ENTROPY BOUNDS, COMPACTNESS AND FINITENESS THEOREMS FOR EMBEDDED SELF-SHRINKERS WITH ROTATIONAL SYMMETRY

JOHN MAN SHUN MA, ALI MUHAMMAD, NIELS MARTIN MØLLER

Abstract. In this work, we study the space of complete embedded rotationally symmetric self-shrinking hypersurfaces in \mathbb{R}^{n+1} . First, using comparison geometry in the context of metric geometry, we derive explicit upper bounds for the entropy of all such self-shrinkers. Second, as an application we prove a smooth compactness theorem on the space of all such shrinkers. We also prove that there are only finitely many such self-shrinkers with an extra reflection symmetry.

1. INTRODUCTION

An *n*-dimensional smooth hypersurface $\Sigma^n \subseteq \mathbb{R}^{n+1}$ is a self-shrinker if

$$
H_{\Sigma}(x) = \frac{1}{2} \langle x, \vec{n} \rangle, \text{ for all } x \in \Sigma.
$$

Here H_{Σ} is the mean curvature of Σ with respect to the outward unit normal \vec{n} . Given a self-shrinker, one obtains by scaling a one parameter family of hypersurfaces $\Sigma_t = \sqrt{-t}\Sigma$, $t \in (-\infty, 0)$ which solves the mean curvature flow (MCF) equation,

(1.1)
$$
\left(\frac{\partial \Sigma_t}{\partial t}\right)^{\perp} = \vec{H}_{\Sigma_t},
$$

where $\vec{H}_{\Sigma} = -H_{\Sigma}\vec{n}$ denotes the mean curvature vector of Σ .

Most importantly, self-shrinkers serve as singularity models for the MCF: under the Type I condition on the singularity, Huisken [\[24\]](#page-19-0) showed that a rescaling of a MCF around a singularity converges locally smoothly subsequentially to a self-shrinker, and proved that closed shrinkers with positive mean curvature are round spheres. Later Ilmanen [\[25\]](#page-19-1) proved the subsequential weak convergence of the tangent flow of any MCF to a self-shrinking solution.

For $n = 1$, all compact immersed self-shrinkers in \mathbb{R}^2 were found in [\[1\]](#page-18-0), the circle being the only embedded example. For $n = 2$, Brendle [\[6\]](#page-18-1) proved the long-standing conjecture that the round sphere of radius 2 is the only closed embedded genus zero self-shrinker in \mathbb{R}^3 . For higher genus, embedded examples are constructed in [\[2\]](#page-18-2), [\[20\]](#page-19-2), [\[27\]](#page-19-3), [\[33\]](#page-19-4), [\[28\]](#page-19-5), [\[31\]](#page-19-6), [\[9\]](#page-18-3). In general, the space of embedded self-shrinkers is not well-understood, even in the case of e.g. topological 2-tori in \mathbb{R}^3 .

The authors were partially supported by DFF Sapere Aude 7027-00110B, by CPH-GEOTOP-DNRF151 and by CF21-0680 from respectively the Independent Research Fund Denmark, the Danish National Research Foundation and the Carlsberg Foundation.

In this paper we direct our attention to the class of complete embedded self-shrinking hypersurfaces in \mathbb{R}^{n+1} with a rotational symmetry, for $n \geq 2$. Using a shooting method in ODEs, Angenent constructed in [\[2\]](#page-18-2) the first nontrivial self-shrinkers in \mathbb{R}^{n+1} besides the round sphere, the generalized cylinders and the plane. These self-shrinkers constructed in [\[2\]](#page-18-2) are rotationally symmetric, embedded, diffeomorphic to $\mathbb{S}^{n-1}\times \mathbb{S}^1$ and are commonly called Angenent doughnuts. They were used in a parabolic maximum principle argument in [\[2\]](#page-18-2) to prove that, when $n \geq 2$, mean curvature flows may develop thin neck-pinch singularities.

In [\[29\]](#page-19-7), Kleene and the third named author proved a partial classification of all complete embedded rotationally symmetric self-shrinkers in any dimension (see also [\[36\]](#page-19-8)). Mramor proved in [\[32\]](#page-19-9) several compactness and finiteness results on the space of all such shrinkers. It is conjectured that, at least in dimension 2, the Angenent doughnut (which when $n = 2$) is topologically a torus) is unique and gives the only embedded self-shrinking torus in \mathbb{R}^3 . This conjecture is still open, even in the rotationally symmetric case [\[18\]](#page-19-10).

The goal of this paper is to study various properties of the space of all complete embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} .

For any hypersurface Σ in \mathbb{R}^{n+1} , let $\lambda(\Sigma)$ be the entropy of Σ defined in [\[14\]](#page-18-4) (see also Section [2](#page-2-0) for the definition).

Theorem 1.1. For each $n \geq 2$, there is a positive number E_n such that

 $1 \leq \lambda(\Sigma) \leq E_n$.

for any complete embedded rotationally symmetric self-shrinker $\Sigma^n \subseteq \mathbb{R}^{n+1}$.

The constants E_n we obtain in Theorem [1.1](#page-1-0) are explicit (see [\(3.1\)](#page-4-0)). For example, when $n = 2$ we have $E_2 \sim 2.24759$, while the 2-dimensional Angenent torus constructed in [\[2\]](#page-18-2) has entropy around 1.85122, as computed numerically in [\[4\]](#page-18-5),[\[22\]](#page-19-11). We remark that for $n = 1$, there is no upper entropy bound for the family of Abresch-Langer immersed selfshrinking curves [\[1\]](#page-18-0). We also remark that if we exclude the stationary plane, the lower bounds can be improved to $\lambda(\mathbb{S}^n)$ ([\[13\]](#page-18-6), [\[3\]](#page-18-7)); if we consider only self shrinking doughnuts, the lower bound can be improved to $\lambda(\mathbb{S}^1) = \sqrt{2\pi/e} \sim 1.52035$, since they all have non-trivial fundamental groups [\[23\]](#page-19-12).

The proof of Theorem [1.1](#page-1-0) will make essential use of the fact that Angenent's Riemannian metrics have Gaussian curvatures bounded below by strictly positive constants (see Section [2](#page-2-0) for the definition of the Angenent metrics).

As an application, our next theorem gives a smooth compactness result for the space of embedded rotationally symmetric self-shrinkers.

Theorem 1.2. For each $n \geq 2$, the space of complete embedded rotationally symmetric self-shrinkers $\Sigma^n \subseteq \mathbb{R}^{n+1}$ is compact in the C_{loc}^{∞} -topology.

Under some extra assumptions on the bounds on entropy and genus, there are already several smooth compactness results for self-shrinkers. Colding and Minicozzi proved in [\[15\]](#page-19-13) the smooth compactness of the set of all complete embedded self-shrinkers $\Sigma^2 \subseteq \mathbb{R}^3$ with bounded genus and Euclidean volume growth (see also [\[16,](#page-19-14) Theorem 1.4]). Later Sun and Wang proved in [\[38\]](#page-19-15) a similar compactness theorem for embedded self-shrinkers

in \mathbb{R}^3 with fixed genus and uniformly bounded entropy. In particular, the $n = 2$ case of Theorem [1.2](#page-1-1) follows from the main theorem in [\[38\]](#page-19-15) and Theorem [1.1.](#page-1-0) We remark that there are more compactness results for two dimensional self-shrinkers in general, even in higher codimension [\[10\]](#page-18-8). In the rotationally symmetric situation, Mramor [\[32\]](#page-19-9) proved several compactness results on the space of compact embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} with various assumptions on n, bounds on entropy and convexity of the profile curves. Theorem [1.2](#page-1-1) is a natural generalization of the results therein.

Theorem [1.2](#page-1-1) has several consequences, which include an index upper bound (Corollary [4.7\)](#page-16-0), finiteness of the set of possible entropy values (Corollary [4.8\)](#page-16-1) and the following finiteness theorem.

Theorem 1.3. For each $n \geq 2$, up to rigid motions there are only finitely many complete embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} which are symmetric with respect to the hyperplane perpendicular to the axis of rotation.

We remark that embeddedness is necessary: there are infinitely many immersed rotationally symmetric self-shrinkers constructed in [\[19\]](#page-19-16) with this extra reflection symmetry.

In Section [2,](#page-2-0) we recall the basic definitions and results needed. In Section [3](#page-4-1) we prove Theorem [1.1.](#page-1-0) In Section [4](#page-8-0) we prove Theorem [1.2](#page-1-1) and [1.3.](#page-2-1)

2. Background

2.1. **Entropy and Self-shrinkers.** We follow the notations in [\[14\]](#page-18-4). Let $\Sigma \subset \mathbb{R}^{n+1}$ be an *n*-dimensional properly embedded hypersurface. For each $x_0 \in \mathbb{R}^{n+1}$, $t_0 > 0$, define the F-functional

$$
F_{x_0,t_0}(\Sigma) := \frac{1}{(4\pi t_0)^{n/2}} \int_{\Sigma} e^{\frac{-|x-x_0|^2}{4t_0}} d\mu,
$$

where $d\mu$ is the volume form of Σ . The entropy of Σ is defined as

$$
\lambda(\Sigma) = \sup_{x_0,t_0} F_{x_0,t_0}(\Sigma).
$$

Using Huisken's monotonicity formula [\[24\]](#page-19-0), it was proved in [\[14\]](#page-18-4) that if $\{\Sigma_t\}_{t\in I}$ satisfies the MCF equation [\(1.1\)](#page-0-0), then the entropy $t \mapsto \lambda(\Sigma_t)$ is non-increasing, and is constant if and only if Σ_t is self-shrinking. We recall the following lemma proved in [\[14,](#page-18-4) Section 7.2].

Proposition 2.1. Let Σ be a properly embedded self-shrinker. Then $\lambda(\Sigma) = F_{0,1}(\Sigma)$.

A hypersurface Σ is a self-shrinker if and only if Σ is critical with respect to the functional $\Sigma \mapsto F_{0,1}(\Sigma)$ [\[14,](#page-18-4) Proposition 3.6]. The second variation of $F_{0,1}$ at a self-shrinker is calculated in [\[14,](#page-18-4) Section 4]: for any normal variation Σ_s of Σ given by $f\vec{n}$, we have

$$
\left. \frac{\partial^2}{\partial s^2} F_{0,1}(\Sigma_s) \right|_{s=0} = -\int_{\Sigma} f L f e^{-\frac{|x|^2}{4}} d\mu_{\Sigma},
$$

where

(2.1)
$$
L = \Delta + |A|^2 - \frac{1}{2} \langle x, \nabla(\cdot) \rangle + \frac{1}{2}
$$

is the stability operator on Σ . It is also shown that all self-shrinkers in \mathbb{R}^{n+1} are Lunstable $([14]$ $([14]$, see also [\[15,](#page-19-13) Theorem 0.5]).

2.2. Rotationally symmetric self-shrinkers; Angenent doughnuts. Let *l* be any line in \mathbb{R}^{n+1} passing through the origin. A hypersurface Σ of \mathbb{R}^{n+1} is rotationally symmetric with respect to l if $R\Sigma = \Sigma$ for all rotations $R \in SO(n+1)$ fixing l. Assume $n \geq 2$ and let $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ be the $(n-1)$ -dimensional unit sphere. We denote the upper half-plane by

$$
\mathbb{H} = \{ (x, r) \in \mathbb{R}^2 : r > 0 \}.
$$

Definition 2.2. Let I be any interval in \mathbb{R} , or $I = \mathbb{S}^1$. Let $\sigma : I \to \mathbb{H}$ be a smooth embedding. Then the embedded hypersurface Σ_{σ} in \mathbb{R}^{n+1} with profile curve $\sigma(s) = (x(s), r(s))$ is given by

$$
\Sigma_{\sigma} = \{ (x(s), \omega r(s)) : \omega \in \mathbb{S}^{n-1}, s \in I \}.
$$

The hypersurface Σ_{σ} is rotationally symmetric with respect to $\ell = \mathbb{R}e_1$.

Proposition 2.3. Given a profile curve σ . Then Σ_{σ} is a self-shrinker if and only if σ is a geodesic in (\mathbb{H}, g_A) , where g_A is the incomplete **Angenent metric** given by

(2.2)
$$
g_A = r^{2(n-1)}e^{-\frac{x^2+r^2}{2}}(dx^2+dr^2).
$$

Direct calculations give

$$
F_{0,1}(\Sigma_{\sigma}) = (4\pi)^{-n/2} \omega_{n-1} L_A(\sigma),
$$

where ω_{n-1} is the surface area of \mathbb{S}^{n-1} and $L_A(\sigma)$ is the length of σ in (\mathbb{H}, g_A) . Hence if Σ_{σ} is a self-shrinker,

(2.3)
$$
\lambda(\Sigma_{\sigma}) = (4\pi)^{-n/2}\omega_{n-1}L_A(\sigma)
$$

by Proposition [2.1.](#page-2-2)

Using a shooting method in ODEs, Angenent constructed in [\[2\]](#page-18-2) a compact embedded rotationally symmetric self-shrinker in \mathbb{R}^{n+1} for each $n \geq 2$. The profile curve of the examples in [\[2\]](#page-18-2) are convex and symmetric with respect to the r-axis. The self-shrinkers so constructed are called Angenent doughnuts. More recently, Drugan and Nguyen in [\[20\]](#page-19-2) used a geometric flow to construct compact embedded rotationally symmetric selfshrinkers in \mathbb{R}^{n+1} with the same property of the profile curve for all $n \geq 2$. It is not known if both constructions in [\[2\]](#page-18-2) and [\[20\]](#page-19-2) resulted in the same self-shrinkers. On the other hand, Kleene and the third named author proved the following partial classification, which we will be making use of in the present paper:

Theorem 2.4. [\[29,](#page-19-7) Theorem 2] Let Σ be a complete embedded rotationally symmetric self-shrinker in \mathbb{R}^{n+1} . Then up to rigid motion, Σ is either

- (i) the hyperplane $\mathcal{P} = \{0\} \times \mathbb{R}^n$,
- (ii) the round sphere S of radius $\sqrt{2n}$,
- (iii) the round cylinder $C = \mathbb{R} \times \mathbb{S}^{n-1}$ with radius $\sqrt{2(n-1)}$, or
- (iv) diffeomorphic to $\mathbb{S}^1 \times \mathbb{S}^{n-1}$.

Note that in the last case of Theorem [2.4,](#page-3-0) $\Sigma = \Sigma_{\sigma}$ for some simple closed geodesic σ in (H, g_A). This theorem is essential to the proof of Theorem [1.2.](#page-1-1) As a first simple but useful consequence, note that if a complete rotationally symmetric self-shrinker is embedded, it is automatically properly embedded.

3. Entropy bound for embedded rotationally symmetric self-shrinkers

In this section we prove Theorem [1.1.](#page-1-0) By a direct calculation using [\(2.2\)](#page-3-1), the Gauss curvature of the Angenent metric g_A is

$$
K = \frac{r^2 + (n-1)}{r^{2n}} e^{\frac{x^2 + r^2}{2}}.
$$

Note that K is strictly positive. For each n, let κ_n be the (positive) minimum of K. By simple calculus, one can find

$$
\kappa_n = \frac{y_n + (n-1)}{y_n^n} e^{\frac{y_n}{2}}, \quad \text{where } y_n = \frac{n-1 + \sqrt{9(n-1)^2 + 8(n-1)}}{2}.
$$

We will prove the following more general result.

Theorem 3.1. Let (M, g) be a 2-dimensional Riemannian manifold so that M is homeomorphic to \mathbb{R}^2 and $K_g \geq \kappa_g > 0$, where K_g is the Gauss curvature of (M, g) . Then every simple closed geodesic in (M, g) has length $\leq 2\pi/\sqrt{\kappa_g}$.

Theorem [1.1](#page-1-0) follows directly from Theorem [3.1](#page-4-2) and (2.3) . Indeed, the constants E_n in Theorem [1.1](#page-1-0) are given by

(3.1)
$$
E_n = \frac{2\pi\omega_{n-1}}{(4\pi)^{n/2}\sqrt{\kappa_n}}.
$$

In the Appendix we show that $2 < E_n \le E_2$ for all $n \ge 2$ and

$$
\lim_{n \to \infty} E_n = \sqrt{\frac{4\pi}{3}} \sim 2.04665.
$$

When M is homeomorphic to \mathbb{S}^2 , Theorem [3.1](#page-4-2) is a classical theorem in comparison geometry, where a proof can be found in [\[30,](#page-19-17) Theorem 3.4.10]; in our situation, (M, g) is non-compact and incomplete, and we are not able to find an exact reference in this generality. As a result, we provide a proof of Theorem [3.1](#page-4-2) using the globalization theorem in metric geometry [\[8\]](#page-18-9).

Let σ be a simple closed geodesic in (M, g) . Since M is homeomorphic to \mathbb{R}^2 , by the Jordan curve theorem σ divides M into two connected components, where exactly one of them has compact closure.

Definition 3.2. Given a simple closed geodesic σ in (M, g) , let Ω be the compact domain in (M, g) with $\partial \Omega = \text{Im}(\sigma)$.

Lemma 3.3. Let $p, q \in \Omega$. Then there is a simple geodesic γ in Ω joining p, q, which is shortest among all piecewise C^1 curves in Ω joining p and q. Moreover,

(i) if one of p, q is in the interior of Ω , γ also lies in the interior of Ω (except possibly at the other end point),

(ii) if both p, q are in $\text{Im}(\sigma) = \partial\Omega$, then either γ lies completely in $\partial\Omega$, or the interior of γ lies inside the interior of Ω .

Proof. Let $p, q \in \Omega$. Since the case $p = q$ is trivial, we assume $p \neq q$. Let

$$
d^{\Omega}(p,q) = \inf_{\gamma} L(\gamma),
$$

where $L(\gamma) = \int \sqrt{g(\dot{\gamma}, \dot{\gamma})}$ is the length of γ and the infimum is taken among all piecewise C¹ curves $\gamma : [0,1] \to \Omega$ so that $\gamma(0) = p$ and $\gamma(1) = q$. It is easy to check that d^{Ω} is a metric on Ω . In particular, $d^{\Omega}(p,q) > 0$.

Let $\gamma_j : [0,1] \to \Omega$ be a sequence of piecewise C^1 curves, parametrized proportional to arc length, joining p, q so that $L(\gamma_j) \to d^{\Omega}(p,q)$ as $j \to \infty$. Since Ω is compact, by passing to a subsequence if necessary, we may assume that (γ_i) converges uniformly in d^{Ω} to a Lipschitz continuous curve $\gamma : [0, 1] \to \Omega$ as $j \to \infty$. If $\gamma(t_0) = q$ for some $t_0 < 1$, we replace γ by $\gamma|_{[0,t_0]}$. Hence we can assume $\gamma^{-1}(\{q\}) = \{1\}.$

First we prove (i). Assume that $p \in \Omega \setminus \partial \Omega$. We claim that

(3.2)
$$
\gamma(t) \notin \partial \Omega, \text{ for all } t < 1.
$$

We argue by contradiction: if not, let t_0 be the infimum of the set $\gamma^{-1}(\partial\Omega \setminus \{q\})$. Since $p \notin \partial\Omega$, we have $t_0 > 0$ and $\gamma([0, t_0))$ lies in the interior of Ω . This together with the definition of γ implies that $\gamma|_{[0,t_0]}$ is a geodesic. Let U be a small geodesically convex neighborhood in (M, g) centered at $\gamma(t_0)$ not containing q. Then there is $\epsilon > 0$ such that $\gamma(t)$ lies in U for all $t \in [t_0 - \epsilon, t_0 + \epsilon]$ and $\gamma(t_0 \pm \epsilon) \neq \gamma(t_0)$. Let σ_0 be the shortest geodesic in U connecting $\gamma(t_0 \pm \epsilon)$. Since $\partial \Omega \cap U$ is a portion of the geodesic σ and U is geodesically convex, σ_0 does not intersect with $\partial\Omega \cap U$ at more than one points. Hence the image of σ_0 except possibly at $\gamma(t_0+\epsilon)$ must lie in the connected component of $U \setminus \partial \Omega$ containing $\gamma(t_0 - \epsilon)$. As a result, the image of σ_0 also lies in $\Omega \cap U$. Since γ is length minimizing, up to reparametrization we have $\gamma|_{[t_0-\epsilon,t_0+\epsilon]} = \sigma_0$ and thus $\gamma|_{[t_0-\epsilon,t_0+\epsilon]}$ is a smooth geodesic. Since γ , σ are tangential at $\gamma(t_0)$ (recall that $\partial\Omega = \text{Im}\sigma$) and both are geodesics, we have $\gamma = \sigma$ locally around $\gamma(t_0)$. This contradicts the choice of t_0 and thus [\(3.2\)](#page-5-0) is shown. This immediately implies (i).

Next we prove (ii). Assume that $p, q \in \partial\Omega$. Let (p_k) be a sequence of points in the interior of Ω converging to p. For each $k \in \mathbb{N}$, let $\gamma_k : [0,1] \to \Omega$ be a shortest geodesic in Ω joining p_k to q constructed in (i). By the smooth dependence of solutions to the geodesic equation and picking a subsequence if necessary, (γ_k) converges smoothly to a geodesic $\gamma : [0, 1] \to \Omega$ joining p, q. Using the triangle inequality

$$
L(\gamma_k) = d^{\Omega}(p_k, q) \le d^{\Omega}(p_k, p) + d^{\Omega}(p, q)
$$

and taking $k \to \infty$, we have $L(\gamma) \leq d^{\Omega}(p, q)$. Thus γ is a length minimizing geodesic in $Ω$. Then either $γ$ lies completely inside the interior of $Ω$ away from the endpoints, or $γ$ touches $\partial\Omega$ at some point in $\partial\Omega \setminus \{p,q\}$, which implies that γ lies completely in $\partial\Omega$ since both of them are geodesics. This finishes the proof of (ii). both of them are geodesics. This finishes the proof of (ii).

We will need some definitions and notations from metric geometry. We use the reference [\[8\]](#page-18-9). For the convenience of the reader, we summarize the basic facts that we shall need to prove Theorem [3.1.](#page-4-2)

A length space (X, d) is a metric space such that the metric d can be obtained as a distance function associated to a length structure (see [\[8\]](#page-18-9) for a definition). The metric d is called an *intrinsic metric* in this case. If every pair of points p, q in X can be joined by a (possibly non-unique) shortest path, then the metric d is called strictly intrinsic. We recall that a shortest path is a curve γ where the length $L(\gamma)$ is given by the distance between the endpoints of γ . A length space whose metric is strictly intrinsic is called a complete length space.

From Lemma [3.3](#page-4-3) we conclude

Proposition 3.4. (Ω, d^{Ω}) is a complete length space.

A triangle Δpqr in (X, d) is a set of points $\{p, q, r\}$ together with three shortest paths $[pq], [qr], [rp]$. The length of a triangle is the sum of the lengths of its sides. For each $\kappa > 0$ and for each triangle Δpqr in X with length $\langle 2\pi/\sqrt{\kappa}$, we can associate a unique (up to an isometry) comparison geodesic triangle $\Delta \bar{p} \bar{q} \bar{r}$ in \mathbb{S}^2 $\frac{2}{1/\sqrt{\kappa}}$ with vertices $\{\bar{p}, \bar{q}, \bar{r}\}\$ such that the corresponding sides of the geodesic triangle $\Delta\bar{p}\bar{q}\bar{r}$ have the same lengths as the sides of the triangle Δpqr . Here \mathbb{S}^2 ²/_{1/ $\sqrt{\kappa}$} is the 2-sphere with radius $1/\sqrt{\kappa}$. For a triangle Δpqr in X, we denote the angles by $\angle p, \angle q, \angle r$, and if confusion arises we will e.g write $\angle pqr$ for the angle $\angle q$. We shall not need the definition of an angle between two shortest paths in a length space, but one can show that on a Riemannian manifold M, if c_1 and c_2 are two geodesics starting at $p = c_1(0) = c_2(0)$, then the angle $\angle p \in [0, \pi]$ between the shortest paths c_1 and c_2 is equal to the usual Riemannian angle between c_1 and c₂. See Corollary 1A.7 in [\[7\]](#page-18-10) for a proof. For the comparison triangle $\Delta \bar{p} \bar{q} \bar{r}$ in \mathbb{S}^2 $\frac{1}{\sqrt{\kappa}}$ we denote the angles by $\angle \bar{p}, \angle \bar{q}, \angle \bar{r}$.

There are several definitions of a *space of curvature* $\geq \kappa$. We shall use the following angle comparison definition.

Definition 3.5. Let X be a complete length space, and let $\kappa > 0$. We say that X is a space of curvature $\geq \kappa$ if for any point $x \in X$ there is a neighborhood U_x of x such that for all triangles $\Delta pqr \subset U_x$ the corresponding angles satisfy the inequalities

(3.3)
$$
\angle p \geq \angle \bar{p}, \quad \angle q \geq \angle \bar{q}, \quad \angle r \geq \angle \bar{r},
$$

for a comparison triangle $\Delta \bar{p}\bar{q}\bar{r}$ in \mathbb{S}^2_1 $\frac{2}{1/\sqrt{\kappa}}$. Furthermore, for any two shortest paths [ab] and [cd] in X where c is an interior point on [ab], the following holds \angle acd + \angle bcd = π .

Remark 3.1. The last statement in Definition [3.5](#page-6-0) above can be summarized as "the sum of adjacent angles equals π ". It is needed to prove the equivalence to other definitions of spaces of curvature $\geq \kappa$. See [\[8,](#page-18-9) Section 4.3].

Proposition 3.6. The complete length space (Ω, d^{Ω}) has curvature $\geq \kappa_q$, where κ_q is the lower bound of the Gaussian curvature of (M, g) in Theorem [3.1.](#page-4-2)

Proof. Let $x \in \Omega$ and let V_x be a geodesically convex neighborhood in (M, g) centered at x. When x is in the interior of Ω , we choose $V_x \subset \Omega$. Let $U_x = \Omega \cap V_x$. For any $p, q \in U_x$, let γ be the unique shortest geodesic in V_x joining p and q. Note that γ must lie inside U_x : this holds when x is in the interior of Ω , since $U_x = V_x$. When $x \in \partial \Omega$, if γ is not in U_x, there are $t_1 < t_2$ such that $\gamma|_{(t_1,t_2)}$ lies in $V_x \setminus U_x$ and $\gamma(t_1), \gamma(t_2)$ are in $\partial\Omega \cap V_x$. Since

 $\partial\Omega \cap V_x$ is also a geodesic passing through $\gamma(t_1), \gamma(t_2)$, this contradicts the fact that V_x is a geodesically convex neighborhood.

In particular, triangles of (Ω, d^{Ω}) in U_x are also triangles of (M, g) in V_x . Together with the assumption that (M, g) has Gaussian curvature $K \ge \kappa_g > 0$, we can argue as in the case for Riemannian manifolds without boundary to show that (Ω, d^{Ω}) has curvature $\geq \kappa_q$; see Theorem 6.5.6. in [\[8\]](#page-18-9).

We will denote the inequalities in (3.3) by the angle comparison condition. If X is a complete length space, then Toponogov's globalization theorem [\[8,](#page-18-9) Theorem 10.3.1] globalizes this local curvature condition to the entirety of X , not only in a neighborhood U_x of each point x:

Theorem 3.7 (Globalization Theorem). Let X be a complete length space of curvature $\geq \kappa$ for some $\kappa > 0$. Then the angle comparison condition [\(3.3\)](#page-6-1) is satisfied for any triangle Δ pqr in X for which there is a unique (up to isometry) comparison triangle $\Delta \bar{p}\bar{q}\bar{r}$ in \mathbb{S}^2_1 $_{1/\sqrt{\kappa}}^{2}.$

Let us remind that when $\kappa > 0$, a comparison triangle only exists if the length of Δpqr does not exceed $2\pi/\sqrt{\kappa}$. As mentioned earlier, we can associate a unique comparison triangle when the length of Δpqr is strictly less than $2\pi/\sqrt{\kappa}$. If the length is equal to $2\pi/\sqrt{\kappa}$, then we have two situations:

- All the sides have length strictly less than $\pi/\sqrt{\kappa}$. Then a comparison triangle is a unique great circle, i.e. all of its angles are equal to π .
- One of the sides has length equal to $\pi/\sqrt{\kappa}$, say [pq]. The sum of lengths of the two other sides $[qr]$ and $[rp]$ is then equal to $\pi/\sqrt{\kappa}$. In this case there does not exist a unique comparison triangle, but we can fix the comparison triangle $\Delta \bar{p} \bar{q} \bar{r}$ where the side $|\bar{p}\bar{q}|$ passes through the point \bar{r} .

Thus, with this convention, the conclusion of Theorem [3.7](#page-7-0) holds for every triangle Δpqr in X with length $\leq 2\pi/\sqrt{\kappa}$. In fact, as a corollary of theorem [3.7](#page-7-0) one can show that any triangle in X has length no greater than $2\pi/\sqrt{\kappa}$, where X is a space of curvature $\geq \kappa$ for some $\kappa > 0$. See [\[8,](#page-18-9) Corollary 10.4.2.]. We shall use this fact in the proof of Theorem [3.1](#page-4-2) below.

Before proving Theorem [3.1,](#page-4-2) we recall the following elementary lemma in spherical geometry. We recall that a convex polygon in the sphere is a polygon so that the interior angle at each vertex is less than or equal to π .

Lemma 3.8. Let P be a convex n-gon in \mathbb{S}^2 $_{1/\sqrt{\kappa}}^{2},$ and denote the length of P by $|P|.$ Then $|P| \leq 2\pi/\sqrt{\kappa}$.

Proof of Theorem [3.1.](#page-4-2) The idea of this proof is similar to the proof presented in [\[30,](#page-19-17) Theorem 3.4.10]. Let Ω be the compact domain bounded by the geodesic $\sigma : [0,1] \to M$ as defined in Definition [3.2,](#page-4-4) so that $\sigma(0) = \sigma(1)$. By Proposition [3.4](#page-6-2) and Proposition [3.6,](#page-6-3) (Ω, d^{Ω}) is a complete length space with curvature $\geq \kappa_g$.

Let $L_0 := L(\sigma)$ be the length of σ . Take numbers $t_1, t_2, t_3, t_4 \in I$, where $t_1 =$ $0, t_4 = 1$, such that each of the subarcs $\sigma|_{[t_1,t_2]}, \sigma|_{[t_2,t_3]}$ and $\sigma|_{[t_3,t_4]}$ has length $L_0/3$. Let $\Delta\sigma(t_1)\sigma(t_2)\sigma(t_3)$ be a triangle in (Ω, d^{Ω}) . By Proposition [3.4](#page-6-2) each of the sides of this triangle either lies completely in $\partial\Omega$ or has interior contained in the interior of Ω . Using [\[8,](#page-18-9) Corollary 10.4.2.] we deduce that the length of this triangle is not greater than $2\pi/\sqrt{\kappa_g}$. If the sides of the triangle $\Delta\sigma(t_1)\sigma(t_2)\sigma(t_3)$ coincide with the images of the subarcs $\sigma|_{[t_1,t_2]}, \sigma|_{[t_2,t_3]}$ and $\sigma|_{[t_3,t_4]}$, then we are done.

If not, then build new triangles

- $\Delta \sigma(t_1) \sigma(t_{1,2}) \sigma(t_2)$,
- $\Delta \sigma(t_2) \sigma(t_2) \dot{\sigma(t_3)}$ and
- $\Delta \sigma(t_3) \sigma(t_{3,1}) \sigma(t_1),$

where $t_{i,k} \in (t_i, t_k)$ are chosen such that $\sigma(t_{i,k})$ define midpoints of the corresponding subarcs. To each one of these triangles, we associate a unique comparison triangle in \mathbb{S}^2 $\frac{2}{1/\sqrt{\kappa_g}}$. We put the triangles together along their common sides and obtain a comparison 6-gon \mathcal{O}_6 in \mathbb{S}_1^2 $\frac{1}{1/\sqrt{\kappa_g}}$. From Theorem [3.7](#page-7-0) we know that the angles in the comparison triangles are not greater than the corresponding angles in Ω . This implies that the angles of the vertices in \mathcal{O}_6 are not bigger than π . Hence \mathcal{O}_6 is convex. By Lemma [3.8,](#page-7-1) the length of \mathcal{O}_6 satisfies $|\mathcal{O}_6| \leq 2\pi/\sqrt{\kappa_g}$. If the sides of the constructed 6-gon in Ω coincide with the arcs of σ , then we are done.

If not, then we continue to construct more triangles as above, and build the corresponding comparison *n*-gons \mathcal{O}_n for increasingly large $n \in \mathbb{N}$. By Lemma [3.8](#page-7-1) again, $|\mathcal{O}_n| \leq 2\pi/\sqrt{\kappa_q}$. For n large enough, the arcs of σ will be the unique shortest paths between the vertices of the constructed n-gon on Ω . This follows from the fact that shortest paths in (Ω, d^{Ω}) are geodesics in (M, g) , and that each side of the n-gon will be contained in a geodesically convex neighborhood in M . In such a neighborhood, every two points are connected by a unique shortest path. This implies the desired bound since for *n* large enough we have $L(\sigma) = |\mathcal{O}_n| \leq 2\pi/\sqrt{\kappa_q}$.

 \Box

4. Compactness and Finiteness of Embedded Self-Shrinkers With Rotational Symmetry

In this section we prove Theorem [1.2,](#page-1-1) [1.3](#page-2-1) and some other related results. We start with the following well known lemma.

Lemma 4.1. Let Σ_1 , Σ_2 be two properly embedded self-shrinkers in \mathbb{R}^{n+1} such that one of them is compact. Then Σ_1 and Σ_2 must intersect.

When both self-shrinkers are compact, this is proved in [\[39,](#page-19-18) Theorem 7.4]. Recently, it was proved in [\[26\]](#page-19-19) that any two properly embedded self-shrinkers that are sufficiently separated at infinity must intersect. See also [\[12,](#page-18-11) Corollary C.4] for the statement for F-stationary varifolds.

Next we restrict attention to complete embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} . By Theorem [2.4,](#page-3-0) there are only two types of non-compact examples - cylinders and hyperplanes through the origin, and all such pairs also intersect. Thus we have

Lemma 4.2. Any two complete embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} intersect.

Now we prove Theorem [1.2.](#page-1-1)

Proof of Theorem [1.2.](#page-1-1) Let (Σ_k) be a sequence of complete embedded rotationally symmetric self-shrinkers in \mathbb{R}^{n+1} each with axis of rotation ℓ_k . After taking a subsequence, a limit axis exists, to which each Σ_k can be rotated, thus it is enough to consider the case where all axes of rotation are identical. By a further rotation, we assume the limit axis is $\ell = \mathbb{R}e_1 \subseteq \mathbb{R}^{n+1}$.

By Theorem [2.4,](#page-3-0) it suffices to assume that Σ_k is a self-shrinking doughnut for each k. Thus $\Sigma_k = \Sigma_{\sigma_k}$, where

$$
\sigma_k : [-d_k, d_k] \to \mathbb{H}
$$

is a unit speed geodesic in (\mathbb{H}, g_A) so that $L(\sigma_k) = 2d_k$ and $\sigma_k(-d_k) = \sigma_k(d_k)$. Since all self-shrinkers have entropy larger than or equal to $\lambda(\mathbb{R}^n) = 1$, we have $d_k \geq d > 0$ for all k by (2.3) for some dimensional constant d.

Let σ_a be the profile curve of the Angenent doughnut constructed in [\[2\]](#page-18-2). By Lemma [4.2,](#page-8-1) each σ_k intersects σ_a . Reparametrizing each σ_k if necessary, we assume that $\sigma_k(0) \in \text{Im } \sigma_a$ for each k. Taking a subsequence if necessary, since $\text{Im}\,\sigma_a$ is compact, we have

$$
\sigma_k(0) \to p, \quad \sigma'_k(0) \to v
$$

as $k \to \infty$. Note that $||v|| = 1$ since each σ_k is of unit speed. Let $\sigma_{\infty} : I \to \mathbb{H}$ be the maximally defined geodesic in (\mathbb{H}, g_A) with $\sigma_\infty(0) = p$, $\sigma'_\infty(0) = v$, where I is an open interval.

For any $R > 1$, let

$$
K_R = [-R, R] \times [R^{-1}, R] \subset \mathbb{H}.
$$

Then there is $R_0 > 1$ so that $\text{Im}(\sigma_a) \subset K_R$ for all $R \geq R_0$. For each $k \in \mathbb{N}$ and $R \geq R_0$, let $I_{k,R}$ be the connected component of $\sigma_k^{-1}(K_R)$ in $[-d_k, d_k]$ containing 0. Since each σ_k is a geodesic in (\mathbb{H}, g_A) , by (4.1) and the smooth dependence on initial data of the ODE within each K_R , $(\sigma_k|_{I_{k,R}})$ converges smoothly to $\sigma_{\infty}|_{I_R}$ in K_R , where I_R is the connected component of $\sigma_{\infty}^{-1}(K_R)$ containing 0.

Since each $\sigma_k|_{I_{k,R}}$ is embedded and $(\sigma_k|_{I_{k,R}})$ converges smoothly to $\sigma_{\infty}|_{I_R}$, it is clear that σ_{∞} does not admit transverse self-intersection. Thus if $\sigma_{\infty}(s) = \sigma_{\infty}(t)$ for some $s \neq t$, then $\sigma'_{\infty}(s) = \pm \sigma'_{\infty}(t)$ and this implies that σ_{∞} is periodic. In this case, σ_{∞} is a simple closed geodesic in (\mathbb{H}, g_A) and (σ_k) converges smoothly to σ_{∞} . Thus we are done.

From now on we may therefore assume that σ_{∞} is not a closed geodesic. Since $(\sigma_k|_{I_{k,R}})$ converges smoothly to σ_{∞} in each K_R and $\cup_{R\geq R_0} K_R = \mathbb{H}$, σ_{∞} is properly immersed. Since σ_{∞} is injective, $\Sigma_{\sigma_{\infty}}$ is a complete embedded rotationally symmetric self-shrinker in \mathbb{R}^{n+1} . By Theorem [2.4,](#page-3-0) $\Sigma_{\sigma_{\infty}}$ is either the plane \mathcal{P} , the sphere $\mathcal S$ or the cylinder \mathcal{C} .

First we assume that $\Sigma_{\sigma_{\infty}}$ is the sphere S and we will derive a contradiction. After that, we consider the hyperplane P and the cylinder C and point out the necessary changes for the contradiction argument.

We split the argument into several lemmas.

Lemma 4.3. There is $R_1 \ge R_0$ so that the following holds: for all $R \ge R_1$, there is $k_1 = k_1(R)$ so that for all $k \geq k_1$, $\text{Im } \sigma_k \cap K_R$ contains a connected component different from $\sigma_k(I_{k,R})$ which intersects K_{R_1} .

Proof of Lemma [4.3.](#page-9-1) First let $s \in (0,1)$ be small so that the scaling of the Angenent doughnut $s\Sigma_{\sigma_a}$ lies completely inside the sphere $\Sigma_{\sigma_\infty} = \mathcal{S}$: that is, $x^2 + r^2 < 2n$ for all $(x, r) \in s \operatorname{Im} \sigma_a$. Let $\delta = d_0(s \Sigma_{\sigma_a}, \Sigma_{\sigma_\infty} \cup \ell)$, where $\ell \subset \mathbb{R}^{n+1}$ is the axis of rotation and d_0 is the Euclidean distance. Note that $\delta > 0$. Let $R_1 = \max\{R_0, 2\delta^{-1}, 3\sqrt{2n}\}\.$ For any $R > R_1$, since $\sigma_k|_{I_{k,R}}$ converges to $\sigma_{\infty}|_{I_R}$ in K_R uniformly, there is $k_1(R) > 0$ so that

(4.2)
$$
d_0(\sigma_k(I_{k,R}), \sigma_\infty(I_R)) < \frac{\delta}{2}
$$

for all $k \geq k_1$. By the choice of δ , we have

$$
\sigma_k(I_{k,R}) \cap s(\operatorname{Im} \sigma_a) = \emptyset, \quad \text{for all } k \geq k_1.
$$

Now we fix $k \geq k_1$ and show that Im $\sigma_k \cap K_R$ has more than one component which intersects K_{R_1} . Assume the contrary, then $\text{Im}\,\sigma_k \cap K_{R_1} = \sigma_k(I_{k,R}) \cap K_{R_1}$. Let

$$
K_{R_1}^{\delta} = K_{R_1} \setminus \{ (x,r) \in \mathbb{H} \mid \sqrt{x^2 + r^2} < \sqrt{2n} + \delta \}.
$$

Note that $K_{R_1}^{\delta}$ is connected. By [\(4.2\)](#page-10-0), $K_{R_1}^{\delta}$ is disjoint from Im σ_k . Since σ_k is a simple closed curve, by the Jordan curve theorem, its image divides H into two connected components, where exactly one of them is compact. There are two cases:

(i) $K_{R_1}^{\delta}$ lies in the compact component (see Figure 1). Let

$$
\tilde{\sigma}_a := s\sigma_a + (2\sqrt{2n}, 0)
$$

be the horizontal translation of $s\sigma_a$ by $(2\sqrt{2n}, 0)$. Since $R_1 > 3\sqrt{2n}$ and $R_1^{-1} < \delta$, $\tilde{\sigma}_a$ lies completely inside $K_{R_1}^{\delta}$ and thus in the compact component (see Figure 1). The translation is horizontal, hence the MCF $\{\Sigma_{\tilde{\sigma}_a}^t\}$ starting at $\Sigma_{\tilde{\sigma}_a}$ is also self-shrinking, centered at $(2\sqrt{2n},0)$. Since $s < 1$, it becomes extinct before the MCF $\{\Sigma_{\sigma_k}^t\}$ starting at Σ_{σ_k} does. This implies that $\Sigma_{\tilde{\sigma}_a}^t$ intersects $\Sigma_{\sigma_k}^t$ for some $t > 0$, which contradicts the maximum principle since $\Sigma_{\tilde{\sigma}_a}$ and Σ_{σ_k} are disjoint.

(ii) $K_{R_1}^{\delta}$ lies in the non-compact component (see Figure 2): let

$$
H_{R_1}^{\delta} = \{ (x, r) : K_{R_1} \mid \sqrt{x^2 + r^2} \le \sqrt{2n} - \delta \}.
$$

Since $H_{R_1}^{\delta}$ is a subset of K_{R_1} and is disjoint from $\sigma_k(I_{k,R}), H_{R_1}^{\delta}$ also lies in a component of $\mathbb{H} \setminus \text{Im}(\sigma_k)$. Since $K_{R_1}^{\delta}$, $H_{R_1}^{\delta}$ are separated by $\sigma_k(\tilde{I}_{k,R})$, then $H_{R_1}^{\delta}$ and $K_{R_1}^{\delta}$ lie in different components of $\mathbb{H} \setminus \text{Im} \sigma_k$. Thus $H_{R_1}^{\delta}$ is in the compact component bounded by Im σ_k . By the choice of R_1 and since $R > R_1$, s Im σ_a lies in the compact component (see Figure 2). By considering the MCF starting at $s\Sigma_{\sigma_a}$ as in (i), we again arrive at a contradiction.

Hence both cases are impossible, and we conclude that $\text{Im}\,\sigma_k \cap K_R$ must contain more than one component which intersects K_{R_1} for all $k \geq k_1$. This finishes the proof of Lemma [4.3.](#page-9-1) \Box

Let $R_1 < R_2 < \ldots < R_k < \ldots$ be any sequence such that $R_k \nearrow +\infty$ as $k \to \infty$. Using Lemma [4.3](#page-9-1) and picking a subsequence of $(\sigma_k)_{k=1}^{\infty}$ if necessary, we may assume that $\text{Im}\,\sigma_k \cap K_{R_k}$ has more than one component which intersects K_{R_1} . For each k, let $N_k \geq 2$ be the number of such connected components. Since there is a uniform positive lower

FIGURE 1. $\tilde{\sigma}_a$ is enclosed by σ_k .

FIGURE 2. $s\sigma_a$ is enclosed by σ_k .

bound on $d_A(K_{R_1}, \mathbb{H} \setminus K_{R_k})$ for all $k \geq 2$, N_k is uniformly bounded by Theorem [1.1](#page-1-0) (see also [\(2.3\)](#page-3-2)). Taking a further subsequence if necessary, we may assume that $N := N_k$ is constant.

Thus for each k, write $\text{Im}\,\sigma_k \cap K_{R_k}$ as the disjoint union

(4.3)
$$
\operatorname{Im} \sigma_k \cap K_{R_k} = \operatorname{Im} \sigma_k^1 \cup \dots \cup \operatorname{Im} \sigma_k^N \cup \sigma_k^C,
$$

where each σ_k^i , $i = 1, \dots, N$ is a simple geodesic arc parametrizing a connected component of Im $\sigma_k \cap K_{R_k}$ which intersects K_{R_1} and σ_k^C is the union of any connected components of Im $\sigma_k \cap K_{R_k}$ which do not intersect K_{R_1} .

For each $k \in \mathbb{N}$, let $i_k \in \{1, \dots, N\}$ be arbitrary. Up to reparametrization, there are $c_k > 0$ such that

$$
\sigma_k^{i_k} : [-c_k, c_k] \to \mathbb{H}
$$

is a unit speed geodesic. Since the image of $\sigma_k^{i_k}$ $\binom{i_k}{k}$ intersects K_{R_1} , one has

$$
2c_k \ge d_A(K_{R_1}, \mathbb{H} \setminus K_{R_2}) > 0,
$$

and there are $\bar{c}_k \in [-c_k, c_k]$ such that $\sigma_k^{i_k}(\bar{c}_k) \in K_{R_1}$. Taking a further subsequence if necessary, by compactness of K_{R_1} , we may assume that

$$
\sigma_k^{i_k}(\bar{c}_k) \to q, \quad (\sigma_k^{i_k})'(\bar{c}_k) \to w
$$

as $k \to \infty$. Let $\tilde{\sigma}_{\infty}: J \to \mathbb{H}$ be the unique complete maximal geodesic with $\tilde{\sigma}_{\infty}(0) = q$ and $\tilde{\sigma}'_{\infty}(0) = w$.

As in the construction of σ_{∞} , $(\sigma_k^{i_k})$ $\binom{n_k}{k}$ converges smoothly to $\tilde{\sigma}_{\infty}$ in K_R for all $R > R_1$ and $\tilde{\sigma}_{\infty}$ is an embedded geodesic in (\mathbb{H}, g_A).

Lemma 4.4. Up to reparametrization, $\tilde{\sigma}_{\infty} = \sigma_{\infty}$.

Proof of Lemma [4.4.](#page-12-0) We may assume that $\sigma_k^{i_k}$ $k_k^{i_k} \neq \sigma_k|_{I_{k,R_k}}$ for all large k, since otherwise we have $\tilde{\sigma}_{\infty} = \sigma_{\infty}$ up to reparametrization. This implies that $\text{Im}(\sigma_k|_{I_{k,R_k}})$ and $\text{Im}\,\sigma_k^{i_k}$ k have empty intersection since both of them are connected components of $\text{Im}\,\sigma_k \cap K_{R_k}$. Assume the contrary, that $\tilde{\sigma}_{\infty} \neq \sigma_{\infty}$. By Lemma [4.2](#page-8-1) and that $\tilde{\sigma}_{\infty}$, σ_{∞} are both complete geodesics, they must intersect transversally. Assume that the intersection is in K_R for some $R > R_1$. Since $\sigma_k|_{I_{k,R}}$ converges locally smoothly to σ_{∞} in K_R , $\sigma_k(I_{k,R})$ also intersects $\tilde{\sigma}_{\infty}$ for k large enough. On the other hand, $\sigma_k^{i_k}$ converges smoothly to $\tilde{\sigma}_{\infty}$, thus $\sigma_k^{i_k}$ also intersects $\sigma_k|_{I_{k,R}}$ in K_R for large k. This contradicts the assumption on $\sigma_k^{i_k}$ and hence the lemma is proved.

Thus the union of subarcs $\text{Im}\,\sigma_k^1\cup\cdots\cup\text{Im}\,\sigma_k^N$ of $\text{Im}\,\sigma_k$ converge as $k\to\infty$ to σ_∞ locally smoothly with multiplicity N. While σ_k^C defined in [\(4.3\)](#page-11-0) might not be empty, we can show that (by passing to a subsequence if necessary) it also stays close to σ_{∞} .

Lemma 4.5. By passing to a subsequence of (σ_k) if necessary,

(4.4)
$$
|\sqrt{x^2 + r^2} - \sqrt{2n}| < R_k^{-1}
$$

for all $(x, r) \in \text{Im}(\sigma_k) \cap K_{R_k}$ and for all $k \in \mathbb{N}$.

Proof of Lemma [4.5.](#page-12-1) Let $k \in \mathbb{N}$ be fixed. First we show that there is n_k so that [\(4.4\)](#page-12-2) holds for all $(x, r) \in \text{Im}(\sigma_i) \cap K_{R_k}$ and for all $i \geq n_k$. To see this we argue by contradiction: if not, then there is a subsequence of (σ_{k_j}) of (σ_k) and $(x_j, r_j) \in \text{Im}(\sigma_{k_j}) \cap K_{R_k}$ so that | $\sqrt{x_j^2 + r_j^2} \sqrt{2n}$ $\ge R_k^{-1}$ for all j. Since K_{R_k} is compact, we may assume that $(x_j, r_j) \to$ $(x_\infty, r_\infty) \in K_{R_k}$, and thus there is a sequence $(\tilde{\sigma}_j)$ of subarcs of (σ_{k_j}) which converges locally smoothly to an embedded complete geodesic in (\mathbb{H}, g_A) , which is different from σ_{∞} since $(x_{\infty}, r_{\infty}) \notin \text{Im}(\sigma_{\infty})$. Arguing similarly as in the proof of Lemma [4.4,](#page-12-0) this is impossible. Lastly, the lemma is proved by passing to a subsequence of (σ_{ι}) . impossible. Lastly, the lemma is proved by passing to a subsequence of (σ_k) .

A priori, as $k \to \infty$ some portions of Im σ_k might escape to infinity (as $x^2 + r^2 \to \infty$) or collapse to the rotational axis $r = 0$. The next lemma shows that this is not the case.

Lemma 4.6. The sequence of self-shrinkers $(\Sigma_{\sigma_k})_{k=1}^{\infty}$ converges in Hausdorff distance to the round sphere $\Sigma_{\sigma_{\infty}}$ as $k \to \infty$.

Proof of Lemma [4.6.](#page-12-3) We use a maximum principle argument similar to that in the proof of Lemma [4.3.](#page-9-1) Let $\epsilon > 0$. Let $t_0 \in (-1, 0)$ be given by

(4.5)
$$
\sqrt{-t_0} = \frac{\sqrt{2n} + 0.5\epsilon}{\sqrt{2n} + \epsilon}.
$$

Let $k_{\epsilon} \in \mathbb{N}$ be large such that $R_{k_{\epsilon}}^{-1} < \epsilon/2$ and the following holds: there are two horizontal translations and scalings of the Angenent torus $\Sigma_{\bar{a}\pm}$ so that

- • the image of \bar{a}_+ and \bar{a}_- lie in $K_{R_{k_\epsilon}}$,
- $\sqrt{2n} + R_{k_{\epsilon}}^{-1} < \sqrt{x^2 + r^2} < \sqrt{2n} + \epsilon$ for all $(x, r) \in \operatorname{Im} \bar{a}_+,$
- $\sqrt{2n} - \epsilon < \sqrt{x^2 + r^2} < \sqrt{2n} - R_{k_{\epsilon}}^{-1}$ for all $(x, r) \in \underline{\text{Im}} \bar{a}$, and
- the MCF starting at $\Sigma_{\bar{a}_{\pm}}$ at $t = -1$ shrinks to $(\sqrt{2n} \pm 0.5\epsilon, 0)$ at time $t_e < t_0$.

By Lemma [4.5,](#page-12-1) \bar{a}_{\pm} do not intersect with σ_k when $k \geq k_{\epsilon}$.

FIGURE 3. Choices of \bar{a}_+ .

Now we claim that

(4.6)
$$
\sqrt{2n} - \epsilon < \sqrt{x^2 + r^2} < \sqrt{2n} + \epsilon, \quad \text{for all } (x, r) \in \text{Im } \sigma_k, \ k \geq k_{\epsilon}.
$$

By Lemma [4.5](#page-12-1) and that $R_{k_{\epsilon}}^{-1} < \epsilon$, it suffices to consider points outside $K_{R_{k_{\epsilon}}}$. Fix any $k \geq k_{\epsilon}$. If there is $(x, r) \in \text{Im}(\sigma_k)$ so that $\sqrt{x^2 + r^2} \geq \sqrt{2n} + \epsilon$, then since each σ_k is connected, there is a subarc β of σ_k connecting (x, r) and $\text{Im}(\sigma_k) \cap K_{R_{k_\epsilon}}$ passing through the region

$$
[\sqrt{2n}, \sqrt{2n} + \epsilon] \times (0, R_{k_{\epsilon}}].
$$

But this is impossible: by [\(4.5\)](#page-13-0), for all $-1 < t < t_0$, $\sqrt{-t\beta}$ contains a point $\{\sqrt{2n} +$ 0.5ϵ } × (0, R_{k_ε). Hence the MCF starting at $\Sigma_{\bar{a}_+}$ (at time −1) would intersect with $\sqrt{-t}\beta$} for some $t \in (-1, t_e)$ and this contradicts the parabolic maximum principle, since Σ_{σ_k} and $\Sigma_{\bar{a}_+}$ are disjoint. Using $\Sigma_{\bar{a}_-}$ and arguing similarly, we conclude that $\sqrt{x^2 + r^2} \leq \sqrt{2n - \epsilon}$ is also impossible.

Thus [\(4.6\)](#page-13-1) is shown and this implies $d_0(p, \Sigma_{\sigma_{\infty}}) < \epsilon$ for all $k \geq k_{\epsilon}$ and $p \in \Sigma_{\sigma_k}$. On the other hand, since $(\sigma_k|_{I_{k,R_k}})$ converges smoothly to σ_{∞} in $K_{2/\epsilon}$, by choosing a larger k_{ϵ} if necessary, we may assume that $d_0(p, \Sigma_{\sigma_k}) < \epsilon/2$ for all $p = (x, r\omega) \in \Sigma_{\sigma_{\infty}}$ with

FIGURE 4. A subarc β of σ_k passing through $\{x = \sqrt{2n} + \epsilon\}$ (left) or ${x = \sqrt{2n} - \epsilon}$ (right).

 $(x, r) \in K_{2/\epsilon}, k \geq k_{\epsilon}$ and $\omega \in \mathbb{S}^{n-1}$. Thus $d_0(p, \Sigma_{\sigma_k}) < \epsilon$ for all $p \in \Sigma_{\sigma_{\infty}}$ and $k \geq k_{\epsilon}$. By definition of the Hausdorff distance d^{μ} ,

$$
d^{\mathcal{H}}(\Sigma_{\sigma_{\infty}}, \Sigma_{\sigma_k}) = \max\{\sup_{p \in \Sigma_{\sigma_k}} d_0(p, \Sigma_{\sigma_{\infty}}), \sup_{p \in \Sigma_{\sigma_{\infty}}} d_0(p, \Sigma_{\sigma_k})\} < \epsilon
$$

for all $k \geq k_{\epsilon}$. Since $\epsilon > 0$ is arbitrary, $(\Sigma_{\sigma_k})_{k=1}^{\infty}$ converges in Hausdorff distance to the round sphere $\Sigma_{\sigma_{\infty}}$ and this finishes the proof of Lemma [4.6.](#page-12-3)

To summarize, we have shown that the sequence of self-shrinkers (Σ_{σ_k}) converges in Hausdorff distance to the sphere $\Sigma_{\sigma_{\infty}}$. Moreover, for all $\epsilon > 0$, the set

$$
\Sigma_{\sigma_k} \setminus (B_{(\sqrt{2n},0)}(\epsilon) \cup (B_{(-\sqrt{2n},0)}(\epsilon))
$$

decomposes into N disjoint graphs $\Sigma_k^1, \cdots \Sigma_k^N$ over Σ_{σ_∞} for large enough k, where $N \geq 2$, and for each $i = 1, \dots, N$, the convergence

$$
\Sigma_k^i \to \Sigma_{\sigma_\infty} \setminus (B_{(\sqrt{2n},0)}(\epsilon) \cup (B_{(-\sqrt{2n},0)}(\epsilon)), \quad \text{as } k \to \infty,
$$

is smooth graphical convergence.

Thus we can apply [\[15,](#page-19-13) Proposition 3.2] to conclude that $\Sigma_{\sigma_{\infty}}$ is L-stable. Although [\[15,](#page-19-13) Proposition 3.2] is stated only for $n = 2$, the same proof, which we now briefly describe, works for all $n \geq 2$ with only notational changes.

For each fixed k, since $\Sigma_k^1, \cdots, \Sigma_k^N$ are disjoint, we can order these N sheets by height (with respect to the outward unit normal of $\Sigma_{\sigma_{\infty}} = S$). Let the top and the bottom layers be represented respectively by two functions w_k^+ k , w_k^- defined on $\Omega_k \subset \mathcal{S} \setminus \{(\pm \sqrt{2n}, 0)\}$ so that $w_k^+ > w_k^-$, $\Omega_k \subset \Omega_{k+1}$ and $\cup_k \Omega_k = \mathcal{S} \setminus \{(\pm \sqrt{2n}, 0)\}\$. Fixing $x_0 \in \mathcal{S}$, then the sequence of functions

$$
u_k = \frac{w_k^+ - w_k^-}{w_k^+(x_0) - w_k^-(x_0)}
$$

converges locally smoothly on $S \setminus \{(\pm \sqrt{2n}, 0)\}\)$ to a smooth function u which satisfies $Lu = 0$, where L is the stability operator [\(2.1\)](#page-2-3). Using the Harnack inequality for linear second order elliptic equations and a maximum principle for minimal hypersurfaces (this is where Lemma [4.6](#page-12-3) is used), one can bound u uniformly. Hence u extends across $\{(\pm\sqrt{2n},0)\}\$ by the removable singularities lemma for *L*-harmonic functions [\[34\]](#page-19-20). Thus we have constructed a positive function on S which satisfies $Lu = 0$ and this is sufficient to conclude that S is L-stable (see [\[15\]](#page-19-13) for more details and [\[21\]](#page-19-21) for a general statement). Since all properly embedded self-shrinkers in \mathbb{R}^{n+1} are *L*-unstable ([\[14\]](#page-18-4), see also [\[15,](#page-19-13) Theorem 0.5]), we have arrived at a contradiction and hence $\Sigma_{\sigma_{\infty}}$ is not the sphere S.

Next we argue by contradiction that $\Sigma_{\sigma_{\infty}}$ is also not the plane P nor the cylinder \mathcal{C} . The arguments are similar to those for S , thus we just point out the differences.

If $\Sigma_{\sigma_{\infty}}$ is the plane \mathcal{P} , then $\sigma_k|_{I_{k,R}}$ converges smoothly to the r-axis in K_R . Hence there is $k_1 = k_1(R) \in \mathbb{N}$ so that $\sqrt{x^2 + r^2} < R^{-1}$ for all $(r, x) \in \text{Im } \sigma_k \cap K_R$ and $k \geq k_1$. For any $R \geq R_1$, let

$$
K_R^{\pm} := \{(x, r) \in K_R : \pm x \ge R^{-1}\}
$$

One can argue that either one of $K_{R_1}^{\pm}$ must intersect the image of σ_k when $k \geq k_1$: if not, then either one of $K_{R_1}^{\pm}$ would lie in the compact region bounded by σ_k . This would lead to a contradiction by putting in suitably scaled and horizontally translated Angenent doughnuts in $K_{R_1}^{\pm}$.

Similar to the previous argument for the sphere S, there is $N \geq 2$ so that for all $k \in \mathbb{N}$, $\text{Im}\,\sigma_k \cap K_{R_k}$ contains N connected components which intersect K_{R_1} , and $\text{Im}\,\sigma_k \cap K_{R_k}$ converges smoothly graphically to the r-axis with multiplicity N.

As in the proof of Lemma [4.6,](#page-12-3) for all $R > R_1$, one can show that Im $\sigma_k \cap K_R$ converges (locally) in Hausdorff distance to $\text{Im}\,\sigma_{\infty} \cap K_R$ as $k \to \infty$ (unlike the case of the sphere, there might be mass loss as $R \to \infty$). This is still sufficient for us to apply [\[15,](#page-19-13) Proposition 3.2 to conclude that the plane is L-stable, which is impossible $[14]$.

For the remaining case for the cylinder C, the curves $\sigma_k|_{I_{k,R}}$ converge smoothly to ${n \choose r} = \sqrt{2(n-1)}$ in K_R . Hence there is $k_1 = k_1(R) \in \mathbb{N}$ such that $|r - \sqrt{2(n-1)}| < R^{-1}$ for all $(x, r) \in \text{Im } \sigma_k \cap K_R$ and $k \geq k_1$. By fitting Angenent doughnuts inside

$$
K_{R_1}^> = \{(x, r) \in K_{R_1} : r > \sqrt{2(n-1)} + R^{-1}\},
$$
\n
$$
K_{R_1}^< = \{(x, r) \in K_{R_1} : r < \sqrt{2(n-1)} - R^{-1}\}
$$

respectively (in $K_{R_1}^>$ we insert a large Angenent doughnut), there is $N \geq 2$ so that Im $\sigma_k \cap K_{R_k}$ converges smoothly graphically to $\{r = \sqrt{2(n-1)}\}$ with multiplicity N. Then again we apply [\[15,](#page-19-13) Proposition 3.2].

To sum up, $\Sigma_{\sigma_{\infty}}$ is neither the plane P nor the cylinder C nor the sphere S. By the classification Theorem [2.4,](#page-3-0) σ_{∞} is an embedded closed geodesic in (H, g_A) and the convergence $\Sigma_{\sigma} \to \Sigma_{\sigma}$ is smooth. This finishes the proof of Theorem 1.2. convergence $\Sigma_{\sigma_k} \to \Sigma_{\sigma_\infty}$ is smooth. This finishes the proof of Theorem [1.2.](#page-1-1)

Remark 4.1. In the proof of Theorem [1.2,](#page-1-1) we argued by contradiction using L-stability that $\Sigma_{\sigma_{\infty}}$ is neither the sphere S, nor the cylinder C nor the plane P. In this remark we give an alternative argument ruling out S and C by instead using the entropy bound $E_n \leq E_2$ (see Lemma [5.1\)](#page-17-0). By Lemma [4.3](#page-9-1) and Lemma [4.4,](#page-12-0) for each k large, one can find N disjoint subarcs $\sigma_k^1 \cdots, \sigma_k^N$ of σ_k so that for each $i = 1, \cdots, N, (\sigma_k^i)$ converges locally smoothly to σ_{∞} as $k \to \infty$. In particular,

$$
\lim_{k \to \infty} L_A(\sigma_k) \geq NL_A(\sigma_\infty).
$$

By (2.3) , we obtain

$$
E_2 \ge E_n \ge \lim_{k \to \infty} \lambda(\Sigma_{\sigma_k}) \ge N\lambda(\Sigma_{\sigma_\infty}).
$$

If $\Sigma_{\sigma_{\infty}} = S$ or C, then $\lambda(\Sigma_{\infty}) > \sqrt{2}$ by [\[37,](#page-19-22) A.4 Lemma]. Thus $N\sqrt{2} \leq E_2$, which is impossible as $N \ge 2$ and $E_2 \sim 2.2476 < 2\sqrt{2}$.

Since $E_n > 2$ and $\lambda(\mathcal{P}) = 1$, this argument is however not sufficient to show $\Sigma_{\sigma_{\infty}} \neq \mathcal{P}$.

Next we prove several consequences of Theorem [1.2.](#page-1-1)

First, a simple contradiction argument using Theorem [1.2](#page-1-1) shows that the profile curve of every embedded rotationally symmetric self-shrinking doughnut must lie in a fixed compact subset in \mathbb{H} . Together with Theorem [1.1](#page-1-0) and [\[5,](#page-18-12) Theorem 1.1], we obtain an upper bound on the index. (for the definition of index of a self-shrinker, see [\[5,](#page-18-12) Definition 2.18].)

Corollary 4.7. There is $I \in \mathbb{N}$ so that for any complete embedded rotationally symmetric self-shrinker Σ in \mathbb{R}^3 , one has $i(\Sigma) \leq I$, where $i(\Sigma)$ is the index of Σ .

Unlike our Theorem [1.1,](#page-1-0) the bound I in the above corollary is not explicit.

The next corollary and Theorem [1.3](#page-2-1) generalize [\[32,](#page-19-9) Theorem 1.3] to any dimension and by allowing rotationally symmetric self-shrinkers with non-convex profile curve and is a consequence of the fact that the Angenent metrics g_A are real analytic. See also [\[11\]](#page-18-13) for a similar theorem for Lagrangian self-shrinking tori in \mathbb{R}^4 .

Corollary 4.8. For each $n \geq 2$, there is a finite set $S_n \subset [1, E_n]$ so that $\lambda(\Sigma) \in S_n$ for all complete embedded rotationally symmetric self-shrinkers Σ in \mathbb{R}^{n+1} .

Proof. By Theorem [2.4,](#page-3-0) it suffices to consider only rotationally symmetric self-shrinkers that are diffeomorphic to $\mathbb{S}^{n-1}\times \mathbb{S}^1$. Let σ be the profile curve of a rotationally symmetric self-shrinking doughnut in \mathbb{R}^{n+1} . Thus σ is a geodesic in (\mathbb{H}, g_A) . Since g_A is real analytic, the mapping

$$
\gamma \mapsto E_A(\gamma) := \frac{1}{2} \int_{\mathbb{S}^1} g_A(\dot{\gamma}, \dot{\gamma})
$$

is a real analytic functional defined on all closed curves γ close to σ . Note that $DE_A(\gamma) =$ 0 if and only if γ is a geodesic, here $DE_A(\gamma)$ is the L^2 -gradient of E_A at γ . By the celebrated Lojasiewicz-Simon gradient inequality [\[35,](#page-19-23) equation (2.2)], there is $C_2 > 0$ and $\theta \in (0, 1/2)$ so that

$$
|E_A(\gamma) - E_A(\sigma)|^{1-\theta} \leq C_2 ||DE_A(\gamma)||_{L^2},
$$

for all γ which are $C^{2,\alpha}$ -close to σ . Hence if γ is another geodesic close to σ , we have $E_A(\gamma) = E_A(\sigma)$. Since geodesics are parametrized by constant length, one has $L_A(\gamma) =$ $L_A(\sigma)$. Thus the length functional $\gamma \mapsto L_A(\gamma)$ is locally constant on the space of all closed geodesics. Since the space of simple closed geodesics is compact by Theorem [1.2,](#page-1-1) the length functional has a finite image. Together with [\(2.3\)](#page-3-2), this finishes the proof of the corollary. \Box

Lastly, we prove Theorem [1.3.](#page-2-1)

Proof of Theorem [1.3.](#page-2-1) In [\[32,](#page-19-9) section 5], Mramor studies the Poincaré map of the (up to renormalization of length) geodesic equation in (\mathbb{H}, g_A)

(4.7)
$$
x' = \cos \theta, \quad r' = \sin \theta, \quad \theta' = \frac{x}{2} \sin \theta + \left(\frac{n-1}{r} - \frac{r}{2}\right) \cos \theta.
$$

For any $R > 0$ and θ , let $(x_{R,\theta}(t), y_{R,\theta}(t), \theta_{R,\theta}(t))$ be the maximal solution to [\(4.7\)](#page-17-1) with initial value $(0, R, \theta)$. Let $T^* = T^*(R, \theta)$ be the second time at which $x_{R,\theta} = 0$ occurs. By $[32, \text{Lemma } 5.1]$, the fixed points of the Poincaré map

$$
P: \mathbb{R} \times (0, \infty) \to \mathbb{R} \times (0, \infty), \quad P(R, \theta) = (r_{R, \theta}(T^*), \theta_{R, \theta}(T^*))
$$

are either isolated points or analytic curves in $\mathbb{R} \times (0, \infty)$. Arguing as in the proof of [\[32,](#page-19-9) Theorem 1.3], the map

$$
P_f: (0, \infty) \to (0, \infty), \quad P_f(R) = r_{R,0}(T^*(R, 0), 0)
$$

has isolated fixed points. Since the profile curve of any embedded self-shrinker with reflectional symmetry must intersect the fixed points of P_f , together with Theorem [1.2,](#page-1-1) Theorem [1.3](#page-2-1) has been proven.

5. APPENDIX: THE SEQUENCE (E_n)

In this appendix, we show the following lemma.

Lemma 5.1. The sequence (E_n) defined in [\(3.1\)](#page-4-0) satisfies $2 < E_n \le E_2$ and

(5.1)
$$
\lim_{n \to \infty} E_n = \sqrt{\frac{4\pi}{3}}.
$$

Proof. It is proved in [\[37,](#page-19-22) A.4 Lemma] that the entropy of the *n*-sphere $\lambda(\mathbb{S}^n)$ satisfies

(5.2)
$$
\lambda(\mathbb{S}^n) = \left(\frac{n}{2\pi e}\right)^{n/2} \omega_n
$$

and the sequence $(\lambda(\mathbb{S}^n))$ is strictly decreasing. Also,

(5.3)
$$
\lim_{n \to \infty} \lambda(\mathbb{S}^n) = \sqrt{2}.
$$

From (3.1) and (5.2) we obtain

(5.4)
$$
E_n = \sqrt{\frac{2\pi}{3} \frac{1+x_n}{1+2x_n/3}} \left(\frac{1}{e} (1+x_n)^{1/x_n}\right)^{a_n/4} \lambda(\mathbb{S}^{n-1}),
$$

where

$$
a_n = y_n - 2(n - 1), \quad x_n = \frac{a_n}{2(n - 1)}.
$$

Direct calculations give

(5.5)
$$
\frac{1}{2} < a_n < \frac{2}{3}, \quad 0 < x_n < 1
$$

and

(5.6)
$$
\lim_{n \to \infty} a_n = \frac{2}{3}, \quad \lim_{n \to \infty} x_n = 0.
$$

Using (5.4) , (5.3) and (5.6) , one obtains (5.1) .

Next we show $2 < E_n \leq E_2$. By the Taylor expansion of $\ln(1+x)$, we have $x + x^3/3$ $ln(1+x) > x - x^2/2$ for all $x \in (0,1)$. Thus

$$
e^{\frac{x^2}{3}} > \frac{1}{e}(1+x)^{1/x} > e^{-\frac{x}{2}},
$$
 for all $x \in (0,1)$.

Together with $\lambda(\mathbb{S}^{n-1}) > \sqrt{2}$, [\(5.5\)](#page-17-7) and [\(5.4\)](#page-17-3),

(5.7)
$$
\frac{\sqrt{2\pi (3 + (n-1)^{-1})}}{3} e^{\frac{1}{162(n-1)^2}} \lambda(\mathbb{S}^{n-1}) > E_n > \sqrt{\frac{4\pi}{3}} e^{-\frac{1}{36(n-1)}}.
$$

Since $\lambda(\mathbb{S}^{n-1})$ is decreasing, the upper bound in [\(5.7\)](#page-18-14) is strictly decreasing in n. Also, the lower bound in [\(5.7\)](#page-18-14) is strictly increasing in n. Plugging in $n = 4$ in the upper and lower bound of [\(5.7\)](#page-18-14) gives

$$
2.21823 \sim \frac{\sqrt{10}\pi}{e^{1093/729}} > E_n > \sqrt{\frac{4\pi}{3}}e^{-\frac{1}{108}} \sim 2.02780, \text{ for all } n \ge 4.
$$

The inequality implies $2 < E_n < E_2$ for all $n \geq 4$. The case $n = 2, 3$ can be checked directly.

 \Box

REFERENCES

- 1. Abresch, U.; Langer, J.: *The normalized curve shortening flow and homothetic solutions*. J. Differential Geom. 23(2): 175-196 (1986).
- 2. Angenent S.B.: *Shrinking Doughnuts*. In: Lloyd N.G., Ni W.M., Peletier L.A., Serrin J. (eds) Nonlinear Diffusion Equations and Their Equilibrium States, 3. Progress in Nonlinear Differential Equations and Their Applications, vol 7. Birkhäuser, Boston, MA.
- 3. Bernstein J.; Wang, L.: *A sharp lower bound for the entropy of closed hypersurfaces up to dimension six*. Invent. Math. 206, 601–627 (2016).
- 4. Berchenko-Kogan, Y: *The entropy of the angenent torus is approximately 1.85122*. Experimental Mathematics, 0(0):1-8, 2019
- 5. Berchenko-Kogan, Y.: *Bounds on the index of rotationally symmetric self-shrinking tori*. Geometriae Dedicata, 213, 82–106, 2021.
- 6. Brendle, S.: *Embedded self-similar shrinkers of genus 0*. Ann. Math. (2) 183, No. 2, 715-728 (2016).
- 7. Bridson, Martin R., and André Haefliger; *Metric spaces of non-positive curvature*. Vol. 319. Springer Science & Business Media, 2013.
- 8. Burago, D.; Burago, Y.; Ivanov, S.: *A course in metric geometry*, Graduate Studies in Mathematics, Volume: 33; 2001
- 9. Buzano, R.; Nguyen, H.T.; Schulz, M.B.: *Noncompact self-shrinkers for mean curvature flow with arbitrary genus*. https://arxiv.org/abs/2110.06027
- 10. Chen, J.; Ma, J.M.S.: *The space of compact self-shrinking solutions to the Lagrangian mean curva*ture flow in \mathbb{C}^2 . J. Reine Angew. Math. 743, 229-244 (2018).
- 11. Chen, J.; Ma, J.M.S.: *Geometry of Lagrangian self-shrinking tori and applications to the Piecewise Lagrangian Mean Curvature Flow*. Am. J. Math. 143, No. 1, 227-264 (2021).
- 12. Chodosh, O.; Choi, K.; Mantoulidis, C.; Schulze, F.: *Mean curvature flow with generic initial data*, https://arxiv.org/abs/2003.14344
- 13. Colding, T.; Ilmanen, T.; Minicozzi, B.; White, B.: *The round sphere minimizes entropy among closed self-shrinkers*. J. Differ. Geom. 95 (2013), no. 1, 53–69.
- 14. Colding, T.; Minicozzi, W.: *Generic mean curvature flow I*. Ann. of Math. 175 (2012), 755-833
- 15. Colding, T.; Minicozzi, W.: *Smooth compactness of self-shrinkers*. Comment. Math. Helv. 87, No. 2, 463-475 (2012).
- 16. Ding, Q.; Xin, Y.L.: *Volume growth eigenvalue and compactness for self-shrinkers*. Asian J. Math. 17, No. 3, 443-456 (2013).
- 17. Drugan, G.: *An immersed* S 2 *self-shrinker*. Trans. Am. Math. Soc. 367, 3139-3159 (2015)
- 18. Drugan, G.; Lee, H.; Nguyen, X.H. *A survey of closed self-shrinkers with symmetry*. Results Math. 73 (2018), no. 1, Paper No. 32, 32 pp.
- 19. Drugan, G., Kleene, S.J.: *Immersed self-shrinkers*. Trans. Am. Math. Soc. 369, 7213-7250 (2017)
- 20. Drugan, G.; Nguyen, X. H.: *Shrinking doughnuts via variational methods*. J. Geom. Anal. 28 (2018), no. 4, 3725-3746.
- 21. Fischer-Colbrie, D.; Schoen, R.: *The structure of complete stable minimal surfaces of non-negative scalar curvature*. Commun. Pure Appl. Math. 33, 199-211 (1980).
- 22. Garcke, H.; N¨urnberg, R.: *Numerical approximation of boundary value problems for curvature flow and elastic flow in Riemannian manifolds*. Numer. Math. 149, No. 2, 375-415 (2021).
- 23. Hershkovits, O.; White, B.: *Sharp entropy bounds for self-shrinkers in mean curvature flow*. Geom. Topol. 23 (2019) 1611–1619.
- 24. Huisken, G.: *Asymptotic behavior for singularities of the mean curvature flow*. J. Differ. Geom. 31, No. 1, 285-299 (1990).
- 25. Ilmanen, T.: *Elliptic regularization and partial regularity for motion by mean curvature*. Mem. Am. Math. Soc. 520, 90 p. (1994).
- 26. Impera, D.; Pigola, S.; Rimoldi, M.: *The Frankel property for self-shrinkers from the viewpoint of elliptic PDEs*. J. Reine Angew. Math. 773, 1-20 (2021).
- 27. Kapouleas, N.; Kleene, S. J.; Møller, N. M. *Mean curvature self-shrinkers of high genus: non-compact examples*. J. Reine Angew. Math. 739, 1-39 (2018).
- 28. Ketover, D.: *Self-shrinking Platonic solids*. https://arxiv.org/abs/1602.07271
- 29. Kleene, S., Møller, N.M.: *Self-shrinkers with a rotational symmetry*, Trans. Amer. Math. Soc. 366 (2014), 3943-3963
- 30. Klingenberg, Wilhelm P.A.. *Riemannian Geometry*, Berlin, New York: De Gruyter, 2011. https://doi.org/10.1515/9783110905120
- 31. Møller, N.M.: *Closed self-shrinking surfaces in* R ³ *via the torus*. [arXiv:1111.7318](http://arxiv.org/abs/1111.7318)
- 32. Mramor, A.: *Compactness and finiteness theorems for rotationally symmetric self shrinkers*. J. Geom. Anal. 31, 5094-5107 (2021)
- 33. Nguyen, X. H. *Construction of complete embedded self-similar surfaces under mean curvature flow, Part III*. Duke Math. J. 163 (2014), no. 11, 2023-2056.
- 34. Serrin, J.: *Removable Singularities of Solutions of Elliptic Equations*, Arch. Rational Mech. Anal. 17, 67–78 (1964). https://doi.org/10.1007/BF00283867
- 35. Simon, L.: *Asymptotics for a class of non-linear evolution equations, with applications to geometric problems*, Ann. Math. 118 (1983) 525-571
- 36. Song, A.: *A maximum principle for self-shrinkers and some consequences*. [arXiv:1412.4755](http://arxiv.org/abs/1412.4755)
- 37. Stone, A.: *A density function and the structure of singularities of the mean curvature flow*. Calc. Var. Partial Differ. Equ. 2, No. 4, 443-480 (1994).
- 38. Sun, A.; Wang, Z.: *Compactness of self-shrinkers in* R ³ *with fixed genus*. Adv. Math. 367, Article ID 107110, 39 p. (2020).
- 39. Wei, G.; Wylie, W.: *Comparison geometry for the Bakry-Emery Ricci tensor*. J. Differ. Geom. 83, No. 2, 377-405 (2009).

John Man Shun Ma Department of Mathematical Science, University of Copenhagen Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark email: jm@math.ku.dk

Ali Muhammad Department of Mathematical Science, University of Copenhagen Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark email: ajhm@math.ku.dk

Niels Martin Møller Department of Mathematical Science, University of Copenhagen Universitetsparken 5, DK-2100 Copenhagen Ø, Denmark email: nmoller@math.ku.dk