Virtual permutations and polymorhisms

YURY A.NERETIN^{[1](#page-0-0)}

There is a natural map from a symmetric group S_n to a smaller symmetric group S_{n-1} , we write a decomposition of a permutation into a product of disjoint cycles and remove the element n from this expression. For this reason there exists the inverse limit $\mathfrak S$ of sets S_n . We equip S_n with the uniform distribution (or more generally with an Ewens distribution) and get a structure of a measure space on \mathfrak{S} (it is called 'virtual permutations' or 'Chinese restaurant process'), a double $S_{\infty} \times S_{\infty}$ of an infinite symmetric group acts on $\mathfrak S$ by left and right 'multiplications'. We discuss the closure of $S_{\infty}\times S_{\infty}$ in the semigroup of polymorphisms (spreading maps with spreaded Radon–Nikodym derivatives) of S. We get formulas for some polymorphisms, in particular for the center of the closure. Expressions are sums of multiple convolutions of Dirichlet distributions, summation sets are certain collections of dessins d'enfant.

1. Introduction. Virtual permutations, chips, and polymorphisms

The topic of the paper is formulas for the action of the Olshanski semigroups of chips on the virtual permutations by polymorphims, in this section we explain these notions and formulate the problem.

1.1. Some notation. Denote by \mathbb{Z}_+ (resp., \mathbb{R}_+) the set of nonnegative integers (resp. reals), by \mathbb{R}° (resp. $\mathbb{R}_{>0}$) the multiplicative group (resp. the set) of positive reals. For a set A we denote by #A the number of its elements. By $A \sqcup B$, $\sqcup_i A_j$ we denote disjoint unions of sets.

We denote a list (v_1, v_2, \dots) by $\{v_i\}$ or by $\{v_i\}_i$ (we apply this notation for both ordered or nonordered lists). For finite nonordered lists $\{k_i\}$ of positive integers we use a notation

(1.1) $\iota_m[\{k_j\}] := \text{number of entries of } m \text{ to a list } \{k_j\}.$

By $\delta_X[[a]]$ we denote the unit atomic measure (delta-function) at a point a of a space X. We denote product-measures by $\mu_1 \times \mu_2$ (to not be confused with the symbol \times of a multiplication).

Denote by S_n the symmetric group, which is considered as the group of permutations of the set

$$
I_n:=\{1,2,\ldots,n\},\
$$

by S_{∞} the group of all permutations of N with finite supports,

$$
S_{\infty} = \lim_{n \to \infty} S_n = \bigcup_{n=1}^{\infty} S_n.
$$

This group is countable and is equipped with the discrete topology.

By \overline{S}_{∞} we denote the group of all permutations of N. This group is continual and is equipped with the unique separable topology compatible with the structure of the group. A sequence $\sigma_j \in \overline{S}_{\infty}$ converges to σ if for any $p \in \mathbb{N}$ we have $\sigma_j(p) = \sigma(p)$ for sufficiently large j .

¹Supported by the grant FWF, Project P31591.

We can regard elements of symmetric groups as diagrams of the form

$$
\begin{array}{c}\n 1 \ 2 \ 3 \ 4 \ 5 \ 6 \\
 \times \ 1 \ \times \dots \\
 1 \ 2 \ 3 \ 4 \ 5 \ 6\n \end{array}
$$
\n
$$
(1.2)
$$

We draw a collection of 'upper circles' enumerated by natural numbers and a collection of 'lower circles' also enumerated by $\mathbb N$. If g sends i to j, then we connect by an arc i -th upper circle with j -th lower circle.

A product of permutations corresponds to a gluing of diagrams

$$
\begin{array}{ccl}\nX & = & X1 \\
\times & \times & 1\n\end{array}
$$

A. Virtual permutations and the bisymmetric group

1.2. The map Υ. Define a canonical map

$$
\Upsilon_{n-1}^n : S_n \to S_{n-1}
$$

in the following way. Let $\sigma \in S_n$.

1) If $\sigma(n) = n$, we set $\Upsilon_{n-1}^n \sigma(j) = \sigma(j)$ for all $j < n$.

2) Let $\sigma(n) = \alpha \neq n$. Then $\sigma^{-1}(n) = \beta$ also $\neq n$. Then we assume $\Upsilon_{n-1}^n \sigma(\beta) =$ α . For $j \neq \beta$ we set $\Upsilon_{n-1}^n \sigma(j) = \sigma(j)$.

Example. For instance,

$$
\Upsilon_4^5 \begin{pmatrix} 1 & 2 & 3 & 4 & 5 \\ 3 & 2 & 4 & 1 & 5 \end{pmatrix} = \begin{pmatrix} 1 & 2 & 3 & 4 \\ 3 & 2 & 4 & 1 \end{pmatrix};
$$

$$
\Upsilon_4^5 \begin{pmatrix} 1 & 2 & \boxed{3} & 4 & 5 \\ 3 & 2 & 5 & 1 & \boxed{4} \end{pmatrix} = \begin{pmatrix} 1 & 2 & \boxed{3} & 4 \\ 3 & 2 & \boxed{4} & 1 \end{pmatrix}
$$

.

 \mathcal{L}

On the language of diagrams we have:

$$
\Upsilon_4^5: \begin{array}{c} 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{4}} 5 \\ 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{4}} 5 \end{array} \rightarrow \begin{array}{c} 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{4}} 5 \\ 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{4}} 5 \end{array} = \begin{array}{c} 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{5}} \\ 1 \stackrel{2}{\cancel{5}} 3 \stackrel{4}{\cancel{4}} \end{array}
$$

Namely, we add an arc connecting the top '5' and the bottom '5'. We get a compound arc $3^{top}-5^{bottom}-5^{top}-4^{bottom}$ and consider it as an arc $3^{top}-4^{bottom}$ (here and below we consider arcs with fixed ends up to isotopies fixing ends). Similarly,

$$
\Upsilon_4^5: \begin{array}{c} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \end{array} \longrightarrow \begin{array}{c} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 3 & 4 & 5 \end{array} = \begin{array}{c} 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \\ 1 & 2 & 3 & 4 \end{array}
$$

We add an arc $5^{top}-5^{bottom}$, get a cycle, and remove it. \boxtimes

The map Υ_{n-1}^n is $S_{n-1} \times S_{n-1}$ -equivariant in the following sense:

(1.3)
$$
\Upsilon_{n-1}^n(h_1^{-1}gh_2) = h_1^{-1}\Upsilon_{n-1}^n(g)h_2, \quad \text{where } g \in S_n, h_1, h_2 \in S_{n-1}.
$$

ANOTHER DESCRIPTION OF THE MAP Υ_{n-1}^n . Decomposing $g \in S_n$ as a product of disjoint cycles,

(1.4)
$$
g = (k_1^1 k_2^1 \dots) (k_1^2 k_2^2 \dots) \dots
$$

and removing n from this expression, we get an element $\Upsilon_{n-1}^n(g)$ defined as product of disjoint cycles.

More generally, for $n \geq m$ we define a canonical map $\Upsilon_m^n : S_n \to S_m$,

$$
\Upsilon_m^n(g) := \Upsilon_{m-1}^m \dots \Upsilon_{n-2}^{n-1} \Upsilon_{n-1}^n(g).
$$

Equivalently, we remove $m + 1, m + 2, \ldots, n$ from expression [\(1.4\)](#page-2-0) and get an element of S_m .

1.3. The inverse limit of the spaces S_n . Thus we have the following chain of surjective maps

$$
(1.5) \qquad \qquad \cdots \stackrel{\Upsilon_{n-1}^n}{\longleftarrow} S_n \stackrel{\Upsilon_{n+1}^{n+1}}{\longleftarrow} S_{n+1} \stackrel{\Upsilon_{n+2}^{n+2}}{\longleftarrow} S_{n+2} \stackrel{\Upsilon_{n+3}^{n+3}}{\longleftarrow} \cdots
$$

So we can consider the inverse limit of sets S_n

$$
\mathfrak{S}=\varprojlim S_n.
$$

By the definition, $\mathfrak S$ consists of sequences

$$
\widetilde{\sigma} = (\sigma_1, \sigma_2, \sigma_3, \dots),
$$
 where $\sigma_k \in S_k$ and $\Upsilon_{k-1}^k \sigma_k = \sigma_{k-1}.$

In particular, we have a canonical map $\Upsilon_n^{\infty} : \mathfrak{S} \to S_n$ defined by $\Upsilon_n^{\infty} \tilde{\sigma} = \sigma_n$.

A point of the space $\mathfrak S$ can be described in the following way. We fix a partition of N into a disjoint union of subsets (*pre-tables*)

(1.6) N = M¹ ⊔ M² ⊔

and fix a cyclic ordering in each subset. A map Υ_n^{∞} is the removing all elements $>n$. Then we get the set $I_n = \{1, 2, ..., n\}$ splitted into a disjoint union of cyclically ordered subsets, i.e., an element of S_n .

The space $\mathfrak S$ is not an inverse limit in the category of groups, since generally

$$
\Upsilon_{n-1}^n(g_1g_2) \neq \Upsilon_{n-1}^n(g_1)\Upsilon_{n-1}^n(g_2).
$$

In particular, $\mathfrak S$ does not have a group structure.

However, for $n > m$ by (1.3) ,

$$
\Upsilon_m^n(h_1^{-1}gh_2) = h_1^{-1}\Upsilon_m^n(g)h_2
$$
, where $g \in S_n$, $h_1, h_2 \in S_m$.

Therefore $S_m \times S_m$ acts on G. Since m is arbitrary, we get an action of the double $S_{\infty} \times S_{\infty}$ on \mathfrak{S} .

It is more natural to consider such inverse limit as a measure space.

1.4. Virtual permutations. For $g \in S_n$ denote by [g] the number of its disjoint cycles. Fix $z > 0$. The Ewens distribution is a probabilistic measure μ_n^z on S_n defined by

$$
\mu_n^z(g) = \frac{z^{[g]}}{z(z+1)\dots(z+n-1)},
$$

for $z = 1$ we get a uniform distribution on S_n .

It easy to show that the pushforward of the Ewens distribution μ_n^z is the Ewens distribution μ_{n-1}^z (i. e., for any subset $K \subset S_{n-1}$ the measure of its preimage

 $(\Upsilon_{n-1}^n)^{-1}(K)$ coincides with the measure of K). We define the space \mathfrak{S}^z of *virtual permutations* as the inverse limit

(1.7)
$$
(\mathfrak{S}^z, \mu^z) := \lim_{\longleftarrow} (S_n, \mu_n^z)
$$

in the category of measure spaces. Namely, we equip $\mathfrak S$ with the measure μ^z defined from the condition: for any n and $g \in S_n$ we set $\mu^z((\Upsilon_n^{\infty})^{-1}g) = \mu_n^z(g)$. By the Carathéodory extension theorem we get a well-defined measure on $\mathfrak{S}.$

REMARK. Apparently, a construction of \mathfrak{S}^z was firstly appeared in Aldous [\[1\]](#page-32-0) (1985), 11.19, with a reference to J. Pitman (see, also, Pitman [\[42\]](#page-34-0)). The action of the group $S_{\infty} \times S_{\infty}$ on G appeared in Kerov, Olshanski, Vershik [\[12\]](#page-33-0). Notice that for a fixed z for each n we have a distribution of lengths of cycles on S_n , i.e., a distribution on the set of partitions of n. So we have a Markov process of growth of such partitions, this process arises to the work of Ewens [\[6\]](#page-32-1), 1972, on population genetics. \boxtimes

1.5. A more convenient description of \mathfrak{S}^z . It is done in two steps.

Step 1. We present the limit distribution of lengths of cycles, see Kingman [\[16\]](#page-33-1). Consider the measure ν on the half-line $x > 0$ defined by

$$
d\nu^z(x) = z x^{-1} e^{-x} dx.
$$

Consider the Poisson measure π^z on the set of (non-ordered) countable subsets (see, e.g., [\[16\]](#page-33-1), [\[24\]](#page-33-2), Sect. X.4) $\{s_1, s_2, \ldots\}$ $\{s_1, s_2, \ldots\}$ $\{s_1, s_2, \ldots\}$ in R defined by the measure² ν^z . It can be easily shown that the series $\sum s_j$ converges a.s. We define a collection $\{\ell_\omega\}$ as $\ell_{\omega} = s_{\omega}/\sum s_j$ and get a measure *(the Poisson–Dirichlet distribution)* on the set of unordered collections $\{\ell_{\omega}\}\$ such that $\sum \ell_{\omega} = 1$.

For a given $\{\ell_{\omega}\}\$ consider a collection U_1, U_2, \ldots of oriented circles of lengths ℓ_1, ℓ_2, \ldots . We imagine them as circles on plane oriented clockwise and call them by *tables*, sets ⊔U^ω we call by *restaurants* ('Chinese restaurants'). Denote the space of all restaurants equipped with the Poisson-Dirichlet distribution by \mathfrak{T}^z .

We can regard points of \mathfrak{T}^z as non-ordered collections of tables $\{U_\omega\}$ or as nonordered collections of lengths $\{\ell_{\omega}\}.$

REMARK. Clearly, the numbers ℓ_{ω} are pairwise distinct a.s. \boxtimes

STEP 2. We chose a random sequence in $\sqcup_{\omega} U_{\omega}$ such that each element is distributed uniformly with respect to the Lebesgue measure on $\sqcup_{\omega} U_{\omega}$. We denote points of a sequence (we call them *guests*) by $\overline{1}$, $\overline{2}$, $\overline{3}$, Configurations of guests on tables are defined upto rotations of tables. A *virtual permutation* or an *occupied restaurant* is a collection $\{\ell_{\omega}\}\$ distributed according the Poisson–Dirichlet law, the corresponding restaurant $\sqcup U_\omega$, and a random sequence $\overline{1}, \overline{2}, \cdots \in \sqcup_\omega U_\omega$. We denote the measure space of all virtual permutations by \mathfrak{S}^z , denote by μ^z the measure on $\mathfrak{S}^z.$

We denote *occupied tables* by \overline{U}_{ω} .

REMARK. Clearly, for almost all points of \mathfrak{S}^z a sequence of guests $\overline{1}, \overline{2}, \ldots$ is dense in $\sqcup_{\omega} U_{\omega}$ and its elements are pairwise distinct. \boxtimes

²For a measurable subset $A \subset \mathbb{R}_{>0}$ denote by $\Omega(A, k)$ the set ('event') of all configurations whose intersections with A have precisely k point. We assume that a propbabibility of $\Omega(A, k)$ is $\exp{-\nu^z(A)}\nu^z(A)^k/k!$ and for pairwise disjoint sets A_j and arbitrary k_j the events $\Omega(A_j, k_j)$ are independent.

REMARK. The set of all tables of a given restaurant can be identified with N, we can order sets of tables according a length ℓ_{ω} or according a minimal number of a guest sitting at the table. We prefer to think that a set of tables of a given restaurant is an abstract countable set and do not identify such sets for different restaurants. ⊠

There is a canonical map $\Upsilon_n^{\infty} : \mathfrak{S}^z \to S_n$. Namely, let us 'forget' all guest $\overline{n+1}$, $\overline{n+2}, \ldots \in \sqcup U_{\omega}$. After this, each table U_{ω} contains a finite collection of guests i_{ω}^1 , $i_{\omega}^2, \ldots, i_{\omega}^{k_{\omega}}$ seating at the table. So we get a partition of $\{1, \ldots, n\}$ into a disjoint union of cyclically ordered sets. Thus we come to a permutation decomposed into a product of disjoint cycles.

The pushforward of the measure μ^z to S_n is the Ewens measure μ_n^z . We also have $\Upsilon_{n-1}^n \circ \Upsilon_n^{\infty} = \Upsilon_{n-1}^{\infty}$.

It remains to explain how pre-tables defined in Subsect. [1.4](#page-2-1) generate tables. Let $\tilde{\sigma}$ be a point of the inverse limit [\(1.7\)](#page-3-1). Consider two points i, j on one pre-table. For each $N \geq i, j$ consider the cycle of $\Upsilon^{\infty}_{N} \widetilde{\sigma}$ containing i, j, say $(... in_1 m_2 ... m_p j ...)$. Denote by $p_{ij}(N)$ the number of elements between i and j. Then (see [\[48\]](#page-34-1), [\[14\]](#page-33-3)) for almost all points of \mathfrak{S}^z for all pairs $\overline{i}, \overline{j}$ lying on one pre-table we define

(1.8) length of arc
$$
[\overline{i}, \overline{j}] := \lim_{N \to \infty} \frac{p_{ij}(N)}{N}
$$
.

The limit exists a.s. for all i, j . Then we get an occupied restasurant.

1.6. The action of $S_{\infty} \times S_{\infty}$ on G. The space \mathfrak{S}^z is not a group, but the group $S_{\infty} \times S_{\infty}$ acts on G by 'left and right multiplications'. Let $\mathfrak{u} \in \mathfrak{S}^z$, let $\{\overline{U}_{\alpha}\}\$ be its occupied tables. Set $u = \Upsilon_n^{\infty} \mathfrak{u}$. For each $i \in I_n$ we denote by $\overline{U}[i]$ the table containing the guest \overline{i} , by $\overline{U}_+[i]$ the arc $[\overline{i}, \overline{u(i)}]$ of $\overline{U}[i]$, by $\overline{U}_-[i]$ the arc $[\overline{u^{-1}(i)}, \overline{i}].$

Fix $g \in S_n$. Cutting tables at points (guests) $\overline{1}, \ldots, \overline{n}$, we get a finite collection of segments ${\{\overline{U}_+[i]\}}_{i\in I_n}$ and a countable number of non-cutted tables. The collection ${\{\overline{U}_-[j]\}}_{j\in I(n)}$ coincides with ${\{\overline{U}_+[i]\}}_{i\in I_n}$ up to a reordering.

To obtain ug, for each $i \in I_n$ we glue the segments \overline{U} -[j] and \overline{U} +[g(j)] identifying points $\overline{j} \in \overline{U}$ = $[j]$ and $\overline{g(j)} \in \overline{U}$ + $[g(j)]$ and putting the guest \overline{j} to the point of gluing. In this way, we get a family of new tables enumerated by disjoint cycles of $ug \in S_n$, and add the collection of non-cutted tables.

To obtain gu we repeat the same steps, but put the guest $\overline{g(j)}$ to the point of gluing. Notice that $gu = g(ug)g^{-1}$, so the restorants of ug and gu have the same tables and the same positions of all guests with numbers $> n$, the guests $\overline{1}, \ldots, \overline{n}$ differ by a permutation q .

REMARK. These rules are simply a rephrasing of a description of multiplication of a permutation defined as map and a permutation defined as a product of cycles. ⊠

1.7. Radon–Nykodim derivatives. Thus, we get an action of the group $S_{\infty} \times S_{\infty}$ on \mathfrak{S}^z given by $\mathfrak{u} \mapsto h_1^{-1} \mathfrak{u} h_2$.

Clearly, on the level of S_n we have

$$
\mu_n^z(h_1^{-1}uh_2) = z^{[h_1^{-1}gh_2]-[g]} \mu_n^z(u).
$$

For $\mathfrak{u} := \mathfrak{S}^z$ the operation of cutting and gluing described above involves only a finite number of tables, so the number

$$
\gamma(h_1, h_2; \mathfrak{u}) := \{ \text{number of tables of } h_1^{-1} \mathfrak{u} h_2 \} - \{ \text{number of tables of } \mathfrak{u} \} :=
$$

$$
:= [h^{-1} \Upsilon_{N}^{\infty}(g) h_2] - [\Upsilon_{N}^{\infty}(g)] \quad \text{for large } N
$$

is well-defined. The Radon-Nikodym derivative^{[3](#page-5-0)} of the transformation $\mathfrak{u} \mapsto h_1^{-1} \mathfrak{u} h_2$ is $z^{\gamma(h_1,h_2;\mathfrak{u})}$.

1.8. The inversion of virtual permutations. Clearly,

$$
\Upsilon_{n-1}^n(g^{-1}) = g^{-1}.
$$

So the inversion $g \mapsto g^{-1}$ defines a measure preserving map $\mathfrak{S}^z \to \mathfrak{S}^z$, we denote it by $\mathfrak{u} \mapsto \mathfrak{u}^{-1}$. This is simply a changing of orientations of all tables from clockwise to counterclockwise. Clearly

$$
(hu g)^{-1} = g^{-1} \mathfrak{u}^{-1} h^{-1}.
$$

1.9. The bisymmetric group. The *bisymmetric group* S is the subgroup in $\overline{S}_{\infty} \times \overline{S}_{\infty}$ consisting of pairs (g_1, g_2) such that $g_1 g_2^{-1} \in S_{\infty}$. Denote by $\mathbb{K} \subset \mathbb{S}$ the diagonal subgroup, i.e., the subgroup consisting of pairs (g, g) , where g ranges in \overline{S}_{∞} . So, $\mathbb{K} \simeq \overline{S}_{\infty}$. We define topology on S assuming that K is an open subgroup equipped with the natural topology. The homogeneous space \mathcal{S}/\mathbb{K} is a countable space with the discrete topology.

The bisymmetric group \Im *acts on* \mathfrak{S}^z . Namely, the subgroup $\mathbb K$ acts by permutations of guests, such permutations are measure-preserving maps. The subgroup $S_{\infty} \times S_{\infty}$ acts as above.

REMARK. 1) The group S was introduced by Olshanski [\[38\]](#page-34-2). It is a type I group, a classification of its irreducible unitary representations is known, see Olshanski [\[38\]](#page-34-2), Okounkov [\[37\]](#page-34-3), see also [\[21\]](#page-33-4). The most important family of irreducible representations are K-spherical representations, i.e., representations having a fixed vector with respect to K (it is automaticaly unique). In fact such representations were classified by Thoma [\[47\]](#page-34-4), 1964, see also [\[50\]](#page-34-5) and [\[38\]](#page-34-2).

2) Decompositions of quasiregular representations of $\mathbb S$ in $L^2(\mathfrak{S}^z)$ are known, see Kerov, Olshanski, Vershik [\[13\]](#page-33-5), Borodin, Olshanski [\[3\]](#page-32-2), [\[4\]](#page-32-3). One of informal aims of the present work is a search of additional possibilities for the analysis on the space $\mathfrak{S}^z.$

B. The bisymmetric group and its train.

1.10. Multiplication of double cosets. See Olshanski [\[38\]](#page-34-2), see also [\[31\]](#page-33-6). For $\alpha \in \mathbb{Z}_+$ we denote by \mathbb{K}_{α} the subgroup in K consisting of (g, g) such that $g \in \overline{S}_{\infty}$ fixes $1, \ldots, \alpha \in \mathbb{N}$. We set $\mathbb{K}_0 := \mathbb{K}$. Denote by

$$
K_{\alpha} := \mathbb{K}_{\alpha} \cap (S_{\infty} \times S_{\infty})
$$

the corresponding subgroup of finitely supported permutations.

 3 See a formal definition below in Subsect. [1.16,](#page-12-0) see also, e.g., [\[2\]](#page-32-4), 9.12.

FIGURE 1. To the definition of a product of double cosets.

For any α , β we consider the double coset space

$$
\mathcal{S}(\alpha,\beta)=\mathbb{K}_{\alpha}\backslash\mathbb{S}/\mathbb{K}_{\beta}.
$$

It is easy to see that each double coset $\mathbb{K}_{\alpha} \cdot g \cdot \mathbb{K}_{\beta}$ has a finitary representative $g' \in S_{\infty} \times S_{\infty}$. Moreover,

(1.9)
$$
\mathbb{K}_{\alpha} \backslash \mathbb{S} / \mathbb{K}_{\beta} \simeq K_{\alpha} \backslash (S_{\infty} \times S_{\infty}) / K_{\beta},
$$

all such quotient spaces are countable and the quotient topologies are discrete.

For any $\alpha, \beta, \gamma \in \mathbb{Z}_+$, there exists a natural multiplication

$$
\mathcal{S}(\alpha,\beta) \times \mathcal{S}(\beta,\gamma) \to \mathcal{S}(\alpha,\gamma)
$$

defined in the following way. For each $\beta \in \mathbb{Z}_+$ we consider the following sequence $\theta^{\beta}[j] \in \mathbb{K}_{\beta}$ defined by

$$
\theta^{\beta}[j](k) = \begin{cases} \alpha & \text{if } k \leqslant \beta; \\ \alpha + j & \text{if } \beta < k \leqslant \beta + j; \\ \alpha - j & \text{if } \beta + j < k \leqslant \beta + 2j; \\ \alpha & \text{if } k > \beta + 2j. \end{cases}
$$

see Fig. [1.](#page-6-0)

Now we take double cosets $\mathfrak{g} = \mathbb{K}_{\alpha} g \mathbb{K}_{\beta}$, $\mathfrak{h} = \mathbb{K}_{\beta} h \mathbb{K}_{\gamma}$, without loss of generality we can think that $g, h \in S_{\infty} \times S_{\infty}$. Consider the sequence

$$
\mathbb{K}_{\alpha} \cdot g \theta^{\beta}[j] h \cdot \mathbb{K}_{\gamma} \in \mathbb{K}_{\alpha} \backslash \mathbb{S} / \mathbb{K}_{\beta}.
$$

It is easy to show that this sequence is eventually constant. We define a product $\mathfrak{g} \circ \mathfrak{h} \in \mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\gamma}$ as the limit (i.e., the stable value) of this sequence. It can be shown that the result does not depend on a choice of representatives $g, h \in S_{\infty} \times S_{\infty}$ and this operation is associative. So we get a category S (*the train of* S) whose objects are nonnegative integers and morphisms $\beta \to \alpha$ are double cosets,

Ob(
$$
\mathcal{S}
$$
) := \mathbb{Z}_+ , Mor _{\mathcal{S}} (β , α) := $\mathcal{S}(\alpha, \beta)$ = $\mathbb{K}_{\alpha} \backslash \mathbb{S} / \mathbb{K}_{\beta}$.

We also define an involution $\mathfrak{g} \mapsto \mathfrak{g}^*$ on the category S. Namely, a map $g \mapsto g^{-1}$ determines bijections $\mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\beta} \to \mathbb{K}_{\beta} \backslash \mathbb{S}/\mathbb{K}_{\alpha}$. Clearly,

$$
(\mathfrak{g}\circ\mathfrak{h})^*=\mathfrak{h}^*\circ\mathfrak{g}^*.
$$

1.11. Chips. Recall Olshanski's description [\[38\]](#page-34-2) of the product of double cosets. We represent an element of the group $S_{\infty} \times S_{\infty}$ as a diagram of the type drawn on Fig. [2.](#page-7-0)a. On Fig. [2.](#page-7-0)b, [2.](#page-7-0)c we define combinatorial data corresponding to a double coset. We get a diagram (a chip^4 chip^4) of the form shown on the Fig. [2.](#page-7-0)c.

⁴We use the term proposed by Kerov [\[11\]](#page-33-7) for elements of the Brauer semigroups. Recall that the Schur–Weyl duality between $GL(N)$ and S_n has a counterpart [\[5\]](#page-32-5), 1936, for orthogonal groups $O(N)$ and symplectic groups $Sp(2N)$, dual objects are certain semigroups of chips (with horisontal arcs but without crosses). Brauer's approach also gives a similar statement for tensors over $GL(N)$

a) An element of S . The symbols l and r in subscripts are abbreviations of 'left' and 'right'. Symbols $+$ and $-$ in superscripts correspond to the top and the bottom of the diagram. The left and the right parts of the diagram are symmetric one to another except a finite number of arcs.

b) On Figure $\beta = 3$, $\alpha = 4$. We add 'horisontal' arcs with crosses.

c) We consider compound arcs on the previous diagram up to isotopies with fixed ends. Also, we get a countable family of circles with two crosses and forget them.

FIGURE 2. The construction of a chip from an element of \mathbb{S} .

It is convenient to think that *each arc has a 'length', which is defined as the half of the number of crosses.*

For figures in our printed text it is more convenient to draw crosses.

We get diagrams of the following type. On the top we have black circles labeled by β_l^+ , ..., 1_l^+ to the left of the dashed line, and 1_r^+ , ..., β_r^+ to the right of the

with covariant and contravariant components, in this case diagrams have a separating dashed line (it separates 'co-' and 'contra-') and horizontal arcs corresponding to convolutions. On the other hand, according Wasserman [\[51\]](#page-34-6) and Olshanski [\[38\]](#page-34-2), representations of S can be realized in certain tensors of infinite order, their decomposition is controlled by certain 'dual' compact groups.

FIGURE 3. The multiplication of chips. We identify corresponding circles of the bottom of the first diagram and on the top of the second diagram.

dashed line. On the bottom we have black circles labeled by $\alpha_{l}^{-},\,\ldots,\,1_{l}^{-}$ on the left and $1^-_r, \ldots, \alpha^-_r$ on the right. Each circle is an end of an arc. There are 3 following types of arcs:

1) arcs $i_r^+ \downarrow i_r^- [\varphi]$ (resp., $i_l^+ \downarrow i_l^- [\varphi]$) of integer length $\varphi \geq 0$ from the top to the bottom located in the right (resp., left) hand side of the diagram;

2) arcs $i_l^+ \frown j_r^+ [\psi]$ (resp., $i_l^- \smile j_r^- [\psi]$) of length $\psi \in (1/2 + \mathbb{Z}_+)$ from left to right on the top (resp., bottom) of the diagram;

3) circles $\bigcirc[k]$ of integer length $k > 0$.

We remove all cycles of length 1 (any diagram obtained by our procedure contains an infinite number of cycles \bigcirc [1], this collection contains no information). So in all cases we get finite objects.

It is easy to show that such diagrams are in one-to-one correspondence with elements of $\mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\beta}$. Multiplication of diagrams is a gluing, see Fig. [3.](#page-8-0) The involution corresponds to the replacement of the top and of the bottom.

REMARK. An element of the semigroup $S(0,0)$ is a collection of cycles $\bigcirc [k_i]$. The product is a union of such collections. In particular, this semigroup is Abelian. ⊠

1.12. The multiplicativity theorem. Let ρ be a unitary representation of the group S in a Hilbert space^{[5](#page-8-1)} H. Denote by H_{α} the subspace of all \mathbb{K}_{α} -fixed vectors, by P_{α} the operator of orthogonal projection to H_{α} . For $g \in \mathbb{S}$ we define the operator

$$
\widetilde{\rho}_{\alpha,\beta}(g):H_{\beta}\to H_{\alpha}
$$

by

$$
\widetilde{\rho}_{\alpha,\beta}(g) := P_\alpha \,\rho(g)\Big|_{H_\beta}.
$$

It is easy to see that $\tilde{\rho}_{\alpha,\beta}(g)$ depends only on the double coset $\mathfrak{g} = \mathbb{K}_{\alpha} g \mathbb{K}_{\beta}$. The following *multiplicativity theorem* holds (see [\[38\]](#page-34-2), [\[31\]](#page-33-6)).

Theorem 1.1. *For any* α , β , $\gamma \in \mathbb{Z}_+$ *and* $\mathfrak{g} \in \mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\beta}$, $\mathfrak{h} \in \mathbb{K}_{\beta} \backslash \mathbb{S}/\mathbb{K}_{\gamma}$ *we have*

(1.10)
$$
\widetilde{\rho}_{\alpha,\beta}(\mathfrak{g})\widetilde{\rho}_{\beta,\gamma}(\mathfrak{h})=\widetilde{\rho}_{\alpha,\gamma}(\mathfrak{g}\circ\mathfrak{h}).
$$

⁵We assume that a Hilbert space is separable by definition.

(1.11)
$$
\widetilde{\rho}(\mathfrak{g}^*) = \widetilde{\rho}(\mathfrak{g})^*
$$

So, for any unitary representation of the group S we get a functor from the category S to the category of Hilbert spaces and bounded operators. This reduces investigation of representations of the group S to representations of the category S.

In particular, this theorem can be applied to quasiregular representations of the bisymmetric group \Im in $L^2(\mathfrak{S}^z)$. There arises a question: *is it possible to describe operators corresponding to chips explicitly?* Below we reformulate this question in terms of polymorphisms.

1.13. Remark. Infinite chips and weak closures of the group S in unitary representations. Denote by $\mathcal{C}(H)$ the semigroup of all operators in a Hilbert space H with norm ≤ 1 (we call such operators *contractive*). It easy to see that $\mathfrak{C}(H)$ is compact and metrizable with respect to the weak operator topology, the multiplication in $\mathcal{C}(H)$ is separately continuous (see, e.g., [\[24\]](#page-33-2), Sect. I.1).

Let G be a group, ρ be its unitary representation in a Hilbert space H. Consider the subset $\rho(G)$ in the space of all operators in H and consider its closure $\rho(G)$ in the semigroup of all contractive operators in the weak operator topology. Clearly, we get a compact separately continuous semigroup, see [\[40\]](#page-34-7), [\[24\]](#page-33-2), Sect. I.1.

A description of such semigroup can be a nontrivial problem even for \mathbb{Z} , i.e., for closure of the set of powers of a given unitary operator U . For a comeagre set of unitary operators this semigroup is the unit ball in $L^{\infty}(S^1, \mu)$, where μ is the spectral measure of U , see [\[20\]](#page-33-8).

For reasonable representations of Lie groups (over real and over local fields) the question usually leads to one-point compactifications, see [\[9\]](#page-33-9). Other types of locally compact groups and discrete non-Abelian groups are not well-understood. For infinite-dimensional groups this question leads to handable and unexpected algebraic structures as it was observed by Olshanski [\[40\]](#page-34-7). Let us describe such compactification for the group S.

The sequence $\theta_{\beta}[j]$ defined above converges to the projector P_{β} in the weak operator topology, see, e.g., [\[24\]](#page-33-2), Theorem VIII.1.4. Therefore the following operators

$$
\widehat{\rho}_{\alpha,\beta}(g) = P_\alpha \,\rho(g) \, P_\beta : H \to H
$$

are contained in $\rho(\mathbb{S})$. It is easy to see that these operators depend only on double coset \boldsymbol{g} containing q and have the following block structure

$$
\widehat{\rho}(\mathfrak{g})=\begin{pmatrix}\widetilde{\rho}(\mathfrak{g})&0\\0&0\end{pmatrix}:\quad H_\beta\oplus H_\beta^\perp\to H_\alpha\oplus H_\alpha^\perp.
$$

Clearly, we have

$$
\widehat{\rho}_{\alpha,\beta}(\mathfrak{g})\,\widehat{\rho}_{\beta,\gamma}(\mathfrak{h})=\widehat{\rho}_{\alpha,\gamma}(\mathfrak{g}\circ\mathfrak{h}),
$$

this identity is a rephrasing of (1.10) . So chips act in the space H itself.

More generally, consider diagrams as on Fig. [2.](#page-7-0)c, where circles on the top and the bottom (from the left and from the right) are enumerated by N. We require the following conditions of finiteness of such diagrams:

- a) the number of cycles $\bigcirc[k]$ is finite (recall that $k \geq 2$);
- b) the number of 'vertical' arcs $i_l^+ \downarrow j_l^- [m], p_r^+ \downarrow q_r^- [m]$ of lengths $m > 0$ is finite;
- c) the number of horizontal arcs $i_l^+ \sim j_r^+[1/2 + k]$, $i_l^- \sim j_r^-[1/2 + k]$ of lengths $1/2 + k > 1/2$ is finite.

d) a diaram is symmetric with respect to the dashed line upto a finite family of arcs.

Denote the semigroup of such chips by $S(\infty, \infty)$. The group of its invertible elements is S.

REMARK. The embedding $\mathcal{S}(n,n) \to \mathcal{S}(\infty,\infty)$ is given by adding arcs $i_l^+ \sim$ $i_r^+[1/2], i_l^- \smile i_r^- [1/2]$ for all $i > n$.

REMARK. The semigroup $\mathcal{S}(\infty,\infty)$ has a center. It consists of chips whose arcs have the form $j_l^+ \downarrow j_l^-$, $j_r^+ \downarrow j_r^-$ for all $j \in \mathbb{N}$ and $\bigcirc [k_j]$, where $k_j \geq 2$. The center is isomorphic to $\mathcal{S}(0,0)$.

For a unitary representation ρ of S denote by Π_{+} the operator of orthogonal projection to the subspace of vectors fixed by the whole group S, by Π[−] the operator of projection to the subspace of vectors v satisfying

$$
\rho(h_1, h_2)v = \text{sgn}(h_1 h_2^{-1})v
$$
, where $(h_1, h_2) \in \mathbb{S}$.

Proposition 1.2. a) *Any unitary representation* ρ *of* S *extends to a representation of the semigroup* $S(\infty, \infty)$ *.*

b) *The closure* $\overline{\rho(S)}$ *consists of the image of* $S(\infty, \infty)$ *and operators* $\Pi_+ + \Pi_$ $and \Pi_+ - \Pi_-.$

REMARKS. a) Representations of the semigroup $\mathcal{S}(\infty,\infty)$ can be constructed in the following way. We split N into a disjoint union of two countable sets $\mathbb{N} = L \sqcup M$. Denote by $\mathbb{K}(\infty)$ the subgroup of K consisting of elements fixing all elements of L. Then $\mathcal{S}(\infty, \infty) \simeq \mathbb{K}(\infty) \backslash \mathbb{S} / \mathbb{K}(\infty)$ and the multiplicativity theorem remains valid with the same proof. The group of invertible elements of $\mathcal{S}(\infty,\infty)$ is isomorphic to $S(\infty, \infty)$. It can be shown that the representation of S in the subspace K(∞)-fixed vectors is equivalent to the initial representation of S.

b) The second statement of the theorem is a relatively easy corollary of results of Olshanski [\[38\]](#page-34-2) and Okounkov [\[37\]](#page-34-3), we omit its proof.

c) We can define sets $S(\infty, m)$ and $S(n, \infty)$ in the same way, and so get a completed category $\overline{\mathcal{S}}$,

$$
\mathrm{Ob}(\mathcal{S}) := \mathbb{Z}_+ \sqcup \infty.
$$

Any unitary representation of S admits a unique extension to a representation of this category. ⊠

C. Polymorphisms

It is natural to reformulate the question about operators in \mathfrak{S}^z corresponding to chips in terms of polymorphisms of the space \mathfrak{S}^z .

1.14. Bistochastic kernels or measure preserving polymorphisms. Recall that a *Lebesgue space* is a measure space, which is equivalent to a union of an interval of a line and a finite, countable or empty set of (atomic) points with nonzero measures (see, $[44]$, $[2]$, §9.4). A measure μ is *probabilistic* if the measure of the whole space is 1. A measure is *continuous* if it has no atomic points. Clearly, the spaces \mathfrak{S}^z of virtual permutations are Lebesgue spaces with continuous measures.

Let (M, μ) be a Lebesgue space with a continuous measure μ . Denote by Ams (M) the group of bijective a.s. measure preserving transformations of M . The group Ams (M) acts in $L^2(M,\mu)$ by unitary operators

$$
(1.12) \t\t T(g)f(m) = f(mg).
$$

Let (M, μ) , (N, ν) be probabilistic Lebesgue measure spaces. A *bistochastic kernel* or a *measure preseving polymorphism*^{[6](#page-11-0)} $\mathfrak{q}: M \twoheadrightarrow N$ is a measure \mathfrak{q} on $M \times N$ such that the pushforward^{[7](#page-11-1)} of q under the projection $M \times N \to M$ coincides with μ and the pushforward under the projection $M \times N \to N$ coincides with ν .

EXAMPLE. Let $g \in \text{Ans}(M)$. Consider the map $M \to M \times M$ given by $m \mapsto$ (m, mg) . Then the pushforward of M under this map is a measure preserving polymorphism. In particular, for the identical transformation $g = 1$ we get an *identical polymorphism.* ⊠

A measure preserving polymorphism $\mathfrak{q}: M \to N$ can be regarded as a 'spreading map' sending points to measures. Namely, consider a partition

$$
M\times N=\coprod_{m\in M}m\times N
$$

and conditional measures^{[8](#page-11-2)} $\varkappa_m(n)$ on N defined for almost all m from the condition

$$
\mathfrak{q}(S) = \int_M \varkappa_m \big((m \times N) \cap S \big) \, d\mu_m(n) \quad \text{for any measurable } S \subset M \times N.
$$

Then *we can regard* q *as a 'spreading map' sending points* $m \in M$ *to measures* $\varkappa_m(n)$ on N.

A *product* of measure preserving polymorphisms corresponds to a double spreading. Namely, let q be a polymorphism $M \to N$ and p be a polymorphism $N \to K$. Let $\varkappa_m(n)$ be the system of conditional measures corresponding to q, and $\pi_n(k)$ be the system of conditional measures corresponding to p. Then the system $\rho_m(k)$ of conditional measures corresponding to the product $\mathfrak{r} = \mathfrak{p} \otimes \mathfrak{q}$ is defined by

$$
\rho_m(k) = \int_N \pi_n(k) \, d\mathcal{K}_m(n).
$$

The map $q \mapsto T(q)$, see [\(1.12\)](#page-11-3), can be extended from Ams(M) to polymorphisms $\mathfrak{q}: M \to N$. Namely, the *operator* $T(\mathfrak{q}): L^2(N) \to L^2(M)$ is given by

$$
T(\mathfrak{q})f(m) = \int_N f(n) \, d\mu_m(n).
$$

Then for $\mathfrak{q}: M \to N$, $\mathfrak{p}: N \to K$ we have

$$
T(\mathfrak{p}\circledast\mathfrak{q})=T(\mathfrak{p})\,T(\mathfrak{q}).
$$

We say that a sequence $\mathfrak{q}^{(j)} : M \to N$ *converges* to **q** if for any measurable $A \subset M, B \subset N$ the sequence $\mathfrak{q}^{(j)}(A \times B)$ converges to $\mathfrak{q}(A \times B)$. This convergence is equivalent to the weak operator convergence $T(\mathfrak{q}^{(j)}) \to T(\mathfrak{q})$.

 6 The notion rises to E. Hopf [\[8\]](#page-33-10), see also [\[36\]](#page-34-9), [\[17\]](#page-33-11), [\[49\]](#page-34-10), such objects are widely used in ergodic theory, see e.g., [\[15\]](#page-33-12), [\[10\]](#page-33-13). We use the term 'polymorphism' proposed in [\[49\]](#page-34-10).

TLet (M, μ) be a measure space, let S be a set, $\psi : M \to S$ a map. We say that a subset $C \subset S$ is measurable if $\psi^{-1}(C)$ is measurable, the pushforward σ of the measure μ is defined by $\sigma(C) = \mu(\psi^{-1}(C)).$

 8 See [\[44\]](#page-34-8), [\[2\]](#page-32-4), Chapter 10.

The product of polymorphisms is separately continuous; the group $\text{Ans}(M)$ is dense in the semigroup of polymorphisms $M \rightarrow M$.

We also have the *involution* $q \mapsto q^*$ on polymorphisms, it sends a measure q on $M \times N$ to the same measure on $N \times M$. We have $T(\mathfrak{q}^*) = T(\mathfrak{q})^*$.

1.15. Quotient spaces and the corresponding measure preserving poly**morphisms.** Let M be a Lebesgue measure space, let $M = \bigcup_{h \in H} S_h$ be a measurable partition, let $\sigma_h(m)$ be the conditional measures on sets S_h (see [\[44\]](#page-34-8)). Consider the quotient space H with the induced measure $\eta(h)$. Then we have the following polymorphisms (see [\[30\]](#page-33-14), Subsect. 3.10):

a) We have the projection $\pi : M \to H$ sending $m \in M$ to the element S_h containing m. We consider the map $M \to M \times H$ given by $m \mapsto (m, \pi(m)),$ we denote by $\mathfrak x$ the image of μ under this map. Then $\mathfrak x$ is a measure-preserving polymorphism $M \rightarrow H$. The operator

$$
T(\mathfrak{x})f(m) = f(\pi(m))
$$

is a canonical isometric embedding $L^2(H) \to L^2(M)$. The image consists of L^2 functions that are constant on sets S_h .

b) We also have a polymorphism \mathfrak{r}^* , the corresponding operator $T(\mathfrak{r}^*) : L^2(M) \to$ $L^2(H)$ is the operator of *conditional expectation*

$$
T(\mathfrak{x}^*) f(h) = \mathcal{E} f(h) := \int_{S_h} f(m) d\sigma_h(m).
$$

c) Notice that $\mathfrak{x} \otimes \mathfrak{x}^*$ is the identical polymorphism $H \to H$.

d) For the polymorphism $\mathfrak{x}^* \otimes \mathfrak{x} : L^2(M) \to L^2(M)$ the corresponding operator is the orthogonal projector to the subspace $L^2(H)$.

1.16. General polymorhisms. Let M be a Lebesgue space with a continuous measure. Recall that a map $g : M \to M$ *leaves a measure* μ *quasiinvariant*^{[9](#page-12-1)} if it is bijective almost sure and there is a function $g'(m)$ (the *Radon–Nikodym derivative*, see, e.g., [\[2\]](#page-32-4), 9.12) such that for any measurable $A \subset M$ we have

$$
\mu(Ag) = \int_A g'(m) \, d\mu(m),
$$

the $g'(m)$ is a natural extension of Jacobians in classical analysis. Denote by Gms(M) the group of transformations of M leaving the measure μ quasiinvariant.

Denote by $\Pi \subset \mathbb{C}$ the strip

$$
(1.13) \t\t 0 \leq \operatorname{Im} \lambda \leq 1.
$$

For $r + is \in \Pi$ we consider the following transformations of the space of functions on M:

(1.14)
$$
T_{r+is}(g) f(m) = f(mg) g'(m)^{r+is},
$$

Then $T_{r+is}(g)$ is an isometric operator $L^{1/r}(M) \to L^{1/r}(M)$.

REMARK. The main topic of our interest is the case $r = 1/2$, when we get a unitary operators $L^2 \to L^2$. However, operators in spaces L^p are used in our argumentation in Theorem [2.1](#page-17-0) below. ⊠

⁹This is also equivalent to the condition: g is bijective a.s. and both g and g^{-1} sent sets of zero measure to sets of zeto measure.

1. The pushforward of $\mathfrak p$ under the projection $M \times N \times \mathbb{R}^{\circ} \to M$ is the measure μ .

2. The pushforward of the measure $t \cdot \mathfrak{p}$ to N coincides with the measure ν .

Denote by $Pol(M, N)$ the space of all polymorphisms $M \rightarrow N$. See [\[23\]](#page-33-15), [\[30\]](#page-33-14).

Example. Measure preserving polymorphisms from Subsect. [1.14](#page-10-0) are polymorphisms, they are supported by the set $M \times N \times \{1\} \subset M \times N \times \mathbb{R}^{\circ}$. ⊠

EXAMPLE. Let $q \in \text{Gms}(M,\mu)$. Consider the map $M \to M \times M \times \mathbb{R}^{\circ}$ defined by

$$
m \mapsto (m, q(m), q'(m)).
$$

Then the pushforward of the measure μ under this map is a polymorphism $M \rightarrow M$. ⊠

EXAMPLE. Denote by $\mathcal{M}^{\nabla}(\mathbb{R}^{\circ})$ the set of all positive finite Borel measures σ on \mathbb{R}° such that $t \cdot \sigma$ also is finite. Clearly, any measurable function $(m, n) \mapsto s_{m,n}$ from $M \times N$ to \mathcal{M}^{∇} determines a certain measure \mathfrak{s} on $M \times N \times \mathbb{R}^{\circ}$. Namely, for measurable sets $A \subset M$, $B \subset N$, $C \subset \mathbb{R}^{\circ}$ we set

$$
\mathfrak{s}(A \times B \times C) = \int_{A \times B} s_{m,n}(C) \, d\mu(m) \, d\nu(n).
$$

We also must assume that

$$
\forall m: \int_N s_{m,n}(\mathbb{R}^\circ) d\nu(n) = 1, \qquad \forall n: \int_M \int_{\mathbb{R}^\circ} t \, ds_{m,n}(t) d\mu(m) = 1,
$$

under these conditions $\mathfrak s$ is contained in Pol(M, N). We say that a polymorphism $\mathfrak s$ obtained in such way is *continuous*. Notice that polymorphisms from the previous example are not continuous. ⊠

1.17. Convergence of polymorphisms. Let $\mathfrak{p}_i \in Pol(M, N)$. Consider measurable subsets $A \subset M$, $B \subset N$. Restrict \mathfrak{p} to $A \times B \times \mathbb{R}^{\circ}$ and take its pushforward under the map $A \times B \times \mathbb{R}^{\circ} \to \mathbb{R}^{\circ}$. Denote the resulting measure on \mathbb{R}° by $\mathfrak{p}[A \times B]$. We say that a sequence $\mathfrak{p}_j \in Pol(M, N)$ converges to \mathfrak{p} , if for any measurable subsets $A \subset M$, $B \subset N$ we have weak convergences of measures

$$
\mathfrak{p}_j[A \times B] \to \mathfrak{p}[A \times B], \qquad t \cdot \mathfrak{p}_j[A \times B] \to t \cdot \mathfrak{p}[A \times B].
$$

It is easy to show (see [\[30\]](#page-33-14), Theorem 5.3) that *the group* Gms(M) *is dense in* $Pol(M, M)$. Continuous polymorphisms $M \to N$ are dense in $Pol(M, N)$.

1.18. Products of polymorphisms. There is an obvious way to multiply continuous polymorphisms. Let $\mathfrak{u} \in \text{Pol}(M, N)$, $\mathfrak{v} \in \text{Pol}(N, K)$ be determined by functions $(m, n) \mapsto u_{m,n}, (n, k) \mapsto v_{n,k}$. Then their product

$$
\mathfrak{w} = \mathfrak{v} \circledast \mathfrak{u} \in \mathrm{Pol}(M, K)
$$

is determined by the function $M \times K \to \mathcal{M}^{\nabla}$ given by

$$
w_{m,k} = \int_N u_{m,n} * v_{n,k} d\nu(n),
$$

where $*$ denotes the convolution on \mathbb{R}° .

The multiplication ⊛ *extends to a separately continuous map*

 $Pol(M, N) \times Pol(N, K) \rightarrow Pol(M, K),$

and moreover this operation is associative (see [\[30\]](#page-33-14), Theorem 5.5, Theorem 5.9). So we get a category Pol whose objects are Lebesgue probabilistic measure spaces and morphisms are polymorphisms.

For different ways to define the multiplication of polymorphisms, see [\[30\]](#page-33-14), however it seems that definitions by continuity (by continuous extension from continuous polymorphisms or from the group $Gms(M)$ are more convenient.

1.19. Polymorphisms as spreading maps. Informally, *polymorphisms are 'maps'* $M \to N$, which spread points of M along N and the Radon–Nikodym deriv*ative also is spread.* Namely, let $q \in Pol(M, N)$. Consider a partition

$$
M \times N \times \mathbb{R}^{\circ} = \coprod_{m \in M} m \times N \times \mathbb{R}^{\circ}
$$

and conditional measures $\varkappa_m(n, t)$ on fibers of the partition. We say that *measures* $\varkappa_m(n,t)$ *are spreaded images of points* m. Below in Section 3 we describe polymorphisms of \mathfrak{S}^z in terms of such conditional measures.

1.20. Involution. Define an involution in the category Pol. Consider the map $(m, n, t) \mapsto (n, m, t^{-1})$. For $\mathfrak{p} \in Pol(M, N)$ we consider its pushforward \mathfrak{p}' under this map and define $\mathfrak{p}^* \in Pol(N, M)$ as the measure $t \cdot \mathfrak{p}'$. Then

$$
(\mathfrak{q}\circledast\mathfrak{p})^{\star}=\mathfrak{p}^{\star}\circledast\mathfrak{q}^{\star}.
$$

1.21. Mellin–Markov transforms of polymorphisms. For each $r + is \in \Pi$ the map $g \mapsto T_{r+is}(g)$ extends to the category of polymorphisms.

Let $\mathfrak{p} \in \text{Pol}(M, N)$. The *Mellin–Markov transform of* $\mathfrak{q}: M \to N$ is a function defined in the strip $r + is \in \Pi$ taking values in operators

$$
T_{r+is}(\mathfrak{q}):L^{1/r}(N)\to L^{1/r}(M)
$$

and determined from the equality

(1.15)
$$
\int_{M} \varphi(m) T_{r+is}(\mathfrak{q}) \psi(m) d\mu(m) = \int_{M \times N \times \mathbb{R}^{\circ}} \varphi(m) \psi(n) t^{r+is} d\mathfrak{q}(m,n,t)
$$

for any $\psi \in L^{1/r}(N)$, $\varphi \in L^{1/(1-r)(M)}$.

The operators $T_{r+is}(\mathfrak{q}) : L^{1/r}(N) \to L^{1/r}(M)$ are bounded and for any $\mathfrak{p} \in$ Pol (M, N) , $\mathfrak{q} \in Pol(M, K)$ we have (see [\[30\]](#page-33-14), Theorem 6.14)

$$
T_{r+is}(\mathfrak{q}) T_{r+is}(\mathfrak{p}) = T_{r+is}(\mathfrak{q} \circledast \mathfrak{p}).
$$

It is more transparent to determine q in terms of conditional measures $\varkappa_n(m, t)$ as in Subsect. [1.19.](#page-14-0) Then

$$
T_{r+is}(\mathfrak{q})f(m)=\int_{N\times\mathbb{R}^\circ}f(n)\,t^{r+is}\,d\varkappa_m(n,t).
$$

1.22. Closures of actions of groups on measure spaces. For a group G acting on a measure space M we get a question about the closure of G in the semigroup Pol(M, M) of polymorphisms. This question for $G = \mathbb{Z}$ acting by measure preserving transformations was a subject of numerous works, see e.g., [\[15\]](#page-33-12), [\[10\]](#page-33-13), [\[46\]](#page-34-11), [\[18\]](#page-33-16), a related topic is intertwiners (joinings) of such actions in class of measure preserving polymorphisms.

For infinite-dimensional groups the first problem of this kind was solved by Nelson [\[22\]](#page-33-17), who examined the action of the infinite-dimensional orthogonal group on the infinite-dimensional space with a Gaussian measure (the resulting semigroup is the semigroup of all contractive operators). For the natural group of symmetries of Poisson measure and for the action of the restricted group GL on a space with a Gaussian measure the problems were discussed in [\[25\]](#page-33-18), [\[28\]](#page-33-19). They are unexpectedly non-trivial and lead to formulas, which are at least unusual. One simple case (measures on the space of infinite Hermitian matrices invariant with respect to unitary groups) was examined in [\[33\]](#page-33-20).

In this paper we consider the action of the bisymmetric group $\mathbb S$ on spaces $\mathfrak S^z$ of virtual permutations.

1.23. The purpose of the paper. First, we show that the categories S and \overline{S} of chips act on the space of virtual permutations by polymorphisms. The category of chips is a representative of a wide zoo of train constructions, in Section 2 we prove a general statement in the following spirit: certain actions of infinite-dimensional group on measure spaces generate actions of their trains by polymorphisms.

In Section 3 we write a formula for the action of the semigroup $\mathcal{S}(0,0)$ on the space of \mathfrak{T}^z of restaurants. In fact, for a given collection $\{\bigcirc[k_j]\}$ of cycles we write spreaded images of points as sums over a certain sets of dessins d'enfant (or equivalently of checker triangulated surfaces), summands are multiple convolutions of Dirichlet distributions (Theorem [3.5\)](#page-23-0).

In Section 4 we get some further statements in this spirit. In Theorem [4.1](#page-31-0) we get formulas for the action of the center $\mathcal{S}(0,0)$ of the semigroup $\mathfrak{S}(\infty,\infty)$ on \mathfrak{S}^z .

In Theorem [3.5](#page-23-0) we write a similar formula for arbitrary chips, whose left sides are trivial (see Fig. [13\)](#page-29-0). Formula involves several types of parameters of combinatorial nature and is longer than for the action of $S(0,0)$ This allows to write explicit formulas in the same spirit for generators of the category S (we omit this). However, the author did not succeed to obtain a general formula for arbitrary chips.

2. TRAINS OF (G, K) -PAIRS AND POLYMORPHISMS.

Train constructions (multiplication of double cosets) and the multiplicativity theorem [1.1](#page-8-3) are relatively usual phenomena for infinite-dimensional groups, including infinite symmetric groups [\[38\]](#page-34-2), [\[29\]](#page-33-21), [\[31\]](#page-33-6), classical groups over reals [?], [\[24\]](#page-33-2), [\[26\]](#page-33-22) (and also over finite fields $[40]$, $[35]$ and p-adic fields $[34]$, $[32]$) groups of transformations of measure spaces [\[23\]](#page-33-15), [\[24\]](#page-33-2) and some exotic cases.

For infinite-dimensional group a lot of quasiinvariant actions (i.e., embeddings to the group $Gms(\cdot)$ is known. In this section, we wish to show that under some conditions such actions automatically generate actions of corresponding train categories by polymorphisms. We prove Theorem [2.1,](#page-17-0) which apparently is sufficient to justify this claim for known zoo of train constructions. Also, the proof is simple, short, and can be easily repeated independently in any explicit case.

We do not pretend to any wider purpose.

2.1. Operators corresponding to double cosets. Let G be a separable topological group, let K be a closed subgroup. Let ρ be a unitary representation of G in a separable Hilbert space H. Denote by $H(K)$ the subspace consisting of vectors fixed by elements of K, by $P(K)$ the orthogonal projection to $H(K)$. For two subgroups K, L and $q \in G$ consider the operator

$$
\widetilde{\rho}_{(K,L)}(g):=P(K)\rho(g)\big|_{H(L)}:\,H(L)\to H(K).
$$

It is easy to verify that the operator-valued function $\tilde{\rho}_{(K,L)}(g)$ is constant on double cosets KgL , so it is a function on $K\backslash G/L$. We define a more coarse equivalence relation on G ,

(2.1)
$$
g \sim g' \quad \text{if} \quad \widetilde{\rho}_{(K,L)}(g) = \widetilde{\rho}_{(K,L)}(g') \quad \text{for all } \rho.
$$

Denote an equivalence class containing g by $[KgL]$, denote the space of equivalence classes by $|K\backslash G/L|$.

2.2. (G, K) -pairs. Now let G be a separable topological group, let K be a subgroup. Let ${K_{\alpha}}_{\alpha\in\mathcal{A}}$ be a family of subgroups in K. Let the set A be partially ordered, and for any $\alpha, \alpha' \in A$ there is $\nu \in A$ such that $\nu \succcurlyeq \alpha, \nu \succcurlyeq \alpha'$. Assume also that $\alpha \succcurlyeq \alpha'$ implies $K_{\alpha'} \supseteq K_{\alpha}$.

A (G, K) -pair is a group and a family of subgroups ${K_{\alpha}}_{\alpha \in A}$ as above satisfying the following conditions:

A. For each α , β , $\gamma \in A$, for each $\mathfrak{g} \in [K_{\alpha} \backslash G / K_{\beta}]$, $\mathfrak{h} \in [K_{\beta} \backslash G / K_{\gamma}]$ there is $\mathfrak{p} \in [K_{\alpha} \backslash G / K_{\gamma}]$ *such that for each unitary representation* ρ *of* G *in a Hilbert space* H *we have*

(2.2)
$$
\widetilde{\rho}_{(K_{\alpha}, K_{\beta})}(\mathfrak{g}) \ \widetilde{\rho}_{(K_{\beta}, K_{\gamma})}(\mathfrak{h}) = \widetilde{\rho}_{(K_{\beta}, K_{\gamma})}(\mathfrak{p}).
$$

B. For each unitary representation, the subspace $\cup_{\alpha \in A} H(K_{\alpha})$ is dense in H. So we get an operation

$$
[K_{\alpha}\backslash G/K_{\beta}] \times [K_{\beta}\backslash G/K_{\gamma}] \rightarrow [K_{\alpha}\backslash G/K_{\gamma}].
$$

Denote it by $\mathfrak{p} = \mathfrak{g} \circ \mathfrak{h}$ or $\mathfrak{p} = g \circ h$. Since the product of linear operators is associative, this operation is also associative. So we get a category – *the train* $\mathcal{T} = \mathcal{T}(G, K)$ of (G, K) . Its objects are enumerated by the set A, morphisms $\beta \to \alpha$ are reduced double cosets $[K_{\alpha}\backslash G/K_{\beta}].$

This category is equipped with the *involution* $\mathfrak{g} \mapsto \mathfrak{g}^*$, which is the map

$$
[K_{\alpha}\backslash G/K_{\beta}] \to [K_{\beta}\backslash G/K_{\alpha}]
$$

induced by the inversion in $G, g \mapsto g^{-1}$.

To simplify notations, denote

$$
H[\alpha] := H(K_{\alpha}), \quad P[\alpha] := P(K_{\alpha}), \quad \widetilde{\rho}_{\alpha,\beta} := \widetilde{\rho}_{(K_{\alpha},K_{\beta})}.
$$

2.3. Actions of (G, K) -pairs by polymorphisms. Consider a (G, K) -pair. Let G act on a Lebesgue probabilistic measure space (M, μ) by transformations leaving the measure quasiinvariant, let K act by measure preserving transformations. In such a case we say that we have an *action of a* (G, K)*-pair on a measure space.*

For each K_{α} consider the σ -algebra $\Sigma[\alpha]$ of all K_{α} -invariant sets. Namely, a set $B \subset M$ is contained in $\Sigma[\alpha]$ if it is measurable and for any $g \in G$ the symmetric sum $B \triangle Bg$ has measure 0; equivalently, the indicator function of B is a K_{α} -fixed element of $L^2(M,\mu)$. This sigma-algebra $\Sigma[\alpha]$ determines a measurable partition^{[10](#page-16-0)}

¹⁰Recall that a partition $M = \bigcup_{h \in H} S_h$ of M is *measurable* if there is a countable family of measurable subsets A_j , which are unions of elements of partitions, such that for each distinct elements S_h , $S_{h'}$ there is A_j such that $S_h \subset A$, $S_{h'} \subset M \setminus A$ or vise versa.

Consider the set $\mathfrak{I}[\alpha] \subset L^2(M)$ of all indicator functions I_C of elements of $\Sigma[\alpha]$. Choose a dense countable subset $\{I_{C_j}\}$ in $\mathbb{I}[\alpha]$. The sigma-algebra generated by C_j and sets of zero measure coincides with $\Sigma[\alpha]$. We say that m, $m' \in M$ are equivalent if for each C_j we have $m, m' \in C_j$ or m, $m' \notin C_j$. By the definition, we get a measurable partition. By the Rohlin theorem [\[44\]](#page-34-8) the quotient is a Lebesgue measure space.

of M, the quotient space is a Lebesgue measure space, denote them $(M[\alpha], \mu[\alpha])$. We have an identification of $L^2(M[\alpha])$ with the space of K_{α} -fixed vectors in $L^2(M)$.

By Subset. [1.15,](#page-12-2) we get a canonical measure preserving polymorphism $\mathfrak{x}[\alpha]$: $M \twoheadrightarrow M[\alpha]$.

Operators $T_{r+is}(\mathfrak{x}[\alpha]) : L^{1/r}(M[\alpha]) \to L^{1/r}(M)$ are operators

$$
\mathcal{E}'_{1/r}[\alpha]\varphi(m) = \varphi(\overline{m}),
$$
 where \overline{m} is element of the partition containing m,

these operators do not depend on s and in fact do not depend on r.

An operator $T_{r+is}(\mathfrak{x}[\alpha])$ is the operator of conditional expectation $\mathcal{E}_{1/r}[\alpha]$: $L^{1/r}(M) \to L^{1/r}(M[\alpha])$. For $r = 1/2$, $T_{1/2+is}(\mathfrak{r}[\alpha]^*)$ is the orthogonal projection to $L^2(M[\alpha])$.

Theorem 2.1. Let a (G, K) -pair act on a measure space (M, μ) . Assign for each $\alpha \in \mathcal{A}$ the measure space $M[\alpha]$. For each $\alpha, \beta \in \mathcal{A}$ and $g \in G$ define the polymor*phism*

(2.3)
$$
\Xi_{\alpha,\beta}(g) := \mathfrak{x}[\alpha]^* g \mathfrak{x}[\beta] \in \mathrm{Pol}(M[\alpha], M[\beta]).
$$

Then the maps $g \mapsto \Xi_{\alpha,\beta}(g)$ *determine a functor from the train* $\mathfrak{T}(G,K)$ *to the category of polymorphisms.*

PROOF. We must verify the identity

(2.4)
$$
\Xi_{\alpha,\beta}([K_{\alpha}gK_{\beta}])\circledast \Xi_{\beta,\gamma}([K_{\beta}hK_{\gamma}])=\Xi_{\alpha,\gamma}(([K_{\alpha}gK_{\beta}]\circ[K_{\beta}hK_{\gamma}])
$$

By [\[30\]](#page-33-14), Theorem 6.12, two polymorphisms are equal if and only if their Mellin– Markov transforms are equal. A Mellin–Markov transform is a holomorphic operatorvalued function in the strip Π, see [\[30\]](#page-33-14), Lemma 6.10, so it is uniquely determined by its values on the line $1/2+is$. Applying functors $T_{1/2+is}$ to both sides of conjectural equality [\(2.4\)](#page-17-1) we come to the equivalent conjectural equality

$$
\left(\mathcal{E}[\alpha]T_{1/2+is}(g)\mathcal{E}'[\beta]\right)\left(\mathcal{E}[\beta]T_{1/2+is}(h)\mathcal{E}'[\gamma]\right)=\mathcal{E}[\alpha]T_{1/2+is}(g\circ h)\mathcal{E}'[\gamma],
$$

these operators act $L^2(M[\gamma]) \to L^2(M[\alpha])$. Since $\mathcal{E}'[\beta]\mathcal{E}[\beta] = P[\beta]$, we can write the left hand side as

$$
P[\alpha] \, T_{1/2+is}(g) \, P[\beta] \, T_{1/2+is}(h) \Big|_{L^2(M[\gamma])}.
$$

By the definition of trains, this equals to

$$
P[\alpha] T_{1/2+is}(g \circ h)\Big|_{L^2(M[\gamma])}
$$

,

i.e., to the right-hand side.

2.4. Application to the bysymmetric group. 1) We take $G = \mathcal{S}, K[\alpha] =$ \mathbb{K}_{α} , $M = \mathfrak{S}^z$, and get an action of the category S.

2) In our case $[\mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\beta}] = \mathbb{K}_{\alpha} \backslash \mathbb{S}/\mathbb{K}_{\beta}$, this is clear from explicit constructions of representations of S, see [\[38\]](#page-34-2). However, in considerations above sets $[K_{\alpha} \backslash G/K_{\beta}]$ are used only for the establishing of the assiciativity, in our case the associativity is obvious. So we can simply repeat the proof of Theorem [2.1](#page-17-0) without a reference to [\[38\]](#page-34-2).

3) If we want to consider the extended category \overline{S} , see Subsect. [1.13,](#page-9-0) then we take a countable subset $\Omega \subset \mathbb{N}$ such that $\mathbb{N} \setminus \Omega$ also is countable. Elements of A are subsets $C \subset \mathbb{N}$ such that $\Omega \setminus C$ are finite. A group $K(C)$ is the subgroup in $\mathbb{K} \simeq \overline{S}_{\infty}$ fixing all elements of C.

FIGURE 4. a) A chip λ_3^6 . b) To Subsect. [3.2.](#page-18-0)

3. ACTION OF THE SEMIGROUP $\mathcal{S}(0,0)$ ON THE SPACE \mathfrak{T}^z OF RESTAURANTS

In the construction of the previous section we set:

 $M = \mathfrak{S}^z$ $G = S$, $K = \mathbb{K}$, $K_n = \mathbb{K}_n$

3.1. Spaces of half-empty restaurants. Denote by \mathfrak{S}_n^z the measure space, whose points are restaurants $\{U_{\omega}\}\in \mathfrak{T}^z$ equipped with guests $\overline{1}, \ldots, \overline{n}$ chosen uniformly, denote such points by $\{U_{\omega}\}, \{\overline{j}\}_{j\in I_n}$. Denote the measure on \mathfrak{S}_n^z by $\overline{\mu}_n^z$. In particular $\mathfrak{S}_0^z = \mathfrak{T}^z$.

We have an obvious forgetting map $\mathfrak{S}^z \to \mathfrak{S}_n^z$, the pushforward of the measure μ^z is $\overline{\mu}_n^z$.

Proposition 3.1.

$$
\mathfrak{S}^z[n]=\mathfrak{S}_n^z.
$$

PROOF. Since \mathfrak{S}_n^z is the quotient space of \mathfrak{S}^z , the space $L^2(\mathfrak{S}_n^z)$ is a canonically defined subspace in $L^2(\mathfrak{S}^z)$ and elements of this subspace are K-fixed. We must show that the space of \mathbb{K}_n -fixed vectors in $L^2(\mathfrak{S}^z)$ is precisely the $L^2(\mathfrak{S}_n^z)$. Notice that $L^2(\mathfrak{S}^z)$ is a direct integral of Hilbert spaces, the base of the integral is the measure space \mathfrak{S}_n^z and a fiber over a point $\{U_\omega\}, \{\overline{j}\}_{j\in I_n}$ is L^2 on a product of countable number of copies of $\sqcup U_\omega$ enumerated by $n+1, n+2, \ldots$. The group \mathbb{K}_n acts trivially on the base $\mathfrak{S}_{n_k}^z$. In each fiber it acts by permutation of factors $\Box U_{\omega}$. So a K_n-fixed vector in $L^2(\mathfrak{S}^z)$ is an integral of K_n-fixed vectors in fibers $L^2((\sqcup U_\omega)^\infty)$. By the zero-one law such a vector is a constant function. \square

REMARK. So the polymorphism $\mathfrak{r}[n]^* : \mathfrak{S}^z \to \mathfrak{S}_n^z$ is the removing of guests $\overline{n+1}$, $\overline{n+2}, \ldots$ The adjoint polymorphism $\mathfrak{r}[n] : \mathfrak{S}_n^z \to \mathfrak{S}^z$ is a random arrangement of new guests $\overline{n+1}$, $\overline{n+2}$, ... (these polymorphisms are measure-preserving).

3.2. Embeddings and projections. Let $m \geq n$. Define the following canonical morphisms $\lambda_n^m \in \mathcal{S}(n,m)$, see Fig. [4.](#page-18-1)a. We take arcs

1) $1_r^+ \downarrow 1_r^-[0], \ldots, n_r^+ \downarrow n_r^-[0]$ in the right hand side;

- 2) $1^+_l \downarrow 1^-_l[0], \ldots, n^+_l \downarrow n^-_l[0]$ in the left hand side;
- 3) $(n+1)⁺_l \cup (n+1)⁺_r[1/2],...,m⁺_l \cup m⁺_r[1/2].$

The following statement immediately follows from the definition:

Proposition 3.2. a The polymorphism $\Xi_{m,n}((\lambda_m^n)^*) : \mathfrak{S}_n^z \to \mathfrak{S}_m^z$ is a forgetting *of guests* $m + 1, \ldots, \overline{n}$.

b) The polymorphism $\Xi(\lambda_n^m) : \mathfrak{S}_m^z \to \mathfrak{S}_n^z$ is a uniform random arrangement of *guests* $\overline{m+1}$, ..., \overline{n} *on a given restaurant* ∟ U_{ω} .

Figure 5. a) A checker surface. b) The corresponding dessin d'enfant.

More formally, let $(LU_{\omega}, {\overline{i}}_{i\leq n}) \in \mathfrak{S}_n^z$. Then its spreaded image under $\mathfrak{r}(\lambda_n^n)$ is the measure on $\mathbb{R}^{\circ} \times \mathfrak{S}_{m}^{z}$ supported by the set consisting of points

$$
1 \times (\sqcup U_{\omega}, \{\overline{i}\}_{i \leq n}, \{\overline{j}\}_{n < j \leq m}),
$$

where the guests $\{\overline{i}\}_{{i \leq n}}$ are the same, and guests $\overline{n+1}$, ..., \overline{m} are uniformly distributed on $\sqcup U_{\omega}$.

REMARK. Proposition [3.2](#page-18-2) also provides us a description of polymorphisms corresponding to diagrams drawn on Fig. [4.](#page-18-1) We forget part of guests and arrange new guests randomly. $□$ ⊠

Our next purpose is to describe the action of the semigroup $\mathcal{S}(0,0)$ on the space $\mathfrak{T}^z = \mathfrak{S}_0^z$. We need to define some additional objects, namely checker surfaces (they also arise in representation theory of infinite symmetric group for other reasons, see $[31]$, $[29]$).

3.3. Checker surfaces. See [\[29\]](#page-33-21), [\[31\]](#page-33-6). Consider an oriented compact twodimensional closed surface (generally speaking it is disconnected). Consider a graph on this surface separating it into triangles. Let triangles be colored black and white in the checker order, i. e., neighbors of a black triangle are white and neighbors of a white triangles are black. Let edges of the graph be colored 3 colors, denote them by a, b, c. Let these colors be arranged clockwise on a perimeter of each white triangles, let they be arranged anticlockwise on perimeters of black triangles. We call such graphs on surfaces by *checker surfaces*. Two surfaces are equivalent if they are isotopic. So such a surface is a pure combinatorial object.

Denote by **?** the set of all checker surfaces, by $\mathbf{P}_n \subset \mathbf{P}$ we denote the set of surfaces with $2n$ triangles (*n* black amd *n* white).

LABELED CHECKER SURFACES. Consider an element of \mathbf{Q}_n . Assign pairwise different labels $1, \ldots, n$ to black black triangles and pairwise different labels $1, \ldots, n$ n to white triangles. We call an object obtained in this way by a *labeled checker surface*. Denote by \mathbf{P}_n^{\bullet} the set of all such surfaces.

Fix an element $\Sigma \in \mathbf{P}_n^{\bullet}$. For each color a, b, c we define an element g_a , g_b , $g_c \in S_n$ in the following way. For each $k = 1, 2, ..., n$ we find a label k on a white triangles. This triangle has a unique edge of color a , this edge is contained in a black triangle with some label, say l. Then g_a sends k to l. In the same way we define permutations g_b and g_c (they are reflections through b-edges and c edges respectively). This determines a *one-to-one correspondence between the set* Ω_n^{\bullet} *and the group* $S_n \times S_n \times S_n$ *.*

THE INVERSE CONSTRUCTION. Fix elements $g_a, g_b, g_c \in S_n$. Consider a col-lection of n white triangles with sides colored a, b, c clockwise, see Fig. [6.](#page-20-0) Assign to these triangles labels 1, \dots , n. Similarly, take a collection of n labeled black

FIGURE 6. Gluing of $\Sigma(g_a, g_b, g_c)$ from triangles.

triangles, whose sides are colored a, b, c counterclockwise. For each $k \leq n$ we glue (according the orientations) the side of type a of k -th white triangle with side of the type a of $g_a(k)$ -th a black triangle. Repeating this operation for g_b and g_c , we get an element of $\mathbf{\hat{S}}_n^{\bullet}$. Denote a checker surface obtained in this way by $\Sigma(g_a, g_b, g_c)$.

Types of vertices. Notice that we have 3 types of vertices. We say that a vertex has a type A if it is adjacent to edges of colors b, c (equivalently, it is opposite to an edge of the type a); a type B if it is adjacent to edges of colors a, c ; a type C if colors of adjacent edges are a and b.

Colors of edges containing a vertex interchange, so valences of vertices of checker surfaces are even. We say that the *order* ord (v) *of a vertex* v is the half of its valence. For $\Sigma(g_a, g_b, g_c) \in \mathbf{P}_n^{\bullet}$, cycles of $g_a^{-1}g_b$ are in one-to-one correspondence with vertices of the type C , etc. Below it is convenient to use a double terminology: *vertices of the type* C = *vertices of the type* (g_a, g_b) , etc.

3.4. Remark. Dessins d'enfant. The set Ω is in one-to-one correspondence with 'dessins d'enfant' in the sense of Grothendieck. Recall that a *dessin d'enfant* is a bipartite graph on an oriented compact two-dimensional surface^{[11](#page-20-1)} such that the complement of the graph is a union of disks^{[12](#page-20-2)}. According the Belyi theorem, dessin d'enfants are in one-to-one correspondence with pairs (R, Φ) , where R is a nonsingular complex curve defined over the algebraic closure \overline{Q} of rational numbers \mathbb{Q} , and Φ is a holomorphic function from R to the Riemann sphere $\overline{\mathbb{C}} = \mathbb{C} \cup \infty$, whose critical values are 0, 1, ∞ . For detailed discussions, applications, and references, see [\[19\]](#page-33-26), [\[45\]](#page-34-12).

Consider a checker surface. Removing edges of colors a, b we get a dessin. Vertices of the type C are in one to-one correspondence with complementary disks.

3.5. Dirichlet measures on simplices. Fix $\ell > 0$. Consider the $(p-1)$ dimensional simplex $\Delta_p(\ell) \subset \mathbb{R}^p$ defined by

$$
x_1 + \dots + x_p = \ell, \qquad \text{where } x_j \geqslant 0.
$$

Let k_1, \ldots, k_p be positive reals. A *Dirichlet distribution* $\Theta_p[k_1, \ldots, k_p; \ell]$ is a (probabilistic) measure on \mathbb{R}^p supported by $\Delta_p(\ell)$ and given by

$$
\frac{\Gamma(k_1+\cdots+k_p)\,\ell^{-p+1}}{\prod_{i=1}^p\Gamma(k_i)}\prod_{i=1}^p x_i^{k_i-1}dx_1\ldots dx_{p-1},
$$

see, e.g., [\[52\]](#page-34-13), 7.7. This is a general definition, actually, we need only integer k_i

¹¹Usually such surfaces are assumed to be connected, we admit disconnected surfaces.

¹²Vertices are colored two colors, and edges can connect only vertices of different colors.

REMARK. In this formula dx_p is absent, but the coordinate x_p is not distinguished, all measures $dx_1 \ldots dx_{j-1} dx_{j+1} \ldots dx_p$ coincide on $\Delta_p(\ell)$. ⊠

If $p = 1$, we define $\Theta_1[k; \ell]$ as the delta-measure on \mathbb{R}^1 supported by $x_1 = \ell$.

More generally, let $m_1, \ldots, m_q \geq 0$. Let I be set of all j such that $m_j > 0$, say $I = \{m_{\alpha_1}, \ldots, m_{\alpha_p}\}.$ Then \mathbb{R}^q splits as a product of the space $\mathbb{R}^{\#I}$ with coordinates x_{β} , where $\beta \in I$, and the space $\mathbb{R}^{q-\#I}$ with coordinates x_{γ} , where $\gamma \notin I$. We define $\Theta_q[m_1,\ldots,m_q;\ell]$ as a product of the measure $\Theta_{\#I}[m_{\alpha_1},\ldots,m_{\alpha_p};\ell]$ on $\mathbb{R}^{\#I}$ and the atomic unit measure on $\mathbb{R}^{q-\#I}$ supported by 0.

We need Dirichlet distributions for the following reason. Let n points be distributed uniformly on the circle of length ℓ . They split the circle into n pieces. Let y_1, \ldots, y_n be their lengths, such collections are elements of the simplex $\Delta_n(\ell)$ distributed according $\Theta_n(1,\ldots,1;\ell)$, i.e., by

$$
(3.1) \t\t (n-1)! \ell^{-n+1} dy_1 \dots dy_{n-1},
$$

this expression is symmetric with respect to permutations of y_1, \ldots, y_n .

Lemma 3.3. *Fix* $k_1, \ldots k_p \in \mathbb{N}, \sum k_j = n$. *Consider the map* $\Delta_n(\ell) \to \Delta_p(\ell)$ *given by*

.

(3.2)
$$
x_1 = \sum_{j=1}^{k_1} y_j, \quad x_2 = \sum_{j=k_1+1}^{k_1+k_2} y_j, \quad \dots, \quad x_p = \sum_{j=k_1+\dots+k_{p-1}+1}^{n}
$$

Then the pushforward of the measure [\(3.1\)](#page-21-0) *is the distribution* $\Theta_p[k_1, \ldots, k_p; \ell]$ *.*

This is a special case of the aggregation property of the Dirichlet distributions, see, e.g., [\[52\]](#page-34-13), Subsect. 7.7.5.

3.6. Laplace transforms of the Dirichlet distributions. Let Ψ be a measure supported by \mathbb{R}^p_+ . Its Laplace transform is

$$
\mathcal{L}\Psi(u_1,\ldots,u_p):=\int_{\mathbb{R}^p_+}e^{-(u_1x_1+\cdots+u_px_p)}\,d\Psi(x_1,\ldots,x_p),
$$

where Re $u_j \geqslant 0$.

Proposition 3.4. *Let* $k_j > 0$. *The Laplace transform of the Dirichlet distribution* $\Theta_p[k_1,\ldots,k_p;a]$ *is given by*

$$
(3.3) \ \mathcal{L}\Theta_p[k_1,\ldots,k_p;a](u_1,\ldots,u_p) = \frac{\Gamma(\sum k_j) a^{-p+1}}{2\pi i} \int_{-i\infty}^{i\infty} e^{az} \prod_{j=1}^p (z+u_j)^{-k_j} dz.
$$

If $k_j \in \mathbb{N}$ *, then this expession equals*

(3.4)
$$
\frac{\left(\sum k_j-1\right)! a^{-p+1}}{\prod_j (k_j-1)!} \prod_{j=1}^p \left(-\frac{\partial}{\partial u_j}\right)^{k_j-1} \left[\sum_m \frac{e^{-au_m}}{\prod_{j:\,j\neq m} (u_j-u_m)}\right].
$$

Below we need (3.4) . The statement (3.3) was obtained in Phillips [\[41\]](#page-34-14), we present proofs of both formulas.

PROOF. Consider functions $f_1(x_1), \ldots, f_p(x_p)$ on \mathbb{R}_+ and an integral

$$
\varphi_a := \int_{\Delta_p(a)} f_1(x_1) \dots f_p(x_p) dx_1 \dots dx_{p-1}.
$$

22

Consider its Laplace transform (Re $z \geqslant 0$)

$$
\int_0^\infty e^{-za} \varphi_a da = \int_{\mathbb{R}^p_+} e^{-z \sum x_j} \prod f_j(x_j) dx_1 \dots dx_p = \prod_{j=1}^p \int_0^\infty e^{-zx} f_j(x) dx.
$$

Applying the inversion formula for the Laplace transform, we get (cf. [\[43\]](#page-34-15), formula $(3.3.4.1)$

(3.5)
$$
\varphi_a = \frac{1}{2\pi i} \int_{-i\infty}^{i\infty} e^{za} \left(\prod_{j=1}^p \int_0^\infty e^{-zx} f_j(x) \, dx \right) dz.
$$

If $f_j(x_j) = x_j^{k_j-1} e^{-u_j x_j}$, then φ_a is (upto a constant factor) the Laplace transform of $\Theta_p[k_1,\ldots,k_p]$. We have

$$
\int_0^{\infty} e^{-zx} f_j(x) dx = \int_0^{\infty} x^{k_j - 1} e^{-(z + u_j)x} dx = \frac{\Gamma(k_j)}{(z + u_j)^{k_j}}.
$$

Applying [\(3.5\)](#page-22-0), we come to the first statement [\(3.3\)](#page-21-2) of the proposition. For general k_i the integral in the right-hand side of [\(3.3\)](#page-21-2) is a kind of multivariate confluent hypergeometric function. For integer k_j this integral can be evaluated by residues.

Now set $k_j = 1$ in [\(3.3\)](#page-21-2). Then

$$
\frac{1}{\prod_{j=1}^{p}(z+u_j)} = \sum_{m} \frac{1}{\prod_{j: j \neq m}(u_j - u_m)} \cdot \frac{1}{z+u_m}.
$$

Keeping in mind the integral

$$
\frac{1}{2\pi i} \operatorname{v.p.} \int_{-i\infty}^{i\infty} \frac{e^{az} dz}{z + u_m} = e^{-au_m},
$$

we come to the identity

$$
\mathcal{L}\Theta_p[1,\ldots,1;a](u_1,\ldots,u_p) =
$$

=
$$
\int\limits_{\Delta_p(a)}\prod_{j=1}^p e^{-u_jx_j} dx_1 \ldots dx_{p-1} = \sum_m \frac{e^{-au_m}}{\prod_{j:\ j\neq m}(u_j-u_m)}.
$$

A multiplication of a distribution by x_j implies a differentiation of its Laplace transform by u_i . This leads to [\(3.4\)](#page-21-1).

3.7. The action of $\mathcal{S}[0,0]$ on \mathfrak{S}_0^z . Consider an element

$$
\{\bigcirc[k_j]\}_j \in \mathcal{S}[0,0],
$$

i.e., a union of cycles of lengths $k_1, \ldots, k_p \geq 2$. To write a formula for the corresponding polymorphism, we need some notation.

Fix a collection of tables $\{U_{\omega}\}\in \mathfrak{S}^z_0$, let ℓ_{ω} be their lengths, recall that they are pairwise different almost sure.

For a given $\{U_{\nu}\}_{\nu} \in \mathfrak{S}_{0}^{z}$ we intend to write the spreaded image of $\{U_{\nu}\}_{\nu}$, i.e., the corresponding conditional measure on the space $\mathbb{R}^{\circ}\times\mathfrak{S}^{z}_{0}$. This measure is supported by points of the type $z^q \times \{V_x\}_\varkappa$, where a set $\{V_x\}_\varkappa$ coincides with $\{U_\nu\}_\nu$ up to a finite number of tables. It is more convenient to speak about collections $\{\ell_{\nu}\}\$ of lengths, which are point configurations on the segment $[0, 1]$. We introduce the operation

$$
\text{Replace}\Big[\{\ell_{\omega_j}\}_j\longrightarrow\Lambda\Big]\{\ell_{\nu}\}
$$

on such configurations, which replaces a finite subcollection $\{\ell_{\omega_j}\}_j$ of the collection $\{\ell_{\nu}\}\$ by a random collection $\subset \mathbb{R}^p_+$ with a given distribution Λ . More generally, we will use such notation for similar transformations of lists of other types.

NOTATION. a) For $\Sigma \in \mathbf{P}$ denote by A_{α} its A-vertices, by B_{β} its B-vertices, by C_{γ} its C-vertices. Denote by $\#\{B_{\beta}\}\$ (resp. $\#\{C_{\gamma}\}\$) the number of B-vertices (resp. C-vertices).

b) For $\Sigma \in \mathbf{Q}$ and $\{U_{\nu}\}_{\nu \in \Omega} \in \mathfrak{S}_{0}^{z}$ we define a *framing* of $\Sigma \in \mathbf{Q}$ as an injective map ω from the set of B-vertices of Σ to the set of tables. In particular, for any Bvertex B_β we assign a number $\ell_{\omega(B_\beta)}$. Denote by $\text{Fr}(\Sigma, {\ell_\nu})$ the set of all framings of Σ.

c) For $\Sigma \in \mathbf{P}$ we denote by $\text{Aut}_B(\Sigma)$ the group of all automorphisms of Σ fixing all B -vertices^{[13](#page-23-1)}.

d) Let $\{k_j\}$ be as above. By $\Gamma[\{k_j\}]$ denote the set of all $\Sigma \in \mathbf{P}_n$ whose A-vertices have orders $\{k_i\}$.

e) Denote by $\iota_m[\{k_i\}]$ the number of entries of m to a list $\{k_i\}$.

Theorem 3.5. *Fix an element* $\{ \bigcirc [k_j] \}_j$ ∈ $\{ \emptyset [0,0]$ *and a restaurant* ⊔U_v *with tables of lengths* $\{\ell_{\nu}\}\$ *. For* $\Sigma \in \Gamma[\{k_j\}]$ *denote by* $m_{\beta\gamma}$ *the number of edges connecting* B_{β} and C_{γ} . Then the polymorphism $\Xi_{0,0}(\{\bigcirc [k_j]\})$ sends the point $\{\ell_{\nu}\}\)$ to the *following measure on* $\mathbb{R}^{\circ} \times \mathfrak{S}_{0}^{z}$:

$$
(3.6) \prod_{j} k_{j} \prod_{m \in \mathbb{N}} \iota_{m}[\{k_{j}\}]! \sum_{\Sigma \in \Gamma[\{k_{j}\}], \omega \in \text{Fr}(\Sigma, \{\ell_{\nu}\})} \frac{1}{\# \text{Aut}_{B}(\Sigma)} \cdot \prod_{B_{\beta}} \frac{\ell_{\omega(B_{\beta})}^{\text{ord}(B_{\beta})}}{(\text{ord}(B_{\beta}) - 1)!} \times \\ \times \left(\delta_{\mathbb{R}^{\circ}}[[z^{\# \{C_{\gamma}\} - \# \{B_{\beta}\} }]] \times \right. \times \text{Replace} \left[\left\{ \ell_{\omega(B_{\beta})} \right\}_{\beta} \longrightarrow \left(\mathbf{\ddot{x}}_{\beta} \Theta_{\# \{C_{\gamma}\}}[\{m_{\beta\gamma}\}_{\gamma}; \ell_{\omega(B_{\beta})}]\right) \right] \left\{ \ell_{\nu} \right\},
$$

where the symbol $\mathbf{\ddot{x}}$ denotes a convolution of Dirichlet distributions on $\mathbb{R}^{\#{C_\gamma}}$.

Notice that a convolution in a summand of the formula is a measure on the simplex $\Delta_{\#(C_\gamma)}(\sum \ell_{\omega(B_\beta)})$, densities of such measures are piecewise polynomial. In any case, by Proposition [3.4,](#page-21-3) Laplace transforms of such measures are explicit elementary functions.

Theorem is proved in Subsect. [3.8–](#page-23-2)[3.12.](#page-27-0)

3.8. A product of two permutations, which are determined in terms of disjoint cycles. Recall that a *ribbon graph* is a graph with fixed cyclic orders of edges at each vertex, see Fig. [7.](#page-24-0) Such a graph can be regarded as an oriented two-dimensional surface consisting of bands along edges. Boundary of this surface is a union of circles, gluing a disk to each component of the boundary we get a compact oriented surface.

A construction of a bipartite ribbon graph by two pemutations. Let $g, h \in S_n$ be represented as products of disjoint cycles

$$
g = (\sigma_1)(\sigma_2)\ldots, \qquad h = (\tau_1)(\tau_2)\ldots
$$

For each cycle (σ_{α}) we draw a *'chamomile'* A_{α} as on Fig. [8](#page-24-1) and enumerate petals

 13 This group is poor, for a connected surface it is cyclic (and usually trivial), for a disconnected surface it is a product of cyclic groups.

FIGURE 7. A ribbon graph and the corresponding band domain on a surface.

Figure 8. A chamomile.

Figure 9. Gluing of chamomiles.

of A_{α} by elements of the cycle (σ_{α}) clockwise. For each cycle (τ_{β}) of h we draw a similar chamomile B_β . Gluing petals of g-chamomiles and petals of h-chamomiles according to the enumeration of petals and orientations (see Fig. [9\)](#page-24-2), we get a bipartite ribbon graph, say $Gr(g, h)$, whose list of vertices is $\{A_{\alpha}\}, \{B_{\beta}\}\$ and edges are enumerated by $1, 2, \ldots, n$.

Each component of the boundary of the ribbon graph $\text{Gr}(q, h)$ can be regarded as a polygonal path with an even number of sides. Passing it counterclockwise we observe edges of two types, with origins in A-vertices and with origins in B-vertices. Labels on edges of the first type form cycles of the permutation hg , labels on edges of the second type form cycles of gh . See Fig. [10.](#page-25-0)a. This correspondence arises at least to Goulden, Jackson [\[7\]](#page-32-6).

Insertions of elements to cycles and ribbon graphs. Such pictures are well-compatible with insertions of additional elements to cycles of h . Namely, take a larger group S_{n+N} . Let $\tilde{g} \in S_{n+N}$ be the trivial extension of g. Choose an

FIGURE 10. a) A product of two permutations g, h in terms of ribbon graphs. b) An insertion of two elements q_1 , q_2 to a cycle of h between j and k . c) A labeled checker surface corresponding to elements $g, h^{-1}, 1 \in S_n$. d) Insertion of two elements to a cycle of h on the language of checker surfaces.

element $h^{\circ N} \in S_{n+N}$ such that

$$
\Upsilon_n^N(h^{\circ N}) = h.
$$

Then the ribbon graph $Gr(\tilde{g}, h^{\circ N})$ can be easily obtained from the ribbon graph $Gr(\tilde{g}, h)$. Now the contract is a set of classes of classes in the contract of classes $Gr(g, h)$. Namely, an insertion of elements $q_1, \ldots, q_r > n$ to a cycle (τ_β) between j and k ,

$$
(\ldots j\,k\ldots)\mapsto(\ldots j\,q_1\ldots q_r k\ldots),
$$

means that we draw at the vertex β edges labeled by q_1, \ldots, q_r between edges j and k. Opposite vertices of these edges have valence 1 (since $\tilde{g}(q_{\mu}) = q_{\mu}$). See, Fig. [10.](#page-25-0)b.

3.9. Checker surfaces and products of permutations. Reformulate this construction in the terms of checker surfaces. For $g, h \in S_n$ we take the surface $\Sigma(g,1,h^{-1}) \in \mathbf{P}_{n}^{\bullet}$. Vertices of this surface have types $(1,g)$, $(h^{-1},1)$, (g,h^{-1}) , they correspond to cycles of permutations g, h, $(hg)^{-1}$ respectively.

For each vertex of (g, h^{-1}) -type, labels in adjacent white triangles passing counterclockwise give a cycle of hg.

Removing vertices of the type (g, h^{-1}) and adjacent edges we get a ribbon graph described above, remaining edges have type (1), labels on both sides of such an

FIGURE 11. To a pass from $\Sigma(g, 1, h^{-1})$ to $\Sigma(\tilde{g}, 1, (h^{\circ N})^{-1})$. Additional drawing on former black triangles.

FIGURE 12. Transformations of a piece of a table with guests.

edge coincide and so these labels can be attributed to edges. We get the ribbon graph $\operatorname{Gr}(g, h^{-1})$.

Insertion of elements to cycles and checker surfaces. Operation of insertion of an element with number $> n$ to a cycle of h is shown on the Fig. [10.](#page-25-0)

So the surface $\Sigma(\tilde{g}, 1, (h^{\circ N})^{-1})$ is obtained from the surface $\Sigma(g, 1, h^{-1})$ by the following operation. We preserve all white triangles with labels, and black triangles are transformed as it is shown on Fig. [11.](#page-26-0)

3.10. The right action of S_n on the space of virtual permutations \mathfrak{S}^z . Let $g \in S_n$, $\mathfrak{u} \in \mathfrak{S}^z$. Denote $u = \Upsilon_n^{\infty} \mathfrak{u}$. Consider the surface $\Sigma(g, 1, u^{-1}) \in \mathbf{P}_n^{\bullet}$. Take a B-vertex, let T be an adjacent black triangle. Let the clockwise white neighbor of T has label j and so the counterclockwise white neighbor has label $u(j)$. We assign to T the interval $(j, u(j))$ of the corresponding occupied table, denote this interval by $\mathcal{O}(T)$, see Fig. [12.](#page-26-1)

For each C-vertex consider a counterclockwise cyclic chain

$$
j_1,\mathcal{O}(T_{i_1}),j_2,\mathcal{O}(T_{i_2}),\ldots
$$

consisting of labels on adjacent white triangles and ordered sets $\mathcal{O}(\cdot)$ on adjacent black triangles. Uniting them, we get a table of ug.

3.11. The action of S_n on \mathfrak{S}_n^z . For an element $g \in S_n$ consider the polymorphism

$$
\Xi_{n,n}(g) = \mathfrak{x}[n]^* g\mathfrak{x}[n]
$$

of \mathfrak{S}_n^z . Consider a point \mathfrak{u} of \mathfrak{S}_n^z . It determines a permutation $u \in S_n$ and a collection of lengths $\{l_j\}$ between points \overline{j} and $\overline{u(j)}$.

Consider the labeled checker surface $\Sigma = \Sigma(g, 1, u^{-1}) \in \mathbf{P}_n^{\bullet}$. For a black triangle of Σ with label j we assign the length of the arc $[\overline{j}, u(j)]$.

Lemma 3.6. *The spreaded image of a point* $\mathfrak{u} \in \mathfrak{S}_n^z$ *under the polymorphism* $g \in S_n$ *is a* δ -measure on $\mathbb{R}^{\circ} \times \mathfrak{S}_n^z$ supported by a point $z^s \times \mathfrak{v}$, where

$$
s = \#\{(g^{-1}, u) \text{-vertices}\} - \#\{(u^{-1}, 1) \text{-vertices}\}
$$

and $\mathfrak{v} \in \mathfrak{S}_n^z$ is defined in the following way. For each (g, u^{-1}) -vertex C_γ of Σ we *draw a table taking labels* $\overline{\mu}_1$, $\overline{\mu}_2$, ... *from white triangles passing counterclockwise* and length of arc $[\overline{\mu}_{i-1}, \overline{\mu}_i]$ from black triangles. This gives us a collection of tables *containing all labels* $\overline{1}, \ldots, \overline{n}$ *. Adding tables of* **u** *that do not contain labels we get* v*.*

In particular, in this case we get a deterministic map $\mathfrak{S}_n^z \to \mathfrak{S}_n^z$, so the group S_n acts on the space \mathfrak{S}_n^z .

The statement follows from the the construction of the previous subsection and formula (1.8) .

3.12. Proof of Theorem [3.5.](#page-23-0) Denote $n := \sum k_j$. Choose an element $g \in$ S_n whose cycles have lengths k_1, \ldots, k_p . Denote by (σ_α) disjoint cycles of g. This choice is noncanonical and further considerations depend on g until the last paragraph of the proof. We extend g to an element \tilde{g} of S_{∞} in a trivial way. Then the double coset $\mathbb{K} \cdot (1,\tilde{g}) \cdot \mathbb{K}$ is $\{\bigcap [k_i]\}\$. We decompose

$$
\{\bigcirc[k_j]\} = (\lambda_0^n)^* g \lambda_0^n.
$$

Recall that the polymorphism $\Xi(\lambda_0^n) : \mathfrak{S}_0^z \to \mathfrak{S}_z^n$ defined in Subsect. [3.2](#page-18-0) is a random uniform arrangement of guests $\overline{1}$, ..., \overline{n} in an empty restaurant. This operation can be described in the following (more complicated) way.

Fix a restaurant, let $\{\ell_{\mu}\}\$ be lengths of tables (recall that they are pairwise distinct a.s).

— First, fix a collection $\{s_{\mu}\}\$ of nonnegative integers such that $\sum s_{\mu} = n$. Assume that a probability of such a choice is

$$
\frac{n! \prod \ell_\mu^{s_\mu}}{\prod s_\mu!}
$$

(products actually are finite, the sum equals 1). Denote $\mathcal{B} = \mathcal{B}(\{s_{\mu}\})$ the set of all μ , for which $s_{\mu} > 0$.

— Second, for each $s_{\mu} > 0$ we take a cyclically ordered set V_{μ} with s_{μ} elements. Choose a bijection $\{1, 2, ..., n\} \rightarrow \sqcup_{\mu \in \mathcal{B}} V_{\mu}$ and assume that all variants have the same probability 1/n!. So we get a collection $\{V_\mu\}_{\mu \in \mathcal{B}}$ of cyclically ordered sets consisting of positive integers $\leq n$. We consider each set up to cyclic ordering, so there are $\prod_{\mu \in \mathcal{B}} s_{\mu}$ ways to obtain the same collection. Thus, we get a permutation $u \in S_n$ defined as a product of cycles.

— Third, for each cycle V_μ of u we insert arcs between elements of the cycle (guests), lengths of arcs are chosen according the uniform probabilistic distribution on the simplex $\Delta_{s_{\mu}}(\ell_{\mu})$ (i.e., the Dirichlet distributon $\Theta_{s_{\mu}}[1,\ldots,1;\ell_{\mu}]).$

So we get an element $\mathfrak u$ of $\mathfrak S_n^z$ depending on $\{s_\mu\}$, $u\in S_n$, and a point of product of simplices $\prod_{\mu \in \mathcal{B}} \Delta_{s_{\mu}}(\ell_{\mu}).$

Keeping in mind Lemma [3.6,](#page-27-1) we can apply $g \in S_n$ to a random element $\mathfrak{u} \in \mathfrak{S}_n^z$.

For each u consider the surface $\Sigma(g, 1, u^{-1})$. By definition, labels on both sides of any edge of the color (1) coincide. Vertices of types A, B, C in formulation of Theorem [3.5](#page-23-0) correspond to types $(1, g)$, $(u^{-1}, 1)$, (g, u^{-1}) . Clockwise cycles of labels

on white triangles about a vertex A_{α} are cycles (σ_{α}) of g. Each surface is equipped with a framing ω , namely B-vertices correspond to tables $\{U_{\mu}\}_{{\mu\in\mathcal{B}}}$ of the initial restaurant $\{U_{\mu}\}\$ (comparatively to the set $\Gamma[\{k_j\}]$ in the formulation of Theorem [3.5,](#page-23-0) we add labels on triangles). Additionally, we assign a positive number $l(j)$ to each black triangle $T(j)$ with label j. Such numbers around a given B-vertex are distributed according the measure $\Theta_{\text{ord}(B_\beta)}[1,\ldots,1;\ell_{\omega(B_\beta)}].$

New tables of a random restaurant $\Xi_{0,n}((\lambda_0^n)^*)\Xi_{n,n}(g)\Xi_{n,0}(\lambda_0^n)$ correspond to vertices C_{γ} , a length $L(\gamma)$ of such a table is sum of real numbers $l(j)$ attributed to black triangles containing C_{γ} ,

$$
L(\gamma) = \sum_{j:\, C_{\gamma} \in T(j)} l(j).
$$

We write this sum in the form

$$
L(\gamma) = \sum_{B_{\beta}} \sum_{j: B_{\beta}, C_{\gamma} \in T(j)} l(j).
$$

By Lemma [3.3,](#page-21-4) for a fixed β a distribution of vectors

$$
\Big\{\sum_{j:\ B_\beta,C_\gamma\in T(j)}l(j)\Big\}_\gamma
$$

is the Dirichlet distribution $\Theta_{\#(C_\gamma)}[\{m_{\beta\gamma}\}_{\gamma};\ell_{\omega(C_\beta)}].$ Therefore the sums $\{L(\gamma)\}_{\gamma}$ are distributed as a convolution $\mathbf{\ddot{x}}_{\beta}$ of such Dirichlet distributions.

Hence the desired conditional measure is

$$
\sum_{\Sigma(g,1,u^{-1}) \text{ with given } g} \sum_{\omega \in \text{Fr}(\Sigma(g,1,u^{-1}),\{\ell_{\nu}\})} \prod_{B_{\beta}} \frac{\ell_{\omega(B_{\beta})}^{\text{ord}(B_{\beta})}}{\text{ord}(B_{\beta})-1)!} \times \times \left(\delta_{\mathbb{R}^{\circ}}[[z^{\# \{C_{\gamma}\} - \# \{ (B_{\beta}) \} }]] \times \right)
$$

$$
\times \left(\text{Replace}\left[\{\ell_{\omega(B_{\beta})}\} \longrightarrow \left(\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{\mathbf{0}}}}}}}}}\right) \left[\{\mathbf{m}_{\beta\gamma}\}_{\gamma};\ell_{\omega(B_{\beta})}\right] \right) \right] \{\ell_{\nu}\}\right)
$$

In fact, the first summation is taken over the subset in \mathbf{P}_n^{\bullet} consisting of labeled surfaces such that:

1) for each c-edge labels on two-sides of the edge coincide;

2) clockwise cycles of labels of white triangles about A-vertices coincide with the disjoint cycles (σ_{α}) of the permutation g.

Forgetting all labels we get an element of the set $\Gamma[\{k_i\}]$ defined in Subsect. [3.7.](#page-22-1) Fix a surface $\Lambda \in \Gamma[\lbrace k_i \rbrace]$ with fixed framing (so we can think that B-vertices of Λ are enumerated) and evaluate the number of different surfaces of the type $\Sigma(g,1,u^{-1})$ over Σ .

1) We must choose a one-to-one correspondence between the set of A-vertices and the set of cycles (σ_{α}) of g (an order of a vertex must coincide with an order of the corresponding cycle). This can be done by $\prod_{m\geqslant 2} \iota_m\big[\{k_j\}\big]$! ways.

2) After this, the collection of labels on white triangles containing a vertex A_{α} consists of elements of the corresponding cycle (σ_{α}) in the same cyclic order. The only freedom is rotation of the cycle, this gives us the factor $\prod \text{ord}(A_{\alpha}) = \prod k_{\alpha}$

3) Labels on black triangles are uniquely determined by labels on white triangles.

.

FIGURE 13. To Subsect. [4.1.](#page-29-1) A chip $\mathfrak b$

4) Two surfaces $\widetilde{\Lambda}_1$, $\widetilde{\Lambda}_2$ obtained by such arrangements (including framings) can be equivalent. In this case we have an automorphism of the surface preserving the framing and identifying two labelings. So we must divide the result by the order of the group of such automorphisms.

We came to formula (3.6) .

4. FURTHER STATEMENTS

4.1. A more general case. Next, let us consider a space \mathfrak{S}_n^z and describe its polymorphism determined by a chip, whose left-hand half is trivial, see Fig. [13.](#page-29-0)

DATA DEFINING A CHIP. Fix a permutation $\sigma \in S_n$, a collection $\varphi_1, \ldots, \varphi_n \in \mathbb{Z}_+$ and a collection of integers $k_1, \ldots, k_p \geq 2$. Define a chip

$$
\mathfrak{b} = \mathfrak{b}\big[\sigma, \{\varphi_i\}, \{k_j\}\big] \in \mathcal{S}(n, n)
$$

consisting of the following arcs:

- 1) $j_l^+ \downarrow j_l^- [0]$, where $j = 1, \ldots, n;$
- 2) $j_r^+ \downarrow \sigma(j)_r^- [\varphi_j]$, where where $j = 1, \ldots, n;$
- 3) $\bigcirc [k_i]$, where $i = 1, \ldots, p$.

DATA DETERMINING A HALF-EMPTY TABLE. Next, fix an element $\tau \in S_n$ represented as a product of disjoint cycles, a collection of positive reals l_1, \ldots, l_n , and a countable collection of positive reals ℓ_{ν} such that $\sum_{i} l_i + \sum_{j} \ell_{j} = 1$. These data define a point

$$
\mathfrak{t} = \mathfrak{t}[\tau, \{l_i\}, \{\ell_\nu\}] \in \mathfrak{T}_n^z.
$$

Namely, for *i*-th cycle of τ we create a table U_i containing points of the cycle as guests (in the same cyclic order), and claim that distances between j and $\tau(j)$ are l_i . Also, we add label-free tables of lengths ℓ_ω .

Now we intend to write the spreaded image of the half-empty restaurant t under the polymorphism b. The result is similar to Theorem [3.5,](#page-23-0) but it must include additional combinatorial data.

THE SET OF SUMMATION. Define the following set $\Gamma[\mathfrak{b};\mathfrak{t}]$ consisting of checker surfaces equipped with following additional structures:

1) An injective map F (*labeling*)

$$
F: \{1, 2, \ldots, n\} \to \{\text{set of white triangles}\}.
$$

2) A bijection between

 $\{$ set of A-vertices of $\Sigma\} \longleftrightarrow$

 \longleftrightarrow {set of cycles \bigcirc | \cup {set of cycles \bigcirc | k_i] of the chip \mathfrak{b} }.

FIGURE 14. a) A cycle of σ (on the figure we have a cycle (273)) and corresponding arcs of a chip. b) The A-vertex corresponding to this cycle σ .

FIGURE 15. A table, the corresponding B-vertex, and sets $T(i)$.

This map must be compatible with the labeling F. Namely, if a vertex A_{α} corresponds to a cycle $\bigcirc [k_i]$ of the chip, then ord $A_\alpha = k_i$ and white triangles containing A_{α} are label-less. If a vertex A_{α} corresponds to a cycle $(\nu_1 \nu_2 \ldots \nu_{\rho})$ of σ , then labeled white triangles containing A are precisely $F(\nu_1), \ldots, F(\nu_\rho)$ in this cyclic order, and for each ν_i there are precisely φ_i label-less white triangles between $F(\nu_i)$ and $F(\nu_{i+1})$, see Fig. [14.](#page-30-0)

3) An injective map ω (framing)

 $\omega: \{\text{set of } B\text{-vertices of } \Sigma\} \to \{\text{set of tables of } t\}.$

Moreover, labels on white triangles containing B_β must coincide with guests on the corresponding table in reversed cyclic order, see Fig. [15.](#page-30-1)

SOME NOTATION. 1) Vertices B_β split into two types. Denote by Φ the set of all B_β such that all white triangles containing B_β are label-less. For $B_\beta \in \Phi$ denote by $T(B_\beta)$ the set of *black* triangles containing B_β .

For $B_\beta \notin \Phi$ consider the counterclockwise cycle of adjacent white triangles, let x_1, \ldots, x_s be labels on labeled white triangles. For each adjacent pair (x_i, x_{i+1}) denote by $T(\varkappa_i)$ the set of *black* triangles between them. Notice that a B-vertex is determined by the number \varkappa_i and we can omit β from the notation, see Fig. [15.](#page-30-1)

2) Similarly, denote by Ψ the set of C-vertices, for which all adjacent triangles are label-less. For $C_\gamma \in \Psi$ denote by $T(C_\gamma)$ the set of *black* triangles containing C. For $C \notin \Psi$ we write counterclockwise labels on adjacent white triangles (μ_1, μ_2, \ldots) . Denote by $R(\mu_i)$ the set of *black* triangles between μ_i and μ_{i+1} .

3) For each vertex of C we consider the counterclockwise cycle of adjacent labeled triangles. Uniting such cycles we get an element ρ of S_n .

4) Denote by $\text{Aut}_B(\Sigma, F)$ the group of all automorphisms of Σ fixing all Bvertices and all labeled triangles. Such an automorphism automatically is trivial on each connected component of Σ containing a labeled triangle.

REMARK. In Theorem [3.5](#page-23-0) parameters of Dirichlet measures $\Theta[\ldots]$ are expressed in the terms of the matrix $m_{\beta\gamma}$ whose elements are number of black triangles with vertices B_β and C_γ . Rows of the matrix are enumerated by B-vertices and columns by C-vertices. In our case, rows of a similar matrix are enumerated by chains $T(i)$ between two labeled triangles at B-vertices $\notin \Phi$ and by vertices $B_\beta \in \Phi$. Columns are enumerated by chains $R(j)$ between labeled triangles at C-vertices $\notin \Psi$ and by vertices $C_{\gamma} \in \Psi$. So analog of the matrix $m_{\beta\gamma}$ contains elements of types

$$
#(T(i) \cap R(j)) \qquad #(T(i) \cap R(C_{\gamma}))
$$

$$
\#(T(B_{\beta}) \cap R(j)) \qquad #(T(B_{\beta}) \cap R(C_{\gamma}))
$$

For this reason an expression for convolutions now is longer, but in fact the structure of the formula is the same. ⊠

Theorem 4.1. *The spreaded image of a point* t *under a polymorphism* b *is the measure on* $\mathbb{R}^{\circ} \times \mathfrak{S}_n$ *given by the formula*

$$
(4.1) \prod k_i \prod \iota_m[\{k_j\}] \sum_{(\Sigma,\omega,F)\in\Gamma[b,t]} \frac{1}{\#G(\Sigma,F)} \prod_{\nu=1}^n \ell_{\nu}^{\#T(\nu)} \prod_{B_{\beta}\in\Phi} \frac{\ell_{\omega(B_{\beta})}^{\text{ord}(B_{\beta})}}{(\text{ord}(B_{\beta})-1)!} \times \\ \times \left(\delta_{\mathbb{R}^{\circ}}\left[\left[z^{\#{C_{\gamma}}\right)-\#{B_{\beta}}\right]\right] \times \\ \times \text{Replace}\left[(\tau;\{l_i\};\{\ell_{\omega(B_{\beta})}\}_{\beta\in\Phi}) \longrightarrow \\ \left(\rho;\underset{i\in I_n}{\star} \Theta_{n+\# \Psi}\left[\{\#(T(i)\cap R(j))\}_{j\in I_n},\{\#(T(i)\cap R(C_{\gamma}))\}_{C_{\gamma}\in\Psi};l_i\right]\right) \star \\ \left(\underset{B_{\beta}\in\Phi}{\star} \Theta_{n+\# \Psi}\left[\{\#(T(B_{\beta})\cap R(j))\}_{j\in I_n},\{\#(T(B_{\beta})\cap R(C_{\gamma}))\}_{C_{\gamma}\in\Psi};\ell_{\omega(B_{\beta})}\right]\right)\right] \qquad (\tau;\{l_i\};\{\ell_{\nu}\})
$$

For a proof we set $N = n + \sum \varphi_i + \sum k_j$ and take an element $g \in S_N$ such that $\mathfrak{b} = (\lambda_n^N)^* g \lambda_n^N.$

Further arguments repeat the proof of Theorem [3.5](#page-23-0) and we omit them.

4.2. Action of the center of $\mathcal{S}(\infty, \infty)$ on \mathfrak{S}^z . Let $k_j \geq 2$. Now consider the element $\mathbf{1}[\{k_j\}] \in \mathcal{S}(\infty, \infty)$ consisting of

1) arcs $i_l^+ \downarrow i_l^-$ [0], $i_r^+ \downarrow i_r^-$ [0] for $i \in \mathbb{N}$ (this collection of arcs defines the unit in $\mathcal{S}(\infty,\infty);$

2) cycles $\bigcirc [k_i]$.

Such elements form the center of the semigroup $\mathcal{S}(\infty,\infty)$.

Let $\mathfrak{u} \in \mathfrak{S}^z$, it is a collection $\{\overline{U_\nu}\}\$ of tables equipped with guests. Consider the set $\Gamma^{\diamond}[\{k_i\}]$ of framed checker surfaces whose A-verices have orders k_j . Fix $(\Sigma, \omega) \in \Gamma^{\circ}[\{k_i\}]$. For each vertex B_{β} denote by $Y(B_{\beta})$ the set of all white triangles containing B_β . For each B_β consider a random map $\eta_\beta: Y(B_\beta) \to U_{\omega(B_\beta)}$ reversing

32

the cyclic order. For each β we equip the set of all η_{β} with the uniform measure $d\eta_{\beta}$, assuming that the total measure is $\ell_{\omega(B)}^{\text{ord }B_{\beta}}$ $\frac{\partial a}{\partial \omega(B)}$ /(# B_β – 1)!. Let Q be a black triangle containing B_β . Let T be its white predecessor according to the cyclic order, T' be the white follower. We assign to Q the arc $\mathcal{J}(Q) := [\eta_{\beta}(T), \eta_{\beta}(T')]$ of the table $U_{\omega(B_\beta)}$ together with guests. Notice that the ends of such arcs are not guests a.s. Define a new collection $\{\zeta_\gamma\} = \{\zeta_\gamma(\{\eta_\beta\})\}$ of occupied tables in the following way: take a vertex C_{γ} , write all black triangles Q_{s_1}, Q_{s_2}, \ldots containing C_{γ} in the counterclockwise order. Then ζ_{γ} is a cyclically ordered set

$$
\zeta_{\beta} := \mathcal{J}(Q_{s_1}) \sqcup \mathcal{J}(Q_{s_2}) \sqcup \ldots
$$

equipped with a natural length. We also have a pushforward $\zeta_*(d\eta)$ of the measure $d\eta = d\eta_1 d\eta_2 \ldots$, this is a new measure on a collection of occupied tables.

Theorem 4.2. *The image of a point* $\mathbf{u} \in \mathfrak{S}^z$ *under the polymorphism* $\mathbf{1}[\{k_i\}]$ *is*

$$
\prod_{j} k_{j}! \prod_{m \geq 2} \iota_{m}[\{k_{j}\}] \sum_{(\Sigma, \omega) \in \Gamma^{\circ}[\{k_{j}\}] } \left(\delta_{\mathbb{R}^{\circ}} \left[\left[z^{\# \{C_{\gamma}\} - \# \{B_{\gamma}\} } \right] \right] \times \text{Replace} \left[\{\overline{U_{\omega(B_{\beta})}} \} \longrightarrow \{\zeta_{*}(d\eta)\} \right] \{\overline{U_{\nu}}\} \right)
$$

To evaluate this image we apply Remark after Proposition [1.2.](#page-10-1) We take the second copy N of N consisting of points $\underline{1}, \underline{2}, \ldots$ Let regard \overline{S}_{∞} as the group $\overline{S}(\mathbb{N}\sqcup\mathbb{N})$ of all permutations of $\mathbb{N} \sqcup \underline{\mathbb{N}}$ and S as a subgroup in $\overline{S}(\mathbb{N} \sqcup \underline{\mathbb{N}}) \times \overline{S}(\mathbb{N} \sqcup \underline{\mathbb{N}})$. Now we can realize $\mathcal{S}(\infty, \infty)$ as the semigroup of double cosets $\overline{S(\mathbb{N})\backslash \mathbb{S}/\overline{S}(\mathbb{N})}$

Let $\sum k_j = n$. Take $g \in \mathbb{S}$ that is trivial on on the left copy of $\mathbb{N} \sqcup \mathbb{N}$, on right copy of N and on the set $\{n+1, n+2, ...\}$. Let also g induces a permutation of the set $\{\underline{1}, \ldots, \underline{n}\}\$ with cycles of lengths k_i .

Further considerations as the same as above. We take an analog of λ_0^n , it corresponds to the chip $\frac{1}{l'}[1/2] \sim \frac{1}{l'}[1/2], \ldots, \frac{n}{l'}[1/2] \sim \frac{n}{l'}[1/2]$ and other arcs are strictly vertical lines of weight 0. It corresponds to putting additional guests $\overline{1}, \ldots$, \overline{n} to the occupied restaurant μ .

We fall to the situation of Subsect. [3.10,](#page-26-2) where the set N is replaced by N ⊔ $\{\overline{1}, \ldots, n\}$ and $g \in S_n$ acts nontrivially only on additional random guests $\overline{1}, \ldots$, \overline{n} . These 'numbers' also are labels $\underline{1}, \ldots, \underline{n}$ on white triangles. We apply the construction of Subsect. [3.10](#page-26-2) and forget random guests.

REFERENCES

- [1] Aldous, D. J. Exchangeability and related topics. École d'été de probabilités de Saint-Flour, XIII–1983, 1-198, Lecture Notes in Math., 1117, Springer, Berlin, 1985.
- [2] Bogachev, V. I. Measure theory. Vol. I, II. Springer-Verlag, Berlin, 2007
- [3] Borodin, A.; Olshanski, G. Point processes and the infinite symmetric group. Math. Res. Lett. 5 (1998), no. 6, 799-816.
- [4] Borodin, A.; Olshanski, G. Representations of the infinite symmetric group. Cambridge University Press, Cambridge, 2017.
- [5] Brauer, R. On algebras which are connected with the semisimple continuous groups. Ann. of Math. (2) 38 (1937), no. 4, 857-872.
- [6] Ewens, W. J. The sampling theory of selectively neutral alleles, Theor. Pop. Biol. 3 (1972), 87-112.
- [7] Goulden, I. P.; Jackson, D. M. The combinatorial relationship between trees, cacti and certain connection coefficients for the symmetric group. European J. Combin. 13 (1992), no. 5, 357- 365.
- [8] Hopf, E. The general temporally discrete Markoff process, J. Rational Mech. Anal., 3 (1954), 13-45.
- [9] Howe, R. E.; Moore, C. C. Asymptotic properties of unitary representations. J. Functional Analysis 32 (1979), no. 1, 72-96.
- [10] Janvresse, E.; Prikhod'ko, A. A.; de la Rue, T.; Ryzhikov, V. V., Weak limits of powers of Chacon's automorphism, Ergodic Theory Dynam. Systems, 35:1 (2015), 128-141.
- [11] Kerov, S. V. Realizations of representations of the Brauer semigroup. (Russian) Zap. Nauchn. Sem. LOMI, 164 (1987), 188-193, 199; English transl. in J. Soviet Math. 47 (1989), no. 2, 2503-2507
- [12] Kerov, S.; Olshanski, G.; Vershik, A. Harmonic analysis on the infinite symmetric group. A deformation of the regular representation. C. R. Acad. Sci. Paris Sér. I Math. 316 (1993), 8, 773-778.
- [13] Kerov, S.; Olshanski, G.; Vershik, A. Harmonic analysis on the infinite symmetric group. Invent. Math. 158 (2004), no. 3, 551-642.
- [14] Kerov, S. V.; Tsilevich, N. V., Stick breaking process generated by virtual permutations with Ewens distribution (Russian) Zap. Nauchn. Sem. POMI, 223, 1995, 162-180; English transl.: J. Math. Sci. (New York), 87:6 (1997), 4082-4093.
- [15] King, J. The commutant is the weak closure of the powers, for rank-1 transformations, Ergodic Theory Dynam. Systems, 6:3 (1986), 363-384.
- [16] Kingman, J. F. C. Poisson processes. Oxford: Oxford University Press, 1993.
- [17] Krengel, U. Ergodic theorems. Walter de Gruyter, Berlin, 1985.
- [18] Kushnir A.Yu., Ryzhikov V.V., Weak closures of ergodic actions, Math. Notes, 101:2 (2017), 277-283.
- [19] Lando, S. K.; Zvonkin, A. K. Graphs on surfaces and their applications. Springer-Verlag, Berlin, 2004.
- [20] Melleray, J.; Tsankov, T. Generic representations of abelian groups and extreme amenability. Israel J. Math. 198 (2013), no. 1, 129-167.
- [21] Méliot, P. L. Representation theory of symmetric groups. CRC Press, Boca Raton, FL, 2017.
- [22] Nelson, E. The free Markoff field, J. Funct. Anal. 12 (1973) 211-227.
- [23] Neretin, Yu. A. Categories of bistochastic measures and representations of some infinitedimensional groups. Russian Acad. Sci. Sb. Math. 75 (1993), 1, 197-219.
- [24] Neretin, Yu. A. Categories of symmetries and infinite-dimensional groups. Lond. Math. Soc. Monogr., New Ser. 16. Oxford: Oxford Univ. Press 1996.
- [25] Neretin, Yu. A. Spreading maps (polymorphisms), symmetries of Poisson processes, and matching summation. Zap. Nauchn. Sem. POMI 292 (2002), 62-91; reprinted in J. Math. Sci. (N.Y.) 126 (2005), 2, 1077-1094.
- [26] Neretin, Yu. A. Multi-operator colligations and multivariate characteristic functions. Anal. Math. Phys. 1 (2011), no. 2-3, 121-138.
- [27] Neretin, Yu. A. Sphericity and multiplication of double cosets for infinite-dimensional classical groups. Funct. Anal. Appl. 45 (2011), 3, 225-239.
- Neretin, Yu. Symmetries of Gaussian measures and operator colligations. J. Funct. Anal. 263 (2012), 3, 782-802.
- [29] Neretin, Yu. Infinite tri-symmetric group, multiplication of double cosets, and checker topological field theories. Int. Math. Res. Not. IMRN 2012, 3, 501-523.
- [30] Neretin, Yu. A. On the boundary of the group of transformations that leave a measure quasiinvariant. Sb. Math. 204 (2013), 7-8, 1161-1194.
- [31] Neretin, Yu. A. Infinite symmetric groups and combinatorial constructions of topological field theory type. Russian Math. Surveys 70 (2015), 4, 715-773.
- [32] Neretin, Yu. A. Infinite-dimensional p-adic groups, semigroups of double cosets, and inner functions on Bruhat-Tits buildings, Izv. Math., 79:3 (2015), 512-553.
- [33] Neretin, Yu. A. Wishart-Pickrell distributions and closures of group actions. Zap. Nauchn. Sem. POMI 448 (2016), 236-245; English translal. in J. Math. Sci. (N.Y.) 224 (2017), 2, 328-334.
- [34] Neretin, Yu. A. On the group of infinite p-adic matrices with integer elements, Zap. Nauchn. Sem. POMI, 468 (2018), 105-125; English transl. in J. Math. Sci. (N. Y.), 240:5 (2019), 572-586.
- [35] Neretin, Yu. A. Groups $GL(\infty)$ over finite fields and multiplications of double cosets. J. Algebra 585 (2021), 370-421.
- [36] Neveu, J. Bases mathématiques du calcul des probabilités. (French) Masson et Cie, Éditeurs, Paris, 1964.
- [37] Okounkov, A. Yu. Thoma's theorem and representations of infinite bisymmetric group. Funct. Anal. Appl. 28 (1994), 101-107.
- [38] Olshanski, G. I. Unitary representations of (G, K) -pairs that are connected with the infinite symmetric group $S(\infty)$, Leningrad Math. J., 1 (1990), 4, 983-1014.
- [39] Olshanski, G. I. Unitary representations of infinite-dimensional pair (G, K) and the formalisms of R. Howe. In: Representations of Lie groups an related topics, 264-465, Gordon and Breach, NY, 1991.
- [40] Olshanski, G. I. On semigroups related to infinite-dimensional groups. In Topics in representation theory, 67-101, Amer. Math. Soc., Providence, RI, 1991.
- [41] Phillips, P. C. B. The characteristic function of the Dirichlet and multivariate F distributions. Cowles Foundation for Research in Economics, 1988 (Discussion Paper 865), 1–17.
- [42] Pitman, J. Combinatorial stochastic processes. Lectures from the 32nd Summer School on Probability Theory held in Saint-Flour, July 7-24, 2002. Lecture Notes in Mathematics, 1875. Springer-Verlag, Berlin, 2006.
- [43] Prudnikov, A. P.; Brychkov, Yu. A.; Marichev, O. I. Integrals and series. Vol. 1. Elementary functions. Gordon & Breach Science Publishers, New York, 1986.
- [44] Rohlin, V. A. On the fundamental ideas of measure theory. (Russian) Mat. Sbornik N.S. 25 (67), (1949), 107-150; English transl. Amer. Math. Soc. Translation (1952), no. 71.
- [45] Shabat, G. Calculating and drawing Belyi pairs, Zap. Nauchn. Sem. POMI, 446 (2016), 182- 220; J. Math. Sci. (N. Y.), 226 (2017), 5, 667-693.
- [46] Solecki, S. Closed subgroups generated by generic measure automorphisms. Ergodic Theory Dynam. Systems 34 (2014), no. 3, 1011-1017.
- [47] Thoma, E. Die unzerlegbaren, positive-definiten Klassenfunktionen der abzählbar unendlichen, symmetrischen Gruppe. Math. Z. 85 (1964), 40-61 .
- [48] Tsilevich, N. V. Distribution of cycle lengths of infinite permutations. (Russian) Zap. Nauchn. Sem. POMI 223 (1995), 148-161, 339; English transl. in J. Math. Sci. (New York) 87 (1997), no. 6, 4072-4081
- [49] Vershik, A. M. Multivalued measure-preserving mappings (polymorphisms) and Markov operators, Zap. Nauchn. Sem. LOMI 72 (1977), 26-61; English transl. in J. Soviet Math., 23:3 (1983), 2243-2266.
- [50] Vershik, A. M.; Kerov, S. V. Characters and factor representations of the infinite symmetric group. Soviet Math. Dokl. 23 (1981), no. 2, 389-392.
- [51] Wassermann, A. J. Automorphic actions of compact groups on operator algebras. Thesis (Ph.D.), University of Pennsylvania, 1981, 176 pp.
- [52] Wilks, S. S. Mathematical statistics. John Wiley & Sons, Inc., New York-London, 1962.

Fakultät für Mathematik, Universität Wien; Institute for Information Transmission Problems, Moscow; Dept. of Mechanics and Mathematics, Moscow State University. e-mail:yurii.neretin@univie.ac.at URL: https://mat.univie.ac.at/ neretin