

# Succinct Amyloid and Non-Amyloid Patterns in Hexapeptides

László Keresztes<sup>a,\*\*</sup>, Evelin Szögi<sup>a,\*\*</sup>, Bálint Varga<sup>a</sup>, Viktor Farkas<sup>c</sup>, András Perczel<sup>c,d</sup>, Vince Grolmusz<sup>a,b,\*</sup>

<sup>a</sup>*PIT Bioinformatics Group, Eötvös University, H-1117 Budapest, Hungary*

<sup>b</sup>*Uratim Ltd., H-1118 Budapest, Hungary*

<sup>c</sup>*MTA-ELTE Protein Modeling Research Group, H-1117 Budapest, Hungary*

<sup>d</sup>*Laboratory of Structural Chemistry and Biology, Eötvös University, H-1117, Budapest, Hungary*

---

## Abstract

Hexapeptides are widely applied as a model system for studying amyloid-forming properties of polypeptides, including proteins. Recently, large experimental databases have become publicly available with amyloidogenic labels. Using these datasets for training and testing purposes, one may build artificial intelligence (AI)-based classifiers for predicting the amyloid state of peptides. In our previous work (Biomolecules, 11(4) 500, (2021)) we described the Support Vector Machine (SVM)-based Budapest Amyloid Predictor (<https://pitgroup.org/bap>). Here we apply the Budapest Amyloid Predictor for discovering numerous amyloidogenic and non-amyloidogenic hexapeptide patterns with accuracy between 80% and 84%, as surprising and succinct novel rules for further understanding the amyloid state of peptides. For example, we have shown that for any independently mutated residue (position marked by “x”), the patterns CxFLWx, FxFLFx, or xxIVIV are predicted to be amyloidogenic, while those of PxDxxx, xxKxEx, and xxPQxx non-amyloidogenic at all. We note that each amyloidogenic pattern with two x’s (e.g., CxFLWx) describes succinctly  $20^2 = 400$  hexapeptides, while the non-amyloidogenic patterns comprising four point mutations (e.g., PxDxxx) gives  $20^4 = 160,000$  hexapeptides in total. We also examine restricted substitutions for positions “x” from subclasses of proteinogenic amino acid residues; for example, if “x” is substituted with hydrophobic amino acids, then there exist patterns containing three x’s, like MxVVxx, predicted to be amyloidogenic. If we can choose for the x positions any hydrophobic amino acids, except the “structure breaker” proline, then we get amyloid patterns with five x positions, e.g., xxxFxx, each corresponding to 32,768 hexapeptides. To our knowledge, no similar applications of artificial intelligence tools or succinct amyloid patterns were described before the present work.

---

\*Corresponding author

\*\*Joint first authors

---

## Introduction

The amyloid formation of proteins and peptides has gained increasing attention in novel areas of medicine and biology in the last months, including the application of amyloidogenic aggregation cores in viral proteins as new antiviral agents [1], or targeting the lethal transthyretin amyloidosis with human *in vivo* CRISPR-Cas9-based gene editing with high success rate [2].

Amyloids are misfolded proteins [3, 4] with well-defined and periodic 3D structure, comprising mostly parallel  $\beta$ -sheets [5, 6]. While amyloids are only seldom present in healthy human tissues [7], they were reported to be connected with several neurodegenerative diseases [8], most importantly with Alzheimer’s disease.

In bioinformatics numerous amyloid-predictors were designed and published in the recent years, including APPNN [9], Zygggregator [10], AGGRESCAN [11], netCSSP [12]; for a recent review of their performance we refer to [13]. These predictors use different machine learning approaches for decision making from training data. In our contribution [14] we applied a powerful but transparent machine learning tool, the linear Support Vector Machine (SVM) [15].

For building an SVM, we needed a training dataset of  $n$ -dimensional vectors  $y^1, y^2, \dots, y^m$ , each labeled with a bit of either 0 or 1. From the training set, we determine a hyperplane, which, in a certain sense, “optimally” separates the 0 and the 1-labeled vectors in a way that most of the 1-labeled vectors are on one side of the hyperplane, and most of the 0-labeled ones are on the other side.

The SVM-predictor now works as follows: If a new vector is situated on the 0-side, then the prediction is “0”, and if it lies on the 1-side, the prediction is “1”.

Recently we built a Support Vector Machine for amyloidogenicity prediction of hexapeptides [14]. The training set was the Waltz dataset [16, 17] of experimentally identified 514 amyloidogenic and 901 non-amyloidogenic hexapeptides. By applying the physico-chemical property dataset of amino acids, AAindex [18], property vectors  $y^1, y^2, \dots, y^m$  were assigned to each hexapeptide, and using this multi-dimensional representation, we have prepared an SVM, called Budapest Amyloid Predictor (abbreviated as BAP), described in detail in [14], and freely available at the address <https://pitgroup.org/bap>.

The accuracy rate of our predictor is 84%, more exactly, ACC = 0.84, TPR=0.75, TNR=0.9, PPV=0.8, NPV=0.86, (that is, accuracy, true positive ratio, true negative ratio, positive predictive value, negative predictive value, resp.). We remark that the accuracy of our SVM is better than or on par with that of APPNN [9], with a simpler, more transparent structure.

In the present contribution, we make use of this transparent structure of the Budapest Amyloid Predictor: we present numerous patterns related to amyloidicity, such that each of those patterns grasps hundreds or even tens of thousands of individual hexapeptides and give predictions of their amyloid-forming properties. For example, we show that for all (independent) substitutions of the

20 amino acids for letter “x”, the hexapeptides CxFLFx, FxFLWx, or xxIVIV are all predicted to be amyloids by the Budapest Amyloid Predictor. Note that each of these patterns describes  $20^2 = 400$  different hexapeptides. We also note that no amyloid-forming patterns exist with three x’s for the predictor. All the 5531 amyloid-forming hexapeptide patterns with two x’s are listed in Table S1 in the Supporting Material.

We also show several patterns, which – by the Budapest Amyloid Predictor – would not form amyloids. For example, the patterns xxDDxx, xxPx Dx, or xxPKxx with any (independently chosen) amino acids for the positions denoted by x are predicted to be non-amyloids by our Budapest Amyloid Predictor. Note that each of these patterns succinctly describes  $20^4 = 160,000$  hexapeptides. We add that non-amyloid patterns with five  $x$  positions do not exist for our tool at <https://pitgroup.org/bap>. All the non-amyloid forming hexapeptide patterns with four  $x$  positions are listed in Table 2 of the present contribution.

We note that these patterns are succinct representations of the predictions of the Budapest Amyloid Predictor (BAP), whose accuracy rate is 84% [14]; that is, we do not state, for example, that all CxFLFx hexapeptides are amyloids, but we state that all of them are predicted to be amyloids by the BAP tool. The transparent linear structure of the Support Vector Machines makes possible the derivation of these intuitive, useful, and well-applicable patterns from an artificial intelligence (AI) tool, as we clarify in the present contribution.

We need to add also that today we are living in the era of the fast-developing AI methods and tools in numerous fields of science and technology. Most of these tools work as follows:

Suppose the tool needs to compute a value  $f(y)$  from another value  $y$ . For constructing such an AI tool, the following steps are applied:

- A large set of previously acquired, correct  $(y, f(y))$  pairs are partitioned into two classes: the training set A and the test set B.
- The training set A is applied to construct a tool, which assigns the predicted value of  $f(y)$ , denoted by  $f_p(y)$ , to each  $y$ ;
- The test set B is used for evaluating the correctness of the tool: the predicted value, produced by the tool,  $f_p(y)$  is compared to the correct, previously known  $f(y)$ .

The AI tool is deemed “good” if it is correct for a large enough portion of the test values.

In general, however, it is difficult to get an insight into the intrinsic decision mechanisms of a typical AI tool; this is especially true for the deep neural networks, which are applied widely today.

In the case of linear Support Vector Machines [15], the decision mechanism is much more transparent, and one can exploit a highly correct SVM for gaining unprecedented scientific information in certain cases [19]. In the present contribution, we show a novel and original method for gaining site-specific amyloid-forming properties of amino acids in hexapeptides and prepar-

ing amyloid-forming and non-amyloid forming patterns for the succinct representation of the SVM prediction results for hundreds (cf. Table S1) or even tens of thousands (cf. Table 2) hexapeptides at the same time.

## Methods

We have introduced the Budapest Amyloid Predictor webserver in the work [14] by applying linear Support Vector Machines as the underlying prediction tool [15], and the Waltz dataset [16, 17] for training and testing purposes. The Waltz dataset consists of 1415 hexapeptides, from which 514 peptides are experimentally labeled as “amyloidogenic” and 901 hexapeptides as “non-amyloidogenic”. The Budapest Amyloid Predictor <https://pitgroup.org/bap> was constructed as follows:

- (i) Each amino acid from the 20 proteinogenic ones was characterized by a 553-dimensional vector, corresponding to its physico-chemical properties published in AAindex [18]. Therefore, a hexapeptide was represented by a length  $6 \times 553 = 3318$  vector  $z$ . We note that this highly redundant representation has given somewhat better predictions than more concise ones [14] and has not caused any difficulties in what follows.
- (ii) By applying a quadratic programming algorithm for SVM computation from the SciKit-learn Python library [20], we have computed a vector  $w$  and a scalar  $b$  such that if the sign of  $w \cdot z + b$  is positive, then the prediction is “amyloidogenic”, otherwise it is “non-amyloidogenic”, with 84% accuracy, for any vector  $z$ , representing a hexapeptide.
- (iii) One can write the dot product  $w \cdot z$ , with  $\ell = 553$ , as

$$w \cdot z = \sum_{i=1}^{6\ell} w_i z_i = \sum_{j=1}^6 \sum_{i=(j-1)\ell+1}^{j\ell} w_i z_i. \quad (1)$$

For any given  $j = 1, 2, \dots, 6$  the  $\ell$   $z_i$ 's are determined by the amino acid at position  $j$  at the hexapeptide. This means that we have only  $6 \times 20 = 120$  second sums in (1) (for six positions and 20 amino acids), and these 120 values can be pre-computed. Table 1 of [14] lists these pre-computed values. Since we need the same table in the present work, we include it also as Table 1 here.

- (iv) Table 1 makes it possible to decide if a hexapeptide is predicted to be amyloidogenic or not, by “hand”; for example, to decide if IVIVIV is amyloidogenic or not, we need to add up the numbers, corresponding to I in the first, to V in the second, to I in the third, to V in the fourth, to I in the fifth and to V in the sixth column, that is

$$-0.06 - 0.14 + 0.26 + 0.14 - 0.06 + 0.01 = 0.15$$

and we need to add to this  $w \cdot z$  value the scalar  $b = 1.083$ , which equals to 1.233, which is a positive number, so IVIVIV is predicted to be amyloidogenic.

	1	2	3	4	5	6
A	-0.26	-0.32	-0.27	-0.14	-0.43	-0.22
R	-0.45	-0.41	-0.46	-0.33	-0.52	-0.35
N	-0.40	-0.34	-0.49	-0.27	-0.46	-0.30
D	-0.49	-0.43	-0.56	-0.41	-0.56	-0.36
C	-0.09	-0.21	0.03	-0.05	-0.17	-0.05
Q	-0.37	-0.30	-0.36	-0.34	-0.48	-0.32
E	-0.51	-0.41	-0.43	-0.30	-0.61	-0.39
G	-0.23	-0.37	-0.46	-0.37	-0.30	-0.33
H	-0.32	-0.26	-0.26	-0.30	-0.35	-0.25
I	-0.06	-0.08	0.26	0.09	-0.06	-0.07
L	-0.10	-0.18	0.02	0.04	-0.22	-0.13
K	-0.39	-0.45	-0.51	-0.35	-0.59	-0.32
M	-0.17	-0.25	-0.02	-0.10	-0.19	-0.18
F	-0.13	-0.11	0.05	-0.03	-0.13	-0.11
P	-0.56	-0.38	-0.56	-0.51	-0.42	-0.45
S	-0.37	-0.35	-0.41	-0.30	-0.48	-0.23
T	-0.34	-0.33	-0.28	-0.23	-0.40	-0.23
W	-0.17	-0.17	-0.09	-0.06	-0.12	-0.16
Y	-0.23	-0.11	-0.13	-0.06	-0.18	-0.15
V	-0.05	-0.14	0.19	0.14	-0.19	0.01

Table 1: The Amyloid Effect Matrix, constructed from the pre-computed values from equation (1). The rows are corresponding to the amino acids, the columns to the positions. The larger numbers show stronger amyloidogenic properties in the given position. Source: [14]. In [14], by ordering the columns of this table, a position-dependent amyloidogenicity order of amino acids are given in a subsequent table.

We refer to Table 1 as the Amyloid Effect Matrix.

As we have demonstrated in paragraph (iv) above, one can simply make the prediction of the SVM by using the values solely from this matrix.

From now on, we would like to exploit the Amyloid Effect Matrix for finding succinct descriptions of amyloidogenic and non-amyloidogenic patterns among the 64 million possible hexapeptides.

### *Patterns of Amyloidicity*

Here we would like to find very characteristic positions and substitutions, which already assure us that all the hexapeptides fitting those patterns are homogeneously either amyloidogenic or non-amyloidogenic. Let us see an example:

**Example 1.** Let us fix the amino acid proline ( $P$  stands for) at positions 3 and 4 and leave all the other four positions free: let us consider the pattern

$$xxPPxx.$$

We state that for all (independent) substitutions for  $x$ 's, the Budapest Amyloid Predictor (abbreviated as BAP) says that the hexapeptide is NOT amyloid. Since we have four  $x$ 's, the pattern  $xxPPxx$  describes exactly  $20^4 = 160000$  hexapeptides; so we state that not one of these 160 000 hexapeptides is predicted to be amyloidogenic.

It is very easy to verify this statement from Table 1. The values corresponding to  $P$ 's in the third and in the fourth positions (-0.56 and -0.51) add up to -1.07. Now, even if we take the largest values of columns 1, 2, 5, and 6, that is, -0.05, -0.08, -0.06, and 0.01, respectively, their sum is -1.25, and adding  $b = 1.083$  to this value, we would still have a result to be a negative number. That is, even the largest values from columns 1,2,5, and 6 could not overweight the large negative sum of -1.07 of the two consecutive proline residues in positions 3 and 4. This means that all hexapeptides, fitting to the pattern of  $xxPPxx$ , are predicted to be non-amyloids by BAP.

**Example 2.** Similarly, one can also find amyloid patterns. For example, we state that all the 400 ( $= 20 \times 20$ ) hexapeptides, fitting to the pattern  $FxFLWx$  are predicted to be amyloids. One can easily verify this statement from Table 1. The  $F$  in position 1 adds -0.13, in position 3 adds 0.05,  $L$  in position 4 adds 0.04, and  $W$  in position 5 contributes -0.12, their sum is -0.16. Now, if we take the smallest values from columns 2 and 6, that is, -0.45 and -0.45, and add  $b = 1.083$  to their sum, we will get  $-0.16 - 0.45 - 0.45 + 1.083 = 0.023$ , that is, a positive number, so independently from the choice of the  $x$ 's,  $FxFLWx$  is predicted to be an amyloid-forming hexapeptide.

#### *Minimal patterns*

In what follows, we will find all the minimal patterns of amyloidicity and non-amyloidicity. These minimal patterns are the most concise representations of the amyloid-forming rules of the BAP predictor.

Here the "minimal" word means that we cannot decrease the number of the fixed amino acids without invalidating the rule. Our goal is to find the patterns with the minimum number of amino acids fixed. From such minimal patterns, one can easily generate valid but non-minimal ones; for example, the  $xxPPxx$  pattern is predicted to be non-amyloidogenic for any substitutions of  $x$ 's. Therefore, e.g.,  $WxPPxx$ , or  $VIPPxx$  are also non-amyloid patterns for any substitutions for  $x$ , but they are not minimal. It is easy to see by observing Table 1 that neither  $xxxPxx$ , nor  $xxPxxx$  are valid non-amyloid patterns, so  $xxPPxx$  is a minimal pattern.

#### *Finding all minimal patterns*

Our goal is to find every hexapeptide pattern, both the amyloidogenic and the non-amyloidogenic ones, as predicted by BAP.

Finding these patterns is straightforward using the Amyloid Effect Matrix (Table 1). Suppose that we intend to generate the minimal amyloid indicating patterns. Finding the non-amyloid patterns is a similar procedure.

Verifying whether a pattern is a valid amyloid indicator is easy: we need to generalize the steps done in the examples: we substitute the minimal amyloid effective amino acids on the free positions (denoted by  $s$ ), and check its score. If the score is already positive, then this least amyloidogenic hexapeptide is already amyloid, then every other hexapeptide from this space is amyloid too.

Finding the rules for hexapeptides could be done by exhaustive search. Let say we want to find all the rules with  $k$  fixed amino acids, where  $k$  is between 1 and 6. In what follows, we call the core of the rule is the number of fixed amino acids (e.g., the core of rule xxPPxx is 2). The positions  $s$  will be referred to as free positions.

For finding all the rules with core  $k$ , our approach is

- (i) generating all the  $\binom{6}{k}$  index subsets;
- (ii) for each index subset, we generate all the  $20^k$  rule candidates by assigning all the possible amino acids to the  $k$  core positions;
- (iii) verify the validity of the pattern by using check each of them as already described;

We remark that this exact exhaustive search is not fast computationally, but it perfectly works for hexapeptides: The number of verifications is

$$\sum_{k=1}^{k=6} \binom{6}{k} 20^k = 21^6 - 1$$

less than 86 million, its running time is several hours in today’s low-end computers.

The amyloid patterns are listed in Table S1 in the Supporting Material, while the non-amyloid patterns are in Table 2 in the next section.

#### *The case of restricted amino acid classes*

Amino acids are frequently characterized and classified by their chemical properties, like polarity, non-polarity, hydrophobicity, hydrophilicity, etc. If we want to find patterns of amyloidicity when for the free positions, denoted by  $x$ , one can choose substitutions only from a given restricted class, then one can have stronger, more specific patterns than in the general case, when  $x$  can be substituted by any of the 20 amino acids.

Finding those patterns in the restricted classes can be done analogously to the general case: the minimum values of Table 1 from the given class need to be considered.

### Statistical analysis

We refer to the work [14] for the statistical accuracy estimations of the Budapest Amyloid Predictor. There we have shown that the predictor has ACC = 0.84, TPR=0.75, TNR=0.9, PPV=0.8, NPV=0.86 (that is, accuracy, true positive ratio, true negative ratio, positive predictive value, negative predictive value, resp.) Figure 1 of [14] also gives ROC (Receiver Operating Characteristics) curve of the tool, with AUC (Area Under Curve) value is 0.89.

### Discussion and Results

Figure 1 visualizes the substitutions into a non-amyloid and an amyloid-pattern.

We have successfully identified all the 5531 minimal amyloid patterns (Table S1) using BAP. For example, xxIYCI, IFIYxx, or CxVVxC are amyloid patterns from Table S1.




Figure 1: Examples for amyloid- and non-amyloid patterns.

We have found that almost all amyloidogenic hexapeptide patterns contain valine (V) and/or isoleucine (I) residues, both of branched and hydrophobic



PxPxxx	PxDxxx	xxPPxx	xxPDxx	xxPGxx	xxPKxx
xxPQxx	xxDPxx	xxDDxx	xxDGxx	xxDKxx	xxDQxx
xxKPxx	xxKDxx	xxNPxx	xxGPxx	xxRPxx	xxPxEx
xxPxKx	xxPxDx	xxDxEx	xxDxKx	xxDxDx	xxKxEx

Table 2: The list of all non-amyloid patterns with four free positions; it contains 24 patterns. Note that each pattern describes  $20^4 = 160000$  hexapeptides succinctly, all of which are predicted to be non-amyloids by the Budapest Amyloid Predictor [14]. From the 24 patterns, only nine do not contain proline in a fixed position. Note also the possibly stabilizing salt-bridge forming [21] amino acid pairs (E-K, D-K) in several patterns without the fixed proline.

sidechains. Among the 5531 patterns identified (Table S1) only 16 patterns are free of both V and I:

CxFLWx, CxFLFx, CxCLWx, CxCLFx, CxLLWx, CxLLFx, LxFLWx, LxFLFx, LxCLWx, LxCLFx, LxLLWx, LxLLFx, FxFLWx, FxFLFx, FxCLWx, CxFLCx.

Furthermore leucine, L, the third branched and hydrophobic side chains is represented one or more times in the above listed 15 patterns. In conclusion, V, I and L residues make hexapeptides intrinsically amyloidogenic.

We remark that no amyloid pattern with three free positions (i.e., x) exists.

Table 2 lists the 24 non-amyloid patterns, each with four free positions. No other non-amyloid patterns exist with four free positions, and no non-amyloid pattern exists with five free positions.

It is interesting to note that from the 24 non-amyloid patterns of Table 2, 15 contain proline in one or two fixed positions. From the 9 remainders, 4 contain E-K or D-K pairs, which can form possibly stabilizing salt bridges [21].

### Results for amino acid subsets

In this subsection, we find amyloid patterns when the  $x$  positions can be substituted only by the members of some specific amino acid classes. The amino acid classes we examine are small non-polar, hydrophobic, and polar amino acids, as classified by [22], and listed in the second column of Table 3.

Class name	Class elements listing	No. of free positions	No. of patterns
small non-polar	GAST	3	411
hydrophobic	CVLIMPFYW	3	43
polar	DENQHKR	3	4
hydrophobic-{P}	CVLIMFYW	5	38
amino acids-{P}	QFYESNCDMLIAHGWRKVT	3	4

Table 3: The amino acid subsets examined. The classification of the residues in the first three rows are by [22]. The last two rows correspond to the classes where we left out proline, a well-known structure-breaker from the hydrophobic set or from all the amino acids. The third column shows the number of free positions we get in the special substitutions, and the fourth column shows the number of patterns found for these special substitutions for “x”.

When the substitutions to the free positions, denoted by  $x$ , can be done only from special subsets, listed in Table 3, we can get amyloid-rules with 3 free positions, in contrast with the unrestricted case, when our rules have 2 free positions (Table S1).

When  $x$  is allowed to be substituted from the small non-polar set, then 411 patterns can be found with three free positions, for example, VIIxxx IIIxxx VxIVxx VxIIxx VxILxx VxIFxx VxICxx VxIWxx VxVVxx VxVIxx. All the existing 411 patterns are listed in Table S2 in the supporting material. No such pattern exists with 4 free positions.

If  $x$  is chosen from the hydrophobic set, then 43 patterns exist with three free positions, listed in Table 4. No such pattern exists with 4 free positions.

VxIVxx	VxIIxx	VxILxx	VxIFxx	VxVVxx	VxVIxx	VxVLxx	IxIVxx
IxIIxx	IxILxx	IxVVxx	IxVIxx	CxIVxx	CxIIxx	CxILxx	CxVVxx
CxVIxx	LxIVxx	LxIIxx	LxILxx	LxVVxx	LxVIxx	FxIVxx	FxIIxx
FxVVxx	MxIVxx	MxIIxx	MxVVxx	WxIVxx	WxIIxx	GxIVxx	YxIVxx
xxIVIx	xxIVxV	xxIVxC	xxIVxI	xxIVxF	xxIIxV	xxIIxC	xxILxV
xxVVxV	xxVVxC	xxVIxV					

Table 4: The list of all the 43 amyloidogenic patterns with three free positions when  $x$  is hydrophobic, i.e., chosen from CVLIMPFYW. Each pattern describes  $9^3 = 729$  hexapeptides.

When  $x$  is chosen from polar amino acids, then the only four patterns with three free positions are xxIVIx xxIVWx xxIIIx xxVVIx.

We note that no pattern exists in these three cases without V and I amino acids; that is, all of the patterns in these three restricted substitutions contain either valine or isoleucine in fixed positions.

If proline is not allowed to be substituted for any  $x$ , but otherwise the remaining 19 amino acids can be chosen for the  $x$  positions, then we have exactly four amyloid patterns with three  $x$  positions: xxIVIx xxIVWx xxIIIx xxVVIx; note that without the restriction to proline, no amyloid pattern exists with three free positions.

These four patterns are exactly the same as in the case of polar residue substitutions, but the set of hexapeptides they represent differ: in the case of polar substitutions, each of the four patterns represent  $7^3 = 343$  hexapeptides, while for the non-proline substitutions  $19^3 = 6859$  hexamers.

If  $x$  could be chosen from hydrophobic amino acids, except proline, the “structure breaker” then we have the “largest” patterns of amyloidicity: 38 patterns exist with just one fixed position, listed in Table 5. Note that each of those patterns describes  $8^5 = 32768$  hexapeptides, such that all of them are predicted to be amyloidogenic.

## Conclusions

Here, we established patterns of amyloidicity and non-amyloidicity in the case of hexapeptides, based on a Support Vector Machine-based predictor, avail-

Vxxxxx	Ixxxxx	Cxxxxx	Lxxxxx	Fxxxxx	Mxxxxx	Wxxxxx	xIxxxx
xFxxxx	xYxxxx	xVxxxx	xWxxxx	xLxxxx	xCxxxx	xxIxxx	xxVxxx
xxFxxx	xxCxxx	xxLxxx	xxMxxx	xxWxxx	xxxVxx	xxxIxx	xxxLxx
xxxFxx	xxxCxx	xxxWxx	xxxYxx	xxxIx	xxxWx	xxxFx	xxxCx
xxxxYx	xxxxxV	xxxxxC	xxxxxI	xxxxxF	xxxxxL		

Table 5: The list of all the 38 amyloidogenic patterns with five free positions when x is hydrophobic, but cannot be proline, i.e., chosen from CVLIMFYW. Each pattern describes  $8^5 = 32768$  hexapeptides.

able at <https://pitgroup.org/bap>. To our knowledge, no machine learning tool was used before to derive succinct chemical knowledge through simple patterns for deep structural properties.

### Data availability

The Budapest Amyloid Predictor webserver is available freely at <https://pitgroup.org/bap>.

### Funding

LK, ES, BV and VG were partially supported by the NKFI-127909 grant of the National Research, Development and Innovation Office of Hungary. LK, ES, AP, VF and VG were partially supported by the ELTE Thematic Excellence Programme (Szint+) subsidized by the Hungarian Ministry for Innovation and Technology.

### Author Contribution

AP, VF and VG have initiated the study and evaluated results, LK and ES constructed the SVM for the prediction, BV constructed the webserver, VG has overseen the work and wrote the paper. AP, VF and VG secured funding.

### Conflicting interest

The authors declare no conflicting interests.

### References

- [1] Emiel Michiels, Kenny Roose, Rodrigo Gallardo, Ladan Khodaparast, Laleh Khodaparast, Rob van der Kant, Maxime Siemons, Bert Houben, Meine Ramakers, Hannah Wilkinson, Patricia Guerreiro, Nikolaos Louros, Suzanne J F Kaptein, Lorena Itati Ibanez, Anouk Smet, Pieter Baatsen, Shu Liu, Ina Vorberg, Guy Bormans, Johan Neyts, Xavier Saelens, Frederic Rousseau, and Joost Schymkowitz. Reverse engineering synthetic antiviral

- amyloids. *Nature Communications*, 11:2832, June 2020. ISSN 2041-1723. doi: 10.1038/s41467-020-16721-8.
- [2] Julian D. Gillmore, Ed Gane, Jorg Taubel, Justin Kao, Marianna Fontana, Michael L. Maitland, Jessica Seitzer, Daniel O’Connell, Kathryn R. Walsh, Kristy Wood, Jonathan Phillips, Yuanxin Xu, Adam Amaral, Adam P. Boyd, Jeffrey E. Cehelsky, Mark D. McKee, Andrew Schiermeier, Olivier Harari, Andrew Murphy, Christos A. Kyriatsous, Brian Zambrowicz, Randy Soltys, David E. Gutstein, John Leonard, Laura Sepp-Lorenzino, and David Lebowitz. Crispr-cas9 in vivo gene editing for transthyretin amyloidosis. *New England Journal of Medicine*, 0(0):null, 0. doi: 10.1056/NEJMoa2107454. URL <https://doi.org/10.1056/NEJMoa2107454>.
- [3] Dániel Horváth, Dóra K Menyhárd, and András Perczel. Protein aggregation in a nutshell: The splendid molecular architecture of the dreaded amyloid fibrils. *Current Protein and Peptide Science*, 20(11):1077–1088, 2019.
- [4] Nora Taricska, Daniel Horvath, Dora K Menyhard, Hanna Akontz-Kiss, Masahiro Noji, Masatomo So, Yuji Goto, Toshimichi Fujiwara, and Andras Perczel. The route from the folded to the amyloid state: Exploring the potential energy surface of a drug-like miniprotein. *Chemistry A European Journal*, 26:1968–1978, February 2020. ISSN 1521-3765. doi: 10.1002/chem.201903826. URL <https://doi.org/10.1002/chem.201903826>.
- [5] Kristóf Takács, Bálint Varga, and Vince Grolmusz. Pdb\_amyloid: an extended live amyloid structure list from the PDB. *FEBS Open Bio*, 9(1): 185–190, 2019.
- [6] Kristof Takacs and Vince Grolmusz. On the border of the amyloidogenic sequences: Prefix analysis of the parallel beta sheets in the PDB\_Amyloid collection. *Journal of Integrative Bioinformatics*, 2021. URL <https://doi.org/10.1515/jib-2020-0043>.
- [7] Samir K Maji, Marilyn H Perrin, Michael R Sawaya, Sebastian Jessberger, Krishna Vadodaria, Robert A Rissman, Praful S Singru, K Peter R Nilsson, Rozalyn Simon, David Schubert, et al. Functional amyloids as natural storage of peptide hormones in pituitary secretory granules. *Science*, 325(5938):328–332, 2009.
- [8] Claudio Soto, Lisbell Estrada, and Joaquín Castilla. Amyloids, prions and the inherent infectious nature of misfolded protein aggregates. *Trends in Biochemical Sciences*, 31(3):150–155, 2006.
- [9] Carlos Familia, Sarah R Dennison, Alexandre Quintas, and David A Phoenix. Prediction of peptide and protein propensity for amyloid formation. *PLoS ONE*, 10:e0134679, 2015. ISSN 1932-6203. doi: 10.1371/journal.pone.0134679.

- [10] Gian Gaetano Tartaglia and Michele Vendruscolo. The zyggregator method for predicting protein aggregation propensities. *Chemical Society reviews*, 37:1395–1401, July 2008. ISSN 0306-0012. doi: 10.1039/b706784b.
- [11] Oscar Conchillo-Sole, Natalia S de Groot, Francesc X Aviles, Josep Vendrell, Xavier Daura, and Salvador Ventura. Aggrescan: a server for the prediction and evaluation of "hot spots" of aggregation in polypeptides. *BMC Bioinformatics*, 8:65, February 2007. ISSN 1471-2105. doi: 10.1186/1471-2105-8-65.
- [12] Changsik Kim, Jiwon Choi, Seong Joon Lee, William J Welsh, and Sukjoon Yoon. Netcssp: web application for predicting chameleon sequences and amyloid fibril formation. *Nucleic acids research*, 37:W469–W473, July 2009. ISSN 1362-4962. doi: 10.1093/nar/gkp351.
- [13] Jaime Santos, Jordi Pujols, Irantzu Pallares, Valentin Iglesias, and Salvador Ventura. Computational prediction of protein aggregation: Advances in proteomics, conformation-specific algorithms and biotechnological applications. *Computational and Structural Biotechnology Journal*, 18:1403–1413, 2020. ISSN 2001-0370. doi: 10.1016/j.csbj.2020.05.026.
- [14] Laszlo Keresztes, Evelin Szogi, Balint Varga, Viktor Farkas, Andras Perczel, and Vince Grolmusz. The Budapest Amyloid Predictor and its applications. *Biomolecules*, 11(4):500. URL <https://doi.org/10.3390/biom11040500>.
- [15] Corinna Cortes and Vladimir Vapnik. Support-vector networks. *Machine Learning*, 20(3):273–297, 1995.
- [16] Jacinte Beerten, Joost Van Durme, Rodrigo Gallardo, Emidio Capriotti, Louise Serpell, Frederic Rousseau, and Joost Schymkowitz. WALTZ-DB: a benchmark database of amyloidogenic hexapeptides. *Bioinformatics (Oxford, England)*, 31:1698–1700, May 2015. ISSN 1367-4811. doi: 10.1093/bioinformatics/btv027.
- [17] Nikolaos Louros, Katerina Konstantoulea, Matthias De Vleeschouwer, Meine Ramakers, Joost Schymkowitz, and Frederic Rousseau. WALTZ-DB 2.0: an updated database containing structural information of experimentally determined amyloid-forming peptides. *Nucleic Acids Research*, 48:D389–D393, January 2020. ISSN 1362-4962. doi: 10.1093/nar/gkz758.
- [18] Shuichi Kawashima, Piotr Pokarowski, Maria Pokarowska, Andrzej Krolinski, Toshiaki Katayama, and Minoru Kanehisa. Aaindex: amino acid index database, progress report 2008. *Nucleic Acids Research*, 36:D202–D205, January 2008. ISSN 1362-4962. doi: 10.1093/nar/gkm998.
- [19] Laszlo Keresztes, Evelin Szogi, Balint Varga, and Vince Grolmusz. Identifying super-feminine, super-masculine and sex-defining connections in the human braingraph. *Cognitive Neurodynamics*, 15(6):949–959, 2021. URL <https://doi.org/10.1007/s11571-021-09687-w>.

- [20] F. Pedregosa, G. Varoquaux, A. Gramfort, V. Michel, B. Thirion, O. Grisel, M. Blondel, P. Prettenhofer, R. Weiss, V. Dubourg, J. Vanderplas, A. Passos, D. Cournapeau, M. Brucher, M. Perrot, and E. Duchesnay. Scikit-learn: Machine learning in Python. *Journal of Machine Learning Research*, 12: 2825–2830, 2011.
- [21] Peter I Nagy and Paul W Erhardt. Theoretical studies of salt-bridge formation by amino acid side chains in low and medium polarity environments. *The Journal of Physical Chemistry B*, 114(49):16436–16442, 2010. URL <https://doi.org/10.1021/jp103313s>.
- [22] Arthur M Lesk. *Introduction to Bioinformatics*. Oxford: Oxford University Press, 2007.

## Supporting Material

**Table S1: Amyloid patterns in hexapeptides**

In the table we list all the existing minimal patterns of amyloid-forming hexapeptides, computed from the SVM model of the Budapest Amyloid Predictor. Each pattern in the Table contains two free “x” positions, where any amino acids can be substituted, and the resulting hexapeptide is predicted to be amyloidogenic. Since the two free positions can be substituted  $20 \times 20 = 400$  ways, each of the 5531 patterns in the Table describes 400 hexapeptides succinctly.

0	1	2	3	4	5	6	7	8	9
VIVxx	VIIxx	VILxx	VIIFxx	VIICxx	VIIWxx	VIIYxx	VIIIMxx	VIIAxx	VIVVxx
VIVIxx	VIVLxx	VIVFxx	VIVCxx	VIVWxx	VIVYxx	VIVVxx	VIVFxx	VIVCxx	VIVCxx
VILVxx	VILIxx	VIMVxx	VFIVxx	VFIIxx	VFILxx	VFIFxx	VFICxx	VFIWxx	VFIIxx
VFIMxx	VFVxx	VFVxx	VFVxx	VFVxx	VFVxx	VFVxx	VFVxx	VFVxx	VFVxx
VYIVxx	VYIIxx	VYILxx	VYIFxx	VYICxx	VYIWxx	VYIYxx	VYIMxx	VYIAxx	VYIVxx
VYVLxx	VYVFxx	VYVCxx	VYFVxx	VYFIxx	VYCVxx	VYLVxx	VYIVxx	VYIIxx	VYILxx
VVIFxx	VVICxx	VVIWxx	VVIYxx	VVVxx	VVVLxx	VVVLxx	VVVFxx	VVFCxx	VVVCxx
VVLVxx	VWIVxx	VWILxx	VWIFxx	VWICxx	VWIXxx	VWIXxx	VWIXxx	VWIXxx	VWIXxx
VWVLxx	VLIVxx	VLIxx	VLIxx	VLIxx	VLIxx	VLIxx	VLIxx	VLIxx	VLIxx
VCIxx	VCIxx	VCVxx	VCVxx	VCVxx	VMIxx	VMIxx	VMIxx	VMIxx	VMIxx
VHIVxx	VHIIxx	VHILxx	VHIFxx	VHICxx	VHIXxx	VHIXxx	VHIXxx	VHIXxx	VHIXxx
VTIVxx	VTIIxx	VNIVxx	VSIxx	VGIxx	IIIVxx	IIIIxx	IIILxx	IIIFxx	IIICxx
IIIVxx	IIIIxx	IIIMxx	IIIAxx	IIVxx	IIVLxx	IIVLxx	IIVFxx	IIVCxx	IIVxx
IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx	IIYVxx
IFILxx	IFIFxx	IFICxx	IFIWxx	IFIYxx	IFIMxx	IFVxx	IFVIxx	IFVLxx	IFVxx
IPFVxx	IFCVxx	IFLVxx	IYIVxx	IYIIxx	IYILxx	IYIFxx	IYICxx	IYIWxx	IYIxx
IYIMxx	IYVxx	IYVxx	IYVxx	IYVxx	IYVxx	IYVxx	IYVxx	IYVxx	IYVxx
IVILxx	IVIFxx	IVICxx	IVIWxx	IVIYxx	IVVxx	IVVIxx	IVVLxx	IVFVxx	IVCVxx
IWIVxx	IWIIxx	IWILxx	IWIFxx	IWICxx	IWVxx	IWVIxx	IWVLxx	ILIVxx	ILIIxx
ILILxx	ILIFxx	ILVVxx	ILVIxx	ILVLxx	ICIVxx	ICIIxx	ICILxx	ICVxx	ICVxx
IMIVxx	IMIIxx	IMILxx	IMVxx	IHIVxx	IHIIxx	IHILxx	IHVxx	IQIVxx	IQIIxx
IQVxx	IAIVxx	ITIVxx	INIVxx	ISIVxx	CIIVxx	CIIIxx	CIILxx	CIIFxx	CIICxx
CIWxx	CIYxx	CIYxx	CIYxx	CIYxx	CIYxx	CIYxx	CIYxx	CIYxx	CIYxx
CICVxx	CILVxx	CFIVxx	CFIIxx	CFILxx	CFIFxx	CFICxx	CFIWxx	CFIYxx	CFVxx
CFVxx	CFVxx	CFVxx	CYIVxx	CYIIxx	CYILxx	CYIFxx	CYICxx	CYIWxx	CYIxx
CYVxx	CYVxx	CYVxx	CYVxx	CYVxx	CYVxx	CYVxx	CYVxx	CYVxx	CYVxx
CVVxx	CVVxx	CVVxx	CVVxx	CVVxx	CVVxx	CVVxx	CVVxx	CVVxx	CVVxx
CLVxx	CLVxx	CCIVxx	CCIIxx	CCILxx	CCVxx	CCVIxx	CMIVxx	CMIIxx	CMVxx
CHIVxx	CHIIxx	CHVxx	CQIVxx	CAIVxx	CTIVxx	LIIVxx	LIIIxx	LILxx	LIIFxx



VxCVWx	VxCVFx	VxCVCx	VxCVYx	VxCVMx	VxCVVx	VxCVLx	VxCVGx	VxCIIx	VxCIWx
VxCIFx	VxCICx	VxCIXx	VxCIMx	VxCIVx	VxCILx	VxCLIx	VxCLWx	VxCLFx	VxCLCx
VxCLYx	VxCLMx	VxCLVx	VxCFIx	VxCFWx	VxCCIx	VxCCWx	VxCWIx	VxCYIx	VxLVIx
VxLVWx	VxLVFx	VxLVCx	VxLVYx	VxLVMx	VxLVVx	VxLVLx	VxLIIx	VxLIWx	VxLIFx
VxLICx	VxLIYx	VxLIMx	VxLIVx	VxLILx	VxLLIx	VxLLWx	VxLLFx	VxLLCx	VxLLYx
VxLFIx	VxLFWx	VxLCIx	VxLWIx	VxLYIx	VxMVIx	VxMVWx	VxMVFx	VxMVCx	VxMZYx
VxMVMx	VxMVVx	VxMVLx	VxMIIx	VxMIWx	VxMIFx	VxMICx	VxMIYx	VxMIMx	VxMIVx
VxMLIx	VxMLWx	VxMLFx	VxMFIx	VxWVIx	VxWVWx	VxWVFx	VxWVCx	VxWVYx	VxWIIx
VxWIWx	VxWLIx	VxYVIx	VxYVWx	VxYVFx	VxYIIx	IxIVIx	IxIVWx	IxIVFx	IxIVCx
IxIVYx	IxIVMx	IxIVVx	IxIVLx	IxIVGx	IxIVHx	IxIVTx	IxIVPx	IxIVAx	IxIVNx
IxIVQx	IxIVSx	IxIIIx	IxIIXx	IxIIFx	IxIICx	IxIIYx	IxIIMx	IxIIVx	IxIILx
IxIIGx	IxIIHx	IxIITx	IxIIPx	IxIIAx	IxIINx	IxILIx	IxILWx	IxILFx	IxILCx
IxILYx	IxILMx	IxILVx	IxILLx	IxILGx	IxILHx	IxILTx	IxIFIx	IxIFWx	IxIFFx
IxIFCx	IxIFYx	IxIFMx	IxIFVx	IxIFLx	IxIFGx	IxIFCx	IxICWx	IxICFx	IxICCx
IxICYx	IxICMx	IxICVx	IxICLx	IxICGx	IxIWIx	IxIWXx	IxIWFx	IxIWCx	IxIWXx
IxIWMx	IxIWWx	IxIWLx	IxIWGx	IxIYIx	IxIYWx	IxIYFx	IxIYCx	IxIYYx	IxIYMx
IxIYVx	IxIYLx	IxIYGx	IxIMIx	IxIMWx	IxIMFx	IxIMCx	IxIMYx	IxIMMx	IxIMVx
IxIMLx	IxIALx	IxIAWx	IxIAFx	IxIACx	IxIAYx	IxIAMx	IxIAVx	IxIALx	IxITIx
IxITWx	IxITFx	IxINIx	IxIHIx	IxIEIx	IxISIx	IxVIIx	IxVVWx	IxVVFx	IxVVCx
IxVVYx	IxVVMx	IxVVVx	IxVVLx	IxVVGx	IxVWHx	IxVVTx	IxVVPx	IxVVAx	IxVIIx
IxVIWx	IxVIFx	IxVICx	IxVIYx	IxVIMx	IxVIVx	IxVILx	IxVIGx	IxVIHx	IxVITx
IxVLIx	IxVLWx	IxVLFx	IxVLCx	IxVLYx	IxVLMx	IxVLVx	IxVLLx	IxVLGx	IxVLHx
IxVFIx	IxVFXx	IxVFFx	IxVFCx	IxVFXx	IxVFMx	IxVFXx	IxVFLx	IxVCIx	IxVCCx
IxVCFx	IxVCCx	IxVCYx	IxVCMx	IxVCVx	IxVCLx	IxVMIx	IxVWx	IxVWFx	IxVWCx
IxVVYx	IxVVMx	IxVVVx	IxVWLx	IxVYIx	IxVYWx	IxVYFx	IxVYCx	IxVYYx	IxVYMx
IxVYVx	IxVYLx	IxVYIx	IxVMWx	IxVMFx	IxVMCx	IxVMYx	IxVMMx	IxVMVx	IxVAIx
IxVAXx	IxVAFx	IxVITx	IxFVIx	IxFVWx	IxFVFXx	IxFVCx	IxFVYx	IxFVMx	IxFVXx
IxFVLx	IxFVGx	IxFIIx	IxFIWXx	IxFIFx	IxFICx	IxFIYx	IxFIMx	IxFIVx	IxFILx
IxFLIx	IxFLWx	IxFLFx	IxFLCx	IxFLYx	IxFLMx	IxFLVx	IxFFIx	IxFFWx	IxFCIx
IxFCWx	IxFWIx	IxFYIx	IxFMIx	IxCVLx	IxCVWx	IxCVFXx	IxCVCx	IxCVYx	IxCVMx
IxCVVx	IxCVLx	IxCIIx	IxCIWx	IxCIFx	IxCICx	IxCIXx	IxCIMx	IxCIVx	IxCILx
IxCLIx	IxCLWx	IxCLFx	IxCLCx	IxCLYx	IxCFIx	IxCFWx	IxCCIx	IxCWIx	IxCIYx
IxLVIx	IxLVWx	IxLVFx	IxLVCx	IxLVYx	IxLVMx	IxLVVx	IxLVLx	IxLIIx	IxLIWx
IxLIFx	IxLICx	IxLIYx	IxLIMx	IxLIVx	IxLILx	IxLLIx	IxLLWx	IxLLFx	IxLLCx
IxLFIx	IxLCIx	IxLWIx	IxLYIx	IxMVIx	IxMVWx	IxMVFXx	IxMVCx	IxMZYx	IxMVMx
IxMVWx	IxMVLx	IxMIIx	IxMIWx	IxMIFx	IxMICx	IxMIYx	IxMLIx	IxMLWx	IxWVIx
IxWVWx	IxWVFx	IxWIIx	IxWIWx	IxYVIx	IxYVWx	IxYIIx	CxIVIx	CxIVWx	CxIVFx
CxIVCx	CxIVYx	CxIVMx	CxIVLx	CxIVGx	CxIVHx	CxIVTx	CxIVPx	CxIVAx	CxIVNx
CxIVQx	CxIVSx	CxIIIx	CxIIWx	CxIIFx	CxIICx	CxIIYx	CxIIMx	CxIIVx	CxIILx
CxIIGx	CxIIHx	CxIITx	CxIIPx	CxIIAx	CxIINx	CxILIx	CxILWx	CxILFx	CxILCx
CxILYx	CxILMx	CxILVx	CxILGx	CxILHx	CxILTx	CxIFIx	CxIFWx	CxIFFx	CxIFCx
CxIFYx	CxIFMx	CxIFVx	CxIFLx	CxIFGx	CxICIx	CxICWx	CxICFx	CxICCx	CxICYx
CxICMx	CxICGx	CxICLx	CxICGx	CxIWXx	CxIWFx	CxIYFx	CxIYCx	CxIYYx	CxIYMx
CxIWXx	CxIWLx	CxIYIx	CxIYWx	CxIYFx	CxIYCx	CxIYVx	CxIYMx	CxIYVx	CxIYLx
CxIMIx	CxIMWx	CxIMFx	CxIMCx	CxIMYx	CxIMMx	CxIMVx	CxIMLx	CxIAIx	CxIAWx
CxIAFx	CxIACx	CxIAYx	CxIAMx	CxIAVx	CxITIx	CxINIx	CxVVIx	CxVVFx	CxVIIx
CxVVCx	CxVVYx	CxVVMx	CxVVVx	CxVVLx	CxVVGx	CxVWHx	CxVVTx	CxVVPx	CxVIIx
CxVIWx	CxVIFx	CxVICx	CxVIYx	CxVIMx	CxVIVx	CxVILx	CxVIGx	CxVIHx	CxVITx
CxVLIx	CxVLFx	CxVLCx	CxVLYx	CxVLMx	CxVLVx	CxVLLx	CxVLGx	CxVFXx	CxVFCx
CxVFIx	CxVFXx	CxVFFx	CxVFCx	CxVFXx	CxVFLx	CxVCIx	CxVCWx	CxVCFx	CxVCCx
CxVCFx	CxVCCx	CxVCYx	CxVCMx	CxVCVx	CxVCLx	CxVMIx	CxVWx	CxVWFx	CxVWCx
CxVVYx	CxVVMx	CxVVVx	CxVWLx	CxVYIx	CxVYWx	CxVYIx	CxVYMx	CxVYVx	CxVYMx
CxVMWx	CxVMFx	CxVMCx	CxVMYx	CxVAIx	CxVAWx	CxVAFx	CxFVIx	CxFVWx	CxFVFXx
CxFVCx	CxFVYx	CxFVMx	CxFVXx	CxFVLx	CxFIIXx	CxFIWXx	CxFIFx	CxFICx	CxFIYx
CxFIMx	CxFIVx	CxFILx	CxFLIx	CxFLWx	CxFLFx	CxFLCx	CxFFIx	CxFCIx	CxFWIx
CxFYIx	CxCVIx	CxCVWx	CxCVFx	CxCVCx	CxCVYx	CxCVMx	CxCVVx	CxCVLx	CxCIIx
CxCIWx	CxCIFx	CxCICx	CxCIXx	CxCIMx	CxCIVx	CxCLIx	CxCLWx	CxCLFx	CxCFIx
CxCCIx	CxLVIx	CxLVWx	CxLVFx	CxLVCx	CxLVYx	CxLVMx	CxLVVx	CxLVLx	CxLIIx
CxLIWx	CxLIFx	CxLICx	CxLIYx	CxLIMx	CxLIVx	CxLLIx	CxLLWx	CxLLFx	CxLFIx
CxMVIx	CxMVWx	CxMVFXx	CxMVCx	CxMVYx	CxMVMx	CxMVVx	CxMILx	CxMIWx	CxMIFx
CxMLIx	CxWVIx	CxWVWx	CxWVFXx	CxWIIx	CxYVIx	LxIVIx	LxIVWx	LxIVFx	LxIVCx
LxIVYx	LxIVMx	LxIVVx	LxIVLx	LxIVGx	LxIVHx	LxIVTx	LxIVPx	LxIVAx	LxIVNx
LxIVQx	LxIIIx	LxIIWx	LxIIFx	LxIICx	LxIIYx	LxIIMx	LxIIVx	LxIILx	LxIIGx
LxIIHx	LxIITx	LxIIPx	LxIIAx	LxILLx	LxILWx	LxILFx	LxILCx	LxILYx	LxILMx
LxILVx	LxILLx	LxILGx	LxILHx	LxIFIx	LxIFWx	LxIFFx	LxIFCx	LxIFYx	LxIFMx
LxIFVx	LxIFLx	LxIFGx	LxICIx	LxICWx	LxICFx	LxICCx	LxICYx	LxICMx	LxICVx
LxICLx	LxIWLx	LxIWXx	LxIWFx	LxIWCx	LxIWXx	LxIWMx	LxIWLx	LxIWIx	LxIYIx
LxIYWx	LxIYFx	LxIYCx	LxIYYx	LxIYMx	LxIYVx	LxIYLx	LxIMIx	LxIMWx	LxIMFx



LxIMCx	LxIMYx	LxIMMx	LxIMVx	LxIMLx	LxIAIx	LxIAWx	LxIAFv	LxIACx	LxIAYx
LxIAMx	LxIAVx	LxITIx	LxINIx	LxVVLx	LxVVVx	LxVVFx	LxVVCx	LxVVYx	LxVVMx
LxVVVx	LxVVLx	LxVVGx	LxVVHx	LxVVTx	LxVIIx	LxVIIx	LxVIFx	LxVICx	LxVIYx
LxVIMx	LxVIVx	LxVILx	LxVIGx	LxVIHx	LxVLIx	LxVLWx	LxVLFx	LxVLCx	LxVLYx
LxVLMx	LxVLVx	LxVLLx	LxVLGx	LxVFIx	LxVFXx	LxVFFx	LxVFCx	LxVFYx	LxVFMx
LxVFXx	LxVFLx	LxVCIx	LxVCWx	LxVCFx	LxVCCx	LxVCYx	LxVCMx	LxVCVx	LxVCLx
LxVVIx	LxVWXx	LxVWFx	LxVWCx	LxVWYx	LxVWMx	LxVWVx	LxVYIx	LxVYVx	LxVYFv
LxVYCx	LxVYYx	LxVYMx	LxVYVx	LxVMLx	LxVMWx	LxVMFv	LxVMCx	LxVAIx	LxVAWx
LxFVIx	LxFVWx	LxFVFx	LxFVCx	LxFVYx	LxFVMx	LxFVVx	LxFVLx	LxFIIx	LxFIWX
LxFIFx	LxFICx	LxFIYx	LxFIMx	LxFIVx	LxFILx	LxFIIX	LxFIWX	LxFIFx	LxFPIx
LxFCIx	LxFWIx	LxFYIx	LxCVIx	LxCVWx	LxCVFx	LxCVCx	LxCVYx	LxCVMx	LxCVVx
LxCVLx	LxCIIx	LxCIWx	LxCIFx	LxCICx	LxCIXx	LxCIMx	LxCIVx	LxCLIx	LxCLWx
LxCLFx	LxCFIx	LxCCIx	LxLVIx	LxLVWx	LxLVFv	LxLVCx	LxLVYx	LxLVMx	LxLVVx
LxLVLx	LxLIIx	LxLIXx	LxLIFx	LxLICx	LxLIYx	LxLIMx	LxLIVx	LxLLIx	LxLLWx
LxLLFx	LxLFIx	LxMVIx	LxMVWx	LxMVFx	LxMVCx	LxMVYx	LxMVMx	LxMVIx	LxMIIx
LxMIWx	LxMIFx	LxMLIx	LxMVIx	LxMWWx	LxMIIx	LxMVIx	FxIVIx	FxIVWx	FxIVFv
FxIVCx	FxIVYx	FxIVMx	FxIVVx	FxIVLx	FxIVGx	FxIVHx	FxIVTx	FxIVPx	FxIVAx
FxIIXx	FxIIXx	FxIIFx	FxIICx	FxIIXx	FxIIXx	FxIIXx	FxIIXx	FxIIXx	FxIIXx
FxIITx	FxILIx	FxILWx	FxILFv	FxILCx	FxILYx	FxILMx	FxILVx	FxILLx	FxILGx
FxIFIx	FxIFWx	FxIFFx	FxIFCx	FxIFYx	FxIFMx	FxIFVx	FxIFLx	FxICIx	FxICWx
FxICFx	FxICCx	FxICYx	FxICMx	FxICYx	FxICLx	FxIWXx	FxIWXx	FxIWFx	FxIWCx
FxIWXx	FxIWMx	FxIWXx	FxIWLx	FxIYIx	FxIYWx	FxIYFv	FxIYCx	FxIYYx	FxIYMx
FxIYVx	FxIYLx	FxIMIx	FxIMWx	FxIMFv	FxIMCx	FxIMYx	FxIMMx	FxIMVx	FxIAIx
FxIAWx	FxIAFv	FxITIx	FxVIIx	FxVWVx	FxVVFx	FxVVCx	FxVVYx	FxVVMx	FxVVVx
FxVVLx	FxVVGx	FxVVHx	FxVIIx	FxVIWx	FxVIFx	FxVICx	FxVIYx	FxVIMx	FxVIVx
FxVILx	FxVIGx	FxVLIx	FxVIXx	FxVLFx	FxVLCx	FxVLYx	FxVLMx	FxVLVx	FxVLLx
FxVFIx	FxVFXx	FxVFFx	FxVFCx	FxVFYx	FxVFMx	FxVFXx	FxVCLx	FxVCMx	FxVCFx
FxVCCx	FxVCYx	FxVCMx	FxVCVx	FxVWXx	FxVWFv	FxVWCx	FxVWYx	FxVYIx	FxVYVx
FxVYVx	FxVYFv	FxVYCx	FxVYVx	FxVMIx	FxVMWx	FxVMFv	FxVMCx	FxVAIx	FxVWVx
FxVFXx	FxVFCx	FxVYVx	FxVFMx	FxVFXx	FxVFLx	FxVIIx	FxVIIx	FxVIIx	FxVIIx
FxFIYx	FxFIMx	FxFIVx	FxFIIX	FxFIWX	FxFIIX	FxFIIX	FxFIIX	FxFIIX	FxFIIX
FxCVXx	FxCVYx	FxCVMx	FxCVXx	FxCVIX	FxCVIX	FxCVIX	FxCVIX	FxCVIX	FxCVIX
FxCLWx	FxLVLx	FxLVWx	FxLVFv	FxLVCx	FxLVYx	FxLVMx	FxLVVx	FxLIIx	FxLIWx
FxLIFx	FxLLIx	FxMVIx	FxMVWx	FxMVFx	FxMIIx	FxMVIx	FxMVIx	MxIVIx	MxIVWx
MxIVFx	MxIVCx	MxIVYx	MxIVVx	MxIVLx	MxIVGx	MxIVHx	MxIVTx	MxIIXx	MxIIXx
MxIIXx	MxIIFx	MxIICx	MxIIXx	MxIIMx	MxIIXx	MxIIXx	MxIIXx	MxIIXx	MxIIXx
MxILWx	MxILFx	MxILCx	MxILYx	MxILMx	MxILVx	MxILLx	MxILGx	MxIFIx	MxIFWx
MxIFFx	MxIFCx	MxIFYx	MxIFMx	MxIFVx	MxIFLx	MxICIx	MxICWx	MxICFx	MxICCx
MxICYx	MxICMx	MxICVx	MxICLx	MxIWLx	MxIWXx	MxIWFx	MxIWCx	MxIWXx	MxIWMx
MxIWXx	MxIYIx	MxIYWx	MxIYFv	MxIYCx	MxIYYx	MxIYMx	MxIYVx	MxIMIx	MxIMWx
MxIMFx	MxIAIx	MxIAWx	MxVIIx	MxVWVx	MxVVFx	MxVVCx	MxVVYx	MxVVMx	MxVVVx
MxVVLx	MxVVGx	MxVVHx	MxVIIx	MxVIWx	MxVIFx	MxVICx	MxVIYx	MxVIMx	MxVIVx
MxVILx	MxVIGx	MxVLIx	MxVIXx	MxVLFx	MxVLCx	MxVLYx	MxVLMx	MxVLVx	MxVLLx
MxVFIx	MxVFXx	MxVFFx	MxVFCx	MxVFXx	MxVCFx	MxVCFx	MxVWLx	MxVWVx	MxVWFx
MxVYIx	MxVYVx	MxVYFv	MxVMIx	MxFVIX	MxFVIX	MxFVIX	MxFVIX	MxFVIX	MxFVIX
MxFVXx	MxFIIX	MxFIWX	MxFIIX	MxFIWX	MxFIIX	MxFIIX	MxFIIX	MxFIIX	MxFIIX
MxCIx	MxCIWx	MxCIFx	MxCLIx	MxLVIx	MxLVWx	MxLVFv	MxLVCx	MxLIIx	MxLIWx
MxLLIx	MxMVIx	MxMVWx	MxMIIx	MxWVIX	WxIVIX	WxIVWx	WxIVFv	WxIVCx	WxIVYx
WxIVMx	WxIVVx	WxIVLx	WxIVGx	WxIVHx	WxIVTx	WxIIXx	WxIIXx	WxIIFx	WxIICx
WxIIXx	WxIIMx	WxIIVx	WxIILx	WxIIGx	WxIIXx	WxIIXx	WxIIXx	WxIIXx	WxIIXx
WxILYx	WxILMx	WxILVx	WxILLx	WxILGx	WxIFIx	WxIFWx	WxIFFx	WxIFCx	WxIFYx
WxIFMx	WxIFVx	WxIFLx	WxICIx	WxICWx	WxICFx	WxICCx	WxICYx	WxICMx	WxICVx
WxICLx	WxIWXx	WxIWXx	WxIWFx	WxIWCx	WxIWXx	WxIWXx	WxIWXx	WxIYIx	WxIYWx
WxIYFx	WxIYCx	WxIYYx	WxIYMx	WxIYVx	WxIMIx	WxIMWx	WxIMFv	WxIAIx	WxIAWx
WxVVIx	WxVWVx	WxVVFx	WxVVCx	WxVVYx	WxVVMx	WxVVVx	WxVVLx	WxVVGx	WxVVHx
WxVIIx	WxVIWx	WxVIFx	WxVICx	WxVIYx	WxVIMx	WxVIVx	WxVILx	WxVIGx	WxVLIx
WxVILx	WxVLFx	WxVLCx	WxVLYx	WxVLMx	WxVLVx	WxVLLx	WxVLFx	WxVFXx	WxVFFx
WxVFCx	WxVCIx	WxVCWx	WxVCFx	WxVWVx	WxVWFx	WxVYIx	WxVYVx	WxVYFv	WxVYMx
WxVMIx	WxFVIX	WxFVIX	WxFVIX	WxFVIX	WxFVIX	WxFVIX	WxFVIX	WxFVIX	WxFVIX
WxFIFx	WxFIIX	WxFIWX	WxFIIX	WxFIWX	WxFIIX	WxFIIX	WxFIIX	WxFIIX	WxFIIX
WxCLIx	WxLVLx	WxLVWx	WxLVFv	WxLVCx	WxLIIx	WxLIWx	WxLLIx	WxMVIx	WxMVWx
WxMIIx	GxIVIX	GxIVWx	GxIVFv	GxIVCx	GxIVYx	GxIVMx	GxIVVx	GxIVLx	GxIVGx
GxIVHx	GxIIIX	GxIIWx	GxIIFx	GxIICx	GxIIXx	GxIIMx	GxIIVx	GxIILx	GxIIGx
GxILLx	GxILWx	GxILFx	GxILCx	GxILYx	GxILMx	GxILVx	GxILLx	GxIFIx	GxIFWx
GxIFFx	GxIFCx	GxICIX	GxICWx	GxICFx	GxIWXx	GxIWXx	GxIWFx	GxIYIx	GxIYWx
GxIYFx	GxIMIx	GxVVIx	GxVWVx	GxVVFx	GxVVCx	GxVVYx	GxVVMx	GxVVVx	GxVVLx
GxVIXx	GxIWXx	GxVIFx	GxVICx	GxVIXx	GxVIMx	GxVIVx	GxVILx	GxVILx	GxVILx
GxVLFx	GxVLCx	GxVFIx	GxVCIx	GxVWIX	GxVYIX	GxFVIX	GxFVIX	GxFVIX	GxFVIX

GxCVIx	GxCVWx	GxCIIx	GxLVIx	GxMVIx	YxIVIx	YxIVWx	YxIVFx	YxIVCx	YxIVYx
YxIVMx	YxIVVx	YxIVLx	YxIVGx	YxIVHx	YxIIIx	YxIIWx	YxIIFx	YxIICx	YxIIXx
YxIIMx	YxIIVx	YxIILx	YxIIGx	YxILLx	YxILWx	YxILFx	YxILCx	YxILYx	YxILMx
YxILVx	YxILLx	YxIFIx	YxIFWx	YxIFFx	YxIFCx	YxICIx	YxICWx	YxICFx	YxIWIx
YxIWXx	YxIWFx	YxIYIx	YxIYWx	YxIYFx	YxIMIx	YxVVIx	YxVWx	YxVVFx	YxVVCx
YxVvYx	YxVvMx	YxVvVx	YxVvLx	YxVvIx	YxVIWx	YxVIFx	YxVICx	YxVIYx	YxVIMx
YxVIVx	YxVILx	YxVLIx	YxVLWx	YxVLFx	YxVLCx	YxVFIx	YxVCIx	YxVVIx	YxVVIx
YxFVIX	YxFVWx	YxFVFx	YxFIIX	YxCVIX	YxCVWx	YxCIIx	YxLVIx	YxMVIx	AxIVIX
AxIVWx	AxIVFx	AxIVCx	AxIVYx	AxIVMx	AxIVVx	AxIVLx	AxIVGx	AxIIIx	AxIIWx
AxIIFx	AxIICx	AxIIXx	AxIIMx	AxIIVx	AxIILx	AxILIx	AxILWx	AxILFx	AxILCx
AxILYx	AxILMx	AxILVx	AxILGx	AxILHx	AxIFWx	AxIFFx	AxICIx	AxICWx	AxIWIx
AxIMIx	AxVIIx	AxVVWx	AxVVFx	AxVVCx	AxVvYx	AxVvMx	AxVvVx	AxVvLx	AxVIIx
AxVIWx	AxVIFx	AxVICx	AxVIYx	AxVIMx	AxVIVx	AxVLIx	AxVLWx	AxVLFx	AxVFIx
AxFVIX	AxCVIX	AxLVIx	HxIVIX	HxIVWx	HxIVFx	HxIVCx	HxIVYx	HxIVMx	HxIVVx
HxIVLx	HxIIIX	HxIIXx	HxIIFx	HxIICx	HxIIXx	HxIIMx	HxIIXx	HxIILx	HxIIXx
HxILFx	HxIFIx	HxICIx	HxVIIx	HxVvWx	HxVvFx	HxVvCx	HxVvYx	HxVvMx	HxVvVx
HxVIIx	HxVIIx	HxVIFx	HxVILx	TxIVIX	TxIVWx	TxIVFx	TxIVCx	TxIVYx	TxIVMx
TxIVVx	TxIVLx	TxIIIX	TxIIXx	TxIIFx	TxIICx	TxIIXx	TxIIMx	TxIIVx	TxIILx
TxILWx	TxILFx	TxIFIx	TxVIIx	TxVvWx	TxVvFx	TxVvCx	TxVvYx	TxVvMx	TxVvVx
SxIVIX	SxIVWx	SxIVFx	SxIVCx	SxIVYx	SxIVMx	SxIVLx	SxIVGx	SxIIIX	SxIIXx
SxILIX	SxVvIX	SxVvVx	SxVvLx	SxVvIx	QxIVIX	QxIVWx	QxIVFx	QxIVCx	QxIVYx
QxIVMx	QxIVVx	QxIIIX	QxIIXx	QxIIFx	QxILIX	QxVvIX	QxVvWx	QxVvFx	QxVIIx
KxIVIX	KxIVWx	KxIVFx	KxIVCx	KxIVYx	KxIVMx	KxIVLx	KxIVGx	KxIIIX	KxIIXx
KxILIX	KxVvIX	KxVvVx	KxVvLx	NxIVIX	NxIVWx	NxIVFx	NxIVCx	NxIVYx	NxIIIX
NxIIXx	NxILIX	NxVIIx	NxVvWx	NxVvIx	RxIVIX	RxIVWx	RxIIIX	RxVvIX	DxIVIX
ExIVIX	VxIVxV	VxIVxW	VxIVxI	VxIVxF	VxIVxL	VxIVxY	VxIVxM	VxIVxN	VxIVxA
VxIVxT	VxIVxS	VxIVxH	VxIVxN	VxIVxQ	VxIVxK	VxIVxG	VxIVxR	VxIVxD	VxIIXV
VxIIXC	VxIIXI	VxIIXF	VxIIXL	VxIIXY	VxIIXW	VxIIXM	VxIIXX	VxIIXZ	VxIIXS
VxIIXH	VxIIXN	VxIIXQ	VxIIXK	VxIIXV	VxIIXL	VxIIXI	VxIIXF	VxIIXL	VxIIXY
VxIIXW	VxIIXM	VxIIXA	VxIIXT	VxIIXS	VxIIXH	VxIIXG	VxIIXR	VxIIXD	VxIIXS
VxIFxL	VxIFxY	VxIFxW	VxIFxC	VxIFxV	VxIFxM	VxIFxI	VxIFxC	VxIFxI	VxIFxY
VxICxW	VxICxM	VxICxV	VxICxI	VxICxY	VxICxL	VxICxI	VxICxF	VxICxI	VxICxY
VxIYxW	VxIYxI	VxIYxF	VxIYxL	VxIYxY	VxIYxW	VxIMxV	VxIMxC	VxIMxI	VxIMxL
VxIMxL	VxIAXV	VxIAXC	VxIAXI	VxITxV	VxVvIX	VxVvX	VxVvN	VxVvI	VxVvL
VxVvY	VxVvW	VxVvM	VxVvA	VxVvT	VxVvS	VxVvH	VxVvX	VxVvI	VxVIX
VxVIXI	VxVIXF	VxVIXL	VxVIXW	VxVIXW	VxVIXM	VxVIXA	VxVIXT	VxVIXS	VxVIXH
VxVLxV	VxVLxC	VxVLxI	VxVLxF	VxVLxL	VxVLxY	VxVLxM	VxVLxX	VxVFX	VxVFX
VxVFXI	VxVFXV	VxVFXL	VxVFXV	VxVFXI	VxVFXI	VxVFXI	VxVFXI	VxVFXI	VxVFXI
VxVWxF	VxVYxV	VxVYxL	VxVYxI	VxVYxF	VxVMxV	VxVMxC	VxVAX	VxVFX	VxVFX
VxVFXI	VxVFXV	VxVFXL	VxVFXI	VxVFXI	VxVFXI	VxVFXI	VxVFXI	VxVFXI	VxVFXI
VxFLxI	VxFPxV	VxCVxV	VxCVxC	VxCVxI	VxCVxF	VxCVxL	VxCVxY	VxCVxI	VxCVxI
VxCIxI	VxCLxV	VxLVxV	VxLVxC	VxLVxI	VxLVxF	VxLVxL	VxLVxV	VxLVxI	VxLVxI
VxLLxV	VxMVxV	VxMVxC	VxMVxI	VxMVxV	VxMVxV	IxIVxV	IxIVxC	IxIVxI	IxIVxF
IxIVxL	IxIVxY	IxIVxW	IxIVxM	IxIVxA	IxIVxT	IxIVxS	IxIVxH	IxIVxN	IxIVxQ
IxIVxK	IxIVxG	IxIVxR	IxIVxD	IxIIXV	IxIIXC	IxIIXI	IxIIXF	IxIIXL	IxIIXY
IxIIXW	IxIIXM	IxIIXA	IxIIXT	IxIIXS	IxIIXH	IxIIXN	IxIIXV	IxIIXL	IxIIXY
IxILxI	IxILxL	IxILxY	IxILxW	IxILxM	IxILxA	IxILxT	IxILxS	IxILxH	IxILxF
IxIFxI	IxIFxI	IxIFxF	IxIFxL	IxIFxY	IxIFxW	IxIFxM	IxICxV	IxICxC	IxICxI
IxICxF	IxICxL	IxICxY	IxICxW	IxIWX	IxIWX	IxIWX	IxIWX	IxIWX	IxIWX
IxIYxV	IxIYxC	IxIYxI	IxIYxF	IxIYxL	IxIYxY	IxIMxV	IxIMxC	IxIMxI	IxIMxL
IxIAXV	IxIAXC	IxIAXI	IxVvX	IxVvxC	IxVvX	IxVvX	IxVvX	IxVvX	IxVvX
IxVvXM	IxVvxA	IxVvXT	IxVvXS	IxVvXH	IxVIX	IxVIX	IxVIX	IxVIX	IxVIX
IxVIxY	IxVIxW	IxVIxM	IxVIxA	IxVIxT	IxVIX	IxVIX	IxVIX	IxVIX	IxVIX
IxVLxF	IxVLxL	IxVLxY	IxVLxW	IxVLxM	IxVFX	IxVFX	IxVFX	IxVFX	IxVFX
IxVCxI	IxVCxI	IxVCxF	IxVWxV	IxVWxC	IxVWxI	IxVYxV	IxVYxC	IxVYxI	IxVYxI
IxVMxC	IxVAX	IxFVxV	IxFVxC	IxFVxI	IxFVxF	IxFVxL	IxFVxY	IxFIxV	IxFIxC
IxFIXI	IxFLxV	IxCVxV	IxCVxC	IxCVxI	IxCVxF	IxCVxL	IxCIxY	IxCIxC	IxCIxI
IxCLxV	IxLVxV	IxLVxC	IxLVxI	IxLVxF	IxLIX	IxLIX	IxLIX	IxLIX	IxLVxV
IxMVxC	IxMVxI	IxMIxV	IxWVxV	CxIVxV	CxIVxC	CxIVxI	CxIVxF	CxIVxL	CxIVxY
CxIVxW	CxIVxM	CxIVxA	CxIVxT	CxIVxS	CxIVxH	CxIVxN	CxIVxQ	CxIVxK	CxIVxG
CxIIXV	CxIIXC	CxIIXI	CxIIXF	CxIIXL	CxIIXY	CxIIXM	CxIIXX	CxIIXZ	CxIIXS
CxIIXS	CxIIXH	CxIIXV	CxIIXC	CxIIXI	CxIIXF	CxIIXL	CxIIXY	CxIIXW	CxIIXM
CxILxA	CxIFxV	CxIFxC	CxIFxI	CxIFxF	CxIFxL	CxIFxY	CxICxC	CxICxC	CxICxI
CxICxF	CxICxL	CxIWX	CxIWX	CxIWX	CxIWX	CxIYxV	CxIYxC	CxIYxI	CxIYxF
CxIMxV	CxIMxC	CxIMxI	CxIAX	CxIAX	CxVvX	CxVvX	CxVvX	CxVvX	CxVvX
CxVvXY	CxVvXW	CxVvXM	CxVvX	CxVvX	CxVvX	CxVvX	CxVIX	CxVIX	CxVIX
CxVIXI	CxVIXL	CxVIXY	CxVIXW	CxVIXM	CxVIX	CxVIX	CxVIX	CxVIX	CxVIX
CxVLxY	CxVFX	CxVFX	CxVFX	CxVFX	CxVFX	CxVFX	CxVFX	CxVFX	CxVFX





xAIVIx	xAIVWx	xAIVFx	xAIIIx	xAVVIx	xTIVIx	xTIVWx	xTIVFx	xTIIIx	xTVVIx
xNIVIx	xNIVWx	xNIIIx	xNIVIx	xSIVIx	xSIVWx	xSIIIx	xGIVIx	xPIVIx	xIIVxV
xIIVxG	xIIVxI	xIIVxF	xIIVxL	xIIVxY	xIIVxW	xIIVxM	xIIVxA	xIIVxT	xIIVxS
xIIIxV	xIIIxC	xIIIxI	xIIIxF	xIIIxL	xIIIxY	xIIIxW	xIIIxM	xIILxV	xIILxG
xIILxI	xIILxF	xIILxL	xIIFxV	xIIFxC	xIICxV	xIICxC	xIIWxV	xIIVxV	xIIMxV
xIVVxV	xIVVxC	xIVVxI	xIVVxF	xIVVxL	xIVVxY	xIVVxW	xIVIxV	xIVIxG	xIVIxI
xIVIxF	xIVLxV	xIVLxC	xIVFxV	xIVFxV	xICVxV	xILVxV	xFIVxV	xFIVxC	xFIVxI
xFIVxF	xFIVxL	xFIVxY	xFIVxW	xFIVxM	xFIIxV	xFIIxC	xFIIxI	xFIIxF	xFIIxL
xFIIxY	xFILxV	xFILxC	xFILxI	xFIFxV	xFICxV	xFIIxV	xFIIxV	xFVxV	xFVxG
xFVvXxI	xFVvXxF	xFVvXxL	xFVIxV	xFVIxC	xFVIxI	xFVLxV	xFFVxV	xYIVxV	xYIVxC
xYIVxI	xYIVxF	xYIVxL	xYIVxY	xYIVxW	xYIVxM	xYIIxV	xYIIxC	xYIIxI	xYIIxF
xYILxL	xYILxY	xYILxV	xYILxC	xYILxI	xYIFxV	xYICxV	xYIWxV	xYIVxV	xYVvXxV
xYVvXxC	xYVvXxI	xYVvXxF	xYVvXxL	xYVIxV	xYVIxC	xYVIxI	xYVLxV	xYFVxV	xVIVxV
xVILxV	xVILxC	xVIFxV	xVIVxL	xVIVxY	xVIVxW	xVIIxV	xVIIxC	xVIIxI	xVIIxF
xWIVxV	xWIVxC	xWIVxI	xWIVxF	xWIVxL	xWIVxY	xWIIxV	xWIIxC	xWIIxI	xWILxV
xWVvXxV	xWVvXxC	xWVvXxI	xWVIxV	xLIVxV	xLIVxC	xLIVxI	xLIVxF	xLIVxL	xLIVxV
xLIIxG	xLIIxI	xLILxV	xLVVxV	xLVVxC	xLVVxI	xLVIxV	xCIVxV	xCIVxC	xCIVxI
xCIVxF	xCIIxV	xCIIxC	xCILxV	xCVVxV	xCVVxC	xCVIxV	xMIVxV	xMIVxC	xMIIxV
xMVVxV	xHIVxV	xHIVxC	xHIIxV	xHVvXxV	xHVvXxC	xQIVxV	xAIVxV	xIIXxI	xIIXxI
xIIXxF	xIIXxL	xIIXxW	xIIXWC	xIIXWI	xIIXFV	xIIXFC	xIIXCV	xIIXV	xIIXMV
xIIXVV	xIVxIC	xIVxW	xIVxW	xIVxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV
xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV	xFIIxV
xYVxIV	xVIxIV	xVIxIC	xVIxW	xVIxV	xVvXxIV	xWIXxIV	xLIXxIV	xCIxIV	xxIVIV
xxIVIC	xxIVII	xxIVIF	xxIVIL	xxIVII	xxIVIV	xxIVIM	xxIVIA	xxIVIT	xxIVIS
xxIVIH	xxIVIN	xxIVIQ	xxIVIK	xxIVIG	xxIVIR	xxIVID	xxIVIE	xxIVV	xxIVWC
xxIVWI	xxIVWF	xxIVWL	xxIVWY	xxIVWW	xxIVWM	xxIVWA	xxIVWT	xxIVWS	xxIVWH
xxIVWN	xxIVWQ	xxIVWK	xxIVWG	xxIVWR	xxIVFS	xxIVFH	xxIVFN	xxIVFQ	xxIVFK
xxIVFY	xxIVFV	xxIVFM	xxIVFA	xxIVFT	xxIVFS	xxIVFH	xxIVFN	xxIVFQ	xxIVFK
xxIVFG	xxIVCV	xxIVCC	xxIVCI	xxIVCF	xxIVCL	xxIVCY	xxIVCW	xxIVCM	xxIVCA
xxIVCT	xxIVCS	xxIVCH	xxIVYV	xxIVYC	xxIVYI	xxIVYF	xxIVYL	xxIVYY	xxIVYW
xxIVYM	xxIVYA	xxIVYT	xxIVYS	xxIVYH	xxIVMV	xxIVMC	xxIVMI	xxIVMF	xxIVML
xxIVMY	xxIVMW	xxIVMM	xxIVMA	xxIVMT	xxIVMS	xxIVMH	xxIVVV	xxIVVC	xxIVVI
xxIVVF	xxIVVL	xxIVVY	xxIVVV	xxIVVM	xxIVVA	xxIVVT	xxIVVS	xxIVVH	xxIVLV
xxIVLC	xxIVLI	xxIVLF	xxIVLL	xxIVLY	xxIVLW	xxIVLM	xxIVLA	xxIVLT	xxIVLS
xxIVLH	xxIVGV	xxIVGC	xxIVGI	xxIVGF	xxIVGL	xxIVGV	xxIVGW	xxIVHV	xxIVHC
xxIVHI	xxIVHF	xxIVTV	xxIVTC	xxIVPV	xxIVPC	xxIVAV	xxIVNV	xxIVQV	xxIIV
xxIIIC	xxIIII	xxIIIF	xxIIIL	xxIIY	xxIIW	xxIIIM	xxIIIA	xxIIIT	xxIIIS
xxIIIH	xxIIIN	xxIIIQ	xxIIIK	xxIIIG	xxIIIR	xxIIWS	xxIIWC	xxIIWI	xxIIF
xxIIVL	xxIIVY	xxIIVW	xxIIVM	xxIIVA	xxIIVT	xxIIVS	xxIIVH	xxIIVV	xxIIFV
xxIIFC	xxIIFI	xxIIFG	xxIIFL	xxIIFY	xxIIFW	xxIIFM	xxIIFA	xxIIFT	xxIIFS
xxIIFH	xxIIFN	xxIIFQ	xxIIFK	xxIIFG	xxIIFL	xxIIFM	xxIIFA	xxIIFT	xxIIFS
xxIICT	xxIICS	xxIICH	xxIICY	xxIICY	xxIICY	xxIICY	xxIICY	xxIICY	xxIICY
xxIIMY	xxIIMX	xxIIMZ	xxIIMW	xxIIMV	xxIIMU	xxIIMI	xxIIMF	xxIIML	xxIIMY
xxIIMW	xxIIMM	xxIIMA	xxIIMT	xxIIMV	xxIIMC	xxIIMI	xxIIMF	xxIIML	xxIIMY
xxIIVW	xxIIVM	xxIIVA	xxIIVT	xxIILV	xxIILC	xxIILI	xxIILF	xxIILL	xxIILY
xxIILW	xxIILM	xxIILG	xxIILC	xxIILY	xxIILV	xxIILW	xxIILX	xxIILI	xxIILF
xxIIPV	xxIIPW	xxIIPX	xxIIPY	xxIIPZ	xxIIPA	xxIIPB	xxIIPC	xxIIPD	xxIIFE
xxILIA	xxILIT	xxILIS	xxILIH	xxILW	xxILV	xxILW	xxILWF	xxILWV	xxILWY
xxILWW	xxILWM	xxILWA	xxILWT	xxILWS	xxILWH	xxILFV	xxILFC	xxILFI	xxILFF
xxILFL	xxILFY	xxILFW	xxILFM	xxILFA	xxILFT	xxILCV	xxILCC	xxILCI	xxILCF
xxILCL	xxILCY	xxILCW	xxILCM	xxILYV	xxILYC	xxILYI	xxILYF	xxILYL	xxILYV
xxILYW	xxILYM	xxILMV	xxILMC	xxILMI	xxILMF	xxILML	xxILMY	xxILMW	xxILV
xxILXV	xxILXI	xxILXW	xxILXW	xxILXW	xxILXW	xxILXW	xxILXW	xxILXW	xxILXW
xxILLL	xxILLY	xxILLV	xxILLV	xxILLV	xxILLV	xxILLV	xxILLV	xxILLV	xxILLV
xxIFII	xxIFIV	xxIFIV	xxIFIV	xxIFIV	xxIFIV	xxIFIV	xxIFIV	xxIFIV	xxIFIV
xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX	xxIFIX
xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY	xxIFWY
xxIFCC	xxIFCI	xxIFCF	xxIFCY	xxIFCY	xxIFCY	xxIFCY	xxIFCY	xxIFCY	xxIFCY
xxIFMF	xxIFVW	xxIFVC	xxIFVI	xxIFVW	xxIFLV	xxIFLC	xxIFLI	xxIFGV	xxICIV
xxICIC	xxICII	xxICIF	xxICIL	xxICIV	xxICIW	xxICIM	xxICW	xxICWC	xxICWI
xxICWF	xxICWL	xxICWY	xxICWZ	xxICFV	xxICFC	xxICFI	xxICFF	xxICFL	xxICFY
xxICCV	xxICCC	xxICCI	xxICCF	xxICVY	xxICYC	xxICYI	xxICMV	xxICMC	xxICMI
xxICVW	xxICVC	xxICVI	xxICLV	xxICLC	xxICLV	xxICLV	xxICLV	xxICLV	xxICLV
xxIWIY	xxIWIW	xxIWIW	xxIWIW	xxIWIW	xxIWIW	xxIWIW	xxIWIW	xxIWIW	xxIWIW
xxIWFC	xxIWFI	xxIWFF	xxIWCV	xxIWCC	xxIWCI	xxIWVY	xxIWYC	xxIWYI	xxIWMV
xxIWMW	xxIWMI	xxIWVW	xxIWCV	xxIWCW	xxIWVI	xxIWLW	xxIYIV	xxIYIC	xxIYII
xxIYIF	xxIYIL	xxIYIY	xxIYIW	xxIYIM	xxIYVW	xxIYWC	xxIYWI	xxIYWF	xxIYWL
xxIYVY	xxIYVW	xxIYFC	xxIYFI	xxIYFF	xxIYCV	xxIYCC	xxIYCI	xxIYVY	xxIYVC

xxIYYI	xxIYMV	xxIYMC	xxIYMI	xxIYVV	xxIYVC	xxIYLV	xxIYLC	xxIMIV	xxIMIC
xxIMII	xxIMIF	xxIMIL	xxIMIY	xxIMIW	xxIMWV	xxIMWC	xxIMWI	xxIMWF	xxIMFV
xxIMFC	xxIMFI	xxIMCV	xxIMCC	xxIMYV	xxIMYC	xxIMMV	xxIMVV	xxIMLV	xxIAIV
xxIAIC	xxIAII	xxIAIF	xxIAWV	xxIAWC	xxIAFV	xxIAFC	xxIACV	xxIAYV	xxIAMV
xxIAVV	xxITIV	xxINIV	xxVVIV	xxVVIC	xxVVII	xxVVIF	xxVVIL	xxVVIY	xxVVIW
xxVVIM	xxVVIA	xxVVIT	xxVVIS	xxVVIH	xxVVIN	xxVVIQ	xxVVIK	xxVVIG	xxVVVV
xxVVWC	xxVVWI	xxVVWF	xxVVWL	xxVVWY	xxVVWV	xxVVWM	xxVVWA	xxVVWT	xxVVWS
xxVVWH	xxVVVF	xxVVFC	xxVVFI	xxVVFF	xxVVFL	xxVVFY	xxVVFW	xxVVFM	xxVVFA
xxVVFT	xxVVFS	xxVVFH	xxVVCV	xxVVCC	xxVVCI	xxVVCF	xxVVCL	xxVVCY	xxVVCW
xxVVCM	xxVVCA	xxVVCT	xxVVCS	xxVVYV	xxVVYC	xxVVYI	xxVVYF	xxVVYL	xxVVYY
xxVVYV	xxVVYM	xxVVYA	xxVVMV	xxVVMC	xxVVMI	xxVVMF	xxVVML	xxVVMY	xxVVMW
xxVVMM	xxVVVV	xxVVVC	xxVVVI	xxVVVF	xxVVVL	xxVVVM	xxVVVW	xxVVVM	xxVVLV
xxVVLC	xxVVLI	xxVVLF	xxVVLL	xxVVLY	xxVVLV	xxVVLM	xxVVGV	xxVVGC	xxVVGI
xxVVGF	xxVVHV	xxVVHC	xxVVTV	xxVVIW	xxVVIC	xxVVIH	xxVVIK	xxVVIY	xxVVIW
xxVVIW	xxVVIH	xxVVIK	xxVVIY	xxVVIW	xxVVIC	xxVVIH	xxVVIK	xxVVIY	xxVVIW
xxVIWL	xxVIWY	xxVIWV	xxVIWM	xxVIWA	xxVIWT	xxVIWS	xxVIVF	xxVIVC	xxVIVF
xxVIFL	xxVIFM	xxVIFV	xxVIFW	xxVIFM	xxVIFA	xxVICV	xxVICC	xxVICI	xxVICF
xxVICL	xxVICY	xxVICW	xxVICM	xxVIYV	xxVIYC	xxVIYI	xxVIYF	xxVIYL	xxVIYY
xxVIYV	xxVIMV	xxVIMC	xxVIMI	xxVIMF	xxVIML	xxVIMY	xxVIMW	xxVIVV	xxVIVC
xxVIVI	xxVIVF	xxVIVL	xxVIVY	xxVIVW	xxVIVL	xxVIVC	xxVILI	xxVILF	xxVILL
xxVIGV	xxVIGC	xxVIGH	xxVLIV	xxVLIC	xxVLII	xxVLIF	xxVLIL	xxVLIY	xxVLIW
xxVLIM	xxVLIA	xxVLIT	xxVLIS	xxVLWV	xxVLWC	xxVLWI	xxVLWF	xxVLWL	xxVLWY
xxVLWV	xxVLWM	xxVLVW	xxVLFC	xxVLFI	xxVLFF	xxVLFL	xxVLFY	xxVLFW	xxVLVC
xxVLCC	xxVLCI	xxVLCF	xxVLYV	xxVLYC	xxVLYI	xxVLYF	xxVLMV	xxVLMC	xxVLCI
xxVLMF	xxVLVV	xxVLVC	xxVLVI	xxVLVF	xxVLLV	xxVLLC	xxVLLI	xxVVLV	xxVVLV
xxVFIC	xxVFII	xxVFIF	xxVFIL	xxVFIV	xxVFIV	xxVFVW	xxVFVC	xxVFVI	xxVFVW
xxVFFV	xxVFFC	xxVFFI	xxVFCV	xxVFCV	xxVFYV	xxVFYC	xxVFMV	xxVFFV	xxVFLV
xxVCIV	xxVCIC	xxVCII	xxVCIF	xxVCIL	xxVCIV	xxVCVW	xxVCWC	xxVCVI	xxVCFV
xxVCCV	xxVCFI	xxVCCV	xxVCYV	xxVCMV	xxVCCV	xxVCLV	xxVWIV	xxVVIC	xxVWII
xxVWIF	xxVWIL	xxVWVW	xxVWCV	xxVWVI	xxVWVW	xxVWFC	xxVWCV	xxVWYV	xxVWVW
xxVWVV	xxVYIV	xxVYIC	xxVYII	xxVYIF	xxVYIL	xxVYVW	xxVYWC	xxVYVI	xxVYFV
xxVYFC	xxVYCV	xxVYVW	xxVYMV	xxVYVW	xxVMIW	xxVMIC	xxVMII	xxVMWV	xxVMWC
xxVMFV	xxVMCV	xxVAIV	xxVAIC	xxVAWV	xxFVIV	xxFVIC	xxFVII	xxFVIF	xxFVIL
xxFVIY	xxFVIW	xxFVIM	xxFVWV	xxFVWC	xxFVWF	xxFVWL	xxFVWV	xxFVWV	xxFVWV
xxFVFC	xxFVFI	xxFVFF	xxFVVC	xxFVCC	xxFVCI	xxFVYV	xxFVYC	xxFVYI	xxFVMV
xxFVMC	xxFVVV	xxFVVC	xxFVLV	xxFIIV	xxFIIIC	xxFIII	xxFIIF	xxFIIIL	xxFIIY
xxFIWV	xxFIWV	xxFIWV	xxFIFV	xxFIFC	xxFIFV	xxFICV	xxFIYV	xxFIMV	xxFIVV
xxFILV	xxFLIV	xxFLIC	xxFLII	xxFLWV	xxFLFV	xxFFIV	xxFCIV	xxFWIV	xxCVIV
xxCVIC	xxCVII	xxCVIF	xxCVIL	xxCVIV	xxCVIW	xxCVVW	xxCVWC	xxCVMI	xxCVWF
xxCVFV	xxCVFC	xxCVFI	xxCVFF	xxCVCV	xxCVCC	xxCVYV	xxCVYC	xxCVVI	xxCVMC
xxCVVV	xxCVVC	xxCVLV	xxCVIV	xxCIIC	xxCIII	xxCIIF	xxCIWV	xxCIWC	xxCIWI
xxCIFV	xxCIFC	xxCICV	xxCIYV	xxCIMV	xxCIVV	xxCLIV	xxCLIC	xxCLWV	xxCLFV
xxCFIV	xxLVIV	xxLVIC	xxLVII	xxLVIF	xxLVIL	xxLVYV	xxLVWV	xxLVWV	xxLVWC
xxLVWI	xxLVWF	xxLVFV	xxLVFC	xxLVFI	xxLVCV	xxLVCC	xxLVYV	xxLVYC	xxLVMV
xxLVVW	xxLVLV	xxLIIV	xxLIIIC	xxLIII	xxLIIF	xxLIWV	xxLIWC	xxLIFV	xxLIFC
xxLICV	xxLIYV	xxLIIV	xxLIVV	xxLLIV	xxLLIC	xxLLWV	xxLLFV	xxLFIV	xxMVIV
xxMVIC	xxMVII	xxMVIF	xxMVWV	xxMVWC	xxMVFV	xxMVFC	xxMVCV	xxMVYV	xxMVWV
xxMVVV	xxMIIV	xxMIIC	xxMIWV	xxMIFV	xxMLIV	xxWVIV	xxWVIC	xxWVVV	xxWVII
xxVVIV									

**Table S2: Amyloid patterns for restricted substitutions with small, non-polar residues**

In the table we list all the existing minimal patterns of amyloid-forming hexapeptides, computed from the SVM model of the Budapest Amyloid Predictor, when the substitutions for the positions, denoted by “x” are allowed by small, non-polar residues, namely G, A, S and T. The table contains 411 patterns.

0	1	2	3	4	5	6	7	8	9
VIIxxx	IIIxxx	VxIVxx	VxIIxx	VxILxx	VxIFxx	VxICxx	VxIWxx	VxIYxx	VxIMxx
VxVVxx	VxVIxx	VxVLxx	VxVFxx	VxVvxx	VxCVxx	VxLVxx	IxIVxx	IxIIxx	IxILxx
IxIFxx	IxICxx	IxIWxx	IxIYxx	IxVvxx	IxVIxx	IxVLxx	IxVFxx	IxFVxx	IxCVxx
IxLVxx	CxIVxx	CxILxx	CxILxx	CxIFxx	CxICxx	CxIWxx	CxIYxx	CxVVxx	CxVLxx
CxVLxx	CxFVxx	LxIVxx	LxIIxx	LxILxx	LxIFxx	LxICxx	LxIWxx	LxVVxx	LxVIxx
LxVLxx	FxIVxx	FxIIxx	FxILxx	FxVvxx	FxVIxx	FxVLxx	MxIVxx	MxIIxx	MxILxx
MxVVxx	MxVIxx	WxIVxx	WxIIxx	WxILxx	WxVvxx	WxVIxx	GxIVxx	GxIIxx	GxVVxx
YxIVxx	YxIIxx	YxVvxx	AxIVxx	VxIxIx	VxIxWx	VxIxFx	VxIxCx	VxIxYx	VxIxMx
VxIxVx	VxIxLx	VxVxIx	VxVxWx	VxVxFx	IxIxIx	IxIxWx	IxIxFx	IxIxCx	IxIxYx
IxIxMx	IxIxVx	IxVxIx	IxVxWx	IxVxFx	CxIxIx	CxIxWx	CxIxFx	CxIxCx	CxIxYx
CxVxIx	LxIxIx	LxIxWx	LxIxFx	LxIxCx	LxVxIx	FxIxIx	FxIxWx	FxIxFx	FxVxIx
MxIxIx	WxIxIx	VxIxxV	VxIxxC	VxVxxV	IxIxxV	IxIxxC	CxIxxV	LxIxxV	xIIVxx
xIIIxx	xIILxx	xIIFxx	xIICxx	xIIWxx	xIIYxx	xIIVxx	xIVIxx	xIVLxx	xIPVxx
xFIVxx	xFIIxx	xFILxx	xFIFxx	xFVvxx	xFVIxx	xFVLxx	xYIVxx	xYIIxx	xYILxx
xYIFxx	xYVvxx	xYVIxx	xYVLxx	xYVvxx	xVIIxx	xVILxx	xVVVxx	xWVvxx	xWIVxx
xWIIxx	xWILxx	xWVvxx	xWVIxx	xLIVxx	xLIIxx	xLILxx	xLVVxx	xLVVxx	xCIVxx
xCIIxx	xCVvxx	xMIVxx	xHIVxx	xIIxIx	xIIxWx	xIIxFx	xIIxCx	xIIxYx	xIVxIx
xIVxWx	xFIxIx	xFIxWx	xFIxFx	xFVxIx	xYIxIx	xYIxWx	xYIxFx	xYVxIx	xVIxIx
xVlxWx	xWlxIx	xLlxIx	xIlxV	xFIxxV	xYIxxV	xxIVIx	xxIVWx	xxIVFx	xxIVCx
xxIVYx	xxIVMx	xxIVVx	xxIVLx	xxIVGx	xxIVHx	xxIVTx	xxIILx	xxIILx	xxIIFx
xxIICx	xxIIXx	xxIIMx	xxIIVx	xxIILx	xxIIGx	xxIIXx	xxILIx	xxILWx	xxILFx
xxILCx	xxILYx	xxILMx	xxILVx	xxILLx	xxILGx	xxIFIx	xxIFWx	xxIFFx	xxIFCx
xxIFYx	xxIFMx	xxIFVx	xxIFLx	xxICIx	xxICWx	xxICFx	xxICCx	xxICYx	xxICMx
xxICVx	xxICLx	xxIWIx	xxIWWx	xxIWFx	xxIWCx	xxIWYx	xxIWMx	xxIWXx	xxIYIx
xxIYWx	xxIYFx	xxIYCx	xxIYYx	xxIYMx	xxIYVx	xxIMIx	xxIMWx	xxIMFx	xxIAIx
xxIAWx	xxVVIx	xxVVWx	xxVVFx	xxVVCx	xxVvYx	xxVVMx	xxVVVx	xxVVLx	xxVVGx
xxVWHx	xxVIIx	xxVIWx	xxVIFx	xxVICx	xxVIYx	xxVIMx	xxVIVx	xxVILx	xxVIGx
xxVLIx	xxVWLx	xxVLFx	xxVLCx	xxVLYx	xxVLMx	xxVLVx	xxVLLx	xxVFIx	xxVFWx
xxVFFx	xxVFCx	xxVCIx	xxVCWx	xxVCFx	xxVVIx	xxVWx	xxVWFx	xxVVIx	xxVVWx
xxVYFx	xxVMIx	xxFVIx	xxFVWx	xxFVFx	xxFVCx	xxFVYx	xxFVMx	xxFVvxx	xxFIIx
xxFIIx	xxFIFx	xxFLIx	xxCVIx	xxCVWx	xxCVFx	xxCVCx	xxCVYx	xxCIIx	xxCIWx
xxCIFx	xxCLIx	xxLVIx	xxLVWx	xxLVFx	xxLVCx	xxLIIx	xxLIWx	xxLLIx	xxMVIx
xxMVWx	xxMIIx	xxWVIx	xxIVxV	xxIVxC	xxIVxI	xxIVxL	xxIVxY	xxIVxW	
xxIVxM	xxIVxA	xxIVxT	xxIVxS	xxIVxH	xxIIXV	xxIIXC	xxIIXI	xxIIXL	
xxIIXY	xxIIXW	xxIIXM	xxIIXV	xxILxC	xxILxI	xxILxF	xxILxL	xxILxY	
xxIFxC	xxIFxI	xxICxV	xxICxC	xxIWxV	xxIWxC	xxIYxV	xxIYxC	xxIMxV	
xxVVxC	xxVVxI	xxVVxF	xxVVxL	xxVVxY	xxVVxM	xxVVxM	xxVIxV	xxVIxC	
xxVIxF	xxVIxL	xxVIxY	xxVLxV	xxVLxC	xxVLxI	xxVFxV	xxVCxV	xxFVxV	
xxFIxV	xxCVxV	xxLVxV	xxIxIV	xxIxIC	xxIxII	xxIxIF	xxIxIL	xxIxIY	
xxIxWV	xxIxWC	xxIxWI	xxIxWF	xxIxFV	xxIxFC	xxIxFI	xxIxCV	xxIxCC	
xxIxYC	xxIxMV	xxIxVV	xxIxLV	xxVxIV	xxVxIC	xxVxII	xxVxWV	xxVxWC	
xxVxCV									