arXiv:2203.00254v1 [quant-ph] 1 Mar 2022

Isolating noise and amplifying signal with quantum Cheshire cat

Soham Sau^{1,2}, Ahana Ghoshal², Debmalya Das^{3,4}, Ujjwal Sen²

¹Central University of Rajasthan, Bandarsindri, Rajasthan, 305 817, India

²Harish-Chandra Research Institute, A CI of Homi Bhabha National Institute, Chhatnag Road, Jhunsi, Prayagraj 211 019, India

³Department of Physics and Astronomy, University of Rochester, Rochester, New York 14627, USA

⁴Center for Coherence and Quantum Optics, University of Rochester, Rochester, New York 14627, USA

The quantum Cheshire cat is a phenomenon in which an object, identified with a "cat", is dissociated from a property of the object, identified with the "grin" of the cat. We propose a thought experiment, similar to this phenomenon, with an interferometric setup, where a property (a component of polarization) of an object (photon) can be separated from the object itself and can simultaneously be amplified when it is already decoupled from its object. We further show that this setup can be used to dissociate two complementary properties, e.g. two orthogonal components of polarization of a photon and identified with the grin and the snarl of a cat, from each other and one of them can be amplified while being detached from the other. Moreover, we extend the work to a noisy scenario, effected by a spin-orbit coupling -like additional interaction term in the Hamiltonian for the measurement process, with the object in this scenario being identified with a "confused Cheshire cat". We devise a gedanken experiment in which such a "confusion" can be successfully dissociated from the system, and we find that the dissociation helps in amplification of signals.

I. INTRODUCTION

In recent times, a technique known as weak value amplification, has been used to amplify small or weak signals [1]. The method relies on extracting information from a system while minimally disturbing it [2–4]. This weak measurement [5, 6] of the system is done using a small coupling strength between the system and a meter, usually described by the Hamiltonian,

$$H_0 = -g(t)\hat{A} \otimes \hat{q},\tag{1}$$

where \hat{q} is a canonical variable of the meter - conjugate to momentum \hat{p} - and g(t) is a time-dependent coupling function with a compact support near the time of measurement (normalized such that its time integral is unity). \hat{A} is the observable to be measured. After the weak measurement of the observable \hat{A} , a strong measurement of a suitable observable B is carried out on the principal system and a quantity called weak value, A_w , is defined by post-selecting an outcome $|\psi_f\rangle$. The mathematical definition of weak value of A [5–7] is given by

$$A_{w} = \frac{\left\langle \psi_{f} \middle| \hat{A} \middle| \psi_{in} \right\rangle}{\left\langle \psi_{f} \middle| \psi_{in} \right\rangle}.$$
(2)

Experimental observations of weak values have been reported in Refs. [8–15].

Although the current work discusses weak values in relation to signal amplification, it may be worthwhile to mention a few other applications of the idea. Weak values can be used in the direct measurement of a photon wavefunction [14, 16], to measure the spin Hall effect [13], in quantum state tomography [17, 18], in the geometric description of quantum states [19] and state visualization [20]. It also finds application in quantum thermometry [21], and measuring the expectation value

of non-Hermitian operators [22, 23]. Weak values have been shown to be acquire complex values [24], and weak values play important roles in the two-state vector formalism [25], a physical understanding of superoscillations [26], and in separating a quantum property from its system [27]. A property of weak values that is of special interest to us is that it can lie outside the eigenvalue spectrum of the observable \hat{A} and can even be very large [5, 6]. This aspect is exploited in weak value amplification.

A significant concern in experiments is the minimizing noise in the relevant signal and designing measurement techniques that achieve the same. A measurement strategy with a greater signal to noise ratio is typically favored by experimentalists [28–33], apart from logistics and dependencies on other constraints imposed by the particular scenario. We consider a situation in which noise may be introduced during the process of weak value amplification if the Hamiltonian in Eq. (1) has extra undesired terms. It becomes necessary, therefore, to eliminate these terms or suppress them to obtain the amplification of the signal alone.

An important ingredient of the strategy we discuss in this paper is a phenomenon known as the quantum Cheshire cat. In this gedanken experiment, based on a modified Mach-Zehnder interferometer, a photon can be detected in one arm while its circular polarization can be detected in the other arm, each being absent in the other arm, by measuring the respective weak values [27]. Thus for a particular combination of preselected and post-selected states, the photon ("cat") can be disembodied from its property ("grin"), leading to the name of the effect being chosen after a magical and enigmatic character in the celebrated literary work, Alice in Wonderland [34]. This counterintuitive phenomenon has been experimentally observed using neutron interferometry in Ref. [8] and using photon interferometry in Refs. [9–11, 35]. The concept has been further expanded upon in Refs. [36–39], and different kinds of manipulations of the photon polarization, independent of the photon, have been achieved in Refs. [40–43].

Typically in a quantum Cheshire cat setup, a photon having horizontal polarization $|H\rangle$ is fed into a 50 : 50 beam splitter BS_1 of a Mach-Zehnder interferometer, creating the pre-selected state

$$|\Psi_{in}\rangle = \frac{1}{\sqrt{2}}(i|L\rangle + |R\rangle)|H\rangle, \qquad (3)$$

where $|L\rangle$ and $|R\rangle$ represent the left and right path degrees of freedom, respectively. In the two arms of the interferometer, weak measurements of the location of the photon and that of the photon's circular polarization are carried out. The photon and its polarization are interacted weakly with appropriate meter states, resulting deflections in the latter. This interaction is of the form defined in Eq. (1). Next, an arrangement of half waveplate *HWP*, phase shifter *PS*, beam splitter *BS*₂, polarization beam splitter *PBS* and detectors D_1 , D_2 and D_3 , elaborated in Fig. 1(a), is used to post-select the state

$$\left|\Psi_{f}\right\rangle = \frac{1}{\sqrt{2}}(\left|L\right\rangle\left|H\right\rangle + \left|R\right\rangle\left|V\right\rangle),\tag{4}$$

where $|V\rangle$ refers to vertical photon polarization. For this particular post-selected state, it is seen that the distributions of the deflected meter states, center around the weak values of the observables being measured in the arms. For tracing the location of the photon, the meters, which are inserted in the left and right arms of the interferometer, measure the projectors $\hat{\Pi}_L = |L\rangle \langle L|$ and $\hat{\Pi}_R = |R\rangle \langle R|$ respectively, and similarly, the polarisation detectors measure the observables $\hat{\sigma}_z^L = \hat{\Pi}_L \otimes \hat{\sigma}_z$ and $\hat{\sigma}_z^R = \hat{\Pi}_R \otimes \hat{\sigma}_z$, where $\hat{\sigma}_z = i(|V\rangle \langle H| - |H\rangle \langle V|)$. The corresponding weak values are

$$(\Pi_L)_w = 1, \quad (\Pi_R)_w = 0 (\hat{\sigma}_z^L)_w = 0, \quad (\hat{\sigma}_z^R)_w = 1.$$
 (5)

This indicates that the photon passed through the left arm but its z component of polarization passed through the right arm.

In this paper, we present a gedanken experiment to amplify a property of a photon at a location, independent of the photon being present at the same location. We also show that the same gedanken setup can amplify one property of a photon independent of another property of the same, where the two properties are complementary and correspond to non-commuting observables. We believe this is a technologically and fundamentally useful amalgamation of the two areas, viz. the quantum Cheshire cat and weak value amplification. As a demonstration, we consider a scenario where the polarization degree of the photon interacts with another degree of freedom, and results in an additional term in the Hamiltonian that governs the measurement. We call the additional term as the "spin-orbit coupling" of the photon (cat), loaning the nomenclature from the spinorbit interaction in the relativistic treatment of an electron's dynamics. This additional term behaves as a noise component that changes the shift of the meter. The meter now gives a deflection proportional to the weak value of an effective observable which is different from the weak value of a polarization component, that was to be measured. To avoid the unwanted disturbances caused by the effect of noise in some amplification techniques, we then formulate a gedanken experiment to separate the required observable from the noisy part. We propose that using this experimental setup, it is possible to reduce the noise effect on an average (as weak value primarily is) and amplify some quantum property by weak value measurement with a certain accuracy.

The rest of the paper is organized as follows. In Sec. II, we present the thought experiment, based on the concept of quantum Cheshire cat, to amplify a property of a photon (z component of polarization) independent of the photon itself. We extend the idea to amplify one property (z component of polarization) independent of the other (x component of polarization) for a noiseless ideal situation. In Sec. III, we evaluate the weak value of the effective observable, resulting from a noisy scenario, where the weak value is obtained from the shift of the meter, weakly coupled with the system, and where the noise is incorporated as a spin-orbit coupling. We also propose an experimental setup useful to amplify the weak value of the effective observable in that section. We present a conclusion in Sec. IV.

II. AMPLIFICATION OF POLARIZATION OF A PHOTON WITHOUT THE PHOTON

Weak values can be used as a tool for amplifying small signals. Further, we have seen how a property can be separated from a quantum system using the technique of the quantum Cheshire cat. Now we aim to achieve the two phenomena simultaneously, namely the separation of a property from the object and amplification of the separated property independent of the object.

The interferometric setup of this gedanken experiment is presented in Fig. 1(b). Let us begin by considering a photon, propagating along a path degree of freedom state denoted by $|L'\rangle$, in a polarization state $\cos \frac{\theta}{2} |H\rangle + \sin \frac{\theta}{2} |V\rangle$. The photon is sent through a polarization beam splitter PBS_1 that transmits horizontally polarized light and reflects the vertically polarized one, leading to the state

$$|\Psi_1\rangle = \cos\frac{\theta}{2} |L\rangle |H\rangle + \sin\frac{\theta}{2} |R\rangle |V\rangle ,$$
 (6)

where $|L\rangle$ and $|R\rangle$ are the two possible photon paths, viz. along the transmitted and reflected paths, respectively, forming the two arms of a Mach-Zehnder interferometer. Note that $|L\rangle$ and $|L'\rangle$ are along the same path after and before the polarization beam-splitter PBS_1 .



FIG. 1. The quantum Cheshire cat without and with amplification. (a) The quantum Cheshire cat setup (without amplification). The areas shaded pink and blue carry out the pre-selection and post-selection respectively. For the latter, only the clicks of detector D_1 are selected. Weak measurements of the photon position and position of polarization are performed by interacting suitable meters weakly, in the two arms of the interferometer. (b) Setup for decoupling the *z* component of polarization of a photon from the photon itself and amplifying it simultaneously. This configuration is also applicable for dissociating the *z* component of polarization of the photon, and amplifying the former simultaneously.

In the right arm of the interferometer, we place a halfwaveplate HWP_1 that converts a vertical polarization into a horizontal one, and vice-versa, followed by a π phase shifter P, which introduces a phase $e^{i\pi}$ in the right path. We now have the state

$$|\Psi_1'\rangle = \left(\cos\frac{\theta}{2}\left|L\right\rangle - i\sin\frac{\theta}{2}\left|R\right\rangle\right)\left|H\right\rangle.$$
(7)

In the parlance of weak values and quantum Cheshire cat, this is the pre-selected state. The post-selection involves a half -waveplate HWP, a phase-shifter PS, a beam-splitter BS, a second polarization beam-splitter PBS and three detectors D_1, D_2 and D_3 . The working principle is the same as discussed for the quantum Cheshire cat scenario without the amplification requirement. Thus the clicking of the detector D_1 can once again be solely selected to obtain the post-selected state

$$\left|\Psi_{f}\right\rangle = \frac{1}{\sqrt{2}}(\left|L\right\rangle\left|H\right\rangle + \left|R\right\rangle\left|V\right\rangle).\tag{8}$$

In the two arms of the interferometric setup, weak measurements of position of the photon and its circular polarization are performed, and the corresponding weak values are obtained. The weak values of the operators Π_L and Π_R , denoting the position of the photon in the left and right arms, and $\sigma_z^{(L)}$ and $\sigma_z^{(R)}$, denoting the positions of *z* components of polarization in the two

arms, are then measured to be

$$(\hat{\Pi}_{L})_{w} = \frac{\left\langle \Psi_{f} \middle| \hat{\Pi}_{L} \middle| \Psi_{1}' \right\rangle}{\left\langle \Psi_{f} \middle| \Psi_{1}' \right\rangle} = 1$$

$$(\hat{\Pi}_{R})_{w} = \frac{\left\langle \Psi_{f} \middle| \hat{\Pi}_{R} \middle| \Psi_{1}' \right\rangle}{\left\langle \Psi_{f} \middle| \Psi_{1}' \right\rangle} = 0$$

$$(\hat{\sigma}_{z}^{L})_{w} = \frac{\left\langle \Psi_{f} \middle| \hat{\sigma}_{z}^{L} \middle| \Psi_{1}' \right\rangle}{\left\langle \Psi_{f} \middle| \Psi_{1}' \right\rangle} = 0$$

$$(\hat{\sigma}_{z}^{R})_{w} = \frac{\left\langle \Psi_{f} \middle| \hat{\sigma}_{z}^{R} \middle| \Psi_{1}' \right\rangle}{\left\langle \Psi_{f} \middle| \Psi_{1}' \right\rangle} = \tan \frac{\theta}{2}$$
(9)

Therefore, the photon is detected in the left arm and the z component of polarization is detected in the right arm with a factor which could be amplified by varying the parameter θ . Thus we have achieved the phenomenon of amplifying a property of an object independent of the object: The photon's polarization component is being amplified in the right arm of the interferometer, and the photon isn't there.

This thought experiment can be further extended by separating two orthogonal components of polarization and then amplifying one component independent of the other. If we consider the two orthogonal components to be the *x* and *z* components of polarization, with $\hat{\sigma}_x = |H\rangle \langle H| - |V\rangle \langle V|$ and $\hat{\sigma}_z = i(|V\rangle \langle H| - |H\rangle \langle V|)$. The operators $\hat{\sigma}_x^L = \hat{\Pi}_L \otimes \hat{\sigma}_x$ and $\hat{\sigma}_x^R = \hat{\Pi}_R \otimes \hat{\sigma}_x$ are measured to detect the *x* component of polarization in the left and right arms of the interferometer respectively.

The corresponding weak values turn out to be

$$(\hat{\sigma}_{x}^{L})_{w} = \frac{\left\langle \phi \mid \hat{\sigma}_{x}^{L} \mid \psi_{1} \right\rangle}{\left\langle \phi \mid \psi_{1} \right\rangle} = 1,$$

$$(\hat{\sigma}_{x}^{R})_{w} = \frac{\left\langle \phi \mid \hat{\sigma}_{x}^{R} \mid \psi_{1} \right\rangle}{\left\langle \phi \mid \psi_{1} \right\rangle} = 0.$$
 (10)

When coupled with the weak values of $\hat{\sigma}_z^L$ and $\hat{\sigma}_z^R$, these indicate that the *z* component of polarization can be amplified independent of the *x* component of polarization. The weak value of $\hat{\sigma}_z^R$ is seen to acquire a value of $\tan \frac{\theta}{2}$, which means that the weak value will result in outcomes beyond the eigenvalue spectrum in the regions $\theta \in (\pi/2, \pi)$ and $\theta \in (-\pi, -\pi/2)$.

For realization of the weak value measurement, the principal system is weakly interacted with a meter. Let us choose the meter to be a qubit system with the computational basis $\{|0\rangle_m, |1\rangle_m\}$. Suppose we intend to measure σ_z^R . The weak interaction between the system and the meter can be effected by a joint unitary $\hat{U}_{\sigma_z^R}$ that couples the *z* component of the photon polarization to the meter. A convenient unitary operator that can serve this purpose, also used in [40, 41], is given by

$$\hat{U}_{\sigma_z^R} = \frac{1}{\sqrt{2}} [(\hat{I} - \hat{\Pi}_R) \otimes \hat{\sigma}_z \otimes \hat{I} + \hat{\Pi}_R \otimes \hat{\sigma}_z \otimes R^{-1}(\theta_g) ZR(\theta(g))],$$
(11)

where $Z = |0\rangle_m \langle 0|_m - |1\rangle_m \langle 1|_m$ and

$$\begin{aligned} R(\theta_g) \left| 0 \right\rangle_m &= \cos(2\theta_g) \left| 0 \right\rangle_m + \sin(2\theta_g) \left| 1 \right\rangle_m, \\ R(\theta_g) \left| 1 \right\rangle_m &= \sin(2\theta_g) \left| 0 \right\rangle_m - \cos(2\theta_g) \left| 1 \right\rangle_m. \end{aligned}$$
(12)

Set $g = 4\theta_g$. For the post-selection given by Eq. (8), the meter goes to the state

$$\left|\Phi\right\rangle_{m} = \left|0\right\rangle + g(\sigma_{Z}^{R})_{w}\left|1\right\rangle,\tag{13}$$

with $(\sigma_Z^R)_w = 1$. Hence, the meter shows a deflection that is directly proportional to the weak value of the measured observable. Further details on the construction of these joint weak unitary evolutions can be found in [12, 44].

III. AMPLIFICATION OF THE POLARIZATION OF A PHOTON IN A NOISY SCENARIO

In the preceding section, we laid out the procedure for amplifying a property of a quantum system in the absence of the system. We now consider a scenario in which the property we are looking to amplify is affected by noise, in a certain way. We first present the situation where the weak value amplification of the noisyaffected observable is carried out. As expected, the amplified quantity will contain a contribution from the unwanted noise. In the second part, we get rid of the noise by separating it from the signal using a setup similar to that of the quantum Cheshire cat and amplify the signal.

We can conveniently take the observable to be weakly measured, as the *z* component of polarization $\hat{\sigma}_z$. We

recall that the measurement of the z component of polarization ideally requires us to set up an interaction of the form

$$H_0 = -g(t)\hat{\sigma_z} \otimes \hat{q},\tag{14}$$

between the polarization degree of freedom and a convenient meter of our choosing, with meter variable \hat{q} . Let us now consider a scenario in which there is "noise" in the interaction Hamiltonian that is analogous to a spin-orbit coupling term $\hat{L}_x \otimes \hat{\sigma}_x$. The total Hamiltonian is now

$$\hat{H} = -g\delta(t)\hat{I}\otimes\hat{\sigma}_z\otimes\hat{q} + g'\hat{L}_x\otimes\hat{\sigma}_x\otimes\hat{I}.$$
 (15)

Here *g* and *g'* are the coupling constants. The coupling between the system and the meter is an instantaneous coupling, while on the contrary, in the second term, there is no dependence of time in spin-orbit coupling. Our aim is to obtain the weak value of $\hat{\sigma}_z$. Due to the noise, the effective observable of the total system, resulting in the deflection in the meter, is different from that in the noiseless case obtained from Eq. (14) in [5, 6].

Let us consider the initial state of the system as $|\Psi_{in}\rangle$ and the post-selected state as $|\Psi_f\rangle$. Let the initial state of the meter be

$$|\Phi_{in}\rangle = \int_{-\infty}^{+\infty} dq \exp\left(-\frac{q^2}{4\Delta^2}\right) |q\rangle.$$
 (16)

After post-selection, the final state of the meter is given by

$$|\Phi_f\rangle = \langle \Psi_f | \hat{U}(t) | \Psi_{in} \rangle | \Phi_{in} \rangle = \langle \Psi_f | e^{-\frac{i}{\hbar} \int_0^t H \, dt} | \Psi_{in} \rangle | \Phi_{in} \rangle$$
(17)

where *t* is the time of measurement. As the Hamiltonians at different times do not commute, the expansion of $\hat{U}(t)$ requires using the Dyson series expansion, leading to

$$\hat{U}(t) = 1 + \sum_{n=1}^{\infty} \frac{1}{n!} \left(\frac{-i}{\hbar}\right)^n \int_0^t dt_1 \int_0^{t_1} dt_2 \cdots \int_0^{t_{n-1}} dt_{n-1} \\ \times \mathcal{J}[H(t_1)H(t_2)\dots H(t_n)].$$
(18)

We re-define the coupling constants g and g', so that we can effectively set $\hbar = 1$. Using time ordering in the expansion, we have

$$1 + (-i) \int_{0}^{t} dt_{1} H(t_{1}) - \frac{1}{2!} (-i)^{2} \int_{0}^{t} dt_{1} \int_{0}^{t_{1}} dt_{2} H(t_{1}) H(t_{2}) - \frac{1}{2!} (-i)^{2} \int_{0}^{t} dt_{2} \int_{0}^{t_{2}} dt_{1} H(t_{2}) H(t_{1}) + \dots$$
(19)

Interchanging t_1 and t_2 in the last (displayed) term, we get

$$1 + (-i) \int_{0}^{t} dt_{1} H(t_{1}) - \int_{0}^{t} dt_{1} \int_{0}^{t_{1}} dt_{2} H(t_{1}) H(t_{2}) + \dots$$
(20)

Substituting the Hamiltonian H from Eq. (15) in the above expression, the form of the unitary turns out to be

$$1 - i(g\hat{I} \otimes \hat{\sigma}_{z} \otimes \hat{q} + g't\hat{L}_{x} \otimes \hat{\sigma}_{x} \otimes \hat{I}) - [g^{2}(\hat{I} \otimes \hat{\sigma}_{z} \otimes \hat{q})^{2} + gg't(\hat{L}_{x} \otimes \hat{\sigma}_{x}\hat{\sigma}_{z} \otimes \hat{q}) + {g'}^{2}\frac{t^{2}}{2}(\hat{L}_{x} \otimes \hat{\sigma}_{x} \otimes \hat{I})^{2}] + \dots (21)$$

The measurement time *t* is chosen from the regime, $\frac{g}{g'} \ll t \ll \frac{\sqrt{g}}{g'}$. Thus we can now assume *g* and *g'* to be sufficiently small to neglect all the terms that are second order in *g* or *g'* and above. On the other hand, we retain the terms containing gg't and g'^2t^2 . Using Eq. (16), the final state of the meter after weak measurement and post-selection reads

$$\begin{split} |\Phi_f\rangle &\simeq \langle \psi_f |\psi_{in}\rangle \int dq \, e^{-\frac{q^2}{4\Delta^2}} [1 + igq(\hat{\sigma}_z)_w \\ &- ig't(\hat{L}_x \otimes \hat{\sigma}_x)_w - gg'tq(\hat{L}_x \otimes \hat{\sigma}_x \hat{\sigma}_z)_w \\ &- g'^2 \frac{t^2}{2} (\hat{L}_x \otimes \hat{\sigma}_x)_w^2] |q\rangle \,. \end{split}$$
(22)

The weak values above are obtained using the definition in Eq. (2). Now, in general, for an observable \hat{A} and post-selection of $|\Psi_f\rangle$, with A_w being the weak value of \hat{A} , the shifted meter state is given by

$$|\Phi_f\rangle \simeq \langle \psi_f | \psi_{in} \rangle \int dq \, e^{iqgA_w} \exp(-\frac{q^2}{4\Delta^2}) | q \rangle$$
. (23)

Comparing Eqs. (22) and (23), we get

$$e^{igqA_w} = 1 + igq(\hat{\sigma}_z)_w - ig't(\hat{L}_x \otimes \hat{\sigma}_z)_w$$
$$-gg'tq(\hat{L}_x \otimes \hat{\sigma}_x \hat{\sigma}_z)_w - g'^2 \frac{t^2}{2} (\hat{L}_x \otimes \hat{\sigma}_z)_w^2.$$
(24)

Let

$$a_w = -ig't(\hat{L}_x \otimes \hat{\sigma}_z)_w - {g'}^2 \frac{t^2}{2} (\hat{L}_x \otimes \hat{\sigma}_z)_w^2,$$

$$A'_w = (\hat{\sigma}_z)_w + ig't(\hat{L}_x \otimes \hat{\sigma}_x \hat{\sigma}_z)_w.$$
(25)

So, the final state of the meter can be rewritten as

$$|\Phi_f\rangle \simeq \langle \psi_f |\psi_{in}\rangle \int dq \, e^{a_w} e^{iqgA'_w} \exp(-\frac{q^2}{4\Delta^2}) |q\rangle$$
 (26)

with the corresponding *p*-representation being,

$$|\Phi_f\rangle \simeq \langle \psi_f | \psi_{in} \rangle \int dp \, e^{a_w} \exp[-\Delta^2 (p - gA'_w)^2] | p \rangle.$$
 (27)

The factor e^{a_w} does not contribute to the shift of the meter. So, the deflection of the meter is proportional to the weak value A'_w . Hence the effective observable, measured in the weak measurement conjured by the noisy Hamiltonian in Eq. (15), is given by

$$A' = \hat{\sigma}_z + ig't\hat{L}_x \otimes \hat{\sigma}_x \hat{\sigma}_z.$$
(28)



FIG. 2. Setup for amplifying the weak value of the effective observable in the noisy scenario. The *L*-splitter adds an orbital angular momentum degree of freedom to the physical system. The *L*'-splitter transmits the component of orbital angular momentum parallel to $|v_a\rangle$ and reflects any component orthogonal to it.

We can also consider a noisy interaction Hamiltonian of a different form. Precisely, we can take the noisy part of the interaction Hamiltonian to be a three-body term so that it is coupled to the meter with a coupling parameter g(t) which has a compact support near the measurement time t:

$$H' = -g(t)[\hat{I} \otimes \hat{\sigma}_z \otimes \hat{q} - \hat{L}_x \otimes \hat{\sigma}_x \otimes \hat{q}].$$
(29)

Using the same method as before, we see that the effective observable resulting in the shift of the meter to be

$$A' = \hat{\sigma}_z - \hat{L}_x \otimes \hat{\sigma}_x. \tag{30}$$

To demonstrate the working principle in either case (i.e., either when the interaction Hamiltonian is given by Eq. (29) or by Eq. (15)), let us consider a set of preselected and post-selected states as follows:

$$\begin{aligned} |\chi_{in}\rangle &= \frac{1}{\sqrt{2}} (|v_a\rangle + i \, |v_b\rangle) \otimes |H\rangle \,, \\ |\chi_f\rangle &= |v_a\rangle \otimes (\cos \alpha \, |H\rangle + \sin \alpha \, |V\rangle). \end{aligned} \tag{31}$$

In these states, the first degree of freedom represents the angular momentum component L_x , while the second degree of freedom represents polarization. We assume that the orbital quantum number l to be conserved, for the system under study, at 1. Correspondingly, the dimension of the space spanned by the eigenvectors of L_x is three-dimensional. For simplicity, we work in a scenario where one of the dimensions in this three-dimensional space is naturally or artificially forbidden. The remaining two-dimensional Hilbert space is spanned by with orthonormal vectors $|v_a\rangle$ and $|v_b\rangle$, with \hat{L}_x being represented as

$$\hat{L}_x = -i(|v_a\rangle \langle v_b| - |v_b\rangle \langle v_a|), \qquad (32)$$

where

$$|v_a\rangle = \frac{1}{\sqrt{2}} \begin{pmatrix} 1\\0\\1 \end{pmatrix}, |v_b\rangle = \begin{pmatrix} 0\\-i\\0 \end{pmatrix}, \tag{33}$$

expressed in the eigenbasis of L_z . To prepare the preselected state, we can send a photon with an initial polarization $|H\rangle$ through an arrangement, that we call the L-splitter in Fig. 2, where it acquires an angular momentum of $\frac{1}{\sqrt{2}}(|v_a\rangle + i |v_b\rangle)$. We intend to measure the weak value of $\hat{\sigma}_z$, and so bring in a meter in the initial state given by Eq. (16) and set up an interaction of the form in Eq. (14). But unknown to us, the form of interaction is actually as in Eq. (15). To get the postselected state, the photon passes through another arrangement, the L'-splitter, that transmits orbital angular momentum $|v_a\rangle$ and reflects any orthogonal component towards the detector D_3 . The transmitted photon is then passed through a polarization beam-splitter *PBS*, that is chosen so that it transmits light of polarization $\cos \alpha |H\rangle + \sin \alpha |V\rangle$ towards a detector D_1 and reflects any orthogonal polarization en route to detector D_2 . Thus by selecting the clicks of D_1 alone, we can post-select the state $|\chi_f\rangle$.

However, due to the inherent noise in the Hamiltonian of the form Eq. (15), as a result of the extra spinorbital interaction, the weak value of the effective observable $A' = \hat{\sigma}_z + ig't\hat{L}_x \otimes \hat{\sigma}_x \hat{\sigma}_z$ is measured (instead of $\hat{\sigma}_z$), and is measured to be

$$A'_w = (g't + i)\tan\alpha. \tag{34}$$

Whereas, if the unknown noise is a three-body interaction as in Eq. (29), the weak value of the effective observable given in Eq. (30) turns out to be

$$A'_w = 1 + i \tan \alpha \tag{35}$$

using the setup schematically demonstrated in Fig. 2.

So, in both the noisy situations (Eqs. (15) and (29)) we can amplify the weak value of the respective effective observables (see Eqs. (28) and (30)) by varying the parameter α using different *PBS*s. However, due to the presence of spin-orbit coupling in the system, we are unable to determine the weak value of $\hat{\sigma}_z$, as instead of $\hat{\sigma}_z$, the deflection of the meter is proportional to the weak value of an observable which contains an unwanted noise along with $\hat{\sigma}_z$. Hence, while attempting to amplify the weak value of the *z*-component of polarization, we unintentionally amplify the weak value of a constituent noise. This, e.g., may cause disadvantages in applications of quantum technologies, that depend on the weak value enhancement of the polarization degree of freedom of the system. It is plausible that presence of the noise effects considered is more probable than the ideal noiseless situation, and therefore it will be beneficial to design an experimental setup that can "disembody" the noise from the required observable.

A. Disembodiment of noise from the ideal system using quantum Cheshire cats

With the general working principle established earlier in Sec. II, we now proceed to get the amplified signal of



FIG. 3. Interferometric setup for separating the "spin-orbit coupling" -like noise, and simultaneously amplifying the signal corresponding to the chosen observable. Details in main text.

the *z*-component of polarization by disassociating, with the help of the Cheshire cat mechanism,

the noise originating from the interaction with the unintended degree of freedom (in this case, L_x) during the weak measurement process. The intended preselected and post-selected states are $|\Psi'_1\rangle$ and $|\Psi_f\rangle$, respectively (see Eqs. (7) and (8)). However, the photon passing through the L-splitter picks up a new degree of freedom, an angular momentum component given by $\frac{1}{\sqrt{2}}(|v_a\rangle + i |v_b\rangle)$, as shown in Fig. 3. So the effective preselected state is

$$|\chi_{in}'\rangle = \left(\cos\frac{\theta}{2}\left|L\right\rangle - i\sin\frac{\theta}{2}\left|R\right\rangle \otimes \frac{1}{\sqrt{2}}(\left|v_{a}\right\rangle + i\left|v_{b}\right\rangle) \otimes \left|H\right\rangle.$$
(36)

To achieve the desired outcome, we are required to carry out post-selection in

$$|\chi_f'\rangle = (\cos\alpha |L\rangle |H\rangle + \sin\alpha |R\rangle |V\rangle) \otimes |v_a\rangle.$$
(37)

To obtain this state, the beam-splitter BS_2 needs to be of a transmission coefficient $\cos^2 \alpha$ and reflection coefficient $\sin^2 \alpha$, rather than being the 50 : 50 type. Also we assume that the state passes through a device, the L'-splitter, that permits only the orbital angular momentum component along $|v_a\rangle$ towards the *PBS* and reflects any orthogonal component towards the detector D_3 , as shown in Fig. 3. The measured weak values are

$$(\hat{\sigma}_{z}^{L})_{w} = 0$$

$$(\hat{\sigma}_{z}^{R})_{w} = \tan \frac{\theta}{2} \tan \alpha$$

$$(\hat{L}_{x} \otimes \hat{\sigma}_{x})_{w}^{L} = 1$$

$$(\hat{L}_{x} \otimes \hat{\sigma}_{x})_{w}^{R} = 0$$
(38)

Therefore, we can conclude that one can amplify the weak value of the observable $\hat{\sigma}_z$ separating it from the noise part $\hat{L}_x \otimes \hat{\sigma}_x$. Moreover, the enhancement of weak value of $\hat{\sigma}_z^R$ can be more than that of the effective observables obtained in the two previous cases (described in Eqs. (34) and (35), using a linear setup, as depicted in Fig. 2), because of the extra tuning parameter θ . Therefore, even if we fix the parameter α , we are still able to amplify the weak value of $\hat{\sigma}_z$, by changing the parameter θ with the use of different *PBS*₁s. So, now we have successfully achieved our goal to amplify the weak value of the required observable by splitting up the noise from the system. And, the enhancement of the weak value of the required observable is greater than that of the effective observable detectable in the noisy situation (compare Eqs. (34) and (35) with Eq. (38)). We need to choose an initial state of suitable polarization in the preparation of the pre-selected state. Also, for measuring the weak values in an experimental procedure, we have to construct a unitary parallel to the one mentioned in the noiseless situation. For measuring $\hat{\sigma}_{z}^{R}$, the constructed unitary is almost the same as in the noiseless scenario, the only difference being that we have to incorporate an identity in the degree of freedom of orbital angular momentum, i.e., we have to use

$$\hat{U}_{\sigma_z^R} = \frac{1}{\sqrt{2}} [(\hat{I} - \hat{\Pi}_R) \otimes \hat{I} \otimes \hat{\sigma}_z \otimes \hat{I} \\ + \hat{\Pi}_R \otimes \hat{I} \otimes \hat{\sigma}_z \otimes R^{-1}(\theta_g) Z R(\theta(g)],$$
(39)

and the same for measuring $(\hat{L}_x \otimes \hat{\sigma}_x)^L$ is

$$\hat{U}_{(L_x \otimes \hat{\sigma}_x)^L} = \frac{1}{\sqrt{2}} [(\hat{I} - \hat{\Pi}_L) \otimes (\hat{L}_x \otimes \hat{\sigma}_x) \otimes \hat{I} \\ + \hat{\Pi}_L \otimes (\hat{L}_x \otimes \hat{\sigma}_x) \otimes R^{-1}(\theta_g) Z R(\theta_g)].$$
(40)

 $R(\theta_q)$ and Z are the same as in the noiseless case.

- P. B. Dixon, D. J. Starling, A. N. Jordan and J. C. Howell, Ultrasensitive Beam Deflection Measurement via Interferometric Weak Value Amplification, Phys. Rev. Lett. 102, 173601 (2009).
- [2] P. Busch and P. J. Lahti. On various joint measurements of position and momentum observables in quantum theory, Phys. Rev. D 29, 1634 (1984).
- [3] A. Barchielli, L. Lanz, and G.M. Prosperi, A model for the macroscopic description and continual observations in quantum mechanics, Il Nuovo Cimento B 72, 79 (1982).

IV. CONCLUSION

To summarize, we have proposed a thought experiment related to the quantum Cheshire cat in which a component of polarization could be amplified independent of the photon, using interferometric arrangements. Furthermore, we extended the setup to a scenario in which, of two complementary polarization components of a photon, one can be amplified while being detached from the other. Moreover, we considered a noisy scenario in which the noise is generated by a spin-orbit coupling -like interaction term in the Hamiltonian governing the measurement process. We analyzed the amplification of a chosen observable in the presence of noise, both with and without the noise term being dissociated, on average, from the object by using a quantum Cheshire cat inspired setup.

It has been pointed out in the literature that the phenomena related to the quantum Cheshire cat are "average" effects. In particular, the weak values indicate average shifts of the meter, conditioned on the pre-selected and the post-selected states, and the object and the property (or two properties of the same object) do not actually travel separately in each arm of the experimental setup, rather they do so only on an average. Nonetheless, just like for the original quantum Cheshire cat, in our setups also, it has been ensured that the weak values observed are not classical averages, but quantum mechanical mean shifts, obtained by considering coupling between the photon or polarization component and a meter. Further discussions on this point can e.g. be found in [41]. Let us also note here that the amplifications reported does not imply that the amplified values could become arbitrarily large - there appear non-linear effects that limit the amplified values [45].

ACKNOWLEDGMENTS

The authors from Harish-Chandra Research Institute acknowledge partial support from the Department of Science and Technology, Government of India through the QuEST grant (grant number DST/ICPS/QUST/Theme-3/2019/120).

- [4] C. M. Caves, Quantum mechanics of measurements distributed in time. A path-integral formulation, Phys. Rev. D 33, 1643 (1986).
- [5] Y. Aharonov, D.Z. Albert and L. Vaidman, *How the result of a measurement of a component of the spin of a spin-1/2 particle can turn out to be 100*, Phys. Rev. Lett. **60**, 1351 (1988).
- [6] I. M. Duck, P. M. Stevenson and E. C. G. Sudarshan, The sense in which a "weak measurement" of a spin-—' particle's spin component yields a value 100, Phys. Rev. D 40, 2112 (1989).

- [7] Y. Aharonov, S. Popescu, and J. Tollaksen, A timesymmetric formulation of quantum mechanics, Phys. Today 63, 27 (2010).
- [8] T. Denkmayr, H. Geppert, S. Sponar, H. Lemmel, A. Matzkin, J. Tollaksen and Y. Hasegawa, Observation of a quantum Cheshire Cat in a matter-wave interferometer experiment, Nat. Commun. 5, 4492 (2014).
- [9] R. Corrêa , M. F. Santos , C. H. Monken and P. Saldanha 'Quantum Cheshire Cat' as simple quantum interference, New J. Phys. 17 053042 (2015).
- [10] S. Sponar, T. Denkmayr, H. Geppert and Y. Hasegawa, Fundamental Features of Quantum Dynamics Studied in Matter-Wave Interferometry—Spin Weak Values and the Quantum Cheshire-Cat, Atoms 4, 11 (2016).
- [11] J. M. Ashby, P. D. Schwarz and M. Schlosshauer, Observation of the quantum paradox of separation of a single photon from one of its properties, Phys.Rev.A 94, 012102 (2016).
- [12] G. J. Pryde, J.L. O'Brien, A. G. White, T. C. Ralph and H. M. Wiseman, *Measurement of Quantum Weak Values of Pho*ton Polarization, Phys.Rev.Lett. 94, 220405 (2005).
- [13] O. Hosten and P. Kwiat, Observation of the Spin Hall Effect of Light via Weak Measurements, Science 319, 787 (2008).
- [14] J. S. Lundeen, B. Sutherland, A. Patel, C. Stewart and C. Bamber, *Direct measurement of the quantum wavefunction*, Nature (London) **474**, 188 (2011).
- [15] M. Cormann, M. Remy, B. Kolaric and Y. Caudano, *Revealing geometric phases in modular and weak values with a quantum eraser*, Phys. Rev. A 93, 042124 (2016).
- [16] J. S. Lundeen and C. Bamber, Procedure for Direct Measurement of General Quantum States Using Weak Measurement, Phys. Rev. Lett. 108, 070402 (2012).
- [17] H. F. Hofmann, Complete characterization of post-selected quantum statistics using weak measurement tomography, Phys. Rev. A 81, 012103 (2010).
- [18] S. Wu, State tomography via weak measurements, Sci. Rep. 3, 1193 (2013).
- [19] M. Cormann and Y. Caudano, Geometric description of modular and weak values in discrete quantum systems using the Majorana representation, J. Phys. A: Math. Theor. 50, 305302 (2017).
- [20] H. Kobayashi, K. Nonaka and Y. Shikano, Stereographical visualization of a polarization state using weak measurements with an optical-vortex beam, Phys. Rev. A 89, 053816 (2014).
- [21] A. K. Pati, C. Mukhopadhyay, S. Chakraborty and S. Ghosh, *Quantum precision thermometry with weak measurements*, Phys. Rev. A 102, 012204 (2020).
- [22] A. K. Pati, U. Singh and U. Sinha, Measuring non-Hermitian operators via weak values, Phys. Rev. A 92, 052120 (2015).
- [23] G. Nirala, S. N. Sahoo, A. K. Pati and U. Sinha, Measuring average of non-Hermitian operator with weak value in a Mach-Zehnder interferometer, Phys. Rev. A 99, 022111 (2019).
- [24] R. Jozsa, Complex weak values in quantum measurement, Phys. Rev. A 76, 044103 (2007).
- [25] C. A. Chatzidimitriou-Dreismann, Weak Values and Two-State Vector Formalism in Elementary Scattering and Reflectivity—A New Effect, Universe 5, 58 (2019).

- [26] M. Nairn, Superoscillations: Realisation of quantum weak values, arXiv:2109.14404.
- [27] Y. Aharonov, S. Popescu, D. Rohrlich and P. Skrzypczyk, *Quantum Cheshire Cats*, New J. Phys. 15, 113015 (2013).
- [28] G. Scala, M. D'Angelo, A. Garuccio, S. Pascazio and F. V. Pepe, Signal-to-noise properties of correlation plenoptic imaging with chaotic light, Phys. Rev. A 99, 053808 (2019).
- [29] L. Li, X. Li, B. Zhang and L. You, Enhancing test precision for local Lorentz-symmetry violation with entanglement, Phys. Rev. A 99, 042118 (2019).
- [30] J.-H. Huang, F.-F. He, X.-Y. Duan, G.-J. Wang, X.-Y. Hu, Improving the precision of weak-value-amplification with two cascaded Michelson interferometers based on Vernier-effect, arXiv:2107.14395
- [31] Y. Jin, J. Yan, S. J. Rahman, X. Yu, and J. Zhang, *Imaging the dipole scattering of an optically levitated dielectric nanoparticle*, Appl. Phys. Lett. **119**, 021106 (2021).
- [32] Y.-Z. Ma, M. Jin, D.-L. Chen, Z.-Quan Zhou, C.-Feng Li and G.-C. Guo, *Elimination of noise in optically rephased pho*ton echoes, Nat. Commun. **12**, 4378 (2021).
- [33] L. Bai, L. Zhang, Y. Yang, R. Chang, Y. Qin, J. He, X. Wen, J. Wang, Enhancement of spin noise spectroscopy of rubidium atomic ensemble by using of the polarization squeezed light, arXiv:2111.09572
- [34] L. Carroll, Alice's Adventures in Wonderland (Macmillan, London, 1865).
- [35] Y. Kim and DG. Im, and YS. Kim, Observing the quantum Cheshire cat effect with noninvasive weak measurement, npj Quantum Inf 7, 13 (2021).
- [36] J.-D. Bancal, Quantum physics: Isolate the subject, Nature Phys. 10, 11 (2014).
- [37] D. P. Atherton, G. Ranjit, A. A. Geraci, J. D. Weinstein, Observation of a classical Cheshire cat in an optical interferometer, Opt. Lett. 40 879 (2015).
- [38] Q. Duprey, S. Kanjilal, U. Sinha, D. Home and A. Matzkin, *The Quantum Cheshire Cat effect: Theoretical basis and obser*vational implications, Ann. Phys. **391** 1 (2018).
- [39] Y. Aharonov, E. Cohen, and S. Popescu, A dynamical quantum Cheshire Cat effect and implications for counterfactual communication, Nat Commun 12, 4770 (2021).
- [40] D. Das and A. K. Pati, Can two quantum Cheshire cats exchange grins?, New J. Phys., 22, 063032 (2020).
- [41] D. Das and U. Sen, Delayed choice of paths in the quantum paradox of separating different properties of a photon, Phys. Rev. A 103, 012228 (2021).
- [42] P. Chowdhury, A. K. Pati, and J.L. Chen, Wave and particle properties can be spatially separated in a quantum entity Photon. Res. 9, 1379 (2021).
- [43] Zheng-Hao Liu, Wei-Wei Pan, Xiao-Ye Xu, Mu Yang, Jie Zhou, Ze-Yu Luo, Kai Sun, Jing-Ling Chen, Jin-Shi Xu, Chuan-Feng Li, Guang-Can Guo, *Experimental exchange* of grins between quantum Cheshire cats, Nat. Commun. 11, 3006 (2020).
- [44] Kim, Y., Kim, YS., Lee, SY. et al. Direct quantum process tomography via measuring sequential weak values of incompatible observables. Nat Commun 9, 192 (2018).
- [45] T. Koike and S. Tanaka, *Limits on amplification by Aharonov-Albert-Vaidman weak measurement*, Phys. Rev. A 84, 062106 (2011).