

LOWER DEFECT GROUPS AND VERTICES OF SIMPLE MODULES

AKIHIKO HIDA AND MASAO KIYOTA

ABSTRACT. We compare lower defect groups associated with p -regular classes and vertices of simple modules for a block of a finite group algebra. We show that lower defect groups are contained in vertices of simple modules after suitable reordering. Moreover, for a block of principal type, we show that a p -regular lower defect group which contains a vertex of simple module is a defect group of the block.

1. INTRODUCTION

Let G be a finite group. Let \mathcal{O} be a complete discrete valuation ring of characteristic 0 such that the residue field $k = \mathcal{O}/J(\mathcal{O})$ is an algebraically closed field of characteristic $p > 0$. Let ZkG be the center of the group algebra kG . We call a primitive idempotent of ZkG a block of kG . In this paper, we consider the relation between two important families of p -subgroups associated with a block b of kG , lower defect groups of b and vertices of simple kGb -modules.

A conjugacy class of G is called p -regular if the order of its element is not divisible by p . The p -regular conjugacy classes of G are distributed to blocks of kG . Let $\text{Cl}(G_{p'})$ be the set of all p -regular classes of G and

$$\text{Cl}(G_{p'}) = \bigcup_b \text{Cl}_{p'}(b)$$

a decomposition of $\text{Cl}(G_{p'})$ into blocks of kG . Let $\text{Cl}_{p'}(b) = \{C_i\}_{1 \leq i \leq l(b)}$ and Q_i a defect group of C_i , namely, Q_i is a Sylow p -subgroup of $C_G(x_i)$ where $x_i \in C_i$. Then Q_i ($1 \leq i \leq l(b)$) are called lower defect groups of b associated with p -regular classes.

Let $\{S_i\}_{1 \leq i \leq l(b)}$ be a representatives of complete set of isomorphism classes of simple kGb -modules. It is known that this set and $\text{Cl}_{p'}(b)$ have the same cardinality. We want to compare Q_i and a vertex $\text{vx}(S_j)$ of S_j . These are both subgroups of some defect group of b . For example, if a defect group P of b is abelian, then $\text{vx}(S_i) = P$ for all i by the theorem of Knörr [9]. Hence in this case, $Q_i \leq_G \text{vx}(S_j)$ for all i, j . Our first result shows that the inclusion $Q_i \leq_G \text{vx}(S_i)$ holds in general for all i if we renumber the indices suitably.

Proposition 1.1. *Let b be a block of kG . Let $\{S_i\}_{1 \leq i \leq l(b)}$ be a set of representatives of isomorphism classes of simple kGb -modules. Let $\{Q_i\}_{1 \leq i \leq l(b)}$ be the lower defect groups of b associated with p -regular classes. Then there exists a permutation σ of $\{1, 2, \dots, l(b)\}$ such that*

$$Q_i \leq_G \text{vx}(S_{\sigma(i)})$$

for all i . In particular,

$$\prod_{i=1}^{l(b)} |Q_i| \leq \prod_{i=1}^{l(b)} |\text{vx}(S_i)|.$$

Next we consider when the equality holds in Proposition 1.1. We show that, under some assumption, if the equality holds then $Q_i = \text{vx}(S_{\sigma(i)})$ is a defect group of b . A block b is of principal type if the image of Brauer homomorphism with respect Q is a block of $kC_G(Q)$ for every p -subgroup Q contained in a defect group of b . For a p -subgroup Q of G , we denote by $m_b^1(Q)$ the multiplicity of Q as a lower defect group of b associated with p -regular classes, that is, $m_b^1(Q)$ is a number of i ($1 \leq i \leq l(b)$) such that $Q_i =_G Q$. The following theorem is our main result.

Theorem 1.2. *Let b be a block of principal type of kG and P a defect group of b . Let S be a simple kGb -module and R a vertex of S . If $R \leq Q < P$, then $m_b^1(Q) = 0$.*

By Proposition 1.1 and Theorem 1.2, we have the following Corollary.

Corollary 1.3. *Let $\{x_i\}_{1 \leq i \leq l}$ be a set of representatives of p -regular conjugacy classes of G . Let Q_i be a Sylow p -subgroup of $C_G(x_i)$. Let $\{S_i\}_{1 \leq i \leq l}$ be a set of representatives of isomorphism classes of simple kG -modules. Then*

$$\prod_{i=1}^l |Q_i| \leq \prod_{i=1}^l |\text{vx}(S_i)|$$

and the equality holds if and only if G is p -nilpotent.

We prove these results in section 3. In section 4, we consider blocks of p -solvable groups. We show a result which is slightly weaker than Theorem 1.2 without the assumption that b is of principal type. Finally, in section 5, we add some results on the complexity of modules.

In this paper, all kG -modules are finite generated right modules. For an indecomposable kG -module M , we denote a vertex of M by $\text{vx}(M)$. We refer to [6], [10], [11] and [12] for modular representations of finite groups.

2. LOWER DEFECT GROUPS OF A BLOCK

Lower defect groups of a block are defined by Brauer [3] and some related results can be found in [5], [8], [13] or [6, V, Section 10], [12, Chapter 5, Section 11]. In this section we quote some known results used in this paper. We consider lower defect groups associated with p -regular classes only.

Let $G_{p'}$ be the set of all p -regular elements of G . We denote by $kG_{p'}$ the k subspace of kG spanned by $G_{p'}$. If C is a conjugacy class of G , we set $\hat{C} = \sum_{x \in C} x \in kG$. Then $\{\hat{C} \mid C \in \text{Cl}(G)\}$ is a basis of the center ZkG of kG . Let $\text{Cl}(G_{p'})$ be the set of all p -regular conjugacy classes of G . Then $\{\hat{C} \mid C \in \text{Cl}(G_{p'})\}$ is a basis of $ZkG_{p'} = ZkG \cap kG_{p'}$. On the other hand,

$$ZkG_{p'} = \bigoplus_b ZkG_{p'}b = \bigoplus_b ZkG'_p \cap ZkGb$$

where b ranges over all blocks of kG and there exists a disjoint decomposition

$$\text{Cl}(G_{p'}) = \bigcup_b \text{Cl}_{p'}(b)$$

such that $\{\hat{C}b \mid C \in \text{Cl}_{p'}(b)\}$ is a basis of $ZkG_{p'}b$ for any block b .

Let $\text{Cl}_{p'}(b) = \{C_i\}_{1 \leq i \leq l(b)}$ and Q_i a defect group of C_i , that is, Q_i is a Sylow p -subgroup of $C_G(x_i)$ where $x_i \in C_i$. We denote by $m_b^1(Q)$ the multiplicity of Q as a lower defect group of b associated with p -regular classes, that is, $m_b^1(Q)$ is a number of i ($1 \leq i \leq l(b)$) such that $Q =_G Q_i$. Here, for subgroups H, K of G , we write $H =_G K$ if H and K are G conjugate.

Let M be a kG -module and $H \subset K$ be subgroups of G . Let M^H be the set of fixed points of H in M . We denote by Tr_H^K the trace map from M^H to M^K defined by

$$\text{Tr}_H^K(m) = \sum_{g \in H \backslash K} mg.$$

We set $M_H^K = \text{Tr}_H^K(M^H)$. The group G acts on kG and $kG_{p'}$ by conjugation and we can define the trace maps

$$\text{Tr}_H^K : (kG)^H \longrightarrow (kG)^K$$

and

$$\text{Tr}_H^K : (kG_{p'})^H \longrightarrow (kG_{p'})^K.$$

The proof of the following basic properties of lower defect groups are found in [6, V, Section 10] and [12, Chapter 5, Section 11].

Proposition 2.1. *Let b be a block of kG and P a defect group of b . Then the following holds.*

- (1) *If Q is a p -subgroup of G , then $m_b^1(Q) = \dim(kG_{p'})_Q^G / \sum_{R < Q} (kG_{p'})_R^G$.*
- (2) *If $m_b^1(Q) > 0$ then Q is conjugate to a subgroup of P and $m_b^1(P) = 1$.*
- (3) *Let $\text{Cl}_{p'}(b) = \{C_i\}_{1 \leq i \leq l(b)}$ and Q_i a defect group of C_i . Then $l(b) = |\text{Cl}_{p'}(b)|$ is the number of isomorphism classes of simple kGb -modules and $\{|Q_i| \mid 1 \leq i \leq l(b)\}$ is the set of elementary divisors of the Cartan matrix of kGb .*
- (4) *Let $m \geq 0$. Then $\sum_Q m_b^1(Q)$ where Q ranges over the set of representatives of conjugacy classes of subgroups of G of order p^m is the multiplicity of p^m as an elementary divisor of the Cartan matrix of kGb .*

We need the following result in Lemma 3.4 and Theorem 3.5.

Proposition 2.2 ([5, Proposition (II) 1.3]). *If Q is a p -subgroup of G then*

$$m_b^1(Q) = \dim(kC_G(Q)_{p'})_Q^{N_G(Q)} \text{Br}_Q(b)$$

where $\text{Br}_Q : ZkG \longrightarrow ZkN_G(Q)$ is the Brauer homomorphism.

3. PROOF OF MAIN RESULTS

The following Proposition is proved in [7, Theorem 10, Corollary]. It is obtained from [4, p.243, Corollaire] or [6, IV, Theorem 2.3] also.

Proposition 3.1. *Let M be an indecomposable kG -module.*

- (1) *Let φ be the Brauer character corresponding to M . Let $x \in G_{p'}$ and $Q \in \text{Syl}_p(C_G(x))$. If*

$$\varphi(x) \notin p\mathcal{O}$$

then $\text{Res}_Q^G M$ has an indecomposable direct summand N such that $\dim N \not\equiv 0 \pmod{p}$.

- (2) *If Q is a p -subgroup of G and $\text{Res}_Q^G M$ has an indecomposable direct summand N such that $\dim N \not\equiv 0 \pmod{p}$, then $Q \leq_G \text{vx}(M)$.*

Proposition 1.1 is immediate from this Proposition.

(Proof of Proposition 1.1)

Let φ_i be the Brauer character of S_i . Let $\text{Cl}_{p'}(b) = \{C_i\}_{1 \leq i \leq l(b)}$ be the p -regular conjugacy classes distributed into b in the block decomposition of $\text{Cl}(G_{p'})$. Fix $x_i \in C_i$ for each i . We may assume that Q_i is a Sylow p -subgroup of $C_G(x_i)$ where $x_i \in C_i$. Then the determinant of the matrix $(\varphi_i(x_j))_{1 \leq i, j \leq l(b)}$ is not contained in $J(\mathcal{O})$ by [12, Chapter 5, Theorem 11.6] and there exists a permutation σ such that

$$\prod_i \varphi_{\sigma(i)}(x_i) \notin J(\mathcal{O}).$$

Since $p\mathcal{O} \subset J(\mathcal{O})$, the result follows from Proposition 3.1.

Next we prove Theorem 1.2. First we study the case that Q is a normal subgroup of G .

Lemma 3.2. *Let $G \triangleright H$. Let b be a block of kH of defect 0. Let $T = T(b)$ be the inertial group of b in G . Assume that $|T : H| \equiv 0 \pmod{p}$. Then $b(kH)_1^G = 0$.*

Proof. By the Mackey decomposition, we have

$$(kH)_1^G \subset \sum_{t \in G/T} ((kH)^t)_1^T = (kH)_1^T.$$

Since b is T -invariant, we have

$$b(kH)_1^T = b \text{Tr}_1^T(kH) = \text{Tr}_1^T(bkH) = \text{Tr}_H^T(\text{Tr}_1^H(bkH)).$$

Then

$$\text{Tr}_1^H(bkH) = Z(bkH) = kb$$

since b is a block of defect 0, and

$$\text{Tr}_H^T(\text{Tr}_1^H(bkH)) = \text{Tr}_H^T(kb) = 0$$

since $|T : H| \equiv 0 \pmod{p}$. □

Lemma 3.3. *Let Q be a normal p -subgroup of G and $\overline{G} = G/Q$. Let $\mu : kG \rightarrow k\overline{G}$ be the surjective algebra homomorphism induced by the natural surjective homomorphism $G \rightarrow \overline{G}$. Let b be a block of kG and $\overline{b} = \mu(b)$. Then μ induces an isomorphism*

$$b \cdot (kC_G(Q)_{p'})_Q^G \cong \overline{b} \cdot (k(\overline{QC_G(Q)})_{p'})_{\overline{1}}^{\overline{G}}$$

Proof. The natural surjective homomorphism $G \rightarrow \overline{G}$ induces a bijection

$$C_G(Q)_{p'} \rightarrow \overline{(QC_G(Q))}_{p'} = (QC_G(Q)/Q)_{p'}.$$

Hence μ induces an isomorphism

$$\mu_0 : kC_G(Q)_{p'} \simeq k(\overline{QC_G(Q)})_{p'}.$$

On the other hand, since

$$\mu(b \cdot (kC_G(Q)_{p'})_Q^G) = \overline{b} \cdot (k(\overline{QC_G(Q)})_{p'})_{\overline{1}}^{\overline{G}}$$

and

$$b \cdot (kC_G(Q)_{p'})_Q^G \subset kC_G(Q)_{p'}, \quad \overline{b} \cdot (k(\overline{QC_G(Q)})_{p'})_{\overline{1}}^{\overline{G}} \subset k(\overline{QC_G(Q)})_{p'},$$

it follows that the restriction of μ_0 induces the desired isomorphism. □

Lemma 3.4. *Let Q be a normal p -subgroup of G and b a block of kG . Let P be a defect group of b . Assume that $Q < P$ and $C_P(Q) = Z(Q)$. Then $b \cdot (kC_G(Q)_{p'})_Q^G = 0$, in particular, $m_b^1(Q) = 0$.*

Proof. Let $H = QC_G(Q)$. The block b is a central idempotent of kH . Let $b = \sum_i b_i$ be the block decomposition of b in kH . Let $T_i = T(b_i)$ be the inertial group of b_i in G . Then there exists a defect group P_i of b such that $P_i \leq T_i$ and $P_i \cap H$ is a defect group of b_i . Since P_i is conjugate to P in G and $C_P(Q) = Z(Q)$,

$$P_i \cap H = P \cap H = Q$$

and it follows that Q is a defect group of b_i . Let $\mu : kG \rightarrow k\overline{G} = k(G/Q)$ be the natural surjective algebra homomorphism. Then $\mu(b_i) = \overline{b}_i$ is a block of $k\overline{H}$ of defect 0 for all i . Since $Q < P$, we have $H \neq HP_i$ and $|T_i : H| \equiv 0 \pmod{p}$. The inertial group of \overline{b}_i in \overline{G} is $\overline{T}_i = T_i/Q$ and $|\overline{T}_i : \overline{H}| \equiv 0 \pmod{p}$. Hence

$$\overline{b}_i(k\overline{H})_{\overline{T}_i}^{\overline{G}} = 0$$

by Lemma 3.2 and

$$b(k\overline{H})_{\overline{T}_i}^{\overline{G}} = \left(\sum_i \overline{b}_i \right) (k\overline{H})_{\overline{T}_i}^{\overline{G}} = 0.$$

Then, by Lemma 3.3,

$$b(kC_G(Q)_{p'})_Q^G \simeq \overline{b}(k\overline{H}_{p'})_{\overline{T}_i}^{\overline{G}} \subset \overline{b}(k\overline{H})_{\overline{T}_i}^{\overline{G}} = 0.$$

Moreover, since Q is a normal subgroup of G , $\text{Br}_Q(b) = b$ and the result follows from Proposition 2.2. \square

In the following, we consider a p -subgroup Q such that $C_P(Q) = Z(Q)$ for any defect group P of b containing Q .

Theorem 3.5. *Let b be a block of kG and Q a p -subgroup of G . Assume that Q is a proper subgroup of a defect group of b . If $C_P(Q) = Z(Q)$ for any defect group P of b containing Q , then $m_b^1(Q) = 0$.*

Proof. Let $N = N_G(Q)$. Let

$$\text{Br}_Q : ZkG \rightarrow ZkN$$

be the Brauer homomorphism and

$$\text{Br}_Q(b) = \sum_i b_i$$

a decomposition into block idempotents of kN . Then $b_i^G = b$ and $Q < D_i$ by Brauer's First Main Theorem where D_i is a defect group of b_i . There exists a defect group P_i of b such that $D_i \leq P_i$. Then

$$C_{D_i}(Q) \leq C_{P_i}(Q) = Z(Q)$$

by the assumption. Hence

$$b_i \cdot (kC_G(Q)_{p'})_Q^N = 0$$

by Lemma 3.4 and

$$(kC_G(Q)_{p'})_Q^N \text{Br}_Q(b) = (kC_G(Q)_{p'})_Q^N \left(\sum_i b_i \right) = 0.$$

Hence the result follows by Proposition 2.2. \square

Lemma 3.6. *Let b be a block of kG and (P, e) a maximal (G, b) -Brauer pair. Let $\mathcal{F} = \mathcal{F}_{(P, e)}(G, b)$ be the fusion system of b with respect to (P, e) . Suppose that $\mathcal{F} = \mathcal{F}_P(G)$. Let Q be a proper subgroup of P . If Q is \mathcal{F} -centric, then $m_b^1(Q) = 0$.*

Proof. Let P_1 is a defect group of b such that $Q < P_1$. Then there exists $g \in G$ such that $P^g = P_1$. Let $Q_1 = {}^gQ$. Then

$$\varphi : Q \longrightarrow Q_1(\leq P), \quad \varphi(u) = gug^{-1}$$

is an isomorphism in $\mathcal{F} = \mathcal{F}_P(G)$. Hence $C_P(Q_1) = Z(Q_1)$ since Q is \mathcal{F} -centric and it follows that $C_{P_1}(Q) = Z(Q)$. Hence b satisfies the assumption of Theorem 3.5. \square

Now we prove Theorem 1.2 and Corollary 1.3. A block b of kG is of principal type if $\text{Br}_Q(b)$ is a block of $kC_G(Q)$ for every p -subgroup Q contained in a defect group of b ([11, Definition 6.3.13]).

(Proof of Theorem 1.2)

There exists a (G, b) -Brauer pair (R, f) such that $Z(R)$ is a defect group of the block f of $kC_G(R)$ by the theorem of Knörr ([9, 3.6 Corollary], [11, Corollary 10.3.2]). Let (P, e) be a maximal (G, b) -Brauer pair such that $(R, f) \leq (P, e)$. Let $\mathcal{F} = \mathcal{F}_{(P, e)}(G, b)$ be the fusion system of b with respect to (P, e) . Then R is an \mathcal{F} -centric subgroup of P by [11, Proposition 8.5.3]. Moreover, since b is a block of principal type, we have $\mathcal{F} = \mathcal{F}_P(G)$ by [11, Proposition 8.5.5]. Since $R \leq Q \leq P$, Q is an \mathcal{F} -centric subgroup by [11, Proposition 8.2.4]. Hence the results follows from Lemma 3.6.

Remark 3.7. In Theorem 1.2, S does not necessarily need to be simple. Let b be a block of $\mathcal{O}G$ with defect group P and S an indecomposable $\mathcal{O}Gb$ -module. If $\underline{\text{End}}_{\mathcal{O}Gb}(S) \cong \mathcal{O}/J(\mathcal{O})^m$ for some $m > 0$, then there is a (G, b) -Brauer pair (R, f) such that $Z(R)$ is a defect group of the block f of $kC_G(R)$ as in the proof of Theorem 1.2 by [11, Corollary 10.3.2]. It follows that if b is of principal type and $R \leq Q < P$, then $m_b^1(Q) = 0$.

(Proof of Corollary 1.3)

The inequality holds by Proposition 1.1. If G is p -nilpotent, then kGb has a unique simple module (up to isomorphism) for every block b of kG . The vertex of the simple kGb -module is a defect group of b and that is a p -regular lower defect group of b . Hence the equality holds. On the other hand, assume that the equality holds. If P is a defect group of a block b of kG , then $m_b^1(P) = 1$. Hence the principal block $b_0(kG)$ of kG has a unique simple module (up to isomorphism) by Proposition 1.1 and Theorem 1.2 since $b_0(kG)$ is a block of principal type by Brauer's Third Main Theorem ([11, Theorem 6.3.14]). Hence G is p -nilpotent.

4. BLOCKS OF p -SOLVABLE GROUPS

Let N be a normal subgroup of G . Let b be a block of kG . Let c be the block of kN such that $bc \neq 0$ and H the inertial group of c in G . Then there exists a block d of kH such that $db = d$ and $\text{Tr}_H^G(d) = b$. The (kGb, kHd) -bimodule $bkGd = kGd$

induces a Morita equivalence between kGb and kHd . In particular, $l(b) = l(d)$. The trace map induces an algebra isomorphism

$$\mathrm{Tr}_H^G : Z(kHd) \longrightarrow Z(kGb).$$

Moreover if P is a defect group of d then P is a defect group of b (cf.[11, Theorem 6.8.3]). With these notations, we have the following two lemmas.

Lemma 4.1. *If Q is a subgroup of H , then Tr_H^G induces an isomorphism*

$$\mathrm{Tr}_H^G : \sum_{t \in G} (kH_{p'})_{Q^t \cap H}^H d \simeq (kG_{p'})_Q^G b.$$

Proof. Tr_H^G induces an injective k -linear map

$$\mathrm{Tr}_H^G : (kH_{p'})_R^H d \longrightarrow (kG_{p'})_R^G b$$

for any subgroup R of H . In particular,

$$\mathrm{Tr}_H^G((kH_{p'})_{Q^t \cap H}^H d) \subset (kG_{p'})_{Q^t}^G b = (kG_{p'})_Q^G b$$

for $t \in G$. On the other hand,

$$(kG_{p'})_Q^G \subset \sum_{t \in G} (kG_{p'})_{Q^t \cap H}^H$$

and

$$\begin{aligned} (kG_{p'})_Q^G d \subset \sum_{t \in G} (kG_{p'})_{Q^t \cap H}^H d &= \sum_{t \in G} ((kH_{p'})_{Q^t \cap H}^H + k(G_{p'} - H_{p'})_{Q^t \cap H}^H) d \\ &= \sum_{t \in G} (kH_{p'})_{Q^t \cap H}^H d \end{aligned}$$

since $gdg = 0$ for any $g \in G - H$. Hence

$$(kG_{p'})_Q^G b = \mathrm{Tr}_H^G((kG_{p'})_Q^G d) \subset \mathrm{Tr}_H^G\left(\sum_{t \in G} (kH_{p'})_{Q^t \cap H}^H d\right).$$

□

Lemma 4.2. *Let U be a p -subgroup of H . Let*

$$\mathcal{U} = \{W \leq H \mid W =_G U\}$$

and let $\{U_j\}$ be a set of representatives of H -conjugacy classes of \mathcal{U} . Then

$$\sum_j m_d^1(U_j) = m_b^1(U).$$

Proof. Let $\{R_{ij}\}_{1 \leq i \leq m, 1 \leq j \leq r(i)}$ be a set of representatives of H -conjugacy classes of subgroups of H of order $|U|$ such that

$$R_{ij} =_G R_{i'j'} \iff i = i'.$$

We may assume $U = R_{11}$ and $\{U_j\} = \{R_{1j}\}$. We set $R_i = R_{i1}$. For each i , Tr_H^G induces a k -linear map

$$\Phi_i : \bigoplus_{1 \leq j \leq r(i)} \left((kH_{p'})_{R_{ij}}^H d / \sum_{R < R_{ij}} (kH_{p'})_R^H d \right) \longrightarrow (kG_{p'})_{R_i}^G b / \sum_{R < R_i} (kG_{p'})_R^G b.$$

We claim that Φ_i is an isomorphism. For $t \in G$, if $R_i^t \leq H$ then $R_i^t =_H R_{ij}$ for some j . On the other hand, if $R_i^t \cap H < R_i^t$ then

$$\mathrm{Tr}_H^G((kH_{p'})_{R_i^t \cap H}^H d) \subset (kG_{p'})_{R_i^t \cap H}^G b = (kG_{p'})_{R_i \cap H}^G b \subset \sum_{R < R_i} (kG_{p'})_R^G b.$$

Hence we have

$$(kG_{p'})_{R_i}^G b = \sum_{t \in G} \mathrm{Tr}_H^G((kH_{p'})_{R_i^t \cap H}^H d) \subset \mathrm{Tr}_H^G \left(\sum_j (kH_{p'})_{R_{ij}}^H d \right) + \sum_{R < R_i} (kG_{p'})_R^G b$$

by Lemma 4.1 and Φ_i is surjective. Hence

$$\sum_{1 \leq j \leq r(i)} m_d^1(R_{ij}) \geq m_b^1(R_i)$$

for each i and

$$\sum_{i,j} m_d^1(R_{ij}) \geq \sum_i m_b^1(R_i).$$

But $\sum_{i,j} m_d^1(R_{ij})$ is the multiplicity of $|U|$ as an elementary divisor of the Cartan matrix C_d of kHd and $\sum_i m_b^1(R_i)$ is that of the Cartan matrix C_b of kGb . Since kGb and kHd are Morita equivalent, $C_d = C_b$ and it follows that

$$\sum_{i,j} m_d^1(R_{ij}) = \sum_i m_b^1(R_i).$$

Hence

$$\sum_{1 \leq j \leq r(i)} m_d^1(R_{ij}) = m_b^1(R_i)$$

and Φ_i is an isomorphism for each i . In particular, for $i = 1$, we have

$$\sum_j m_d^1(U_j) = m_b^1(U).$$

□

The following theorem is the main result of this section. If the block b is of principal type, then this is a consequence of Theorem 1.2.

Theorem 4.3. *Let G be a p -solvable group. Let b be a block of kG . Let $\mathrm{Cl}_{p'}(b) = \{C_i\}_{1 \leq i \leq l(b)}$ and Q_i a defect group of C_i . Let $\{S_i\}_{1 \leq i \leq l(b)}$ be a set of representatives of isomorphism classes of simple kGb -modules. Then there exists a permutation σ of $\{1, \dots, l(b)\}$ such that*

$$Q_i \leq_G \mathrm{vx}(S_{\sigma(i)})$$

for all i and

$$Q_i <_G \mathrm{vx}(S_{\sigma(i)})$$

unless Q_i is a defect group of b .

Proof. Let c be a block of $kO_{p'}(G)$ such that $bc \neq 0$ and H the inertial group of c in G . Then there exists a block d of kH such that $db = d$ and $\mathrm{Tr}_H^G(d) = b$. Moreover (kGb, kHd) -bimodule $bkGd = kGd$ induces a Morita equivalence between kGb and kHd . In particular, $l(b) = l(d)$. Moreover if P is a defect group of d then P is a defect group of b .

If $G = H$, then the result holds by Proposition 1.1 and Theorem 1.2 since $b = d$ is a block of principal type by [11, Lemma 10.6.5].

Suppose that $G > H$. Let $\text{Cl}_{p'}(b) = \{C_i\}$ and $\text{Cl}_{p'}(d) = \{\tilde{C}_i\}$. Let Q_i (resp. \tilde{Q}_i) be a defect group of C_i (resp. \tilde{C}_i). If Q is a p -subgroup of P ,

$$|\{1 \leq j \leq l(d) \mid \tilde{Q}_j =_G Q\}| = m_b^1(Q) = |\{1 \leq i \leq l(b) \mid Q_i =_G Q\}|$$

by Lemma 4.2. Hence we may assume $Q_i = \tilde{Q}_i$ for every $1 \leq i \leq l(b)$. Let $\tilde{S}_i = S_i \otimes_{kGb} kGd$ be the simple kHd -module corresponding to S_i . Then $\text{vx}(\tilde{S}_i) =_G \text{vx}(S_i)$. By induction there exists a permutation σ of $\{1, \dots, l(d)\}$ such that

$$Q_i \leq_H \text{vx}(\tilde{S}_{\sigma(i)})$$

for all i and

$$Q_i <_H \text{vx}(\tilde{S}_{\sigma(i)})$$

if $Q_i <_H P$. Since P is a defect group of b , it follows that

$$Q_i \leq_G \text{vx}(S_{\sigma(i)})$$

for all i and

$$Q_i <_G \text{vx}(S_{\sigma(i)})$$

unless Q_i is a defect group of b . \square

The following corollary is a block version of Corollary 1.3 for p -solvable groups.

Corollary 4.4. *Let G be a p -solvable group. Let b be a block of kG . Let $\{S_i\}_{1 \leq i \leq l(b)}$ be a set of representatives of isomorphism classes of simple kGb -modules. Then*

$$\det C_b \leq \prod_{i=1}^{l(b)} |\text{vx}(S_i)|$$

where C_b is the Cartan matrix of kGb and the equality holds if and only if $l(b) = 1$.

5. COMPLEXITY OF MODULES

Let Q be a p -subgroup of G and M an indecomposable kG -module. Suppose that $\text{Res}_Q^G M$ has a direct summand N such that $\dim N \not\equiv 0 \pmod{p}$. We set $|\text{vx}(M)| = p^{v(M)}$ and $|Q| = p^a$. Then by Proposition 3.1, $Q \leq_G \text{vx}(M)$ and in particular $a \leq v(M)$. If the dimension of a source of M is divisible by p , then proper inequality $a < v(M)$ holds. We consider another information on this inequality related to the complexity of M . For the complexity of a module, we refer to [1] and [2, Section 5].

Let $c(M)$ be the complexity of M and $r(M)$ the p -rank of $\text{vx}(M)$. Since M is a direct summand of $\text{Ind}_{\text{vx}(M)}^G \text{Res}_{\text{vx}(M)}^G M$ and $c(\text{Res}_{\text{vx}(M)}^G M) \leq r(M)$, we have

$$c(M) \leq r(M) \leq v(M).$$

Proposition 5.1. *Let Q be a p -subgroup of G and M an indecomposable kG -module. Suppose that $\text{Res}_Q^G M$ has a direct summand N such that $\dim N \not\equiv 0 \pmod{p}$. If $|Q| = p^a$ then*

$$a \leq v(M) + c(M) - r(M).$$

Proof. Since N is a direct summand of $\text{Res}_Q^G M$ and $\dim N \not\equiv 0 \pmod{p}$,

$$\text{rank}(Q) = c(N) \leq c(\text{res}_Q^G M) \leq c(M)$$

and

$$\text{rank}(Q) + v(M) - r(M) \leq v(M) + c(M) - r(M)$$

where $\text{rank}(Q)$ is the p -rank of Q . We may assume that $Q \leq \text{vx}(M)$ by Proposition 3.1(2). Let E be an elementary abelian p -subgroup of $\text{vx}(M)$ of maximal rank and F an elementary abelian p -subgroup of Q of maximal rank. The class of elementary abelian p -groups satisfies the condition in Lemma 5.2 below. Hence $|Q|/|F| \leq |\text{vx}(M)|/|E|$ by Lemma 5.2 and we have

$$a \leq \text{rank}(Q) + v(M) - r(M).$$

□

Lemma 5.2. *Let \mathcal{X} be a class of finite p -groups which satisfies the following property:*

$$P \in \mathcal{X}, P \triangleright R \Rightarrow R \in \mathcal{X}.$$

Let P be a p -group and Q be a subgroup of P . Suppose $E \leq P$, $F \leq Q$ and $E, F \in \mathcal{X}$. If $|L| \leq |F|$ for any subgroup $L \leq Q$ such that $L \in \mathcal{X}$, then

$$|Q : F| \leq |P : E|.$$

Proof. We proceed by induction on $|P : Q|$. If $P = Q$ then $|E| \leq |F|$ by the assumption. Hence $|Q : F| \leq |P : E|$. Assume that $P > Q$ and let R be a maximal subgroup of P such that $Q \leq R < P$. Then $R \cap E \in \mathcal{X}$ since $P \triangleright R$ and $E \triangleright R \cap E$. If $R \not\geq E$, then $P = RE$ and $P/R = RE/R \cong E/R \cap E$. It follows that $|P|/|E| = |R|/|R \cap E|$ and $|R|/|R \cap E| \geq |Q|/|F|$ by induction. If $R \geq E$, then $|P|/|E| > |R|/|E|$ and it follows that $|R|/|E| \geq |Q|/|F|$ by induction. □

REFERENCES

- [1] J. L. Alperin, L. Evens, Representations, resolutions and Quillen's dimension theorem, J. Pure Appl. Algebra 22 (1981) 1-9.
- [2] D. J. Benson, Representations and cohomology II: Cohomology of groups and modules, Cambridge studies in advanced mathematics 31, Cambridge University Press, 1991.
- [3] R. Brauer, Defect groups in the theory of representations of finite groups, Illinois J. Math. 13 (1969) 53-73.
- [4] M. Broué, Sur l'induction des modules indécomposables et la projectivité relative, Math. Z. 149 (1976) 227-245.
- [5] M. Broué, Brauer coefficients of p -subgroups associated with a p -block of a finite group, J. Algebra 56 (1979) 365-383.
- [6] W. Feit, The representation theory of finite groups, North-Holland, 1982.
- [7] J. A. Green, On the indecomposable representations of a finite group, Math. Z. 70 (1958/59) 430-445.
- [8] K. Iizuka, A note on blocks of characters of a finite group, J. Algebra 20 (1972) 196-201.
- [9] R. Knörr, On the vertices of irreducible modules, Ann. of Math. 110 (1979) 487-499.
- [10] M. Linckelmann, The block theory of finite group algebras, Vol. 1, London Math. Soc. Student Texts 91, Cambridge University Press, 2018.
- [11] M. Linckelmann, The block theory of finite group algebras, Vol. 2, London Math. Soc. Student Texts 92, Cambridge University Press, 2018.
- [12] H. Nagao, Y. Tsushima, Representations of finite groups, Academic Press, 1989.
- [13] J. B. Olsson, Lower defect groups, Comm. Algebra 8 (1980) 261-288.

AKHIKO HIDA, FACULTY OF EDUCATION, SAITAMA UNIVERSITY, SHIMO-OKUBO 255, SAKURA-KU, SAITAMA-CITY, SAITAMA, 338-8570, JAPAN
Email address: ahida@mail.saitama-u.ac.jp

MASAO KIYOTA, COLLEGE OF LIBERAL ARTS AND SCIENCES, TOKYO MEDICAL AND DENTAL UNIVERSITY, KOHNODAI 2-8-30, ICHIKAWA, CHIBA, 272-0827, JAPAN
Email address: kiyota.las@tmd.ac.jp