AFFINE RSK CORRESPONDENCE AND CRYSTALS OF LEVEL ZERO EXTREMAL WEIGHT MODULES

JAE-HOON KWON AND HYUNSE LEE

ABSTRACT. We give an affine analogue of the Robison-Schensted-Knuth (RSK) correspondence, which generalizes the affine Robinson-Schensted correspondence by Chmutov-Pylyavskyy-Yudovina. The affine RSK map sends a generalized affine permutation of period (m, n) to a pair of tableaux (P, Q) of the same shape, where P belongs to a tensor product of level one perfect Kirillov-Reshetikhin crystals of type $A_{m-1}^{(1)}$, and Q belongs to a crystal of extremal weight module of type $A_{n-1}^{(1)}$ when $m, n \ge 2$. We consider two affine crystal structures of types $A_{m-1}^{(1)}$ and $A_{n-1}^{(1)}$ on the set of generalized affine permutations, and show that the affine RSK map preserves the crystal equivalence. We also give a dual affine Robison-Schensted-Knuth correspondence.

1. INTRODUCTION

The Robinson-Schensted-Knuth (simply RSK) correspondence [12] is a fundamental algorithm in algebraic combinatorics with rich applications and generalizations in various areas. In representation theory, we may regard it as a combinatorial aspect of Howe duality [6], where the pair of groups (GL_m , GL_n) acts as mutual centralizers on the symmetric or exterior algebra generated by the tensor product of their natural representations. From a view point of crystal base theory, the bijection can be further understood as an isomorphism of ($\mathfrak{gl}_m, \mathfrak{gl}_n$)-bicrystals [14].

The Robinson-Schensted (simply RS) correspondence, a special case of RSK correspondence restricted to $n \times n$ permutation matrices, gives a bijection from the symmetric group \mathfrak{S}_n on n letters to the set of pairs of standard tableaux of the same shape, and it provides a combinatorial tool to study the left and right cells of \mathfrak{S}_n in Kazhdan-Lusztig theory [11].

An affine analogue of the RS correspondence has been introduced by Shi [20, 21] in the study of affine Kazhdan-Lusztig cells, which maps an affine permutation w to a pair of tabloids (P(w), Q(w)), but not necessarily in a injective way. Recently in [3], Chmutov-Pylyavskyy-Yudovina construct a bijection from the set of extended affine permutations w to the set of triples $(P(w), Q(w), \rho)$, which recovers the Shi's algorithm and characterizes each fiber of (P(w), Q(w)), where ρ is an integral vector satisfying a condition so called dominance. The main ingredient of the affine RS correspondence in [3] is the *affine matrix-ball construction*, which generalizes the matrix-ball construction in the usual RSK algorithm [4].

²⁰¹⁰ Mathematics Subject Classification. 17B37, 22E46, 05E10.

This work is supported by the National Research Foundation of Korea(NRF) grant funded by the Korea government(MSIT) (No. 2019R1A2C1084833 and 2020R1A5A1016126).

In this paper we show that the affine RS correspondence in [3] can be naturally extended to a bijection on the set of generalized affine permutations, which also admits an isomorphism of affine crystals.

Let m and n be positive integers greater than 1. Let $\widehat{\mathcal{M}}_{m \times n}$ be the set of matrices $A = (a_{ij})_{i,j \in \mathbb{Z}}$ with non-negative integral entries such that $a_{i+mj+n} = a_{ij}$ for all $i, j \in \mathbb{Z}$, and for each j, $a_{ij} = 0$ except for finitely many *i*'s. We show that there exists a bijection

(1.1)
$$\kappa: \ \widehat{\mathcal{M}}_{m \times n} \longrightarrow \bigsqcup_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} CSST_{[m]}(\lambda) \times \mathcal{B}_n(\lambda) ,$$
$$A \longmapsto (P_0, Q)$$

where the union is over the partitions λ with length no more than $\min\{m, n\}$, $CSST_{[m]}(\lambda)$ is the set of column semistandard tableaux of shape λ with entries from 1 to m, and $\mathcal{B}_n(\lambda)$ is the crystal of an extremal weight module of type $A_{n-1}^{(1)}$ with extremal weight corresponding to λ [9]. The bijection is also well-defined when m = 1 or n = 1.

The bijection κ recovers the affine RS correspondence in [3] after a suitable standardization. One key observation is that the dominance condition on ρ in [3] fits nicely into the description of connected components in $\mathcal{B}_n(\lambda)$, and hence gives the image of κ as in (1.1).

We further show that κ is compatible with affine crystal structure. Note that $CSST_{[m]}(\lambda)$ is a finite affine crystal of type $A_{m-1}^{(1)}$ with classical weights, which is isomorphic to the tensor product of Kirillov-Reshetikhin crystals corresponding to fundamental representations. We consider two affine crystal structures of types $A_{m-1}^{(1)}$ and $A_{n-1}^{(1)}$ on $\widehat{\mathcal{M}}_{m \times n}$, where the Kashiwara operators for $A_{m-1}^{(1)}$ commute with those for $A_{n-1}^{(1)}$. Then we show that the bijection κ preserves the crystal equivalence for these crystal structures. Indeed, we show that κ commutes with the Kashiwara operators for both A_{m-1} and $A_{n-1}^{(1)}$, while the operators \widetilde{e}_0 and \widetilde{f}_0 on A for $A_{m-1}^{(1)}$ may change the second component Q in (1.1) under κ .

As a corollary, we have an isomorphism of $A_{n-1}^{(1)}$ -crystals with affine weights of level 0

$$\widehat{\mathcal{M}}_{m \times n} \cong \bigoplus_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} \mathcal{B}_n(\lambda)^{\oplus m_\lambda}$$

where $m_{\lambda} = |CSST_{[m]}(\lambda)|$ and \oplus means a disjoint union of crystals. We expect that these affine crystal structures on $\widehat{\mathcal{M}}_{m \times n}$ coincide with the ones in [15]. We also have a dual affine RSK correspondence, where $\widehat{\mathcal{M}}_{m \times n}$ is replaced by the set of binary matrices and $CSST_{[m]}(\lambda)$ is replaced by the set of row semistandard tableaux of the conjugate shape λ' .

Finally, we remark that an affine RSK correspondence is also given in [7], where the algorithm is given by using dynamics of Sagan-Stanley's skew RSK correspondence [19], but its algorithm looks different from the one by affine matrix-ball construction. It would be interesting to compare these two algorithms.

The paper is organized as follows: In Section 2, we introduce necessary background and notions for the affine RSK algorithm, which are analogous to those introduced in [3]. In Section 3, we construct the affine RSK correspondence κ in (1.1) (Theorem 3.9), where the proof of its well-definedness and bijectivity is given in Section 5. In Section 4, we discuss

affine crystal structure on both sides of (1.1), and its compatibility with κ (Theorem 4.14). The proof is given in Section 6. In Section 7, we give a dual affine RSK correspondence.

Acknowledgement The authors would like to thank D. Kim for kind explanation on the work [3].

2. Preliminaries

Let us introduce some necessary notations and terminologies following [3] with slight modification and generalization.

2.1. Generalized affine permutations. Throughout the paper, let m and n denote the positive integers. Let \mathbb{Z}_+ denote the set of non-negative integers and $[k] = \{1, 2, ..., k\}$ for $k \ge 1$ with the usual linear order.

Let

$$\widehat{\mathcal{M}}_{m \times n} = \left\{ A = (a_{ij})_{i,j \in \mathbb{Z}} \middle| \begin{array}{c} (1) \ a_{ij} \in \mathbb{Z}_+ \text{ and } a_{i+mj+n} = a_{ij} \text{ for all } i, j \in \mathbb{Z}, \\ (2) \text{ for each } j, \ a_{ij} = 0 \text{ except for finitely many } i\text{'s} \end{array} \right\}$$

We call $\widehat{\mathcal{M}}_{m \times n}$ the set of generalized affine permutations of period (m, n). For $A \in \widehat{\mathcal{M}}_{m \times n}$, let

$$supp(A) = \{ c = (i, j) \in \mathbb{Z}^2 \mid a_{ij} > 0 \}$$

be the support of $A \in \widehat{\mathcal{M}}_{m \times n}$. We denote by \mathbb{O} the zero matrix.

Note that $A \in \widehat{\mathcal{M}}_{m \times n}$ can be viewed as an *extended affine permutation*, an element of the extended affine Weyl group of type $A_{n-1}^{(1)}$ when $m = n \ge 2$ and $\sum_{j \in \mathbb{Z}} a_{ij} = \sum_{i \in \mathbb{Z}} a_{ij} = 1$ for each *i* and *j*. Following [3], we call *A* a *partial (extended affine) permutation* if $\sum_{j \in \mathbb{Z}} a_{ij} = \sum_{i \in \mathbb{Z}} a_{ij} \le 1$ for each *i* and *j*.

An element c = (i, j) in the lattice \mathbb{Z}^2 will be often called a *cell*. We regard each cell as a position of the entry a_{ij} in $A = (a_{ij}) \in \widehat{\mathcal{M}}_{m \times n}$, where the row index *i* (resp. the column index *j*) is increasing from top to bottom (resp. from left to right). With this convention, we define partial orders $>_{\text{NW}}, \ge_{\text{nw}}$ and \leq_{ne} on \mathbb{Z}^2 as follows:

- (1) $c_1 >_{NW} c_2$ if and only if $i_1 < i_2$ and $j_1 < j_2$,
- (2) $c_1 \ge_{nw} c_2$ if and only if $i_1 \le i_2$ and $j_1 \le j_2$,
- (3) $c_1 \leq_{ne} c_2$ if and only if $i_1 \geq i_2$ and $j_1 \leq j_2$,

for $c_1 = (i_1, j_1), c_2 = (i_2, j_2) \in \mathbb{Z}^2$. By convention, we use N (or E, W, S) to emphasize strict inequality, while n (or $\mathbf{e}, \mathbf{w}, \mathbf{s}$) allows equality of the co-ordinates of cells.

Let $\tau = \tau_{m,n}$ denote a bijection on \mathbb{Z}^2 given by

$$\tau(i,j) = (i+m,j+n) \quad ((i,j) \in \mathbb{Z}^2).$$

Example 2.1. The following is an example of a generalized affine permutation in $\mathcal{M}_{4\times 5}$.



The co-ordinate of * is (1, 1) and hence the co-ordinates of the cells with red entries are (-1, -6), (3, -1), (7, 4), (11, 9) from northwest to southeast, while those with blue entries are (-2, 3), (2, 8), (6, 13), (10, 18).

2.2. Standardization. Let $M = (m_1, \ldots, m_s) \in \mathbb{Z}_+^s$ be a matrix with a single row and let $r = m_1 + \cdots + m_s$. Suppose that r > 0. Let $R = R_1 \sqcup \cdots \sqcup R_s$ be an index set with r elements, where each R_k is a linearly ordered set with m_k elements. We assume that R_k is an empty set if $m_k = 0$. We assume that there is a linear order on R, where x > y if and only if $x \in R_i$ and $y \in R_j$ with i < j, for x and y in different R_k 's. Then we define $R \times [s]$ -matrix $M^\circ = (m_{ij}^\circ)_{i \in R, j \in [s]}$ by

(2.1)
$$m_{ij}^{\circ} = \begin{cases} 1 & \text{if } i \in R_j, \\ 0 & \text{otherwise} \end{cases}$$

If r = 0, then we define $M^{\circ} = M$. Note that $\sum_{j=1}^{s} m_{ij}^{\circ} = 1$ for $i \in R$ and $\sum_{i \in R} m_{ij}^{\circ} = m_j$ for $1 \leq j \leq s$. For example, if M = (1, 1, 0, 3), then

$$M^{\circ} = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix},$$

which is a 5 × 4-matrix with $R = [5], R_1 = \{5\}, R_2 = \{4\}, R_3 = \emptyset$, and $R_4 = \{1, 2, 3\}$.

Given linearly ordered sets ${\mathcal A}$ and ${\mathcal B},$ let

$$\mathcal{M}_{\mathcal{A}\times\mathcal{B}} = \left\{ M = (m_{ab})_{a\in\mathcal{A},b\in\mathcal{B}} \mid m_{ab}\in\mathbb{Z}_+, \ \sum_b m_{ab} < \infty \ (a\in\mathcal{A}), \ \sum_a m_{ab} < \infty \ (b\in\mathcal{B}) \right\}.$$

Let $M \in \mathcal{M}_{\mathcal{A} \times \mathcal{B}}$ be given. We define M° to be the matrix obtained from M by replacing each row $M_a = (m_{ab})_{b \in \mathcal{B}}$ with M_a° for $a \in \mathcal{A}$ as in (2.1). We may assume that M° is a matrix in $\mathcal{M}_{\mathcal{A}^{\circ} \times \mathcal{B}}$, where \mathcal{A}° is a set with a linear order < satisfying

- (1°) \mathcal{A}° has a partition parametrized by \mathcal{A} , that is, $\mathcal{A}^{\circ} = \bigsqcup_{a \in \mathcal{A}} \mathcal{A}_{a}^{\circ}$,
- (2°) $|\mathcal{A}_a^{\circ}| = \max\{\sum_b m_{ab}, 1\}$ for each $a \in \mathcal{A}$,
- (3°) $x_1 < x_2$ for $x_1 \in \mathcal{A}_{a_1}^\circ$ and $x_2 \in \mathcal{A}_{a_2}^\circ$ with $a_1 < a_2$.

We define $M^{\circ'}$ in a similar way with respect to the columns of M, that is, $M^{\circ'} = ((M^t)^{\circ})^t$, where M^t denotes the transpose of M. Then we define the *standardization of* M to be

$$M^{\mathtt{st}} = (M^{\circ})^{\circ'}.$$

By definition, we have $M^{\mathtt{st}} = (M^{\circ})^{\circ'} = (M^{\circ'})^{\circ}$, and $M^{\mathtt{st}} \in \mathfrak{M}_{\mathcal{A}^{\circ} \times \mathfrak{B}^{\circ}}$, where \mathfrak{B}° is a linearly ordered set satisfying the properties $(1^{\circ})-(3^{\circ})$ with respect to M^{t} .

Remark 2.2. Let $M = (m_1, \ldots, m_s) \in \mathbb{Z}_+^s$ and $R = R_1 \sqcup \cdots \sqcup R_s$ be given as above. We define an $R \times [s]$ matrix M^{\bullet} in the same way as in M° by reversing a linear order on R, that is, x < y if and only if $x \in R_i$ and $y \in R_j$ with i < j, for x and y in different R_k 's. Then M^{\bullet} is given by rearranging the rows of M° in a reverse way. For M = (1, 1, 0, 3), we have

$$M^{\bullet} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

In general, for $M \in \mathfrak{M}_{\mathcal{A}\times\mathfrak{B}}$, we define M^{\bullet} and $M^{\bullet'} = ((M^t)^{\bullet})^t$. We may define another versions of standardization of M using \circ, \circ', \bullet and \bullet' .

2.3. Standardization of generalized affine permutations. Suppose that $A = (a_{ij}) \in \widehat{\mathcal{M}}_{m \times n} \subset \mathcal{M}_{\mathbb{Z} \times \mathbb{Z}}$ is given, and consider A^{st} . We assume that the subsets [m] and [n] in the row and column indices of A are replaced by

$$I = I_1 \sqcup \cdots \sqcup I_m, \quad J = J_1 \sqcup \cdots \sqcup J_n,$$

in A^{st} , where we have $x_1 < x_2$ for $x_1 \in X_{i_1}$ and $x_2 \in X_{i_2}$ with $i_1 < i_2$ (X = I, J). Note that the entry 1 of A^{st} at $(k, l) \in I_i \times J_j$ is obtained from the non-zero entry a_{ij} at $(i, j) \in \text{supp}(A)$ by applying \circ and \circ' .

Let M = |I| and N = |J|. Identifying I and J with [M] and [N], respectively, we put

(2.2)
$$I_{i+kM} = I_i + kM, \quad J_{j+kN} = J_j + kN \quad (i \in [m], \ j \in [n], \ k \in \mathbb{Z}).$$

Then we have

$$\mathbb{Z} = \bigsqcup_{i \in \mathbb{Z}} I_i = \bigsqcup_{j \in \mathbb{Z}} J_j.$$

With this identification, we have $A^{st} \in \widehat{\mathcal{M}}_{M \times N}$. If M < N (resp. M > N), then we may add additional N - M (resp. M - N) elements, for example, to I_m (resp. J_n) and then the trivial rows (resp. columns) to A^{st} corresponding to those indices and their translates in (2.2). So we may assume that

where $L = \max\{M, N\}$. In particular A^{st} is a partial extended affine permutation.

Definition 2.3. For $A \in \widehat{\mathcal{M}}_{m \times n}$, we call A^{st} in (2.3) a standardization of A. For $c = (i, j) \in$ supp(A), we denote by A_c^{st} the matrix in $\mathcal{M}_{\mathbb{Z} \times \mathbb{Z}}$, which is equal to A^{st} at the positions of $(k, l) \in I_i \times J_j$ and has zero entries elsewhere.

Remark 2.4. Note that A_c^{st} has an $a_{ij} \times a_{ij}$ block submatrix at $I_i \times J_j$ with 1 on the antidiagonal, and zero entries elsewhere.

The following lemmas can be checked easily.

Lemma 2.5. Let $c_1, c_2 \in \text{supp}(A^{st})$ be given with $c_1 = (i_1, j_1)$ and $c_2 = (i_2, j_2)$.

- (1) If $i_1 < i_2$ and $c_1, c_2 \in I_i \times \mathbb{Z}$ for some $i \in \mathbb{Z}$, then $c_2 \leq_{ne} c_1$.
- (2) If $j_1 < j_2$ and $c_1, c_2 \in \mathbb{Z} \times J_j$ for some $j \in \mathbb{Z}$, then $c_1 \leq_{ne} c_2$.

Lemma 2.6. Let $c_1, c_2 \in \text{supp}(A^{\text{st}})$ be given with $c_i \in \text{supp}\left(A_{c'_i}^{\text{st}}\right)$ for some $c'_i \in \text{supp}(A)$ (i = 1, 2). Then we have

(1) $c_2 >_{\text{NW}} c_1$ if and only if $c'_2 >_{\text{NW}} c'_1$, (2) $c_2 = \tau_{L,L}(c_1)$ implies $c'_2 = \tau_{m,n}(c'_1)$.



Example 2.7. Let m = 3, n = 4. If $A \in \widehat{\mathfrak{M}}_{3 \times 4}$ is a generalized affine permutation given by



2.4. **Proper numberings.** Suppose that a non-zero $A \in \widehat{\mathcal{M}}_{m \times n}$ is given.

Definition 2.8. A numbering on A is a function $d : \operatorname{supp}(A) \longrightarrow \mathbb{Z}$. It is called *proper* if

- (1) $d(c_2) < d(c_1)$ for $c_1, c_2 \in \text{supp}(A)$ with $c_2 >_{NW} c_1$,
- (2) for $c_1 \in \text{supp}(A)$, there exists $c_2 \in \text{supp}(A)$ with $c_2 >_{\text{NW}} c_1$ and $d(c_2) = d(c_1) 1$.

The conditions (1) and (2) are called *monotone* and *continuous*, respectively.

The notion of proper numbering on a partial permutation was introduced in [3]. It also plays an important role when we describe an affine analogue for the matrix-ball construction of the RSK correspondence in this paper.

Let d be a numbering on A. Let d^{st} be a numbering on $supp(A^{st})$ given by

$$d^{\mathtt{st}}(c) = d(c')$$
 if $c \in \operatorname{supp}(A_{c'}^{\mathtt{st}})$ for some $c' \in \operatorname{supp}(A)$,

for $c \in \operatorname{supp}(A^{\mathtt{st}})$.

Lemma 2.9. We have the following:

- (1) d is a proper numbering on A if and only if d^{st} is a proper numbering on A^{st} ,
- (2) any proper numbering on A^{st} is given by d^{st} for a unique proper numbering d on A.

Proof. (1) Since no two cells corresponding to non-zero entries in A_c^{st} ($c \in \text{supp}(A)$) are comparable with respect to $>_{NW}$ (see Remark 2.4), it follows from Lemma 2.6(1) that d satisfies the conditions Definition 2.8(1) and (2) if and only if d^{st} does so.

(2) Let d' be a proper numbering on A^{st} . We claim that $d' = d^{st}$ for some proper numbering d on A. By Lemma 2.6(1), it suffices to show that d' is constant on A_c^{st} for

 $\overline{7}$

each $c \in \operatorname{supp}(A)$. Suppose that it does not hold. Then there exist $c_1, c_2 \in A_c^{st}$ for some $c \in \operatorname{supp}(A)$ such that $d'(c_1) < d'(c_2)$. By Definition 2.8(1), there exists $c_3 \in A_{c'}^{st}$ for some $c' \in \operatorname{supp}(A)$ such that $c' >_{NW} c$ and $d'(c_3) = d'(c_1)$. Since $c_3 >_{NW} c_1$ by Lemma 2.6, it is a contradiction. This proves the claim.

Lemma 2.10 (cf. [3]). Under the above hypothesis, the following hold.

- (1) For any proper numbering d on A, there exists a positive integer ℓ , which we call period, such that $d(\tau(c)) = d(c) + \ell$ for $c \in \text{supp}(A)$.
- (2) If ℓ and ℓ' are the periods of any two proper numberings d and d' on A, respectively, then we have $\ell = \ell'$, which we call the width of A.

Proof. The assertions are proved in [3, Proposition 3.4], when A is a partial permutation. Therefore for arbitrary A, (1) and (2) follow from Lemmas 2.6(2), 2.9 and [3, Proposition 3.4].

2.5. Streams and channels. Let us verify the existence of a proper numbering on $A \in \widehat{\mathcal{M}}_{m \times n}$. This can be done by using standardization and the result in case of partial permutations in [3].

Definition 2.11. A stream is an infinite collection of cells $\mathbf{s} = \{c_i\}_{i \in \mathbb{Z}}$, which is invariant under τ and forms a chain with respect to $>_{\mathbb{NW}}$, that is, $c_i >_{\mathbb{NW}} c_{i+1}$ for $i \in \mathbb{Z}$. A flow of a stream \mathbf{s} is a positive integer l such that $\tau(c_i) = c_{i+l}$ for $i \in \mathbb{Z}$. A defining data of a stream $\mathbf{s} = \{c_i = (a_i, b_i)\}_{i \in \mathbb{Z}}$ of flow l is a triple $(\mathbf{a}, \mathbf{b}, r)$, where

- (1) $\mathbf{a} = (a_{1+r_1}, \dots, a_{l+r_1})$ with $1 \leq a_{1+r_1} < \dots < a_{l+r_1} \leq m$,
- (2) $\mathbf{b} = (b_{1+r_2}, \dots, b_{l+r_2})$ with $1 \leq b_{1+r_2} < \dots < b_{l+r_2} \leq n$,
- (3) $r = r_1 r_2$.

Suppose that a non-zero $A \in \widehat{\mathcal{M}}_{m \times n}$ is given.

Definition 2.12. A stream **s** is called a *stream of* A if $\mathbf{s} \subset \text{supp}(A)$. A stream **s** of A is called a *channel of* A if its flow is maximal among the streams of A.

Let $C = \{c_i\}_{i \in \mathbb{Z}}$ be a channel of A and let $c \in \text{supp}(A)$. Let k be the maximal integer such that $c_k >_{\text{NW}} c$. The maximal property of channel forces that $c >_{\text{NW}} c_{k+1}$. This implies either $c_{k+1} \leq_{\text{ne}} c$ or $c \leq_{\text{ne}} c_{k+1}$. In other words, we have

$$\operatorname{supp}(A) = C_{\operatorname{ne}} \cup C_{\operatorname{sw}},$$

where

(2.4)
$$C_{ne} = \{ c \in \operatorname{supp}(A) \mid c' \leq_{ne} c \text{ for some } c' \in C \},$$

$$C_{sw} = \{ c \in \operatorname{supp}(A) \mid c \leq_{ne} c' \text{ for some } c' \in C \},$$

and $C_{ne} \cap C_{sw} = C$. Let \mathcal{C}_A denote the set of channels of A, which is finite. For $C_1, C_2 \in \mathcal{C}_A$, we define

(2.5)
$$C_1 \geq_{sw} C_2$$
 if and only if $C_1 \subset (C_2)_{sw}$.

Then \geq_{sw} is a partial order on \mathcal{C}_A .

Let $\mathbf{s} = \{c_i\}_{i \in \mathbb{Z}}$ be a stream of A. Let $\mathbf{s}^{\mathtt{st}} = \{c_i^{\mathtt{st}}\}_{i \in \mathbb{Z}}$ be a collection of cells such that $c_i^{\mathtt{st}}$ is the cell corresponding to the most southwest 1 in the block submatrix $A_{c_i}^{\mathtt{st}}$ for $i \in \mathbb{Z}$.

On the other hand, let $\mathbf{t} = \{c'_i\}_{i \in \mathbb{Z}}$ be a stream of $A^{\mathtt{st}}$. Let \mathbf{t}' be a collection of cells obtained from \mathbf{t} by replacing each c'_i by the cell corresponding to the most southwest nonzero entry in $A_c^{\mathtt{st}}$ if $c'_i \in \mathrm{supp}(A_c^{\mathtt{st}})$ for $c \in \mathrm{supp}(A)$. Also let $\overline{\mathbf{t}} = \{\overline{c}_i\}_{i \in \mathbb{Z}}$ be a collection of cells in $\mathrm{supp}(A)$ such that $c_i \in \mathrm{supp}(A_{\overline{c}i}^{\mathtt{st}})$ for $i \in \mathbb{Z}$.

Lemma 2.13. Under the above hypothesis, we have the following:

- (1) $\mathbf{s}^{\mathtt{st}}$ is a stream of $A^{\mathtt{st}}$ with the same flow as \mathbf{s} ,
- (2) \mathbf{t}^{\checkmark} and $\overline{\mathbf{t}}$ are streams of $A^{\mathtt{st}}$ and A with the same flow as \mathbf{t} , respectively,
- (3) **t** is a channel if and only if \mathbf{t}' is a channel, in which case we have $\mathbf{t}' \geq_{sw} \mathbf{t}$,
- (4) **s** is a channel if and only if \mathbf{s}^{st} is a channel,
- (5) for any channel \mathbf{s}' of A, $\mathbf{s} \geq_{sw} \mathbf{s}'$ if and only if $\mathbf{s}^{st} \geq_{sw} \mathbf{s}'^{st}$.

Proof. It is straightforward to check (1) and (2). The assertion (3) follows from the fact that no two cells of non-zero entries in A_c^{st} for $c \in \text{supp}(A)$ belong to a stream of A^{st} . The assertion (4) follows from the fact that for any stream **t** of A^{st} , there exists a unique stream **s** of A such that $\mathbf{s}^{st} = \mathbf{t}^{\checkmark}$. This implies (5) immediately.

Proposition 2.14 (cf. [3]). The set \mathcal{C}_A has a greatest element with respect to \geq_{sw} , which we denote by C_A^{sw} .

Proof. As in the proof of Lemma 2.10, we reduce the proof to the case when A is a partial permutation. Let C_1 and C_2 be two channels of A. By Lemma 2.13, C_1^{st} and C_2^{st} be two channels of A^{st} . By [3, Proposition 3.13], there exists a channel C of A^{st} such that $C \geq_{sw} C_1^{st}, C_2^{st}$. Let C_0 be a channel of A such that $C_0^{st} = C^{\checkmark}$. Hence we have $C_0^{st} = C^{\checkmark} \geq_{sw} C_1^{st}, C_2^{st}$, which implies that $C_0 \geq_{sw} C_1, C_2$ by Lemma 2.13. This completes the proof.

Definition 2.15. We call C_A^{sw} the southwest channel of A.

Corollary 2.16. Under the above hypothesis, we have

$$\left(C_A^{\texttt{sw}}\right)^{\texttt{st}} = C_{A^{\texttt{st}}}^{\texttt{sw}}$$

Proof. It follows from Lemma 2.13(4) and (5).

Let $C = \{c_i\}_{i \in \mathbb{Z}}$ be a channel of A and let d_0 be a numbering on C given by $d_0(c_i) = i$ for $i \in \mathbb{Z}$. For $c \in \text{supp}(A)$, we define

$$(2.6) d_A^C(c) = \sup \left\{ \begin{array}{c} d_0(c'_k) + k \\ k \ge 0 \end{array} \middle| \begin{array}{c} c'_k >_{\text{NW}} \cdots >_{\text{NW}} c'_0 \text{ is a chain in } \operatorname{supp}(A) \\ (k \ge 0) \text{ such that } c'_0 = c \text{ and } c'_k \in C \end{array} \right\}.$$

Proposition 2.17 (cf. [3]). The numbering d_A^C on A is a well-defined proper numbering. Moreover, we have $d_A^C(c) = d_0(c)$ for $c \in C$.

Proof. Put $d = d_A^C$. We claim that

$$(2.7) d^{st} = d_{A^{st}}^{C^{st}}$$

where $d_{A^{\text{st}}}^{C^{\text{st}}}$ is defined in the same way as in (2.6) with respect to C^{st} .

Let $c \in \operatorname{supp}(A^{\operatorname{st}})$ be given. Let $\overline{c} \in \operatorname{supp}(A)$ such that $c \in \operatorname{supp}(A_{\overline{c}}^{\operatorname{st}})$. Suppose that we are given a chain $c_k >_{\operatorname{NW}} \cdots >_{\operatorname{NW}} c_0$ in $\operatorname{supp}(A)$ from $c_0 = \overline{c}$ to $c_k \in C$. Let c'_i be a cell in $\operatorname{supp}(A_{c_i}^{\operatorname{st}})$ $(1 \leq i \leq k)$ such that $c'_k \in C^{\operatorname{st}}$. Then we have a chain $c'_k >_{\operatorname{NW}} \cdots >_{\operatorname{NW}} c'_0$ in $\operatorname{supp}(A^{\operatorname{st}})$ from $c'_0 = c$ to $c_k \in C^{\operatorname{st}}$. Conversely, if we have a chain of length k + 1 in $\operatorname{supp}(A^{\operatorname{st}})$ from c to C^{st} , then we can associate a chain of length k + 1 in $\operatorname{supp}(A)$ from \overline{c} to C by taking representatives of the block submatrix to which each cell in the chain belongs to. This implies that $d^{\operatorname{st}}(c) = d_{A^{\operatorname{st}}}^{\operatorname{Cst}}(c)$, which proves the claim. Hence, d is also proper by Lemma 2.9(1) since $d_{A^{\operatorname{st}}}^{\operatorname{Cst}}$ is proper by [3, Proposition 3.10]. It follows from $d^{\operatorname{st}} = d_{A^{\operatorname{st}}}^{\operatorname{Cst}}$ and [3, Proposition 11.3] that $d(c) = d_0(c)$ $(c \in C)$.

Remark 2.18. Note that the width of A or the period of the channel numbering d_A^C is equal to the flow of a channel C of A.

Definition 2.19. When $C = C_A^{sw}$, we write $d_A^{sw} = d_A^{C_A^{sw}}$ for short, and call it the *southwest* channel numbering on A.

Example 2.20. Let A be the generalized affine permutation in Example 2.1. The southwest channel of A is given by

$$C_A^{\text{sw}} = \left\{ \cdots >_{\text{NW}} \tau^{-1}(4,7) >_{\text{NW}} (2,3) >_{\text{NW}} (3,6) >_{\text{NW}} (4,7) >_{\text{NW}} \tau(2,3) >_{\text{NW}} \cdots \right\},$$

which has flow 3. Then the southwest channel numbering d_A^{sw} is represented as follows:



Here each circle represents a cell in supp(A) and the blue circles forms the southwest channel C_A^{sw} . The number in each circle is the numbering d_A^{sw} of that cell, where \overline{k} means -k. For example, $d_A^{sw}(2,8) = d_A^{sw}(3,6) = d_A^{sw}(7,4) = 1$.

Remark 2.21. Consider the southwest channel $C_{A^t}^{sw}$ of A^t . Then

$$C_A^{\texttt{ne}} = \{\,(i,j) \,|\, (j,i) \in C_{A^t}^{\texttt{sw}}\,\}$$

is the minimal element in \mathcal{C}_A with respect to \geq_{sw} , which is called the *northeast channel of* A. Let d_A^{ne} be the channel numbering corresponding to C_A^{ne} . Then it follows from definition that $d_{A^t}^{sw}(j,i) = d_A^{ne}(i,j)$ for $(j,i) \in \text{supp}(A^t)$.

The following lemma gives a characterization of a channel numbering (cf. [3, Remark 11.8]).

Lemma 2.22. Let C be a channel of A. Let d be a proper numbering on A such that $d(c) = d_A^C(c)$ for $c \in C$. Then the following are equivalent:

- (1) $d = d_A^C$,
- (2) for $c \in \operatorname{supp}(A)$, there exists a chain $c_k >_{NW} \cdots >_{NW} c_0$ in $\operatorname{supp}(A)$ such that $c_0 = c$, $c_k \in C$ and $d(c_i) = d(c) - i$ for $0 \leq i \leq k$,
- (3) if d' is a proper numbering such that $d'(c) = d_A^C(c)$ for $c \in C$, then we have $d(c) \leq d'(c)$ for every $c \in \text{supp}(A)$.

Proof. Suppose that (1) holds. Let $c_k >_{NW} \cdots >_{NW} c_0 = c$ be a chain which gives the maximum value $d_0(c_k) + k$ in (2.6). Since d is monotone, we have

(2.8)
$$d(c_k) + k \leq d(c_{k-1}) + k - 1 \leq \dots \leq d(c) = d_0(c_k) + k.$$

Since $c_k \in C$, we have $d(c_k) = d_0(c_k)$ by Proposition 2.17. Thus all the inequalities in (2.8) are in fact equalities and hence, $d(c_i) = d(c) - i$ for $0 \le i \le k$. This implies (2).

Suppose that (2) holds. For $c \in \text{supp}(A)$, let $c_k >_{NW} \cdots >_{NW} c_0 = c$ be a chain satisfying the condition in (2). Let d' be a proper numbering such that $d' = d_A^C$ on C. Along this chain, we have

$$d'(c_k) + k \leq d'(c_{k-1}) + k - 1 \leq \dots \leq d'(c).$$

from the monotonicity of d'. Since $d'(c_k) = d(c_k)$ we conclude that $d(c) = d(c_k) + k = d'(c_k) + k \leq d'(c)$. This implies (3).

Suppose that (3) holds. Then, in particular, we have $d(c) \leq d_A^C(c)$ for $c \in \text{supp}(A)$ by letting $d' = d_A^C$. Let $c_k >_{NW} \cdots >_{NW} c_0 = c$ be a chain which gives the maximal value $d_A^C(c)$. We have $d(c_k) + k \leq d(c_{k-1}) + k - 1 \leq \cdots \leq d(c)$ from the monotonicity of d. Then we see that $d_A^C(c) = d_A^C(c_k) + k = d(c_k) + k \leq d(c)$. Hence $d(c) = d_A^C(c)$.

Remark 2.23. By Lemma 2.22, we may understand that the channel numbering d_A^C is the proper numbering with minimal values among the proper numberings d which coincide with d_0 on C.

Corollary 2.24. Let C_1 , C_2 be channels of A such that $C_1 \cap C_2 \neq \emptyset$. If $d_A^{C_1} = d_A^{C_2}$ on $C_1 \cap C_2$, then we have $d_A^{C_1} = d_A^{C_2}$.

Proof. Let $C_1 = \{\cdots >_{NW} c_0 >_{NW} \cdots >_{NW} c_l = \tau(c_0) >_{NW} \cdots \}$, where *l* is the common flow of C_1 and C_2 . We may assume that $c_0 \in C_1 \cap C_2$. Along C_1 , we have

$$d_A^{C_2}(c_0) + l \leq d_A^{C_2}(c_1) + (l-1) \leq \dots \leq d_A^{C_2}(c_l).$$

By Lemma 2.10, we see that $d_A^{C_2}(c_l) = d_A^{C_2}(c_0) + l$. Hence all the inequalities above are equalities, and for $0 \leq i \leq l$

$$d_A^{C_2}(c_i) = d_A^{C_2}(c_0) + i = d_A^{C_1}(c_0) + i = d_A^{C_1}(c_i).$$

By Lemma 2.22(3), we have $d_A^{C_1}(c) \leq d_A^{C_2}(c)$ for $c \in \text{supp}(A)$. The reverse inequality can be obtained in the same way. Hence we obtain $d_A^{C_1} = d_A^{C_2}$.

2.6. **Tableaux.** Let \mathscr{P} be the set of partitions or Young diagrams $\lambda = (\lambda_1, \lambda_2, \cdots)$. Let $\lambda' = (\lambda'_1, \lambda'_2, \cdots)$ be the conjugate of λ , $\ell(\lambda)$ the length of λ , and $|\lambda| = \sum_{i \ge 1} \lambda_i$. For $n \ge 1$, let $\mathscr{P}_n = \{\lambda \in \mathscr{P} | \ell(\lambda) \le n\}$.

Let \mathcal{A} be a linearly ordered set. For example, \mathbb{Z} and [k] $(k \ge 1)$ with the usual linear order. For $\lambda \in \mathscr{P}$, let $SST_{\mathcal{A}}(\lambda)$ be the set of semistandard (or \mathcal{A} -semistandard) tableaux of shape λ , that is, tableaux with entries in \mathcal{A} such that (1) the entries in each row are weakly increasing from left to right, (2) the entries in each column are strictly increasing from top to bottom.

We denote by $CSST_{\mathcal{A}}(\lambda)$ the set of column semistandard tableaux of shape λ , that is, tableaux of shape λ with entries in \mathcal{A} which satisfy the condition (2). We also let $CST_{\mathcal{A}}(\lambda)$ be the set of $T \in CSST_{\mathcal{A}}(\lambda)$ such that all the entries in T are distinct.

Similarly, we let $RSST_{\mathcal{A}}(\lambda)$ be the set of row semistandard tableaux of shape λ and let $RST_{\mathcal{A}}(\lambda)$ be the set of $T \in RSST_{\mathcal{A}}(\lambda)$ such that all the entries in T are distinct.

3. Affine RSK correspondence

3.1. Proper numberings and zig-zags.

Definition 3.1. Let $\mathbf{z} = \{c_i\}_{i \in \mathbb{Z}}$ an infinite collection of cells such that c_{i+1} is adjacent to c_i for $i \in \mathbb{Z}$.

- (1) We say that \mathbf{z} is a *zig-zag* when it satisfies the following:
 - (i) c_{i+1} is located to the east or north of c_i for all i,
 - (ii) c_{i+1} is located to the east of c_i for $i \gg 0$,
 - (iii) c_{i-1} is located to the south of c_i for $i \ll 0$.
- (2) A cell c_i in a zig-zag \mathbf{z} is called an *inner corner* if c_{i-1} is located to the south and c_{i+1} is located to the east of c_i , and called an *outer corner* if c_{i-1} is located to the west and c_{i+1} is located to the north of c_i . Note that a zig-zag \mathbf{z} has at least one inner corner.
- (3) A cell $c = (i_l, j_k)$ is called a *back-post corner* of a zig-zag **z** when $c_k = (i_k, j_k)$ and $c_l = (i_l, j_l)$ are the leftmost and the rightmost inner corners of **z**, respectively.

Let d be a proper numbering on a non-zero $A \in \widehat{\mathcal{M}}_{m \times n}$. For each $k \in \mathbb{Z}$, the level set $d^{-1}(k)$ forms a chain with respect to \leq_{ne} , by monotonicity of d (Definition 2.8).

We associate a set of zig-zags $Z_d = {\mathbf{z}_k}_{k \in \mathbb{Z}}$ to d, where the inner corners of \mathbf{z}_k are the set of maximal elements in $d^{-1}(k)$ with respect to \geq_{nw} . It is straightforward to see that ${\mathbf{z}_k}_{k \in \mathbb{Z}}$ satisfies

- (z.1) the inner corners of each \mathbf{z}_k are contained in supp(A),
- (z.2) \mathbf{z}_k 's are mutually disjoint and $\operatorname{supp}(A) \subseteq \bigsqcup_{k \in \mathbb{Z}} \mathbf{z}_k$,
- (z.3) \mathbf{z}_k is located to the southeast of \mathbf{z}_{k-1} for $k \in \mathbb{Z}$ in the sense that

(3.1) for each
$$c_1 \in \mathbf{z}_k$$
, there exists $c_2 \in \mathbf{z}_{k-1}$ such that $c_2 >_{NW} c_1$

Conversely, a set of zig-zags $Z = {\mathbf{z}_k}_{k \in \mathbb{Z}}$ satisfying (z.1)-(z.3) determines a unique proper numbering d^Z on A given by

(3.2)
$$d^{Z}(c) = k \quad \text{if } c \in \text{supp}(A) \cap \mathbf{z}_{k}$$

whose associated zig-zags is Z.

3.2. Matrix-ball construction. Now suppose that a non-zero $A \in \widehat{\mathcal{M}}_{m \times n}$ is given and let

• $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$: the set of zig-zags associated to d_A^{sw} .

Note that $\tau(\mathbf{z}_k) = \mathbf{z}_{k+\ell}$ $(k \in \mathbb{Z})$ if ℓ is the period of d_A^{sw} (cf. Lemma 2.10).

Example 3.2. Let A and d_A^{sw} be as in Example 2.20. The period of d_A^{sw} is 3. The zig-zags \mathbf{z}_k corresponding to the level sets $(d_A^{sw})^{-1}(k)$ for k = 1, 2, 3 are given as red lines below.



Let

- A^{\flat} : the matrix in $\widehat{\mathcal{M}}_{m \times n}$ obtained from A by
 - (i) subtracting one at the inner corners in \mathbf{z}_k $(k \in \mathbb{Z})$,
 - (ii) adding one at the outer corners in \mathbf{z}_k $(k \in \mathbb{Z})$,
- $A^{(t)}$ $(t \ge 0)$: the matrices in $\widehat{\mathcal{M}}_{m \times n}$ defined inductively as follows:

$$A^{(0)} = A, \quad A^{(t)} = \left(A^{(t-1)}\right)^{\flat}.$$

We have $A^{(t)} = \mathbb{O}$ for a sufficiently large t, since the number of outer corners in each zig-zag \mathbf{z}_k are strictly smaller than those of inner corners. Suppose that

$$A^{(s-1)} \neq \mathbb{O}, \quad A^{(s)} = \mathbb{O}$$

for some $s \ge 1$. For $1 \le t \le s$, we consider

- $\{\mathbf{z}_{k}^{(t)}\}_{k\in\mathbb{Z}}$: the set of zig-zags associated to $d_{A^{(t-1)}}^{sw}$,
- $\mathbf{s}^{(t)}$: the stream consisting of the back-post corners of $\mathbf{z}_k^{(t)}$ $(k \in \mathbb{Z})$,
- μ_t : the flow of $\mathbf{s}^{(t)}$ or the width of $A^{(t-1)}$,
- $(\mathbf{a}_t, \mathbf{b}_t, \rho_t)$: the defining data of $\mathbf{s}^{(t)}$ $(1 \le t \le s)$.

Lemma 3.3 (cf. [3]). The flows of $\mathbf{s}^{(1)}, \ldots, \mathbf{s}^{(s)}$ is a weakly decreasing sequence of positive integers, that is, $\mu_1 \ge \cdots \ge \mu_s > 0$.

Proof. Let ℓ and ℓ' be the widths of A and A^{\flat} , respectively. Consider the zig-zags \mathbf{z}_k $(k \in \mathbb{Z})$ associated to d_A^{sw} . By definition, $\mathrm{supp}(A^{\flat})$ is contained in the union of \mathbf{z}_k 's. Suppose that $\ell < \ell'$. Let $C = \{c_i\}_{i\in\mathbb{Z}}$ be a channel of A^{\flat} . Note that each cell c_i belong to a different zig-zag \mathbf{z}_k . But the width of A is ℓ , which implies that at least two cells among ℓ' cells $c_1 >_{\mathrm{NW}} c_2 >_{\mathrm{NW}} \cdots >_{\mathrm{NW}} c_{\ell'} = \tau(c_0)$, belong to the same zig-zag \mathbf{z}_k for some k. This is a contradiction.

Now we let

- $\lambda = \mu'$: the conjugate partition of $\mu = (\mu_1, \dots, \mu_s) \in \mathscr{P}_s$,
- P_0 : the tableau of shape λ , whose t-th column from the left is \mathbf{a}_t $(1 \leq t \leq s)$,
- Q_0 : the tableau of shape λ , whose t-th column from the left is \mathbf{b}_t $(1 \leq t \leq s)$,
- $\rho = (\rho_1, \ldots, \rho_s) \in \mathbb{Z}^s$.

Here we understand that the entries of the *t*-th column in P_0 and Q_0 are given by those in \mathbf{a}_t and \mathbf{b}_t respectively. Note that $\lambda \in \mathscr{P}_m \cap \mathscr{P}_n$, and P_0, Q_0 are column semistandard tableaux.

Summarizing, we have a map

(3.3)
$$\kappa_{0}: \widehat{\mathcal{M}}_{m \times n} \longrightarrow \bigsqcup_{\lambda \in \mathscr{P}_{m} \cap \mathscr{P}_{n}} CSST_{[m]}(\lambda) \times CSST_{[n]}(\lambda) \times \mathbb{Z}^{\lambda_{1}},$$
$$A \longmapsto (P_{0}, Q_{0}, \rho)$$

where we assume $\kappa_0(\mathbb{O}) = (\emptyset, \emptyset, (0, \dots, 0))$ with \emptyset the empty tableau.

Example 3.4. Let A be the generalized affine permutation in Example 2.1.



where the red lines denote the zig-zags in Example 3.2. The stream consisting of the backpost corners of \mathbf{z}_k ($k \in \mathbb{Z}$) is

$$\mathbf{s}^{(1)} = \{ \dots >_{\mathsf{NW}} (2,4) >_{\mathsf{NW}} (3,6) >_{\mathsf{NW}} (4,7) >_{\mathsf{NW}} \dots \}.$$

By subtracting one at each inner corner and adding one at each outer corner of \mathbf{z}_k , we obtain A^{\flat} as follows:



Repeating this process, we see that



with $A^{(5)} = \mathbb{O}$, and

$$\begin{split} \mathbf{s}^{(2)} &= \{ \cdots >_{\text{NW}} (1,3) >_{\text{NW}} (2,6) >_{\text{NW}} (4,7) >_{\text{NW}} \cdots \}, \\ \mathbf{s}^{(3)} &= \{ \cdots >_{\text{NW}} (2,0) >_{\text{NW}} (3,2) >_{\text{NW}} \cdots \}, \\ \mathbf{s}^{(4)} &= \{ \cdots >_{\text{NW}} (1,-1) >_{\text{NW}} (3,3) >_{\text{NW}} \cdots \}, \\ \mathbf{s}^{(5)} &= \{ \cdots >_{\text{NW}} (2,3) >_{\text{NW}} (3,4) >_{\text{NW}} \cdots \}. \end{split}$$

Hence $\kappa_0(A) = (P_0, Q_0, \rho)$, where

Remark 3.5. We have shown that $(d_A^{sw})^{st} = d_{A^{st}}^{sw}$ in (2.7). This implies that

$$(A^{\mathtt{st}})^{\flat} = (A^{\flat})^{\mathtt{st}}.$$

More precisely, it should be understood that the righthand side is obtained by removing some zero rows and columns on the lefthand side (cf. Section 2.2 for the convention of row and column indices of A^{st}).

3.3. Semistandard tableaux of rectangular shape. Let $\lambda \in \mathscr{P}$ be given. For $T \in SST_{\mathbb{Z}}(\lambda)$, let us write $T = (T^s, \ldots, T^2, T^1)$, where T^j denotes the *j*-th column of T from the right. For any $1 \leq j < k \leq s$, we may regard $(T^k, \ldots, T^{j+1}, T^j)$ as a semistandard subtableau of T with the columns $T^k, \ldots, T^{j+1}, T^j$.

For $a \ge 1$ and $1 \le b \le n$, let

$$R = (a^b) = (\underbrace{a, \dots, a}_{b}) \in \mathscr{P}_n$$

be a Young diagram of rectangular shape. Let

$$\mathcal{B}(R) = \left\{ T \in SST_{\mathbb{Z}}(R) \mid T^{j}(b) - T^{j}(1) < n \ (1 \leq j \leq a) \right\},\$$

where $T^{j}(i)$ denotes the entry of T^{j} at the *i*-th row from the top. Let u_{R} denote the unique element of $\mathcal{B}(R)$ such that the *i*-th row is filled with *i* for $1 \leq i \leq b$.

For $T \in \mathcal{B}((1^b))$, let $\tau_n(T) = \tau(T)$ be the column semistandard tableau obtained from T by replacing its entries $T(1) < T(2) < \cdots < T(b)$ with $T(2) < T(3) < \cdots < T(b) < T(1) + n$. Hence τ defines a bijection on $\mathcal{B}((1^b))$ and τ^{-1} denotes its inverse. In general, for $\alpha = (\alpha_1, \ldots, \alpha_a) \in \mathbb{Z}^a$ and $(T_1, \ldots, T_a) \in \mathcal{B}((1^b)) \times \cdots \times \mathcal{B}((1^b))$, we define

(3.4)
$$\tau^{\alpha}(T_1,\ldots,T_a) = \left(\tau^{\alpha_1}(T_1),\ldots,\tau^{\alpha_a}(T_a)\right),$$

which gives a bijection on $\mathcal{B}((1^b)) \times \cdots \times \mathcal{B}((1^b))$. Regarding $\mathcal{B}(R) \subset \mathcal{B}((1^b)) \times \cdots \times \mathcal{B}((1^b))$, we may define τ^{α} on $\mathcal{B}(R)$. Let

$$\mathcal{B}(R)_0 = \left\{ T \in \mathcal{B}(R) \mid (T^{j+1}, \tau^{-1}(T^j)) \text{ is not semistandard for } 1 \le j \le a-1 \right\}.$$

Lemma 3.6. We have a bijection

$$\begin{aligned} &\mathcal{B}(R)_0 \times \mathscr{P}_{a-1} \longrightarrow &\mathcal{B}(R) \\ &(T,\nu) \longmapsto & \tau^{\nu_{\rm rev}}(T) \end{aligned}$$

for $T = (T^a, \dots, T^1)$ and $\nu = (\nu_1, \dots, \nu_{a-1}) \in \mathscr{P}_{a-1}$, where $\nu_{rev} = (0, \nu_{a-1}, \dots, \nu_2, \nu_1)$.

Let us also regard $CSST_{[n]}(R) \subset \mathcal{B}((1^b)) \times \cdots \times \mathcal{B}((1^b))$, and define $\tau^{\alpha} : CSST_{[n]}(R) \longrightarrow \mathcal{B}((1^b))^a$ for $\alpha = (\alpha_1, \ldots, \alpha_a) \in \mathbb{Z}^a$ as in (3.4).

Definition 3.7. Let $T \in CSST_{[n]}(R)$ be given. For $1 \leq j \leq a - 1$, let r_j be the minimal non-negative integer such that

$$(T^{j+1}, \tau^{r_j}(T^j))$$

is Z-semistandard, and put $\eta_j = r_j + r_{j+1} + \cdots + r_{a-1}$. We call (r_1, \ldots, r_{a-1}) the offset vector and $\eta = (\eta_1, \ldots, \eta_{a-1})$ the symmetrized offset vector of T.

The symmetrized offset vector is the unique $\eta \in \mathscr{P}_{a-1}$ such that $\tau^{\eta_{\text{rev}}}(T) \in \mathcal{B}(R)_0$. The following is immediate from the definition of η .

Lemma 3.8. Let $T \in CSST_{[n]}(R)$ and $\alpha = (\alpha_1, \ldots, \alpha_a) \in \mathbb{Z}^a$ be given. Then $\tau^{\alpha_{rev}}(T) \in \mathcal{B}(R)$ with $\alpha_{rev} = (\alpha_a, \ldots, \alpha_1)$ if and only if

$$\alpha_a \leqslant \alpha_{a-1} - \eta_{a-1} \leqslant \cdots \leqslant \alpha_1 - \eta_1.$$

By Lemma 3.8, we have a bijection

$$(3.5) \qquad CSST_{[n]}(R) \times \mathcal{P}_a \longrightarrow \mathcal{B}(R)$$
$$(T,\nu) \longmapsto \tau^{\nu_{\text{rev}} + \eta_{\text{rev}}}(T)$$

where $\mathcal{P}_a = \{ \nu = (\nu_1, \dots, \nu_a) \in \mathbb{Z}^a \mid \nu_1 \ge \dots \ge \nu_a \}$ is the set of generalized partitions of length a and η is the symmetrized offset vector of T. Let

(3.6)
$$\tau : \mathcal{B}(R) \longrightarrow \mathcal{B}(R)$$
$$T = (T^a, \dots, T^1) \longmapsto (\tau(T^a), \dots, \tau(T^1))$$

be a bijection given by applying τ to each column of the tableaux in $\mathcal{B}(R)$, which induces a \mathbb{Z} -action on $\mathcal{B}(R)$ and $\mathcal{B}(R)_0$. Let $\mathcal{B}(R)_0/\mathbb{Z}$ denote the set of equivalence classes under this \mathbb{Z} -action. We may identify $\mathcal{B}(R)_0/\mathbb{Z}$ with the set of $T \in \mathcal{B}(R)_0$ such that the first column have entries in [n]. Hence, we have another bijection

$$(3.7) \qquad CSST_{[n]}(R) \longrightarrow \mathcal{B}(R)_0/\mathbb{Z} ,$$
$$T \longmapsto [\tau^{\eta_{rev}}(T)]$$

where [T] denotes the equivalence class of T and η is the symmetrized offset vector of T...

3.4. Rectangular decomposition. Let $\lambda \in \mathscr{P}_n$ be given. We regard λ (its Young diagram) as the one obtained by gluing the Young diagrams of rectangular shapes:

(3.8)
$$R_i = (\underbrace{m_i, \dots, m_i}_{i}) \quad (1 \le i \le l),$$

where m_i is the number of occurrences of i in λ' and $l = \ell(\lambda)$. Here we assume that R_i is empty when $m_i = 0$. Let

(3.9)
$$\begin{aligned} & \mathcal{B}_n(\lambda)_0 = \mathcal{B}(\lambda)_0 = \mathcal{B}(R_l)_0 \times \dots \times \mathcal{B}(R_1)_0, \\ & \mathcal{B}_n(\lambda) = \mathcal{B}(\lambda) = \mathcal{B}(R_l) \times \dots \times \mathcal{B}(R_1). \end{aligned}$$

If we put $\mathscr{P}(\lambda) = \mathscr{P}_{m_l-1} \times \cdots \times \mathscr{P}_{m_2-1} \times \mathscr{P}_{m_1-1}$, where we take the product over $m_i \ge 1$, then we have a bijection

$$\begin{array}{ccc} \mathcal{B}(\lambda)_0 \times \mathscr{P}(\lambda) & \longrightarrow & \mathcal{B}(\lambda) \\ \\ \left(\left(T^{(i)} \right)_{1 \leqslant i \leqslant l}, (\nu^{(i)})_{1 \leqslant i \leqslant l} \right) & \longmapsto & \left(\tau^{\nu^{(i)}_{rev}}(T^{(i)}) \right)_{1 \leqslant i \leqslant l} \end{array}$$

by applying Lemma 3.6 to each component, where $T^{(i)} \in \mathcal{B}(R_i)_0$ and $\nu^{(i)} \in \mathscr{P}_{m_i-1}$.

Similarly, if we let $\mathcal{P}(\lambda) = \mathcal{P}_{m_l} \times \cdots \times \mathcal{P}_{m_1}$, where we take the product over $m_i \ge 1$, and regard

$$CSST_{[n]}(\lambda) = CSST_{[n]}(R_l) \times \cdots \times CSST_{[n]}(R_1),$$

then by (3.6) we have a bijection

$$(3.10) \qquad CSST_{[n]}(\lambda) \times \mathcal{P}(\lambda) \longrightarrow \mathcal{B}(\lambda)$$
$$\left(\left(T^{(i)} \right)_{1 \leq i \leq l}, (\nu^{(i)})_{1 \leq i \leq l} \right) \longmapsto \left(\tau^{\nu_{\mathsf{rev}}^{(i)} + \eta_{\mathsf{rev}}^{(i)}}(T^{(i)}) \right)_{1 \leq i \leq l}$$

where $\eta^{(i)} \in \mathscr{P}_{m_i-1}$ is the symmetrized offset vector of $T^{(i)}$.

Let

$$\mathcal{B}(\lambda)_0/\mathbb{Z}^l = \mathcal{B}(R_l)_0/\mathbb{Z} \times \cdots \times \mathcal{B}(R_1)_0/\mathbb{Z},$$

where each $\mathcal{B}(R_i)_0/\mathbb{Z}$ is the set of equivalence classes under the \mathbb{Z} -action (3.6). Then we also have a bijection

$$CSST_{[n]}(\lambda) \longrightarrow \mathcal{B}(\lambda)_0/\mathbb{Z}^l$$
$$(T^{(i)})_{1 \leq i \leq l} \longmapsto ([\tau^{\eta^{(i)}_{rev}}(T^{(i)})])_{1 \leq i \leq l}$$

where $\eta^{(i)} \in \mathscr{P}_{m_i-1}$ is the symmetrized offset vector of $T^{(i)}$.

3.5. Affine RSK correspondence. Suppose that $A \in \widehat{\mathcal{M}}_{m \times n}$ is given. We keep the notations in Sections 3.2 and 3.4.

Let (P_0, Q_0, ρ) be given as in (3.3). For $1 \leq i \leq l$ with $m_i \geq 1$, let

- $P_0^{(i)}, Q_0^{(i)}$: the subtableaux of P_0 and Q_0 corresponding to R_i , respectively,
- $\rho^{(i)} \in \mathbb{Z}^{m_i}$: the subsequence of ρ corresponding to the columns of R_i ,
- $\eta^{(i)} \in \mathscr{P}_{m_i-1}$: the symmetrized offset vector of $P_0^{(i)}$.

Then we define

(3.11)
$$Q = \left(\tau^{\rho^{(l)} + \eta^{(l)}_{\text{rev}}}(Q_0^{(l)}), \dots, \tau^{\rho^{(1)} + \eta^{(1)}_{\text{rev}}}(Q_0^{(1)})\right) = \left(Q^{(l)}, \dots, Q^{(1)}\right).$$

Note that the action of τ on Q_0 should be understood as τ_n . The following is one of the main results in this paper. The proof will be given in Section 5.

,

,

,

Theorem 3.9. We have a bijection

Remark 3.10. Applying the bijection (3.10) to $\mathcal{B}_n(\lambda)$, we have a bijection

$$\kappa^{\diamond}: \quad \widehat{\mathcal{M}}_{m \times n} \quad \longrightarrow \quad \bigsqcup_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} CSST_{[m]}(\lambda) \times CSST_{[n]}(\lambda) \times \mathcal{P}(\lambda)$$

Again applying the inverse of (3.10) to $CSST_{[m]}(\lambda) \times \mathcal{P}(\lambda)$, we have a bijection

$$\kappa': \ \widehat{\mathcal{M}}_{m \times n} \longrightarrow \bigsqcup_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} \mathcal{B}_m(\lambda) \times CSST_{[n]}(\lambda) \ .$$

Example 3.11. Let $\kappa_0(A) = (P_0, Q_0, \rho)$ be as in Example 3.4. The rectangular decompositions of P_0 , Q_0 and ρ are

$$P_0^{(2)} = \boxed{2 \ 1 \ 2}_{3 \ 3 \ 3}, \qquad Q_0^{(2)} = \boxed{2 \ 3 \ 3}_{5 \ 4 \ 4}, \qquad \rho^{(2)} = (-1, -1, 0)$$

$$P_0^{(3)} = \boxed{2 \ 1}_{3 \ 2 \ 4 \ 4}, \qquad Q_0^{(3)} = \boxed{1 \ 1}_{2 \ 2 \ 4 \ 3}, \qquad \rho^{(3)} = (2, 2),$$

where R_2 and R_3 are the only non-trivial rectangles in this decomposition. The symmetrized offset vectors of $P_0^{(2)}$ and $P_0^{(3)}$ are

$$\eta^{(2)}_{\rm rev} = (0, 1, 1), \quad \eta^{(3)}_{\rm rev} = (0, 1),$$

and hence

$$Q = \frac{4 \ 6 \ 0 \ 3 \ 4}{6 \ 7 \ 2 \ 4 \ 8}.$$

4. Affine crystals

4.1. Crystals. Let us give a brief review on crystals (see [5] for more details). Let \mathfrak{g} be the Kac-Moody algebra associated to a symmetrizable generalized Cartan matrix $A = (a_{ij})_{i,j\in I}$ indexed by a set I. Let P^{\vee} be the dual weight lattice, $P = \operatorname{Hom}_{\mathbb{Z}}(P^{\vee}, \mathbb{Z})$ the weight lattice, $\Pi^{\vee} = \{h_i \mid i \in I\} \subset P^{\vee}$ the set of simple coroots, and $\Pi = \{\alpha_i \mid i \in I\} \subset P$ the set of simple roots of \mathfrak{g} such that $\langle \alpha_j, h_i \rangle = a_{ij}$ for $i, j \in I$.

A \mathfrak{g} -crystal (or crystal if there is no confusion on \mathfrak{g}) is a set B together with the maps wt : $B \longrightarrow P$, $\varepsilon_i, \varphi_i : B \longrightarrow \mathbb{Z} \cup \{-\infty\}$ and $\tilde{e}_i, \tilde{f}_i : B \longrightarrow B \cup \{\mathbf{0}\}$ for $i \in I$ such that for $b \in B$ and $i \in I$

(1)
$$\varphi_i(b) = \langle \operatorname{wt}(b), h_i \rangle + \varepsilon_i(b),$$

- (2) $\varepsilon_i(\widetilde{e}_i b) = \varepsilon_i(b) 1, \ \varphi_i(\widetilde{e}_i b) = \varphi_i(b) + 1, \ \operatorname{wt}(\widetilde{e}_i b) = \operatorname{wt}(b) + \alpha_i \ \operatorname{if} \ \widetilde{e}_i b \neq \mathbf{0},$
- (3) $\varepsilon_i(\widetilde{f}_i b) = \varepsilon_i(b) + 1, \ \varphi_i(\widetilde{f}_i b) = \varphi_i(b) 1, \ \mathrm{wt}(\widetilde{f}_i b) = \mathrm{wt}(b) \alpha_i \ \mathrm{if} \ \widetilde{f}_i b \neq \mathbf{0},$
- (4) $\widetilde{f}_i b = b'$ if and only if $b = \widetilde{e}_i b'$ for $b, b' \in B$,
- (5) $\widetilde{e}_i b = \widetilde{f}_i b = \mathbf{0}$ if $\varphi_i(b) = -\infty$,

where **0** is a formal symbol. Here we assume that $-\infty + n = -\infty$ for all $n \in \mathbb{Z}$. A crystal B can be viewed as an I-colored graph where $b \xrightarrow{i} b'$ if and only if $\tilde{f}_i b = b'$ for $b, b' \in B$ and $i \in I$. A crystal B is called *normal* if $\varepsilon_i(b) = \max\{k \mid \tilde{e}_i^k b \neq \mathbf{0}\}$ and $\varphi_i(b) = \max\{k \mid \tilde{f}_i^k b \neq \mathbf{0}\}$ for $b \in B$ and $i \in I$.

Let B_1 and B_2 be crystals. A morphism $\psi : B_1 \longrightarrow B_2$ is a map from $B_1 \cup \{0\}$ to $B_2 \cup \{0\}$ such that

- (1) $\psi(0) = 0$,
- (2) wt($\psi(b)$) = wt(b), $\varepsilon_i(\psi(b)) = \varepsilon_i(b)$, and $\varphi_i(\psi(b)) = \varphi_i(b)$ if $\psi(b) \neq \mathbf{0}$,
- (3) $\psi(\tilde{e}_i b) = \tilde{e}_i \psi(b)$ if $\psi(b) \neq \mathbf{0}$ and $\psi(\tilde{e}_i b) \neq \mathbf{0}$,
- (4) $\psi(\widetilde{f}_i b) = \widetilde{f}_i \psi(b)$ if $\psi(b) \neq \mathbf{0}$ and $\psi(\widetilde{f}_i b) \neq \mathbf{0}$,

for $b \in B_1$ and $i \in I$. We call ψ an *embedding* when ψ is injective, and call ψ strict if $\psi : B_1 \cup \{\mathbf{0}\} \longrightarrow B_2 \cup \{\mathbf{0}\}$ commutes with \tilde{e}_i and \tilde{f}_i for all $i \in I$, where we assume that $\tilde{e}_i \mathbf{0} = \tilde{f}_i \mathbf{0} = \mathbf{0}$. When ψ is a bijection, it is called an isomorphism.

Given $b_1 \in B_1$ and $b_2 \in B_2$, we say that b_1 is \mathfrak{g} -crystal equivalent to b_2 if there is an isomorphism of crystals $C(b_1) \longrightarrow C(b_2)$ mapping b_1 to b_2 , where $C(b_i)$ denotes the connected component of b_i in B_i for i = 1, 2.

The tensor product $B_1 \otimes B_2$ is defined to be $B_1 \times B_2$ as a set with elements denoted by $b_1 \otimes b_2$, where

$$(4.1)$$

$$wt(b_1 \otimes b_2) = wt(b_1) + wt(b_2),$$

$$\varepsilon_i(b_1 \otimes b_2) = \max\{\varepsilon_i(b_1), \varepsilon_i(b_2) - \langle wt(b_1), h_i \rangle\},$$

$$\varphi_i(b_1 \otimes b_2) = \max\{\varphi_i(b_1) + \langle wt(b_2), h_i \rangle, \varphi_i(b_2)\},$$

$$\tilde{e}_i(b_1 \otimes b_2) = \begin{cases} \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \ge \varepsilon_i(b_2), \\ b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2), \end{cases}$$

$$\tilde{f}_i(b_1 \otimes b_2) = \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \ge \varepsilon_i(b_2), \\ b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \end{cases}$$

for $i \in I$. Here, we assume that $\mathbf{0} \otimes b_2 = b_1 \otimes \mathbf{0} = \mathbf{0}$. Then $B_1 \otimes B_2$ is a crystal.

Suppose that $\sigma = (\ldots, \sigma_{-2}, \sigma_{-1}, \sigma_0, \sigma_1, \sigma_2, \ldots)$ is a sequence (not necessarily finite) where $\sigma_k \in \{+, -, \cdot\}$ and $\sigma_k = \cdot$ except for finitely many k. In the sequence σ , we replace a pair $(\sigma_s, \sigma_{s'}) = (+, -)$, where s < s' and $\sigma_t = \cdot$ for s < t < s', with (\cdot, \cdot) , and repeat this process as far as possible until we get a sequence with no – placed to the right of +. We denote the resulting reduced sequence by $\tilde{\sigma}$, where no + is placed to the left of any – in the sequence. It is often used to describe a crystal structure on the tensor product of crystals.

4.2. $A_{n-1}^{(1)}$ -crystals. Suppose that \mathfrak{g} is of type $A_{n-1}^{(1)}$ for $n \ge 2$. We assume that $I = \{0, 1, \ldots, n-1\}, P^{\vee} = \bigoplus_{i \in I} \mathbb{Z}h_i \oplus \mathbb{Z}d$, and $P = \bigoplus_{i \in I} \mathbb{Z}\Lambda_i \oplus \mathbb{Z}\delta$, where $\delta = \sum_{i \in I} \alpha_i$ is the null root such that $\langle \delta, h_j \rangle = 0$ and $\langle \delta, d \rangle = 1$, and Λ_i is the *i*-th fundamental weight such

that $\langle \Lambda_i, h_j \rangle = \delta_{ij}$ and $\langle \Lambda_i, d \rangle = 0$. We let P_+ be the set of dominant integral weights and let

$$P^{0} = \mathbb{Z}\varpi_{1} \oplus \cdots \oplus \mathbb{Z}\varpi_{n-1} \oplus \mathbb{Z}\delta, \quad P_{cl} = P/\mathbb{Z}\delta, \quad P_{cl}^{0} = P^{0}/\mathbb{Z}\delta,$$

where $\varpi_i = \Lambda_i - \Lambda_0$ is the *i*-th level zero fundamental weight for $i \in \{1, \ldots, n-1\}$. Let cl : $P \longrightarrow P_{cl}$ be the canonical projection. Define $\epsilon_1 = \varpi_1$ and $\epsilon_i = \varpi_i - \varpi_{i-1}$ for $i \in \{2, \ldots, n-1\}$. Put $\epsilon_n = -\epsilon_1 - \cdots - \epsilon_{m-1}$. We have $\alpha_i = \epsilon_i - \epsilon_{i+1}$ for $i \in \{1, \ldots, n-1\}$ and $\alpha_0 = \epsilon_n - \epsilon_1 + \delta$.

Let X be one of P^0 and P^0_{cl} . We define an $A^{(1)}_{n-1}$ -crystal B with X-weights in the same manner, where wt : $B \longrightarrow X$. We often denote wt by wt⁰ and wt⁰_{cl} when X is P^0 and P^0_{cl} , respectively.

For $\Lambda \in P$, let $B(\Lambda)$ denote the crystal of the *extremal weight module* generated by an extremal vector of weight Λ [9].

Example 4.1. Let \mathcal{F} be the set of sequences $\mathbf{v} = (v_k)_{k \in \mathbb{Z}}$ such that $v_k \in \{0, 1\}, v_k = 1$ for $k \ll 0$ and $v_k = 0$ for $k \gg 0$. For $i \in I$, consider a sequence

$$\sigma_{\mathbf{v},i} = (\ldots, \sigma_{i-n}, \sigma_{i+1-n}, \sigma_i, \sigma_{i+1}, \sigma_{i+n}, \sigma_{i+1+n}, \ldots),$$

where we assign for $k \equiv i \pmod{n}$

$$(\sigma_k, \sigma_{k+1}) = \begin{cases} (+, \cdot) & \text{if } (v_k, v_{k+1}) = (1, 0), \\ (\cdot, -) & \text{if } (v_k, v_{k+1}) = (0, 1), \\ (\cdot, \cdot) & \text{if } (v_k, v_{k+1}) = (0, 0) \text{ or } (1, 1) \end{cases}$$

We define $\tilde{e}_i \mathbf{v}$ to be the sequence obtained from \mathbf{v} by replacing $(v_k, v_{k+1}) = (0, 1)$ with (1,0), where $k + 1 \equiv i + 1 \pmod{n}$ is the index corresponding to the rightmost – in the reduced sequence $\widetilde{\sigma_{\mathbf{v},i}}$. We define $\tilde{e}_i \mathbf{v} = \mathbf{0}$ if there is no such k. Similarly, we define $\tilde{f}_i \mathbf{v}$ to be the sequence obtained from \mathbf{v} by replacing $(v_k, v_{k+1}) = (1,0)$ with (0,1), where $k \equiv i \pmod{n}$ is the index corresponding to the leftmost + in $\widetilde{\sigma_{\mathbf{v},i}}$. We define $\tilde{f}_i \mathbf{v} = \mathbf{0}$ if there is no such k.

For $j \in I$, let $\mathbf{v}_{\Lambda_j} = (a_k)_{k \in \mathbb{Z}}$ be such that $v_k = 1$ if and only if $k \leq j$. It is well-known that the connected component of \mathbf{v}_{Λ_j} in \mathcal{F} with respect to \tilde{e}_i and \tilde{f}_i for $i \in I$ is isomorphic to $B(\Lambda_j)$, where we put $\operatorname{wt}(\mathbf{v}_{\Lambda_j}) = \Lambda_j$ [16]. Moreover, we have $\mathcal{F} \cong \bigoplus_{i \in \mathbb{Z}} B(\Lambda_j)$.

The dual crystal can be described similarly. Let \mathcal{F}^{\vee} be the set of sequences $\mathbf{v} = (v_k)_{k \in \mathbb{Z}}$ such that $v_k \in \{0, 1\}$, $v_k = 0$ for $k \ll 0$ and $v_k = 1$ for $k \gg 0$. Let $\mathbf{v}_{-\Lambda_j} = (v_k)_{k \in \mathbb{Z}}$ be such that $a_k = 0$ if and only if $k \leq j$. Then the connected component of $\mathbf{v}_{-\Lambda_j}$ in \mathcal{F}^{\vee} with respect to \tilde{e}_i and \tilde{f}_i for $i \in I$ is isomorphic to $B(-\Lambda_j)$, where $\operatorname{wt}(\mathbf{v}_{-\Lambda_j}) = -\Lambda_j$.

4.3. Bicrystal structure on $\widehat{\mathcal{M}}_{m \times n}$. In this subsection, we define two crystal structures on $\widehat{\mathcal{M}}_{m \times n}$ of type $A_{m-1}^{(1)}$ and $A_{n-1}^{(1)}$.

For $i, j \in \mathbb{Z}$, let E_{ij} denote the elementary matrix with 1 at the (i, j)-position and 0 elsewhere and put

$$\widehat{E}_{ij} = \sum_{k \in \mathbb{Z}} E_{i+km\,j+kn} \in \widehat{\mathcal{M}}_{m \times n}.$$

Let us first describe an $A_{m-1}^{(1)}$ -crystal structure on $\widehat{\mathcal{M}}_{m \times n}$ for $m \geq 2$. Suppose that $A = (a_{ij})_{i,j \in \mathbb{Z}} \in \widehat{\mathcal{M}}_{m \times n}$ is given. For $i \in \{0, 1, \ldots, m-1\}$, we define $\widetilde{e}_i A$ and $\widetilde{f}_i A$ as follows:

(1) Let σ be a sequence of $\{+, -\}$ given by

$$\sigma = (\cdots, \underbrace{-\cdots}_{a_{i+1\,j}}, \underbrace{+\cdots+}_{a_{i\,j}}, \underbrace{-\cdots}_{a_{i+1\,j+1}}, \underbrace{+\cdots+}_{a_{i\,j+1}}, \cdots)$$

and let $\tilde{\sigma}$ be the reduced one, which is well-defined since σ has only finitely many +'s and -'s.

(2) If $\tilde{\sigma}$ has at least one –, then we define

$$\widetilde{e}_i A = A + \widehat{E}_{ij_0} - \widehat{E}_{i+1j_0},$$

where j_0 is the column index of A corresponding the rightmost - in $\tilde{\sigma}$. If $\tilde{\sigma}$ has no -, then we define $\tilde{e}_i A = \mathbf{0}$. Similarly, if $\tilde{\sigma}$ has at least one +, then we define

$$\widetilde{f}_i A = A - \widehat{E}_{ij_1} + \widehat{E}_{i+1j_1},$$

where j_1 is the column index of A corresponding the leftmost + in $\tilde{\sigma}$. If $\tilde{\sigma}$ has no +, then we define $\tilde{f}_i A = \mathbf{0}$.

Put

(4.2)
$$\operatorname{wt}^{0}(A) = \sum_{k \in \mathbb{Z}} \left(\sum_{j=1}^{n} a_{1+km \, j} \right) (\epsilon_{1} - k\delta) + \dots + \sum_{k \in \mathbb{Z}} \left(\sum_{j=1}^{n} a_{m+km \, j} \right) (\epsilon_{m} - k\delta) \in P^{0},$$

(4.3)
$$\operatorname{wt}^{0}_{\operatorname{cl}}(A) = \operatorname{cl}(\operatorname{wt}^{0}(A)) = \left(\sum_{j \in \mathbb{Z}} a_{1j}\right) \operatorname{cl}(\epsilon_{1}) + \dots + \left(\sum_{j \in \mathbb{Z}} a_{mj}\right) \operatorname{cl}(\epsilon_{m}) \in P^{0}_{\operatorname{cl}}.$$

Example 4.2. Let A be the generalized affine permutation in Example 2.1. For i = 2, the associated sequence σ and its reduced one $\tilde{\sigma}$ are

$$\sigma = (+, -, -, +, -, +, +, -), \\ \widetilde{\sigma} = (\cdot, \cdot, -, \cdot, \cdot, +, \cdot, \cdot).$$

Hence the cell (2,8) is the position corresponding to the leftmost + in $\tilde{\sigma}$, and $\tilde{f}_i A = A - \hat{E}_{28} + \hat{E}_{38}$.



Here we present the submatrices of A and $\tilde{f}_2 A$ with the row indices in [4].

Lemma 4.3. The set $\widehat{\mathcal{M}}_{m \times n}$ is a normal $A_{m-1}^{(1)}$ -crystal with P^0 and P_{cl}^0 -weights with respect to wt, \widetilde{e}_i , \widetilde{f}_i for $i \in \{0, 1, \ldots, m-1\}$, where wt = wt⁰ and wt⁰_{cl}, respectively.

Next, we define an $A_{n-1}^{(1)}$ -crystal structure on $\widehat{\mathcal{M}}_{m \times n}$ for $n \ge 2$, say wt^t, $\varepsilon_i^t, \varphi_i^t, \widetilde{e}_i^t, \widetilde{f}_i^t$ for $i \in \{0, 1, \dots, n-1\}$, by applying the $A_{n-1}^{(1)}$ -crystal structure on $\widehat{\mathcal{M}}_{n \times m}$ to the transpose A^t of $A \in \widehat{\mathcal{M}}_{m \times n}$

$$\begin{aligned} &\operatorname{wt}^{t}(A) = \operatorname{wt}(A^{t}),\\ &\widetilde{e}_{i}^{t}A = (\widetilde{e}_{i}A^{t})^{t}, \quad \widetilde{f}_{i}^{t}A = (\widetilde{f}_{i}A^{t})^{t},\\ &\varepsilon_{i}^{t}(A) = \varepsilon_{i}(A^{t}), \quad \varphi_{i}^{t}(A) = \varphi_{i}(A^{t}). \end{aligned}$$

Proposition 4.4. The operators \tilde{e}_i and \tilde{f}_i for $i \in \{0, 1, \ldots, m-1\}$ commute with \tilde{e}_j^t and \tilde{f}_i^t for $j \in \{0, 1, \ldots, n-1\}$ on $\hat{\mathcal{M}}_{m \times n} \cup \{\mathbf{0}\}$.

Proof. It is straightforward to check that $\tilde{x}_i \tilde{y}_j^t = \tilde{y}_j^t \tilde{x}_i$ for $i \in \{0, 1, \dots, m-1\}, j \in \{0, 1, \dots, n-1\}$ and $x, y \in \{e, f\}$. We leave the details to the reader (see for example [13]).

Remark 4.5. If we assume that

$$\operatorname{wt}(A) = \operatorname{wt}_{\operatorname{cl}}^{0}(A) \quad \operatorname{wt}^{t}(A) = \operatorname{wt}_{\operatorname{cl}}^{0}(A^{t}),$$

for $A \in \widehat{\mathcal{M}}_{m \times n}$, then $\widehat{\mathcal{M}}_{m \times n}$ is an $(A_{m-1}^{(1)}, A_{n-1}^{(1)})$ -bicrystal. In other words, \widetilde{x}_i and \widetilde{y}_j^t on $\widehat{\mathcal{M}}_{m \times n} \cup \{\mathbf{0}\}$ are strict morphisms of $A_{n-1}^{(1)}$ -crystals and $A_{m-1}^{(1)}$ -crystals, respectively for $x, y \in \{e, f\}, i \in \{0, 1, \dots, m-1\}$, and $j \in \{0, 1, \dots, n-1\}$. On the other hand, if we assume that wt = wt⁰ on $\widehat{\mathcal{M}}_{m \times n}$ as an $A_{m-1}^{(1)}$ -crystal, then $\widetilde{e}_j^t, \widetilde{f}_j^t$ preserves the weights only when $j \neq 0$. In case of $\widetilde{e}_0^t, \widetilde{f}_0^t$, we have

$$\operatorname{wt}(\widetilde{e}_0^t A) = \operatorname{wt}(A) - \delta, \quad \operatorname{wt}(\widetilde{f}_0^t A) = \operatorname{wt}(A) + \delta,$$

for $A \in \widehat{\mathcal{M}}_{m \times n}$. The same holds for \widetilde{e}_i and \widetilde{f}_i .

4.4. Crystals of level zero extremal weight modules. Let $n \ge 2$. In this subsection, we describe a structure of $A_{n-1}^{(1)}$ -crystal with P^0 -weights on $\mathcal{B}(\lambda) = \mathcal{B}_n(\lambda)$ for $\lambda \in \mathscr{P}_n$.

First, consider $\mathcal{B}((1)) = SST_{\mathbb{Z}}((1))$, which can be identified with \mathbb{Z} . For $k \in \mathcal{B}((1))$ and $i \in I$, we define

$$\operatorname{wt}^{0}(\overline{k}) = \epsilon_{r} - q\delta, \quad \text{where } k = nq + r \text{ for } q \in \mathbb{Z} \text{ and } 1 \leqslant r \leqslant n,$$
$$\widetilde{e}_{i}[k] = \begin{cases} \boxed{k-1} & \text{if } k \equiv i+1 \pmod{n}, \\ \mathbf{0} & \text{otherwise,} \end{cases} \qquad \widetilde{f}_{i}[k] = \begin{cases} \boxed{k+1} & \text{if } k \equiv i \pmod{n}, \\ \mathbf{0} & \text{otherwise.} \end{cases}$$

Then $\mathcal{B}((1))$ is a normal $A_{n-1}^{(1)}$ -crystal with respect to wt⁰, \tilde{e}_i, \tilde{f}_i for $i \in I$.

Next, consider $\mathcal{B}((1^b))$ for $2 \leq b \leq n$. Let $T \in \mathcal{B}((1^b))$ be given with entries $T(1) < T(2) < \cdots < T(b)$. For $i \in I$, we define $\tilde{e}_i T$ as follows:

- (1) If $T(k) \equiv i+1 \pmod{n}$ and T(k-1) < T(k) 1 for some $2 \leq k \leq b$, then we define $\tilde{e}_i T$ to be the tableau obtained by replacing T(k) with T(k) 1.
- (2) If $T(1) \equiv i + 1 \pmod{n}$ and T(b) < T(1) 1 + n, then we define $\tilde{e}_i T$ to be the tableau obtained by replacing T(1) with T(1) 1.
- (3) Otherwise, we define $\tilde{e}_i T = \mathbf{0}$.

We define $\tilde{f}_i T$ in a similar way. We put $\mathrm{wt}^0(T) = \sum_{k=1}^b \mathrm{wt}^0(T(k))$, where we regard T(k) as an element in $\mathcal{B}((1))$.

Proposition 4.6. For $1 \leq b \leq n$, $\mathcal{B}((1^b))$ is a connected normal $A_{n-1}^{(1)}$ -crystal with P^0 -weights. For $1 \leq b \leq n-1$, $\mathcal{B}((1^b))$ is isomorphic to $B(\varpi_b)$.

Proof. It is easy to see that $\mathcal{B}((1^b))$ is a connected normal $A_{n-1}^{(1)}$ -crystal. In particular, $\mathcal{B}((1^n))$ is a crystal of single element of weight 0.

Suppose that $1 \leq b \leq n-1$. Let us keep the notations in Example 4.1. For $i \in \mathbb{Z}$, let $\mathbf{e}_i = (\delta_{ik})_{k \in \mathbb{Z}}$. Let $T \in \mathcal{B}((1^b))$ be given with entries $T(1) < \cdots < T(b)$. Suppose that

$$T(1) < \dots < T(k) \le 0 < T(k+1) < \dots < T(b)$$

for some $1 \le k \le b-1$. We assume that k = 0 if 0 < T(1) and k = b if $T(b) \le 0$. Let

$$\mathbf{v}_1 = \mathbf{v}_{\Lambda_k} + \mathbf{e}_{T(k+1)} + \dots + \mathbf{e}_{T(b)},$$

$$\mathbf{v}_2 = \mathbf{v}_{-\Lambda_k} + \mathbf{e}_{T(1)} + \dots + \mathbf{e}_{T(k)}.$$

Then the map sending T to $\mathbf{v}_1 \otimes \mathbf{v}_2$ gives a strict embedding

$$\mathcal{B}((1^b)) \longrightarrow B(\Lambda_b) \otimes B(-\Lambda_0) ,$$

where $u_{(1^b)}$ maps to $\mathbf{v}_{\Lambda_b} \otimes \mathbf{v}_{-\Lambda_0}$. This implies that $\mathcal{B}((1^b))$ is isomorphic to the connected component of $\mathbf{v}_{\Lambda_b} \otimes \mathbf{v}_{-\Lambda_0}$ in $B(\Lambda_b) \otimes B(-\Lambda_0)$. Hence $\mathcal{B}((1^b))$ is isomorphic to $B(\varpi_b)$ by [9] and [10, Proposition 5.4].

Corollary 4.7. For $T \in \mathcal{B}((1^b))$, we have $\tau(\tilde{e}_i T) = \tilde{e}_i \tau(T)$ and $\tau(\tilde{f}_i T) = \tilde{f}_i \tau(T)$ for $i \in I$, where $\mathrm{wt}^0(\tau(T)) = \mathrm{wt}^0(T) - \delta$.

Proof. It follows directly from the crystal structure on $\mathcal{B}((1^b))$.

Suppose that $R = (a^b) \in \mathscr{P}_n$ is given for $a \ge 1$ and $1 \le b \le n$. Define a map

$$\psi: \ \mathcal{B}(R) \longrightarrow \mathcal{B}((1^b))^{\otimes a}$$
$$T = (T^a, \dots, T^1) \longmapsto T^1 \otimes \dots \otimes T^a$$

Lemma 4.8. Under the above hypothesis, we have the following.

- (1) The image of $\mathfrak{B}(R)$ under ψ is invariant under \tilde{e}_i and \tilde{f}_i for $i \in I$. Hence $\mathfrak{B}(R)$ is an $A_{n-1}^{(1)}$ -crystal with P^0 -weights such that ψ is a strict embedding.
- (2) The connected component of u_R in $\mathcal{B}(R)$ is $\mathcal{B}(R)_0$ when b < n.

(3) The map

$$\mathcal{B}(R)_0 \times \mathscr{P}_{a-1} \longrightarrow \mathcal{B}(R)$$

in Lemma 3.6 is an isomorphism of $A_{n-1}^{(1)}$ -crystals with P^0 -weights. Here we understand $\mathcal{B}(R)_0 \times \mathscr{P}_{a-1}$ as $\mathcal{B}(R)_0 \otimes \mathscr{P}_{a-1}$ and \mathscr{P}_{a-1} as an $A_{n-1}^{(1)}$ -crystal, where $\operatorname{wt}(\nu) = -|\nu|\delta, \ \tilde{e}_i\nu = \tilde{f}_i\nu = \mathbf{0}$, and $\varepsilon_i(\nu) = \varphi_i(\nu) = 0$ for $\nu \in \mathscr{P}_{a-1}$ and $i \in I$.

Proof. (1) Since $\mathcal{B}((1^b))$ is invariant under \tilde{e}_i and \tilde{f}_i for $i \in I$, it is enough to show that $\tilde{e}_i T$ and $\tilde{f}_i T$ are semistandard. The proof is similar to the case of type A_{n-1} (cf. [5]).

(2) Let $T = (T^a, \ldots, T^1) \in \mathcal{B}(R)_0$ be given.

Suppose that $\tilde{e}_i T \notin \mathcal{B}(R)_0 \cup \{\mathbf{0}\}$ for some *i*. Then $\tilde{e}_i T = (\dots, \tilde{e}_i T^j, \dots)$ for some $1 \leq j \leq a$, and $\tilde{e}_i T^j$ is obtained from T^j by replacing $T^j(k)$ with $T^j(k) - 1$ for some $1 \leq k \leq b$. If $(\tilde{e}_i T^j, \tau^{-1}(T^{j-1}))$ is semistandard, then it is not difficult to see that $(T^j, \tau^{-1}(T^{j-1}))$ is also semistandard, which is a contradiction. For the other cases, we have similar contradiction. By the same arguments, we also have $\tilde{f}_i T \in \mathcal{B}(R)_0 \cup \{\mathbf{0}\}$. Hence $\mathcal{B}(R)_0$ is invariant under \tilde{e}_i and \tilde{f}_i for $i \in I$.

Conversely, we claim that any $T \in \mathcal{B}(R)_0$ is connected to u_R . Let $t = T^a(1)$. We first show that $T \in \mathcal{B}(R)_0$ is connected to $T_0 = \tau^t(u_R)$. Suppose that $T \neq T_0$. Let $d(T) = \sum_{j=1}^a \sum_{k=1}^b (T^j(k) - T_0^j(k)) \ge 0$. Since $T \in \mathcal{B}(R)_0$ and $\mathcal{B}(R)_0$ is invariant under \tilde{e}_i and \tilde{f}_i for $i \in I$, we have $0 \le d(\tilde{e}_i T) < d(T)$ for i such that $\tilde{e}_i T \neq \mathbf{0}$. Note that there exists at least one i such that $\tilde{e}_i T \neq \mathbf{0}$. By induction on d(T), we conclude that T is connected to T_0 .

Next, we have

$$\begin{cases} \tilde{e}^{a}_{t+b-2} \dots \tilde{e}^{a}_{t} \tilde{e}^{a}_{t-1} T_{0} = \tau^{t-1}(u_{R}) & (t > 1), \\ \tilde{f}^{a}_{t} \dots \tilde{f}^{a}_{t+b-2} \tilde{f}^{a}_{t+b-1} T_{0} = \tau^{t+1}(u_{R}) & (t < 1). \end{cases}$$

Repeating this step, we conclude that T_0 is connected to u_R . Therefore, $\mathcal{B}(R)_0$ is the connected component of u_R in $\mathcal{B}(R)$.

(3) It follows from Corollary 4.7.

Remark 4.9. Suppose that b = n. In this case, $\mathcal{B}(R)_0 = \{\tau^t(u_R) | t \in \mathbb{Z}\}$, where each $\{\tau^t(u_R)\}$ forms a trivial crystal of weight $-at\delta$.

Proposition 4.10. For $R = (a^b)$ with $a \ge 1$ and $1 \le b < n$, $\mathcal{B}(R)$ is isomorphic to $B(a\varpi_b)$ as an $A_{n-1}^{(1)}$ -crystal with P^0 -weights.

Proof. It follows from Proposition 4.6, Lemma 4.8, and [1, Theorem 4.16(a)] (see also [17, Theorem 2]. \Box

Let $\lambda \in \mathscr{P}_n$ be given and let $\mathscr{B}(\lambda)$ as given in (3.9). We regard $\mathscr{B}(\lambda)$ as an $A_{n-1}^{(1)}$ -crystal with P^0 -weights by identifying

$$\mathcal{B}(\lambda) = \mathcal{B}(R_1) \otimes \cdots \otimes \mathcal{B}(R_l).$$

Let

$$\varpi_{\lambda} = m_1 \varpi_1 + \dots + m_{n-1} \varpi_{n-1}$$

where m_i is the multiplicity of i in λ . By [1, Theorem 4.16] and [10, Conjectures 13.1, 13.2] (see also [1, Remark 4.17]), we have the following.

Proposition 4.11. For $\lambda \in \mathscr{P}_{n-1}$, $\mathfrak{B}(\lambda)$ is isomorphic to $B(\varpi_{\lambda})$.

Remark 4.12. (1) Let $u_{\lambda} = u_{R_1} \otimes \ldots \otimes u_{R_l}$. Then $\mathcal{B}(\lambda)_0 = \mathcal{B}(R_1)_0 \otimes \cdots \otimes \mathcal{B}(R_l)_0$ is the connected component of u_{λ} in $\mathcal{B}(\lambda)$ (see [1, Remark 4.17]). In case of $A_{n-1}^{(1)}$, we can prove it directly using the crystal structure described here.

(2) Suppose that $\lambda \in \mathscr{P}_n$ with $n = \ell(\lambda)$. We have $\mathscr{B}(\lambda) = \mathscr{B}(\mu) \otimes \mathscr{B}(R_n)$, where $\mu = (\lambda_1, \ldots, \lambda_{n-1})$ (see Remark 4.9).

(3) We may also regard $B(\lambda)$ as an $A_{n-1}^{(1)}$ -crystal with P_{cl}^{0} -weights by using $wt_{cl}^{0} = cl(wt_{cl}^{0})$.

Remark 4.13. Let $R = (a^b)$. Then the set $\mathcal{B}(R)_0/\mathbb{Z}$ is invariant under \tilde{e}_i and \tilde{f}_i for $i \in I$ since τ commutes with \tilde{e}_i and \tilde{f}_i , and $\mathrm{wt}_{\mathrm{cl}}^0$ is constant on any equivalence class. Hence by (3.7), we may assume that $CSST_{[n]}(R)$ is a normal $A_{n-1}^{(1)}$ -crystal with P_{cl}^0 -weights. Then $CSST_{[n]}(R)$ is isomorphic to $SST_{[n]}((1^b))^{\otimes a}$. Note that $SST_{[n]}((1^b)) = CSST_{[n]}((1^b))$ and it is isomorphic to the crystal of Kirillov-Reshetikhin module or level one perfect crystal associated to $\mathrm{cl}(\varpi_b)$ [8].

Furthermore, if (r_1, \ldots, r_{a-1}) is the offset vector of $T \in CSST_{[n]}(R)$, then $-r_j$ is equal to the value of the local energy function on $SST_{[n]}((1^b))^{\otimes 2}$ at (T^{j+1}, T^j) or $T^j \otimes T^{j+1}$ for $1 \leq j \leq a-1$ (see [18, Section 3]).

In general, for $\lambda \in \mathscr{P}_{n-1}$ we regard as an $A_{n-1}^{(1)}\text{-crystal}$

$$CSST_{[n]}(\lambda) = CSST_{[n]}(R_1) \otimes \cdots \otimes CSST_{[n]}(R_l).$$

Hence if we regard $\mathcal{B}(\lambda)$ as an $A_{n-1}^{(1)}$ -crystal with P_{cl}^{0} -weights, then the bijection (3.10)

$$CSST_{[n]}(\lambda) \times \mathcal{P}(\lambda) \longrightarrow \mathcal{B}(\lambda)$$

is an isomorphism.

4.5. Isomorphism of crystals. Let

(4.4)
$$\mathfrak{T}_{m \times n} = \bigsqcup_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} CSST_{[m]}(\lambda) \times \mathfrak{B}_n(\lambda).$$

We first assume that $\mathcal{T}_{m \times n}$ is a normal $A_{m-1}^{(1)}$ -crystal with P_{cl}^{0} -weights with respect to \tilde{e}_i, \tilde{f}_i for $i \in \{0, 1, \ldots, m-1\}$ for $m \ge 2$, where \tilde{e}_i, \tilde{f}_i are the Kashiwara operators on $CSST_{[m]}(\lambda)$ (see Remark 4.13). Also we assume that $\mathcal{T}_{m \times n}$ is a normal $A_{n-1}^{(1)}$ -crystal with P^0 -weights with respect to $\tilde{e}_j^t, \tilde{f}_j^t$ for $j \in \{0, 1, \ldots, n-1\}$ for $n \ge 2$, where $\tilde{e}_j^t, \tilde{f}_j^t$ are the Kashiwara operators on $\mathcal{B}_n(\lambda)$.

The following is the second main result in this paper. The proof is given in Section 6.

Theorem 4.14. The bijection

 $\kappa: \hat{\mathfrak{M}}_{m \times n} \longrightarrow \mathfrak{T}_{m \times n}$

commutes with \tilde{e}_i , \tilde{f}_i for $i \in \{1, \ldots, m-1\}$ and \tilde{e}_j^t , \tilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$.

We remark that the map κ does not commute with \tilde{e}_0 and \tilde{f}_0 , but $\kappa_1 := \pi_1 \circ \kappa$ does, where π_1 is the projection of $\mathcal{T}_{m \times n}$ along the first component (see Remark 6.9). Since wt⁰_{cl}(A) = wt⁰_{cl}(P₀), κ_1 induces the following.

Corollary 4.15. A generalized affine permutation $A \in \widehat{\mathcal{M}}_{m \times n}$ is $A_{m-1}^{(1)}$ -crystal equivalent to P_0 , where $\kappa(A) = (P_0, Q)$.

Note that κ is an isomorphism of $A_{n-1}^{(1)}$ -crystals with P_{cl}^{0} -weights, but it does not preserve P^{0} -weights. More precisely, for $A \in \widehat{\mathcal{M}}_{m \times n}$ with $\kappa(A) = (P_0, Q)$, we see from the definition of κ that

$$\operatorname{wt}^{t}(A) = \operatorname{wt}^{0}(A^{t}) = \operatorname{wt}^{0}(Q) - \left(\sum_{i=1}^{l} |\eta^{(i)}|\right) \delta,$$

where $\eta^{(i)}$ is the symmetrized offset vectors of $P_0^{(i)}$ in Section 3.5. So in order to have a morphism of $A_{n-1}^{(1)}$ crystals with P^0 -weights, we may modify the weight on $\widehat{\mathcal{M}}_{m \times n}$ in P^0 by

(4.5)
$$\mathbf{wt}^t(A) = \mathrm{wt}^0(A^t) + H_m(A)\delta,$$

where $H_m(A) = \sum_{i=1}^l |\eta^{(i)}|$. Then we have the following.

Corollary 4.16. The set $\widehat{\mathbb{M}}_{m \times n}$ is an $A_{n-1}^{(1)}$ -crystal with P^0 -weights with respect to \mathbf{wt}^t , \widetilde{e}_j^t , \widetilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$, and κ is an isomorphism of $A_{n-1}^{(1)}$ -crystals with P^0 -weights. In particular, a generalized affine permutation $A \in \widehat{\mathbb{M}}_{m \times n}$ is $A_{n-1}^{(1)}$ -crystal equivalent to Q, where $\kappa(A) = (P_0, Q)$. Moreover, $\widehat{\mathbb{M}}_{m \times n}$ decomposes into a disjoint union of crystals of extremal weight modules as follows:

$$\widehat{\mathcal{M}}_{m \times n} \cong \bigoplus_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} \mathcal{B}_n(\lambda)^{\bigoplus m_\lambda},$$

where $m_{\lambda} = |CSST_{[m]}(\lambda)|$, and \oplus means a disjoint union of crystals.

We remark that not both m and n need to be greater than 1 for Theorem 4.14 and its corollaries. In particular, Corollary 4.15 holds for n = 1 and Corollary 4.16 holds for m = 1.

Remark 4.17. The function $H_m(\cdot)$ in (4.5) is related to the intrinsic energy function on $A_{m-1}^{(1)}$ -crystals with P_{cl}^{0} -weights as follows. Let $T \in CSST_{[m]}(R)$ be given where $R = (a^b)$. Let $r = (r_1, \ldots, r_{a-1})$ be the offset vector and $\eta = (\eta_1, \ldots, \eta_{a-1})$ the symmetrized offset vector of T. Then we have

$$\mathcal{H}_m(T) = |\eta| - a|r|,$$

where $|r| = r_1 + \cdots + r_{a-1}$ and $\mathcal{H}_m(\cdot)$ is the intrinsic energy function on $CSST_{[n]}(R)$ with $\mathcal{H}_m(u_R) = 0$ (cf. [22] for its definition). Hence for $A \in \mathcal{M}_{m \times n}$ with $\kappa(A) = (P_0, Q)$, we have

$$H_m(A) = \sum_{i=1}^{l} \left(\mathcal{H}_m(P_0^{(i)}) + m_i |r^{(i)}| \right),$$

where $R_i = (a_i^{m_i})$ is the shape and $r^{(i)}$ is the offset vector of $P_0^{(i)}$ in Section 3.5, respectively.

5. Proof of Theorem 3.9

5.1. Standardization and κ_0 . In this subsection, we show the map κ_0 in (3.3) is compatible with the standardization (2.3).

Let $A = (a_{ij})_{i,j \in \mathbb{Z}} \in \widehat{\mathcal{M}}_{m \times n}$ be given. Let

$$\operatorname{row}(A) = \left(\sum_{j \in \mathbb{Z}} a_{1j}, \dots, \sum_{j \in \mathbb{Z}} a_{mj}\right), \quad \operatorname{col}(A) = \left(\sum_{i \in \mathbb{Z}} a_{i1}, \dots, \sum_{i \in \mathbb{Z}} a_{in}\right)$$

be the row and column contents of A, respectively.

Let $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{Z}_+^m$ and $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{Z}_+^n$ be given such that $|\alpha| = |\beta|$, where $|\alpha| = \alpha_1 + \cdots + \alpha_m$. Put $\operatorname{supp}(\alpha) = \{i \in [m] \mid \alpha_i \neq 0\}$, $\operatorname{supp}(\beta) = \{j \in [n] \mid \beta_j \neq 0\}$, and $K = |\alpha|$. Let

$$\widehat{\mathcal{M}}_{m \times n}(\alpha, \beta) = \{ A \mid A \in \widehat{\mathcal{M}}_{m \times n}, \text{ row}(A) = \alpha, \text{ col}(A) = \beta \}.$$

We have $\widehat{\mathfrak{M}}_{m \times n} = \bigsqcup_{\alpha, \beta} \widehat{\mathfrak{M}}_{m \times n}(\alpha, \beta).$

Suppose that $A \in \widehat{\mathcal{M}}_{m \times n}(\alpha, \beta)$ is given. Let A^{st} be the standardization of A in (2.3), which is a partial extended affine permutation. By removing the rows and columns in A^{st} with zero entries only, we may obtain an extended affine permutation, which we denote by A^{perm} .

Let us describe more explicitly the index sets for the rows and columns in A^{perm} . We may assume that K > 1 since A^{perm} is trivial in case of K = 0, 1.

For $i \in \text{supp}(\alpha)$ and $j \in \text{supp}(\beta)$, let

$$\begin{aligned} \mathfrak{I}_i &= \{ k \in [K] \mid \alpha_0 + \alpha_1 + \dots + \alpha_{i-1} < k \leq \alpha_1 + \dots + \alpha_{i-1} + \alpha_i \}, \\ \mathfrak{J}_j &= \{ l \in [K] \mid \beta_0 + \beta_1 + \dots + \beta_{j-1} < l \leq \beta_1 + \dots + \beta_{j-1} + \beta_j \}, \end{aligned}$$

where $\alpha_0 = \beta_0 = 0$, and let

$$\mathcal{I}_{i+sm} = \mathcal{I}_i + sK, \quad \mathcal{J}_{j+sn} = \mathcal{J}_j + sK \quad (s \in \mathbb{Z}).$$

Then we have

(5.1)
$$[K] = \bigsqcup_{i \in \operatorname{supp}(\alpha)} \mathfrak{I}_{i} = \bigsqcup_{j \in \operatorname{supp}(\beta)} \mathfrak{J}_{j}$$
$$\mathbb{Z} = \bigsqcup_{i \in \mathfrak{Z}_{\alpha}} \mathfrak{I}_{i} = \bigsqcup_{j \in \mathfrak{Z}_{\beta}} \mathfrak{J}_{j},$$

where $\mathcal{Z}_{\alpha} = \operatorname{supp}(\alpha) + \mathbb{Z}m$ and $\mathcal{Z}_{\beta} = \operatorname{supp}(\beta) + \mathbb{Z}n$. Note that $\tau_K(\mathcal{I}_i \times \mathcal{J}_j) = \mathcal{I}_{i+m} \times \mathcal{J}_{j+n}$ for $(i, j) \in \mathcal{Z}_{\alpha} \times \mathcal{Z}_{\beta}$. Hence we may regard

where the row and column indices are partitioned as in (5.1).

Remark 5.1. For $(i, j) \in \mathbb{Z}_{\alpha} \times \mathbb{Z}_{\beta}$, the submatrix of A^{perm} corresponding to $\mathcal{J}_i \times \mathcal{J}_j$, which is an $\alpha_i \times \beta_j$ matrix, includes an $a_{ij} \times a_{ij}$ matrix with 1 on the antidiagonal and 0 elsewhere, and has no other non-zero entry.

For K > 1, let $\widetilde{W}_K \subset \widehat{\mathcal{M}}_{K \times K}$ denote the set of extended affine permutations of type $A_{K-1}^{(1)}$. Let $B \in \widetilde{W}_K$ be given. We say that $i \in [K]$ is a *descent* of B if $(i, j), (i + 1, j') \in \operatorname{supp}(B)$ and j > j'. Let $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{Z}_+^m$ and $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{Z}_+^n$ be given with $|\alpha| = |\beta| = K$. We say that $B \in \widetilde{W}_K$ is α -descending if for any $k \in \operatorname{supp}(\alpha)$ and i with

$$\alpha_0 + \alpha_1 + \dots + \alpha_{k-1} < i < \alpha_1 + \dots + \alpha_{k-1} + \alpha_k,$$

i is a descent of *B*, where $\alpha_0 = 0$. We say that *B* is (α, β) -descending if *B* is α -descending and B^t is β -descending. Let $\widetilde{W}_{K,(\alpha,\beta)}$ denote the set of extended affine permutations which are (α, β) -descending.

Lemma 5.2. Under the above hypothesis, we have a bijection

$$\widehat{\mathcal{M}}_{m \times n}(\alpha, \beta) \longrightarrow \widetilde{W}_{K,(\alpha,\beta)} \\
A \longmapsto A^{\operatorname{perm}}$$

Proof. For $A \in \widehat{\mathcal{M}}_{m \times n}(\alpha, \beta)$, we have $A^{\operatorname{perm}} \in \widetilde{W}_{K,(\alpha,\beta)}$ by Lemma 2.5, where we replace I_i and J_j by \mathfrak{I}_i and \mathfrak{J}_j , respectively. Hence the map is well-defined.

Let us prove the existence of its inverse. Let $B = (b_{kl})_{k,l \in \mathbb{Z}} \in \widetilde{W}_{K,(\alpha,\beta)}$ be given. Consider the partition $\{\mathcal{I}_i \times \mathcal{J}_j\}_{(i,j)\in \mathbb{Z}_{\alpha}\times \mathbb{Z}_{\beta}}$ of $\mathbb{Z} \times \mathbb{Z}$ associated to α and β in (5.1). Then we define $A = (a_{ij})_{i,j \in \mathbb{Z}}$ to be a $\mathbb{Z} \times \mathbb{Z}$ matrix where

$$a_{ij} = \begin{cases} \sum_{(k,l)\in\mathcal{I}_i\times\mathcal{J}_j} b_{kl} & \text{if } (i,j)\in\mathcal{Z}_\alpha\times\mathcal{Z}_\beta, \\ 0 & \text{otherwise.} \end{cases}$$

Since $B \in \widetilde{W}_{K,(\alpha,\beta)}$, we see from the definition that $\mathcal{I}_i \times \mathcal{J}_j$ -submatrix of B contains an $a_{ij} \times a_{ij}$ matrix with 1 on the antidiagonal and 0 elsewhere. Then it is easy to see that A is the unique matrix in $\widehat{\mathcal{M}}_{m \times n}(\alpha,\beta)$ such that $A^{\text{perm}} = B$. This proves the bijectivity. \Box

Now, let us consider standardization on the side of tableaux in (3.3).

Let $\lambda \in \mathscr{P}_n$ and $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}^n_+$ be given. Let $CSST_{[n]}(\lambda)_{\alpha}$ be the set of $T \in CSST_{[n]}(\lambda)$ with content α , that is, α_k is the number of occurrence of $k \in [n]$ in T. For $T \in CSST_{[n]}(\lambda)_{\alpha}$, we define T^{st} to be a tableau obtained from T by replacing each $k \in [n]$ in T with $\alpha_k \neq 0$ by the consecutive numbers

 $\alpha_0 + \alpha_1 + \dots + \alpha_{k-1} + 1 < \dots < \alpha_1 + \dots + \alpha_{k-1} + \alpha_k$

from left to right, where $\alpha_0 = 0$. Let $K = |\lambda|$. By definition, we have

$$T^{\mathtt{st}} \in CST_{[K]}(\lambda),$$

(cf. Section 2.6). We call T^{st} the standardization of T. The following lemma implies that the symmetrized offset vector of T is invariant under standardization.

Lemma 5.3. Let $T \in CSST_{[n]}(R)$ be given where $R = (a^b) \in \mathscr{P}_n$ for some $a, b \ge 1$. For $\nu \in \mathscr{P}_{a-1}$, we have

$$\tau_n^{\nu_{\text{rev}}}(T) \in \mathcal{B}_n(R)_0 \quad \text{if and only if} \quad \tau_K^{\nu_{\text{rev}}}(T^{\text{st}}) \in \mathcal{B}_K(R)_0,$$

where K = ab.

Proof. Let $T = (T^a, \ldots, T^1)$ and $T^{st} = S = (S^a, \ldots, S^1)$. For $1 \leq j \leq a - 1$, let r_j be the smallest integer satisfying $(T^{j+1}, \tau_n^{r_j}(T^j))$ is semistandard. It suffices to show that $(S^{j+1}, \tau_K^{r_j}(S^j))$ is semistandard but $(S^{j+1}, \tau_K^{r_j-1}(S^j))$ is not. It is straightforward to see from the definition of T^{st} .

Let $T \in CST_{[K]}(\lambda)$ be given with $|\lambda| = K$. We say that $i \in [K]$ is a *(column) descent* of T if the letter i + 1 appears to the right of i in T (i + 1) lies in the column with a smaller index than that of i). For $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}^n_+$ with $K = |\alpha|$, we say that $T \in CST_{[K]}(\lambda)$ is α -descending if for any $k \in [n]$ and i with

$$\alpha_0 + \alpha_1 + \dots + \alpha_{k-1} < i < \alpha_1 + \dots + \alpha_{k-1} + \alpha_k,$$

i is a descent of *T*, where $\alpha_0 = 0$. Let $CST_{[K],\alpha}(\lambda)$ be the set of $T \in CST_{[K]}(\lambda)$ which are α -descending. Then we can check the following.

Lemma 5.4. Under the above hypothesis, we have a bijection

$$CSST_{[n]}(\lambda)_{\alpha} \longrightarrow CST_{[K],\alpha}(\lambda)$$
$$T \longmapsto T^{st}$$

The following generalizes a well-known property of the usual RS correspondence (cf. [4, Section 1.1]).

Lemma 5.5. Let $B \in \widetilde{W}_K$ be given with $\kappa_0(B) = (P_0, Q_0, \rho)$. Then $i \in [K]$ is a descent of B if and only if i is a descent of P_0 . Similarly, $j \in [K]$ is a descent of B^t if and only if j is a descent of Q_0 .

Proof. Let $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ be the set of zig-zags associated to $d = d_B^{sw}$. Let $i \in [K]$ be given. Let $c_1 = (i, j), c_2 = (i + 1, j') \in \operatorname{supp}(B)$ and let $d(c_1) = k_1, d(c_2) = k_2$.

Suppose that *i* is a descent of *B*, that is, j' < j or $c_2 \leq_{ne} c_1$. Since *d* is continuous, there exists $c_3 \in \text{supp}(B)$ such that $c_3 >_{NW} c_2$ with $d(c_3) = k_2 - 1$. Since c_1 is the unique non-zero cell which lies in the *i*-th row of *B*, we have $c_3 >_{NW} c_1$. Hence we have $k_2 - 1 < k_1$ or $k_2 \leq k_1$, which implies that $\mathbf{z}_{k_1} = \mathbf{z}_{k_2}$ or \mathbf{z}_{k_1} is located to the southwest of \mathbf{z}_{k_2} in the sense of (3.1). So we conclude that c_2 is not the rightmost inner corner of a zigzag, and hence i + 1 does not occur in the first column of P_0 (from the left). In particular, if *i* occurs in the first column of P_0 . Let c_4 and c_5 be the unique outer corners in the *i*-th and (i + 1)-th rows in \mathbf{z}_{k_1} and \mathbf{z}_{k_2} , respectively. Note that $c_5 \leq_{ne} c_4$ since $k_2 \leq k_1$, and $c_4 \neq c_1$ since *i* does not occur in the first column of P_0 . Hence c_4 and c_5 are the inner corners of B^{\flat} , which implies that *i* is a descent of $\sum_{i \in [m]} \sum_{j \in \mathbb{Z}} b_{ij}$ where $B = (b_{ij})$, we conclude that *i* is a descent of P_0 .

Conversely, suppose that *i* is not a descent of *B*, that is, j < j'. Since $c_1 >_{NW} c_2$, we see that $k_1 < k_2$ and \mathbf{z}_{k_2} is located to the southeast of \mathbf{z}_{k_1} . Hence $c_1^* >_{NW} c_2^*$, where c_1^* and c_2^* are the back-post corners of \mathbf{z}_{k_1} and \mathbf{z}_{k_2} respectively. If *i* appears in the first column of P_0 ,

then i + 1 also appears in the first column of P_0 since $c_1^* >_{NW} c_2^*$. If both i and i + 1 do not appear in the first column of P_0 , then by similar arguments in the previous paragraph we see that i is not a descent of B^{\flat} . Therefore, we conclude that i is not a descent of P_0 .

The proof for B^t are almost the same.

Corollary 5.6. Let $B \in \widetilde{W}_K$ be given with $\kappa_0(B) = (P_0, Q_0, \rho)$, where $P_0, Q_0 \in CST_{[K]}(\lambda)$ and $\rho \in \mathbb{Z}_+^{\lambda_1}$ with $\lambda \in \mathscr{P}_m \cap \mathscr{P}_n$. For $\alpha \in \mathbb{Z}_+^m$ and $\beta \in \mathbb{Z}_+^n$ with $|\alpha| = |\beta| = K$, the following are equivalent:

- (1) $B \in \widetilde{W}_{K,(\alpha,\beta)},$
- (2) $P_0 \in CST_{[K],\alpha}(\lambda)$ and $Q_0 \in CST_{[K],\beta}(\lambda)$.

It is clear that κ_0 in (3.3) preserves contents, that is, if $A \in \widehat{\mathcal{M}}_{m \times n}(\alpha, \beta)$ with $\kappa_0(A) = (P_0, Q_0, \rho)$, then $P_0 \in CSST_{[m]}(\lambda)_{\alpha}$ and $Q_0 \in CSST_{[n]}(\lambda)_{\beta}$. The following proposition shows that κ_0 is compatible with standardizations.

Proposition 5.7. For $A \in \widehat{\mathcal{M}}_{m \times n}$ with $\kappa_0(A) = (P_0, Q_0, \rho)$, we have

$$\kappa_0(A^{\texttt{perm}}) = (P_0^{\texttt{st}}, Q_0^{\texttt{st}}, \rho)$$

Proof. Assume that $A \in \widehat{\mathcal{M}}_{m \times n}(\alpha, \beta)$ for some $\alpha \in \mathbb{Z}_{+}^{m}$ and $\beta \in \mathbb{Z}_{+}^{n}$ with $|\alpha| = |\beta| = K$. Let $\kappa_{0}(A^{\text{perm}}) = (P_{1}, Q_{1}, \varrho)$. By Lemma 5.2 and Corollary 5.6, we have

$$P_1 \in CST_{[K],\alpha}(\lambda), \quad Q_1 \in CST_{[K],\beta}(\lambda)$$

for some $\lambda \in \mathscr{P}$. Since A^{perm} is obtained from A^{st} by removing the rows and columns with no positive entries, we have $\kappa_0(A^{\text{st}}) = \kappa_0(A^{\text{perm}})$, where we identify the indices of remaining rows and columns in A^{st} with the those of A^{perm} .

We claim that P_0 (resp. Q_0) is the image of P_1 (resp. Q_1) under the inverse of the bijection in Lemma 5.4 and $\rho = \rho$. Let $\{\mathcal{J}_i \times \mathcal{J}_j\}_{(i,j) \in \mathcal{Z}_\alpha \times \mathcal{Z}_\beta}$ be the partition of $\mathbb{Z} \times \mathbb{Z}$ associated to α and β in (5.1). Let \mathbf{s} and \mathbf{s}^{st} be the stream given from the the back-post corners of the zig-zags $\{\mathbf{z}_k\}_{k \in \mathbb{Z}}$ and $\{\mathbf{z}_k^{st}\}_{k \in \mathbb{Z}}$ associated to d_A^{sw} and $d_{A^{st}}^{sw}$, respectively. Since we have $(d_A^{sw})^{st} = d_{A^{st}}^{sw}$ by (2.7), the flow of \mathbf{s} is equal to that of \mathbf{s}^{st} , and \mathbf{s} is obtained from \mathbf{s}^{st} by replacing each (k, l) in \mathbf{s}^{st} with (i, j) when $(k, l) \in \mathcal{I}_i \times \mathcal{J}_j$. Since $(A^{\flat})^{st} = (A^{st})^{\flat}$ (see Remark 3.5), we use induction on K to prove the claim. Hence we have $P_0^{st} = P_1$ and $Q_0^{st} = Q_1$.

5.2. Dominance condition and κ . In this subsection, we characterize the image of κ_0 , and then prove Theorem 3.9.

5.2.1. Offset constants. Let $\lambda = (2^l)$ be of rectangular shape with two columns for $1 \leq l \leq \min\{m, n\}$.

Suppose that $P_0 = (P_0^2, P_0^1) \in CSST_{[m]}(\lambda)$ and $Q_0 = (Q_0^2, Q_0^1) \in CSST_{[n]}(\lambda)$ are given. Let $(0, \eta)$ and $(0, \theta)$ be the offset vectors for P_0 and Q_0 , that is,

(5.3)
$$\tau^{(0,\eta)}(P_0) = \left(P_0^2, \tau^{\eta}(P_0^1)\right) \in \mathcal{B}_m(\lambda)_0, \quad \tau^{(0,\theta)}(Q_0) = \left(Q_0^2, \tau^{\theta}(Q_0^1)\right) \in \mathcal{B}_n(\lambda)_0,$$

and put

(5.4)
$$r = \theta - \eta$$

We claim that r is equal to the offset constant introduced in [3, Section 5.2], which plays an important role in characterizing the image of κ_0 in case of extended affine permutations.

Write $P_0 = (\mathbf{a}_1, \mathbf{a}_2)$ and $Q_0 = (\mathbf{b}_1, \mathbf{b}_2)$. For $(\rho_1, \rho_2) \in \mathbb{Z}^2$, let

$$\mathbf{s} = \{ c_i = (a_i, b_i) \}_{i \in \mathbb{Z}}, \quad \mathbf{s}' = \{ c'_i = (a'_i, b'_i) \}_{i \in \mathbb{Z}}$$

be the streams of flow l, whose defining data are $(\mathbf{a}_1, \mathbf{b}_1, \rho_1)$ and $(\mathbf{a}_2, \mathbf{b}_2, \rho_2)$, respectively. We may assume that

(5.5)
$$\mathbf{a}_{1} = (a_{1}, \dots, a_{l}), \quad \mathbf{b}_{1} = (b_{1-\rho_{1}}, \dots, b_{l-\rho_{1}}), \mathbf{a}_{2} = (a'_{1}, \dots, a'_{l}), \quad \mathbf{b}_{2} = (b'_{1-\rho_{2}}, \dots, b'_{l-\rho_{2}}).$$

Let k be the smallest integer such that $c_i \geq_{nw} c'_{i+k}$ for all $i \in \mathbb{Z}$, and then consider the following two streams

$$\mathbf{t} = \{ d_i = (a'_{i+k}, b_i) \}_{i \in \mathbb{Z}}, \quad \mathbf{t}' = \{ d'_i = (a_i, b'_{i+k}) \}_{i \in \mathbb{Z}}.$$

Let $A = (a_{ij})_{i,j \in \mathbb{Z}} \in \widehat{\mathcal{M}}_{m \times n}$, where a_{ij} is the number of occurrence of (i, j) in $\mathbf{t} \cup \mathbf{t'}$. It is easy to observe that l is the width of A and \mathbf{t} is the southwest channel of A. Let d be a proper numbering on A defined by $d(d_i) = d(d'_i) = i$ for $i \in \mathbb{Z}$.

Lemma 5.8. Under the above hypothesis, we have $\rho_2 - \rho_1 \ge r$ if and only if $d = d_A^{sw}$.

Proof. We observe that in order to show $d = d_A^{sw}$, it suffices to find *i* such that $d_i >_{NW} d'_{i+1}$ by the condition (2) in Lemma 2.22. Furthermore, $d_{i-1} >_{NW} d'_i$ for some *i* is equivalent to $a'_{i-1+k} < a_i$ since $b_{i-1} < b'_{i+k}$ is redundant by our choice of *k*.

By (5.3), the constants η and θ are the smallest integers such that

$$a_i \leqslant a'_{i+\eta}, \quad b_i \leqslant b'_{i+\rho_1-\rho_2+\theta} \quad (i \in \mathbb{Z}),$$

respectively. By the minimality of η , θ , we have $\eta \leq k$ and $\theta \leq \rho_2 - \rho_1 + k$.

Suppose that $d = d_A^{sw}$. Then by the above observation, there exists *i* with $a_i > a'_{i+k-1}$. In this case, we have $k - 1 < \eta \leq k$ by the minimality of η . Hence $\eta = k$ and

$$\rho_2 - \rho_1 \ge \theta - k = \theta - \eta = r.$$

Note that we have $\eta \ge \rho_1 - \rho_2 + \theta$, which implies that $b_i \le b'_{i+\eta}$.

Conversely, suppose that $\rho_2 - \rho_1 \ge \theta - \eta$. Since $\rho_2 - \rho_1 + \eta \ge \theta$, we have

(5.6)
$$b_i \leq b'_{i+\rho_1-\rho_2+(\rho_2-\rho_1+\eta)} = b'_{i+\eta} \quad (i \in \mathbb{Z}).$$

Since $a_i \leq a'_{i+\eta}$, we have $k \leq \eta$ by the minimality of k, which implies $a'_{i+k-1} < a_i$ for some i. Hence $d = d_A^{sw}$.

The following is another characterization of r.

Corollary 5.9. Under the above hypothesis, we have $\rho_2 - \rho_1 \ge r$ if and only if $\tau^{(\rho_1,\rho_2+\eta)}(Q_0) \in \mathcal{B}_n(\lambda).$

Proof. We have shown in the proof of Lemma 5.8 that $\rho_2 - \rho_1 \ge r$ is equivalent to $b_i \le b'_{i+\eta}$ (5.6). On the other hand, by (5.5) it is equivalent to

$$\tau^{(\rho_1,\rho_2+\eta)}(Q_0) = (\tau^{\rho_1}\mathbf{b}_1,\tau^{\rho_2+\eta}\mathbf{b}_2) \in \mathcal{B}_n(\lambda).$$

Remark 5.10. It is not difficult to see that the constants η and θ are equal to the *local* charges of P_0 and Q_0 , respectively [2, Definition 5.3]. Hence by [2, Theorem 5.10], the constant $r = \theta - \eta$ in (5.4) is equal to the offset constant of (P_0, Q_0) in [3] when $P_0, Q_0 \in CST_{[K]}((2^l))$ for some K.

5.2.2. Proof of Theorem 3.9. We keep the notations in Section 3.5. Suppose that $(P_0, Q_0, \rho) \in CSST_{[m]}(\lambda) \times CSST_{[n]}(\lambda) \times \mathbb{Z}^{\lambda_1}$ is given for $\lambda \in \mathscr{P}_m \cap \mathscr{P}_n$.

For $1 \leq i \leq l$ with $m_i \geq 1$, we let

- $\rho^{(i)} \in \mathbb{Z}^{m_i}$: the subsequence of ρ corresponding to the columns of R_i ,
- $\eta^{(i)} \in \mathscr{P}_{m_i-1}$: the symmetrized offset vector of $P_0^{(i)}$,
- $\theta^{(i)} \in \mathscr{P}_{m_i-1}$: the symmetrized offset vector of $Q_0^{(i)}$,
- $\zeta^{(i)} = \theta^{(i)} \eta^{(i)} \in \mathbb{Z}^{m_i}$ with the last component being zero.

We write $\zeta = (\zeta^{(1)}, ..., \zeta^{(l)}).$

Definition 5.11. Under the above hypothesis, we say that (P_0, Q_0, ρ) is *dominant* if

$$\rho_{\text{rev}} - \zeta \in \mathcal{P}(\lambda).$$

Lemma 5.12. Under the above hypothesis, we have (P_0, Q_0, ρ) is dominant if and only if $(P_0^{st}, Q_0^{st}, \rho)$ is dominant.

Proof. For $1 \leq i \leq l$ with $m_i \geq 1$, let $\eta_{\mathtt{st}}^{(i)}$ and $\theta_{\mathtt{st}}^{(i)}$ be the symmetrized offset vectors of $(P_0^{\mathtt{st}})^{(i)}$ and $(Q_0^{\mathtt{st}})^{(i)}$ respectively. It is immediate from Lemma 5.3 that $\eta^{(i)} = \eta_{\mathtt{st}}^{(i)}$ and $\theta^{(i)} = \theta_{\mathtt{st}}^{(i)}$. Hence the assertion follows.

Remark 5.13. By Remark 5.10, we see that $(P_0^{st}, Q_0^{st}, \rho)$ is dominant if and only if ρ is dominant in the sense of [3, Definition 5.8].

Now, we can describe the image of κ_0 as follows.

Proposition 5.14. Let

$$\Omega_{\rm dom} = \left\{ \begin{array}{c|c} (P_0, Q_0, \rho) \\ (P_0, Q_0, \rho) \end{array} \middle| \begin{array}{c} (1) \ P_0 \in CSST_{[m]}(\lambda), \ Q_0 \in CSST_{[n]}(\lambda) & (\lambda \in \mathscr{P}_m \cap \mathscr{P}_n), \\ (2) \ \rho \in \mathbb{Z}^{\lambda_1}, \\ (3) \ (P_0, Q_0, \rho) \ is \ dominant. \end{array} \right\}.$$

Then the map κ_0 gives a bijection

$$\widehat{\mathfrak{M}}_{m \times n} \longrightarrow \Omega_{\mathrm{dom}}$$
.

Proof. Suppose that $A \in \widehat{\mathcal{M}}_{m \times n}$ is given. We have $\kappa_0(A) = (P_0, Q_0, \rho)$ for some $P_0 \in CSST_{[m]}(\lambda), Q_0 \in CSST_{[n]}(\lambda)$ with $\lambda \in \mathscr{P}_m \cap \mathscr{P}_n$, and $\rho \in \mathbb{Z}^{\lambda_1}$.

Then we have $\kappa_0(A^{\text{perm}}) = (P_0^{\text{st}}, Q^{\text{st}}, \rho)$ by Proposition 5.7. Since A^{perm} is an extended affine permutation, $(P_0^{\text{st}}, Q^{\text{st}}, \rho)$ is dominant by [3, Theorem 5.11] (see Remark 5.13), and hence $\kappa_0(A) = (P_0, Q_0, \rho)$ is dominant by Lemma 5.12.

Conversely, suppose that $(P_0, Q_0, \rho) \in \Omega_{\text{dom}}$ is given. We claim that there exists unique $A \in \widehat{\mathcal{M}}_{m \times n}$ with $\kappa_0(A) = (P_0, Q_0, \rho)$.

Let α and β be the contents of P_0 and Q_0 respectively, and let $K = |\alpha| = |\beta| = |\lambda|$. By Lemma 5.12, $(P_0^{st}, Q_0^{st}, \rho)$ is dominant. Then it follows from [3, Theorem 5.1, Theorem 5.12] that there exists a unique extended affine permutation $B \in \widehat{\mathcal{M}}_{K \times K}$ such that $\kappa_0(B) = (P_0^{st}, Q_0^{st}, \rho)$. By Corollary 5.6, B is (α, β) -descending, and hence by Lemma 5.2 there exists a unique $A \in \widehat{\mathcal{M}}_{m \times n}(\alpha, \beta)$ such that $A^{\text{perm}} = B$. Finally, we have $\kappa_0(A) = (P_0, Q_0, \rho)$ by Lemma 5.4.

Corollary 5.15. The map κ is well-defined.

Proof. It follows from Lemma 5.9.

Lemma 5.16. For $\lambda \in \mathscr{P}_m \cap \mathscr{P}_n$, let $\Omega_{\text{dom}}(\lambda)$ denote the set of $(P_0, Q_0, \rho) \in \Omega_{\text{dom}}$ such that the shape of P_0 and Q_0 is λ . Then we have a bijection

$$\Omega_{\text{dom}}(\lambda) \longrightarrow CSST_{[m]}(\lambda) \times \mathcal{B}_n(\lambda)$$
$$(P_0, Q_0, \rho) \longmapsto (P_0, Q)$$

where Q is given as in (3.11).

Proof. Consider a bijection

(5.7)
$$\Omega_{\text{dom}}(\lambda) \longrightarrow CSST_{[m]}(\lambda) \times CSST_{[n]}(\lambda) \times \mathcal{P}(\lambda) .$$
$$(P_0, Q_0, \rho) \longmapsto (P_0, Q_0, \rho_{\text{rev}} - \zeta)$$

Then the map $(P_0, Q_0, \rho) \mapsto (P_0, Q)$ is a composition of (5.7) followed by the bijection (3.10) to the last two component respectively. Hence the bijectivity follows. \Box

Since $\Omega_{\text{dom}} = \bigsqcup_{\lambda \in \mathscr{P}_m \cap \mathscr{P}_n} \Omega_{\text{dom}}(\lambda)$, Theorem 3.9 follows from Proposition 5.14 and Lemma 5.16.

Remark 5.17. The well-definedness of κ also follows directly from Proposition 5.14 and Lemma 5.16.

6. Proof of Theorem 4.14

In this section, we prove

(6.1) $\kappa(\widetilde{x}_i A) = \widetilde{x}_i \kappa(A), \quad \kappa(\widetilde{y}_i^t A) = \widetilde{y}_i^t \kappa(A),$

for $A \in \widehat{\mathcal{M}}_{m \times n}$, $i \in \{1, \dots, m-1\}, j \in \{0, 1, \dots, n-1\}$ and $x, y \in \{e, f\}$.

6.1. Notations. Let **s** be a stream of flow l with defining data $(\mathbf{a}, \mathbf{b}, r)$. We regard **s** as an element of $\mathcal{T}_{m \times n}$ in (4.4), which is both an $A_{m-1}^{(1)}$ and $A_{n-1}^{(1)}$ crystal with P_{cl}^{0} -weights.

$$\mathbf{s} = (\mathbf{a}, \tau^r \mathbf{b}) \in CSST_{[m]}((1^l)) \times \mathcal{B}_n((1^l)) \subset \mathcal{T}_{m \times n}.$$

For $A \in \widehat{\mathcal{M}}_{m \times n}$ with the corresponding streams $\mathbf{s}^{(1)}, \ldots, \mathbf{s}^{(s)}$ in Section 3.2, we identify $\kappa(A)$ with $\mathbf{s}^{(s)} \otimes \cdots \otimes \mathbf{s}^{(1)}$.

Define a map

$$\Psi: \ \widehat{\mathcal{M}}_{m \times n} \longrightarrow \bigsqcup_{l \ge 0} \ \widehat{\mathcal{M}}_{m \times n} \otimes \left(CSST_{[m]}((1^l)) \times \mathcal{B}_n((1^l)) \right)$$
$$A \longmapsto A^{\flat} \otimes \mathbf{s}^{(1)}$$

Since

(6.2)
$$\left(\left(\Psi \otimes \mathrm{id}^{\otimes s-1}\right) \circ \cdots \circ \left(\Psi \otimes \mathrm{id}\right) \circ \Psi\right)(A) = \mathbb{O} \otimes \mathbf{s}^{(s)} \otimes \cdots \otimes \mathbf{s}^{(1)} = \mathbb{O} \otimes \kappa(A),$$

where id is the identity morphism, it suffices to show that Ψ commutes with \tilde{x}_i and \tilde{y}_j^t for the proof of (6.1).

In order to simplify the description of \tilde{x}_i and \tilde{y}_j^t on $A^{\flat} \otimes \mathbf{s}$ (see (6.6)), let us introduce some additional notations and conventions. Let $\mathbb{Z}^* = \mathbb{Z} \cup \{\infty\}$, where we understand that $a < \infty$ and $a + \infty = \infty$ for $a \in \mathbb{Z}$. Let $A \in \widehat{\mathcal{M}}_{m \times n}$ be given. Let $\Psi(A) = A^{\flat} \otimes \mathbf{s}^{(1)}$, where $A^{\flat} = (a_{ij}^{\flat})_{i,j\in\mathbb{Z}}$. We define $A^* = (a_{ij}^*)_{i,j\in\mathbb{Z}^*}$ by

$$a_{ij}^* = \begin{cases} a_{ij}^\flat & \text{if } (i,j) \in \mathbb{Z} \times \mathbb{Z}, \\ 1 & \text{if } i = \infty \text{ and } (k,j) \in \mathbf{s}^{(1)} \text{ for some } k \in \mathbb{Z}, \\ 1 & \text{if } j = \infty \text{ and } (i,k) \in \mathbf{s}^{(1)} \text{ for some } k \in \mathbb{Z}, \\ 0 & \text{otherwise.} \end{cases}$$

In other words, A^* is an augmented matrix obtained from A^{\flat} by

$$A^* = A^{\flat} + \sum_{(i,j)\in\mathbf{s}^{(1)}} (E_{i\infty} + E_{\infty j}).$$

Note that A^* satisfies $a_{i+mj+n}^* = a_{ij}^*$ for $(i, j) \in \mathbb{Z}^* \times \mathbb{Z}^*$.

Let \mathbf{z} be a zig-zag of A with the back-post corner (i, j). Let $\mathbf{z}^* = \mathbf{z} \cup \{(\infty, j), (i, \infty)\}$ and regard (∞, j) and (i, ∞) as outer corners of \mathbf{z}^* . Then we may understand $\Psi(A) = A^*$ as a $\mathbb{Z}^* \times \mathbb{Z}^*$ -matrix obtained by

- identifying $A = (a_{ij})_{i,j\in\mathbb{Z}}$ with $(a_{ij})_{i,j\in\mathbb{Z}}$ where $a_{\infty j} = a_{i\infty} = 0$ for $i, j \in \mathbb{Z}^*$,
- applying the same rules for \flat in Section 3.2 to A along \mathbf{z}_k^* instead of \mathbf{z}_k ,

where $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ is the set of zig-zags associated to d_A^{sw} . Note that we can recover A^{\flat} and $\mathbf{s}^{(1)}$ from A^* and $\{\mathbf{z}_k^*\}_{k\in\mathbb{Z}}$. From now on, we assume that a matrix is a $\mathbb{Z}^* \times \mathbb{Z}^*$ -matrix and a zig-zag is of the form \mathbf{z}^* .

Example 6.1. Let A be the generalized affine permutation in Example 2.1. We regard A as a $\mathbb{Z}^* \times \mathbb{Z}^*$ matrix as follows.



where the red lines denote the zig-zags $\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3$ associated to d_A^{sw} . Then A^* is given as follows.



6.2. Tensor product rule. From now on, we fix $A \in \widehat{\mathcal{M}}_{m \times n}$ and $j \in [n]$. If there is no confusion, let us write \widetilde{f}_j , ε_j , and φ_j instead of \widetilde{f}_j^t , ε_j^t , and φ_j^t for simplicity. In the remaining of this section, we will focus on the proof of

(6.3)
$$\widetilde{f}_{j}\Psi(A) = \Psi(\widetilde{f}_{j}A).$$

Let

(6.4)
$$\sigma = (\cdots, \underbrace{-\cdots}_{a_{i-1j+1}}, \underbrace{+\cdots+}_{a_{i-1j}}, \underbrace{-\cdots}_{a_{ij+1}}, \underbrace{+\cdots+}_{a_{ij}}, \cdots),$$
$$\sigma^* = (\cdots, \underbrace{-\cdots}_{a_{i-1j+1}^*}, \underbrace{+\cdots+}_{a_{i-1j}^*}, \underbrace{-\cdots}_{a_{ij+1}^*}, \underbrace{+\cdots+}_{a_{ij}^*}, \cdots) \cdot (\underbrace{+}_{a_{\infty,j}^*}, \underbrace{-}_{a_{\infty j+1}^*})$$

where σ^* is a concatenation of two sequences. By tensor product rule (4.1), we see that (6.5)

$$\widetilde{f}_{j}\left(A^{\flat}\otimes\mathbf{s}\right) = \begin{cases} A^{\flat}\otimes\widetilde{f}_{j}\mathbf{s} & \text{if the leftmost} + \text{in } \widetilde{\sigma^{*}} \text{ corresponds to } (\infty,j), \\ \left(\widetilde{f}_{j}A^{\flat}\right)\otimes\mathbf{s} & \text{if the leftmost} + \text{in } \widetilde{\sigma^{*}} \text{ corresponds to } (i,j) \text{ for some } i < \infty, \\ \mathbf{0} & \text{if } \widetilde{\sigma^{*}} \text{ has no } +. \end{cases}$$

In terms of A^* , this can be simplified as

(6.6)
$$\widetilde{f}_{j}A^{*} = \begin{cases} A^{*} - \widehat{E}_{ij} + \widehat{E}_{ij+1} & \text{if the leftmost} + \text{in } \widetilde{\sigma^{*}} \text{ corresponds to } (i, j), \\ \mathbf{0} & \text{if } \widetilde{\sigma^{*}} \text{ has no } +, \end{cases}$$

where we assume that $\hat{E}_{\infty j} = \sum_{k \in \mathbb{Z}} E_{\infty j+kn}$.

Lemma 6.2. We have $\tilde{f}_j A \neq \mathbf{0}$ if and only if $\tilde{f}_j \Psi(A) \neq \mathbf{0}$.

Proof. We may assume that there exists a non-zero cell in the *j*-th column. Otherwise, we have $\tilde{f}_j A = \tilde{f}_j \Psi(A) = \mathbf{0}$.

Let $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ be the set of zig-zags associated to d_A^{sw} . Let k_0 (resp. k_1) be the minimal (resp. maximal) value of d_A^{sw} in the *j*-th column. For $k_0 \leq k \leq k_1$, let i_k be the minimal row index with $(i_k, j) \in \mathbf{z}_k$. Note that $(i_k, j+1) \in \mathbf{z}_k$.

Put

$$\sigma_k = (\underbrace{+\cdots+}_{a_{i_k j}}, \underbrace{-\cdots-}_{a_{i_k+1 j+1}}, \cdots, \underbrace{+\cdots+}_{a_{i_{k+1}-1 j}}, \underbrace{-\cdots-}_{a_{i_{k+1} j+1}}),$$

$$\sigma_k^* = (\underbrace{+\cdots+}_{a_{i_k j}^*}, \underbrace{-\cdots-}_{a_{i_{k+1} j+1}^*}, \cdots, \underbrace{+\cdots+}_{a_{i_{k+1}-1 j}^*}, \underbrace{-\cdots-}_{a_{i_{k+1} j+1}^*}),$$

for $k_0 \leq k < k_1$, and

$$\sigma_{-\infty} = (\cdots, \underbrace{-\cdots}_{a_{i_{k_0}-1j+1}}, \underbrace{-\cdots}_{a_{i_{k_0}j+1}}), \quad \sigma_{\infty} = (\underbrace{+\cdots}_{a_{i_{k_1}j}}, \underbrace{-\cdots}_{a_{i_{k_1}+1j+1}}, \cdots),$$

$$\sigma_{-\infty}^* = (\cdots, \underbrace{-\cdots}_{a_{i_{k_0}-1j+1}}, \underbrace{-\cdots}_{a_{i_{k_0}j+1}}), \quad \sigma_{\infty}^* = (\underbrace{+\cdots}_{a_{i_{k_1}j}}, \underbrace{-\cdots}_{a_{i_{k_1}+1j+1}}, \cdots) \cdot (\underbrace{+}_{a_{\infty}^*_{j}}, \underbrace{-}_{a_{\infty}^*_{j+1}}).$$

Then σ and σ^* in (6.4) decompose as follows:

$$\sigma = \sigma_{-\infty} \cdot \sigma_{k_0} \cdot \cdots \cdot \sigma_{k_1-1} \cdot \sigma_{\infty},$$

$$\sigma^* = \sigma^*_{-\infty} \cdot \sigma^*_{k_0} \cdot \cdots \cdot \sigma^*_{k_1-1} \cdot \sigma^*_{\infty}.$$

Suppose that k is given with $k_0 \leq k < k_1$. Let u be the maximal row index with $(u, j) \in \mathbf{z}_k$, and let v be the minimal row index with $(v, j + 1) \in \mathbf{z}_{k+1}$. Note that $i_k \leq u$ and

 $v \leq i_{k+1}$. Suppose first that u < v. Then we have

$$\sigma_k = (\underbrace{+\cdots+}_{a_{i_k j}}, +\cdots+, \underbrace{+\cdots+}_{a_{u j}}, \underbrace{-\cdots-}_{a_{v j+1}}, -\cdots-, \underbrace{-\cdots-}_{a_{i_{k+1} j+1}}),$$

$$\sigma_k^* = (\underbrace{+\cdots+}_{a_{i_k j}-1}, +\cdots+, \underbrace{+\cdots+}_{a_{u j}+1}, \underbrace{-\cdots-}_{a_{v j+1}-1}, -\cdots-, \underbrace{-\cdots-}_{a_{i_{k+1} j+1}+1}) = \sigma_k,$$

and hence $\widetilde{\sigma_k} = \widetilde{\sigma_k^*}$. Next, suppose that $u \ge v$. In this case, we have $i_k < v \le u < i_{k+1}$. Therefore, (i_k, j) , (v, j+1) are inner corners, and (u, j), $(i_{k+1}, j+1)$ are outer corners. Then we have

$$\sigma_{k} = (\underbrace{+\cdots+}_{a_{i_{k}j}}, +\cdots+, \underbrace{-\cdots-}_{a_{v\,j+1}}, \cdots, \underbrace{+\cdots+}_{a_{u,j}}, -\cdots-, \underbrace{-\cdots-}_{a_{i_{k+1}j+1}}),$$

$$\sigma_{k}^{*} = (\underbrace{+\cdots+}_{a_{i_{k}j}-1}, +\cdots+, \underbrace{-\cdots-}_{a_{v\,j+1}-1}, \cdots, \underbrace{+\cdots+}_{a_{u,j}+1}, -\cdots-, \underbrace{-\cdots-}_{a_{i_{k+1}j+1}+1}).$$

Since one cancelling pair (+, -) of σ_k in

$$(\underbrace{+\cdots+}_{a_{i_k j}}, +\cdots+, \underbrace{-\cdots-}_{a_{v j+1}})$$

is moved to a pair (+,-) of σ_k^* in

$$(\underbrace{+\cdots+}_{a_{u\,j}+1}, -\cdots-, \underbrace{-\cdots-}_{a_{i_{k+1}\,j+1}+1}),$$

we conclude that $\widetilde{\sigma_k} = \widetilde{\sigma_k^*}$. By similar argument, we see that $(\sigma_{\pm \infty})^{\sim} = (\sigma_{\pm \infty}^*)^{\sim}$. Hence we have

(6.7)
$$\widetilde{\sigma_k} = \widetilde{\sigma_k^*} \quad (-\infty \leqslant k \leqslant \infty).$$

Since reducing a sequence does not depend on the order of cancelling (+, -), we have

$$\widetilde{\sigma} = (\sigma_{-\infty} \cdots \sigma_{\infty})^{\sim} = ((\sigma_{-\infty})^{\sim} \cdots (\sigma_{\infty})^{\sim})^{\sim}$$
$$= ((\sigma_{-\infty}^{*})^{\sim} \cdots (\sigma_{\infty}^{*})^{\sim})^{\sim} = (\sigma_{-\infty}^{*} \cdots \sigma_{\infty}^{*})^{\sim} = \widetilde{\sigma^{*}}.$$

This shows that $\varphi_j(A) = \varphi_j(\Psi(A))$ and $\varepsilon_j(A) = \varepsilon_j(\Psi(A))$, and hence the lemma follows. \Box

From now on, we assume that $\tilde{f}_j A \neq \mathbf{0}$ and $\tilde{f}_j \Psi(A) \neq \mathbf{0}$. We also assume the following notations:

- u: the row index corresponding to the leftmost + in σ,
 u*: the row index corresponding to the leftmost + in σ*,
- $s = d_A^{sw}(u, j),$ $\widetilde{A} = \widetilde{F} A$

•
$$A = f_j A = (\widetilde{a}_{ij})_{i,j \in \mathbb{Z}^*}.$$

We have

(6.8)
$$\widetilde{A} = \widetilde{f}_j A = A - \widehat{E}_{uj} + \widehat{E}_{uj+1}, \quad \widetilde{f}_j A^* = A^* - \widehat{E}_{u^*j} + \widehat{E}_{u^*j+1}.$$

Note that the leftmost + in $\widetilde{\sigma^*}$ also appears in $(\sigma_s^*)^\sim$ by (6.7). More explicitly, u^* is the minimal row index such that $u \leq u^*$ and $a_{u^*i}^* \neq 0$. We have

$$\begin{cases} u < u^* & \text{if } (u, j) \text{ is an inner corner of } \mathbf{z}_s \text{ with } a_{uj} = 1, \\ u = u^* & \text{otherwise.} \end{cases}$$

6.3. Southwest channel numberings on A and \tilde{A} . In this subsection, we discuss the relation between the southwest channel numberings on A and \tilde{A} .

For
$$(x, y) \in \mathbb{Z}^2$$
, let $(x, y)^{\wedge} = \{ \tau^k(x, y) \mid k \in \mathbb{Z} \}$. We have

$$\operatorname{supp}(A) - \operatorname{supp}(A) \subseteq (u, j+1)^{\wedge}, \quad \operatorname{supp}(A) - \operatorname{supp}(A) \subseteq (u, j)^{\wedge},$$

where the equalities hold when $a_{uj+1} = 0$ and $a_{uj} = 1$ respectively.

Lemma 6.3. We have the following.

(1) Let d be a proper numbering on A with the associated zig-zags $Z = \{\mathbf{z}_k\}_{k \in \mathbb{Z}}$ and d(u, j) = s. Then there exists a proper numbering d^- on \widetilde{A} satisfying

$$d^{-}(c) = d(c) \quad \text{if } c \in \text{supp}(A) \cap \text{supp}(A),$$
$$d^{-}(u, j+1) = \begin{cases} d(u, j) & \text{if } u \text{ is minimal such that } (u, j) \in \mathbf{z}_s, \\ d(u, j) + 1 & \text{otherwise.} \end{cases}$$

(2) Let d be a proper numbering on \widetilde{A} with the associated zig-zags $Z = \{\mathbf{z}_k\}_{k \in \mathbb{Z}}$ and d(u, j + 1) = t. Then there exists a proper numbering d^+ on A satisfying

$$d^{+}(c) = d(c) \quad \text{if } c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A}),$$
$$d^{+}(u,j) = \begin{cases} d(u,j) - 1 & \text{if } (u,j) \in \mathbf{z}_{t-1}, \\ d(u,j) & \text{otherwise.} \end{cases}$$

In particular, the widths of A and \widetilde{A} are the same.

Proof. (1) We construct a set of zig-zags Z^- (by adjusting Z) which satisfy the conditions (z.1)-(z.3) in Section 3.2 with respect to \widetilde{A} and hence gives a proper numbering d^- on \widetilde{A} .

Let i_s be the minimal row index with $(i_s, j) \in \mathbf{z}_s$. If $i_s = u$, then $(u, j+1) \in \mathbf{z}_s$ by definition of zig-zag. Suppose that $i_s < u$. Then (i_s, j) is an inner corner of \mathbf{z}_s , and $a_{i_s j} > 0$. Consider a subsequence of σ in (6.4)

(6.9)
$$(\underbrace{+\cdots+}_{a_{i_sj}}, \underbrace{-\cdots-}_{a_{i_s+1j+1}}, \cdots, \underbrace{+\cdots+}_{a_{u-1j}}, \underbrace{-\cdots-}_{a_{u_jj+1}}).$$

Since (u, j) is the cell corresponding to the leftmost + in $\tilde{\sigma}$, there exists no + in the reduced form of (6.9). This implies that there exists some $a_{vj+1} > 0$ for some v with $i_s < v \leq u$ so that + in the cell (i_s, j) is paired with - in (v, j+1). It is easy to see that d(v, j+1) = s+1, and hence $(u, j+1) \in \mathbf{z}_{s+1}$.

Hence we see that $\operatorname{supp}(\widetilde{A}) \subseteq \bigsqcup_{k \in \mathbb{Z}} \mathbf{z}_k$ and Z satisfies the conditions in (z.2) and (z.3) for \widetilde{A} . Note that the condition (z.1) fails if and only if (u, j) is an inner corner of \mathbf{z}_s with $a_{uj} = 1$.

$$d^{-}(u, j+1) = \begin{cases} s & \text{if } u = i_s, \\ s+1 & \text{if } i_s < u, \end{cases}$$

and $d^{-}(c) = d(c)$ for $c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A})$.

Case 2. Suppose that (u, j) is an inner corner of \mathbf{z}_s with $a_{uj} = 1$. In this case, the condition (z.1) fails since the inner corner (u, j) of \mathbf{z}_s does not lie in $\operatorname{supp}(\widetilde{A})$. Now let us modify \mathbf{z}_s as follows: Let (v, j) be an outer corner of \mathbf{z}_s and let

$$w = \min\{i \in \mathbb{Z} \mid u < i \leq v \text{ and } a_{ij} > 0\}.$$

Note that $w = u^*$ when $d = d_A^{sw}$. Consider a subsequence of σ

(6.10)
$$\left(\underbrace{+}_{a_{uj}}, \underbrace{-\cdots-}_{a_{u+1\,j+1}}, \cdots \underbrace{+\cdots+}_{a_{w-1\,j}}, \underbrace{-\cdots-}_{a_{w\,j+1}}\right)$$

We see that $a_{ij} = 0$ for $u < i \le w - 1$ by definition of w, and moreover $a_{ij+1} = 0$ for $u < i \le w$ since the reduced form of (6.10) is (+). Indeed the sequence in (6.10) is (+). According to this observation, we define a zig-zag \mathbf{z}_s^- by replacing the cells

$$(w-1, j), \cdots, (u+1, j), (u, j)$$

in \mathbf{z}_s with the following cells

$$(w, j + 1), (w - 1, j + 1), \dots, (u + 1, j + 1).$$

Then each inner corner of \mathbf{z}_s^- lies in $\operatorname{supp}(\widetilde{A})$.

Let Z^- be the set of zig-zags obtained from Z by replacing $\{\tau^k \mathbf{z}_s \mid k \in \mathbb{Z}\}$ with $\{\tau^k \mathbf{z}_s^- \mid k \in \mathbb{Z}\}$. \mathbb{Z} . Then Z^- satisfies the conditions (z.1)-(z.3) for \widetilde{A} , and hence induces a proper numbering d^- on \widetilde{A} . It is easy to see that $d^-(u, j + 1) = s$ since $u = i_s$, and that $d^-(c) = d(c)$ for $c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A})$.

By definition, the proper numberings d and d^- have the same flow, which implies that A and \widetilde{A} have the same width. This proves (1).

(2) As in (1), we construct a set of zig-zags from Z to which a proper numbering d^+ on A is associated.

Case 1. Suppose first that $(u, j) \in \mathbf{z}_{t-1}$. Then there exists an inner corner (v, j) of \mathbf{z}_{t-1} and an inner corner (w, j + 1) of \mathbf{z}_t with $v < w \leq u$. In particular, we have $\tilde{a}_{vj} > 0$. Consider a subsequence of σ

$$(\underbrace{+\cdots+}_{\widetilde{a}_{vi}},\underbrace{-\cdots-}_{\widetilde{a}_{v-1i+1}},\cdots,\underbrace{+\cdots+}_{\widetilde{a}_{u-1i}},\underbrace{-\cdots-}_{\widetilde{a}_{ui+1}}).$$

Since $\tilde{a}_{vj} > 0$ and (u, j + 1) is the cell corresponding to the rightmost -, we have

$$\widetilde{a}_{wj+1} + \widetilde{a}_{w+1\,j+1} + \dots + \widetilde{a}_{uj+1} \ge 2.$$

This shows that w < u and the inner corner (w, j + 1) of \mathbf{z}_t lies in $\operatorname{supp}(A)$. Hence $Z^+ := Z$ satisfies the conditions (z.1)-(z.3), and it induces a proper numbering d^+ on A. It is obvious that $d^+(u, j) = t - 1$ and $d^+(c) = d(c)$ for $c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A})$.

Case 2. Suppose that $(u, j) \notin \mathbf{z}_{t-1}$. Let i_t be the maximal row index with $(i_t, j+1) \in \mathbf{z}_t$. Since $(u, j) \notin \mathbf{z}_{t-1}$, we have $\tilde{a}_{ij} = 0$ for $u \leq i < i_t$. Consider a subsequence of σ

(6.11)
$$(\underbrace{-\cdots-}_{\widetilde{a}_{u\,j+1}}, \underbrace{+\cdots+}_{\widetilde{a}_{uj}}, \cdots, \underbrace{-\cdots-}_{\widetilde{a}_{i_tj+1}}) = (\underbrace{-\cdots-}_{\widetilde{a}_{u\,j+1}}, \cdots, \underbrace{-\cdots-}_{\widetilde{a}_{i_t\,j+1}}).$$

Since the rightmost - in $\tilde{\sigma}$ corresponding to position (u, j + 1) is the one in (6.11), we see that $\tilde{a}_{ij+1} = 0$ for $u < i \leq i_t$. We define \mathbf{z}_t^+ to be a zig-zag by replacing the cells

$$(i_t, j+1), (i_t-1, j+1), \cdots, (u+1, j+1)$$

in \mathbf{z}_t with the following cells

$$(i_t - 1, j), \cdots, (u - 1, j), (u, j).$$

Then (u, j) is an inner corner of \mathbf{z}_t^+ , and $(u, j) \in \operatorname{supp}(A)$. Let Z^+ be the set of zig-zags obtained from Z by replacing $\{\tau^k \mathbf{z}_t \mid k \in \mathbb{Z}\}$ with $\{\tau^k \mathbf{z}_t^+ \mid k \in \mathbb{Z}\}$. Then Z^+ satisfies the conditions (z.1)-(z.3) for A, and it induce a proper numbering d^+ on A. We have $d^+(u, j) = t$ and $d^+(c) = d(c)$ for $c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A})$. This proves (2).

Remark 6.4. Let d_1 be a proper numbering on A. If follows from the construction of d^{\pm} in the proof of Lemma 6.3 that

- (1) $(d_1^-)^+ = d_1,$
- (2) if d_2 is another proper numbering on A such that $d_1(c) \leq d_2(c)$ for $c \in \text{supp}(A)$, then $d_1^-(c) \leq d_2^-(c)$ for $c \in \text{supp}(\widetilde{A})$.

The similar properties hold for a proper numbering on \tilde{A} .

Lemma 6.5. We have the following.

(1) Let C be a channel of A. Then there exists a channel C^- of \widetilde{A} given by

$$\begin{cases} \left(C - (u, j)^{\wedge}\right) \cup (u, j+1)^{\wedge} & \text{if } (u, j) \in C, \ a_{uj} = 1 \text{ and } (i, j+1) \notin C \text{ for all } i, \\ \left(C - (u, j)^{\wedge}\right) \cup (v, j)^{\wedge} & \text{if } (u, j) \in C, \ a_{uj} = 1 \text{ and } (i, j+1) \in C \text{ for some } i, \\ C & \text{otherwise,} \end{cases}$$

where v is the minimal row index such that u < v and $a_{vj} > 0$ if it exists. (2) Let C be a channel of \widetilde{A} . Then there exists a channel C^+ of A given by

$$\begin{cases} \left(C - (u, j+1)^{\wedge}\right) \cup (u, j)^{\wedge} & \text{if } (u, j+1) \in C, \ a_{u\,j+1} = 1 \ and \ (i, j) \notin C \ for \ all \ i, \\ \left(C - (u, j+1)^{\wedge}\right) \cup (w, j+1)^{\wedge} & \text{if } (u, j+1) \in C, \ a_{u\,j+1} = 1 \ and \ (i, j) \in C \ for \ some \ i, \\ C & \text{otherwise}, \end{cases}$$

where w is the maximal row index such that w < u and $a_{w j+1} > 0$ if it exists.

Proof. Let us prove (1) only, since the proof of (2) is similar.

Case 1. Suppose $(u, j) \notin C$ or $a_{uj} > 1$. Then $C \subseteq \operatorname{supp}(\widetilde{A})$ and C is a channel of \widetilde{A} . We put $C^- = C$ in this case.

Now suppose that $(u, j) \in C$ and $a_{uj} = 1$. Then we have $C \not\subseteq \operatorname{supp}(\widetilde{A})$. Let us write

$$C = \{ \dots >_{\text{NW}} c_{s-1} >_{\text{NW}} c_s >_{\text{NW}} c_{s+1} >_{\text{NW}} \dots \}$$

with $c_s = (u, j)$.

Case 2. Suppose that $(i, j) \notin C$ for all i. Let C^- be a set obtained from C by replacing $c_s^{\wedge} = (u, j)^{\wedge} \subset C$ with $(u, j + 1)^{\wedge}$, which is clearly a stream of \widetilde{A} by assumption.

Case 3. Suppose that $(i, j + 1) \in C$ for some *i*. Then we have $c_{s+1} = (u', j + 1)$ for some u' > u. Consider a subsequence of σ in (6.4)

(6.12)
$$(\underbrace{+}_{a_{uj}}, \underbrace{-\cdots-}_{a_{u+1\,j+1}}, \cdots, \underbrace{+\cdots+}_{a_{u'-1j}}, \underbrace{-\cdots-}_{a_{u'\,j+1}}).$$

Since - in c_{s+1} is paired with + in (6.12) other than + in $c_s = (u, j)$, we have $a_{ij} > 0$ for some u < i < u'. Let v be the minimal such one. Then we have

(6.13)
$$c_{s-1} >_{\text{NW}} (v, j+1) >_{\text{NW}} c_{s+1}.$$

Let C^- be a set obtained from C by replacing $c_s^{\wedge} = (u, j)^{\wedge} \subset C$ with $(v, j + 1)^{\wedge}$, which is a stream of \widetilde{A} by (6.13).

By definition, C^- has the same flow as C. Since A and \widetilde{A} have the same width by Lemma 6.3, C^- is a stream of maximal flow, and hence a channel of \widetilde{A} (cf. Remark 2.18).

Remark 6.6. Let C be channel of A. Under the above hypothesis, we have

$$(C^{-})^{+} - C = \begin{cases} (v, j) & \text{if } (u, j) \in C, \ a_{uj} = 1 \text{ and } (i, j+1) \in C \text{ for some } i, \\ (u, j+1) & \text{if } (u, j) \in C, \ a_{uj} = 1, \ a_{u\,j+1} > 0 \text{ and } (i, j+1) \notin C \text{ for all } i, \\ \emptyset & \text{otherwise.} \end{cases}$$

Note that there exists no cell in supp(A) between (u, j) and (v, j) and between (u, j) and (u, j + 1). Hence it follows that

$$C \geq_{\mathsf{sw}} (C^-)^+$$
 or $(C^-)^+ \geq_{\mathsf{sw}} C$,

and there exists no other channel between C and $(C^{-})^{+}$. It is also easy to check that if C' is another channel of A with $C \geq_{sw} C'$, then we have

$$(6.14) C^- \geqslant_{\mathsf{sw}} (C')^-.$$

The similar properties also hold with respect to channels of \widetilde{A} .

Lemma 6.7. We have the following.

- Let C be a channel of A. If d is the channel numbering on A associated to C, then d⁻ is the channel numbering on A associated to C⁻.
- (2) Let C be a channel of A. If d is the channel numbering on A associated to C, then d⁺ is the channel numbering on A associated to C⁺.

Proof. Let us prove (1) only since the proof of (2) is similar.

Let d' be the channel numbering on \widetilde{A} associated to C^- . Since the widths of A and \widetilde{A} coincide, we may assume that d^- coincides with d' on the channel C^- by adding a constant to d^- . Hence we have $d' \leq d^-$ by Lemma 2.22, and $(d')^+ \leq (d^-)^+ = d$ by Remark 6.4.

Let ℓ be the common width of A and \widetilde{A} . If $\ell > 1$, then we see from Lemma 6.5(1) that there exists $c \in C \cap C^-$ such that $c \in \operatorname{supp}(A) \cap \operatorname{supp}(\widetilde{A})$. We have

$$d(c) = d^{-}(c) = d'(c) = (d')^{+}(c).$$

So $(d')^+$ and d also coincide on C (cf. Remark 6.6), and $d \leq (d')^+$ by Lemma 2.22. Therefore, we have $d = (d')^+$, and $d^- = d'$ by Remark 6.4.

If $\ell = 1$, then it is not possible to have $d^-(u, j+1) = d(u, j) + 1$ or $C^- = (C - (u, j)^{\wedge}) \cup (v, j)^{\wedge}$ since we must have another cell (i, j+1) with $(u, j) >_{NW} (i, j+1) >_{NW} (u, j) + (m, n)$. Hence we see directly that $(d')^+(u, j) = d(u, j) = d^-(u, j+1) = d'(u, j+1)$. By similar arguments as in the above case, we conclude that $d^- = d'$.

Now we can describe the southwest channel numbering on \widetilde{A} in terms of the one on A.

Proposition 6.8. Let d be the southwest channel numbering on A. Then d^- is the southwest channel numbering on \widetilde{A} . Equivalently, let d be the southwest channel numbering on \widetilde{A} . Then d^+ is the southwest channel numbering on A.

Proof. Let $C_1 = C_A^{sw}$ and $C_2 = C_{\widetilde{A}}^{sw}$. Let d' be the southwest channel numbering on \widetilde{A} . By Lemma 6.7, we have

$$d^- = d_{\widetilde{A}}^{C_1^-}, \quad (d')^+ = d_A^{C_2^+}.$$

Thus it suffices to show that either $C_1 = C_2^+$ or $C_1^- = C_2$, which implies that $d = (d')^+$ or $d^- = d'$, respectively (see also Remark 6.4).

Since C_1 and C_2 are the southwest channels, we have

$$(6.15) C_1 \geqslant_{\mathsf{sw}} C_2^+, \quad C_2 \geqslant_{\mathsf{sw}} C_1^-.$$

By Remark 6.6 (cf. (6.14)), we get from (6.15)

$$(6.16) C_1 \geq_{\mathsf{sw}} C_2^+ \geq_{\mathsf{sw}} (C_1^-)^+, \quad C_2 \geq_{\mathsf{sw}} C_1^- \geq_{\mathsf{sw}} (C_2^+)^-$$

We claim that $C_1 = C_2^+$ if $C_2 >_{sw} C_1^-$. By (6.16), we have $C_2 >_{sw} (C_2^+)^-$. By Remark 6.6, we see that $C_2 >_{sw} (C_2^+)^-$ occurs only when

$$C_2^+ = (C_2 - (u, j+1)^{\wedge}) \cup (w, j+1)^{\wedge}.$$

Then we have $C_2^+ = (C_2^+)^-$.

On the other hand, we have $C_1^- = (C_2^+)^-$ since there is no other channel between C_2 and $(C_2^+)^-$. Hence we get $C_1^- = (C_2^+)^- = C_2^+$, and in particular $(w, j + 1) \in C_1^-$. Since we have $(u, j) \notin C_1$ by Lemma 6.5(1), it follows that $C_1 = C_1^- = C_2^+$. This proves the claim. \Box

6.4. **Proof of** (6.3). Now we are in a position to prove (6.3). Let $d = d_A^{sw}$ and let $Z = \{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ be the set of zig-zags associated to d. Let Z^- be the set of zig-zags associated to d^- (see the proof of Lemma 6.3(1)). Note that $d^- = d_{\tilde{A}}^{sw}$ by Proposition 6.8.

Case 1. Suppose that (u, j) is not an inner corner of \mathbf{z}_s or (u, j) is an inner corner of \mathbf{z}_s with $a_{uj} > 1$. Since $Z = Z^-$ and the cells corresponding the leftmost + in A and A* coincide in this case, we have $\widetilde{A^*} = (\widetilde{A})^*$.

Case 2. Suppose that (u, j) is an inner corner of \mathbf{z}_s with $a_{uj} = 1$.

Let us first compare \mathbf{z}_s with the modified zig-zag \mathbf{z}_s^- . Let (v_0, j) be an outer corner of \mathbf{z}_s and let v_1 be the minimal row index with $(v_1, j+1) \in \mathbf{z}_s$. Note that the inner and outer corners of Z and Z^- always coincide other than the following cells (more precisely, their orbits under $\tau^{\pm 1}$):

$$(6.17) (u,j), (u^*,j), (v_0,j), (v_1,j+1), (u,j+1), (u^*,j+1),$$

where $v_1 \leq u < u^* \leq v_0 \leq \infty$. For the reader's convenience, we summarize the positions of the cells in (6.17) as follows:

	\mathbf{Z}_{S}	\mathbf{z}_s^-
(u,j)	inner corner	
(u^*, j)	outer corner if $u^* = v_0$	inner corner if $u^* < v_0$
(v_0, j)	outer corner	outer corner if $u^* < v_0$
$(v_1, j+1)$	inner corner if $v_1 < u$	inner corner
(u, j+1)	outer corner if $v_1 < u$	inner corner if $v_1 = u$
$(u^*, j+1)$		outer corner

Hence we may write

(6.18)
$$A^* = A - \hat{E}_{uj} + \hat{E}_{v_0j} - \hat{E}_{v_1j+1} + \hat{E}_{u\,j+1} + B,$$
$$(\widetilde{A})^* = \widetilde{A} - \hat{E}_{u*j} + \hat{E}_{v_0j} - \hat{E}_{v_1j+1} + \hat{E}_{u*\,j+1} + B,$$

where B is a finite linear combination of $\hat{E}_{k,l}$'s over the cells (k, l) not belonging to (6.17). Combining (6.8) and (6.18), we have

$$\begin{aligned} \widehat{A^*} &= A^* - \widehat{E}_{u^*j} + \widehat{E}_{u^*j+1} \\ &= \left(A - \widehat{E}_{uj} + \widehat{E}_{v_0j} - \widehat{E}_{v_1j+1} + \widehat{E}_{uj+1} + B\right) - \widehat{E}_{u^*j} + \widehat{E}_{u^*j+1} \\ &= \left(A - \widehat{E}_{uj} + \widehat{E}_{uj+1}\right) - \widehat{E}_{u^*j} + \widehat{E}_{v_0j} - \widehat{E}_{v_1j+1} + \widehat{E}_{u^*j+1} + B \\ &= \widetilde{A} - \widehat{E}_{u^*j} + \widehat{E}_{v_0j} - \widehat{E}_{v_1j+1} + \widehat{E}_{u^*j+1} + B \\ &= (\widetilde{A})^*. \end{aligned}$$

By Case 1 and Case 2, we have $\widetilde{A^*} = (\widetilde{A})^*$. Let us write $\Psi(\widetilde{f}_j A) = (\widetilde{f}_j A)^{\flat} \otimes \mathbf{s}'$. From $\widetilde{A^*} = (\widetilde{A})^*$ and Proposition 6.8, we see that

$$(\widetilde{f}_j A)^{\flat} \otimes \mathbf{s}' = \begin{cases} A^{\flat} \otimes \widetilde{f}_j \mathbf{s} & \text{if } u^* = \infty, \\ (\widetilde{f}_j A^{\flat}) \otimes \mathbf{s} & \text{if } u^* < \infty. \end{cases}$$

Comparing this with (6.5), we have (6.3).

By (6.2), κ commutes with \widetilde{f}_j , and hence κ commutes with \widetilde{e}_j for $j \in \{0, 1, \dots, n-1\}$.

6.5. **Proof of Theorem 4.14.** We have proved that κ commutes with \tilde{e}_j^t and \tilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$. Let us finish the proof of Theorem 4.14 by showing that κ commutes with \tilde{e}_i and \tilde{f}_i for $i \in \{1, \ldots, m-1\}$.

First, it is not difficult to see that Proposition 6.8 still holds if we replace the southwest channel numberings with the northeast channel numberings (cf. Remark 2.21). Hence by the same arguments as Section 6.4 we have

(6.19)
$$\widetilde{x}_i A^* = (\widetilde{x}_i A)^*$$

for $i \in \{0, 1, ..., m-1\}$ and $x \in \{e, f\}$. This implies that Ψ commutes with \tilde{e}_i and \tilde{f}_i for $i \in \{1, ..., m-1\}$ (see Remark 6.9 for i = 0). Hence κ commutes with \tilde{e}_i and \tilde{f}_i for $i \in \{1, ..., m-1\}$. This completes the proof of Theorem 4.14.

Remark 6.9. We should remark that \widetilde{e}_0 and \widetilde{f}_0 may not commute with Ψ . Let $A \in \widehat{\mathcal{M}}_{m \times n}$ be given such that $\widetilde{x}_0 A \neq \mathbf{0}$ for $x \in \{e, f\}$. Suppose that $\Psi(A) = A^{\flat} \otimes \mathbf{s}$ and $\mathbf{s} = (\mathbf{a}, \mathbf{b}, r)$.

If $\tilde{x}_0(A^{\flat} \otimes \mathbf{s}) = A^{\flat} \otimes \tilde{x}_0 \mathbf{s}$, then it follows from (6.19) that $\Psi(\tilde{x}_0 A) = A^{\flat} \otimes \mathbf{s}'$ and $\mathbf{s}' = (\tilde{x}_0 \mathbf{a}, \mathbf{b}, r')$, where r' = r+1 (resp. r-1) if x = e (resp. x = f). Since $\tilde{x}_0 \mathbf{s} = (\tilde{x}_0 \mathbf{a}, \mathbf{b}, r) \neq \mathbf{s}'$, we have $\Psi(\tilde{x}_0 A) \neq \tilde{x}_0 \Psi(A)$. If $\kappa(A) = (P_0, Q)$, then by applying Ψ repeatedly we have

$$\kappa(\widetilde{x}_0 A) = (\widetilde{x}_0 P_0, Q'),$$

for some $Q' \in \mathcal{B}_n(\lambda)$ with $Q' \neq Q$.

7. DUAL AFFINE RSK CORRESPONDENCE

In this section, we construct a dual analogue of Theorem 3.9.

7.1. Generalized dual affine permutations. Let

$$\widehat{\mathbb{N}}_{m \times n} = \left\{ A = (a_{ij})_{i,j \in \mathbb{Z}} \middle| \begin{array}{c} (1) \ a_{ij} \in \{0,1\} \text{ and } a_{i+mj+n} = a_{ij} \text{ for all } i, j \in \mathbb{Z}, \\ (2) \text{ for each } j, \ a_{ij} = 0 \text{ except for finitely many } i\text{'s.} \end{array} \right\}$$

For $A \in \widehat{\mathcal{N}}_{m \times n}$, we define a *(dual) standardization of A* to be

$$A^{\mathtt{st}'} = (A^{\bullet})^{\circ'},$$

(see Remark 2.2 for the definition of A^{\bullet}). We may adopt the same notations in Section 2.3 for $A^{st'}$. Note that a cell c in supp(A) corresponds to a unique cell in supp $(A_c^{st'})$.

Example 7.1. Let $A \in \widehat{\mathcal{N}}_{3 \times 4}$ be given as follows.



We define partial orders $>_{nW}$ and $<_{Ne}$ on \mathbb{Z}^2 as follows:

- (1) $c_1 >_{\mathbf{n} \mathbf{W}} c_2$ if and only if $i_1 \leq i_2$ and $j_1 < j_2$,
- (2) $c_1 <_{\text{Ne}} c_2$ if and only if $i_1 > i_2$ and $j_1 \leq j_2$,

for $c_1 = (i_1, j_1)$ and $c_2 = (i_2, j_2)$ in \mathbb{Z}^2 .

With respect to these partial orders, we have natural dual analogues of the notions and their properties in Section 2, the proofs of which are almost parallel to those in the case of $\widehat{\mathcal{M}}_{m \times n}$. Let us summarize them as follows: suppose that $A \in \widehat{\mathcal{N}}_{m \times n}$ is given.

• A proper numbering d on A is defined as in Definition 2.8 with respect to $>_{nW}$ instead of $>_{NW}$. Let $d^{st'}$ denote the proper numbering on $A^{st'}$ which naturally correspond to d. Lemmas 2.9 and 2.10 hold for proper numberings on A.

- A stream s and a channel C of A are defined in the same way as in Definition 2.12 with respect to $>_{nW}$. Let $s^{st'}$ and $C^{st'}$ denote the stream and channel of $A^{st'}$ which naturally correspond to s and C, respectively.
- Let \mathcal{C}'_A be the set of channels of A. We define a partial order \geq_{sw} on \mathcal{C}'_A as in (2.5) by using \leq_{Ne} instead of \leq_{ne} (see also (2.4)). Then \mathcal{C}'_A has the greatest and the smallest elements with respect to \geq_{sw} . It can be proved by the same arguments as in Proposition 2.14 using $A^{st'}$.
- Let $C_A^{\mathbf{sw}'}$ be the greatest element in \mathcal{C}'_A , which we call the southwest channel of A. We have $(C_A^{\mathbf{sw}'})^{\mathbf{st}'} = C_{A^{\mathbf{st}'}}^{\mathbf{sw}'}$.
- For $C \in \mathcal{C}'_A$, we define a channel numbering d^C_A as in (2.6) with respect to $>_{nW}$. Then d^C_A is a well-defined proper numbering on A (cf. Proposition 2.17), and

$$\left(d^C_A\right)^{\rm st'} = d^{C^{\rm st'}}_{A^{\rm st'}}.$$

• Let $d_A^{ss'} = d_A^{C_A^{ss'}}$ be the southwest channel numbering on A. Lemma 2.22 also holds.

7.2. Row semistandard tableaux and offset vectors. For $a \ge 1$, let $\mathcal{B}'_n(a) = \mathcal{B}'(a)$ be the set of $T \in SST_{\mathbb{Z}}((a))$ such that T(a) - T(1) < n, where T(j) is the *j*-th entry of *T* from the left for $1 \le j \le a$. Let $\tau_n = \tau$ be a bijection on $\mathcal{B}'(a)$ where $\tau(T)$ is the row semistandard tableau obtained from $T \in \mathcal{B}'(a)$ by replacing its entries $T(1) \le T(2) \le \cdots \le T(a)$ with $T(2) \le T(3) \le \cdots \le T(a) \le T(1) + n$.

Let $b \ge 1$. For $\alpha = (\alpha_1, \ldots, \alpha_b) \in \mathbb{Z}^b$ and $(T_1, \ldots, T_b) \in \mathcal{B}'(a) \times \cdots \times \mathcal{B}'(a)$, we define

$$\tau^{\alpha}(T_1,\ldots,T_b)=(\tau^{\alpha_1}(T_1),\ldots,\tau^{\alpha_a}(T_b)).$$

Let $R = (a^b)$. Let us regard $T \in RSST_{[n]}(R)$ as $T = (T^b, \ldots, T^1) \in \mathcal{B}'(a) \times \cdots \times \mathcal{B}'(a)$, where T^i is the *i*-th row of T from the bottom. For $1 \leq i \leq b-1$, let r_i be the minimal non-negative integer such that

$$(T^{i+1}, \tau^{r_i}(T^i))$$

is a \mathbb{Z} -semistandard tableau of shape (a^2) , and put $\eta_i = r_i + r_{i+1} + \cdots + r_{b-1}$. We call $r = (r_1, \ldots, r_{b-1})$ an offset vector and $\eta = (\eta_1, \ldots, \eta_{b-1})$ a symmetrized offset vector of T. Note that $\tau^{\eta_{\text{rev}}}(T) \in SST_{\mathbb{Z}}(R)$.

7.3. Dual affine RSK correspondence. Let d be a proper numbering on $A \in \hat{\mathbb{N}}_{m \times n}$. Note that each level set $d^{-1}(k)$ forms a chain with respect to $<_{Ne}$.

Let $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ be the set of zig-zags associated to d, where the inner corners of \mathbf{z}_k are the set of elements in $d^{-1}(k)$ maximal with respect to $>_{nW}$ (cf. Section 3.2). Then $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$ satisfies

- (z'.1) the inner corners of each \mathbf{z}_k are contained in supp(A),
- $(\mathbf{z}'.2) \operatorname{supp}(A) \subseteq \bigcup_{k \in \mathbb{Z}} \mathbf{z}_k,$
- (z'.3) \mathbf{z}_k is located to the southeast of \mathbf{z}_{k-1} for $k \in \mathbb{Z}$ in the sense of (3.1) with respect to $>_{nW}$.

Remark 7.2. We should remark that no outer cell of \mathbf{z}_k belongs to $\operatorname{supp}(A)$, and \mathbf{z}_k 's are not always mutually disjoint. More precisely, two horizontal lines (or line segments) in \mathbf{z}_k

and \mathbf{z}_l (k < l) may have non-trivial intersection, while vertical lines (or line segmenents) in \mathbf{z}_k and \mathbf{z}_l (k < l) are always disjoint.

Suppose that a non-zero $A \in \widehat{\mathcal{N}}_{m \times n}$ is given and let

- $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$: the set of zig-zags associated to $d_A^{\mathbf{sw}'}$,
- $A^{\flat'}$: the matrix obtained from A by the same rule as in A^{\flat} with respect to $\{\mathbf{z}_k\}_{k\in\mathbb{Z}}$,
- $A^{(t)}$: the matrices in $\widehat{\mathcal{N}}_{m \times n}$ defined inductively by

$$A^{(0)} = A, \quad A^{(t)} = \left(A^{(t-1)}\right)^{\flat'} \quad (t \ge 0).$$

Note that $A^{(s-1)} \neq \mathbb{O}$ and $A^{(s)} = \mathbb{O}$ for some $s \ge 1$. For $1 \le t \le s$, we let

- $\{\mathbf{z}_k^{(t)}\}_{k\in\mathbb{Z}}$: the set of zig-zags associated to $d_{A^{(t-1)}}^{\mathbf{sw}'}$,
- $\mathbf{s}^{(t)} = (\mathbf{a}_t, \mathbf{b}_t, \rho_t)$: the stream of the back-post corners of $\{\mathbf{z}_k^{(t)}\}_{k \in \mathbb{Z}}$ with flow μ_t ,

where we can check that $\mu = (\mu_1, \ldots, \mu_s) \in \mathscr{P}_s$ as in Lemma 3.3. Now let

- P_0, Q_0 : the tableau of shape $\lambda = \mu'$ defined as in Section 3.2,
- P_0^t : the tableau of shape μ obtained by flipping P_0 with respect to the main diagonal,
- $\rho = (\rho_1, \ldots, \rho_s) \in \mathbb{Z}^s.$

Lemma 7.3. Under the above hypothesis, P_0^t is row semistandard, while Q_0 is column semistandard.

Proof. It follows immediately from Remark 7.2.

Let $l = \ell(\lambda)$ and let m_i be the number of occurrences of i in λ' for $1 \leq i \leq l$. Let

- R_i : the Young diagram given in (3.8),
- $P_0^{(i)}, Q_0^{(i)}$: the subtableaux of P_0 and Q_0 corresponding to R_i , respectively,
- $\rho^{(i)} \in \mathbb{Z}^{m_i}$: the subsequence of ρ corresponding to the columns of R_i ,
- $\eta^{(i)} \in \mathscr{P}_{m_i-1}$: the symmetrized offset vector of $\left(P_0^{(i)}\right)^{\iota}$.

for $1 \leq i \leq l$ with $m_i \geq 1$, and

• Q: the tableau defined as in (3.11).

Then we have the following, which is a dual analogue of Theorem 3.9.

Theorem 7.4. We have a bijection

$$\kappa': \quad \widehat{\mathcal{N}}_{m \times n} \longrightarrow \bigsqcup_{\lambda \in \mathscr{P}_n} RSST_{[m]}(\lambda') \times \mathcal{B}_n(\lambda) ,$$
$$A \longmapsto (P_0^t, Q)$$

where $\kappa'(\mathbb{O}) = (\emptyset, \emptyset)$.

Example 7.5. Let A be the generalized permutation in Example 7.1. Then



where the red lines denote the zig-zag's associated to $d_{A^{(t-1)}}^{\mathbf{s}\mathbf{w}'}$ for $1\leqslant t\leqslant 4,$ and



In this case, R_1 and R_3 are the only non-trivial rectangles in the decomposition of the shape of P_0 and Q_0 . It is easy to see that $\eta_{\text{rev}}^{(3)} = (0,2)$, $\eta_{\text{rev}}^{(1)} = (0,0)$, and hence



7.4. **Proof of Theorem 7.4.** The proof of Theorem 7.4 is almost parallel to that of Theorem 3.9 in Section 5. So we give an outline of its proof and leave the details to the reader.

Let $\alpha = (\alpha_1, \ldots, \alpha_m) \in \mathbb{Z}_+^m$ and $\beta = (\beta_1, \ldots, \beta_n) \in \mathbb{Z}_+^n$ be given. Let

$$\widehat{\mathbb{N}}_{m \times n}(\alpha, \beta) = \{ A \mid A \in \widehat{\mathbb{N}}_{m \times n}, \text{ row}(A) = \alpha, \text{ col}(A) = \beta \},\$$

where row(A) and col(A) are given as in (5.1). For $A \in \widehat{\mathbb{N}}_{m \times n}(\alpha, \beta)$, we denote by $A^{\operatorname{perm}'}$ the extended affine permutation in \widetilde{W}_K obtained from $A^{\operatorname{st}'}$ in the same way as in (5.2), where $|\alpha| = |\beta| = K$.

Assume that K > 1. We say that $i \in [K]$ is an *ascent* of $B \in \widetilde{W}_K$ if $(i, j), (i + 1, j') \in$ supp(B) and j < j'. we say that $B \in \widetilde{W}_K$ is α -ascending if for any $k \in$ supp (α) and i with

 $\alpha_0 + \alpha_1 + \dots + \alpha_{k-1} < i < \alpha_1 + \dots + \alpha_{k-1} + \alpha_k,$

i is an ascent of *B*, where $\alpha_0 = 0$. Let $\widetilde{W}_{K,[\alpha,\beta]}$ denote the set of $B \in \widetilde{W}_K$ such that *B* is α -ascending and B^t is β -descending. As in Lemma 5.2, we have a bijection

$$\widehat{\mathbb{N}}_{m \times n}(\alpha, \beta) \longrightarrow \widetilde{W}_{K,[\alpha,\beta]} .$$

$$A \longmapsto A^{\operatorname{perm}'}$$

Let $\lambda \in \mathscr{P}$ be given with $|\lambda| = K$. Let $RSST_{[m]}(\lambda)_{\alpha}$ be the set of $T \in RSST_{[m]}(\lambda)$ with content α . For $T \in RSST_{[m]}(\lambda)_{\alpha}$, we define $T^{st'}$ to be a tableau obtained from T by replacing each $k \in [m]$ in T with $\alpha_k \neq 0$ by the consecutive numbers

$$\alpha_0 + \alpha_1 + \dots + \alpha_{k-1} + 1 < \dots < \alpha_1 + \dots + \alpha_{k-1} + \alpha_k$$

from the bottom row to the top one, and from left to right in each row, where $\alpha_0 = 0$. Then we have

$$T^{\mathsf{st}'} \in RST_{[K]}(\lambda),$$

We call $T^{st'}$ the standardization of T. Suppose that $R = (a^b) \in \mathscr{P}$ for some $a, b \ge 1$. As in Lemma 5.3, we have

for $T \in RSST_{[m]}(R)$ and $\nu \in \mathscr{P}_{a-1}$. In particular, the symmetrized offset vector of T is the same as that of $T^{st'}$.

Let $S \in RST_{[K]}(\lambda)$ be given. We say that $i \in [K]$ is a *(row) ascent* of S if the letter i + 1 does not appear below i in T. We define $S \in RST_{[K]}(\lambda)$ to be α -ascending in the same way as in the case of $CST_{[K]}(\lambda)$ with respect to ascent. Let $RST_{[K],\alpha}(\lambda)$ be the set of α -ascending tableaux in $RST_{[K]}(\lambda)$. As in Lemma 5.4, we have a bijection

$$RSST_{[m]}(\lambda)_{\alpha} \longrightarrow RST_{[K],\alpha}(\lambda)$$
$$T \longmapsto T^{st'}$$

Let $B \in \widetilde{W}_K$ be given with $\kappa_0(B) = (P_0, Q_0, \rho)$, where κ_0 is the map in (3.3). Suppose that the shape of P_0 and Q_0 is λ so that $P_0^t \in RST_{[K]}(\lambda'), Q_0 \in CST_{[K]}(\lambda)$. By Lemma 5.5, we have as in Corollary 5.6

(7.2)
$$B \in \widetilde{W}_{K,[\alpha,\beta]}$$
 if and only if $P_0^t \in RST_{[K],\alpha}(\lambda')$ and $Q_0 \in CST_{[K],\beta}(\lambda)$.

We also have the following analogue of Proposition 5.7. For $A \in \widehat{\mathcal{N}}_{m \times n}$, let $\kappa'_0(A) = (P_0, Q_0, \rho)$ be given in Section 7.3. Then

(7.3)
$$\kappa_0(A^{\texttt{perm}'}) = (P_0^{\texttt{st}'}, Q_0^{\texttt{st}}, \rho),$$

where we understand $P_0^{\mathsf{st}'}$ as $((P_0^t)^{\mathsf{st}'})^t$.

Let (P_0, Q_0, ρ) be a triple such that $(P_0^t, Q_0, \rho) \in RSST_{[m]}(\lambda) \times CSST_{[n]}(\lambda) \times \mathbb{Z}^{\lambda_1}$ for some $\lambda \in \mathscr{P}_n$. Keeping the notations in Section 7.3, we let

- $\theta^{(i)} \in \mathscr{P}_{m_i-1}$: the symmetrized offset vector of $Q_0^{(i)}$,
- $\zeta^{(i)} = \theta^{(i)} \eta^{(i)} \in \mathbb{Z}^{m_i}$ with the last component being zero,

for $1 \leq i \leq l$ with $m_i \geq 1$. We say that (P_0, Q_0, ρ) is *dominant* if $\rho_{rev} - \zeta \in \mathcal{P}(\lambda)$, where $\zeta = (\zeta^{(l)}, \ldots, \zeta^{(1)}) \in \mathbb{Z}^{m_l} \times \cdots \times \mathbb{Z}^{m_1}$. Then as in Lemma 5.12, it follows from (7.1) that

(7.4)
$$(P_0, Q_0, \rho)$$
 is dominant if and only if $(P_0^{st'}, Q_0^{st}, \rho)$ is dominant.

Furthermore, by almost the same arguments as in Section 5.2, we can check that $(P_0^{\text{st}'}, Q_0^{\text{st}}, \rho)$ is dominant if and only if ρ is dominant in the sense of [3, Definition 5.8].

Let Ω'_{dom} be the set of (P_0, Q_0, ρ) which are dominant. Then it follows from (7.2), (7.3), (7.4) and [3, Theorem 5.1, Theorem 5.11] that we have a bijection

$$\kappa'_0: \widehat{\mathbb{N}}_{m \times n} \longrightarrow \Omega'_{\mathrm{dom}} .$$

Finally, by the same arguments as in 5.16, we conclude that

$$\kappa': \quad \widehat{\mathcal{N}}_{m \times n} \longrightarrow \bigsqcup_{\lambda \in \mathscr{P}_n} RSST_{[m]}(\lambda') \times \mathcal{B}_n(\lambda) ,$$
$$A \longmapsto (P_0^t, Q)$$

is a well-defined bijection. This proves Theorem 7.4.

7.5. Affine crystal morphisms. We first remark that $\widehat{N}_{m \times n}$ has a normal $A_{m-1}^{(1)}$ -crystal structure with P^0 and P_{cl}^0 -weights with respect to wt, $\widetilde{e}_i, \widetilde{f}_i$ for $i \in \{0, 1, \ldots, m-1\}$ for $m \ge 2$, where wt = wt⁰ in (4.2) and wt_{cl}⁰ in (4.3) respectively, and $\widetilde{f}_i A$ is given as follows:

• Let

$$\sigma = (\cdots, \underbrace{+}_{a_{i\,j+1}}, \underbrace{-}_{a_{i+1\,j+1}}, \underbrace{+}_{a_{i\,j}}, \underbrace{-}_{a_{i+1\,j}}, \cdots).$$

If $\tilde{\sigma}$ has at least one +, then we define $\tilde{f}_i A = A - \hat{E}_{ij_1} + \hat{E}_{i+1j_1}$, where j_1 is the column index of A corresponding to the leftmost + in $\tilde{\sigma}$. If $\tilde{\sigma}$ has no +, then we define $\tilde{f}_i A = \mathbf{0}$.

Similarly, $\widehat{\mathcal{N}}_{m \times n}$ has a normal $A_{n-1}^{(1)}$ -crystal with P^0 and P_{cl}^0 -weights with respect to $\operatorname{wt}^t, \widetilde{e}_j^t, \widetilde{f}_j^t$ for $j \in \{0, 1, \ldots, n-1\}$ for $n \ge 2$, where $\operatorname{wt}^t(A) = \operatorname{wt}(A^t)$, and $\widetilde{f}_j^t A$ is given as follows:

• Let

$$\sigma' = (\cdots, \underbrace{+}_{a_{ij}}, \underbrace{-}_{a_{ij+1}}, \underbrace{+}_{a_{i+1j}}, \underbrace{-}_{a_{i+1j+1}}, \cdots).$$

If $\widetilde{\sigma'}$ has at least one +, then we define $\widetilde{f}_j^t A = A - \widehat{E}_{i_1 j} + \widehat{E}_{i_1 j+1}$, where i_1 is the index corresponding to the leftmost + in $\widetilde{\sigma'}$. If $\widetilde{\sigma'}$ has no +, then we define $\widetilde{f}_j^t A = \mathbf{0}$.

The following is an analogue of Proposition 4.4, which is straightforward to check.

Proposition 7.6. The operators \tilde{e}_i and \tilde{f}_i for $i \in \{0, 1, \ldots, m-1\}$ commute with \tilde{e}_j^t and \tilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$ on $\hat{\mathbb{N}}_{m \times n} \cup \{\mathbf{0}\}$.

For $r \ge 1$, the set $SST_{[m]}((r))$ has a normal $A_{m-1}^{(1)}$ -crystal structure with P_{cl}^{0} -weights with respect to $\operatorname{wt}_{cl}^{0}$, \tilde{e}_{i} , \tilde{f}_{i} for $i \in \{0, 1, \ldots, m-1\}$, where for $T \in SST_{[m]}((r))$

- (1) wt⁰_{cl}(T) = $\sum_{k=j}^{r} \operatorname{cl}(\epsilon_{T(j)}),$
- (2) $\tilde{f}_i T$ is the row semistandard tableau obtained by replacing an entry i in T with $i + 1 \pmod{n}$ if exists, and **0** otherwise.

Recall that $SST_{[m]}((r))$ is isomorphic to the crystal of the Kirillov-Reshetikhin module corresponding to $r \operatorname{cl}(\varpi_1)$ [8].

In general, for $\mu = (\mu_1, \ldots, \mu_l) \in \mathscr{P}$, we regard $RSST_{[m]}(\mu)$ as an $A_{m-1}^{(1)}$ -crystal with P_{cl}^0 by identifying $T \in RSST_{[m]}(\mu)$ with $T^{\ell} \otimes \ldots \otimes T^1 \in SST_{[m]}(\mu_1) \otimes \ldots \otimes SST_{[m]}(\mu_l)$, where T^i is the *i*-th row of T from the bottom.

Let

$$\mathbb{S}_{m \times n} = \bigsqcup_{\lambda \in \mathscr{P}_n} RSST_{[m]}(\lambda') \times \mathbb{B}_n(\lambda).$$

We assume the normal $A_{m-1}^{(1)}$ -crystal structure with P_{cl}^{0} -weights for $m \ge 2$ and $A_{n-1}^{(1)}$ -crystal structure with P^{0} -weights for $n \ge 2$ on $\mathcal{S}_{m \times n}$ in the same manner as in $\mathcal{T}_{m \times n}$.

Theorem 7.7. The bijection

$$\kappa': \widehat{\mathcal{N}}_{m \times n} \longrightarrow \mathcal{S}_{m \times n}$$

commutes with \tilde{e}_i , \tilde{f}_i for $i \in \{1, \ldots, m-1\}$ and \tilde{e}_j^t , \tilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$.

Proof. The proof is almost parallel to that of Theorem 4.14, where we need to modify the arguments in Section 6 with respect to the notions in Section 7.1. We leave the details to the reader. \Box

Let $\kappa'_1 = \pi_1 \circ \kappa'$, where π_1 is the projection of $S_{m \times n}$ along the first component. Then κ'_1 commutes with \tilde{e}_i and \tilde{f}_i for $i \in \{0, 1, \ldots, m-1\}$, and preserves $\mathrm{wt}^0_{\mathrm{cl}}$. Hence we have the following.

Corollary 7.8. A generalized dual affine permutation $A \in \widehat{\mathbb{N}}_{m \times n}$ is $A_{m-1}^{(1)}$ -crystal equivalent to P_0 , where $\kappa'(A) = (P_0, Q)$.

Moreover, if we define \mathbf{wt}^t on $\widehat{\mathbb{N}}_{m \times n}$ in the same way as in (4.5) with respect to κ' , then we have the following analogue of Corollary 4.16.

Corollary 7.9. The set $\widehat{\mathbb{N}}_{m \times n}$ is an $A_{n-1}^{(1)}$ -crystal with P^0 -weights with respect to \mathbf{wt}^t , \widetilde{e}_j^t , \widetilde{f}_j^t for $j \in \{0, 1, \ldots, n-1\}$, and κ' is an isomorphism of $A_{n-1}^{(1)}$ -crystals with P^0 -weights. In particular, a generalized dual affine permutation $A \in \widehat{\mathbb{N}}_{m \times n}$ is $A_{n-1}^{(1)}$ -crystal equivalent to Q, where $\kappa'(A) = (P_0, Q)$. Moreover, $\widehat{\mathbb{N}}_{m \times n}$ decomposes into a disjoint union of crystals of extremal weight modules as follows:

$$\widehat{\mathbb{N}}_{m \times n} \cong \bigoplus_{\lambda \in \mathscr{P}_n} \mathcal{B}_n(\lambda)^{\bigoplus m'_\lambda}$$

where $m'_{\lambda} = |RSST_{[m]}(\lambda')|.$

We remark that not both m and n need to be greater than 1 in Theorem 7.7 and its corollaries. In particular, when m = 1 we have the following multiplicity-free decomposition

$$\widehat{\mathbb{N}}_{1 \times n} \cong \bigoplus_{\lambda \in \mathscr{P}_n} \mathcal{B}_n(\lambda),$$

since $m'_{\lambda} = 1$ for all $\lambda \in \mathscr{P}_n$.

References

- J. Beck, H. Nakajima, Crystal bases and two-sided cells of quantum affine algebras, Duke Math. J. 123 (2004) 335–402.
- [2] M. Chmutov, J. Brewster Lewis, P Pylyavskyy, Monodromy in Kazhdan-Lusztig cells in affine type A, preprint (2017), arXiv:1706.00471.
- [3] M. Chmutov, P. Pylyavskyy, E. Yudovina, Matrix-ball construction of affine Robinson-Schensted correspondence Selecta Math. (N.S.) 24 (2018) 667–750.
- [4] W. Fulton, Young tableaux, with Application to Representation theory and Geometry, Cambridge Univ. Press, 1997.
- [5] J. Hong, S.-J. Kang, Introduction to Quantum Groups and Crystal Bases, Graduate Studies in Mathematics 42, Amer. Math. Soc., 2002.
- [6] R. Howe, Remarks on classical invariant theory, Trans. Amer. Math. Soc. 313 (1989) 539–570.
- [7] T. Imamura, M. Mucciconi, T. Sasamoto, Skew RSK dynamics: Greene invariants, affine crystals and applications to q-Whittaker polynomials, preprint (2021), arXiv:2106.11922.
- [8] S-J. Kang, M. Kashiwara, K. C. Misra, T. Miwa, T. Nakashima, A. Nakayashiki, Perfect crystals of quantum affine Lie algebras, Duke Math. J. 68 (1992) 499–607.
- [9] M. Kashiwara, Crystal bases of modified quantized enveloping algebra, Duke Math. J. 73 (1994) 383–413.
- [10] M. Kashiwara, On level-zero representations of quantized affine algebras, Duke Math. J. 112 (2002) 117–175.
- [11] D. Kazhdan, G. Lusztig, Representations of Coxeter groups and Hecke algebras, Invent. Math. 53 (1979) 165–184.
- [12] D. Knuth, Permutations, matrices, and the generalized Young tableaux, Pacific J. Math. 34 (1970) 709–727.
- [13] J.-H. Kwon, Crystal graphs for Lie superalgebras and Cauchy decompositions, J. Algebraic Combin. 25 (2007) 57–100.
- [14] A. Lascoux, Double crystal graphs, Studies in Memory of Issai Schur, Progress in Math. 210, Birkhäuser (2003) 95–114.
- [15] G. Lusztig, Aperiodicity of quantum affine $\mathfrak{gl}_n,$ Asian J. Math. 3 (1999) 147–178.

- [16] K. Misra, T. Miwa, Crystal base for the basic representation of $U_q(\hat{\mathfrak{sl}}(n))$, Comm. Math. Phys. 134 (1990) 79–88.
- [17] S. Naito, D. Sagaki, Path model for a level-zero extremal weight module over a quantum affine algebra II, Adv. Math. 200 (2006) 102–124.
- [18] A. Nakayashiki, Y. Yamada, Kostka polynomials and energy functions in solvable lattice models, Selecta Math. (N.S.) 3 (1997) 547–599.
- [19] B. Sagan, R. Stanley, Robinson-Schensted algorithms for skew tableaux, J. Combin. Theory, Series A, 55 (1990) 161–193.
- [20] J. Y. Shi, Kazhdan-Lusztig cells of certain affine Weyl groups, Volume 1179 of Lecture Notes in Mathematics. Springer (1986).
- [21] J. Y. Shi, The generalized Robinson-Schensted algorithm on the affine Weyl group of type A_{n-1} , J. Algebra **139** (1991) 364–394.
- [22] M. Shimozono, Crystals for Dummies, http://www.aimath.org/WWN/kostka/crysdumb.pdf, 2005.

DEPARTMENT OF MATHEMATICAL SCIENCES AND RIM, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KO-REA

Email address: jaehoonkw@snu.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA Email address: cestlavie813@snu.ac.kr