

# The SKA as a prebiotic molecule detector

Izaskun Jiménez-Serra<sup>1,\*</sup>, Jesús Martín-Pintado<sup>1</sup>, Aran Insausti<sup>2,3</sup>, Elena R. Alonso<sup>2,3</sup>, Emilio J. Cocinero<sup>2,3</sup> and Tyler L. Bourke<sup>4</sup>

<sup>1</sup>Centro de Astrobiología (CSIC/INTA), Ctra. de Torrejón a Ajalvir km 4, E-28850, Torrejón de Ardoz, Spain

<sup>2</sup>Departamento de Química Física, Facultad de Ciencia y Tecnología, Universidad del País Vasco, (UPV-EHU), 48080 Bilbao, Spain

<sup>3</sup>Biofisika Institute (CSIC, UPV/EHU), 48940 Leioa, Spain

<sup>4</sup>SKA Observatory, Jodrell Bank, Macclesfield SK11 9FT, United Kingdom

Correspondence\*:

Izaskun Jiménez-Serra

ijimenez@cab.inta-csic.es

## ABSTRACT

One of the theories for the origin of life proposes that a significant fraction of prebiotic material could have arrived to Earth from outer space between 4.1 and 3.8 billion years ago. This suggests that those prebiotic compounds could have originated in interstellar space, to be later on incorporated to small Solar-system bodies and planetesimals. The recent discovery of prebiotic molecules such as hydroxylamine and ethanolamine in the interstellar medium, strongly supports this hypothesis. However, some species such as sugars, key for the synthesis of ribonucleotides and for metabolic processes, remain to be discovered in space. The unmatched sensitivity of the Square Kilometer Array (SKA) at centimeter wavelengths will be able to detect even more complex and heavier prebiotic molecules than existing instrumentation. In this contribution, we illustrate the potential of the SKA to detect simple sugars with three and four carbon atoms, using a moderate investment of observing time.

**Keywords:** Square Kilometer Array (SKA), complex organic molecules, prebiotic chemistry, interstellar medium, astrochemistry

## 1 INTRODUCTION

The question of the origin of life has attracted the interest of researchers for centuries. Two main lines of thought have been proposed: i) Life may have emerged endogenously so that the building blocks of life could have formed "in situ" on Earth; or ii) Life could have originated somewhere else in the Universe and been transported to Earth in small Solar-system bodies. Alternatively, an intermediate theory contemplates the possibility that a fraction of the prebiotic material essential for the origin of life could have originated exogenously and been transferred to a young Earth via planetesimal impact on its surface (Anders, 1989; Chyba and Sagan, 1992). Prebiotic molecules such as amino acids, nucleobases and sugars have indeed been detected in meteorites (Cooper et al., 2001; Pizzarello et al., 2006; Glavin et al., 2020; Callahan et al., 2011; Furukawa et al., 2019) and in comets (Altwegg et al., 2016), which supports the latter hypothesis.

In the past decade, it has become clear that the interstellar medium (ISM) is an extraordinary chemical factory. About 250 molecules, including ringed-molecules (see e.g. Cernicharo et al., 2021;

McGuire et al., 2021), have so far been reported in the ISM. In addition, the pace at which new molecules are detected not only seems steady but accelerating (McGuire, 2021). In particular, the so-called complex organic molecules (or COMs)<sup>1</sup> have attracted great interest in recent years since a subset of them could have been involved in the first biochemical reactions leading to the origin of life. This sub-set of COMs are typically called *prebiotic*. Some examples of prebiotic COMs include urea and hydroxylamine (Belloche et al., 2019; Jiménez-Serra et al., 2020; Rivilla et al., 2020) since they are possible precursors of ribonucleotides (see e.g. Becker et al., 2019; Menor Salvan et al., 2020); ethanolamine and n-propanol because they could have triggered the formation of phospholipids (Rivilla et al., 2021; Jimenez-Serra et al., 2021); or amino acetonitrile, vinyl amine and ethyl amine since they are considered precursors of amino acids (Belloche et al., 2008; Zeng et al., 2021).

One of the most extended theories about the origin of life is the primordial RNA world. In this scenario, early forms of life relied solely on RNA to store genetic information and to catalyze chemical reactions. The basic units of RNA are ribonucleotides, which are composed of a phosphate group, a nitrogenous base, and a ribose sugar (a C<sub>5</sub> sugar with five carbon atoms). Interestingly, only small precursors of sugars such as glycolonitrile (HOCH<sub>2</sub>CN; see Patel et al., 2015) or the simplest C<sub>2</sub> sugar molecule, glycolaldehyde (CH<sub>2</sub>OHCHO), have been reported in the ISM (Hollis et al., 2000; Beltrán et al., 2009; Jørgensen et al., 2012; Zeng et al., 2019). Indeed, searches of C<sub>3</sub> sugars such as glyceraldehyde (CHOCHOHCH<sub>2</sub>OH) or dihydroxyacetone (DHA, with the chemical formula CH<sub>2</sub>OHCOCH<sub>2</sub>OH), have not yielded any robust detection (Hollis et al., 2004; Widicus Weaver and Blake, 2005; Apponi et al., 2006; Jiménez-Serra et al., 2020). In contrast, larger sugar-like species such as C<sub>3</sub> sugars and ribose have been found in meteorites (de Marcellus et al., 2015; Furukawa et al., 2019), which opens the possibility that these species form in interstellar space.

The search of sugars in the ISM has partly been hindered by the lack of spectroscopic rotational data since these species are thermolabile and hygroscopic. This has triggered the development of new sample preparation techniques as well as the use of ultrafast laser vaporization sources to avoid their decomposition (Cocinero et al., 2012; Calabrese et al., 2020). Recently, the rotational spectroscopy of C<sub>4</sub> and C<sub>5</sub> sugars such as erythrulose and ribose has been obtained in the laboratory (Cocinero et al., 2012; Insausti et al., 2021), which enables their search in interstellar space.

## 2 SEARCHING FOR SUGARS IN SOURCES WITH LOW EXCITATION TEMPERATURES ( $T_{EX}$ )

The first searches in the ISM of glyceraldehyde and dihydroxyacetone (DHA) targeted the massive star-forming region SgrB2 N-LMH. While an upper limit to the abundance of glyceraldehyde of  $\leq 2.4\text{--}5.7 \times 10^{-11}$  was obtained by Hollis et al. (2004), Widicus Weaver and Blake (2005) reported a tentative detection of DHA toward this source with an abundance of  $1.2 \times 10^{-9}$ . The latter detection, however, was never confirmed (Apponi et al., 2006). For low-mass star-forming regions, Jiménez-Serra et al. (2020) also reported upper limits to the abundance of both species in the range  $\leq 0.6\text{--}4 \times 10^{-10}$  toward the low-mass hot corino IRAS16293-2422 B.

Massive hot cores and low-mass hot corinos are among the most chemically rich sources in the Galaxy and, traditionally, they have been the selected targets for searches of new prebiotic COMs in the ISM (see e.g. Jørgensen et al., 2012; Belloche et al., 2014; Jørgensen et al., 2016; Belloche et al., 2019). Their disadvantage, however, lies in the fact that their observed rotational spectra present hundreds (even

<sup>1</sup> COMs are defined as carbon-based molecules with  $\geq 6$  atoms in their structure (see e.g. methanol or CH<sub>3</sub>OH; Herbst and van Dishoeck, 2009).

thousands) of different molecular rotational transitions as a result of the high excitation temperatures (of  $T_{ex}=100-300$  K). These high temperatures populate a wide number of energy levels of a COM since its partition function is large. In addition, at temperatures of 100-300 K COM rotational spectra present their peak emission shifted towards millimeter and sub-millimeter wavelengths, which are heavily contaminated by the emission from smaller and lighter species such as CO. Consequently, the COM spectra observed in massive hot cores and low-mass hot corinos not only largely suffer from line blending and line confusion but also present weak lines due to the large partition functions expected at high excitation temperatures.

Alternatively, sources where COMs show low excitation temperatures ( $T_{ex}$ ) represent better targets for the search and discovery of new large prebiotic species in the ISM (Jimenez-Serra et al., 2014)<sup>2</sup>. As a result of the low  $T_{ex}$ , the emission peak of the COM observed spectra is shifted towards lower frequencies, which are cleaner from the contribution from lighter molecules. In addition, the line intensities increase since only the lowest energy levels of the COMs can be populated at such low  $T_{ex}$  and so, the number of rotational transitions present in the measured spectra is significantly smaller than in hotter sources. The frequency span for the transitions between the lowest energy levels is also much larger than for those between the higher energy levels, which helps reducing significantly line blending and line confusion in the observed spectra (Jimenez-Serra et al., 2014).

Giant Molecular Clouds (GMCs) located in the Galactic Center such as the molecular cloud G+0.693-0.027 (hereafter G+0.693) have proven to be not only efficient chemical factories for the formation of complex organics (Requena-Torres et al., 2006, 2008; Zeng et al., 2018), but also excellent targets for the discovery of new prebiotic species (see e.g. Jiménez-Serra et al., 2020; Rivilla et al., 2020, 2021; Zeng et al., 2021). These clouds show low  $H_2$  gas densities of  $\sim 10^4$   $cm^{-3}$  and their gas and dust temperatures are decoupled (while  $T_{dust} \leq 20$  K,  $T_{gas} \sim 70-150$  K; see Rodríguez-Fernández et al., 2000; Zeng et al., 2018). The low  $H_2$  gas densities also imply that for high-dipole moment molecules (such as COMs), and despite the high gas kinetic temperatures and broad linewidths of their emission (of  $\sim 20$   $km\ s^{-1}$ ), their  $T_{ex}$  is low (between 5 and 20 K) and thus their observed spectra will be less affected by line blending and line confusion. Recent searches of prebiotic molecules toward the Galactic Center molecular cloud G+0.693 have yielded the discovery of several of these species for the first time in the ISM such as hydroxylamine ( $NH_2OH$ ; Rivilla et al., 2020), ethanolamine ( $NH_2CH_2CH_2OH$ ; Rivilla et al., 2021), thioformic acid ( $HC(O)SH$ ; Rodríguez-Almeida et al., 2021) or *n*-propanol ( $n-C_3H_7OH$ ; Jimenez-Serra et al., 2021). As a result, Galactic Center GMCs are prime targets for the search and discovery of large  $C_3$  and  $C_4$  sugars in the ISM.

### 3 THE POTENTIAL OF THE SKA TO DETECT NEW PREBIOTIC SPECIES IN THE ISM: THE CASE OF SUGARS

The Square Kilometre Array (SKA) will be the largest radio telescope in the world operating at centimeter and meter wavelengths. The Phase 1 of this observatory will consist of two radio interferometers located at two different sites: SKA1 LOW in Western Australia and SKA1 MID in South Africa (Braun et al., 2019). SKA1 LOW will have 512 stations of 256 log periodic dipole antennas each operating within the frequency range between 50 and 350 MHz, while SKA1 MID will have a total of 197 antennas (that include the 64 antennas of the MeerKAT array) operating at frequencies between 350 MHz and 15 GHz (Braun et al., 2019). Given the frequency coverage and how dramatically the strength of the rotational

<sup>2</sup> In massive hot cores and hot corinos,  $T_{ex} \sim T_{kin}$  because of the high  $H_2$  gas densities ( $\geq 10^6$   $cm^{-3}$ ) found in these star-forming regions.

transitions of high-dipole molecules drops for frequencies  $\leq 5$  GHz, SKA1 MID, and in particular its Band 5 receivers covering the frequency range between 4.8 and 15.3 GHz, are well suited for searches of prebiotic molecules in the ISM. On top of the advantages of observing at centimeter wavelengths noted in Section 2, it is important to stress that radio interferometers such as the SKA1 filter out any extended emission in the line of sight of the targeted source, significantly decreasing the linewidths of the observed molecular line emission and reducing the level of line blending and line confusion. In Section 3.1, we evaluate the feasibility of detecting large sugar-like species such as glyceraldehyde, dihydroxyacetone and erythrose in the ISM using the Band 5 receivers of SKA1 MID.

### 3.1 SKA1 simulations for C<sub>3</sub> and C<sub>4</sub> sugars in Galactic Center GMCs: glyceraldehyde, dihydroxyacetone and erythrose

The emission from COMs in the Galactic Center GMCs is known to be extended (Martín-Pintado et al., 2001; Li et al., 2017, 2020). This is particularly interesting for absorption studies in which COMs can be searched for against a bright continuum background source. This technique may allow the detection of low-abundance COMs whose emission spectra is expected to be very weak. Indeed, in the presence of a bright background continuum source, the predicted absorption line intensity ( $T_L$ ) is given by (see e.g. Martín et al., 2019):

$$T_L = (T_{ex} - T_c - T_{bg}) \times [1 - \exp(-\tau_\nu)] \quad (1)$$

where  $T_c$  is the temperature of a continuum source and  $T_{bg}$  is the temperature of the cosmic microwave background radiation ( $T_{bg}=2.73$  K). If  $T_c \gg T_{ex}$ , this will largely enhance the observed absorption line intensity over the emission for the  $T_{ex}$  expected to be close to  $T_{bg}$ .

Therefore, we simulate the case of absorption spectra of glyceraldehyde, dihydroxyacetone and erythrose (C<sub>4</sub>H<sub>8</sub>O<sub>4</sub>) using the SLIM tool within the MADCUBA software package<sup>3</sup> (see Martín et al., 2019). We focus our simulations on C<sub>3</sub> and C<sub>4</sub> sugars because, as discussed in Section 3.2, C<sub>5</sub> sugars such as ribose are not expected to be abundant enough to be detected with SKA1 in a reasonable amount of observing time. We assume the physical conditions of the G+0.693 molecular cloud with a typical  $T_{ex}$  of  $\leq 10$  K, linewidths of  $\Delta_v=20$  km s<sup>-1</sup> and extended morphology across the SKA1 beam for the COMs emission (Zeng et al., 2020). As background source, we consider the L source located in the SgrB2 N complex, which is a bright HII region found nearby the emission peak of G+0.693, and whose measured flux at 23 GHz is  $\sim 300$  mJy within a beam of  $\sim 1''$  (de Pree et al., 1995). We have assumed that the emission from the HII region is optically thin and thus, that it shows an almost flat spectral energy distribution between 23 GHz and 10 GHz. The simulations have been obtained considering a beam of  $1''$ , which will easily be reached by the SKA1 at the frequencies of the Band 5 receivers of SKA1 MID (see Braun et al., 2019). The spectroscopic data for glyceraldehyde have been extracted from the CDMS catalogue (entry 090501; Endres et al., 2016), while the data for dihydroxyacetone is available in the JPL molecular catalogue (entry 90002; Pickett et al., 1998). The spectroscopic information for erythrose can be found in Insausti et al. (2021).

As column densities, we have assumed that glyceraldehyde and dihydroxyacetone have column densities of  $10^{13}$  cm<sup>-2</sup> and  $6.8 \times 10^{12}$  cm<sup>-2</sup>, respectively. These column densities correspond to the upper limits obtained toward G+0.693 in previous spectral surveys (see Table 2 in Jiménez-Serra et al., 2020). For

<sup>3</sup> MADCUBA, or MAdrid Data CUBe Analysis, is a software developed at the Center of Astrobiology in Madrid: <https://cab.inta-csic.es/madcuba/index.html>.

**Table 1.** List of transitions of glyceraldehyde, dihydroxyacetone and erythrulose that present the deepest features in the simulated spectra

Molecule	Formula	Frequency (MHz)	Transition	log [Int(300K)]	E <sub>1</sub> (cm <sup>-1</sup> )	I <sub>depth</sub> (mJy)
Glyceraldehyde	CHOCHOHCH <sub>2</sub> OH	4955.4998	6 <sub>2,5</sub> →6 <sub>1,5</sub>	-7.7002	3.8458	-0.488
		5907.2556	5 <sub>2,4</sub> →5 <sub>1,4</sub>	-7.6715	2.7805	-0.360
		6762.4103	4 <sub>2,3</sub> →4 <sub>1,3</sub>	-7.6931	1.8888	-0.260
		4835.6480	12 <sub>3,10</sub> →12 <sub>2,10</sub>	-7.4495	14.2538	-0.223
		6240.1302	11 <sub>3,9</sub> →11 <sub>2,9</sub>	-7.3127	12.1285	-0.186
Dihydroxyacetone	CH <sub>2</sub> OHCOCH <sub>2</sub> OH	9605.9342	4 <sub>1,3</sub> →4 <sub>0,4</sub>	-7.6869	1.2584	-0.172
		10540.6391	5 <sub>1,4</sub> →5 <sub>0,5</sub>	-7.5446	1.8837	-0.166
		8898.3733	3 <sub>1,2</sub> →3 <sub>0,3</sub>	-7.8427	0.7563	-0.161
		11731.7585	6 <sub>1,5</sub> →6 <sub>0,6</sub>	-7.4104	2.6308	-0.147
		8391.9655	2 <sub>1,1</sub> →2 <sub>0,2</sub>	-8.0253	0.3786	-0.132
Erythrulose	C <sub>4</sub> H <sub>8</sub> O <sub>4</sub>	4609.6719	10 <sub>3,8</sub> →10 <sub>2,8</sub>	-8.3205	5.1592	-0.010
		5583.4870	9 <sub>3,7</sub> →9 <sub>2,7</sub>	-8.2442	4.2624	-0.0075
		5131.3265	2 <sub>2,1</sub> →2 <sub>1,1</sub>	-9.0513	0.3329	-0.0071 <sup>a</sup>
		5131.4115	2 <sub>0,2</sub> →1 <sub>0,1</sub>	-8.8061	0.0861	-0.0071 <sup>a</sup>
		6508.3240	8 <sub>3,6</sub> →8 <sub>2,6</sub>	-8.199	3.4522	-0.0061

<sup>a</sup> Intensity of the feature resulting from the blending of the two indicated transitions.

erythrulose, we consider that this C<sub>4</sub> sugar is a factor of 10 less abundant than the C<sub>3</sub> sugar glyceraldehyde, with an assumed column density of 10<sup>12</sup> cm<sup>-2</sup> for erythrulose. We note that this decrease by one order of magnitude in the column density of molecules from the same family when one carbon atom is added to the molecular structure, has been reported for alcohols (Jimenez-Serra et al., 2021) and thiols (Rodríguez-Almeida et al., 2021). The assumed spectral resolution considered in our simulations is 79.3 kHz (equivalent to a velocity resolution between ~1.6 and 5.0 km s<sup>-1</sup>) across the full Band 5 frequency coverage.

Figure 1 presents the predicted absorption spectra of glyceraldehyde, dihydroxyacetone and erythrulose obtained for the extended GMC G+0.693 against the background source L. Depending on the structure of the molecule, the transitions that show the larger absorption depths lie in the range between 4.8-7.0 GHz for glyceraldehyde and erythrulose, and 8-14 GHz for dihydroxyacetone. The maximum bandwidth that will be covered simultaneously by the Band 5 receivers of SKA1 MID in its Phase 1 is 4 GHz at frequencies between 4.6 and 8.5 GHz, and 5 GHz for frequencies between 8.3 and 15.3 GHz. Given that the deepest absorption lines appear clustered within frequency ranges of ≤4-5 GHz-width, future detection experiments of these lines with SKA1 MID could be carried out in just two observing runs: one to simultaneously cover all the deepest features of glyceraldehyde and erythrulose, and a second one to simultaneously cover all the transitions of dihydroxyacetone. We stress that Figure 1 includes all possible transitions of these C<sub>3</sub> and C<sub>4</sub> sugars available within the frequency range covered by the Band 5 receivers of SKA1 MID. The intensities of the deepest absorption features in our simulated spectra reach values of -0.47 mJy for glyceraldehyde, of -0.17 mJy for dihydroxyacetone, and -0.01 mJy for erythrulose. Table 1 lists the spectroscopic information of the deepest features found in our predicted spectra for each molecule. In Section 3.2, we evaluate the observing time required to perform these detection experiments with SKA1 MID during its Phase 1 of operations, which will have only 133 antennas equipped with Band 5 receivers.



### 3.2 Time estimates and Key Science Projects

As seen from Table 1, the predicted intensities for the deepest absorption features lie in the range between  $-0.19$  and  $-0.5$  mJy for glyceraldehyde,  $-0.13$  and  $-0.2$  mJy for dihydroxyacetone, and  $-0.006$  and  $-0.01$  mJy for erythrulose. According to Braun et al. (2019), a line sensitivity of  $90 \mu\text{Jy}$  will be achieved in one hour of observing time with the Band 5 receivers of SKA1 MID at frequencies of 4.6-8.5 GHz, and of  $85 \mu\text{Jy}$  at frequencies of 8.5-15.3 GHz, for a velocity resolution of  $\sim 30 \text{ km s}^{-1}$ .

Taking these numbers into account, the deepest absorption line of glyceraldehyde at 4955 MHz could be detected with a  $S/N \geq 5$  in just 8.5 hrs assuming a velocity resolution of  $3 \text{ km s}^{-1}$ . The weakest transitions of glyceraldehyde at 4835 MHz and at 6240 MHz, however, would require about 50 hrs of observing time for a similar velocity resolution and  $S/N$ . This is also the case of dihydroxyacetone, for which a total of  $\geq 100$  hrs of integration time on-source would be needed to detect its weakest transitions at 8391 MHz and 11731 MHz with a  $S/N \geq 5$  and a velocity resolution of  $3 \text{ km s}^{-1}$ . A more time-consuming experiment would have to be performed for the discovery of the deepest absorption features of erythrulose. In order to detect the erythrulose transition at 4609 MHz with a  $S/N \geq 3$ , and considering a velocity resolution of  $20 \text{ km s}^{-1}$  (i.e. the linewidth of the molecular emission in G+0.693; see Zeng et al., 2018), a total of  $\geq 1100$  hrs of observing time on-source would be required.

The time estimates to perform the detection experiments of glyceraldehyde, dihydroxyacetone and erythrulose proposed above, may seem excessive. However, we note that these long integration times are contemplated within the future Key Science Projects (KSPs) scheme that will be carried out by the SKA1. The KSPs are major surveys targeting ground-breaking transformational science in Astrophysics and Astrobiology that require considerable amount of time. Therefore, the expected integration time requests for these projects are typically of a few thousands of hours, which guarantees the detection of at least  $C_3$  sugars such as glyceraldehyde and dihydroxyacetone, if present in the ISM. As discussed in Section 4, the discovery of  $C_4$  sugars such as erythrulose may require the development of future instrumentation for SKA1 MID such as the high-frequency Band 6 receivers.

We note that, when performing deep integrations as proposed here, it is expected that the level of line confusion increases even at the low frequencies covered by the SKA1 as a consequence of weak features of low-abundance molecules becoming bright in the spectra. However, this potential problem is likely not an issue for the discovery of intermediate-sized species such as  $C_3$  and  $C_4$  sugars. As recently found by Jiménez-Serra et al. (2021) and by Rodríguez-Almeida et al. (2021), the addition of a  $-\text{CH}_2$  group to the structure of alcohols, thiols, isocyanates, and carboxylic acids implies a decrease in their abundance by at least one order of magnitude (see also Section 3.1). This decrease in the abundance would imply an increase in the required SKA1 observing time by a factor of 100 and thus, for a reasonable integration time within the context of SKA1 Key Science Projects, the only sugars for which absorption features could be detected with SKA1 are the  $C_3$  and the  $C_4$  sugars together with their lowest-energy conformers. Indeed, for the case of *n*-propanol, only the *Ga* and *Aa* conformers with relative energies of  $E=0$  K and  $E=40$  K, have been found toward G+0.693 (the rest of conformers with energies  $>50$  K do not show any detected features; see Jiménez-Serra et al., 2021). Therefore, although the conformers of  $C_3$  and  $C_4$  sugars as well as even larger sugars may be present in Galactic Center Giant Molecular Clouds such as G+0.693, these species likely do not confuse much the observed SKA1 spectrum because either their abundance is too low (as for  $C_5$  sugars such as ribose) or their low-energy levels are not populated at the low excitation temperatures measured in these clouds (as for high-energy conformers).

## 4 FUTURE EXPANSION OF THE SKA: THE BAND 6 RECEIVERS

From Section 3.2, it is clear that, while relatively small molecules such as glyceraldehyde or dihydroxyacetone could be detected with a moderate investment of SKA1 MID observing time, larger prebiotic COMs such as erythrose would still be well below the limit of what the SKA1 will be able to detect in its Phase 1. For this reason, one possible future expansion of SKA1 MID contemplates the development of higher-frequency receiver(s), the Band 6 receivers, which will increase the frequency coverage of SKA1 MID to higher frequencies from 15.3 up to 50 GHz. As reported in Memo 20-01 of SKA1 titled "SKA1 Beyond 15GHz: The Science case for Band 6" (see Conway et al., 2021), high-dipole moment molecules such as COMs present rotational transitions at frequencies  $\geq 20$  GHz that can be factors  $\geq 10$  brighter than those found at frequencies  $\leq 15$  GHz (see Section 3.3.1 in this Memo). This is due to the fact that the Einstein  $A_{ul}$  coefficients increase as  $\nu^3$  with  $\nu$  being the frequency, making them one order of magnitude higher at 36 GHz than at 12 GHz. This implies that observing times about  $\sim 100$  times shorter would be needed to detect the  $C_4$  sugar erythrose in the ISM, reducing the total observing time needed for the discovery of this species from  $\geq 1100$  hrs to a few tens of hours (or at most, to a few hundreds of hours). Therefore, if these receivers were finally included in the development program of the SKA1, they would make this observatory an unbeatable machine for the discovery of large prebiotic compounds in space.

## 5 CONCLUSIONS

In this contribution, we evaluate the feasibility of detecting small sugars in the ISM using the Band 5 receivers of SKA1 in its Phase 1. Our simulations show that Galactic Center Giant Molecular Clouds such as G+0.693 represent prime targets for future searches of these key prebiotic species in space. As shown in Section 3.1, the detection of small sugars could be achieved by carrying out broad spectroscopic surveys between 5 and 14GHz in absorption against a bright continuum background source. By taking as template the L HII region source located in the SgrB2 N molecular complex, we estimate that the detection of  $C_3$  sugars could be achieved in a few hundreds of hours. Larger  $C_4$  sugars such as erythrose would require thousands of hours of observing time. Future developments of the SKA such as the Band 6 receivers (which will increase the frequency coverage of SKA1 MID up to, at least, 24 GHz), will enable the search of these large sugars and other prebiotic COMs in just a few hundreds of hours.


## 6 FIGURES

### AUTHOR CONTRIBUTIONS

I.J.-S. has written the first version of the manuscript. I.J.-S. and J.M.-P. have produced the MADCUBA-SLIM simulations of the spectra of glyceraldehyde, dihydroxyacetone and erythrose reported in Section 3.1. All authors have contributed to the discussion of the results and have provided comments on the text.

### FUNDING

I.J.-S. and J.M.-P. acknowledge partial support from the Spanish State Research Agency (AEI) through project numbers PID2019-105552RB-C41 and MDM-2017-0737 Unidad de Excelencia *María de Maeztu*–Centro de Astrobiología (CSIC-INTA). E.J.C. thanks the support received from the MINECO (PID2020-117892RB-I00), the Basque Government (IT1162-19 and PIBA 2018-11), the UPV/EHU



**Figure 1.** Predicted absorption spectra of the  $C_3$  and  $C_4$  sugars glycerinaldehyde, dihydroxyacetone and erythrulose obtained with the SLIM tool of the MADCUBA package considering the physical conditions of the G+0.693 molecular cloud ( $T_{ex}=10$  K, linewidths of  $\Delta v=20$  km s $^{-1}$  and extended emission across the beam) and a background source similar to the L source in the SgrB2 N molecular complex (measured flux of  $\sim 300$  mJy within a beam of  $1''$  at 10 GHz). The assumed column densities for glycerinaldehyde and dihydroxyacetone are  $10^{13}$  cm $^{-2}$  and  $6.8 \times 10^{12}$  cm $^{-2}$ , respectively, while for erythrulose we assume a column density one order of magnitude lower than that of glycerinaldehyde (of  $10^{12}$  cm $^{-2}$ ; see Section 3.1).

(PPG17/10, GIU18/207), and CSIC (PIC2018, LINKA20249). Computational resources and laser facilities of the UPV/EHU (SGIker) and CESGA were used in this work.

## ACKNOWLEDGMENTS

We would like to thank an anonymous referee for his/her positive and constructive comments on an earlier version of the manuscript.

## REFERENCES

- Altwegg, K., Balsiger, H., Bar-Nun, A., Berthelier, J. J., Bieler, A., Bochslers, P., et al. (2016). Prebiotic chemicals—amino acid and phosphorus—in the coma of comet 67P/Churyumov-Gerasimenko. *Science Advances* 2, e1600285–e1600285. doi:10.1126/sciadv.1600285
- Anders, E. (1989). Pre-biotic organic matter from comets and asteroids. *Nature* 342, 255–257. doi:10.1038/342255a0



- Apponi, A. J., Halfen, D. T., Ziurys, L. M., Hollis, J. M., Remijan, A. J., and Lovas, F. J. (2006). Investigating the Limits of Chemical Complexity in Sagittarius B2(N): A Rigorous Attempt to Confirm 1,3-Dihydroxyacetone. *The Astrophysical Journal Letters* 643, L29–L32. doi:10.1086/504979
- Becker, S., Feldmann, J., Wiedemann, S., Okamura, H., Schneider, C., Iwan, K., et al. (2019). Unified prebiotically plausible synthesis of pyrimidine and purine rna ribonucleotides. *Science* 366, 76–82. doi:10.1126/science.aax2747
- Belloche, A., Garrod, R. T., Müller, H. S. P., and Menten, K. M. (2014). Detection of a branched alkyl molecule in the interstellar medium: iso-propyl cyanide. *Science* 345, 1584–1587. doi:10.1126/science.1256678
- Belloche, A., Garrod, R. T., Müller, H. S. P., Menten, K. M., Medvedev, I., Thomas, J., et al. (2019). Re-exploring Molecular Complexity with ALMA (ReMoCA): interstellar detection of urea. *Astronomy & Astrophysics* 628, A10. doi:10.1051/0004-6361/201935428
- Belloche, A., Menten, K. M., Comito, C., Müller, H. S. P., Schilke, P., Ott, J., et al. (2008). Detection of amino acetonitrile in Sgr B2(N). *Astronomy & Astrophysics* 482, 179–196. doi:10.1051/0004-6361:20079203
- Beltrán, M. T., Codella, C., Viti, S., Neri, R., and Cesaroni, R. (2009). First Detection of Glycolaldehyde Outside the Galactic Center. *The Astrophysical Journal Letters* 690, L93–L96. doi:10.1088/0004-637X/690/2/L93
- Braun, R., Bonaldi, A., Bourke, T., Keane, E., and Wagg, J. (2019). Anticipated Performance of the Square Kilometre Array – Phase 1 (SKA1). *arXiv e-prints*, arXiv:1912.12699
- Calabrese, C., Uriarte, I., Insausti, A., Vallejo-López, M., Basterretxea, F. J., Cochrane, S. A., et al. (2020). Observation of the unbiased conformers of putative dna-scaffold ribosugars. *ACS Central Science* 6, 293–303. doi:10.1021/acscentsci.9b01277. PMID: 32123748
- Callahan, M. P., Smith, K. E., Cleaves, H. J., Ruzicka, J., Stern, J. C., Glavin, D. P., et al. (2011). Carbonaceous meteorites contain a wide range of extraterrestrial nucleobases. *Proceedings of the National Academy of Science* 108, 13995–13998. doi:10.1073/pnas.1106493108
- Cernicharo, J., Agúndez, M., Cabezas, C., Tercero, B., Marcelino, N., Pardo, J. R., et al. (2021). Pure hydrocarbon cycles in TMC-1: Discovery of ethynyl cyclopropenylidene, cyclopentadiene, and indene. *Astronomy & Astrophysics* 649, L15. doi:10.1051/0004-6361/202141156
- Chyba, C. and Sagan, C. (1992). Endogenous production, exogenous delivery and impact-shock synthesis of organic molecules: an inventory for the origins of life. *Nature* 355, 125–132. doi:10.1038/355125a0
- Cocinero, E. J., Lesarri, A., Écija, P., Basterretxea, F. J., Grabow, J.-U., Fernández, J. A., et al. (2012). Ribose found in the gas phase. *Angewandte Chemie International Edition* 51, 3119–3124. doi:https://doi.org/10.1002/anie.201107973
- Conway, J., Beswick, R., Bourke, T., Coriat, M., Ferrari, C., Jiménez-Serra, I., et al. (2021). *Memo 20-01, SKA1 Beyond 15GHz: The Science case for Band 6* (Square Kilometre Array)
- Cooper, G., Kimmich, N., Belisle, W., Sarinana, J., Brabham, K., and Garrel, L. (2001). Carbonaceous meteorites as a source of sugar-related organic compounds for the early Earth. *Nature*, 879–883doi: https://doi.org/10.1038/414879a
- de Marcellus, P., Meinert, C., Myrgorodska, I., Nahon, L., Buhse, T., d’Hendecourt, L. L. S., et al. (2015). Aldehydes and sugars from evolved precometary ice analogs: Importance of ices in astrochemical and prebiotic evolution. *Proceedings of the National Academy of Science* 112, 965–970. doi:10.1073/pnas.1418602112

- de Pree, C. G., Gaume, R. A., Goss, W. M., and Claussen, M. J. (1995). The Sagittarius B2 Star-forming Region. II. High-Resolution H66 alpha Observations of Sagittarius B2 North. *The Astrophysical Journal* 451, 284. doi:10.1086/176218
- Endres, C. P., Schlemmer, S., Schilke, P., Stutzki, J., and Müller, H. S. P. (2016). The Cologne Database for Molecular Spectroscopy, CDMS, in the Virtual Atomic and Molecular Data Centre, VAMDC. *Journal of Molecular Spectroscopy* 327, 95–104. doi:10.1016/j.jms.2016.03.005
- Furukawa, Y., Chikaraishi, Y., Ohkouchi, N., Ogawa, N. O., Glavin, D. P., Dworkin, J. P., et al. (2019). Extraterrestrial ribose and other sugars in primitive meteorites. *Proceedings of the National Academy of Science* 116, 24440–24445. doi:10.1073/pnas.1907169116
- Glavin, D. P., McLain, H. L., Dworkin, J. P., Parker, E. T., Elsila, J. E., Aponte, J. C., et al. (2020). Abundant extraterrestrial amino acids in the primitive cm carbonaceous chondrite asuka 12236. *Meteoritics & Planetary Science* 55, 1979–2006. doi:https://doi.org/10.1111/maps.13560
- Herbst, E. and van Dishoeck, E. F. (2009). Complex Organic Interstellar Molecules. *Annual Review of Astronomy & Astrophysics* 47, 427–480. doi:10.1146/annurev-astro-082708-101654
- Hollis, J. M., Jewell, P. R., Lovas, F. J., and Remijan, A. (2004). Green Bank Telescope Observations of Interstellar Glycolaldehyde: Low-Temperature Sugar. *The Astrophysical Journal Letters* 613, L45–L48. doi:10.1086/424927
- Hollis, J. M., Lovas, F. J., and Jewell, P. R. (2000). Interstellar glycolaldehyde: The first sugar. *The Astrophysical Journal* 540, L107–L110. doi:10.1086/312881
- Insausti, A., Alonso, E. R., Tercero, B., Santos, J. I., Calabrese, C., Vogt, N., et al. (2021). Laboratory Observation of, Astrochemical Search for, and Structure of Elusive Erythrulose in the Interstellar Medium. *The Journal of Physical Chemistry Letters* 12, 1352–1359. doi:10.1021/acs.jpcclett.0c03050
- Jiménez-Serra, I., Martín-Pintado, J., Rivilla, V. M., Rodríguez-Almeida, L., Alonso, E. R., Zeng, S., et al. (2020). Toward the RNA-World in the Interstellar Medium—Detection of Urea and Search of 2-Amino-oxazole and Simple Sugars. *Astrobiology* 20, 1048–1066. doi:10.1089/ast.2019.2125
- Jimenez-Serra, I., Rodríguez-Almeida, L. F., Martín-Pintado, J., Rivilla, V. M., Melosso, M., Zeng, S., et al. (2021). Precursors of fatty alcohols in the ISM: Discovery of n-propanol. *Astronomy & Astrophysics*
- Jimenez-Serra, I., Testi, L., Caselli, P., and Viti, S. (2014). Detectability of Glycine in Solar-type System Precursors. *The Astrophysical Journal Letters* 787, L33. doi:10.1088/2041-8205/787/2/L33
- Jørgensen, J. K., Favre, C., Bisschop, S. E., Bourke, T. L., van Dishoeck, E. F., and Schmalzl, M. (2012). Detection of the Simplest Sugar, Glycolaldehyde, in a Solar-type Protostar with ALMA. *The Astrophysical Journal Letters* 757, L4. doi:10.1088/2041-8205/757/1/L4
- Jørgensen, J. K., van der Wiel, M. H. D., Coutens, A., Lykke, J. M., Müller, H. S. P., van Dishoeck, E. F., et al. (2016). The ALMA Protostellar Interferometric Line Survey (PILS). First results from an unbiased submillimeter wavelength line survey of the Class 0 protostellar binary IRAS 16293-2422 with ALMA. *Astronomy & Astrophysics* 595, A117. doi:10.1051/0004-6361/201628648
- Li, J., Shen, Z., Wang, J., Chen, X., Li, D., Wu, Y., et al. (2017). Widespread Presence of Glycolaldehyde and Ethylene Glycol around Sagittarius B2. *The Astrophysical Journal Letters* 849, 115. doi:10.3847/1538-4357/aa9069
- Li, J., Wang, J., Qiao, H., Quan, D., Fang, M., Du, F., et al. (2020). Mapping observations of complex organic molecules around Sagittarius B2 with the ARO 12 m telescope. *Monthly Notices of the Royal Astronomical Society* 492, 556–565. doi:10.1093/mnras/stz3337
- Martín, S., Martín-Pintado, J., Blanco-Sánchez, C., Rivilla, V. M., Rodríguez-Franco, A., and Rico-Villas, F. (2019). Spectral Line Identification and Modelling (SLIM) in the MAdrid Data CUBe Analysis

- (MADCUBA) package. Interactive software for data cube analysis. *Astronomy & Astrophysics* 631, A159. doi:10.1051/0004-6361/201936144
- Martín-Pintado, J., Rizzo, J. R., de Vicente, P., Rodríguez-Fernández, N. J., and Fuente, A. (2001). Large-Scale Grain Mantle Disruption in the Galactic Center. *The Astrophysical Journal Letters* 548, L65–L68. doi:10.1086/318937
- McGuire, B. A. (2021). 2021 Census of Interstellar, Circumstellar, Extragalactic, Protoplanetary Disk, and Exoplanetary Molecules. *arXiv e-prints*, arXiv:2109.13848
- McGuire, B. A., Loomis, R. A., Burkhardt, A. M., Lee, K. L. K., Shingledecker, C. N., Charnley, S. B., et al. (2021). Detection of two interstellar polycyclic aromatic hydrocarbons via spectral matched filtering. *Science* 371, 1265–1269. doi:10.1126/science.abb7535
- Menor Salvan, C., Bouza, M., Fialho, D. M., Burcar, B. T., Fernández, F. M., and Hud, N. V. (2020). Prebiotic origin of pre-rna building blocks in a urea “warm little pond” scenario. *ChemBioChem* 21, 3504–3510. doi:https://doi.org/10.1002/cbic.202000510
- Patel, B. H., Percivalle, C., Ritson, D. J., Duffy, C. D., and Sutherland, J. D. (2015). Common origins of RNA, protein and lipid precursors in a cyanosulfidic protometabolism. *Nature Chemistry*, 301–307doi: https://doi.org/10.1038/nchem.2202
- Pickett, H. M., Poynter, R. L., Cohen, E. A., Delitsky, M. L., Pearson, J. C., and Müller, H. S. P. (1998). Submillimeter, millimeter and microwave spectral line catalog. *Journal of Quantitative Spectroscopy and Radiative Transfer* 60, 883–890. doi:10.1016/S0022-4073(98)00091-0
- Pizzarello, S., Cooper, G. W., and Flynn, G. J. (2006). The Nature and Distribution of the Organic Material in Carbonaceous Chondrites and Interplanetary Dust Particles. In *Meteorites and the Early Solar System II*, eds. D. S. Lauretta and H. Y. McSween. 625
- Requena-Torres, M. A., Martín-Pintado, J., Martín, S., and Morris, M. R. (2008). The Galactic Center: The Largest Oxygen-bearing Organic Molecule Repository. *The Astrophysical Journal* 672, 352–360. doi:10.1086/523627
- Requena-Torres, M. A., Martín-Pintado, J., Rodríguez-Franco, A., Martín, S., Rodríguez-Fernández, N. J., and de Vicente, P. (2006). Organic molecules in the Galactic center. Hot core chemistry without hot cores. *Astronomy & Astrophysics* 455, 971–985. doi:10.1051/0004-6361:20065190
- Rivilla, V. M., Jiménez-Serra, I., Martín-Pintado, J., Briones, C., Rodríguez-Almeida, L. F., Rico-Villas, F., et al. (2021). Discovery in space of ethanolamine, the simplest phospholipid head group. *Proceedings of the National Academy of Science* 118, 2101314118. doi:10.1073/pnas.2101314118
- Rivilla, V. M., Martín-Pintado, J., Jiménez-Serra, I., Martín, S., Rodríguez-Almeida, L. F., Requena-Torres, M. A., et al. (2020). Prebiotic Precursors of the Primordial RNA World in Space: Detection of NH<sub>2</sub>OH. *The Astrophysical Journal Letters* 899, L28. doi:10.3847/2041-8213/abac55
- Rodríguez-Almeida, L. F., Jiménez-Serra, I., Rivilla, V. M., Martín-Pintado, J., Zeng, S., Tercero, B., et al. (2021). Thiols in the Interstellar Medium: First Detection of HC(O)SH and Confirmation of C<sub>2</sub>H<sub>5</sub>SH. *The Astrophysical Journal Letters* 912, L11. doi:10.3847/2041-8213/abf7cb
- Rodríguez-Fernández, N. J., Martín-Pintado, J., de Vicente, P., Fuente, A., Hüttemeister, S., Wilson, T. L., et al. (2000). Non-equilibrium H<sub>2</sub> ortho-to-para ratio in two molecular clouds of the Galactic Center. *Astronomy & Astrophysics* 356, 695–704
- Widicus Weaver, S. L. and Blake, G. A. (2005). 1,3-Dihydroxyacetone in Sagittarius B2(N-LMH): The First Interstellar Ketose. *The Astrophysical Journal Letters* 624, L33–L36. doi:10.1086/430407
- Zeng, S., Jiménez-Serra, I., Rivilla, V. M., Martín, S., Martín-Pintado, J., Requena-Torres, M. A., et al. (2018). Complex organic molecules in the Galactic Centre: the N-bearing family. *Monthly Notices of the Royal Astronomical Society* 478, 2962–2975. doi:10.1093/mnras/sty1174

- Zeng, S., Jiménez-Serra, I., Rivilla, V. M., Martín-Pintado, J., Rodríguez-Almeida, L. F., Tercero, B., et al. (2021). Probing the Chemical Complexity of Amines in the ISM: Detection of Vinylamine ( $C_2H_3NH_2$ ) and Tentative Detection of Ethylamine ( $C_2H_5NH_2$ ). *The Astrophysical Journal Letters* 920, L27. doi:10.3847/2041-8213/ac2c7e
- Zeng, S., Quénard, D., Jiménez-Serra, I., Martín-Pintado, J., Rivilla, V. M., Testi, L., et al. (2019). First detection of the pre-biotic molecule glycolonitrile ( $HOCH_2CN$ ) in the interstellar medium. *Monthly Notices of the Royal Astronomical Society* 484, L43–L48. doi:10.1093/mnrasl/slz002
- Zeng, S., Zhang, Q., Jiménez-Serra, I., Tercero, B., Lu, X., Martín-Pintado, J., et al. (2020). Cloud-cloud collision as drivers of the chemical complexity in Galactic Centre molecular clouds. *Monthly Notices of the Royal Astronomical Society* 497, 4896–4909. doi:10.1093/mnras/staa2187