

The case of equality in geometric instances of Barthe's reverse Brascamp-Lieb inequality

Karoly J. Boroczok*, Pavlos Kalantzopoulos†, Dongmeng Xi‡

November 16, 2022

Abstract

The works of Bennett, Carbery, Christ, Tao and of Valdimarsson have clarified when equality holds in the Brascamp-Lieb inequality. Here we characterize the case of equality in the Geometric case of Barthe's reverse Brascamp-Lieb inequality.

1 Introduction

For a proper linear subspace E of \mathbb{R}^n ($E \neq \mathbb{R}^n$ and $E \neq \{0\}$), let P_E denote the orthogonal projection into E . We say that the subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ form a Geometric Brascamp-Lieb data if they satisfy

$$\sum_{i=1}^k c_i P_{E_i} = I_n. \quad (1)$$

The name ‘‘Geometric Brascamp-Lieb data’’ coined by Bennett, Carbery, Christ, Tao [17] comes from the following theorem, originating in the work of Brascamp, Lieb [22] and Ball [3, 4] in the rank one case ($\dim E_i = 1$ for $i = 1, \dots, k$), and Lieb [75] and Barthe [8] in the general case. In the rank one case, the Geometric Brascamp-Lieb data is known as Parseval frame in coding theory and computer science (see for example Casazza, Tran, Tremain [33]).

Theorem 1 (Brascamp-Lieb, Ball, Barthe) *For the linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), and for non-negative $f_i \in L_1(E_i)$, we have*

$$\int_{\mathbb{R}^n} \prod_{i=1}^k f_i(P_{E_i}x)^{c_i} dx \leq \prod_{i=1}^k \left(\int_{E_i} f_i \right)^{c_i} \quad (2)$$

*Alfred Renyi Institute of Mathematics, Realtanoda utca 13-15, 1053, Budapest, Hungary, Supported by NKFIH 132002

†Central European University, Nador utca 9, 1051, Budapest, Hungary

‡Department of Mathematics, Shanghai University, Shanghai 200444, China, Supported by National Natural Science Foundation of China (12071277).

Remark This is Hölder's inequality if $E_1 = \dots = E_k = \mathbb{R}^n$ and $B_i = I_n$, and hence $\sum_{i=1}^k c_i = 1$.

We note that equality holds in Theorem 1 if $f_i(x) = e^{-\pi\|x\|^2}$ for $i = 1, \dots, k$; and hence, each f_i is a Gaussian density. Actually, Theorem 1 is an important special case discovered by Ball [4, 5] in the rank one case and by Barthe [8] in the general case of the general Brascamp-Lieb inequality Theorem 5.

After partial results by Barthe [8], Carlen, Lieb, Loss [31] and Bennett, Carbery, Christ, Tao [17], it was Valdimarsson [93] who characterized equality in the Geometric Brascamp-Lieb inequality. In order to state his result, we need some notation. Let E_1, \dots, E_k the proper linear subspaces of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfy (1). In order to understand extremizers in (5), following Carlen, Lieb, Loss [31] and Bennett, Carbery, Christ, Tao [17], we say that a non-zero linear subspace V is a critical subspace if

$$\sum_{i=1}^k c_i \dim(E_i \cap V) = \dim V,$$

which is turn equivalent saying that

$$E_i = (E_i \cap V) + (E_i \cap V^\perp) \text{ for } i = 1, \dots, k$$

according to [17] (see also Lemma 18). We say that a critical subspace V is indecomposable if V has no proper critical linear subspace.

Valdimarsson [93] introduced the so called independent subspaces and the dependent space. We write J to denote the set of 2^k functions $\{1, \dots, k\} \rightarrow \{0, 1\}$. If $\varepsilon \in J$, then let $F_{(\varepsilon)} = \bigcap_{i=1}^k E_i^{(\varepsilon(i))}$ where $E_i^{(0)} = E_i$ and $E_i^{(1)} = E_i^\perp$ for $i = 1, \dots, k$. We write J_0 to denote the subset of $\varepsilon \in J$ such that $\dim F_{(\varepsilon)} \geq 1$, and such an $F_{(\varepsilon)}$ is called independent following Valdimarsson [93]. Readily $F_{(\varepsilon)}$ and $F_{(\tilde{\varepsilon})}$ are orthogonal if $\varepsilon \neq \tilde{\varepsilon}$ for $\varepsilon, \tilde{\varepsilon} \in J_0$. In addition, we write F_{dep} to denote the orthogonal component of $\bigoplus_{\varepsilon \in J_0} F_{(\varepsilon)}$. In particular, \mathbb{R}^n can be written as a direct sum of pairwise orthogonal linear subspaces in the form

$$\mathbb{R}^n = \left(\bigoplus_{\varepsilon \in J_0} F_{(\varepsilon)} \right) \oplus F_{\text{dep}}. \quad (3)$$

Here it is possible that $J_0 = \emptyset$, and hence $\mathbb{R}^n = F_{\text{dep}}$, or $F_{\text{dep}} = \{0\}$, and hence $\mathbb{R}^n = \bigoplus_{\varepsilon \in J_0} F_{(\varepsilon)}$ in that case.

For a non-zero linear subspace $L \subset \mathbb{R}^n$, we say that a linear transformation $A : L \rightarrow L$ is positive definite if $\langle Ax, y \rangle = \langle x, Ay \rangle$ and $\langle x, Ax \rangle > 0$ for any $x, y \in L \setminus \{0\}$.

Theorem 2 (Valdimarsson) *For the proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), let us assume that equality holds in the Brascamp-Lieb inequality (2) for non-negative $f_i \in L_1(E_i)$, $i = 1, \dots, k$. If $F_{\text{dep}} \neq \mathbb{R}^n$, then let F_1, \dots, F_ℓ be the independent subspaces, and if $F_{\text{dep}} = \mathbb{R}^n$, then let $\ell = 1$ and $F_1 = \{0\}$. There exist $b \in F_{\text{dep}}$ and $\theta_i > 0$ for*

$i = 1, \dots, k$, integrable non-negative $h_j : F_j \rightarrow [0, \infty)$ for $j = 1, \dots, \ell$, and a positive definite matrix $A : F_{\text{dep}} \rightarrow F_{\text{dep}}$ such that the eigenspaces of A are critical subspaces and

$$f_i(x) = \theta_i e^{-\langle AP_{F_{\text{dep}}}x, P_{F_{\text{dep}}}x - b \rangle} \prod_{F_j \subset E_i} h_j(P_{F_j}(x)) \quad \text{for Lebesgue a.e. } x \in E_i. \quad (4)$$

On the other hand, if for any $i = 1, \dots, k$, f_i is of the form as in (4), then equality holds in (2) for f_1, \dots, f_k .

Theorem 2 explains the term "independent subspaces" because the functions h_j on F_j are chosen freely and independently from each other.

A reverse form of the Geometric Brascamp-Lieb inequality was proved by Barthe [8]. We write $\int_{\mathbb{R}^n}^*$ to denote the outer integral for a possibly non-integrable function $\varphi : \mathbb{R}^n \rightarrow [0, \infty)$; namely, the infimum (actually minimum) of $\int_{\mathbb{R}^n} \psi$ where $\psi \geq \varphi$ is Lebesgue measurable.

Theorem 3 (Barthe) For the non-trivial linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), and for non-negative $f_i \in L_1(E_i)$, we have

$$\int_{\mathbb{R}^n}^* \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k f_i(x_i)^{c_i} dx \geq \prod_{i=1}^k \left(\int_{E_i} f_i \right)^{c_i}. \quad (5)$$

Remark This is the Prékopa-Leindler inequality Theorem 28 if $E_1 = \dots = E_k = \mathbb{R}^n$ and $B_i = I_n$, and hence $\sum_{i=1}^k c_i = 1$.

We say that a function $h : \mathbb{R}^n \rightarrow [0, \infty)$ is log-concave if $h((1-\lambda)x + \lambda y) \geq h(x)^{1-\lambda} h(y)^\lambda$ for any $x, y \in \mathbb{R}^n$ and $\lambda \in (0, 1)$; or in other words, $h = e^{-W}$ for a convex function $W : \mathbb{R}^n \rightarrow (-\infty, \infty]$. Our main result is the following characterization of equality in the Geometric Barthe's inequality (5).

Theorem 4 For linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), if $F_{\text{dep}} \neq \mathbb{R}^n$, then let F_1, \dots, F_ℓ be the independent subspaces, and if $F_{\text{dep}} = \mathbb{R}^n$, then let $\ell = 1$ and $F_1 = \{0\}$.

If equality holds in the Geometric Barthe's inequality (5) for non-negative $f_i \in L_1(E_i)$ with $\int_{E_i} f_i > 0$, $i = 1, \dots, k$, then

$$f_i(x) = \theta_i e^{-\langle AP_{F_{\text{dep}}}x, P_{F_{\text{dep}}}x - b_i \rangle} \prod_{F_j \subset E_i} h_j(P_{F_j}(x - w_i)) \quad \text{for Lebesgue a.e. } x \in E_i \quad (6)$$

where

- $\theta_i > 0$, $b_i \in E_i \cap F_{\text{dep}}$ and $w_i \in E_i$ for $i = 1, \dots, k$,
- $h_j \in L_1(F_j)$ is non-negative for $j = 1, \dots, \ell$, and in addition, h_j is log-concave if there exist $\alpha \neq \beta$ with $F_j \subset E_\alpha \cap E_\beta$,

- $A : F_{\text{dep}} \rightarrow F_{\text{dep}}$ is a positive definite matrix such that the eigenspaces of A are critical subspaces.

On the other hand, if for any $i = 1, \dots, k$, f_i is of the form as in (6) and equality holds for all $x \in E_i$ in (6), then equality holds in (5) for f_1, \dots, f_k .

In particular, if for any $\alpha = 1, \dots, k$, $\{E_i\}_{i \neq \alpha}$ spans \mathbb{R}^n in Theorem 4, then any extremizer of the Geometric Barthe's inequality is log-concave.

The explanation for the phenomenon concerning the log-concavity of h_j in Theorem 4 is as follows (see the proof of Proposition 29). Let $\ell \geq 1$ and $j \in \{1, \dots, \ell\}$, and hence $\sum_{E_i \supset F_j} c_i = 1$. If f_1, \dots, f_k are of the form (6), then equality in Barthe's inequality (5) yields

$$\int_{F_j}^* \sup_{\substack{x = \sum_{E_i \supset F_j} c_i x_i \\ x_i \in F_j}} h_j(x_i - P_{F_j} w_i)^{c_i} dx = \prod_{E_i \supset F_j} \left(\int_{F_j} h_j(x - P_{F_j} w_i) dx \right)^{c_i} \left(= \int_{F_j} h_j(x) dx \right).$$

Therefore, if there exist $\alpha \neq \beta$ with $F_j \subset E_\alpha \cap E_\beta$, then the equality conditions in the Prékopa-Leindler inequality Proposition 28 imply that h_j is log-concave. On the other hand, if there exists $\alpha \in \{1, \dots, k\}$ such that $F_j \subset E_\beta^\perp$ for $\beta \neq \alpha$, then we do not have any condition on h_j , and $c_\alpha = 1$.

For completeness, let us state and discuss the general Brascamp-Lieb inequality and its reverse form due to Barthe. The following was proved by Brascamp, Lieb [22] in the rank one case and Lieb [75] in general.

Theorem 5 (Brascamp-Lieb Inequality) *Let $B_i : \mathbb{R}^n \rightarrow H_i$ be surjective linear maps where H_i is n_i -dimensional Euclidean space, $n_i \geq 1$, for $i = 1, \dots, k$, and let $c_1, \dots, c_k > 0$ satisfy $\sum_{i=1}^k c_i n_i = n$. For non-negative $f_i \in L_1(H_i)$, we have*

$$\int_{\mathbb{R}^n} \prod_{i=1}^k f_i(B_i x)^{c_i} dx \leq C \prod_{i=1}^k \left(\int_{H_i} f_i \right)^{c_i} \quad (7)$$

where C is determined by choosing centered Gaussians $f_i(x) = e^{-\langle A_i x, x \rangle}$, A_i positive definite.

Remark The Geometric Brascamp-Lieb Inequality is readily a special case of (7). We note that (7) is Hölder's inequality if $H_1 = \dots = H_k = \mathbb{R}^n$ and each $B_i = I_n$, and hence $C = 1$ and $\sum_{i=1}^k c_i = 1$ in that case.

We say that two Brascamp-Lieb data $\{(B_i, c_i)\}_{i=1, \dots, k}$ and $\{(B'_i, c'_i)\}_{i=1, \dots, k'}$ as in Theorem 5 are called equivalent if $k' = k$, $c'_i = c_i$, and there exists linear isomorphism $\Phi_i : H_i \rightarrow H'_i$ for $i = 1, \dots, k$ such that $B'_i = \Phi_i \circ B_i$. It was proved by Carlen, Lieb, Loss [31] in the rank one case, and by Bennett, Carbery, Christ, Tao [17] in general that there exists a set of extremizers f_1, \dots, f_k for (7) if and only if the Brascamp-Lieb data $\{(B_i, c_i)\}_{i=1, \dots, k}$ is equivalent

to some Geometric Brascamp-Lieb data. Therefore, Valdimarsson's Theorem 2 provides a full characterization of the equality case in Theorem 5, as well.

The following reverse version of the Brascamp-Lieb inequality was proved by Barthe in [7] in the rank one case, and in [8] in general.

Theorem 6 (Barthe's Inequality) *Let $B_i : \mathbb{R}^n \rightarrow H_i$ be surjective linear maps where H_i is n_i -dimensional Euclidean space, $n_i \geq 1$, for $i = 1, \dots, k$, and let $c_1, \dots, c_k > 0$ satisfy $\sum_{i=1}^k c_i n_i = n$. For non-negative $f_i \in L_1(H_i)$, we have*

$$\int_{\mathbb{R}^n}^* \sup_{x=\sum_{i=1}^k c_i B_i^* x_i, x_i \in H_i} \prod_{i=1}^k f_i(x_i)^{c_i} dx \geq D \prod_{i=1}^k \left(\int_{H_i} f_i \right)^{c_i} \quad (8)$$

where D is determined by choosing centered Gaussians $f_i(x) = e^{-\langle A_i x, x \rangle}$, A_i positive definite.

Remark The Geometric Barthe's Inequality is readily a special case of (8). We note that (8) is the Prékopa-Leindler inequality if $H_1 = \dots = H_k = \mathbb{R}^n$ and each $B_i = I_n$, and hence $D = 1$ and $\sum_{i=1}^k c_i = 1$ in that case.

Concerning extremals in Theorem 6, Lehec [69] proved that if there exists some Gaussian extremizers for Barthe's Inequality (8), then the corresponding Brascamp-Lieb data $\{(B_i, c_i)\}_{i=1, \dots, k}$ is equivalent to some Geometric Brascamp-Lieb data; therefore, the equality case of (8) can be understood via Theorem 4 in that case.

However, it is still not known whether having any extremizers in Barthe's Inequality (8) yields the existence of Gaussian extremizers. One possible approach is to use iterated convolutions and renormalizations as in Bennett, Carbery, Christ, Tao [17] in the case of Brascamp-Lieb inequality.

There are three main methods of proofs that work for proving both the Brascamp-Lieb Inequality and its reverse form, Barthe's inequality. The paper Barthe [8] used optimal transportation to prove Barthe's Inequality ("the Reverse Brascamp-Lieb inequality") and reprove the Brascamp-Lieb Inequality simultaneously. A heat equation argument was provided in the rank one case by Carlen, Lieb, Loss [31] for the Brascamp-Lieb Inequality and by Barthe, Cordero-Erausquin [10] for Barthe's inequality. The general versions of both inequalities are proved via the heat equation approach by Barthe, Huet [12]. Finally, simultaneous probabilistic arguments for the two inequalities are due to Lehec [69].

We note that Chen, Dafnis, Paouris [34] and Courtade, Liu [36], as well, deal systematically with finiteness conditions in Brascamp-Lieb and Barthe's inequalities. The importance of the Brascamp-Lieb inequality is shown by the fact that besides harmonic analysis, probability and convex geometry, it has been also even applied in number theory, see eg. Guo, Zhang [58]. Various versions of the Brascamp-Lieb inequality and its reverse form have been obtained by Balogh, Kristaly [6] Barthe [9], Barthe, Cordero-Erausquin [10], Barthe, Cordero-Erausquin, Ledoux, Maurey [11], Barthe, Wolff [13, 14], Bennett, Bez, Flock, Lee [15], Bennett, Bez, Buschenhenke, Cowling, Flock [16], Bobkov, Colesanti, Fragalà [19], Bueno, Pivarov [26], Chen, Dafnis, Paouris [34], Courtade, Liu [36], Duncan [40], Ghilli, Salani [46], Kolesnikov, Milman [68],

Livshyts [72, 73], Lutwak, Yang, Zhang [77, 78], Maldague [79], Marsiglietti [80], Rossi, Salani [89, 90].

Concerning the proof of Theorem 4, we discuss the structure theory of a Brascamp-Lieb data, Barthe's crucial determinantal inequality (*cf.* Proposition 22) and the extremality of Gaussians (*cf.* Proposition 25) in Sections 3, 4 and 5. Section 6 explains how Barthe's proof of his inequality using optimal transportation in [8] yields the splitting along independent and dependent subspaces in the case of equality in Barthe's inequality for positive C^1 probability densities f_1, \dots, f_k , and how the equality case of the Prékopa-Leindler inequality leads to the log-concavity of certain functions involved. However, one still needs to produce suitably smooth extremizers given any extremizers of Barthe's inequality. In order to achieve this, we discuss that convolution and suitable products of extremizers are also extremizers in Section 7. To show that extremizers are Gaussians on the dependent subspace, we use a version of Caffarelli's Contraction Principle in Section 8. Finally, all ingredients are pieced together to prove Theorem 4 in Section 9.

As applications of the understanding the equality case of the Brascamp-Lieb and Barthe's inequalities, we discuss the Bollobas-Thomason inequality and in its dual version in Section 2, and provide the characterization of the equality cases in Section 10.

2 Some applications: Equality in the Bollobas-Thomason inequality and in its dual

We write e_1, \dots, e_n to denote an orthonormal basis of \mathbb{R}^n . For a compact set $K \subset \mathbb{R}^n$ with $\dim \text{aff } K = m$, we write $|K|$ to denote the m -dimensional Lebesgue measure of K .

The starting point of this section is the classical Loomis-Whitney inequality [74].

Theorem 7 (Loomis, Whitney) *If $K \subset \mathbb{R}^n$ is compact and affinely spans \mathbb{R}^n , then*

$$|K|^{n-1} \leq \prod_{i=1}^k |P_{e_i^\perp} K|, \quad (9)$$

with equality if and only if $K = \bigoplus_{i=1}^n K_i$ where $\text{aff } K_i$ is a line parallel to e_i .

Meyer [83] provided a dual form of the Loomis-Whitney inequality where equality holds for affine crosspolytopes.

Theorem 8 (Meyer) *If $K \subset \mathbb{R}^n$ is compact convex with $o \in \text{int } K$, then*

$$|K|^{n-1} \geq \frac{n!}{n^n} \prod_{i=1}^k |K \cap e_i^\perp|, \quad (10)$$

with equality if and only if $K = \text{conv}\{\pm \lambda_i e_i\}_{i=1}^n$ for $\lambda_i > 0$, $i = 1, \dots, n$.

We note that various Reverse and dual Loomis-Whitney type inequalities are proved by Campi, Gardner, Gronchi [76], Brazitikos *et al* [24, 25], Alonso-Gutiérrez *et al* [1, 2].

To consider a generalization of the Loomis-Whitney inequality and its dual form, we set $[n] := \{1, \dots, n\}$, and for a non-empty proper subset $\sigma \subset [n]$, we define $E_\sigma = \text{lin}\{e_i\}_{i \in \sigma}$. For $s \geq 1$, we say that the not necessarily distinct proper non-empty subsets $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ if each $j \in [n]$ is contained in exactly s of $\sigma_1, \dots, \sigma_k$.

The Bollobas-Thomason inequality [18] reads as follows.

Theorem 9 (Bollobas, Thomason) *If $K \subset \mathbb{R}^n$ is compact and affinely spans \mathbb{R}^n , and $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$, then*

$$|K|^s \leq \prod_{i=1}^k |P_{E_{\sigma_i}} K|. \quad (11)$$

We note that additional the case when $k = n$, $s = n - 1$, and hence when we may assume that $\sigma_i = [n] \setminus e_i$, is the Loomis-Whitney inequality Theorem 7.

Liakopoulos [71] managed to prove a dual form of the Bollobas-Thomason inequality. For a finite set σ , we write $|\sigma|$ to denote its cardinality.

Theorem 10 (Liakopoulos) *If $K \subset \mathbb{R}^n$ is compact convex with $o \in \text{int}K$, and $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$, then*

$$|K|^s \geq \frac{\prod_{i=1}^k |\sigma_i|!}{(n!)^s} \cdot \prod_{i=1}^k |K \cap E_{\sigma_i}|. \quad (12)$$

The equality case of the Bollobas-Thomason inequality Theorem 9 based on Valdimarsson [93] has been known to the experts but we present this argument in order to have a written account. Let $s \geq 1$, and let $\sigma_1, \dots, \sigma_k \subset [n]$ be an s -uniform cover of $[n]$. We say that $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l \subset [n]$ form a 1-uniform cover of $[n]$ induced by the s -uniform cover $\sigma_1, \dots, \sigma_k$ if $\{\tilde{\sigma}_1, \dots, \tilde{\sigma}_l\}$ consists of all non-empty distinct subsets of $[n]$ of the form $\bigcap_{i=1}^k \sigma_i^{\varepsilon(i)}$ where $\varepsilon(i) \in \{0, 1\}$ and $\sigma_i^0 = \sigma_i$ and $\sigma_i^1 = [n] \setminus \sigma_i$. We observe that $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l \subset [n]$ actually form a 1-uniform cover of $[n]$; namely, $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ is a partition of $[n]$.

Theorem 11 (Folklore) *Let $K \subset \mathbb{R}^n$ be compact and affinely span \mathbb{R}^n , and let $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$. Then equality holds in (11) if and only if $K = \bigoplus_{i=1}^l P_{E_{\tilde{\sigma}_i}} K$ where $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ is the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$.*

Our main result in this section is the characterization of the equality case of the dual Bollobas-Thomason inequality Theorem 10 relating it to the Geometric Barthe's inequality.

Theorem 12 *Let $K \subset \mathbb{R}^n$ be compact convex with $o \in \text{int}K$, and let $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$. Then equality holds in (12) if and only if $K = \text{conv}\{K \cap F_{\tilde{\sigma}_i}\}_{i=1}^l$ where $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ is the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$.*

3 The determinantal inequality and structure theory for rank one Geometric Brascamp-Lieb data

We first discuss the basic properties of a set of vectors $u_1, \dots, u_k \in S^{n-1}$ and constants $c_1, \dots, c_k > 0$ occurring in the Geometric Brascamp-Lieb inequality; namely, satisfying

$$\sum_{i=1}^k c_i u_i \otimes u_i = I_n. \quad (13)$$

This section just retells the story of Section 2 of Barthe [8] in the language of Carlen, Lieb, Loss [31] and Bennett, Carbery, Christ, Tao [17].

Lemma 13 *For $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ satisfying (13), we have*

- (i) $\sum_{i=1}^k c_i = n$;
- (ii) $\sum_{i=1}^k c_i \langle u_i, x \rangle^2 = \|x\|^2$ for all $x \in \mathbb{R}^n$;
- (iii) $c_i \leq 1$ for $i = 1, \dots, k$ with equality if and only if $u_j \in u_i^\perp$ for $j \neq i$;
- (iv) u_1, \dots, u_k spans \mathbb{R}^n , and $k = n$ if and only if u_1, \dots, u_n is an orthonormal basis of \mathbb{R}^n and $c_1 = \dots = c_n = 1$;
- (v) if L is a proper linear subspace of \mathbb{R}^n , then

$$\sum_{u_i \in L} c_i \leq \dim L,$$

with equality if and only if $\{u_1, \dots, u_k\} \subset L \cup L^\perp$.

Remark If $\sum_{u_i \in L} c_i = \dim L$ in (v), then $\text{lin}\{u_i : u_i \in L\} = L$ and $\text{lin}\{u_i : u_i \in L^\perp\} = L^\perp$.

Proof: Here (i) follows from comparing the traces of the two sides of (13), and (ii) is just an equivalent form of (13). To prove $c_j \leq 1$ with the characterization of equality, we substitute $x = u_j$ into (ii).

Turning to (iv), u_1, \dots, u_k spans \mathbb{R}^n by (ii). Next, let us assume that $u_1, \dots, u_n \in S^{n-1}$ and $c_1, \dots, c_n > 0$ satisfy (13). We consider $w_j \in S^{n-1}$ for $j = 1, \dots, n$ such that $\langle w_j, u_i \rangle = 0$ if $i \neq j$, and hence (ii) shows that $u_j = \pm w_j$ and $c_j = 1$.

For (v), if $u_i \notin L$, then we consider the unit vector

$$\tilde{u}_i = \frac{P_{L^\perp} u_i}{\|P_{L^\perp} u_i\|} \in L^\perp.$$

We deduce that if $x \in L^\perp$, then

$$\|x\|^2 = \sum_{i=1}^k c_i \langle u_i, x \rangle^2 = \sum_{u_i \notin L} c_i \langle P_{L^\perp} u_i, x \rangle^2 = \sum_{u_i \notin L} c_i \|P_{L^\perp} u_i\|^2 \langle \tilde{u}_i, x \rangle^2.$$

It follows from (i) and (ii) applied to $\{\tilde{u}_i : u_i \notin L\}$ in L^\perp that

$$\dim L^\perp = \sum_{u_i \notin L} c_i \|P_{L^\perp} u_i\|^2 \leq \sum_{u_i \notin L} c_i.$$

In turn, we conclude the inequality in (v) by (i). Equality holds in (v) if and only if $\|P_{L^\perp} u_i\| = 1$ whenever $u_i \notin L$; therefore, $u_1, \dots, u_k \subset L \cup L^\perp$. \square

Let $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ satisfy (13). Following Bennett, Carbery, Christ, Tao [17], we say that a non-zero linear subspace V is a critical subspace with respect to u_1, \dots, u_k and c_1, \dots, c_k if

$$\sum_{u_i \in V} c_i = \dim V.$$

In particular, \mathbb{R}^n is a critical subspace according to Lemma 13. We say that a non-empty subset $\mathcal{U} \subset \{u_1, \dots, u_k\}$ is indecomposable if $\text{lin } \mathcal{U}$ is an indecomposable critical subspace.

In order to understand the equality case of the rank one Brascamp-Lieb inequality, Barthe [8] indicated an equivalence relation on $\{u_1, \dots, u_k\}$. We say that a subset $\mathcal{D} \subset \{u_1, \dots, u_k\}$ is minimally dependent if \mathcal{D} is dependent and no proper subset of \mathcal{D} is dependent. The following is folklore in matroid theory, was known most probably already to Tutte (see for example Theorem 7.3.6 in Recski [88]). For the convenience of the reader, we provide an argument.

Lemma 14 *Given non-zero v_1, \dots, v_k spanning \mathbb{R}^n , $n \geq 1$, we write $v_i \bowtie v_j$ if either $v_i = v_j$, or there exists a minimal dependent set $\mathcal{D} \subset \{v_1, \dots, v_k\}$ satisfying $v_i, v_j \in \mathcal{D}$.*

- (i) $v_i \bowtie v_j$ if and only if there exists a subset $\mathcal{U} \subset \{v_1, \dots, v_k\}$ of cardinality $n - 1$ such that both $\{v_i\} \cup \mathcal{U}$ and $\{v_j\} \cup \mathcal{U}$ are independent;
- (ii) \bowtie is an equivalence relation on $\{v_1, \dots, v_k\}$;
- (iii) if V_1, \dots, V_m are the linear hulls of the equivalence classes with respect to \bowtie , then they span \mathbb{R}^n and $V_i \cap V_j = \{0\}$ for $i \neq j$.

Proof: We prove the lemma by induction on $n \geq 1$ where the case $n = 1$ readily holds. Therefore, we assume that $n \geq 2$.

We may readily assume that

$$\{v_1, \dots, v_k\} \cap \text{lin } \{v_i\} = \{v_i\} \quad \text{for } i = 1, \dots, k. \quad (14)$$

For (i), if \mathcal{D} is a minimal dependent set with $v_i, v_j \in \mathcal{D}$, then adding some $\mathcal{V} \subset \{v_1, \dots, v_k\}$ to $\mathcal{D} \setminus \{v_i, v_j\}$, we obtain a basis of \mathbb{R}^n , and we may choose $\mathcal{U} = \mathcal{V} \cup (\mathcal{D} \setminus \{v_i, v_j\})$. On the other hand, if the suitable \mathcal{U} of cardinality $n - 1$ exists such that both $\{v_i\} \cup \mathcal{U}$ and $\{v_j\} \cup \mathcal{U}$ are independent, then any dependent subset of $\mathcal{U} \cup \{v_i, v_j\}$ contains v_i and v_j .

For (ii) and (iii), we call a non-zero linear subspace $W \subset \mathbb{R}^n$ unsplitable with respect to $\{v_1, \dots, v_k\}$ if W is spanned by $W \cap \{v_1, \dots, v_k\}$, but there exist no non-zero complementary linear subspaces $A, B \subset W$ with $\{v_1, \dots, v_k\} \cap W \subset A \cup B$. Readily, there exist pairwise

complementary unsplitable linear subspaces $W_1, \dots, W_m \subset \mathbb{R}^n$ such that $\{v_1, \dots, v_k\} \subset W_1 \cup \dots \cup W_m$.

On the one hand, if $v_i \in W_\alpha$ and $v_j \in W_\beta$ for $\alpha \neq \beta$, then trivially $v_i \not\propto v_j$. Therefore all we need to prove is that if $v_i, v_j \in W_\alpha$, then $v_i \propto v_j$. By the induction on n , we may assume that $m = 1$ and $W_\alpha = \mathbb{R}^n$. We may also assume that $i = 1$ and $j = 2$.

The final part of argument is indirect; therefore, we suppose that

$$v_1 \not\propto v_2, \quad (15)$$

and seek a contradiction.

(15) implies that v_1 and v_2 are independent, and hence $v_1 \not\propto v_2$ and (14) yield that $L = \text{lin}\{v_1, v_2\}$ satisfies

$$\{v_1, \dots, v_k\} \cap L = \{v_1, v_2\}. \quad (16)$$

Now \mathbb{R}^n is unsplitable, thus $n \geq 3$.

Since v_1, \dots, v_k span \mathbb{R}^n , we may assume that v_1, \dots, v_n form a basis of \mathbb{R}^n . Let $L_0 = \text{lin}\{v_3, \dots, v_n\}$, and $L_t = \text{lin}\{v_t, L_0\}$ for $t = 1, 2$. We may also assume that v_1, \dots, v_n is an orthonormal basis.

For any $l > n$, (i) and $v_1 \not\propto v_2$ yield that

$$\text{either } v_l \in L_1, \text{ or } v_l \in L_2. \quad (17)$$

Since \mathbb{R}^n is unsplitable, there exist $p, q > n$ such that

$$v_p \in L_1 \setminus L_0 \text{ and } v_q \in L_2 \setminus L_0. \quad (18)$$

For any $w \notin L$, we write

$$\text{supp } w = \{v_l : l \in \{3, \dots, n\} \ \& \ \langle w, v_l \rangle \neq 0\};$$

namely, the basis vectors where the corresponding coordinate of $w|_L = 0$ is non-zero.

Case 1 *There exist $v_p \in L_1 \setminus L_0$ and $v_q \in L_2 \setminus L_0$, $p, q > n$, such that $(\text{supp } v_p) \cap (\text{supp } v_q) \neq \emptyset$*

Let $v_s \in (\text{supp } v_p) \cap (\text{supp } v_q)$. Now the $n + 1$ element set

$$\{v_1, v_p, v_2, v_q\} \cup \{v_l : l \in \{3, \dots, n\} \setminus \{s\}\}$$

is dependent, and considering the 1^{st} , 2^{nd} and s^{th} coordinates show that both v_1 and v_2 lie in any dependent subset. This fact contradicts (15).

Case 2 *$(\text{supp } v_p) \cap (\text{supp } v_q) = \emptyset$ for any $v_p \in L_1 \setminus L_0$ and $v_q \in L_2 \setminus L_0$ with $p, q > n$*

Let $\mathcal{U}_t = \cup\{\text{supp } v_p : p > n \ \& \ v_p \in L_t \setminus L_0\}$ for $t = 1, 2$. It follows that $\mathcal{U}_1 \cap \mathcal{U}_2 = \emptyset$, thus $n \geq 4$. For any partition $\mathcal{U}'_1 \cup \mathcal{U}'_2 = \{v_3, \dots, v_n\}$ (and hence $\mathcal{U}'_1 \cap \mathcal{U}'_2 = \emptyset$) such that $\mathcal{U}_1 \subset \mathcal{U}'_1$ and $\mathcal{U}_2 \subset \mathcal{U}'_2$, there exists some $v_l \in L_0$ that is contained neither in $\text{lin}(\mathcal{U}'_1 \cup \{v_1\})$ nor in $\text{lin}(\mathcal{U}'_2 \cup \{v_2\})$ because \mathbb{R}^n is unsplitable. In turn we deduce that we may reindex the vectors v_3, \dots, v_n on the one hand, and the vectors v_{n+1}, \dots, v_k on the other hand to ensure the following properties:

- $v_{n+1} \in L_1 \setminus L_0$ and $v_{n+2} \in L_2 \setminus L_0$;
- there exist $\alpha \in \{3, \dots, n-1\}$ and $\beta \in \{n+3, \dots, k\}$ such that $\text{supp } v_l \subset \{v_\alpha, \dots, v_n\}$ for $l \in \{n+1, \dots, \beta\}$, and $v_l \in L_0$ if $n+3 \leq l \leq \beta$;
- for any partition $\mathcal{W}_1 \cup \mathcal{W}_2 = \{v_\alpha, \dots, v_n\}$ into non-empty sets, there exist $l \in \{n+1, \dots, \beta\}$ such that $\text{supp } v_l$ intersects both \mathcal{W}_1 and \mathcal{W}_2 .

We observe that $\tilde{L}_0 = \text{lin}\{v_\alpha, \dots, v_n\}$ is unsplitable with respect to

$$\{v_\alpha, \dots, v_n, v_{n+1}|L_0, v_{n+2}|L_0, v_{n+3}, \dots, v_\beta\}.$$

Therefore, this last set contains a minimal dependent subset $\tilde{\mathcal{D}}$ with $v_{n+1}|L_0, v_{n+2}|L_0 \in \tilde{\mathcal{D}}$ by induction; namely, the elements of $\tilde{\mathcal{D}}$ different from $v_{n+1}|L_0, v_{n+2}|L_0$ are vectors of the form v_l that lie in L_0 . We conclude that

$$\mathcal{D} = \{v_1, v_2, v_{n+1}, v_{n+2}\} \cup \left(\tilde{\mathcal{D}} \setminus \{v_{n+1}|L_0, v_{n+2}|L_0\} \right)$$

is a minimal dependent set, contradicting (15), and proving Lemma 14. \square

Lemma 15 For $u_1, \dots, u_k \in S^{m-1}$ and $c_1, \dots, c_k > 0$ satisfying (13), we have

- (i) a proper linear subspace $V \subset \mathbb{R}^n$ is critical if and only if $\{u_1, \dots, u_k\} \subset V \cup V^\perp$;
- (ii) if V, W are proper critical subspaces with $V \cap W \neq \{0\}$, then $V^\perp, V \cap W$ and $V + W$ are critical subspaces;
- (iii) the equivalence classes with respect to the relation \bowtie in Lemma 14 are the indecomposable subsets of $\{u_1, \dots, u_k\}$;
- (iv) the proper indecomposable critical subspaces are pairwise orthogonal, and any critical subspace is the sum of some indecomposable critical subspaces.

Proof: (i) directly follows from Lemma 13 (v), and in turn (i) yields (ii).

We prove (iii) and first half of (iv) simultaneously. Let V_1, \dots, V_m be the linear hulls of the equivalence classes of u_1, \dots, u_k with respect to the \bowtie of Lemma 14. We deduce from Lemma 13 (v) that each V_i is a critical subspace, and if $i \neq j$, then V_i and V_j are orthogonal.

Next let $\mathcal{U} \subset \{u_1, \dots, u_k\}$ be an indecomposable set, and let $V = \text{lin } \mathcal{U}$. We write $I \subset \{1, \dots, m\}$ to denote the set of indices i such that $V_i \cap \mathcal{U} \neq \emptyset$. Since V is a critical subspace, we deduce from Lemma 13 (v) that $V_i \cap V$ is a critical subspace for $i \in I$, as well; therefore, I consists of a unique index p as \mathcal{U} is indecomposable. In particular, $V = V_p$.

It follows from Lemma 13 (v) that $\{u_1, \dots, u_k\} \subset V \cup V^\perp$; therefore, there exists no minimally dependent subset of $\{u_1, \dots, u_k\}$ intersecting both \mathcal{U} and its complement. We conclude that $V = V_p$.

Finally, the second half of (iv) follows from (i) and (ii). \square

The following is the main result of this section, where the inequality is proved by Ball [3, 4], and the equality case is clarified by Barthe [8].

Proposition 16 (Ball-Barthe Lemma) For $u_1, \dots, u_k \in S^{n-1}$ and $c_1, \dots, c_k > 0$ satisfying (13), if $t_i > 0$ for $i = 1, \dots, k$, then

$$\det \left(\sum_{i=1}^k c_i t_i u_i \otimes u_i \right) \geq \prod_{i=1}^k t_i^{c_i}. \quad (19)$$

Equality holds in (19) if and only if $t_i = t_j$ for any u_i and u_j lying in the same indecomposable subset of $\{u_1, \dots, u_k\}$.

Proof: To simplify expressions, let $v_i = \sqrt{c_i} u_i$ for $i = 1, \dots, k$.

In this argument, I always denotes some subset of $\{1, \dots, k\}$ of cardinality n . For $I = \{i_1, \dots, i_n\}$, we define

$$d_I := \det[v_{i_1}, \dots, v_{i_n}]^2 \quad \text{and} \quad t_I := t_{i_1} \cdots t_{i_n}.$$

For the $n \times k$ matrices $M = [v_1, \dots, v_k]$ and $\widetilde{M} = [\sqrt{t_1} v_1, \dots, \sqrt{t_k} v_k]$, we have

$$MM^T = I_n \quad \text{and} \quad \widetilde{M}\widetilde{M}^T = \sum_{i=1}^k t_i v_i \otimes v_i. \quad (20)$$

It follows from the Cauchy-Binet formula that

$$\sum_I d_I = 1 \quad \text{and} \quad \det \left(\sum_{i=1}^k t_i v_i \otimes v_i \right) = \sum_I t_I d_I,$$

where the summations extend over all sets $I \subset \{1, \dots, k\}$ of cardinality n . It follows that the discrete measure μ on the n element subsets of $\{1, \dots, k\}$ defined by $\mu(\{I\}) = d_I$ is a probability measure. We deduce from inequality between the arithmetic and geometric mean that

$$\det \left(\sum_{i=1}^k t_i v_i \otimes v_i \right) = \sum_I t_I d_I \geq \prod_I t_I^{d_I}. \quad (21)$$

The factor t_i occurs in $\prod_I t_I^{d_I}$ exactly $\sum_{I, i \in I} d_I$ times. Moreover, the Cauchy-Binet formula applied to the vectors $v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k$ implies

$$\begin{aligned} \sum_{I, i \in I} d_I &= \sum_I d_I - \sum_{I, i \notin I} d_I = 1 - \det \left(\sum_{j \neq i} v_j \otimes v_j \right) \\ &= 1 - \det(\text{Id}_n - v_i \otimes v_i) = \langle v_i, v_i \rangle = c_i. \end{aligned}$$

Substituting this into (21) yields (19).

We now assume that equality holds in (19). Since equality holds in (21) when applying arithmetic and geometric mean, all the t_I are the same for any subset I of $\{1, \dots, k\}$ of cardinality

n with $d_I \neq 0$. It follows that $t_i = t_j$ whenever $u_i \bowtie u_j$, and in turn we deduce that $t_i = t_j$ whenever u_i and u_j lie in the same indecomposable set by Lemma 15 (i).

On the other hand, Lemma 15 (ii) yields that if $t_i = t_j$ whenever u_i and u_j lie in the same indecomposable set, then equality holds in (19). \square

Combining Lemma 15 and Proposition 16 leads to the following:

Corollary 17 *For $u_i \in S^{n-1}$ and $c_i, t_i > 0$, $i = 1, \dots, k$ satisfying (13), equality holds in (19) if and only if there exist pairwise orthogonal linear subspaces V_1, \dots, V_m , $m \geq 1$, such that $\{u_1, \dots, u_k\} \subset V_1 \cup \dots \cup V_m$ and $t_i = t_j$ whenever u_i and u_j lie in the same V_p for some $p \in \{1, \dots, m\}$.*

4 Structure theory of a Brascamp-Lieb data and the determinantal inequality corresponding to the higher rank case

We build a structural theory for a Brascamp-Lieb data based on results proved or indicated in Barthe [8], Bennett, Carbery, Christ, Tao [17] and Valdimarsson [93].

For non-zero linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying the Geometric Brascamp-Lieb condition

$$\sum_{i=1}^k c_i P_{E_i} = I_n, \quad (22)$$

we connect (22) to (13). For $i = 1, \dots, k$, let $\dim E_i = n_i$ and let $u_1^{(i)}, \dots, u_{n_i}^{(i)}$ be any orthonormal basis of E_i . In addition, for $i = 1, \dots, k$, we consider the $n \times n_i$ matrix $M_i = \sqrt{c_i}[u_1^{(i)}, \dots, u_{n_i}^{(i)}]$. We deduce that

$$c_i P_{E_i} = M_i M_i^T = \sum_{j=1}^{n_i} c_i u_j^{(i)} \otimes u_j^{(i)} \quad \text{for } i = 1, \dots, k; \quad (23)$$

$$I_n = \sum_{i=1}^k c_i P_{E_i} = \sum_{i=1}^k \sum_{j=1}^{n_i} c_i u_j^{(i)} \otimes u_j^{(i)} = \sum_{i=1}^k \sum_{j=1}^{n_i} c_j^{(i)} u_j^{(i)} \otimes u_j^{(i)} \quad (24)$$

and hence $u_j^{(i)} \in S^{n-1}$ and $c_j^{(i)} = c_i > 0$ for $i = 1, \dots, k$ and $j = 1, \dots, n_i$ form a Geometric Brascamp-Lieb data like in (13).

Lemma 18 *For linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22),*

(i) *if $x \in \mathbb{R}^n$, then $\sum_{i=1}^k c_i \|P_{E_i} x\|^2 = \|x\|^2$;*

(ii) *if $V \subset \mathbb{R}^n$ is a proper linear subspace, then*

$$\sum_{i=1}^k c_i \dim(E_i \cap V) \leq \dim V \quad (25)$$

where equality holds if and only if $E_i = (E_i \cap V) + (E_i \cap V^\perp)$ for $i = 1, \dots, k$; or equivalently, when $V = (E_i \cap V) + (E_i^\perp \cap V)$ for $i = 1, \dots, k$.

Proof: For $i = 1, \dots, k$, let $\dim E_i = n_i$ and let $u_1^{(i)}, \dots, u_{n_i}^{(i)}$ be any orthonormal basis of E_i such that if $V \cap E_i \neq \{0\}$, then $u_1^{(i)}, \dots, u_{m_i}^{(i)}$ is any orthonormal basis of $V \cap E_i$ where $m_i \leq n_i$.

For any $x \in \mathbb{R}^n$ and $i = 1, \dots, k$, we have $\|P_{E_i}x\|^2 = \sum_{j=1}^{n_i} \langle u_j^{(i)}, x \rangle^2$, thus Lemma 13 (ii) yields (i).

Concerning (ii), Lemma 13 (v) yields (25). On the other hand, if equality holds in (25), then V is a critical subspace for the rank one Geometric Brascamp-Lieb data $u_j^{(i)} \in S^{n-1}$ and $c_j^{(i)} = c_i > 0$ for $i = 1, \dots, k$ and $j = 1, \dots, n_i$ satisfying (24). Thus Lemma 18 (ii) follows from Lemma 13 (v). \square

We say that a non-zero linear subspace V is a critical subspace with respect to the proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22) if

$$\sum_{i=1}^k c_i \dim(E_i \cap V) = \dim V.$$

In particular, \mathbb{R}^n is a critical subspace by calculating traces of both sides of (22). For a proper linear subspace $V \subset \mathbb{R}^n$, Lemma 18 yields that V is critical if and only if V^\perp is critical, which is turn equivalent saying that

$$E_i = (E_i \cap V) + (E_i \cap V^\perp) \text{ for } i = 1, \dots, k; \quad (26)$$

or in other words,

$$V = (E_i \cap V) + (E_i^\perp \cap V) \text{ for } i = 1, \dots, k. \quad (27)$$

We observe that (26) has the following consequence: If V_1 and V_2 are orthogonal critical subspaces, then

$$E_i \cap (V_1 + V_2) = (E_i \cap V_1) + (E_i \cap V_2) \text{ for } i = 1, \dots, k. \quad (28)$$

We recall that a critical subspace V is indecomposable if V has no proper critical linear subspace.

Lemma 19 *If E_1, \dots, E_k are linear subspaces of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), and V, W are proper critical subspaces, then V^\perp and $V + W$ are critical subspaces, and even $V \cap W$ is critical provided that $V \cap W \neq \{0\}$.*

Proof: We may assume that $\dim E_i \geq 1$ for $i = 1, \dots, k$.

The fact that V^\perp is also critical follows directly from (26).

Concerning $V \cap W$ when $V \cap W \neq \{0\}$, we need to prove that if $i = 1, \dots, k$, then

$$(V \cap W) \cap E_i + (V \cap W)^\perp \cap E_i = E_i. \quad (29)$$

For a linear subspace $L \subset E_i$, we write $L^{\perp_i} = L^{\perp} \cap E_i$ to denote the orthogonal complement within E_i . We observe that as V and W are critical subspaces, we have $(V \cap E_i)^{\perp_i} = V^{\perp} \cap E_i$ and $(W \cap E_i)^{\perp_i} = W^{\perp} \cap E_i$. It follows from the identity $(V \cap W)^{\perp} = V^{\perp} + W^{\perp}$ that

$$\begin{aligned}
E_i &\supset (V \cap W) \cap E_i + (V \cap W)^{\perp} \cap E_i = (V \cap E_i) \cap (W \cap E_i) + (V^{\perp} + W^{\perp}) \cap E_i \\
&\supset (V \cap E_i) \cap (W \cap E_i) + (V^{\perp} \cap E_i) + (W^{\perp} \cap E_i) \\
&= (V \cap E_i) \cap (W \cap E_i) + (V \cap E_i)^{\perp_i} + (W \cap E_i)^{\perp_i} \\
&= (V \cap E_i) \cap (W \cap E_i) + [(V \cap E_i) \cap (W \cap E_i)]^{\perp_i} = E_i,
\end{aligned}$$

yielding (29).

Finally, $V + W$ is also critical as $V + W = (V^{\perp} \cap W^{\perp})^{\perp}$. \square

We deduce from Lemma 19 that any critical subspace can be decomposed into indecomposable ones.

Corollary 20 *If E_1, \dots, E_k are proper linear subspaces of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfy (22), and W is a critical subspace or $W = \mathbb{R}^n$, then there exist pairwise orthogonal indecomposable critical subspaces V_1, \dots, V_m , $m \geq 1$, such that $W = V_1 + \dots + V_m$ (possibly $m = 1$ and $W = V_1$).*

We note that the decomposition of \mathbb{R}^n into indecomposable critical subspaces is not unique in general for a Geometric Brascamp-Lieb data. Valdimarsson [93] provides some examples, and in addition, we provide an example where we have a continuous family of indecomposable critical subspaces.

Example 21 (Continuous family of indecomposable critical subspaces) *In \mathbb{R}^4 , let us consider the following six unit vectors: $u_1(1, 0, 0, 0)$, $u_2(\frac{1}{2}, \frac{\sqrt{3}}{2}, 0, 0)$, $u_3(\frac{-1}{2}, \frac{\sqrt{3}}{2}, 0, 0)$, $v_1(0, 0, 1, 0)$, $v_2(0, 0, \frac{1}{2}, \frac{\sqrt{3}}{2})$, $v_3(0, 0, \frac{-1}{2}, \frac{\sqrt{3}}{2})$, which satisfy $u_2 = u_1 + u_3$ and $v_2 = v_1 + v_3$.*

For any $x \in \mathbb{R}^4$, we have

$$\|x\|^2 = \sum_{i=1}^3 \frac{2}{3} \cdot (\langle x, u_i \rangle^2 + \langle x, v_i \rangle^2)$$

Therefore, we define the Geometric Brascamp-Lieb Data $E_i = \text{lin}\{u_i, v_i\}$ and $c_i = \frac{2}{3}$ for $i = 1, 2, 3$ satisfying (1). In this case, $F_{\text{dep}} = \mathbb{R}^4$.

For any angle $t \in \mathbb{R}$, we have a two-dimensional indecomposable critical subspace

$$V_t = \text{lin}\{(\cos t)u_1 + (\sin t)v_1, (\cos t)u_2 + (\sin t)v_2, (\cos t)u_3 + (\sin t)v_3\}.$$

Next we prove the crucial determinantal inequality. Its proof is kindly provided by Franck Barthe.

Proposition 22 (Barthe) For linear subspaces E_1, \dots, E_k of \mathbb{R}^n , $n \geq 1$ and $c_1, \dots, c_k > 0$ satisfying (22), if $A_i : E_i \rightarrow E_i$ is a positive definite linear transformation for $i = 1, \dots, k$, then

$$\det \left(\sum_{i=1}^k c_i A_i P_{E_i} \right) \geq \prod_{i=1}^k (\det A_i)^{c_i}. \quad (30)$$

Equality holds in (30) if and only if there exist linear subspaces V_1, \dots, V_m where $V_1 = \mathbb{R}^n$ if $m = 1$ and V_1, \dots, V_m are pairwise orthogonal indecomposable critical subspaces spanning \mathbb{R}^n if $m \geq 2$, and a positive definite $n \times n$ matrix Φ such that V_1, \dots, V_m are eigenspaces of Φ and $\Phi|_{E_i} = A_i$ for $i = 1, \dots, k$. In addition, $\Phi = \sum_{i=1}^k c_i A_i P_{E_i}$ in the case of equality.

Proof: We may assume that $\dim E_i \geq 1$ for $i = 1, \dots, k$.

For $i = 1, \dots, k$, let $\dim E_i = n_i$, let $u_1^{(i)}, \dots, u_{n_i}^{(i)}$ be an orthonormal basis of E_i consisting of eigenvectors of A_i , and let $\lambda_j^{(i)} > 0$ be the eigenvalue of A_i corresponding to $u_j^{(i)}$. In particular $\det A_i = \prod_{j=1}^{n_i} \lambda_j^{(i)}$ for $i = 1, \dots, k$. In addition, for $i = 1, \dots, k$, we set $M_i = \sqrt{c_i} [u_1^{(i)}, \dots, u_{n_i}^{(i)}]$ and B_i to be the positive definite transformation with $A_i = B_i B_i$, and hence

$$c_i A_i P_{E_i} = (M_i B_i)(M_i B_i)^T = \sum_{j=1}^{n_i} c_i \lambda_j^{(i)} u_j^{(i)} \otimes u_j^{(i)}.$$

We deduce from Lemma 16 and (24) that

$$\begin{aligned} \det \left(\sum_{i=1}^k c_i A_i P_{E_i} \right) &= \det \left(\sum_{i=1}^k \sum_{j=1}^{n_i} c_i \lambda_j^{(i)} u_j^{(i)} \otimes u_j^{(i)} \right) \\ &\geq \prod_{i=1}^k \left(\prod_{j=1}^{n_i} \lambda_j^{(i)} \right)^{c_i} = \prod_{i=1}^k (\det A_i)^{c_i}. \end{aligned} \quad (31)$$

If we have equality in (30), and hence also in (31), then Corollary 17 implies that there exist pairwise orthogonal critical subspaces V_1, \dots, V_m , $m \geq 1$ spanning \mathbb{R}^n and $\lambda_1, \dots, \lambda_m > 0$ (where $V_1 = \mathbb{R}^n$ if $m = 1$) such that if $E_i \cap V_j \neq \{0\}$, then $E_i \cap V_j$ is an eigenspace of A_i with eigenvalue λ_j . We conclude from (26) that each V_j is a critical subspace, and from Corollary 20 that each V_j can be assumed to be indecomposable. Finally, (28) yields that each E_i is spanned by the subspaces $E_i \cap V_j$ for $j = 1, \dots, m$.

To show that each V_j is an eigenspace for the positive definite linear transform $\sum_{i=1}^k c_i A_i P_{E_i}$ of \mathbb{R}^n with eigenvalue λ_j , we observe that

$$A_i P_{E_i} x = \lambda_j P_{E_i} x$$

for any $i = 1, \dots, k$ and $x \in V_j$. It follows that if $x \in V_j$, then

$$\sum_{i=1}^k c_i A_i P_{E_i} x = \lambda_j \sum_{i=1}^k c_i P_{E_i} x = \lambda_j x,$$

proving that we can choose $\Phi = \sum_{i=1}^k c_i A_i P_{E_i}$.

On the other hand, let us assume that there exists a positive definite $n \times n$ matrix Θ whose eigenspaces W_1, \dots, W_l are critical subspaces (or $l = 1$ and $W_1 = \mathbb{R}^n$) and $\Theta|_{E_i} = A_i$ for $i = 1, \dots, k$. In this case, for any $i = 1, \dots, k$, we may choose the orthonormal basis $u_1^{(i)}, \dots, u_{n_i}^{(i)}$ of E_i in a way such that $u_1^{(i)}, \dots, u_{n_i}^{(i)} \subset W_1 \cup \dots \cup W_l$, and hence Corollary 17 yields that equality holds in (30). \square

Remark While Proposition 22 has a crucial role in proving both the Brascamp-Lieb inequality (2) and Barthe's inequality (5) and their equality cases, Proposition 22 can be actually derived from say (2). In the Brascamp-Lieb inequality, choose $f_i(z) = e^{-\pi \langle A_i z, z \rangle}$ for $z \in E_i$ and $i = 1, \dots, k$, and hence $\int_{E_i} f_i = (\det A_i)^{-\frac{1}{2}}$. On the other hand, if $x \in \mathbb{R}^n$, then

$$\prod_{i=1}^k f_i(P_{E_i} x)^{c_i} = e^{-\pi \sum_{i=1}^k c_i \langle A_i P_{E_i} x, P_{E_i} x \rangle} = e^{-\pi \sum_{i=1}^k c_i \langle A_i P_{E_i} x, x \rangle} = e^{-\pi \langle \sum_{i=1}^k c_i A_i P_{E_i} x, x \rangle},$$

therefore, the Brascamp-Lieb inequality (2) yields

$$\left(\det \sum_{i=1}^k c_i A_i P_{E_i} \right)^{-\frac{1}{2}} \leq \prod_{i=1}^k (\det A_i)^{-\frac{c_i}{2}}.$$

In addition, the equality conditions in Proposition 22 can be derived from Valdimarsson's Theorem 2.

Let us show why indecomposability of the critical subspaces in Proposition 22 is useful.

Lemma 23 *Let the linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfy (22), let $F_{\text{dep}} \neq \mathbb{R}^n$, and let F_1, \dots, F_l be the independent subspaces, $l \geq 1$. If V is an indecomposable critical subspace, then either $V \subset F_{\text{dep}}$, or there exists an independent subspace F_j , $j \in \{1, \dots, l\}$ such that $V \subset F_j$.*

Proof: It is equivalent to prove that if V is an indecomposable critical subspace and $j \in \{1, \dots, l\}$, then

$$V \not\subset F_j \text{ implies } F_j \subset V^\perp. \quad (32)$$

We deduce that $V \cap F_j = \{0\}$ from the facts that V is indecomposable and F_j is a critical subspace, thus $F_j \cap V$ is a critical subspace or $\{0\}$. There exists a partition $M \cup N = \{1, \dots, k\}$ with $M \cap N = \emptyset$ such that

$$F_j = \left(\bigcap_{i \in M} E_i \right) \cap \left(\bigcap_{i \in N} E_i^\perp \right).$$

Let $y \in F_j$. Since V is a critical subspace, we conclude that $P_V y \in E_i$ for $i \in M$ and $P_V y \in E_i^\perp$ for $i \in N$, and hence $P_V y \in V \cap \left(\bigcap_{i \in M} E_i \right) \cap \left(\bigcap_{i \in N} E_i^\perp \right) = \{0\}$. Therefore, $y \in V^\perp$. \square

5 Typical Gaussian extremizers for some Geometric Brascamp-Lieb data

This section continues to build on work done in Barthe [8], Bennett, Carbery, Christ, Tao [17] and Valdimarsson [93].

For linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), we deduce from Lemma 18 (i) and (27) that if V is a critical subspace, then writing $P_{E_i \cap V}^{(V)}$ to denote the restriction of $P_{E_i \cap V}$ onto V , we have

$$\sum_{E_i \cap V \neq \{0\}} c_i P_{E_i \cap V}^{(V)} = I_V \quad (33)$$

where I_V denotes the identity transformation on V .

The equality case of Proposition 22 indicates why Lemma 24 is important.

Lemma 24 *For linear subspaces E_1, \dots, E_k of \mathbb{R}^n , $n \geq 1$ and $c_1, \dots, c_k > 0$ satisfying (22), if Φ is a positive definite linear transform whose eigenspaces are critical subspaces, then for any $x \in \mathbb{R}^n$, we have*

$$\|\Phi x\|^2 = \min_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \sum_{i=1}^k c_i \|\Phi x_i\|^2. \quad (34)$$

Proof: We may assume that $\dim E_i \geq 1$ for $i = 1, \dots, k$.

As the eigenspaces of Φ are critical subspaces, we deduce that

$$\Phi(E_i) = E_i \text{ and } \Phi(E_i^\perp) = E_i^\perp. \quad (35)$$

For any $x \in \mathbb{R}^n$, we have $\Phi P_{E_i} x = P_{E_i} \Phi x$ for $i = 1, \dots, k$ by (35); therefore, Lemma 18 (i) yields

$$\langle \Phi x, \Phi x \rangle = \sum_{i=1}^k c_i \|P_{E_i} \Phi x\|^2 = \sum_{i=1}^k c_i \|\Phi P_{E_i} x\|^2. \quad (36)$$

Since $x = \sum_{i=1}^k c_i P_{E_i} x$ by (22), we may choose $x_i = P_{E_i} x$ in (34), and we have equality in (34) in this case. Therefore, Lemma 24 is equivalent to proving that if $x = \sum_{i=1}^k c_i x_i$ for $x_i \in E_i$, $i = 1, \dots, k$, then

$$\|\Phi x\|^2 \leq \sum_{i=1}^k c_i \|\Phi x_i\|^2. \quad (37)$$

Case 1 $\dim E_i = 1$ for $i = 1, \dots, k$ and $\Phi = I_n$

Let $E_i = \mathbb{R}u_i$ for $u_i \in S^{n-1}$. If $x \in \mathbb{R}^n$, then $P_{E_i} x = \langle u_i, x \rangle u_i$ for $i = 1, \dots, k$, and (36) yields that

$$\langle x, x \rangle = \sum_{i=1}^k c_i \langle u_i, x \rangle^2.$$

In addition, any $x_i \in E_i$ is of the form $x_i = t_i u_i$ for $i = 1, \dots, k$ where $\|x_i\|^2 = t_i^2$. If $x = \sum_{i=1}^k c_i t_i u_i$, then the Hölder inequality yields

$$\langle x, x \rangle = \left\langle x, \sum_{i=1}^k c_i t_i u_i \right\rangle = \sum_{i=1}^k c_i t_i \langle x, u_i \rangle \leq \sqrt{\sum_{i=1}^k c_i t_i^2} \cdot \sqrt{\sum_{i=1}^k c_i \langle x, u_i \rangle^2} = \sqrt{\sum_{i=1}^k c_i t_i^2} \cdot \sqrt{\langle x, x \rangle},$$

proving (37) in this case.

Case 2 The general case, E_1, \dots, E_k and Φ are as in Lemma 24

Let V_1, \dots, V_m , $m \geq 1$, be the eigenspaces of Φ corresponding to the eigenvalues $\lambda_1, \dots, \lambda_m$. As V_1, \dots, V_m are orthogonal critical subspaces and $\mathbb{R}^n = \bigoplus_{j=1}^m V_j$. As V_1, \dots, V_m are orthogonal critical subspaces and $\mathbb{R}^n = \bigoplus_{j=1}^m V_j$, we deduce that $x_{ij} = P_{V_j} x_i \in E_i \cap V_j$ for any $i = 1, \dots, k$ and $j = 1, \dots, m$, and $x_i = \sum_{j=1}^m x_{ij}$ for any $i = 1, \dots, k$. It follows that

$$x = \sum_{j=1}^m \left(\sum_{E_i \cap V_j \neq \{0\}} c_i x_{ij} \right) \text{ where}$$

$$P_{V_j} x = \sum_{E_i \cap V_j \neq \{0\}} c_i x_{ij}. \quad (38)$$

For any $i = 1, \dots, k$, the vectors $\Phi x_{ij} = \lambda_j x_{ij}$ are pairwise orthogonal for $j = 1, \dots, m$, thus

$$\sum_{i=1}^k c_i \|\Phi x_i\|^2 = \sum_{i=1}^k \left(\sum_{j=1}^m c_i \|\Phi x_{ij}\|^2 \right) = \sum_{j=1}^m \left(\sum_{E_i \cap V_j \neq \{0\}} c_i \|\Phi x_{ij}\|^2 \right).$$

Since $\|\Phi x\|^2 = \sum_{j=1}^m \|P_{V_j} \Phi x\|^2 = \sum_{j=1}^m \|\Phi P_{V_j} x\|^2$, (37) follows if for any $j = 1, \dots, m$, we have

$$\|\Phi P_{V_j} x\|^2 \leq \sum_{E_i \cap V_j \neq \{0\}} c_i \|\Phi x_{ij}\|^2. \quad (39)$$

To prove (39), if $E_i \cap V_j \neq \{0\}$, then let $\dim(E_i \cap V_j) = n_{ij}$, and let $u_1^{(ij)}, \dots, u_{n_{ij}}^{(ij)}$ be an orthonormal basis of $E_i \cap V_j$. Since V_j is a critical subspace (see (33)), if $z \in V_j$, then

$$z = \sum_{i=1}^k c_i P_{E_i} z = \sum_{E_i \cap V_j \neq \{0\}} c_i P_{E_i \cap V_j} z = \sum_{E_i \cap V_j \neq \{0\}} \sum_{\alpha=1}^{n_{ij}} c_i \langle u_\alpha^{(ij)}, z \rangle u_\alpha^{(ij)}. \quad (40)$$

(40) shows that the system of all $u_1^{(ij)}, \dots, u_{n_{ij}}^{(ij)}$ when $E_i \cap V_j \neq \{0\}$ form a rank one Brascamp-Lieb data where the coefficient corresponding to $u_\alpha^{(ij)}$ is c_i .

According to (38), we have

$$P_{V_j} x = \sum_{E_i \cap V_j \neq \{0\}} \sum_{\alpha=1}^{n_{ij}} c_i \langle u_\alpha^{(ij)}, x_{ij} \rangle u_\alpha^{(ij)}.$$

We deduce from Case 1 applying to $P_{V_j}x$ to the rank one Brascamp-Lieb data in V_j above that

$$\begin{aligned} \|\Phi P_{V_j}x\|^2 &= \lambda_j^2 \|P_{V_j}x\|^2 \leq \lambda_j^2 \sum_{E_i \cap V_j \neq \{0\}} \sum_{\alpha=1}^{n_{ij}} c_i \langle u_\alpha^{(ij)}, x_{ij} \rangle^2 \\ &= \lambda_j^2 \sum_{E_i \cap V_j \neq \{0\}} c_i \|x_{ij}\|^2 = \sum_{E_i \cap V_j \neq \{0\}} c_i \|\Phi x_{ij}\|^2, \end{aligned}$$

proving (39), and in turn (37) that is equivalent to Lemma 24. \square

We now use Proposition 22 and Lemma 24 to exhibit the basic type of Gaussian extremizers of Barthe's inequality.

Proposition 25 *For linear subspaces E_1, \dots, E_k of \mathbb{R}^n , $n \geq 1$ and $c_1, \dots, c_k > 0$ satisfying (22), if Φ is a positive definite linear transform whose eigenspaces are critical subspaces, then*

$$\int_{\mathbb{R}^n}^* \left(\sup_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k e^{-c_i \|\Phi x_i\|^2} \right) dx = \prod_{i=1}^k \left(\int_{E_i} e^{-\|\Phi x_i\|^2} dx_i \right)^{c_i}.$$

Proof: Let $\tilde{\Phi} = \pi^{-\frac{1}{2}} \Phi$. For $i = 1, \dots, k$, let $A_i = \tilde{\Phi}|_{E_i}$, and hence $A_i : E_i \rightarrow E_i$ as the eigenspaces of $\tilde{\Phi}$ are critical subspaces. We deduce first using Lemma 24, and then the equality case of Proposition 22 that

$$\begin{aligned} \int_{\mathbb{R}^n}^* \left(\sup_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k e^{-c_i \|\Phi x_i\|^2} \right) dx &= \int_{\mathbb{R}^n} e^{-\pi \|\tilde{\Phi}x\|^2} dx = (\det \tilde{\Phi})^{-1} = \prod_{i=1}^k (\det A_i)^{-c_i} \\ &= \prod_{i=1}^k \left(\int_{E_i} e^{-\pi \|\tilde{\Phi}x_i\|^2} dx_i \right)^{c_i} = \prod_{i=1}^k \left(\int_{E_i} e^{-\|\Phi x_i\|^2} dx_i \right)^{c_i}, \end{aligned}$$

proving Proposition 25. \square

6 Splitting smooth extremizers along independent and dependent subspaces

Optimal transportation as a tool proving geometric inequalities was introduced by Gromov in his Appendix to [84] in the case of the Brunn-Minkowski inequality. Actually, Barthe's inequality in [8] was one of the first inequalities in probability, analysis or geometry that was obtained via optimal transportation.

We write $\nabla\Theta$ to denote the first derivative of a C^1 vector valued function Θ defined on an open subset of \mathbb{R}^n , and $\nabla^2\varphi$ to denote the Hessian of a real C^2 function φ . We recall that a vector valued function Θ on an open set $U \subset \mathbb{R}^n$ is C^α for $\alpha \in (0, 1)$ if for any $x_0 \in U$ there exist an open neighbourhood U_0 of x_0 and a $c_0 > 0$ such that $\|\Theta(x) - \Theta(y)\| \leq c_0\|x - y\|^\alpha$ for $x, y \in U_0$. In addition, a real function φ is $C^{2,\alpha}$ if φ is C^2 and $\nabla^2\varphi$ is C^α .

Combining Corollary 2.30, Corollary 2.32, Theorem 4.10 and Theorem 4.13 in Villani [94] on the Brenier map based on McCann [81, 82] for the first two, and on Caffarelli [27, 28, 29] for the last two theorems, we deduce the following:

Theorem 26 (Brenier, McCann, Caffarelli) *If f and g are positive C^α probability density functions on \mathbb{R}^n , $n \geq 1$, for $\alpha \in (0, 1)$, then there exists a $C^{2,\alpha}$ convex function φ on \mathbb{R}^n (unique up to additive constant) such that $T = \nabla\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is bijective and*

$$g(x) = f(T(x)) \cdot \det \nabla T(x) \text{ for } x \in \mathbb{R}^n. \quad (41)$$

Remarks The derivative $T = \nabla\varphi$ is the Brenier (transportation) map pushing forward the measure on \mathbb{R}^n induced by g to the measure associated to f ; namely, $\int_{T(X)} f = \int_X g$ for any measurable $X \subset \mathbb{R}^n$.

In addition, $\nabla T = \nabla^2\varphi$ is a positive definite symmetric matrix in Theorem 26, and if f and g are C^k for $k \geq 1$, then T is C^{k+1} .

Sometimes it is practical to consider the case $n = 0$, when we set $T : \{0\} \rightarrow \{0\}$ to be the trivial map.

Proof of Theorem 3 based on Barthe [8]. First we assume that each f_i is a C^1 positive probability density function on \mathbb{R}^n , and let us consider the Gaussian density $g_i(x) = e^{-\pi\|x\|^2}$ for $x \in E_i$. According to Theorem 26, if $i = 1, \dots, k$, then there exists a C^3 convex function φ_i on E_i such that for the C^2 Brenier map $T_i = \nabla\varphi_i$, we have

$$g_i(x) = \det \nabla T_i(x) \cdot f_i(T_i(x)) \text{ for all } x \in E_i. \quad (42)$$

It follows from the Remark after Theorem 26 that $\nabla T_i = \nabla^2\varphi_i(x)$ is positive definite symmetric matrix for all $x \in E_i$. For the C^2 transformation $\Theta : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$\Theta(y) = \sum_{i=1}^k c_i T_i(P_{E_i}y), \quad y \in \mathbb{R}^n, \quad (43)$$

its differential

$$\nabla\Theta(y) = \sum_{i=1}^k c_i \nabla T_i(P_{E_i}y)$$

is positive definite by Proposition 22. It follows that $\Theta : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is injective (see [8]), and

actually a diffeomorphism. Therefore Proposition 22, (42) and Lemma 18 (i) imply

$$\begin{aligned}
& \int_{\mathbb{R}^n}^* \sup_{x=\sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k f_i(x_i)^{c_i} dx \\
& \geq \int_{\mathbb{R}^n}^* \left(\sup_{\Theta(y)=\sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k f_i(x_i)^{c_i} \right) \det(\nabla\Theta(y)) dy \\
& \geq \int_{\mathbb{R}^n} \left(\prod_{i=1}^k f_i(T_i(P_{E_i}y))^{c_i} \right) \det\left(\sum_{i=1}^k c_i \nabla T_i(P_{E_i}y)\right) dy \\
& \geq \int_{\mathbb{R}^n} \left(\prod_{i=1}^k f_i(T_i(P_{E_i}y))^{c_i} \right) \prod_{i=1}^k (\det \nabla T_i(P_{E_i}y))^{c_i} dy \tag{44} \\
& = \int_{\mathbb{R}^n} \left(\prod_{i=1}^k g_i(P_{E_i}y)^{c_i} \right) dy = \int_{\mathbb{R}^n} e^{-\pi\|y\|^2} dy = 1.
\end{aligned}$$

Finally, Barthe's inequality (5) for arbitrary non-negative integrable functions f_i follows by scaling and approximation (see Barthe [8]). \square

We now prove that if equality holds in Barthe's inequality (5), then the diffeomorphism Θ in (43) in the proof of Barthe's inequality splits along the independent subspaces and the dependent subspace. First we explain how Barthe's inequality behaves under the shifts of the functions involved. Given proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), first we discuss in what sense Barthe's inequality is translation invariant. For non-negative integrable function f_i on E_i , $i = 1, \dots, k$, let us define

$$F(x) = \sup_{x=\sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k f_i(x_i)^{c_i}.$$

We observe that for any $e_i \in E_i$, defining $\tilde{f}_i(x) = f_i(x + e_i)$ for $x \in E_i$, $i = 1, \dots, k$, we have

$$\tilde{F}(x) = \sup_{x=\sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k \tilde{f}_i(x_i)^{c_i} = F\left(x + \sum_{i=1}^k c_i e_i\right). \tag{45}$$

Proposition 27 *For non-trivial linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), we write F_1, \dots, F_l to denote the independent subspaces (if exist), and F_0 to denote the dependent subspace (possibly $F_0 = \{0\}$). Let us assume that equality holds in (5) for positive C^1 probability densities f_i on E_i , $i = 1, \dots, k$, let $g_i(x) = e^{-\pi\|x\|^2}$ for $x \in E_i$, let $T_i : E_i \rightarrow E_i$ be the C^2 Brenier map satisfying*

$$g_i(x) = \det \nabla T_i(x) \cdot f_i(T_i(x)) \text{ for all } x \in E_i, \tag{46}$$

and let

$$\Theta(y) = \sum_{i=1}^k c_i T_i(P_{E_i}y), \quad y \in \mathbb{R}^n.$$

(i) For any $i \in \{1, \dots, k\}$ there exists positive C^1 integrable $h_{i0} : F_0 \cap E_i \rightarrow [0, \infty)$ (where $h_{i0}(0) = 1$ if $F_0 \cap E_i = \{0\}$), and for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$, there exists positive C^1 integrable $h_{ij} : F_j \rightarrow [0, \infty)$ such that

$$f_i(x) = h_{i0}(P_{F_0}x) \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} h_{ij}(P_{F_j}x) \quad \text{for } x \in E_i.$$

(ii) For $i = 1, \dots, k$, $T_i(E_i \cap F_p) = E_i \cap F_p$ whenever $E_i \cap F_p \neq \{0\}$ for $p \in \{0, \dots, l\}$, and if $x \in E_i$, then

$$T_i(x) = \bigoplus_{\substack{E_i \cap F_p \neq \{0\} \\ p \geq 0}} T_i(P_{F_p}x).$$

(iii) For $i = 1, \dots, k$, there exist C^2 functions $\Omega_i : E_i \rightarrow E_i$ and $\Gamma_i : E_i^\perp \rightarrow E_i^\perp$ such that

$$\Theta(y) = \Omega_i(P_{E_i}y) + \Gamma_i(P_{E_i^\perp}y) \quad \text{for } y \in \mathbb{R}^n.$$

(iv) If $y \in \mathbb{R}^n$, then the eigenspaces of the positive definite matrix $\nabla\Theta(y)$ are critical subspaces, and $\nabla T_i(P_{E_i}y) = \nabla\Theta(y)|_{E_i}$ for $i = 1, \dots, k$.

Proof: According to (45), we may assume that

$$T_i(0) = 0 \quad \text{for } i = 1, \dots, k, \tag{47}$$

If equality holds in (5), then equality holds in the determinantal inequality in (44) in the proof of Barthe's inequality; therefore, we apply the equality case of Proposition 22. In particular, for any $x \in \mathbb{R}^n$, there exist $m_x \geq 1$ and linear subspaces $V_{1,x}, \dots, V_{m_x,x}$ where $V_{1,x} = \mathbb{R}^n$ if $m_x = 1$, and $V_{1,x}, \dots, V_{m_x,x}$ are pairwise orthogonal indecomposable critical subspaces spanning \mathbb{R}^n if $m_x \geq 2$, and there exist $\lambda_{1,x}, \dots, \lambda_{m_x,x} > 0$ such that if $E_i \cap V_{j,x} \neq \{0\}$, then

$$\nabla T_i(P_{E_i}x)|_{E_i \cap V_{j,x}} = \lambda_{j,x} I_{E_i \cap V_{j,x}}; \tag{48}$$

and in addition, each E_i satisfies (cf. (28))

$$E_i = \bigoplus_{E_i \cap V_{j,x} \neq \{0\}} E_i \cap V_{j,x}. \tag{49}$$

Let us consider a fixed E_i , $i \in \{1, \dots, k\}$. First we claim that if $y \in E_i$, then

$$\begin{aligned} \nabla T_i(y)(F_p) &= F_p && \text{if } p \geq 1 \text{ and } E_i \cap F_p \neq \{0\} \\ \nabla T_i(y)(F_0 \cap E_i) &= F_0 \cap E_i. \end{aligned} \tag{50}$$

To prove (50), we take $y = x$ in (48). If $p \geq 1$ and $E_i \cap F_p \neq \{0\}$, then $F_p \subset E_i$, and Lemma 23 yields that

$$\begin{aligned}\bigoplus_{F_p \cap V_{j,y} \neq \{0\}} V_{j,y} &\subset F_p \\ \bigoplus_{F_p \cap V_{j,y} = \{0\}} V_{j,y} &\subset F_p^\perp.\end{aligned}$$

Since the subspaces $V_{j,y}$ span \mathbb{R}^n , we have

$$F_p = \bigoplus_{\substack{E_i \cap V_{j,y} \neq \{0\} \\ V_{j,y} \subset F_p}} V_{j,y};$$

therefore, (48) implies (50) if $p \geq 1$.

For the case of F_0 in (50), it follows from (49) and Lemma 23 that if $E_i \cap F_0 \neq \{0\}$, then

$$E_i \cap F_0 = \bigoplus_{\substack{E_i \cap V_{j,y} \neq \{0\} \\ V_{j,y} \subset F_0}} E_i \cap V_{j,y}. \quad (51)$$

Therefore, (48) completes the proof of (50).

It follows from (50) that if $E_i \cap F_p \neq \{0\}$, $y \in E_i$, $v \in E_i \cap F_p \cap S^{n-1}$ and $w \in E_i \cap F_p^\perp \cap S^{n-1}$, then

$$\left\langle v, \frac{\partial}{\partial t} T_i(y + tw) \Big|_{t=0} \right\rangle = 0. \quad (52)$$

In turn, (50), (52) and $T_i(0) = 0$ (cf. (47)) imply that if $y \in E_i$, then

$$T_i(E_i \cap F_p) = E_i \cap F_p \text{ whenever } E_i \cap F_p \neq \{0\} \text{ for } p \geq 0, \quad (53)$$

$$T_i(y) = \bigoplus_{\substack{E_i \cap F_p \neq \{0\} \\ p \geq 0}} T_i(P_{F_p} y). \quad (54)$$

We deduce from (54) that if $y \in E_i$, then

$$\det \nabla T_i(y) = \prod_{\substack{E_i \cap F_p \neq \{0\} \\ p \geq 0}} \det (\nabla T_i(P_{F_p} y)|_{F_p}). \quad (55)$$

We conclude (i) from (52), (53), (54), and (55) as (46) yields that if $y \in E_i$, then

$$f_i(T_i(y)) = \prod_{\substack{E_i \cap F_p \neq \{0\} \\ p \geq 0}} \frac{e^{-\pi \|P_{F_p} y\|^2}}{\det (\nabla T_i(P_{F_p} y)|_{F_p})}.$$

We deduce (ii) from (53) and (54).

For (iii), it follows from Proposition 22 that for any $x \in \mathbb{R}^n$, the spaces $V_{j,x}$ are eigenspaces for $\nabla \Theta(x)$ and span \mathbb{R}^n ; therefore, (27) implies that if $x \in \mathbb{R}^n$ and $i \in \{1, \dots, k\}$, then

$$\nabla \Theta(x) = \nabla \Theta(x)|_{E_i} \oplus \nabla \Theta(x)|_{E_i^\perp}.$$

Since $\Theta(0) = 0$ by (47), for fixed $i \in \{1, \dots, k\}$, we conclude

$$\begin{aligned}\Theta(E_i) &= E_i; \\ \Theta(x) &= \Theta(P_{E_i}x)|_{E_i} \oplus \Theta\left(P_{E_i^\perp}x\right)\Big|_{E_i^\perp} \quad \text{if } x \in \mathbb{R}^n.\end{aligned}$$

Finally, (iv) directly follows from Proposition 22, completing the proof of Proposition 27. \square

Next we show that if the extremizers f_1, \dots, f_k in Proposition 27 are of the form as in (i), then for any given $F_j \neq \{0\}$, the functions h_{ij} on F_j for all i with $E_i \cap F_j \neq \{0\}$ are also extremizers. We also need the Prékopa-Leindler inequality Theorem 28 (proved in various forms by Prékopa [86, 87], Leindler [70] and Borell [20]) whose equality case was clarified by Dubuc [38] (see the survey Gardner [45]). In turn, the Prékopa-Leindler inequality (56) is of the very similar structure like Barthe's inequality (5).

Theorem 28 (Prékopa, Leindler, Dubuc) *For $m \geq 2$, $\lambda_1, \dots, \lambda_m \in (0, 1)$ with $\lambda_1 + \dots + \lambda_m = 1$ and integrable $\varphi_1, \dots, \varphi_m : \mathbb{R}^n \rightarrow [0, \infty)$, we have*

$$\int_{\mathbb{R}^n} \sup_{x = \sum_{i=1}^m \lambda_i x_i, x_i \in \mathbb{R}^n} \prod_{i=1}^m \varphi_i(x_i)^{\lambda_i} dx \geq \prod_{i=1}^m \left(\int_{\mathbb{R}^n} \varphi_i \right)^{\lambda_i}, \quad (56)$$

and if equality holds and the left hand side is positive and finite, then there exist a log-concave function φ and $a_i > 0$ and $b_i \in \mathbb{R}^n$ for $i = 1, \dots, m$ such that

$$\varphi_i(x) = a_i \varphi(x - b_i)$$

for Lebesgue a.e. $x \in \mathbb{R}^n$, $i = 1, \dots, m$.

For linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), we assume that $F_{\text{dep}} \neq \mathbb{R}^n$, and write F_1, \dots, F_l to denote the independent subspaces. We verify that if $j \in \{1, \dots, l\}$, then

$$\sum_{E_i \supset F_j} c_i = 1. \quad (57)$$

For this, let $x \in F_j \setminus \{0\}$. We observe that for any E_i , either $F_j \subset E_i$, and hence $P_{E_i}x = x$, or $F_j \subset E_i^\perp$, and hence $P_{E_i}x = o$. We deduce from (1) that

$$x = \sum_{i=1}^k c_i P_{E_i}x = \left(\sum_{F_j \subset E_i} c_i \right) \cdot x,$$

which formula in turn implies (57).

Proposition 29 *For linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (1), we write F_1, \dots, F_l to denote the independent subspaces (if exist), and F_0 denote the dependent*

subspace (possibly $F_0 = \{0\}$). Let us assume that equality holds in Barthe's inequality (5) for probability densities f_i on E_i , $i = 1, \dots, k$, and for any $i \in \{1, \dots, k\}$ there exists integrable $h_{i0} : F_0 \cap E_i \rightarrow [0, \infty)$ (where $h_{i0}(0) = 1$ if $F_0 \cap E_i = \{0\}$), and for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$, there exists non-negative integrable $h_{ij} : F_j \rightarrow [0, \infty)$ such that

$$f_i(x) = h_{i0}(P_{F_0}x) \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} h_{ij}(P_{F_j}x) \quad \text{for } x \in E_i. \quad (58)$$

(i) If $F_0 \neq \{0\}$, then $\sum_{E_i \cap F_0 \neq \{0\}} c_i P_{E_i \cap F_0} = \text{Id}_{F_0}$ and

$$\int_{F_0}^* \sup_{x = \sum \{c_i x_i : x_i \in E_i \cap F_0 \text{ \& } E_i \cap F_0 \neq \{0\}\}} \prod_{E_i \cap F_0 \neq \{0\}} h_{i0}(x_i)^{c_i} dx = \prod_{E_i \cap F_0 \neq \{0\}} \left(\int_{E_i \cap F_0} h_{i0} \right)^{c_i}.$$

(ii) If $F_0 \neq \mathbb{R}^n$, then there exist integrable $\psi_j : F_j \rightarrow [0, \infty)$ for $j = 1, \dots, l$ where ψ_j is log-concave whenever $F_j \subset E_\alpha \cap E_\beta$ for $\alpha \neq \beta$, and there exist $a_{ij} > 0$ and $b_{ij} \in F_j$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$ such that $h_{ij}(x) = a_{ij} \cdot \psi_j(x - b_{ij})$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$.

Proof: We only present the argument in the case $F_0 \neq \mathbb{R}^n$ and $F_0 \neq \{0\}$. If $F_0 = \mathbb{R}^n$, then the same argument works ignoring the parts involving F_1, \dots, F_l , and if $F_0 = \{0\}$, then the same argument works ignoring the parts involving F_0 .

Since $F_0 \oplus F_1 \oplus \dots \oplus F_l = \mathbb{R}^n$ and F_0, \dots, F_l are critical subspaces, (28) yields for $i = 1, \dots, k$ that

$$E_i = (E_i \cap F_0) \oplus \bigoplus_{\substack{F_j \subset E_i \\ j \geq 1}} F_j; \quad (59)$$

therefore, the Fubini theorem and (58) imply that

$$\int_{E_i} f_i = \left(\int_{E_i \cap F_0} h_{i0} \right) \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} \int_{F_j} h_{ij}. \quad (60)$$

On the other hand, using again $F_0 \oplus F_1 \oplus \dots \oplus F_l = \mathbb{R}^n$, we deduce that if $x = \sum_{j=0}^l z_j$ where $z_j \in F_j$ for $j \geq 0$, then $z_j = P_{F_j}x$. It follows from (59) that for any $x \in \mathbb{R}^n$, we have

$$\begin{aligned} \sup_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i)^{c_i} &= \left(\sup_{\substack{P_{F_0}x = \sum_{i=1}^k c_i x_{0i} \\ x_{0i} \in E_i \cap F_0}} \prod_{i=1}^k h_{i0}(x_{0i}) \right) \times \\ &\quad \times \prod_{j=1}^l \left(\sup_{\substack{P_{F_j}x = \sum_{F_j \subset E_i} c_i x_{ji} \\ x_{ji} \in F_j}} \prod_{F_j \subset E_i} h_{ij}(x_{ji})^{c_i} \right), \end{aligned}$$

and hence

$$\int_{\mathbb{R}^n}^* \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i)^{c_i} dx = \left(\int_{F_0}^* \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i \cap F_0}} \prod_{i=1}^k h_{i0}(x_i) dx \right) \times \quad (61)$$

$$\times \prod_{j=1}^l \left(\int_{F_j}^* \sup_{\substack{x=\sum_{F_j \subset E_i} c_i x_i, \\ x_i \in F_j}} \prod_{F_j \subset E_i} h_{ij}(x_i)^{c_i} dx \right).$$

As F_0 is a critical subspace, we have

$$\sum_{i=1}^k c_i P_{E_i \cap F_0} = \text{Id}_{F_0},$$

and hence Barthe's inequality (5) yields

$$\int_{F_0}^* \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i \cap F_0}} \prod_{i=1}^k h_{i0}(x_i) dx \geq \prod_{i=1}^k \left(\int_{E_i \cap F_0} h_{i0} \right)^{c_i}. \quad (62)$$

We deduce from (57) and the Prékopa-Leindler inequality (56) that if $j = 1, \dots, l$, then

$$\int_{F_j}^* \sup_{\substack{x=\sum_{F_j \subset E_i} c_i x_i, \\ x_i \in F_j}} \prod_{F_j \subset E_i} h_{ij}(x_i)^{c_i} dx \geq \prod_{E_i \supset F_j} \left(\int_{F_j} h_{ij} \right)^{c_i}. \quad (63)$$

Combining (60), (61), (62) and (63) with the fact that f_1, \dots, f_k are extremizers for Barthe's inequality (5) implies that if $j = 1, \dots, l$, then

$$\int_{F_0}^* \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i \cap F_0}} \prod_{i=1}^k h_{i0}(x_i) dx = \prod_{i=1}^k \left(\int_{E_i \cap F_0} h_{i0} \right)^{c_i} \quad (64)$$

$$\int_{F_j}^* \sup_{\substack{x=\sum_{F_j \subset E_i} c_i x_i, \\ x_i \in F_j}} \prod_{F_j \subset E_i} h_{ij}(x_i)^{c_i} dx = \prod_{E_i \supset F_j} \left(\int_{F_j} h_{ij} \right)^{c_i}. \quad (65)$$

We observe that (64) is just (i). In addition, (ii) follows from the equality conditions in the Prékopa-Leindler inequality (see Theorem 28). \square

7 Convolution and product of extremizers

Given proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), we say that the non-negative integrable functions f_1, \dots, f_k with positive integrals are extremizers if

equality holds in (5). In order to deal with positive smooth functions, we use convolutions. More precisely, Lemma 2 in Barthe [8] states the following.

Lemma 30 *Given proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), if f_1, \dots, f_k and g_1, \dots, g_k are extremizers in Barthe's inequality (5), then $f_1 * g_1, \dots, f_k * g_k$ are also extremizers.*

Proof: We define

$$F(x) = \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k f_i(x_i)^{c_i}$$

$$G(y) = \sup_{y = \sum_{i=1}^k c_i y_i, y_i \in E_i} \prod_{i=1}^k g_i(y_i)^{c_i}.$$

Possibly F and G are not measurable but as f_1, \dots, f_k and g_1, \dots, g_k are extremizers, there exist measurable $\tilde{F} \geq F$ and $\tilde{G} \geq G$ such that $\int_{\mathbb{R}^n} \tilde{F}(x) dx = \int_{\mathbb{R}^n} \tilde{G}(x) dx = 1$. We deduce that

$$\begin{aligned} \int_{\mathbb{R}^n} \tilde{F} * \tilde{G}(x) dx &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \tilde{F}(x-y) \tilde{G}(y) dy dx = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \tilde{F}(x-y) \tilde{G}(y) dx dy \\ &= \int_{\mathbb{R}^n} \tilde{G}(y) \left(\int_{\mathbb{R}^n} \tilde{F}(x-y) dx \right) dy = \int_{\mathbb{R}^n} \tilde{G}(y) \cdot 1 dy = 1. \end{aligned} \quad (66)$$

We deduce writing $x_i = z_i + y_i$ in (67) for $i = 1, \dots, k$ and using Barthe's inequality in (68) that

$$\begin{aligned} 1 &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \tilde{F}(x-y) \tilde{G}(y) dy dx \\ &\geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{x-y = \sum_{i=1}^k c_i z_i, z_i \in E_i} \prod_{i=1}^k f_i(z_i)^{c_i} \sup_{y = \sum_{i=1}^k c_i y_i, y_i \in E_i} \prod_{i=1}^k g_i(y_i)^{c_i} dy dx \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{x-y = \sum_{i=1}^k c_i z_i, z_i \in E_i} \sup_{y = \sum_{i=1}^k c_i y_i, y_i \in E_i} \prod_{i=1}^k f_i(z_i)^{c_i} \prod_{i=1}^k g_i(y_i)^{c_i} dy dx \end{aligned} \quad (67)$$

$$\begin{aligned} &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \sup_{y = \sum_{i=1}^k c_i y_i, y_i \in E_i} \prod_{i=1}^k f_i(x_i - y_i)^{c_i} \prod_{i=1}^k g_i(y_i)^{c_i} dy dx \\ &\geq \int_{\mathbb{R}^n} \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \int_{\mathbb{R}^n} \sup_{y = \sum_{i=1}^k c_i y_i, y_i \in E_i} \prod_{i=1}^k (f_i(x_i - y_i) g_i(y_i))^{c_i} dy dx \end{aligned} \quad (68)$$

$$\begin{aligned} &\geq \int_{\mathbb{R}^n} \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k \left(\int_{E_i} f_i(x_i - y_i) g_i(y_i) dy_i \right)^{c_i} dx \\ &= \int_{\mathbb{R}^n} \sup_{x = \sum_{i=1}^k c_i x_i, x_i \in E_i} \prod_{i=1}^k (f_i * g_i(x_i))^{c_i} dx \end{aligned}$$

Since for $i = 1, \dots, k$, $\int_{E_i} f_i * g_i = 1$ can be proved similarly to (66), we conclude that $f_i * g_i$, $i = 1, \dots, k$, is also an extremizer. \square

Since in a certain case we want to work with Lebesgue integral instead of outer integrals, we use the following statement that can be proved via compactness argument.

Lemma 31 *Given proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), if h_i is a positive continuous functions satisfying $\lim_{x \rightarrow \infty} h_i(x) = 0$ for $i = 1, \dots, k$, then the function*

$$h(x) = \sup_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k h_i(x_i)^{c_i}$$

of $x \in \mathbb{R}^n$ is continuous.

Next we show that the product of a shift of a smooth extremizer and a Gaussian is also an extremizer for Barthe's inequality.

Lemma 32 *Given proper linear subspaces E_1, \dots, E_k of \mathbb{R}^n and $c_1, \dots, c_k > 0$ satisfying (22), if f_1, \dots, f_k are positive bounded C^1 are extremizers in Barthe's inequality (5), and $g_i(x) = e^{-\pi\|x\|^2}$ for $x \in E_i$, then there exist $z_i \in E_i$, $i = 1, \dots, k$, such that the functions $y \mapsto f_i(y - z_i)g_i(y)$ of $y \in E_i$, $i = 1, \dots, k$, are also extremizers for (5).*

Proof: We may assume that f_1, \dots, f_k are probability densities.

Readily the functions $\tilde{f}_1, \dots, \tilde{f}_k$ defined by $\tilde{f}_i(y) = f_i(-y)$ for $y \in E_i$ and $i = 1, \dots, k$ are also extremizers. We deduce from Lemma 30 that the functions $\tilde{f}_i * g_i$ for $i = 1, \dots, k$ are also extremizers where each $\tilde{f}_i * g_i$ is a probability density on E_i . According to Theorem 26, if $i = 1, \dots, k$, then there exists a C^2 Brenier map $S_i : E_i \rightarrow E_i$ such that

$$g_i(x) = \det \nabla S_i(x) \cdot (\tilde{f}_i * g_i)(S_i(x)) \text{ for all } x \in E_i,$$

and $\nabla S_i(x)$ is a positive definite symmetric matrix for all $x \in E_i$. As in the proof of Theorem 3 above, we consider the C^2 diffeomorphism $\Theta : \mathbb{R}^n \rightarrow \mathbb{R}^n$ given by

$$\Theta(y) = \sum_{i=1}^k c_i S_i(P_{E_i} y), \quad y \in \mathbb{R}^n.$$

whose positive definite differential is

$$\nabla \Theta(y) = \sum_{i=1}^k c_i \nabla S_i(P_{E_i} y).$$

On the one hand, we note that if $x = \sum_{i=1}^k c_i x_i$ for $x_i \in E_i$, then

$$\|x\|^2 \leq \sum_{i=1}^k c_i \|x_i\|^2$$

holds according to Barthe [8]; or equivalently,

$$\prod_{i=1}^k g_i(x_i)^{c_i} \leq e^{-\pi\|x\|^2}.$$

Since f_i is positive, bounded, continuous and in $L_1(E_i)$ for $i = 1, \dots, k$, we observe that the function

$$z \mapsto \int_{\mathbb{R}^n} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - S_i(P_{E_i} \Theta^{-1} z))^{c_i} g_i(x_i)^{c_i} dx \quad (69)$$

of $z \in \mathbb{R}^n$ is continuous.

Using also that $\tilde{f}_1, \dots, \tilde{f}_k$ are extremizers and probability density functions, we have

$$\begin{aligned} & \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{\substack{z=\sum_{i=1}^k c_i z_i, \\ z_i \in E_i}} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - z_i)^{c_i} g_i(x_i)^{c_i} dx dz \\ &= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \left(\prod_{i=1}^k g_i(x_i)^{c_i} \right) \sup_{\substack{z=\sum_{i=1}^k c_i z_i, \\ z_i \in E_i}} \prod_{i=1}^k f_i(x_i - z_i)^{c_i} dz dx \\ &\leq \int_{\mathbb{R}^n} e^{-\pi\|x\|^2} \int_{\mathbb{R}^n} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \sup_{\substack{z=\sum_{i=1}^k c_i z_i, \\ z_i \in E_i}} \prod_{i=1}^k f_i(x_i - z_i)^{c_i} dz dx \\ &= \int_{\mathbb{R}^n} e^{-\pi\|x\|^2} \int_{\mathbb{R}^n} \sup_{\substack{z-x=\sum_{i=1}^k c_i y_i, \\ y_i \in E_i}} \prod_{i=1}^k \tilde{f}_i(y_i)^{c_i} dz dx \\ &= \int_{\mathbb{R}^n} e^{-\pi\|x\|^2} \int_{\mathbb{R}^n} \sup_{\substack{w=\sum_{i=1}^k c_i y_i, \\ y_i \in E_i}} \prod_{i=1}^k \tilde{f}_i(y_i)^{c_i} dw dx \\ &= \int_{\mathbb{R}^n} e^{-\pi\|x\|^2} dx = 1. \end{aligned}$$

Using Lemma 31 and (69) in (70), Barthe's inequality (5) in (71) and Proposition 22 in (72),

we deduce that

$$\begin{aligned}
1 &\geq \int_{\mathbb{R}^n}^* \int_{\mathbb{R}^n}^* \sup_{\substack{z=\sum_{i=1}^k c_i z_i \\ z_i \in E_i}} \sup_{\substack{x=\sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - z_i)^{c_i} g_i(x_i)^{c_i} dx dz \\
&\geq \int_{\mathbb{R}^n}^* \int_{\mathbb{R}^n}^* \sup_{\substack{x=\sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - S_i(P_{E_i}\Theta^{-1}z))^{c_i} g_i(x_i)^{c_i} dx dz \tag{70}
\end{aligned}$$

$$= \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - S_i(P_{E_i}\Theta^{-1}z))^{c_i} g_i(x_i)^{c_i} dx dz \tag{71}$$

$$\begin{aligned}
&\geq \int_{\mathbb{R}^n} \prod_{i=1}^k \left(\int_{E_i} f_i(x_i - S_i(P_{E_i}\Theta^{-1}z)) g_i(x_i) dx_i \right)^{c_i} dz \\
&= \int_{\mathbb{R}^n} \prod_{i=1}^k (\tilde{f}_i * g_i)(S_i(P_{E_i}\Theta^{-1}z))^{c_i} dz \\
&= \int_{\mathbb{R}^n} \left(\prod_{i=1}^k (\tilde{f}_i * g_i)(S_i(P_{E_i}y))^{c_i} \right) \det(\nabla\Theta(y)) dy \\
&= \int_{\mathbb{R}^n} \left(\prod_{i=1}^k (\tilde{f}_i * g_i)(S_i(P_{E_i}y))^{c_i} \right) \det\left(\sum_{i=1}^k c_i \nabla S_i(P_{E_i}y)\right) dy \tag{72} \\
&\geq \int_{\mathbb{R}^n} \left(\prod_{i=1}^k (\tilde{f}_i * g_i)(S_i(P_{E_i}y))^{c_i} \right) \prod_{i=1}^k (\det \nabla S_i(P_{E_i}y))^{c_i} dy \\
&= \int_{\mathbb{R}^n} \left(\prod_{i=1}^k g_i(P_{E_i}y)^{c_i} \right) dy = \int_{\mathbb{R}^n} e^{-\pi\|y\|^2} dy = 1.
\end{aligned}$$

In particular, we conclude that

$$\begin{aligned}
1 &\geq \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \sup_{\substack{x=\sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - S_i(P_{E_i}\Theta^{-1}z))^{c_i} g_i(x_i)^{c_i} dx dz \\
&\geq \int_{\mathbb{R}^n} \prod_{i=1}^k \left(\int_{E_i} f_i(x_i - S_i(P_{E_i}\Theta^{-1}z)) g_i(x_i) dx_i \right)^{c_i} dz \geq 1.
\end{aligned}$$

Because of Barthe's inequality (5), it follows from (69) that

$$\begin{aligned} & \int_{\mathbb{R}^n} \sup_{\substack{x = \sum_{i=1}^k c_i x_i, \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i - S_i(P_{E_i} \Theta^{-1} z))^{c_i} g_i(x_i)^{c_i} dx \\ &= \prod_{i=1}^k \left(\int_{E_i} f_i(x_i - S_i(P_{E_i} \Theta^{-1} z)) g_i(x_i) dx_i \right)^{c_i} \end{aligned}$$

for any $z \in \mathbb{R}^n$; therefore, we may choose $z_i = S_i(0)$ for $i = 1, \dots, k$ in Lemma 32. \square

8 h_{i0} is Gaussian in Proposition 27

For positive C^α probability density functions f and g on \mathbb{R}^n for $\alpha \in (0, 1)$, the C^1 Brenier map $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ in Theorem 26 pushing forward the the measure on \mathbb{R}^n induced by g to the measure associated to f satisfies that ∇T is positive definite. We deduce that

$$\langle T(y) - T(x), y - x \rangle = \int_0^1 \langle \nabla T(x + t(y - x)) \cdot (y - x), y - x \rangle dt \geq 0 \quad \text{for any } x, y \in \mathbb{R}^n. \quad (73)$$

We say that a continuous function $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ has linear growth if there exists a positive constant $c > 0$ such that

$$\|T(x)\| \leq c\sqrt{1 + \|x\|^2}$$

for $x \in \mathbb{R}^n$. It is equivalent saying that

$$\limsup_{\|x\| \rightarrow \infty} \frac{\|T(x)\|}{\|x\|} < \infty. \quad (74)$$

In general, T has polynomial growth, if there exists $k \geq 1$ such that

$$\limsup_{\|x\| \rightarrow \infty} \frac{\|T(x)\|}{\|x\|^k} < \infty.$$

Proposition 33 related to Caffarelli Contraction Principle in Caffarelli [30] was proved by Emanuel Milman, see for example Colombo, Fathi [35], De Philippis, Figalli [37], Fathi, Gozlan, Prod'homme [41], Y.-H. Kim, E. Milman [62], Klartag, Putterman [66], Kolesnikov [67], Livshyts [72] for relevant results.

Proposition 33 (Emanuel Milman) *If a Gaussian probability density g and a positive C^α , $\alpha \in (0, 1)$, probability density f on \mathbb{R}^n satisfy $f \leq c \cdot g$ for some positive constant $c > 0$, then the Brenier map $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ pushing forward the measure on \mathbb{R}^n induced by g to the measure associated to f has linear growth.*

Proof: We may assume that $g(x) = e^{-\pi\|x\|^2}$.

We observe that $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is bijective as both f and g are positive. Let S be the inverse of T ; namely, $S : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the bijective Brenier map pushing forward the measure on \mathbb{R}^n induced by f to the measure associated to g . In particular, any Borel $X \subset \mathbb{R}^n$ satisfies

$$\int_{S(X)} g = \int_X f. \quad (75)$$

We note that (74), and hence Proposition 33 is equivalent saying that

$$\liminf_{x \rightarrow \infty} \frac{\|S(x)\|}{\|x\|} > 0. \quad (76)$$

The main idea of the argument is the following observation. For any unit vector u and $\theta \in (0, \pi)$, we consider

$$\Xi(u, \theta) = \{y : \langle y, u \rangle \geq \|y\| \cdot \cos \theta\}.$$

Since S is surjective, and $\langle S(z) - S(w), z - w \rangle \geq 0$ for any $z, w \in \mathbb{R}^n$ according to (73), we deduce that

$$S(w) + \Xi(u, \theta) \subset S\left(w + \Xi\left(u, \theta + \frac{\pi}{2}\right)\right) \quad (77)$$

for any $u \in S^{n-1}$ and $\theta \in (0, \frac{\pi}{2})$.

We suppose that T does not have linear growth, and seek a contradiction. According to (76), there exists a sequence $\{x_k\}$ of points of $\mathbb{R}^n \setminus \{0\}$ tending to infinity such that

$$\lim_{k \rightarrow \infty} \|x_k\| = \infty \quad \text{and} \quad \lim_{k \rightarrow \infty} \frac{\|S(x_k)\|}{\|x_k\|} = 0.$$

In particular, we may assume that

$$\|S(x_k)\| < \frac{\|x_k\|}{8}. \quad (78)$$

For any k , we consider the unit vector $e_k = x_k / \|x_k\|$. We observe that $X_k = x_k + \Xi(e_k, \frac{3\pi}{4})$ avoids the interior of the ball $\frac{\|x_k\|}{\sqrt{2}} B^n$; therefore, if k is large, then

$$\int_{X_k} f \leq c \cdot n \kappa_n \int_{\|x_k\|/\sqrt{2}}^{\infty} r^{n-1} e^{-\pi r^2} dr < \int_{\|x_k\|/\sqrt{2}}^{\infty} e^{-2r^2} \sqrt{2} r dr = e^{-\|x_k\|^2} \quad (79)$$

On the other hand, $S(x_k) + \Xi(e_k, \frac{\pi}{4})$ contains the ball

$$\tilde{B} = S(x_k) + \frac{x_k}{8} + \frac{\|x_k\|}{8\sqrt{2}} B^n \subset \frac{\|x_k\|}{2} B^n$$

where we have used (78). It follows from (75) and (77) that if k is large, then

$$\int_{X_k} f = \int_{S(X_k)} g \geq \int_{\tilde{B}} g \geq \kappa_n \left(\frac{\|x_k\|}{8\sqrt{2}}\right)^n e^{-\pi(\|x_k\|/2)^2} > e^{-\|x_k\|^2}.$$

This inequality contradicts (79), and in turn proves (76). \square

Proposition 36 shows that if the whole space is the dependent subspace and the Brenier maps corresponding to the extremizers f_1, \dots, f_k in Proposition 27 have at most linear growth, then each f_i is actually Gaussian. The proof of Proposition 36 uses classical Fourier analysis, and we refer to Grafakos [50] for the main properties. For our purposes, we need only the action of a tempered distribution on the space of $C_0^\infty(\mathbb{R}^m)$ of C^∞ functions with compact support, do not need to consider the space of Schwarz functions in general. We recall that if u is a tempered distribution on Schwarz functions on \mathbb{R}^n , then the support $\text{supp } u$ is the intersection of all closed sets K such that if $\varphi \in C_0^\infty(\mathbb{R}^n)$ with $\text{supp } \varphi \subset \mathbb{R}^n \setminus K$, then $\langle u, \varphi \rangle = 0$. We write \hat{u} to denote the Fourier transform of a u . In particular, if θ is a function of polynomial growth and $\varphi \in C_0^\infty(\mathbb{R}^n)$, then

$$\langle \hat{\theta}, \varphi \rangle = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \theta(x) \varphi(y) e^{-2\pi i \langle x, y \rangle} dx dy. \quad (80)$$

We consider the two well-known statements Lemma 34 and Lemma 35 about the support of a Fourier transform to prepare the proof of Proposition 36.

Lemma 34 *If θ is a measurable function of polynomial growth on \mathbb{R}^n , and there exist linear subspace E with $1 \leq \dim E \leq n - 1$ and function ω on E such that $\theta(x) = \omega(P_E x)$, then $\text{supp } \hat{\theta} \subset E$.*

Proof: We write a $z \in \mathbb{R}^n$ in the form $z = (z_1, z_2)$ with $z_1 \in E$ and $z_2 \in E^\perp$. Let $\varphi \in C_0^\infty(\mathbb{R}^n)$ satisfy that $\text{supp } \varphi \subset \mathbb{R}^n \setminus E$, and hence $\varphi(x_1, 0) = 0$ for $x_1 \in E$, and the Fourier Integral Theorem in E^\perp implies

$$\varphi(x_1, z) = \int_{E^\perp} \int_{E^\perp} \varphi(x_1, x_2) e^{2\pi i \langle z - x_2, y_2 \rangle} dx_2 dy_2$$

for $x_1 \in E$ and $z \in E^\perp$. It follows from (80) that

$$\begin{aligned} \langle \hat{\theta}, \varphi \rangle &= \int_{E^\perp} \int_E \int_{E^\perp} \int_E \omega(x_1) \varphi(x_1, x_2) e^{-2\pi i \langle x_1, y_1 \rangle} e^{-2\pi i \langle x_2, y_2 \rangle} dx_1 dx_2 dy_1 dy_2 \\ &= \int_E \int_E \omega(x_1) e^{-2\pi i \langle x_1, y_1 \rangle} \left(\int_{E^\perp} \int_{E^\perp} \varphi(x_1, x_2) e^{2\pi i \langle -x_2, y_2 \rangle} dx_2 dy_2 \right) dy_1 dx_1 \\ &= \int_E \int_E \omega(x_1) e^{-2\pi i \langle x_1, y_1 \rangle} \varphi(x_1, 0) dy_1 dx_1 = 0. \quad \square \end{aligned}$$

Next, Lemma 35 directly follows from Proposition 2.4.1 in Grafakos [50].

Lemma 35 *If θ is a continuous function of polynomial growth on \mathbb{R}^n and $\text{supp } \hat{\theta} \subset \{0\}$, then θ is a polynomial.*

Proposition 36 For linear subspaces E_1, \dots, E_k of \mathbb{R}^m and $c_1, \dots, c_k > 0$ satisfying (1), we assume that

$$\bigcap_{i=1}^k (E_i \cup E_i^\perp) = \{0\}. \quad (81)$$

Let $g_i(x) = e^{-\pi\|x\|^2}$ for $i = 1, \dots, k$ and $x \in E_i$, let equality hold in (5) for positive C^1 probability densities f_i on E_i , $i = 1, \dots, k$, and let $T_i : E_i \rightarrow E_i$ be the C^2 Brenier map satisfying

$$g_i(x) = \det \nabla T_i(x) \cdot f_i(T_i(x)) \text{ for all } x \in E_i. \quad (82)$$

If each T_i , $i = 1, \dots, k$, has linear growth, then there exist a positive definite matrix $A : \mathbb{R}^m \rightarrow \mathbb{R}^m$ whose eigenspaces are critical subspaces, and $a_i > 0$ and $b_i \in E_i$, $i = 1, \dots, k$, such that

$$f_i(x) = a_i e^{-\langle Ax, x + b_i \rangle} \text{ for } x \in E_i.$$

Proof: We may assume that each linear subspace is non-zero.

We note that the condition (81) is equivalent saying that \mathbb{R}^m itself is the dependent subspace with respect to the Brascamp-Lieb data. We may assume that for some $1 \leq l \leq k$, we have $1 \leq \dim E_i \leq m - 1$ if $i = 1, \dots, l$, and still

$$\bigcap_{i=1}^l (E_i \cup E_i^\perp) = \{0\}. \quad (83)$$

We use the diffeomorphism $\Theta : \mathbb{R}^m \rightarrow \mathbb{R}^m$ of Proposition 27 defined by

$$\Theta(y) = \sum_{i=1}^k c_i T_i(P_{E_i} y), \quad y \in \mathbb{R}^m.$$

It follows from (45) that we may assume

$$T_i(0) = 0 \text{ for } i = 1, \dots, k, \text{ and hence } \Theta(0) = 0. \quad (84)$$

We claim that there exists a positive definite matrix $B : \mathbb{R}^m \rightarrow \mathbb{R}^m$ whose eigenspaces are critical subspaces, and

$$\nabla \Theta(y) = B \text{ for } y \in \mathbb{R}^m. \quad (85)$$

Let $\Theta(y) = (\theta_1(y), \dots, \theta_m(y))$ for $y \in \mathbb{R}^m$ and $\theta_j \in C^2(\mathbb{R}^m)$, $j = 1, \dots, m$. Since each T_i , $i = 1, \dots, k$ has linear growth, it follows that Θ has linear growth, and in turn each θ_j , $j = 1, \dots, m$, has linear growth.

According to Proposition 27 (iii), there exist C^2 functions $\Omega_i : E_i \rightarrow E_i$ and $\Gamma_i : E_i^\perp \rightarrow E_i^\perp$ such that

$$\Theta(y) = \Omega_i(P_{E_i} y) + \Gamma_i(P_{E_i^\perp} y)$$

for $i = 1, \dots, k$ and $y \in \mathbb{R}^n$. We write $\Omega_i(x) = (\omega_{i1}(x), \dots, \omega_{im}(x))$ and $\Gamma_i(x) = (\gamma_{i1}(x), \dots, \gamma_{im}(x))$; therefore,

$$\theta_j(y) = \omega_{ij}(P_{E_i} y) + \gamma_{ij}(P_{E_i^\perp} y) \quad (86)$$

for $j = 1, \dots, m$ and $i = 1, \dots, k$.

Fix a $j \in \{1, \dots, m\}$. It follows from Lemma 34 and (86) that

$$\text{supp } \hat{\theta}_j \subset E_i \cup E_i^\perp$$

for $i = 1, \dots, l$. Thus (83) yields that

$$\text{supp } \hat{\theta}_j \subset \{0\},$$

and in turn we deduce from Lemma 35 that θ_j is a polynomial. Given that θ_j has linear growth, it follows that there exist $w_j \in \mathbb{R}^m$ and $\alpha_j \in \mathbb{R}$ such that $\theta_j(y) = \langle w_j, y \rangle + \alpha_j$. We deduce from $\theta_j(o) = 0$ (cf. (84)) that $\alpha_j = 0$.

The argument so far yields that there exists an $m \times m$ matrix B such that $\Theta(y) = By$ for $y \in \mathbb{R}^m$. As $\nabla\Theta(y) = B$ is positive definite and its eigenspaces are critical subspaces, we conclude the claim (85).

Since $\nabla T_i(P_{E_i}y) = \nabla\Theta(y)|_{E_i}$ for $i = 1, \dots, k$ and $y \in \mathbb{R}^m$ by Proposition 27 (iv), we deduce that $T_i^{-1} = B^{-1}|_{E_i}$ for $i = 1, \dots, k$. It follows from (82) that

$$f_i(x) = e^{-\pi\|B^{-1}x\|^2} \cdot \det(B^{-1}|_{E_i}) \quad \text{for } x \in E_i$$

for $i = 1, \dots, k$. Therefore, we can choose $A = \pi B^{-2}$. \square

9 Proof of Theorem 4

We may assume that each linear subspace E_i is non-zero in Theorem 4. Let f_i be a probability density on E_i in a way such that equality holds for f_1, \dots, f_k in (5). For $i = 1, \dots, k$ and $x \in E_i$, let $g_i(x) = e^{-\pi\|x\|^2}$, and hence g_i is a probability distribution on E_i , and g_1, \dots, g_k are extremizers in Barthe's inequality (5).

It follows from Lemma 30 that the convolutions $f_1 * g_1, \dots, f_k * g_k$ are also extremizers for (5). We observe that for $i = 1, \dots, k$, $f_i * g_i$ is a bounded positive C^∞ probability density on E_i . Next we deduce from Lemma 32 that there exist $z_i \in E_i$ and $\gamma_i > 0$ for $i = 1, \dots, k$ such that defining

$$\tilde{f}_i(x) = \gamma_i \cdot g_i(x) \cdot (f_i * g_i)(x - z_i) \quad \text{for } x \in E_i,$$

$\tilde{f}_1, \dots, \tilde{f}_k$ are probability densities that are extremizers for (5). We note that if $i = 1, \dots, k$, then \tilde{f}_i is positive and C^∞ , and there exists $c > 1$ satisfying

$$\tilde{f}_i \leq c \cdot g_i. \tag{87}$$

Let $\tilde{T}_i : E_i \rightarrow E_i$ be the C^∞ Brenier map satisfying

$$g_i(x) = \det \nabla \tilde{T}_i(x) \cdot \tilde{f}_i(\tilde{T}_i(x)) \quad \text{for all } x \in E_i, \tag{88}$$

We deduce from (87) and Proposition 33 that \tilde{T}_i has linear growth.

For $i = 1, \dots, k$ and $x \in F_0 \cap E_i$, let $g_{i0}(x) = e^{-\pi\|x\|^2}$. It follows from Proposition 27 (i) that for $i \in \{1, \dots, k\}$, there exists positive C^1 integrable $h_{i0} : F_0 \cap E_i \rightarrow [0, \infty)$ (where $h_{i0}(o) = 1$ if $F_0 \cap E_i = \{0\}$), and for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$, there exists positive C^1 integrable $\tilde{h}_{ij} : F_j \rightarrow [0, \infty)$ such that

$$\tilde{f}_i(x) = \tilde{h}_{i0}(P_{F_0}x) \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} \tilde{h}_{ij}(P_{F_j}x) \quad \text{for } x \in E_i.$$

We deduce from Proposition 27 (ii) that $\tilde{T}_{i0} = \tilde{T}_i|_{F_0 \cap E_i}$ is the Brenier map pushing forward the measure on $F_0 \cap E_i$ determined g_{i0} onto the measure determined by \tilde{h}_{i0} . Since \tilde{T}_i has linear growth, \tilde{T}_{i0} has linear growth, as well, for $i = 1, \dots, k$.

We deduce from Proposition 29 (i) that $\sum_{i=1}^k c_i P_{E_i \cap F_0} = \text{Id}_{F_0}$, the Geometric Brascamp Lieb data $E_1 \cap F_0, \dots, E_k \cap F_0$ in F_0 has no independent subspaces, and $\tilde{h}_{10}, \dots, \tilde{h}_{k0}$ are extremizers in Barthe's inequality for this data in F_0 .

As \tilde{T}_{i0} has linear growth for $i = 1, \dots, k$, Proposition 36 yields the existence of a positive definite matrix $\tilde{A} : F_0 \rightarrow F_0$ whose eigenspaces are critical subspaces, and $\tilde{a}_i > 0$ and $\tilde{b}_i \in F_0 \cap E_i$ for $i = 1, \dots, k$, such that

$$\tilde{f}_i(x) = \tilde{a}_i e^{-\langle \tilde{A}x, x + \tilde{b}_i \rangle} \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} \tilde{h}_{ij}(P_{F_j}x) \quad \text{for } x \in E_i.$$

Dividing by g_i and shifting, we deduce that there exist a symmetric matrix $\bar{A} : F_0 \rightarrow F_0$ whose eigenspaces are critical subspaces, and $\bar{a}_i > 0$ and $\bar{b}_i \in F_0 \cap E_i$ for $i = 1, \dots, k$, and for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$, there exists positive C^1 $\bar{h}_{ij} : F_j \rightarrow [0, \infty)$ such that

$$f_i * g_i(x) = \bar{a}_i e^{-\langle \bar{A}x, x + \bar{b}_i \rangle} \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} \bar{h}_{ij}(P_{F_j}x) \quad \text{for } x \in E_i.$$

Since $f_i * g_i$ is a probability density on E_i , it follows that \bar{A} is positive definite and $\bar{h}_{ij} \in L_1(E_i \cap F_j)$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$.

For any $i = 1, \dots, k$, we write $\hat{\varrho}$ for the Fourier transform of a function $\varrho \in L_1(E_i)$, thus we can take the inverse Fourier transform in the sense that ϱ is a.e. the L_1 limit of

$$x \mapsto \int_{\mathbb{R}^n} \hat{\varrho}(\xi) e^{-a|\xi|^2} e^{2\pi i \langle \xi, x \rangle} d\xi$$

as $a > 0$ tends to zero. For $i = 1, \dots, k$, using that $\widehat{f_i * g_i} = \hat{f}_i \cdot \hat{g}_i$, we deduce that the restriction of \hat{f}_i to $F_0 \cap E_i$ is the quotient of two Gaussian densities. Since f_i is bounded and zero at infinity, we deduce that the restriction of \hat{f}_i to $F_0 \cap E_i$ is a Gaussian density for $i = 1, \dots, k$, as well, with the symmetric matrix involved being positive definite. We conclude using the inverse Fourier transform above and the fact that the linear subspaces F_j , $j = 0, \dots, l$, are pairwise orthogonal that there exist a symmetric matrix $A : F_0 \rightarrow F_0$ whose eigenspaces are critical subspaces, and

$a_i > 0$ and $b_i \in F_0 \cap E_i$ for $i = 1, \dots, k$, and for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$, there exists $h_{ij} : F_j \rightarrow [0, \infty)$ such that

$$f_i(x) = a_i e^{-\langle Ax, x + b_i \rangle} \cdot \prod_{\substack{F_j \subset E_i \\ j \geq 1}} h_{ij}(P_{F_j}x) \quad \text{for a.e. } x \in E_i.$$

Since f_i is a probability density on E_i , it follows that A is positive definite and each h_{ij} is non-negative and integrable. Finally, Proposition 29 (ii) yields that there exist integrable $\psi_j : F_j \rightarrow [0, \infty)$ for $j = 1, \dots, l$ where ψ_j is log-concave whenever $F_j \subset E_\alpha \cap E_\beta$ for $\alpha \neq \beta$, and there exist $a_{ij} > 0$ and $b_{ij} \in F_j$ for any $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$ such that $h_{ij}(x) = a_{ij} \cdot \psi_j(x - b_{ij})$ for $i \in \{1, \dots, k\}$ and $j \in \{1, \dots, l\}$ with $F_j \subset E_i$.

Finally, we assume that f_1, \dots, f_k are of the form as described in (6) and equality holds for all $x \in E_i$ in (6). According to (45), we may assume that there exist a positive definite matrix $\Phi : F_0 \rightarrow F_0$ whose proper eigenspaces are critical subspaces and a $\tilde{\theta}_i > 0$ for $i = 1, \dots, k$ such that

$$f_i(x) = \tilde{\theta}_i e^{-\|\Phi P_{F_0}x\|^2} \prod_{F_j \subset E_i} h_j(P_{F_j}(x)) \quad \text{for } x \in E_i. \quad (89)$$

We recall that according to (57), if $j \in \{1, \dots, l\}$, then

$$\sum_{E_i \supset F_j} c_i = 1. \quad (90)$$

We set $\theta = \prod_{i=1}^k \tilde{\theta}_i^{c_i}$ and $h_0(x) = e^{-\|\Phi x\|^2}$ for $x \in F_0$. On the left hand side of Barthe's inequality (5), we use first (90) and the log-concavity of h_j whenever $j \geq 1$ and $F_j \subset E_\alpha \cap E_\beta$ for $\alpha \neq \beta$, secondly Proposition 25, thirdly (90), fourth the Fubini Theorem, and finally (90) again to prove that

$$\begin{aligned} \int_{\mathbb{R}^n}^* \sup_{\substack{x = \sum_{i=1}^k c_i x_i \\ x_i \in E_i}} \prod_{i=1}^k f_i(x_i)^{c_i} dx &= \theta \int_{\mathbb{R}^n}^* \sup_{\substack{x = \sum_{i=1}^k \sum_{j=0}^l c_i x_{ij} \\ x_{ij} \in E_i \cap F_j}} \prod_{j=0}^l \prod_{i=1}^k h_j(x_{ij})^{c_i} dx \\ &= \theta \int_{\mathbb{R}^n}^* \prod_{j=0}^l \sup_{\substack{P_{F_j}x = \sum_{i=1}^k c_i x_{ij} \\ x_{ij} \in E_i \cap F_j}} \prod_{i=1}^k h_j(x_{ij})^{c_i} dx \\ &= \theta \int_{\mathbb{R}^n}^* \left(\sup_{\substack{P_{F_0}x = \sum_{i=1}^k c_i x_{i0} \\ x_{i0} \in E_i \cap F_0}} \prod_{i=1}^k e^{-c_i \|\Phi x_{i0}\|^2} \right) \times \prod_{j=1}^l h_j(P_{F_j}x) dx \\ &= \theta \left(\prod_{i=1}^k \left(\int_{F_0 \cap E_i} e^{-\|\Phi y\|^2} dy \right)^{c_i} \right) \times \prod_{j=1}^l \int_{F_j} h_j \\ &= \prod_{i=1}^k \left(\int_{E_i} f_i \right)^{c_i}, \end{aligned}$$

completing the proof of Theorem 4. \square

10 Equality in the Bollobas-Thomason inequality and in its dual

We fix an orthonormal basis e_1, \dots, e_n of \mathbb{R}^n for the whole section. We set $\sigma_i^0 = \sigma_i$ and $\sigma_i^1 = [n] \setminus \sigma_i$. When we write $\tilde{\sigma}_1, \dots, \tilde{\sigma}_\ell$ for the induced cover from $\sigma_1, \dots, \sigma_k$, we assume that the sets $\tilde{\sigma}_1, \dots, \tilde{\sigma}_\ell$ are pairwise distinct.

Lemma 37 *For $s \geq 1$, let $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$, and let $\tilde{\sigma}_1, \dots, \tilde{\sigma}_\ell$ be the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$. Then*

(i) *the subspaces $E_{\sigma_i} := \text{lin}\{e_j : j \in \sigma_i\}$ satisfy*

$$\sum_{i=1}^k \frac{1}{s} P_{E_{\sigma_i}} = I_n \quad (91)$$

i.e. form a Geometric Brascamp Lieb data;

(ii) *For $r \in \tilde{\sigma}_j$, $j = 1, \dots, \ell$, we have*

$$\tilde{\sigma}_j := \bigcap_{r \in \sigma_i} \sigma_i^0 \cap \bigcap_{r \notin \sigma_i} \sigma_i^1; \quad (92)$$

(iii) *the subspaces $F_{\tilde{\sigma}_j} := \text{lin}\{e_r : r \in \tilde{\sigma}_j\}$ are the independent subspaces of the Geometric Brascamp Lieb data (91) and $F_{\text{dep}} = \{0\}$.*

Proof: Since $\sigma_1, \dots, \sigma_k$ form a s -uniform cover, every $e_i \in \mathbb{R}^n$ is contained in exactly s of $E_{\sigma_1}, \dots, E_{\sigma_k}$, yielding (i).

For (ii), the definition of $\tilde{\sigma}_j$ directly implies (92).

For (iii), the linear subspaces $F_{\tilde{\sigma}_1}, \dots, F_{\tilde{\sigma}_\ell}$ are pairwise orthogonal because $\sigma_i^0 \cap \sigma_i^1 = \emptyset$ for $i = 1, \dots, k$. On the other hand, for any $r \in [n]$, $r \in \bigcap_{i=1}^n \sigma_i^{\varepsilon(i)}$ where $\varepsilon(i) = 0$ if $r \in \sigma_i$, and $\varepsilon(i) = 1$ if $r \notin \sigma_i$; therefore, $F_{\tilde{\sigma}_1}, \dots, F_{\tilde{\sigma}_\ell}$ span \mathbb{R}^n . In particular, $F_{\text{dep}} = \{0\}$. \square

Let us introduce the notation that we use when handling both the Bollobas-Thomason inequality and its dual. Let $\sigma_1, \dots, \sigma_k$ be the s cover of $[n]$ occurring in Theorem 11 and Theorem 12, and hence $E_i = E_{\sigma_i}$, $i = 1, \dots, k$, satisfies

$$\frac{1}{s} \sum_{i=1}^k P_{E_{\sigma_i}} = I_n. \quad (93)$$

Let $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ be the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$. It follows that

$$F_j = E_{\tilde{\sigma}_j} \text{ for } j = 1, \dots, l \text{ are the independent subspaces,} \quad (94)$$

$$F_{\text{dep}} = \{0\}. \quad (95)$$

For any $i \in \{1, \dots, k\}$, we set

$$I_i = \{j \in \{1, \dots, l\} : F_j \subset E_i\},$$

and for any $j \in \{1, \dots, l\}$, we set

$$J_j = \{i \in \{1, \dots, k\} : F_j \subset E_i\}.$$

For the reader's convenience, we restate Theorem 9 and Theorem 11 as Theorem 38, and Theorem 10 and Theorem 12 as Theorem 39.

Theorem 38 *If $K \subset \mathbb{R}^n$ is compact and affinely spans \mathbb{R}^n , and $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$, then*

$$|K|^s \leq \prod_{i=1}^k |P_{E_{\sigma_i}} K|. \quad (96)$$

Equality holds if and only if $K = \bigoplus_{i=1}^l P_{F_{\tilde{\sigma}_i}} K$ where $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ is the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$ and $F_{\tilde{\sigma}_i}$ is the linear hull of the e_i 's with indices from $\tilde{\sigma}_i$.

Proof: We set $E_i := E_{\sigma_i}$ which subspaces compose a geometric data according to Lemma 37. We start with a proof of Bollobas-Thomason inequality. It follows directly from the Brascamp-Lieb inequality as

$$\begin{aligned} |K| &= \int_{\mathbb{R}^n} 1_K(x) dx \leq \int_{\mathbb{R}^n} \prod_{i=1}^k 1_{P_{E_i}(K)}(P_{E_i}(x))^{\frac{1}{s}} dx \\ &\leq \prod_{i=1}^k \left(\int_{E_i} 1_{P_{E_i}(K)} \right)^{\frac{1}{s}} = \prod_{i=1}^k |P_{E_i}(K)|^{\frac{1}{s}} \end{aligned} \quad (97)$$

where the first inequality is from the monotonicity of the integral while the second is Brascamp-Lieb inequality Theorem 1. Now, if equality holds in (97), then on the one hand,

$$1_K(x) = \prod_{i=1}^k 1_{P_{E_i}(K)}(P_{E_i}(x))$$

and on the other hand, if F_1, \dots, F_l are the independent subspaces of the data, then they span \mathbb{R}^n according to Lemma 37; namely, $F_{\text{dep}} = \{0\}$. It follows from Theorem 2 that there exist integrable functions $h_j : F_j \rightarrow \mathbb{R}$, such that, for Lebesgue a.e. $x_i \in E_i$

$$1_{P_{E_i}K}(x_i) = \theta_i \prod_{j \in I_i} h_j(P_{F_j}(x_i))$$

Therefore from the previous two, we have for $x \in \mathbb{R}^n$

$$1_K(x) = \prod_{i=1}^k \theta_i \prod_{j \in I_i} h_j(P_{F_j}(P_{E_i}(x)))$$

Now, since for $j \in I_i$ we have $F_j \subset E_i$ we can delete the P_{E_i} on the above product. Thus, for $\theta = \prod_{i=1}^k \theta_i$, we have for Lebesgue a.e. $x \in \mathbb{R}^n$

$$1_K(x) = \theta \prod_{i=1}^k \prod_{j \in I_i} h_j(P_{F_j}(x)) = \theta \prod_{j=1}^l h_j(P_{F_j}(x))^{|J_j|}. \quad (98)$$

Now, for $x \in K$ the last product on above is constant, so

$$\theta = \frac{1}{\prod_{i=1}^l h_j(P_{F_j}(x_0))^{|J_j|}} \quad (99)$$

for some $x_0 \in K$. For $j = 1, \dots, l$ we set $\varphi_j : F_j \rightarrow \mathbb{R}^n$, by

$$\varphi_j(x) = \frac{h_j(x + P_{F_j}(x_0))^{|J_j|}}{h_j(P_{F_j}(x_0))^{|J_j|}}.$$

We see that $\varphi_j(o) = 1$ and also (98) and (99) yields

$$1_{K-x_0}(x) = \prod_{j=1}^l \varphi_j(P_{F_j}(x)) \quad (100)$$

For $m \in \{1, \dots, l\}$, taking $x \in F_m$ in (100) (and hence $\varphi_j(P_{F_j}(x)) = 1$ for $j \neq m$) shows that

$$1_{K-x_0}(y) = \varphi_m(y),$$

for Lebesgue a.e. $y \in F_m$. Therefore (100) and the orthogonality of the F_j 's,

$$K - x_0 = \bigcap_{j=1}^l P_{F_j}^{-1}(P_{F_j}(K - x_0)) = \bigoplus_{j=1}^l P_{F_j}(K - x_0),$$

completing the proof of Theorem 38. □

To prove Theorem 39, we use two small observations. First if M is any convex body with $o \in \text{int } M$, then

$$\int_{\mathbb{R}^n} e^{-\|x\|_M} dx = \int_0^\infty e^{-r} n r^{n-1} |M| dr = n! |M|. \quad (101)$$

Secondly, if F_j are pairwise orthogonal subspaces and $M = \text{conv} \{M_1, \dots, M_l\}$ where $M_j \subset F_j$ is a $\dim F_j$ -dimensional compact convex set with $o \in \text{relint } M_j$, then for any $x \in \mathbb{R}^n$

$$\|x\|_M = \sum_{i=1}^l \|P_{F_j} x\|_{M_j}. \quad (102)$$

In addition, we often use the fact, for a subspace F of \mathbb{R}^n and $x \in F$, then $\|x\|_K = \|x\|_{K \cap F}$.

Theorem 39 *If $K \subset \mathbb{R}^n$ is compact convex with $o \in \text{int}K$, and $\sigma_1, \dots, \sigma_k \subset [n]$ form an s -uniform cover of $[n]$ for $s \geq 1$, then*

$$|K|^s \geq \frac{\prod_{i=1}^k |\sigma_i|!}{(n!)^s} \cdot \prod_{i=1}^k |K \cap E_{\sigma_i}|. \quad (103)$$

Equality holds if and only if $K = \text{conv}\{E_{\tilde{\sigma}_i} \cap K\}_{i=1}^l$ where $\tilde{\sigma}_1, \dots, \tilde{\sigma}_l$ is the 1-uniform cover of $[n]$ induced by $\sigma_1, \dots, \sigma_k$.

Proof: We define

$$f(x) = e^{-\|x\|_K}, \quad (104)$$

which is a log-concave function with $f(o) = 1$, and satisfying (cf (101))

$$\int_{\mathbb{R}^n} f(y)^n dy = \int_{\mathbb{R}^n} e^{-n\|y\|_K} dy = \int_{\mathbb{R}^n} e^{-\|y\|_{\frac{1}{n}K}} dy = n! \left| \frac{1}{n}K \right| = \frac{n!}{n^n} \cdot |K|. \quad (105)$$

We claim that

$$n^n \int_{\mathbb{R}^n} f(y)^n dy \geq \prod_{i=1}^k \left(\int_{E_i} f(x_i) dx_i \right)^{1/s}. \quad (106)$$

Equating the traces of the two sides of (91), we deduce that, $d_i := |\sigma_i| = \dim E_i$

$$\sum_{i=1}^k \frac{d_i}{sn} = 1. \quad (107)$$

For $z = \sum_{i=1}^k \frac{1}{s} x_i$ with $x_i \in E_i$, the log-concavity of f and its definition (104), imply

$$f(z/n) \geq \prod_{i=1}^k f(x_i/d_i)^{\frac{d_i}{ns}} = \prod_{i=1}^k f(x_i)^{\frac{1}{ns}}. \quad (108)$$

Now, the monotonicity of the integral and Barthe's inequality yield

$$\int_{\mathbb{R}^n} f(z/n)^n dz \geq \int_{\mathbb{R}^n}^* \sup_{z = \sum_{i=1}^k \frac{1}{s} x_i, x_i \in E_i} \prod_{i=1}^k f(x_i)^{1/s} dz \geq \prod_{i=1}^k \left(\int_{E_i} f(x_i) dx_i \right)^{1/s}. \quad (109)$$

Making the change of variable $y = z/n$ we conclude to (106). Computing the right hand side of (106), we have

$$\int_{E_i} f(x_i) dx_i = \int_{E_i} e^{-\|x_i\|_K} dx_i = \int_{E_i} e^{-\|x_i\|_{K \cap E_i}} dx_i = d_i! |K \cap E_i|. \quad (110)$$

Therefore, (105), (106) and (110) yield (103).

Let us assume that equality holds in (103), and hence we have two equalities in (109). We set

$$M = \text{conv}\{K \cap F_j\}_{1 \leq j \leq l}.$$

Clearly, $K \supseteq M$. For the other inclusion, we start with $z \in \text{int}K$, namely $\|z\|_K < 1$. Equality in the first inequality in (109) means,

$$(e^{-\|z/n\|_K})^n = \sup_{z = \sum_{i=1}^k \frac{1}{s} x_i, x_i \in E_i} \prod_{i=1}^k e^{-\|x_i\|_K 1/s},$$

or in other words,

$$\|z\|_K = \frac{1}{s} \cdot \inf_{z = \sum_{i=1}^k \frac{1}{s} x_i, x_i \in E_i} \sum_{i=1}^k \|x_i\|_K = \inf_{z = \sum_{i=1}^k y_i, y_i \in E_i} \sum_{i=1}^k \|y_i\|_K. \quad (111)$$

We deduce that there exist $y_i \in E_i$, $i = 1, \dots, k$ such that

$$z = \sum_{i=1}^k y_i \quad \text{and} \quad \sum_{i=1}^k \|y_i\|_K < 1, \quad (112)$$

Therefore, from (112), then (102) and after the triangle inequality for $\|\cdot\|_{K \cap F_j}$, we have

$$\|z\|_M = \left\| \sum_{i=1}^k \sum_{j \in I_i} P_{F_j} y_i \right\|_M = \sum_{i=1}^k \left\| \sum_{j \in I_i} P_{F_j} y_i \right\|_{K \cap F_j} \leq \sum_{i=1}^k \sum_{j \in I_i} \|P_{F_j} y_i\|_{K \cap F_j}. \quad (113)$$

It suffices to show that

$$K \cap E_i = \text{conv}\{K \cap F_j\}_{j \in I_i} \quad (114)$$

because then, from (113), applying (102) and (112), we have

$$\|z\|_M \leq \sum_{j=1}^l \sum_{i \in J_j} \|P_{F_j} y_i\|_{K \cap F_j} = \sum_{i=1}^k \|y_i\|_{K \cap E_i} < 1,$$

which means $z \in M$. Now, to show (114), we start with the equality case of Barthe's inequality which has been applied in (109). From Theorem 4, there exist $\theta_i > 0$ and $w_i \in E_i$ and log-concave $h_j : F_j \rightarrow [0, \infty)$, namely $h_j = e^{-\varphi_j}$ for a convex function φ_j , such that

$$e^{-\|x_i\|_{K \cap E_i}} = \theta_i \prod_{j \in I_i} h_j(P_{F_j}(x_i - w_i)). \quad (115)$$

for Lebesgue a.e. $x_i \in E_i$. For $i \in [k]$ and $j \in I_i$ we set, $\psi_{ij} : F_j \rightarrow \mathbb{R}$ by

$$\psi_{ij}(x) = \varphi_j(x - P_{F_j} w_i) - \varphi_j(-P_{F_j} w_i) + \frac{\ln \theta_i}{|I_i|}.$$

We see

$$\psi_{ij}(o) = 0 \text{ and } \psi_{ij} \text{ is convex on } F_j. \quad (116)$$

and also (115) yields, for $x \in E_i$

$$e^{-\|x\|_{K \cap E_i}} = \exp \left(- \sum_{j \in I_i} \psi_{ij}(P_{F_j}x) \right). \quad (117)$$

For $x \in F_j$, we apply λx to (117) with $\lambda > 0$, and we have from $\psi_{im}(o) = 0$ for $m \in I_i \setminus \{j\}$ that

$$\psi_{ij}(\lambda x) = \lambda \psi_{ij}(x) \text{ and } \psi_{ij}(x) > 0. \quad (118)$$

We deduce from (116) and (118) that ψ_{ij} is a norm. Therefore, $\psi_{ij}(x) = \|x\|_{C_{ij}}$ for some $(\dim F_j)$ -dimensional compact convex set $C_{ij} \subset F_j$ with $o \in \text{relint } C_{ij}$. Now (117) becomes,

$$\|x\|_{K \cap E_i} = \sum_{j \in I_i} \|P_{F_j}x\|_{C_{ij}}$$

and hence by (102) we conclude to

$$K \cap E_i = \text{conv} \{C_{ij}\}_{j \in I_i}.$$

In particular, if $i \in [k]$ and $j \in I_i$, then $C_{ij} = (K \cap E_i) \cap F_j = K \cap F_j$, completing the proof of (114), and in turn yielding Theorem 12. \square

Acknowledgements We thank Alessio Figalli, Greg Kuperberg and Christos Saroglou for helpful discussions. We are especially grateful to Emanuel Milman for providing the proof of Proposition 33, and for Franck Barthe for providing the proof of Proposition 22 and insight on the history of the subject, and for further ideas and extremely helpful discussions. We thank the referee for correcting a mistake in Theorem 4 and significantly improving the presentation of the whole paper.

The first named author is also grateful for the hospitality and excellent working environment provided by University of California, Davis and by ETH Zürich during various parts of this project.

References

- [1] D. Alonso-Gutiérrez, J. Bernués, S. Brazitikos, A. Carbery: On affine invariant and local Loomis-Whitney type inequalities. arXiv:2002.05794
- [2] D. Alonso-Gutiérrez, S. Brazitikos: Reverse Loomis-Whitney inequalities via isotropicity. arXiv:2001.11876

- [3] K.M. Ball: Volumes of sections of cubes and related problems. In: J. Lindenstrauss and V.D. Milman (ed), Israel seminar on Geometric Aspects of Functional Analysis 1376, Lectures Notes in Mathematics. Springer-Verlag, 1989.
- [4] K.M. Ball: Volume ratios and a reverse isoperimetric inequality. *J. London Math. Soc.* 44 (1991), 351–359
- [5] K.M. Ball: Convex geometry and functional analysis. In: W B. Johnson, L. Lindenstrauss (eds), Handbook of the geometry of Banach spaces, 1, (2003), 161–194.
- [6] Z. Balogh, A. Kristály: Equality in Borell-Brascamp-Lieb inequalities on curved spaces. *Adv. Math.* 339 (2018), 453-494.
- [7] F. Barthe: Inégalités de Brascamp-Lieb et convexité. *C. R. Acad. Sci. Paris* 324 (1997), 885–888.
- [8] F. Barthe: On a reverse form of the Brascamp-Lieb inequality. *Invent. Math.* 134 (1998), 335–361.
- [9] F. Barthe: A continuous version of the Brascamp-Lieb inequalities. *Geometric Aspects of Functional Analysis, Lecture Notes in Mathematics Volume 1850*, 2004, 53–63.
- [10] F. Barthe, D. Cordero-Erausquin: Inverse Brascamp-Lieb inequalities along the heat equation. *Geometric aspects of functional analysis*, 65-71, *Lecture Notes in Math.*, 1850, Springer, Berlin, 2004.
- [11] F. Barthe, D. Cordero-Erausquin, M. Ledoux, B. Maurey: Correlation and Brascamp-Lieb inequalities for Markov semigroups. *Int. Math. Res. Not.* 10 (2011), 2177–2216.
- [12] F. Barthe, N. Huet: On Gaussian Brunn-Minkowski inequalities. *Studia Math.* 191 (2009), 283–304.
- [13] F. Barthe, P. Wolff: Positivity improvement and Gaussian kernels. *C. R. Math. Acad. Sci. Paris*, 352 (2014), 1017–1021.
- [14] F. Barthe, P. Wolff: Positive Gaussian Kernels also Have Gaussian Minimizers. *Mem. Amer. Math. Soc.*, 276 (2022), no. 1359, iii+90 pp.
- [15] J. Bennett, N. Bez, T.C. Flock, S. Lee: Stability of the Brascamp–Lieb constant and applications. *Am. J. Math.* 140(2) (2018), 543-569.
- [16] J. Bennett, N. Bez, S. Buschenhenke, M.G. Cowling, T.C. Flock: On the nonlinear Brascamp-Lieb inequality. *Duke Math. J.* 169(17) (2020), 3291-3338
- [17] J. Bennett, T. Carbery, M. Christ, T. Tao: The Brascamp–Lieb Inequalities: Finiteness, Structure and Extremals. *Geom. Funct. Anal.* 17 (2008), 1343–1415.

- [18] B. Bollobas, A. Thomason: Projections of bodies and hereditary properties of hypergraphs. *Bull. Lond. Math. Soc.* 27, (1995), 417–424.
- [19] S.G. Bobkov, A. Colesanti, I. Fragalà: Quermassintegrals of quasi-concave functions and generalized Prékopa-Leindler inequalities. *Manuscripta Math.*, 143 (2014), 131-169.
- [20] C. Borell: The Brunn-Minkowski inequality in Gauss spaces. *Invent. Math.* 30 (1975), 207–216.
- [21] K.J. Böröczky, M. Henk: Cone volume measure and stability. [arXiv:1407.7272](https://arxiv.org/abs/1407.7272).
- [22] H.J. Brascamp, E.H. Lieb: Best constants in Young’s inequality, its converse, and its generalization to more than three functions. *Adv. Math.* 20 (1976), 151-173.
- [23] S. Brazitikos, A. Giannopoulos, P. Valettas, B.-H. Vritsiou: *Geometry of isotropic convex bodies. Mathematical Surveys and Monographs 196*, American Mathematical Society, Providence, RI, 2014.
- [24] S. Brazitikos, S. Dann, A. Giannopoulos, A. Koldobsky: On the average volume of sections of convex bodies. *Israel J. Math.* 222 (2017), 921–947.
- [25] S. Brazitikos, A. Giannopoulos, D-M. Liakopoulos: Uniform cover inequalities for the volume of coordinate sections and projections of convex bodies. *Adv. Geom.* 18 (2018), 345–354.
- [26] J.R. Bueno, P. Pivarov: A stochastic Prékopa-Leindler inequality for log-concave functions. *Commun. Contemp. Math.*, 23 (2021), no. 2, Paper No. 2050019, 17 pp.
- [27] L.A. Caffarelli: A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity. *Ann. of Math. (2)* 131, (1990), 129-134.
- [28] L.A. Caffarelli: Interior $W^{2,p}$ estimates for solutions of the Monge-Ampère equation. *Ann. of Math. (2)*, 131 (1990), 135-150.
- [29] L.A. Caffarelli: The regularity of mappings with a convex potential. *J. Amer. Math. Soc.*, 5 (1992), 99-104.
- [30] L.A. Caffarelli: Monotonicity properties of optimal transportation and the FKG and related inequalities. *Comm. Math. Phys.* 214 (2000), no. 3, 547-563.
- [31] E. Carlen, E.H. Lieb, M. Loss: A sharp analog of Young’s inequality on S^N and related entropy inequalities. *J. Geom. Anal.*, 14 (2004), 487-520.
- [32] E. Carlen, D. Cordero-Erausquin: Subadditivity of the entropy and its relation to Brascamp-Lieb type inequalities. *Geom. Funct. Anal.*, 19 (2009), 373-405.
- [33] P.G. Casazza, T.T. Tran, J.C. Tremain: Regular two-distance sets. *J. Fourier Anal. Appl.*, 26 (2020), no. 3, Paper No. 49, 32 pp.

- [34] W-K. Chen, N. Dafnis, G. Paouris: Improved Hölder and reverse Hölder inequalities for Gaussian random vectors. *Adv. Math.* 280 (2015), 643–689.
- [35] M. Colombo, M. Fathi: Bounds on optimal transport maps onto log-concave measures. *J. Differential Equations* 271 (2021), 1007-1022.
- [36] T.A. Courtade, J. Liu: Euclidean forward-reverse Brascamp-Lieb inequalities: finiteness, structure, and extremals. *J. Geom. Anal.*, 31 (2021), 3300–3350.
- [37] G. De Philippis, A. Figalli: Rigidity and stability of Caffarelli’s log-concave perturbation theorem. *Nonlinear Anal.* 154 (2017), 59-70.
- [38] S. Dubuc: Critères de convexité et inégalités intégrales. *Ann. Inst. Fourier Grenoble*, 27 (1) (1977), 135–165.
- [39] L. Dümbgen: Bounding standard Gaussian tail probabilities. arXiv:1012.2063v3
- [40] J. Duncan: An algebraic Brascamp-Lieb inequality. *J. Geom. Anal.* 31 (2021), 10136-10163.
- [41] M. Fathi, N. Gozlan, M. Prod’homme: A proof of the Caffarelli contraction theorem via entropic regularization. *Calc. Var. Partial Differential Equations* 59 (2020), no. 3, Paper No. 96, 18 pp.
- [42] A. Figalli, F. Maggi, A. Pratelli: A refined Brunn-Minkowski inequality for convex sets. *Annales de IHP (C) Non Linear Analysis* 26 (2009), 2511–2519.
- [43] A. Figalli, F. Maggi, A. Pratelli: A mass transportation approach to quantitative isoperimetric inequalities. *Invent. Math.* 182 (2010), 167–211.
- [44] N. Fusco, F. Maggi, A. Pratelli: The sharp quantitative isoperimetric inequality. *Ann. of Math.* 168 (2008), 941–980.
- [45] R. Gardner: The Brunn-Minkowski inequality. *Bull. Amer. Math. Soc.* 39 (2002), 355–405.
- [46] D. Ghilli, P. Salani: Quantitative Borell-Brascamp-Lieb inequalities for power concave functions. *J. Convex Anal.* 24 (2017), 857-888.
- [47] A. Giannopoulos, V. Milman: Extremal problems and isotropic positions of convex bodies. *Israel J. Math.* 117 (2000), 29–60.
- [48] A. Giannopoulos, V. Milman: Euclidean structure in finite dimensional normed spaces. *Handbook of the geometry of Banach spaces*, Vol. I, 707-779, North-Holland, Amsterdam, 2001.
- [49] A. Giannopoulos, M. Papadimitrakis: Isotropic surface area measures. *Mathematika* 46 (1999), 1-13

- [50] L. Grafakos: Classical Fourier analysis. Graduate Texts in Mathematics, 249. Springer, 2014.
- [51] H. Groemer: Stability properties of geometric inequalities. *Amer. Math. Monthly* 97 (1990), no. 5, 382–394.
- [52] H. Groemer: Stability of geometric inequalities. *Handbook of convex geometry*, Vol. A, B, 125–150, North-Holland, Amsterdam, 1993.
- [53] H. Groemer, R. Schneider: Stability estimates for some geometric inequalities. *Bull. London Math. Soc.* 23 (1991), no. 1, 67–74.
- [54] P.M. Gruber: Convex and discrete geometry. *Grundlehren der Mathematischen Wissenschaften*, Springer, Berlin, 2007.
- [55] P.M. Gruber, F.E. Schuster: An arithmetic proof of John’s ellipsoid theorem. *Arch. Math.* 85 (2005), 82–88.
- [56] B. Grünbaum: Partitions of mass-distributions and of convex bodies by hyperplanes. *Pacific J. Math.* 10 (1960), 1257–1261.
- [57] O. Guedon, E. Milman: Interpolating thin-shell and sharp large-deviation estimates for isotropic log-concave measures. *Geom. Funct. Anal.* 21 (2011), 1043–1068.
- [58] S. Guo, R. Zhang: On integer solutions of Parsell-Vinogradov systems. *Invent. Math.* 218 (2019), 1–81.
- [59] W. Gustin: An isoperimetric minimax. *Pacific J. Math.* 3 (1953), 403–405.
- [60] F. John: Polar correspondence with respect to a convex region. *Duke Math. J.* 3 (1937), 355–369.
- [61] R. Kannan, L. Lovász, M. Simonovits: Isoperimetric problems for convex bodies and a localization lemma. *Discrete Comput. Geom.* 13 (1995), 541–559.
- [62] Y.-H. Kim, E. Milman: A generalization of Caffarelli’s contraction theorem via (reverse) heat flow. *Math. Ann.* 354 (2012), no. 3, 827–862.
- [63] B. Klartag: A Berry-Esseen type inequality for convex bodies with an unconditional basis. *Probab. Theory Related Fields* **145** (2009), 1–33.
- [64] B. Klartag: On nearly radial marginals of high-dimensional probability measures. *J. Eur. Math. Soc.* 12 (2010), 723–754.
- [65] B. Klartag, E. Milman: Centroid bodies and the logarithmic Laplace transform—a unified approach. *J. Funct. Anal.* 262 (2012), 10–34.

- [66] B. Klartag, E. Putterman: Spectral monotonicity under Gaussian convolution. arXiv:2107.09496
- [67] A.V. Kolesnikov: On Sobolev regularity of mass transport and transportation inequalities. *Theory Probab. Appl.* 57 (2013), no. 2, 243-264.
- [68] A.V. Kolesnikov, E. Milman: Local L_p -Brunn-Minkowski inequalities for $p < 1$. *Memoirs AMS*, accepted. arXiv:1711.01089
- [69] J. Lehec: Short probabilistic proof of the Brascamp-Lieb and Barthe theorems. *Canad. Math. Bull.*, 57 (2014), 585-597.
- [70] L. Leindler: On a certain converse of Hölder's inequality. II. *Acta Sci. Math. (Szeged)* 33 (1972), 217–223.
- [71] Liakopoulos, D.-M.: Reverse Brascamp-Lieb inequality and the dual Bollobás-Thomason inequality. *Arch. Math. (Basel)* 112 (2019), 293–304.
- [72] G.V. Livshyts: Some remarks about the maximal perimeter of convex sets with respect to probability measures. *Commun. Contemp. Math.* 23 (2021), no. 5, Paper No. 2050037, 19 pp.
- [73] G.V. Livshyts: On a conjectural symmetric version of Ehrhard's inequality. arXiv:2103.11433
- [74] L.H. Loomis, H. Whitney: An inequality related to the isoperimetric inequality, *Bull. Amer. Math. Soc.*, 55 (1949), 961–962.
- [75] E.H. Lieb: Gaussian kernels have only Gaussian maximizers. *Invent. Math.* 102 (1990), 179–208.
- [76] S. Campi, R. Gardner, P. Gronchi: Reverse and dual Loomis-Whitney-type inequalities. *Trans. Amer. Math. Soc.*, 368 (2016), 5093–5124.
- [77] E. Lutwak, D. Yang, G. Zhang: Volume inequalities for subspaces of L_p . *J. Diff. Geom.*, 68 (2004), 159-184.
- [78] E. Lutwak, D. Yang, G. Zhang: Volume inequalities for isotropic measures. *Amer. J. Math.*, 129 (2007), 1711-1723.
- [79] D. Maldague: Regularized Brascamp–lieb Inequalities And An Application. *The Quarterly Journal of Mathematics*. <https://doi.org/10.1093/qmath/haab032>
- [80] A. Marsiglietti: Borell's generalized Prékopa-Leindler inequality: a simple proof. *J. Convex Anal.* 24 (2017), 807-817.
- [81] R.J. McCann: Existence and uniqueness of monotone measure-preserving maps. *Duke Math. J.*, 80 (1995), 309-323.

- [82] R.J. McCann: A convexity principle for interacting gases. *Adv. Math.* 128 (1997), 153-179.
- [83] M. Meyer: A volume inequality concerning sections of convex sets. *Bull. Lond. Math. Soc.*, 20 (1988), 15-155.
- [84] V.D. Milman, G. Schechtman: *Asymptotic theory of finite-dimensional normed spaces. With an appendix by M. Gromov.* Springer-Verlag, Berlin, 1986.
- [85] C.M. Petty: Surface area of a convex body under affine transformations. *Proc. Amer. Math. Soc.* 12 (1961), 824–828,
- [86] A. Prékopa: Logarithmic concave measures with application to stochastic programming. *Acta Sci. Math. (Szeged)* 32 (1971), 301–316.
- [87] A. Prékopa: On logarithmic concave measures and functions. *Acta Sci. Math. (Szeged)* 34 (1973), 335–343.
- [88] A. Recski: *Matroid Theory and its Applications in Electric Network Theory and in Statics.* Springer, 1989.
- [89] A. Rossi, P. Salani: Stability for Borell-Brascamp-Lieb inequalities. *Geometric aspects of functional analysis, Lecture Notes in Math.*, 2169, Springer, Cham, (2017), 339-363.
- [90] A. Rossi, P. Salani: Stability for a strengthened Borell-Brascamp-Lieb inequality. *Appl. Anal.*, 98 (2019), 1773-1784.
- [91] R. Schneider: *Convex bodies: the Brunn-Minkowski Theory.* Cambridge University Press, Cambridge, 1993, Second expanded edition, 2014.
- [92] S.I. Valdimarsson: Geometric Brascamp-Lieb has the optimal best constant. *J. Geom. Anal.* 21 (2011), 1036-1043.
- [93] S.I. Valdimarsson: Optimisers for the Brascamp-Lieb inequality. *Israel J. Math.* 168 (2008), 253-274.
- [94] C. Villani: *Topics in optimal transportation.* AMS, Providence, RI, 2003.