Quantum Entanglement in the Sachdev-Ye-Kitaev Model and its Generalizations

Pengfei Zhang¹

¹Institute for Quantum Information and Matter and Walter Burke Institute for Theoretical Physics, California Institute of Technology, Pasadena, CA 91125, USA (Dated: March 4, 2022)

Entanglement is one of the most important concepts in quantum physics. We review recent progress in understanding the quantum entanglement in many-body systems using large-N solvable models: the Sachdev-Ye-Kitaev (SYK) model and its generalizations. We present the study of entanglement entropy in the original SYK Model using three different approaches: the exact diagonalization, the eigenstate thermalization hypothesis, and the path-integral representation. For coupled SYK models, the entanglement entropy shows linear growth and saturation at the thermal value. The saturation is related to replica wormholes in gravity. Finally, we consider the steady-state entanglement entropy of quantum many-body systems under repeated measurements. The traditional symmetry breaking in the enlarged replica space leads to the measurement-induced entanglement phase transition

18

CONTENTS

1.	introduction	1
Π.	The SYK model and its generalizations	2
	A. Large- <i>N</i> expansion	2
	B. Low-energy limit and holography	2 3
	C. Generalizations of the SYK model	4
III.	Entanglement in the SYK model	5
	A. The exact diagonalization	5
	B. The eigenstate thermalization hypothesis	6
	C. The path-integral approach	7
	D. Exact results for SYK ₂	9
IV.	Entanglement in coupled SYK models	9
	A. The set-up and G - Σ action	10
	B. Early-time linear growth	11
	C. Geometric interpretation for $N_2 = N_1$	12
	D. Thermalization and replica wormholes	13
	E. Entanglement dynamics of SYK chains	14
V.	Entanglement in non-unitary SYK models	14
	A. Emergent replica conformal symmetry	15
	B. Entanglement phase transitions	16
	C. Effects of long-range couplings	17
VI.	Outlook	17

I. INTRODUCTION

References

Quantum entanglement, a fascinating property of quantum states, has been widely studied in quantum information, condensed matter, and high-energy physics. For example, in quantum information, entanglement is crucial for pure-state quantum algorithms to offer a speed-up over classical computation [1–3]. In condensed matter physics, entanglement entropy can be used to distinguish thermalized/localized systems [4–8], as well as to characterize topologically ordered

phases [9, 10]. In high energy physics, the discovery of the Ryu-Takayanagi (RT) formula [11–16], which relates the fine-grained entropy of a subsystem to geometric properties, has considerably deepened our understanding of gravity, and recently led to recent progress in the information paradox [17–22]. However, the entanglement entropy is notoriously difficult to calculate due to the lack of efficient algorithms, in particular, for strongly interacting many-body systems. Most previous studies focus on systems with conformal symmetry [23], holographic duality [24], or well-defined large-*N* expansions [25–27].

The Sachdev-Ye-Kitaev (SYK) model, proposed by Kitaev [28] and related to the early work by Sachdev and Ye [29], describes N Majorana fermions with random interactions. The SYK model is solvable under large-N expansion [30]. For example, to the leading order of 1/N, the Schwinger-Dyson equation for the Green's function can be derived in a closed form, the solution of which also determines thermodynamics. In the low-temperature limit, the SYK model shows emergent conformal symmetry. Its low-energy dynamics have a holographic dual representation by the 2D Jackiw-Teitelboim gravity [31, 32], which implies the system is maximally chaotic [33]. There are also studies on quantum dynamics of the SYK model, which reveals it exhibits quantum thermalization [34]. Later, different generalizations of the SYK model have been proposed to study various physical problems [35], including models with charge conservation [36– 39], Brownian interaction [40, 41], or multiple sites [42–51].

In this review, we summarize recent progress in the quantum entanglement in the SYK model and its generalizations. In the original SYK model, the entanglement entropy has been studied using three different approaches [52–58]: the exact diagonalization (ED), the eigenstate thermalization hypothesis (ETH), and the path-integral representation. Using the ED approach [52, 53], existing numerical analysis give results of the ground state entanglement of the SYK model, up a system size N=44 [52]. On the other hand, the ETH states that in thermalized systems, small subsystems can be approximated by thermal ensembles with an effective temperature determined by the energy density [7, 8]. In particular, the entanglement entropy is equal to the corresponding thermal entropy. For systems with local interaction, the boundary effect

is small, and the effective temperature is independent of subsystem size. Whereas for systems with non-local interaction, the effective temperature depends on the subsystem size [54]. One can also express the *n*-th Rényi entropy as a path integral and perform the large-*N* expansion as for thermodynamics [55–57]. This leads to results exact in the large-*N* limit. Results from these three approaches match to good precision.

For generalized SYK models that contain multiple sites, with some of the sites described by the SYK model, there is no simple approach to employ the ETH analysis, and the ED is limited to small system size N. Consequently, previous studies rely on the path-integral approach [22, 59–69]. We are interested in the entanglement dynamics of the system prepared in some short-range entangled state. In the simplest case, we consider two SYK sites with N_1 and N_2 Majorana modes, coupled by random pair hopping [22, 59-61]. In the earlytime regime, the entanglement entropy shows linear growth in time [22, 59–63]. In the long-time limit, when $N_1 = N_2$, it is found that in the canonical ensemble the Rényi entanglement entropy does not saturate to the thermal value due to the existence of soft modes [22, 59, 60]. On the contrary, if $N_2 \gg N_1$, the second site serves as a heat bath, and the Rényi entanglement entropy of the first site always matches the thermal entropy [60, 61]. When the entanglement entropy approaches the thermal value, a first-order transition occurs. This is a direct analog of the appearance of replica wormholes [21, 22] in the entanglement calculations of systems with gravity, which is an important breakthrough in high-energy physics that resolves the information paradox [70]. Alternatively, one may instead consider replacing the static SYK interaction with the Brownian counterpart [64]. Analytical results can then be derived under certain assumptions. There are also attempts on understanding the von Neumann entanglement entropy [63].

In addition to unitary evolutions where thermalization generally takes place, the evolution of quantum many-body systems can also become non-unitary when measurements are performed [71]. Recent studies show that if we follow the quantum trajectories of systems with repeated measurements, the steady-state can exhibit volume law, logarithmic, or area law entanglement [72–98]. This mechanism works for either projective or general measurements, regardless of whether the outcome is post-selected or not. SYK-like models are ideal platforms for revealing the underlying mechanism of these transitions [92–98]. In the last part of this review, we summarize recent progress in understanding the steady-state entanglement entropy of non-unitary SYK models. For small measurement rates, when the system is non-interacting, an emergent replica conformal symmetry shows up due to the existence of Goldstone modes after introducing multiple replicas [93, 94]. This leads to logarithmic entanglement entropy. After adding interactions, the Goldstone modes become massive [94], and the system becomes volume law entangled. For large measurement rates, the steady-state of the system may become area law entangled, and the transition is a traditional symmetry-breaking transition on the enlarged replica space [94, 98].

This review is organized as follows: In section II, we give a brief introduction to the SYK model and its various gen-




FIG. 1. Schematics of the SYK_4 model, which describes Majorana fermions with random 4-body interactions.

eralizations. Then, in section III, we summarize existing results for the entanglement entropy of the original SYK model. The entanglement entropy for coupled SYK models and non-unitary SYK models are discussed separately in sections IV and V. Finally, we conclude our review by a few outlooks in section VI.

II. THE SYK MODEL AND ITS GENERALIZATIONS

In this section, we give a brief introduction to the original SYK model [28, 30]. The Hamiltonian of the SYK $_q$ model reads

$$H_{\text{SYK}_q} = \sum_{1 \le i_1 \le i_2 \le \dots \le i_n \le N} i^{q/2} J_{i_1 i_2 \dots i_q} \chi_{i_1} \chi_{i_2} \dots \chi_{i_q}. \tag{1}$$

Here q is an even integer. It describes q-body random interactions between N Majorana modes χ_i . We choose the convention that $\{\chi_i,\chi_j\} = \delta_{ij}$. $J_{i_1i_2...i_q}$ with different set of indices are independent Gaussian variables, with expectations and variances given by

$$\overline{J_{i_1 i_2 \dots i_q}} = 0, \quad \overline{J_{i_1 i_2 \dots i_q}^2} = \frac{(q-1)! J^2}{N^{q-1}} = \frac{2^{q-1} (q-1)! \mathcal{J}^2}{q N^{q-1}}.$$
 (2)

Here the scaling of N is chosen such that the energy is extensive. \mathcal{J} is introduced for convenience in the large-q analysis [30], which should be fixed as a constant as $q \to \infty$. The model describes Majorana fermions with random hopping for q = 2, and becomes randomly interacting for $q \ge 4$.

A. Large-N expansion

The large-N expansion is a standard approach for analyzing quantum field theories that contain large number of fields [99]. Here we explain the large-N expansion in the SYK model by considering the partition function $Z = \operatorname{tr} e^{-\beta H_{\text{SYK}_q}}$. The path-integral representation takes the form of $Z = \int \mathcal{D}\chi \exp\left(-S\left[\chi,J\right]\right)$, with

$$S[\chi, J] = \int d\tau \left[\sum_{i} \frac{1}{2} \chi_{i} \partial_{\tau} \chi_{i} + H_{\text{SYK}_{q}}[\chi] \right].$$
 (3)

By definition, the free-energy after the disorder average is given by $F(\beta) = -\beta^{-1} \overline{\log Z}$. For general systems, the computation of $F(\beta)$ then requires introducing n disorder replicas

and using $\log Z = \lim_{n\to 0} (Z^n - 1)/n$ [100]. However, for the SYK model, it is known that replica off-diagonal solutions are sub-leading in N [32], and one can approximate $\log Z \approx \log \overline{Z}$. Physically, this indicates the thermodynamical properties of the SYK model are self-averaging. Performing the Gaussian integral over $J_{i_1i_2...i_q}$, we have

$$\overline{Z} = \int \mathcal{D}\chi \ e^{-\frac{1}{2} \int d\tau \ \sum_i \chi_i(\tau) \partial_\tau \chi_i(\tau) + \frac{\jmath^2}{2qN^{q-1}} \int d\tau d\tau' \ [\sum_i \chi_i(\tau) \chi_i(\tau')]^q}. \tag{4}$$

The combination $\sum_i \chi_i(\tau) \chi_i(\tau')/N$ takes the form of averaging over multiple variables. By the central limit theorem [101], its variance should vanish in the large-N limit. This motivates the introduction of bilocal fields $\tilde{G}(\tau,\tau') = \sum_i \chi_i(\tau) \chi_i(\tau')/N$, which should be imposed by an additional Lagrangian multiplier field $\tilde{\Sigma}(\tau,\tau')$

$$1 = \int \mathcal{D}\tilde{G}\mathcal{D}\tilde{\Sigma} \ e^{\frac{1}{2}\int d\tau d\tau' \ \tilde{\Sigma}(\tau,\tau')\left(\sum_{i}\chi_{i}(\tau)\chi_{i}(\tau') - N\tilde{G}(\tau,\tau')\right)}.$$
 (5)

Here the integral over Σ is along the imaginary axis. Inserting (5) into (4) and replacing $\sum_i \chi_i(\tau) \chi_i(\tau') / N$ with $G(\tau, \tau')$, we find

$$\overline{Z} = \int \mathcal{D}\chi \mathcal{D}\tilde{G}\mathcal{D}\tilde{\Sigma} e^{-\int d\tau d\tau' \left[\frac{1}{2}\sum_{i}\chi_{i}(\partial_{\tau}-\tilde{\Sigma})\chi_{i}+\frac{N}{2}(\tilde{\Sigma}\tilde{G}-\frac{j^{2}}{q}\tilde{G}^{q})\right]}. \quad (6)$$

Now the fermion fields become bilinear and thus can be integrated out straightforwardly

$$\overline{Z} = \int \mathcal{D}\tilde{G}\mathcal{D}\tilde{\Sigma} \exp\left(-S_{G\Sigma}[\tilde{G},\tilde{\Sigma}]\right).$$

$$S_{G\Sigma} = -\frac{N}{2}\log\det\left(\partial_{\tau} - \tilde{\Sigma}\right) + \frac{N}{2}\int d\tau d\tau' \left(\tilde{\Sigma}\tilde{G} - \frac{J^{2}}{q}\tilde{G}^{q}\right).$$
(7)

 $S_{G\Sigma}$ is known as the G- Σ action for the SYK model. Importantly, $S_{G\Sigma}$ is proportional to N. As a result, we can apply the saddle-point approximation for the integration over \tilde{G} and $\tilde{\Sigma}$. This is consistent with our previous arguments using the central limit theorem. The saddle-point equations read

Here (G, Σ) is the saddle-point value of $(\tilde{G}, \tilde{\Sigma})$, which is only a function of the time difference due to the time translation symmetry. The first equation can also be written in short as $(\partial_{\tau} - \Sigma) \circ G = I$. This is equivalent to the Schwinger-Dyson equation of Majorana fermions with melon diagrams. In the diagram, we take q = 4 as an example. The solid lines are Green's functions for Majorana fermions, and the dashed line represents the disorder average. The numerical algorithm for solving (8) can be found in [30]. After obtaining the solution, we can determine the free energy as $\beta F = I = S_{G\Sigma}[G, \Sigma]$, or explicitly

$$\frac{2\beta F}{N} = -\sum_{n} \log(-i\omega_n - \Sigma(i\omega_n)) + \frac{(q-1)\beta}{q} \int d\tau \, \Sigma G, \quad (9)$$

where $\omega_n = 2\pi(n + 1/2)/\beta$ with $n \in \mathbb{Z}$ is the Matsubara frequency.

In the large-q limit, the Green's function can be expanded in terms of 1/q and the free energy can be obtained in the closed form [30]

$$-\frac{\beta F}{N} = \frac{\log 2}{2} + \frac{\pi v}{q^2} \left[\tan \frac{\pi v}{2} - \frac{\pi v}{4} \right],\tag{10}$$

with $\beta \mathcal{J} = \pi v / \cos(\pi v / 2)$. Under the Legendre transformation, this gives the thermal entropy

$$\frac{S_{\text{th}}(E)}{N} = \frac{\log 2}{2} - \frac{1}{q^2}\arcsin^2\left(\frac{E}{E_0}\right). \tag{11}$$

We have defined the ground state energy $E_0 = -\mathcal{J}/q^2$. In the zero-temperature limit, there is a residue entropy $S_0/N = \log 2/2 - \pi^2/4q^2$. Such a finite zero-temperature entropy is a general phenomenon for the SYK model. It is known that for finite q [30, 37, 102],

$$\frac{S_0}{N} = \int_{1/q}^{1/2} dx \, \pi \left(\frac{1}{2} - x\right) \tan \pi x. \tag{12}$$

This indicates the existence of exponentially many low-energy states. Although (11) is derived in the large-q limit, it turns out to be a good approximation for finite q away from $|E/E_0| = 1$. It has also been derived using a different approach in [103].

One can go further by considering the fluctuation around saddle-point solutions. The quadratic order expansion determines the fluctuation $\langle (\tilde{G} - G)(\tilde{G} - G) \rangle$, which corresponds to a connected four-point function of Majorana fermions. Detailed analysis can be found in [30].

B. Low-energy limit and holography

The SYK model shows emergent conformal symmetry and holographic duality in the low-temperature limit $\beta \mathcal{J} \gg 1$. Let us first perform the scaling analysis for the kinetic term $\chi \partial \chi$ and the interaction $J\chi^q$ in (3). If we assume the kinetic term is marginal, this gives the scaling dimension $[\chi]=0$, and the interaction term is relevant. This is not a stable fixed point. Consequently, we should have assumed the interaction term is marginal, which gives $[\chi] \equiv \Delta = 1/q$. This implies in the low-energy limit $\tau \mathcal{J} \gg 1$, we can neglect the ∂_{τ} term in the Schwinger-Dyson equation

$$-J^{2} \int d\tau' \ G^{q-1}(\tau - \tau') G(\tau') = \delta(\tau), \tag{13}$$

and the solution reads

$$G(\tau) = b \left(\frac{\pi}{\beta \sin \frac{\pi \tau}{\beta}} \right)^{2\Delta}, \qquad J^2 b^q \pi = \left(\frac{1}{2} - \Delta \right) \tan \pi \Delta.$$
 (14)

This is the standard form of Green's functions in the conformal field theory (CFT), which indicates the emergence of

conformal symmetry in the SYK model. Indeed, after neglecting the kinetic term, the G- Σ action (7) is invariant under the reparametrization (or conformal transformation in 1D) $\tau \to f(\tau)$, with

$$\tilde{G}(\tau_1, \tau_2) = (f(\tau_1)' f(\tau_2)')^{\Delta} \tilde{G}(f(\tau_1), f(\tau_2)). \tag{15}$$

In particular, this indicates that fluctuations

$$\tilde{G}_f(\tau_1, \tau_2) = (f(\tau_1)' f(\tau_2)')^{\Delta} G(f(\tau_1) - f(\tau_2))$$

cost no energy. These are called reparametrization modes, which governs the SYK low-energy dynamics. After adding back contributions from the kinetic term ∂_{τ} , a non-vanishing action for these reparametrization modes can be derived [30, 32], which is known as the Schwarzian action

$$S_{\text{Sch}} = -\frac{N\alpha_S}{\mathcal{J}} \int d\tau \left\{ \tan \frac{\pi f(\tau)}{\beta}, \tau \right\}.$$
 (16)

Here $\{f, \tau\} = f'''/f' - 3(f''/f')^2/2$ is the Schwarzian derivative and α_S is some numerical factor. There are extensive discussions of the exact solution of the Schwarzian action from different perspectives [104–107].

The holography dual of the SYK low-energy sector refers to the fact that the Schwarzian action can also be obtained from an independent gravity calculation [31]. The gravity theory is the Jackiw-Teitelboim (JT) gravity in two-dimensional Anti-de-Sitter (AdS_2) space. The action reads

$$S_{\rm JT} = -\frac{1}{16\pi G} \left[\int dx^2 \sqrt{g} \phi(R+2) + 2 \int_{\rm bdy} \phi_b K \right].$$
 (17)

The first/second term is a bulk/boundary term. ϕ is the dilaton field, which can be integrated out as a Lagrangian multiplier to impose R = -2. In 2D, this fixes all degrees of freedom in bulk. We choose the bulk to be a Poincaré disk, with metric

$$ds^2 = d\rho^2 + \sinh^2 \rho \ d\theta^2. \tag{18}$$

The only remaining dynamical degrees of freedom is the fluctuation of the boundary curve. We fix the boundary condition that $g_{\tau\tau}=1/\epsilon^2\gg 1$ with boundary time τ , this gives

$$\sinh \rho(\tau) \, \theta'(\tau) \approx \epsilon^{-1}.$$
(19)

As a result, the shape of the boundary curve can be parametrized by a single function $\theta(\tau)$, and the energy cost is determined by the extrinsic curvature term (17). Explicit calculation shows the result is exactly the Schwarzian action (16), with $2\pi f(\tau) = \beta \theta(\tau)$. In particular, the saddle-point solution $f(\tau) = \tau$ corresponds to a circle on the Poincaré disk. This completes the discussion of the holographic dual of the SYK low-energy sector.

C. Generalizations of the SYK model

We have introduced the original SYK model in previous subsections. Here we turn to its generalizations. The simplest generalization includes adding SYK terms with different q into a single Hamiltonian $H = \sum_q c_q H_{\text{SYK}_q}$, with $c_q \neq 0$ for some q. In addition, there are mainly three routes for generalizing the SYK model: adding symmetry, making coupling time-dependent, and introducing multiple sites. In this subsection, we illustrate these ideas using a few examples.

Adding Symmetry Symmetry plays an important role in modern physics. The original SYK model only exhibits energy conservation, and there are several attempts to add symmetries for the SYK model. The simplest generalization is to replace Majorana fermions with complex fermions [36–39] and impose the U(1) charge conservation. This leads to the complex SYK (or cSYK) model. The Hamiltonian of the cSYK_a model reads

$$H_{\text{cSYK}_q} = \sum_{\substack{i_1 < \dots < i_{q/2} \\ j_1 < \dots < j_{q/2}}} J_{i_1 i_2 \dots i_{q/2} j_1 j_2 \dots j_{q/2}} c_{i_1}^{\dagger} \dots c_{i_{q/2}}^{\dagger} c_{j_1} \dots c_{j_{q/2}}, \quad (20)$$

where the Gaussian distributions of random couplings $J_{i_1i_2...i_{q/2}j_1j_2...j_{q/2}}$ satisfy

$$\overline{J_{i_1 i_2 \dots i_{q/2} j_1 j_2 \dots j_{q/2}}} = 0, \qquad \overline{J_{i_1 i_2 \dots i_{q/2} j_1 j_2 \dots j_{q/2}}^2} = \frac{\frac{q}{2}! (\frac{q}{2} - 1)! J^2}{N^{q-1}}. \tag{21}$$

In the grand canonical ensemble, an chemical potential μ can be added to tune the charge filling $Q = \sum_i c_i^{\dagger} c_i$.

The large-*N* expansion of the cSYK model is similar to that of the Majorana SYK model. For example, the Schwinger-Dyson equation can be written as [29, 36]

$$\delta(\tau) = (\partial_{\tau} - \mu)G(\tau) - \int d\tau' \ \Sigma(\tau - \tau')G(\tau'),$$

$$\Sigma(\tau) = J^{2}G^{q/2}(\tau)[-G(-\tau)]^{q/2-1} =$$
(22)

and the free energy can also be expressed as a functional of G and Σ . In the low-energy limit, where is an additional low-energy parameter θ , which is related to the charge filling Q and characterizes the asymmetry of the spectral function between particles and holes [36]. There is also an additional soft mode $\lambda(\tau)$, which is associated with the phase fluctuation [36, 37].

Time-dependent Coupling The Hamiltonian of the original SYK model is static. Later, there are studies in which couplings between Majorana fermions are also random in time. This is known as the Brownian SYK (or BSYK) model [40, 41]. The Hamiltonian of the BSYK model is time-dependent:

$$H_{\text{BSYK}_q}(t) = \sum_{1 \le i_1 < i_2 < \dots < i_q \le N} i^{q/2} J_{i_1 i_2 \dots i_q}(t) \chi_{i_1} \chi_{i_2} \dots \chi_{i_q}, \qquad (23)$$

 $J_{i_1i_2...i_q}(t)$ are Gaussian variables with zero expectations and

$$\overline{J_{i_1 i_2 \dots i_q}(t) J_{i_1 i_2 \dots i_q}(t')} = \frac{(q-1)! J}{N^{q-1}} \delta(t-t'). \tag{24}$$

Since there is no energy conservation, only the infinite temperature ensemble is well-defined. Instead of the imaginary-time path-integral, we consider the Keldysh contour which contains one forward and one backward evolution [108]. The partition function $Z = \text{tr } e^{iHT} e^{-iHT}$ with $T \to \infty$ reads

$$Z = \int \mathcal{D}\chi^{+} \mathcal{D}\chi^{-} e^{-\int_{-\infty}^{\infty} dt \sum_{\pm} \left[\sum_{i} \pm \frac{1}{2} \chi_{i}^{\pm} \partial_{i} \chi_{i}^{\pm} \pm i H_{\text{BSYK}_{q}} [\chi^{\pm}] \right]}.$$
 (25)

 χ_i^{\pm} are Majorana fields on the forward/backward evolution contour. One can follow derivation in section II A to integrate over the random interactions, introduce the $(\tilde{G}, \tilde{\Sigma})$ fields, and derive the saddle-point equations [109]. The result reads

$$\begin{bmatrix} \begin{pmatrix} \partial_t & 0 \\ 0 & -\partial_t \end{pmatrix} - \begin{pmatrix} \Sigma^{++} & \Sigma^{+-} \\ \Sigma^{-+} & \Sigma^{--} \end{pmatrix} \end{bmatrix} \circ \begin{pmatrix} G^{++} & G^{+-} \\ G^{-+} & G^{--} \end{pmatrix} = I,$$
(26)

with self-energy $\Sigma^{ss'}(t,t') = -Jss'\delta(t-t')G^{ss'}(t,t')^{q-1}$, which is non-zero only when two time variables coincide. This is the consequence of having Brownian couplings.

Multiple Sites By introducing different copies of the SYK model and adding interactions between them, we can construct SYK-like models with more than one site [42–51]. As an example, we consider coupling two SYK₄ models using SYK₄ random pair hopping:

$$H = \sum_{\{i_n\}} J^{\chi}_{i_1 i_2 i_3 i_4} \chi_{i_1} \chi_{i_2} \chi_{i_3} \chi_{i_4} + \sum_{\{a_n\}} J^{\psi}_{a_1 a_2 a_3 a_4} \psi_{a_1} \psi_{a_2} \psi_{a_3} \psi_{a_4}$$

$$+ \sum_{i_1 < i_2, a_1 < a_2} V_{i_1 i_2 a_1 a_2} \chi_{i_1} \chi_{i_2} \psi_{a_1} \psi_{a_2}.$$

$$(27)$$

Here we have N_1 χ fermions $(1 \le i_1 < ... < i_4 \le N_1)$ and N_2 ψ fermions $(1 \le a_1 < ... < a_4 \le N_2)$. We take both $N_1, N_2 \to \infty$ with their ratio $r = N_2/N_1$ fixed. We consider that both $J_{i_1 i_2 i_3 i_4}^{\chi}$ and $J_{a_1 a_2 a_3 a_4}^{\psi}$ are SYK random interactions with strength J. The random pair hopping strengths satisfy

$$\overline{V_{i_1 i_2 a_1 a_2}} = 0, \qquad \overline{V_{i_1 i_2 a_1 a_2}^2} = \frac{2V^2}{N_1 N_2^2}.$$
 (28)

Now the self-energies become

$$\begin{split} & \Sigma_{\chi}(\tau) = J^2 G_{\chi}(\tau)^3 + V^2 G_{\psi}(\tau)^2 G_{\chi}(\tau), \\ & \Sigma_{\psi}(\tau) = J^2 G_{\psi}(\tau)^3 + V^2 \frac{N_1}{N_2} G_{\psi}(\tau) G_{\chi}(\tau)^2. \end{split} \tag{29}$$

In the low-energy limit, the emergent conformal symmetry and the soft reparametrization mode $f(\tau)$ still exist, with a Schwarzian effective action. Following the same spirit, one can introduce a chain of SYK sites with random pair hopping. This leads to higher-dimensional SYK lattices [42], in which the low-energy butterfly velocity and the energy diffusion constant has been studied.

III. ENTANGLEMENT IN THE SYK MODEL

After introducing the SYK model in section II, we now review existing results for its entanglement entropy. We divide




FIG. 2. The von Neumann entanglement entropy for the ground state of the SYK₄+SYK₂ model using ED. The result is averaged over 10 disorder realizations. M = m is the subsystem size of the cSYK model, while for the Majorana SYK the subsystem size is M = 2m. The system contains N = 22 complex fermions with filling fraction $\kappa = Q/N$ or N = 44 Majorana fermions. Adapted from [52].

the system into A and B, where A contains M Majorana modes and B contains N-M Majorana modes. We fix $M/N = \lambda$ when taking the large-N limit. The definition of A and B subsystems in the cSYK model is similar. Given the density matrix of the total system ρ , the reduced density matrix of A can be computed as $\rho_A = \operatorname{tr}_B \rho$. The n-th Rényi entropy $S_A^{(n)}$ and the von Neumann entropy S_A are defined as

$$S_A^{(n)} = -\frac{1}{n-1} \log \operatorname{tr}_A \rho_A^n, \quad S_A = -\operatorname{tr}_A \rho_A \log \rho_A.$$
 (30)

It is straightforward to show that the von Neumann entropy is a special limit of the Rényi entropy $S_A = S_A^{(1)}$.

There are three different approaches for studying either $S_A^{(n)}$ or S_A in the SYK model: the ED, the ETH assumption, and the path-integral representation. All of them have some advantages and disadvantages. The ED can give exact numerical results for both $S_A^{(n)}$ and S_A of any state, but is limited to small system size N and small q. The ETH analysis can lead to large-N results for S_A , but it requires further justification and knowledge about thermodynamics. The path-integral approach works in the large-N limit and can be easily extended to other models, but in most cases is restricted to $S_A^{(n)}$ with small integers $n \geq 2$. In the following subsections, we summarize results obtained using these approaches separately, and show their consistency.

A. The exact diagonalization

We first review results obtained from the exact diagonalization in [52]. In this work, authors numerically study the ground-state von Neumann entropy S_A for a complex fermion

model with Hamiltonian

$$H_{\rm c} = H_{\rm cSYK_4} + \frac{g}{N} H_{\rm cSYK_2},\tag{31}$$

or a Majorana fermion model with H_{cSYK_q} replaced by its Majorana counterpart H_{SYK_q} . Using the sparse matrix method, authors study systems with N=22 complex fermions or N=44 Majorana fermions. For complex fermions, one can further fix the charge filling $\kappa=Q/N$. The result is shown in Figure 2.

For $M \ll N$, the entanglement entropy is maximal for any g. For the Majorana fermion model, this corresponds to $S_A = M \log 2/2$. For the complex fermion model, it depends on the filling fraction κ :

$$S_A = -M \left(\kappa \log \kappa + (1 - \kappa) \log(1 - \kappa) \right). \tag{32}$$

For $M/N \sim O(1)$, the entanglement entropy is lowered when g is increased. For $g \to \infty$, the entanglement entropy matches the SYK₂ result, as will be discussed in section (III D). The authors also compare the numerical results with the Page curve [110], which is the von Neumann entanglement entropy of a random pure state

$$S_A^{\text{Page}} = M \log 2 - 2^{2M-N-1}.$$
 (33)

Here we take the complex fermion model as an example. For the Majorana model, one should make the replacement of $(N, M) \rightarrow (N, M)/2$. The first term is the maximal entropy, and the second term is a small correction that vanishes in the large-N limit. As shown in Figure 2, the numerical result of the SYK₄ model is smaller than the Page value. The difference is expected to become smaller when we consider models with larger q.

B. The eigenstate thermalization hypothesis

The entanglement entropy of the SYK $_q$ model can also be studied using the ETH. The ETH states that an isolated quantum system can serve as its own bath. More precisely, let us consider a eigenstate $|\psi\rangle$ of some local chaotic Hamiltonian H with energy $E=\langle\psi|H|\psi\rangle$. If we measure some local observable O in the subsystem A, the result will match the prediction from a thermal ensemble

$$\langle \psi | O | \psi \rangle = \frac{\text{tr } O e^{-\beta H}}{\text{tr } e^{-\beta H}}.$$
 (34)

Here β is fixed by matching the energy with O = H. This can be understood as proposing the density matrix of the subsystem A is almost thermal. Consequently, the entanglement entropy of the subsystem A matches the corresponding thermal entropy $S_A = MS_{th}(E)/N$.

The SYK model describes Majorana fermions with non-local interactions. As a result, this analysis should be modified following the discussions in [54]. In this subsection, we outline the basic idea. For simplicity, we choose the subsystem A to contain Majorana modes χ_i with i = 1, 2, ..., M. To understand the reduced density matrix of subsystem A, we first




FIG. 3. A comparison between different approaches for the entangelement entropy for the SYK₄ model. In the path-integral approach, we consider $S_A^{(2)}$ set $\beta J = 50$ [55, 56]. The ED data is copied from Figure (2). The analytical ETH result is plotted according to (39) and the numerical ETH result is adapted from [55].

define its Hamiltonian H_A by restricting the summation over indices in (1):

$$H_A = \sum_{1 \le i_1 < i_2 < \dots < i_q \le M} i^{q/2} J_{i_1 i_2 \dots i_q} \chi_{i_1} \chi_{i_2} \dots \chi_{i_q}.$$
 (35)

Comparing to the original Hamiltonian, which contains $\sim N^q$ terms, here the summation is restricted to $\sim M^q$ terms. As a result, to the leading order of 1/N, we have

$$E_A = \langle \psi | H_A | \psi \rangle = \frac{M^q}{N^q} \langle \psi | H | \psi \rangle = \lambda^q E.$$
 (36)

Here we have assumed the state is symmetric under the permutation of indices. This is very different from the case with local Hamiltonians, where the energy density will not change.

In the spirit of ETH, we assume the reduced density matrix ρ_A is a thermal density matrix of H_A for $\lambda < 1/2$, and the entanglement entropy S_A matches thermal entropy. To determine the entropy density $s = S_A/M$, we need additional knowledge about H_A . Assume the ground state energy of the SYK model with N Majorana fermions with coupling J is $E_0 = Nc_0J$, the ground state energy $E_{A,0}$ of H_A is determined by realizing

$$\overline{J_{i_1 i_2 \dots i_q}^2} = \frac{(q-1)!J^2}{N^{q-1}} = \frac{(q-1)!(\lambda^{(q-1)/2}J)^2}{M^{q-1}}.$$
 (37)

As a result, H_A is an SYK model that contains M Majorana fermions with coupling strength $J\lambda^{(q-1)/2}$. This gives $E_{A,0} = Mc_0\lambda^{(q-1)/2} = \lambda^{(q+1)/2}E_0$. We have

$$\frac{E_A}{E_{A,0}} = \lambda^{\frac{q-1}{2}} \frac{E}{E_0},\tag{38}$$

Physically, this means that for non-local Hamiltonians, the effective temperature changes when varying the subsystem size. The entanglement entropy is then determined by the thermal entropy with the corresponding energy density. In the limit of

 $\lambda \to 0$, the energy density vanishes, and the thermal ensemble is at infinite temperature. This explains the observed maximal entropy behavior for small subsystem size in ED.

A simple analytical approximation of S_A can be derived using (11):

$$\frac{S_A}{N} = \lambda \left(\frac{\log 2}{2} - \frac{1}{q^2} \arcsin^2 \left(\lambda^{\frac{q-1}{2}} \frac{E}{E_0} \right) \right). \tag{39}$$

In particular, for the ground state of the SYK₄ model (E/E_0 = 1), this gives S_A = 0.338M at λ = 1/2. On the other hand, one can also perform the numerical study to derive the equation of state as explained in (II A). The numerical results are shown in Figure 3, which match results of the path-integral approach (explained in the next subsection) to good precision.

C. The path-integral approach

The Rényi entropy in the SYK model can also be studied using the large-N expansion. This is only possible for certain states where the path-integral takes a simple form. In this section, we consider the SYK model prepared in either a thermal ensemble or a Kourkoulou-Maldacena (KM) pure state [111]. We focus on the second Rényi entropy $S_A^{(2)}$, while a generalization to higher orders is straightforward. When the total system is in a thermal ensemble, the result contains both entanglement contributions and thermal contributions. However, as we will show later, the subsystem entropy of a thermal ensemble matches the steady-state entanglement entropy of the KM pure-state with the same energy for $\lambda < 1/2$, and can be directly compared to the ED and the ETH results.

Thermal Ensemble We first consider the subsystem entropy $S_A^{(2)}$ of a thermal ensemble $\rho = e^{-\beta H}/Z$ [55]. A pictorial representation of the density matrix ρ and the reduced density matrix ρ_A read

$$Z\rho = \beta \left(\begin{array}{c} B \\ A \end{array} \right)^0, \quad Z\rho_A = \beta \left(\begin{array}{c} B \\ A \end{array} \right)^0. \quad (40)$$

Here the solid lines represent the imaginary-time evolutions, and the dashed lines represent the interaction between subsystems. The black dot indicates the anti-periodic boundary condition for fermion fields, and the free ends can be assigned with quantum states to give matrix elements of density matrices. To compute $S_A^{(2)}$, we sew two ρ_A together:

$$\exp\left(-S_A^{(2)}\right) = \operatorname{tr} \rho_A^2 = \frac{1}{Z^2} \times \left[\beta \left(\begin{array}{c} \vdots \\ B_2 \\ \vdots \\ B_1 \\ A \end{array}\right) \right]. \tag{41}$$

The contribution in the square bracket is very similar to the path-integral of a thermal ensemble with inverse temperature 2β , which reads

$$Z^{(2)} = \int_{\text{b.c.}} \mathcal{D}\chi \ e^{-\int_0^{2\beta} d\tau \left[\sum_i \frac{1}{2} \chi_i \partial_\tau \chi_i + H_{\text{SYK}_q}[\chi]\right]}. \tag{42}$$

The only difference is the boundary condition. Here we should impose

$$\chi_i(0^+) = -\chi_i(2\beta^-), \quad \chi_i(\beta^+) = \chi_i(\beta^-), \quad \text{if} \quad i \in A;
\chi_i(0^+) = -\chi_i(\beta^-), \quad \chi_i(\beta^+) = -\chi_i(2\beta^-), \quad \text{if} \quad i \in B.$$
(43)

These boundary conditions break the time translation symmetry and the Green's functions depend on both time variables. In [57], this boundary condition is expressed in an equivalent form with an integral over auxiliary fermion modes.

To proceed, we need to average over the disorder realizations as $\overline{S_A^{(2)}} = -\log Z^{(2)}/Z^2$. Similar to the calculation of the free energy, here we assume the disorder replica is diagonal, and thus $\overline{S_A^{(2)}} = -\log \overline{Z^{(2)}}/\overline{Z}^2$ with

$$\overline{Z^{(2)}} = \int dJ P(J) \int_{\mathbb{R}^{C}} \mathcal{D}\chi \ e^{-\int_{0}^{2\beta} d\tau \left[\sum_{i} \frac{1}{2} \chi_{i} \partial_{\tau} \chi_{i} + H_{\text{SYK}_{q}}[\chi]\right]}. \tag{44}$$

The calculation of $\overline{Z^{(2)}}$ is parallel to that of \overline{Z} . One can integrate over disorder realizations, introduce bilocal fields

$$\tilde{G}_{A}(\tau, \tau') = \frac{1}{M} \sum_{i \in A} \chi_{i}(\tau) \chi_{i}(\tau'),$$

$$\tilde{G}_{B}(\tau, \tau') = \frac{1}{N - M} \sum_{i \in R} \chi_{i}(\tau) \chi_{i}(\tau'),$$
(45)

with Lagrangian multipliers $\tilde{\Sigma}_{A/B}$, and make the saddle-point approximation after integrating out Majorana modes. Skipping the details, the saddle-point equations read

$$G_{A} = (\partial_{\tau} - \Sigma_{A})_{A}^{-1}, \qquad G_{B} = (\partial_{\tau} - \Sigma_{B})_{B}^{-1},$$

$$\Sigma_{A} = \Sigma_{B} = J^{2} (\lambda G_{A} + (1 - \lambda) G_{B})^{q-1}.$$
(46)

These self-energies can again be understood by melon diagrams, with each internal line being G_A or G_B with probability λ or $1-\lambda$. We add subscripts A/B as a reminder that the inversion should be performed under the corresponding boundary condition (43). As a result, although the self-energy for fermions in A/B systems are the same, Green's functions are different. Generalizations to the cSYK model or with multiple flavors are straightforward. After solving (46), the on-shell action $I^{(2)} = -\log \overline{Z^{(2)}}$ reads

$$\frac{I^{(2)}}{N} = \frac{\lambda}{2} \log \det G_A + \frac{\lambda(q-1)}{2q} \int ds ds' G_A \Sigma_A$$

$$+ (\lambda \to 1 - \lambda, A \to B).$$
(47)

Recall that we have $\overline{Z} = e^{-I}$, the subsystem Rényi entropy is given by $\overline{S_A^{(2)}} = I^{(2)} - 2I$. Generalizations to the cSYK model and coupled SYK models are straightforward. In particular, for cSYK model at half-filling, the large-N entropy is only different from the Majorana case by a factor of 2 [55].

For $\lambda=1$, the calculation is reduced to the thermal Rényi entropy. For $\lambda\ll 1$, one can relate the partition function to the normalization of KM pure states and prove that the subsystem entropy is maximal [55], consistent with previous discussions.

In more general cases, one needs to perform a numerical study by solving the saddle-point equation (46). Compared to the thermal equilibrium calculation, this requires additional efforts since Green's function depends on two-time variables. Fortunately, Green's function of the SYK model is smooth, and a small discretization is enough. Then we can store each Green's functions as matrices and directly perform iterations as explained in [55]. As an example, we show results with q=4 and $\beta J=50$ in Figure 3 for $\lambda<1/2$. It clearly shows the subsystem entropy is smaller than the maximal value for $\lambda\sim O(1)$. Interestingly, this subsystem entropy matches the entanglement entropy of thermalized pure states. This will be explained below.

KM Pure States Since the entanglement entropy is messed up with the thermal entropy if the full system is in a mixed state, it is natural to ask whether some pure-state entanglement entropy can be computed using the path-integral approach for the SYK model. As discussed in [56], such a calculation is possible for KM pure states [111].

The KM pure states are defined similarly to the thermofield double (TFD) state [112] (also see section IV). We first prepare a maximally entangled state between Majorana modes. Without loss of generality, we choose to entangle fermions with even and odd indices. Defining $c_j = \frac{\chi_{2j-1} + i \chi_{2j}}{\sqrt{2}}$, we introduce the (unnormalized) KM pure states

$$|KM(\{s\},\beta)\rangle = e^{-\frac{\beta H}{2}}|\{s\}\rangle, \qquad (1-2n_j)|\{s\}\rangle = s_j|\{s\}\rangle.$$
 (48)

Here $n_j = c_j^{\dagger} c_j$ and $s_j = \pm 1$. Under transformation $\chi_{2j} \rightarrow -\chi_{2j}$, s_j changes sign. As a result, we are free to average over all $\{s_j = \pm 1\}$ when computing the energy. Due to the completeness of $|\{s\}\rangle$, we find

$$\frac{\langle \text{KM}(\{s\},\beta)|H|\text{KM}(\{s\},\beta)\rangle}{\langle \text{KM}(\{s\},\beta)|\text{KM}(\{s\},\beta)\rangle} = \frac{1}{Z} \text{tr } He^{-\beta H}, \tag{49}$$

and the energy density matches the corresponding thermal ensemble. A holographic interpretation for KM states has also been proposed in [111]. The KM pure states are generally non-thermal. In particular, they are product states when $\beta=0$. To make the states thermal, we add additional real-time evolutions, and the states become

$$|KM(\lbrace s\rbrace, \beta, t)\rangle = e^{-iHt} e^{-\frac{\beta H}{2}} |\lbrace s\rbrace\rangle,. \tag{50}$$

The path-integral representation of KM pure states can be pictorially drawn as

$$|KM(\lbrace s\rbrace, \beta)\rangle = e^{-\frac{\beta H}{2}}|\lbrace 1\rbrace \rangle = \chi_1 \bigcup_{0}^{\frac{\beta}{2}} \chi_2$$

$$\downarrow_0 \bigcup_{|\lbrace s\rbrace \rangle} \chi_2$$
(51)

Here χ_1/χ_2 represents Majorana modes with odd/even indices. The solid line represents an imaginary-time evolution $\tau \in [0, \beta/2]$. The black dot represents the boundary condition that $(\chi_{2j-1} + is_j\chi_{2j})|\{s\}\rangle = 0$, which can be solved by introducing a single $\chi_j(s)$ field with $s \in [0, \beta]$, where

$$\chi_{j}(s) = \begin{cases} \chi_{2j-1}(\beta/2 - s) & \text{for } s \in [0, \beta/2), \\ -is_{j}\chi_{2j}(s - \beta/2) & \text{for } s \in [\beta/2, \beta). \end{cases}$$
(52)




FIG. 4. The Rényi entanglement entropy $S_A^{(2)}(t)$ on KM pure state as a function of subsystem size λ for q=4 and $\beta=0$. The black dashed lines are $\lambda \log(2)/2$ and $(1-\lambda)\log(2)/2$. Adapted from [56].

This reduces the original problem for N Majorana fields with boundary conditions to N/2 Majorana fields with the smooth condition and doubled the contour length. Choosing the A subsystem containing M/2 pair of Majorana fermions, we can express $S_A^{(2)}(t)$ as

$$e^{-S_A^{(2)}(t)} = \frac{1}{Z_{KM}^2} \times A \qquad A$$
, (53)

Here we have defined the normalization of KM states $Z_{\rm KM} = \langle {\rm KM}|{\rm KM}\rangle$. The evolution contour involves both imaginary-time and real-time evolution, with a direction indicated by the arrows. Now we can parametrize the contour by a single real variable $s \in [0, 4(\beta+2t)]$. The saddle-point equation can be derived as

$$G_A = (\partial_s - \Sigma_A)_A^{-1}, \qquad G_B = (\partial_s - \Sigma_B)_B^{-1},$$

$$\Sigma_A = \Sigma_B = J^2 (\lambda G_A + (1 - \lambda) G_B)^{q-1} \tilde{P}.$$
(54)

Comparing to the thermal ensemble case (46), the differences are two-folds. Firstly, the boundary condition is now different. We have two separate contours for both subsystems. Secondly, there is an additional factor $\tilde{P}(s,s') = P(s,s')f(s)f(s')$. P(s,s') = 1 if $\chi_j(s)$ and $\chi_j(s')$ both correspond to original fields with even/odd indices and otherwise zero. This is due to the fact that random interactions for χ_{2j-1} and χ_{2j} are independent. The factor f(s) = 1 for imaginary-time evolutions, and $f(s) = \pm i$ for forward/backward real-time evolutions. The detailed derivation can be found in [56]. With the saddle-point solution, we can compute

$$\frac{I_{\text{KM}}^{(2)}(\lambda)}{N} = \frac{\lambda}{4} \log \det G_A + \frac{\lambda(q-1)}{4q} \int ds ds' G_A \Sigma_A$$

$$+ (\lambda \to 1 - \lambda, A \to B).$$
(55)

Comparing to (47), there is an additional factor of 1/2. This is because we are now counting Majorana modes in terms of N/2

pairs. The Rényi entanglement entropy is given by $S_A^{(2)}(t) = I_{KM}^{(2)}(\lambda) - I_{KM}^{(2)}(0)$.

As an example, we show numerical results for the $S_A^{(2)}(t)$ at $\beta = 0$ for different time tJ. In the short-time regime, the entanglement entropy is a smooth function of λ . However, at the longer time regime, there are two saddle-point solutions near $\lambda = 1/2$, with a first-order transition at $\lambda = 1/2$. (These solutions take the form of "replica wormholes", which will be explained in more details in section IV.) The final result of $S_A^{(2)}(t)$ is determined by the solution with smaller on-shell action $I_{\text{KM}}^{(2)}$. This leads to a singularity of $S_A^{(2)}$ at $\lambda = 1/2$. In the long-time limit, $S_A^{(2)}$ saturates to the maximal value, while each saddle-point gives a curve with entropy $\lambda \log(2)/2$ or $(1 - \lambda) \log(2)/2$. These values corresponds to the entropy of subsystem A or B when the total system is prepared in an thermal ensemble, which has been discussed above. This can be understood as follows [56]: Let us consider a general pure state $|\psi(t)\rangle = \sum_{c} c_n e^{-iE_n t} |E_n\rangle$, the entanglement entropy of A can be written as

$$e^{-S_A^{(2)}(t)} = \operatorname{tr}\left[\hat{S}^A|\psi(t)\rangle\langle\psi(t)|\otimes|\psi(t)\rangle\langle\psi(t)|\right]. \tag{56}$$

In the long-time limit, the eigenstates should pair up to cancel the fast oscillation phase factor. As a result, we find

$$e^{-S_A^{(2)}(t)} \approx \sum_{m,n} |c_m c_n|^2 \text{tr} \left[(\hat{S}^A + \hat{S}^B) |E_m\rangle \langle E_m| \otimes |E_n\rangle \langle E_n| \right]$$

$$= \text{tr} \left[(\hat{S}^A + \hat{S}^B) \rho_{\text{th}} \otimes \rho_{\text{th}} \right] = e^{-S_{A,th}^{(2)}} + e^{-S_{B,th}^{(2)}}.$$
(57)

Here we have assumed the diagonal ensemble looks thermal $\sum_m |c_m|^2 |E_m\rangle\langle E_m| = \rho_{\text{th}}$. In the path-integral representation, each of the term comes from a separate saddle point, and there is a first-order transition in the large-N limit. The equivalence between the subsystem entropy of a thermal ensemble and the long-time entanglement of KM pure state has also been directly proved in [56] using the path-integral approach.

D. Exact results for SYK₂

Before ending this section, we review exact results for the entanglement entropy in the SYK_2 and $cSYK_2$ random hopping models [52, 55, 113]. We would focus on the $cSYK_2$ model since the path-integral approach suggests the large-N entanglement entropy of the SYK_q model is given by one half of the $cSYK_q$ model result at half-filling.

The idea is to use the correlation matrix approach for the entanglement entropy of the free particles [114]. The ground-state correlation function of free fermion Hamiltonians satisfies the wick theorem, even if restricted to arbitrary subsystems. The validity of the wick theorem indicates the reduced density matrix takes the form of Gaussian states

$$\rho_A = e^{-K_A}/\text{tr}[e^{-K_A}], \qquad K_A = \sum_{ij \in A} K_{ij} c_i^{\dagger} c_j.$$
(58)

In other words, ρ_A is a thermal ensemble of an auxiliary quadratic Hamiltonian K_A , which is known as the Modular Hamiltonian or the entanglement Hamiltonian. The parameter K_{ij} can be fixed by matching the correlation matrix

 $C_{ij} = \langle c_i^{\dagger} c_j \rangle$ with $i, j \in A$. The simplest way is to transform into the diagonal basis of $C_{\eta\xi} = \sum_{ij} U_{\eta i}^* U_{\xi j} C_{ij}$, where $C_{\eta\xi} = C_{\eta}\delta_{\eta\xi}$. In the same basis, we expect K_A is also diagonal. Writing $K_A = K_{\eta}\delta_{\eta\xi}$, we have

$$C_{\eta} = \frac{e^{-K_{\eta}}}{1 + e^{-K_{\eta}}} = \frac{1}{e^{K_{\eta}} + 1}.$$
 (59)

The entanglement entropy of the system is then given by $S_A = -\text{tr } \rho_A \log \rho_A$, which reads

$$S_A = -\sum_{\eta} \left(C_{\eta} \log C_{\eta} + (1 - C_{\eta}) \log \left(1 - C_{\eta} \right) \right). \tag{60}$$

In the continuum limit, if we define a normalized distribution f(x) of eigenvalue $x = C_{\eta}$, we have

$$S_A = -M \int dx \left(x \log x + (1 - x) \log(1 - x) \right) f(x).$$

$$S_A^{(n)} = -\frac{M}{n - 1} \int dx \log \left(x^n + (1 - x)^n \right) f(x).$$
(61)

The only remaining task is to determine f(x) for the cSYK₂ model. Since the single-particle Hamiltonian is a random Hermitian matrix, its eigenstates form a Haar random unitary matrix U [52]. For a filling fraction $\kappa = Q/N$, first Q states are occupied. We parametrize U as [52]

$$U = \begin{pmatrix} V_{Q \times M} & V'_{Q \times (N-M)} \\ W_{(N-Q) \times M} & W'_{(N-Q) \times (N-M)} \end{pmatrix}^T, \tag{62}$$

and the correlation matrix restricted to the subsystem A is given by $C_{ij} = V^{\dagger}V$. It is known that C then forms a β -Jacobi ensemble with $\beta = 2$ [115]. The eigenvalue distribution f(x) is known to satisfy the Wachter law [116]:

$$f(x) = \frac{1}{2\pi\lambda} \frac{\sqrt{(\lambda_{+} - x)(x - \lambda_{-})}}{x(1 - x)} \Theta \left[(\lambda_{+} - x)(x - \lambda_{-}) \right] + \left(1 - \frac{\kappa}{\lambda} \right) \Theta(\lambda - \kappa) \delta(x).$$
(63)

with $\lambda_{\pm} = [\sqrt{\kappa(1-\lambda)} \pm \sqrt{\lambda(1-\kappa)}]^2$. For $\lambda \to 0$, (61) immediately predicts the maximal entropy (32), consistent with all previous discussions. For general λ , the entropy can be analytically obtained for n=1 and all even n [55, 113], and also for finite N [117]. They show that the entanglement entropy is much smaller than the Page value, as expected for non-interacting systems.

IV. ENTANGLEMENT IN COUPLED SYK MODELS

Having summarized results for the SYK/cSYK model, in this section, we review studies of the entanglement dynamics in coupled SYK models. We focus on the coupled $\chi-\psi$ system with the Hamiltonian (27) for different $r=N_2/N_1$. We copy the Hamiltonian here for convenience:

$$H = \sum_{\{i_n\}} J^{\chi}_{i_1 i_2 i_3 i_4} \chi_{i_1} \chi_{i_2} \chi_{i_3} \chi_{i_4} + \sum_{\{a_n\}} J^{\psi}_{a_1 a_2 a_3 a_4} \psi_{a_1} \psi_{a_2} \psi_{a_3} \psi_{a_4}$$

$$+ \sum_{i_1 < i_2, a_1 < a_2} V_{i_1 i_2 a_1 a_2} \chi_{i_1} \chi_{i_2} \psi_{a_1} \psi_{a_2}.$$

$$(64)$$

Unlike the single SYK model, where many approaches can be employed, here the path-integral approach is the only efficient method. Motivated by gravity calculations [20], which would be reviewed in section IV D, the coupled system is prepared in a TFD state with an auxiliary system and evolves under the coupled Hamiltonian H [118]. In the early-time regime, the entanglement entropy grows linearly for both Rényi and von Neumann entropies. In the long-time limit, Rényi entropy saturates to a non-thermal value for r = 1. On the other hand, for $r \gg 1$, the ψ system contains large degrees of freedom and serves as a heat bath [48, 119, 120]. The coupled system is then qualitatively similar to [61]

$$H = H_{\text{SYK}_4}[\chi] + i \sum_{x,i} \frac{\Lambda}{2} \psi_{i,x} \psi_{i,x+1} + \sum_i i V \sqrt{\Lambda} \chi_i \psi_{i,0}, \quad (65)$$

which is closely related to its gravity counterpart in AdS_2 spacetime [20]. As a result, a replica wormhole solution appears in the long-time limit, and the entanglement entropy approaches the thermal value [66]. These results will be explained in details in the following subsections.

A. The set-up and $G-\Sigma$ action

We begin with the setup of the problem. TFD states can be defined generally by replicating the system [59], with arbitrariness in choosing the basis. Here we focus on a convenient convention for Majorana fermions. We consider a system that contains N Majorana modes χ_i with Hamiltonian $H[\chi]$. We introduce 2N fermion modes χ_i^L, χ_i^R . Similar to the definition of KM pure states, we pair up the left and the right fermions by defining $c_i = \frac{\chi_i^L + i \chi_i^R}{\sqrt{2}}$. The unnormalized TFD state reads

$$|\text{TFD}\rangle = e^{-\frac{\beta H_L}{2}}|I\rangle, \quad c_i|I\rangle = 0.$$
 (66)

Here we choose a single maximal entangled state $|I\rangle$ with no filling of c_i , and keep the β dependence of $|\text{TFD}\rangle$ implicit. We have defined the Hamiltonian for the left system as $H_L = H[\chi^L]$. We could also introduce the Hamiltonian for the right system as $H_R = H[-i\chi^R]$, which satisfies $H_R|I\rangle = H_L|I\rangle$, and the TFD state can be equivalently prepared by evolving $|I\rangle$ with $H_{\text{tot}} = H_L + H_R$ for an imaginary time $\beta/4$, or with H_R for an imaginary time $\beta/2$.

We add a few comments. Firstly, any maximally entangled state can be written as $|I\rangle=2^{-N/4}\sum_{i=1}^{2^{N/2}}|i\rangle_L|i'\rangle_R$, where both $\{|i\rangle_L\}$ and $\{|i'\rangle_R\}$ are orthonormal bases in the corresponding Hilbert space. As a result, the reduced density matrix of a single side is thermal $\rho^\alpha=e^{-\beta H^\alpha}/Z$ with $\alpha=L,R$. Secondly, for small β , $|\text{TFD}\rangle$ mainly contains entanglement between left and right copies, while the entanglement between Majorana modes in the same copy is weak. This indicates the TFD state is not thermal, similar to the KM pure states. After long-time evolution, we expect the system thermalize and $S_A^{(2)}=\min\{2S_{A,\text{th}}^{(2)},2S_{B,\text{th}}^{(2)}\}$. Here $S_{A/B,\text{th}}^{(2)}$ is the subsystem Rényi entropy of thermal density matrices. Thirdly, under holographic duality, the TFD states are described by the

Hartle-Hawking states in the two-sided black holes [112], in which a version of the information paradox exists [20].

To study the entanglement dynamics, we further evolve the coupled system with H_{tot} . This gives


$$|\text{TFD}(t)\rangle = e^{-iH_{\text{tot}}t}|\text{TFD}\rangle = e^{-2iH_Lt}|\text{TFD}\rangle.$$
 (67)

Then one can compute the reduced density matrix ρ_A and Rényi entanglement entropy $S_A^{(n)}$.

In this section, we focus on coupled SYK models with both χ and ψ fermions. After doubling the system, we have four species of fermions χ^L , ψ^L , χ^R , and ψ^R . The Hamiltonian H_α contains interactions between χ^α and ψ^α , and there is no interaction between left and right copies. The TFD state after the real-time evolution can be represented pictorially

$$|\text{TFD}\rangle = \chi$$
 (68)

Here the horizontal solid lines represent real-time evolutions, while the arcs represent imaginary-time evolutions. We choose the subsystem A as $\psi^L \cup \psi^R$, which is in a pure state when $\beta = t = 0$. The purity tr ρ_A^2 then takes two equivalent forms [61]



The left graph can be understood as inserting a pair of twist operators $T_{\chi,L}/T_{\chi,R}$ to the χ system:

$$Z^{2}\operatorname{tr}\rho_{A}^{2} = \langle \operatorname{TFD}(t)|^{\otimes 2}T_{\chi,L}T_{\chi,R}|\operatorname{TFD}(t)\rangle^{\otimes 2}.$$
 (70)

The right graph is more convenient for the path-integral approach. Following the discussions in section II A by introducing bilocal fields $G_{\chi/\psi}^{mn}$ and $\Sigma_{\chi/\psi}^{mn}$ with $m,n\in\{1,2\}$, one find the G- Σ action reads

$$\frac{S_{\text{TFD}}^{(2)}}{N_{1}} = -\frac{1}{2} \log \det(\partial_{\tau} - \tilde{\Sigma}_{\chi}) - \frac{r}{2} \log \det(\partial_{\tau} - \tilde{\Sigma}_{\psi}) \\
+ \frac{1}{2} \int_{\tau\tau'} \left[\tilde{\Sigma}_{\chi}^{mn} \tilde{G}_{\chi}^{mn} - \frac{J^{2}}{4} (\tilde{G}_{\chi}^{mn})^{4} + r \tilde{\Sigma}_{\psi}^{mn} \tilde{G}_{\psi}^{mn} \right. (71) \\
- \frac{J^{2}r}{4} (\tilde{G}_{\psi}^{mn})^{4} - \frac{V^{2}}{2} (\tilde{G}_{\chi}^{mn})^{2} (\tilde{G}_{\psi}^{m'n'})^{2} c_{mm'} c_{nn'} \right].$$

Here the integrals over τ and τ' are over the contour $C_1 \cup C_2$ on the complex plane. An additional phase factor appears if we use a single real parameter as we did in section III C. We have defined $c_{mm'}(\tau)$ and $c_{nn'}(\tau')$ as a permutation between two replicas [59, 98]:

$$c(\tau) = \begin{cases} \mathbf{I} & \text{for } \tau \in C_1, \\ \boldsymbol{\sigma}_{\boldsymbol{x}} & \text{for } \tau \in C_2. \end{cases}$$
 (72)

The saddle-point equations of (71) read

$$G_{\chi} = (\partial_{\tau} - \Sigma_{\chi})^{-1}, \qquad G_{\psi} = (\partial_{\tau} - \Sigma_{\psi})^{-1},$$

$$\Sigma_{\chi}^{mn} = J^{2} (G_{\chi}^{mn})^{3} + V^{2} c_{mm'} c_{nn'} (G_{\psi}^{m'n'})^{2} G_{\chi}^{mn},$$

$$\Sigma_{\psi}^{mn} = J^{2} (G_{\psi}^{mn})^{3} + V^{2} c_{mm'} c_{nn'} (G_{\chi}^{m'n'})^{2} G_{\psi}^{mn}/r,$$
(73)

and the on shell action $I_{\text{TFD}}^{(2)} = S_{\text{TFD}}^{(2)}[G, \Sigma]$. The second Rényi entropy is given by $S_A^{(2)} = I_{\text{TFD}}^{(2)} - 2I$, with I being the onshell action of the thermal ensemble. Equivalently, 2I can be computed by setting $c(\tau) = \mathbf{I}$ in (71).

B. Early-time linear growth

We begin to analyze the entanglement dynamics of the coupled system. An important observation [59] is that the saddlepoint equations (73) admit solutions that is "replica diagonal":

$$G_{\psi/\chi}^{mn}(\tau,\tau') = \delta^{mn} G_{\psi/\chi}(\tau,\tau'). \tag{74}$$

The arguments contain two steps: Without any interaction V=0, the solution of $G_{\psi/\chi}$ satisfies the replica diagonal condition. When we turn on a finite V, new terms in the self-energy is again replica diagonal due to the presence of $G_{\psi/\chi}$. As a result, the replica diagonal solution is consistent with the saddle-point equation. It turns out that this is the correct saddle-point solution in the early-time regime.

Now let's assume V is small, and perform a perturbative calculation of the entanglement entropy near the saddle-point solution with V = 0 [22, 59, 62, 63, 66]. We have

$$\frac{\delta I_{\text{TFD}}^{(n)}}{nN_1} = -\frac{V^2}{4} \left(\int_{\tau, \tau' \in C_1} + \int_{\tau, \tau' \in C_2} \right) G_{\chi}(\tau, \tau')^2 G_{\psi}(\tau, \tau')^2. \tag{75}$$

Here we consider the *n*-th Rényi entropy. On the other hand, there is a similar contribution for *I*:

$$\frac{\delta I}{N_1} = -\frac{V^2}{4} \int_{\tau, \tau' \in C_1 \cup C_2} G_{\chi}(\tau, \tau')^2 G_{\psi}(\tau, \tau')^2. \tag{76}$$

Combining these results, we have

$$S_A^{(n)} = \frac{nV^2 N_1}{2(n-1)} \int_{\tau \in C_1} \int_{\tau' \in C_2} G_{\chi}(\tau, \tau')^2 G_{\psi}(\tau, \tau')^2.$$
 (77)

For V=0, the Green's functions only depend on time differences. When the evolution time $t \ge t_{\rm dis}$, with the dissipation time $t_{\rm dis}$ charactering the decay of two-point functions, the main time-dependent contribution $\Delta S_A^{(n)}(t) = S_A^{(n)}(t) - S_A^{(n)}(0)$ comes from the integral over real-time evolutions. This gives

$$\Delta S_A^{(n)}(t) \approx \frac{nV^2 N_1}{2(n-1)} \int_0^{2t} \int_0^{2t} dt_1 dt_2 G_\chi^{>}(t_{12})^2 G_\psi^{>}(t_{12})^2$$

$$\approx \left[\frac{nV^2 N_1}{n-1} \int_{-\infty}^{\infty} dt' G_\chi^{>}(t')^2 G_\psi^{>}(t')^2 \right] t \equiv \kappa t.$$
(78)

Here $G_{\chi}^{>}(t) = -i \langle \chi(t) \chi(0) \rangle$ is the real-time Green's function [108]. We find at early-time the entanglement entropy grows linearly.




FIG. 5. An illustration of the geometric problem on the Poincaré disk for the dynamics of Rényi entropy with $N_1 = N_2 = N$ in the low-energy limit. The saddle-point solution is given by joining two arcs. Adapted from [59].

Such an linear-growth behavior is valid beyond SYK-like models and the replica diagonal consistency is not necessary. As an illustration, we consider some arbitrary χ and ψ systems, with a general coupling $H_{\text{int}} = V \sum_{i=1}^{N} O_{\chi}^{i} O_{\psi}^{i}$ [63]. The perturbative analysis can still be carried out diagrammatically. For n-th Rényi entropy with $n \geq 2$, the leading order time-dependent contribution comes from the diagram

$$\psi^{1} \qquad \psi^{2} = -\frac{nV^{2}N}{n-1} \operatorname{tr} \left[G_{O_{\chi}}^{>} \circ G_{O_{\psi}}^{>,T} \right].$$
(79)

Here we treat Green's functions as matrices in the time domain. This leads to a linear growth

$$\Delta S_A^{(n)}(t) \approx \left[-\frac{2nV^2N}{n-1} \int \frac{d\omega}{2\pi} G_{O_\chi}^{>}(\omega) G_{O_\psi}^{>}(-\omega) \right] t. \tag{80}$$

However, above calculations do not have a well-defined limit $n \to 1$, in which the Rényi entropy $S_A^{(n)}$ reduces to the von Neumann entropy S_A . As noticed by authors [63], this is because an additional diagram contributes for $n \to 1$:

$$\psi^{1} = -\frac{V^{2n}N}{n-1} \left[-G_{O_{\chi}}^{>} \circ G_{O_{\psi}}^{>,T} \right]^{n}.$$
 (81)

Although this contribution is at a higher order of V for any n > 1, It becomes important in the limit of $n \to 1$. Summing up (79) and (81), we get a finite expression for the von Neumann entropy, which predicts a slope

$$\frac{\Delta S_A}{N} \approx \left[2 \int \frac{d\omega}{2\pi} g(\omega) \left(1 - \log[g(\omega)] \right) \right] t, \tag{82}$$

with $g(\omega) \equiv -V^2 G_{O_\chi}^>(\omega) G_{O_\psi}^>(-\omega)$. Interestingly, this shows that the linear growth rate for the von Neumann entropy is proportional to $-V^2 \log V$ for small V, parametrically faster than any n-th Rényi entropy with n > 1.

C. Geometric interpretation for $N_2 = N_1$

Now we turn to the entanglement dynamics beyond the early-time regime. Generally, this requires direct numerical studies of the saddle-point equations (73). In this subsection, we focus on the special limit with $N_1 = N_2 = N$, $V \ll J$, and $\beta J \rightarrow \gg 1$, where analytical results exist [59]. The χ and ψ systems now become symmetric. Following the discussions in [59], we further assume the replica diagonal solution also works at late times [121]. This has been verified in numerics [60, 61].

Since $V \ll J$, the perturbation due to the coupling between two sites are small. As a result, one can assume the changes of Green's functions are restricted to the low-energy manifold. In other words, we can use the Schwarzian action (16) to determine the entanglement dynamics. Similar to the previous subsection, the effective action contains two parts $S_{\rm eff} = S_{\rm Sch} + S_{\rm int}$, with

$$\frac{S_{\text{Sch}}}{n} = -\frac{N\alpha_S}{\mathcal{J}} \int_{C_1 \cup C_2} d\tau \left\{ \tan \frac{\pi f(\tau)}{\beta}, \tau \right\},$$

$$\frac{S_{\text{int}}}{n} = \frac{V^2}{2} \int_{C_1} d\tau_1 \int_{C_2} d\tau_2 \, \tilde{G}_f(\tau_1, \tau_2)^4.$$
(83)

We have assumed the only relevant mode satisfies $\tilde{G}_{\psi}(\tau_1,\tau_2)=\tilde{G}_{\chi}(\tau_1,\tau_2)=\tilde{G}_{f}(\tau_1,\tau_2)$. The integral is along a contour on the complex plane. To avoid this, we perform the analytical continuation $t\to -i\tau$, and then the integral is purely on the imaginary time. We choose $C_1\in [\tau,\beta/2-\tau)$ and $C_2\in [0,\tau)\cup [\beta/2-\tau,\beta)$. A second analytical continuation back to the real time $\tau\to it$ is performed at the end of the calculation. This gives

$$\frac{S_{\text{int}}}{n} = \frac{\gamma}{2} \log \frac{\beta^2 \sin^2 \frac{\pi}{\beta} (f(\beta/2 - \tau) - f(\tau))}{\pi^2 \epsilon^2 f'(\beta/2 - \tau) f'(\tau)}.$$
 (84)

Here $\gamma = V^2/4\pi J^2$ and $\epsilon \sim 1/\mathcal{J}$ is a cutoff.

The problem of solving (83) and (84) can be mapped to a geometric problem on a Poincaré disk thanks to the holographic duality [59]. For a curve $(\rho(\tau), \theta(\tau))$ on a Poincaré disk with $\theta(\tau) = 2\pi f(\tau)/\beta$ and $g_{\tau\tau}(\tau) = \mathcal{J}^2$, the Schwarzian action can be represented by

$$\frac{S_{\text{sch}}}{nN} = \alpha_S(\beta \mathcal{J} - A - 2\pi). \tag{85}$$

Here A is the area enclosed by the curve. This is a consequence of (17) and the Gauss-Bonnet theorem [122], and a direct verification is given in [59]. On the other hand, S_{int} represents an attractive force between two boundary points

$$\frac{S_{\text{int}}}{nN} = \frac{\gamma}{2} \log \cosh D \left(\frac{\beta}{2} - \tau, \tau \right). \tag{86}$$

Here $D(\tau_1, \tau_2)$ is the geodesic distance between two points $(\rho(\tau_1), \theta(\tau_1))$ and $(\rho(\tau_2), \theta(\tau_2))$ on the curve. Intuitively, this can be understood by realizing that the combination in (84) takes the form of the two-point function, which can be approximated by $e^{-D(\tau_1, \tau_2)}$.




FIG. 6. Numerical results for the entanglement dynamics of coupled SYK models. (a). 2-nd Rényi entropy for Hamiltonian (64) with $\beta J=4, r\to\infty, N_1=N$ and V/J=1; (b). 2-nd Rényi entropy for Hamiltonian (65) with $\beta J=4, V^2/J=0.05, N_1=N$ and $\Lambda/J=5$; (c). Typical Green's function Re G_χ in the early-time regime; (d). Typical Green's function Re G_χ in the late-time regime. (a), (c), and (d) are adapted from [61].

The entanglement dynamics is determined by minimizing $S_{\rm sch} + S_{\rm int}$ with respect to the shape of the curve $f(\tau)$. Away from τ and $\beta/2 - \tau$, the equation of motion is the same as V = 0, under which the solution is a circle. As a result, the solution takes the form of joining two arcs (up to an SL(2,R) transformation on the Poincaré disk). The minimization has been worked out in [59] by expressing both terms through the radius and open-angle of each arc. In the long-time limit, the authors find

$$\frac{n-1}{nN}S^{(n)} \approx \frac{4\pi\alpha_S}{\beta\mathcal{J}} + \gamma \left(1 + \log\frac{8\sqrt{2}\alpha_S}{\gamma}\right). \tag{87}$$

If we take $\gamma \to 0$, this reduces to the excitation energy of the SYK model, which does not contain the zero-temperature entropy S_0 . This may lead to the conclusion that the coupled SYK model fail to thermalize for $N_1 = N_2$. However, this may be an artifact of the large-N limit for Rényi entropies, and the von Neumann entropy may show parametrically different behaviors as we have seen in the last subsection. For example [123], if we consider a toy density matrix

$$\rho = \frac{P}{P+Q}|E_1\rangle\langle E_1| + \frac{1}{P+Q}\mathbf{I}_{Q\times Q},$$

when $P, Q \gg 1$, We have

$$S^{(n)} = -\frac{1}{n-1} \log \frac{P^n + Q}{(P+Q)^n} \approx \frac{n}{n-1} \log \frac{P+Q}{P},$$

$$S = \frac{P}{P+Q} \log \frac{P+Q}{P} + \frac{Q}{P+Q} \log(P+Q).$$
(88)




FIG. 7. An illustration of the gravity calculation of the von Neumann entropy: (a). The RT surface without an island, and (b). The RT surface with an island outside the horizon.

Here we have assumed $P^2 \gg Q$. $|E_1\rangle$ is an analog of the excited state, and $\mathbf{I}_{\mathcal{Q}\times\mathcal{Q}}$ is an analog of the SYK ground state ensemble. We see that although $S^{(n)}$ only contains the contribution from $|E_1\rangle$ if we take the large P, Q limit, the von Neumann entropy S contains both contributions. It is also known that one can get rid of the non-thermal behavior by considering the microcanonical ensemble [22].

One could also ask about results at higher temperatures. In this case, the quantum dynamics is no longer restricted to the low-energy manifold, and the numerical study [60, 61] shows that the entanglement entropy saturates to the thermal value through a first-order transition. We will review similar physics in the next subsection.

D. Thermalization and replica wormholes

Now we turn to the situation where the ψ system contains much more degrees of freedom $r \gg 1$. In this case, the Green's function of ψ receives no correction from V (73) and we can fix the G_{ψ} by thermal two-point functions. Moreover, we can neglect all \tilde{G}_{ψ} and $\tilde{\Sigma}_{\psi}$ terms in the G- Σ action as

$$\frac{S_{\text{TFD}}^{(2)}}{N} = -\frac{1}{2} \log \det(\partial_{\tau} - \tilde{\Sigma}_{\chi}) + \frac{1}{2} \int_{\tau\tau'} \left[\tilde{\Sigma}_{\chi}^{mn} \tilde{G}_{\chi}^{mn} - \frac{J^{2}}{4} (\tilde{G}_{\chi}^{mn})^{4} - \frac{V^{2}}{2} (\tilde{G}_{\chi}^{mn})^{2} (G_{\psi}^{m'n'})^{2} c_{mm'} c_{nn'} \right].$$
(89)

This is because they cancel out with corresponding terms in 21. Here we set $N_1 = N$ for conciseness. (89) can be understood as firstly integrating out the ψ system and focus on the response of χ fermions. For the model with a bath described by a Majorana chain (65), the only difference is to replace the last term in the square bracket by [61]

$$-V^2\Lambda \tilde{G}_{\chi}^{mn}G_{\psi}^{m'n'}c_{mm'}c_{nn'}, \qquad (90)$$

with $G_{\psi}^{m'n'}$ computed using the free Majorana chain. The similarity between $G\text{-}\Sigma$ actions indicates two models show qualitatively the same dynamics of the entanglement entropy. The analysis in section (IVB) shows the entanglement entropies of both models grow linearly under perturbation. When perturbation theory fails, one should perform a numerical study of Schwinger-Dyson equations. We parametrize the

contour using a single real parameter $s \in [0, 4(2t + \beta/2)]$, similar to the study of KM pure states. The results are shown in Figure 6 (a) for the SYK₄ bath and (b) for the chain bath. The results are qualitatively the same. In the early-time regime, the entanglement entropy grows following a smooth curve with a nearly constant slope. This persists to a very long time even when $S_A^{(2)} > 2S_{A,\text{th}}^{(2)}$, which leads to a negative (second Rényi) mutual information $S_{\psi^L}^{(2)} + S_{\psi^R}^{(2)} - S_{\psi^L \cup \psi^R}^{(2)} = 2S_{A,\text{th}}^{(2)} - S_A^{(2)} < 0$. Such a violation of quantum information bounds is known as the information paradox [70]. Fortunately, an additional saddle-point solution appears in the intermediate time regime, which dominates the path-integral and resolves the paradox in the long-time limit by a first-order transition. Similar transitions also show up in Brownian models [64, 92].

To understand the difference between two saddle-point solutions, we plot the typical Green's functions G_{ν} in Figure 6 (c) and (d). Here we use the convention illustrated in (69), where replica diagonal solutions of G_{ν} and G_{ψ} take the form

$$G_{\chi}^{\mathrm{diag}}:$$
 $G_{\psi}^{\mathrm{diag}}:$ (91)

Since G_{ψ} does not depends on V, it is always replica diagonal and we only need to focus on G_{χ} . On the early-time saddle, we expect the Green's functions take a similar form as V = 0. For the SYK₄ bath, this suggests that G_{ψ} is strictly replica diagonal on the early-time saddle. For the Majorana chain bath, the replica diagonal assumption is not consistent with a finite V. As a result, G_{ψ} is only approximately replica diagonal, as shown in Figure 6 (c).

However, G_{ψ} takes a complete different form on the longtime saddle point, as shown in Figure 6 (d). The solution approximately takes the form of G_{ψ}^{diag} . This suggests that although the contour of $s \in [0, \beta/2 + 2t)$ is connected with $s \in [3(\beta/2 + 2t), 4(\beta/2 + 2t))$, the solution favors the pairing between $s \in [0, \beta/2+2t)$ and $s \in [\beta/2+2t, 2(\beta/2+2t))$. Due to the reasons explained below, we call this a replica wormhole solution. As we mentioned in section III C, similar transitions also present when considering the entanglement dynamics of KM pure states. In [61], authors explains the replica wormhole solutions can be understood as a factorization of twist operators, which gives

$$e^{-S_A^{(2)}} = \langle T_{\chi,L} T_{\chi,R} \rangle \approx \langle T_{\chi,L} \rangle \langle T_{\chi,R} \rangle = e^{-2S_{A,th}^{(2)}}.$$
 (92)

The replica wormholes were introduced in the entanglement calculation of holographic systems [21, 22]. Towards a gravity analog of the SYK4 fermions coupled to a Majorana chain, we can replace the SYK model with a JT gravity in the AdS₂ spacetime. The Majorana chain corresponds to a bath in flat spacetime without gravity dynamics. An SYK₄ fermion has a scaling dimension of $\Delta = 1/4$, which is dual to a massive bulk field. However, massive fields make the analytical calculation hard. In the gravity calculation [20], authors assume there is a conformal field theory in both regions, with transparent boundary conditions. The system is prepared in

the TFD state, which is dual to a two-sided black hole, and evolved under the total Hamiltonian H^{tot} . An illustration of the geometry is shown in Figure 7.

The von Neumann entanglement entropy of boundary region A can be computed using the RT formula [11–16]:

$$S_A = \min \left[\text{ext}_{\gamma_A} \left(\frac{\text{Area}(\gamma_A)}{4G_N} + S_{\text{bulk}} \right) \right].$$
 (93)

The first term is the geometric term, and the second term is the entanglement of bulk quantum fields. Here we should consider all possible bulk extremal surfaces γ_A homologous to A, and choose the minimal one. In our case, the system is 1+1-D, and the RT surface is a set of points. The area term is replaced by the expectation of the dilaton field [13]. The naive RT surface only contains points that separate gravity systems and baths, as illustrated in Figure 7 (a). Since there is no gravity dynamics in the bath, only S_{bulk} contributes. A CFT calculation shows the result grows linearly in the long-time limit [20], due to the establish of the entanglement of bulk fields. Similar to the SYK calculation, this gives rise to an information paradox.

The resolution is to add the contribution in Figure 7 (b). Here the RT surface contains additional points in the gravity region, which forms an island. Without bulk quantum fields, islands always lead to an increase of (93) because the area increases. However, in the current calculation, this considerably reduces the entanglement of bulk fields. The explicit calculation shows the long-time limit is then a constant, which approaches $2S_{A,\text{th}}^{(2)}$. The presence of the island can be dated back to the gravity path-integral with multiple replicas [21, 22]. For the naive saddle point, the bath is twisted while the gravity region is untwisted. This indicates the way of pairing different contours is different for the gravity system and the bath. The island can be viewed as emergent dynamical twist operators. After the island appears, most of spacetime regions are twisted, and the contours for the gravity system and the bath are paired similarly. This is consistent with the SYK result. These emergent twist operators connect gravity systems in different replicas through wormholes, which are known as "replica wormholes" [21, 22]. When the island presents, the entanglement wedge reconstruction [124] implies part of the gravity region will be accessible to the bath. Different concrete protocols to extract information have been proposed in [22] and [125].

E. Entanglement dynamics of SYK chains

We finally extend the discussions for the coupled χ - ψ systems to SYK chains [42]. On each site, there is an SYK₄ model, and they are coupled by SYK-like random pair hopping terms. The Hamiltonian reads

$$H = \sum_{x} \left(\sum_{i_{1} < i_{2} < i_{3} < i_{4}} J^{x}_{i_{1}i_{2}i_{3}i_{4}} \chi_{x,i_{1}} \chi_{x,i_{2}} \chi_{x,i_{3}} \chi_{x,i_{4}} \right.$$

$$+ \sum_{i_{1} < i_{2},j_{1} < j_{2}} V^{x}_{i_{1}i_{2}j_{1}j_{2}} \chi_{x,i_{1}} \chi_{x,i_{2}} \chi_{x+1,j_{1}} \chi_{x+1,j_{2}} \right).$$

$$(94)$$

Here $x \in [1, L]$ with periodic boundary condition. We prepare the system in a TFD state with an auxiliary system. Entanglement dynamics of KM states is similar. We choose the subsystem A as the first $L_A \le L/2$ sites.


The first contribution to the Rényi entropy is the (approximate) replica diagonal solution. Following the discussions in section IV B, the entanglement entropy will grow linearly for $L_A \geq 1$, with, a slope v_E independent of subsystem size. Choosing the same convention as the last subsection, the Green's functions G_x look like G_χ^{diag} if $x \in A$, and G_ψ^{diag} if $x \in B$. Moreover, Green's functions in the bulk of A/B system approach two copies of the thermal Green's function.

The second contribution is the replica wormhole solution, where all Green's functions G_x look like G_ψ^{diag} . However, due to the mismatch between connectivity of the contour and the pairing indicated by the Green function, small deviations from G_ψ^{diag} always exist near the insertion time of twist operators (as in Figure 6 (d)), even in the bulk of subsystem A. As a result, the replica wormhole solution leads to a volume law phase with $S_A^{(2)} = s_0 L_A$. Putting both contributions together, the entanglement dynamics satisfies

$$S_A^{(2)}(t, L_A) = \begin{cases} v_E t & \text{if} \quad t < s_0 L_A / v_E \\ s_0 L_A & \text{if} \quad t \ge s_0 L_A / v_E \end{cases}$$
(95)

This is consistent with numerical results in [92].

Interestingly, this calculation can formulated as a classical spin problem. We can treat the choice of the Green's functions as a spin degree of freedom. To be concrete, we label G_{ψ}^{diag} as 1 and G_{χ}^{diag} as σ . The boundary condition is fixed by twist operators: (69) suggests we should impose 1 for $x \in B$ and σ for $x \in A$ near s = 0 and $\beta/2 + 2t$. The replica diagonal and replica wormhole solution can then be illustrated as



The replica diagonal solution corresponds to domain walls along the time-direction, the energy cost of which grows linearly. The replica wormhole solution corresponds to having domain walls parallel to the spatial direction, which indicates a volume law entanglement entropy. Similar picture was proposed in random tensor networks [126] and random circuits [127, 128]. As we will see in the next subsection, such a picture gives a natural understanding of measurement induced phase transitions.

V. ENTANGLEMENT IN NON-UNITARY SYK MODELS

Recent years, it was realized novel quantum entanglement dynamics can be achieved for "hybrid" quantum systems, in which evolutions involve both unitary evolutions and measurements [72–98]. When tunning the measurement rate, the steady states can show entanglement phase transitions between volume law entangled phases and area law entangled phases, separated by a critical point with logarithmic entanglement entropy if the transition is second order.

Since SYK-like models are solvable, they are ideal platforms for understanding the entanglement phase transitions [92–98]. In this section, we review the studies of non-unitary SYK models, focusing on continuous forced measurements which are equivalent to non-Hermitian Hamiltonian dynamics. A general measurement is described by a set of Kraus operators $\{M_{\xi}\}$ that satisfies $\sum_{\xi} M_{\xi}^{\dagger} M_{\xi} = I$. Let us assume the system is prepared in some state $|\psi_0\rangle$. After performing the measurement, the probability p_{ξ} of getting the result ξ and the corresponding post-selected state $|\psi_{\xi}\rangle$ read

$$p_{\xi} = \langle \psi_0 | M_{\xi}^{\dagger} M_{\xi} | \psi_0 \rangle, \qquad | \psi_{\xi} \rangle = \frac{1}{\sqrt{p_{\xi}}} M_{\xi} | \psi_0 \rangle. \tag{97}$$

Now we focus on the special case where $\xi = 1, 2$ with $M_1 \approx I - \epsilon H_1$ and $M_2 \approx \sqrt{2\epsilon} H_1$, with Hermitian and positive semidefinite H_1 and $\epsilon \ll 1$. We now post select the measurement outcome with $\xi = 1$. This leads to

$$|\psi_{\mathcal{E}}\rangle \propto (1 - \epsilon H_{\rm I}) |\psi_0\rangle \approx e^{-\epsilon H_{\rm I}} |\psi_0\rangle.$$
 (98)

This is equivalent to an imaginary-time evolution of the initial state. Taking the continuum limit by setting $\epsilon = dt$ and adding the unitary evolution part with Hamiltonian H_R , we find

$$|\psi(T)\rangle \propto e^{-iHT}|\psi_0\rangle, \qquad H = H_R - iH_I.$$
 (99)

We choose the subsystem A as first L_A sites. The n-th Rényi entropy reads

$$e^{-(n-1)S_A^{(n)}} = \frac{\operatorname{tr}_A \left(\operatorname{tr}_B e^{-iHT} |\psi_0\rangle \langle \psi_0| e^{iH^{\dagger}T} \right)^n}{\left(\langle \psi_0| e^{iH^{\dagger}T} e^{-iHT} |\psi_0\rangle \right)^n}.$$
 (100)

Main previous studies focus on n = 2 [92–97]. The von Neumann entropy limit $n \to 1$ has been studied in [98] with further assumptions.

In the following subsections, we will consider different non-Hermitian SYK chains and derive effective actions that govern the entanglement dynamics of the 2-nd Rényi entropy. An important ingredient is the emergence of conformal symmetry in replicated non-interacting systems for small measurement rates, which leads to logarithmic entanglement entropy. After adding interactions, the gapless modes become gapped, and the volume law entanglement replaces the logarithmic law entanglement. We can drive transitions to area law phases in certain setups for a larger measurement rate. We will also discuss the consequence of adding long-range couplings.

A. Emergent replica conformal symmetry

The non-unitary dynamics of free fermions has been studied numerically [90], and the result shows logarithmic entanglement entropy with emergent conformal symmetry. In this




FIG. 8. (a-b). A sketch of the path-integral contour without/with twist operators for subsystems A. (c). The configuration of θ when computing the entanglement entropy in non-Hermitian SYK₂ chains. (d). The configuration of θ for the purification dynamics in non-Hermitian SYK₂ chains. (c), (d) are adapted from [95].

subsection, we review the progress of understanding these results using SYK solvable models. We consider the non-interacting SYK₂ chains [93]. The Hamiltonian reads [129]

$$H_{R} = i \sum_{x} \left[\sum_{j,k} J_{jk}^{x}(t) \chi_{x,j} \chi_{x+1,k} + \sum_{j < k} \tilde{J}_{jk}^{x}(t) \chi_{x,j} \chi_{x,k} \right],$$

$$H_{I} = i \sum_{x} \left[\sum_{i,k} V_{jk}^{x}(t) \chi_{x,j} \chi_{x+1,k} + \sum_{i < k} \tilde{V}_{jk}^{x}(t) \chi_{x,j} \chi_{x,k} \right].$$
(101)

For simplicity, we choose couplings to be Brownian:

$$\overline{J_{ij}^{x}(t)J_{ij}^{x}(0)} = \frac{J_{1}}{2N}\delta(t), \quad \overline{J_{ij}^{x}(t)J_{ij}^{x}(0)} = \frac{J_{0}}{N}\delta(t),
\overline{V_{ij}^{x}(t)V_{ij}^{x}(0)} = \frac{V_{1}}{2N}\delta(t), \quad \overline{V_{ij}^{x}(t)V_{ij}^{x}(0)} = \frac{V_{0}}{N}\delta(t).$$
(102)

The static coupling case has been analyzed similarly in [93].

In section IV E, we have seen that the entanglement entropy calculation in interacting systems with unitary evolution corresponds to the energy of domain walls. We expect similar pictures work here: the twist operator creates some defects which excite low-energy excitations, and the entanglement entropy is given by the corresponding energy. To understand the possible low-energy excitations, we start with the path-integral without any twists $\lim_{T\to\infty} (\operatorname{tr} e^{iHT} e^{iHT})^2$. We label four evolution contours by 1, 2, 3, 4, where the evolution is forward/backward on $\{1,3\}/\{2,4\}$, as illustrated in Figure 8 (a). Following the derivation in section II A, the saddle-point equation reads

$$\left[(-1)^{a+1} \delta^{ac} \partial_t - \Sigma_x^{ac} \right] \circ G_x^{cb} = I^{ab}, \tag{103}$$

and

$$\Sigma_{x}^{ab}(t,t') = \left[(V_{1}^{2} - (-1)^{a+b} J_{1}^{2}) \frac{G_{x+1}^{ab}(t,t) + G_{x-1}^{ab}(t,t)}{2} + (V_{0}^{2} - (-1)^{a+b} J_{0}^{2}) G_{x}^{ab}(t,t) \right] \delta(t-t').$$
(104)

Interestingly, there is a $O(2) \otimes O(2)$ symmetry: $G(t,t') \rightarrow OG(t,t')O^T$ with $O = \exp(-\gamma_{13}\theta_{13} - \gamma_{24}\theta_{24})$ left the equation invariant. Here $(\gamma_{cd})^{ab} = \delta_{ac}\delta_{bd} - \delta_{bc}\delta_{ad}$ is a 4×4 symmetry generator in the contour space. This can be understood as the promotion of the permutation symmetry $(1 \leftrightarrow 3), (2 \leftrightarrow 4)$ for quadratic actions.

This symmetry is broken by the saddle-point solution. From Figure 8 (a), we see that contour (1,2) is disconnected from the contour (3,4). As a result, we expect G_s^{ab} to be block diagonal $G_s^{13} = G_s^{14} = G_s^{23} = G_s^{24} = 0$, and satisfying $G_s^{ab} = G_s^{a+2,b+2}$ for $a,b \in \{1,2\}$. Explicitly, on steady states, the solution of (103) and (104) reads

$$G_s^{11}(\omega) = \frac{i\omega}{\omega^2 + \Gamma^2/4}, \quad G_s^{12}(\omega) = -\frac{\Gamma/2}{\omega^2 + \Gamma^2/4}.$$
 (105)

Here we have defined $\Gamma = V + J = V_0 + V_1 + J_0 + J_1$. The solution (105) is only invariant under the transformation if we have $\gamma_{13} = \gamma_{24}$, and the residue symmetry group is O(2). Since the relative rotation symmetry is broken, there is an associated Goldstone mode living in the coset space $O(2) \otimes O(2)/O(2) = O(2)$. We will use a $\theta \in [0, 2\pi)$ field for this Goldstone mode. The effective theory for θ can be derived by considering fluctuations around the saddle point. The result reads [93]

$$\frac{S_{\theta}}{N} = \frac{1}{2} \int_{\omega k} \left(\frac{J_1 + V_1}{4} k^2 + \frac{1}{4V} \omega^2 \right) |\theta(\omega, k)|^2.$$
 (106)

This is the 2D XY model, with dynamical exponent z = 1. It is also a representative conformal field theory, which explains the conformal symmetry of steady states observed in numerical studies. Since this conformal symmetry only emerges after introducing replicas, the authors call it an emergent replica conformal symmetry [93].

The entanglement entropy can be computed using the effective action (106). The important observation is that the Goldstone mode θ is exactly the degree of freedom that labels the pairing between contours. For $\theta=0$, contours (1, 2) and (3, 4) are paired, as shown in (105). This is an analog of I in section IV E. After performing a relative rotation θ , the Green's functions become $G^{12}=\cos\theta\,G_s^{12}$ and $G^{14}=\sin\theta\,G_s^{12}$. Consequently, $\theta=\pi/2$ corresponds to pairing (1, 4) and (2, 3). This is an analog of σ in section IV E. Consequently, when computing the entanglement entropy, we should impose $\theta=0$ near the boundary of the contour for sites in subsystem B, and $\theta=\pi/2$ for sites in subsystem A. This is illustrated in Figure 8 (b). For the XY model, (106), this boundary condition excites a pair of half-vortexes, with a energy logarithmic in their distance:

$$S_A^{(2)} \propto \rho_s \log L_A = \sqrt{\frac{J_1 + V_1}{V}} N \log L_A.$$
 (107)

Here ρ_s is known as the superfluid density [130]. (107) has been verified by comparing to numerical results in [93].

We finally comment on interaction effects. The absence of interactions is essential for the presence of $O(2) \otimes O(2)$ symmetry, and thus the emergence of replica conformal symmetry. When interactions are introduced, for example an on-site Brownian SYK₄ term, the $O(2) \otimes O(2)$ symmetry is broken

explicitly back to $Z_4 \otimes Z_4$, where θ is fixed to be $n\pi/2$, as a combination of $\chi^a \to -\chi^a$ and permutations between contours. More explanation is given in the next subsection. The Goldstone modes gain mass and a smooth rotation of θ is no longer allowed. As a result, the boundary condition of $S_A^{(2)}$ will excites a domain wall instead of a half-vortex pair. This leads to a volume law phase similar to section IV E. This concludes that no entanglement phase transition shows up in this simple Hamiltonian (101).

B. Entanglement phase transitions

A non-Hermitian SYK chain with entanglement phase transition was proposed in [94]. Comparing to the model in the last subsection, it contains two copy of SYK chains, labeled by L and R. The Hamiltonian reads

$$H_{R} = \sum_{x,a} \left(\sum_{jk} i J_{jk}^{x,a}(t) \psi_{x,a,j} \psi_{x+1,a,k} + \sum_{j_{1} < \dots < j_{n}} i^{q/2} U_{j_{1}\dots j_{q}}^{x,a}(t) \psi_{x,a,j_{1}} \dots \psi_{x,a,j_{q}} \right),$$
(108)

with a = L, R, and

$$H_{\rm I} = -\frac{\mu}{2} \int dt \sum_{x,i} i \psi_{x,L,i} \psi_{x,R,i}. \tag{109}$$

The model can be understood as performing continuous forced measurements on the inter-chain fermion parity. We assume $q \ge 4$ and U = 0 corresponds to the non-interacting limit.

Following the discussions in the last subsection, we first consider the saddle-point solutions without twist operators. For U = 0, the system still exhibits the $O(2) \otimes O(2)$ symmetry. The solution of saddle-point equations reads [94]

$$G = \begin{cases} \frac{e^{-\frac{J|t|}{2}}}{2} (\operatorname{sgn}(t)\sigma^{z} - \sqrt{1 - \tilde{\mu}^{2}}i\sigma^{y} + \tilde{\mu}\tau^{y}), & \tilde{\mu} < 1\\ \frac{e^{-\frac{J|t|}{2}}}{2} (\operatorname{sgn}(t)\sigma^{z} + \tau^{y}), & \tilde{\mu} \ge 1 \end{cases}$$
(110)

Here $\tilde{\mu}=\mu/J$. σ_a (τ_a) are the Pauli matrices in the 1, 2 (L,R) space. We find $G^{12}\neq 0$ for $\mu < J$, where the forward evolution pairs with the backward evolution, while $G^{12}>0$ for $\mu \geq J$, where the pairing is between two copies of the SYK chain. When the pairing between forward and backward evolution presents, the $O(2)\otimes O(2)$ symmetry is broken to O(2), and the Goldstone mode exists. On the other hand, when $G^{12}=0$, the saddle-point solution preserves the full symmetry, and the system is gapped. Such a transition can be formulated as an effective action [94]:

$$\frac{S_{\phi}}{N} = \int dt dx \left(\frac{1}{2} (\partial \vec{\phi})^2 + r \vec{\phi}^2 + \lambda \vec{\phi}^4 \right), \tag{111}$$

Here $\lambda > 0$ and $r \propto \frac{\mu - J}{2}$ determines whether the pairing field $\vec{\phi} \sim (G^{12} + G^{34}, G^{14} + G^{23})$ condenses or not. This suggests the entanglement phase transition for non-interacting systems




FIG. 9. The phase diagram of the non-Hermitian SYK₂ chain with long-range couplings. Adapted from [95].

with large local Hilbert space dimension is a large-N O(2) symmetry breaking transition. For $\mu < J$, the Goldstone modes lead to logarithmic entanglement, as in the previous subsection. For $\mu > J$, the system is disordered with respect to boundary conditions. As a result, the insertion of twist operators will not excite any defect, and the entanglement entropy is area law.

The relation between G^{12} and the scaling of the entanglement entropy can also be understood from the perturbation theory in section IV B [92]. By identifying $G^{21} = iG^{>}$, the early-time contribution (79) always grows linearly. As explained in section IV E, this is the area law contribution. In the long-time limit, it becomes larger than the replica wormhole contribution, and the saddle point is dominated by the latter. On the contrary, $G^{12} = 0$, the perturbative calculation predicts a saturation value for the area law entanglement. Consequently, it will dominate the entanglement entropy in the long-time limit and lead to an area law entangled phase.

Now we consider small interaction $U/J \ll 1$. Similar to (7), this introduces an additional term G^4 into the action, which contains a contribution $(\phi_1^4 + \phi_2^4)$. Consequently, the revised effective action reads

$$\frac{S_{\phi}}{N} = \int dt dx \, \left(\frac{1}{2} (\partial \vec{\phi})^2 + r \vec{\phi}^2 + \lambda \vec{\phi}^4, + \lambda' (\phi_1^4 + \phi_2^4) \right), \quad (112)$$

with $\lambda' \propto U$. The additional term breaks the O(2) symmetry, and the residue group is Z_4 . The transition at r=0 becomes a large-N Z_4 symmetry breaking transition. The disordered phase is similar to the non-interacting case, while the ordered phase exhibits volume law entanglement entropy. In [97], authors also use effective actions to study the entanglement phase transition from the quantum error correction perspective. The extension to phase transitions of the von Neumann entropy is studied in [98].

C. Effects of long-range couplings

The long-range coupling presents in most of the state-ofthe-art experimental platforms for quantum dynamics, including NMR systems and cold atom systems. In this subsection, we review efforts to understand the non-unitary dynamics using SYK chains. We focus on the non-interacting case studied in [95]. The effect of long-range couplings in interacting SYK-like models has been studied in [96].

We consider adding long-range hopping terms to the

Hamiltonian (101) as

$$H_R = i \sum_{x} \left[\sum_{j,k,r} \frac{J_{jk}^{x,r}(t)}{r^{\alpha}} \chi_{x,j} \chi_{x+r,k} + \sum_{j < k} \tilde{J}_{jk}^{x}(t) \chi_{x,j} \chi_{x,k} \right]. \tag{113}$$

Here $r \in \{1,2,...\}$ labels the range of hopping. For $\alpha \leq 0.5$, the decay of random hopping is too slow, and the system is effectively all-to-all connected. We focus on the $\alpha > 0.5$, where the system has a well-defined thermodynamical limit. The solution of the saddle-point equation is still given by (105), with $\Gamma = V_0 + V_1 + J_0 + \sum_r J_1/r^{2\alpha}$.

Comparing to the short-range hopping case, there is no difference from the symmetry perspective. To determine the scaling of entanglement entropy, we should derive the effective action for Goldstone modes. The k^2 term in (106) can be understood as the kinetic energy of a Majorana fermion pair $(V_1 + J_1)k^2/2 = (V_1 + J_1)(1 - \cos k)$. After adding long-range coupling, we replace it with

$$\epsilon(k) = V_1(1 - \cos(k)) + \sum_{r=1}^{\infty} \frac{J_1}{r^{2\alpha}} (1 - \cos(rk)).$$
 (114)

The low-energy physics is determined by the small k expansion of $\epsilon(k)$, which depends on α as

$$\epsilon(k) \propto
\begin{cases}
k^2/2 & \text{for } \alpha > 1.5, \\
k^{2\alpha - 1} & \text{for } \alpha \le 1.5.
\end{cases}$$
(115)

As a result, for $\alpha \geq 1.5$, there is no difference compared to models with short-range hopping. We have dynamical exponent z=1 and logarithmic entanglement entropy. For $\alpha < 1.5$, the dynamical exponent depends on α as $z=\frac{2\alpha-1}{2}$. Moreover, the entanglement entropy is no longer equal to the energy of a half vortex pair, and a direct calculation using dual fields of θ shows $S_A^{(2)} \propto L_A^{1-z}$ [95]. This is known as a fractal phase [131]. Implications on the purification dynamics are also discussed in [95].

VI. OUTLOOK

This review summarizes recent studies of the quantum entanglement in the SYK model and its generalizations. The entanglement entropy of the single-site SYK model can be studied using the ED, the ETH, and the path-integral approach. Results from different methods match to good precision. For coupled SYK models, the Rényi entanglement entropy generally shows first-order transitions, where the long-time saddle is an analog of replica wormholes. Non-unitary SYK chains can also be used to study measurement induced entanglement phase transitions for the 2-nd Rényi entropy, which can be understood as conventional symmetry breaking transitions on replicated Hilbert space.

There are several interesting open questions:

1. Von Neumann Entropy. In gravity calculations, the von Neumann entropy is much easier to compute compared with Rényi entropies [132], which receive additional backreactions. However, in the SYK model and

its generalizations, it's hard to take the limit $n \to 1$ and derive results for the von Neumann entropy, except for the Brownian models [98], or perturbative calculations [63]. It is interesting to develop the methodology for the path-integral approach for the von Neumann entropy for interacting quantum systems.

- 2. General Measurements. In section V, we focus on forced measurements, under which the evolution is described by non-Hermitian Hamiltonian dynamics. It is interesting to consider more general measurements, in particular, to understand the consequence of random outcomes. It is also inspiring to reveal the gravity picture of the measurement induced phase transitions.
- 3. General Many-body Systems. The SYK model and

its generalizations are solvable in the large-N limit, and consequently, we can understand their entanglement dynamics using the path-integral representation. However, there is no simple saddle-point description in general quantum systems with strong interactions. It is of vital importance to develop approximation methods for entanglement dynamics using motivations from SYK-like models.

Acknowledgment. We acknowledge helpful discussions with Xiao Chen, Yiming Chen, Yingfei Gu, and Chunxiao Liu. We thank Yingfei Gu, Yiming Chen, Chunxiao Liu, and Ning Sun for carefully reading the manuscript and giving valuable suggestions. PZ acknowledges support from the Walter Burke Institute for Theoretical Physics at Caltech.

- [1] R. Jozsa, arXiv preprint quant-ph/9707034 (1997).
- [2] R. Jozsa and N. Linden, Proceedings of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences 459, 2011 (2003).
- [3] S. Ding and Z. Jin, Chinese Science Bulletin 52, 2161 (2007).
- [4] A. Pal and D. A. Huse, Physical review b 82, 174411 (2010).
- [5] R. Nandkishore and D. A. Huse, Annu. Rev. Condens. Matter Phys. 6, 15 (2015).
- [6] D. A. Abanin, E. Altman, I. Bloch, and M. Serbyn, Reviews of Modern Physics 91, 021001 (2019).
- [7] J. M. Deutsch, Physical review a 43, 2046 (1991).
- [8] M. Srednicki, Physical review e **50**, 888 (1994).
- [9] A. Kitaev and J. Preskill, Physical review letters 96, 110404 (2006).
- [10] M. Levin and X.-G. Wen, Physical review letters 96, 110405 (2006).
- [11] S. Ryu and T. Takayanagi, Physical review letters **96**, 181602 (2006).
- [12] S. Ryu and T. Takayanagi, JHEP 08, 045 (2006).
- [13] A. Lewkowycz and J. Maldacena, JHEP 08, 090 (2013).
- [14] V. E. Hubeny, M. Rangamani, and T. Takayanagi, JHEP 07, 062 (2007).
- [15] T. Faulkner, A. Lewkowycz, and J. Maldacena, JHEP 11, 074 (2013).
- [16] N. Engelhardt and A. C. Wall, JHEP 01, 073 (2015).
- [17] G. Penington, Journal of High Energy Physics 2020, 1 (2020).
- [18] A. Almheiri, N. Engelhardt, D. Marolf, and H. Maxfield, Journal of High Energy Physics **2019**, 1 (2019).
- [19] A. Almheiri, R. Mahajan, J. Maldacena, and Y. Zhao, arXiv preprint arXiv:1908.10996 (2019).
- [20] A. Almheiri, R. Mahajan, and J. Maldacena, arXiv preprint arXiv:1910.11077 (2019).
- [21] A. Almheiri, T. Hartman, J. Maldacena, E. Shaghoulian, and A. Tajdini, arXiv preprint arXiv:1911.12333 (2019).
- [22] G. Penington, S. H. Shenker, D. Stanford, and Z. Yang, arXiv preprint arXiv:1911.11977 (2019).
- [23] P. Calabrese and J. Cardy, Journal of physics a: mathematical and theoretical 42, 504005 (2009).
- [24] M. Rangamani and T. Takayanagi, in *Holographic Entangle-ment Entropy* (Springer, 2017) pp. 35–47.
- [25] M. A. Metlitski, C. A. Fuertes, and S. Sachdev, Physical Review B 80, 115122 (2009).
- [26] S. Whitsitt, W. Witczak-Krempa, and S. Sachdev, Physical

- Review B 95, 045148 (2017).
- [27] W. Donnelly, S. Timmerman, and N. Valdés-Meller, arXiv preprint arXiv:1911.09302 (2019).
- [28] A. Kitaev, in Talk given at the Fundamental Physics Prize Symposium, Vol. 10 (2014).
- [29] S. Sachdev and J. Ye, Physical review letters **70**, 3339 (1993).
- [30] J. Maldacena and D. Stanford, Physical Review D 94, 106002 (2016).
- [31] J. Maldacena, D. Stanford, and Z. Yang, Progress of Theoretical and Experimental Physics 2016 (2016).
- [32] A. Kitaev and S. J. Suh, Journal of High Energy Physics **2018**, 1 (2018).
- [33] J. Maldacena, S. H. Shenker, and D. Stanford, Journal of High Energy Physics 2016, 1 (2016).
- [34] A. Eberlein, V. Kasper, S. Sachdev, and J. Steinberg, Physical Review B **96**, 205123 (2017).
- [35] D. Chowdhury, A. Georges, O. Parcollet, and S. Sachdev, arXiv preprint arXiv:2109.05037 (2021).
- [36] R. A. Davison, W. Fu, A. Georges, Y. Gu, K. Jensen, and S. Sachdev, Physical Review B 95, 155131 (2017).
- [37] Y. Gu, A. Kitaev, S. Sachdev, and G. Tarnopolsky, arXiv preprint arXiv:1910.14099 (2019).
- [38] P. Chaturvedi, Y. Gu, W. Song, and B. Yu, Journal of High Energy Physics **2018**, 1 (2018).
- [39] K. Bulycheva, Journal of High Energy Physics 2017, 1 (2017).
- [40] P. Saad, S. H. Shenker, and D. Stanford, arXiv preprint arXiv:1806.06840 (2018).
- [41] C. Sünderhauf, L. Piroli, X.-L. Qi, N. Schuch, and J. I. Cirac, Journal of High Energy Physics 2019, 1 (2019).
- [42] Y. Gu, X.-L. Qi, and D. Stanford, Journal of High Energy Physics 2017, 1 (2017).
- [43] Y. Gu, A. Lucas, and X.-L. Qi, SciPost Phys 2, 018 (2017).
- [44] S. Banerjee and E. Altman, Physical Review B **95**, 134302 (2017)
- [45] X. Chen, R. Fan, Y. Chen, H. Zhai, and P. Zhang, Physical review letters **119**, 207603 (2017).
- [46] X.-Y. Song, C.-M. Jian, and L. Balents, Physical review letters 119, 216601 (2017).
- [47] S.-K. Jian and H. Yao, Physical review letters 119, 206602 (2017).
- [48] Y. Chen, H. Zhai, and P. Zhang, Journal of High Energy Physics **2017**, 1 (2017).
- [49] P. Zhang, Physical Review B 96, 205138 (2017).

- [50] Z. Bi, C.-M. Jian, Y.-Z. You, K. A. Pawlak, and C. Xu, Physical Review B 95, 205105 (2017).
- [51] P. Narayan and J. Yoon, Journal of High Energy Physics 2017, 1 (2017).
- [52] C. Liu, X. Chen, and L. Balents, Physical Review B 97, 245126 (2018).
- [53] W. Fu and S. Sachdev, Physical Review B 94, 035135 (2016).
- [54] Y. Huang, Y. Gu, et al., Physical Review D 100, 041901 (2019).
- [55] P. Zhang, C. Liu, and X. Chen, SciPost Physics 8, Art (2020).
- [56] P. Zhang, Journal of High Energy Physics 2020, 1 (2020).
- [57] A. Haldar, S. Bera, and S. Banerjee, Physical Review Research 2, 033505 (2020).
- [58] J. Kudler-Flam, R. Sohal, and L. Nie, arXiv preprint arXiv:2107.04043 (2021).
- [59] Y. Gu, A. Lucas, and X.-L. Qi, Journal of High Energy Physics 2017, 1 (2017).
- [60] R. Sohal, L. Nie, X.-Q. Sun, and E. Fradkin, Journal of Statistical Mechanics: Theory and Experiment 2022, 013103 (2022).
- [61] Y. Chen, X.-L. Qi, and P. Zhang, Journal of High Energy Physics 2020, 1 (2020).
- [62] Y. Chen, Journal of High Energy Physics 2021, 1 (2021).
- [63] P. Dadras and A. Kitaev, Journal of High Energy Physics 2021, 1 (2021).
- [64] S.-K. Jian and B. Swingle, Journal of High Energy Physics 2021, 1 (2021).
- [65] S.-K. Jian and B. Swingle, arXiv preprint arXiv:2109.03825 (2021).
- [66] S. Kaixiang, P. Zhang, and H. Zhai, Journal of High Energy Physics 2021 (2021).
- [67] D. L. Nedel, Physics Letters B 817, 136340 (2021).
- [68] Y. Chen and P. Zhang, Journal of High Energy Physics 2019, 1 (2019).
- [69] D. Pouria, Journal of High Energy Physics **2022** (2022).
- [70] S. W. Hawking, Physical Review D 14, 2460 (1976).
- [71] M. A. Nielsen and I. Chuang, "Quantum computation and quantum information," (2002).
- [72] Y. Li, X. Chen, and M. P. Fisher, Physical Review B 98, 205136 (2018).
- [73] X. Cao, A. Tilloy, and A. D. Luca, SciPost Phys. 7, 24 (2019).
- [74] Y. Li, X. Chen, and M. P. Fisher, Physical Review B 100, 134306 (2019).
- [75] B. Skinner, J. Ruhman, and A. Nahum, Physical Review X 9, 031009 (2019).
- [76] A. Chan, R. M. Nandkishore, M. Pretko, and G. Smith, Physical Review B 99, 224307 (2019).
- [77] Y. Bao, S. Choi, and E. Altman, Physical Review B 101, 104301 (2020).
- [78] S. Choi, Y. Bao, X.-L. Qi, and E. Altman, Physical Review Letters 125, 030505 (2020).
- [79] M. J. Gullans and D. A. Huse, Physical Review X 10, 041020 (2020).
- [80] M. J. Gullans and D. A. Huse, Physical review letters 125, 070606 (2020).
- [81] C.-M. Jian, Y.-Z. You, R. Vasseur, and A. W. Ludwig, Physical Review B 101, 104302 (2020).
- [82] M. Szyniszewski, A. Romito, and H. Schomerus, Physical Review B 100, 064204 (2019).
- [83] A. Zabalo, M. J. Gullans, J. H. Wilson, S. Gopalakrishnan, D. A. Huse, and J. Pixley, Physical Review B 101, 060301 (2020).
- [84] Q. Tang and W. Zhu, Physical Review Research 2, 013022 (2020).

- [85] L. Zhang, J. A. Reyes, S. Kourtis, C. Chamon, E. R. Mucciolo, and A. E. Ruckenstein, Physical Review B 101, 235104 (2020).
- [86] S. Goto and I. Danshita, Physical Review A 102, 033316 (2020).
- [87] C.-M. Jian, B. Bauer, A. Keselman, and A. W. Ludwig, arXiv preprint arXiv:2012.04666 (2020).
- [88] Y. Bao, S. Choi, and E. Altman, arXiv preprint arXiv:2102.09164 (2021).
- [89] O. Alberton, M. Buchhold, and S. Diehl, arXiv preprint arXiv:2005.09722 (2020).
- [90] X. Chen, Y. Li, M. P. A. Fisher, and A. Lucas, Physical Review Research 2 (2020).
- [91] A. Nahum and B. Skinner, Physical Review Research 2
- [92] C. Liu, P. Zhang, and X. Chen, SciPost Physics 10, Art (2021).
- [93] P. Zhang, S.-K. Jian, C. Liu, and X. Chen, Quantum 5, 579 (2021).
- [94] S.-K. Jian, C. Liu, X. Chen, B. Swingle, and P. Zhang, Physical Review Letters 127, 140601 (2021).
- [95] P. Zhang, C. Liu, S.-K. Jian, and X. Chen, arXiv preprint arXiv:2105.08895 (2021).
- [96] S. Sahu, S.-K. Jian, G. Bentsen, and B. Swingle, arXiv preprint arXiv:2109.00013 (2021).
- [97] S.-K. Jian, C. Liu, X. Chen, B. Swingle, and P. Zhang, arXiv preprint arXiv:2106.09635 (2021).
- [98] S.-K. Jian and B. Swingle, arXiv preprint arXiv:2108.11973 (2021).
- [99] S. Coleman, Aspects of symmetry: selected Erice lectures (Cambridge University Press, 1988).
- [100] A. Altland and B. D. Simons, *Condensed matter field theory* (Cambridge university press, 2010).
- [101] C. M. Grinstead and J. L. Snell, *Introduction to probability* (American Mathematical Soc., 1997).
- [102] A. Georges, O. Parcollet, and S. Sachdev, Physical Review B 63, 134406 (2001).
- [103] A. M. García-García and J. J. Verbaarschot, Physical Review D 96, 066012 (2017).
- [104] D. Bagrets, A. Altland, and A. Kamenev, Nuclear Physics B 911, 191 (2016).
- [105] D. Stanford and E. Witten, Journal of High Energy Physics 2017, 1 (2017).
- [106] T. G. Mertens, G. J. Turiaci, and H. L. Verlinde, Journal of High Energy Physics 2017, 1 (2017).
- [107] Z. Yang, Journal of High Energy Physics **2019**, 1 (2019).
- [108] A. Kamenev, Field theory of non-equilibrium systems (Cambridge University Press, 2011).
- [109] P. Zhang, Y. Gu, and A. Kitaev, Journal of High Energy Physics 2021, 1 (2021).
- [110] D. N. Page, Physical review letters **71**, 1291 (1993).
- [111] I. Kourkoulou and J. Maldacena, arXiv preprint arXiv:1707.02325 (2017).
- [112] W. Israel, Physics Letters A 57, 107 (1976).
- [113] P. Łydżba, M. Rigol, and L. Vidmar, Physical review letters 125, 180604 (2020).
- [114] H. Casini and M. Huerta, Journal of Physics A: Mathematical and Theoretical **42**, 504007 (2009).
- [115] P. J. Forrester, Journal of Physics A: Mathematical and General 39, 6861 (2006).
- [116] P. J. Forrester, *Log-gases and random matrices (LMS-34)* (Princeton University Press, 2010).
- [117] E. Bianchi, L. Hackl, and M. Kieburg, Physical Review B 103, L241118 (2021).
- [118] There are also studies on subsystem entropy of systems pre-

- pared in thermal ensembles and coupled to a bath [66].
- [119] P. Zhang, Physical Review B 100, 245104 (2019).
- [120] A. Almheiri, A. Milekhin, and B. Swingle, arXiv preprint arXiv:1912.04912 (2019).
- [121] Similar calculations has been carried out in [59] for SYK chains.
- [122] M. P. Do Carmo, Differential geometry of curves and surfaces: revised and updated second edition (Courier Dover Publications, 2016).
- [123] We thank Yingfei Gu for explaining this example.
- [124] X. Dong, D. Harlow, and A. C. Wall, Physical review letters 117, 021601 (2016).
- [125] Y. Chen, Journal of High Energy Physics **2020**, 1 (2020).

- [126] P. Hayden, S. Nezami, X.-L. Qi, N. Thomas, M. Walter, and Z. Yang, Journal of High Energy Physics 2016, 1 (2016).
- [127] A. Nahum, S. Vijay, and J. Haah, Physical Review X **8**, 021014 (2018).
- [128] C. Von Keyserlingk, T. Rakovszky, F. Pollmann, and S. L. Sondhi, Physical Review X 8, 021013 (2018).
- [129] Here H_I is not positive semidefinite. However, we can always make it positive semidefinite by shifting a large enough constant.
- [130] H. Zhai, *Ultracold Atomic Physics* (Cambridge University Press, 2021).
- [131] M. Ippoliti, T. Rakovszky, and V. Khemani, arXiv preprint arXiv:2103.06873 (2021).
- [132] X. Dong, Nature communications 7, 1 (2016).