On signed graphs whose spectral radius does not exceed $\sqrt{2+\sqrt{5}}$

Dijian Wang^a, Wenkuan Dong^a, Yaoping Hou^b, Deqiong Li^c

^aSchool of Science, Zhejiang University of Science and Technology,

Hangzhou, Zhejiang, 310023, P. R. China

^bCollege of Mathematics and Statistics, Hunan Normal University,

Changsha, Hunan, 410081, P. R. China

^cSchool of Mathematics and Statistics, Hunan University of Technology and Business,

Changsha, Hunan, 410205, P. R. China

Abstract

The Hoffman program with respect to any real or complex square matrix M associated to a graph G stems from Hoffman's pioneering work on the limit points for the spectral radius of adjacency matrices of graphs does not exceed $\sqrt{2+\sqrt{5}}$. A signed graph $\dot{G}=(G,\sigma)$ is a pair (G,σ) , where G=(V(G),E(G)) is a simple graph and $\sigma:E(G)\to\{+1,-1\}$ is the sign function. In this paper, we study the Hoffman program of signed graphs. Here, all signed graphs whose spectral radius does not exceed $\sqrt{2+\sqrt{5}}$ will be identified.

AMS classification: 05C50

Keywords: Signed graphs; Spectral radius; Hoffman program.

1 Introduction

All graphs considered here are simple, undirected and finite. For a graph G = (V(G), E(G)), let A(G) denote the adjacency matrix of G, the eigenvalues of A(G) are all real and the spectral radius of G is equal to the largest eigenvalue of A(G).

The Hoffman program is the identification of connected graphs whose spectral radius does not exceed some special limit points established by Hoffman [8]. The smallest limit point for the spectral radius of G is 2, that is, identifies all connected graphs whose spectral radius does not exceed 2. This problem has already been completely solved by Smith [13]. They are known as Smith graphs. After that, the problem jumps to the next significant limit point, which is $\lambda^* := \sqrt{2 + \sqrt{5}} = \tau^{\frac{1}{2}} + \tau^{-\frac{1}{2}}$ (≈ 2.05817), where τ

^{*}Corresponding author: yphou@hunnu.edu.cn





Figure 1: The signed graphs \dot{S}_{14} , \dot{S}_{16} and \dot{T}_{2k} .

is the golden mean. In [5], Cvetković, Doob and Gutman determined the structures of graphs with spectral radius between 2 and $\sqrt{2+\sqrt{5}}$. Their description was completed by Brouwer and Neumaier [2]. In 2020, Jiang and Polyanskii [9] studied the forbidden subgraphs characterization for \mathcal{G}^{λ} (where \mathcal{G}^{λ} denotes the family of connected graphs of spectral radius $\leq \lambda$) with applications to estimate the maximum cardinality of equiangular lines in the n-dimensional Euclidean space \mathbb{R}^n . For more on the results between Hoffman program and equiangular lines, see [9,11,13].

A signed graph $\dot{G} = (G, \sigma)$ is a pair (G, σ) , where G is a simple graph with n vertices and m edges, called the underlying graph, and $\sigma : E(G) \to \{+1, -1\}$ is the sign function. An edge $v_i v_j$ is called positive (negative) if $\sigma(v_i v_j) = +1$ (resp. $\sigma(v_i v_j) = -1$) and denoted by $v_i \stackrel{+}{\sim} v_j$ (resp. $v_i \stackrel{-}{\sim} v_j$). An unsigned graph (or a graph) is a signed graph without negative edges. Given a signed graph \dot{G} , its adjacency matrix is defined by $A(\dot{G}) = (\sigma_{ij})$, where $\sigma_{ij} = \sigma(v_i v_j)$ if $v_i \sim v_j$, and $\sigma_{ij} = 0$ otherwise. The eigenvalues, denoted by $\lambda_1(\dot{G}) \geq \cdots \geq \lambda_n(\dot{G})$, of $A(\dot{G})$ are defined as the eigenvalues of \dot{G} ; they are all real since $A(\dot{G})$ is a real symmetric matrix. The spectral radius of \dot{G} is defined by $\rho(\dot{G}) = \max\{|\lambda_i(\dot{G})| : 1 \leq i \leq n\} = \max\{\lambda_1(\dot{G}), -\lambda_n(\dot{G})\}$.

The theory of limit points for the spectral radius of graph sequences studied by Hoffman is still valid in the context of signed graphs. Some interesting results on the Hoffman program of signed graphs can be found in [1,7,10,14]. In [10], Jiang and Polyanskii studied the forbidden subgraphs characterization for families of signed graphs with eigenvalues bounded from below with applications to determine the maximum cardinality of a spherical two-distance set with two fixed angles in high dimensions. The smallest limit point for the spectral radius of \dot{G} is also 2, see [14, Propostion 6.1]. Those signed graphs have been identified by McKee and Smyth [12]. Therefore, the natural next step is the next limit point: $\lambda^* = \sqrt{2 + \sqrt{5}}$, that is, identifies all connected signed graphs whose spectral radius does not exceed λ^* . In this paper, we study the Hoffman program of signed graphs and all signed graphs whose spectral radius does not exceed λ^* will be identified.



In this paper, positive edges are depicted as bold lines and negative edges are depicted as dashed lines. The rest of the paper is organized as follows. Our contribution is reported in Section 2. In Section 3, we give some preliminaries lemmas and theorems. In Section 4, we will prove the Theorem 2.4.

2 Main results

Let $T_{a,b,c}$ be the graph with a + b + c + 1 vertices consisting of three paths with a, b, and c edges, respectively, where these paths have one end vertex in common, and let $Q_{a,b,c}$ be the graph with a + b + c + 3 vertices consisting of a path with a + b + c edges and two extra edges affixed at v_0 and w_0 . See Fig. 2.

For any fixed $\rho > 0$, the symbol \mathcal{G}^{ρ} (resp. \mathcal{G}^{ρ}_{S}) denotes the set of the connected (resp. signed) graphs whose spectral radius does not exceed ρ .

Theorem 2.1. [2, 5, 13] Let the graphs $T_{a,b,c}$ and $Q_{a,b,c}$ be depicted in Fig. 2. Then

(i)
$$\mathcal{G}^2 = \{C_n, K_{1,4}, T_{2,2,2}, T_{1,3,3}, P_n, T_{1,1,n-1}, T_{1,2,2}, T_{1,2,3}, T_{1,2,4}, Q_{1,n-5,1}\}.$$

(ii) $\mathcal{G}^{\lambda^*} = \mathcal{G}^2 \cup \{T_{a,b,c} | a = 1, b = 2, c > 5; or \ a = 1, b > 2, c > 3; or \ a = b = 2, c > 2; or \ a = 2, b = c = 3\} \cup \{Q_{a,b,c} | (a,b,c) \in \mathcal{S}; or \ c \geq a > 0, b \geq b^*(a,c), (a,c) \neq (1,1)\}$ where

$$\mathcal{S} = \{(1,1,2), (2,4,2), (2,5,3), (3,7,3), (3,8,4)\}$$

and

$$b^{*}(a,c) = \begin{cases} a+c+2 & \text{for } a > 2; \\ c+3 & \text{for } a = 2; \\ c & \text{for } a = 1. \end{cases}$$

Theorem 2.2. [12] Signed graphs in \mathcal{G}_S^2 are the induced subgraphs of \dot{S}_{14} , \dot{S}_{16} or \dot{T}_{2k} .

Definition 2.3. (1) Let \dot{G} be a connected signed graph, and let v be a vertex in \dot{G} .

- Denote (\dot{G}, v, k) the signed graph obtained from \dot{G} by appending a \dot{T}'''_{2k} at v.
- Denote $[\dot{G}, v, k]$ the signed graph obtained from \dot{G} by appending a \dot{T}_{2k}'' at v.




Figure 3: The signed graphs $\dot{T}'_{2k},\,\dot{T}''_{2k}$ and \dot{T}'''_{2k} .




Figure 4: The signed graphs in Definition 2.3.

- (2) Let \dot{G}_1 , \dot{G}_2 be two connected disjoint signed graphs, and let v, u be vertices in \dot{G}_1 , \dot{G}_2 respectively.
- Define $(\dot{G}_1, v, k, u, \dot{G}_2)$ to be the signed graph obtained from \dot{G}_1 and \dot{G}_2 by joining them with a path of k vertices connecting v and u.
- Define $[\dot{G}_1, v, k, u, \dot{G}_2]$ to be the signed graph obtained from \dot{G}_1 and \dot{G}_2 by joining them with a \dot{T}_{2k}''' connecting v and u.

The above mentioned signed graphs are depicted in Figs. 3 and 4.

The main results of this paper are presented as following.

Theorem 2.4. Signed graphs in $\mathcal{G}_S^{\lambda^*}$ are the induced subgraphs (up to switching isomorphic) of

- $(i) \dot{S}_{14}, \dot{S}_{16}, \dot{T}_{2k},$
- (ii) $\dot{C}_4[2,2,2,2], \dot{H}_1 \dot{H}_8, \dot{G}_6^1 \dot{G}_6^7, \dot{G}_8^0 \dot{G}_8^3, \dot{\Theta}_{8,2,0}, \dot{G}_{10},$
- (iii) $\dot{C}_k^{1,\frac{k}{2}+1}$ (k is even), \dot{G}_0^k (k ≥ 12 and k is even), $\dot{U}_6^{n_1,n_2}$ ($n_1 \geq 1$ and $n_2 \geq 1$), $\dot{C}_4[n_1,1,n_3,1]$ ($n_1 \geq 1$ and $n_3 \geq 1$), \dot{S}_1^n ($n \geq 8$), \dot{S}_2^n ($n \geq 10$),
- (iv) $[\dot{G}, v, s, u, \dot{H}]$ ($s \ge 3$), where (\dot{G}, v) , $(\dot{H}, u) \in \{(\dot{G}_4^{12}, v_{12}), (\dot{G}_2^9, v_9), (T_{a,1,a-1}, v_{a-1}), (\dot{G}_{11}^{\prime\prime}, v_{n_1}), (\dot{G}_5^{10}, v_{10}), (P_4, v_2)\}$, $a \ge 3$ and $n_1 \ge 1$,
- $(v) \ (\dot{Q}'_{1,0},v_1,2,v_2,P_4), \ (\dot{Q}'_{a_4,a_4},v_{a_4},2,v_2,P_4), \ (\dot{Q}'_{b,b},v_b,2,v_{a_3-1},T_{a_3,1,a_3-1}) \ (b\geq a_3-1), \\ (\dot{Q}'_{1,0},v_1,2,v_{a_3-1},T_{a_3,1,a_3-1}), \ (\dot{Q}'_{a_1,a_1},v_{a_1},2,v_1,\dot{Q}'_{1,0}), \ (\dot{Q}'_{a_3,a_3},v_{a_3},2,v_{a_2},\dot{Q}'_{a_2,a_2}), \ (\dot{G}^{10}_5,v_{10},2,v_1,\dot{Q}'_{1,0}), \ (\dot{G}^{10}_5,v_{10},2,v_{a_4},\dot{Q}'_{a_4,a_4}), \ (\dot{G}^{9}_2,v_9,2,v_2,P_4), \ (\dot{G}^{9}_2,v_9,2,v_2,T_{3,1,2}), \ (\dot{G}^{9}_2,v_9,2,v_3,T_{4,1,3}), \\ (\dot{G}^{9}_2,v_9,2,v_1,\dot{Q}'_{1,0}), \ (\dot{G}^{9}_2,v_9,2,v_{a_2},\dot{Q}'_{a_2,a_2}), \ (\dot{G}^{11}_3,v_{11},2,v_2,T_{3,1,2}), \ (\dot{G}^{11}_3,v_{11},2,v_3,T_{4,1,3}), \ (\dot{G}^{11}_3,v_{11},2,v_2,\dot{Q}'_{a_2,a_2}), \ (\dot{G}^{12}_4,v_{12},2,v_1,\dot{Q}'_{1,0}), \ (\dot{G}^{12}_4,v_{12},2,v_1,\dot{Q}'_{1,0}), \ (\dot{G}^{12}_4,v_{12},2,v_1,\dot{Q}'_{1,0}), \ (\dot{G}^{12}_4,v_{12},2,v_{12},2,v_{13},\dot{Q}'_{a_4,a_4}), \ where \ a_i\geq i \ for \ i=1,2,3,4. \end{cases}$

The above mentioned signed graphs are depicted in Figs. 2, 5, 6 and 7.




Figure 5: The signed graphs in Theorem 2.4 (ii). The number denotes the spectral radius of the corresponding signed graph.




Figure 6: The signed graphs in Theorem 2.4.

Remark 2.5. (1) In Appendix A, we will prove that each signed graph of Theorem 2.4 (iii) and (iv) has spectral radius $\rho(\dot{G}) < \lambda^*$. See Lemma 4.21.

(2) Each signed graph of Theorem 2.4 (v) belongs to Lemma 3.5.

3 Preliminaries

The sign of a cycle \dot{C} of \dot{G} is $\sigma(\dot{C}) = \prod_{e \in \dot{C}} \sigma(e)$, whose sign is +1 (resp. -1) is called positive (resp. negative) and denoted by \dot{C}_n^+ (resp. \dot{C}_n^-). A signed graph \dot{G} is called balanced if all its cycles are positive; otherwise it is called unbalanced. If there is a vertex subset $S \subseteq V(\dot{G})$, such that \dot{G}^S is obtained by reversing the sign of every edge with one end in S and the other in $V(\dot{G}) \setminus S$, then we say that \dot{G} and \dot{G}^S are switching equivalent and write $\dot{G} \sim \dot{G}^S$. If \dot{G} is isomorphic to a signed graph switching equivalent to \dot{G}_1 , we say \dot{G} is switching isomorphic to \dot{G}_1 , denoted by $\dot{G} \simeq \dot{G}_1$. Switching isomorphic signed




Figure 7: The graph P_n and the signed graphs \dot{G}_1^n , \dot{G}_2^n , \dot{G}_3^n , \dot{G}_4^n and \dot{G}_5^n .

graphs share the same spectrum. More basic results on the signed graphs, see [15].

As usual, d_u is the degree of a vertex u and $\Delta(\dot{G}) = \max_{u \in V(\dot{G})} d_u$. Let $N_X(u)$ be the set of neighbours of a vertex u in a set X and $d_X(u)$ be the cardinality of $N_X(u)$. For $U \subset V(\dot{G})$, $\dot{G}[U]$ denotes the induced subgraph of \dot{G} respect to the U. The signed graph \dot{C}_k^1 (resp. \dot{C}_k^1) is obtained by a balanced cycle \dot{C}_k^+ (resp. an unbalanced cycle \dot{C}_k^-) with one pendent edge. The signed graph $\dot{C}_k^{i,j}$ ($i \neq j$) is obtained by an unbalanced cycle \dot{C}_k^- with one pendent edge at vertex v_i (in \dot{C}_k^-) and one pendent edge at vertex v_j (in \dot{C}_k^-).

A signed graph is called *maximal* with respect to some property \mathcal{P} if it is not a proper induced subgraph of some other signed graph satisfying \mathcal{P} . In this context, we call a signed graph is *maximal* if it is not a proper induced subgraph of some other signed graphs whose spectral radius does not exceed λ^* .

Given a signed graph \dot{H} and $\rho(\dot{H}) \leq 2$, a subset U of $V(\dot{H})$, and a vertex w not in $V(\dot{H})$, the signed graph $\dot{H}_U(w)$ is obtained by inserting the edges (the sign of edge is any) between w and all vertices of U. We call that w is a good vertex if $2 < \rho(\dot{H}_U(w)) \leq \lambda^*$.

The first lemma is interlacing theorem.

Lemma 3.1. [3] Let A be a symmetric matrix of order n with eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_n$ and B a principal submatrix of A of order m with eigenvalues $\mu_1 \geq \mu_2 \geq \cdots \geq \mu_m$. Then the eigenvalues of B interlace the eigenvalues of A, that is, $\lambda_i \geq \mu_i \geq \lambda_{n-m+i}$ for $i = 1, \ldots, m$.

Define $\overline{\mathcal{G}}_S^{\lambda^*}$ to be the set of the connected unbalanced signed graphs with spectral radius $2 < \rho(\dot{G}) \le \lambda^*$. Note that the balanced signed graph and unsigned graph share the same spectrum, thus the main goal of this paper is to determine all signed graphs of the set $\overline{\mathcal{G}}_S^{\lambda^*}$. Some properties of the signed graph \dot{G} of the set $\overline{\mathcal{G}}_S^{\lambda^*}$ are listed as following.

Lemma 3.2. Let $\dot{G} \in \mathcal{G}_S^{\lambda^*}$. Then $\Delta(\dot{G}) \leq 4$.

Proof. Note that
$$\rho(\dot{G})^2 = \max\{\lambda_i(A(\dot{G})^2)|i=1,\ldots,n\}$$
, then $\rho(\dot{G})^2 \geq \max\{(A(\dot{G})^2)_{ii}|i=1,\ldots,n\}$ = $\Delta(\dot{G})$. Since $\Delta(\dot{G}) \leq \rho(\dot{G})^2 \leq (\lambda^*)^2 < 5$, then $\Delta(\dot{G}) \leq 4$.




Figure 8: Forbidden signed graphs $\dot{F}_1 - \dot{F}_{11}$ (up to switching equivalence). The number denotes the spectral radius of corresponding signed graph.

We designate that a signed graph \dot{H} is an induced subgraph of \dot{G} by writing $\dot{H} \subset \dot{G}$. If \dot{H} is not an induced subgraph of \dot{G} , then we say that \dot{G} is \dot{H} -free.

Lemma 3.3. Let $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$. Then \dot{G} is \dot{C}_3 -free.

Proof. Suppose on the contrary that $\dot{C}_3 \subset \dot{G}$. By the table of the spectra of signed graphs with at most six vertices [4], we find that there is no signed graph $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ of order $n \leq 6$. Next we consider $n \geq 7$. Then \dot{G} must contain an induced subgraph \dot{G}_1 with order $|V(\dot{G}_1)| = 6$ and $\rho(\dot{G}_1) \leq \lambda^*$. By [4, page 19–40], we have $\dot{G}_1 \sim \dot{T}_6$. Since \dot{T}_6 is 4-regular, then $\Delta(\dot{G}) \geq 5$, which contradicts to Lemma 3.2. Hence, \dot{G} is \dot{C}_3 -free.

We say that \dot{H} is forbidden if $\rho(\dot{H}) > \lambda^*$. Evidently, if \dot{H} is forbidden, then \dot{G} is \dot{H} -free.

Lemma 3.4. All (unsigned) graphs expect the graphs in the set \mathcal{G}^{λ^*} , the signed graph $\dot{\mathcal{C}}^1_{2\ell+1}$ and the signed graphs \dot{F}_i ($i=1,\ldots,11$) of Fig. 8 are forbidden.

Proof. It is easily to see that all (unsigned) graphs expect the graphs in the set \mathcal{G}^{λ^*} , the signed graph $\dot{\mathcal{C}}^1_{2\ell+1}$ (since $\rho(\dot{\mathcal{C}}^1_{2\ell+1}) = \rho(\dot{\mathcal{C}}^1_{2\ell+1})$) and the signed graphs $\dot{F}_1 - \dot{F}_{11}$ of Fig. 8 have spectral radius $\rho(\dot{G}) > \lambda^*$. So all of them are forbidden by interlacing.

The next lemma plays an important role in this paper. The proof is presented in Appendix A.

Lemma 3.5. Let $\dot{\mathcal{A}}_1^{n_1,n_2,n_3,n_4}$, $\dot{\mathcal{A}}_2^{n_1,n_2,n_3}$, $\dot{\mathcal{A}}_3^{n_1,n_2,n_3}$ $(n_1 \geq n_2)$, $\dot{\mathcal{A}}_i^{n_1,n_2}$ $(i = 4, \ldots, 14)$, $\dot{\mathcal{A}}_i^{n_1}$ $(i = 15, \ldots, 19)$ be the signed graphs depicted in Fig. 9. Then

- $(1) \ if \ \rho(\dot{\mathcal{A}}_{1}^{n_{1},n_{2},n_{3},n_{4}}) \leq \lambda^{*}, \ then \ \dot{\mathcal{A}}_{1}^{n_{1},n_{2},n_{3},n_{4}} \ is \ an \ induced \ subgraph \ of \ [P_{4},v_{2},s,v_{2},P_{4}], \\ [T_{a_{3},1,a_{3}-1},v_{a_{3}-1},s_{3},v_{2},P_{4}] \ or \ [T_{a_{3},1,a_{3}-1},v_{a_{3}-1},s,v_{b_{3}-1},T_{b_{3},1,b_{3}-1}],$
- $(2) \ if \ \rho(\dot{\mathcal{A}}_{2}^{n_{1},n_{2},n_{3}}) \leq \lambda^{*}, \ then \ \dot{\mathcal{A}}_{2}^{n_{1},n_{2},n_{3}} \ is \ an \ induced \ subgraph \ of \ (\dot{Q}_{1,0}',v_{1},2,v_{2},P_{4}), \\ (\dot{Q}_{a_{4},a_{4}}',v_{a_{4}},2,v_{2},P_{4}), \ (\dot{Q}_{1,0}',v_{1},2,v_{a_{3}-1},T_{a_{3},1,a_{3}-1}), \ (\dot{Q}_{a,a}',v_{a},2,v_{a_{3}-1},T_{a_{3},1,a_{3}-1}) \ (where \ a \geq a_{3}-1), \ [\dot{Q}_{a_{1},a_{1}}',v_{a_{1}},s,v_{2},P_{4}] \ or \ [\dot{Q}_{a_{1},a_{1}}',v_{a_{1}},s,v_{a_{3}-1},T_{a_{3},1,a_{3}-1}],$




Figure 9: The signed graphs in Lemma 3.5.

- (3) if $\rho(\dot{\mathcal{A}}_{3}^{n_{1},n_{2},n_{3}}) \leq \lambda^{*}$, then $\dot{\mathcal{A}}_{3}^{n_{1},n_{2},n_{3}}$ is an induced subgraph of $(\dot{Q}'_{a_{1},a_{1}},v_{a_{1}},2,v_{1},\dot{Q}'_{1,0})$, $(\dot{Q}'_{a_{3},a_{3}},v_{a_{3}},2,v_{a_{2}},\dot{Q}'_{a_{2},a_{2}})$ or $[\dot{Q}'_{a_{1},a_{1}},v_{a_{1}},s,v_{b_{1}},\dot{Q}'_{b_{1},b_{1}}]$,
- (4) if $\rho(\dot{\mathcal{A}}_{4}^{n_{1},n_{2}}) \leq \lambda^{*}$, then $\dot{\mathcal{A}}_{4}^{n_{1},n_{2}}$ is an induced subgraph of $[\dot{G}_{5}^{10},v_{10},s,v_{2},P_{4}]$ or $[\dot{G}_{5}^{10},v_{10},s,v_{a_{3}-1},T_{a_{3},1,a_{3}-1}]$,
- (5) if $\rho(\dot{\mathcal{A}}_{5}^{n_{1},n_{2}}) \leq \lambda^{*}$, then $\dot{\mathcal{A}}_{5}^{n_{1},n_{2}}$ is an induced subgraph of $(\dot{G}_{5}^{10},v_{10},2,v_{1},\dot{Q}'_{1,0})$, $(\dot{G}_{5}^{10},v_{10},2,v_{a_{4}},\dot{Q}'_{a_{4},a_{4}})$ or $[\dot{G}_{5}^{10},v_{10},s,v_{a_{1}},\dot{Q}'_{a_{1},a_{1}}]$,
- $(6) \ if \ \rho(\dot{\mathcal{A}}_{i}^{n_{1},n_{2}}) \leq \lambda^{*} \ (i=6,8), \ then \ \dot{\mathcal{A}}_{i}^{n_{1},n_{2}} \ is \ an \ induced \ subgraph \ of \ (\dot{G}_{2}^{9},v_{9},2,v_{2},P_{4}), \\ [\dot{G}_{2}^{9},v_{9},s,v_{2},P_{4}], \ (\dot{G}_{2}^{9},v_{9},2,v_{2},T_{3,1,2}), \ (\dot{G}_{2}^{9},v_{9},2,v_{3},T_{4,1,3}) \ \ or \ [\dot{G}_{2}^{9},v_{9},s,v_{a_{3}-1},T_{a_{3},1,a_{3}-1}], \\$
- (7) if $\rho(\dot{\mathcal{A}}_{i}^{n_{1},n_{2}}) \leq \lambda^{*}$ (i=7,9), then $\dot{\mathcal{A}}_{i}^{n_{1},n_{2}}$ is an induced subgraph of $(\dot{G}_{2}^{9},v_{9},2,v_{1},\dot{Q}'_{1,0})$, $(\dot{G}_{2}^{9},v_{9},2,v_{a_{2}},\dot{Q}'_{a_{2},a_{2}})$ or $[\dot{G}_{2}^{9},v_{9},s,v_{a_{1}},\dot{Q}'_{a_{1},a_{1}}]$,
- (8) if $\rho(\dot{\mathcal{A}}_{10}^{n_1,n_2}) \leq \lambda^*$, then $\dot{\mathcal{A}}_{10}^{n_1,n_2}$ is an induced subgraph of $[\dot{G}_3^{11}, v_{11}, s, v_2, P_4]$, $(\dot{G}_3^{11}, v_{11}, s, v_2, T_{4,1,3})$, $(\dot{G}_3^{11}, v_{11}, s, v_{4,1,3})$ or $[\dot{G}_3^{11}, v_{11}, s, v_{4,1,3}, T_{4,1,3}, v_{4,1,3}, v_{4,$
- (9) if $\rho(\dot{\mathcal{A}}_{11}^{n_1,n_2}) \leq \lambda^*$, then $\dot{\mathcal{A}}_{11}^{n_1,n_2}$ is an induced subgraph of $(\dot{G}_3^{11},v_{11},2,v_1,\dot{Q}'_{1,0})$, $(\dot{G}_3^{11},v_{11},2,v_{a_2},\dot{Q}'_{a_2,a_2})$ or $[\dot{G}_3^{11},v_{11},s,v_{a_1},\dot{Q}'_{a_1,a_1}]$,
- (10) if $\rho(\dot{\mathcal{A}}_{12}^{n_1,n_2}) \leq \lambda^*$, then $\dot{\mathcal{A}}_{12}^{n_1,n_2}$ is an induced subgraph of $[\dot{G}_4^{12},v_{12},s,v_2,P_4]$ or $[\dot{G}_4^{12},v_{12},s,v_{a_3-1},T_{a_3,1,a_3-1}]$,
- (11) if $\rho(\dot{A}_{13}^{n_1,n_2}) \leq \lambda^*$, then $\dot{A}_{13}^{n_1,n_2}$ is an induced subgraph of $(\dot{G}_4^{12}, v_{12}, 2, v_1, \dot{Q}'_{1,0})$, $(\dot{G}_4^{12}, v_{12}, 2, v_{a_4}, \dot{Q}'_{a_4,a_4})$ or $[\dot{G}_4^{12}, v_{12}, s, v_{a_1}, \dot{Q}'_{a_1,a_1}]$,
 - $(12) \ \ \textit{if} \ \rho(\dot{\mathcal{A}}_{14}^{n_1,n_2}) \leq \lambda^*, \ \ \textit{then} \ \ \dot{\mathcal{A}}_{14}^{n_1,n_2} \subset [P_4,v_2,s] \ \ \textit{or} \ \ \dot{\mathcal{A}}_{14}^{n_1,n_2} \subset [T_{a_3,1,a_3-1},v_{a_3-1},s].$
- (13) if $\rho(\dot{\mathcal{A}}_i^{n_1}) \leq \lambda^*$ (i = 15, 16, 17, 18, 19), then $\dot{\mathcal{A}}_i^{n_1}$ is an induced subgraph of $[\dot{G}, v, s, u, \dot{H}]$, where (\dot{G}, v) , $(\dot{H}, u) \in \{(\dot{G}_4^{12}, v_{12}), (\dot{G}_2^{9}, v_{9}), (\dot{G}_5^{10}, v_{10})\}$,

where $b_1 \ge 1$, $b_3 \ge 1$, $a_i \ge i$ for i = 1, 2, 3, 4 and $s \ge 3$.

Remark 3.6. Each signed graph in Lemma 3.5 is one of the signed graphs of Theorem 2.4 (v) or an induced subgraph of the signed graphs of Theorem 2.4 (iv).

Now we pay attention to the uncyclic and bicyclic signed graphs of the set $\mathcal{G}_S^{\lambda^*}$.

Lemma 3.7. Let $\dot{G} = (G, \sigma) \in \overline{\mathcal{G}}_{S}^{\lambda^{*}}$ $(G \neq C_{n})$ be a uncyclic signed graph with a k-cycle. Then $\dot{G} \subset \dot{\mathcal{C}}_{k}^{1,\frac{k}{2}+1}, \dot{\mathcal{C}}_{10}^{1,5}, \dot{\mathcal{C}}_{8}^{1,4}, \dot{\mathcal{G}}_{8}^{0}, \dot{U}_{6}, \dot{U}_{6}^{n_{1},n_{2}}, \dot{\mathcal{A}}_{1}^{n_{1},n_{2},n_{3},n_{4}}, \dot{\mathcal{A}}_{2}^{n_{1},n_{2},n_{3}}, \dot{\mathcal{A}}_{4}^{n_{1},n_{2}}, \dot{\mathcal{A}}_{14}^{n_{1},n_{2}}, \dot{\mathcal{C}}_{4}[2,2,2,2],$ $\dot{\mathcal{C}}_{4}[4,2,1,0], \, \dot{\mathcal{C}}_{4}[2,3,1,0], \, \dot{\mathcal{C}}_{4}[5,3,0,0], \, \dot{\mathcal{C}}_{4}[n_{1},1,n_{3},1].$ See Figs. 5, 6, 9 and 10.

Proof. Let $V(\dot{C}_k) = \{v_1, v_2, \dots, v_k\}$ and let $d_{v_1} \geq 3$ and u_1 be the new neighbor of v_1 . By forbidding \dot{C}_{ℓ}^1 and $\dot{C}_{2\ell+1}^1$, then the cycle \dot{C}_k is unbalanced and k is even. In addition, by forbidding \dot{F}_1 , if $k \geq 6$, then $d_{v_i} \leq 3$ for all $i = 1, \dots, k$.

Case 1. $k \geq 14$. Then $d_{v_i} = 2$ for $i = 2, 3, \ldots, \frac{k}{2}$, otherwise $d_{v_i} \geq 3$ and let $u_i \sim v_i$, then $Q_{2,i-1,\frac{k}{2}-3} \subset \dot{G} - v_{i+3}$ and $\rho(\dot{G}) \geq \rho(Q_{2,i-1,\frac{k}{2}-3}) > \lambda^*$, which is a contradiction. Because of symmetry, we have $d_{v_j} = 2$ for $j = \frac{k}{2} + 2, \ldots, k$. If $d_{u_1} \geq 2$, let $u_2 \stackrel{+}{\sim} u_1$, then $\{v_1, v_2, v_3, v_4, v_5, v_k, v_{k-1}, v_{k-2}, u_1, u_2\}$ induces the $T_{2,3,4}$ and $\rho(\dot{G}) \geq \rho(T_{2,3,4}) > \lambda^*$, which is a contradiction. So $d_{u_1} = 1$. Therefore, $\dot{G} \subset \dot{C}_k^{1,\frac{k}{2}+1}$.

Case 2. k=10 or 12. Then $d_{u_1}=1$, otherwise $d_{u_1} \geq 2$ and let $u_2 \stackrel{+}{\sim} u_1$, then $\{v_1, v_2, v_3, v_4, v_5, v_k, v_{k-1}, v_{k-2}, u_1, u_2\}$ induces the $T_{2,3,4}$ and $\rho(\dot{G}) \geq \rho(T_{2,3,4}) > \lambda^*$, which is a contradiction. So all vertices of $V(\dot{G}) \setminus V(\dot{C}_k)$ are within distance 1 of the cycle \dot{C}_k . By directed calculations, it is not hard to get that $\dot{G} \sim \dot{C}_{10}^{1,5}, \, \dot{C}_{10}^{1,6}, \, \dot{C}_{12}^{1,7}$.

Case 3. k=8 or 6. If all vertices of $V(\dot{G}) \setminus V(\dot{C}_k)$ are within distance 1 of the cycle \dot{C}_k , a directed calculation leads that $\dot{G} \subset \dot{C}_6^{1,4}, \dot{C}_8^{1,4}, \dot{C}_8^{1,5}$ or \dot{U}_6 . Otherwise, we assume that $d_{u_1} \geq 2$ and let u_2 be the new neighbor of u_1 . If k=8, by forbidding $T_{3,3,3}$ and \dot{F}_2 , then $d_{u_2}=1$ and $d_{u_1}=d_{v_i}=2$ for i=2,3,4,6,7,8, respectively. So $\dot{G} \subset \dot{G}_8^0$. If k=6, then $d_{v_2}=d_{v_6}=2$ (by forbidding \dot{F}_2). If $d_{v_3}=3$ or $d_{v_5}=3$, let u_3 (resp. u_5) be the new neighbor of v_3 (resp. v_5), by forbidding \dot{F}_2 , $Q_{1,2,3}$, $Q_{2,2,2}$, \dot{F}_1 and $Q_{1,1,4}$, we have $d_{u_1}=2$, $d_{u_2}=d_{u_3}=d_{u_5}=1$ and $d_{v_4}=2$. Then $\dot{G} \subset \dot{U}_6$. See Fig. 10. Next we consider that $d_{v_3}=d_{v_5}=2$, then $\dot{U}_6^{n_1,n_2}\subset \dot{G}$. See Fig. 6. Since \dot{G} is \dot{F}_2 -free, then each vertex of $\dot{U}_6^{n_1,n_2}$ expect the vertices of $V(\dot{C}_6)$ has degree at most 2. Hence, $\dot{G} \sim \dot{U}_6^{n_1,n_2}$.

Case 4. k = 4. Then $\dot{C}_4[n_1, n_2, n_3, n_4] \subset \dot{G}$. Let $n_4 = min\{n_1, n_2, n_3, n_4\}$. If $d_{v_1} = 4$, then $n_1 = 1$, $d_{v_2} = d_{v_4} = 2$ (by forbidding \dot{F}_1) and at most one vertex of $\{t_1, \ldots, t_{n_3}\}$ has degree 3 (by forbidding \dot{F}_3). So, $\dot{G} \sim \dot{C}_4'$ if $d_{v_3} = 4$ (where $\rho(\dot{C}_4') = 2$) or $\dot{G} \subset \dot{A}_{14}^{n_1, n_2}$ if




Figure 10: The signed graphs \dot{U}_6 , $\dot{C}_4[n_1, n_2, n_3, n_4]$ and \dot{C}'_4 .

 $d_{v_3} \leq 3$. See Figs. 9 and 10. Next suppose that $d_{v_i} \leq 3$ for i=1,2,3,4. If there is one vertex expect v_1, v_2, v_3, v_4 having degree greater than 2, without loss of generality, assume that $d_{w_i} \geq 3$ $(1 \leq i \leq n_1 - 1)$. By forbidding \dot{F}_1 , \dot{F}_2 and \dot{F}_3 , then $d_{w_i} = 3$ and at most one vertex of $V(\dot{G}) \setminus \{v_1, v_3, w_1, \ldots, w_{n_1}\}$ has degree 3. If $n_2 \geq 3$, then $\rho_1(\dot{G}) \geq \lambda_1(\dot{F}_8) > \lambda^*$ (if $n_1 \geq 6$), $\rho_1(\dot{G}) \geq \lambda_1(\dot{F}_5) > \lambda^*$ (if $n_1 = 5$) and $\rho_1(\dot{G}) \geq \lambda_1(Q_{1,3,4}) > \lambda^*$ (if $2 \leq n_1 \leq 4$), which is a contradiction. So $n_2 \leq 2$. Furthermore, if $n_2 = 2$, then $n_3 = 0$ (by forbidding \dot{F}_2). Therefore, $\dot{G} \subset \dot{\mathcal{A}}_1^{n_1,n_2,n_3,n_4}$ (if $n_2 = 0$), $\dot{G} \subset \dot{\mathcal{A}}_2^{n_1,n_2,n_3}$ (if $n_2 = 1$) or $\dot{G} \subset \dot{\mathcal{A}}_4^{n_1,n_2}$ (if $n_2 = 2$). If all vertices expect v_1, v_2, v_3, v_4 have degree at most 2, then \dot{G} is $\dot{\mathcal{C}}_4[n_1, n_2, n_3, n_4]$. If $n_i \geq 2$ for i = 1, 2, 3, then $n_1 = n_2 = n_3 = 2$ by forbidding $T_{3,3,3}$ and $T_{2,3,4}$. So $\dot{G} \subset \dot{\mathcal{C}}_4[2, 2, 2, 2]$. Otherwise, $n_2 \leq 1$ or $n_3 \leq 1$. If $n_2 \leq 1$, then $\dot{G} \subset \dot{\mathcal{C}}_4[n_1, 1, n_3, 1]$. If $n_3 \leq 1$ and $n_2 \geq 2$, by forbidding \dot{F}_4 , \dot{F}_6 , \dot{F}_7 , \dot{F}_8 and $T_{2,3,4}$, we have $\dot{G} \subset \dot{\mathcal{C}}_4[n_1, n_2, n_3, n_4]$, where $(n_1, n_2, n_3, n_4) \in \{(2, 2, 2, 2), (4, 2, 1, 0), (2, 3, 1, 0), (5, 3, 0, 0), (n_1, 2, 0, 0)\}$.

It is known that there are three types of bicyclic graphs in term of their base graph as described next. A bicyclic graph is said to be a *bicyclic base graph* if contains no pendent vertices.

The type $\theta_{p,q,r}$ is the union of three internally disjoint paths P_{p+2} , P_{q+2} , and P_{r+2} which have the same two distinct end vertices, where $p \ge q \ge r \ge 0$.

The type $B_r^{a,b}$ consists of two vertex disjoint cycles C_a and C_b joined by a path P_r having only its end vertices in common with the cycles, where $a \geq 3$, $b \geq 3$ and $r \geq 2$.

The type $B_0^{a,b}$ is the union of two cycles C_a and C_b with precisely one vertex in common, where $a \geq 3$ and $b \geq 3$.

Lemma 3.8. Let $\dot{G} \in \mathcal{G}_S^{\lambda^*}$ be a bicyclic signed base graph. Then \dot{G} is switching equivalent to one of the $\dot{\mathcal{B}}_0^{4,4}$, $\dot{\mathcal{B}}_r^{4,4}$, $\dot{\Theta}_{3,3,1}$, $\dot{\Theta}_{p,1,1}$, $\dot{\Theta}_{2,2,0}$, $\dot{\Theta}_{4,2,0}$, $\dot{\Theta}_{6,2,0}$ or $\dot{\Theta}_{8,2,0}$. See Fig. 11.

Proof. For types $B_0^{a,b}$ and $B_r^{a,b}$, by forbidding \dot{C}_ℓ^1 , $\dot{C}_{2\ell+1}^1$, \dot{F}_1 and \dot{F}_2 , then $\sigma(\dot{C}_a) = \sigma(\dot{C}_b) = -1$ and a = b = 4. So $\dot{G} \sim \dot{\mathcal{B}}_0^{4,4}$ or $\dot{G} \sim \dot{\mathcal{B}}_r^{4,4}$. For type $\theta_{p,q,r}$, by forbidding \dot{C}_ℓ^1 and $\dot{C}_{2\ell+1}^1$, then $\dot{G} \sim \dot{\Theta}_{p,q,1}$ (if q > 1, then p and q are odd) or $\dot{G} \sim \dot{\Theta}_{p,q,0}$ (p and q are even).




Figure 11: The signed graphs $\dot{\mathcal{B}}_0^{4,4}$, $\dot{\mathcal{B}}_r^{4,4}$, $\dot{\Theta}_{p,q,1}$ and $\dot{\Theta}_{p,q,0}$.




Figure 12: The graphs G_1, G_2 and the signed graphs $\dot{G}_1, \dot{G}_2, \dot{\mathcal{X}}_1, \dot{\mathcal{X}}_2$ and $\dot{\mathcal{X}}_3$.

Case 1. $\dot{G} \sim \dot{\Theta}_{p,q,1}$. If $q \geq 3$ and $p \geq 5$, then $Q_{2,2,2} \subset \dot{G}$, which contradicts to Lemma 3.4. Therefore, p = q = 3 or q = 1.

Case 2. $\dot{G} \sim \dot{\Theta}_{p,q,0}$. If $q \geq 4$ (resp. $p \geq 10$), then $Q_{2,1,2} \subset \dot{G}$ (resp. $\dot{F}_4 \subset \dot{G}$), which contradicts to Lemma 3.4. Therefore, q = 2 and $p \in \{2, 4, 6, 8\}$.

Hence,
$$\dot{G} \sim \dot{\Theta}_{3,3,1}, \, \dot{\Theta}_{\nu,1,1}, \, \dot{\Theta}_{2,2,0}, \, \dot{\Theta}_{4,2,0}, \, \dot{\Theta}_{6,2,0} \text{ or } \dot{\Theta}_{8,2,0}.$$

Let $\dot{G} \in \mathcal{G}_S^{\lambda^*}$ be a signed graph with $m \geq n+1$ and \dot{C}_k be a cycle in \dot{G} . If each vertex outside of \dot{C}_k is adjacent to at most one vertex of \dot{C}_k , then $\dot{C}_k \subset \dot{\mathcal{B}}_r^{4,4}$ or $\dot{C}_k \subset \dot{\mathcal{B}}_0^{4,4}$ (by Lemma 3.8). So k=4. Therefore, if $k \geq 5$, then there is a vertex outside of \dot{C}_k adjacent to at least two vertices of \dot{C}_k . Then we have

Lemma 3.9. Let $\dot{G} = (G, \sigma)$ be a signed graph obtained by a cycle \dot{C}_k and a vertex u not in \dot{C}_k such that u is adjacent to at least two vertices of \dot{C}_k . If $\rho(\dot{G}) \leq \lambda^*$, then \dot{G} is switching equivalent to one of the $\dot{\Theta}_{3,3,1}, \dot{\Theta}_{k-3,1,1}, \dot{\mathcal{X}}_1, \dot{\mathcal{X}}_2$ or $\dot{\mathcal{X}}_3$. See Figs. 11 and 12.

Proof. If $d_{\dot{C}_k}(u)=2$, then \dot{G} is bicyclic. By Lemma 3.8, then $\dot{G}\sim\dot{\Theta}_{3,3,1}$ or $\dot{G}\sim\dot{\Theta}_{k-3,1,1}$. Next assume that $3\leq d_{\dot{C}_k}(u)\leq 4$, then the underlying graph G of \dot{G} is G_1 or G_2 . See Fig. 12. By forbidding \dot{C}_{ℓ}^1 and $\dot{C}_{2\ell+1}^1$, then the cycles C_1,C_2,C_3,C_4 in G_1 and G_2 are even and unbalanced. If $\sigma(\dot{C}_k)=+1$ and $d_{\dot{C}_k}(u)=3$, or $\sigma(\dot{C}_k)=-1$ and $d_{\dot{C}_k}(u)=4$, then at least one of the cycles C_1,C_2,C_3 and C_4 is balanced, which is a contradiction. So, either $\sigma(\dot{C}_k)=+1$ and $d_{\dot{C}_k}(u)=4$, or $\sigma(\dot{C}_k)=-1$ and $d_{\dot{C}_k}(u)=3$. Then $\dot{G}\sim\dot{G}_1$ or $\dot{G}\sim\dot{G}_2$. If $\dot{G}\sim\dot{G}_1$, then $n_1=n_2=n_3=n_4=1$ (by forbidding $\dot{C}_{2\ell+1}^1$ and \dot{F}_1). If $\dot{G}\sim\dot{G}_2$, then $n_i\in\{1,3\}$ for i=1,2,3 and at most one of n_1,n_2 and n_3 is equal to 3 (by forbidding $\dot{C}_{2\ell+1}^1,C_{2,2,2}$ and \dot{F}_2). Hence, $\dot{G}\sim\dot{X}_i$ for i=1,2 or 3. See Fig. 12.

In the final of this section, we give an algorithm for searching the signed graph \dot{G} with spectral radius $2 < \rho(\dot{G}) \le \lambda^*$.

Algorithm 1:

Input: The adjacency matrix $A_0 = A(\dot{G}) = (a_{ij})_{n \times n}$ of a signed graph with order n, and a positive integer number k.

Output: The set of the matrices $S_k = \{A_k = A(\dot{G}_k) = (a_{ij})_{(n+k)\times(n+k)} \mid 2 < \rho(\dot{G}_k) \leq \lambda^* \}$. Step 1. For A_0 , constructing a vector set $R_1 = \{\mathbf{r}_1 = (r_1, r_2, \dots, r_n) \mid r_j \in \{-1, 0, 1\}, j = \{-1, 0, 1$

 $1, \ldots, n$. (Restrict that $\sum_{i=1}^{n} |r_i| \le 4$ by Lemma 3.2).

- (1.1) Traverse each vector \mathbf{r}_1 of R_1 and construct a matrix $A_1 = \begin{pmatrix} A_0 & \mathbf{r}_1^T \\ \mathbf{r}_1 & 0 \end{pmatrix}$,
- $(1.2) \ a_1 \leftarrow \max\{|\lambda_i(A_1)| \mid i = 1, n\},\$
- (1.3) Traverse each matrix A_1 , and add A_1 to the set S_1 if the sum of each row of the matrix $|A_1|$ is less than or equal to 4 (by Lemma 3.2) and $a_1 \leq \lambda^*$,

Step 2. For each matrix A_1 of S_1 , constructing a vector set $R_2 = \{\mathbf{r}_2 = (r_1, r_2, \dots, r_{n+1}) \mid r_j \in \{-1, 0, 1\}, j = 1, \dots, n+1\}$. (Restrict that $\sum_{i=1}^{n+1} |r_i| \le 4$ by Lemma 3.2).

- (2.1) Traverse each vector \mathbf{r}_2 of R_2 and construct a matrix $A_2 = \begin{pmatrix} A_1 & \mathbf{r}_2^T \\ \mathbf{r}_2 & 0 \end{pmatrix}$,
- $(2.2) \ a_2 \leftarrow \max\{|\lambda_i(A_2)| \ | \ i = 1, n\},\$
- (2.3) Traverse each matrix A_2 , and add A_2 to the set S_2 if the sum of each row of the matrix $|A_2|$ is less than or equal to 4 (by Lemma 3.2) and $a_2 \leq \lambda^*$,

.

Step k. For each matrix A_{k-1} of S_{k-1} , constructing a vector set $R_k = \{\mathbf{r}_k = (r_1, r_2, ..., r_{n+k-1}) \mid r_j \in \{-1, 0, 1\}, j = 1, ..., n+k-1\}$. (Restrict that $\sum_{i=1}^{n+k-1} |r_i| \le 4$ by Lemma 3.2).

- (k.1) Traverse each vector \mathbf{r}_k of R_k and construct a matrix $A_k = \begin{pmatrix} A_{k-1} & \mathbf{r}_k^T \\ \mathbf{r}_k & 0 \end{pmatrix}$,
- $(k.2) \ a_k \leftarrow \max\{|\lambda_i(A_k)| \mid i = 1, n\},\$
- (k.3) Traverse each matrix A_k , and add A_k to the set S_k if the sum of each row of the matrix $|A_k|$ is less than or equal to 4 (by Lemma 3.2) and $2 < a_k \le \lambda^*$,
 - (k.4) Output the S_k .

The whole algorithm is over.

Corollary 3.10. Applying algorithm 1 to each signed graph of Theorem 2.4 (ii), we can check that all of them are maximal.




Figure 13: The signed graphs $\dot{X}_1 - \dot{X}_8$ where k is odd or $k \ge 10$.

4 Signed graph whose spectral radius does not exceed $\sqrt{2+\sqrt{5}}$

In this section, we identify all signed graphs whose spectral radius does not exceed $\sqrt{2+\sqrt{5}}$. By Lemma 3.7, we assume that $m \ge n+1$. We break into four subsections.

4.1 $\dot{C}_k \subset \dot{G}$ where k is odd or $k \geq 10$

Firstly we consider that $\dot{C}_k \subset \dot{G}$ where k is odd or $k \geq 10$.

Lemma 4.1. Let \dot{G} be one of the signed graphs $\dot{X}_1, \ldots, \dot{X}_8$ where k is odd or $k \geq 10$, see Fig. 13. Then $\rho(\dot{G}) \leq \lambda^*$ if and only if $\dot{G} \sim \dot{X}_3$ and k = 10.

Proof. If $\dot{G} \sim \dot{X}_3$ and k = 10, then $\dot{G} \subset \dot{G}_{10}$. Note that $\dot{C}^1_{\ell} \subset \dot{X}_1$ $(4 \leq \ell < k)$, $Q_{2,2,2} \subset \dot{X}_2$ or $\dot{C}^1_{2\ell+1} \subset \dot{X}_2$, $\dot{F}_9 \subset \dot{X}_3$ $(k \geq 12 \text{ and } k \text{ is even})$, $\dot{C}^1_k \subset \dot{X}_3$ (k is odd), $\dot{C}^1_4 \subset \dot{X}_4$, $T_{2,3,4} \subset \dot{X}_5$ and $\dot{C}^1_k \subset \dot{X}_j$ for j = 6, 7, 8. By Lemma 3.4, then $\rho(\dot{X}_i) > \lambda^*$ for $i = 1, \ldots, 8$ expect the signed graph \dot{X}_3 where k = 10.

Lemma 4.2. Let $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ be a signed graph with $m \geq n+1$. If $\dot{C}_k \subset \dot{G}$ where k is odd or $k \geq 10$, then $\dot{G} \subset \dot{\Theta}_{8,2,0}, \dot{G}_{10}$ or \dot{G}_0^k $(k \geq 12 \text{ and } k \text{ is even})$. See Figs. 5 and 6.

Proof. By Lemma 3.9, then $\dot{\Theta}_{8,2,0} \subset \dot{G}$ or $\dot{\Theta}_{k-3,1,1} \subset \dot{G}$. By Corollary 3.10, we know that $\dot{\Theta}_{8,2,0}$ is maximal. Then we consider that $\dot{\Theta}_{k-3,1,1} \subset \dot{G}$. Let \dot{H} be the induced subgraph of \dot{G} such that $\dot{\Theta}_{k-3,1,1} \subset \dot{H}$ and $\rho(\dot{H}) \leq 2$. Clearly, $\dot{H} \subset \dot{T}_{2k}$. We use the vertex label of Fig. 1 and let $V(\dot{\Theta}_{k-3,1,1}) = \{v_1, \ldots, v_k, u_1\}$. Let w be a good vertex in $V(\dot{G}) \setminus V(\dot{H})$.

Claim 1. If $w \sim v_i$ for one i, then k = 10 and $\dot{H}_U(w) \sim \dot{X}_3$.

Proof. By Lemmas 3.4 and 3.9, then $w \stackrel{+}{\sim} v_{i-1}$ and $w \stackrel{-}{\sim} v_{i+1}$ for one $i \pmod k$. It is not hard to get that $i \neq 1$ and $u_i \notin V(\dot{H})$ (otherwise $\dot{C}_4^1 \subset \dot{G}$ and $\rho(\dot{G}) \geq \rho(\dot{C}_4^1) > \lambda^*$). Since $\dot{H}_U(w)$ is not an induced subgraph of \dot{T}_{2k} , then one of the followings happens:

- $w \sim u_j$ for one $j \neq i \pm 1$, then $\dot{X}_1 \subset \dot{G}$ or $\dot{X}_2 \subset \dot{G}$;
- $w \nsim u_j$ for one $j \in \{i \pm 1\}$, then $\dot{X}_3 \subset \dot{G}$;
- $w \stackrel{+}{\sim} u_j$ for one $j \in \{i \pm 1\}$, then $\dot{X}_4 \subset \dot{G}$.

By Lemma 4.1, then $\rho(\dot{H}_U(w)) \leq \lambda^*$ if and only if k = 10 and $\dot{H}_U(w) \sim \dot{X}_3$.

Claim 2. If $w \not\sim v_i$ for all i = 1, ..., k, then $w \sim u_1$. Moreover, $\dot{H}_U(w) \sim \dot{G}_0^k$.

Proof. If $w \not\sim u_1$, then there is a path from the vertex w to the vertex u_i of \dot{H} . And now $\dot{X}_i \subset \dot{G}$ for one $i \in \{5,6,7,8\}$ and $\rho(\dot{G}) > \lambda^*$ (by Lemma 4.1), which is a contradiction. So $w \sim u_1$. If $u_i \in V(\dot{H})$ for $i \neq 1$, we also get that $\dot{X}_i \subset \dot{G}$ for one $i \in \{5,6,7,8\}$, which is a contradiction. Hence, $\dot{H}_U(w) \sim \dot{G}_0^k$. See Fig. 6.

If k = 10, then $\dot{H}_U(w) \sim \dot{X}_3$ or $\dot{H}_U(w) \sim \dot{G}_0^{10}$. Applying algorithm 1 to $A(\dot{H}_U(w))$, we obtain that $\dot{G} \subset \dot{G}_{10}$.

If k is odd, or k is even and $k \geq 12$, then $\dot{H}_U(w) \sim \dot{G}_0^k$. If k is odd, then $\dot{C}_k^1 \subset \dot{G}$, which contradicts to Lemma 3.4. If k is even and $k \geq 12$, then \dot{G}_0^k is maximal, otherwise there is another vertex w' in $V(\dot{G}) \setminus V(\dot{H})$, then $w' \sim u_1$ (by Claims 1 and 2) and $\dot{F}_1 \subset \dot{G}$, which contradicts to Lemma 3.4. Hence, $\dot{G} \sim \dot{G}_0^k$.

Remark 4.3. By Lemma 4.2, we next state that the signed graph \dot{G} is bipartite.

4.2 $\dot{C}_8 \subset \dot{G}$

Secondly we consider that \dot{G} is \dot{C}_k -free where $k \geq 10$ and $\dot{C}_8 \subset \dot{G}$. By Lemma 3.9, then $\dot{\Theta}_{5,1,1}, \dot{\Theta}_{6,2,0}, \dot{\Theta}_{3,3,1}, \dot{\mathcal{X}}_2$ or $\dot{\mathcal{X}}_3$ is an induced subgraph of \dot{G} . Let \dot{H} be the induced subgraph of \dot{G} such that $\dot{\Theta}_{5,1,1} \subset \dot{H}, \, \dot{\Theta}_{3,3,1} \subset \dot{H}, \, \dot{\mathcal{X}}_2 \subset \dot{H}$ or $\dot{\mathcal{X}}_3 \subset \dot{H}$ and $\rho(\dot{H}) = 2$.

Lemma 4.4. Let $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ be a \dot{C}_k -free $(k \geq 10)$ bipartite signed graph with $m \geq n + 1$. If $\dot{C}_8 \subset \dot{G}$, then $\dot{G} \subset \dot{G}_8^i$ for i = 1, 2, 3. See Fig. 5.

Proof. Applying algorithm 1 to $A(\dot{\Theta}_{6,2,0})$, we obtain that $\dot{G} \subset \dot{G}_8^1$ or $\dot{G} \subset \dot{G}_8^2$. We next suppose that $\dot{\Theta}_{5,1,1} \subset \dot{G}, \dot{\Theta}_{3,3,1} \subset \dot{G}, \dot{\mathcal{X}}_2 \subset \dot{G}$ or $\dot{\mathcal{X}}_3 \subset \dot{G}$. If $\dot{\Theta}_{5,1,1} \subset \dot{G}$, then $\dot{\Theta}_{5,1,1} \subset \dot{H}$. So, $\dot{H} \subset \dot{S}_{16}$ or $\dot{H} \subset \dot{T}_{16}$. Then $9 \leq |V(\dot{H})| \leq 15$. By Fig. 1, if $|V(\dot{H})| = 9$, then \dot{H} is $\dot{\Theta}_{5,1,1}$; if $|V(\dot{H})| = 10$, then $\dot{H} \sim \dot{\Theta}_{5,1,1}^i$ for $i = 1, \ldots, 10$ (see Fig. 14); if $|V(\dot{H})| = 14$ or 15, then \dot{H} is switching isomorphic to one of the signed graphs of Fig. 15. Moreover, if




Figure 14: The signed graphs $\dot{\Theta}_{5,1,1}^1 - \dot{\Theta}_{5,1,1}^{10}$.




Figure 15: The signed graphs in the proof of Lemma 4.4.

 $11 \leq |V(\dot{H})| \leq 13$, then \dot{H} contains one $\dot{\Theta}^i_{5,1,1}$ $(i=1,\ldots,10)$ as an induced subgraph. Now applying algorithm 1 to $A(\dot{H})$ where \dot{H} is $\dot{\Theta}_{5,1,1}$ or one of the signed graphs of Fig. 15, we get that there is no signed graph \dot{G} with order $|V(\dot{H})| + 1$ and $2 < \rho(\dot{G}) \leq \lambda^*$, and applying algorithm 1 to $A(\dot{\Theta}^i_{5,1,1})$ for $i=1,\ldots,10$, we get that $\dot{G} \subset \dot{G}^3_8$. If $\dot{\Theta}_{3,3,1} \subset \dot{G}$, $\dot{\mathcal{X}}_2 \subset \dot{G}$ or $\dot{\mathcal{X}}_3 \subset \dot{G}$, we can similarly get that $\dot{G} \subset \dot{G}^3_8$.

4.3 $\dot{C}_6 \subset \dot{G}$

Thirdly we consider that \dot{G} is \dot{C}_k -free where $k \geq 8$ and $\dot{C}_6 \subset \dot{G}$. By Lemma 3.9, then $\dot{\Theta}_{4,2,0} \subset \dot{G}$, $\dot{\Theta}_{3,1,1} \subset \dot{G}$ or $\dot{\mathcal{X}}_1 \subset \dot{G}$. Let \dot{H} be the induced subgraph of \dot{G} such that $\dot{\Theta}_{4,2,0} \subset \dot{H}$, $\dot{\Theta}_{3,1,1} \subset \dot{H}$ or $\dot{\mathcal{X}}_1 \subset \dot{H}$ and $\rho(\dot{H}) \leq 2$.

Lemma 4.5. Let $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ be a \dot{C}_k -free $(k \geq 8)$ bipartite signed graph with $m \geq n+1$. If $\dot{C}_6 \subset \dot{G}$, then $\dot{G} \subset \dot{G}_6^1, \dot{G}_6^2, \dot{G}_6^3, \dot{G}_6^4, \dot{G}_6^5, \dot{G}_6^6, \dot{G}_6^7, \dot{G}_8^3, \dot{S}_1^n$ or \dot{S}_2^n . See Figs. 5 and 6.

Proof. If $\dot{\Theta}_{4,2,0} \subset \dot{H}$, then $\dot{H} \subset \dot{S}_{14}$ or $\dot{H} \subset \dot{S}_{16}$. Since \dot{G} is \dot{C}_k -free $(k \geq 8)$, by Fig. 1, then $\dot{H} \subset \dot{\Theta}^i_{4,2,0}$ for i = 1, 2, 3, 4. See Fig. 16. Applying algorithm 1 to $A(\dot{H})$ where $\dot{H} \subset \dot{\Theta}^i_{4,2,0}$ for i = 1, 2, 3, 4, we get that $\dot{G} \subset \dot{G}^3_8$ or $\dot{G} \subset \dot{G}^i_6$ for i = 1, 2, 3, 4, 5, 6.

If $\dot{\Theta}_{3,1,1} \subset \dot{H}$, then $\dot{H} \subset \dot{S}_{14}$ or $\dot{H} \subset \dot{T}_{12}$. So, $7 \leq |V(\dot{H})| \leq 13$. By Fig. 1, if n = 7, then \dot{H} is $\dot{\Theta}_{3,1,1}$; if $|V(\dot{H})| = 8$, then $\dot{H} \sim \dot{\Theta}^i_{3,1,1}$ for $i = 1, \ldots, 6$; if $|V(\dot{H})| = 12$ or 13, then $\dot{H} \sim \dot{S}^i_{14}$ for i = 1, 2, 3, 4. See Fig. 16. Moreover, if $9 \leq |V(\dot{H})| \leq 11$, then \dot{H} contains one $\dot{\Theta}^i_{3,1,1}$ ($i = 1, \ldots, 6$) as an induced subgraph. Now applying algorithm 1 to $A(\dot{\Theta}_{3,1,1})$ and $A(\dot{S}^i_{14})$ for i = 1, 2, 3, 4, then there is no signed graph \dot{G} with order




Figure 16: The signed graphs $\dot{\Theta}^1_{4,2,0} - \dot{\Theta}^4_{4,2,0}$, \dot{G}'_6 , \dot{G}''_6 , $\dot{\Theta}^1_{3,1,1} - \dot{\Theta}^6_{3,1,1}$, $\dot{S}^1_{14} - \dot{S}^4_{14}$.




Figure 17: The signed graphs $H_1^n - H_3^n$.

 $|V(\dot{H})|+1$ and $2<\rho(\dot{G})\leq\lambda^*$, and applying algorithm 1 to $A(\dot{\Theta}^i_{3,1,1})$ for $i=1,\ldots,6$, we can get all signed graphs $\dot{G}\in\overline{\mathcal{G}}^{\lambda^*}_S$ (where $\dot{\Theta}^i_{3,1,1}\subset\dot{G}$) of order $n\leq12$, which are the induced subgraphs of \dot{G}^7_6 or \dot{S}^{12}_1 . Notice that $|V(\dot{G}^7_6)|=11$. See Figs. 5 and 6. If $n\geq13$, then $\dot{S}^{12}_1\subset\dot{G}$. Set $V(\dot{G})\setminus V(\dot{S}^{12}_1)=\{v_{13},\ldots,v_n\}$. Since \dot{S}^{12}_1 is the unique signed graph of order 12 such that $\dot{\Theta}_{3,1,1}\subset\dot{G}$ and $2<\rho(\dot{G})\leq\lambda^*$, then $v_j\not\sim v_i$ for each $j\geq13$ and all $i\leq11$. By forbidding \dot{F}_2 , then $d_{v_i}\leq2$ for all $i\geq12$. Hence, $\dot{G}\sim\dot{S}^n_1$ and \dot{S}^n_1 is maximal.

If $\dot{\mathcal{X}}_1 \subset \dot{H}$, then $\dot{H} \subset \dot{S}_{14}$ or $\dot{H} \subset \dot{S}_{16}$. From above discussions, we can suppose that \dot{H} is $\{\dot{\Theta}_{4,2,0}, \dot{\Theta}_{3,1,1}\}$ -free. By Fig. 1, then $\dot{H} \subset \dot{G}'_6$, $\dot{H} \subset \dot{G}''_6$ or $\dot{H} \subset \dot{S}^{11}_2$. See Figs. 16 and 6. Similarly, applying algorithm 1 to $A(\dot{H})$, then all signed graphs $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ of order $n \leq 14$ can be found, which are the induced subgraphs of \dot{H}_i^{14} (i = 1, 2, 3) or \dot{S}_2^{14} . See Figs. 17 and 6. If $n \geq 15$, then $\dot{S}_2^{14} \subset \dot{G}$ or $\dot{H}_i^{14} \subset \dot{G}$ (i = 1, 2, 3).

Case 1. $\dot{S}_{2}^{14} \subset \dot{G}$. Set $V(\dot{G}) \setminus V(\dot{S}_{2}^{14}) = \{v_{15}, \dots, v_{n}\}$. Since $\dot{H}_{1}^{14}, \dot{H}_{2}^{14}, \dot{H}_{3}^{14}$ and \dot{S}_{2}^{14} are the only four signed graphs of order n = 14 such that $\dot{\mathcal{X}}_{1} \subset \dot{G}$ and $2 < \rho(\dot{G}) \leq \lambda^{*}$, then $v_{j} \not\sim v_{i}$ for each $j \geq 15$ and all $i \leq 9$. By forbidding \dot{F}_{2} , then $d_{v_{i}} \leq 2$ for all $i \geq 10$. Hence, $\dot{G} \sim \dot{S}_{2}^{n}$ and \dot{S}_{2}^{n} is maximal.

Case 2. $\dot{H}_i^{14} \subset \dot{G}$ for i = 1, 2 or 3. We will prove that $\dot{G} \subset \dot{S}_2^n$. Set $V(\dot{G}) \setminus V(\dot{H}_3^{14}) = \{v_{15}, \ldots, v_n\}$. If there is a vertex v_j $(j \geq 15)$ adjacent to some vertices of $\{v_1, \ldots, v_{13}\}$,




Figure 19: The signed graphs $\dot{B}_1 - \dot{B}_{19}$.

then $\dot{G}[\{v_1,\ldots,v_{13},v_j\}] \simeq \dot{S}_2^{14}$ and $\dot{S}_2^{14} \subset \dot{G}$. By case 1, then $\dot{G} \subset \dot{S}_2^n$. Otherwise, $v_j \not\sim v_i$ for each $j \geq 15$ and all $i \leq 13$. By forbidding \dot{F}_2 , then $d_{v_i} \leq 2$ for all $i \geq 9$. Hence, $\dot{G} \subset \dot{H}_3^n$ and $\dot{G} \subset \dot{S}_2^n$. The proofs of the cases $\dot{H}_1^{14} \subset \dot{G}$ or $\dot{H}_2^{14} \subset \dot{G}$ are similar. \square

4.4 \dot{G} is \dot{C}_k -free where k=3 or $k\geq 5$

In last subsection we shall consider that \dot{G} is \dot{C}_k -free where k=3 or $k\geq 5$, i.e., $k\neq 4$. For convenience, let \dot{H} be the induced subgraph of \dot{G} such that $\rho(\dot{H})\leq 2$. Since \dot{H} may not be unique, we choose one that the order of \dot{H} is maximal.

4.4.1 $\dot{\Theta}_{2,2,0} \subset \dot{G}$

First of all, we focus on that the order of \dot{G} is less than or equal to 14.

Lemma 4.6. Let $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ be a \dot{C}_k -free (where k = 3 or $k \geq 5$) bipartite signed graph of order $n \leq 14$. If $\dot{\Theta}_{2,2,0} \subset \dot{G}$, then \dot{G} is the induced subgraph of $\dot{H}_1, \ldots, \dot{H}_5, \dot{B}_1, \ldots, \dot{B}_{19}$ (up to switching equivalence). See Figs. 5 and 19.

Proof. If $\dot{\Theta}_{2,2,0} \subset \dot{G}$, then $\dot{\Theta}_{2,2,0} \subset \dot{H}$. Obviously, $\dot{H} \subset \dot{S}_{14}$ or $\dot{H} \subset \dot{S}_{16}$. By Fig. 1, then $\dot{H} \subset \dot{A}_i$ for $i = 1, \ldots, 10$ and $|V(\dot{H})| \leq 10$. See Fig. 18. Applying algorithm 1 to $A(\dot{H})$, then all signed graphs $\dot{G} \in \overline{\mathcal{G}}_S^{\lambda^*}$ of order $n \leq 14$ can be found, which are the induced subgraphs of $\dot{H}_1, \ldots, \dot{H}_5, \dot{B}_1, \ldots, \dot{B}_{19}$ (up to switching equivalence).

Remark 4.7. (1) By algorithm 1, we can check that the signed graphs \dot{B}_1 , \dot{B}_2 , \dot{B}_3 and \dot{B}_4 are maximal \dot{C}_k -free (where k=3 or $k \geq 5$) signed graphs.

- (2) $\dot{B}_1 \subset \dot{G}_8^3$, $\dot{B}_2 \subset \dot{G}_8^3$, $\dot{B}_3 \subset \dot{G}_6^6$, $\dot{B}_4 \subset \dot{G}_{10}$, $\dot{B}_i \subset [\dot{G}_4^{12}, v_{12}, s]$ for $i = 5, \dots, 8$ and $\dot{B}_i \subset [\dot{G}_2^9, v_9, s]$ for $i = 9, \dots, 19$.
- (3) If $n \geq 15$, then \dot{G} is $\{\dot{H}_1, \ldots, \dot{H}_5, \dot{B}_1, \ldots, \dot{B}_4\}$ -free. Furthermore, let V' be the vertex subset of $V(\dot{G}) \setminus V(\dot{\Theta}_{2,2,0})$ with order 8, then $\dot{G}[V(\dot{\Theta}_{2,2,0}) \cup V']$ is disconnected or switching equivalent to \dot{B}_i for one $i = 5, \ldots, 19$.

Secondly, we consider that $n \geq 15$. By Lemma 4.6, then $\dot{G}_i^{m_i}$ (i = 1, 2, 3 or 4) is an induced subgraph of \dot{G} , where $m_1 \geq 9$, $m_2 \geq 10$, $m_3 \geq 12$ and $m_4 \geq 14$. See Fig. 6. Without loss of generality, assume that the choice of m_i is largest. Let

$$V(\dot{G}_{1}^{m_{1}}) = V_{1} \cup V_{2}, \text{ where } V_{1} = \{v_{1}, \dots, v_{8}\} \text{ and } V_{2} = \{v_{9}, \dots, v_{m_{1}}\},$$

$$V(\dot{G}_{2}^{m_{2}}) = V_{1} \cup V_{2}, \text{ where } V_{1} = \{v_{1}, \dots, v_{9}\} \text{ and } V_{2} = \{v_{10}, \dots, v_{m_{2}}\},$$

$$V(\dot{G}_{3}^{m_{3}}) = V_{1} \cup V_{2}, \text{ where } V_{1} = \{v_{1}, \dots, v_{11}\} \text{ and } V_{2} = \{v_{12}, \dots, v_{m_{3}}\},$$

$$V(\dot{G}_{4}^{m_{4}}) = V_{1} \cup V_{2}, \text{ where } V_{1} = \{v_{1}, \dots, v_{12}\} \text{ and } V_{2} = \{v_{13}, \dots, v_{m_{4}}\}.$$

$$(4.1)$$

Lemma 4.8. Let x_1 and x_2 be two vertices in $V(\dot{G}) \setminus V(\dot{G}_i^{m_i})$ (i = 1, 2, 3 or 4). Then $d_{V_1}(x_1) = 0$ or $d_{V_2}(x_1) = 0$. Furthermore, if $x_1 \sim v_i$ and $x_2 \sim v_j$ where $v_i \in V_1$ and $v_j \in V_2$, then $x_1 \not\sim x_2$.

Proof. Suppose on the contrary that $d_{V_1}(x_1) \geq 1$ and $d_{V_2}(x_1) \geq 1$. Then $x_1 \sim v_i$ and $x_1 \sim v_j$ where $v_i \in V_1$ and $v_j \in V_2$. If $j \geq 15$, then $\{v_i, v_{i+1}, \ldots, v_j, x_1\}$ induces a cycle \dot{C}_{ℓ} ($\ell \geq 5$), which is a contradiction. If $j \leq 14$, then \dot{G} contains a \dot{C}_{ℓ} ($\ell \geq 5$) or $\dot{G}[\{v_1, \ldots, v_{13}, x_1\}]$ is not switching equivalent to \dot{B}_i for $i = 5, \ldots, 19$, which is a contradiction. So $d_{V_1}(x_1) = 0$ or $d_{V_2}(x_1) = 0$. Furthermore, if $x_1 \sim v_i$ and $x_2 \sim v_j$ where $v_i \in V_1$ and $v_j \in V_2$, and if $x_1 \sim x_2$, then $\dot{G}[\{v_1, \ldots, v_{12}, x_1, x_2\}]$ contains a \dot{C}_{ℓ} ($\ell \geq 5$) or is not switching equivalent to \dot{B}_i for $i = 5, \ldots, 19$, which is a contradiction. So $x_1 \not\sim x_2$. \square

Then we let

$$V(\dot{G}) \setminus V(\dot{G}_i^{m_i}) = U_0 \cup U_1 \cup U_2$$
, where $i = 1, 2, 3 \text{ or } 4$, (4.2)

where each vertex in U_0 is adjacent to no vertex of $V(\dot{G}_i^{m_i})$, each vertex in U_1 is adjacent to some vertices of V_1 and each vertex in U_2 is adjacent to some vertices of V_2 .

Set $|U_i| = a_i$ for i = 0, 1, 2, then we have




Figure 20: The signed graph $\dot{\Theta}_{2,2,0}^t$.

Lemma 4.9. (1) If U_0 is nonempty, then $a_0 = 1$ and there is a vertex w_2 in U_2 such that $\dot{G}[V(\dot{G}_i^{m_i}) \cup \{w_2\} \cup U_0] \sim \dot{\mathcal{A}}_{2i+5}^{n_1,n_2}$ for i = 1, 2, 3, 4. See Fig. 9.

(2) If $\dot{G}_{i}^{m_{i}} \subset \dot{G}$ for i = 1, 2, then $\dot{G}[V_{1} \cup U_{1}] \subset \dot{G}_{4}^{10}$; if $\dot{G}_{i}^{m_{i}} \subset \dot{G}$ for i = 3, 4, then $\dot{G}[V_{1} \cup U_{1}] \subset \dot{G}_{4}^{12}$.

Proof. (1) Let w_1 be a vertex in U_0 . Then there is a shortest path $P_{w_1v_j} = w_1 \dots w_\ell v_j$ ($\ell \geq 2$) from w_1 to a vertex $v_j \in V(\dot{G}_i^{m_i})$. If $v_j \in V_1$, we can find a connected induced subgraph \dot{G}' of \dot{G} with order $|V(\dot{G}')| = 14$ and is not switching equivalent to \dot{B}_i for $i = 5, \dots, 19$, which contradicts to Remark 4.7 (3). Then $v_j \in V_2$. Since \dot{G} is $\{\dot{F}_1, \dot{F}_2, \dot{G}_i^{m_i+1}\}$ -free, then $\ell = 2$ and $d_{V_2}(w_\ell) = 2$. Therefore, $\dot{G}[V(\dot{G}_i^{m_i}) \cup \{w_1, w_2\}] \sim \dot{\mathcal{A}}_{2i+5}^{n_1, n_2}$ for i = 1, 2, 3, 4. If $a_0 \geq 2$, then there is another vertex $w'_1 \neq w_1$ in U_0 and a vertex w'_2 in U_2 such that $\dot{G}[V(\dot{G}_i^{m_i}) \cup \{w'_1, w'_2\}] \sim \dot{\mathcal{A}}_{2i+5}^{n_1, n_2}$. And now we can check that $\dot{F}_1 \subset \dot{G}$, $\dot{F}_2 \subset \dot{G}$ or $\dot{F}_3 \subset \dot{G}$, which contradicts to Lemma 3.4. Hence, $a_0 = 1$.

(2) Choose $14 - (|V_1| + a_1)$ vertices (say vertex set V_1') from $V_2 \cup U_0 \cup U_2$ such that $\dot{G}[V_1 \cup V_1']$ is connected. By Remark 4.7 (3), then $\dot{G}[V_1 \cup V_1' \cup U_1] \sim \dot{B}_i$ for one $i = 5, \ldots, 19$. It is not hard to see that if $\dot{G}_i^{m_i} \subset \dot{G}$ for i = 1, 2, then $\dot{G}[V_1 \cup U_1] \subset \dot{G}_4^{10}$; if $\dot{G}_i^{m_i} \subset \dot{G}$ for i = 3, 4, then $\dot{G}[V_1 \cup U_1] \subset \dot{G}_4^{12}$.

The next lemma deals with the case that $\dot{\Theta}_{2,2,0}^t \subset \dot{G}$. See Fig. 20. Note that $\dot{\mathcal{A}}_6^{3,t} \sim \dot{\Theta}_{2,2,0}^t - u_2$. By Lemma 3.5 (6), we have $t \geq 6$. Then we let

$$V(\dot{G}) \setminus V(\dot{\Theta}_{2,2,0}^t) = X_0 \cup X_1 \cup X_2 \cup X_3,$$

where each vertex in X_0 is adjacent to no vertices of $V(\dot{\Theta}_{2,2,0}^t)$, each vertex in X_1 is adjacent to some vertices of $\{v_1,\ldots,v_6,w_1,w_2\}$, each vertex in X_2 is adjacent to some vertices of $\{u_1,\ldots,u_6,w_{t-1},w_t\}$ and each vertex in X_3 is adjacent to some vertices of $\{w_3,\ldots,w_{t-2}\}$. By Lemma 4.8, then $X_1\cap X_2=\emptyset$, $X_1\cap X_3=\emptyset$, $X_2\cap X_3=\emptyset$ and there is no edge between three parts X_1 , X_2 and X_3 . If $X_0\neq\emptyset$, let $u_0\in X_0$, Lemma 4.9 (1) implies that there is a path $P_{u_0z_i}=u_0u_1w_i$ (where $u_1\in X_3$) from the vertex u_0 to the vertex w_i . Then $\dot{F}_3\subset\dot{G}$, which contradicts to Lemma 3.4. So $X_0=\emptyset$.

Lemma 4.10. Let $\dot{G} \in \overline{\mathcal{G}}_{S}^{\lambda^{*}}$ be a \dot{C}_{k} -free $(k \neq 4)$ bipartite signed graph of order $n \geq 15$. If $\dot{\Theta}_{2,2,0}^{t} \subset \dot{G}$, then \dot{G} is the induced subgraph of \dot{H}_{6} , \dot{H}_{7} , \dot{H}_{8} , $[\dot{G}_{2}^{9}, v_{9}, s, v_{9}, \dot{G}_{2}^{9}]$, $[\dot{G}_{4}^{12}, v_{12}, s, v_{9}, \dot{G}_{2}^{9}]$ or $[\dot{G}_{4}^{12}, v_{12}, s, v_{12}, \dot{G}_{4}^{12}]$. See Figs. 5 and 7.

Proof. By Lemma 4.9 (2), then $\dot{G}[V(\dot{\Theta}_{2,2,0}^t) \cup X_1 \cup X_2] \sim (\dot{G}_i^{m_i}, v_{m_i}, s_1, v_{m_j}, \dot{G}_j^{m_j})$, where $(\dot{G}_i^{m_i}, v_{m_i}), (\dot{G}_j^{m_j}, v_{m_j}) \in \{(\dot{G}_1^8, v_8), (\dot{G}_2^9, v_9), (\dot{G}_3^{11}, v_{11}), (\dot{G}_4^{12}, v_{12})\}, s_1 \geq 3 \text{ if } i = 4 \text{ or } j = 4 \text{ (by forbidding } \dot{F}_{10} \text{ and } \dot{F}_{11}) \text{ and } s_1 \geq 2 \text{ otherwise (by Lemma 3.5 (6) and (8))}.$

Let $X_3 = \{x_{i_1}, \dots, x_{i_\ell}\}$, then $d_{\dot{\Theta}_{2,2,0}^t}(x_{i_j}) = 2$ (by forbidding \dot{F}_3). Up to switching equivalence, let $x_j \stackrel{+}{\sim} w_j$ and $x_j \stackrel{-}{\sim} w_{j+2}$ for $j = i_1, \dots, i_\ell$, where $j \geq 3$ and $j + 2 \leq t - 2$. Since \dot{G} is $\{\dot{F}_3, \dot{C}_k, \dot{C}_4^1\}$ -free $(k \neq 4)$, then $i_{j_1} \neq i_{j_2}$ if $j_1 \neq j_2, x_{i_{j_1}} \stackrel{-}{\sim} x_{i_{j_2}}$ if $|i_{j_1} - i_{j_2}| = 1$ and $x_{i_{j_1}} \not\sim x_{i_{j_2}}$ if $|i_{j_1} - i_{j_2}| \geq 2$. So, $\dot{G}[V(\dot{\Theta}_{2,2,0}^t) \cup X_3] \subset [\dot{G}_1^8, v_8, s_2, v_8, \dot{G}_1^8]$ $(s_2 \geq 3)$.

If $s_1 = 2$, then $\dot{G} \subset \dot{H}_i$ for i = 6, 7, 8. If $s_1 \geq 3$, then $\dot{G} \subset [\dot{G}_i^{m_i}, v_{m_i}, s, v_{m_j}, \dot{G}_j^{m_j}]$ $(s \geq 3)$, where $i, j \in \{1, 2, 3, 4\}$. By Lemma 3.5 (6), (8) and (10), then $m_i, m_j \geq 8$ if i, j = 1; $m_i, m_j \geq 9$ if i, j = 2; $m_i, m_j \geq 11$ if i, j = 3; $m_i, m_j \geq 12$ if i, j = 4. Hence, \dot{G} is the induced subgraph of $[\dot{G}_2^9, v_9, s, v_9, \dot{G}_2^9]$, $[\dot{G}_4^{12}, v_{12}, s, v_9, \dot{G}_2^9]$ or $[\dot{G}_4^{12}, v_{12}, s, v_{12}, \dot{G}_4^{12}]$. \square

Next we assume that \dot{G} is $\dot{\Theta}_{2,2,0}^t$ -free.

Lemma 4.11. Let $\dot{G} \in \overline{\mathcal{G}}_{S}^{\lambda^{*}}$ be a $\{\dot{C}_{k}, \dot{\Theta}_{2,2,0}^{t}\}$ -free $(k \neq 4)$ bipartite signed graph of order $n \geq 15$. If $\dot{\Theta}_{2,2,0} \subset \dot{G}$, then \dot{G} is the induced subgraph of $\dot{\mathcal{A}}_{i}^{n_{1},n_{2}}$ (i = 8, 9, 10, 11, 12, 13), $\dot{\mathcal{A}}_{i}^{n_{1}}$ (i = 16, 17, 18, 19) or $[\dot{G}, v, s, u, \dot{H}]$, where $(\dot{G}, v) \in \{(\dot{G}_{4}^{12}, v_{12}), (\dot{G}_{2}^{9}, v_{9})\}$ and $(\dot{H}, u) \in \{(T_{a,1,a-1}, v_{a-1}), (\dot{\mathcal{Q}}_{n_{1},n_{1}}^{\prime}, v_{n_{1}}), (\dot{G}_{5}^{10}, v_{10}), (P_{4}, v_{2})\}$, $s \geq 3$, $a \geq 3$ and $n_{1} \geq 1$.

Proof. Recall that $\dot{G}_{i}^{m_{i}}$ (i=1,2,3 or 4) is an induced subgraph of \dot{G} . Here, we only consider that $\dot{G}_{4}^{m_{4}} \subset \dot{G}$. The proofs of the cases $\dot{G}_{i}^{m_{i}} \subset \dot{G}$ for i=1,2,3 are similar. We use the vertex partitions of Eqs. (4.1) and (4.2). By Lemma 4.9 (2), then $U_{1} = \emptyset$. Since \dot{G} is $\dot{\Theta}_{2,2,0}^{t}$ -free, then each vertex in U_{2} is adjacent to at most two vertices of V_{2} . Then let $U_{2} = U_{21} \cup U_{22}$, where each vertex in U_{2i} (i=1 or 2) is adjacent to i vertices of V_{2} . Furthermore, set $U_{21} = \{u_{j_{1}}, \ldots, u_{j_{p}}\}$ $(13 \leq j_{1} \leq \cdots \leq j_{p} \leq m_{4} - 1)$ where $u_{j} \stackrel{+}{\sim} v_{j}$ for $j=j_{1},\ldots,j_{p}$, and $U_{22} = \{u'_{i_{1}},\ldots,u'_{i_{q}}\}$ $(13 \leq i_{1} \leq \cdots \leq i_{q} \leq m_{4} - 2)$ where $u'_{i} \stackrel{+}{\sim} v_{i}$ and $u'_{i} \stackrel{-}{\sim} v_{i+2}$ for $i=i_{1},\ldots,i_{q}$. Since \dot{G} is $\{\dot{\Theta}_{2,2,0}^{t}, \dot{F}_{3}, \dot{C}_{k}, \dot{C}_{4}^{t}, \dot{G}_{4}^{m_{4}+1}\}$ -free $(k \neq 4)$, we have $(\mathbf{A1}) \ \dot{G}[U_{21}]$ is K_{1} or K_{2}^{-} . If $\dot{G}[U_{21}]$ is K_{2}^{-} , then $1 \leq m_{4} - j_{2} \leq 2$ (by forbidding \dot{F}_{8}), $(\mathbf{A2}) \ i_{j_{1}} \neq i_{j_{2}}$ if $j_{1} \neq j_{2}$, $u'_{i_{j_{1}}} \stackrel{-}{\sim} u'_{i_{j_{2}}}$ if $|i_{j_{1}} - i_{j_{2}}| = 1$ and $u'_{i_{j_{1}}} \not\sim u'_{i_{j_{2}}}$ if $|i_{j_{1}} - i_{j_{2}}| \geq 2$. Then $\dot{G}[V(\dot{G}_{4}^{m_{4}}) \cup U_{21}] \sim \dot{A}_{12}^{n_{1},n_{2}}$, $\dot{A}_{13}^{0,n_{2}}$ or $\dot{A}_{19}^{n_{1}}$ and $\dot{G}[V(\dot{G}_{4}^{m_{4}}) \cup U_{22}] \subset [\dot{G}_{4}^{12}, v_{12}, s]$.

Case 1. U_0 is empty. If U_{21} or U_{22} is empty, then we are done. If U_{21} and U_{22} are not empty, according to $(\mathbf{A1})$, we break into two subcases:

Subcase 1.1. $\dot{G}[U_{21}] = K_1$. Then $U_{21} = \{u_{j_1}\}, i_q \leq j_1 - 1 \text{ and } i_q \neq j_1 - 2 \text{ (otherwise } \dot{F}_1 \subset \dot{G} \text{ or } \dot{F}_3 \subset \dot{G}, \text{ which contradicts to Lemma 3.4).}$

Subsubcase 1.1.1. $i_q = j_1 - 3$. By (**A2**), then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s_1, v_2, P_\ell]$. By forbidding $Q_{1,1,3}$, then $\ell \leq 4$. So, $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_2, P_4]$.

Subsubcase 1.1.2. $i_q \leq j_1 - 4$. By (**A2**), then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s_1, v_c, T_{c,1,a}]$. If a = 1 or 2, then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_2, P_4]$. If $a \geq 3$, by forbidding $Q_{1,c+1,a}$ (where $c \leq a - 2$), then $c \geq a - 1$. So, $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{a-1}, T_{a-1,1,a}]$.

Subsubcase 1.1.3. $i_q = j_1 - 1$. Then $u'_{i_q} \stackrel{+}{\sim} v_{j_1-1}$ and $u'_{i_q} \stackrel{-}{\sim} v_{j_1+1}$. If $u'_{i_q} \sim u_{j_1}$, then $u'_{i_q} \stackrel{-}{\sim} u_{j_1}$ and $j_1 = m_4 - 1$ (by forbidding \dot{C}_4^1). By (**A2**), then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s]$. If $u'_{i_q} \not\sim u_{j_1}$, by forbidding \dot{C}_4^1 and \dot{F}_1 , then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s_1, v_{n_2}, \dot{Q}'_{n_1, n_2}]$. By Lemma 3.5 (2), then $n_2 \geq 1$ if $n_1 = 0$ and $n_2 \geq n_1$ if $n_1 \geq 1$. So, $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{n_1}, \dot{Q}'_{n_1, n_1}]$ $(n_1 \geq 1)$.

Subcase 1.2. $\dot{G}[U_{21}] = K_2^-$. Then $U_{21} = \{u_{j_1}, u_{j_2}\}, \ j_2 = j_1 + 1 \text{ and } u_{j_1} \sim u_{j_2}$. Since \dot{G} is $\{\dot{\Theta}_{2,2,0}^t, \dot{F}_1\}$ -free, then $i_q \leq j_1 - 3$. If $j_2 = m_4 - 1$, then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{n_2}, \dot{Q}'_{0,n_2}]$, which has been done in subsubcase 1.1.3. If $j_2 = m_4 - 2$, then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{n_1}, \dot{G}_5^{n_1}]$. By Lemma 3.5 (4), then $n_1 \geq 10$. So, $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{10}, \dot{G}_5^{10}]$ $(s \geq 3)$.

Case 2. U_0 is nonempty. Then $|U_{21}| = 0$ as \dot{G} is $\{\dot{F}_1, \dot{F}_3\}$ -free. Similarly, by (A2) and forbidding $\dot{\Theta}_{2,2,0}^t, \dot{F}_1, \dot{C}_4^1$, then $\dot{G} \subset [\dot{G}_4^{12}, v_{12}, s, v_{n_2}, \dot{Q}'_{n_1,n_2}]$, which has been done in subsubcase 1.1.3.

This completes the proof.

4.4.2 \dot{G} is $\dot{\Theta}_{2,2,0}$ -free

Lastly it remains that \dot{G} is $\dot{\Theta}_{2,2,0}$ -free. First we suppose that $\dot{H} \subset \dot{T}_{2k}$. Notice that if $\dot{H} \subset \dot{T}'_{2k}$, then $\dot{H} \subset \dot{T}'_{2n}$ for all $n \geq k$. Without loss of generality, we assume that $\dot{H} \subset \dot{T}'_{2t}$, where t is maximal. Up to switching isomorphic, let $V(\dot{H}) = V_1 \cup U_1$ where $V_1 = \{v_1, v_2, \dots, v_t\}$ and $U_1 = \{u_{k_1}, \dots, u_{k_\ell}\}$ $(1 \leq k_1 < \dots < k_\ell \leq t)$. Let w be a good vertex in $V(\dot{G}) \setminus V(\dot{H})$. If t = 3 or 4, applying algorithm 1 to $A(\dot{H})$, then there is no signed graph \dot{G} with order $|V(\dot{H})| + 1$ and $2 < \rho(\dot{G}) \leq \lambda^*$. Next we let $t \geq 5$.

Before giving the main result of this subsubsection, we need some preliminary lemmas.

Lemma 4.12. $d_{V_1}(w) \leq 1$ and $d_{U_1}(w) \leq 1$.

Proof. Since \dot{G} is $\{\dot{\Theta}_{2,2,0}, \dot{C}_{4}^{1}\}$ -free, then $d_{V_{1}}(w) \leq 2$. If $d_{V_{1}}(w) = 2$, then $w \stackrel{+}{\sim} v_{i-1}$ and

 $w \stackrel{-}{\sim} v_{i+1}$ (switching at w if necessary). Since \dot{G} is $\{\dot{C}_4^1, \dot{\Theta}_{2,2,0}, \dot{C}_k\}$ -free $(k \neq 4)$, then $u_i \notin U_1, \ w \not\sim u_j$ for all $j \neq i \pm 1$ and $w \stackrel{-}{\sim} u_{i\pm 1}$ if $u_{i\pm 1} \in U_1$. So $\dot{H}_U(w) \simeq \dot{G}[V(\dot{H}) \cup \{u_i\}]$ and $\dot{H}_U(w) \subset \dot{T}'_{2t}$, which is a contradiction. Hence, $d_{V_1}(w) \leq 1$. Similarly, we have $d_{U_1}(w) \leq 1$.

Then we let

$$V(\dot{G}) \setminus V(\dot{H}) = V_0 \cup V_1' \cup U_1' \cup Y,$$

where each vertex in V_0 is adjacent to no vertex of $V_1 \cup U_1$, each vertex in V'_1 is adjacent to exactly one vertex of V_1 and no vertex of U_1 , each vertex in U'_1 is adjacent to exactly one vertex of U_1 and no vertex of V_1 , each vertex in Y is adjacent to exactly one vertex of V_1 and exactly one vertex of U_1 .

Lemma 4.13. *If* $w \in Y$, then

- (i) $w \stackrel{+}{\sim} v_{t-2}$, $w \stackrel{-}{\sim} u_t$ and $u_i \notin U_1$ for i = t 1, t 2, t 3, t 4, or
- (ii) $w \stackrel{+}{\sim} v_3$, $w \stackrel{+}{\sim} u_1$ and $u_i \notin U_1$ for i = 2, 3, 4, 5.

Proof. Let $w \sim v_i$ and $w \sim u_j$. Since \dot{G} is \dot{C}_k -free $(k \neq 4)$, then j = i or $j = i \pm 2$. If i = j, then $w \stackrel{+}{\sim} v_i$ and $w \stackrel{-}{\sim} u_i$ (switching at w if necessary). By forbidding \dot{C}_4^1 , then i = t - 1. If $u_t \not\in U_1$ (resp. $u_t \in U_1$), then $\dot{H}_U(w) \subset \dot{T}'_{2t}$ (resp. $\dot{G}[\{v_{t-1}, v_t, u_{t-1}, u_t, w\}] \sim \theta_{1,1,1}$), which is a contradiction. So $j = i \pm 2$. If j = i + 2, since \dot{G} is $\{\dot{C}_4^1, \dot{\Theta}_{2,2,0}, \dot{F}_1\}$ -free, then t = i + 2, $w \stackrel{+}{\sim} v_{t-2}$, $w \stackrel{-}{\sim} u_t$ and $u_i \not\in U_1$ for i = t - 1, t - 2, t - 3. If $u_{t-4} \in U_1$, then $\dot{G}[v_t, v_{t-1}, v_{t-2}, v_{t-3}, v_{t-4}, u_t, u_{t-4}, w] \sim \dot{A}_2^{0,1,0}$. By Lemma 3.5 (2), then $\rho(\dot{G}) \geq \rho(\dot{A}_2^{0,1,0}) > \lambda^*$, which is a contradiction. So $u_{t-4} \not\in U_1$. If j = i - 2, because of symmetry, then i = 3, $w \stackrel{+}{\sim} v_3$, $w \stackrel{+}{\sim} u_1$ and $u_i \not\in U_1$ for i = 2, 3, 4, 5.

Lemma 4.14. V_0 is empty.

Proof. For a contradiction, assume that $w \in V_0$. Then there is a shortest path $P_{wx_{i+1}} = ww_1 \dots w_\ell x_{i+1}$ ($\ell \geq 1$) from w to $x_{i+1} \in V(\dot{H})$ where x = v or u. Since t is maximal, then $\ell + 1 \leq i$ and $\ell + 1 \leq t - (i+1)$. By forbidding $T_{2,3,4}$, then $\ell = 1$, i = 2 and $t \geq 5$; or $\ell = 1$, i = t - 3 and $t \geq 5$; or $\ell = 1$, i = 3 and t = 7. By forbidding \dot{F}_1 and \dot{F}_2 , if x = v, then $V_0 = \{w\}$, $U_1 = \emptyset$ or $U_1 = \{u_{i+1}\}$, $V'_1 = \{w_1\}$, $Y = \emptyset$ and $|U'_1| \leq 1$; if x = u, then $V_0 = \{w\}$, $U_1 = \{u_{i+1}\}$, $U'_1 = \{w_1\}$, $Y = \emptyset$ and $|V'_1| \leq 1$. Therefore, \dot{G} is a tree or unicyclic, which has been done in Theorem 2.1 and Lemma 3.7. Hence, V_0 is empty. \Box

Lemma 4.15. If $w \in V_1'$, then

- (i) if $w \sim v_2$, then $u_2 \in U_1$ and $u_i \notin U_1$ for i = 1, 3, 4, 5,
- (ii) if $w \sim v_{t-1}$, then $u_{t-1} \in U_1$ and $u_i \notin U_1$ for i = t, t-2, t-3, t-4,
- (ii) if $w \sim v_i$ (3 \le i \le t-2), then $u_{i\pm 1} \notin U_1$ and one of the following holds: (1) $k_1 = i$ and $k_2 \geq i+2$; (2) $k_1 \geq i+2$; (3) $k_\ell = i$ and $k_{\ell-1} \leq i-2$; (4) $k_\ell \leq i-2$.
- Proof. (i) By forbidding \dot{F}_1 , then $u_1 \notin U_1$. Since $\dot{H}_U(w)$ is not an induced subgraph of \dot{T}'_{2t} , then $u_2 \in U_1$. By forbidding \dot{F}_1 and \dot{C}^1_4 , then $u_4 \notin U_1$ and $u_3 \notin U_1$. If $u_5 \in U_1$, then $\dot{G}[v_1, v_2, v_3, v_4, v_5, u_2, u_5, w] \sim \dot{A}^{0,1,0}_2$. By Lemma 3.5 (2), then $\rho(\dot{G}) \geq \rho(\dot{A}^{0,1,0}_2) > \lambda^*$, which is a contradiction. Hence, $u_5 \notin U_1$.
 - (ii) The proof is similar to (i).
- (iii) By forbidding \dot{F}_1 , then $u_{i\pm 1} \notin U_1$. By forbidding \dot{F}_3 , it is impossible that $k_{j_1} \leq i-2$ and $k_{j_2} \geq i+2$ for $1 \leq j_1 < j_2 \leq \ell$. So, $k_1 \geq i$ or $k_{\ell} \leq i$. If $k_1 = i$, then $k_2 \geq i+2$. Otherwise, $k_1 \geq i+2$. If $k_{\ell} = i$, then $k_{\ell-1} \leq i-2$. Otherwise, $k_{\ell} \leq i-2$.

Now we are going to give the main result of this subsubsection. Set $V'_1 = \{v'_{i_1}, \ldots, v'_{i_p}\}$ where $v'_i \stackrel{+}{\sim} v_i$ for each $i = i_1, \ldots, i_p$, and $U'_1 = \{u'_{j_1}, \ldots, u'_{j_q}\}$ where $u'_j \stackrel{+}{\sim} u_j$ for each $j = j_1, \ldots, j_q$. If $i_{s_1} = j_{s_2}$ for one $1 \leq s_1 \leq p$ and one $1 \leq s_2 \leq q$, by forbidding \dot{F}_1 , \dot{F}_2 and \dot{C}_4^1 , we have $|U_1| = |V'_1| = |U'_1| = 1$ and |Y| = 0. So, $\dot{G} \sim \dot{C}_4[n_1, 1, t - n_1, 1]$ (see Lemma 3.7). Otherwise, $i_{s_1} \neq j_{s_2}$ for all $s_1 = 1, \ldots, p$ and all $s_2 = 1, \ldots, q$. Then there is a switching isomorphic of \dot{G} such that $u'_j \not\sim u_j$ and $u'_j \stackrel{+}{\sim} v_j$ for all $j = j_1, \ldots, j_q$, that is, U'_1 becomes empty and $V'_1 = \{v'_{i_1}, \ldots, v'_{i_p}, u'_{j_1}, \ldots, u'_{j_q}\}$.

Lemma 4.16. Let $\dot{G} \in \overline{\mathcal{G}}_{S}^{\lambda^{*}}$ be a $\{\dot{C}_{k}, \dot{\Theta}_{2,2,0}\}$ -free $(k \neq 4)$ bipartite signed graph with $m \geq n+1$. Then \dot{G} is an induced subgraph of $\dot{\mathcal{A}}_{3}^{n_{1},n_{2},n_{3}}, \dot{\mathcal{A}}_{5}^{n_{1},n_{2}}$ or $[\dot{G},v,s,u,\dot{H}]$, where $(\dot{G},v),(\dot{H},u) \in \{(P_{4},v_{2}),(T_{a,1,a-1},v_{a-1}),(\dot{Q}'_{n_{1},n_{1}},v_{n_{1}}),(\dot{G}_{5}^{10},v_{10})\}, s \geq 3, a \geq 3 \text{ and } n_{1} \geq 1.$

Proof. Since \dot{G} is $\{\dot{\Theta}_{2,2,0}, \dot{C}_k\}$ -free $(k \neq 4)$, then $\dot{G}[V_1'] = t_1 K_2^- \cup t_2 K_1$, where $t_1 + t_2 \leq 2$ (by forbidding \dot{F}_3). By Lemma 4.13, we break into three cases.

Case 1. |Y| = 0.

Subcase 1.1. $\dot{G}[V_1'] = K_1$. Then $V_1' = \{v_{i_1}'\}$. Since t is maximal, then $2 \le i_1 \le t - 1$. Recall that $|V(\dot{H})|$ is maximal. By Lemma 4.15, then we have

Subsubcase 1.1.1. $i_1 = 2$ or $i_1 = t - 1$. Then $\dot{G} \subset [\dot{Q}'_{1,0}, v_1, s] \subset [\dot{Q}'_{1,1}, v_1, s]$.

Subsubcase 1.1.2. $3 \le i_1 \le t - 2$. If $k_1 = i_1$ or $k_\ell = i_1$, then $\dot{G} \subset [\dot{Q}'_{n_1,n_2}, v_{n_2}, s]$ $(n_1 \ge 1)$, where $n_2 = n_1$ (by Lemma 3.5 (2)). If $k_1 = i_1 + 2$ or $k_\ell = i_1 - 2$, then $\dot{G} \subset [P_\ell, v_2, s]$, where $\ell = 4$ (by forbidding $Q_{1,1,3}$). If $k_1 \ge i_1 + 3$ or $k_\ell \le i_1 - 3$, then $\dot{G} \subset [T_{a,1,b}, v_b, s]$ $(a \ge 3$ and $b \ge 1)$, where b = a - 1 (by Lemma 3.5 (1)).

Subcase 1.2. $\dot{G}[V_1'] = 2K_1$. Then $V_1' = \{v_{i_1}', v_{i_2}'\}$. Similar to subcase 1.1, then $\dot{G} \subset \dot{\mathcal{A}}_3^{n_1, n_2, n_3}$ or $\dot{G} \subset [\dot{G}, v, s, u, \dot{H}]$, where $(\dot{G}, v), (\dot{H}, u) \in \{(T_{a,1,a-1}, v_{a-1}), (\dot{Q}'_{n_1,n_1}, v_{n_1}), (P_4, v_2)\}$ $(a \geq 3 \text{ and } n_1 \geq 1)$.

Subcase 1.3. $\dot{G}[V_1'] = K_2^-$. Then $V_1' = \{v_{i_1}', v_{i_2}'\}$, $i_2 = i_1 + 1$ and $v_{i_1}' - v_{i_2}'$. Without loss of generality, we may assume that $i_1 \leq \lfloor \frac{t}{2} \rfloor$. Since $|V(\dot{H})|$ is maximal, then $i_2 \geq 4$. By Lemma 4.15, then $k_1 \geq i_2 + 2$. So, $Q_{i_2-1,k_1-i_2-1,1} \subset \dot{G}$. By forbidding $Q_{i_2-1,b,1}$ (where $b \leq i_2 - 2$), then $k_1 \geq 2i_2$. If $i_2 = 5$ or $i_2 \geq 6$, then $\dot{F}_8 \subset \dot{G}$ or $\dot{F}_4 \subset \dot{G}$, respectively, which contradicts to Lemma 3.4. So, $i_2 = 4$ and $\dot{G} \subset [G_5^{10}, v_{10}, s]$.

Subcase 1.4. $\dot{G}[V_1'] = K_2^- \cup K_1$. Then $V_1' = \{v_{i_1}', v_{i_2}', v_{i_3}'\}$, where $i_2 = i_1 + 1$ and $v_{i_1}' - v_{i_2}'$. By subcases 1.1 and 1.3, then $\dot{G} \subset \dot{\mathcal{A}}_5^{n_1, n_2}$ or $\dot{G} \subset [G_5^{10}, v_{10}, s, v, \dot{G}]$, where $(\dot{G}, v) \in \{(P_4, v_2), (T_{a,1,a-1}, v_{a-1}), (\dot{Q}'_{n_1,n_1}, v_{n_1})\}$ $(a \geq 3 \text{ and } n_1 \geq 1)$.

Subcase 1.5. $\dot{G}[V_1'] = 2K_2^-$. Similar to subcase 1.3, then $\dot{G} \subset [G_5^{10}, v_{10}, s, v_{10}, G_5^{10}]$.

Case 2. |Y| = 1. By forbidding \dot{F}_1 and \dot{F}_3 , then $t_1 + t_2 \leq 1$.

Subcase 2.1. $|V_1'| = 0$. By Lemma 4.13, then $\dot{G} \subset [\dot{Q}'_{1,0}, v_1, s] \subset [\dot{Q}'_{1,1}, v_1, s]$.

Subcase 2.2. $\dot{G}[V_1'] = K_1$. By subcases 1.1 and 2.1, then $\dot{G} \subset \dot{\mathcal{A}}_3^{0,n_2,n_3}$ or $\dot{G} \subset [\dot{Q}_{1,1}', v_1, s, v, \dot{G}]$, where $(\dot{G}, v) \in \{(T_{a,1,a-1}, v_{a-1}), (\dot{Q}_{n_1,n_1}', v_{n_1}), (P_4, v_2)\}$ $(a \geq 3 \text{ and } n_1 \geq 1)$.

Subcase 2.3. $\dot{G}[V_1'] = K_2^-$. By subcases 1.3 and 2.1, then $\dot{G} \subset \dot{\mathcal{A}}_5^{0,n_2}$ or $\dot{G} \subset [\dot{Q}_{1,1}', v_1, s, v_{10}, \dot{G}_5^{10}]$.

Case 3. |Y| = 2. Then $k_1 = 1$ and $k_{\ell} = t$. By forbidding \dot{F}_1 and \dot{F}_3 , then V'_1 is empty. If $|U_1| = 2$, then $\dot{G} \sim \dot{\mathcal{A}}_3^{0,0,n_3}$. If $|U_1| \geq 3$, then $6 \leq k_2 < \cdots < k_{\ell-1} \leq t-5$ (by Lemma 4.13). So $\dot{G} \subset [\dot{Q}'_{1,0}, v_1, s, v_1, \dot{Q}'_{1,0}] \subset [\dot{Q}'_{1,1}, v_1, s, v_1, \dot{Q}'_{1,1}]$.

This completes the proof.

Remark 4.17. Finally, it remains the case that \dot{H} is an induced subgraph of \dot{S}_{14} or \dot{S}_{16} but not an induced subgraph of \dot{T}_{2k} . Since $m \geq n+1$, then $\dot{\mathcal{B}}_r^{4,4}$, $\dot{\mathcal{B}}_0^{4,4}$ or $\dot{\Theta}_{1,1,1}$ is an induced subgraph of \dot{G} . More importantly, $\dot{\mathcal{B}}_r^{4,4}$, $\dot{\mathcal{B}}_0^{4,4}$ or $\dot{\Theta}_{1,1,1}$ is also an induced subgraph of \dot{H} . From Fig. 1, it is not too hard to get that there is no such signed graph \dot{H} .

Proof of Theorem 2.4. Summarizing with all results of Theorems 2.1 and 2.2, Lemmas

3.5, 3.7, 4.2, 4.4, 4.5, 4.6, 4.10, 4.11, 4.16 and 4.21, we complete the proof of Theorem 2.4. \Box

Acknowledgments

This project is supported by the National Natural Science Foundation of China (No.119 71164, 12001185, 12100557) and the Zhejiang Provincial Natural Science Foundation of China (LQ21A010004).

References

- [1] F. Belardo, S. Cioabă, J. Koolen, J. F. Wang, Open problems in the spectral theory of signed graphs, *Art Discrete Appl. Math.* 1 (2018) #P2.10.
- [2] A. E. Brouwer, A. Neumaier, The graphs with spectral radius between 2 and $\sqrt{2+\sqrt{5}}$, Linear Algebra Appl. 114/115 (1989) 273–276.
- [3] A. E. Brouwer, W. H. Haemers, Spectra of graphs, Springer, 2011.
- [4] F. C. Bussemaker, P. J. Cameron, J. J. Seidel, S. V. Tsaranov, Tables of signed graphs, Eut Report 91-WSK-01, Eindhoven, 1991.
- [5] D. Cvetković, M. Doob, I. Gutman, On graphs whose spectral radius does not exceed $\sqrt{2+\sqrt{5}}$, Ars Combinat. 14 (1982) 225–239.
- [6] M. K. Gill, B. D. Acharya, A recurrence formula for computing the characteristic polynomial of a sigraph, J. Comb. Inf. Syst. Sci. 5 (1980) 68–72.
- [7] G. Greaves, J. Koolen, A. Munemasa, Y. Sano, T. Taniguchi, Edge-signed graphs with smallest eigenvalue greater than -2, J. Combin. Theory Ser. B 110 (2015) 90–111.
- [8] A. J. Hoffman, On limit points of spectral radii of non-negative symmetric integral matrices, in: Y. Alavi, et al. (Eds.), Lecture Notes Math, vol. 303, Springer-Verlag, Berlin, 1972, pp. 165–172.
- [9] Z. L. Jiang, A. Polyanskii, Forbidden subgraphs of bounded spectral radius with applications to equiangular lines, *Israel J. Math.* 236 (2020) 393–421.

- [10] Z. L. Jiang, A. Polyanskii, Forbidden induced subgraphs of graphs and signed graphs with eigenvalues bounded from below, arXiv: 2111.10366v1.
- [11] P. W. H. Lemmens, J. J. Seidel, Equiangular lines, J. Algebra 24 (1973) 494–512.
- [12] J. McKee, C. Smyth, Integer symmetric matrices having all their eigenvalues in the interval [-2, 2], J. Algebra 317 (2007) 260–290.
- [13] J. H. Smith, Some properties of the spectrum of a graph, Combinatorial Structures and their Applications, Gordon and Breach, New York, 1970, pp. 403–406.
- [14] J. F. Wang, J. Wang, M. Brunetti, The hoffman program of graphs: old and new, arXiv: 2012.13079v1.
- [15] T. Zaslavsky, Signed graphs, Discrete Appl. Math. 4 (1982) 47–74.

Appendix A

Gill and Acharya [6] obtained the following recurrence formula for the characteristic polynomial of a signed graph.

Lemma 4.18. [6] Let \dot{G} be a signed graph and v be its arbitrary vertex. Then

$$\phi_{\dot{G}}(x) = x\phi_{\dot{G}-v}(x) - \sum_{vu \in E(\dot{G})} \phi_{\dot{G}-v-u}(x) - 2\sum_{\dot{C} \in \dot{C}(v)} \sigma(\dot{C})\phi_{\dot{G}-\dot{C}}(x),$$

where $\dot{C}(v)$ denotes the set of signed cycles passing through v, and $\dot{G} - \dot{C}$ denotes the signed graph obtained from \dot{G} by deleting \dot{C} .

Denote p_n and q_n the characteristic polynomials of P_n and C_n , respectively. Then

$$p_0(x) = 1$$
, $p_1(x) = x$, $p_n(x) = xp_{n-1}(x) - p_{n-2}(x)$, $q_{n+1}(x) = p_{n+1}(x) - p_{n-1}(x) - 2$,

for all $n \geq 2$. Moreover, the recursion gives

$$p_n(x) = \frac{\theta^{2n+2} - 1}{\theta^{n+2} - \theta^n}, \quad q_n(x) = \theta^n + \theta^{-n} - 2, \quad \text{where} \quad \theta = \theta(x) := \frac{x + \sqrt{x^2 - 4}}{2}.$$
 (4.3)

Then

$$\phi_{T_{a,1,b}}(x) = xp_{a+b+1} - p_a p_b,$$

$$\phi_{Q_{a,b,c}}(x) = x^2 p_{a+b+c+1} - xp_{a+b} p_c - xp_a p_{b+c} + p_a p_{b-1} p_c.$$
(4.4)

Let \dot{T}_{2k} be the signed graph depicted in Fig. 1 with the adjacency matrix A_{σ} , and let $V(\dot{T}_{2k}) = V_1 \cup V_2$ be a partition of the vertex set. Then

$$A_{\sigma} = \begin{bmatrix} A_1 & B \\ B^T & A_2 \end{bmatrix}$$
, where A_i $(i = 1, 2)$ is the adjacency matrix of $\dot{T}_{2k}[V_i]$.

It is known that the spectrum of \dot{T}_{2k} is $\{2^k, -2^k\}$ (see [12, Theorem 1]). Then

$$(A_{\sigma})^2 = \begin{bmatrix} (A_1)^2 + BB^T & A_1B + BA_2 \\ B^T A_1 + A_2 B^T & B^T B + (A_2)^2 \end{bmatrix} = 4I_{2k} \text{ and } B^T A_1 + A_2 B^T = \mathbf{0}.$$

Lemma 4.19. If $\mu \neq \pm 2$ is an eigenvalue of $\dot{T}_{2k}[V_1]$, then $-\mu$ is an eigenvalue of $\dot{T}_{2k}[V_2]$.

Proof. Let β be the eigenvector corresponding to the eigenvalue $\mu \neq \pm 2$ of $T_{2k}[V_1]$. If $B^T\beta = \mathbf{0}$, then

$$A_{\sigma} \begin{bmatrix} \beta \\ \mathbf{0} \end{bmatrix} = \begin{bmatrix} A_1 & B \\ B^T & A_2 \end{bmatrix} \begin{bmatrix} \beta \\ \mathbf{0} \end{bmatrix} = \begin{bmatrix} A_1 \beta \\ B^T \beta \end{bmatrix} = \begin{bmatrix} A_1 \beta \\ \mathbf{0} \end{bmatrix} = \mu \begin{bmatrix} \beta \\ \mathbf{0} \end{bmatrix}.$$

This gives that μ is an eigenvalue of A_{σ} and $\mu = \pm 2$, which contradicts to hypothesis. So $B^T \beta \neq \mathbf{0}$. Since $B^T A_1 + A_2 B^T = \mathbf{0}$, then $A_2 B^T \beta = -B^T A_1 \beta = -B^T \mu \beta = -\mu B^T \beta$. Hence, $-\mu$ is an eigenvalue of $\dot{T}_{2k}[V_2]$.

Corollary 4.20. (i)
$$\phi_{\dot{T}'_{2s}}(x) = x^2(x \pm 2)^{s-1}$$
, (ii) $\phi_{\dot{T}''_{2s}}(x) = x(x \pm \sqrt{2})(x \pm 2)^{s-2}$, (iii) $\phi_{\dot{T}''_{2s}}(x) = (x \pm \sqrt{2})^2(x \pm 2)^{s-3}$.

Proof. Note that $V(\dot{T}_{2(s+1)}) = V(\dot{T}'_{2s}) \cup V(2K_1)$, $V(\dot{T}_{2(s+1)}) = V(\dot{T}''_{2s}) \cup V(P_3, \sigma)$ and $V(\dot{T}_{2(s+1)}) = V(\dot{T}'''_{2s}) \cup V(\dot{C}_4)$. By Lemma 3.1, then ± 2 is an eigenvalue of \dot{T}'_{2s} (resp. \dot{T}'''_{2s}) with multiplicity at least s-1 (resp. s-2, s-3).

(i) It is known that the spectrum of $2K_1$ is $\{0^2\}$. By Lemma 4.19, then 0 is an eigenvalue of \dot{T}'_{2s} with multiplicity 2. Therefore, $\phi_{\dot{T}'_{2s}}(x) = x^2(x \pm 2)^{s-1}$.

The proofs of (ii) and (iii) are similar since the spectrum of (P_3, σ) is $\{\pm\sqrt{2}, 0\}$ and the spectrum of \dot{C}_4^- is $\{\pm\sqrt{2}^2\}$.

Let the pair (\dot{G}, v) be defined in Theorem 2.4 (iv) and (v), then

$$\phi_{[\dot{G},v,s]}(x) = x\phi_{\dot{T}_{2s}^{"}}(x)\phi_{\dot{G}-v}(x) - \phi_{\dot{T}_{2(s-1)}^{"}}(x)\phi_{\dot{G}-v}(x) - \phi_{\dot{T}_{2s}^{"}}(x)\sum_{uv\in E(\dot{G})}\phi_{\dot{G}-v-u}(x)$$

$$= x^{2}(x\pm\sqrt{2})(x\pm2)^{s-2}\phi_{\dot{G}-v}(x) - x^{2}(x\pm2)^{s-2}\phi_{\dot{G}-v}(x)$$

$$- x(x\pm\sqrt{2})(x\pm2)^{s-2}\sum_{uv\in E(\dot{G})}\phi_{\dot{G}-v-u}(x) \text{ (by Corollary 4.20)}$$

$$= (x\pm2)^{s-2}(x^{2}(x\pm\sqrt{2})\phi_{\dot{G}-v}(x) - x^{2}\phi_{\dot{G}-v}(x) - x(x\pm\sqrt{2})\sum_{uv\in E(\dot{G})}\phi_{\dot{G}-v-u}(x))$$

$$= (x\pm2)^{s-2}\phi_{[\dot{G},v,2]}(x),$$

$$(4.5)$$

and $\phi_{[\dot{G},v,s]}(\lambda^*) = (\sqrt{5} - 2)^{s-2}\phi_{[\dot{G},v,2]}(\lambda^*)$. Similarly, we have $\phi_{(\dot{G},v,s)}(\lambda^*) = (\sqrt{5} - 2)^{s-3}\phi_{(\dot{G},v,3)}(\lambda^*),$

$$\phi_{(\dot{G},v,s)}(\lambda^*) = (\sqrt{5} - 2)^{s-3}\phi_{(\dot{G},v,3)}(\lambda^*),$$

$$\phi_{[\dot{G},v,s,u,\dot{H}]}(\lambda^*) = (\sqrt{5} - 2)^{s-3}\phi_{[\dot{G},v,3,u,\dot{H}]}(\lambda^*).$$
(4.6)

By Lemma 4.18 and Eqs. (4.3)–(4.6), we can obtain the characteristic polynomials of all signed graphs \dot{G} of Theorem 2.4 (*iii*) and (*iv*). More importantly, by computations (using Wolfram Mathematica 12), we get that $\phi_{\dot{G}}(\lambda^*) > 0$. See Tables 1 and 2. (In Table 1, $a_1 = 2\sqrt{\sqrt{5}-1}$ and $a_2 = \sqrt{\sqrt{5}-2}$). Then we have

Lemma 4.21. Let \dot{G} be one of the signed graphs of Theorem 2.4 (iii) and (iv) with n vertices. Then $\rho(\dot{G}) < \lambda^*$.

Proof. Let \dot{G} be one of the signed graphs of Theorem 2.4 (iii) and (iv), then we can get that \dot{G} is bipartite and \dot{G} contains an induced subgraph \dot{G}_1 with order $|V(\dot{G}_1)| = n-1$ and $\lambda_1(\dot{G}_1) < \lambda^*$. (For example, if \dot{G} is $[\dot{G}_4^{12}, v_{12}, s, v_{12}, \dot{G}_4^{12}]$, let $\dot{G}_1 = \dot{G} - v_{12}$, then $\lambda_1(\dot{G}_1) = \lambda_1(\dot{G}_4^{12}, v_{12}, s) = \lambda_1(\dot{G}_4^{12}, v_{12}, 3) < \lambda^*$.) Then $\lambda_2(\dot{G}) \leq \lambda_1(\dot{G}_1) < \lambda^*$ by interlacing theorem. Since $\phi_{\dot{G}}(\lambda^*) > 0$ (by Tables 1 and 2), then $\rho(\dot{G}) = \lambda_1(\dot{G}) < \lambda^*$.

Finally we are going to give the proof of Lemma 3.5.

Proof of Lemma 3.5. Let \dot{G} be a signed graph in Lemma 3.5. We can check that \dot{G} is bipartite and \dot{G} contains an induced subgraph \dot{G}_1 with order $|V(\dot{G}_1)| = n - 1$ and $\lambda_1(\dot{G}_1) < \lambda^*$. Then $\lambda_2(\dot{G}) \le \lambda_1(\dot{G}_1) < \lambda^*$ by interlacing theorem. So if $\phi_{\dot{G}}(\lambda^*) > 0$, then $\rho(\dot{G}) < \lambda^*$. Now we just need to prove that $\phi_{\dot{G}}(\lambda^*) > 0$. Obviously, if \dot{G} is an induced subgraph of the signed graphs of Theorem 2.4 (iv), then $\rho(\dot{G}) < \lambda^*$ by Lemma 4.21.

- (1) If $n_1 \geq 3$, by forbidding $Q_{n_1,n_2+1,1}$ (where $n_2 \leq n_1 2$), then $n_2 \geq n_1 1$. Similarly, if $n_4 \geq 3$, then $n_3 \geq n_4 1$. Therefore, if $n_1 \leq 2$ and $n_4 \leq 2$, then $\dot{\mathcal{A}}_1^{n_1,n_2,n_3,n_4} \subset [P_4, v_2, s, v_2, P_4]$; if $n_1 \geq 3$ and $n_4 \geq 3$, then $\dot{\mathcal{A}}_1^{n_1,n_2,n_3,n_4} \subset [T_{a_3,1,a_3-1}, v_{a_3-1}, s, v_{b_3-1}, T_{b_3,1,b_3-1}]$; if $n_1 \leq 2$ and $n_4 \geq 3$, or $n_1 \geq 3$ and $n_4 \leq 2$, then $\dot{\mathcal{A}}_1^{n_1,n_2,n_3,n_4} \subset [T_{a_3,1,a_3-1}, v_{a_3-1}, s, v_2, P_4]$, where $s, a_3, b_3 \geq 3$.
 - (2) By Lemma 4.18 and Eq. (4.4), then

$$\phi_{\dot{\mathcal{A}}_{2}^{n_{1},n_{2},n_{3}}}(x) = x^{3} p_{n_{1}+n_{2}+n_{3}+4} - x^{2} (p_{n_{1}+n_{3}+3} p_{n_{2}} + p_{n_{1}+1} p_{n_{2}+n_{3}+2} - 2 p_{n_{1}} p_{n_{2}+n_{3}+1})$$

$$+ x (p_{n_{1}+1} p_{n_{3}+1} p_{n_{2}} - p_{n_{1}} p_{n_{2}+n_{3}+4} - p_{n_{1}+3} p_{n_{2}+n_{3}+1} - 2 p_{n_{1}} p_{n_{2}} p_{n_{3}})$$

$$+ p_{n_{1}} p_{n_{2}} p_{n_{3}+3} + p_{n_{1}+3} p_{n_{2}} p_{n_{3}}.$$

$$(4.7)$$

If $n_3 \ge n_1 + n_2 + 2$ and $n_2 \le 2$, then $\dot{\mathcal{A}}_2^{n_1, n_2, n_3} \subset [\dot{Q}'_{a_1, a_1}, v_{a_1}, s, v_2, P_4]$ $(a_1 \ge 1)$.

If $n_3 \geq n_1 + n_2 + 2$ and $n_2 \geq 3$, and if $n_1 = 0$ and $n_3 = n_2 + 2$, then $\dot{\mathcal{A}}_2^{0,n_2,n_2+2} \sim (\dot{Q}'_{1,0}, v_1, 2, v_{a_3-1}, T_{a_3,1,a_3-1})$ (where $\rho(\dot{\mathcal{A}}_2^{0,n_2,n_2+2}) < \lambda^*$ by Table 3 (3)); otherwise, $\dot{\mathcal{A}}_2^{n_1,n_2,n_3} \subset [\dot{Q}'_{a_1,a_1}, v_{a_1}, s, v_{a_3-1}, T_{a_3,1,a_3-1}]$, where $a_1 \geq 1$, $a_3 \geq 3$ and $s \geq 3$.

Next we consider that $n_3 \leq n_1 + n_2 + 1$.

Case 1. $n_2 = 1$.

Subcase 1.1. $n_1 \leq 1$. If $n_3 = 0$, by computations, we have $\phi_{\dot{\mathcal{A}}_2^{0,1,0}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_2^{1,1,0}}(\lambda^*) < 0$. Then $n_3 \geq 1$ and $\dot{\mathcal{A}}_2^{n_1,1,n_3} \subset [\dot{Q}'_{1,1},v_1,s]$.

Subcase 1.2. $n_1 \geq 2$. By forbidding $Q_{n_1+1,n_3+2,1}$ (where $n_3 \leq n_1-2$), then $n_3 \geq n_1-1$. By Table 3 (1), we have $\phi_{\dot{\mathcal{A}}_2^{n_1,1,n_1-1}}(\lambda^*) < 0$. Then $n_3 \geq n_1$ and $\dot{\mathcal{A}}_2^{n_1,1,n_3} \subset [\dot{Q}'_{n_1,n_1}, v_{n_1}, s]$.

Case 2. $n_2 = 2$. Then $n_3 \le n_1 + 3$. Furthermore, by forbidding $Q_{2,n_3+1,2}$ (where $n_3 \le 2$), then $n_3 \ge 3$.

Subcase 2.1. $n_1 = 0$. Then $n_3 = 3$. Note that $\rho(\dot{A}_2^{0,2,3}) = 2.057 < \lambda^*$, then $\dot{A}_2^{0,2,3} \sim (\dot{Q}'_{1,0}, v_1, 2, v_2, P_4)$.

Subcase 2.2. $1 \le n_1 \le 3$. If $n_3 \le n_1 + 2$, by computations, we have $\phi_{\dot{\mathcal{A}}_2^{n_1,2,n_3}}(\lambda^*) < 0$. Then $n_3 \ge n_1 + 3$. So $\dot{\mathcal{A}}_2^{n_1,2,n_3} \subset [\dot{Q}'_{n_1,n_1}, v_{n_1}, s, v_2, P_4]$ where $s \ge 3$.

Subcase 2.3. $n_1 \geq 4$. By forbidding $Q_{n_1+1,n_3+2,2}$ (where $n_3 \leq n_1+1$), then $n_3 \geq n_1+2$. If $n_3=n_1+2$, then $\phi_{\dot{\mathcal{A}}_2^{n_1,2,n_1+2}}(\lambda^*)>0$ (by Table 3 (2)) and $\dot{\mathcal{A}}_2^{n_1,2,n_1+2}\sim (\dot{Q}'_{n_1,n_1},v_{n_1},2,v_2,P_4)$. If $n_3 \geq n_1+3$, then $\dot{\mathcal{A}}_2^{n_1,2,n_3}\subset [\dot{Q}'_{n_1,n_1},v_{n_1},s,v_2,P_4]$ where $s\geq 3$.

Case 3. $n_2 \geq 3$. By forbidding the graph $Q_{a,b,c} \notin \mathcal{G}^{\lambda^*}$, we obtain that if $n_2 = 3$ and $1 \leq n_1 \leq 3$, or $n_2 = 4$ and $n_1 = 2$, then $n_3 \geq n_1 + n_2$; otherwise, $n_3 \geq n_1 + n_2 + 1$.

Subcase 3.1. $n_2 = 3$ and $n_1 = 1$. Then $4 \le n_3 \le 5$. By computations, we have $\phi_{\dot{\mathcal{A}}_{2}^{1,3,4}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_{2}^{1,3,5}}(\lambda^*) < 0$.

Subcase 3.2. $n_2 = 3$ and $n_1 = 2$. Then $5 \le n_3 \le 6$. By computations, we have $\phi_{\dot{\mathcal{A}}_2^{2,3,5}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_2^{2,3,6}}(\lambda^*) > 0$. So $\dot{\mathcal{A}}_2^{2,3,6} \sim (\dot{Q}'_{2,2}, v_2, 2, v_2, T_{3,1,2})$.

Subcase 3.3. $n_2 = 3$ and $n_1 = 3$. Then $6 \le n_3 \le 7$. By computations, we have $\phi_{\dot{\mathcal{A}}_2^{3,3,6}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_2^{3,3,7}}(\lambda^*) > 0$. So $\dot{\mathcal{A}}_2^{3,3,7} \sim (\dot{Q}_{3,3}', v_3, 2, v_2, T_{3,1,2})$.

Subcase 3.4. $n_2 = 4$ and $n_1 = 2$. Then $6 \le n_3 \le 7$. By computations, we have $\phi_{\dot{\mathcal{A}}_2^{2,4,6}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_2^{2,4,7}}(\lambda^*) < 0$.

Subcase 3.5. $n_2 \ge 3$ and $n_3 = n_1 + n_2 + 1$. Let $n_1 = n_2 + k$. Then

$$\phi_{\dot{\mathcal{A}}_2^{n_2+k,n_2,2n_2+k+1}}(\lambda^*) = 2^{n_2}(\sqrt{5}+1)^{-k-2n_2}f(k,n_2),$$

where $f(k, n_2) = 2^k (\sqrt{5} + 1)^{n_2+1} ((\frac{\sqrt{5}+3}{\sqrt{5}+1})(\frac{\sqrt{5}+1}{2})^k + 2(\frac{2}{\sqrt{5}+1})^{n_2+1} - 1)$. If $k \le -2$, then $f(k, n_2) < 0$ and $\phi_{\dot{\mathcal{A}}_2^{n_2+k, n_2, 2n_2+k+1}}(\lambda^*) < 0$ as $(\frac{\sqrt{5}+3}{\sqrt{5}+1})(\frac{\sqrt{5}+1}{2})^k \le (\frac{\sqrt{5}+3}{\sqrt{5}+1})(\frac{\sqrt{5}+1}{2})^{-2} \approx 0.618$ and

 $2(\frac{2}{\sqrt{5}+1})^{n_2+1} \leq 2(\frac{2}{\sqrt{5}+1})^4 \approx 0.2918. \text{ If } k \geq -1, \text{ then } f(k,n_2) > 0 \text{ and } \phi_{\dot{\mathcal{A}}_2^{n_2+k},n_2,2n_2+k+1}(\lambda^*) > 0 \text{ as } (\frac{\sqrt{5}+3}{\sqrt{5}+1})(\frac{\sqrt{5}+1}{2})^k \geq (\frac{\sqrt{5}+3}{\sqrt{5}+1})(\frac{\sqrt{5}+1}{2})^{-1} = 1. \text{ So, } \dot{\mathcal{A}}_2^{n_1,n_2,n_3} \subset (\dot{Q}'_{n_1,n_1},v_{n_1},2,v_{n_2-1},T_{n_2,1,n_2-1}), \text{ where } n_1 \geq n_2 - 1. \text{ We complete the proof of (2).}$

(3) Let $n_1 = n_2 + k$ where $k \ge 0$. By forbidding $Q_{2,b,2}$ (where $b \le 3$), we have $n_3 \ge 3$.

Case 1. $n_1 = n_2 = 0$. By computations, we have $\phi_{\dot{\mathcal{A}}_3^{0,0,3}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{0,0,4}}(\lambda^*) > 0$. Then $n_3 \geq 4$. So $\dot{\mathcal{A}}_3^{0,0,4} \sim (\dot{Q}'_{1,0}, v_1, 2, v_1, \dot{Q}'_{1,0}) \subset (\dot{Q}'_{1,1}, v_1, 2, v_1, \dot{Q}'_{1,0})$ (if $n_3 = 4$) or $\dot{\mathcal{A}}_3^{0,0,n_3} \subset [\dot{Q}'_{1,0}, v_1, s, v_1, \dot{Q}'_{1,0}] \subset [\dot{Q}'_{1,1}, v_1, s, v_1, \dot{Q}'_{1,1}]$ where $s \geq 3$ (if $n_3 \geq 5$).

Case 2. $n_1 = 1$.

Subcase 2.1. $n_2 = 0$. By computations, we have $\phi_{\dot{\mathcal{A}}_3^{1,0,3}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{1,0,4}}(\lambda^*) > 0$. Then $n_3 \geq 4$. So $\dot{\mathcal{A}}_3^{1,0,4} \sim (\dot{Q}'_{1,1}, v_1, 2, v_1, \dot{Q}'_{1,0})$ (if $n_3 = 4$) or $\dot{\mathcal{A}}_3^{1,0,n_3} \subset [\dot{Q}'_{1,1}, v_1, s, v_1, \dot{Q}'_{1,0}] \subset [\dot{Q}'_{1,1}, v_1, s, v_1, \dot{Q}'_{1,1}]$ where $s \geq 3$ (if $n_3 \geq 5$).

Subcase 2.2. $n_2 = 1$. By computations, we have $\phi_{\dot{\mathcal{A}}_3^{1,1,3}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{1,1,4}}(\lambda^*) < 0$. Then $n_3 \geq 5$. So $\dot{\mathcal{A}}_3^{1,1,n_3} \subset [\dot{Q}_{1,1}', v_1, s, v_1, \dot{Q}_{1,1}']$ where $s \geq 3$.

Case 3. $n_1 = 2$.

Subcase 3.1. $n_2 = 0$. By computations, we have $\phi_{\dot{\mathcal{A}}_3^{2,0,3}}(\lambda^*) < 0$, $\phi_{\dot{\mathcal{A}}_3^{2,0,4}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{2,0,5}}(\lambda^*) > 0$. Then $n_3 \geq 5$. So $\dot{\mathcal{A}}_3^{2,0,5} \sim (\dot{Q}'_{2,2}, v_2, 2, v_1, \dot{Q}'_{1,0})$ (if $n_3 = 5$) or $\dot{\mathcal{A}}_3^{2,0,n_3} \subset [\dot{Q}'_{2,2}, v_2, s, v_1, \dot{Q}'_{1,0}] \subset [\dot{Q}'_{2,2}, v_2, s, v_1, \dot{Q}'_{1,1}]$ where $s \geq 3$ (if $n_3 \geq 6$).

Subcase 3.2. $1 \le n_2 \le 2$. By computations, if $n_3 \le n_1 + n_2 + 2$, then $\phi_{\dot{\mathcal{A}}_3^{2,n_2,n_3}}(\lambda^*) < 0$. Then $n_3 \ge n_1 + n_2 + 3$. So $\dot{\mathcal{A}}_3^{2,n_2,n_3} \subset [\dot{Q}'_{2,2}, v_2, s, v_{n_2}, \dot{Q}'_{n_2,n_2}]$ where $s \ge 3$.

Case 4. $n_1 \ge 3$.

Subcase 4.1. $n_2 = 0$. By forbidding $Q_{n_1+1,n_3+2,2}$ (where $n_3 \leq n_1 + 1$), we have $n_3 \geq n_1 + 2$. Since $\phi_{\dot{\mathcal{A}}_3^{n_1,0,n_1+2}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{n_1,0,n_1+3}}(\lambda^*) > 0$ (by Table 3 (4) and (5)), then $n_3 \geq n_1 + 3$. So $\dot{\mathcal{A}}_3^{n_1,0,n_1+3} \sim (\dot{Q}'_{n_1,n_1}, v_{n_1}, 2, v_1, \dot{Q}'_{1,0})$ (if $n_3 = n_1 + 3$) or $\dot{\mathcal{A}}_3^{n_1,0,n_3} \subset [\dot{Q}'_{n_1,n_1}, v_{n_1}, s, v_1, \dot{Q}'_{1,0}] \subset [\dot{Q}'_{n_1,n_1}, v_{n_1}, s, v_1, \dot{Q}'_{1,1}]$ where $s \geq 3$ (if $n_3 \geq n_1 + 4$).

Subcase 4.2. $n_2 = 1$. By forbidding $Q_{n_1+1,n_3+3,2}$ (where $n_3 \leq n_1 + 2$), we have $n_3 \geq n_1 + 3$. Since $\phi_{\dot{\mathcal{A}}_3^{n_1,1,n_1+3}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{n_1,1,n_1+4}}(\lambda^*) > 0$ (by Table 3 (6) and (7)), then $n_3 \geq n_1 + 4$. So $\dot{\mathcal{A}}_3^{n_1,1,n_3} \subset [\dot{Q}'_{n_1,n_1}, v_{n_1}, s, v_1, \dot{Q}'_{1,1}]$ where $s \geq 3$.

Subcase 4.3. $n_2 \geq 2$. If $n_1 = 3$ and $n_2 = 2$, by forbidding $Q_{4,n_3+3,3}$ (where $n_3 \leq 4$), we have $n_3 \geq 5$. By computations, we have $\phi_{\dot{\mathcal{A}}_3^{3,2,5}}(\lambda^*) < 0, \phi_{\dot{\mathcal{A}}_3^{3,2,6}}(\lambda^*) < 0$ and $\phi_{\dot{\mathcal{A}}_3^{3,2,7}}(\lambda^*) > 0$. Then $n_3 \geq 7$. So $\dot{\mathcal{A}}_3^{3,2,7} \sim (\dot{Q}_{3,3}', v_3, 2, v_2, \dot{Q}_{2,2}')$ (if $n_3 = 7$) or $\dot{\mathcal{A}}_3^{3,2,n_3} \subset [\dot{Q}_{3,3}', v_3, s, v_2, \dot{Q}_{2,2}']$ where $s \geq 3$ (if $n_3 \geq 8$). Otherwise, either $n_1 \geq 4$ and $n_2 = 2$, or $n_1 \geq 3$ and $n_2 \geq 3$. By forbidding Q_{n_1+1,n_3+3,n_2+1} (where $n_3 \leq n_1+n_2$), we have $n_3 \geq n_1+n_2+1$. If $n_3 = n_1+n_2+1$,

then $\phi_{\dot{\mathcal{A}}_3^{n_1,n_2,n_1+n_2+1}}(\lambda^*) = -2\sqrt{2}(\sqrt{5}+1)^{-n_1-n_2-\frac{1}{2}}((\sqrt{5}+1)^{n_1}2^{n_2}+2^{n_1}(\sqrt{5}+1)^{n_2}+(3-\sqrt{5})2^{n_1+n_2}) < 0$. Therefore, $n_3 \ge n_1 + n_2 + 2$.

If $n_3 = n_1 + n_2 + 2$, then $\phi_{\dot{\mathcal{A}}_3^{n_1, n_2, n_1 + n_2 + 2}}(\lambda^*) = (\sqrt{5} + 1)^{-k - 2n_2 - 1} f(n_2, k)$, where $f(n_2, k) = (\sqrt{5} + 1)^{k + 2n_2 + 1} + (\sqrt{5} - 3)2^{2n_2 + k + 2} - 2^{n_2 + 2}(\sqrt{5} + 1)^{n_2}((\sqrt{5} + 1)^k + 2^k)$. Since

$$f(n_2, k) > (\sqrt{5} + 1)^{k+2n_2+1} - (\sqrt{5} + 1)^{n_2} 2^{k+n_2+2} - 2^{n_2+2} (\sqrt{5} + 1)^{k+n_2} - 2^{k+2n_2+2}$$
$$> (\sqrt{5} + 1)^{n_2} ((\sqrt{5} + 1)^{k+n_2+1} - ((\sqrt{5} + 1)^k + 2^{k+1}) 2^{n_2+2}) = (\sqrt{5} + 1)^{n_2} g(n_2, k),$$

then $\phi_{\dot{A}_{3}^{n_{1},n_{2},n_{1}+n_{2}+2}}(\lambda^{*}) > (\sqrt{5}+1)^{-k-n_{2}-1}g(n_{2},k)$. If k=0, then $n_{1}=n_{2}\geq 3$ and $g(n_{2},0)=(\sqrt{5}+1)^{n_{2}+1}-3\cdot 2^{n_{2}+2}>0$. If $k\geq 1$, then $g(n_{2},k)>(\sqrt{5}+1)^{k}((\sqrt{5}+1)^{n_{2}+1}-2^{n_{2}+3})>0$ for $n_{2}\geq 2$. So, $\phi_{\dot{A}_{2}^{n_{1},n_{2},n_{1}+n_{2}+2}}(\lambda^{*})>0$.

Hence, $\dot{\mathcal{A}}_{3}^{n_{1},n_{2},n_{1}+n_{2}+2} \sim (\dot{Q}'_{n_{1},n_{1}},v_{n_{1}},2,v_{n_{2}},\dot{Q}'_{n_{2},n_{2}})$ (if $n_{3}=n_{1}+n_{2}+2$) or $\dot{\mathcal{A}}_{3}^{n_{1},n_{2},n_{3}} \subset [\dot{Q}'_{n_{1},n_{1}},v_{n_{1}},s,v_{n_{2}},\dot{Q}'_{n_{2},n_{2}}]$ where $s \geq 3$ (if $n_{3} \geq n_{1}+n_{2}+3$). We complete the proof of (3).

The proofs of (4) - (11) are similar, and all values $\phi_{\mathcal{A}_i^{n_1,n_2}}(\lambda^*)$ ($i=4,\ldots,13$) are presented in Table 3, where $a_3=\sqrt{\sqrt{5}-1}, a_4=\sqrt{17\sqrt{5}+22}, a_5=\sqrt{73\sqrt{5}+151},$ $a_6=\sqrt{17\sqrt{5}-22}, \ a_7=\sqrt{5\sqrt{5}-11}, a_8=\sqrt{73\sqrt{5}-151}, a_9=\sqrt{5\sqrt{5}-11}, a_{10}=\sqrt{185\sqrt{5}-409}, a_{11}=\frac{2}{\sqrt{13\sqrt{5}+29}}, a_{12}=\frac{31}{\sqrt{337\sqrt{5}+751}}.$

- (12) If $n_1 \geq 3$, by forbidding Q_{1,n_2+1,n_1} (where $n_2 \leq n_1 2$), then $n_2 \geq n_1 1$. So $\dot{\mathcal{A}}_{14}^{n_1,n_2} \subset [P_4, v_2, s]$ (if $n_1 \leq 2$) or $\dot{\mathcal{A}}_{14}^{n_1,n_2} \subset [T_{n_1,1,n_1-1}, v_{n_1-1}, s]$ (if $n_1 \geq 3$), where $s \geq 3$.
- (13) By computations, we have $\phi_{\dot{A}_{16}^{n_1}}(\lambda^*) < 0$ and $\phi_{\dot{A}_{17}^{n_1}}(\lambda^*) < 0$ for $n_1 \leq 8$ and $\phi_{\dot{A}_{15}^{n_1}}(\lambda^*) < 0$, $\phi_{\dot{A}_{18}^{n_1}}(\lambda^*) < 0$ and $\phi_{\dot{A}_{19}^{n_1}}(\lambda^*) < 0$ for $n_1 \leq 10$. Therefore, $\dot{A}_{15}^{n_1}$ is an induced subgraph of $[\dot{G}_5^{10}, v_{10}, s, v_{10}, \dot{G}_5^{10}]$, $\dot{A}_i^{n_1}$ (i = 16, 17) is an induced subgraph of $[\dot{G}_2^{12}, v_9, s, v_{10}, \dot{G}_5^{10}]$, $\dot{A}_i^{n_1}$ (i = 18, 19) is an induced subgraph of $[\dot{G}_4^{12}, v_{12}, s, v_{10}, \dot{G}_5^{10}]$.

Table 1: $\phi_{\dot{G}}(x)$ and $\phi_{\dot{G}}(\lambda^*)$ where \dot{G} belongs to Theorem 2.4 (iii) and (iv)

$$\begin{split} \phi_{Q_{n_1,n_2}}(x) &= xp_{n_1+n_2+3} - p_{n_1+2}p_{n_2} - p_{n_1}p_{n_2+2} + 2p_{n_1}p_{n_2} \\ \phi_{Q_{n_1,n_2}}(x) &= x\phi_{Q_{n_1,n_2}}(x) - p_{n_1+n_2+3} \\ \phi_{C_k^1,\frac{k}{2}+1}(x) &= x(x(p_k-p_{k-2}+2)-p_{k-1}) - \phi_{T_{\frac{k}{2}-1,1,\frac{k}{2}-1}}(x) \\ \phi_{C_k^1,\frac{k}{2}+1}(\lambda^*) &= (\sqrt{5}+3)2^{\frac{k}{2}}(\sqrt{5}+1)^{1-\frac{k}{2}} > 0 \\ \phi_{C_4(n_1,1,n_3,1]}(x) &= x\phi_{Q_{n_1,n_3}}(x) - \phi_{T_{n_1+1,1,n_3+1}}(x) \\ \phi_{C_4(n_1,1,n_3,1]}(\lambda^*) &= 2^{\frac{k}{2}(n_1+n_3+2)}(\sqrt{5}+1)^{-\frac{n_1}{2}-\frac{n_2}{2}+1} > 0 \\ \phi_{C_6^{n_1,n_2}}(x) &= x\phi_{T_{n_2,1,n_1+3}}(x) - p_{n_1+n_2+4} - \phi_{T_{n_2,1,2}}(x)p_{n_1} + 2p_{n_1}p_{n_2} \\ \phi_{C_6^{n_1,n_2}}(\lambda^*) &= 2^{\frac{k}{2}(n_1+n_2+2)}(\sqrt{5}+1)^{-\frac{n_1}{2}-\frac{n_2}{2}+1} > 0 \\ \phi_{G_6^{n_0}}(x) &= x^2q_k - (q_k+2xp_{k-1}) + 2x^2 + 2xp_{k-3} \\ \phi_{G_6^{n_0}}(x) &= x^2(q_k-2xp_{k-1}) + 2x^2 + 2xp_{k-3} \\ \phi_{G_6^{n_0}}(x) &= x(2n_1-\frac{k}{2}) > 0 \quad (k \text{ is even}) \\ \phi_{S_1^{n_1}}(x) &= xp_{n-8}\phi_{S_1^{n_1}}(x) - q_6p_{n-8} - p_{n-9}\phi_{S_1^{n_1}}(x) \\ \phi_{S_1^{n_1}}(x) &= xp_{n-8}\phi_{S_1^{n_1}}(x) - q_6p_{n-8} - p_{n-12}\phi_{S_2^{n_2}}(x) \\ \phi_{S_2^{n_1}}(x) &= xp_{n-11}\phi_{S_2^{n_1}}(x) - p_{n-11}\phi_{S_2^{n_1}}(x) - p_{n-12}\phi_{S_2^{n_1}}(x) \\ \phi_{S_2^{n_1}}(x) &= xp_{n-1}\phi_{S_2^{n_1}}(x) - p_{n-11}\phi_{S_2^{n_1}}(x) - p_{n-12}\phi_{S_2^{n_1}}(x) \\ \phi_{S_2^{n_1}}(x) &= xp_{n-11}\phi_{S_2^{n_1}}(x) - p_{n-12}\phi_{S_2^{n_1}}(x) \\ \phi_{S_2^{n_1}}(x) &= 2^{\frac{n}{2}+2}(\sqrt{5}+1)^{\frac{n}{2}-\frac{n}{2}} > 0 \\ \phi_{C_3^{n_1},v_{2,3},v_{n_1},\dot{Q}_{n_1,n_1}}(\lambda^*) &= \frac{3(194\sqrt{5}-7142)(\sqrt{5}+1)^{n_1+1}(1292\sqrt{5}-2889)2^{n_1+3}}{(\sqrt{5}+1)^{n_1+1}} > 0 \\ \phi_{[C_3^{n_1},v_{2,3},v_{n_1},\dot{Q}_{n_1,n_1}]}(\lambda^*) &= \frac{2(\sqrt{5}+1)^{n_1+1}}{3(305\sqrt{5}+682)^{1/2}(\sqrt{5}+1)^{n_1+1}} > 0 \\ \phi_{[C_3^{n_1},v_{2,3},v_{n_1},\dot{Q}_{n_1,n_1}]}(\lambda^*) &= \frac{2(\sqrt{5}+1)^{n_1+1}}{3(35\sqrt{5}+682)^{1/2}(\sqrt{5}+1)^{n_1+1}} > 0 \\ \phi_{[C_{3}^{n_1},v_{3},v_{n_1},\dot{Q}_{n_1,n_1}]}(\lambda^*) &= \frac{2(\sqrt{5}+1)^{n_1+1}-(\sqrt{5}-2)2^{n_1+3}}{(\sqrt{5}+1)^{n_1+1}} > 0 \\ \phi_{[C_3^{n_1},v_{3,3},v_{n_1},\dot{Q}_{n_1,n_1}]}(\lambda^*) &= \frac{2(\sqrt{5}+1)^{n_1+1}-(\sqrt{5}-2)2^{n_1+3}}{(\sqrt{5}+1)^{n_1+1}} > 0 \\ \phi_{[C_3^{n_1},v_{3,3},v_{n_1},\dot{Q}_{n_1,n$$

Table 2: $\phi_{\dot{G}}(\lambda^*)$ where \dot{G} belongs to Theorem 2.4 (iv)

$\phi_{[\dot{G}_{4}^{12}, v_{12}, 3, v_{12}, \dot{G}_{4}^{12}]}(\lambda^*) \approx 0.0007$	$\phi_{[\dot{G}_{2}^{9}, v_{9}, 3, v_{9}, \dot{G}_{2}^{9}]}(\lambda^{*}) \approx 0.02$	$\phi_{[\dot{G}_5^{10}, v_{10}, 3, v_2, P_4]}(\lambda^*) \approx 0.03$
$\phi_{[\dot{G}_{4}^{12}, v_{12}, 3, v_{9}, \dot{G}_{2}^{9}]}(\lambda^{*}) \approx 0.004$	$\phi_{[\dot{G}_{2}^{9},v_{9},3,v_{10},\dot{G}_{5}^{10}]}(\lambda^{*}) \approx 0.015$	$\phi_{[P_4,v_2,3,v_2,P_4]}(\lambda^*) \approx 0.05$
$\phi_{[\dot{G}_{4}^{12}, v_{12}, 3, v_{10}, \dot{G}_{5}^{10}]}(\lambda^*) \approx 0.003$	$\phi_{[\dot{G}_{2}^{9}, v_{9}, 3, v_{2}, P_{4}]}(\lambda^{*}) \approx 0.033$	
$\phi_{[\dot{G}_{4}^{12}, v_{12}, 3, v_{2}, P_{4}]}(\lambda^{*}) \approx 0.006$	$\phi_{[\dot{G}_5^{10}, v_{10}, 3, v_{10}, \dot{G}_5^{10}]}(\lambda^*) \approx 0.01$	

Table 3: $\phi_{\dot{G}}(\lambda^*)$ where \dot{G} belongs to Lemma 3.5

$$\begin{array}{l} (1) \ \phi_{\mathcal{A}_{1}^{n_{1},1,n_{1}-1}}(\lambda^{*}) = -(\sqrt{5}-1)^{\frac{1}{2}}2^{n_{1}+\frac{1}{2}}\left(\sqrt{5}+1\right)^{-n_{1}} < 0 \\ (2) \ \phi_{\mathcal{A}_{2}^{n_{1},2,n_{1}+2}}(\lambda^{*}) = 2\sqrt{2}(\sqrt{5}+1)^{-n_{1}-\frac{3}{2}}((\sqrt{5}+1)^{n_{1}}-\sqrt{5}\cdot2^{n_{1}+1}) > 0 \ (n_{1}\geq 4) \\ (3) \ \phi_{\mathcal{A}_{2}^{0,n_{2},n_{2}+2}}(\lambda^{*}) = \frac{(\sqrt{5}-1)^{\frac{1}{2}}2^{n_{2}+\frac{3}{2}}}{(\sqrt{5}+1)^{n_{2}}} + 3(\sqrt{5}+2)^{\frac{1}{2}}-(2(\sqrt{5}+1))^{\frac{1}{2}} > 0 \\ (4) \ \phi_{\mathcal{A}_{3}^{n_{1},0,n_{1}+2}}(\lambda^{*}) = -(2\sqrt{5}+3)2^{n_{1}+5}(\sqrt{5}+1)^{-n_{1}-4} - \sqrt{5}+2 < 0 \ (n_{1}\geq 3) \\ (5) \ \phi_{\mathcal{A}_{3}^{n_{1},0,n_{1}+3}}(\lambda^{*}) = (\sqrt{5}+1)^{-n_{1}-\frac{3}{2}}(2\sqrt{10}(\sqrt{5}+1)^{n_{1}}+(\sqrt{5}-4)2^{n_{1}+\frac{5}{2}}) > 0 \\ (6) \ \phi_{\mathcal{A}_{3}^{n_{1},1,n_{1}+3}}(\lambda^{*}) = (5\sqrt{5}-13)(\frac{2}{\sqrt{5}+1})^{n_{1}} + \sqrt{5}-2 < 0 \\ (7) \ \phi_{\mathcal{A}_{3}^{n_{1},1,n_{1}+4}}(\lambda^{*}) = (\sqrt{5}-7)(\frac{1}{2}(\sqrt{5}+1))^{-n_{1}-\frac{5}{2}} + 3\sqrt{\sqrt{5}-2} > 0 \\ (8) \ \phi_{\mathcal{A}_{4}^{n_{1},n_{2}}}(\lambda^{*}) = \frac{a_{3}\cdot2^{n_{1}}(\sqrt{5}+1)^{n_{2}+a_{4}}\cdot2^{n_{1}+n_{2}+\frac{3}{2}}-a_{5}(\sqrt{5}+1)^{n_{1}}2^{n_{2}}}{2^{\frac{1}{2}(n_{1}+n_{2}+2)}(\sqrt{5}+1)^{\frac{1}{2}(n_{1}+n_{2}-1)}} \\ (9) \ \phi_{\mathcal{A}_{4}^{n_{1},n_{2}}}(\lambda^{*}) = \frac{((4\sqrt{5}+1)^{n_{1}+1}+(\sqrt{5}+1)^{n_{1}+1}+\frac{3}{2}}{(\sqrt{5}-1)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{1}+n_{2}-1)}}} \\ (10) \ \phi_{\mathcal{A}_{6}^{n_{1},n_{2}}}(\lambda^{*}) = \frac{((\sqrt{5}-1)2^{n_{1}+1}}(\sqrt{5}+1)^{n_{1}+1}+3)((\sqrt{5}-1)2^{n_{1}-1}(\sqrt{5}+1)^{n_{1}})2^{n_{2}+2}}}{(\sqrt{5}-1)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{1}+n_{2}+3)}}} \\ (11) \ \phi_{\mathcal{A}_{7}^{n_{1},n_{2}}}}(\lambda^{*}) = \frac{3(\sqrt{5}-1)^{n_{1}+1}(\sqrt{5}+1)^{n_{2}+3}((\sqrt{5}-1)2^{n_{1}+n_{2}+3})}{(\sqrt{5}-3)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{1}+n_{2}+3)}}} \\ (12) \ \phi_{\mathcal{A}_{8}^{n_{1},n_{2}}}}(\lambda^{*}) = \frac{a_{3}\cdot2^{n_{1}}(\sqrt{5}+1)^{n_{2}+3}(\sqrt{5}-1)^{n_{1}+1}(\sqrt{5}+1)^{n_{2}+3}+2^{n_{2}+3}}}{(\sqrt{5}-3)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{1}+n_{2}+3)}}} \\ (13) \ \phi_{\mathcal{A}_{9}^{n_{1},n_{2}}}}(\lambda^{*}) = \frac{a_{3}\cdot2^{n_{1}}(\sqrt{5}+1)^{n_{2}+3}(\sqrt{5}-2)^{n_{1}+1}(\sqrt{5}+1)^{n_{2}+3}+2^{n_{2}+3}}}{(\sqrt{5}-3)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{1}+n_{2}+3)}}} \\ (14) \ \phi_{\mathcal{A}_{10}^{n_{1},n_{2}}}(\lambda^{*}) = \frac{a_{3}\cdot2^{n_{1}}(\sqrt{5}+1)^{n_{2}}(\sqrt{5}-2)(\sqrt{5}+1)^{\frac{1}{2}(n_{1}+n_{2}+3)}}}{(\sqrt{5}-3)(2(\sqrt{5}+1))^{\frac{1}{2}(n_{$$