

Emergent space-time supersymmetry at disorder quantum critical point

Xue-Jia Yu,¹ Peng-Lu Zhao,^{2,3,*} Shao-Kai Jian,^{4,†} and Zhiming Pan^{5,‡}

¹*International Center for Quantum Materials, School of Physics, Peking University, Beijing 100871, China*

²*Shenzhen Institute for Quantum Science and Engineering and Department of Physics, Southern University of Science and Technology, Shenzhen*

³*Shenzhen Key Laboratory of Quantum Science and Engineering, Shenzhen 518055, China*

⁴*Department of Physics, Brandeis University, Waltham, Massachusetts 02453, USA*

⁵*Institute for Theoretical Sciences, WestLake University, Hangzhou 310024, China*

(Dated: March 4, 2022)

We study the effect of disorder on the spacetime supersymmetry that is proposed to emerge at the quantum critical point of pair density wave transition in (2+1)D Dirac semimetals and (3+1)D Weyl semimetals. In the (2+1)D Dirac semimetal, we consider three types of disorder, including random scalar potential, random vector potential and random mass potential, while the random mass disorder is absent in the (3+1)D Weyl semimetal. Via a systematic renormalization group analysis, we find that any type of weak random disorder is irrelevant due to the couplings between the disorder potential and the Yukawa vertex. The emergent supersymmetry is thus stable against weak random potentials. Our work will pave the way for exploration supersymmetry in realistic condensed matter systems.

I. INTRODUCTION

About five decades ago, the spacetime supersymmetry (SUSY) was proposed as a possible way of solving the hierarchy problem of the standard model [1–4] and the cosmological constant problem[5]. Later, some supersymmetric theories have been studied as toy models to understand strong coupling physics rigorously [6, 7]. Due to these attractive features, SUSY has been studied intensively in past fifty years and there is some expectation before that SUSY may be revealed in the large hadron collider (LHC). Unfortunately, the recent experiments at the LHC have found no evidence of SUSY and/or its spontaneous breaking in particle physics.

Three-dimensional (3D) Weyl fermions [8–10] in non-centrosymmetric materials [11–15] provide an opportunity to test and investigate important concepts developed in the context of high-energy physics in realistic condensed matter systems. It has been suggested that SUSY can emerge in the low-energy limit of a number of non-supersymmetric models [16–24]. In particular, SUSY is proposed to emerge at quantum critical points (QCPs) in Bose-Fermi lattice models [25, 26], in the (2+1)D surface states of topological insulators [27–31], as well as at multicritical points in some low-dimensional systems [32–34]. Moreover, an interesting recent suggestion [35] is that SUSY can be realized at certain pair-density-wave (PDW) superconducting quantum critical points of ideal Weyl semimetals [36, 37](WSMs).

The realization of SUSY at QCPs relies crucially on the fact that the infrared fixed point is stable against small perturbations. In particular, for the emergent SUSY to

be realized, it must be robust when the fermions are subject to small perturbations from quenched disorder and other dissipation effects. Here, we are particularly interested in the impact of quenched disorder on the emergent SUSY, because disorder unavoidably exists in all realistic materials. It is well known that disorder plays an essential role in condensed matter systems [38–45] and may lead to a plenty of prominent phenomena, such as Anderson localization and metal-insulator transition. In graphene-like Dirac semimetals (DSMs), depending on the specific type, disorder can either enhance or reduce the effective Coulomb interaction strength [46–52], which in turn drastically modifies the phase diagram obtained in the clean limit [46–52]. Moreover, disorder may have a significant impact on the low-temperature properties of various Dirac or Weyl semimetals, such as the conductivity of graphene [47, 53, 54], the optical conductivity of WSMs [55], and the low-energy spectral, thermodynamic, and transport behaviors of *d*-wave cuprate superconductors [43, 56–59]. Disorder also plays a vital role in quantum Hall systems [60–63] and topological insulators [27, 28].

In this paper, we investigate the stability of emergent SUSY against the disorder scattering. We focus on the disorder-induced unusual renormalization of the fermion velocity [49, 59, 64], and examine whether such a renormalization effect causes a substantial difference between the velocities of fermions and bosons at low energies, and ruins the emergent SUSY. Based on this analysis, we are able to identify the influence of non-magnetic disorder on the *particular* fixed point that is argued to display an emergent SUSY at the QCP of pair density wave (PDW) transition in (3+1)D WSMs and (2+1)D DSMs [35]. In the case of (2+1)D DSMs, we consider three types of disorder, including random scalar potential (RSP), random vector potential (RVP), and random mass (RM). In (2+1)D, our systematic RG analysis reveals that weak RVP, RMP and RSP are irrelevant at the QCPs where

* zhaoplu@gmail.com

† skjian@brandeis.edu

‡ Corresponding author: panzhiming@westlake.edu.cn




FIG. 1. Schematic RG flow for (3+1)D lattice model [35] with two Weyl fermions as a function the Hubbard attraction U . In the clean system, the $\mathcal{N} = 2$ space-time SUSY in (3+1) D emerges in low energy and long distance at the PDW critical point $U = U_c$. Moreover, the emergent SUSY is stable for weak random quench disorders.

fermion velocity and boson velocity flow into same value under renormalization, which certainly does not breaks the emergent SUSY. In (3+1)D WSMs, the disorder potential becomes more irrelevant and the effective SUSY is robust against weak disorder. The schematic RG flow diagram for emergent SUSY is shown in Fig.1

This paper is structured as follows. In Sec. II, we present the effective model for (2+1)D disordered DSMs and perform the RG calculations. In Sec. III, the same analysis is carried out in (3+1)D WSMs. We briefly summarize the results of this work in Sec. IV. Further RG details for our calculations are provided in Appendix.

II. (2+1)D DIRAC SEMIMETALS

As demonstrated in Ref. 35, a spactime SUSY could emerge in the low-energy region at the PDW QCP of (2+1)D DSMs only when the number of massless Dirac fermions is $N_f = 2$. In this case, the low-energy effective action at the PDW criticality in a clean system is given by

$$S = S_f + S_b + S_I, \quad (1)$$

$$S_f = \int d^2x d\tau \sum_{n=\pm} \psi_n^\dagger \left[\partial_\tau + iv_f \sum_{j=1}^2 \gamma_j \partial_j \right] \psi_n, \quad (2)$$

$$S_b = \int d^2x d\tau \left\{ \sum_{n=\pm} \left[|\partial_\tau \phi_n|^2 + v_b^2 \sum_{j=1}^2 |\partial_j \phi_n|^2 \right. \right. \quad (3)$$

$$\left. \left. + r|\phi_n|^2 + u|\phi_n|^4 \right] + u_{+-} |\phi_1|^2 |\phi_2|^2 \right\},$$

$$S_I = \int d^2x d\tau g \sum_{n=\pm} \left[\phi_n \psi_n^T \sigma_y \psi_n + h.c. \right], \quad (4)$$

where $\gamma_j = (\sigma_x, \sigma_y)$. $\sigma_\alpha, \alpha = x, y$ is Pauli matrix with spin indices. S_f corresponds to the action for two non-interacting two-component Dirac fermions ψ_\pm at two

Dirac points Q_\pm [65, 66], with quartic and higher order self-coupling terms being irrelevant [25] at low energies. S_b describes the quantum fluctuation and the self-coupling of the PDW order parameter ϕ_n near the QCP, where ϕ_\pm is the superconducting order with momentum $2Q_\pm$, respectively. Terms with higher powers of ϕ_n are all irrelevant, whereas $\phi_n^* \partial_\tau \phi_n$ is excluded by particle-hole symmetry [35]. S_I represents the Yukawa coupling between Dirac fermions and bosons. The terms of the form $\phi_-^* \psi_+ \sigma_y \psi_+$ and $\phi_-^* \psi_+ \sigma_y \psi_-$ are not allowed because they do not satisfy momentum conservation [25]. Therefore, the effective action given above is of the most general form. It has been shown through renormalization group analysis that an emergent spacetime SUSY occurs at the low energy limit. A necessary condition for the emergent SUSY is that velocities of fermions and bosons flow to the same value under RG, which renders the emergent Lorentz symmetry. It was claimed that such an emergent Lorentz symmetry can be naturally realized in a number of correlated electron systems [25–31, 35].




FIG. 2. Feynman diagrams for all the relevant one-loop diagrams that survive within replica limit. Here, the solid line represents the free fermion propagator, the wavy line the free boson propagator, and the dashed line the disorder.

The aim of the present work is to examine whether the emergent SUSY is robust against disorder scattering. For this purpose, we now introduce a direct fermion-disorder coupling term to the system via the standard form, also see Appendix A [43, 46–49, 58, 60, 67],

$$S_{\text{dis}} = \int d^2x d\tau \sum_{n=\pm} \psi_n^\dagger \left(\sum_{\Gamma} V_{\Gamma}(\mathbf{x}) \Gamma \right) \psi_n, \quad (5)$$

where $V_{\Gamma}(\mathbf{x})$ stands for the random potential and Γ labels

the type of the disorder potential. We assume $V_\Gamma(\mathbf{x})$ to be a quenched, Gaussian white noise potential characterized by the following identities:

$$\langle V_\Gamma(\mathbf{x}) \rangle = 0, \quad \langle V_\Gamma(\mathbf{x}) V_{\Gamma'}(\mathbf{x}') \rangle = \Delta_\Gamma \delta_{\Gamma\Gamma'} \delta(\mathbf{x} - \mathbf{x}') \quad (6)$$

where $\langle \dots \rangle$ denotes average over disorder distribution and Δ_Γ is introduced to characterize the strength of random potential.

We consider three different types of disorder classified by the different matrices Γ . In particular, $\Gamma = \mathbb{I}_{2 \times 2}$ for RSP, $\Gamma = \sigma_z$ for RM, and $\Gamma = (\sigma_x, \sigma_y)$ for RVP. These

three types are most frequently studied in the literature and they can be induced by some specific mechanisms in realistic materials [67–73]. These three types of random potential might exist individually, or coexist in the same material. To be general, we assume that they coexist in the system and analyze their impact by performing RG calculations.

The random potential $V(\mathbf{x})$ can be properly averaged by employing the replica trick [39, 51, 52, 55, 74, 75], which leads us to an interacting effective action of short-range fermion-fermion interaction:

$$\begin{aligned} S_{\text{dis}} = & -\frac{1}{2} \int d^2x d\tau d\tau' \left\{ \sum_{n=\pm} \left[\Delta_S (\psi_n^{\dagger\alpha} \psi_n^\alpha)_x (\psi_n^{\dagger\beta} \psi_n^\beta)_{x'} + \Delta_M (\psi_n^{\dagger\alpha} \sigma_z \psi_n^\alpha)_x (\psi_n^{\dagger\beta} \sigma_z \psi_n^\beta)_{x'} \right. \right. \\ & + \Delta_V \sum_j (\psi_n^{\dagger\alpha} \sigma_j \psi_n^\alpha)_x (\psi_n^{\dagger\beta} \sigma_j \psi_n^\beta)_{x'} \left. \right] + 2\Delta'_S (\psi_+^{\dagger\alpha} \psi_+^\alpha)_x (\psi_-^{\dagger\beta} \psi_-^\beta)_{x'} \\ & + 2\Delta'_M (\psi_+^{\dagger\alpha} \sigma_z \psi_+^\alpha)_x (\psi_-^{\dagger\beta} \sigma_z \psi_-^\beta)_{x'} + 2\Delta'_V \sum_j (\psi_+^{\dagger\alpha} \sigma_j \psi_+^\alpha)_x (\psi_-^{\dagger\beta} \sigma_j \psi_-^\beta)_{x'} \left. \right\} \quad (7) \end{aligned}$$

where $j = (x, y)$, α and β are the replica indices, $x \equiv (\mathbf{x}, \tau)$ and $x' \equiv (\mathbf{x}, \tau')$ are the space-time coordinate. The repeated indices α and β are summed automatically. In the replica theory, the replica limit $\sum_\alpha = N \rightarrow 0$ is implemented in the following RG calculation. Three parameters Δ_S , Δ_M , and Δ_V characterize the effective strength of quartic couplings of Dirac fermions induced by averaging over RM, RVP, and RSP, respectively. The two pieces of Dirac fermions share the same random potential. Three cross terms characterized by Δ'_S , Δ'_M , and Δ'_V are induced in the replica limit. In the RG analysis, the bare values of the parameters are the same, $\Delta_S^0 = \Delta_S$, $\Delta_M^0 = \Delta_M$, and $\Delta_V^0 = \Delta_V$, moreover, the RG equations for Δ_Γ and $\Delta_{\Gamma'}$ are the same (see Appendix B for details), so we focus on Δ_Γ in the following.

As shown in previous calculations [49, 59, 64], disorder can strongly affect the RG flow of fermion velocity as the energy is lowered. If the disorder coupling is relevant that flow to a finite value at low energy limit, it will drive the fermion velocity to vanish at sufficiently low energies, which then spoils the Lorentz symmetry for

the fermion sector, but not for the boson sector. As a result, the emergent Lorentz symmetry, and thus the emergent SUSY will be ruined by disorder. However, whether this takes place relies crucially on the scale dependence of disorder coupling parameter. In the case of (2+1)D DSMs, naive power-counting, according to Eqs. (2) and (7), shows that disorder is marginal. A careful analysis of the marginal disorder effect is helpful to tell us whether a irrelevant and a relevant coupling need to investigate further.

To this end, we carry out a detailed RG analysis starting from the critical action with $r = 0$, represented by Eq. (1) along with Eq. (7), by considering the leading order of the ϵ -expansion, where $\epsilon = 4 - D = 3 - d$, D and d are the spacetime dimension and the spatial dimension, respectively. The pertinent one-loop Feynman diagrams are shown in Fig. 2. After integrating out the fast modes defined within the momentum shell $e^{-\Lambda} < |\mathbf{p}| < \Lambda$ and then performing RG transformations [76], we obtain the following RG equations (the detailed results are presented in appendix)

$$\frac{dv_f}{dl} = v_f \left[g^2 (G_1 - G_0) - 2 \sum_\Gamma \Delta_\Gamma \right], \quad (8)$$

$$\frac{da}{dl} = g^2 \left(\frac{1 - a^2}{2a} + a(G_0 - G_1) \right) + 2a \sum_\Gamma \Delta_\Gamma, \quad (9)$$

$$\frac{dg^2}{dl} = \epsilon g^2 - g^4 (1 - G_0 + 3G_1) + (4\Delta_S + 2 \sum_\Gamma \Delta_\Gamma) g^2, \quad (10)$$

$$\frac{d\Delta_S}{dl} = (\epsilon - 1)\Delta_S + 2\Delta_S(\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta_M\Delta_V + (G_0 - 3G_2 - 2G_1)\Delta_S g^2, \quad (11)$$

$$\frac{d\Delta_M}{dl} = (\epsilon - 1)\Delta_M - 2\Delta_M(\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta_S\Delta_V - (G_0 + 3G_2 + 2G_1)\Delta_M g^2, \quad (12)$$

$$\frac{d\Delta_V}{dl} = (\epsilon - 1)\Delta_V + 2\Delta_M\Delta_S - (G_0 + 2G_1)\Delta_V g^2, \quad (13)$$

where $a = v_b/v_f$, $G_0 = \frac{4}{a(a+1)^2}$, $G_1 = \frac{4(2a+1)}{3a(a+1)^2}$, and $G_2 = \frac{4(2+a)}{3a(a+1)^2}$. In the above calculations, we have rescaled all the couplings as follows: $g^2\Lambda^{-\epsilon}S_{D-1}/[2(2\pi)^{D-1}v_f^{D-1}] \rightarrow g^2$, and $\Delta_\Gamma\Lambda^{1-\epsilon}S_{D-1}/[(2\pi)^{D-1}v_f^2] \rightarrow \Delta_\Gamma$, with $S_d = 2\pi^{d/2}/\Gamma(d/2)$ as the area of the unit sphere in d dimensions. By setting all $\Delta_\Gamma = 0$, Eqs. (9) - (10) recover the RG equations for v_f , a , and g^2 previously obtained in Refs. 25 and 35. In the case of disordered Dirac fermion systems with $g = 0$, our RG results for Δ_Γ are in accordance with that previously obtained in [53, 54, 77]. The RG equations for u and u_{+-} , which are not shown here, are exactly the same as those presented in Refs. 25 and 35, since there is no direct coupling between boson and fermion disorder potential. In the case of clean system, as demonstrated in Refs. 25 and 35, $a^* = 1$ is the only stable infrared fixed point for a , which means that the bosons and fermions have the same velocity at low energies. Moreover, the coupling constant g , u and u_{+-} will flow to a strongly coupled fixed point that preserves SUSY. In the following, we first analyze the effects of single disorder, and then consider the interplay between different types of disorder.

We now consider the case in which RVP exists by itself by taking $\Delta_M = \Delta_S = 0$. Noting that the physical case of (2+1)D corresponds to $\epsilon \rightarrow 1$, Eq. (13) becomes

$$\frac{d\Delta_V}{dl} = -(G_0 + 2G_1)\Delta_V g^2 \quad (14)$$

Thus the effective coupling strength for RVP, namely Δ_V , is irrelevant and flows to zero. Without the fermion-boson coupling g , RVP is marginal, which originates from the existence of a time-independent gauge transformation that ensures RVP unrenormalized and is valid at any order of loop expansion [47, 48, 60, 64, 77]. Nevertheless, near the emergent SUSY fixed point, where the coupling constant g remains finite, RVP is irrelevant, and thus the emergent SUSY is stable.

We then assume that RM exists alone, which means $\Delta_S = \Delta_V = 0$ in Eq. (12), and we have

$$\frac{d\Delta_M}{dl} = -2\Delta_M^2 - (G_0 + 3G_2 + 2G_1)\Delta_M g^2. \quad (15)$$

From the RG function, we see that Δ_M is always irrelevant. We thus can infer that the emergent SUSY is also robust against RM.

The RSP can be similarly analyzed. The simplified RG equations for RSP are

$$\frac{da}{dl} = g^2 \left(\frac{1-a^2}{2a} + a(G_0 - G_1) \right) + 2a\Delta_S, \quad (16)$$




FIG. 3. Flow diagram in the $\Delta_S - a$ plane when the system contains only RSP. There is an stable fixed point (red point) $(\Delta_S^*, a^*) = (0, 1)$.

$$\frac{dg^2}{dl} = g^2 - g^4(1 - G_0 + 3G_1) + 6g^2\Delta_S, \quad (17)$$

$$\frac{d\Delta_S}{dl} = 2\Delta_S^2 + (G_0 - 3G_2 - 2G_1)\Delta_S g^2, \quad (18)$$

There is a stable fixed point g_c^2 for g^2 ,

$$g_c^2 = \frac{1 + 6\Delta_S}{1 - G_0 + 3G_1}, \quad (19)$$

the RG function of Δ_S near $\Delta_S = 0$ is negative ($G_0 - 3G_2 - 2G_1 < 0$), and we expect small Δ_S is irrelevant. The RG flow of the equations within the critical plane $g^2 = g_c^2$ is shown in fig. (3). We find that SUSY is stable: an arbitrarily strong RSP flows to the weak coupling regime in the lowest energy limit. This means that the RSP is an irrelevant perturbation, which is consistent with the one-loop RG result in [79]. Previous results show that the RSP is marginally relevant for Dirac fermions, and will induce an instability of the system, leading to a diffusive motion of the Dirac fermions [40, 43, 51-54, 77]. However, our result shows that, the RSP is rendered irrelevant by the critical fluctuations at the PDW QCP, and there exists no diffusive behaviors.

When more than one type of disorder exist, from the properties of the RG equation, Eqs. (12)-(14) that the coexistence of any two types disorder dynamically generate the third one. Thus we need to analyze the full set of RG equations given by Eqs. (8)-(13). It is easy to see that there is a fixed point given by $(v_f, a^*, g^{2*}, \Delta_S^*, \Delta_M^*, \Delta_V^*) = (v_f, 1, \frac{\epsilon}{3}, 0, 0, 0)$, where v_f

can take any values. Combining with the RG equation for coupling constants u and u_{+-} , it turns out that this is the SUSY fixed point. Now we examine whether this fixed point is stable by expanding the RG equation at the fixed point, and calculate eigenvalues of the stability matrix (see Appendix C). The eigenvalues of the stability matrix are all negative except for one marginal direction at v_f . So we can conclude that the emergent SUSY is robust against weak disorders.

III. (3+1)D WEYL SEMIMETALS

In this section, we examine the disorder effects on the emergent SUSY in (3+1)D WSMs [35]. The effective action of the disordered system in the vicinity of PDW QCP is given by

$$S = S_f + S_b + S_I + S_{\text{dis}}, \quad (20)$$

$$S_f = \int d^4x \sum_{n=\pm} \left[\psi_n^\dagger \partial_\tau \psi_n + \sum_{j=1}^3 i v_{fj} \psi_n^\dagger \gamma_n^j \partial_j \psi_n \right], \quad (21)$$

$$S_b = \int d^4x \left\{ \sum_{n=\pm} \left[|\partial_\tau \phi_n|^2 + \sum_{j=1}^3 v_{bj}^2 |\partial_j \phi_n|^2 + r |\phi_n|^2 + u |\phi_n|^4 \right] + u_{+-} |\phi_+|^2 |\phi_-|^2 \right\}, \quad (22)$$

$$S_I = \int d^4x g \sum_{n=\pm} \left[\phi_n \psi_n \sigma^y \psi_n + h.c. \right], \quad (23)$$

$$S_{\text{dis}} = -\frac{1}{2} \int d^3x d\tau d\tau' \left[\Delta_S (\psi_\alpha^\dagger \sigma_0 \psi_\alpha)_x (\psi_\beta^\dagger \sigma_0 \psi_\beta)_{x'} + \sum_{i=x,y,z} \Delta_i (\psi_\alpha^\dagger \sigma_i \psi_\alpha)_x (\psi_\beta^\dagger \sigma_i \psi_\beta)_{x'} \right], \quad (24)$$

where $\gamma_\pm^j = (\sigma^x, \sigma^y, \pm\sigma^z)$. Now, ψ_\pm denotes the two-component Weyl fermions at two Weyl points \mathbf{Q}_\pm , and ϕ_\pm is the superconducting order with momentum $2\mathbf{Q}_\pm$, respectively. In (3+1)D Weyl semimetal, RVP has three components, i.e., RM becomes the third component [80, 81]. In general, the fermion velocity is anisotropic, with unequal values along different directions. As shown in Ref. 35, even in the extremely anisotropic case, an emergent Lorentz symmetry can be established in the lowest energy limit. Our current concern is whether this emergent Lorentz symmetry can be broken by quenched disorder, thus it suffices to consider the isotropic case. We now can assume that $v_{fx} = v_{fy} = v_{fz} = v_f$, and also make the same assumption for the bosonic field. We employ the same symbol a to identify the ratio between boson and fermion, and the same definition of G_0, G_1 and the rescaled couplings in Sec. II. Calculating the same diagrams in Fig. 2, we obtain the following RG equations:

$$\frac{d \ln v_f}{dl} = g^2 (G_1 - G_0) - 2 \left(\Delta_S + \sum_i \Delta_i \right), \quad (25)$$

$$\frac{d \ln a^2}{dl} = \frac{g^2 (1 - a^2)}{a^2} - 2g^2 (G_1 - G_0) + 4 \left(\Delta_S + \sum_i \Delta_i \right), \quad (26)$$

$$\frac{d \ln g^2}{dl} = \epsilon - g^2 (3G_1 - G_0 + 1) + \left(6\Delta_S + 2 \sum_i \Delta_i \right), \quad (27)$$

$$\frac{d \Delta_S}{dl} = (\epsilon - 1) \Delta_S + 2\Delta_S \left(\Delta_S + \sum_i \Delta_i \right) + \frac{2}{3} \sum_i \sum_{j \neq i} \Delta_i \Delta_j + g^2 (G_0 - 3G_2 - 2G_1) \Delta_S, \quad (28)$$

$$\frac{d \Delta_i}{dl} = (\epsilon - 1) \Delta_i + \frac{4}{3} \sum_{j \neq i} \Delta_S \Delta_j - \frac{2}{3} \Delta_i \left(\Delta_S + 2\Delta_i - \sum_j \Delta_j \right) - \Delta_i g^2 (G_0 + G_2 + 2G_1), \quad (29)$$

in these equations, the index i is summed over x, y, z . The analysis of these RG equations follows similarly as done in Sec. II. Firstly, we consider only a single vector component disorder exists, which means

$$\Delta_i \neq 0, \quad \Delta_{j \neq i} = 0, \quad \Delta_S = 0, \quad (30)$$

by substituting this conditions to Eqs. (25-29), three simplified RG equations for a, g^2 and Δ_i are obtained, we just exhibit the result for disorder coupling as

$$\frac{d \Delta_i}{dl} = (\epsilon - 1) \Delta_i - \frac{2}{3} \Delta_i^2$$

$$- \Delta_i g^2 (G_0 + G_2 + 2G_1). \quad (31)$$

Therefore, for an exact (3+1)D system corresponding to $\epsilon = 0$, any component of RVP is irrelevant. Therefore, in (3+1)D WSMs, the emergent SUSY is robust against any single component of RVP.

For there is only RSP in the system, we have $\Delta_i = 0$. Now Eqs. (25) - (29) are simplified to

$$\frac{d \ln a^2}{dl} = \frac{g^2 (1 - a^2)}{a^2} - 2g^2 (G_1 - G_0) + 4\Delta_S, \quad (32)$$

$$\frac{d \ln g^2}{dl} = \epsilon - g^2(3G_1 - G_0 + 1) + 6\Delta_S, \quad (33)$$

$$\begin{aligned} \frac{d\Delta_S}{dl} &= (\epsilon - 1)\Delta_S + 2\Delta_S^2 \\ &\quad + g^2(G_0 - 3G_2 - 2G_1)\Delta_S, \end{aligned} \quad (34)$$

according to Eq. (33), and noting the fact $3G_1 - G_0 + 1 > 0$, the stable fixed point for g^2 is located at $g_c^2 = 6\Delta_S/(3G_1 - G_0 + 1)$. Weak RSP itself is irrelevant as indicated in because $G_0 - 3G_2 - 2G_1 < 0$. Whereas for strong RSP, the scenario is similar to the case of (2+1) DSMs, namely, the RSP becomes irrelevant due to the interplay between the Weyl fermion and the PDW order parameter. We thus conclude that the emergent SUSY is stable against RSP.

Next, we consider the coexisting case. According to Eq. (28), the coexistence of two components of RVP can dynamically generate RSP even when RSP does not exist at the beginning. We also learn from Eq. (29) that the coexistence of RSP and any component of RVP produce the other two components. Therefore, we need to consider the generic case in which all three components of RVP coexist with RSP. Now the disorder effects should be analyzed by solving the complete set of equations given by Eq. (25) - (29). It is hard to solve these coupled equations, but fortunately, it is simple to show that in the weak disorder regime the SUSY fixed point is stable against all random potentials, similar to the case of (2+1) DSMs.

IV. SUMMARY AND DISCUSSION

In summary, we perform a standard perturbative RG to study the effect of disorder on the emergent SUSY in (2+1)D DSMs [25, 35] and (3+1)D WSMs [35]. According to our RG results, the effective SUSY fixed point is robust against any weak disorder irrespective of the type of disorder potentials. Our RG analysis of the disorder effects on the emergent SUSY appearing in (2+1)D DSMs [25, 35] and (3+1)D WSMs [35] can be directly extended to other analogous models, which may more realistic to detect emergent SUSY in quantum materials. In our one-loop RG analysis, we have omitted new vertex that could be generated from the disorder potential and the Yukawa coupling. It will be interesting to include their effects although at tree-level they are irrelevant. Our work can shed new light on the understanding and exploring the emergent SUSY in realistic condensed matter systems.

ACKNOWLEDGMENTS

This work was supported by National Natural Science Foundation of China (No. 12147104). The work of SKJ is also supported in part by the Simons Foundation via the It From Qubit Collaboration.

APPENDIX A: DISORDER POTENTIALS

In this section, we briefly consider the possible disorder potential in the two-component Dirac (Weyl) system. The original spinful fermion annihilation operator ψ can be expand around two Dirac (Weyl) point \mathbf{Q}_\pm ,

$$\psi(\mathbf{x}) = e^{-i\mathbf{Q}_+ \cdot \mathbf{x}} \psi_+(\mathbf{x}) + e^{-i\mathbf{Q}_- \cdot \mathbf{x}} \psi_-(\mathbf{x}), \quad (35)$$

where ψ_\pm corresponds to the two low-energy Dirac (Weyl) fermion. For the fermion system ψ , a general disorder potential takes the form,

$$H_{\text{dis}} = \int d^d \mathbf{x} \psi^\dagger(\mathbf{x}) \left(\sum_{\Gamma} V_{\Gamma}(\mathbf{x}) \Gamma \right) \psi(\mathbf{x}), \quad (36)$$

where the $V_{\Gamma}(\mathbf{x})$ stands for the randomly distributed potential and $\Gamma = I, \sigma_\mu$ ($\mu = x, y, z$) labels type of the disorder potential. We focus on the quenched disorder potential $V_{\Gamma}(\mathbf{x})$, with Gaussian white noise potential characterized by the following identities:

$$\langle V_{\Gamma}(\mathbf{x}) \rangle = 0, \quad \langle V_{\Gamma}(\mathbf{x}) V_{\Gamma'}(\mathbf{x}') \rangle = \Delta_{\Gamma} \delta_{\Gamma\Gamma'} \delta^d(\mathbf{x} - \mathbf{x}') \quad (37)$$

Substituting the fermion field Eq. (35) into the disordered Hamiltonian H_{dis} , we can arrive at the disorder Hamiltonian for the two low-energy Dirac (Weyl) fermion ψ_\pm . Notice that for the coupling between the two Dirac field, e.g. $\psi_+^\dagger(\mathbf{x}) \Gamma \psi_-(\mathbf{x})$, there is an overall oscillating factor $e^{\pm i(\mathbf{Q}_+ + \mathbf{Q}_-) \cdot \mathbf{x}}$. Such disorder potential coupled the two Dirac (Weyl) fermion ψ_\pm will be small in general. We only need to consider the disorder Hamiltonian Eq. (5) with respect to a piece of Dirac fermion. It should be emphasized that the two pieces of Dirac (Weyl) fermion share the same statistical distribution of random potential.

APPENDIX B: RG DETAILS

We present here the detailed calculation of Fig. 2(a)-Fig. 2(h) as well as the RG equations in (2+1)D, the calculation for (3+1)D is directly followed, which is not detailed shown here.

From the free action of fermions and bosons,

$$S_{f0} = \int d^2 x d\tau \sum_{n=\pm} \psi_n^\dagger \left[\partial_\tau + iv_f \sum_{j=1}^2 \gamma_j \partial_j \right] \psi_n, \quad (38)$$

$$S_{b0} = \int d^2 x d\tau \left\{ \sum_{n=\pm} \left[|\partial_\tau \phi_n|^2 + v_b^2 \sum_{j=1}^2 |\partial_j \phi_n|^2 \right] \right\}, \quad (39)$$

the free propagator for fermions and bosons are

$$\begin{aligned} G_0(k) &= \frac{1}{ik_\tau - v_f \boldsymbol{\gamma} \cdot \mathbf{k}}; \\ D_0(k) &= \frac{1}{k_\tau^2 + v_b^2 \mathbf{k}^2}. \end{aligned}$$

with $k = (k_\tau, \mathbf{k})$ in the momentum space through replacement $(\partial_\tau, \partial_i) \rightarrow (ik_\tau, ik_i)$. The free propagators for the two pieces of Dirac fermion take the same form. For Fig. 2(a), it corresponds

$$\begin{aligned} \Pi(k) &= -2g^2 \int_p \text{Tr} [\sigma^y G_0^T(p) \sigma^y G_0(-p-k)] \\ &= \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} l \left(k_\tau^2 + \left(2 - \frac{3}{d}\right) v_f^2 \mathbf{k}^2 \right) \end{aligned} \quad (40)$$

where $\int_p \equiv \int dp_\tau d^d \mathbf{p} / (2\pi)^D$ is the $D = d+1$ dimensional momentum integral and $S_d = 2\pi^{d/2} / \Gamma(d/2)$ is the area of the unit sphere in d dimensions. For Fig. 2(c), it gives

$$\begin{aligned} \Sigma_f^{(b)}(k) &= -4g^2(-) \int_p \sigma^y G_0^T(p) \sigma^y D_0(-k-p) \\ &= \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} l \left(\frac{4(ik_\tau)}{a(a+1)^2} - \frac{4(2a+1)}{ad(a+1)^2} (v_f \boldsymbol{\gamma} \cdot \mathbf{k}) \right) \\ &= \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} l \left(G_0(ik_\tau) - G_1(v_f \boldsymbol{\gamma} \cdot \mathbf{k}) \right) \end{aligned} \quad (41)$$

The diagram of Fig. 2(d) is ,

$$\begin{aligned} \Sigma_f^{(c)}(k) &= - \sum_\Gamma (\Delta_\Gamma + \Delta'_\Gamma) \int \frac{d^d \mathbf{k}}{(2\pi)^d} \Gamma G_0(k) \Gamma \\ &= \sum_\Gamma (\Delta_\Gamma + \Delta'_\Gamma) \frac{\Lambda^{1-\epsilon} S_d}{(2\pi)^d v_f^2} (ik_\tau) \ell, \end{aligned} \quad (42)$$

which only contribute to the velocity renormalization at one-loop. The diagram of Fig. 2(e) is given by

$$\delta \Delta_\Gamma^{(d)} = -8\Delta_\Gamma g^2 \int_k \sigma^y D(-k) G^T(k) \Gamma^T G^T(k) \sigma^y \quad (43)$$

$$\delta \Delta'_\Gamma^{(d)} = -8\Delta'_\Gamma g^2 \int_k \sigma^y D(-k) G^T(k) \Gamma^T G^T(k) \sigma^y \quad (44)$$

Calculating out these integrals for different Γ_a one by one, we have

$$\delta \Delta_S^{(d)} = \Delta_S \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} (G_0 - dG_2), \quad (45)$$

$$\delta \Delta_M^{(d)} = \Delta_M \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} (-G_0 - dG_2), \quad (46)$$

$$\delta \Delta_V^{(d)} = \Delta_V \frac{g^2 S_d \Lambda^{-\epsilon}}{2(2\pi)^d v_f^3} (-G_0), \quad (47)$$

and same for $\Delta \rightarrow \Delta'$. The diagram of Fig. 2(b) involves Δ_γ -vertex and is given by

$$\begin{aligned} \delta g^{(h)} &= g \sum_\Gamma \Delta_\Gamma \int_k \Gamma^T G_+^T(k) \sigma^y G_+(-k) \Gamma \\ &= gl \sum_\Gamma \frac{\Delta_\Gamma \Lambda^{1-\epsilon} S_d}{(2\pi)^d v_f^2} \Gamma^T \sigma^y \Gamma \end{aligned} \quad (48)$$

The remaining diagrams Fig. 2(f,g,h) correspond to pure coupling between disorder potentials [60], which can be obtained as follows,

$$\delta \Delta_S^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[+2\Delta_S (\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta_M \Delta_V \right] \quad (49)$$

$$\delta \Delta_M^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[-2\Delta_M (\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta_S \Delta_V \right] \quad (50)$$

$$\delta \Delta_V^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[2\Delta_M \Delta_S \right] \quad (51)$$

and similarly for the other three terms,

$$\delta \Delta'_S{}^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[+2\Delta'_S (\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta'_M \Delta'_V \right] \quad (52)$$

$$\delta \Delta'_M{}^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[-2\Delta'_M (\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta'_S \Delta'_V \right] \quad (53)$$

$$\delta \Delta'_V{}^{(e,f,g)} = \left(\frac{S_d}{(2\pi)^d v_f^2 \Lambda^{2-d}} \right) l \left[2\Delta'_M \Delta'_S \right] \quad (54)$$

According to above results, label and rescale the couplings as follows:

$$G_0 = \frac{4}{a(1+a)^2}, \quad G_1 = \frac{4(1+2a)}{da(1+a)^2}, \quad G_2 = \frac{4(2+a)}{da(1+a)^2}, \quad \frac{g^2 \Lambda^{-\epsilon} S_d}{2(2\pi)^d v_f^d} \rightarrow g^2, \quad \frac{\Delta_\Gamma \Lambda^{1-\epsilon} S_d}{(2\pi)^d v_f^2} \rightarrow \Delta_\Gamma. \quad (55)$$

Then, the results obtained for Fig. 2(a)-Fig. 2(h) can be simplified, leading to the one-loop quantum corrections of

the action. These one-loop results produce the RG equations

$$\frac{dv_f}{dl} = v_f \left[g^2(G_1 - G_0) - \sum_{\Gamma} \Delta_{\Gamma} - \sum_{\Gamma} \Delta'_{\Gamma} \right], \quad (56)$$

$$\frac{da}{dl} = g^2 \left(\frac{1-a^2}{2a} + a(G_0 - G_1) \right) + a \sum_{\Gamma} \Delta_{\Gamma} + a \sum_{\Gamma} \Delta'_{\Gamma}, \quad (57)$$

$$\frac{dg^2}{dl} = \epsilon g^2 - g^4(1 - G_0 + 3G_1) + (4\Delta_S - \sum_{\Gamma} \Delta_{\Gamma} + 3 \sum_{\Gamma} \Delta'_{\Gamma})g^2, \quad (58)$$

$$\frac{d\Delta_S}{dl} = (\epsilon - 1)\Delta_S + 2\Delta_S(\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta_M\Delta_V + (G_0 - 3G_2 - 2G_1)\Delta_S g^2, \quad (59)$$

$$\frac{d\Delta_M}{dl} = (\epsilon - 1)\Delta_M - 2\Delta_M(\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta_S\Delta_V - (G_0 + 3G_2 + 2G_1)\Delta_M g^2, \quad (60)$$

$$\frac{d\Delta_V}{dl} = (\epsilon - 1)\Delta_V + 2\Delta_M\Delta_S - (G_0 + 2G_1)\Delta_V g^2, \quad (61)$$

$$\frac{d\Delta'_S}{dl} = (\epsilon - 1)\Delta'_S + 2\Delta'_S(\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta'_M\Delta'_V + (G_0 - 3G_2 - 2G_1)\Delta'_S g^2, \quad (62)$$

$$\frac{d\Delta'_M}{dl} = (\epsilon - 1)\Delta'_M - 2\Delta'_M(\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta'_S\Delta'_V - (G_0 + 3G_2 + 2G_1)\Delta'_M g^2, \quad (63)$$

$$\frac{d\Delta'_V}{dl} = (\epsilon - 1)\Delta'_V + 2\Delta'_M\Delta'_S - (G_0 + 2G_1)\Delta'_V g^2, \quad (64)$$

Since the initial values of the parameters are $\Delta_{\Gamma}^0 = \Delta'_{\Gamma}$, Δ_{Γ} and Δ'_{Γ} will flow in the same way. We can simplify the equations by taken $\Delta'_{\Gamma} = \Delta_{\Gamma}$ and produce the Eqs. (8) - (13).

do not show details here.

For the case of $(3+1)D$, due to the change of disorder types, the one-loop corrections of disorder couplings need to recalculate, the extension is direct for which we

APPENDIX C: GENERAL DISORDER CASE

In this section, we study the case where all the disorder potential is appeared. We focus on the regime near the Lorentz symmetric fixed point, $a = 1 + \delta a$ with small $\delta a \ll 1$. The RG equations in 2D now become,

$$\frac{dv_f}{dl} = v_f \left[g^2 \frac{2}{3} \delta a - 2 \sum_{\Gamma} \Delta_{\mu} \right], \quad (65)$$

$$\frac{d\delta a}{dl} = \left(-\frac{5}{3}g^2 + 2 \sum_{\Gamma} \Delta_{\Gamma} \right) \delta a, \quad (66)$$

$$\frac{dg^2}{dl} = \epsilon g^2 - g^4(3 - 2\delta a) + (4\Delta_S + 2 \sum_{\Gamma} \Delta_{\Gamma})g^2, \quad (67)$$

$$\frac{d\Delta_S}{dl} = (\epsilon - 1)\Delta_S + 2\Delta_S(\Delta_S + \Delta_M + 2\Delta_V) + 4\Delta_M\Delta_V + \left(-4 + \frac{17}{3}\delta a\right)\Delta_S g^2, \quad (68)$$

$$\frac{d\Delta_M}{dl} = (\epsilon - 1)\Delta_M - 2\Delta_M(\Delta_S + \Delta_M - 2\Delta_V) + 4\Delta_S\Delta_V - \left(6 - \frac{29}{3}\delta a\right)\Delta_M g^2, \quad (69)$$

$$\frac{d\Delta_V}{dl} = (\epsilon - 1)\Delta_V + 2\Delta_M\Delta_S - \left(3 - \frac{14}{3}\delta a\right)\Delta_V g^2, \quad (70)$$

There is a fixed point given by $(v_f, a^*, g^{2*}, \Delta_S^*, \Delta_M^*, \Delta_V^*) = (v_f, 1, \frac{\epsilon}{3}, 0, 0, 0)$. For small disorder case, the stability matrix at this fixed

point is

$$\begin{pmatrix} 0 & \frac{2}{9}\epsilon v_f & 0 & -2v_f & -2v_f & -2v_f \\ 0 & -\frac{5}{9}\epsilon & 0 & 2 & 2 & 2 \\ 0 & \frac{2}{9}\epsilon^2 & -\epsilon & 2\epsilon & \frac{2}{3}\epsilon & \frac{2}{3}\epsilon \\ 0 & 0 & 0 & -1 - \frac{\epsilon}{3} & 0 & 0 \\ 0 & 0 & 0 & 0 & -1 - \epsilon & 0 \\ 0 & 0 & 0 & 0 & 0 & -1 \end{pmatrix}. \quad (71)$$

The eigenvalues of the stability matrix are all negative except for one marginal direction at v_f . We can conclude that the coupling between Yukawa potential and disorder will suppress weak random disorder potential. Similarly argument can also be applied to three dimensional case.

-
- [1] S. Weinberg, *The Quantum Theory of Fields: Supersymmetry*, Cambridge University Press, Cambridge (2000).
- [2] J. L. Gervais and B. Sakita, Nucl. Phys. B **34**, 632 (1971).
- [3] J. Wess and B. Zumino, Nucl. Phys. B **70**, 39 (1974).
- [4] S. Dimopoulos and H. Georgi, Nucl. Phys. B **193**, 150 (1981).
- [5] E. Cremmer, S. Ferrara, C. Kounnas, and D. V. Nanopoulos, Phys. Lett. B **133**, 61 (1983).
- [6] N. Seiberg and E. Witten, Nucl. Phys. B **426**, 19 (1994).
- [7] N. Seiberg, Nucl. Phys. B **435**, 129 (1995).
- [8] X. Wan, A. M. Turner, A. Vishwanath, and S. Y. Savrasov, Phys. Rev. B **83**, 205101 (2011).
- [9] G. Xu, H. Weng, Z. Wang, X. Dai, and Z. Fang, Phys. Rev. Lett. **107**, 186806 (2011).
- [10] A. A. Burkov and L. Balents, Phys. Rev. Lett. **107**, 127205 (2011).
- [11] H. Weng, C. Fang, Z. Fang, B. A. Bernevig, and X. Dai, Phys. Rev. X **5**, 011029 (2015).
- [12] S.-M. Huang, S.-Y. Xu, I. Belopolski, C.-C. Lee, G. Chang, B. Wang, N. Alidoust, G. Bian, M. Neupane, C. Zhang, S. Jia, A. Bansil, H. Lin, and M. Z. Hasan, Nat. Commun. **6**, 7373 (2015).
- [13] S.-Y. Xu, I. Belopolski, N. Alidoust, M. Neupane, G. Bian, C. Zhang, R. Sankar, G. Chang, Z. Yuan, C.-C. Lee, S.-M. Huang, H. Zheng, J. Ma, D. S. Sanchez, B. Wang, A. Bansil, F. Chou, P. P. Shibayev, H. Lin, S. Jia, and M. Z. Hasan, Science **349**, 613 (2015).
- [14] B. Q. Lv, H. M. Weng, B. B. Fu, X. P. Wang, H. Miao, J. Ma, P. Richard, X. C. Huang, L. X. Zhao, G. F. Chen, Z. Fang, X. Dai, T. Qian, and H. Ding, Phys. Rev. X **5**, 031013 (2015).
- [15] L. X. Yang, Z. K. Liu, Y. Sun, H. Peng, H. F. Yang, T. Zhang, B. Zhou, Y. Zhang, Y. F. Guo, M. Rahn, D. Prabhakaran, Z. Hussain, S. K. Mo, C. Felser, B. Yan, and Y. L. Chen, Nat. Phys. **11**, 728 (2015).
- [16] G. Curci and G. Veneziano, Nucl. Phys. B **292**, 555 (1987).
- [17] H.-S. Goh, M. A. Luth and S.-P. Ng, J. High Energy Phys. **01**, 40 (2005).
- [18] S. Thomas, *Emergent Supersymmetry*, KITP talk (2005).
- [19] Prakash, Abhishodh, and Juven Wang, Physical Review Letters **126**.23 (2021): 236802.
- [20] Li, Zi-Xiang, et al, Science advances **4**.11 (2018): eaau1463.
- [21] Li, Zi-Xiang, and Hong Yao, Physical Review B **96**.24 (2017): 241101.
- [22] Li, Zi-Xiang, Yi-Fan Jiang, and Hong Yao, Physical Review Letters **119**.10 (2017): 107202.
- [23] Jian, Shao-Kai, et al, Physical review letters **118**.16 (2017): 166802.
- [24] Yu, Jiabin, et al, Physical Review B **100**.7 (2019): 075153.
- [25] S.-S. Lee, Phys. Rev. B **76**, 075103 (2007).
- [26] Y. Yu and K. Yang, Phys. Rev. Lett. **105**, 150605 (2010).
- [27] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
- [28] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
- [29] T. Grover, D. N. Sheng, and A. Vishwanath, Science **344**, 208 (2014).
- [30] P. Ponte and S.-S. Lee, New J. Phys. **16**, 013044 (2014).
- [31] N. Zerf, C.-H. Lin, and J. Maciejko, Phys. Rev. B **94**, 205106 (2016).
- [32] D. Friedan, Z. Qiu, and S. Shenker, Phys. Rev. Lett. **52**, 1575 (1984).
- [33] O. Foda, Nucl. Phys. B **300**, 611 (1988).
- [34] L. Huijse, B. Bauer, and E. Berg, Phys. Rev. Lett. **114**, 090404 (2015).
- [35] S.-K. Jian, Y.-F. Jiang, and H. Yao, Phys. Rev. Lett. **114**, 237001 (2015).
- [36] J. Ruan, S.-K. Jian, H. Yao, H. Zhang, S.-C. Zhang, and D. Xing, Nat. Commun. **7**, 11136 (2016).
- [37] J. Ruan, S.-K. Jian, D. Zhang, H. Yao, H. Zhang, S.-C. Zhang, and D. Xing, Phys. Rev. Lett. **116**, 226801 (2016).
- [38] B. L. Altshuler and A. G. Aronov, in *Electron-Electron Interaction in Disordered Systems*, edited by M. Pollak and A. L. Efros (North-Holland, Amsterdam, 1984).
- [39] P. A. Lee and T. V. Ramakrishnan, Rev. Mod. Phys. **57**, 287 (1985).
- [40] E. Fradkin, Phys. Rev. B **33**, 3257 (1986); **33**, 3263 (1986).
- [41] D. Belitz and T. R. Kirkpatrick, Rev. Mod. Phys. **66**, 261 (1994).
- [42] E. Abrahams, S. V. Kravchenko, and M. R. Sarachik, Rev. Mod. Phys. **73**, 251 (2001).
- [43] A. Altland, B. D. Simons, and M. R. Zirnbauer, Phys. Rep. **359**, 283 (2002).
- [44] S. D. Sarma, S. Adam, E. H. Hwang, and E. Rossi, Rev. Mod. Phys. **83**, 407 (2011).

- [45] V. N. Kotov, B. Uchoa, V. M. Pereira, F. Guinea, and A. H. Castro Neto, *Rev. Mod. Phys.* **84**, 1067 (2012).
- [46] T. Stauber, F. Guinea, and M. A. H. Vozmediano, *Phys. Rev. B* **71**, 041406(R) (2005).
- [47] I. F. Herbut, V. Juričić, and O. Vafek, *Phys. Rev. Lett.* **100**, 046403 (2008).
- [48] O. Vafek and M. J. Case, *Phys. Rev. B* **77**, 033410 (2008).
- [49] J.-R. Wang and G.-Z. Liu, *Phys. Rev. B* **89**, 195404 (2014).
- [50] P.-L. Zhao, J.-R. Wang, A.-M. Wang, and G.-Z. Liu, *Phys. Rev. B* **94**, 195114 (2016).
- [51] P. Goswami and S. Chakravarty, *Phys. Rev. Lett.* **107**, 196803 (2011).
- [52] B. Roy and S. Das Sarma, *Phys. Rev. B* **90**, 241112(R) (2014).
- [53] P. M. Ostrovsky, I. V. Gornyi, and A. D. Mirlin, *Phys. Rev. B* **74**, 235443 (2006).
- [54] M. S. Foster and I. L. Aleiner, *Phys. Rev. B* **77**, 195413 (2008).
- [55] B. Roy, V. Juričić, and S. Das Sarma, *Sci. Rep.* **6**, 32446 (2016).
- [56] D. H. Kim, P. A. Lee, and X.-G. Wen, *Phys. Rev. Lett.* **79**, 2109 (1997).
- [57] P. A. Lee, N. Nagaosa, and X.-G. Wen, *Rev. Mod. Phys.* **78**, 17 (2006).
- [58] J. Wang, G.-Z. Liu, and H. Kleinert, *Phys. Rev. B* **83**, 214503 (2011).
- [59] J. Wang, *Phys. Rev. B* **87**, 054511 (2013).
- [60] A. W. W. Ludwig, M. P. A. Fisher, R. Shankar, and G. Grinstein, *Phys. Rev. B* **50**, 7526 (1994).
- [61] J. E. Furneaux, S. V. Kravchenko, W. E. Mason, G. E. Bowker, and V. M. Pudalov, *Phys. Rev. B* **51**, 17227 (1995).
- [62] J. Ye and S. Sachdev, *Phys. Rev. Lett.* **80**, 5409 (1998).
- [63] J. Ye, *Phys. Rev. B* **60**, 8290 (1999).
- [64] P.-L. Zhao, A.-M. Wang, and G.-Z. Liu, arXiv:1701.08529.
- [65] J. González, F. Guinea, and M. A. H. Vozmediano, *Nucl. Phys. B* **406**[FS], 771 (1993).
- [66] J. González, F. Guinea, and M. A. H. Vozmediano, *Nucl. Phys. B* **424**[FS], 595 (1994).
- [67] A. A. Nersesyan, A. M. Tsvetik, and F. Wenger, *Nucl. Phys. B* **438**, 561 (1995).
- [68] A. H. Castro Neto, F. Guinea, N. M. R. Peres, K. S. Novoselov, and A. K. Geim, *Rev. Mod. Phys.* **81**, 109 (2009).
- [69] N. M. R. Peres, *Rev. Mod. Phys.* **82**, 2673 (2010).
- [70] E. R. Mucciolo and C. H. Lewenkopf, *J. Phys. Condens. Matter* **22**, 273201 (2010).
- [71] J. C. Meyer, A. K. Geim, M. I. Katsnelson, K. S. Novoselov, T. J. Booth, and S. Roth, *Nature* **446**, 60 (2007).
- [72] T. Champel and S. Florens, *Phys. Rev. B* **82**, 045421 (2010).
- [73] S. V. Kusminskiy, D. K. Campbell, A. H. Castro Neto, and F. Guinea, *Phys. Rev. B* **83**, 165405 (2011).
- [74] I. V. Lerner, arXiv:cond-mat/0307471.
- [75] B. Roy and S. Das Sarma, *Phys. Rev. B* **94**, 115137 (2016).
- [76] R. Shankar, *Rev. Mod. Phys.* **66**, 129 (1994).
- [77] M. S. Foster, *Phys. Rev. B* **85**, 085122 (2012).
- [78] J. Wang, P.-L. Zhao, J.-R. Wang, and G.-Z. Liu, *Phys. Rev. B* **95**, 054507 (2017).
- [79] R. Nandkishore, J. Maciejko, D. A. Huse, and S. L. Sondhi, *Phys. Rev. B* **87**, 174511 (2013).
- [80] B. Sbierski, K. S. C. Decker, and P. W. Brouwer, *Phys. Rev. B* **94**, 220202(R) (2016).
- [81] S. V. Syzranov, L. Radzihovsky, arXiv:1609.05694