On Data Augmentation for Models Involving Reciprocal Gamma Functions

Yasuyuki Hamura^{*}, Kaoru Irie[†], and Shonosuke Sugasawa[‡]

August 29, 2022

Abstract

In this paper, we introduce a new and efficient data augmentation approach to the posterior inference of the models with shape parameters when the reciprocal gamma function appears in full conditional densities. Our approach is to approximate full conditional densities of shape parameters by using Gauss's multiplication formula and Stirling's formula for the gamma function, where the approximation error can be made arbitrarily small. We use the techniques to construct efficient Gibbs and Metropolis-Hastings algorithms for a variety of models that involve the gamma distribution, Student's *t*-distribution, the Dirichlet distribution, the negative binomial distribution, and the Wishart distribution. The proposed sampling method is numerically demonstrated through simulation studies.

Key words and phrases: Gauss's multiplication formula, Markov chain Monte Carlo, reciprocal gamma function, Stirling's formula.

1 Introduction

Markov chain Monte Carlo (MCMC) algorithms are now widely adopted in Bayesian posterior computation, where parameters are iteratively sampled from their respective conditional distributions. However, when the models of interest involve the gamma and related distributions, it is computationally costly to sample the shape parameters from their full conditional posteriors. The main difficulty here is that the full conditional densities of the shape parameters involve the reciprocal gamma function, $1/\Gamma(\xi)$, $\xi > 0$, and are not any well-known distributions. Thus, it is not straightforward to construct efficient MCMC algorithms when the shape parameters are also estimated.

Several sampling strategies that have been proposed in the literature are customized for each class of distributions. For gamma distributions, Miller (2019) provided an accurate approximation of the full conditional distribution of the shape parameter. For Student's t-distributions,

^{*}Corresponding author. Graduate School of Economics, Kyoto University, Yoshida-Honmachi, Sakyo-ku, Kyoto, 606-8501, JAPAN.

E-Mail: yasu.stat@gmail.com

[†]Faculty of Economics, The University of Tokyo.

E-Mail: irie@e.u-tokyo.ac.jp

[‡]Center for Spatial Information Science, The University of Tokyo. E-Mail: sugasawa@csis.u-tokyo.ac.jp

Fonseca et al. (2008) considered the unknown degrees of freedom, at the cost of the complication of the priors. For Dirichlet-multinomial and negative-binomial models, sampling algorithms for the shape parameters have been proposed by Nandram (1998) and Zhou and Carin (2015), respectively.

Rather than focusing on a particular class of distributions, it is also possible to devise the sampling methods that are applicable to the general class of models with shape parameters, at the cost of efficiency and computational time. For example, the approximation of log-concave densities (Gilks and Wild, 1992; Devroye, 2012) and the MH acceptance-rejection method (Tierney, 1994; Chib and Greenberg, 1995) can be used for the posterior inference for models with the reciprocal gamma functions. The latter needs to be further customized to each model, as practiced for Student's t-models in Watanabe (2001). Another approach is the data augmentation scheme, where several latent variables are introduced to simplify the full conditionals of the model. He et al. (2021) proposed a general and efficient data augmentation for models with reciprocal gamma functions, where the simulation from power truncated normal (PTN) distributions become necessary. In this paper, we also take the data-augmentation approach, but propose a new augmentation where we only need to simulate from well-known distributions.

Our strategy for deriving an augmented model is twofold: (i) using Gauss's multiplication formula for the gamma function to introduce conditionally beta-distributed latent variables and (ii) approximating the augmented densities by Stirling's formula. The full conditionals of the shape parameters and latent variables of the resulting model are all well-known distributions, such as gamma and beta distributions, from which it is easy and fast to simulate. Finally, the accept/reject step is added to justify the sampling algorithm as an independent Metropolis-Hastings (MH) method.

To assess the efficiency of the sampling algorithm based on the proposed augmentation, we evaluate the upper and lower bounds of the approximation error and show that, in many cases, the acceptance probability is close to one. Due to its simplicity, our augmentation scheme can be applied directly to many models with reciprocal gamma functions, including the Student's t-distribution, Dirichlet-multinomial distribution, negative binomial distribution and Wishart distribution.

The remainder of the paper is organized as follows. In Section 2, we develop a new data augmentation and approximation of the reciprocal gamma function and illustrate our approach using a simple gamma model. For simplicity, we consider only proper priors for shape parameters as well as other variables, which ensures that full conditional distributions are always proper. In Section 3, we use our approach for a model based on Student's *t*-distribution. In Section 4, we consider a Dirichlet-multinomial model and apply a generic method. Some concluding remarks are given in Section 5. Proofs and additional results are provided in the Supplementary Material.

2 Beta Data Augmentation

2.1 General ideas

The most important result for our method is the following integral expression, which is based on Gauss's multiplication formula for the gamma function. **Theorem 1.** Let $m \in \mathbb{N}$. Then we have

$$\frac{1}{\{\Gamma(\xi)\}^m} = C_m \frac{1}{\xi^{m\xi}} \xi^{m+1/2-1} e^{m\xi} \Big\{ \prod_{j=2}^m \int_0^1 \rho_j^{\xi+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j \Big\} \frac{(m\xi)^{m\xi-1/2}}{\Gamma(m\xi)e^{m\xi}},$$

for all $\xi > 0$, where $C_m = 1/\{(2\pi)^{(m-1)/2} \prod_{j=2}^m \Gamma((m-j+1)/m)\}.$

The proof is given in the Supplementary Material. By Theorem 1, we can rewrite the *m*th power of the reciprocal gamma function, $1/{\{\Gamma(\xi)\}}^m$, by using integrals of m-1 beta densities, such that the reciprocal gamma function appears only once in the right-hand side.

Suppose that the target distribution, or the posterior distribution, is the joint density of shape parameter ξ and other variables ϑ of the form,

$$p(\xi, \boldsymbol{\vartheta}) \propto f(\xi, \boldsymbol{\vartheta}) \frac{1}{\{\Gamma(\xi)\}^m},$$

where typically $f(\xi, \vartheta) \stackrel{\xi}{\propto} \operatorname{Ga}(\xi|a_1, b_1)$ for some $a_1, b_1 > 0$. This framework covers, for example, the case of n independent observations from a gamma distribution, $x_1, \ldots, x_n \sim \operatorname{Ga}(\alpha, \beta)$; in this case, m = n, $(\xi, \vartheta) = (\alpha, \beta)$, and $p(\alpha, \beta)$ is the posterior of (α, β) given x_1, \ldots, x_n , or $p(\alpha, \beta) \propto \pi(\alpha, \beta) \times \beta^{n\alpha} (\prod_{i=1}^n x_i)^{\alpha} e^{-\beta \sum_{i=1}^n x_i}$, where $\pi(\alpha, \beta)$ is a prior density (see Section 2.2). In general, some of the variables ϑ may be latent variables introduced based on data augmentation. We are interested in the repeated sampling from the conditional distributions, $p(\xi|\vartheta)$ and $p(\vartheta|\xi)$. We assume that it is relatively easy to sample ϑ from $p(\vartheta|\xi)$, and we focus on the problem of sampling ξ from $p(\xi|\vartheta)$ in the following.

The derivation of the augmented model is a three-step process. First, we rewrite $p(\xi, \vartheta)$ as

$$p(\xi, \boldsymbol{\vartheta}) \propto \frac{f(\xi, \boldsymbol{\vartheta})}{\xi^{m\xi}} \xi^{m+1/2-1} e^{m\xi} \Big\{ \prod_{j=2}^{m} \int_{0}^{1} \rho_{j}^{\xi+(j-1)/m-1} (1-\rho_{j})^{(m-j+1)/m-1} d\rho_{j} \Big\} \frac{(m\xi)^{m\xi-1/2}}{\Gamma(m\xi) e^{m\xi}},$$

by using Theorem 1. The *m*th power, $1/{\{\Gamma(\xi)\}}^m$, is simplified to a single reciprocal gamma function, $1/\Gamma(m\xi)$, which we further evaluate in the following steps. A set of additional latent variables, $\rho = (\rho_2, \ldots, \rho_m) \in (0, 1)^{m-1}$, has the full conditional of the simple form, $\prod_{j=2}^m \text{Beta}(\rho_j|\xi + (j-1)/m, (m-j+1)/m)$, from which we can easily sample.

Second, the conditional density of (ξ, ϑ) given ρ is

$$p(\xi, \boldsymbol{\vartheta} | \boldsymbol{\rho}) \propto \frac{f(\xi, \boldsymbol{\vartheta})}{\xi^{m\xi}} \xi^{a_2 - 1} e^{-b_2 \xi} \frac{(m\xi)^{m\xi - 1/2}}{\Gamma(m\xi) e^{m\xi}},$$

where $a_2 = m+1/2$ and $b_2 = -m + \sum_{j=2}^m \log(1/\rho_j)$. In the above expression, there are two factors that make it difficult to sample ξ from the full conditional: $1/\xi^{m\xi}$ and $(m\xi)^{m\xi-1/2}/\{\Gamma(m\xi)e^{m\xi}\}$. Here, in order to eliminate $1/\xi^{m\xi}$, we assume that we can make the change of variables $\tilde{\boldsymbol{\vartheta}} = \varphi(\boldsymbol{\vartheta};\xi)$ with Jacobian $\xi^{m\xi}$, so that $f(\xi,\boldsymbol{\vartheta})d(\xi,\boldsymbol{\vartheta}) = \xi^{m\xi}\tilde{f}(\xi,\tilde{\boldsymbol{\vartheta}})d(\xi,\tilde{\boldsymbol{\vartheta}})$, and the density of interest becomes

$$p(\xi, \widetilde{\boldsymbol{\vartheta}} | \boldsymbol{\rho}) \propto \widetilde{f}(\xi, \widetilde{\boldsymbol{\vartheta}}) \xi^{a_2 - 1} e^{-b_2 \xi} \frac{(m\xi)^{m\xi - 1/2}}{\Gamma(m\xi) e^{m\xi}}$$

This change-of-variable is available for many models, including the gamma model of Section 2.2. The models for which there is no such change-of-variable, including the Dirichlet-multinomial model of Section 4, are discussed in Section 2.3.

Third, we use the above expression to construct an independent MH algorithm. Let ξ^{old} be a current value of ξ . To generate a new value ξ^{new} , we first sample a proposal ξ^* from the approximate full conditional density proportional to $\tilde{f}(\xi, \tilde{\vartheta})\xi^{a_2-1}e^{-b_2\xi}$ and compute

$$p = \min \Big\{ 1, \frac{(m\xi^*)^{m\xi^* - 1/2}}{\Gamma(m\xi^*)e^{m\xi^*}} / \frac{(m\xi^{\text{old}})^{m\xi^{\text{old}} - 1/2}}{\Gamma(m\xi^{\text{old}})e^{m\xi^{\text{old}}}} \Big\}.$$

Then we set $\xi^{\text{new}} = \xi^*$ with probability p, otherwise $\xi^{\text{new}} = \xi^{\text{old}}$. We note that in all the models considered in this paper, proposal distributions corresponding to $\tilde{f}(\xi, \tilde{\vartheta})\xi^{a_2-1}e^{-b_2\xi}$ are easy to sample from. The factor dropped in the approximate distribution can be evaluated as

$$\frac{e^{-1/(12\xi)}}{(2\pi)^{1/2}} < \frac{\xi^{\xi-1/2}}{\Gamma(\xi)e^{\xi}} < \frac{1}{(2\pi)^{1/2}},\tag{1}$$

for any $\xi > 0$ by Stirling's formula. This expression shows that the factor is almost constant when ξ is not extremely small, and that the acceptance probability p is close to one. This can be confirmed by bounding the acceptance probability below as $p \ge e^{-1/(12m\xi^*)} \ge 1 - 1/(12m\xi^*)$, where the lower bound is almost unity unless ξ^* is extremely small.

2.2 An illustration using a gamma model

Here, we consider a simple gamma model for illustration. For this model, several methods for posterior inference are available (e.g. Gilks and Wild 1992). In particular, the method of Miller (2019) is customized for this model and highly efficient.

Suppose that observations $\boldsymbol{x} = (x_1, \ldots, x_n)$ have been independently generated from a gamma distribution $Ga(\alpha, \beta)$. We assume the independent gamma prior distributions for α and β : Ga(a, b) and Ga(c, d), respectively. Then the posterior of (α, β) is

$$p(\alpha,\beta|\mathbf{x}) \propto \operatorname{Ga}(\alpha|a,b)\beta^{c-1}e^{-d\beta}\frac{\beta^{n\alpha}}{\{\Gamma(\alpha)\}^n} \Big(\prod_{i=1}^n x_i\Big)^{\alpha}e^{-\beta\sum_{i=1}^n x_i}.$$

Using Theorem 1, we can rewrite the above posterior density as

$$p(\alpha,\beta|\boldsymbol{x}) \propto \operatorname{Ga}(\alpha|a,b)\beta^{c-1}e^{-d\beta}\beta^{n\alpha} \Big(\prod_{i=1}^{n} x_i\Big)^{\alpha} e^{-\beta\sum_{i=1}^{n} x_i} \\ \times \frac{1}{\alpha^{n\alpha}}\alpha^{n+1/2-1}e^{n\alpha} \Big\{\prod_{i=2}^{n} \int_0^1 \rho_i^{\alpha+(i-1)/n-1} (1-\rho_i)^{(n-i+1)/n-1} d\rho_i\Big\} \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}}.$$

Now, we consider $\boldsymbol{\rho} = (\rho_2, \dots, \rho_n) \in (0, 1)^{n-1}$ as a set of additional latent variables. Then the conditional distribution of $(\alpha, \beta, \boldsymbol{\rho})$ given \boldsymbol{x} is

$$p(\alpha, \beta, \boldsymbol{\rho} | \boldsymbol{x}) \propto \operatorname{Ga}(\alpha | a, b) \beta^{c-1} e^{-d\beta} \beta^{n\alpha} \Big(\prod_{i=1}^{n} x_i \Big)^{\alpha} e^{-\beta \sum_{i=1}^{n} x_i} \\ \times \frac{1}{\alpha^{n\alpha}} \alpha^{n+1/2-1} e^{n\alpha} \Big[\prod_{i=2}^{n} \{ \rho_i^{\alpha+(i-1)/n-1} (1-\rho_i)^{(n-i+1)/n-1} \} \Big] \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha) e^{n\alpha}}.$$

In order to obtain MCMC samples of $(\alpha, \beta, \rho)|\mathbf{x}$, we can use the MH within Gibbs sampler. It is easy to sample ρ from its full conditional distribution since $p(\rho|\alpha, \beta, \mathbf{x}) = \prod_{i=2}^{n} \text{Beta}(\rho_i|\alpha + (i-1)/n, (n-i+1)/n)$. Meanwhile, the full conditional of (α, β) is

$$p(\alpha,\beta|\boldsymbol{\rho},\boldsymbol{x}) \propto \frac{1}{\alpha^{n\alpha}} \alpha^{n-1/2+a-1} \exp\left\{-\alpha\left(-\sum_{i=1}^{n}\log x_{i} + \sum_{i=2}^{n}\log\frac{1}{\rho_{i}} - n + b\right)\right\}$$
$$\times \beta^{n\alpha+c-1} \exp\left\{-\beta\left(\sum_{i=1}^{n}x_{i} + d\right)\right\} \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}}.$$

Although the full conditional of β is a gamma distribution, the full conditional density of α does not have a standard form because of the two factors: $g_1(\alpha) = 1/\alpha^{n\alpha}$ and $g_2(\alpha) = (n\alpha)^{n\alpha-1/2}/{\{\Gamma(n\alpha)e^{n\alpha}\}}$.

First, in order to eliminate $g_1(\alpha)$ from the above expression, we make the change of variables $\gamma = \beta/\alpha$. Then

$$p(\alpha, \gamma | \boldsymbol{\rho}, \boldsymbol{x}) \propto \alpha^{n-1/2+c+a-1} \exp\left\{-\alpha \left(-\sum_{i=1}^{n} \log x_i + \sum_{i=2}^{n} \log \frac{1}{\rho_i} - n + b\right)\right\}$$
$$\times \gamma^{n\alpha+c-1} \exp\left\{-\alpha \gamma \left(\sum_{i=1}^{n} x_i + d\right)\right\} g_2(\alpha).$$

The full conditional of $\gamma = \beta/\alpha$ is given by $\operatorname{Ga}(\gamma | n\alpha + c, \alpha (\sum_{i=1}^{n} x_i + d))$ and tractable similar to that of the original parameter β .

Next, we use the MH algorithm to update α . The full conditional density of α is given by $p(\alpha|\gamma, \boldsymbol{\rho}, \boldsymbol{x}) \propto \operatorname{Ga}(\alpha|A, B)g_2(\alpha)$, where A = n - 1/2 + c + a and $B = -\sum_{i=1}^n \log x_i + \sum_{i=2}^n \log(1/\rho_i) - n - n \log \gamma + \gamma (\sum_{i=1}^n x_i + d) + b$. We sample a proposal α^* from $\operatorname{Ga}(\alpha|A, B)$. We accept α^* if an independent standard uniform variable U is less than or equal to $g_2(\alpha^*)/g_2(\alpha^{\text{old}})$, where α^{old} denotes the current value of α . The new value of α , or α^{new} , is set to α^* if α^* is accepted, and to α^{old} otherwise.

The MH within Gibbs sampler is summarized as follows.

Algorithm 1. The variables α , γ , and ρ are updated in the following way.

- Sample $\gamma^* \sim \operatorname{Ga}(n\alpha + c, \alpha(\sum_{i=1}^n x_i + d)).$
- Sample $\rho^* = (\rho_2^*, \dots, \rho_n^*) \sim \prod_{i=2}^n \text{Beta}(\alpha + (i-1)/n, (n-i+1)/n).$
- Sample $\alpha^* \sim \operatorname{Ga}(A, B)$, where A = n 1/2 + c + a and

$$B = -\sum_{i=1}^{n} \log x_i + \sum_{i=2}^{n} \log \frac{1}{\rho_i^*} - n - n \log \gamma^* + \gamma^* \left(\sum_{i=1}^{n} x_i + d\right) + b,$$

and accept α^* with probability

$$\min\left\{1,\frac{(n\alpha^*)^{n\alpha^*-1/2}}{\Gamma(n\alpha^*)e^{n\alpha^*}}/\frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}}\right\}$$

The accuracy of approximation, or the acceptance probability, has already been evaluated in (1). The acceptance probability is, at least, $1 - 1/(12n\alpha^*)$.

2.3 PTN data augmentation

The key to the augmentation strategy of Section 2.1 is to find suitable changes of variables $\tilde{\vartheta} = \varphi(\vartheta; \xi)$ to eliminate the factor $1/\xi^{m\xi}$ in the second step. Because this is not always straightforward, an alternative method is developed in this section. We modify the proposed method of Section 2.1 by introducing additional latent variables. The main tool is the integral expression in the following lemma.

Lemma 1. Let $m \in \mathbb{N}$. Then

$$\frac{1}{\xi^{m\xi}} = (m\xi)^{1/2} e^{m\xi} \frac{(m\xi)^{m\xi-1/2}}{\Gamma(m\xi)e^{m\xi}} \int_0^\infty w^{m\xi-1} e^{-wm\xi^2} dw$$

for all $\xi > 0$.

We assume that $f(\xi, \vartheta) \stackrel{\xi}{\propto} \operatorname{Ga}(\xi|a_1, b_1)$ for simplicity and consider the conditional density

$$p(\xi|\boldsymbol{
ho},\boldsymbol{\vartheta}) \propto \frac{1}{\xi^{m\xi}} \xi^{a_3-1} e^{-b_3\xi} \frac{(m\xi)^{m\xi-1/2}}{\Gamma(m\xi)e^{m\xi}},$$

where $a_3 = a_1 + m - 1/2$ and $b_3 = b_1 - m + \sum_{j=2}^m \log(1/\rho_j)$. Using Lemma 1, we see that $p(\xi|\boldsymbol{\rho},\boldsymbol{\vartheta})$ is the marginal density of

$$p(\xi, w | \boldsymbol{\rho}, \boldsymbol{\vartheta}) \propto \xi^{a_3 - 1/2} e^{-(b_3 - m)\xi} \left\{ \frac{(m\xi)^{m\xi - 1/2}}{\Gamma(m\xi) e^{m\xi}} \right\}^2 w^{m\xi - 1} e^{-wm\xi^2},$$

where $w \in (0, \infty)$ is an additional latent variable. Clearly, $p(w|\xi, \rho, \vartheta) = \text{Ga}(w|m\xi, m\xi^2)$. On the other hand,

$$p(\xi|w,\boldsymbol{\rho},\boldsymbol{\vartheta}) / \left\{ \frac{(m\xi)^{m\xi-1/2}}{\Gamma(m\xi)e^{m\xi}} \right\}^2 \propto \xi^{c-1} e^{-a\xi^2 + b\xi}, \tag{2}$$

where $c = a_3 + 1/2$, a = mw, and $b = m \log w + m - b_3$. The right-hand side is proportional to the power truncated normal (PTN) distribution (He et al., 2021) with parameters c, a, and b, which is denoted by PTN(c, a, b). Since the denominator of the left-hand side in (2) is almost constant as seen in (1), the conditional density $p(\xi|w, \rho, \vartheta)$ is approximated by $PTN(\xi|c, a, b)$. Then, we generate a proposal, $\xi^* \sim PTN(c, a, b)$, and accept it with probability

$$\min\left\{1, \left\{\frac{(m\xi^*)^{m\xi^*-1/2}}{\Gamma(m\xi^*)e^{m\xi^*}} / \frac{(m\xi^{\text{old}})^{m\xi^{\text{old}}-1/2}}{\Gamma(m\xi^{\text{old}})e^{m\xi^{\text{old}}}}\right\}^2\right\},\$$

where ξ^{old} is the current state of ξ .

2.4 Additional data augmentation for the PTN distribution

In order to sample from the PTN distribution (2), one can use the accept/reject algorithm described by He et al. (2021). In this paper, we consider other approaches so that we do not necessarily need to use accept/reject algorithms. Our approaches also have potential flexibility that they are easily extended to the case where $f(\xi, \vartheta)$ is proportional to a generalized-inverse-Gaussian density as a function of ξ .

Let M > 0 be a constant possibly dependent on w, ρ , and ϑ such that M > b. (A convenient choice is $M = 1 + \max\{0, b\}$.) Then, the PTN density is written as

$$PTN(\xi|c,a,b) \propto \xi^{c-1} e^{-a\xi^2 - b'\xi} e^{M\xi},$$

where c, a, b' = M - b, and M are all positive.

The exponential term $e^{M\xi}$ can be augmented in two ways. The first approach is based on the following expression:

$$\begin{aligned} \operatorname{PTN}(\xi|c,a,b) &\propto \xi^{c-1} e^{-a\xi^2 - b'\xi} \sum_{\zeta=0}^{\infty} \frac{M^{\zeta}\xi^{\zeta}}{\zeta!} \\ &= \sum_{\zeta=0}^{\infty} \frac{M^{\zeta}\xi^{\zeta}}{\zeta!} \xi^{c-1} e^{-a\xi^2} \int_0^{\infty} \frac{1}{\sqrt{2\pi}} \eta^{1/2-1} e^{-\eta/2} e^{-(b')^2\xi^2/(2\eta)} d\eta \end{aligned}$$

where we consider $\zeta \in \mathbb{N}_0 = \{0, 1, 2, ...\}$ and $\eta \in (0, \infty)$ as additional latent variables. Then the full conditional distributions of ζ and η are $\operatorname{Po}(\zeta | M\xi)$ and $\operatorname{GIG}(\eta | 1/2, 1, (b')^2 \xi^2)$, respectively. The full conditional density of ξ divided by $\{(m\xi)^{m\xi-1/2}/\Gamma(m\xi)e^{m\xi}\}^2$ is proportional to

$$\xi^{\zeta+c-1}e^{-\{a+(b')^2/(2\eta)\}\xi^2}.$$

We can easily sample from the above distribution since it is simply the square root of a gamma variable.

The second approach utilizes the integral expression based on the normal density.

Lemma 2. For all $\xi > 0$, we have

$$e^{\xi} = \int_{-\infty}^{\infty} \frac{1}{\sqrt{2\pi}} \frac{1}{\sqrt{2\xi}} e^{-\theta^2/(4\xi) + \theta} d\theta$$

By this lemma, we have

$$\begin{aligned} \operatorname{PTN}(\xi|c,a,b) &\propto \xi^{c-1} e^{-a\xi^2 - b'\xi} \int_{-\infty}^{\infty} \xi^{1/2 - 1} e^{-\theta^2/(4M\xi) + \theta} d\theta \\ &\propto \int_{-\infty}^{\infty} e^{\theta} \xi^{c-1/2 - 1} e^{-a\xi^2} \Big[\int_{0}^{\infty} \eta^{1/2 - 1} e^{-\eta/2} e^{-\{b'\xi + \theta^2/(4M\xi)\}^2/(2\eta)} d\eta \Big] d\theta, \end{aligned}$$

where we consider $\theta \in (-\infty, \infty)$ and $\eta \in (0, \infty)$ as additional latent variables. Sampling from the full conditional of (η, θ) can be done in a compositional way; we sample θ (with η marginalized out) from N($\theta|2M\xi, 2M\xi$), then $(\eta|\theta)$ from GIG $(\eta|1/2, 1, \{b'\xi + \theta^2/(4M\xi)\}^2)$. The full conditional density of ξ divided by $\{(m\xi)^{m\xi-1/2}/\Gamma(m\xi)e^{m\xi}\}^2$ is proportional to

$$\xi^{c-1/2-1}e^{-\{a+(b')^2/(2\eta)\}\xi^2}e^{-\{\theta^4/(32M^2\eta)\}/\xi^2},$$

which is the square root of a generalized-inverse-Gaussian distribution.

3 Student's *t*-Distribution

3.1 Sampling algorithm

Student's t-distribution is widely adopted in Bayesian inference to handle outliers in samples or heavy-tailed properties of data generating processes (e.g. Geweke, 1993; Fonseca et al., 2008; Villa and Rubio, 2018; da Silva et al., 2020). A typical problem in using Student's t-distribution is that the posterior inference of the degrees of freedom is not straightforward since its full conditional distribution has a complicated form. However, we can use our data-augmentation approach. We here consider the simplest case where the means of all observations are the same, and we use the normal-scale-mixture representation of Student's t-distribution, under which the degrees-of-freedom parameter is regarded as the shape parameter in the gamma distribution.

Suppose that for $i = 1, \ldots, n$,

$$x_i \sim t(x_i | (\theta, \tau), 2\alpha) = \frac{\Gamma(\alpha + 1/2)}{\sqrt{2\pi\tau^{1/2}\alpha^{1/2}\Gamma(\alpha)}} / \left\{ 1 + \frac{(x_i - \theta)^2}{2\tau\alpha} \right\}^{\alpha + 1/2},$$

$$(\theta, \tau) \sim p(\theta, \tau), \qquad \alpha \sim p(\alpha),$$

where $x_i \in \mathbb{R}$, $\theta \in \mathbb{R}$, and $\tau \in (0, \infty)$. Then the posterior distribution $p(\theta, \tau, \alpha | \boldsymbol{x})$ is obtained as the marginal distribution of

$$p(\theta, \tau, \alpha, \boldsymbol{w}, \boldsymbol{\rho} | \boldsymbol{x}) \propto \frac{p(\theta, \tau)}{\tau^{n/2}} p(\alpha) \alpha^{n+1/2-1} e^{n\alpha} \Big(\prod_{i=1}^{n} [w_i^{\alpha+1/2-1} e^{-w_i \{\alpha + (x_i - \theta)^2/(2\tau)\}}] \Big) \\ \times \Big[\prod_{i=2}^{n} \{\rho_i^{\alpha + (i-1)/n - 1} (1 - \rho_i)^{(n-i+1)/n - 1} \} \Big] \frac{(n\alpha)^{n\alpha - 1/2}}{\Gamma(n\alpha) e^{n\alpha}},$$
(3)

where $\boldsymbol{w} = (w_1, \ldots, w_n) \in (0, \infty)^n$ and $\boldsymbol{\rho} = (\rho_2, \ldots, \rho_n) \in (0, 1)^{n-1}$ are additional latent variables. The above expression is derived in Section S6 of the Supplementary Material by using Theorem 1.

If we use the priors $p(\theta, \tau) = N(\theta|b, \tau/a)IG(\tau|c, d)$ and $p(\alpha) = Ga(\alpha|a_0, b_0)$ for $a, c, d \in (0, \infty)$ and $b \in \mathbb{R}$ and $a_0, b_0 \in (0, \infty)$, we can use the following algorithm to generate posterior samples.

Algorithm 2. The variables θ , τ , α , w, and ρ are updated in the following way.

- Sample $\tau^* \sim \text{IG}(c', d')$, where c' = n/2 + c and

$$d' = \frac{1}{2} \left\{ ab^2 + \sum_{i=1}^n w_i x_i^2 - \frac{\left(ab + \sum_{i=1}^n w_i x_i\right)^2}{a + \sum_{i=1}^n w_i} \right\} + d.$$

- Sample $\theta^* \sim N(b', \tau^*/a')$, where $a' = a + \sum_{i=1}^n w_i$ and

$$b' = \frac{ab + \sum_{i=1}^{n} w_i x_i}{a + \sum_{i=1}^{n} w_i}$$

- Sample $\mathbf{w}^* = (w_1^*, \dots, w_n^*) \sim \prod_{i=1}^n \operatorname{Ga}(\alpha + 1/2, \alpha + (x_i \theta^*)^2/(2\tau^*)).$
- Sample $\rho^* = (\rho_2^*, \dots, \rho_n^*) \sim \prod_{i=2}^n \text{Beta}(\alpha + (i-1)/n, (n-i+1)/n).$

- Sample $\alpha^* \sim \text{Ga}(a_0', b_0')$, where $a_0' = a_0 + n - 1/2$ and

$$b_0' = b_0 - n + \sum_{i=1}^n (w_i^* - \log w_i^*) + \sum_{i=2}^n \log \frac{1}{\rho_i^*},$$

and accept α^* with probability

$$\min\left\{1,\frac{(n\alpha^*)^{n\alpha^*-1/2}}{\Gamma(n\alpha^*)e^{n\alpha^*}}/\frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}}\right\}.$$

Since we introduce the additional latent variables ρ_2, \ldots, ρ_n , our method is less efficient than an alternative method in terms of the effective sample size for an MCMC sequence of a fixed number of parameter values. However, since we do not need to use numerical approximation, our method takes less time. These are confirmed in Section 3.2.

We remark that our method is flexible and we can use many other types of priors. For example, we can use a scale mixture of gamma distributions as a prior for α . We can use a truncated gamma prior for α and this case is considered in the second half of Section 3.2. Also, for $a_0, b_0, c_0 \in (0, \infty)$, we can use the beta-type prior $p(\alpha) \propto \alpha^{a_0-1}(1-\alpha/c_0)^{b_0-1}\chi_{(0,c_0)}(\alpha)$.

3.2 Simulation study

Here, we compare the performance of our method based on data augmentation (DA) with the performance of an alternative method based on the approximation proposed by Miller (2019) (A-MH). See Section S7 of the Supplementary Material for details of the A-MH method.

First, we set either n = 10, n = 30, or n = 100 and use the conjugate prior $p(\theta, \tau) = N(\theta|0, \tau/(1/10)) \times IG(\tau|1/10, 1/10)$ and the gamma prior $p(\alpha) = Ga(\alpha|1/10, 1/10)$. We generate x_i from $t(x_i|(3, 1), 2\alpha_0)$. We consider the cases $2\alpha_0 = 1/10$, $2\alpha_0 = 1$, and $2\alpha_0 = 10$. Then, for each of the two methods, we generate 4,000 posterior samples after discarding the first 1,000 samples. We use $(\varepsilon, M) = (10^{-8}, 10)$ for the convergence tolerance and the maximum number of iterations for the A-MH method as recommended in Miller (2019). We repeat this simulation 100 times.

Boxplots of the ratios of the effective sample sizes for α , τ , and θ to the computation times for the two methods are shown in Figure 1 for n = 10. (For the boxplots for n = 30 and n = 100, see Figures S1 and S2 of the Supplementary Material.) Table 1 reports the averages over the simulations of the ratios (sESS) and the original effective sample sizes (ESS), as well as the mean squared error (MSE) ratios of the estimators of α , τ , and θ , where the MSE ratio is defined as the MSE of the alternative method divided by that of our proposed method. In terms of MSE, there is little difference between the two methods in many cases including those in the Supplementary Material. In terms of sESS, our method is better especially for θ and τ when n = 10. When n = 30, the alternative method becomes better in terms of α and competitive in terms of τ and θ . When n = 100, the alternative method is clearly better than ours. This increase of sESS of the DA method for large n is most likely due to the increased number of latent parameters $\rho_{2:n}$, affecting both efficiency and computational time. For example, the ESSs of center parameter θ are almost unchanged (or even improve) when n increases from 30 to 100, hence the decrease of the sESSs for θ is mainly due to the increased computational time.

Thus, when n is large, our method benefits rather from its simplicity and applicability to more complicated models. To see this point, we consider additional scenarios where a truncated



Figure 1: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for n = 10.

Table 1: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH) by Miller (2019), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method.

				ESS			sESS		М	MSE ratio		
n	$2\alpha_0$	method	θ	au	α	θ	au	α	θ	au	α	
10	0.1	DA	874	382	1972	7148	3121	16058	_	_	_	
10	0.1	A-MH	805	361	2097	5887	2643	15408	1.14	8.87	1.06	
10	1	DA	1506	686	466	12749	5815	3926	—	—	_	
10	1	A-MH	1514	755	581	11252	5648	4285	1.00	1.05	0.93	
10	10	DA	2276	1862	288	19406	15875	2458	—	—	_	
10	10	A-MH	2316	1989	505	17634	15193	3854	1.00	1.01	0.84	
30	0.1	DA	796	188	2068	4836	1135	12525	_	_	_	
30	0.1	A-MH	834	178	2103	5115	1095	12914	1.10	0.79	1.02	
30	1	DA	1408	444	510	9019	2837	3252	—	—	_	
30	1	A-MH	1440	473	645	9114	2991	4072	1.00	1.00	1.00	
30	10	DA	2371	715	122	15235	4570	782	_	_	_	
30	10	A-MH	2448	921	220	16451	6235	1479	1.02	1.02	1.03	
100	0.1	DA	904	109	1770	3200	384	6270	_	-	_	
100	0.1	A-MH	895	105	1802	4040	472	8120	1.00	0.77	1.00	
100	1	DA	1359	386	527	4889	1388	1888	_	_	_	
100	1	A-MH	1364	391	640	6251	1794	2926	1.02	1.01	1.03	
100	10	DA	2711	284	56	10380	1093	214	_	_	_	
100	10	A-MH	2721	407	100	13793	2075	507	1.01	1.04	0.98	

gamma prior is used for the shape parameter; $p(\alpha) \propto \text{Ga}(\alpha|1/10, 1/10)\chi_{(\alpha,\infty)}(\alpha)$, where $\alpha > 0$. With this truncated priors, the method of Miller (2019) must evaluate the expected values of truncated gamma distributions, taking longer time for posterior computation. In contrast, no complication is needed for our method to use the truncated prior, except that we now need to sample from truncated distributions. We set $2\alpha_0 = 10$ and conduct the same simulation study for the truncated gamma prior with $2\alpha = 1, 3$.

Boxplots of sESSs are shown in Figure 2 for n = 10 (and in Figures S3 and S4 for n = 30 and n = 100, respectively), and Table 2 lists the averages of ESSs, sESSs and the ratios of MSEs computed in this experiment. In these scenarios, our method becomes more competitive even for large n. In particular, our method outperforms the A-MH method in terms of sESS for θ and τ when n = 10 and n = 30, and for θ when n = 100.

4 The Dirichlet-Multinomial Distribution

4.1 Sampling algorithm

Dirichlet-multinomial distribution is useful for modeling multi-label variables, as used in topic modeling (e.g. Blei et al., 2003). Since the full conditional distribution of the shape parameters of the Dirichlet distribution includes the reciprocal gamma function, their posterior sampling is



Figure 2: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for $2\underline{\alpha} = 1,3$ for n = 10.

Table 2: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method for $2\alpha = 1, 3$.

				ESS			sESS			MSE ratio			
n	$2\underline{\alpha}$	method	θ	au	α	θ	au	α	θ	au	α		
10	1	DA	2376	2033	293	7029	6013	867	_	_	_		
10	1	A-MH	2371	2092	533	3866	3419	868	1.01	0.98	0.96		
10	3	DA	2885	2735	396	8963	8494	1230	—	-	-		
10	3	A-MH	2893	2761	695	4877	4660	1171	1.00	1.00	0.99		
30	1	DA	2341	601	126	6354	1627	342	_	_	_		
30	1	A-MH	2371	767	217	4074	1325	373	1.01	0.99	0.89		
30	3	DA	2710	1057	141	7376	2871	383	—	-	-		
30	3	A-MH	2702	1206	251	4567	2051	424	1.00	1.03	1.03		
100	1	DA	2660	247	59	5595	521	123	_	_	_		
100	1	A-MH	2678	346	106	4278	552	168	0.99	1.00	0.94		
100	3	DA	2679	283	63	5670	600	134	_	-	-		
100	3	A-MH	2719	381	108	4300	604	170	1.00	1.03	1.04		

typically not straightforward (e.g. Nandram, 1998). Although in this section our focus is the estimation of the shape parameters of the Dirichlet-multinomial distribution, our result is also relevant in the context of finite mixture modeling (e.g. Frühwirth-Schnatter, 2006).

Suppose that for $i = 1, \ldots, n$,

$$\boldsymbol{x}_{i} \sim \operatorname{Multin}_{L}(\boldsymbol{x}_{i}|N_{i},\boldsymbol{p}_{i}) = \frac{N_{i}!}{\prod_{l=0}^{L}(x_{i,l}!)} \prod_{l=0}^{L} p_{i,l}^{x_{i,l}},$$
$$\boldsymbol{p}_{i} \sim \operatorname{Dir}_{L}(\boldsymbol{p}_{i}|\boldsymbol{\alpha}) = \frac{\Gamma\left(\sum_{l=0}^{L} \alpha_{l}\right)}{\prod_{l=0}^{L} \Gamma(\alpha_{l})} \prod_{l=0}^{L} p_{i,l}^{\alpha_{l}-1},$$
$$\boldsymbol{\alpha} \sim p(\boldsymbol{\alpha}),$$

where $\boldsymbol{x}_i = (x_{i,1}, \ldots, x_{i,L}), \ x_{i,0} = N_i - \sum_{l=1}^L x_{i,l}, \ \boldsymbol{p}_i = (p_{i,1}, \ldots, p_{i,L}) \in D_L = \{(\tilde{p}_1, \ldots, \tilde{p}_L) \in (0,1)^L | \tilde{p}_1, \ldots, \tilde{p}_L > 0, \sum_{l=1}^L \tilde{p}_l < 1\}, \ p_{i,0} = 1 - \sum_{l=1}^L p_{i,l}, \ \text{and} \ \boldsymbol{\alpha} = (\alpha_0, \ldots, \alpha_L).$ Let $\boldsymbol{x} = (\boldsymbol{x}_1, \ldots, \boldsymbol{x}_n)$ and $\boldsymbol{p} = (\boldsymbol{p}_1, \ldots, \boldsymbol{p}_n)$. Since we have been unable to find good changes of variables to perform the second step of Section 2.1, we use the flexible method of Section 2.3. In Section S6 of the Supplementary Material, we prove that the posterior distribution $p(\boldsymbol{p}, \boldsymbol{\alpha} | \boldsymbol{x})$ is obtained

as the marginal distribution of

$$p(\boldsymbol{p}, \boldsymbol{\alpha}, \boldsymbol{z}, \boldsymbol{w}, \boldsymbol{\rho} | \boldsymbol{x}) \propto p(\boldsymbol{\alpha}) \Big\{ \prod_{l=0}^{L} (\alpha_l^n e^{2n\alpha_l}) \Big\} \Big(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l} x_{i,l} + \alpha_l - 1 \Big) \Big\{ \prod_{i=1}^{n} (z_i \sum_{l=0}^{L} \alpha_l - 1 e^{-z_i}) \Big\} \\ \times \Big\{ \prod_{l=0}^{L} (w_l^{n\alpha_l - 1} e^{-w_l n\alpha_l^2}) \Big\} \Big[\prod_{i=2}^{n} \prod_{l=0}^{L} \{\rho_{i,l} \alpha_l + (i-1)/n - 1 (1 - \rho_{i,l})^{(n-i+1)/n - 1} \} \Big] \\ \times \prod_{l=0}^{L} \Big\{ \frac{(n\alpha_l)^{n\alpha_l - 1/2}}{\Gamma(n\alpha_l) e^{n\alpha_l}} \Big\}^2,$$
(4)

where $\boldsymbol{z} = (z_1, \dots, z_n) \in (0, \infty)^n$, $\boldsymbol{w} = (w_0, \dots, w_L) \in (0, \infty)^{L+1}$, and $\boldsymbol{\rho} = (\boldsymbol{\rho}_2, \dots, \boldsymbol{\rho}_n) = ((\rho_{2,0}, \dots, \rho_{2,L}), \dots, (\rho_{n,0}, \dots, \rho_{n,L})) \in (0, 1)^{(n-1)(L+1)}$ are additional latent variables.

If we use the prior $p(\boldsymbol{\alpha}) = \prod_{l=0}^{L} \operatorname{Ga}(\alpha_{l}|a, b)$ for example, we can use the following algorithm to generate posterior samples.

Algorithm 3. The variables p, α , z, w, and ρ are updated in the following way.

- Sample $p^* = ((p_{1,0}^*, \dots, p_{1,L}^*), \dots, (p_{n,0}^*, \dots, p_{n,L}^*)) \sim \prod_{i=1}^n \operatorname{Dir}_L(x_i + \alpha).$
- Sample $\mathbf{z}^* = (z_1^*, \dots, z_n^*) \sim \{ \operatorname{Ga}(\sum_{l=0}^L \alpha_l, 1) \}^n$.
- Sample $\boldsymbol{w}^* = (w_0^*, \dots, w_L^*) \sim \prod_{l=0}^L \operatorname{Ga}(n\alpha_l, n\alpha_l^2).$
- Sample $\boldsymbol{\rho}^* = ((\rho_{2,0}^*, \dots, \rho_{2,L}^*), \dots, (\rho_{n,0}^*, \dots, \rho_{n,L}^*)) \sim \prod_{i=2}^n \prod_{l=0}^L \operatorname{Beta}(\alpha_l + (i-1)/n, (n-i+1)/n).$
- For l = 0, ..., L, let $c_l = n + a$, $a_l = nw_l^*$, and

$$b_{l} = -\sum_{i=1}^{n} \log \frac{1}{p_{i,l}^{*}} + \sum_{i=1}^{n} \log z_{i}^{*} + 2n + n \log w_{l}^{*} - \sum_{i=2}^{n} \log \frac{1}{\rho_{i,l}^{*}} - b$$

and sample α_l^* in one of the following three ways and accept it with probability

$$\min\left\{1,\left\{\frac{(n\alpha_l^*)^{n\alpha_l^*-1/2}}{\Gamma(n\alpha_l^*)e^{n\alpha_l^*}}\right\}^2/\left\{\frac{(n\alpha_l)^{n\alpha_l-1/2}}{\Gamma(n\alpha_l)e^{n\alpha_l}}\right\}^2\right\}$$

- (i) Sample $\alpha_l^* \sim \text{PTN}(c_l, a_l, b_l)$ by using the PTN sampler developed by He et al. (2021).
- (ii) Let $M_l = 1 + \max\{0, b_l\}$ and $b_l' = M_l b_l$.
 - Sample $\zeta_l^* \sim \operatorname{Po}(M_l \alpha_l).$
 - Sample $\eta_l^* \sim \text{GIG}(1/2, 1, (b_l')^2 \alpha_l^2).$
 - Sample $\tilde{\alpha}_{l}^{*} \sim \text{Ga}((\zeta_{l}^{*} + c_{l})/2, a_{l} + (b_{l}')^{2}/(2\eta_{l}^{*}))$ and set $\alpha_{l}^{*} = (\tilde{\alpha}_{l}^{*})^{1/2}$.
- (iii) Let $M_l = 1 + \max\{0, b_l\}$ and $b_l' = M_l b_l$.
 - Sample $\theta_l^* \sim N(2M_l\alpha_l, 2M_l\alpha_l)$.
 - Sample $\eta_l^* \sim \text{GIG}(1/2, 1, \{b_l' \alpha_l + \theta_l^2/(4M_l \alpha_l)\}^2).$
 - $Sample \tilde{\alpha}_l^* \sim \text{GIG}(c_l/2 1/4, 2a_l + (b_l')^2/\eta_l^*, (\theta_l^*)^4/(16M_l^2\eta_l^*)) \text{ and set } \alpha_l^* = (\tilde{\alpha}_l^*)^{1/2}.$

4.2 Simulation study

In this section, we conduct a simulation study– the posterior inference of Dirichlet shape parameters– to compare our method and the method of He et al. (2021). Both methods are based on data augmentation but in different ways. Many other standard methods, including one by Miller (2019), are not directly applicable.

Following He et al. (2021), we set L + 1 = 10 and $N_1 = \cdots = N_n = 500$ and use the prior $p(\alpha) = \prod_{j=0}^{9} \operatorname{Ga}(\alpha_l | b/10, b)$ with b = 1. We generate p_i from $\operatorname{Dir}_9(\alpha_0)$ and then x_i from $\operatorname{Multin}_9(500, p_i)$. We consider the cases n = 100 and n = 1000. For each of these cases, we consider two scenarios: (I) $\alpha_0 = (1/10, \ldots, 1/10)$ (equal case) and (II) $\alpha_0 = (1/10, 2/10, \ldots, 10/10)$. Other scenarios are also considered and reported in the Supplementary Material. We generate 4,000 posterior samples after discarding the first 1,000 samples. We repeat this simulation 100 times. The method of He et al. (2021) requires sampling from the exponential reciprocal gamma (ERG) distribution, for which they gave three methods. We use the first method because it is the easiest to implement. Setting N equal to a large value in (16) of He et al. (2021) makes their approximation accurate. We set N = 3, so that their approximation is sufficiently accurate.

We consider the proposed method based on (iii), (ii) and (i) of Algorithm 3 (DA-N, DA-P and DA-PT, respectively), as well as the method of He et al. (2021) (ERG). Using these methods, we calculate the averages over the simulations of the means of the effective sample sizes for $\alpha_0, \ldots, \alpha_9$ (ESS), the averages over the simulations of the computation times (CT), and the averages over the simulations of the ratios of the means of the effective sample sizes to the computation times (sESS). We also calculate the mean squared errors (MSE) of the estimators of $\alpha_0, \ldots, \alpha_9$.

The results are reported in Table 3. In all scenarios, the ERG method has the largest ESS but the longest CT. In contrast, the DA-N, DA-P and DA-PT methods are less competitive than the ERG method in ESSs, but significantly outperform it in computational time. Consequently, all of our methods have much larger sESSs than the state-of-the-art ERG method. Among the proposed methods, the DA-PT method has the best sESS. The other two methods cost computational efficiency for the simplicity of their algorithms, as noted in Section 2.3. In terms of MSE, no significant difference can be seen in the four methods.

Thus, the difference of the method of He et al. (2021) and ours in computational efficiency critically depends on the computational time. For the fairness of comparison, it should be noted that the computation by the ERG method can speed-up by using parallelization, and could be competitive as our methods in some computational environments that enable such parallelization. Other than the efficiency, the advantage of our method to be emphasized is its simplicity; no explicit parallelization is needed in implementing our method. In addition, our method is tuning parameter free, while the ERG method requires tuning N.

5 Concluding Remarks

The data augmentation approach proposed in this paper can be applicable to any posterior inference if the conditional posterior involves the reciprocal gamma functions. Examples of such models include the one-parameter Dirichlet, negative binomial and Wishart models, in addition to the gamma, Student's t and Dirichlet-multinomial models considered in the previous sections. The sampling algorithms for those models can be derived straightforwardly and are provided in Section S1 of the Supplementary Material.

Table 3: The average effective sample size (ESS), the average computation time (CT), the standardized effective sample size by the computation time (sESS), and the mean squared error (MSE) for the proposed data-augmentation method with normal latent variables (DA-N), Poisson latent variables (DA-P), and the PTN sampler of He et al. (2021) (DA-PT) and the original method proposed by He et al. (2021) (ERG). These values are averaged over $\alpha_0, \ldots, \alpha_9$. MSE values under n = 100 and n = 1000 are multiplied by 10^3 and 10^4 , respectively.

n	Scenario	method	ESS	CT	sESS	MSE
100	(I)	DA-N	863	2.1	414	0.82
		DA-P	856	1.8	480	0.83
		DA-PT	1199	1.8	667	0.82
		ERG	1808	81.4	22	0.82
100	(II)	DA-N	580	2.0	287	5.70
		DA-P	668	1.7	389	5.68
		DA-PT	846	1.8	478	5.73
		ERG	1315	79.5	17	5.66
1000	(I)	DA-N	846	7.7	110	0.80
		DA-P	846	7.4	115	0.80
		DA-PT	1192	7.4	162	0.80
		ERG	1800	819.1	2	0.80
1000	(II)	DA-N	582	7.6	76	5.06
		DA-P	671	7.2	93	5.18
		DA-PT	834	7.3	115	5.15
		ERG	1317	835.4	2	5.17

A remaining issue related to the proposed approach is that our method is likely to be less efficient for extremely small n. In that case, the data augmentation in Theorem S1 should be customized for the model of interest. For example, if n = 1 and $0 < \alpha_l \ll 1$ in the Dirichlet-multinomial model, we could improve the proposed augmentation; see Section S5 of the Supplementary Material.

Acknowledgments

Research of the authors was supported in part by JSPS KAKENHI Grant Number 20J10427, 19K11852, 17K17659, and 21H00699 from Japan Society for the Promotion of Science.

References

- Blei, D.M., Ng, A.Y. and Jordan, M.I. (2003). Latent Dirichlet allocation. Journal of Machine Learning Research, 3, 993–1022.
- [2] Chib, S. and Greenberg, E. (1995). Understanding the Metropolis-Hastings algorithm. The American Statistician, 49, 327–335.

- [3] da Silva, N.B., Prates, M.O. and Goncalves, F.B. (2020). Bayesian linear regression models with flexible error distributions. *Journal of Statistical Computation and Simulation*, 90, 2571–2591.
- [4] Devroye, L. (2012). A note on generating random variables with log-concave densities. Statistics and Probability Letters, 82, 1035–1039.
- [5] Fonseca, T.C., Ferreira, M.A. and Migon, H.S. (2008). Objective Bayesian analysis for the Student-t regression model. *Biometrika*, 95, 325–333.
- [6] Frühwirth-Schnatter, S. (2006). Finite mixture and Markov switching models. Springer-Verlag, New York.
- [7] Geweke, J. (1993). Bayesian treatment of the independent Student-t linear model. Journal of Applied Econometrics, 8, 19–40.
- [8] Gilks, W.R. and Wild, P. (1992). Adaptive rejection sampling for Gibbs sampling. Journal of the Royal Statistical Society. Series C (Applied Statistics), 41, 337–348.
- [9] He, J., Polson, N. and Xu, J. (2021). Bayesian inference for gamma models. arXiv preprint arXiv:2106.01906.
- [10] Miller, J.W. (2019). Fast and accurate approximation of the full conditional for gamma shape parameters. *Journal of Computational and Graphical Statistics*, 28, 476–480.
- [11] Nandram, B. (1998). A bayesian analysis of the three-stage hierarchical multinomial model. Journal of Statistical Computation and Simulation, 61, 97–126.
- [12] Polson, N.G., Scott, J.G. and Windle, J. (2013). Bayesian inference for logistic models using Polya-gamma latent variables. *Journal of the American Statistical Association*, 108, 1339–1349.
- [13] Tierney, L. (1994). Markov Chains for Exploring Posterior Distributions (with discussion). The Annals of Statistics, 22, 1701–1728.
- [14] van Dyk, D.A. and Jiao, X. (2015). Metropolis-Hastings within partially collapsed Gibbs samplers. Journal of Computational and Graphical Statistics, 24, 301–327.
- [15] van Dyk, D.A. and Park, T. (2008). Partially collapsed Gibbs samplers. Journal of the American Statistical Association, 103, 790–796.
- [16] Villa, C. and Rubio, F.J. (2018). Objective priors for the number of degrees of freedom of a multivariate t distribution and the t-copula. Computational Statistics and Data Analysis, 124, 197–219.
- [17] Watanabe, T. (2001). On sampling the degree-of-freedom of Student's-t disturbances. Statistics & Probability Letters, 52, 177–181.
- [18] Xiao, Y., Kub, Y.-C., Bloomfield, P. and Ghosh, S.K. (2015). On the degrees of freedom in MCMC-based Wishart models for time series data. *Statistics and Probability Letters*, 98, 59–64.

[19] Zhou, M. and Carin, L. (2015). Negative binomial process count and mixture modeling. IEEE Transactions on Pattern Analysis and Machine Intelligence, 37, 307–320.

Supplementary Material for "On Data Augmentation for Models Involving Reciprocal Gamma Functions"

In Section S1, we briefly discuss how we could use our method for three models not considered in the main text. In Section S2, we prove Theorem 1 of the main text. In Section S3, we prove two lemmas. In Section S4, we prove Theorem S1, from which Theorem 1 and the theorems of Section S5 follow. In Section S5, we consider extensions of Theorem 1. In Section S6, we derive the expressions (3) and (4) of the main text and the expressions (S1), (S2), (S3), and (S4) of this Supplementary Material. In Section S7, details of the alternative method considered in Section 3.2 of the main text are given. Additional results for the simulation study of Section 3.2 of the main text are in Section S8.

S1 Other examples

Here, we consider three additional models to which our approach is relevant. In Sections S1.1, S1.2, and S1.3, we consider models based on the one-parameter Dirichlet distribution, the negative binomial distribution, and the Wishart distribution, respectively. The expressions (S1), (S2), (S3), and (S4) are derived in Section S6 of this Supplementary Material.

S1.1 The one-parameter Dirichlet prior distribution

Here, we first consider a simple model based on the Dirichlet distribution with a single shape parameter. Suppose that for i = 1, ..., n,

$$\begin{aligned} \boldsymbol{x}_i &\sim p_i(\boldsymbol{x}_i | \boldsymbol{p}_i), \\ \boldsymbol{p}_i &\sim \operatorname{Dir}_L(\boldsymbol{p}_i | \alpha, \dots, \alpha) = \frac{\Gamma((L+1)\alpha)}{\{\Gamma(\alpha)\}^{L+1}} \prod_{l=0}^L p_{i,l} \alpha^{\alpha-1}, \\ \alpha &\sim p(\alpha), \end{aligned}$$

where $\boldsymbol{x}_i \in \mathcal{X}_i$, $\boldsymbol{p}_i = (p_{i,1}, \ldots, p_{i,L}) \in D_L = \{ (\tilde{p}_1, \ldots, \tilde{p}_L) \in (0, 1)^L | \tilde{p}_1, \ldots, \tilde{p}_L > 0, \sum_{l=1}^L \tilde{p}_l < 1 \}$, and $p_{i,0} = 1 - \sum_{l=1}^L p_{i,l}$. Then the posterior distribution $p(\boldsymbol{p}, \alpha | \boldsymbol{x})$ is obtained as the marginal distribution of

$$p(\boldsymbol{p}, \alpha, \boldsymbol{\rho} | \boldsymbol{x}) \propto p(\alpha) \alpha^{nL} \exp\left(-\alpha \sum_{i=1}^{n} \left[\sum_{l=1}^{L} \log \frac{1}{\rho_{i,l}} + \sum_{l=0}^{L} \left\{\log \frac{1}{p_{i,l}} - \log(L+1)\right\}\right]\right) \\ \times \left[\prod_{i=1}^{n} \prod_{l=1}^{L} \{\rho_{i,l}^{l/(L+1)-1} (1-\rho_{i,l})^{(L-l+1)/(L+1)-1}\}\right] \left(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{-1}\right) \prod_{i=1}^{n} p_{i}(\boldsymbol{x}_{i} | \boldsymbol{p}_{i}),$$
(S1)

where $\boldsymbol{\rho} = (\boldsymbol{\rho}_1, \dots, \boldsymbol{\rho}_n) = ((\rho_{1,1}, \dots, \rho_{1,L}), \dots, (\rho_{n,1}, \dots, \rho_{n,L})) \in (0,1)^{nL}$ is a set of additional latent variables.

By (S1), we can construct a Gibbs sampler without a MH step, for example, if $p(\boldsymbol{x}_i) =$ Multin_L $(\boldsymbol{x}_i|N_i, \boldsymbol{p}_i) = \{(N_i!)/\prod_{l=0}^L (x_{i,l}!)\}\prod_{l=0}^L p_{i,l}x_{i,l}$ for $i = 1, \ldots, n$ and $p(\alpha) = \text{Ga}(\alpha|a, b)$ for a, b > 0, where $(x_{i,1}, \ldots, x_{i,L}) = \mathbf{x}_i$ and $x_{i,0} = N_i - \sum_{l=1}^L x_{i,l}$ for $i = 1, \ldots, n$. The full conditional distributions are as follows.

- The full conditional distributions of p_i are $\text{Dir}_L(p_i|x_{i,0} + \alpha, \dots, x_{i,L} + \alpha)$.
- The full conditional distributions of $\rho_{i,l}$ are $\text{Beta}(\rho_{i,l}|\alpha + l/(L+1), (L-l+1)/(L+1))$.
- The full conditional distribution of α is

$$Ga\left(\alpha \left| nL + a, \sum_{i=1}^{n} \left[\sum_{l=1}^{L} \log \frac{1}{\rho_{i,l}} + \sum_{l=0}^{L} \left\{ \log \frac{1}{p_{i,l}} - \log(L+1) \right\} \right] + b \right),$$

where for all i = 1, ..., n, we have $\sum_{l=0}^{L} \{ \log(1/p_{i,l}) - \log(L+1) \} > 0$ by Jensen's inequality.

S1.2 The negative binomial distribution

Here, we consider the estimation of a negative binomial shape parameter. Suppose that for i = 1, ..., n,

$$y_i \sim \text{NB}(y_i | \alpha, p_i) = \frac{\Gamma(\alpha + y_i)}{y_i! \Gamma(\alpha)} p_i^{\alpha} (1 - p_i)^{y_i},$$

 $\alpha \sim p(\alpha).$

We assume that $p_1, \ldots, p_n \in (0, 1)$ are known and fixed for simplicity. However, we can consider the famous negative binomial regression model by setting $p_i = \mathbf{x}_i^{\top} \boldsymbol{\beta}$ for known $\mathbf{x}_i \in \mathbb{R}^p$ and unknown $\boldsymbol{\beta} \sim \pi(\boldsymbol{\beta})$ and using the data augmentation scheme of Polson et al. (2013); see He et al. (2021).

Let $\boldsymbol{y} = (y_1, \ldots, y_n)$. The posterior distribution $p(\alpha | \boldsymbol{y})$ is obtained as the marginal distribution of

$$p(\alpha, \boldsymbol{z}, w, \boldsymbol{\rho} | \boldsymbol{y}) \propto p(\alpha) \alpha^{n} e^{2n\alpha} \Big(\prod_{i=1}^{n} p_{i} \Big)^{\alpha} \Big\{ \prod_{i=1}^{n} (z_{i}^{\alpha+y_{i}-1} e^{-z_{i}}) \Big\} w^{n\alpha-1} e^{-wn\alpha^{2}} \\ \times \Big[\prod_{i=2}^{n} \{ \rho_{i}^{\alpha+(i-1)/n-1} (1-\rho_{i})^{(n-i+1)/n-1} \} \Big] \Big\{ \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}} \Big\}^{2}, \qquad (S2)$$

where $\boldsymbol{z} = (z_1, \ldots, z_n) \in (0, \infty)^n$, $w \in (0, \infty)$, and $\boldsymbol{\rho} = (\rho_2, \ldots, \rho_n) \in (0, 1)^{n-1}$ are additional latent variables. The approaches outlined in Section 2.4 could be useful when we use (S2). An alternative approach would be to combine Theorem 1 with the result of Zhou and Carin (2015).

S1.3 The Wishart distribution

Estimating the shape parameter of the Wishart distribution is also an important problem. For example, Xiao et al. (2015) discussed the importance of considering a fractional shape parameter in the context of analyzing time series data.

Consider the simple model where for $i = 1, \ldots, n$,

$$\begin{aligned} \boldsymbol{x}_{i} &\sim \mathcal{N}_{p}(\boldsymbol{x}_{i}|\boldsymbol{0}_{p}, \Psi^{-1}) = \frac{1}{(2\pi)^{p/2}} |\Psi|^{1/2} e^{-\boldsymbol{x}_{i}^{\top} \boldsymbol{\Psi} \boldsymbol{x}_{i}/2}, \\ \boldsymbol{\Psi} &\sim \mathcal{W}_{p}(\boldsymbol{\Psi}|2\alpha + p - 1, (\beta \boldsymbol{I}_{p})^{-1}) = \frac{\beta^{p\{\alpha + (p-1)/2\}}}{2^{p\{\alpha + (p-1)/2\}} \Gamma_{p}(\alpha + (p-1)/2)} |\Psi|^{\alpha - 1} e^{-\beta \operatorname{tr} \boldsymbol{\Psi}/2}, \\ (\alpha, \beta) &\sim \operatorname{Ga}(\alpha|a, b) \operatorname{Ga}(\beta|c, d) g(\alpha, \beta). \end{aligned}$$

In this case, although Theorem 1 is not applicable to $1/\Gamma_p(\alpha + (p-1)/2)$, we can derive a convenient expression based on a similar idea; see Lemma S5 of Section S6. Let $\gamma = \beta/\alpha$. Our results are the following.

(i) If p = 2m for $m \in \mathbb{N}$, then the posterior distribution $p(\Psi, \alpha, \gamma | \boldsymbol{x})$ is obtained as the marginal distribution of

$$p(\Psi, \alpha, \gamma, \rho | \boldsymbol{x}) \propto g(\alpha, \alpha \gamma) \frac{e^{2m\alpha}}{\alpha^{-1/2}} \frac{\alpha^{p(p-1)/2 + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} \\ \times |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \Big[\prod_{j=2}^m \{\rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} \} \Big] \frac{(2m\alpha)^{2m\alpha - 1/2}}{\Gamma(2m\alpha) e^{2m\alpha}}.$$
(S3)

where $\boldsymbol{\rho} = (\rho_2, \dots, \rho_m) \in (0, 1)^{m-1}$ is a set of additional latent variables.

(ii) If p = 2m - 1 for $m \in \mathbb{N}$, then the posterior distribution $p(\Psi, \alpha, \gamma | \boldsymbol{x})$ is obtained as the marginal distribution of

$$p(\Psi, \alpha, \gamma, \rho, z | \boldsymbol{x}) \propto g(\alpha, \alpha \gamma) \frac{e^{2m\alpha}}{\alpha^{1/2}} \frac{\alpha^{p(p-1)/2 + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} \\ \times |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \Big[\prod_{j=2}^m \{ \rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} \} \Big] z^{\alpha + m - 1/2 - 1} e^{-z} \frac{(2m\alpha)^{2m\alpha - 1/2}}{\Gamma(2m\alpha) e^{2m\alpha}},$$
(S4)

where ρ and $z \in (0, \infty)$ are additional latent variables.

For simplicity, here we consider only the case of even p. Let $m = p/2 \in \mathbb{N}$. The full conditional distributions under the gamma prior $p(\alpha, \beta) = \operatorname{Ga}(\alpha|a, b)\operatorname{Ga}(\beta|c, d)$ are as follows.

- The full conditional of $\boldsymbol{\Psi}$ is $W_p(\boldsymbol{\Psi}|n+2\alpha+p-1,(\alpha\gamma \boldsymbol{I}_p+\sum_{i=1}^n \boldsymbol{x}_i\boldsymbol{x}_i^{\top})^{-1}).$
- The full conditional distributions of ρ_j are $\text{Beta}(\rho_j | 2\alpha + (j-1)/m, (2-1/m)(j-1))$.
- The full conditional of γ is $\operatorname{Ga}(\gamma | p\{\alpha + (p-1)/2\} + c, (\operatorname{tr} \Psi/2 + d)\alpha)$.
- The full conditional of α is

$$p(\alpha|\Psi,\gamma,\boldsymbol{\rho},\boldsymbol{x}) \propto \operatorname{Ga}\left(\alpha\Big|\frac{1}{2} + \frac{p(p-1)}{2} + c + a, B
ight) \frac{(p\alpha)^{p\alpha-1/2}}{\Gamma(p\alpha)e^{p\alpha}},$$

where

$$B = b + d\gamma + \frac{\gamma}{2} \operatorname{tr} \Psi - p - \left(p \log \frac{\gamma}{2} + \log |\Psi| \right) + \sum_{j=2}^{m} 2 \log \frac{1}{\rho_j}$$

is positive since the full conditional must be proper and since the factor $(p\alpha)^{p\alpha-1/2}/{\{\Gamma(p\alpha)e^{p\alpha}\}}$ is bounded as in (1).

S2 Proof of Theorem 1

Here, we prove Theorem 1 of the main text.

Proof of Theorem 1. We first write $1/{\{\Gamma(\xi)\}^m}$ as

$$\frac{1}{\{\Gamma(\xi)\}^m} = \frac{1}{\prod_{j=0}^{m-1} \Gamma(\xi+j/m)} \prod_{j=1}^m \frac{\Gamma(\xi+(j-1)/m)}{\Gamma(\xi)}.$$

By Gauss's multiplication formula, we have

$$\frac{1}{\prod_{j=0}^{m-1} \Gamma(\xi+j/m)} = \frac{(2\pi)^{(1-m)/2} m^{m\xi-1/2}}{\Gamma(m\xi)}.$$

On the other hand, for all $j = 1, \ldots, m$,

$$\frac{\Gamma(\xi + (j-1)/m)}{\Gamma(\xi)} = \frac{\Gamma(\xi + (j-1)/m)}{\Gamma(\xi)} \int_0^1 \text{Beta}\Big(\rho_j\Big|\xi + \frac{j-1}{m}, \frac{m-j+1}{m}\Big)d\rho_j$$
$$= \frac{\xi}{\Gamma((m-j+1)/m)} \int_0^1 \rho_j^{\xi + (j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j.$$

Therefore,

$$\frac{1}{\{\Gamma(\xi)\}^m} = \frac{(2\pi)^{(1-m)/2} m^{m\xi-1/2}}{\Gamma(m\xi)} \prod_{j=2}^m \frac{\xi}{\Gamma((m-j+1)/m)} \int_0^1 \rho_j^{\xi+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j,$$

and the result follows.

S3 Lemmas

Here, we present two lemmas.

Lemma S1. Let $M \in \mathbb{N}_0$. Then

$$\frac{1}{\Gamma(\xi)} = \frac{(M+1)^{-1/2}}{(2\pi)^{M/2}} \frac{(M+1)^{(M+1)\xi}}{\Gamma((M+1)\xi)} \prod_{j=2}^{M+1} \int_0^\infty t_j \xi^{+(j-1)/(M+1)-1} e^{-t_j} dt_j$$

for all $\xi > 0$.

Proof. By Gauss's multiplication formula,

$$\begin{split} \frac{1}{\Gamma(\xi)} &= \frac{1}{\prod_{j=0}^{M} \Gamma(\xi+j/(M+1))} \prod_{j=2}^{M+1} \Gamma(\xi+(j-1)/(M+1)) \\ &= \frac{(2\pi)^{-M/2} (M+1)^{(M+1)\xi-1/2}}{\Gamma((M+1)\xi)} \prod_{j=2}^{M+1} \int_{0}^{\infty} t_{j}^{\xi+(j-1)/(M+1)-1} e^{-t_{j}} dt_{j}, \end{split}$$

which is the desired result.

Lemma S2. Let $K \in \mathbb{N}_0$. Let $M_1, \ldots, M_K \in \mathbb{N}_0$. Then

$$\frac{1}{\Gamma(\xi)} = \left\{ \prod_{k=1}^{K} \frac{(M_k+1)^{-1/2}}{(2\pi)^{M_k/2}} \right\} \frac{\prod_{k=1}^{K} (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)\xi}}{\Gamma((M_K+1)\cdots(M_1+1)\xi)} \\ \times \prod_{k=1}^{K} \prod_{j=2}^{M_k+1} \int_0^\infty t_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)\xi+(j-1)/(M_k+1)-1} e^{-t_{k,j}} dt_{k,j}$$

for all $\xi > 0$.

Proof. By Lemma S1, we have

$$\begin{aligned} \frac{1}{\Gamma(\xi)} &= \frac{(M_1+1)^{-1/2}}{(2\pi)^{M_1/2}} \frac{(M_1+1)^{(M_1+1)\xi}}{\Gamma((M_1+1)\xi)} \prod_{j=2}^{M_1+1} \int_0^\infty t_{1,j}^{\xi+(j-1)/(M_1+1)-1} e^{-t_{1,j}} dt_{1,j}, \\ \frac{1}{\Gamma((M_1+1)\xi)} &= \frac{(M_2+1)^{-1/2}}{(2\pi)^{M_2/2}} \frac{(M_2+1)^{(M_2+1)(M_1+1)\xi}}{\Gamma((M_2+1)(M_1+1)\xi)} \prod_{j=2}^{M_2+1} \int_0^\infty t_{2,j}^{(M_1+1)\xi+(j-1)/(M_2+1)-1} e^{-t_{2,j}} dt_{2,j}, \\ \vdots \end{aligned}$$

$$\frac{1}{\Gamma((M_{K-1}+1)\cdots(M_1+1)\xi)} = \frac{(M_K+1)^{-1/2}}{(2\pi)^{M_K/2}} \frac{(M_K+1)^{(M_K+1)(M_{K-1}+1)\cdots(M_1+1)\xi}}{\Gamma((M_K+1)(M_{K-1}+1)\cdots(M_1+1)\xi)} \times \prod_{j=2}^{M_K+1} \int_0^\infty t_{K,j} (M_{K-1}+1)\cdots(M_1+1)\xi + (j-1)/(M_K+1) - 1} e^{-t_{K,j}} dt_{K,j}.$$

Therefore,

$$\begin{aligned} \frac{1}{\Gamma(\xi)} &= \Big\{ \prod_{k=1}^{K} \frac{(M_k+1)^{-1/2}}{(2\pi)^{M_k/2}} \Big\} \frac{\prod_{k=1}^{K} (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)\xi}}{\Gamma((M_K+1)\cdots(M_1+1)\xi)} \\ &\times \prod_{k=1}^{K} \prod_{j=2}^{M_k+1} \int_0^\infty t_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)\xi+(j-1)/(M_k+1)-1} e^{-t_{k,j}} dt_{k,j}, \end{aligned}$$

which is the desired result.

S4 General explicit expressions for $1/\{\Gamma(\xi)\}^m$ for beta-gamma data augmentation

Theorem 1 follows from part (i) of the following theorem.

Theorem S1. Let $m \in \mathbb{N}$. Let $K \in \mathbb{N}_0$ and let $M_1, \ldots, M_K \in \mathbb{N}_0$.

(i) For all $\xi > 0$, we have

$$\begin{split} \frac{1}{\{\Gamma(\xi)\}^m} &= \frac{\{(M_K+1)\cdots(M_1+1)m/(2\pi)\}^{\sum_{k=1}^K M_k/2}}{(2\pi)^{(m-1)/2}\prod_{j=2}^m \Gamma((m-j+1)/m)} \frac{\xi^{m+\sum_{k=1}^K M_k/2+1/2-1}e^{(M_K+1)\cdots(M_1+1)m\xi}}{\{(M_K+1)\cdots(M_1+1)\}^{m\xi}\xi^{m\xi}} \\ &\times \Big\{\prod_{k=1}^K (M_k+1)^{(M_k+1)\cdots(M_1+1)m\xi}\Big\}\Big\{\prod_{j=2}^m \int_0^1 \rho_j^{\xi+(j-1)/m-1}(1-\rho_j)^{(m-j+1)/m-1}d\rho_j\Big\} \\ &\times \Big\{\prod_{k=1}^K \prod_{j=2}^{M_k+1} \int_0^\infty t_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi+(j-1)/(M_k+1)-1}e^{-t_{k,j}(M_K+1)\cdots(M_1+1)m\xi}dt_{k,j}\Big\} \\ &\times \frac{\{(M_K+1)\cdots(M_1+1)m\xi\}^{(M_K+1)\cdots(M_1+1)m\xi-1/2}}{\Gamma((M_K+1)\cdots(M_1+1)m\xi)e^{(M_K+1)\cdots(M_1+1)m\xi}}. \end{split}$$

(ii) For all $\xi > 0$, we have

$$\frac{1}{\{\Gamma(\xi)\}^m} = \frac{\{(M_K+1)\cdots(M_1+1)m/(2\pi)\}^{\sum_{k=1}^K M_k/2}}{(2\pi)^{(m-1)/2}\prod_{j=2}^m \Gamma((m-j+1)/m)} \xi^m \frac{(\xi+1)^{\sum_{k=1}^K M_k/2+1/2-1}e^{(M_K+1)\cdots(M_1+1)m(\xi+1)}}{\{(M_K+1)\cdots(M_1+1)\}^{m(\xi+1)}(\xi+1)^{m\xi}} \\
\times \left\{\prod_{k=1}^K (M_k+1)^{(M_k+1)\cdots(M_1+1)m(\xi+1)}\right\} \left\{\prod_{j=2}^m \int_0^1 \rho_j^{\xi+1+(j-1)/m-1}(1-\rho_j)^{(m-j+1)/m-1}d\rho_j\right\} \\
\times \left\{\prod_{k=1}^K \prod_{j=2}^{M_k+1} \int_0^\infty t_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m(\xi+1)+(j-1)/(M_k+1)-1}e^{-t_{k,j}(M_K+1)\cdots(M_1+1)m(\xi+1)}dt_{k,j}\right\} \\
\times \frac{\{(M_K+1)\cdots(M_1+1)m(\xi+1)\}^{(M_K+1)\cdots(M_1+1)m(\xi+1)-1/2}}{\Gamma((M_K+1)\cdots(M_1+1)m(\xi+1))e^{(M_K+1)\cdots(M_1+1)m(\xi+1)}}.$$

Proof. Part (ii) follows from part (i). For part (i), we have, by Theorem 1,

$$\frac{1}{\{\Gamma(\xi)\}^m} = \frac{(2\pi)^{(1-m)/2}m^{-1/2}}{\prod_{j=2}^m \Gamma((m-j+1)/m)} \frac{\xi^{m-1}m^{m\xi}}{\Gamma(m\xi)} \prod_{j=2}^m \int_0^1 \rho_j^{\xi+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j.$$

By Lemma S2,

$$\frac{1}{\Gamma(m\xi)} = \left\{ \prod_{k=1}^{K} \frac{(M_k+1)^{-1/2}}{(2\pi)^{M_k/2}} \right\} \frac{\prod_{k=1}^{K} (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)m\xi}}{\Gamma((M_K+1)\cdots(M_1+1)m\xi)} \\ \times \prod_{k=1}^{K} \prod_{j=2}^{M_k+1} \int_0^\infty t_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi+(j-1)/(M_k+1)-1} e^{-t_{k,j}} dt_{k,j}.$$

By making the change of variables $t_{k,j} = \tilde{t}_{k,j}(M_K + 1)\cdots(M_1 + 1)m\xi$,

$$\frac{1}{\Gamma(m\xi)} = \left\{ \prod_{k=1}^{K} \frac{(M_k+1)^{-1/2}}{(2\pi)^{M_k/2}} \right\} \frac{\prod_{k=1}^{K} (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)m\xi}}{\Gamma((M_K+1)\cdots(M_1+1)m\xi)} \\ \times \left\{ (M_K+1)\cdots(M_1+1)m\xi \right\}^{(M_K+1)\cdots(M_1+1)m\xi-m\xi+\sum_{k=1}^{K} M_k/2} \\ \times \prod_{k=1}^{K} \prod_{j=2}^{M_k+1} \int_0^\infty \tilde{t}_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi+(j-1)/(M_k+1)-1} e^{-\tilde{t}_{k,j}(M_K+1)\cdots(M_1+1)m\xi} d\tilde{t}_{k,j}.$$

Therefore,

$$\begin{split} \frac{1}{\{\Gamma(\xi)\}^m} &= \frac{(2\pi)^{(1-m)/2}m^{-1/2}}{\prod_{j=2}^m \Gamma((m-j+1)/m)} \xi^{m-1} m^m \xi \Big\{ \prod_{j=2}^m \int_0^1 \rho_j \xi^{+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j \Big\} \\ &\times \Big\{ \prod_{k=1}^K \frac{(M_k+1)^{-1/2}}{(2\pi)^{M_k/2}} \Big\} \frac{e^{(M_K+1)\cdots(M_1+1)m\xi} \prod_{k=1}^K (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)m\xi}}{\{(M_K+1)\cdots(M_1+1)m\xi\}^{m\xi-\sum_{k=1}^K M_k/2-1/2}} \\ &\times \frac{\{(M_K+1)\cdots(M_1+1)m\xi\}^{(M_K+1)\cdots(M_1+1)m\xi^{-1/2}}}{\Gamma((M_K+1)\cdots(M_1+1)m\xi)e^{(M_K+1)\cdots(M_1+1)m\xi}} \\ &\times \prod_{k=1}^K \prod_{j=2}^M \int_0^\infty \tilde{t}_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi^{+(j-1)/(M_k+1)-1}} e^{-\tilde{t}_{k,j}(M_K+1)\cdots(M_1+1)m\xi} d\tilde{t}_{k,j} \\ &= \frac{\prod_{k=1}^K \{(M_K+1)\cdots(M_1+1)m/(2\pi)\}^{M_k/2}}{(2\pi)^{(m-1)/2} \prod_{j=2}^m \Gamma((m-j+1)/m)} \frac{\xi^{m+\sum_{k=1}^K M_k/2+1/2-1}e^{(M_K+1)\cdots(M_1+1)m\xi}}{\{(M_K+1)\cdots(M_1+1)\xi\}^{m\xi}} \\ &\times \Big\{ \prod_{k=1}^K (M_k+1)^{(M_k+1)(M_{k-1}+1)\cdots(M_1+1)m\xi} \Big\} \Big\{ \prod_{j=2}^m \int_0^1 \rho_j \xi^{+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j \Big\} \\ &\times \prod_{k=1}^K \prod_{j=2}^{M_k+1} \int_0^\infty \tilde{t}_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi+(j-1)/(M_k+1)-1} e^{-\tilde{t}_{k,j}(M_K+1)\cdots(M_1+1)m\xi} d\tilde{t}_{k,j} \\ &\times \prod_{k=1}^K \prod_{j=2}^{M_k+1} \int_0^\infty \tilde{t}_{k,j}^{(M_{k-1}+1)\cdots(M_1+1)m\xi+(j-1)/(M_k+1)-1} e^{-\tilde{t}_{k,j}(M_K+1)\cdots(M_1+1)m\xi} d\tilde{t}_{k,j} \\ &\times \frac{\{(M_K+1)\cdots(M_1+1)m\xi\}^{(M_K+1)\cdots(M_1+1)m\xi-1/2}}{\Gamma((M_K+1)\cdots(M_1+1)m\xi)e^{(M_K+1)\cdots(M_1+1)m\xi-1/2}}. \end{split}$$

This proves part (i).

S5 Extensions of Theorem 1

Parts (i) and (ii) of Theorem S2 follow from parts (i) and (ii) of Theorem S1, respectively. Theorem S2 can be used in a manner similar to that for Theorem 1.

Theorem S2. Let $m \in \mathbb{N}$ and $K \in \mathbb{N}_0$.

(i) For all $\xi > 0$, we have

$$\begin{split} \frac{1}{\{\Gamma(\xi)\}^m} &= C_{m,K} \frac{1}{\xi^{m\xi}} \xi^{m+K/2+1/2-1} e^{2^K m\xi} \\ &\quad \times 2^{\{2(2^K-1)-K\}m\xi} \Big\{ \prod_{j=2}^m \int_0^1 \rho_j ^{\xi+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j \Big\} \\ &\quad \times \Big\{ \prod_{k=1}^K \int_0^\infty t_k ^{2^{k-1}m\xi+1/2-1} e^{-t_k 2^K m\xi} dt_k \Big\} \frac{(2^K m\xi)^{2^K m\xi-1/2}}{\Gamma(2^K m\xi) e^{2^K m\xi}}, \end{split}$$
where $C_{m,K} = \{2^K m/(2\pi)\}^{K/2} / \{ (2\pi)^{(m-1)/2} \prod_{j=2}^m \Gamma((m-j+1)/m) \}.$

(ii) For all $\xi > 0$, we have

$$\begin{split} \frac{1}{\{\Gamma(\xi)\}^m} &= C_{m,K} \frac{(\xi+1)^{K/2+1/2-1}}{(\xi+1)^{m\xi}} \xi^m e^{2^K m(\xi+1)} \\ &\quad \times 2^{\{2(2^K-1)-K\}m(\xi+1)} \Big\{ \prod_{j=2}^m \int_0^1 \rho_j ^{\xi+1+(j-1)/m-1} (1-\rho_j)^{(m-j+1)/m-1} d\rho_j \Big\} \\ &\quad \times \Big\{ \prod_{k=1}^K \int_0^\infty t_k ^{2^{k-1}m(\xi+1)+1/2-1} e^{-t_k 2^K m(\xi+1)} dt_k \Big\} \frac{\{2^K m(\xi+1)\}^{2^K m(\xi+1)-1/2}}{\Gamma(2^K m(\xi+1)) e^{2^K m(\xi+1)}}, \end{split}$$
where $C_{m,K} = \{2^K m/(2\pi)\}^{K/2} / \Big\{ (2\pi)^{(m-1)/2} \prod_{j=2}^m \Gamma((m-j+1)/m) \Big\}.$

As discussed in Section 2.1, when we use Theorem 1, we consider ρ_j as additional latent variables, the full conditional distributions of ρ_j are beta, and we can easily sample ρ_j in a MCMC algorithm. When we use Theorem S2, we consider $\mathbf{t} = (t_1, \ldots, t_K) \in (0, \infty)^K$ as a set of latent variables in addition to $\boldsymbol{\rho}$. The full conditional distributions of t_k are $\operatorname{Ga}(t_k|2^{k-1}m\xi+1/2, 2^Km\xi)$ if we use part (i) and $\operatorname{Ga}(t_k|2^{k-1}m(\xi+1)+1/2, 2^Km(\xi+1))$ if we use part (ii).

Although there are more latent variables, the approximation to the full conditional of ξ becomes more accurate if we use Theorem S2 instead of Theorem 1. For example, in the Ga(α, β) case considered in Section 2.2, we have better lower bounds for the acceptance probability. That is, we have

$$\exp\left\{-\frac{1}{12(2^{K}n\alpha^{*})}\right\} \ge 1 - \frac{1}{12(2^{K}n\alpha^{*})}$$

for part (i) and

$$\exp\left[-\frac{1}{12\{2^{K}n(\alpha^{*}+1)\}}\right] \ge 1 - \frac{1}{12\{2^{K}n(\alpha^{*}+1)\}} \ge 1 - \frac{1}{12(2^{K}n)}$$

for part (ii). The lower bounds are clearly increasing functions of K and n. As $K \to \infty$, they converge to 1 exponentially fast. The last bound is independent of all the variables.

When we use part (ii), we have a factor of the form $(\xi + 1)^c$. It is rewritten in the following way.

(a) If c < 0,

$$(\xi+1)^c = \int_0^\infty \frac{z^{-c-1}}{\Gamma(-c)} e^{-z(\xi+1)} dz.$$

(b) If c = 0,

$$(\xi + 1)^c = 1.$$

(c) If c > 0,

$$(\xi+1)^c = \xi^{c+\varepsilon} \sum_{\zeta=0}^{\infty} \binom{c+\varepsilon+\zeta-1}{\zeta} \int_0^\infty \frac{\eta^{\zeta+\varepsilon-1}}{\Gamma(\zeta+\varepsilon)} e^{-\eta(\xi+1)} d\eta$$

for any $\varepsilon > 0$ by Lemma S3 below.

In case (a), we introduce an additional latent variable $z \in (0, \infty)$. Its full conditional distribution is $\operatorname{Ga}(z|-c,\xi+1)$. In case (c), we introduce two latent variables $\zeta \in \mathbb{N}_0$ and $\eta \in (0,\infty)$. The variable η is marginalized out except when we sample ξ and η (see van Dyk and Park (2008) and van Dyk and Jiao (2015) for more details). The full conditional of η is $\operatorname{Ga}(\eta|\zeta+\varepsilon,\xi+1)$. The full conditional of ζ after marginalizing out η is proportional to

$$\binom{C+\varepsilon+\zeta-1}{\zeta}\frac{1}{(\xi+1)^{\zeta+\varepsilon}} \propto \binom{C+\varepsilon+\zeta-1}{\zeta} \left(\frac{\xi}{\xi+1}\right)^{C+\varepsilon} \left(\frac{1}{\xi+1}\right)^{\zeta} = \operatorname{NB}\left(\zeta \middle| C+\varepsilon, \frac{\xi}{\xi+1}\right).$$

Lemma S3. Let $\varepsilon > 0$ and $C > -\varepsilon$. Then

$$\begin{aligned} (\xi_1 + \xi_2)^C &= \xi_1^{C+\varepsilon} \sum_{\zeta=0}^{\infty} \binom{C+\varepsilon+\zeta-1}{\zeta} \frac{\xi_2^{\zeta}}{(\xi_1 + \xi_2)^{\zeta+\varepsilon}} \\ &= \xi_1^{C+\varepsilon} \sum_{\zeta=0}^{\infty} \binom{C+\varepsilon+\zeta-1}{\zeta} \xi_2^{\zeta} \int_0^{\infty} \frac{\eta^{\zeta+\varepsilon-1}}{\Gamma(\zeta+\varepsilon)} e^{-\eta(\xi_1+\xi_2)} d\eta \end{aligned}$$

for all $\xi > 0$.

Finally, we can also use Lemma S4 in case (c).

Lemma S4. Let C > -1/2. Then

$$\begin{aligned} (\xi_1 + \xi_2)^C &= \int_0^\infty \frac{\eta^{C+1/2-1}}{\Gamma(C+1/2)} \frac{e^{-\eta/(\xi_1 + \xi_2)}}{(\xi_1 + \xi_2)^{1/2}} d\eta \\ &= \int_0^\infty \frac{\eta^{C+1/2-1}}{\Gamma(C+1/2)} \Big[\int_{-\infty}^\infty \frac{1}{\sqrt{2\pi}} \xi_1^{-1/2} \xi_2^{-1/2} e^{-\theta^2/(2\xi_1) - \{\theta - (2\eta)^{1/2}\}^2/(2\xi_2)} d\theta \Big] d\eta \\ &= \int_0^\infty \frac{\eta^{C+1/2-1}}{\Gamma(C+1/2)} \Big\{ \int_{-\infty}^\infty \frac{\eta^{1/2}}{\sqrt{\pi}} \xi_1^{-1/2} \xi_2^{-1/2} e^{-\eta\tilde{\theta}^2/\xi_1 - \eta(\tilde{\theta} - 1)^2/\xi_2} d\tilde{\theta} \Big\} d\eta \end{aligned}$$

for all $\xi_1, \xi_2 > 0$.

S6 Proofs of the expressions (3), (4), (S1), (S2), (S3), and (S4)

Here, we derive the expressions (3) and (4) of the main text and the expressions (S1), (S2), (S3), and (S4) of this Supplementary Material.

Proof of (3). The posterior of θ , τ , and α is

$$p(\theta,\tau,\alpha|\boldsymbol{x}) \propto p(\alpha) \frac{p(\theta,\tau)}{\tau^{n/2}} \prod_{i=1}^{n} \left[\alpha^{\alpha} \frac{\Gamma(\alpha+1/2)}{\Gamma(\alpha)} / \left\{ \alpha + \frac{(x_i-\theta)^2}{2\tau} \right\}^{\alpha+1/2} \right]$$
$$= \frac{p(\theta,\tau)}{\tau^{n/2}} p(\alpha) \frac{\alpha^{n\alpha}}{\{\Gamma(\alpha)\}^n} \prod_{i=1}^{n} \int_0^\infty w_i^{\alpha+1/2-1} e^{-w_i \{\alpha + (x_i-\theta)^2/(2\tau)\}} dw_i.$$

Then, by Theorem 1, we have

$$p(\theta,\tau,\alpha|\mathbf{x}) \propto \frac{p(\theta,\tau)}{\tau^{n/2}} p(\alpha) \alpha^{n+1/2-1} e^{n\alpha} \Big[\prod_{i=1}^n \int_0^\infty w_i^{\alpha+1/2-1} e^{-w_i \{\alpha+(x_i-\theta)^2/(2\tau)\}} dw_i \Big] \\ \times \Big\{ \prod_{i=2}^n \int_0^1 \rho_i^{\alpha+(i-1)/n-1} (1-\rho_i)^{(n-i+1)/n-1} d\rho_i \Big\} \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}},$$

and this completes the proof.

Proof of (4). The posterior of p and α is

$$p(\boldsymbol{p}, \boldsymbol{\alpha} | \boldsymbol{x}) \propto p(\boldsymbol{\alpha}) \Big\{ \frac{\Gamma\left(\sum_{l=0}^{L} \alpha_{l}\right)}{\prod_{l=0}^{L} \Gamma(\alpha_{l})} \Big\}^{n} \prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{x_{i,l}+\alpha_{l}-1} = p(\boldsymbol{\alpha}) \Big(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{x_{i,l}+\alpha_{l}-1} \Big) \Big[\prod_{l=0}^{L} \frac{1}{\{\Gamma(\alpha_{l})\}^{n}} \Big] \Big(\prod_{i=1}^{n} \int_{0}^{\infty} z_{i} \sum_{l=0}^{L} \alpha_{l} - 1 e^{-z_{i}} dz_{i} \Big).$$

By Theorem 1 and Lemma 1,

$$\frac{1}{\{\Gamma(\alpha_l)\}^n} \propto \alpha_l^n e^{2n\alpha_l} \left\{ \prod_{i=2}^n \int_0^1 \rho_{i,l}^{\alpha_l + (i-1)/n - 1} (1 - \rho_{i,l})^{(n-i+1)/n - 1} d\rho_{i,l} \right\} \\ \times \left\{ \frac{(n\alpha_l)^{n\alpha_l - 1/2}}{\Gamma(n\alpha_l) e^{n\alpha_l}} \right\}^2 \int_0^\infty w_l^{n\alpha_l - 1} e^{-w_l n\alpha_l^2} dw_l.$$

Thus,

$$p(\boldsymbol{p}, \boldsymbol{\alpha} | \boldsymbol{x}) \propto p(\boldsymbol{\alpha}) \Big(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l} x_{i,l} + \alpha_{l} - 1 \Big) \Big(\prod_{i=1}^{n} \int_{0}^{\infty} z_{i} \sum_{l=0}^{L} \alpha_{l} - 1 e^{-z_{i}} dz_{i} \Big) \\ \times \prod_{l=0}^{L} \Big[\alpha_{l}^{n} e^{2n\alpha_{l}} \Big\{ \prod_{i=2}^{n} \int_{0}^{1} \rho_{i,l} \alpha_{l} + (i-1)/n - 1 (1 - \rho_{i,l})^{(n-i+1)/n - 1} d\rho_{i,l} \Big\} \\ \times \Big\{ \frac{(n\alpha_{l})^{n\alpha_{l} - 1/2}}{\Gamma(n\alpha_{l}) e^{n\alpha_{l}}} \Big\}^{2} \int_{0}^{\infty} w_{l}^{n\alpha_{l} - 1} e^{-w_{l}n\alpha_{l}^{2}} dw_{l} \Big],$$

and the result follows.

Proof of (S1). The posterior of p and α is

$$p(\boldsymbol{p}, \alpha | \boldsymbol{x}) \propto p(\alpha) \prod_{i=1}^{n} \left[\frac{\Gamma((L+1)\alpha)}{\{\Gamma(\alpha)\}^{L+1}} \Big(\prod_{l=0}^{L} p_{i,l}^{\alpha-1} \Big) p(\boldsymbol{x}_{i} | \boldsymbol{p}_{i}) \right].$$
(S5)

By Theorem 1, we have

$$\begin{split} \frac{\Gamma((L+1)\alpha)}{\{\Gamma(\alpha)\}^{L+1}} &\propto \Gamma((L+1)\alpha) \frac{\alpha^{L+1/2} e^{(L+1)\alpha}}{\alpha^{(L+1)\alpha}} \\ &\times \Big\{ \prod_{l=1}^{L} \int_{0}^{1} \rho_{l}^{\alpha+l/(L+1)-1} (1-\rho_{l})^{(L-l+1)/(L+1)-1} d\rho_{l} \Big\} \frac{\{(L+1)\alpha\}^{(L+1)\alpha-1/2}}{\Gamma((L+1)\alpha) e^{(L+1)\alpha}} \\ &\propto \alpha^{L} (L+1)^{(L+1)\alpha} \prod_{l=1}^{L} \int_{0}^{1} \rho_{l}^{\alpha+l/(L+1)-1} (1-\rho_{l})^{(L-l+1)/(L+1)-1} d\rho_{l}. \end{split}$$

Therefore,

$$\prod_{i=1}^{n} \frac{\Gamma((L+1)\alpha)}{\{\Gamma(\alpha)\}^{L+1}} \propto \int_{(0,1)^{nL}} \alpha^{nL} (L+1)^{n(L+1)\alpha} \Big(\prod_{i=1}^{n} \prod_{l=1}^{L} \rho_{i,l}^{\alpha} \Big) \Big[\prod_{i=1}^{n} \prod_{l=1}^{L} \{\rho_{i,l}^{l/(L+1)-1} (1-\rho_{i,l})^{(L-l+1)/(L+1)-1} \} \Big] d\rho.$$

Substituting this into (S5) gives

$$p(\boldsymbol{p}, \alpha | \boldsymbol{x}) \propto \int_{(0,1)^{nL}} \left(p(\alpha) \alpha^{nL} (L+1)^{n(L+1)\alpha} \left(\prod_{i=1}^{n} \prod_{l=1}^{L} \rho_{i,l}^{\alpha} \right) \left(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{\alpha} \right) \right. \\ \times \left[\prod_{i=1}^{n} \prod_{l=1}^{L} \left\{ \rho_{i,l}^{l/(L+1)-1} (1-\rho_{i,l})^{(L-l+1)/(L+1)-1} \right\} \right] \left(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{-1} \right) \prod_{i=1}^{n} p(\boldsymbol{x}_{i} | \boldsymbol{p}_{i}) \right) d\boldsymbol{\rho},$$

where

$$(L+1)^{n(L+1)\alpha} \Big(\prod_{i=1}^{n} \prod_{l=1}^{L} \rho_{i,l}^{\alpha} \Big) \Big(\prod_{i=1}^{n} \prod_{l=0}^{L} p_{i,l}^{\alpha} \Big) = \exp\Big(-\alpha \sum_{i=1}^{n} \Big[\sum_{l=1}^{L} \log \frac{1}{\rho_{i,l}} + \sum_{l=0}^{L} \Big\{ \log \frac{1}{p_{i,l}} - \log(L+1) \Big\} \Big] \Big),$$

and this completes the proof.

Proof of (S2). The posterior of α is

$$p(\alpha|\mathbf{y}) \propto p(\alpha) \frac{1}{\{\Gamma(\alpha)\}^n} \Big\{ \prod_{i=1}^n \Gamma(\alpha+y_i) \Big\} \Big(\prod_{i=1}^n p_i \Big)^{\alpha} \\ = p(\alpha) \frac{1}{\{\Gamma(\alpha)\}^n} \Big(\prod_{i=1}^n \int_0^\infty z_i^{\alpha+y_i-1} e^{-z_i} dz_i \Big) \Big(\prod_{i=1}^n p_i \Big)^{\alpha}.$$

By Theorem 1 and Lemma 1,

$$\frac{1}{\{\Gamma(\alpha)\}^n} \propto \alpha^n e^{2n\alpha} \Big\{ \prod_{i=2}^n \int_0^1 \rho_i^{\alpha+(i-1)/n-1} (1-\rho_i)^{(n-i+1)/n-1} d\rho_i \Big\} \\ \times \Big\{ \frac{(n\alpha)^{n\alpha-1/2}}{\Gamma(n\alpha)e^{n\alpha}} \Big\}^2 \int_0^\infty w^{n\alpha-1} e^{-wn\alpha^2} dw.$$

The desired result follows from the above two expressions.

Proof of (S3) and (S4). The posterior of Ψ , α , and β is

$$p(\boldsymbol{\Psi}, \boldsymbol{\alpha}, \boldsymbol{\beta} | \boldsymbol{x}) \propto g(\boldsymbol{\alpha}, \boldsymbol{\beta}) \frac{\alpha^{a-1} e^{-b\alpha} \beta^{p\{\boldsymbol{\alpha}+(p-1)/2\}+c-1} e^{-d\beta}}{2^{p\alpha} \Gamma_p(\boldsymbol{\alpha}+(p-1)/2)} |\boldsymbol{\Psi}|^{n/2+\alpha-1} e^{-\operatorname{tr}(\boldsymbol{\beta} \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^{\top}) \boldsymbol{\Psi}/2}$$

By making the change of variables $\gamma = \beta / \alpha$, we have

$$\frac{p(\boldsymbol{\Psi}, \boldsymbol{\alpha}, \boldsymbol{\gamma} | \boldsymbol{x})}{g(\boldsymbol{\alpha}, \boldsymbol{\alpha} \boldsymbol{\gamma})} \propto \frac{\alpha^{p\{\boldsymbol{\alpha} + (p-1)/2\} + c + a - 1} e^{-b\alpha} \boldsymbol{\gamma}^{p\{\boldsymbol{\alpha} + (p-1)/2\} + c - 1} e^{-d\alpha \boldsymbol{\gamma}}}{2^{p\alpha} \Gamma_p(\boldsymbol{\alpha} + (p-1)/2)} |\boldsymbol{\Psi}|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\boldsymbol{\alpha} \boldsymbol{\gamma} \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \boldsymbol{\Psi}/2}.$$

Suppose first that p = 2m for $m \in \mathbb{N}$. Then, by Lemma S5 below,

$$\frac{p(\Psi, \alpha, \gamma | \boldsymbol{x})}{g(\alpha, \alpha \gamma)} \propto \frac{\alpha^{p\{\alpha + (p-1)/2\} + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} d\rho_j \\ \propto \frac{e^{2m\alpha}}{\alpha^{-1/2}} \frac{\alpha^{p(p-1)/2 + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \frac{(2m\alpha)^{2m\alpha - 1/2}}{\Gamma(2m\alpha) e^{2m\alpha}} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} d\rho_j.$$

Next, suppose that p = 2m - 1 for $m \in \mathbb{N}$. Then, by Lemma S5 below,

$$\frac{p(\Psi, \alpha, \gamma | \boldsymbol{x})}{g(\alpha, \alpha \gamma)} \propto \frac{\alpha^{p\{\alpha + (p-1)/2\} + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} d\rho_j \int_0^\infty z^{\alpha + m - 1/2 - 1} e^{-z} dz \\ \propto \frac{e^{2m\alpha}}{\alpha^{1/2}} \frac{\alpha^{p(p-1)/2 + c + a - 1} e^{-b\alpha} \gamma^{p\{\alpha + (p-1)/2\} + c - 1} e^{-d\alpha \gamma}}{2^{p\alpha}} |\Psi|^{n/2 + \alpha - 1} e^{-\operatorname{tr}(\alpha \gamma \boldsymbol{I}_p + \sum_{i=1}^n \boldsymbol{x}_i \boldsymbol{x}_i^\top) \Psi/2} \\ \times \frac{(2m\alpha)^{2m\alpha - 1/2}}{\Gamma(2m\alpha)e^{2m\alpha}} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha + (j-1)/m - 1} (1 - \rho_j)^{(2-1/m)(j-1) - 1} d\rho_j \int_0^\infty z^{\alpha + m - 1/2 - 1} e^{-z} dz$$

This completes the proof.

Lemma S5. Let $p, m \in \mathbb{N}$ and $\alpha > 0$. Then

$$\begin{split} &\frac{\pi^{p(p-1)/4}}{\Gamma_p(\alpha+(p-1)/2)} / \frac{\left\{\prod_{j=1}^m 2^{2(j-1)-1}\right\}(2\pi)^{(1-m)/2}m^{-1/2}}{\pi^{m/2}\prod_{j=2}^m \Gamma((2-1/m)(j-1))} \\ &= \begin{cases} \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha+(j-1)/m-1}(1-\rho_j)^{(2-1/m)(j-1)-1}d\rho_j, & \text{if } p = 2m, \\ \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha+(j-1)/m-1}(1-\rho_j)^{(2-1/m)(j-1)-1}d\rho_j \int_0^\infty z^{\alpha+m-1/2-1}e^{-z}dz, & \text{if } p = 2m-1. \end{cases}$$

Proof. By the definition of the multivariate gamma function,

$$\frac{\pi^{p(p-1)/4}}{\Gamma_p(\alpha + (p-1)/2)} = \frac{1}{\prod_{j=1}^p \Gamma(\alpha + (p-1)/2 - (j-1)/2)}$$
$$= \frac{1}{\prod_{j=1}^p \Gamma(\alpha + (p-j)/2)} = \frac{1}{\prod_{j=1}^p \Gamma(\alpha + (j-1)/2)}.$$

By the duplication formula for the gamma function,

$$\prod_{j=1}^{2m} \Gamma\left(\alpha + \frac{j-1}{2}\right) = \prod_{j=1}^{m} \left\{ \Gamma(\alpha + j - 1) \Gamma\left(\alpha + j - \frac{1}{2}\right) \right\} = \pi^{m/2} \prod_{j=1}^{m} \frac{\Gamma(2(\alpha + j - 1))}{2^{2(\alpha + j - 1) - 1}}.$$
 (S6)

Therefore,

$$\frac{\pi^{m/2} / \prod_{j=1}^{m} 2^{2(\alpha+j-1)-1}}{\prod_{j=1}^{2m} \Gamma(\alpha+(j-1)/2)} = \prod_{j=1}^{m} \frac{1}{\Gamma(2\alpha+2(j-1))}$$

$$= \frac{1}{\Gamma(2\alpha)} \prod_{j=2}^{m} \left\{ \frac{1}{\Gamma(2\alpha+2(j-1))} \int_{0}^{1} \operatorname{Beta}\left(\rho_{j} \left| 2\alpha+\frac{j-1}{m}, \left(2-\frac{1}{m}\right)(j-1)\right) d\rho_{j} \right\}$$

$$= \frac{1}{\Gamma(2\alpha)} \prod_{j=2}^{m} \left\{ \frac{1}{\Gamma(2\alpha+(j-1)/m)\Gamma((2-1/m)(j-1))} \int_{0}^{1} \rho_{j}^{2\alpha+(j-1)/m-1}(1-\rho_{j})^{(2-1/m)(j-1)-1} d\rho_{j} \right\}.$$
(S7)

Then, by Gauss's multiplication formula for the gamma function, it follows that

$$\frac{\pi^{m/2} / \prod_{j=1}^{m} 2^{2(j-1)-1}}{\prod_{j=1}^{2m} \Gamma(\alpha + (j-1)/2)} \prod_{j=2}^{m} \Gamma((2-1/m)(j-1))
= \frac{2^{2m\alpha}}{\Gamma(2\alpha)} \prod_{j=2}^{m} \left\{ \frac{1}{\Gamma(2\alpha + (j-1)/m)} \int_{0}^{1} \rho_{j}^{2\alpha + (j-1)/m-1} (1-\rho_{j})^{(2-1/m)(j-1)-1} d\rho_{j} \right\}
= 2^{2m\alpha} \frac{(2\pi)^{(1-m)/2} m^{2m\alpha-1/2}}{\Gamma(2m\alpha)} \prod_{j=2}^{m} \int_{0}^{1} \rho_{j}^{2\alpha + (j-1)/m-1} (1-\rho_{j})^{(2-1/m)(j-1)-1} d\rho_{j}.$$
(S8)

Thus, if p = 2m,

$$\begin{aligned} &\frac{\pi^{p(p-1)/4}}{\Gamma_p(\alpha+(p-1)/2)} = \frac{1}{\prod_{j=1}^p \Gamma(\alpha+(j-1)/2)} \\ &= \frac{\left\{\prod_{j=1}^m 2^{2(j-1)-1}\right\}(2\pi)^{(1-m)/2}m^{-1/2}}{\pi^{m/2}\prod_{j=2}^m \Gamma((2-1/m)(j-1))} \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha+(j-1)/m-1}(1-\rho_j)^{(2-1/m)(j-1)-1} d\rho_j. \end{aligned}$$

On the other hand, if p = 2m - 1, we have, by the duplication formula,

$$\begin{split} &\frac{\pi^{p(p-1)/4}}{\Gamma_p(\alpha+(p-1)/2)} = \frac{1}{\prod_{j=1}^{2m-1} \Gamma(\alpha+(j-1)/2)} \\ &= \frac{1}{\prod_{j=1}^{2m-2} \Gamma(\alpha+(j-1)/2)} \frac{\Gamma(\alpha+m-1/2)}{\Gamma(\alpha+m-1)\Gamma(\alpha+m-1/2)} \\ &= \frac{1}{\prod_{j=1}^{2(m-1)} \Gamma(\alpha+(j-1)/2)} \frac{\Gamma(\alpha+m-1/2)}{\pi^{1/2}} \frac{2^{2(\alpha+m-1)-1}}{\Gamma(2(\alpha+m-1))} \\ &= \frac{1}{\pi^{(m-1)/2}} \Big\{ \prod_{j=1}^{m-1} \frac{2^{2(\alpha+j-1)-1}}{\Gamma(2(\alpha+j-1))} \Big\} \frac{\Gamma(\alpha+m-1/2)}{\pi^{1/2}} \frac{2^{2(\alpha+m-1)-1}}{\Gamma(2(\alpha+m-1))} \\ &= \frac{1}{\pi^{m/2}} \Big\{ \prod_{j=1}^{m} \frac{2^{2(\alpha+j-1)-1}}{\Gamma(2(\alpha+j-1))} \Big\} \int_{0}^{\infty} z^{\alpha+m-1/2-1} e^{-z} dz, \end{split}$$

where the fourth equality follows from (S6). Then, since, by (S7) and (S8),

$$2^{2m\alpha} \prod_{j=1}^{m} \frac{1}{\Gamma(2\alpha + 2(j-1))} \prod_{j=2}^{m} \Gamma((2-1/m)(j-1))$$

= $2^{2m\alpha} \frac{(2\pi)^{(1-m)/2} m^{2m\alpha-1/2}}{\Gamma(2m\alpha)} \prod_{j=2}^{m} \int_{0}^{1} \rho_{j}^{2\alpha + (j-1)/m-1} (1-\rho_{j})^{(2-1/m)(j-1)-1} d\rho_{j},$

we obtain

$$\begin{split} & \frac{\pi^{p(p-1)/4}}{\Gamma_p(\alpha+(p-1)/2)} \\ &= \frac{1}{\pi^{m/2}} \frac{\prod_{j=1}^m 2^{2(j-1)-1}}{\prod_{j=2}^m \Gamma((2-1/m)(j-1))} \int_0^\infty z^{\alpha+m-1/2-1} e^{-z} dz \\ &\quad \times 2^{2m\alpha} \frac{(2\pi)^{(1-m)/2} m^{2m\alpha-1/2}}{\Gamma(2m\alpha)} \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha+(j-1)/m-1} (1-\rho_j)^{(2-1/m)(j-1)-1} d\rho_j \\ &= \frac{\left\{ \prod_{j=1}^m 2^{2(j-1)-1} \right\} (2\pi)^{(1-m)/2} m^{-1/2}}{\pi^{m/2} \prod_{j=2}^m \Gamma((2-1/m)(j-1))} \\ &\quad \times \frac{(2m)^{2m\alpha}}{\Gamma(2m\alpha)} \left\{ \prod_{j=2}^m \int_0^1 \rho_j^{2\alpha+(j-1)/m-1} (1-\rho_j)^{(2-1/m)(j-1)-1} d\rho_j \right\} \int_0^\infty z^{\alpha+m-1/2-1} e^{-z} dz. \end{split}$$

This completes the proof.

S7 An alternative Metropolis-Hastings algorithm for the model of Section 3

For the model of Section 3, we can use the method of Miller (2019) and construct a MH algorithm. The posterior of θ , τ , and α is

$$p(\theta,\tau,\alpha|\mathbf{x}) \propto p(\alpha) \frac{p(\theta,\tau)}{\tau^{n/2}} \prod_{i=1}^{n} \left[\alpha^{\alpha} \frac{\Gamma(\alpha+1/2)}{\Gamma(\alpha)} / \left\{ \alpha + \frac{(x_i-\theta)^2}{2\tau} \right\}^{\alpha+1/2} \right]$$
$$= \frac{p(\theta,\tau)}{\tau^{n/2}} p(\alpha) \frac{\alpha^{n\alpha}}{\{\Gamma(\alpha)\}^n} \prod_{i=1}^{n} \int_0^\infty w_i^{\alpha+1/2-1} e^{-w_i \{\alpha + (x_i-\theta)^2/(2\tau)\}} dw_i$$

Then the full conditional distributions of (θ, τ) and w_i are as given in Section 3.1. The full conditional distribution of α under the prior $p(\alpha) = \text{Ga}(\alpha|a, b)$ can be written as

$$\operatorname{Ga}(\alpha|a,b) \underbrace{\prod_{i=1}^{n} \operatorname{Ga}(w_{i}|\alpha,\alpha)}_{\text{``likelihood''}}.$$

Therefore, we can apply the method of Miller (2019) to find numbers A, B > 0 such that $Ga(\alpha|A, B)$ is a good approximation to the full conditional density.

S8 Additional results for the simulation studies of Sections 3.2 and 4.2 of the main text

Additional results for the simulation studies of Sections 3.2 and 4.2 of the main text are in this section. First, Figures S1 and S3 correspond to the case of n = 30, whereas Figures S2 and S4 correspond to the case of n = 100, and these are as mentioned in the main text. Second, Tables S1 and S2 correspond to Table 1 of the main text and show results when we generate x_i from $t(x_i|(3,4), 2\alpha_0)$ and $t(x_i|(6,1), 2\alpha_0)$, whereas Tables S3 and S4 correspond to Table 2 of the main text. As in the main text, our method is better in terms of sESS for θ if n is small or if the prior for α is truncated. (Corresponding figures such as Figures 1 and 2 of the main text are omitted because they will not be so different.) Finally, Table S5 corresponds to Table 3 of the main text and shows results for two additional scenarios: (III) $\alpha_0 = (1, \ldots, 1)$ and (IV) $\alpha_0 = ((1/2, \ldots, 1/2), (1, \ldots, 1)) \in \mathbb{R}^5 \times \mathbb{R}^5$. It can be seen that our methods show good performance in terms of sESS.



Figure S1: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for n = 30.



Figure S2: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for n = 100.



Figure S3: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for $2\underline{\alpha} = 1,3$ for n = 30.



Figure S4: Boxplots of the effective sample sizes standardized by the computation times for the proposed method (DA) and the alternative method (A-MH) for $2\underline{\alpha} = 1,3$ for n = 100.

Table S1: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH) by Miller (2019), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method. Here, we consider the case of $(\theta, \tau) = (3, 4)$.

				ESS			sESS		М	SE rat	tio
n	$2\alpha_0$	method	θ	au	α	θ	au	α	θ	au	α
10	0.1	DA	832	362	1750	6828	2958	14213	_	_	—
10	0.1	A-MH	802	338	1860	6046	2522	13975	1.04	2.36	0.97
10	1	DA	1312	500	401	11263	4315	3457	_	-	-
10	1	A-MH	1314	523	516	10355	4133	4030	0.97	0.90	1.00
10	10	DA	1785	1053	263	15706	9261	2312	_	-	-
10	10	A-MH	1796	1234	444	14571	10034	3608	1.00	1.00	0.98
30	0.1	DA	641	150	1646	3998	936	10197	_	-	_
30	0.1	A-MH	645	148	1628	4121	950	10427	1.06	1.19	1.01
30	1	DA	1256	368	449	7938	2323	2833	_	_	_
30	1	A-MH	1274	400	583	8295	2612	3800	0.98	1.00	0.92
30	10	DA	2122	565	118	13852	3696	768	_	_	_
30	10	A-MH	2226	706	208	15259	4856	1421	1.01	1.00	0.90
100	0.1	DA	778	87	1298	2777	310	4622	_	_	_
100	0.1	A-MH	779	88	1315	3621	411	6118	1.00	1.02	0.99
100	1	DA	1346	359	489	4922	1309	1784	_	_	_
100	1	A-MH	1340	374	609	6311	1762	2860	1.00	1.00	1.01
100	10	DA	2646	221	57	10114	849	217	_	-	-
100	10	A-MH	2692	337	102	13679	1724	516	1.01	1.00	0.80

Table S2: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH) by Miller (2019), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method. Here, we consider the case of $(\theta, \tau) = (6, 1)$.

				ESS			sESS			ISE rat	io
n	$2\alpha_0$	method	θ	au	α	θ	au	α	θ	au	α
10	0.1	DA	1000	404	2166	8477	3416	18278	_	_	_
10	0.1	A-MH	894	425	2233	6922	3268	17200	0.68	35.33	1.00
10	1	DA	1823	1010	471	15689	8705	4030	—	—	—
10	1	A-MH	1878	1081	626	14862	8569	4931	1.01	1.03	1.10
10	10	DA	2745	3004	313	23912	26157	2728	—	—	—
10	10	A-MH	2780	3067	575	22710	25109	4711	1.01	1.02	1.00
30	0.1	DA	1100	243	2542	6856	1520	15867	_	—	_
30	0.1	A-MH	1107	237	2631	7147	1531	16981	0.97	0.70	1.02
30	1	DA	1645	577	585	10488	3674	3727	_	_	_
30	1	A-MH	1666	591	753	10886	3848	4889	1.01	1.07	0.97
30	10	DA	2618	1122	127	17030	7263	827	_	—	—
30	10	A-MH	2674	1310	232	18308	9004	1588	1.01	0.98	0.95
100	0.1	DA	1007	140	2399	3608	500	8585	_	_	_
100	0.1	A-MH	995	130	2448	4621	604	11335	1.01	0.94	0.99
100	1	DA	1415	449	581	5227	1652	2132	_	_	_
100	1	A-MH	1427	454	742	6727	2143	3493	0.99	0.99	0.97
100	10	DA	2794	349	56	10786	1354	218	_	_	_
100	10	A-MH	2857	477	101	14736	2466	520	1.01	1.00	0.91

Table S3: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method for $2\underline{\alpha} = 1, 3$. Here, we consider the case of $(\theta, \tau) = (3, 4)$.

				ESS			sESS		Μ	SE rat	tio
n	$2\underline{\alpha}$	method	θ	au	α	θ	au	α	θ	au	α
10	1	DA	1985	1287	275	6088	3946	842	_	-	-
10	1	A-MH	2046	1445	467	3432	2438	784	0.99	1.00	0.90
10	3	DA	2702	2286	370	8343	7052	1142	_	-	-
10	3	A-MH	2690	2309	660	4610	3962	1130	0.99	1.00	0.94
30	1	DA	2167	505	119	5952	1391	327	_	_	_
30	1	A-MH	2147	623	209	3686	1077	357	0.99	1.00	0.92
30	3	DA	2516	979	136	6908	2689	374	_	-	-
30	3	A-MH	2611	1153	245	4426	1960	415	1.01	1.00	1.04
100	1	DA	2559	211	56	5462	450	118	_	-	-
100	1	A-MH	2582	316	96	4125	506	153	0.99	1.00	0.94
100	3	DA	2729	306	59	5803	649	124	_	_	_
100	3	A-MH	2752	436	106	4357	692	168	0.99	1.00	0.98

Table S4: The averages of the effective sample sizes (ESS) for the proposed method (DA) and the alternative method (A-MH), the averages of those standardized by computation time (sESS), and the ratios of the mean squared errors (MSE) of the A-MH method to those of the DA method for $2\underline{\alpha} = 1, 3$. Here, we consider the case of $(\theta, \tau) = (6, 1)$.

				ESS			sESS		М	SE rat	tio
n	$2\underline{\alpha}$	method	θ	au	α	θ	au	α	θ	au	α
10	1	DA	2871	3015	325	8880	9324	1004	_	_	_
10	1	A-MH	2852	2982	594	4905	5134	1019	1.01	0.99	0.93
10	3	DA	3121	3228	393	9704	10043	1221	_	_	-
10	3	A-MH	3142	3211	718	5379	5500	1229	1.00	1.00	0.96
30	1	DA	2725	1272	130	7561	3531	359	_	-	-
30	1	A-MH	2735	1534	237	4798	2696	415	1.00	1.01	1.06
30	3	DA	2868	1658	141	8025	4638	394	_	_	-
30	3	A-MH	2895	1902	262	5091	3356	460	0.99	1.00	1.00
100	1	DA	2808	333	56	6088	724	122	_	-	-
100	1	A-MH	2793	437	101	4597	719	166	1.00	1.03	0.89
100	3	DA	2833	336	60	6114	726	130	_	-	-
100	3	A-MH	2836	502	104	4629	823	168	1.00	0.99	1.01

Table S5: The average effective sample size (ESS), the average computation time (CT), the standardized effective sample size by the computation time (sESS), and the mean squared error (MSE) for the proposed data-augmentation method with normal latent variables (DA-N), Poisson latent variables (DA-P), and the PTN sampler of He et al. (2021) (DA-PT) and the original method proposed by He et al. (2021) (ERG). These values are averaged over $\alpha_0, \ldots, \alpha_9$. MSE values under n = 100 and n = 1000 are multiplied by 10^3 and 10^4 , respectively.

\overline{n}	Scenario	method	ESS	CT	sESS	MSE
100	(III)	DA-N	311	2.1	151	10.37
		DA-P	448	1.7	257	10.17
		DA-PT	593	1.8	323	10.19
		ERG	920	82.4	11	10.16
100	(IV)	DA-N	564	2.1	272	6.40
		DA-P	672	1.8	381	6.51
		DA-PT	790	1.8	431	6.32
		ERG	1256	82.4	15	6.35
1000	(III)	DA-N	308	7.5	41	9.65
		DA-P	444	7.1	63	9.66
		DA-PT	598	7.1	84	9.66
		ERG	918	856.7	1	9.63
1000	(IV)	DA-N	560	7.5	74	6.20
		DA-P	670	7.2	93	6.12
		DA-PT	751	7.2	104	6.18
		ERG	1231	834.8	1	6.15