

TIME EVOLUTION OF THE CMB QUADRUPOLE

ABRAHAM LOEB¹

¹*Astronomy Department, Harvard University, 60 Garden St., Cambridge, MA 02138, USA*

ABSTRACT

I show that the quadrupole of the Cosmic Microwave Background (CMB) evolves more rapidly than previously expected, as a result of the acceleration of the Sun towards the Galactic center. The acceleration, measured most recently by *Gaia* EDR3, implies a fractional change in the quadrupole of $\sim 10^{-9}$ per year, an order of magnitude larger than expected from the evolution in the last scattering surface of the CMB.

INTRODUCTION

Within a timescale of order the age of the Universe, $\sim 10^{10}$ yr, the large-scale anisotropies of the Cosmic Microwave Background (CMB) are expected to change as a result of the change in the location of the last scattering surface (Lange & Page 2007; Zibin et al. 2007; Moss et al. 2008). Naively, one would expect anisotropies on the largest scales - such as the quadrupole moment, to change most slowly. However, here we show this not to be true as a result of the acceleration of the Sun towards the center of the Milky Way. This acceleration was measured most recently by *Gaia* EDR3 to be, $\dot{v} = 2.32(\pm 0.16) \times 10^{-8}$ cm s $^{-2}$, towards the Galactic center (Gaia Collaboration et al. 2021).

CALCULATION

The velocity of the Sun relative to the cosmic frame of reference, \vec{v} , results in a CMB dipole moment, which is routinely removed from CMB anisotropy maps. But to second-order in $\vec{\beta} \equiv (\vec{v}/c)$, the motion also leads to a quadrupole anisotropy with an angular dependence of $(\cos^2 \theta - 1/3)$, where θ is the angle between the velocity vector \vec{v} and the photon direction.

The fractional change in the CMB intensity as a result of the kinematic quadrupole depends on photon frequency (Kamionkowski & Knox 2003),

$$\left(\frac{\Delta I_\nu}{I_\nu}\right)_Q = F(x)\beta^2, \quad (1)$$

with $F(x) = [xe^x/(e^x - 1)][(x/2) \coth(x/2)]$. Here, $x = (h\nu/kT)$ with $I_\nu = [2(kT)^3/(hc)^2]x^3/(e^x - 1)$ being the CMB blackbody intensity at a frequency ν and the mean CMB temperature $T = 2.725$ K (Fixsen 2009). At the low frequencies of the Rayleigh-Jeans regime where $x \ll 1$, we get $F(x) \approx 1$ and $(\Delta I_\nu/I_\nu) = (\Delta T/T)$.

CONCLUSIONS

The time derivative of equation (1) yields a kinematic variation in the quadrupole, which for $x \ll 1$ is given by,

$$\left(\frac{\dot{T}}{T}\right)_Q = 2\vec{\beta} \cdot \dot{\vec{\beta}}, \quad (2)$$

where from the *Gaia* measurement, $\dot{\beta} = 2.44 \times 10^{-11}$ yr $^{-1}$, and from the CMB dipole $\beta \approx 1.23 \times 10^{-3}$ (Kogut et al. 1993; Fixsen et al. 1994). The dot-product of the acceleration vector towards the Galactic center and the CMB dipole velocity vector yields a geometric projection factor of 0.065 (Loeb & Narayan 2008).

Given the quadrupole moment measured by the Planck satellite, $Q \equiv (\Delta T/T)_Q \sim 4.5 \times 10^{-6}$ (Notari & Quartin 2015), the fractional time derivative of the CMB quadrupole, as a result of the acceleration of the Sun towards the Galactic center, is

given by,

$$\left(\frac{\dot{Q}}{Q}\right) \approx 10^{-9} \text{ yr}^{-1}. \quad (3)$$

This rate of evolution is over an order of magnitude larger than expected from the change in the last scattering surface of the CMB (Lange & Page 2007; Zibin et al. 2007; Moss et al. 2008).

The expected time evolution of the CMB quadrupole spectrum, $F(x)$, in equation (1), could potentially be measured - especially for $x \gg 1$ where $F(x) \gg 1$, with future CMB spectral distortion experiments (Chluba et al. 2021).

ACKNOWLEDGEMENTS

This work was supported in part by Harvard's *Black Hole Initiative*, which is funded by grants from JFT and GBMF.

REFERENCES

- Chluba, J., Abitbol, M. H., Aghanim, N., et al. 2021, *Experimental Astronomy*, 51, 1515, doi: [10.1007/s10686-021-09729-5](https://doi.org/10.1007/s10686-021-09729-5)
- Fixsen, D. J. 2009, *ApJ*, 707, 916, doi: [10.1088/0004-637X/707/2/916](https://doi.org/10.1088/0004-637X/707/2/916)
- Fixsen, D. J., Cheng, E. S., Cottingham, D. A., et al. 1994, *ApJ*, 420, 445, doi: [10.1086/173575](https://doi.org/10.1086/173575)
- Gaia Collaboration, Klioner, S. A., Mignard, F., et al. 2021, *A&A*, 649, A9, doi: [10.1051/0004-6361/202039734](https://doi.org/10.1051/0004-6361/202039734)
- Kamionkowski, M., & Knox, L. 2003, *PhRvD*, 67, 063001, doi: [10.1103/PhysRevD.67.063001](https://doi.org/10.1103/PhysRevD.67.063001)
- Kogut, A., Lineweaver, C., Smoot, G. F., et al. 1993, *ApJ*, 419, 1, doi: [10.1086/173453](https://doi.org/10.1086/173453)
- Lange, S., & Page, L. 2007, *ApJ*, 671, 1075, doi: [10.1086/523097](https://doi.org/10.1086/523097)
- Loeb, A., & Narayan, R. 2008, *MNRAS*, 386, 2221, doi: [10.1111/j.1365-2966.2008.13187.x](https://doi.org/10.1111/j.1365-2966.2008.13187.x)
- Moss, A., Zibin, J. P., & Scott, D. 2008, *PhRvD*, 77, 043505, doi: [10.1103/PhysRevD.77.043505](https://doi.org/10.1103/PhysRevD.77.043505)
- Notari, A., & Quartin, M. 2015, *JCAP*, 2015, 047, doi: [10.1088/1475-7516/2015/06/047](https://doi.org/10.1088/1475-7516/2015/06/047)
- Zibin, J. P., Moss, A., & Scott, D. 2007, *PhRvD*, 76, 123010, doi: [10.1103/PhysRevD.76.123010](https://doi.org/10.1103/PhysRevD.76.123010)