

# Anisotropic compact stars in $D \rightarrow 4$ limit of Gauss-Bonnet gravity

G.G.L. Nashed,<sup>1, 2,\*</sup> S.D. Odintsov,<sup>3, 4, †</sup> and V.K. Oikonomou<sup>5, 2, ‡</sup>

<sup>1</sup>*Centre for Theoretical Physics, The British University,  
P.O. Box 43, El Sherouk City, Cairo 11837, Egypt*

<sup>2</sup>*Laboratory for Theoretical Cosmology, Tomsk State University of  
Control Systems and Radioelectronics (TUSUR), 634050 Tomsk, Russia*

<sup>3</sup>*Institut de Ciències de l’Espai (ICE-CSIC/IEEC),  
Campus, c. Can Magrans s/n, 08193, Barcelona, Spain*

<sup>4</sup>*Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain*

<sup>5</sup>*Department of Physics, Aristotle University of Thessaloniki, Thessaloniki 54124, Greece*  
(Dated: March 7, 2022)

In the frame of Gauss-Bonnet gravity and in the limit of  $D \rightarrow 4$ , based on the fact that spherically symmetric solution derived using any of regularization schemes will be the same form as the original theory [1–4], we derive a new interior spherically symmetric solution assuming specific forms of the metric potentials that have two constants. Using the junction condition we determine these two constants. By using the data of the star EXO 1785 – 248, whose mass is  $M = 1.3 \pm 0.2 M_{\odot}$  and radius  $l = 8.849 \pm 0.4 \text{ km}$ , we calculate the numerical values of these constants, in terms of the dimensionful coupling parameter of the Gauss-Bonnet term, and eventually, we get real values for these constants. In this regard, we show that the components of the energy-momentum tensor has a finite value at the center of the star as well as a smaller value to the surface of the star. Moreover, we show that the equations of the state behave in a non-linear way due to the impact of the Gauss-Bonnet term. Using the Tolman-Oppenheimer-Volkoff equation, the adiabatic index, and stability in the static state we show that the model under consideration is always stable. Finally, the solution of this study is matched with observational data of other pulsars showing satisfactory results.

PACS numbers: 11.30.-j; 04.50.Kd; 97.60.Jd.

Keywords: Modified gravity; neutron stars; stellar structure

## I. INTRODUCTION

To date a considerable amount of observational data of compact stellar objects data, such as neutron stars and spherically symmetric black holes, are available from gravitational-wave (GW) detectors, like X-ray observations, advanced LIGO [5], Kagra [6, 7], Advanced Virgo [8], and from the networks of radio telescopes, like the event horizon telescope [9], from the NICER mission [10] and there is much activity in the study of such objects also, i.e., neutron stars in  $F(R)$  [11, 12]. In the frame of current observational uncertainties, all data are consistent with Einstein’s general relativity (GR) and despite this. However, there is a venue for a extended gravity theories, which might be useful to future observations which require alternative descriptions than that of Einstein’s GR theory, like for example the curious GW190814 event [13]. Regardless of the direct tests of Einstein GR, there are also pending questions to be clarified, like the maximum mass of neutron stars, the high-density equation of state (EoS), and so on. The observations obtained recently by the compact object merger GW190814 [13] shows that the low-mass component of the binary, possesses a mass of  $2.59^{+0.08}_{-0.09} M_{\odot}$ , which directly puts it in the present observational mass gap between neutron stars and black holes. Moreover, among the gravitational waves, there are three events/candidates involving the merger of one neutron star or light mass black hole with another compact object. If this event is described solely by GR, the lower mass component of the binary can either be a neutron star with an unexpectedly stiff (or exotic) EoS, a black hole with an unexpectedly small mass, or a neutron star with an unexpectedly rapid rotation [14–34]. On the other hand, an natural and non-exotic description of the lower mass component of the event GW190814 can be given by extended gravity [35–37]. In the frame of modified gravitational theories there are many studied tackled the problem of anisotropic compact stars for example in the frame of  $f(R, G)$ , where  $R$  is the Ricci scalar and  $G$  is the Gauss-Bonnet term [38], in the frame of mimetic theory [39], in the frame of  $f(R)$  [40–43], in the frame of teleparallel gravity [44, 45], and in the frame of scalar Gauss-Bonnet theory [46].

To date, two GW outcomes have been verified as binary neutron star (BNS) mergers, GW170817 [47] and GW190425 [48], and extra outcomes are anticipated in the near future [49]. Revealing of the GW from the inspiral phase

---

\* [nashed@bue.edu.eg](mailto:nashed@bue.edu.eg)

† [odintsov@ieec.uab.es](mailto:odintsov@ieec.uab.es)

‡ [v.k.oikonomou1979@gmail.com](mailto:v.k.oikonomou1979@gmail.com)

of GW170817, in connection with observations of its kilonova electromagnetic counterpart [50–52], have put new constraints on the dimensionless tidal deformability of neutron stars and hence on their EoS [53–59]. Such EoS constraints are expected to be more precise in the next years, via the combination of a larger number of observations [60–64]. Despite the sensitivity of the Advanced LIGO and Advanced Virgo detectors, these were not sensitive enough in order to determine the post-merger phase in GW170817 [47, 65]. However, this post-merger phase may be revealed in the near future, by several experiments and collaborations, like [49], or by third generation detectors [66, 67], or even with high-frequency detectors [68, 69]. The observation of GW in the post-merger phase of a BNS merger would then offer another opportunity to probe the high-density EoS of NSs, see [70–81].

In view of the future observational data which might indicate deviations from GR, it is important to study compact objects in the context of extended theories of gravity. In the present work, we will study compact objects in the context of Einstein-Gauss-Bonnet gravity (EGB) [82, 83] which provides a number of attractive analytic solutions. The action of this theory is given as:

$$S = \frac{1}{2\kappa} \int d^4x \sqrt{-g} \{R + \beta G\} + S_m, \quad (1)$$

with  $\kappa = 8\pi G/c^4$ , and  $G = R^2 - 4R_{\mu\nu}R^{\mu\nu} + R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$  is the Gauss-Bonnet scalar while  $S_m$  is the matter Lagrangian. Here  $\beta$  has a dimension of length squared. We will then formulate models of neutron stars, and by using numerical methods we will study several solutions for neutron stars.

Lately, Glavan and Lin [84] provided a new general covariant modified model of the EGB theory that elucidates multiple non-trivial issues taking place in 4-dimensional spacetime. Note that such scenario to get consistent 2-dimensional GR from D-dimensional GR in the limit of  $D \rightarrow 2$  was proposed in [85, 86]. The GB expression ( $G$ ) in 4-dimensions is a total derivative term and hence does not impact the gravitational dynamics. However, on a higher dimension, i.e.,  $D > 4$ , it involves an exterior result of the total derivative. When  $D > 4$ , Torii et al. [87] and, Mardones et al. [88] showed that the support of the GB term is proportional to a vanishing factor in 4-dimensions. Moreover, Glavan and Lin [84] rescaled the dimensional parameter  $\beta$ , i.e.,  $\beta \rightarrow \frac{\beta}{(D-4)}$  showing a very much identical to GR behavior, as it preserves equal degrees of freedom in all dimensions and sets aside the Ostrogradsky instability [89]. Using the regularization of the GB expression at  $D = 4$ , one can derive a non-trivial impact in the four-dimensional dynamics [90].

Here, we stress the fact that the novel 4D EGB theory [84] has got several quibbles. The criticisms on the correctness by taking the limit  $D \rightarrow 4$  have been discussed in [91, 92]. Nevertheless, it was shown in Refs. [93–95] that in 4-dimensions spacetime the output field equations are not regular and no regular action which creates the proposed regularized field equations can be exist [96]. Kaluza-Klein-reduction method of the limit  $D \rightarrow 4$  yields a special class of scalar-tensor theories in the Horndeski family [97, 98]. Similar method was used in [2, 99, 100] by adding a counter term in D-dimensions and then taking  $D \rightarrow 4$  limit. Furthermore, many projects have been analyzed the previous issues coming from the new 4-dimensions EGB gravity. Depending on the selection of the “regularization method”, several theories have been presented with various degrees of freedom and various characteristic. Using the procedure of [4, 91], the handling was shown to be symmetrical by ignoring the 4-dimensions diffeomorphism invariance. Nevertheless, it is interesting to note that spherically symmetric 4-dimensions solutions are valid in these regularized theories [1]. Actually, black hole solutions through the rescaling method [84] still correct in these regularized theories [2, 3]. Therefore, it becomes clear that the spherically symmetric solution derived using any of these regularization schemes will be identical to the authentic theory. In this frame, the interior spherically symmetric star solution is significant to study. Therefore, we decide to obtain the field equations starting with outlines depending on the new 4-dimensions EGB gravity. Moreover, one of the motivation of the present study is to constrains the numerical value of Gauss-Bonnet parameter in the frame of anisotropic model so that we get a realizable astrophysical model. Additionally, we want to test the behavior of the equation of state in this model and compared it with the Einstein general relativity.

The GB theory has various applications in different topics of gravity and therefore, it appeared to be quite unconventional to scientists in the field of cosmology and astrophysics. Glavan and Lin [84] in the frame of GB theory derived a static spherically symmetric vacuum black hole which was proved to be quite different from Schwarzschild black holes. Gurses et. al [95] showed that EGB theory does not support any intrinsic 4-dimensional solution in terms of the metric, thus its viability of the black hole solutions is appropriate for the case  $D > 4$ . More applications of EGB theory were used to calculate the thermodynamics of non-rotating and rotating black holes and evaluate the thermodynamical quantities as entropy, mass, temperature [101–104]. Moreover, EGB theory is used for the study of various black hole related issues, like thermodynamics of AdS black holes [105], quasi-normal modes [106], black gravitational Lensing of strong and weak types [101], Hawking radiation [107], geodesic motions for spinning test particles [108], and wormhole solutions [109]. Aside from the issues of the black hole, EGB theory is applied for the study of neutron stars, which is a subject of continuous research interest. Recent observations of neutron stars have further constrained the equation of state of nuclear matter, but the inner structure of neutron stars still remains a

mystery. An investigation of the photon sphere and the shadow observed by a distant observer and the exploration of the effects of black hole parameters have been studied in the frame of EGB [110].

The determination of the inner core matter is quite non-trivial as the composition varies with the nature of interaction and quark matter, hyperon matter, Bose-Einstein condensate, strange mesons are all considered to be the core matter. But the invention of  $2M_{\odot}$  Neutron stars like PSR J1614-2230 [ $1.97 \pm 0.04M_{\odot}$ ] [14] and J0348+0432 [ $2.01 \pm 0.04M_{\odot}$ ] [111] brought criterion on the nature of the core matter. Quantum chromodynamics [QCD], supports the conversion of hadronic matter into deconfined quarks inside the neutron stars. This invention makes the Bodmer-Witten hypothesis quite important to the context as it already predicted for a quark-matter component in the cosmic rays detected from neutron stars [112, 113]. This strengthens the presence of quark stars.

The present study aims to derive an interior solutions for EGB theory, using the fact that spherically symmetric solution obtained from any regularization methods will be identical with the original theory [1–3], and then check the physical viability of such a solution. According to the procedure, it is possible to construct self-consistent compact stellar models capable of describing anisotropic neutron stars in the context of EGB gravity. The layout of the present paper is as follows: In Sec. II, a summary of EGB theory is reported. Also in Sec. II, we apply the field equation of EGB to  $D$ -dimensional spherically symmetric spacetime and write them in the limit  $D \rightarrow 4$  and solve the resulting differential equations assuming specific forms of the metric potentials. In Sec. III, we use the junction condition, i.e. match the interior solution with the exterior solution derived [84], and fix the two constants that characterize our solution. In Sec. IV, we discuss the physical conditions that any real compact star must satisfy and show that the energy components of our model have finite values at the center of the star and decrease towards the star of the surface. Moreover, we show that the model under consideration has a positive anisotropy which means that the tangential pressure is greater than the radial one. Also, we prove that our model satisfies the causality condition and the energy-momentum conditions. Finally, we study the equation of state, (EoS), and show that the impact of Gauss-Bonnet expression makes its pattern has a non-linear form. In Sec. V we show that our model is stable under various conditions, like TOV equation, adiabatic index, and the static state. We confront our results with observational data using and we tabulated our results in Tables I and II. Finally, the conclusions follow in the end of the paper<sup>1</sup>.

## II. COMPACT STAR IN GAUSS-BONNET THEORY

The Lagrangian of Einstein-Gauss-Bonnet, in  $D$ -dimensional spacetime has the form:

$$\mathcal{L}_{EGB} = \mathcal{R} + \frac{\beta}{D-4} G + \mathcal{L}_m, \quad (2)$$

whose corresponding action takes the form:

$$S_{EGB} = \frac{1}{16\pi} \int d^D x \sqrt{-g} \left( \mathcal{R} + \frac{\beta}{D-4} G \right) + S_{matter}, \quad (3)$$

where  $\mathcal{R}$  is the Ricci scalar curvature and  $G$  is the Gauss-Bonnet invariant associated with the  $D$ -dimensional spacetime. The dimensionful parameter  $\beta$  is the Gauss-Bonnet coupling parameter and  $S_{matter}$  is the action contributed from the matter perfect fluid. The Gauss-Bonnet term,  $G$ , takes the form:

$$G = \mathcal{R}^2 - 4\mathcal{R}_{\mu\nu}\mathcal{R}^{\mu\nu} + \mathcal{R}_{\mu\nu\kappa\lambda}\mathcal{R}^{\mu\nu\kappa\lambda}, \quad (4)$$

where  $R_{\mu\nu}$  and  $\mathcal{R}_{\mu\nu\kappa\lambda}$  are the Ricci curvature and Riemann curvature tensors, respectively. Variation of the action (3) with respect to the metric tensor yields:

$$\mathcal{G}_{\mu\nu} + \frac{\beta}{N-4} \mathcal{H}_{\mu\nu} = 8\pi \mathcal{T}_{\mu\nu}, \quad (5)$$

where the Einstein tensor  $\mathcal{G}_{\mu\nu}$ , the Lanczos tensor  $\mathcal{H}_{\mu\nu}$  and the stress tensor  $\mathcal{T}_{\mu\nu}$  are defined respectively as:

$$\mathcal{G}_{\mu\nu} = \mathcal{R}_{\mu\nu} - \frac{1}{2} \mathcal{R} g_{\mu\nu}, \quad (6)$$

$$\mathcal{H}_{\mu\nu} = 2 \left( \mathcal{R}\mathcal{R}_{\mu\nu} - 2\mathcal{R}_{\mu\lambda}\mathcal{R}_{\nu}^{\lambda} - 2\mathcal{R}_{\mu\lambda\nu\rho}\mathcal{R}^{\lambda\rho} - \mathcal{R}_{\mu\alpha\beta\gamma}\mathcal{R}_{\nu}^{\alpha\beta\gamma} \right) - \frac{1}{2} g_{\mu\nu} G, \quad (7)$$

$$\mathcal{T}_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta(\sqrt{-g} S_m)}{\delta g^{\mu\nu}}. \quad (8)$$

---

<sup>1</sup> In this study, we are going to use the geometrized units units.

The field equation (5) has no meaning when  $D = 4$ . In spite of this, Glavan et. al., [84, 114] show that using a re-scale of the parameter  $\beta$  as  $\beta \rightarrow \beta/(D-4)$  and by specifying spacetimes of curvature scale  $K$  which are maximally symmetric that set the variation of GB contribution as:

$$\frac{g_{\mu\lambda}}{\sqrt{-g}} \frac{\delta G}{\delta g_{\nu\lambda}} = \frac{\beta(D-2)(D-3)}{2(D-1)} K^2 \delta_\mu^\nu, \quad (9)$$

that shows in a clear way the non-vanishing of  $G$  when  $D = 4$ .

Now we consider a spacetime of  $D$ -dimensions that has the form:

$$ds^2 = -e^{2h(r)} dt^2 + e^{2h_1(r)} dr^2 + r^2 d\Omega_{D-2}^2, \quad (10)$$

where,  $d\Omega_{D-2}^2$  refers to the  $(D-2)$ -dimensional surface of a unit sphere.

As shown in [84], the effective 4-dimensional theory is obtained in the singular limit  $D \rightarrow 4$  with  $\beta = \beta/(D-4)$ . Using the limit  $D \rightarrow 4$  we derive the dimensionally reduced field equations as:

$$8\pi\rho = \frac{\beta(1-e^{-2h_1})}{r^3} \left[ 4h'_1 e^{-2h_1} - \frac{(1-e^{-2h_1})}{r} \right] + e^{-2h_1} \left( \frac{2h'_1}{r} - \frac{1}{r^2} \right) + \frac{1}{r^2}, \quad (11)$$

$$8\pi p_r = \frac{\beta(1-e^{-2h_1})}{r^3} \left[ 4h' e^{-2h_1} + \frac{(1-e^{-2h_1})}{r} \right] + e^{-2h_1} \left( \frac{2h'}{r} + \frac{1}{r^2} \right) - \frac{1}{r^2}, \quad (12)$$

$$\begin{aligned} 8\pi p_\theta &= e^{-2h_1} \left[ h'' + h'^2 + \frac{1}{r}(h' - h'_1) + h'h'_1 \left( \frac{8\beta e^{-2h_1}}{r^2} - 1 \right) - \frac{2\beta(1-e^{-2h_1})}{r^2} \right. \\ &\quad \times \left. \left\{ \frac{1}{r}(h' - h'_1) - 2(h'' + h'^2 - h'h'_1) + \frac{1}{r^2}(e^{2h_1} - 1) \right\} \right], \end{aligned} \quad (13)$$

where the “prime” indicates differentiation with respect to the radial coordinate.

Moreover, we suppose that the stress energy-momentum tensor of anisotropic fluid has the form:

$$\mathcal{T}_{\mu\nu} = (\rho + p_\theta) u_\mu u_\nu + p_\theta g_{\mu\nu} + (p_r - p_\theta) \xi_\nu \xi^\mu, \quad (14)$$

where  $\rho$  is the energy-density,  $p_r$  and  $p_\theta$  are the radial and tangential pressures, where  $u^\mu$  is the time-like vector defined as  $u^\mu = [1, 0, 0, 0]$  and  $\xi_\mu$  is the unit radial vector with its spacelike property, defined by  $\xi_\mu = [0, 1, 0, 0]$  such that  $u^\mu u_\mu = -1$  and  $\xi^\mu \xi_\mu = 1$ . In the present study, we shall consider the matter content described by the energy density  $\rho$ , radial and tangential pressures,  $p_r$  and  $p_\theta$  respectively<sup>2</sup>. quantities

The above system of differential equations, (11)–(13), constitutes three differential equations with five variables, thus we need two extra conditions. In this study we are going to assume the metric potentials,  $g_{tt}$  and  $g_{rr}$  to have special forms so that the resulting model can be consistent with a real compact star. Here we assume

$$h(r) = -4 \ln(c_1 - c_0 r^2), \quad h_1(r) = -4 \ln(1 + c_0 r^2), \quad (15)$$

where  $c_0$  and  $c_1$  are constants that will be fixed from the junction conditions. We assume the ansatz given by Eq. (15) so that we can get non-singular values of energy-density, radial and tangential pressures and also finite values of these quantities at the center of the star as we will show in the following sections. Using Eq. (15) in Eqs. (11)–(13) we get the form of energy-density, radial and tangential pressures which are quite lengthy to quote them here and therefore, we present these in Appendix A. In the next section we are going to fix the two constants using the junction conditions, i.e., we will match the interior solution with exact exterior solution.

### III. JUNCTION CONDITIONS

The vacuum spherically symmetric solution in Gauss-Bonnet gravitational theory is derived in [84] and has the following form in the limit  $D \rightarrow 4$ :

$$ds^2 = -A(r) dt^2 + \frac{dr^2}{A(r)} + r^2 d\Omega_2^2, \quad (16)$$

---

<sup>2</sup> Throughout this study we shall use geometrized units in which  $G = c = 1$ .

where, the unknown function  $A(r)$  has the form

$$A(r) = 1 + \frac{r^2}{32\pi\beta} \left[ 1 \pm \left\{ 1 + \frac{128\pi\beta m}{r^3} \right\}^{1/2} \right]. \quad (17)$$

In this study, we consider the negative branch in Eq. (17) that yields<sup>3</sup>:

$$A(r) = 1 + \frac{r^2}{32\pi\beta} \left[ 1 - \left\{ 1 + \frac{128\pi\beta M}{r^3} \right\}^{1/2} \right] \approx 1 - \frac{2M}{r} + \frac{64\pi\beta M^2}{r^4}. \quad (18)$$

Equation (18) shows that the direct effect of the GB parameter will be of order  $\mathcal{O}(r^{-4})$  and in case we neglect it we will return to the usual Schwarzschild solution. Now matching the interior and exterior spacetimes at the boundary  $r = l$ , i.e.,  $e^{2h(l)} = e^{-2h_1(l)} = A(l)$  yields the value of the two constants  $c_0$  and  $c_1$  which are:

$$c_0 = \frac{\sqrt{2} \sqrt[4]{\frac{2(32\pi\beta + l^2 - l^2\sqrt{\frac{l^3 + 128\pi\beta M}{l^3}})}{\pi\alpha}} - 4}{4l^2}, \quad (19)$$

$$c_1 = \frac{1}{2\sqrt[4]{\pi\beta} \parallel \mathbb{B} \parallel [2\sqrt[4]{2} \parallel \mathbb{B} \parallel -4\sqrt[4]{\pi}\sqrt[4]{\beta}]} \left\{ \sqrt{2} \parallel \mathbb{B}^3 \parallel -8\sqrt[4]{2\pi\beta}\mathbb{B}^2 + 8\sqrt{\pi\beta}\mathbb{B} \right. \\ \left. - 8\sqrt{\pi\beta}\sqrt[4]{[8\sqrt{\pi\beta} - 8\sqrt[4]{2\pi\beta} \parallel \mathbb{B} \parallel + \sqrt{2}\mathbb{B}^2][2\sqrt[4]{2} \parallel \mathbb{B} \parallel -4\sqrt[4]{\pi\beta}]} \right\}, \quad (20)$$

where  $\mathbb{B} = \sqrt[4]{\sqrt{l}\sqrt{l^3 + 128\pi\beta M} - 32\pi\beta - l^2}$ . Now we are going to calculate the derivative of the metric potentials (15) and the exterior solution given by Eq. (18) and get:

$$g'_{tt}(r) = \frac{dg_{tt}(r)}{dr} = \frac{8rc_0}{(c_1 - c_0 r^2)^5}, \quad g'_{rr}(r) = \frac{dg_{rr}(r)}{dr} = -\frac{8rc_0}{(1 + c_0 r^2)^5}, \\ A'(r) = \frac{dA(r)}{dr} = \frac{r^{3/2}\sqrt{r^3 + 128\pi\beta M} + r^3 + 32\pi\beta M}{\pi\beta\sqrt{r^4 + 128r\pi\beta M}}. \quad (21)$$

The second junction condition should be checked as well which deals with the first derivative of the metric with respect to the radial coordinate. Having the results in Eq. (21), one can check the second junction condition to see whether it trivially satisfies or one needs to consider a shell with an appropriate energy-momentum tensor.

#### IV. PHYSICS OF THE SOLUTION GIVEN IN APPENDIX A

Now, we are ready to examine if the interior solution which is given in Appendix A can describe a realistic star. For such a purpose, we are going to derive some necessary conditions which must be satisfied in order to have a real star.

##### A. Energy-momentum tensor

For any real interior solution, we must have a positive value of all the components of the energy-momentum tensor, i.e., the energy-density, and the radial, and transverse pressures should have positive well-defined values. Additionally, all these components must be finite at the center of the star and should decrease in the radial direction towards the surface of the star. Finally, the tangential pressure should exceed the radial one at the center of the star. In the present study we shall consider the stellar model of the observed pulsar EXO 1785–248 whose mass is  $M = 1.3 \pm 0.2 M_\odot$  and its radius  $l = 8.849 \pm 0.4 \text{ km}$  for which the constants  $c_0$  and  $c_1$  can be calculated numerically. We must stress that the model solution given in Appendix A cannot be reduced to the GR solution because the parameter  $\beta$  is not allowed to take zero values. We depict the components of energy-momentum in Fig. 1. The numerical values of the parameters

<sup>3</sup> The matching condition of the positive branch of Eq. (17) yields unphysical quantities, that is, negative values of density, radial and tangential pressures. So we will not consider this case in this study.




Figure 1. Plot of the components of the energy-momentum tensor and anisotropic force where we put  $\beta = -0.0001$  and the constants  $c_0$  and  $c_1$  take the numerical values,  $-0.001763358117$  and  $-0.9329227851$  respectively.

$c_0$  and  $c_1$  used in Fig. 1 are  $c_0 = -0.001763358117$ ,  $c_1 = -0.9329227851$ . Moreover,  $\rho(r \rightarrow 0) = 0.00168473468$ ,  $p_r(r \rightarrow 0) = 0.0005755839216$ , and  $p_\theta(r \rightarrow 0) = 0.0005755799012$ . Figure 1 ensures that the energy-density, radial and tangential pressures are decreasing towards the surface stellar. Also, in Fig. 1 (d), we present the behavior of the anisotropy parameter which is defined as  $\Delta(r) = p_\theta - p_r$ . Moreover, in Fig. 1 (d) we present the anisotropic force that has a positive behavior which means that we have a repulsive force, due to the fact  $p_\theta \geq p_r$ .

## B. Causality

To test the behavior of the sound velocities, we evaluate the derivative of the energy-density, and the radial and tangential pressures, the final expressions of which we present in the Appendix B. The equations given in Appendix B do not inform us if the gradients of the components of the energy-momentum tensor have a positive or negative behavior, therefore, we plot them in Fig. 2 (a) and from the plots it is ensured that the gradients of the energy-momentum tensor are negative, as it is required for any realistic physical stellar. To verify the causality conditions we must prove that the radial and transverse sound speeds,  $v_r^2$  and  $v_\perp^2$ , have values less than the speed of light. We evaluate these quantities,  $v_r^2$  and  $v_\perp^2$ , and we present their functional forms in the Appendix C : We plot the equations given in Appendix C in Fig. 2 (b) to ensure the validity of the conditions  $1 > v_r^2 > 0$  and  $1 > v_\perp^2 > 0$ .

The appearance of non-vanishing radial force with different signs in different regions of the fluid is called *gravitational cracking*. This happen when the radial force is directed inwards in the inner part of the sphere for all values of the




Figure 2. The plot of the gradients of the energy-momentum components where we put  $\beta = -0.0001$  and the constants  $c_0$  and  $c_1$  take the numerical values,  $-0.001763358117$  and  $-0.9329227851$  respectively.

radial coordinate  $r$  between the center, and some value beyond which the force reverses its direction [115]. It is shown in Ref. [116], that a simple requirement to avoid gravitational cracking is  $0 > v_\theta^2 - v_r^2 > -1$ . In Fig. 2 (b), we show that the solution given in Appendix C is stable against cracking.

### C. Energy conditions

Energy conditions are considered as important tests for non-vacuum solutions. To satisfy the dominant energy condition (DEC), we have to prove that  $\rho - p_r > 0$  &  $\rho - p_\theta > 0$ . As shown in Fig. 3 (a) the DEC is satisfied for suitable choices of the parameters  $\beta$ ,  $c_0$ , and  $c_1$ . Also, in Fig. 3 (b), (c) and (d) we show that the Weak Energy Condition (WEC), the Null Energy Condition (NEC), and the Strong Energy Condition (SEC) are all satisfied. It is interesting to note that the problem of energy conditions in the frame of  $f(G)$  has been studied and the results presented in this study are consistent with the results presented in [117].

### D. Mass-Radius relation

The compactification factor,  $u(r)$ , is the one defined as the ratio between the mass and radius. It has an important role towards revealing the physical properties of compact objects. Starting from the solution given in Appendix A, we define the gravitational mass by the following expression:

$$M(r) = \int_0^r \rho(\xi) \xi^2 d\xi = \frac{c_0 r^3}{16\pi} (c_0 r^2 + 2) (c_0^2 r^4 + 2 + 2 c_0 r^2) (c_0^4 r^8 + 4 c_0^3 r^6 + 6 c_0^2 r^4 + 4 c_0 r^2 + 2) \\ \times (\beta c_0^8 r^{14} + 8 \beta c_0^7 r^{12} + 28 \beta c_0^6 r^{10} + 56 \beta c_0^5 r^8 + 70 \beta c_0^4 r^6 + 56 \beta c_0^3 r^4 + 28 \beta c_0^2 r^2 + 8 \beta c_0 - 1) , \quad (22)$$

where we substitute the value of energy-density from the first equation presented in Appendix A. The compactification factor  $C(r)$  is then defined as:

$$C(r) = \frac{M(r)}{l} . \quad (23)$$

Substituting Eq. (22) into (23), one can get the explicit form of the compactification factor. The behavior of the gravitational mass and the compactification factor are plotted in Fig. 4 which indicates in a clear way that the gravitational mass and the compactification factor are directly proportional to the radial coordinate.




Figure 3. DEC, WEC, NEC and SEC for  $\beta = -0.0001$ ,  $c_0 = -0.001763358117$  and  $c_1 = -0.9329227851$ .

### E. Equation of State

Finally, let us study the equation of state (EoS) of a compact stellar object as presented in [118] which uses a linear EoS. In the present case, we show that the EoS is not linear due to the contribution of the GB term. To show this, we define the radial and transverse EoS as:

$$\omega_r = \frac{p_r(r)}{\rho(r)}, \quad \omega_t = \frac{p_\theta(r)}{\rho(r)}, \quad (24)$$

with  $\omega_r$  and  $\omega_t$  being the radial and transverse EoS parameters. Using the form of energy-density, radial and tangential pressures presented in Appendix A we get the EoSs of our model as presented in Appendix D. As it can be seen in Fig. 4 (b), the EoS is non-linear because of the contribution of the GB term.

Now, we are going to calculate the EoS as  $P(\rho) = \omega\rho$ , for such purpose we write  $r = r(\rho)$ . Using the form of energy-density given in Appendix A we get the form of the radial, tangential and pressure components in Appendix D. The behavior of the EoS as a function of the energy-density is presented in Fig. 4 (c), (d) and (e). As it can be seen in Figs. 4 (c), (d) and (e), the radial and tangential EoS parameters are positive, while the total EoS, i.e.,  $P = \frac{p_r+2p_\theta}{3}$  is negative.

### V. STABILITY OF THE MODEL

We use two different procedures to study the stability problem, the first is the Tolman-Oppenheimer-Volkoff (TOV) equation and the second is to study the adiabatic index. Now we are going to discuss the TOV equation in the frame




Figure 4. Schematic plots of (a) the gravitational mass and compactification factor of the solution given by Eqs. (22), and (23); (b) the components of radial and tangential EoS parameters as function of the radial coordinate; (c) the radial EoS as a function of the energy-density; (d) the tangential EoS as a function of the energy-density and; (e) the EoS as a function of the energy-density. The values the GB parameter and the constants  $c_0$  and  $c_1$  are taken as  $\beta = -0.0001$ ,  $c_0 = -0.001763358117$  and  $c_1 = -0.9329227851$  in these plots.

of our solution in presented in Appendix A.

### A. The Tolman-Oppenheimer-Volkoff equation

To study the stability of our solution displayed in Appendix A, we are going to suppose the hydrostatic equilibrium by using the TOV equation [119, 120] as given in [121]:

$$\frac{2[p_\theta - p_r]}{r} - \frac{M_g(r)[\rho(r) + p_r(r)]e^{[h(r)-h_1(r)]/2}}{r} - \frac{dp_r(r)}{dr} = 0. \quad (25)$$

Here  $M_g(r)$  represents the gravitational mass evaluated at the radius  $r$  and has the following form:

$$M_g(r) = 4\pi \int_0^r \left( \mathcal{T}_t^t - \mathcal{T}_r^r - \mathcal{T}_\theta^\theta - \mathcal{T}_\phi^\phi \right) y^2 e^{[h(y)+h_1(y)]} dy = \frac{rh' e^{[h_1(r)-h(r)]/2}}{2}, \quad (26)$$

Using Eq. (26) in Eq. (25) gives:

$$\frac{2(p_\theta - p_r)}{r} - \frac{dp_r}{dr} - \frac{h'[\rho(r) + p_r(r)]}{2} = F_g + F_a + F_h = 0. \quad (27)$$




Figure 5. Schematic plots of (a) which represents the different forces acting on the model under consideration; (b) which represents the adiabatic index of the model under consideration; (c) which represents the variation of the mass (which is a function of the central density) via the central density; (d) which represents the stability of the model in the static state. The numerical values of the GB and the constants  $c_0$  and  $c_1$  used in these plots are given as  $\beta = -0.0001$ ,  $c_0 = -0.001763358117$  and  $c_1 = -0.9329227851$  in this figure.

Here  $F_g = -\frac{h'[\rho+p_r]}{2}$ ,  $F_a = \frac{2(p_\theta-p_r)}{r}$  and  $F_h = -\frac{dp_r(r)}{dr}$  are the gravitational mass, the anisotropic and the hydrostatic forces, respectively. The explicit forms of these forces can be found in Appendix E. We plot the different forces presented in Appendix E in Fig. 5 (a), and it can be seen, it is ensured that the hydrostatic and anisotropic forces are positive and dominate over the gravitational force, which has a negative value, hence opposite direction, and the system is kept in static equilibrium.

## B. The Adiabatic Index

The adiabatic index connects the structure of a spherically symmetric static object to the EoS of the interior solution. One can use the adiabatic index to discuss the stability of the interior solution [122]. For any interior solution, it becomes stable if its adiabatic index is greater than  $4/3$  [123] however, if  $\gamma = \frac{4}{3}$ , then the isotropic sphere is in neutral equilibrium. Following Chan et al. [124], the stability condition of a relativistic anisotropic sphere,  $\gamma > \Gamma$ , must be satisfied. Here  $\Gamma$  is defined as

$$\Gamma = \frac{4}{3} - \left\{ \frac{4(p_r - p_\theta)}{3|p'_r|} \right\}_{max}, \quad (28)$$

Table I. Values of model parameters [128]

Pulsar	Mass ( $M_{\odot}$ )	Radius (km)	$c_0$	$c_1$
4U 1724-207	$1.81^{+0.25}_{-0.37}$	$12.2^{+1.4}_{-1.4}$	$\approx -0.75 \times 10^{-3}$	$\approx -0.9$
4U 1820-30	$1.46^{+0.21}_{-0.21}$	$11.1^{+1.8}_{-1.8}$	$\approx -0.7 \times 10^{-3}$	$\approx -1.5$
SAX J1748.9-2021	$1.81^{+0.25}_{-0.37}$	$11.7^{+1.7}_{-1.7}$	$\approx -0.2 \times 10^{-2}$	$\approx -1.3$
EXO 1745-268	$1.65^{+0.21}_{-0.31}$	$10.5^{+1.6}_{-1.6}$	$\approx -0.1 \times 10^{-2}$	$\approx -1.4$
4U 1608-52	$1.57^{+0.30}_{-0.29}$	$9.8^{+1.8}_{-1.8}$	$\approx -0.1 \times 10^{-2}$	$\approx -1.4$
KS 1731-260	$1.61^{+0.35}_{-0.37}$	$10.0^{+2.2}_{-2.2}$	$\approx -0.1 \times 10^{-2}$	$\approx -2.4$

Table II. Values of physical quantities

Pulsar	$\rho _0$	$\rho _l$	$\frac{dp_r}{d\rho} _0$	$\frac{dp_r}{d\rho} _l$	$\frac{dp_\theta}{d\rho} _0$	$\frac{dp_\theta}{d\rho} _l$	$(\rho - p_r - 2p_\theta) _0$	$(\rho - p_r - 2p_\theta) _l$	$z _l$
4U 1724-207	$\approx 0.71 \times 10^{-3}$	$\approx 0.32 \times 10^{-3}$	$\approx 0.68$	$\approx 0.57$	$\approx 0.28$	$0.43$	$\approx 0.42 \times 10^{-3}$	$\approx 0.41 \times 10^{-4}$	$0.014$
4U 1820-30	$\approx 0.65 \times 10^{-3}$	$\approx 0.34 \times 10^{-3}$	$\approx 0.36$	$\approx 0.26$	$\approx 0.21$	$\approx 0.15$	$\approx 0.58 \times 10^{-3}$	$\approx 0.28 \times 10^{-3}$	$0.012$
SAX J1748.9-2021	$\approx 0.2 \times 10^{-2}$	$\approx 0.6 \times 10^{-3}$	$\approx 0.45$	$\approx 0.24$	$\approx 0.26$	$\approx 0.2$	$\approx 0.16 \times 10^{-2}$	$\approx 0.35 \times 10^{-3}$	$\approx 0.017$
EXO 1745-268	$\approx 0.1 \times 10^{-2}$	$\approx 0.4 \times 10^{-3}$	$\approx 0.4$	$\approx 0.27$	$\approx 0.23$	$\approx 0.17$	$\approx 0.8 \times 10^{-3}$	$\approx 0.3 \times 10^{-3}$	$\approx 0.014$
4U 1608-52	$\approx 0.1 \times 10^{-2}$	$\approx 0.44 \times 10^{-3}$	$\approx 0.38$	$\approx 0.25$	$\approx 0.22$	$\approx 0.15$	$\approx 0.92 \times 10^{-3}$	$\approx 0.34 \times 10^{-3}$	$\approx 0.0148$
KS 1731-260	$\approx 0.1 \times 10^{-2}$	$\approx 0.4 \times 10^{-3}$	$\approx 0.15$	$\approx 0.034$	$\approx 0.048$	$\approx 0$	$\approx 0.1 \times 10^{-2}$	$\approx 0.4 \times 10^{-3}$	$\approx 0.0147$

where the collapse happens according to the nature of the anisotropy. When  $p_r > p_\theta$  one can have  $\gamma < 4/3$  and the system still stable and when  $p_\theta > p_r$  one can have unstable system even if  $\gamma > 4/3$ . The behavior of the adiabatic index is shown in Fig. 5 (b) which ensures the stability condition of our solution.

### C. Stability in the Static State

Finally, let us study the stability through the procedure given by Harrison, Zeldovich and Novikov [125–127] who showed that for stable compact stars the mass, (which is a function of the central density) must be positive and increasing and also the derivative of the mass with respect to the central density must have a positive value, i.e.,  $\frac{\partial M}{\partial \rho_0} > 0$ . Applying this condition to our model we get the density at the center of the star:

$$\rho_0(r \rightarrow 0) = \frac{3(8\beta c_0 - 1)c_0}{\pi} \implies c_0 = \frac{3 \pm \sqrt{9 + 96\pi\beta\rho_0}}{48\beta}. \quad (29)$$

Using Eq. (29) in Eq. (22) we get the form of mass in terms of the central density which is written in Appendix F. The pattern of the derivative of mass with respect to the central density given by the equation presented in Appendix  $F_1$  and it is plotted in Fig. 5 (d) which ensures the stability of our model. In addition to EXO1785 – 248, a similar analysis can be developed for other pulsars. In tables I and II we report the results for other observed pulsars.

## VI. DISCUSSION AND CONCLUSIONS

Among the higher curvature theories, the EGB theory offers new possibilities for 4D gravity. This new proposal has received much interest because the GB invariant may share Einstein's field equations through the redefinition of the GB constant to be  $\beta \rightarrow \frac{\beta}{D-4}$  in  $D$ -dimensions and taking the limit  $D \rightarrow 4$ . Inspired by this new proposal of EGB theory, in this work, we rigorously explained a static and spherically symmetric interior solution. Assuming a specific form of the metric potentials, we derived in this new proposal of EGB a new interior solution. In this study we showed that the GB parameter,  $\beta$ , must take a tiny negative value otherwise we will have imaginary quantities for the components of pressures. As we showed the effect of the constant  $\beta$  acting on the EoS, yielded its behavior to behave in a non-linear form, unlike the Einstein GR.

We checked the impact of the anisotropy regime and showed that it is always positive which means that we have a repulsive force. Moreover, we listed the physical conditions that any real star must satisfy and showed that our model yields:

i-A well-known behavior of the energy-density, radial and tangential pressures at the center as well as at the surface of the star as shown in Fig. 1 (a), (b) and (c).

ii-The causality condition is satisfied by showing that the gradient of the energy momentum components have negative value as shown in Fig. 2 (a). Moreover, in Fig. 2 (b) we showed that the speed of sound of radial and tangential components are less than one, as required for any realistic star.

iii- In Fig. 3 we showed that the energy conditions which must be satisfied for any realistic star, are satisfied.

iv- Using different techniques, TOV, adiabatic index and stability of static state, we studied the stability of our model and we showed that it is stable as Fig. 5 (a), (b) and (d) show. The results of this study can be applied to the observational data of various stars showing consistent results as shown in Tables I and II. For these objects, it is possible to calculate the density at the center and at the surface, the radial and tangential speed, at the center and the surface of the star, the SEC at the center and the surface of the star, and the redshift at the surface of the star. It is interesting to see that all the results of the different stars of our solution are compatible with observations. From the above discussions we can say that our results presented in this study are in agreement with that presented in [129] from the viewpoint of the structure of compact star.

In summary, we derived a new interior solution in the framework of EGB in the limit  $D \rightarrow 4$ , assuming physically motivated metric potentials. This solution has a non-trivial form of the Ricci scalar as well as the GB term. The internal solution can be matched with the external in the limit  $D \rightarrow 4$ . In a forthcoming paper, we will consider a similar situation assuming a charged interior solution.

#### ACKNOWLEDGEMENTS

The Authors would like to thank the referees for their valuable comments. This work was supported by MINECO (Spain), project PID2019-104397GB-I00.

### Appendix A: Energy density, radial and tangential pressures.

Now, let us present the quantities related to the interior solution discussed in the present study and write them as:

$$\begin{aligned}
8\pi\rho(r) &= \left\{ 192\beta c_0 - 140c_0 r^2 - 392c_0^2 r^4 - 630c_0^3 r^6 - 616c_0^4 r^8 - 364c_0^5 r^{10} - 120c_0^6 r^{12} - 17c_0^7 r^{14} - 24 + 11760\beta c_0^3 r^4 \right. \\
&\quad + 38304\beta c_0^4 r^6 + 87472\beta c_0^5 r^8 + 148512\beta c_0^6 r^{10} + 193020\beta c_0^7 r^{12} + 194480\beta c_0^8 r^{14} + 152152\beta c_0^9 r^{16} + 91728\beta c_0^{10} r^{18} \\
&\quad \left. + 41860\beta c_0^{11} r^{20} + 14000\beta c_0^{12} r^{22} + 3240\beta c_0^{13} r^{24} + 464\beta c_0^{14} r^{26} + 31\beta c_0^{15} r^{28} + 2240\beta c_0^2 r^2 \right\} c_0, \\
8\pi p_r(r) &= \frac{c_0}{c_0 r^2 - c_1} \left\{ -16 - 420c_0^2 r^4 - 840c_0^3 r^6 - 1050c_0^4 r^8 - 840c_0^5 r^{10} - 420c_0^6 r^{12} - 120c_0^7 r^{14} - c_0^7 r^{14} c_1 \right. \\
&\quad - 120c_0 r^2 + 256\beta c_0 - 70c_0^3 r^6 c_1 - \beta c_0^{15} r^{28} c_1 - 448\beta c_0^2 r^2 c_1 - 11440\beta c_0^8 r^{14} c_1 - 8008\beta c_0^9 r^{16} c_1 - 1680\beta c_0^3 r^4 c_1 \\
&\quad - 4256\beta c_0^4 r^6 c_1 - 7952\beta c_0^5 r^8 c_1 - 11424\beta c_0^6 r^{10} c_1 - 12868\beta c_0^7 r^{12} c_1 - 4368\beta c_0^{10} r^{18} c_1 - 1820\beta c_0^{11} r^{20} c_1 - 560\beta c_0^{12} r^{22} c_1 \\
&\quad - 120\beta c_0^{13} r^{24} c_1 - 16\beta c_0^{14} r^{26} c_1 + 33\beta c_0^{16} r^{30} - 28c_0^5 r^{10} c_1 - 56c_0^4 r^8 c_1 - 64\beta c_0 c_1 - 8c_0^6 r^{12} c_1 - 28c_0 r^2 c_1 - 56c_0^2 r^4 c_1 \\
&\quad + 3008\beta c_0^2 r^2 + 528\beta c_0^{15} r^{28} + 3960\beta c_0^{14} r^{26} + 60060\beta c_0^{12} r^{22} + 18480\beta c_0^{13} r^{24} + 264264\beta c_0^{10} r^{18} + 144144\beta c_0^{11} r^{20} \\
&\quad + 377248\beta c_0^7 r^{12} + 263312\beta c_0^6 r^{10} + 377520\beta c_0^9 r^{16} + 424676\beta c_0^8 r^{14} + 16576\beta c_0^3 r^4 + 57680\beta c_0^4 r^6 + 142240\beta c_0^5 r^8 \\
&\quad \left. - 8c_1 - 15c_0^8 r^{16} \right\}, \\
8\pi p_\theta(r) &= \frac{2c_0}{(c_0 r^2 - c_1)^2} \left\{ 228c_0^2 r^4 + 700c_0^3 r^6 + 1204c_0^4 r^8 + 1260c_0^5 r^{10} + 812c_0^6 r^{12} + 308c_0^7 r^{14} + 28c_1^2 c_0 r^2 + 232c_0^7 r^{14} c_1 \right. \\
&\quad + 32c_0 r^2 + 952c_0^3 r^6 c_1 + 224\beta c_0^2 c_1^2 r^2 + 1680\beta c_0^3 c_1^2 r^4 + 6384\beta c_0^4 c_1^2 r^6 + 15904\beta c_0^5 c_1^2 r^8 + 28560\beta c_0^6 c_1^2 r^{10} \\
&\quad + 38604\beta c_0^7 c_1^2 r^{12} + 40040\beta c_0^8 c_1^2 r^{14} + 32032\beta c_0^9 r^{16} c_1^2 + 19656\beta c_0^{10} r^{18} c_1^2 + 9100\beta c_0^{11} r^{20} c_1^2 + 3080\beta c_0^{12} r^{22} c_1^2 \\
&\quad + 720\beta c_0^{13} r^{24} c_1^2 + 104\beta c_0^{14} r^{26} c_1^2 + 7\beta c_0^{15} r^{28} c_1^2 - 406\beta c_0^{16} r^{30} c_1 - 6096\beta c_0^{15} r^{28} c_1 - 5600\beta c_0^2 r^2 c_1 \\
&\quad - 2651072\beta c_0^8 r^{14} c_1 - 2642640\beta c_0^9 r^{16} c_1 - 42112\beta c_0^3 r^4 c_1 - 187600\beta c_0^4 r^6 c_1 - 566944\beta c_0^5 r^8 c_1 - 1246112\beta c_0^6 r^{10} c_1 \\
&\quad - 2069600\beta c_0^7 r^{12} c_1 - 2050048\beta c_0^{10} r^{18} c_1 - 1227408\beta c_0^{11} r^{20} c_1 - 556920\beta c_0^{12} r^{22} c_1 - 185360\beta c_0^{13} r^{24} c_1 \\
&\quad - 42720\beta c_0^{14} r^{26} c_1 + 3432\beta c_0^{16} r^{30} + 1288c_0^5 r^{10} c_1 + 1400c_0^4 r^8 c_1 - 320\beta c_0 c_1 + 728c_0^6 r^{12} c_1 + 88c_0 r^2 c_1 + 392c_0^2 r^4 c_1 \\
&\quad - 960\beta c_0^2 r^2 + 22800\beta c_0^{15} r^{28} + 92680\beta c_0^{14} r^{26} + 508872\beta c_0^{12} r^{22} + 256620\beta c_0^{13} r^{24} + 772200\beta c_0^{10} r^{18} + 736736\beta c_0^{11} r^{20} \\
&\quad - 46592\beta c_0^7 r^{12} - 129360\beta c_0^6 r^{10} + 553428\beta c_0^9 r^{16} + 211920\beta c_0^8 r^{14} - 9120\beta c_0^3 r^4 - 38752\beta c_0^4 r^6 - 94080\beta c_0^5 r^8 \\
&\quad + 84c_1^2 c_0^5 r^{10} + 140c_1^2 c_0^4 r^8 + 4c_1^2 c_0^7 r^{14} + 28c_1^2 c_0^6 r^{12} + 239\beta c_0^{17} r^{32} + 140c_1^2 c_0^3 r^6 + 84c_1^2 c_0^2 r^4 + 32c_0^8 r^{16} c_1 + 8c_1 \\
&\quad \left. + 60c_0^8 r^{16} + 4c_0^9 r^{18} + 4c_1^2 \right\}, \\
8\pi\Delta(r) &= \frac{c_0^2}{(c_0 r^2 - c_1)^2} \left\{ 80r^2 + 1120c_0^2 r^6 c_1 + 6336\beta c_0^{15} r^{30} + 352c_0^6 r^{14} c_1 + 1064c_0^5 r^{12} c_1 + 1792c_0^4 r^{10} c_1 + 1820c_0^3 r^8 c_1 \right. \\
&\quad - 400960\beta c_0^5 r^{10} + 682180\beta c_0^8 r^{16} + 1680\beta c_0^2 c_1^2 r^4 + 8512\beta c_0^3 c_1^2 r^6 + 23856\beta c_0^4 c_1^2 r^8 + 45696\beta c_0^5 c_1^2 r^{10} \\
&\quad + 64340\beta c_0^6 c_1^2 r^{12} + 68640\beta c_0^7 c_1^2 r^{14} + 56056\beta c_0^8 r^{16} c_1^2 + 34944\beta c_0^9 r^{18} c_1^2 + 16380\beta c_0^{10} r^{20} c_1^2 + 5600\beta c_0^{11} r^{22} c_1^2 \\
&\quad + 1320\beta c_0^{12} r^{24} c_1^2 + 192\beta c_0^{13} r^{26} c_1^2 + 13\beta c_0^{14} r^{28} c_1^2 - 778\beta c_0^{15} r^{30} c_1 - 11648\beta c_0^{14} r^{28} c_1 - 8128\beta c_0 r^2 c_1 \\
&\quad - 4864600\beta c_0^7 r^{14} c_1 - 4896320\beta c_0^8 r^{16} c_1 - 67200\beta c_0^2 r^4 c_1 - 315840\beta c_0^3 r^6 c_1 - 987392\beta c_0^4 r^8 c_1 - 2220960\beta c_0^5 r^{10} c_1 \\
&\quad - 3750528\beta c_0^6 r^{12} c_1 - 3827824\beta c_0^9 r^{18} c_1 - 2306304\beta c_0^{10} r^{20} c_1 - 1051960\beta c_0^{11} r^{22} c_1 - 351680\beta c_0^{12} r^{24} c_1 - 81360\beta c_0^{13} r^{26} c_1 \\
&\quad - 384\beta c_1 + 64r^2 c_1 - 64\beta c_1^2 + 28r^2 c_1^2 + 1209208\beta c_0^{10} r^{20} - 356496\beta c_0^6 r^{12} - 94080\beta c_0^3 r^6 - 245840\beta c_0^4 r^8 \\
&\quad + 46592\beta c_0^7 r^{14} - 21248\beta c_0^2 r^4 - 2176\beta c_0 r^2 + 41640\beta c_0^{14} r^{28} + 392c_0 r^4 c_1 + 453180\beta c_0^{12} r^{24} + 1166880\beta c_0^9 r^{18} \\
&\quad + 166880\beta c_0^{13} r^{26} + 873600\beta c_0^{11} r^{22} + 140c_1^2 c_0^4 r^{10} + 224c_1^2 c_0^3 r^8 + 7c_1^2 c_0^6 r^{14} + 48c_1^2 c_0^5 r^{12} + 445\beta c_0^{16} r^{32} \\
&\quad + 210c_1^2 c_0^2 r^6 + 112c_1^2 c_0 r^4 + 50c_0^7 r^{16} c_1 + 576c_0 r^4 + 1820c_0^2 r^6 + 3248c_0^3 r^8 + 3570c_0^4 r^{10} + 2464c_0^5 r^{12} + 1036c_0^6 r^{14} \\
&\quad \left. + 240c_0^7 r^{16} + 23c_0^8 r^{18} \right\}. \tag{A}
\end{aligned}$$

where  $\Delta$  is the anisotropy defined as  $\Delta = p_\theta - p_r$ .

## Appendix B: The gradients of energy-density, radial and transverse pressures.

The gradients of the components of the energy-momentum tensor given in the Appendix A take the following form:

$$\begin{aligned}
\rho' &= \frac{d\rho}{dr} = \frac{c_0^2 r}{4\pi} \left\{ 23520 \beta c_0^2 r^2 - 140 - 784 c_0 r^2 - 1890 c_0^2 r^4 - 2464 c_0^3 r^6 - 1820 c_0^4 r^8 - 720 c_0^5 r^{10} - 119 c_0^6 r^{12} \right. \\
&\quad + 114912 \beta c_0^3 r^4 + 349888 \beta c_0^4 r^6 + 742560 \beta c_0^5 r^8 + 1158120 \beta c_0^6 r^{10} + 1361360 \beta c_0^7 r^{12} + 1217216 \beta c_0^8 r^{14} + 2240 \beta c_0 \\
&\quad \left. + 825552 \beta c_0^9 r^{16} + 418600 \beta c_0^{10} r^{18} + 154000 \beta c_0^{11} r^{20} + 38880 \beta c_0^{12} r^{22} + 6032 \beta c_0^{13} r^{24} + 434 \beta c_0^{14} r^{26} \right\}, \\
p'_r &= \frac{c_0^2 r}{4\pi (c_0 r^2 - c_1)^2} \left\{ 128 c_1 + 3360 \beta c_0^2 c_1^2 r^2 + 12768 \beta c_0^3 c_1^2 r^4 + 31808 \beta c_0^4 c_1^2 r^6 + 57120 \beta c_0^5 c_1^2 r^8 + 77208 \beta c_0^6 c_1^2 r^{10} \right. \\
&\quad + 80080 \beta c_0^7 c_1^2 r^{12} + 64064 \beta c_0^8 c_1^2 r^{14} + 39312 \beta c_0^9 r^{16} c_1^2 + 18200 \beta c_0^{10} r^{18} c_1^2 + 6160 \beta c_0^{11} r^{20} c_1^2 + 1440 \beta c_0^{12} r^{22} c_1^2 \\
&\quad + 208 \beta c_0^{13} r^{24} c_1^2 + 14 \beta c_0^{14} r^{26} c_1^2 - 508 \beta c_0^{15} r^{28} c_1 - 33152 \beta c_0^2 r^2 c_1 - 3088800 \beta c_0^8 r^{14} c_1 - 2434432 \beta c_0^9 r^{16} c_1 \\
&\quad - 577472 \beta c_0^4 r^6 c_1 - 1340416 \beta c_0^5 r^8 c_1 - 2309184 \beta c_0^6 r^{10} c_1 - 3037072 \beta c_0^7 r^{12} c_1 - 1476384 \beta c_0^{10} r^{18} c_1 - 174720 \beta c_0^3 r^4 c_1 \\
&\quad + 16 - 677040 \beta c_0^{11} r^{20} c_1 - 227360 \beta c_0^{12} r^{22} c_1 - 52800 \beta c_0^{13} r^{24} c_1 - 7584 \beta c_0^{14} r^{26} c_1 - 256 \beta c_0 - 420 c_0^2 r^4 - 1680 c_0^3 r^6 \\
&\quad - 3150 c_0^4 r^8 - 3360 c_0^5 r^{10} - 2100 c_0^6 r^{12} - 720 c_0^7 r^{14} - 105 c_0^8 r^{16} + 28 c_1^2 + 16576 \beta c_0^3 r^4 + 115360 \beta c_0^4 r^6 \\
&\quad + 426720 \beta c_0^5 r^8 + 1053248 \beta c_0^6 r^{10} + 1886240 \beta c_0^7 r^{12} + 2548056 \beta c_0^8 r^{14} + 2642640 \beta c_0^9 r^{16} + 2114112 \beta c_0^{10} r^{18} \\
&\quad + 1297296 \beta c_0^{11} r^{20} + 600600 \beta c_0^{12} r^{22} + 203280 \beta c_0^{13} r^{24} + 47520 \beta c_0^{14} r^{26} + 6864 \beta c_0^{15} r^{28} + 114 c_0^7 r^{14} c_1 + 800 c_0^6 r^{12} c_1 \\
&\quad + 840 c_0 r^2 c_1 + 2464 c_0^2 r^4 c_1 + 4060 c_0^3 r^6 c_1 + 4032 c_0^4 r^8 c_1 + 2408 c_0^5 r^{10} c_1 - 2944 \beta c_0 c_1 + 462 \beta c_0^{16} r^{30} + 112 c_1^2 c_0 r^2 \\
&\quad \left. + 210 c_1^2 c_0^2 r^4 + 224 c_1^2 c_0^3 r^6 + 140 c_1^2 c_0^4 r^8 + 48 c_1^2 c_0^5 r^{10} + 7 c_1^2 c_0^6 r^{12} + 448 \beta c_0 c_1^2 \right\}, \\
p'_\theta &= \frac{c_0^2 r}{\pi (-c_1 + c_0 r^2)^3} \left\{ 42000 \beta c_0^2 c_1^2 r^2 - 24 c_1 + 281400 \beta c_0^3 c_1^2 r^4 + 1137080 \beta c_0^4 c_1^2 r^6 + 3131184 \beta c_0^5 c_1^2 r^8 \right. \\
&\quad + 6251640 \beta c_0^6 c_1^2 r^{10} + 9355960 \beta c_0^7 c_1^2 r^{12} + 10670660 \beta c_0^8 c_1^2 r^{14} + 9321312 \beta c_0^9 r^{16} c_1^2 + 6205836 \beta c_0^{10} r^{18} c_1^2 \\
&\quad + 3099460 \beta c_0^{11} r^{20} c_1^2 + 1126020 \beta c_0^{12} r^{22} c_1^2 + 281280 \beta c_0^{13} r^{24} c_1^2 + 43244 \beta c_0^{14} r^{26} c_1^2 + 3087 \beta c_0^{15} r^{28} c_1^2 \\
&\quad - 4551 \beta c_0^{16} r^{30} c_1 - 16 c_0 r^2 - 62316 \beta c_0^{15} r^{28} c_1 + 11920 \beta c_0^2 r^2 c_1 - 8841392 \beta c_0^8 r^{14} c_1 - 11402820 \beta c_0^9 r^{16} c_1 \\
&\quad + 58128 \beta c_0^3 r^4 c_1 + 94360 \beta c_0^4 r^6 c_1 - 243544 \beta c_0^5 r^8 c_1 - 1729392 \beta c_0^6 r^{10} c_1 - 4880920 \beta c_0^7 r^{12} c_1 - 10858848 \beta c_0^{10} r^{18} c_1 \\
&\quad - 7708428 \beta c_0^{11} r^{20} c_1 - 4045860 \beta c_0^{12} r^{22} c_1 - 1529220 \beta c_0^{13} r^{24} c_1 - 394560 \beta c_0^{14} r^{26} c_1 + 350 c_0^3 r^6 + 1204 c_0^4 r^8 \\
&\quad + 1890 c_0^5 r^{10} + 1624 c_0^6 r^{12} + 770 c_0^7 r^{14} + 180 c_0^8 r^{16} + 14 c_0^9 r^{18} - 48 c_1^2 - 19376 \beta c_0^4 r^6 - 94080 \beta c_0^5 r^8 \\
&\quad - 194040 \beta c_0^6 r^{10} - 93184 \beta c_0^7 r^{12} + 529800 \beta c_0^8 r^{14} + 1660284 \beta c_0^9 r^{16} + 2702700 \beta c_0^{10} r^{18} + 2946944 \beta c_0^{11} r^{20} \\
&\quad + 2289924 \beta c_0^{12} r^{22} + 1283100 \beta c_0^{13} r^{24} + 509740 \beta c_0^{14} r^{26} + 136800 \beta c_0^{15} r^{28} + 480 \beta c_0^2 r^2 + 340 c_0^7 r^{14} c_1 \\
&\quad + 378 c_0^6 r^{12} c_1 - 272 c_0 r^2 c_1 - 1050 c_0^2 r^4 c_1 - 1932 c_0^3 r^6 c_1 - 1750 c_0^4 r^8 c_1 - 504 c_0^5 r^{10} c_1 + 800 \beta c_0 c_1 + 22308 \beta c_0^{16} r^{30} \\
&\quad + 78 c_0^8 r^{16} c_1 + 1673 \beta c_0^{17} r^{32} - 406 c_1^2 c_0 r^2 - 1428 c_1^2 c_0^2 r^4 - 2730 c_1^2 c_0^3 r^6 - 3080 c_1^2 c_0^4 r^8 - 2058 c_1^2 c_0^5 r^{10} \\
&\quad - 756 c_1^2 c_0^6 r^{12} - 118 c_1^2 c_0^7 r^{14} + 2800 \beta c_0 c_1^2 - 14 c_1^3 - 1680 \beta c_0^2 c_1^3 r^2 - 9576 \beta c_0^3 c_1^3 r^4 - 31808 \beta c_0^4 c_1^3 r^6 \\
&\quad - 71400 \beta c_0^5 c_1^3 r^8 - 115812 \beta c_0^6 c_1^3 r^{10} - 140140 \beta c_0^7 c_1^3 r^{12} - 128128 \beta c_0^8 c_1^3 r^{14} - 88452 \beta c_0^9 r^{16} c_1^3 - 45500 \beta c_0^{10} r^{18} c_1^3 \\
&\quad - 16940 \beta c_0^{11} r^{20} c_1^3 - 4320 \beta c_0^{12} r^{22} c_1^3 - 676 \beta c_0^{13} r^{24} c_1^3 - 49 \beta c_0^{14} r^{26} c_1^3 - 84 c_1^3 c_0 r^2 \\
&\quad \left. - 210 c_1^3 c_0^2 r^4 - 280 c_1^3 c_0^3 r^6 - 210 c_1^3 c_0^4 r^8 - 84 c_1^3 c_0^5 r^{10} - 14 c_1^3 c_0^6 r^{12} - 112 \beta c_0 c_1^3 \right\}, \tag{B}
\end{aligned}$$

where  $\rho' = \frac{d\rho}{dr}$ ,  $p_r' = \frac{dp_r}{dr}$  and  $p_\theta' = \frac{dp_\theta}{dr}$ .

### Appendix C: Derivation of the radial and tangential speeds of sound

Using the equations of the gradient of energy-momentum components presented in Appendix B we get:

$$\begin{aligned}
v_r^2 &= \frac{p'_r}{\rho'} = \left\{ 16 - 256 \beta c_0 - 420 c_0^2 r^4 - 1680 c_0^3 r^6 - 3150 c_0^4 r^8 - 3360 c_0^5 r^{10} - 2100 c_0^6 r^{12} - 720 c_0^7 r^{14} - 105 c_0^8 r^{16} \right. \\
&\quad + 28 c_1^2 + 426720 \beta c_0^5 r^8 + 115360 \beta c_0^4 r^6 + 1053248 \beta c_0^6 r^{10} + 16576 \beta c_0^3 r^4 + 1886240 \beta c_0^7 r^{12} + 2548056 \beta c_0^8 r^{14} \\
&\quad + 2642640 \beta c_0^9 r^{16} + 2114112 \beta c_0^{10} r^{18} + 1297296 \beta c_0^{11} r^{20} + 600600 \beta c_0^{12} r^{22} + 203280 \beta c_0^{13} r^{24} + 47520 \beta c_0^{14} r^{26} \\
&\quad + 6864 \beta c_0^{15} r^{28} + 114 c_0^7 r^{14} c_1 + 800 c_0^6 r^{12} c_1 + 840 c_0^5 r^2 c_1 + 2464 c_0^2 r^4 c_1 + 4060 c_0^3 r^6 c_1 + 4032 c_0^4 r^8 c_1 + 2408 c_0^5 r^{10} c_1 \\
&\quad - 2944 \beta c_0 c_1 + 462 \beta c_0^{16} r^{30} + 112 c_1^2 c_0 r^2 + 210 c_1^2 c_0^2 r^4 + 224 c_1^2 c_0^3 r^6 + 140 c_1^2 c_0^4 r^8 + 48 c_1^2 c_0^5 r^{10} + 7 c_1^2 c_0^6 r^{12} \\
&\quad + 448 \beta c_0 c_1^2 + 3360 \beta c_0^2 c_1^2 r^2 + 12768 \beta c_0^3 c_1^2 r^4 + 31808 \beta c_0^4 c_1^2 r^6 + 57120 \beta c_0^5 c_1^2 r^8 + 77208 \beta c_0^6 c_1^2 r^{10} \\
&\quad + 80080 \beta c_0^7 c_1^2 r^{12} + 64064 \beta c_0^8 c_1^2 r^{14} + 39312 \beta c_0^9 r^{16} c_1^2 + 18200 \beta c_0^{10} r^{18} c_1^2 + 6160 \beta c_0^{11} r^{20} c_1^2 + 1440 \beta c_0^{12} r^{22} c_1^2 \\
&\quad + 208 \beta c_0^{13} r^{24} c_1^2 + 14 \beta c_0^{14} r^{26} c_1^2 - 508 \beta c_0^{15} r^{28} c_1 - 33152 \beta c_0^{16} r^{30} c_1 - 3088800 \beta c_0^8 r^{14} c_1 - 2434432 \beta c_0^9 r^{16} c_1 \\
&\quad - 174720 \beta c_0^3 r^4 c_1 - 577472 \beta c_0^4 r^6 c_1 - 1340416 \beta c_0^5 r^8 c_1 - 2309184 \beta c_0^6 r^{10} c_1 - 3037072 \beta c_0^7 r^{12} c_1 - 1476384 \beta c_0^{10} r^{18} c_1 \\
&\quad - 677040 \beta c_0^{11} r^{20} c_1 - 227360 \beta c_0^{12} r^{22} c_1 - 52800 \beta c_0^{13} r^{24} c_1 - 7584 \beta c_0^{14} r^{26} c_1 + 128 c_1 \Big\} \left[ (c_0 r^2 - c_1)^2 \left\{ 23520 \beta c_0^2 r^2 \right. \right. \\
&\quad 140 - 784 c_0 r^2 - 1890 c_0^2 r^4 - 2464 c_0^3 r^6 - 1820 c_0^4 r^8 - 720 c_0^5 r^{10} - 119 c_0^6 r^{12} + 114912 \beta c_0^3 r^4 + 349888 \beta c_0^4 r^6 \\
&\quad + 742560 \beta c_0^5 r^8 + 1158120 \beta c_0^6 r^{10} + 1361360 \beta c_0^7 r^{12} + 1217216 \beta c_0^8 r^{14} + 825552 \beta c_0^9 r^{16} + 418600 \beta c_0^{10} r^{18} \\
&\quad \left. \left. + 154000 \beta c_0^{11} r^{20} + 38880 \beta c_0^{12} r^{22} + 6032 \beta c_0^{13} r^{24} + 434 \beta c_0^{14} r^{26} + 2240 \beta c_0 \right\} \right]^{-1}, \\
v_t^2 &= \frac{p'_\theta}{\rho'} = -4 \left\{ 770 c_0^7 r^{14} - 280 c_1^3 c_0^3 r^6 + 78 c_0^8 r^{16} c_1 + 180 c_0^8 r^{16} + 350 c_0^3 r^6 + 1204 c_0^4 r^8 + 1890 c_0^5 r^{10} + 1624 c_0^6 r^{12} \right. \\
&\quad - 16 c_0 r^2 - 48 c_1^2 - 94080 \beta c_0^5 r^8 - 19376 \beta c_0^4 r^6 - 194040 \beta c_0^6 r^{10} - 93184 \beta c_0^7 r^{12} + 529800 \beta c_0^8 r^{14} + 1660284 \beta c_0^9 r^{16} \\
&\quad + 2702700 \beta c_0^{10} r^{18} + 2946944 \beta c_0^{11} r^{20} + 2289924 \beta c_0^{12} r^{22} + 1283100 \beta c_0^{13} r^{24} + 509740 \beta c_0^{14} r^{26} + 136800 \beta c_0^{15} r^{28} \\
&\quad + 340 c_0^7 r^{14} c_1 + 378 c_0^6 r^{12} c_1 - 272 c_0 r^2 c_1 - 1050 c_0^2 r^4 c_1 - 1932 c_0^3 r^6 c_1 - 1750 c_0^4 r^8 c_1 - 504 c_0^5 r^{10} c_1 + 800 \beta c_0 c_1 \\
&\quad + 22308 \beta c_0^{16} r^{30} - 406 c_1^2 c_0 r^2 - 1428 c_1^2 c_0^2 r^4 - 2730 c_1^2 c_0^3 r^6 - 3080 c_1^2 c_0^4 r^8 - 2058 c_1^2 c_0^5 r^{10} - 756 c_1^2 c_0^6 r^{12} \\
&\quad + 2800 \beta c_0 c_1^2 + 480 \beta c_0^2 r^2 + 42000 \beta c_0^2 c_1^2 r^2 + 281400 \beta c_0^3 c_1^2 r^4 + 1137080 \beta c_0^4 c_1^2 r^6 + 3131184 \beta c_0^5 c_1^2 r^8 \\
&\quad + 6251640 \beta c_0^6 c_1^2 r^{10} + 9355960 \beta c_0^7 c_1^2 r^{12} + 10670660 \beta c_0^8 c_1^2 r^{14} + 9321312 \beta c_0^9 r^{16} c_1^2 + 6205836 \beta c_0^{10} r^{18} c_1^2 \\
&\quad + 3099460 \beta c_0^{11} r^{20} c_1^2 + 1126020 \beta c_0^{12} r^{22} c_1^2 + 281280 \beta c_0^{13} r^{24} c_1^2 + 43244 \beta c_0^{14} r^{26} c_1^2 - 62316 \beta c_0^{15} r^{28} c_1 \\
&\quad + 11920 \beta c_0^2 r^2 c_1 - 8841392 \beta c_0^8 r^{14} c_1 - 11402820 \beta c_0^9 r^{16} c_1 + 58128 \beta c_0^3 r^4 c_1 + 94360 \beta c_0^4 r^6 c_1 - 243544 \beta c_0^5 r^8 c_1 \\
&\quad - 1729392 \beta c_0^6 r^{10} c_1 - 4880920 \beta c_0^7 r^{12} c_1 - 10858848 \beta c_0^{10} r^{18} c_1 - 7708428 \beta c_0^{11} r^{20} c_1 - 4045860 \beta c_0^{12} r^{22} c_1 \\
&\quad - 1529220 \beta c_0^{13} r^{24} c_1 - 394560 \beta c_0^{14} r^{26} c_1 - 24 c_1 - 112 \beta c_0 c_1^3 - 210 c_1^3 c_0^4 r^8 - 84 c_1^3 c_0^5 r^{10} - 210 c_1^3 c_0^2 r^4 - 14 c_1^3 c_0^6 r^{12} \\
&\quad - 118 c_1^2 c_0^7 r^{14} + 1673 \beta c_0^{17} r^{32} + 3087 \beta c_0^{15} r^{28} c_1^2 - 4551 \beta c_0^{16} r^{30} c_1 - 1680 \beta c_0^2 c_1^3 r^2 - 9576 \beta c_0^3 c_1^3 r^4 - 31808 \beta c_0^4 c_1^3 r^6 \\
&\quad - 71400 \beta c_0^5 c_1^3 r^8 - 115812 \beta c_0^6 c_1^3 r^{10} - 140140 \beta c_0^7 c_1^3 r^{12} - 128128 \beta c_0^8 c_1^3 r^{14} - 88452 \beta c_0^9 r^{16} c_1^3 - 45500 \beta c_0^{10} r^{18} c_1^3 \\
&\quad - 16940 \beta c_0^{11} r^{20} c_1^3 - 4320 \beta c_0^{12} r^{22} c_1^3 - 676 \beta c_0^{13} r^{24} c_1^3 - 49 \beta c_0^{14} r^{26} c_1^3 - 84 c_1^3 c_0 r^2 + 14 c_0^9 r^{18} - 14 c_1^3 \Big\} \\
&\quad \times \left[ (-c_1 + c_0 r^2)^3 \left[ -140 - 784 c_0 r^2 - 1890 c_0^2 r^4 - 2464 c_0^3 r^6 - 1820 c_0^4 r^8 - 720 c_0^5 r^{10} - 119 c_0^6 r^{12} + 23520 \beta c_0^2 r^2 \right. \right. \\
&\quad + 114912 \beta c_0^3 r^4 + 349888 \beta c_0^4 r^6 + 742560 \beta c_0^5 r^8 + 1158120 \beta c_0^6 r^{10} + 1361360 \beta c_0^7 r^{12} + 1217216 \beta c_0^8 r^{14} \\
&\quad \left. \left. + 825552 \beta c_0^9 r^{16} + 418600 \beta c_0^{10} r^{18} + 154000 \beta c_0^{11} r^{20} + 38880 \beta c_0^{12} r^{22} + 6032 \beta c_0^{13} r^{24} + 434 \beta c_0^{14} r^{26} + 2240 \beta c_0 \right\} \right]^{-1}. \tag{C}
\end{aligned}$$

## Appendix D: Derivation of the EoS's

Using the form of energy-density, radial and tangential pressures presented in Appendix A, we get the EoS as:

$$\begin{aligned}
\omega_r = & \left\{ -16 - 120 c_0 r^2 + 256 \beta c_0 - 420 c_0^2 r^4 - 840 c_0^3 r^6 - 1050 c_0^4 r^8 - 840 c_0^5 r^{10} - 420 c_0^6 r^{12} - 120 c_0^7 r^{14} - 15 c_0^8 r^{16} \right. \\
& + 142240 \beta c_0^5 r^8 + 57680 \beta c_0^4 r^6 + 263312 \beta c_0^6 r^{10} + 16576 \beta c_0^3 r^4 + 377248 \beta c_0^7 r^{12} + 424676 \beta c_0^8 r^{14} + 377520 \beta c_0^9 r^{16} \\
& + 264264 \beta c_0^{10} r^{18} + 144144 \beta c_0^{11} r^{20} + 60060 \beta c_0^{12} r^{22} + 18480 \beta c_0^{13} r^{24} + 3960 \beta c_0^{14} r^{26} + 528 \beta c_0^{15} r^{28} - c_0^7 r^{14} c_1 \\
& - 8 c_0^6 r^{12} c_1 - 28 c_0^2 r^2 c_1 - 56 c_0^2 r^4 c_1 - 70 c_0^3 r^6 c_1 - 56 c_0^4 r^8 c_1 - 28 c_0^5 r^{10} c_1 - 64 \beta c_0 c_1 + 33 \beta c_0^{16} r^{30} + 3008 \beta c_0^2 r^2 \\
& - \beta c_0^{15} r^{28} c_1 - 448 \beta c_0^2 r^2 c_1 - 11440 \beta c_0^8 r^{14} c_1 - 8008 \beta c_0^9 r^{16} c_1 - 1680 \beta c_0^3 r^4 c_1 - 4256 \beta c_0^4 r^6 c_1 - 7952 \beta c_0^5 r^8 c_1 \\
& - 11424 \beta c_0^6 r^{10} c_1 - 12868 \beta c_0^7 r^{12} c_1 - 4368 \beta c_0^{10} r^{18} c_1 - 1820 \beta c_0^{11} r^{20} c_1 - 560 \beta c_0^{12} r^{22} c_1 - 120 \beta c_0^{13} r^{24} c_1 \\
& \left. - 16 \beta c_0^{14} r^{26} c_1 - 8 c_1 \right\} \left[ (-c_1 + c_0 r^2) \left\{ 192 \beta c_0 - 140 c_0 r^2 - 392 c_0^2 r^4 - 630 c_0^3 r^6 - 616 c_0^4 r^8 - 364 c_0^5 r^{10} - 120 c_0^6 r^{12} \right. \right. \\
& - 17 c_0^7 r^{14} - 24 + 11760 \beta c_0^3 r^4 + 38304 \beta c_0^4 r^6 + 87472 \beta c_0^5 r^8 + 148512 \beta c_0^6 r^{10} + 193020 \beta c_0^7 r^{12} + 194480 \beta c_0^8 r^{14} \\
& + 152152 \beta c_0^9 r^{16} + 91728 \beta c_0^{10} r^{18} + 41860 \beta c_0^{11} r^{20} + 14000 \beta c_0^{12} r^{22} + 3240 \beta c_0^{13} r^{24} + 464 \beta c_0^{14} r^{26} + 31 \beta c_0^{15} r^{28} \\
& \left. \left. + 2240 \beta c_0^2 r^2 \right\} \right]^{-1}, \\
\omega_\theta = & 2 \left\{ 32 c_0^8 r^{16} c_1 + 32 c_0 r^2 + 228 c_0^2 r^4 + 700 c_0^3 r^6 + 1204 c_0^4 r^8 + 1260 c_0^5 r^{10} + 812 c_0^6 r^{12} + 308 c_0^7 r^{14} + 60 c_0^8 r^{16} \right. \\
& - 94080 \beta c_0^5 r^8 - 38752 \beta c_0^4 r^6 - 129360 \beta c_0^6 r^{10} - 9120 \beta c_0^3 r^4 - 46592 \beta c_0^7 r^{12} + 211920 \beta c_0^8 r^{14} + 553428 \beta c_0^9 r^{16} \\
& + 4 c_1^2 + 772200 \beta c_0^{10} r^{18} + 736736 \beta c_0^{11} r^{20} + 508872 \beta c_0^{12} r^{22} + 256620 \beta c_0^{13} r^{24} + 92680 \beta c_0^{14} r^{26} + 22800 \beta c_0^{15} r^{28} \\
& + 232 c_0^7 r^{14} c_1 + 728 c_0^6 r^{12} c_1 + 88 c_0^2 r^2 c_1 + 392 c_0^2 r^4 c_1 + 952 c_0^3 r^6 c_1 + 1400 c_0^4 r^8 c_1 + 1288 c_0^5 r^{10} c_1 - 320 \beta c_0 c_1 \\
& + 3432 \beta c_0^{16} r^{30} + 28 c_1^2 c_0 r^2 + 84 c_1^2 c_0^2 r^4 + 140 c_1^2 c_0^3 r^6 + 140 c_1^2 c_0^4 r^8 + 84 c_1^2 c_0^5 r^{10} + 28 c_1^2 c_0^6 r^{12} - 960 \beta c_0^2 r^2 \\
& + 224 \beta c_0^2 c_1^2 r^2 + 1680 \beta c_0^3 c_1^2 r^4 + 6384 \beta c_0^4 c_1^2 r^6 + 15904 \beta c_0^5 c_1^2 r^8 + 28560 \beta c_0^6 c_1^2 r^{10} + 38604 \beta c_0^7 c_1^2 r^{12} \\
& + 40040 \beta c_0^8 c_1^2 r^{14} + 32032 \beta c_0^9 r^{16} c_1^2 + 19656 \beta c_0^{10} r^{18} c_1^2 + 9100 \beta c_0^{11} r^{20} c_1^2 + 3080 \beta c_0^{12} r^{22} c_1^2 + 720 \beta c_0^{13} r^{24} c_1^2 \\
& + 104 \beta c_0^{14} r^{26} c_1^2 - 6096 \beta c_0^{15} r^{28} c_1 - 5600 \beta c_0^2 r^2 c_1 - 2651072 \beta c_0^8 r^{14} c_1 - 2642640 \beta c_0^9 r^{16} c_1 - 42112 \beta c_0^3 r^4 c_1 \\
& - 187600 \beta c_0^4 r^6 c_1 - 566944 \beta c_0^5 r^8 c_1 - 1246112 \beta c_0^6 r^{10} c_1 - 2069600 \beta c_0^7 r^{12} c_1 - 2050048 \beta c_0^{10} r^{18} c_1 - 1227408 \beta c_0^{11} r^{20} c_1 \\
& - 556920 \beta c_0^{12} r^{22} c_1 - 185360 \beta c_0^{13} r^{24} c_1 - 42720 \beta c_0^{14} r^{26} c_1 + 8 c_1 + 4 c_1^2 c_0^7 r^{14} + 239 \beta c_0^{17} r^{32} + 7 \beta c_0^{15} r^{28} c_1^2 \\
& \left. - 406 \beta c_0^{16} r^{30} c_1 + 4 c_0^9 r^{18} \right\} \left[ (-c_1 + c_0 r^2)^2 \left( 192 \beta c_0 - 140 c_0 r^2 - 392 c_0^2 r^4 - 630 c_0^3 r^6 - 616 c_0^4 r^8 - 364 c_0^5 r^{10} \right. \right. \\
& - 120 c_0^6 r^{12} - 17 c_0^7 r^{14} - 24 + 11760 \beta c_0^3 r^4 + 38304 \beta c_0^4 r^6 + 87472 \beta c_0^5 r^8 + 148512 \beta c_0^6 r^{10} + 193020 \beta c_0^7 r^{12} \\
& \left. \left. + 194480 \beta c_0^8 r^{14} + 152152 \beta c_0^9 r^{16} + 91728 \beta c_0^{10} r^{18} + 41860 \beta c_0^{11} r^{20} + 14000 \beta c_0^{12} r^{22} + 3240 \beta c_0^{13} r^{24} + 464 \beta c_0^{14} r^{26} \right. \right. \\
& \left. \left. + 31 \beta c_0^{15} r^{28} + 2240 \beta c_0^2 r^2 \right) \right]. \tag{D}
\end{aligned}$$

Using the approximate form of the energy-density listed in Appendix A we get:

$$\rho(r) \approx (192 \beta c_0 - 24) c_0 + (-140 c_0 + 2240 \beta c_0^2) c_0 r^2 + (-392 c_0^2 + 11760 \beta c_0^3) c_0 r^4 + \mathcal{O}(r^{-6})$$

$$r = \frac{\sqrt{7} \sqrt{\left( 5 c_0 - 80 \beta c_0^2 + \sqrt{-5120 \beta^2 c_0^4 + 1024 \beta c_0^3 - 23 c_0^2 + 480 \beta c_0^2 \pi \rho - 16 c_0 \pi \rho} \right)}}{14 c_0 \sqrt{30 \beta c_0 - 1}}, \tag{D_1}$$

where the first equation of Eq. (D<sub>1</sub>) has four roots and we write only the real one given by the second equation of Eq. (D<sub>1</sub>). The second equation of (D<sub>1</sub>) ensures that the constant  $c_0$  must not equal zero and  $\beta \neq \frac{1}{30 c_0}$ . Using Eq.

(D<sub>1</sub>) in the form of  $p_r$  and  $p_\theta$  displayed in Appendix A we get:

$$p_r(\rho) = \frac{1}{2c_1^3\pi} \left\{ 332416\pi\rho\beta c_0 - 12983040\pi\rho\beta^2 c_0^2 - 87808c_1^2\pi\rho - 25088c_1\pi\rho - 10976c_1^3\pi\rho - 1551032\Upsilon\beta c_0 c_1 - 3136\pi\rho + 252073920\Upsilon\beta^2 c_0^2 c_1^2 - 2857680\beta c_1^3\beta^2 c_0^2 + 56560000\Upsilon\beta^2 c_0^2 c_1 - 6134688\Upsilon\beta c_0 c_1^2 - 314874\Upsilon c_1^3\beta c_0 - 4465036800\Upsilon\beta^3 c_0^3 c_1^2 + 284004000\Upsilon\beta^3 c_0^3 c_1^3 - 896582400\Upsilon\beta^3 c_0^3 c_1 + 28449792000\Upsilon\beta^4 c_0^4 c_1^2 - 2963520000\Upsilon\beta^4 c_0^4 c_1^3 + 5193216000\Upsilon\beta^4 c_0^4 c_1 - 1354752000\beta^4 c_0^4 \pi\rho + 220147200\beta^3 c_0^3 \pi\rho - 85349376000\beta^4 c_0^4 \pi\rho c_1^2 + 8890560000\beta^4 c_0^4 \pi\rho c_1^3 - 15579648000\beta^4 c_0^4 \pi\rho c_1 + 10905753600\beta^3 c_0^3 \pi\rho c_1^2 - 592704000\beta^3 c_0^3 \pi\rho c_1^3 + 2235340800\beta^3 c_0^3 \pi\rho c_1 + 1960\Upsilon - 3612672000\beta^5 c_0^6 - 227598336000\beta^5 c_0^6 c_1^2 + 23708160000\beta^5 c_0^6 c_1^3 - 41545728000\beta^5 c_0^6 c_1 - 165580800\beta^4 c_0^5 - 2694451200\beta^4 c_0^5 c_1 - 18334310400\beta^4 c_0^5 c_1^2 + 3358656000\beta^4 c_0^5 c_1^3 + 451584000\Upsilon\beta^4 c_0^4 - 86553600\Upsilon\beta^3 c_0^3 + 6028960\Upsilon\beta^2 c_0^2 - 180320\Upsilon\beta c_0 + 15652\Upsilon c_1 + 54656\Upsilon c_1^2 + 6811\Upsilon c_1^3 + 658560c_1^3\beta c_0 \pi\rho - 119669760\beta^2 c_0^2 c_1 \pi\rho + 2834944\beta c_0 c_1 \pi\rho - 521579520\beta^2 c_0^2 c_1^2 \pi\rho + 11063808\beta c_0 c_1^2 \pi\rho - 129477600c_1^3\beta^3 c_0^4 - 79125312\beta^2 c_0^3 c_1^2 + 570384640\beta^3 c_0^4 c_1 + 119870c_1^3\beta c_0^2 - 20165376\beta^2 c_0^3 c_1 + 2744071680\beta^3 c_0^4 c_1^2 - 4903920c_1^3\beta^2 c_0^3 + 99624\beta c_0^2 c_1 + 252832c_1^2\beta c_0^2 + 392c_0 + 14112\beta c_0^2 - 2356704\beta^2 c_0^3 + 56949760\beta^3 c_0^4 + 2996c_0 c_1 + 9860c_1^2 c_0 + 1129c_1^3 c_0 \right\},$$

$$p_\theta(\rho) = -\frac{1}{392c_1^3\pi(30\beta c_0 - 1)^2} \left\{ 172800\pi\rho\beta^2 c_0^2 + 23080\beta c_0 c_1 \Upsilon + 31360\beta c_0 c_1^2 \Upsilon - 238400\beta^2 c_0^2 c_1 \Upsilon - 245280\beta^2 c_0^2 c_1^2 \Upsilon - 2576c_1^3\beta c_0 \Upsilon + 10080c_1^3\beta^2 c_0^2 \Upsilon + 5360\beta c_0 \Upsilon - 57600\beta^2 c_0^2 \Upsilon - 110\Upsilon - 352c_1 \Upsilon - 224c_1^2 \Upsilon - 7c_1^3 \Upsilon - 11040\pi\rho\beta c_0 + 176\pi\rho + 896c_1^2\pi\rho + 832c_1\pi\rho + 168c_1^3\pi\rho - 100800c_1^3\beta^2 c_0^2 \pi\rho - 1680c_1^3\beta c_0 \pi\rho + 1219200\beta^2 c_0^2 c_1 \pi\rho - 65600\beta c_0 c_1 \pi\rho + 2499840\beta^2 c_0^2 c_1^2 \pi\rho - 110208\beta c_0 c_1^2 \pi\rho + 1612800c_1^3\beta^3 c_0^4 + 3822112\beta^2 c_0^3 c_1^2 - 10188800\beta^3 c_0^4 c_1 + 2128c_1^3\beta c_0^2 + 2193600\beta^2 c_0^3 c_1 - 12149760\beta^3 c_0^4 c_1^2 - 358400c_1^3\beta^2 c_0^3 - 73208\beta c_0^2 c_1 - 99008c_1^2\beta c_0^2 - 22c_0 - 1744\beta c_0^2 + 66560\beta^2 c_0^3 + 460800\beta^3 c_0^4 + 736c_0 c_1 + 784c_1^2 c_0 + 77c_1^3 c_0 \right\},$$

$$p(\rho) = \left\{ c_0 \left( \frac{752}{7} \frac{\beta \Upsilon_1}{\Upsilon_2} - 16 - \frac{\alpha \Upsilon_1^{14} c_1}{182059119829942534144 c_0^{13} \Upsilon_2^{14}} - \frac{715\beta \Upsilon_1^7 c_1}{843308032 c_0^6 \Upsilon_2^7} - \frac{143\beta \Upsilon_1^8 c_1}{6746464256 c_0^7 \Upsilon_2^8} - \frac{15\beta \Upsilon_1^2 c_1}{7c_0 \Upsilon_2^2} \right. \right. \\ - \frac{71\beta \Upsilon_1^4 c_1}{5488 c_0^3 \Upsilon_2^4} - \frac{51\beta \Upsilon_1^5 c_1}{76832 c_0^4 \Upsilon_2^5} - \frac{3217\beta \Upsilon_1^6 c_1}{120472576 c_0^5 \Upsilon_2^6} - \frac{39\beta \Upsilon_1^9 c_1}{94450499584 c_0^8 \Upsilon_2^9} - \frac{65\beta \Upsilon_1^{10} c_1}{10578455953408 c_0^9 \Upsilon_2^{10}} - \frac{5\beta \Upsilon_1^{11} c_1}{74049191673856 c_0^{10} \Upsilon_2^{11}} \\ - 64\beta c_0 c_1 - \frac{19}{98} \frac{\beta \Upsilon_1^3 c_1}{c_0^2 \Upsilon_3^3} - \frac{15}{29027283136151552} \frac{\beta \Upsilon_1^{12} c_1}{c_0^{11} \Upsilon_2^{12}} - \frac{1}{406381963906121728} \frac{\beta \Upsilon_1^{13} c_1}{c_0^{12} \Upsilon_2^{13}} - \frac{30}{7} \frac{\Upsilon_1}{c_0 \Upsilon_2} - \frac{15}{28} \frac{\Upsilon_1^2}{c_0^2 \Upsilon_2^2} \\ - 8c_1 - \frac{15\Upsilon_1^3}{392c_0^3 \Upsilon_2^3} - \frac{75\Upsilon_1^4}{43904c_0^4 \Upsilon_2^4} - \frac{15\Upsilon_1^5}{307328c_0^5 \Upsilon_2^5} - \frac{15\Upsilon_1^6}{17210368c_0^6 \Upsilon_2^6} - \frac{15\Upsilon_1^7}{1686616064c_0^7 \Upsilon_2^7} - \frac{15\Upsilon_1^8}{377801998336c_0^8 \Upsilon_2^8} - \frac{16\beta \Upsilon_1 c_1}{\Upsilon_2} \\ + \frac{148}{7} \frac{\beta^2}{c_0 \Upsilon_2^2} + \frac{515}{196} \frac{\beta \Upsilon_1^3}{c_0^2 \Upsilon_2^3} + \frac{635}{2744} \frac{\beta \Upsilon_1^4}{c_0^3 \Upsilon_2^4} + \frac{2351}{153664} \frac{\beta \Upsilon_1^5}{c_0^4 \Upsilon_2^5} + \frac{11789}{15059072} \frac{\beta \Upsilon_1^6}{c_0^5 \Upsilon_2^6} + \frac{15167}{481890304} \frac{\beta \Upsilon_1^7}{c_0^6 \Upsilon_2^7} + \frac{23595}{23612624896} \frac{\beta \Upsilon_1^8}{c_0^7 \Upsilon_2^8} \\ + \frac{4719}{188900999168} \frac{\beta \Upsilon_1^9}{c_0^8 \Upsilon_2^9} + \frac{1287}{2644613988352} \frac{\beta \Upsilon_1^{10}}{c_0^9 \Upsilon_2^{10}} + \frac{2145}{296196766695424} \frac{\beta \Upsilon_1^{11}}{c_0^{10} \Upsilon_2^{11}} + \frac{165}{2073377366867968} \frac{\beta \Upsilon_1^{12}}{c_0^{11} \Upsilon_2^{12}} \\ + \frac{495}{812763927812243456} \frac{\beta \Upsilon_1^{13}}{c_0^{12} \Upsilon_2^{13}} + \frac{33}{11378694989371408384} \frac{\beta \Upsilon_1^{14}}{c_0^{13} \Upsilon_2^{14}} - \frac{1}{13492928512} \frac{\Upsilon_1^7 c_1}{c_0^7 \Upsilon_2^7} - \frac{1}{60236288} \frac{\Upsilon_1^6 c_1}{c_0^6 \Upsilon_2^6} - \frac{\Upsilon_1 c_1}{c_0 \Upsilon} \\ \left. \left. - \frac{\Upsilon_1^2 c_1}{14c_0^2 \Upsilon^2} - \frac{5}{1568} \frac{\Upsilon_1^3 c_1}{c_0^3 \Upsilon^3} - \frac{1}{10976} \frac{\Upsilon_1^4 c_1}{c_0^4 \Upsilon_2^4} - \frac{1}{614656} \frac{\Upsilon_1^5 c_1}{c_0^5 \Upsilon_2^5} + \frac{33}{5097655355238390956032} \frac{\beta \Upsilon_1^{15}}{c_0^{14} \Upsilon_2^{15}} + 256\beta c_0 \right) \right\} \\ \left[ 24\pi \left( \frac{5c_0 - 80\beta c_0^2 + \Upsilon}{28c_0 \Upsilon} - c_1 \right) \right]^{-1} + \left\{ c_0 \left( - \frac{381}{11378694989371408384} \frac{\beta \Upsilon_1^{14} c_1}{c_0^{13} \Upsilon^{14}} - \frac{41423}{210827008} \frac{\beta \Upsilon_1^7 c_1}{c_0^6 \Upsilon_2^7} - \frac{23595}{3373232128} \frac{\beta \Upsilon_1^8 c_1}{c_0^7 \Upsilon_2^8} \right. \right. \\ - \frac{376}{7} \frac{\beta \Upsilon_1^2 c_1}{c_0 \Upsilon_2^2} - \frac{1675}{196} \frac{\beta \Upsilon_1^3 c_1}{c_0^2 \Upsilon_2^3} - \frac{2531}{2744} \frac{\beta \Upsilon_1^4 c_1}{c_0^3 \Upsilon_2^4} - \frac{5563}{76832} \frac{\beta \Upsilon_1^5 c_1}{c_0^4 \Upsilon_2^5} - \frac{64675}{15059072} \frac{\beta \Upsilon_1^6 c_1}{c_0^5 \Upsilon_2^6} - \frac{143}{737894528} \frac{\beta \Upsilon_1^9 c_1}{c_0^8 \Upsilon^9} \\ - \frac{10959}{2644613988352} \frac{\beta \Upsilon_1^{10} c_1}{c_0^9 \Upsilon_2^{10}} - \frac{9945}{148098383347712} \frac{\beta \Upsilon_1^{11} c_1}{c_0^{10} \Upsilon_2^{11}} - \frac{1655}{2073377366867968} \frac{\beta \Upsilon_1^{12} c_1}{c_0^{11} \Upsilon_2^{12}} - \frac{1335}{203190981953060864} \frac{\beta \Upsilon_1^{13} c_1}{c_0^{12} \Upsilon_2^{13}} \\ \left. \left. + \frac{15}{7} \frac{\beta c_1^2 \Upsilon_1^2}{c_0 \Upsilon^2} + \frac{57}{196} \frac{\beta c_1^2 \Upsilon_1^3}{c_0^2 \Upsilon_2^3} + \frac{71}{2744} \frac{\beta c_1^2 \Upsilon_1^4}{c_0^3 \Upsilon_2^4} + \frac{255}{153664} \frac{\beta c_1^2 \Upsilon_1^5}{c_0^4 \Upsilon_2^5} + \frac{9651}{120472576} \frac{\beta c_1^2 \Upsilon_1^6}{c_0^5 \Upsilon_2^6} + \frac{715}{240945152} \frac{\beta c_1^2 \Upsilon_1^7}{c_0^6 \Upsilon_2^7} \right) \right\}$$

$$\begin{aligned}
& -320 \beta c_0 c_1 + \frac{143}{1686616064} \frac{\beta \Upsilon_1^8 c_1^2}{c_0^7 \Upsilon_2^8} + \frac{351}{188900999168} \frac{\beta \Upsilon_1^9 c_1^2}{c_0^8 \Upsilon_2^9} + \frac{325}{10578455953408} \frac{\beta \Upsilon_1^{10} c_1^2}{c_0^9 \Upsilon_2^{10}} + \frac{55}{148098383347712} \frac{\beta \Upsilon_1^{11} c_1^2}{c_0^{10} \Upsilon_2^{11}} \\
& + 8 c_1 + \frac{45}{14513641568075776} \frac{\beta \Upsilon_1^{12} c_1^2}{c_0^{11} \Upsilon_2^{12}} + \frac{13}{812763927812243456} \frac{\beta \Upsilon_1^{13} c_1^2}{c_0^{12} \Upsilon_2^{13}} + \frac{1}{26008445689991790592} \frac{\beta \Upsilon_1^{14} c_1^2}{c_0^{13} \Upsilon_2^{14}} \\
& - \frac{29}{364118239659885068288} \frac{\beta \Upsilon_1^{15} c_1}{c_0^{14} \Upsilon_2^{15}} - \frac{240}{7} \frac{\beta \Upsilon_1}{\Upsilon} + \frac{8}{7} \frac{-80 \beta c_0^2 + 5 c_0 + \Upsilon}{c_0 \Upsilon_2} + \frac{57}{196} \frac{\Upsilon_1^2}{c_0^2 \Upsilon_2^2} + \frac{25}{784} \frac{\Upsilon_1^3}{c_0^3 \Upsilon_2^3} + \frac{43}{21952} \frac{\Upsilon_1^4}{c_0^4 \Upsilon_2^4} \\
& + \frac{45}{614656} \frac{\Upsilon_1^5}{c_0^5 \Upsilon_2^5} + \frac{29}{17210368} \frac{\Upsilon_1^6}{c_0^6 \Upsilon_2^6} + \frac{11}{481890304} \frac{\Upsilon_1^7}{c_0^7 \Upsilon_2^7} + \frac{15}{94450499584} \frac{\Upsilon_1^8}{c_0^8 \Upsilon_2^8} + \frac{1}{2644613988352} \frac{\Upsilon_1^9}{c_0^9 \Upsilon_2^9} - \frac{200 \beta \Upsilon_1 c_1}{\Upsilon_2} \\
& - \frac{570}{49} \frac{\beta \Upsilon_1^2}{c_0 \Upsilon_2^2} - \frac{173}{98} \frac{\beta \Upsilon_1^3}{c_0^2 \Upsilon_2^3} - \frac{15}{98} \frac{\beta \Upsilon_1^4}{c_0^3 \Upsilon_2^4} - \frac{165}{21952} \frac{\beta \Upsilon_1^5}{c_0^4 \Upsilon_2^5} - \frac{13}{134456} \frac{\beta \Upsilon_1^6}{c_0^5 \Upsilon_2^6} + \frac{13245}{843308032} \frac{\beta \Upsilon_1^7}{c_0^6 \Upsilon_2^7} + \frac{138357}{94450499584} \frac{\beta \Upsilon_1^8}{c_0^7 \Upsilon_2^8} \\
& + \frac{96525}{1322306994176} \frac{\beta \Upsilon_1^9}{c_0^8 \Upsilon_2^9} + \frac{3289}{1322306994176} \frac{\beta \Upsilon_1^{10}}{c_0^9 \Upsilon_2^{10}} + \frac{9087}{148098383347712} \frac{\beta \Upsilon_1^{11}}{c_0^{10} \Upsilon_2^{11}} + \frac{9165}{8293509467471872} \frac{\beta \Upsilon_1^{12}}{c_0^{11} \Upsilon_2^{12}} \\
& + \frac{1655}{116109132544606208} \frac{\beta \Upsilon_1^{13}}{c_0^{12} \Upsilon_2^{13}} + \frac{1425}{11378694989371408384} \frac{\beta \Upsilon_1^{14}}{c_0^{13} \Upsilon_2^{14}} + \frac{29}{1686616064} \frac{\Upsilon_1^7 c_1}{c_0^7 \Upsilon_2^7} + \frac{13}{8605184} \frac{\Upsilon_1^6 c_1}{c_0^6 \Upsilon_2^6} \\
& + \frac{22}{7} \frac{\Upsilon_1 c_1}{c_0 \Upsilon_2} + \frac{\Upsilon_1^2 c_1}{2 c_0^2 \Upsilon_2^2} + \frac{17}{392} \frac{\Upsilon_1^3 c_1}{c_0^3 \Upsilon_2^3} + \frac{25}{10976} \frac{\Upsilon_1^4 c_1}{c_0^4 \Upsilon_2^4} + \frac{23}{307328} \frac{\Upsilon_1^5 c_1}{c_0^5 \Upsilon_2^5} + \frac{429}{637206919404798869504} \frac{\beta \Upsilon_1^{15}}{c_0^{14} \Upsilon_2^{15}} \\
& + \frac{1}{11806312448} \frac{\Upsilon_1^8 c_1}{c_0^8 \Upsilon_2^8} + \frac{239}{142734349946674946768896} \frac{\beta \Upsilon_1^{16}}{c_0^{15} \Upsilon_2^{16}} + \frac{c_1^2 \Upsilon_1}{c_0 \Upsilon_2} + \frac{3}{28} \frac{c_1^2 \Upsilon_1^2}{c_0^2 \Upsilon_2^2} + \frac{5}{784} \frac{c_1^2 \Upsilon_1^3}{c_0^3 \Upsilon_2^3} + \frac{5}{21952} \frac{c_1^2 \Upsilon_1^4}{c_0^4 \Upsilon_2^4} \\
& + \frac{3}{614656} \frac{c_1^2 \Upsilon_1^5}{c_0^5 \Upsilon_2^5} + \frac{1}{17210368} \frac{c_1^2 \Upsilon_1^6}{c_0^6 \Upsilon_2^6} + \frac{1}{3373232128} \frac{c_1^2 \Upsilon_1^7}{c_0^7 \Upsilon_2^7} + 8 \frac{\beta c_1^2 \Upsilon_1}{\Upsilon_2} \Big) \Big\} \left[ 6\pi \left( \frac{\Upsilon_1}{28 c_0 \Upsilon_2} - c_1 \right)^2 \right]^{-1}, \quad (\mathbf{D}_2)
\end{aligned}$$

where  $\Upsilon = \sqrt{c_0 (480 \pi \rho \beta c_0 - 5120 \beta^2 c_0^3 + 1024 \beta c_0^2 - 23 c_0 - 16 \pi \rho)}$ ,  $\Upsilon_1 = (-80 \beta c_0^2 + 5 c_0 + \Upsilon)$ ,  $\Upsilon_2 = (-1 + 30 \beta c_0)$ .

### Appendix E: The different forces of on the model.

The forces that act on the model presented in Appendix A have the following forms:

$$\begin{aligned}
F_g &= \frac{c_0^2 r}{16 (c_0 r^2 - c_1)^6 \pi} \left( 21 c_0^5 r^{10} c_1 - 9 c_0 r^2 - 2 \beta c_0^{15} r^{28} c_1 - 168 \beta c_0^2 r^2 c_1 - 12870 \beta c_0^8 r^{14} c_1 - 10010 \beta c_0^9 r^{16} c_1 \right. \\
&\quad - 840 \beta c_0^3 r^4 c_1 - 2660 \beta c_0^4 r^6 c_1 - 5964 \beta c_0^5 r^8 c_1 - 9996 \beta c_0^6 r^{10} c_1 - 12868 \beta c_0^7 r^{12} c_1 - 6006 \beta c_0^{10} r^{18} c_1 - 2730 \beta c_0^{11} r^{20} c_1 \\
&\quad - 910 \beta c_0^{12} r^{22} c_1 - 210 \beta c_0^{13} r^{24} c_1 - 30 \beta c_0^{14} r^{26} c_1 + c_1 + 1176 \beta c_0^3 r^4 + 4340 \beta c_0^4 r^6 + 11284 \beta c_0^5 r^8 + 21924 \beta c_0^6 r^{10} \\
&\quad + 32860 \beta c_0^7 r^{12} + 38606 \beta c_0^8 r^{14} + 35750 \beta c_0^9 r^{16} + 26026 \beta c_0^{10} r^{18} + 14742 \beta c_0^{11} r^{20} + 6370 \beta c_0^{12} r^{22} + 2030 \beta c_0^{13} r^{24} \\
&\quad + 450 \beta c_0^{14} r^{26} + 62 \beta c_0^{15} r^{28} + 200 \beta c_0^2 r^2 + c_0^7 r^{14} c_1 + 7 c_0^6 r^{12} c_1 + 7 c_0 r^2 c_1 + 21 c_0^2 r^4 c_1 + 35 c_0^3 r^6 c_1 + 35 c_0^4 r^8 c_1 \\
&\quad \left. - 16 \beta c_0 c_1 + 4 \beta c_0^{16} r^{30} + 16 \beta c_0 - 35 c_0^2 r^4 - 77 c_0^3 r^6 - 105 c_0^4 r^8 - 91 c_0^5 r^{10} - 49 c_0^6 r^{12} - 15 c_0^7 r^{14} - 2 c_0^8 r^{16} - 1 \right), \\
F_a &= \frac{c_0^2}{4r\pi (c_0 r^2 - c_1)^2} \left\{ 80 r^2 - 11648 \beta c_0^{14} r^{28} c_1 - 8128 \beta c_0 r^2 c_1 - 4864600 \beta c_0^7 r^{14} c_1 - 4896320 \beta c_0^8 r^{16} c_1 \right. \\
&\quad - 67200 \beta c_0^2 r^4 c_1 - 315840 \beta c_0^3 r^6 c_1 - 987392 \beta c_0^4 r^8 c_1 - 2220960 \beta c_0^5 r^{10} c_1 - 3750528 \beta c_0^6 r^{12} c_1 - 3827824 \beta c_0^9 r^{18} c_1 \\
&\quad - 2306304 \beta c_0^{10} r^{20} c_1 - 1051960 \beta c_0^{11} r^{22} c_1 - 351680 \beta c_0^{12} r^{24} c_1 - 81360 \beta c_0^{13} r^{26} c_1 + 1680 \beta c_0^2 c_1^2 r^4 + 8512 \beta c_0^3 c_1^2 r^6 \\
&\quad + 23856 \beta c_0^4 c_1^2 r^8 + 45696 \beta c_0^5 c_1^2 r^{10} + 64340 \beta c_0^6 c_1^2 r^{12} + 68640 \beta c_0^7 c_1^2 r^{14} + 56056 \beta c_0^8 r^{16} c_1^2 + 34944 \beta c_0^9 r^{18} c_1^2 \\
&\quad + 16380 \beta c_0^{10} r^{20} c_1^2 + 5600 \beta c_0^{11} r^{22} c_1^2 + 1320 \beta c_0^{12} r^{24} c_1^2 + 192 \beta c_0^{13} r^{26} c_1^2 + 13 \beta c_0^{14} r^{28} c_1^2 - 778 \beta c_0^{15} r^{30} c_1 \\
&\quad + 1209208 \beta c_0^{10} r^{20} + 1166880 \beta c_0^9 r^{18} + 682180 \beta c_0^8 r^{16} + 46592 \beta c_0^7 r^{14} - 400960 \beta c_0^5 r^{10} - 356496 \beta c_0^6 r^{12} \\
&\quad - 94080 \beta c_0^3 r^6 - 245840 \beta c_0^4 r^8 + 352 c_0^6 r^{14} c_1 - 2176 \beta c_0 r^2 + 392 c_0 r^4 c_1 + 1064 c_0^5 r^{12} c_1 + 873600 \beta c_0^{11} r^{22} \\
&\quad + 453180 \beta c_0^{12} r^{24} + 166880 \beta c_0^{13} r^{26} + 41640 \beta c_0^{14} r^{28} + 6336 \beta c_0^{15} r^{30} + 1792 c_0^4 r^{10} c_1 + 1820 c_0^3 r^8 c_1 + 1120 c_0^2 r^6 c_1 \\
&\quad + 112 c_1^2 c_0 r^4 + 210 c_1^2 c_0^2 r^6 + 50 c_0^7 r^{16} c_1 + 445 \beta c_0^{16} r^{32} + 140 c_1^2 c_0^4 r^{10} + 224 c_1^2 c_0^3 r^8 + 7 c_1^2 c_0^6 r^{14} + 48 c_1^2 c_0^5 r^{12} \\
&\quad - 21248 \beta c_0^2 r^4 + 64 r^2 c_1 - 384 \beta c_1^2 + 28 r^2 c_1^2 + 576 c_0 r^4 + 1820 c_0^2 r^6 + 3248 c_0^3 r^8 + 3570 c_0^4 r^{10} \\
&\quad \left. + 2464 c_0^5 r^{12} + 1036 c_0^6 r^{14} + 240 c_0^7 r^{16} + 23 c_0^8 r^{18} \right\}, \\
F_h &= \frac{c_0}{8\pi (c_1 - c_0 r^2)} \left\{ - 28 \beta c_0^{15} r^{27} c_1 - 896 \beta c_0^2 r c_1 - 160160 \beta c_0^8 r^{13} c_1 - 128128 \beta c_0^9 r^{15} c_1 - 6720 \beta c_0^3 r^3 c_1 \right. \\
&\quad - 25536 \beta c_0^4 r^5 c_1 - 63616 \beta c_0^5 r^7 c_1 - 114240 \beta c_0^6 r^9 c_1 - 154416 \beta c_0^7 r^{11} c_1 - 78624 \beta c_0^{10} r^{17} c_1 - 36400 \beta c_0^{11} r^{19} c_1 \\
&\quad - 12320 \beta c_0^{12} r^{21} c_1 - 2880 \beta c_0^{13} r^{23} c_1 - 416 \beta c_0^{14} r^{25} c_1 - 240 c_0^8 r^{15} - 240 c_0 r - 1680 c_0^2 r^3 - 5040 c_0^3 r^5 - 8400 c_0^4 r^7 \\
&\quad - 8400 c_0^5 r^9 - 5040 c_0^6 r^{11} - 1680 c_0^7 r^{13} + 66304 \beta c_0^3 r^3 + 346080 \beta c_0^4 r^5 + 1137920 \beta c_0^5 r^7 + 2633120 \beta c_0^6 r^9 \\
&\quad + 4526976 \beta c_0^7 r^{11} + 5945464 \beta c_0^8 r^{13} + 6040320 \beta c_0^9 r^{15} + 4756752 \beta c_0^{10} r^{17} + 2882880 \beta c_0^{11} r^{19} + 1321320 \beta c_0^{12} r^{21} \\
&\quad + 443520 \beta c_0^{13} r^{23} + 102960 \beta c_0^{14} r^{25} + 14784 \beta c_0^{15} r^{27} + 6016 \beta c_0^2 r - 14 c_0^7 r^{13} c_1 - 96 c_0^6 r^{11} c_1 - 56 c_0 r c_1 - 224 c_0^2 r^3 c_1 \\
&\quad \left. - 420 c_0^3 r^5 c_1 - 448 c_0^4 r^7 c_1 - 280 c_0^5 r^9 c_1 + 990 \beta c_0^{16} r^{29} \right\}, \quad (\text{E})
\end{aligned}$$

Appendix F: The form of the mass in terms of the central density at the surface of the star

The form of the mass in terms of the central density at the surface of the star takes the form:

$$\begin{aligned}
M(\rho_0) = & \frac{1}{2836619246897071128576\beta^{15}\pi} \left\{ (3 + \Upsilon_3) l^3 (3l^2 + l^2\Upsilon_3 + 96\beta) \left( 3l^4 + l^4\Upsilon_3 + 16l^4\beta\rho_0\pi + 768\beta^2 + 48l^2\beta \right. \right. \\
& + 16l^2\beta\Upsilon_3 \Big) \times \left( 9l^8 + 3l^8\Upsilon_3 + 96l^8\beta\rho_0\pi + 16l^8\Upsilon_3\beta\rho_0\pi + 128l^8\beta^2\rho_0^2\pi^2 + 288l^6\beta + 96l^6\beta\Upsilon_3 + 2304l^6\beta^2\rho_0\pi \right. \\
& + 256l^6\beta^2\Upsilon_3\rho_0\pi + 3456l^4\beta^2 + 1152l^4\beta^2\Upsilon_3 + 18432l^4\beta^3\rho_0\pi + 18432l^2\beta^3 + 6144l^2\beta^3\Upsilon_3 + 147456\beta^4 \Big) \left( -1358954496\beta^7 \right. \\
& + 81l^{14} + 6193152l^8\beta^4\Upsilon_3\rho_0\pi + 5505024l^8\beta^5\Upsilon_3\rho_0^2\pi^2 + 41287680l^6\beta^5\sqrt{9+96\beta\rho_0\pi}\rho_0\pi + 132120576l^4\beta^6\Upsilon_3\rho_0\pi \\
& + 23040l^{12}\beta^2\Upsilon_3\rho_0\pi + 73728l^{12}\beta^3\Upsilon_3\rho_0^2\pi^2 + 32768l^{12}\beta^4\Upsilon_3\rho_0^3\pi^3 + 145152l^{10}\beta^2 + 23224320l^6\beta^4 + 148635648l^4\beta^5 \\
& + 5184l^{12}\beta + 2322432l^8\beta^3 + 594542592l^2\beta^6 + 516096l^{10}\beta^3\Upsilon_3\rho_0\pi + 1032192l^{10}\beta^4\Upsilon_3\rho_0^2\pi^2 + 432l^{14}\Upsilon_3\beta\rho_0\pi \\
& + 1920l^{14}\Upsilon_3\beta^2\rho_0^2\pi^2 + 2048l^{14}\Upsilon_3\beta^3\rho_0^3\pi^3 + 96768l^{12}\beta^2\rho_0\pi + 516096l^{12}\beta^3\rho_0^2\pi^2 + 688128l^{12}\beta^4\rho_0^3\pi^3 + 2322432l^{10}\beta^3\rho_0\pi \\
& + 9289728l^{10}\beta^4\rho_0^2\pi^2 + 5505024l^{10}\beta^5\rho_0^3\pi^3 + 30965760l^8\beta^4\rho_0\pi + 82575360l^8\beta^5\rho_0^2\pi^2 + 247726080l^6\beta^5\rho_0\pi \\
& + 330301440l^6\beta^6\rho_0^2\pi^2 + 1189085184l^4\beta^6\rho_0\pi + 3170893824l^2\beta^7\rho_0\pi + 1728l^{14}\beta\rho_0\pi + 11520l^{14}\beta^2\rho_0^2\pi^2 \\
& + 24576l^{14}\beta^3\rho_0^3\pi^3 + 8192l^{14}\beta^4\rho_0^4\pi^4 + 452984832\Upsilon_3\beta^7 + 27l^{14}\Upsilon_3 + 1728l^{12}\beta\Upsilon_3 + 48384l^{10}\beta^2\Upsilon_3 + 774144l^8\beta^3\Upsilon_3 \\
& \left. \left. + 49545216l^4\beta^5\Upsilon_3 + 7741440l^6\beta^4\Upsilon_3 + 198180864l^2\beta^6\Upsilon_3 \right) \right\}, \tag{F1}
\end{aligned}$$

$\Upsilon_3 = \sqrt{9 + 96\beta\rho_0\pi}$ . The form of the derivative of  $M(\rho_0)$  w.r.t. the central density takes the form

$$\begin{aligned}
\frac{\partial M(\rho_0)}{\partial \rho_0} = & \frac{l^3}{461689330549653504 \beta^{14} \Upsilon_3} \left\{ 50497270528868352 \alpha^{12} l^4 + 215205838848 l^{18} \alpha^5 \Upsilon_3 + 509483520 l^{22} \beta^3 \Upsilon_3 \right. \\
& + 262440 l^{26} \beta \Upsilon_3 + 12227604480 l^{20} \beta^4 \Upsilon_3 + 6673792914948096 \beta^{10} l^8 \Upsilon_3 + 22355562473717760 \beta^{11} l^6 \Upsilon_3 \\
& + 1442426916962304 \beta^9 l^{10} \Upsilon_3 + 2869411184640 l^{16} \beta^6 \Upsilon_3 + 67329694038491136 \beta^{13} l^2 \Upsilon_3 + 14696640 l^{24} \beta^2 \Upsilon_3 \\
& + 29495597727744 \beta^7 l^{14} \Upsilon_3 + 235047487537152 \beta^8 l^{12} \Upsilon_3 + 88486793183232 \beta^7 l^{14} + 4327280750886912 \beta^9 l^{10} \\
& + 67066687421153280 \beta^{11} l^6 + 151491811586605056 \beta^{12} l^4 + 705142462611456 \beta^8 l^{12} + 201989082115473408 \beta^{13} l^2 \\
& + 44089920 l^{24} \beta^2 + 36682813440 l^{20} \beta^4 + 645617516544 l^{18} \beta^5 + 787320 l^{26} \beta + 1528450560 l^{22} \beta^3 + 8608233553920 l^{16} \beta^6 \\
& + 20021378744844288 \beta^{10} l^8 + 76948221758275584 \beta^{14} \Upsilon_3 + 2187 l^{28} \Upsilon_3 + 81618807029760 l^{18} \beta^8 \Upsilon_3 \rho_0^3 \pi^3 \\
& + 8115506380800 l^{20} \beta^7 \Upsilon_3 \rho_0^3 \pi^3 + 1672151040 l^{26} \beta^5 \Upsilon_3 \rho_0^4 \pi^4 + 901722931200 l^{22} \beta^7 \Upsilon_3 \rho_0^4 \pi^4 + 23224320 l^{28} \beta^4 \rho_0^4 \pi^4 \Upsilon_3 \\
& + 1981808640 l^{26} \beta^6 \rho_0^5 \pi^5 \Upsilon_3 + 52022476800 l^{24} \beta^6 \rho_0^4 \pi^4 \Upsilon_3 + 37158912 l^{28} \beta^5 \rho_0^5 \pi^5 \Upsilon_3 + 660602880 l^{26} \beta^7 \rho_0^6 \pi^6 \Upsilon_3 \\
& + 41617981440 l^{24} \beta^7 \rho_0^5 \pi^5 \Upsilon_3 + 412216197120 l^{22} \beta^8 \rho_0^5 \pi^5 \Upsilon_3 + 9274864435200 l^{20} \beta^8 \rho_0^4 \pi^4 \Upsilon_3 + 54412538019840 l^{18} \beta^9 \rho_0^4 \pi^4 \Upsilon_3 \\
& + 22020096 l^{28} \beta^6 \rho_0^6 \pi^6 \Upsilon_3 + 5284823040 l^{24} \beta^8 \rho_0^6 \pi^6 \Upsilon_3 + 1648864788480 l^{20} \beta^9 \rho_0^5 \pi^5 \Upsilon_3 + 2097152 l^{28} \Upsilon_3 \beta^7 \rho_0^7 \pi^7 \\
& + 53562342113280 l^{16} \beta^7 \Upsilon_3 \rho_0 \pi + 306070526361600 l^{16} \beta^8 \Upsilon_3 \rho_0^2 \pi^2 + 544125380198400 l^{16} \beta^9 \Upsilon_3 \rho_0^3 \pi^3 \\
& + 145100101386240 l^{16} \beta^{10} \Upsilon_3 \rho_0^4 \pi^4 + 471978486005760 \beta^8 l^{14} \Upsilon_3 \rho_0 \pi + 2097986565832704 \beta^9 l^{14} \Upsilon_3 \rho_0^2 \pi^2 \\
& + 2238339386179584 \beta^{10} l^{14} \Upsilon_3 \rho_0^3 \pi^3 + 3136706152759296 \beta^9 l^{12} \Upsilon_3 \rho_0 \pi + 10050610019696640 \beta^{10} l^{12} \Upsilon_3 \rho_0^2 \pi^2 \\
& + 4476678772359168 \beta^{11} l^{12} \Upsilon_3 \rho_0^3 \pi^3 + 15451638768599040 \beta^{10} l^{10} \Upsilon_3 \rho_0 \pi + 31122449718312960 \beta^{11} l^{10} \Upsilon_3 \rho_0^2 \pi^2 \\
& + 54267032044044288 \beta^{11} l^{8} \Upsilon_3 \rho_0 \pi + 49795919549300736 \beta^{12} l^{8} \Upsilon_3 \rho_0^2 \pi^2 + 126243176322170880 \beta^{12} l^6 \Upsilon_3 \rho_0 \pi \\
& + 168324235096227840 \beta^{13} l^4 \Upsilon_3 \rho_0 \pi + 13586227200 l^{22} \beta^4 \Upsilon_3 \rho_0 \pi + 8398080 l^{26} \beta^2 \Upsilon_3 \rho_0 \pi + 102643200 l^{26} \beta^3 \Upsilon_3 \rho_0^2 \pi^2 \\
& + 597196800 l^{26} \beta^4 \Upsilon_3 \rho_0^3 \pi^3 + 431101440 l^{24} \beta^3 \Upsilon_3 \rho_0 \pi + 4702924800 l^{24} \beta^4 \Upsilon_3 \rho_0^2 \pi^2 + 130427781120 l^{22} \beta^5 \Upsilon_3 \rho_0^2 \pi^2 \\
& + 293462507520 l^{20} \beta^5 \Upsilon_3 \rho_0 \pi + 4591057895424 l^{18} \beta^6 \Upsilon_3 \rho_0 \pi + 75816 l^{28} \Upsilon_3 \beta \rho_0 \pi + 1026432 l^{28} \Upsilon_3 \beta^2 \rho_0^2 \pi^2 \\
& + 6842880 l^{28} \Upsilon_3 \beta^3 \rho_0^3 \pi^3 + 541033758720 l^{22} \beta^6 \Upsilon_3 \rho_0^3 \pi^3 + 2434651914240 l^{20} \beta^6 \Upsilon_3 \rho_0^2 \pi^2 + 32137405267968 l^{18} \beta^7 \Upsilon_3 \rho_0^2 \pi^2 \\
& + 23410114560 l^{24} \beta^5 \Upsilon_3 \rho_0^3 \pi^3 + 6561 l^{28} + 1887865021661184 \beta^8 l^{14} \rho_0 \pi + 12587136637206528 \beta^9 l^{14} \rho_0^2 \pi^2 \\
& + 26856941602996224 \beta^{10} l^{14} \rho_0^3 \pi^3 + 8954749114122240 \beta^{11} l^{14} \rho_0^4 \pi^4 + 13170878258872320 \beta^9 l^{12} \rho_0 \pi \\
& + 70310435701653504 \beta^{10} l^{12} \rho_0^2 \pi^2 + 93893362389614592 \beta^{11} l^{12} \rho_0^3 \pi^3 + 69433746977193984 \beta^{10} l^{10} \rho_0 \pi \\
& + 279050020995465216 \beta^{11} l^{10} \rho_0^2 \pi^2 + 166921533137092608 \beta^{12} l^{10} \rho_0^3 \pi^3 + 269581782771302400 \beta^{11} l^8 \rho_0 \pi \\
& + 732911773648158720 \beta^{12} l^8 \rho_0^2 \pi^2 + 736418528545996800 \beta^{12} l^6 \rho_0 \pi + 1066053488942776320 \beta^{13} l^6 \rho_0^2 \pi^2 \\
& + 1312929033750577152 \beta^{13} l^4 \rho_0 \pi + 1615912656923787264 \beta^{14} l^2 \rho_0 \pi + 262440 l^{28} \beta \rho_0 \pi + 4199040 l^{28} \beta^2 \rho_0^2 \pi^2 \\
& + 34214400 l^{28} \beta^3 \rho_0^3 \pi^3 + 149299200 l^{28} \beta^4 \rho_0^4 \pi^4 + 29393280 l^{26} \beta^2 \rho_0 \pi + 431101440 l^{26} \beta^3 \rho_0^2 \pi^2 + 3135283200 l^{26} \beta^4 \rho_0^3 \pi^3 \\
& + 1528450560 l^{24} \beta^3 \rho_0 \pi + 20379340800 l^{24} \beta^4 \rho_0^2 \pi^2 + 130427781120 l^{24} \beta^5 \rho_0^3 \pi^3 + 48910417920 l^{22} \beta^4 \rho_0 \pi \\
& + 586925015040 l^{22} \beta^5 \rho_0^2 \pi^2 + 1076029194240 l^{20} \beta^5 \rho_0 \pi + 11477644738560 l^{20} \beta^6 \rho_0^2 \pi^2 + 17216467107840 l^{18} \beta^6 \rho_0 \pi \\
& + 206597605294080 l^{16} \beta^7 \rho_0 \pi + 405775319040 l^{24} \beta^6 \rho_0^4 \pi^4 + 3246202552320 l^{22} \beta^6 \rho_0^3 \pi^3 + 53562342113280 l^{20} \beta^7 \rho_0^3 \pi^3 \\
& + 160687026339840 l^{18} \beta^7 \rho_0^2 \pi^2 + 1652780842352640 l^{16} \beta^8 \rho_0^2 \pi^2 + 102023508787200 l^{20} \beta^8 \rho_0^4 \pi^4 \\
& + 612141052723200 l^{18} \beta^8 \rho_0^3 \pi^3 + 4897128421785600 l^{16} \beta^9 \rho_0^3 \pi^3 + 816188070297600 l^{18} \beta^9 \rho_0^4 \pi^4 + 11705057280 l^{26} \beta^5 \rho_0^4 \pi^4 \\
& + 8115506380800 l^{22} \beta^7 \rho_0^4 \pi^4 + 3917702737428480 l^{16} \beta^{10} \rho_0^4 \pi^4 + 334430208 l^{28} \beta^5 \rho_0^5 \pi^5 + 330301440 l^{28} \beta^6 \rho_0^6 \pi^6 \\
& + 20808990720 l^{26} \beta^6 \rho_0^5 \pi^5 + 541033758720 l^{24} \beta^7 \rho_0^5 \pi^5 + 94371840 l^{28} \beta^7 \rho_0^7 \pi^7 + 13872660480 l^{26} \beta^7 \rho_0^6 \pi^6 \\
& + 206108098560 l^{24} \beta^8 \rho_0^6 \pi^6 + 7419891548160 l^{22} \beta^8 \rho_0^5 \pi^5 + 54412538019840 l^{20} \beta^9 \rho_0^5 \pi^5 + 1509949440 l^{26} \beta^8 \rho_0^7 \pi^7 \\
& \left. + 1099243192320 l^{22} \beta^9 \rho_0^6 \pi^6 + 174120121663488 l^{18} \beta^{10} \rho_0^5 \pi^5 \right\}. \tag{F2}
\end{aligned}$$

- 
- [1] A. Banerjee, T. Tangphati, and P. Channuie, *Astrophys. J.* **909**, 14 (2021), arXiv:2006.00479 [gr-qc].  
[2] R. A. Hennigar, D. Kubiznák, R. B. Mann, and C. Pollack, *JHEP* **07**, 027 (2020), arXiv:2004.09472 [gr-qc].  
[3] A. Casalino, A. Colleaux, M. Rinaldi, and S. Vicentini, *Phys. Dark Univ.* **31**, 100770 (2021), arXiv:2003.07068 [gr-qc].  
[4] K. Aoki, M. A. Gorji, and S. Mukohyama, *Phys. Lett. B* **810**, 135843 (2020), arXiv:2005.03859 [gr-qc].  
[5] J. Aasi *et al.* (LIGO Scientific), *Class. Quant. Grav.* **32**, 074001 (2015), arXiv:1411.4547 [gr-qc].

- [6] K. Somiya (KAGRA), *Class. Quant. Grav.* **29**, 124007 (2012), arXiv:1111.7185 [gr-qc].
- [7] Y. Aso, Y. Michimura, K. Somiya, M. Ando, O. Miyakawa, T. Sekiguchi, D. Tatsumi, and H. Yamamoto (KAGRA), *Phys. Rev. D* **88**, 043007 (2013), arXiv:1306.6747 [gr-qc].
- [8] F. Acernese *et al.* (VIRGO), *Class. Quant. Grav.* **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
- [9] D. Psaltis *et al.* (Event Horizon Telescope), *Phys. Rev. Lett.* **125**, 141104 (2020), arXiv:2010.01055 [gr-qc].
- [10] G. Raaijmakers *et al.*, *Astrophys. J. Lett.* **893**, L21 (2020), arXiv:1912.11031 [astro-ph.HE].
- [11] A. V. Astashenok, S. Capozziello, and S. D. Odintsov, *JCAP* **12**, 040 (2013), arXiv:1309.1978 [gr-qc].
- [12] A. V. Astashenok, S. Capozziello, and S. D. Odintsov, *Phys. Rev. D* **89**, 103509 (2014), arXiv:1401.4546 [gr-qc].
- [13] R. Abbott *et al.* (LIGO Scientific, Virgo), *Astrophys. J. Lett.* **896**, L44 (2020), arXiv:2006.12611 [astro-ph.HE].
- [14] K. Huang, J. Hu, Y. Zhang, and H. Shen, *Astrophys. J.* **904**, 39 (2020), arXiv:2008.04491 [nucl-th].
- [15] I. Bombaci, A. Drago, D. Logoteta, G. Pagliara, and I. Vidaña, *Phys. Rev. Lett.* **126**, 162702 (2021), arXiv:2010.01509 [nucl-th].
- [16] Z. Roupas, G. Panotopoulos, and I. Lopes, *Phys. Rev. D* **103**, 083015 (2021), arXiv:2010.11020 [astro-ph.HE].
- [17] X. Zhou, A. Li, and B.-A. Li, *Astrophys. J.* **910**, 62 (2021), arXiv:2011.11934 [astro-ph.HE].
- [18] A. Awad and G. Nashed, *JCAP* **1702**, 046 (2017), arXiv:1701.06899 [gr-qc].
- [19] E. R. Most, L. J. Papenfort, L. R. Weih, and L. Rezzolla, *Mon. Not. Roy. Astron. Soc.* **499**, L82 (2020), arXiv:2006.14601 [astro-ph.HE].
- [20] G. G. L. Nashed, *EPL* **105**, 10001 (2014), arXiv:1501.00974 [gr-qc].
- [21] G. G. L. Nashed, *Int. J. Mod. Phys. A* **21**, 3181 (2006), arXiv:gr-qc/0501002.
- [22] H. Tan, J. Noronha-Hostler, and N. Yunes, *Phys. Rev. Lett.* **125**, 261104 (2020), arXiv:2006.16296 [astro-ph.HE].
- [23] K. Vattis, I. S. Goldstein, and S. M. Koushiappas, *Phys. Rev. D* **102**, 061301 (2020), arXiv:2006.15675 [astro-ph.HE].
- [24] N.-B. Zhang and B.-A. Li, *Astrophys. J.* **902**, 38 (2020), arXiv:2007.02513 [astro-ph.HE].
- [25] F. J. Fattoyev, C. J. Horowitz, J. Piekarewicz, and B. Reed, *Phys. Rev. C* **102**, 065805 (2020), arXiv:2007.03799 [nucl-th].
- [26] T. Shirafuji, G. G. L. Nashed, and Y. Kobayashi, *Prog. Theor. Phys.* **96**, 933 (1996), arXiv:gr-qc/9609060.
- [27] A. Tsokaros, M. Ruiz, and S. L. Shapiro, *Astrophys. J.* **905**, 48 (2020), arXiv:2007.05526 [astro-ph.HE].
- [28] I. Tews, P. T. H. Pang, T. Dietrich, M. W. Coughlin, S. Antier, M. Bulla, J. Heinzel, and L. Issa, *Astrophys. J. Lett.* **908**, L1 (2021), arXiv:2007.06057 [astro-ph.HE].
- [29] V. Dexheimer, R. O. Gomes, T. Klähn, S. Han, and M. Salinas, *Phys. Rev. C* **103**, 025808 (2021), arXiv:2007.08493 [astro-ph.HE].
- [30] D. A. Godzieba, D. Radice, and S. Bernuzzi, *Astrophys. J.* **908**, 122 (2021), arXiv:2007.10999 [astro-ph.HE].
- [31] A. Kanakis-Pegios, P. S. Koliogiannis, and C. C. Moustakidis, *Symmetry* **13**, 183 (2021), arXiv:2012.09580 [astro-ph.HE].
- [32] A. Nathanail, E. R. Most, and L. Rezzolla, *Astrophys. J. Lett.* **908**, L28 (2021), arXiv:2101.01735 [astro-ph.HE].
- [33] Z. Roupas, *Astrophys. Space Sci.* **366**, 9 (2021), arXiv:2007.10679 [gr-qc].
- [34] B. Biswas, R. Nandi, P. Char, S. Bose, and N. Stergioulas, *mnras* **505**, 1600 (2021), arXiv:2010.02090 [astro-ph.HE].
- [35] R. C. Nunes, J. G. Coelho, and J. C. N. de Araujo, *Eur. Phys. J. C* **80**, 1115 (2020), arXiv:2008.10395 [astro-ph.HE].
- [36] A. V. Astashenok, S. Capozziello, S. D. Odintsov, and V. K. Oikonomou, *Phys. Lett. B* **811**, 135910 (2020), arXiv:2008.10884 [gr-qc].
- [37] A. V. Astashenok, S. Capozziello, S. D. Odintsov, and V. K. Oikonomou, *Phys. Lett. B* **816**, 136222 (2021), arXiv:2103.04144 [gr-qc].
- [38] G. Mustafa, M. F. Shamir, and X. Tie-Cheng, *Phys. Rev. D* **101**, 104013 (2020), arXiv:2005.03997 [gr-qc].
- [39] G. G. L. Nashed, *Astrophys. J.* **919**, 113 (2021), arXiv:2108.04060 [gr-qc].
- [40] G. G. L. Nashed, S. D. Odintsov, and V. K. Oikonomou, *Eur. Phys. J. C* **81**, 528 (2021), arXiv:2106.13607 [gr-qc].
- [41] G. G. L. Nashed and S. Capozziello, *Eur. Phys. J. C* **81**, 481 (2021), arXiv:2105.11975 [gr-qc].
- [42] A. V. Astashenok, S. Capozziello, S. D. Odintsov, and V. K. Oikonomou, (2021), 10.1209/0295-5075/ac3d6c, arXiv:2111.14179 [gr-qc].
- [43] A. V. Astashenok, S. Capozziello, S. D. Odintsov, and V. K. Oikonomou, *EPL* **134**, 59001 (2021), arXiv:2106.01234 [gr-qc].
- [44] G. G. L. Nashed, A. Abebe, and K. Bamba, *Eur. Phys. J. C* **80**, 1109 (2020).
- [45] G. G. L. Nashed and S. Capozziello, *Eur. Phys. J. C* **80**, 969 (2020), arXiv:2010.06355 [gr-qc].
- [46] A. Övgün, *Phys. Lett. B* **820**, 136517 (2021), arXiv:2105.05035 [gr-qc].
- [47] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [48] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Astrophys. J. Lett.* **892**, L3 (2020), arXiv:2001.01761 [astro-ph.HE].
- [49] B. P. Abbott *et al.* (KAGRA, LIGO Scientific, Virgo, VIRGO), *Living Rev. Rel.* **21**, 3 (2018), arXiv:1304.0670 [gr-qc].
- [50] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi GBM, INTEGRAL, IceCube, AstroSat Cadmium Zinc Telluride Imager Team, IPN, Insight-Hxmt, ANTARES, Swift, AGILE Team, 1M2H Team, Dark Energy Camera GW-EM, DES, DLT40, GRAWITA, Fermi-LAT, ATCA, ASKAP, Las Cumbres Observatory Group, OzGrav, DWF (Deeper Wider Faster Program), AST3, CAASTRO, VINROUGE, MASTER, J-GEM, GROWTH, JAGWAR, CaltechNRAO, TTU-NRAO, NuSTAR, Pan-STARRS, MAXI Team, TZAC Consortium, KU, Nordic Optical Telescope, ePESSTO, GROND, Texas Tech University, SALT Group, TOROS, BOOTES, MWA, CALET, IKI-GW Follow-up, H.E.S.S., LOFAR, LWA, HAWC, Pierre Auger, ALMA, Euro VLBI Team, Pi of Sky, Chandra Team at McGill University, DFN, ATLAS Telescopes, High Time Resolution Universe Survey, RIMAS, RATIR, SKA South Africa/MeerKAT), *Astrophys. J. Lett.* **848**, L12 (2017), arXiv:1710.05833 [astro-ph.HE].
- [51] B. P. Abbott *et al.* (LIGO Scientific, Virgo, Fermi-GBM, INTEGRAL), *Astrophys. J. Lett.* **848**, L13 (2017),

- arXiv:1710.05834 [astro-ph.HE].
- [52] A. Goldstein *et al.*, *Astrophys. J. Lett.* **848**, L14 (2017), arXiv:1710.05446 [astro-ph.HE].
- [53] A. Bauswein, O. Just, H.-T. Janka, and N. Stergioulas, *Astrophys. J. Lett.* **850**, L34 (2017), arXiv:1710.06843 [astro-ph.HE].
- [54] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Phys. Rev. Lett.* **121**, 161101 (2018), arXiv:1805.11581 [gr-qc].
- [55] C. D. Capano, I. Tews, S. M. Brown, B. Margalit, S. De, S. Kumar, D. A. Brown, B. Krishnan, and S. Reddy, *Nature Astron.* **4**, 625 (2020), arXiv:1908.10352 [astro-ph.HE].
- [56] V. B. Thapa, A. Kumar, and M. Sinha, *Mon. Not. Roy. Astron. Soc.* **507**, 2 (2021), arXiv:2108.04318 [astro-ph.HE].
- [57] T. Dietrich, M. W. Coughlin, P. T. H. Pang, M. Bulla, J. Heinzel, L. Issa, I. Tews, and S. Antier, *Science* **370**, 1450 (2020), arXiv:2002.11355 [astro-ph.HE].
- [58] M. Breschi, A. Perego, S. Bernuzzi, W. Del Pozzo, V. Nedora, D. Radice, and D. Vescovi, *Mon. Not. Roy. Astron. Soc.* **505**, 1661 (2021), arXiv:2101.01201 [astro-ph.HE].
- [59] K. Chatzioannou, *Gen. Rel. Grav.* **52**, 109 (2020), arXiv:2006.03168 [gr-qc].
- [60] W. Del Pozzo, T. G. F. Li, M. Agathos, C. Van Den Broeck, and S. Vitale, *Phys. Rev. Lett.* **111**, 071101 (2013), arXiv:1307.8338 [gr-qc].
- [61] K. Chatzioannou, K. Yagi, A. Klein, N. Cornish, and N. Yunes, *Phys. Rev. D* **92**, 104008 (2015), arXiv:1508.02062 [gr-qc].
- [62] B. D. Lackey and L. Wade, *Phys. Rev. D* **91**, 043002 (2015), arXiv:1410.8866 [gr-qc].
- [63] F. Hernandez Vivanco, R. Smith, E. Thrane, P. D. Lasky, C. Talbot, and V. Raymond, *Phys. Rev. D* **100**, 103009 (2019), arXiv:1909.02698 [gr-qc].
- [64] K. Chatzioannou and S. Han, *Phys. Rev. D* **101**, 044019 (2020), arXiv:1911.07091 [gr-qc].
- [65] B. P. Abbott *et al.* (LIGO Scientific, Virgo), *Astrophys. J. Lett.* **851**, L16 (2017), arXiv:1710.09320 [astro-ph.HE].
- [66] B. P. Abbott *et al.* (LIGO Scientific), *Class. Quant. Grav.* **34**, 044001 (2017), arXiv:1607.08697 [astro-ph.IM].
- [67] M. Maggiore *et al.*, *JCAP* **03**, 050 (2020), arXiv:1912.02622 [astro-ph.CO].
- [68] D. Ganapathy, L. McCuller, J. Graef Rollins, E. D. Hall, L. Barsotti, and M. Evans, *Phys. Rev. D* **103**, 022002 (2021), arXiv:2010.15735 [astro-ph.IM].
- [69] M. A. Page *et al.*, (2020), arXiv:2007.08766 [physics.optics].
- [70] F. A. Rasio and S. L. Shapiro, *Astrophys. J.* **401**, 226 (1992).
- [71] M. Shibata, *Phys. Rev. Lett.* **94**, 201101 (2005), arXiv:gr-qc/0504082.
- [72] E. Elizalde, G. G. L. Nashed, S. Nojiri, and S. D. Odintsov, *Eur. Phys. J. C* **80**, 109 (2020), arXiv:2001.11357 [gr-qc].
- [73] A. Bauswein and H. T. Janka, *Phys. Rev. Lett.* **108**, 011101 (2012), arXiv:1106.1616 [astro-ph.SR].
- [74] J. Clark, A. Bauswein, L. Cadonati, H. T. Janka, C. Pankow, and N. Stergioulas, *Phys. Rev. D* **90**, 062004 (2014), arXiv:1406.5444 [astro-ph.HE].
- [75] L. Rezzolla and K. Takami, *Phys. Rev. D* **93**, 124051 (2016), arXiv:1604.00246 [gr-qc].
- [76] A. Bauswein and N. Stergioulas, *J. Phys. G* **46**, 113002 (2019), arXiv:1901.06969 [gr-qc].
- [77] M. Breschi, S. Bernuzzi, F. Zappa, M. Agathos, A. Perego, D. Radice, and A. Nagar, *Phys. Rev. D* **100**, 104029 (2019), arXiv:1908.11418 [gr-qc].
- [78] K. W. Tsang, T. Dietrich, and C. Van Den Broeck, *Phys. Rev. D* **100**, 044047 (2019), arXiv:1907.02424 [gr-qc].
- [79] S. Vretinaris, N. Stergioulas, and A. Bauswein, *Phys. Rev. D* **101**, 084039 (2020), arXiv:1910.10856 [gr-qc].
- [80] P. J. Easter, S. Ghonge, P. D. Lasky, A. R. Casey, J. A. Clark, F. H. Vivanco, and K. Chatzioannou, *Phys. Rev. D* **102**, 043011 (2020), arXiv:2006.04396 [astro-ph.HE].
- [81] J. L. Friedman and N. Stergioulas, *Int. J. Mod. Phys. D* **29**, 2041015 (2020), arXiv:2005.14135 [astro-ph.HE].
- [82] B. Zwiebach, *Physics Letters B* **156**, 315 (1985).
- [83] D. J. Gross and J. H. Sloan, *Nuclear Physics B* **291**, 41 (1987).
- [84] D. Glavan and C. Lin, *Phys. Rev. Lett.* **124**, 081301 (2020), arXiv:1905.03601 [gr-qc].
- [85] R. B. Mann and S. F. Ross, *Class. Quant. Grav.* **10**, 1405 (1993), arXiv:gr-qc/9208004.
- [86] S. Nojiri and S. D. Odintsov, *EPL* **130**, 10004 (2020), arXiv:2004.01404 [hep-th].
- [87] T. Torii and H.-a. Shinkai, *Phys. Rev. D* **78**, 084037 (2008), arXiv:0810.1790 [gr-qc].
- [88] A. Mardones and J. Zanelli, *Classical and Quantum Gravity* **8**, 1545 (1991).
- [89] R. P. Woodard, *Scholarpedia* **10**, 32243 (2015), arXiv:1506.02210 [hep-th].
- [90] Y. Tomozawa, (2011), arXiv:1107.1424 [gr-qc].
- [91] K. Aoki, M. A. Gorji, S. Mizuno, and S. Mukohyama, *JCAP* **01**, 054 (2021), arXiv:2010.03973 [gr-qc].
- [92] J. Bonifacio, K. Hinterbichler, and L. A. Johnson, *Phys. Rev. D* **102**, 024029 (2020), arXiv:2004.10716 [hep-th].
- [93] W.-Y. Ai, *Commun. Theor. Phys.* **72**, 095402 (2020), arXiv:2004.02858 [gr-qc].
- [94] S. Mahapatra, *Eur. Phys. J. C* **80**, 992 (2020), arXiv:2004.09214 [gr-qc].
- [95] M. Gürses, T. c. Sışman, and B. Tekin, *Eur. Phys. J. C* **80**, 647 (2020), arXiv:2004.03390 [gr-qc].
- [96] M. Hohmann, C. Pfeifer, and N. Voicu, *The European Physical Journal Plus* **136**, 1 (2021).
- [97] L. Ma and H. Lu, *Eur. Phys. J. C* **80**, 1209 (2020), arXiv:2004.14738 [gr-qc].
- [98] H. Lu and Y. Pang, *Phys. Lett. B* **809**, 135717 (2020), arXiv:2003.11552 [gr-qc].
- [99] P. G. S. Fernandes, P. Carrilho, T. Clifton, and D. J. Mulryne, *Phys. Rev. D* **102**, 024025 (2020), arXiv:2004.08362 [gr-qc].
- [100] A. Banerjee, S. Hansraj, and L. Moodly, *Eur. Phys. J. C* **81**, 790 (2021).
- [101] S. G. Ghosh and R. Kumar, *Class. Quant. Grav.* **37**, 245008 (2020), arXiv:2003.12291 [gr-qc].
- [102] A. Kumar and S. G. Ghosh, (2020), arXiv:2004.01131 [gr-qc].

- [103] A. Kumar and R. Kumar, (2020), arXiv:2003.13104 [gr-qc].
- [104] C.-Y. Zhang, S.-J. Zhang, P.-C. Li, and M. Guo, JHEP **08**, 105 (2020), arXiv:2004.03141 [gr-qc].
- [105] S. A. Hosseini Mansoori, Phys. Dark Univ. **31**, 100776 (2021), arXiv:2003.13382 [gr-qc].
- [106] A. K. Mishra, Gen. Rel. Grav. **52**, 106 (2020), arXiv:2004.01243 [gr-qc].
- [107] C.-Y. Zhang, P.-C. Li, and M. Guo, Eur. Phys. J. C **80**, 874 (2020), arXiv:2003.13068 [hep-th].
- [108] Y.-P. Zhang, S.-W. Wei, and Y.-X. Liu, Universe **6**, 103 (2020), arXiv:2003.10960 [gr-qc].
- [109] K. Jusufi, A. Banerjee, and S. G. Ghosh, Eur. Phys. J. C **80**, 698 (2020), arXiv:2004.10750 [gr-qc].
- [110] B. Eslam Panah, K. Jafarzade, and S. H. Hendi, Nucl. Phys. B **961**, 115269 (2020), arXiv:2004.04058 [hep-th].
- [111] J. Antoniadis *et al.*, Science **340**, 6131 (2013), arXiv:1304.6875 [astro-ph.HE].
- [112] A. Bodmer, Physical Review D **4**, 1601 (1971).
- [113] E. Witten, Phys. Rev. D **30**, 272 (1984).
- [114] S. G. Ghosh and S. D. Maharaj, Phys. Dark Univ. **30**, 100687 (2020), arXiv:2003.09841 [gr-qc].
- [115] L. Herrera, Physics Letters A **188**, 402 (1994).
- [116] H. Abreu, H. Hernandez, and L. A. Nunez, Class. Quant. Grav. **24**, 4631 (2007), arXiv:0706.3452 [gr-qc].
- [117] K. Bamba, M. Ilyas, M. Z. Bhatti, and Z. Yousaf, Gen. Rel. Grav. **49**, 112 (2017), arXiv:1707.07386 [gr-qc].
- [118] S. Das, F. Rahaman, and L. Baskey, Eur. Phys. J. C **79**, 853 (2019).
- [119] R. C. Tolman, Phys. Rev. **55**, 364 (1939).
- [120] J. R. Oppenheimer and G. M. Volkoff, Phys. Rev. **55**, 374 (1939).
- [121] J. Ponce de Leon, General Relativity and Gravitation **25**, 1123 (1993).
- [122] C. C. Moustakidis, Gen. Rel. Grav. **49**, 68 (2017), arXiv:1612.01726 [gr-qc].
- [123] H. Heintzmann and W. Hillebrandt, aap **38**, 51 (1975).
- [124] R. Chan, L. Herrera, and N. O. Santos, Monthly Notices of the Royal Astronomical Society **265**, 533 (1993), <http://oup.prod.sis.lan/mnras/article-pdf/265/3/533/3807712/mnras265-0533.pdf>.
- [125] B. K. Harrison, K. S. Thorne, M. Wakano, and J. A. Wheeler, *Gravitation Theory and Gravitational Collapse* (1965).
- [126] Y. B. Zeldovich and I. D. Novikov, *Relativistic astrophysics. Vol.1: Stars and relativity* (1971).
- [127] I. B. Zeldovich and I. D. Novikov, *Relativistic astrophysics. Vol.2: The structure and evolution of the universe* (1983).
- [128] F. Özel, D. Psaltis, T. Güver, G. Baym, C. Heinke, and S. Guillot, apj **820**, 28 (2016), arXiv:1505.05155 [astro-ph.HE].
- [129] M. Z.-u.-H. Bhatti, M. Sharif, Z. Yousaf, and M. Ilyas, Int. J. Mod. Phys. D **27**, 1850044 (2017).

