# **GAMES ON BASE MATRICES**

#### VERA FISCHER, MARLENE KOELBING, AND WOLFGANG WOHOFSKY

ABSTRACT. Using a game characterization of distributivity, we show that base matrices for  $\mathcal{P}(\omega)$ /fin of regular height larger than  $\mathfrak{h}$  necessarily have maximal branches which are not cofinal.

### 1. INTRODUCTION

A forcing  $\mathbb{P}$  is  $\delta$ -distributive if any system of  $\delta$  many maximal antichains has a common refinement. The distributivity of a forcing notion  $\mathbb{P}$ , denoted by  $\mathfrak{h}(\mathbb{P})$ , is the least  $\lambda$  such that  $\mathbb{P}$  is not  $\lambda$ -distributive. In particular,  $\mathfrak{h}(\mathcal{P}(\omega)/\mathfrak{fin})$  is the classical cardinal characteristic  $\mathfrak{h}$ . Note that  $\mathfrak{h}(\mathbb{P})$  is actually the least  $\lambda$  such that there is a system of  $\lambda$  many *refining* maximal antichains without common refinement, which gives rise to the following definition:

**Definition 1.1.** We say that  $\mathcal{A} = \{A_{\xi} \mid \xi < \lambda\}$  is a *distributivity matrix for*  $\mathbb{P}$  *of height*  $\lambda$  if

- (1)  $A_{\xi}$  is a maximal antichain in  $\mathbb{P}$ , for each  $\xi < \lambda$ ,
- (2)  $A_{\eta}$  refines  $A_{\xi}$  whenever  $\eta \ge \xi$ , i.e., for each  $b \in A_{\eta}$  there exists  $a \in A_{\xi}$  such that  $b \le a$ , and
- (3) there is no common refinement, i.e., there is no maximal antichain *B* which refines every  $A_{\xi}$ .

A special sort of distributivity matrices have been considered in the seminal paper [BPS80] of Balcar, Pelant, and Simon, where h has been introduced:

**Definition 1.2.** A distributivity matrix  $\{A_{\xi} | \xi < \lambda\}$  for  $\mathbb{P}$  is a *base matrix* if  $\bigcup_{\xi < \lambda} A_{\xi}$  is dense in  $\mathbb{P}$ , i.e., for each  $p \in \mathbb{P}$  there is  $\xi < \lambda$  and  $a \in A_{\xi}$  such that  $a \le p$ .

In [BPS80], the famous base matrix theorem has been shown: there exists a base matrix for  $\mathcal{P}(\omega)/\text{fin}$  of height b. A more general version for a wider class of forcings has been given in [BDH15, Theorem 2.1].

Due to its refining structure, a distributivity matrix  $\{A_{\xi} | \xi < \lambda\}$  can be viewed as a tree, with level  $\xi$  being  $A_{\xi}$ . Let us say that  $\langle a_{\xi} | \xi < \delta \rangle$  is a *branch of the distributivity matrix*  $\{A_{\xi} | \xi < \lambda\}$  if  $a_{\xi} \in A_{\xi}$  for each  $\xi < \delta$ , and  $a_{\eta} \leq a_{\xi}$  for each

<sup>2010</sup> Mathematics Subject Classification. 03E05, 03E17.

*Key words and phrases.* base matrices; distributivity game; distributivity matrices; cardinal characteristics.

*Acknowledgments.* The authors would like to thank the Austrian Science Fund (FWF) for the generous support through grants Y1012, I4039 (Fischer, Wohofsky) and P28420 (Koelbing). The second author is also grateful for the support by the ÖAW Doc fellowship.

 $\xi \le \eta < \delta$ . We say that the branch is *maximal* if there is no branch of  $\{A_{\xi} | \xi < \lambda\}$  strictly extending it. If  $\delta = \lambda$ , the branch  $\langle a_{\xi} | \xi < \delta \rangle$  is called *cofinal* in  $\{A_{\xi} | \xi < \lambda\}$ .

The structure of base matrices for  $\mathcal{P}(\omega)/\text{fin}$  has been investigated in the literature. Note that each maximal branch of a distributivity matrix for  $\mathcal{P}(\omega)/\text{fin}$  which is not cofinal is a tower. So if there are no towers of length strictly less than b, i.e., if t = b, all maximal branches of a distributivity matrix of height b are cofinal. On the other hand, Dow showed that in the Mathias model, there exists a base matrix of height b without cofinal branches (see [Dow89, Lemma 2.17]). It is actually consistent that *no* base matrix of height b has cofinal branches. This was proved by Dordal by constructing a model in which b does not belong to the tower spectrum (see [Dor87] or<sup>1</sup> [Dor89, Corollary 2.6]).

In [FKW], the authors of this paper have shown that consistently there exists a distributivity matrix for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than  $\mathfrak{h}$  in which all maximal branches are cofinal.

In [Bre], Brendle has shown that if  $\lambda \leq c$  is regular and greater or equal than the splitting number  $\mathfrak{s}$  (or, alternatively, there exists no strictly  $\subseteq^*$ -decreasing sequence of length  $\lambda$ ), then there exists a base matrix for  $\mathcal{P}(\omega)/\text{fin of height } \lambda$ . In particular, there always exists a base matrix of height  $\mathfrak{c}$  provided that  $\mathfrak{c}$  is regular. He mentions that in the Cohen and random models base matrices of height larger than  $\mathfrak{h}$  necessarily have maximal branches which are not cofinal (in fact, there are no strictly  $\subseteq^*$ -decreasing sequences of length larger than  $\omega_1$ ).

We will show below that, in ZFC, any base matrix for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than h has maximal branches which are not cofinal.

# 2. MAIN RESULT

In the proof of Theorem 2.3, we will use a game characterization of being  $\lambda$ -distributive. It generalizes the game characterization for being  $\omega$ -distributive which can be found in [Jec03, Lemma 30.23].

**Definition 2.1.** Let  $\mathbb{P}$  be a forcing notion. Let  $G_{\lambda}(\mathbb{P})$  denote the  $\lambda$ -distributivity game (which has length  $\lambda$ ):

Ι	$a_0$		$a_1$		•••		$a_{\mu+1}$		•••	
II		$b_0$		$b_1$		 $b_{\mu}$		$b_{\mu+1}$		

The players alternately pick conditions in  $\mathbb{P}$  such that the resulting sequence is decreasing, i.e.,  $b_j \leq a_i$  and  $a_{i+1} \leq b_i$  for every  $i \leq j < \lambda$ . Player I starts the game, and at limits  $\mu$ , Player II has to play. If Player II cannot play at limits (because the sequence played till then has no lower bound), the game ends and Player I wins immediately. If the game continuous for  $\lambda$  many steps, Player II wins if and only if there exists a  $b \in \mathbb{P}$  with  $b \leq a_i$  for every successor  $i < \lambda$ .

<sup>&</sup>lt;sup>1</sup>Dordal's original model (in which  $c = \omega_2$ ) is presented in [Dor87], whereas [Dor89, Corollary 2.6] is a more general result which also gives models satisfying  $\mathfrak{h} = \mathfrak{c} > \omega_2$  (but is, interestingly enough, easier to prove).

Recall that, by definition, a forcing  $\mathbb{P}$  is  $\leq \lambda$ -strategically closed if Player II has a winning strategy in  $G_{\lambda}(\mathbb{P})$ . A slightly weaker property turns out to be equivalent to being  $\lambda$ -distributive; this was shown by Foreman in [For83, Theorem on page 718]:

### **Theorem 2.2.** *The following are equivalent:*

- (1) Player I has no winning strategy in the game  $G_{\lambda}(\mathbb{P})$ .
- (2)  $\mathbb{P}$  is  $\lambda$ -distributive.

We now show that a base matrix for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than h necessarily has (through every node) branches which are dying out (see Corollary 2.5). In fact, we show the following more general theorem:

**Theorem 2.3.** Let  $\mathcal{A} = \{A_{\xi} \mid \xi < \lambda\}$  be a base matrix for  $\mathbb{P}$  of regular height  $\lambda > \mathfrak{h}(\mathbb{P})$ . Then there is a maximal branch of  $\mathcal{A}$  which is not cofinal.

*Proof.* Assume towards contradiction that every maximal branch is cofinal. By definition,  $\mathbb{P}$  is *not*  $\mathfrak{h}(\mathbb{P})$ -distributive. Therefore, using the game characterization of distributivity (see Theorem 2.2), Player I has a winning strategy  $\sigma$  in the game  $G_{\mathfrak{h}(\mathbb{P})}(\mathbb{P})$ .

Consider the following run of the game of (full) length  $\mathfrak{h}(\mathbb{P})$ :

where Player I plays according to  $\sigma$  (i.e.,  $b_0 = \sigma(\langle \rangle)$  and  $b_{i+1} = \sigma(\langle b_0, a_0, \dots, a_i \rangle)$ for each  $i < \mathfrak{h}(\mathbb{P})$ ), and Player II plays as follows (where the  $a_i$  are going to be in the matrix for each  $i < \mathfrak{h}(\mathbb{P})$ ). For successors i (and for i = 0), let  $a_i \le b_i$  with  $a_i$ in the matrix; this is possible, because it is a base matrix. For limit  $\mu \le \mathfrak{h}(\mathbb{P})$ , the following holds by induction:  $\langle a_i | i < \mu \rangle$  is a  $\le$ -decreasing sequence such that  $a_i$  is in the matrix for each  $i < \mu$ . So (for  $\mu < \mathfrak{h}(\mathbb{P})$ ) Player II can play a lower bound  $a_{\mu}$ in the matrix, by the following claim.

## **Claim.** The sequence $\langle a_i | i < \mu \rangle$ has a lower bound in the matrix.

*Proof.* We can assume that the sequence is not eventually constant. Moreover, we can assume that it is strictly decreasing. It is easy to check that there is a strictly increasing sequence  $\langle \xi_i \mid i < \mu \rangle \subseteq \lambda$  with  $a_i \in A_{\xi_i}$  for each  $i < \mu$ . Then  $\sup(\{\xi_i \mid i < \mu\}) < \lambda$ , because  $\mu \le \mathfrak{h}(\mathbb{P}) < \lambda$  and  $\lambda$  is regular. So the corresponding branch is not cofinal in the matrix, hence it is not maximal by assumption. Consequently, there exists an *a* in the matrix such that  $a \le a_i$  for each  $i < \mu$ .

Finally, for  $\mu = \mathfrak{h}(\mathbb{P})$ , the claim yields a lower bound of  $\langle a_i | i < \mathfrak{h}(\mathbb{P}) \rangle$ , witnessing that Player II wins this run of the game. This contradicts that  $\sigma$  is a winning strategy for Player I.

**Remark 2.4.** In the above theorem, the assumption that  $\lambda > \mathfrak{h}(\mathbb{P})$  can be replaced by the weaker assumption that  $\mathbb{P}$  is, for some  $\nu < \lambda$ , not  $\leq \nu$ -strategically closed. In the proof, one can turn (using a well-order on the base matrix) the description of the moves of Player II into a strategy for Player II, which is then a winning strategy.

Also, note that the proof still works for singular  $\lambda$  as long as  $cf(\lambda) > h(\mathbb{P})$  (or if  $\mathbb{P}$  is not  $\leq v$ -strategically closed for some  $v < cf(\lambda)$ ).

For the important case of  $\mathcal{P}(\omega)/\text{fin}$ , we can now derive the following:

**Corollary 2.5.** Let  $\mathcal{A} = \{A_{\xi} \mid \xi < \lambda\}$  be a base matrix for  $\mathcal{P}(\omega)/fin$  of regular height  $\lambda > \mathfrak{h}$ . Then for every  $a \in \bigcup_{\xi < \lambda} A_{\xi}$ , there is a maximal branch of  $\mathcal{A}$  containing a which is not cofinal.

*Proof.* Fix *a* in the matrix (i.e.,  $a \in \bigcup_{\xi < \lambda} A_{\xi}$ ). Let  $\mathbb{P} := \{b \mid b \subseteq^* a\}$  be the part of  $\mathcal{P}(\omega)/\text{fin}$  below *a*. Recall that  $\mathcal{P}(\omega)/\text{fin}$  is homogenous, hence  $\mathfrak{h}(\mathbb{P}) = \mathfrak{h}(\mathcal{P}(\omega)/\text{fin}) = \mathfrak{h}$ . Note that the part of  $\mathcal{A}$  below *a* is a base matrix for  $\mathbb{P}$  of height  $\lambda$ . Therefore, by Theorem 2.3, there is a maximal branch of  $\mathcal{A}$  containing *a* which is not cofinal.

So the above theorem actually says that distributivity matrices for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than h cannot simultaneously have only cofinal maximal branches and be a base matrix. Therefore, Brendle's theorem from [Bre] together with the above theorem shows that there are distributivity matrices for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than h with maximal branches which are not cofinal (at least if c > h is regular or s < c). On the other hand, the above theorem shows that the generic distributivity matrix of regular height larger than b from [FKW] cannot be a base matrix because all its maximal branches are cofinal (this can also be seen by analyzing the forcing construction, see the end of [FKW, Section 7.1]).

Further note that in the model from [FKW], there are both kinds of distributivity matrices for  $\mathcal{P}(\omega)/\text{fin}$  of regular height larger than  $\mathfrak{h}$ : matrices all whose maximal branches are cofinal, and matrices with maximal branches which are not cofinal.

#### References

- [BDH15] Bohuslav Balcar, Michal Doucha, and Michael Hrušák. Base tree property. *Order*, 32(1):69–81, 2015.
- [BPS80] Bohuslav Balcar, Jan Pelant, and Petr Simon. The space of ultrafilters on N covered by nowhere dense sets. *Fund. Math.*, 110(1):11–24, 1980.
- [Bre] Jörg Brendle. Base matrices of various heights. http://arxiv.org/abs/2202.00897.
- [Dor87] Peter Lars Dordal. A model in which the base-matrix tree cannot have cofinal branches. J. Symbolic Logic, 52(3):651–664, 1987.
- [Dor89] Peter Lars Dordal. Towers in  $[\omega]^{\omega}$  and  ${}^{\omega}\omega$ . Ann. Pure Appl. Logic, 45(3):247–276, 1989.
- [Dow89] Alan Dow. Tree  $\pi$ -bases for  $\beta$ N N in various models. *Topology Appl.*, 33(1):3–19, 1989.
- [FKW] Vera Fischer, Marlene Koelbing, and Wolfgang Wohofsky. On heights of distributivity matrices. *Submitted*. http://arxiv.org/abs/2202.09255.
- [For83] Matthew Foreman. Games played on Boolean algebras. *J. Symbolic Logic*, 48(3):714–723, 1983.
- [Jec03] Thomas Jech. *Set theory*. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2003. The third millennium edition, revised and expanded.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF VIENNA, KOLINGASSE 14–16, 1090 WIEN, AUSTRIA *Email address*: vera.fischer@univie.ac.at

INSTITUTE OF MATHEMATICS, UNIVERSITY OF VIENNA, KOLINGASSE 14–16, 1090 WIEN, AUSTRIA *Email address*: marlenekoelbing@web.de

INSTITUTE OF MATHEMATICS, UNIVERSITY OF VIENNA, KOLINGASSE 14–16, 1090 WIEN, AUSTRIA *Email address*: wolfgang.wohofsky@gmx.at