

# Estimating the circumference of a graph in terms of its leaf number

Jingru Yan \*

Department of Mathematics, East China Normal University, Shanghai 200241, China

## Abstract

Let  $\mathcal{T}$  be the set of spanning trees of  $G$  and let  $L(T)$  be the number of leaves in a tree  $T$ . The leaf number  $L(G)$  of  $G$  is defined as  $L(G) = \max\{L(T)|T \in \mathcal{T}\}$ . Let  $G$  be a connected graph of order  $n$  and minimum degree  $\delta$  such that  $L(G) \leq 2\delta - 1$ . We show that the circumference of  $G$  is at least  $n - 1$ , and that if  $G$  is regular then  $G$  is hamiltonian.

**Keywords.** Leaf number, circumference, hamiltonian

**Mathematics Subject Classification.** 05C38, 05C45

## 1 Introduction

We will deal with only finite nontrivial simple graphs. Let  $G$  be a graph with vertex set  $V(G)$  and edge set  $E(G)$ . The order and size of a graph  $G$  are its number of vertices and edges, respectively. The notations  $N_G(v)$  and  $N_G[v]$  denote the neighborhood and closed neighborhood of  $v \in V(G)$ , respectively. The degree of  $v$  is  $d_G(v) = |N_G(v)|$ .  $\delta(G)$  and  $\Delta(G)$  denote the minimum and maximum degree of a graph  $G$ , respectively. If the graph  $G$  is clear from the context, we will omit it as subscript. For terminology and notations not explicitly described in this paper, the reader is referred the books [1, 19].

Let  $\mathcal{T}$  be the set of spanning trees of  $G$ .  $L(T)$  denotes the number of leaves in a tree  $T$ , where a leaf means a vertex of degree 1. Then the leaf number  $L(G) = \max\{L(T)|T \in \mathcal{T}\}$ . Many researchers have estimated the circumference of graphs by various invariants. The purpose of this paper is to estimate the circumference of a connected graph  $G$  by the two invariants  $\delta(G)$  and  $L(G)$ .

DeLaViña's computer program, Graffiti.pc, posed attractive conjectures [3] and some of the conjectures speculate sufficient conditions for traceability based on the minimum degree and leaf number. In 2013, Mukwembi gave a partial solution to the Graffiti.pc 190a. He [14] showed that if  $G$  is a finite connected graph with minimum degree  $\delta(G) \geq 5$ , and leaf

---

\*E-mail address: mathyjr@163.com

number  $L(G)$  such that  $\delta(G) \geq L(G) - 1$ , then  $G$  is hamiltonian and thus traceable. In the same year, he [16] relaxed the condition  $\delta(G) \geq 5$  to  $\delta(G) \geq 3$ . After that, Mukwembi [15] proved that if  $G$  is a connected claw-free graph with  $\delta(G) \geq (L(G) + 1)/2$ , then  $G$  is hamiltonian. In recent years, several authors reported on sufficient conditions for a graph to be hamiltonian or traceable based on minimum degree and leaf number, see [9–13].

We state the following results, some of which will be used later in this paper.

**Theorem 1.** [12] *If  $G$  is a connected graph with  $\delta(G) \geq (L(G)+2)/2$ , then  $G$  is hamiltonian.*

**Theorem 2.** [11] *If  $G$  is a connected graph with  $\delta(G) \geq (L(G) + 1)/2$ , then  $G$  is traceable.*

**Theorem 3.** [13] *Let  $G$  be a connected triangle-free graph with  $L(G) \leq 2\delta(G) - 1$ . Then  $G$  is either hamiltonian or  $G \in \mathcal{F}_2$ , where  $\mathcal{F}_2$  is the class of non-hamiltonian graphs with leaf number  $2\delta(G) - 1$ .*

Let  $p(G)$  and  $c(G)$  be the order of a longest path and a longest cycle in a graph  $G$ , respectively. Note that  $c(G)$  is equal to the circumference of a graph  $G$ . Many researchers have investigated the relation between  $p(G)$  and  $c(G)$  ([5],[8],[17],[18]). Motivated by Theorem 2, we obtain the following main result.

**Theorem 4.** *Let  $G$  be a connected graph of order  $n$ . If  $L(G) \leq 2\delta(G) - 1$ , then  $c(G) \geq n - 1$ . The bound is sharp and the condition cannot be relaxed.*

We also consider regular graphs.

## 2 Main results

We start with some lemmas that will be used repeatedly.

**Lemma 5.** [15] *Let  $G$  be a connected graph of order  $n$ . If  $L(G) \leq 2\delta(G) - 1$ , then  $n \leq \max\{2\delta(G) + 6, 3\delta(G)\}$ .*

**Lemma 6.** [15] *Let  $G$  be a connected graph with  $L(G) \leq 2\delta(G) - 1$ . Then  $G$  is 2-connected.*

For a graph  $G$ ,  $\kappa(G)$  and  $\alpha(G)$  denote the connectivity and independent number of  $G$ , respectively. Let  $\sigma_k(G)$  be the minimum degree sum of  $k$  independent vertices of  $G$  if  $\alpha(G) \geq k$ .  $K_n$  stands for the complete graph of order  $n$ .

**Lemma 7.** [2] Let  $G$  be a connected graph. If  $\kappa(G) \geq \alpha(G)$ , then  $G$  is hamiltonian except for  $G = K_2$ .

**Lemma 8.** [5] Let  $G$  be a 2-connected graph of order  $n$ . If  $\sigma_3(G) \geq n + 2$ , then  $c(G) \geq p(G) - 1$ .

Now we first show that the result of Theorem 4 is true when  $n \leq 3\delta(G)$ .

**Lemma 9.** [16] Let  $G$  be a connected graph of order  $n$ . If  $\delta(G) = 2$  and  $L(G) \leq 3$ , then  $c(G) \geq n - 1$ .

Given graphs  $G$  and  $H$ , the notation  $G + H$  means the *disjoint union* of  $G$  and  $H$ . Then  $tG$  denotes the disjoint union of  $t$  copies of  $G$ . The notation  $G \vee H$  means the *joint* of  $G$  and  $H$ . For graphs we will use equality up to isomorphism, so  $G = H$  means that  $G$  and  $H$  are isomorphic.

For any graph  $G$ ,  $G[S]$  denotes the subgraph of  $G$  induced by  $S \subseteq V(G)$ . Let  $A, B \subseteq V(G)$  and  $A \cap B = \emptyset$ . Denote by  $E(A, B)$  the set of edges of  $G$  with one end in  $A$  and the other end in  $B$  and  $e(A, B) = |E(A, B)|$ .

**Lemma 10.** [5] Let  $G$  be a connected graph with order  $n \geq 3$ . If  $\sigma_3(G) \geq n$ , then  $G$  satisfies  $c(G) \geq p(G) - 1$  or  $G \in \mathcal{F}(n)$ , where  $\mathcal{F}(n)$  is the class of graphs defined below.

$\mathcal{F}(n)$  consists six subclasses:

$$\mathcal{F}(n) = \mathcal{F}_1(n) \cup \mathcal{F}_2(n) \cup \mathcal{F}_3(n) \cup \mathcal{F}_4(n) \cup \mathcal{F}_5(n) \cup \mathcal{F}_6(n).$$

For any graph  $G \in \mathcal{F}(n)$ , we have  $|V(G)| = n$  and  $\sigma_3(G) \geq n$ . The subclasses are defined as follows (more details can be found in [5]):

$\mathcal{F}_1(n)$ :  $G \in \mathcal{F}_1(n)$  if  $V(G) = A \cup B$  with  $A \cap B = \emptyset$ ,  $G[A]$  and  $G[B]$  are hamiltonian or isomorphic to  $K_2$ , and  $e(A, B) = 1$ .

$\mathcal{F}_2(n)$ :  $G \in \mathcal{F}_2(n)$  if  $V(G) = A \cup B$  with  $A \cap B = \{x\}$ ,  $G[A]$  and  $G[B]$  are both hamiltonian or both isomorphic to  $K_2$ , and  $e(A \setminus \{x\}, B \setminus \{x\}) = 0$ .

$\mathcal{F}_3(n)$ :  $G \in \mathcal{F}_3(n)$  if  $G$  is a 2-connected spanning subgraph of  $K_2 \vee (K_a + K_b + K_c)$  with  $a, b, c \geq 2$  ( $n = a + b + c + 2$ ).

$\mathcal{F}_4(n)$ :  $G \in \mathcal{F}_4(n)$  if  $G$  is a 2-connected spanning subgraph of  $K_3 \vee (aK_2 + bK_3)$  with  $a, b \geq 0$  and  $a + b = 4$  ( $n = 2a + 3b + 3$ ,  $11 \leq n \leq 15$ ).

$\mathcal{F}_5(n)$ :  $G \in \mathcal{F}_5(n)$  if  $G$  is a 2-connected spanning subgraph of  $K_s \vee (sK_2 + K_3)$  with  $s \geq 4$  ( $n = 3s + 3$ ).

$\mathcal{F}_6(n)$ :  $G \in \mathcal{F}_6(n)$  if  $G$  is a 2-connected spanning subgraph of  $K_s \vee (s + 1)K_2$  with  $s \geq 4$  ( $n = 3s + 2$ ).

**Theorem 11.** *Let  $G$  be a connected graph with order  $n \leq 3\delta(G)$ . If  $L(G) \leq 2\delta(G) - 1$ , then  $c(G) \geq n - 1$ .*

*Proof.* Let  $G$  be a connected graph with order  $n \leq 3\delta(G)$  and  $L(G) \leq 2\delta(G) - 1$ . By Lemma 6,  $G$  is 2-connected. If  $\alpha(G) \leq 2$ , by Lemma 7, then  $G$  is hamiltonian and hence  $c(G) \geq n - 1$ . Clearly,  $\delta(G) \neq 1$ . By Lemma 9, the result holds true for  $\delta(G) = 2$ . Now, it suffices to consider the case of  $\alpha(G) \geq 3$  and  $\delta(G) = \delta \geq 3$ . Note that  $G$  is a connected graph with order  $n > 3$ , by Theorem 2 and Lemma 10,  $c(G) \geq p(G) - 1 = n - 1$  or  $G \in \mathcal{F}(n)$ . Suppose to the contrary that  $G \in \mathcal{F}(n)$ .

Recall that  $G$  is 2-connected. This implies that  $G \notin \mathcal{F}_1(n) \cup \mathcal{F}_2(n)$ . First suppose  $G \in \mathcal{F}_3(n)$ . For any vertex  $x$  of  $V(K_a)$ ,  $d_{K_a}(x) \geq \delta - 2$  in  $G$  and hence  $|V(K_a)| \geq \delta - 1$ . Similarly,  $|V(K_b)| \geq \delta - 1$  and  $|V(K_c)| \geq \delta - 1$ . Then

$$|V(K_a)| + |V(K_b)| + |V(K_c)| + 2 \geq 3(\delta - 1) + 2 = 3\delta - 1.$$

It implies that either  $n = 3\delta - 1$  or  $n = 3\delta$ . For the first case,  $|V(K_a)| = |V(K_b)| = |V(K_c)| = \delta - 1$  and hence  $G = K_2 \vee (K_a + K_b + K_c)$ . It can easily be shown that  $G$  contains a spanning tree with leaf number at least  $2\delta$ , a contradiction. For the second case, exactly one of  $|V(K_a)|$ ,  $|V(K_b)|$  and  $|V(K_c)|$  is equal to  $\delta$ , and the rest are equal to  $\delta - 1$ . Without loss of generality, suppose that  $|V(K_a)| = \delta$  and  $|V(K_b)| = |V(K_c)| = \delta - 1$ . Then  $G[G - V(K_a)] = K_2 \vee (K_b + K_c)$ . The subgraph induced by the vertex set of  $G[G - V(K_a)]$  with one vertex of  $V(K_a)$  has a spanning tree with leaf number  $2\delta$ , contradicting  $L(G) \leq 2\delta - 1$ . Thus  $G \notin \mathcal{F}_3(n)$ .

Next assume that  $G \in \mathcal{F}_4(n)$ . For  $n \leq 3\delta - 2$ , by Lemmas 6 and 8,  $c(G) \geq p(G) - 1$  since  $n \leq \sigma_3(G) - 2$ . For  $3\delta - 1 \leq n \leq 3\delta$ ,  $\delta = 4$  or  $5$  since  $11 \leq n \leq 15$ . Note that  $a + b = 4$  and  $n = a + b + c + 2$ . If  $\delta = 4$ ,  $n = 11, a = 4, b = 0$  or  $n = 12, a = 3, b = 1$ . It is easy to check that  $L(G) \geq 8 > 2\delta - 1$  in both cases. If  $\delta = 5$ ,  $n = 14, a = 1, b = 3$  or  $n = 15, a = 0, b = 4$ . Since  $\delta = 5$ , the first case is not allowed. For  $n = 15, a = 0, b = 4$ , we have  $L(G) \geq 10 > 2\delta - 1$ . Thus  $G \notin \mathcal{F}_4(n)$ .

Now assume that  $G \in \mathcal{F}_5(n)$ . For any vertex  $x$  of  $V(sK_2)$ ,  $d_{K_s}(x) \geq \delta - 1$  in  $G$  and hence

$|V(K_s)| \geq \delta - 1$ . Then  $n = 3s + 3 \geq 3(\delta - 1) + 3 = 3\delta$ . Since  $n \leq 3\delta$ , then  $s = \delta - 1$ . It implies that  $G[V(K_s) \cup V(sK_2)]$  contains  $(\delta - 1)K_1 \vee (\delta - 1)K_2$  as a subgraph. Note that the subgraph induced by  $V(K_3)$  contains no isolated vertex in  $G$ . Then we can split this into two cases. For  $G[V(K_3)] = K_3$ , it is easy to check that  $G$  contains a spanning tree with leaf number at least  $2\delta$ , a contradiction. For  $G[V(K_3)] = P_3$ , let  $w_1, w_2 \in V(G[V(K_3)])$  and  $d_{G[V(K_3)]}(w_1) = d_{G[V(K_3)]}(w_2) = 1$ . Then  $d_{K_s}(w_1) = d_{K_s}(w_2) = \delta - 1$ . We also obtain  $G$  has a spanning tree with leaf number at least  $2\delta$ , contradicting  $L(G) \leq 2\delta - 1$ . Thus  $G \notin \mathcal{F}_5(n)$ .

It follows that  $G \in \mathcal{F}_6(n)$ . Since  $n = 3s + 2 \leq 3\delta$ , we have  $s \leq \delta - 1$ . For any vertex  $x$  of  $V((s + 1)K_2)$ ,  $d_{K_s}(x) \geq \delta - 1$  in  $G$  since  $d(x) \geq \delta$ . Then  $s = \delta - 1$  and  $n = 3s + 2 = 3\delta - 1$ . Further,  $G$  contains  $K_1 \vee \delta K_2$  as a subgraph. Thus,  $G$  has a spanning tree with leaf number at least  $2\delta$ , a contradiction. This completes the proof of Theorem 11.  $\square$

Before giving the proof of the main theorem, we prove a conclusion about regular graphs.

**Lemma 12.** [7] *Every 2-connected  $k$ -regular ( $k \geq 3$ ) graph of order at most  $3k + 3$  is hamiltonian except the Petersen graph  $P$  and the graph obtained from  $P$  by replacing one vertex of  $P$  by a triangle.*

Denote by  $P^\Delta$  be the graph obtained from  $P$  by replacing one vertex of  $P$  by a triangle.




Fig. 1. The Petersen graph  $P$




Fig. 2. The graph  $P^\Delta$

**Theorem 13.** *Let  $G$  be a  $k$ -regular connected graph. If  $L(G) \leq 2k - 1$ , then  $G$  is hamiltonian and the condition cannot be relaxed.*

*Proof.* It is easy to verify that  $L(P) = 6$  and  $L(P^\Delta) = 7$  (see Fig 1 and Fig 2). The Petersen graph  $P$  is non-hamiltonian but satisfies  $L(P) = 6 = 2k$ , so the condition cannot be relaxed.

Let  $G$  be a  $k$ -regular connected graph of order  $n$  with  $L(G) \leq 2k - 1$ . Since  $L(G) \geq 2$ , then  $k \geq 2$ . Clearly,  $G$  is hamiltonian when  $k = 2$ . Next suppose that  $k \geq 3$ . By Lemma 5,

we have  $n \leq \max\{2k+6, 3k\}$ . Note that  $3k+3 \geq \max\{2k+6, 3k\}$  when  $k \geq 3$ . By Lemmas 6 and 12,  $G$  is hamiltonian. This completes the proof of Theorem 13.  $\square$

The following lemmas play the key role in the proof of Theorem 4.

**Lemma 14.** *Let  $G$  be a connected graph with  $L(G) \leq 2\delta(G) - 1$ . For  $\delta(G) \geq 3$ , if there is one vertex  $x \in V(G)$  with degree  $2\delta(G) - 1$ , then  $|V(G) \setminus N[x]| \leq 2$ .*

*Proof.* Let  $G$  be a connected graph with  $L(G) \leq 2\delta(G) - 1$ . Since  $L(G) \leq 2\delta(G) - 1$ , each vertex of  $N(x)$  has at most one neighbour in  $V(G) \setminus N[x]$ . By Lemma 6,  $G$  is 2-connected. Then there are two vertices  $y_1, y_2 \in V(G) \setminus N[x]$  have neighbors in  $N(x)$ . Similarly, each vertex of  $\{y_1, y_2\}$  has at most one neighbour in  $V(G) \setminus N[x]$  and hence at least  $\delta(G) - 1$  neighbors in  $N(x)$ . Suppose that  $|V(G) \setminus N[x]| \geq 3$ . There exists one vertex  $y_3 \in V(G) \setminus (N[x] \cup \{y_1, y_2\})$  and  $y_3$  is adjacent to  $y_1$  or  $y_2$ . Clearly,  $N(y_3) \cap N(x) = \emptyset$ . Without loss of generality, assume that  $y_3$  is adjacent to  $y_1$ .  $G[N[x] \cup \{y_1\} \cup N(y_3)]$  contains a tree with leaf number  $3\delta(G) - 3$ . Further, since  $\delta(G) \geq 3$ , we have  $3\delta(G) - 3 > 2\delta(G) - 1$ , a contradiction. This completes the proof of Lemma 14.  $\square$

**Lemma 15.** [4] *Let  $G$  be a 2-connected graph of order  $n$  and let  $C$  be a longest cycle in  $G$ . Then  $|V(C)| \geq \min\{n, 2\delta(G)\}$ .*

**Lemma 16.** [6] *Let  $G$  be a connected graph of order  $n$ .*

(1) *If  $\delta(G) \geq 4$ , then  $L(G) \geq \frac{2n+8}{5}$ .*

(2) *If  $\delta(G) \geq 5$ , then  $L(G) \geq \frac{n}{2} + 2$ .*

**Lemma 17.** *Let  $G$  be a connected graph of order  $n$  and let  $C = c_1, c_2, \dots, c_k, c_1$  be a longest cycle in  $G$ . The subscripts of the vertices  $c_i$  are taken modulo  $k$ .*

(1) *The vertices  $c_i$  and  $c_{i+1}$  have no common neighbor in  $V(G) \setminus V(C)$ .*

(2) *Let  $x, y \in V(G) \setminus V(C)$ . If  $c_i, c_j \in N_C(x)$ , then  $c_{i+1}$  and  $c_{j+1}$  cannot both belong to  $N(y)$ .*

(3) *Let  $P_C = p_1, p_2, \dots, p_s$  be a longest path in  $G - V(C)$ . If the vertices  $p_1$  and  $p_s$  have distinct neighbors in  $V(C)$ , then  $s \leq \lfloor \frac{k}{2} \rfloor - 1$ .*

*Proof.* It is easy to show that the results of Lemma 17, so we omit them.  $\square$

Finally, we show that the proof of Theorem 4.

*Proof.* Let  $G$  be a connected graph of order  $n$ . For  $n \leq 3\delta(G)$ , by Theorem 11,  $c(G) \geq n - 1$ . For  $n \geq 3\delta(G) + 1$ , by Lemma 5, we have  $\delta(G) \leq 5$ . Clearly,  $\delta(G) \neq 1$ . By Lemma 9, the

result is true when  $\delta(G) = 2$ . Denote by  $\delta(G) = \delta$ . Now, it suffices to consider the case of  $3\delta + 1 \leq n \leq 2\delta + 6$  and  $3 \leq \delta \leq 5$ .

*Case 1.*  $\delta = 3$

Note that  $10 \leq n \leq 12$ . Since  $L(G) \leq 2\delta - 1 = 5$ , we have  $3 \leq \Delta(G) \leq 5$ . If  $\Delta(G) = 3$ , by Theorem 13,  $G$  is hamiltonian and hence  $c(G) \geq n - 1$ . If  $\Delta(G) = 5$ , by Lemma 14,  $n \leq 6 + 2 = 8 < 10$ , a contradiction. Next suppose that  $\Delta(G) = 4$ . We discuss it in two Subcases according to the order of  $G$ .

*Subcase 1.1.* Consider  $n = 10$ . Let  $C = c_1, c_2, \dots, c_k, c_1$  be a longest cycle in  $G$  and let  $P_C$  be a longest path in  $G - V(C)$ . By Lemmas 6 and 15,  $k \geq 6$ . Now we show that  $k \geq 9$ . Suppose to the contrary that  $6 \leq k \leq 8$ .

For  $k = 6$ , by Lemma 17 (3),  $|V(P_C)| \leq 2$ . Recall that  $\delta = 3$  and  $\Delta(G) = 4$ , by Lemma 17 (1) and (2), we obtain at most two isolated vertices in  $G - V(C)$ . Hence  $|V(P_C)| = 2$  and  $G[V(G) \setminus V(C)] = 2K_1 + K_2$  or  $2K_2$ . Let  $x, y \in V(G) \setminus V(C)$  and  $x$  is adjacent to  $y$ . Since  $k = 6$ , we have  $d(x) = d(y) = 3$  and  $N(x) \setminus \{y\} = N(y) \setminus \{x\}$ . Clearly,  $N(x) \setminus \{y\} = \{c_1, c_4\}$  or  $\{c_2, c_5\}$  or  $\{c_3, c_6\}$ . It is not difficult to see that the proof methods for the above three cases are similar. So let us just consider the first case. Note that  $|V(G) \setminus V(C)| = 4$  and  $\Delta(G) = 4$ . There is one vertex  $z \in V(G) \setminus V(C)$  is adjacent to at least one of  $\{c_2, c_3, c_5, c_6\}$ . Suppose  $z$  is adjacent to  $c_2$ . The subgraph induced by  $\{c_6, c_1, x, y, c_2, c_3, z\}$  contains a tree with leaf number 5. By Lemma 14,  $n \leq 7 + 2 = 9$ , a contradiction. The remaining cases can be proved in the same way.

For  $k = 7$ , by Lemma 17 (3),  $|V(P_C)| \leq 2$ . If  $|V(P_C)| = 2$ , then  $G[V(G) \setminus V(C)] = K_1 + K_2$ . Let  $x, y, z \in V(G) \setminus V(C)$  and  $x$  is adjacent to  $y$ . It is easy to check that  $d(x) = d(y) = 3$  and  $N(x) \setminus \{y\} = N(y) \setminus \{x\}$ . Without loss of generality, suppose that  $N(x) \setminus \{y\} = \{c_1, c_4\}$ . Recall that  $\delta = 3$  and  $\Delta(G) = 4$ . Then  $z$  is adjacent to at least three vertices in  $V(C) \setminus \{c_1, c_4\}$ . If  $z$  is adjacent to  $c_2$ , the subgraph induced by  $\{c_6, c_1, x, y, c_2, c_3, z\}$  contains a tree with leaf number 5. By Lemma 14,  $n \leq 7 + 2 = 9$ , a contradiction. Using a similar argument as above, we deduce that  $z$  is not adjacent to  $c_3, c_5$  and  $c_6$ , contradicting  $d(z) \geq 3$ . Next suppose  $|V(P_C)| = 1$ . Then  $G[V(G) \setminus V(C)] = 3K_1$ . Let  $x, y, z \in V(G) \setminus V(C)$ . By Lemma 17 (1) and (2),  $d(x) = d(y) = d(z) = 3$  and  $N(x) = N(y) = N(z)$ . Then there is one vertex of  $N(x)$  has degree at least 5, a contradiction.

For  $k = 8$ ,  $|V(P_C)| \leq 2$  since  $n = 10$ . If  $|V(P_C)| = 2$ , then  $G[V(G) \setminus V(C)] = K_2$ . Let  $x, y \in V(G) \setminus V(C)$ . Without loss of generality, suppose  $x$  is adjacent to  $c_1$ . Since  $C$  is a

longest cycle in  $G$ , then  $y$  is adjacent to  $c_4$ ,  $c_5$  or  $c_6$ . Obviously, we can split this into two cases. The first case is where  $y$  is adjacent to  $c_4$ . One can easily show that  $d(x) = d(y) = 3$  and  $N(x) \setminus \{y\} = N(y) \setminus \{x\} = \{c_1, c_4\}$ . Consider the vertex  $c_2$ . If  $c_2$  is not adjacent to  $c_8$ , the subgraph induced by  $N(c_1) \cup N(c_2)$  contains a tree with leaf number 5 since  $d(c_2) \geq 3$ . By Lemma 14,  $n \leq 7 + 2 = 9 < 10$ , a contradiction. If  $c_2$  is adjacent to  $c_8$ ,  $G$  contains a cycle  $c_2, c_8, c_7, c_6, c_5, c_4, y, x, c_1, c_2$  with length 9 (see Fig. 3), contradicting to  $k = 8$ . The second case is where  $y$  is adjacent to  $c_5$ . Similarly, we have  $d(x) = d(y) = 3$  and  $N(x) \setminus \{y\} = N(y) \setminus \{x\} = \{c_1, c_5\}$ . The following results which are derived from the above proof:  $c_2$  is adjacent to  $c_8$  and  $c_4$  is adjacent to  $c_6$ . Then  $G$  contains a cycle  $c_2, c_8, c_7, c_6, c_4, c_5, y, x, c_1, c_2$  with length 9, a contradiction. Next suppose  $|V(P_C)| = 1$  and




Fig. 3. The subgraph of  $G$  when  $c_2$  is adjacent to  $c_8$




Fig. 4. The subgraph of  $G$  when  $y$  is adjacent to  $c_1$

hence  $G[V(G) \setminus V(C)] = 2K_1$ . Let  $x, y \in V(G) \setminus V(C)$ . Since  $C$  is a longest cycle in  $G$  and  $L(G) \leq 5$ , then  $d(x) = d(y) = 3$ . Without loss of generality, suppose that  $N(x) = \{c_1, c_3, c_5\}$  or  $\{c_1, c_3, c_6\}$ . For the first case, assert that  $N(y) \cap N(x) = \emptyset$ . Otherwise,  $G$  contains a tree with leaf number at least 6 if  $y$  is adjacent to  $c_1$  or  $c_5$  (see Fig.4), a contradiction. And if  $y$  is adjacent to  $c_3$ , the subgraph induced by the vertex set  $\{c_1, c_2, c_3, c_4, c_5, x, y\}$  contains a tree with leaf number 5. Then, by Lemma 14,  $n \leq 9$ , a contradiction. So,  $N(y) \subseteq V(C) \setminus N(x)$  and  $|N(y)| = 3$ , by Lemma 17 (1) and (2), which is not allowed. For the second case, the proof method is similar to the first case, and will not be repeated here.

*Subcase 1.2.* Consider  $n = 11$  or  $12$ . Let  $x \in V(G)$  with  $d(x) = 4$ . Set  $N(x) = \{x_1, x_2, x_3, x_4\}$ . Assert that any vertex of  $N(x)$  has at most one neighbor in  $V(G) \setminus N[x]$ . Since  $L(G) \leq 5$ , we have  $d_{G-N[x]}(x_i) \leq 2$  for  $i \in \{1, 2, 3, 4\}$ . If there is one vertex of  $N(x)$  has exactly two neighbors in  $V(G) \setminus N[x]$ , by Lemma 14,  $n \leq 5 + 2 + 2 = 9 < 11$ , a contradiction. Hence,  $e(N(x), V(G) \setminus N[x]) \leq 4$ . Let  $N_2(x) \subseteq V(G) \setminus N[x]$  and each vertex of  $N_2(x)$  has neighbor in  $N(x)$ . Similarly, by Lemma 14, we can show that each vertex of  $N_2(x)$  has at most one neighbour in  $V(G) \setminus N[x]$ . Then each vertex of  $N_2(x)$  has at least two neighbours in



$N(x)$ , since  $\delta = 3$ . By Lemma 6,  $G$  is 2-connected and hence  $|N_2(x)| \geq 2$ . Then  $|N_2(x)| = 2$  and  $e(N(x), V(G) \setminus N[x]) = 4$ . Set  $N_2(x) = \{y_1, y_2\}$ . Without loss of generality, suppose that  $N(y_1) \cap N(x) = \{x_1, x_2\}$  and  $N(y_2) \cap N(x) = \{x_3, x_4\}$ . It is easy to check that  $y_1$  is not adjacent to  $y_2$ , since  $n \geq 11$ . Let  $z_1 = N(y_1) \setminus \{x_1, x_2\}$  and  $z_2 = N(y_2) \setminus \{x_3, x_4\}$ . Since  $G$  is 2-connected, then  $z_1 \neq z_2$ . Note that  $G[N(x)]$  contains  $2K_2$ . Then  $G$  contains a path of length 8 with endpoints  $z_1$  and  $z_2$ . For  $n = 11$ , it remains two vertices  $w_1$  and  $w_2$ . Obviously,  $d(w_1) = d(w_2) = 3$  and  $N(w_1) = \{w_2, z_1, z_2\}$ ,  $N(w_2) = \{w_1, z_1, z_2\}$ . Thus  $c(G) = n$ . For  $n = 12$ , it remains three vertices  $w_1, w_2$  and  $w_3$ . One can easy show that  $c(G) \geq n - 1$ . So Case 1 is proven.

*Case 2.*  $\delta = 4$

Note that  $13 \leq n \leq 14$ . For  $n = 14$ , by Lemma 16 (1),  $L(G) \geq \frac{2n+8}{5} = \frac{36}{5} > 7$ , contradicting  $L(G) \leq 2\delta - 1 = 7$ . Then we only need to consider  $n = 13$ . Suppose  $n = 13$ . Since  $L(G) \leq 7$ , we have  $4 \leq \Delta(G) \leq 7$ . By Lemma 14,  $\Delta(G) \neq 7$ . For  $\Delta(G) = 6$ , let  $x \in V(G)$  with  $d(x) = 6$ . Then any vertex of  $N(x)$  has at most two neighbors in  $V(G) \setminus N[x]$ . Let  $N_2(x) \subseteq V(G) \setminus N[x]$  and each vertex of  $N_2(x)$  has neighbor in  $N(x)$ . By Lemma 14, any vertex of  $N(x) \cup N_2(x)$  has at most one neighbor in  $V(G) \setminus N[x]$ . This implies that  $e(N(x), N_2(x)) \leq 6$ . Since  $\delta = 4$ , then  $|N_2(x)| \leq 2$ . By Lemma 6,  $G$  is 2-connected and hence  $|N_2(x)| = 2$ . Clearly,  $y_1$  is not adjacent to  $y_2$  since  $n = 13$ . Let  $z_1 \in N(y_1)$  and  $z_2 \in N(y_2)$ . Recall that  $G$  is 2-connected.  $z_1 \neq z_2$ . Hence  $|N(z_1) \setminus \{y_1\}| \geq 3$ . From Fig.5, we obtain  $G$  contains a tree with leaf number 8, a contradiction. For  $\Delta(G) = 4$ , by Theorem 13,  $c(G) = n \geq n - 1$ .




Fig. 5. The subgraph of  $G$  in Case 2 when  $\Delta(G)=6$

It remains the case of  $\Delta(G) = 5$ . Let  $C = c_1, c_2, \dots, c_k, c_1$  be a longest cycle in  $G$  and let  $P_C$  be a longest path in  $G - V(C)$ . By Lemmas 6 and 15,  $k \geq 8$ . Now we show that  $k \geq 12$ . Suppose to the contrary that  $8 \leq k \leq 11$ .

For  $k = 8$ , by Lemma 17 (3),  $|V(P_C)| \leq 3$ . Since  $\Delta(G) = 5$ , by Lemma 17 (1) and (2), there are at most three isolated vertices in  $G - V(C)$ . Then  $2 \leq |V(P_C)| \leq 3$ . If  $|V(P_C)| = 2$ , then  $G - V(C) = 3K_1 + K_2$  or  $K_1 + 2K_2$ . Let  $x, y \in V(G) \setminus V(C)$  and  $x$  is adjacent to  $y$ . Without loss of generality, suppose that  $c_1 \in N(x)$ . By Lemma 17 (1) and (2),  $N(y) \subseteq \{x, c_1, c_4, c_5, c_6\}$ . Then  $N(y) = \{x, c_1, c_4, c_6\}$ , since  $C$  is a longest cycle and  $\delta = 4$ . Further, we have  $N(x) = \{c_1, y\}$ , contradicting  $d(x) \geq 4$ . Next suppose  $|V(P_C)| = 3$ . Let  $P_C = x, y, z$ . It follows that  $N_C(x) = N_C(z)$  and  $|N_C(x)| = |N_C(z)| = 2$ . Recall that  $C$  is a longest cycle in  $G$ . Without loss of generality, suppose that  $N_C(x) = N_C(z) = \{c_1, c_5\}$ . Since  $\delta(G) = 4$  and  $|V(P_C)| = 3$ , we have  $x$  is adjacent to  $z$ . Note that we have a new path  $P'_C = y, x, z$  in  $G - V(C)$ . Similarly,  $N_C(y) = N_C(z)$  and  $|N_C(y)| = |N_C(z)| = 2$ . So,  $N_C(x) = N_C(y) = N_C(z) = \{c_1, c_5\}$  and  $d(x) = d(y) = d(z) = 4$ . Then  $G[V(G) \setminus V(C)] = 2K_1 + K_3$ . Let  $\{u, v\} = V(G) \setminus (V(C) \cup \{x, y, z\})$ . Clearly,  $N(u) = N(v) = \{c_2, c_4, c_6, c_8\}$ . The subgraph induced by  $\{c_3, c_1, c_2, c_3, x, y, z, u, v\}$  contains a tree with leaf number 7 (see Fig.6). By Lemma 14,  $n \leq 9 + 2 = 11 < 13$ , a contradiction.




Fig. 6. The subgraph of  $G$  in Case 2 when  $k=8$  and  $|V(P_C)|=3$

For  $k = 9$ , by Lemma 17,  $|V(P_C)| \leq 3$  and there are at most two isolated vertices in  $G - V(C)$ . Then  $2 \leq |V(P_C)| \leq 3$ . If  $|V(P_C)| = 2$ , then  $G[V(G) \setminus V(C)] = 2K_1 + K_2$  or  $2K_2$ . Let  $x, y \in V(G) \setminus V(C)$  and  $x$  is adjacent to  $y$ . One can easy show that  $d(x) = d(y) = 4$  and  $N(x) \cap N(y) = \{c_1, c_4, c_7\}$  or  $\{c_2, c_5, c_8\}$  or  $\{c_3, c_6, c_9\}$ . Without loss of generality, suppose that  $N(x) \cap N(y) = \{c_1, c_4, c_7\}$ . Since  $L(G) \leq 7$ , any vertex of  $V(G) \setminus (V(C) \cup \{x, y\})$  has no neighbor in  $\{c_1, c_4, c_7\}$ . Further, by Lemma 17 (1) and (2),  $G[V(G) \setminus V(C)] \neq 2K_1 + K_2$  and  $G[V(G) \setminus V(C)] \neq 2K_2$ , since  $\delta = 4$ . Next suppose  $|V(P_C)| = 3$ . Let  $P_C = x, y, z$ . Using the same method as the case of  $k = 8$  and  $|V(P_C)| = 3$ , we obtain  $G[V(G) \setminus V(C)] = K_1 + K_3$  and  $N_C(x) = N_C(y) = N_C(z)$ . Without loss of generality, suppose that  $N_C(x) = N_C(y) = N_C(z) = \{c_1, c_5\}$ . Consider the vertex  $c_2$ . If  $c_2$  is adjacent to  $c_9$ ,  $G$  contains a cycle  $c_1, c_2, c_9, c_8, c_7, c_6, c_5, z, y, x, c_1$  with length 10, a contradiction. If  $c_2$  is

not adjacent to  $c_9$ , the subgraph induced by  $N(c_1) \cup N(c_2)$  contains a tree with leaf number 7. By Lemma 14,  $n \leq 10 + 2 = 12 < 13$ , a contradiction.

For  $k = 10$ ,  $|V(P_C)| \leq 3$ . For  $|V(P_C)| = 3$ , let  $V(G) \setminus V(C) = \{x, y, z\}$ . Similarly, one can easily show that  $G[V(G) \setminus V(C)] = K_3$  and  $N_C(x) = N_C(y) = N_C(z)$  and  $|N_C(x)| = |N_C(y)| = |N_C(z)| = 2$ . Without loss of generality, suppose that  $N_C(x) = \{c_1, c_5\}$  or  $\{c_1, c_6\}$ . If  $N_C(x) = \{c_1, c_5\}$ , we consider the vertex  $c_3$ . Note that  $d(c_1) = d(c_5) = 5$ . Then  $|N(c_3) \cap (V(C) \setminus \{c_1, c_2, c_4, c_5\})| \geq 2$ , since  $\delta = 4$ . If  $c_3$  is adjacent to  $c_6$  (see Fig.7a) or  $c_{10}$ , then  $G$  contains a cycle  $c_5, c_4, c_3, c_6, c_7, c_8, c_9, c_{10}, c_1, x, y, z, c_5$  with length 12 or a cycle  $c_1, c_2, c_3, c_{10}, c_9, c_8, c_7, c_6, c_5, z, y, x, c_1$  with length 12, a contradiction. If  $c_3$  is adjacent to  $c_7$  (see Fig.7b) or  $c_9$ , then  $G$  contains a cycle  $c_5, c_4, c_3, c_7, c_8, c_9, c_{10}, c_1, x, y, z, c_5$  with length 11 or a cycle  $c_1, c_2, c_3, c_9, c_8, c_7, c_6, c_5, z, y, x, c_1$  with length 11, a contradiction. Hence we have  $N(c_3) \subseteq \{c_2, c_4, c_8\}$ , contradicting  $\delta = 4$ . If  $N_C(x) = \{c_1, c_6\}$ , using the same method, we




Fig. 7a. The subgraph of  $G$  when  $k=10$  and  $c_3$  is adjacent to  $c_6$




Fig. 7b. The subgraph of  $G$  when  $k=10$  and  $c_3$  is adjacent to  $c_7$

have  $c_7, c_{10} \notin N(c_3)$  and  $c_7, c_{10} \notin N(c_4)$ . Now we show that  $c_3$  and  $c_4$  are both adjacent to exactly one of  $\{c_8, c_9\}$ . Suppose to the contrary that  $c_3$  and  $c_4$  are adjacent to  $c_8$  and  $c_9$  or  $c_9$  and  $c_8$ , respectively. Then  $G$  contains a cycle  $c_1, c_2, c_3, c_8, c_9, c_4, c_5, c_6, z, y, x, c_1$  with length 11 or a cycle  $c_1, c_2, c_3, c_9, c_8, c_4, c_5, c_6, z, y, x, c_1$  with length 11, contradicting  $k = 10$ . Without loss of generality, suppose  $c_9 \in N(c_3) \cap N(c_4)$ . Since  $\delta = 4$ ,  $c_4$  is adjacent to  $c_2$ . Then  $G$  contains a cycle  $c_1, c_{10}, c_9, c_3, c_2, c_4, c_5, c_6, z, y, x, c_1$  with length 11, a contradiction. For  $|V(P_C)| \leq 2$ , it implies that there exists at least one isolated vertex in  $G - V(C)$ . Assert that any isolated vertex of  $G - V(C)$  has degree 4. Otherwise, suppose  $d_{G-V(C)}(x) = 0$  and  $d(x) = 5$ . Then  $N(x) = \{c_1, c_3, c_5, c_7, c_9\}$  or  $\{c_2, c_4, c_6, c_8, c_{10}\}$ . We show that the first case, the second can be proved by same method. Note the subgraph induced by  $\{c_9, c_{10}, c_1, c_2, c_3, c_4, c_5, c_6, c_7, x\}$  contains a tree with leaf number 7. By Lemma 14,  $n \leq 10 + 2 = 12 < 13$ , a contradiction. Let  $V(G) \setminus V(C) = \{x, y, z\}$  and  $d_{G-V(C)}(x) = 0$ . Then, by Lemma 14,  $N(y) \cap N(x) = \emptyset$  and  $N(z) \cap N(x) = \emptyset$ . By Lemma 17 (1) and (2),  $|V(P_C)| > 1$  since  $\delta = 4$ . Hence  $|V(P_C)| = 2$

and  $y$  is adjacent to  $z$ . Since  $d(x) = 4$ , then the neighbors of  $x$  divide  $C$  into four parts. The lengths of the four parts of  $C$  are 2,2,2,4 or 2,2,3,3 or 2,3,2,3 (see Fig.8). By Lemma 17 (1) and (2), it is easy to show that in each case there is a contradiction, so we omit it.




Fig. 8. The subgraph of  $G$  when  $k=10$  and  $|V(P_C)|=2$

For  $k = 11$ ,  $|V(P_C)| \leq 2$ . Let  $x, y \in V(G) \setminus V(P)$ . Since  $n = 13$ , by Lemma 14,  $d(x) = d(y) = 4$ . For  $|V(P_C)| = 2$ ,  $G[V(G) \setminus V(C)] = K_2$ . Note that the neighbors of  $x$  in  $V(C)$  divide  $C$  into three parts. Then we have five cases and the lengths of the three parts of  $C$  are 2,2,7 or 2,3,6 or 2,4,5 or 3,3,5 or 3,4,4. The proof methods for the first three cases are similar. Since  $C$  is a longest cycle in  $G$ , one can easy show that  $d(y) \leq 3 < 4$ , a contradiction. The proofs for the latter two cases are similar, so we only give the proof for one of them here. Without loss of generality, suppose  $N(x) = \{c_1, c_4, c_7\}$ . Consider the vertex  $c_3$ . We assert that  $N(c_3) \cap N(x) = \{c_4\}$ . Otherwise, if  $c_3$  is adjacent to  $c_1$ , the subgraph induced by  $\{c_{11}, c_1, c_2, c_3, x, y, c_4, c_6, c_7, c_8\}$  contains a tree with leaf number 7 and hence by Lemma 14,  $n \leq 12 < 13$ , a contradiction. Similarly, we can show that  $c_3$  is not adjacent to  $c_7$ . If  $c_3$  is adjacent to  $c_5$ ,  $G$  contains a cycle  $c_4, c_3, c_5, c_6, c_7, c_8, c_9, c_{10}, c_{11}, c_1, x, y, c_4$  of length 12, contradicting  $k = 11$ . Hence  $N(c_3) \cap \{x, y, c_1, c_5, c_7\} = \emptyset$ . Note that the subgraph induced by  $N[c_3] \cup \{x, y, c_1, c_5, c_7\}$  contains a tree with leaf number 7, since  $d(c_3) \geq 4$ . Then, by Lemma 14,  $n \leq 12 < 13$ , a contradiction. For  $|V(P_C)| = 1$ ,  $G[V(G) \setminus V(C)] = 2K_1$ . Recall that  $d(x) = d(y) = 4$ . We assert that  $N(x) \cap N(y) = \emptyset$ . Otherwise, by Lemma 14, we will have  $n < 13$ . Note that the neighbors of  $x$  in  $V(C)$  divide  $C$  into four parts and hence we have four cases. The lengths of each parts of  $C$  are 2,2,2,5 or 2,2,3,4 or 2,3,2,4 or 2,3,3,3. Similarly, by Lemma 17 (1) and (2), one can easy show that in each case there is a contradiction. Then Case 2 is proven.

*Case 3.*  $\delta = 5$

Note that  $n = 16$ . By Lemma 16 (2),  $L(G) \geq \frac{n}{2} + 2 = 10$ , contradicting  $L(G) \leq 2\delta - 1$ . So Case 3 is proven.

For the sharpness, consider the following graph. The graph  $G_1$  of order  $n$  is formed by taking the cycle  $C_{n-1} = v_1, v_2, \dots, v_{n-2}, v_{n-1}, v_1$  and add one vertex  $v_n$  together with edges  $v_1v_n, v_3v_n$ . Note that  $\delta(G_1) = 2$  and  $L(G_1) = 3$ . Then  $G_1$  satisfying  $L(G_1) \leq 2\delta(G_1) - 1$  and  $c(G_1) = n - 1$ .

The condition  $L(G) \leq 2\delta(G) - 1$  cannot be relaxed. The graph  $G_2$  with order  $n \geq 8$  is formed by taking the cycle  $C_{n-2} = v_1, v_2, \dots, v_{n-2}, v_1$  and add two vertices  $v_{n-1}$  and  $v_n$  together with edges  $v_1v_{n-1}, v_3v_{n-1}, v_{n-5}v_n, v_{n-3}v_n$ . Clearly,  $\delta(G_2) = 2$  and  $L(G_2) = 4$ . Then  $G_2$  satisfying  $L(G_2) \leq 2\delta(G_2)$  but  $c(G_2) = n - 2$ . This completes the proof of Theorem 4.  $\square$

**Acknowledgement** This research was supported Science and Technology Commission of Shanghai Municipality (STCSM) grant 18dz2271000.

## References

- [1] J.A. Bondy, U.S.R. Murty, Graph Theory, in: GTM, vol. 244, Springer, 2008.
- [2] V. Chvátal, P. Erdős, A note on Hamiltonian circuits. *Discrete Math.* 2 (1972), 111–113.
- [3] E. DeLaViña, Written on the Wall II (Conjectures of Graffiti.pc). <http://cms.dt.uh.edu/faculty/delavinae/research/wowII/>.
- [4] G. A. Dirac, Some theorems on abstract graphs, *Proc. London Math. Soc.* (3) 2 (1952), 69–81.
- [5] H. Enomoto, J. van den Heuvel, A. Kaneko, A. Saito, Relative length of long paths and cycles in graphs with large degree sums, *J. Graph Theory* 20 (1995), no. 2, 213–225.
- [6] J.R. Griggs, M. Wu, Spanning trees in graphs of minimum degree 4 or 5. *Discrete Math.* 104 (1992), no. 2, 167–183.
- [7] F. Hilbig, Kantenstrukturen in Nichthamiltonschen Graphen (PHD Thesis). Technische Universität (1986).
- [8] S.C. Locke, Relative lengths of paths and cycles in  $k$ -connected graphs, *J. Combin. Theory Ser. B* 32 (1982), no. 2, 206–222.

- [9] P. Mafuta, Leaf number and Hamiltonian  $C_4$ -free graphs, *Afr. Mat.* 28 (2017), no. 7-8, 1067–1074.
- [10] P. Mafuta, S. Mukwembi, On minimum degree, leaf number, traceability and Hamiltonicity in graphs, *Discrete Appl. Math.* 221 (2017), 89–94.
- [11] P. Mafuta, S. Mukwembi, S. Munyira, Spanning paths in graphs, *Discrete Appl. Math.* 255 (2019), 278–282.
- [12] P. Mafuta, S. Mukwembi, S. Munyira, T. Vetrík, Hamiltonicity, minimum degree and leaf number, *Acta Math. Hungar.* 152 (2017), no. 1, 217–226.
- [13] P. Mafuta, J. Mushanyu, Spanning paths and cycles in triangle-free graphs, *Quaest. Math.* 43 (2020), no. 12, 1737–1747.
- [14] S. Mukwembi, Minimum degree, leaf number and Hamiltonicity, *Amer. Math. Monthly* 120 (2) (2013) 115.
- [15] S. Mukwembi, Minimum degree, leaf number and traceability, *Czechoslovak Math. J.* 63(138) (2013), no. 2, 539–545.
- [16] S. Mukwembi, On spanning cycles, paths and trees, *Discrete Appl. Math.* 161 (2013), no. 13-14, 2217–2222.
- [17] K. Ozeki, T. Yamashita, Length of longest cycles in a graph whose relative length is at least two, *Graphs Combin.* 28 (2012), no. 6, 859–868.
- [18] D. Paulusma, K. Yoshimoto, Relative length of longest paths and longest cycles in triangle-free graphs, *Discrete Math.* 308 (2008), no. 7, 1222–1229.
- [19] D.B. West, *Introduction to Graph Theory*, Prentice Hall, Inc., Upper Saddle River, NJ, (1996).