

# On the Maximum $F_5$ -free Subhypergraphs of a Random Hypergraph

Igor Araujo\*

József Balogh†

Haoran Luo‡

## Abstract

Denote by  $F_5$  the 3-uniform hypergraph on vertex set  $\{1, 2, 3, 4, 5\}$  with hyperedges  $\{123, 124, 345\}$ . Balogh, Butterfield, Hu, and Lenz proved that if  $p > K \log n/n$  for some large constant  $K$ , then every maximum  $F_5$ -free subhypergraph of  $G^3(n, p)$  is tripartite with high probability, and showed if  $p_0 = 0.1\sqrt{\log n}/n$ , then with high probability there exists a maximum  $F_5$ -free subhypergraph of  $G^3(n, p_0)$  that is not tripartite. In this paper, we sharpen the upper bound to be best possible up to a constant factor. We prove that when  $p > C\sqrt{\log n}/n$  for some large constant  $C$ , then every maximum  $F_5$ -free subhypergraph of  $G^3(n, p)$  is tripartite with high probability.

## 1 Introduction

The first theorem in extremal graph theory is Mantel's Theorem [10], which states that every triangle-free graph on  $n$  vertices has at most  $\lfloor n^2/4 \rfloor$  edges. Additionally, the complete bipartite graph whose part sizes differ by at most one is the unique maximum triangle-free graph. Later, Turán [14] generalized Mantel's Theorem for complete graphs. Denote by  $K_s$  the complete graph on  $s$  vertices and by  $T_s(n)$  the complete  $s$ -partite graph on  $n$  vertices where the sizes of the parts differ by at most 1. Turán's Theorem states that  $T_{s-1}(n)$  is the unique maximum  $K_s$ -free graph on  $n$  vertices. Turán's Theorem can also be understood as a property of  $K_n$ . Namely, it confirms that every maximum  $K_s$ -free subgraph of  $K_n$  is  $(s-1)$ -partite.

Let  $G(n, p)$  be the standard binomial model of random graphs, where each edge in  $K_n$  is chosen independently with probability  $p$ . We say an event occurs *with high probability*

---

\*Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. Email: [igoraa2@illinois.edu](mailto:igoraa2@illinois.edu).

†Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. E-mail: [jbal@illinois.edu](mailto:jbal@illinois.edu). Research is partially supported by NSF Grant DMS-1764123 and RTG DMS-1937241, Arnold O. Beckman Research Award (UIUC Campus Research Board RB 22000), the Langan Scholar Fund (UIUC), and the Simons Fellowship.

‡Department of Mathematics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA. E-mail: [haoranl8@illinois.edu](mailto:haoranl8@illinois.edu).

(w.h.p.) if its probability goes to 1 as  $n$  goes to infinity. A question related to Turán's Theorem arises when  $G(n, p)$  replaces the role of  $K_n$ , that is for what  $p = p(n)$  we have that every maximum  $K_s$ -free subgraph of  $G(n, p)$  is  $(s - 1)$ -partite with high probability. This question was first considered by Babai, Simonovits and Spencer [1], who studied the case when  $p = \frac{1}{2}$  and  $s = 3$ .

DeMarco and Kahn [6] determined the correct order of  $p$  for  $s = 3$ . They [6] showed that if  $p > K\sqrt{\log n/n}$  for some large constant  $K$ , then every maximum triangle-free subgraph of  $G(n, p)$  is w.h.p. bipartite, while if  $p = 0.1\sqrt{\log n/n}$ , then this does not hold w.h.p., where here and throughout the paper, the logarithms are in natural base. Finally, DeMarco and Kahn [7] answered this question for every  $s \geq 3$ .

Figure 1: The hypergraph  $F_5$ .





Figure 2: The hypergraph  $K_4^-$ .



Similar problems can also be considered for hypergraphs. Denote by  $K_4^-$  the hypergraph obtained from the complete 3-uniform hypergraph on 4 vertices by removing one hyperedge. Let  $F_5$ , which sometimes is called the *generalized triangle*, be the hypergraph on vertex set  $\{1, 2, 3, 4, 5\}$  with hyperedges  $\{123, 124, 345\}$ . Denote by  $S(n)$  the complete 3-partite 3-uniform hypergraph on  $n$  vertices whose parts have sizes  $\lfloor n/3 \rfloor$ ,  $\lfloor (n+1)/3 \rfloor$ , and  $\lfloor (n+2)/3 \rfloor$ , and let  $s(n) = \lfloor n/3 \rfloor \cdot \lfloor (n+1)/3 \rfloor \cdot \lfloor (n+2)/3 \rfloor$  be the number of hyperedges in  $S(n)$ . Bollobás [4] proved that  $S(n)$  is the unique maximum  $\{K_4^-, F_5\}$ -free  $n$ -vertex hypergraph. Frankl and Füredi [8] proved that, for  $n \geq 3000$ , the maximum number of hyperedges in an  $n$ -vertex  $F_5$ -free 3-uniform hypergraph is  $s(n)$ . Later, this result was improved by Keevash and Mubayi [9], who proved that if  $n \geq 33$ , the unique maximum  $n$ -vertex  $F_5$ -free 3-uniform hypergraph is  $S(n)$ .

The random version of this theorem was obtained by Balogh, Butterfield, Hu and Lenz [2]. Let  $G^3(n, p)$  be the random 3-uniform hypergraph on vertex set  $[n] := \{1, \dots, n\}$ , where each triple in  $\binom{[n]}{3}$  is included in  $G^3(n, p)$  with probability  $p$  independently of each other. Note that when  $p$  is very small, then  $G^3(n, p)$  itself is tripartite and hence  $F_5$ -free with high probability. Therefore, the interesting case is when  $p$  is large enough. In [2], it was proved that when  $p > K \log n/n$  for some large constant  $K$ , every maximum  $F_5$ -free subhypergraph of  $G^3(n, p)$  is w.h.p. tripartite, and it was conjectured that it should be enough to require

$p > C\sqrt{\log n}/n$  for some large constant  $C$ . In this paper, we verify this conjecture. Note that this is best possible up to the constant factor, as it was also shown in [2] that when  $p = 0.1\sqrt{\log n}/n$ , then w.h.p. there is a maximum  $F_5$ -free subhypergraph of  $G^3(n, p)$  that is not tripartite.

**Theorem 1.** *There exists a constant  $C > 0$  such that, for every  $p > C\sqrt{\log n}/n$ , w.h.p. every maximum  $F_5$ -free subhypergraph of  $G^3(n, p)$  is tripartite.*

Our approach will follow the general structure of [2]. Several key lemmas are improved and adapted for this smaller  $p$ . We trust that the new ideas used in our proof could be useful for solving other problems when one of the tasks in the proof is to beat the union bound. In Section 2, we will introduce the notation and lemmas needed. In Section 3, we give the proof of Theorem 1.

## 2 Preliminaries

To improve readability, as it is standard in the literature, we will usually pretend that large numbers are integers to avoid using essentially irrelevant floor and ceiling symbols. For a set  $S$ , we use  $\binom{S}{k}$  to stand for the collection of all the subsets of  $S$  with size  $k$ . We will often use the standard upper bound  $\binom{n}{k} \leq (\frac{en}{k})^k$  for binomial coefficients. We write  $x = (1 \pm c)y$  for  $(1 - c)y \leq x \leq (1 + c)y$ . We use  $G$  for  $G^3(n, p)$  hereinafter and denote by  $t(G)$  the number of hyperedges in a maximum tripartite subhypergraph of  $G$ . Let  $C$  be a large enough constant such that all assumptions in the lemmas in this section are satisfied and the proof in Section 3 works.

We will always assume that the hypergraphs are on vertex set  $[n] = \{1, \dots, n\}$ , so we can identify a hypergraph  $H$  by its hyperedges and  $|H|$  stands for the number of hyperedges of  $H$ . We use  $\pi = (V_1, V_2, V_3)$  to stand for a 3-partition of  $[n]$ . We say a 3-partition  $\pi$  is *balanced* if every part has size  $(1 \pm 10^{-10})n/3$ . Denote by  $K_\pi$  the set of triples with exactly one vertex in each part of  $\pi$ . Let  $G_\pi := G \cap K_\pi$ . For a hypergraph  $H \subseteq G$ , let  $H_i := \{e \in H : |e \cap V_i| \geq 2\}$  for  $i = 1, 2, 3$ . Let  $H_\pi := H \cap K_\pi$  and  $\bar{H}_\pi := G_\pi \setminus H_\pi$ . We will call hyperedges in  $H_\pi$  the *crossing hyperedges* of  $H$ , and the hyperedges in  $\bar{H}_\pi$  the *missing crossing hyperedges* of  $H$ .

For a hypergraph  $H$ , a partition  $\pi = (V_1, V_2, V_3)$ ,  $v, v' \in [n]$ , and  $S, T \subseteq [n]$ , let  $N_{S,T}^H(v) := \{\{y, z\} : y \in S, z \in T, vyz \in H\}$  be the *link graph* in  $H$  of  $v$  between  $S$  and  $T$ ,  $d_{S,T}^H(v) := |N_{S,T}^H(v)|$  be the *degree* in  $H$  of  $v$  between  $S$  and  $T$ ,  $N_S^H(v, v') := \{z : z \in S, vv'z \in H\}$  be the *neighbors* of  $v$  and  $v'$  in  $S$ , and  $d_S^H(v, v') := |N_S^H(v, v')|$  be the *codegree* of  $v$  and  $v'$  in  $S$ . Let  $L_{S,T}^H(v, v') := N_{S,T}^H(v) \cap N_{S,T}^H(v')$  be the *common neighborhood* of  $v$  and  $v'$  between  $S$  and  $T$  in  $H$ . When  $S$  or  $T$  is  $[n]$  or  $H = G$ , we omit to write  $S, T$  or  $G$  when there is no ambiguity. When  $S$  or  $T$  is  $V_i$ , we often just use  $i$  in the subscript. For example,  $d(x)$  is just the degree of  $x$  in  $G$  and  $N_1^H(y, z)$  is the neighbors of  $y$  and  $z$  in  $V_1$  in  $H$ . We will often use  $xy$  to stand for  $\{x, y\}$  and  $xyz$  to stand for  $\{x, y, z\}$ . Finally, define

$$Q(\pi) := \{xy \subset V_1 : |L_{2,3}(x, y)| < 0.8p^2n^2/9\}.$$

The first proposition we need from [2] is the following stability result, which is a special case of a general transference result of Conlon and Gowers [5], and as Samotij observed [11] of Schacht [13]. It was also proved by the hypergraph container method [3, 12].

**Proposition 2.** *For every  $\delta > 0$ , there exist  $\varepsilon > 0$  and  $C > 0$  such that if  $p > C/n$ , the follow statement is true. Let  $H$  be a maximum  $F_5$ -free subhypergraph of  $G$  and  $\pi$  be a 3-partition maximizing  $|H_\pi|$ . Then  $|H| \geq (2/9 - \varepsilon)p\binom{n}{3}$ ,  $\pi$  is balanced and  $|H \setminus H_\pi| \leq \delta pn^3$ .*

The following concentration results are also in [2]. Lemma 3 is the standard Chernoff's bound. Lemma 4 and Lemma 5 are standard properties of random hypergraphs.

**Lemma 3.** *Let  $Y$  be the sum of mutually independent indicator random variables, and let  $\mu = \mathbb{E}[Y]$ . For every  $\varepsilon > 0$ , we have*

$$\mathbb{P}[|Y - \mu| > \varepsilon \mu] < 2e^{-c_\varepsilon \mu},$$

where  $c_\varepsilon = \min \{-\log(e^\varepsilon(1 + \varepsilon)^{-(1+\varepsilon)}), \varepsilon^2/2\}$ .

**Lemma 4.** *For every  $\varepsilon > 0$ , there exists a positive constant  $C$  such that if  $p > C \log n/n^2$ , then w.h.p. for every vertex  $v$ , we have  $d(v) = (1 \pm \varepsilon)pn^2/2$ .*

**Lemma 5.** *For every  $\varepsilon > 0$ , there exists a positive constant  $C$  such that if  $p > C/n$ , then w.h.p. for every 3-partition  $\pi = (V_1, V_2, V_3)$  with  $|V_2|, |V_3| \geq n/20$  and every vertex  $v \in V_1$ , we have  $d_{2,3}(v) = (1 \pm \varepsilon)p|V_2||V_3|$ .*

We also have the following concentration results.

**Lemma 6.** *There exists a positive constant  $C$  such that if  $p > C\sqrt{\log n}/n$ , then w.h.p. for every pair of vertices  $x, y$ , we have  $d(x, y) \leq pn\sqrt{\log n}/\log \log n$ .*

*Proof.* For every pair of vertices  $xy$ , the probability that  $d(x, y) \geq \frac{pn\sqrt{\log n}}{\log \log n}$  is at most  $\left(\frac{n}{pn\sqrt{\log n}}\right)p^{\frac{pn\sqrt{\log n}}{\log \log n}}$ . There are  $\binom{n}{2}$  pairs of vertices, so by using a union bound, the probability that there exists a pair of vertices  $xy$  such that  $d(x, y) \geq \frac{pn\sqrt{\log n}}{\log \log n}$  is at most

$$\begin{aligned} \binom{n}{2} \left(\frac{n}{pn\sqrt{\log n}}\right)p^{\frac{pn\sqrt{\log n}}{\log \log n}} &\leq n^2 \left(\frac{enp}{pn\sqrt{\log n}}\right)^{\frac{pn\sqrt{\log n}}{\log \log n}} \leq n^2 \left(\frac{e \log \log n}{\sqrt{\log n}}\right)^{\frac{C \log n}{\log \log n}} \\ &\leq \exp \left(2 \log n + \frac{C \log n}{\log \log n} \log \frac{e \log \log n}{\sqrt{\log n}}\right) = \exp \left(\left(2 + \frac{C(1 + \log \log \log n)}{\log \log n} - \frac{C}{2}\right) \log n\right), \end{aligned}$$

where the last expression is  $o(1)$  for sufficiently large  $C$ .  $\square$

**Remark 7.** Lemma 6 shows one of the major differences in the proof compared to the case when  $p$  is only at least  $C\sqrt{\log n}/n$ , whereas in [2] it is proved that  $d(x, y) \leq 2pn$  with high probability when  $p > K \log n/n$ . Also note that with a direct application of the Chernoff's inequality we can only conclude that  $d(x, y) \leq pn\sqrt{\log n}$ , without the  $\log \log n$  factor. As we will see in Section 3, this  $\log \log n$  factor plays a vital role in the proof of Theorem 1 (see Remark 19).

**Lemma 8.** *There exists a constant  $C > 0$  such that if  $p > C\sqrt{\log n}/n$ , then w.h.p. for every subset  $S \subseteq [n]$ ,*

$$\left| \left\{ \{x, y\} \in \binom{[n] \setminus S}{2} : d_S(x, y) \geq 3pn \right\} \right| \leq n^2 e^{-\sqrt{\log n}}.$$

*Proof.* First, consider a fixed set  $S$ . For every pair of vertices  $\{x, y\} \in \binom{[n] \setminus S}{2}$ , the probability that  $d_S(x, y) > 3pn$  is at most  $\binom{|S|}{3pn} p^{3pn}$ . For different pairs  $\{x_1, y_1\}$  and  $\{x_2, y_2\}$  in  $\binom{[n] \setminus S}{2}$ ,  $d_S(x_1, y_1)$  and  $d_S(x_2, y_2)$  are independent. Hence, given a family of  $n^2 e^{-\sqrt{\log n}}$  pairs of vertices in  $\binom{[n] \setminus S}{2}$ , the probability that  $d_S(x, y) \geq 3pn$  for every pair  $\{x, y\}$  in this family is at most  $\left( \binom{|S|}{3pn} p^{3pn} \right)^{n^2 e^{-\sqrt{\log n}}}$ . Then, by a union bound over all the families of size  $n^2 e^{-c\sqrt{\log n}}$  of pairs of vertices not in  $S$ , we get

$$\begin{aligned} \mathbb{P} \left( \left| \left\{ \{x, y\} \in \binom{[n] \setminus S}{2} : d_S(x, y) \geq 3pn \right\} \right| > n^2 e^{-\sqrt{\log n}} \right) \\ \leq \binom{n^2}{n^2 e^{-\sqrt{\log n}}} \left( \binom{|S|}{3pn} p^{3pn} \right)^{n^2 e^{-\sqrt{\log n}}}. \end{aligned}$$

Finally, using a union bound over all the choices of  $S$ , the probability of failure is at most

$$2^n \binom{n^2}{n^2 e^{-\sqrt{\log n}}} \left( \binom{n}{3pn} p^{3pn} \right)^{n^2 e^{-\sqrt{\log n}}} \leq 2^n \left( e^{1+\sqrt{\log n}} \left( \frac{e}{3} \right)^{3pn} \right)^{n^2 e^{-\sqrt{\log n}}} = o(1). \quad \square$$

Let  $s = s(n, p)$ ,  $r = r(n, p)$ ,  $i = i(n, p)$  be positive integers depending on  $n$  and  $p$ , where  $s \leq n$ ,  $r \leq \binom{n}{2}$  and  $i \leq n$ . Let  $E_{s,r,i}$  be the event that for every vertex  $v \in [n]$  and vertex set  $S \subseteq [n]$ , where  $v \notin S$  and  $|S| = s$ , there are at most  $r$  pairs of vertices  $\{y_k, z_k\} \in \binom{[n] \setminus S}{2}$  such that  $vy_k z_k \in G$  and  $d_S(y_k, z_k) \geq i$  for every  $k$ , see Figure 3. Define  $g(p, s, r, i) := n \binom{n}{s} \binom{n^2}{r} (p \binom{s}{i} p^i)^r$ , and let  $O$  be the set of  $(s, r, i)$  such that  $g(p, s, r, i) = o(n^{-5})$  given  $p > C\sqrt{\log n}/n$ . Let  $E$  be the event  $\bigcap_{(s,r,i) \in O} E_{s,r,i}$ .

**Lemma 9.**  *$E$  happens with high probability.*


*Proof.* First consider  $\bar{E}_{s,r,i}$ , the complement of the event  $E_{s,r,i}$ . Fix distinct vertices  $v, y, z \in [n]$  and vertex set  $S \subset [n]$  where  $v, y, z \notin S$  and  $|S| = s$ , then

$$\mathbb{P}(vyz \in G) = p, \quad \mathbb{P}(d_S(y, z) \geq i) \leq \binom{s}{i} p^i.$$

Note that these two events are independent since  $v \notin S$ . Therefore, the probability that there are at least  $r$  pairs of vertices  $(y_k, z_k) \in \binom{[n] \setminus S}{2}$  where  $vy_k z_k \in G$  and  $d_S(y_k, z_k) \geq i$  is at most  $\binom{n^2}{r} (p \binom{s}{i} p^i)^r$ . Then, using a union bound over all choices of  $v \in [n]$  and  $S \subset [n]$  of size  $s$ , we get that  $\mathbb{P}(\bar{E}_{s,r,i}) \leq g(p, s, r, i) = o(n^{-5})$ . Then we have

$$\mathbb{P}(\bar{E}) = \mathbb{P} \left( \bigcup_{(s,r,i) \in O} \bar{E}_{s,r,i} \right) \leq \sum_{(s,r,i) \in O} \mathbb{P}(\bar{E}_{s,r,i}) \leq n \binom{n}{2} n \cdot o(n^{-5}) = o(1). \quad \square$$

Figure 3: The event  $E_{s,r,i}$  is that there are at most  $r$  pairs of vertices  $\{y_k, z_k\} \in \binom{[n] \setminus S}{2}$  such that  $vy_k z_k \in G$  and  $d_S(y_k, z_k) \geq i$  for every  $k$ .



In our proof of Theorem 1, we will repeatedly use Lemma 9 to show that for some given vertex  $v$  and vertex set  $S$ , the number of pairs that are in the link graph of  $v$  and have a large neighborhood in  $S$  is small.

**Remark 10.** Lemma 8 and Lemma 9 are similar and one may hope that we can still keep the  $e^{-\sqrt{\log n}}$  factor by proving  $(s, pn^2 e^{-\sqrt{\log n}}, 3ps) \in O$  for every  $1 \leq s \leq n$ . Unfortunately, this is not necessary true. In the proof of Lemma 8, we have  $n^2 e^{-\sqrt{\log n}}$  in the exponent, which can beat the number of choices of  $S$ , whereas  $pn^2 e^{-\sqrt{\log n}}$  is not larger than  $n$ . Instead, we have  $(s, pn^2 / \log n, 3pn) \in O$ , see Claim 25.

For fixed vertex  $v$ , vertex set  $A \subseteq [n] \setminus \{v\}$ , subset  $T$  of  $N_{[n] \setminus A, [n] \setminus A}(v)$  and subset  $E$  of  $\{vxy \in G : x \in A\}$ , define

$$K(v, E, A, T) := \{xyz : x \in A, yz \in T, \text{ there exists } e \in E \text{ such that } x \in e, y \notin e, z \notin e\},$$

and  $G(v, E, A, T) := K(v, E, A, T) \cap G$ . We have that  $\mathbb{E}(|G(v, E, A, T)|) = p|K(v, E, A, T)|$ ; note that here the randomness in this expectation is of the hyperedges in  $\{xyz : x \in A, yz \in T\}$  being or not in  $G$ . Also note that the events of the hyperedges in  $\{xyz : x \in A, yz \in T\}$  being or not in  $G$  are independent from the events from hyperedges containing  $v$  being in  $G$ . We will condition on which hyperedges containing  $v$  are present in  $G$  in the argument that follows, and we will do so repeatedly without further notice. Also note that for every  $xyz \in G(v, E, A, T)$  with  $x \in A, yz \in T$ , we can find an  $F_5 = \{yzv, yzx, vxw\}$  in  $G$  where  $vxw \in E$ , see Figure 4. The condition  $y \notin e, z \notin e$  in the definition of  $K(v, E, A, T)$  guarantees that we indeed find an  $F_5$  instead of an  $K_4^-$ .

**Lemma 11.** *For any constants  $\varepsilon, \varepsilon_1, \varepsilon_2 > 0$ , there exists a constant  $C > 0$  such that if  $p > C\sqrt{\log n}/n$ , then with high probability for every choice of  $v, E, A, T$  as above with  $|A| \geq \varepsilon_1 n / \sqrt{\log n}$  and  $|T| \geq \varepsilon_2 pn^2$ , we have  $|G(v, E, A, T)| = (1 \pm \varepsilon)p|A||T|$ .*

The proof of Lemma 11 follows the same lines of the proof of Proposition 9 in [2].




Figure 4: A copy of  $F_5$  from  $xyz \in G(v, E, A, T)$ .

*Proof.* Here we reveal the randomness into two phases. We first reveal the hyperedges containing  $v$ ; given this choice we can fix  $A$ ,  $T$  and  $E$ . We have at most  $n$  choices for  $v$ . For sets  $A, T$  with  $|A| = a$  and  $|T| = t$ , there are at most  $\binom{n}{a}$  choices for  $A$ , and with high probability there are at most  $2^{apn \frac{\sqrt{\log n}}{\log \log n}}$  choices for  $E$  (by Lemma 6) and at most  $\binom{pn^2}{t}$  choices for  $T$  (by Lemma 4).

For vertex  $x \in A$ , let  $[x, T] := \{xyz : yz \in T\}$ ,  $d^E(x) := |\{e \in E : x \in e\}|$  and  $T_x := \{yz \in T | vxy \in E \text{ or } vxz \in E\}$ . Let  $[A, T] := \{xyz : x \in A, yz \in T\}$ . Note that  $[A, T] \supseteq K(v, E, A, T)$ . If  $d^E(x) \geq 3$ , choose  $v x w_1, v x w_2, v x w_3$  arbitrarily from  $E$ . For every  $yz \in T$ , there exists some  $i \in \{1, 2, 3\}$  such that  $y \neq w_i$  and  $z \neq w_i$ . Hence, by the definition of  $K(v, E, A, T)$ , we have  $[x, T] \subseteq K(v, E, A, T)$ . If  $d^E(x) \leq 2$ , by Lemma 6, we have  $|T_x| \leq 2 \cdot \frac{pn\sqrt{\log n}}{\log \log n}$  and  $[x, T \setminus T_x] = \{xyz : yz \in T \setminus T_x\} \subseteq K(v, E, A, T)$ . Therefore,

$$|[A, T]| - |K(v, E, A, T)| \leq \sum_{x: x \in A, d^E(x) \leq 2} |T_x| \leq |A| \cdot 2pn \frac{\sqrt{\log n}}{\log \log n}.$$

We have  $|[A, T]| = |A||T| \geq |A|\varepsilon_2 pn^2$ , so  $|K(v, E, A, T)| = (1 - o(1))|A||T|$  with high probability.

Now, we reveal the hyperedges in  $[A, T]$ , Let  $\mu = \mathbb{E}(G(v, E, A, T)) = p|K(v, E, A, T)| = (1 - o(1))p|A||T|$ . By Lemma 3, we have

$$\mathbb{P}(|G(v, E, A, T)| - \mu > \varepsilon \mu) < 2e^{-c_\varepsilon \mu}.$$

Then by the union bound, the probability that the statement in the lemma does not hold is

at most

$$\begin{aligned}
& \sum_{a \geq \frac{\varepsilon_1 n}{\sqrt{\log n}}} \sum_{t \geq \varepsilon_2 p n^2} n \binom{n}{a} 2^{apn \frac{\sqrt{\log n}}{\log \log n}} \binom{pn^2}{t} \cdot 2e^{-c_\varepsilon \cdot atp/2} \\
& \leq \sum_{a \geq \frac{\varepsilon_1 n}{\sqrt{\log n}}} \sum_{t \geq \varepsilon_2 p n^2} n \exp\left(a \log \frac{en}{a}\right) 2^{apn \frac{\sqrt{\log n}}{\log \log n}} \exp\left(t \log \frac{pn^2}{t}\right) \cdot 2e^{-c_\varepsilon \cdot atp/2} \\
& \leq 2 \sum_{a \geq \frac{\varepsilon_1 n}{\sqrt{\log n}}} \sum_{t \geq \varepsilon_2 p n^2} \exp\left(\log n + a \log\left(\frac{e}{\varepsilon_1} \sqrt{\log n}\right) + apn \frac{\sqrt{\log n} \log 2}{\log \log n} + t \log \frac{1}{\varepsilon_2} - c_\varepsilon atp/2\right) \\
& \leq 2 \sum_{a \geq \frac{\varepsilon_1 n}{\sqrt{\log n}}} \sum_{t \geq \varepsilon_2 p n^2} \exp(-c_\varepsilon atp/4) \leq 2n^3 \exp(-n\sqrt{\log n}) = o(1). \quad \square
\end{aligned}$$

### 3 Proof of the Main Theorem

Recall that for a given partition  $\pi = (V_1, V_2, V_3)$ ,

$$\begin{aligned}
\bar{H}_\pi &= G_\pi \setminus H_\pi, \\
H_i &= \{e \in H : |e \cap V_i| \geq 2\} \quad \text{for } i \in \{1, 2, 3\}, \quad \text{and} \\
Q(\pi) &= \{\{u, v\} \subset V_1 : |L_{2,3}(u, v)| < 0.8p^2n^2/9\}.
\end{aligned}$$

For a 3-uniform hypergraph  $H$ , the *shadow graph* of  $H$  is the graph on the same vertex set, where  $xy$  is an edge if and only if there exist another vertex  $z$  such that  $xyz$  is a hyperedge in  $H$ .

**Proposition 12.** *Let  $H$  be an  $F_5$ -free subhypergraph of  $G$  and  $\pi = (V_1, V_2, V_3)$  be a balanced partition maximizing  $|H_\pi|$ . Then there exist positive constants  $C$  and  $\delta$  such that if  $p > C\sqrt{\log n}/n$  and if the following conditions hold:*

1.  $\sum_{i=1}^3 |H_i| \leq \delta p n^3$ ,
2. the shadow graph of  $H_1$  is disjoint from  $Q(\pi)$ ,

*then with high probability  $|\bar{H}_\pi| \geq 3|H_1|$ , where equality is possible only if  $H$  is tripartite.*

**Proposition 13.** *There exist positive constants  $C$  and  $\delta$  such that if  $p > C\sqrt{\log n}/n$  and the 3-partition  $\pi$  is balanced, then with high probability*

$$t(G) \geq |G_\pi| + |Q(\pi)|\delta n^2 p^2,$$

*where equality is possible only if  $Q(\pi) = \emptyset$ .*

Based on Propositions 12 and 13, Theorem 1 easily follows, whose proof is similar to the proof of the main theorem in [2].



*Proof of Theorem 1.* Let  $H$  be a maximum  $F_5$ -free subhypergraph of  $G$ . We have  $|H| \geq t(G)$ , since  $F_5$  is not tripartite. Let  $\pi = (V_1, V_2, V_3)$  be a 3-partition of  $[n]$  maximizing  $|H_\pi|$ . Without loss of generality, we may assume  $|H_1| \geq |H_2|, |H_3|$ . Besides, by Proposition 2, we know that  $\pi$  is balanced and  $\sum_{i=1}^3 |H_i| \leq \delta p n^3$ , where  $\delta$  is the constant in Proposition 12. Now let

$$B(\pi) := \{e \in G : \text{there exists } \{u, v\} \in Q(\pi) \text{ such that } \{u, v\} \subset e\}$$

and  $H' := H \setminus B(\pi)$ , so the shadow graph of  $H'_1$  is disjoint from  $Q(\pi)$ . Since  $B(\pi)$  consists of only non-crossing hyperedges of  $H$ ,  $\pi$  is still a partition maximizing  $|H'_\pi|$  and  $\sum_{i=1}^3 |H'_i| \leq \sum_{i=1}^3 |H_i| \leq \delta p n^3$ . Hence,  $H'$  satisfies the assumptions in Proposition 12. Now, we have w.h.p.

$$\begin{aligned} |H| &\leq |H_\pi| + 3|H_1| = |H'_\pi| + 3|H'_1| + 3|H \cap B(\pi)| \\ &\leq |H'_\pi| + 3|H'_1| + 3|B(\pi)| \\ &\leq |G_\pi| + 3|B(\pi)| \end{aligned} \tag{1}$$

$$\leq |G_\pi| + 3|Q(\pi)|pn\sqrt{\log n}/\log \log n \tag{2}$$

$$\leq |G_\pi| + |Q(\pi)|\delta p^2 n^2 \tag{3}$$

$$\leq t(G). \tag{4}$$

Here we use Proposition 12 for (1), Lemma 6 for (2), the assumption  $p > C\sqrt{\log n}/n$  for (3), and Proposition 13 for (4).  $|H|$  cannot be strictly smaller than  $t(G)$ , so all the inequalities must hold with equality. By Propositions 12 and 13,  $H'$  is tripartite and  $Q(\pi)$  is empty. Then, we conclude that  $H = H'$  is tripartite.  $\square$

The proof of Proposition 13 is exactly the same as the one in [2], where it is required that  $p > K \log n/n$ . It can be easily checked that all the arguments still work verbatim with  $p > C\sqrt{\log n}/n$ . It remains to prove Proposition 12. Clearly, we can assume that  $|\bar{H}_\pi| \leq 3|H_1|$  and  $|H_1| \geq |H_2|, |H_3|$ . Let  $\delta$  be small enough so that the following arguments work, and fix three small positive constants  $\varepsilon_1, \varepsilon_2$  and  $\varepsilon_3$  such that

$$\frac{1}{72\varepsilon_1} \geq 30, \quad \frac{1}{20} \cdot \frac{1}{2\varepsilon_2} \geq 10, \quad \frac{1}{10} - \varepsilon_2 \geq \frac{1}{20}, \quad \frac{100\delta}{\varepsilon_1} \leq \varepsilon_3 \leq \varepsilon_1, \quad \frac{\frac{1}{4}\varepsilon_1\varepsilon_2}{3\varepsilon_3} \geq 20.$$


For example, we can set

$$\delta = 10^{-100}, \quad \varepsilon_1 = \frac{1}{3000}, \quad \varepsilon_2 = \frac{1}{400}, \quad \varepsilon_3 = 10^{-10}.$$

Denote by  $J$  the induced subgraph of the shadow graph of  $H_1$  on the vertex set  $V_1$  and use  $N^J(x)$ ,  $d^J(x)$  for neighborhood and degree of  $x$  in graph  $J$ , respectively. Call a 4-set  $\{w_1, w_2, y, z\}$  an  $\hat{F}_5$  if  $w_1yz, w_2yz \in G_\pi$  and there exists  $e \in H$  such that  $w_1, w_2 \in e \cap V_1$ ,  $y, z \notin e$ . Note that  $\{w_1yz, w_2yz, e\}$  forms a copy of  $F_5$  in  $G$ , so at least one of  $w_1yz$  and  $w_2yz$  has to be in  $\bar{H}_\pi$ .

We next count copies of  $\hat{F}_5$  to lower bound the number of missing crossing hyperedges  $|\bar{H}_\pi|$ , based on the size of (a subgraph of)  $J$ . First, we prove the following claim, which will be used in the proofs of the next few lemmas.

Figure 5: The set  $\{w_1, w_2, y, z\}$  is an  $\hat{F}_5$  if there exists an edge  $e \in H_1$  as below.



**Claim 14.** *If  $p > C\sqrt{\log n}/n$ , then with high probability for every edge  $x_1x_2 \in J \setminus Q(\pi)$ , there are at least  $p^2n^2/12$  choices of  $(y, z)$  where  $y \in V_2, z \in V_3$  such that  $\{x_1, x_2, y, z\}$  spans an  $\hat{F}_5$ .*

*Proof.* Since  $x_1x_2$  is in  $J$ , there exists  $v_0$  such that  $x_1x_2v_0 \in H$ . Since  $x_1x_2$  is not in  $Q(\pi)$ , there exists at least  $0.8p^2n^2/9$  choices of  $(y, z)$  such that  $y \in V_2, z \in V_3$ , and  $x_1yz$  and  $x_2yz$  are both in  $G_\pi$ . By Lemma 6, we know  $d(x_1, v_0)$  is at most  $pn \frac{\sqrt{\log n}}{\log \log n}$ , so there can be at most  $pn \frac{\sqrt{\log n}}{\log \log n}$  vertices  $v$  such that  $x_1v_0v \in G_\pi$ . Therefore, there are at least  $0.8p^2n^2/9 - pn \frac{\sqrt{\log n}}{\log \log n} \geq p^2n^2/12$  choices of  $(y, z)$  such that  $x_1yz, x_2yz \in G_\pi$  and  $v_0 \notin \{y, z\}$ . For every such  $(y, z)$ ,  $\{x_1, x_2, y, z\}$  forms an  $\hat{F}_5$ .  $\square$

**Lemma 15.** *Suppose the assumptions of Proposition 12 hold. Let  $J'$  be a subgraph of  $J$  and denote by  $\Delta(J')$  the maximum degree of  $J'$ . If  $\Delta(J') \leq \varepsilon_1 n$ , then with high probability*

$$|\bar{H}_\pi| \geq 30pn|J'|.$$

*If further  $\Delta(J') \leq \varepsilon_1 n / \sqrt{\log n}$ , then with high probability*

$$|\bar{H}_\pi| \geq 20pn|J'| \frac{\sqrt{\log n}}{\log \log n}.$$

*Proof.* For each  $wx \in J'$ , we get  $wx \notin Q(\pi)$  by the assumption of Proposition 12, so there are at least  $p^2n^2/12$  choices of  $(y, z)$  such that  $\{w, x, y, z\}$  spans an  $\hat{F}_5$ , by Claim 14. Then there are at least  $\frac{1}{12}|J'|p^2n^2 = \frac{1}{2} \sum_{x \in V_1} d^{J'}(x) \frac{p^2n^2}{12}$  copies of  $\hat{F}_5$  in total.

Now consider missing crossing hyperedges  $xyz \in \bar{H}_\pi$  with  $x \in V_1$ . Call  $xyz \in \bar{H}_\pi$  bad if

$$d^{J'}(x) \leq \frac{\varepsilon_1 n}{\sqrt{\log n}} \quad \text{and} \quad |N(y, z) \cap N^{J'}(x)| \geq \frac{pn \log \log n}{500\sqrt{\log n}}, \quad \text{or}$$

$$\frac{\varepsilon_1 n}{\sqrt{\log n}} < d^{J'}(x) \leq \varepsilon_1 n \quad \text{and} \quad |N(y, z) \cap N^{J'}(x)| \geq 3\varepsilon_1 pn.$$

Otherwise, call  $xyz$  *good*. We will show that the number of copies of  $\hat{F}_5$  that contain a good hyperedge from  $\bar{H}_\pi$  is at least  $\frac{1}{4} \sum_{x \in V_1} d^{J'}(x) \frac{p^2 n^2}{12}$ . For  $x \in V_1$ , let  $n_x$  be the number of copies of  $\hat{F}_5$  that contain vertex  $x$  and a bad missing crossing hyperedge, and let  $r_x$  be the number of  $(y, z)$  such that  $y \in V_2, z \in V_3$ , and  $xyz$  is bad. We will repeatedly use Lemma 9 for some choices of  $(s, r, i) \in O$  to give upper bounds for  $r_x$  and then obtain upper bounds for  $n_x$ . For every  $x \in V_1$ , we consider the following two cases.

- **Case 1:** If  $d^{J'}(x) \leq \frac{\varepsilon_1 n}{\sqrt{\log n}}$ . If  $d^{J'}(x) = 0$ , then  $n_x$  is trivially 0. For every positive integer  $s \leq \frac{\varepsilon_1 n}{\sqrt{\log n}}$ ,  $r = \frac{pn}{\sqrt{\log n}} s$ , and  $i = \frac{pn \log \log n}{500 \sqrt{\log n}}$ , we have  $(s, r, i) \in O$  (see Claim 23 in the Appendix for the proof). Hence, by Lemma 9, we get  $r_x \leq \frac{pn}{\sqrt{\log n}} d^{J'}(x)$ . By Lemma 6, we can assume every pair of vertices has codegree at most  $pn \frac{\sqrt{\log n}}{\log \log n}$  in  $G$ . We conclude

$$n_x \leq pn \frac{\sqrt{\log n}}{\log \log n} \cdot \frac{pn}{\sqrt{\log n}} d^{J'}(x) \leq \frac{p^2 n^2}{100} d^{J'}(x).$$

- **Case 2:** If  $\frac{\varepsilon_1 n}{\sqrt{\log n}} < d^{J'}(x) \leq \varepsilon_1 n$ . For every integer  $s \in (\frac{\varepsilon_1 n}{\sqrt{\log n}}, \varepsilon_1 n]$ ,  $r = \frac{pn}{500} s$ , and  $i = 3\varepsilon_1 pn$ , we have  $(s, r, i) \in O$  (see Claim 24 in the Appendix for the proof). Hence, by Lemma 9, we have  $r_x \leq \frac{pn}{500} d^{J'}(x)$ . By Lemma 6, we can assume every pair of vertices has codegree at most  $pn \frac{\sqrt{\log n}}{\log \log n}$  in  $G$ . Besides, we have  $(s, pn^2 / \log n, 3pn) \in O$  for every  $1 \leq s \leq n$  (see Claim 25 in the Appendix for the proof), so there are at most  $pn^2 / \log n$  pairs  $(y, z)$  where  $y \in V_2, z \in V_3$  such that  $|N(y, z) \cap N^{J'}(x)| \geq 3pn$ . We conclude

$$n_x \leq \frac{pn^2}{\log n} \cdot pn \frac{\sqrt{\log n}}{\log \log n} + 3r_x pn \leq \frac{p^2 n^2}{\log \log n} \cdot \frac{n}{\sqrt{\log n}} + \frac{3p^2 n^2}{500} d^{J'}(x) \leq \frac{p^2 n^2}{100} d^{J'}(x).$$

Thus, the number of copies of  $\hat{F}_5$  that contain a good hyperedge from  $\bar{H}_\pi$  is at least

$$\frac{1}{2} \sum_{x \in V_1} d^{J'}(x) \frac{p^2 n^2}{12} - \sum_{x \in V_1} \frac{p^2 n^2}{100} d^{J'}(x) \geq \frac{1}{4} \sum_{x \in V_1} d^{J'}(x) \frac{p^2 n^2}{12} = \frac{1}{24} |J'| p^2 n^2. \quad (5)$$

If  $\Delta(J') \leq \varepsilon_1 n$ , every good missing crossing hyperedge is in at most  $3\varepsilon_1 pn$  copies of  $\hat{F}_5$  estimated in (5), so

$$|\bar{H}_\pi| \geq \frac{\frac{1}{24} |J'| p^2 n^2}{3\varepsilon_1 pn} \geq \frac{pn |J'|}{72\varepsilon_1} \geq 30pn |J'|.$$

If further  $\Delta(J') \leq \varepsilon_1 n / \sqrt{\log n}$ , then every good missing crossing hyperedge is in at most  $\frac{pn \log \log n}{500 \sqrt{\log n}}$  copies of  $\hat{F}_5$  estimated in (5), so

$$|\bar{H}_\pi| \geq \frac{\frac{1}{24} |J'| p^2 n^2}{pn \log \log n / (500 \sqrt{\log n})} \geq \frac{20pn |J'| \sqrt{\log n}}{\log \log n}. \quad \square$$

Now, we divide the rest of the proof into two cases according to the size of  $\bar{H}_\pi$ . The notations and calculations are similar in both cases.

### 3.1 $|\bar{H}_\pi| \leq \delta p n^3 / \log n$

Let  $S = \{x \in V_1 : d^J(x) \geq \varepsilon_1 n / \sqrt{\log n}\}$ ,  $S_1 = \{x \in S : d_{2,3}^H(x) \geq \varepsilon_2 p n^2\}$  and  $S_2 = S \setminus S_1$ . For the following lemmas, we always assume that  $|\bar{H}_\pi| \leq \delta p n^3 / \log n$  and the assumptions of Proposition 12 hold.

**Lemma 16.** *With high probability  $|S| \leq \varepsilon_3 n / \sqrt{\log n}$ .*

*Proof.* For each  $wx \in J'$ , we get  $wx \notin Q(\pi)$  by the assumption of Proposition 12, so there are at least  $p^2 n^2 / 12$  choices of  $(y, z)$  such that  $\{w, x, y, z\}$  spans an  $\hat{F}_5$ , by Claim 14. Then there are at least  $\frac{1}{12} |J'| p^2 n^2$  copies of  $\hat{F}_5$  in total. On the other hand, at least one of  $wyz$  and  $xyz$  must be in  $\bar{H}_\pi$ . For  $xyz$  where  $d(y, z) \leq 3pn$ ,  $xyz$  can be in at most  $3pn$  copies of  $\hat{F}_5$ . Hence the number of copies of  $\hat{F}_5$  containing a pair  $(y, z) \in V_2 \times V_3$  with  $d(y, z) \leq 3pn$  is at most  $3|\bar{H}_\pi|pn$ . By Lemma 8, there can be at most  $n^2 e^{-\sqrt{\log n}}$  pairs  $(y, z)$  with  $d(y, z) \geq 3pn$ , and by Lemma 6, for every pair  $(y, z)$ , we can assume  $d(y, z) \leq pn \frac{\sqrt{\log n}}{\log \log n}$ . Hence the number of copies of  $\hat{F}_5$  containing a pair  $(y, z) \in V_2 \times V_3$  with  $d(y, z) \geq 3pn$  is at most  $n^2 e^{-\sqrt{\log n}} \left( pn \frac{\sqrt{\log n}}{\log \log n} \right)^2$ . Therefore, we get

$$n^2 e^{-\sqrt{\log n}} \left( pn \frac{\sqrt{\log n}}{\log \log n} \right)^2 + |\bar{H}_\pi| \cdot 3pn \geq \frac{|J| p^2 n^2}{12}. \quad (6)$$

Note that  $n^2 e^{-\sqrt{\log n}} \left( pn \frac{\sqrt{\log n}}{\log \log n} \right)^2 \leq \frac{\delta p^2 n^4}{\log n}$  and by the assumption  $|\bar{H}_\pi| < \frac{\delta p n^3}{\log n}$  so  $|\bar{H}_\pi| \cdot 3pn \leq \frac{3\delta p^2 n^4}{\log n}$ . We get  $|J| \leq \frac{48\delta n^2}{\log n}$ . Every vertex in  $S$  has degree at least  $\frac{\varepsilon_1 n}{\sqrt{\log n}}$  in  $J$ , so  $\frac{\varepsilon_1 n}{\sqrt{\log n}} |S| \leq 2|J| \leq \frac{96\delta n^2}{\log n}$ , which confirms that  $|S| \leq \frac{100\delta}{\varepsilon_1} \cdot \frac{n}{\sqrt{\log n}} \leq \frac{\varepsilon_3 n}{\sqrt{\log n}}$ .  $\square$

**Lemma 17.** *With high probability  $|\bar{H}_\pi| \geq 20pn^2|S_1|$ .*

*Proof.* We can assume that  $|S_1| > 0$ , since otherwise this inequality is trivial. For every vertex  $x \in S_1$ , define

$$T_x := \{yz : y \in V_2, z \in V_3, xyz \in H, d_{S_1}(y, z) \leq 3\varepsilon_3 pn / \sqrt{\log n}\}.$$

By Lemma 16, we know  $|S_1| \leq |S| \leq \varepsilon_3 n / \sqrt{\log n}$ . For every  $s \leq \varepsilon_3 n / \sqrt{\log n}$ , we have  $(s, \varepsilon_2 p n^2 / 2, 3\varepsilon_3 pn / \sqrt{\log n}) \in O$  (see Claim 26 in the Appendix for the proof), so there can be at most  $\varepsilon_2 p n^2 / 2$  pairs of  $(y, z)$  such that  $xyz \in G$  and  $d_{S_1}(y, z) \geq 3\varepsilon_3 pn / \sqrt{\log n}$ . By the definition of  $S_1$ , we have  $d_{2,3}^H(x) \geq \varepsilon_2 p n^2$  for every  $x \in S_1$ . Thus, we get  $|T_x| \geq \varepsilon_2 p n^2 / 2$ .

Now, we count the copies of  $\hat{F}_5 = \{x, w, y, z\}$  where  $x \in S_1$ ,  $w \in N^J(x)$ , and  $\{y, z\} \in T_x$ . Note that  $wyz$  must be in  $\bar{H}_\pi$ , since  $xyz \in H$ . We have that  $|N^J(x)| \geq \varepsilon_1 n / \sqrt{\log n}$  by the definition of  $S$ , and we just confirm  $|T_x| \geq \varepsilon_2 p n^2 / 2$ . Applying Lemma 11 by setting  $v = x$ ,  $E = H_1$ ,  $A = N^J(x)$ , and  $T = T_x$ , we get the number of such copies of  $\hat{F}_5$  is at least

$$\sum_{x \in S_1} \frac{1}{2} p |N^J(x)| |T_x| \geq \sum_{x \in S_1} \frac{1}{2} p \cdot \frac{\varepsilon_1 n}{\sqrt{\log n}} \cdot \frac{1}{2} \varepsilon_2 p n^2 \geq \frac{\varepsilon_1 \varepsilon_2 p^2 n^3 |S_1|}{4\sqrt{\log n}}. \quad (7)$$

Every  $wyz \in \bar{H}_\pi$  can be in at most  $3\varepsilon_3 pn / \sqrt{\log n}$  copies of  $\hat{F}_5$  evaluated in (7), because  $x$  is assumed to be in  $S_1$  and  $d_{S_1}(y, z) \leq 3\varepsilon_3 pn / \sqrt{\log n}$ . Therefore,

$$|\bar{H}_\pi| \geq \frac{\frac{1}{4}\varepsilon_1\varepsilon_2 p^2 n^3 |S_1| / \sqrt{\log n}}{3\varepsilon_3 pn / \sqrt{\log n}} \geq 20pn^2 |S_1|. \quad \square$$

**Lemma 18.** *With high probability  $|\bar{H}_\pi| \geq \frac{1}{20}pn^2 |S_2|$ .*

*Proof.* For every vertex  $x \in S_2$ , we have  $d_{2,3}^H(x) < \varepsilon_2 pn^2$  by the definition of  $S_2$ , but by Lemma 5,  $d_{2,3}(x) \geq pn^2/10$ . Thus, there are at least  $pn^2/20$  hyperedges in  $\bar{H}_\pi$  containing  $x$ , so  $|\bar{H}_\pi| \geq |S_2|pn^2/20$ .  $\square$

Finally, we deduce Proposition 12 assuming  $|\bar{H}_\pi| \leq \delta pn^3 / \log n$ .

*Proof of Proposition 12.* We will show that with high probability  $|\bar{H}_\pi| \geq 3|H_1|$  by using the lower bounds in Lemmas 15, 17 and 18. Partition  $H_1$  into the following three sets.

- $H_1(1) = \{e \in H_1 : |e \cap S| \geq 2 \text{ or } |e \cap V_1 \setminus S| \geq 2\}$ .
- $H_1(2) = \{e \in H_1 \setminus H_1(1) : |e \cap S_1| = 1\}$ . Hence,  $H_1(2)$  contains those hyperedges in  $H_1$  with exactly one vertex in  $S_1$ , one vertex in  $V_1 \setminus S$ , and one vertex in  $[n] \setminus V_1$ .
- $H_1(3) = H_1 \setminus (H_1(1) \cup H_1(2))$ . Hence,  $H_1(3)$  contains those hyperedges in  $H_1$  with exactly one vertex in  $S_2$ , one vertex in  $V_1 \setminus S$ , and one vertex in  $[n] \setminus V_1$ .

There are three cases needed to be handled.

- **Case 1:** If  $3|H_1(1)| \geq |H_1|$ .

Let  $J' := J[S] \cup J[V_1 \setminus S]$ , where  $J[S]$  and  $J[V_1 \setminus S]$  are the induced subgraph of  $J$  on  $S$  and  $V_1 \setminus S$  separately. By Lemma 6,  $|H_1(1)| \leq |J'|pn\sqrt{\log n} / \log \log n$ . For every vertex  $x \in S$ ,  $d^{J'}(x)$ , the degree of  $x$  in  $J'$ , is at most  $|S| - 1 \leq \varepsilon_3 n / \sqrt{\log n} \leq \varepsilon_1 n / \sqrt{\log n}$ , by Lemma 16. For every vertex  $x \in V \setminus S$ , we have  $d^{J'}(x) \leq d^J(x) \leq \varepsilon_1 n / \sqrt{\log n}$ , by the definition of  $S$ . Hence,  $\Delta(J') \leq \varepsilon_1 n / \sqrt{\log n}$ . Then, by Lemma 15,

$$|\bar{H}_\pi| \geq \frac{20pn|J'|\sqrt{\log n}}{\log \log n} \geq 20|H_1(1)| \geq 3|H_1|.$$

- **Case 2:** If  $3|H_1(2)| \geq |H_1|$ .

For every vertex  $x \in S_1$ , there are at most  $2pn^2$  hyperedges in  $H_1 \setminus H_1(1)$  containing it, by Lemma 4. Hence,  $|H_1(2)| \leq 2pn^2 |S_1|$ . Then, by Lemma 17,

$$|\bar{H}_\pi| \geq 20pn^2 |S_1| \geq 10|H_1(2)| \geq 3|H_1|.$$

- **Case 3:** If  $3|H_1(3)| \geq |H_1|$ .

By the definition of  $S_2$ , every vertex  $x \in S_2$  has  $d_{2,3}^H(x) \leq \varepsilon_2 p n^2$ . Recall that  $\pi$  maximizes  $|H_\pi|$ , so  $d_{1,2}^H(x), d_{1,3}^H(x) \leq \varepsilon_2 p n^2$ . Hence,  $|H_1(3)| \leq 2\varepsilon_2 p n^2 |S_2|$ . Then, by Lemma 18,

$$|\bar{H}_\pi| \geq \frac{1}{20} p n^2 |S_2| \geq \frac{1}{20} \cdot \frac{|H_1(3)|}{2\varepsilon_2} \geq 10|H_1(3)| \geq 3|H_1|. \quad \square$$

**Remark 19.** The  $\log \log n$  factor in Lemma 6 plays an important role in the case  $3|H_1(1)| \geq |H_1|$ . Without this factor, we would need to conclude  $|\bar{H}_\pi| \geq 20pn|J'|\sqrt{\log n}$  from Lemma 15, which we can only obtain when  $\Delta(J') \leq \varepsilon_1 n / \log^{1/2+c} n$  for some constant  $c > 0$ . Then, we would need to modify the definition of  $S$  to be  $\{x \in V_1 : d^J(x) \geq \varepsilon_1 n / \log^{1/2+c} n\}$ . However, the assumption of Lemma 11 is no longer valid for sets  $A$  of smaller size. Therefore, we would not be able to use Lemma 11 in the proof of Lemma 17.

### 3.2 $|\bar{H}_\pi| > \delta p n^3 / \log n$

All the notations and theorems here are similar to those in the previous subsection, so we will just point out the necessary modifications. Note that by the assumption in Proposition 12, we have  $|\bar{H}_\pi| \leq \sum_{i=1}^3 |H_i| \leq \delta p n^3$ . Let  $S' = \{x \in V_1 : d^J(x) \geq \varepsilon_1 n\}$ ,  $S'_1 = \{x \in S' : d_{2,3}^H(x) \geq \varepsilon_2 p n^2\}$  and  $S'_2 = S' \setminus S'_1$ . For the following lemmas, we always assume that  $|\bar{H}_\pi| > \delta p n^3 / \log n$  and the assumptions of Proposition 12 hold.

**Lemma 20.** *With high probability  $|S'| \leq \varepsilon_3 n$ .*

*Proof.* Inequality (6) still holds and now we get  $|J| \leq 48\delta n^2$ . Every vertex in  $S'$  has degree at least  $\varepsilon_1 n$  in  $J$ , so  $\varepsilon_1 n |S'| \leq 2 \cdot 48\delta n^2$ , which gives  $|S'| \leq \frac{100\delta}{\varepsilon_1} n = \varepsilon_3 n$ .  $\square$

**Lemma 21.** *With high probability  $|\bar{H}_\pi| \geq 20pn^2 |S'_1|$ .*

*Proof.* We can assume that  $|S'_1| \geq 1$ , otherwise this inequality is trivial. For each  $x \in S'_1$ , define

$$T'_x := \{yz : y \in V_2, z \in V_3, xyz \in H, d_{S'_1}(y, z) \leq 3\varepsilon_3 p n\}.$$

By Lemma 20,  $|S'_1| \leq |S'| \leq \varepsilon_3 n$ . For every  $s \leq \varepsilon_3 n$ , we have  $(s, \varepsilon_2 p n^2 / 2, 3\varepsilon_3 p n) \in O$  (see Claim 27 in the Appendix for the proof). Then by Lemma 9 and the definition of  $S'_1$ , we get that  $|T'_x| \geq \frac{1}{2} \varepsilon_2 p n^2$ . Now we count those copies of  $\hat{F}_5 = \{x, w, y, z\}$  where  $x \in S'_1$ ,  $w \in N^J(x)$ , and  $\{y, z\} \in T'_x$ . By Lemma 11, the number of such copies of  $\hat{F}_5$  is at least  $\frac{1}{4} \varepsilon_1 \varepsilon_2 p^2 n^3 |C_1|$ . Every  $wyz \in \bar{H}_\pi$  can be in at most  $3\varepsilon_3 p n$  such copies of  $\hat{F}_5$ , because  $x$  is assumed to be in  $S'_1$  and  $d_{S'_1}(y, z) \leq 3\varepsilon_3 p n$ . Therefore,

$$|\bar{H}_\pi| \geq \frac{\frac{1}{4} \varepsilon_1 \varepsilon_2 p^2 n^3 |S'_1|}{3\varepsilon_3 p n} \geq 20pn^2 |S'_1|. \quad \square$$

**Lemma 22.** *With high probability  $|\bar{H}_\pi| \geq \frac{1}{20} p n^2 |S'_2|$ .*

*Proof.* Exactly the same as the proof of Lemma 18.  $\square$

We are now able to conclude the proof of Proposition 12 for the remaining case that  $\frac{\delta pn^3}{\log n} < |\bar{H}_\pi| \leq \delta pn^3$ .

*Proof of Proposition 12.* Similarly to the proof in the previous subsection, we define  $H'_1(1) = \{e \in H_1 : |e \cap S'| \geq 2 \text{ or } |e \cap V_1 \setminus S'| \geq 2\}$ ,  $H'_1(2) = \{e \in H_1 \setminus H'_1(1) : |e \cap S'_1| = 1\}$  and  $H'_1(3) = H'_1 \setminus (H'_1(1) \cup H'_1(2))$ .

We still split the proof into three cases. The cases  $3|H'_1(2)| \geq |H_1|$  and  $3|H'_1(3)| \geq |H_1|$  follow with the same proof as in the previous subsection. For the case  $3|H'_1(1)| \geq |H_1|$ , now let  $J' = J[S'] \cup J[V_1 \setminus S']$ . By Lemmas 6 and 8,

$$|H'_1(1)| \leq |J'| \cdot 3pn + n^2 e^{-\sqrt{\log n}} \cdot pn \frac{\sqrt{\log n}}{\log \log n}. \quad (8)$$

Recall we have the assumptions that  $3|H_1| \geq |\bar{H}_\pi|$ , so  $3|H'_1(1)| \geq |H_1| \geq \delta pn^3 / (3 \log n)$ . Hence by (8), we have that  $0.9|H'_1(1)| \leq |J'| \cdot 3pn$ . Then, by Lemma 20 and the definition of  $S'_1$ , we get that  $\Delta(J') \leq \varepsilon_1 n$ . Finally, by Lemma 15,  $|\bar{H}_\pi| \geq 30pn|J'| \geq 9|H'_1(1)| \geq 3|H_1|$ .  $\square$

## References

- [1] László Babai, Miklós Simonovits, and Joel Spencer, *Extremal subgraphs of random graphs*, Journal of Graph Theory **14** (1990), no. 5, 599–622.
- [2] József Balogh, Jane Butterfield, Ping Hu, and John Lenz, *Mantel’s theorem for random hypergraphs*, Random Structures & Algorithms **48** (2016), no. 4, 641–654.
- [3] József Balogh, Robert Morris, and Wojciech Samotij, *Independent sets in hypergraphs*, Journal of the American Mathematical Society **28** (2015), no. 3, 669–709.
- [4] Béla Bollobás, *Three-graphs without two triples whose symmetric difference is contained in a third*, Discrete Mathematics **8** (1974), no. 1, 21–24.
- [5] David Conlon and Timothy Gowers, *Combinatorial theorems in sparse random sets*, Annals of Mathematics (2016), 367–454.
- [6] Bobby DeMarco and Jeff Kahn, *Mantel’s theorem for random graphs*, Random Structures & Algorithms **47** (2015), no. 1, 59–72.
- [7] ———, *Turán’s theorem for random graphs*, arXiv:1501.01340 (2015).
- [8] Peter Frankl and Zoltán Füredi, *A new generalization of the Erdős-Ko-Rado theorem*, Combinatorica **3** (1983), no. 3, 341–349.
- [9] Peter Keevash and Dhruv Mubayi, *Stability theorems for cancellative hypergraphs*, Journal of Combinatorial Theory, Series B **92** (2004), no. 1, 163–175.
- [10] Willem Mantel, *Problem 28*, Wiskundige Opgaven (1907), 60–61.
- [11] Wojciech Samotij, *Stability results for random discrete structures*, Random Structures & Algorithms **44** (2014), no. 3, 269–289.
- [12] David Saxton and Andrew Thomason, *Hypergraph containers*, Inventiones mathematicae **201** (2015), no. 3, 925–992.

- [13] Mathias Schacht, *Extremal results for random discrete structures*, Annals of Mathematics (2016), 333–365.
- [14] Paul Turán, *Eine Extremalaufgabe aus der Graphentheorie*, Mat. Fiz. Lapok **48** (1941), no. 61, 436–452.

## Appendix: Final computations

Recall  $g(p, s, r, i) = n \binom{n}{s} \binom{n^2}{r} (p \binom{s}{i} p^i)^r$  and  $O$  is the set of  $(s, r, i)$  such that  $g(p, s, r, i) = o(n^{-5})$ , given  $p > C\sqrt{\log n}/n$ . In this appendix, we prove the following claims.

**Claim 23.** *For every positive integer  $s \leq \frac{\varepsilon_1 n}{\sqrt{\log n}}$ ,  $r = \frac{pn}{\sqrt{\log n}}$ , and  $i = \frac{pn \log \log n}{500\sqrt{\log n}}$ , we have  $(s, r, i) \in O$ .*

*Proof.* We have

$$\begin{aligned}
g(p, s, r, i) &= n \binom{n}{s} \binom{n^2}{\frac{pn}{\sqrt{\log n}} s} \left( p \binom{s}{\frac{pn \log \log n}{500\sqrt{\log n}}} p^{\frac{pn \log \log n}{500\sqrt{\log n}}} \right)^{\frac{pn}{\sqrt{\log n}} s} \\
&\leq n \left( \frac{en}{s} \right)^s \left( \frac{epn^2 \sqrt{\log n}}{pns} \left( \frac{500eps \sqrt{\log n}}{pn \log \log n} \right)^{\frac{pn \log \log n}{500\sqrt{\log n}}} \right)^{\frac{pn}{\sqrt{\log n}} s} \\
&\leq n \left( \frac{en}{s} \right)^s \left( \frac{en \sqrt{\log n}}{s} \left( \frac{500es \sqrt{\log n}}{n \log \log n} \right)^{\frac{pn \log \log n}{500\sqrt{\log n}}} \right)^{\frac{pn}{\sqrt{\log n}} s}. \tag{9}
\end{aligned}$$

There are three cases depending on  $s$ .

- If  $1 \leq s \leq \sqrt{n}$ .

In this case, (9) is at most

$$n(en)^s \left( en \sqrt{\log n} \left( \frac{\sqrt{\log n}}{\sqrt{n}} \right)^{\frac{C \log \log n}{500}} \right)^{Cs} \leq n(en)^s \left( \frac{1}{n} \right)^{Cs} = o(n^{-5}).$$

- If  $\sqrt{n} \leq s \leq \frac{n}{\log n}$ .

In this case, (9) is at most

$$\begin{aligned}
n \left( \frac{en}{s} \right)^s \left( \frac{en \sqrt{\log n}}{s} \left( \frac{s}{n} \right)^{100} \right)^{\frac{pn}{\sqrt{\log n}} s} &\leq n \left( \frac{en}{s} \right)^s \left( \frac{s}{n} \right)^{\frac{pn}{\sqrt{\log n}} s} \\
&\leq n \left( \frac{en}{s} \right)^s \left( \frac{s}{n} \right)^{Cs} \leq n \left( \frac{s}{n} \right)^s \leq n \left( \frac{1}{\log n} \right)^{\sqrt{n}} = o(n^{-5}).
\end{aligned}$$



- If  $\frac{n}{\log n} \leq s \leq \frac{\varepsilon_1 n}{\sqrt{\log n}}$ . In this case, (9) is at most

$$\begin{aligned} n(e \log n)^s \left( e \log^{3/2} n \left( \frac{500e\varepsilon_1}{\log \log n} \right)^{\frac{C \log \log n}{500}} \right)^{\frac{pn}{\sqrt{\log n}} s} &\leq n(e \log n)^s \left( \left( \frac{1}{\log \log n} \right)^{\log \log n} \right)^{Cs} \\ &\leq n \left( \frac{1}{\log n} \right)^s \leq n \left( \frac{1}{\log n} \right)^{\frac{n}{\log n}} = o(n^{-5}). \quad \square \end{aligned}$$

**Claim 24.** For every positive integer  $s \in (\frac{\varepsilon_1 n}{\sqrt{\log n}}, \varepsilon_1 n]$ ,  $r = \frac{pn}{500}s$ , and  $i = 3\varepsilon_1 pn$ , we have  $(s, r, i) \in O$ .

*Proof.* We have that  $g(p, s, r, i)$  is

$$\begin{aligned} n \binom{n}{s} \binom{n^2}{\frac{pn}{500}s} \left( p \binom{s}{3\varepsilon_1 pn} p^{3\varepsilon_1 pn} \right)^{\frac{pn}{500}s} &\leq n \left( \frac{en}{s} \right)^s \left( \frac{500pn^2}{pns} \left( \frac{eps}{3\varepsilon_1 pn} \right)^{3\varepsilon_1 pn} \right)^{\frac{pn}{500}s} \\ &= n \left( \frac{en}{s} \right)^s \left( \frac{500n}{s} \left( \frac{es}{3\varepsilon_1 n} \right)^{3\varepsilon_1 pn} \right)^{\frac{pn}{500}s} \leq n \left( \frac{e\sqrt{\log n}}{\varepsilon_1} \right)^s \left( \frac{500\sqrt{\log n}}{\varepsilon_1} \left( \frac{e}{3} \right)^{3\varepsilon_1 C\sqrt{\log n}} \right)^{Cs\sqrt{\log n}} \\ &\leq n \left( \frac{e\sqrt{\log n}}{\varepsilon_1} \right)^s \left( 2^{-\sqrt{\log n}} \right)^{Cs\sqrt{\log n}} \leq n2^{-s} \leq n2^{-\frac{\varepsilon_1 n}{\sqrt{\log n}}} = o(n^{-5}). \quad \square \end{aligned}$$

**Claim 25.** For every positive integer  $s \leq n$ , we have  $(s, \frac{pn^2}{\log n}, 3pn) \in O$ .

*Proof.* We have that  $g(p, s, r, i)$  is

$$\begin{aligned} n \binom{n}{s} \binom{n^2}{\frac{pn^2}{\log n}} \left( p \binom{s}{3pn} p^{3pn} \right)^{\frac{pn^2}{\log n}} &\leq n2^n \binom{n^2}{\frac{pn^2}{\log n}} \left( p \binom{n}{3pn} p^{3pn} \right)^{\frac{pn^2}{\log n}} \\ &\leq n2^n \left( \frac{epn^2 \log n}{pn^2} \left( \frac{enp}{3pn} \right)^{3pn} \right)^{\frac{pn^2}{\log n}} \leq n2^n \left( e \log n \left( \frac{e}{3} \right)^{3pn} \right)^{\frac{pn^2}{\log n}} \\ &\leq n2^n \left( e \log n \left( \frac{e}{3} \right)^{3C\sqrt{\log n}} \right)^{\frac{Cn}{\sqrt{\log n}}} \leq n2^n \left( \left( \frac{e}{3} \right)^{\sqrt{\log n}} \right)^{\frac{Cn}{\sqrt{\log n}}} = n2^n \left( \frac{e}{3} \right)^{Cn} = o(n^{-5}). \quad \square \end{aligned}$$

**Claim 26.** For every positive integer  $s \leq \frac{\varepsilon_3 n}{\sqrt{\log n}}$ , we have  $(s, \frac{\varepsilon_2 pn^2}{2}, \frac{3\varepsilon_3 pn}{\sqrt{\log n}}) \in O$ .

*Proof.* We have that  $g(p, s, r, i)$  is

$$\begin{aligned}
& n \binom{n}{s} \binom{n^2}{\frac{1}{2}\varepsilon_2 p n^2} \left( p \binom{s}{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} p^{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \leq n \binom{n}{\frac{\varepsilon_3 n}{\sqrt{\log n}}} \binom{n^2}{\frac{1}{2}\varepsilon_2 p n^2} \left( p \binom{\frac{\varepsilon_3 n}{\sqrt{\log n}}}{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} p^{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \\
& \leq n \left( \frac{en\sqrt{\log n}}{\varepsilon_3 n} \right)^{\frac{\varepsilon_3 n}{\sqrt{\log n}}} \left( \frac{2epn^2}{\varepsilon_2 p n^2} \left( \frac{e\varepsilon_3 p n}{3\varepsilon_3 p n} \right)^{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \\
& = n \left( \frac{e\sqrt{\log n}}{\varepsilon_3} \right)^{\frac{\varepsilon_3 n}{\sqrt{\log n}}} \left( \frac{2e}{\varepsilon_2} \left( \frac{e}{3} \right)^{\frac{3\varepsilon_3 p n}{\sqrt{\log n}}} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \\
& \leq n \left( \frac{e\sqrt{\log n}}{\varepsilon_3} \right)^{\frac{\varepsilon_3 n}{\sqrt{\log n}}} 2^{-\frac{1}{2}\varepsilon_2 p n^2} \leq n \left( \frac{e\sqrt{\log n}}{\varepsilon_3} \right)^{\frac{\varepsilon_3 n}{\sqrt{\log n}}} 2^{-\frac{1}{2}\varepsilon_2 C n \sqrt{\log n}} = o(n^{-5}). \quad \square
\end{aligned}$$

**Claim 27.** For every positive integer  $s \leq \varepsilon_3 n$ , we have  $\left(s, \frac{\varepsilon_2 p n^2}{2}, 3\varepsilon_3 p n\right) \in O$ .

*Proof.* We have that  $g(p, s, r, i)$  is

$$\begin{aligned}
& n \binom{n}{s} \binom{n^2}{\frac{1}{2}\varepsilon_2 p n^2} \left( p \binom{s}{3\varepsilon_3 p n} p^{3\varepsilon_3 p n} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \leq n \binom{n}{\varepsilon_3 n} \binom{n^2}{\frac{1}{2}\varepsilon_2 p n^2} \left( p \binom{\varepsilon_3 n}{3\varepsilon_3 p n} p^{3\varepsilon_3 p n} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \\
& \leq n \left( \frac{en}{\varepsilon_3 n} \right)^{\varepsilon_3 n} \left( \frac{2epn^2}{\varepsilon_2 p n^2} \left( \frac{e\varepsilon_3 p n}{3\varepsilon_3 p n} \right)^{3\varepsilon_3 p n} \right)^{\frac{1}{2}\varepsilon_2 p n^2} = n \left( \frac{e}{\varepsilon_3} \right)^{\varepsilon_3 n} \left( \frac{2e}{\varepsilon_2} \left( \frac{e}{3} \right)^{3\varepsilon_3 p n} \right)^{\frac{1}{2}\varepsilon_2 p n^2} \\
& \leq n 2^{-n\sqrt{\log n}} = o(n^{-5}). \quad \square
\end{aligned}$$