

A new infinite family of irregular algebraic surfaces with canonical map of degree 8

NGUYEN BIN

April 22, 2022

Abstract

In this note we construct an unlimited family of irregular algebraic surfaces of general type with canonical map of degree 8, irregularity 1 and arbitrarily large geometric genus such that the image of the canonical map is not a surface of minimal degree.

1 Introduction

Let X be a minimal smooth complex surface of general type and denote by $\varphi_{|K_X|} : X \dashrightarrow \mathbb{P}^{p_g(X)-1}$ the canonical map of X , where K_X is the canonical divisor of X and $p_g(X) = \dim H^0(X, K_X)$ is the geometric genus. The existence of surfaces of general type with non-birational canonical map has been studied intensively by many authors in the last decades. This problem is motivated by the work of A. Beauville [1]. We refer to the recent preprint by M. Mendes Lopes and R. Pardini [6] on the subject. We restrict our attention to the existence of surfaces of general type with canonical map of degree 8. It is known that for surfaces of general type, the degree d of the canonical map is at most 9 if the holomorphic Euler-Poincaré characteristic $\chi(\mathcal{O}_X)$ is bigger than or equal to 31. This was proven by A. Beauville in [1]. In 1986, G. Xiao improved this result by showing that the degree of the canonical map is less than or equal to 8 if the geometric genus of the surface is bigger than 132 [11]. The first unlimited family of surfaces with canonical map of degree 8 was found by A. Beauville [8]. These examples were constructed as double covers of the product surface of a non-hyperelliptic curve of genus 3 and the rational curve \mathbb{P}^1 . In [8], some unlimited families of surfaces with canonical map of degree 8 were found by as \mathbb{Z}_2^3 -covers of the first Hirzebruch surface \mathbb{F}_1 or its blow-up in a point. In these all known examples, the image of the canonical map is a surface of

Mathematics Subject Classification (2010): 14J29.

Key words: Surfaces of general type, Canonical maps, Abelian covers.

minimal degree. In this note we construct an unlimited family of algebraic surfaces with $d = 8$ and arbitrarily large p_g such that the image of the canonical map is not of minimal degree.

Theorem 1. *Let n be an integer with $n \geq 3$. There exist minimal surfaces of general type X satisfying*

K_X^2	$p_g(X)$	$q(X)$	$\deg(\text{im } \varphi_{ K_X })$	$ K_X $
$16n$	$2n$	1	$2n$	is base point free

such that the canonical map $\varphi_{|K_X|}$ has degree 8.

In the above theorem, $q(X) = \dim H^1(X, K_X)$ is the irregularity of X . These surfaces are constructed as \mathbb{Z}_2^3 -covers of the product surface of a smooth elliptic curve C and the rational curve \mathbb{P}^1 . The building data $\{L_\chi, D_\sigma\}_{\chi, \sigma}$ of \mathbb{Z}_2^3 -covers (see Section 2) are chosen such that there is a character χ' of \mathbb{Z}_2^3 with arbitrarily large $h^0(L_{\chi'} + K_{\mathbb{P}^1 \times C})$ and that $h^0(L_\chi + K_{\mathbb{P}^1 \times C})$ vanishes for all other characters χ of \mathbb{Z}_2^3 . From the decomposition of the space of 2-forms of the surfaces (see Proposition 2)

$$H^0(X, K_X) = H^0(\mathbb{P}^1 \times C, K_{\mathbb{P}^1 \times C}) \oplus \bigoplus_{\chi \neq \chi_{000}} H^0(\mathbb{P}^1 \times C, K_{\mathbb{P}^1 \times C} + L_\chi),$$

such a choice of the building data allows to conclude that the canonical map of X factors through the \mathbb{Z}_2^3 -cover. Furthermore, we choose the divisor $L_{\chi'}$ in $\mathbb{P}^1 \times C$ such that $L_{\chi'} + K_{\mathbb{P}^1 \times C} \equiv E + \sum_{i=1}^n F_{ii}$, where E is a general elliptic fiber and F_{ii} are distinct rational fibres of $\mathbb{P}^1 \times C$. This leads to the fact that the map $\varphi_{|E + \sum_{i=1}^n F_{ii}|}$ is of degree one for all $n \geq 3$. Thus, the canonical map of X is of degree 8. We notice that the covers of such a product guarantees the irregularity of the result by pulling back 1-forms.

In our construction, if $n = 2$, we obtain a smooth minimal surface of general type X with $K_X^2 = 32, p_g(X) = 4, q(X) = 1$ and $d = 16$ since the linear system $|E + F_{11} + F_{22}|$ is a map of degree 2 onto $\mathbb{P}^1 \times \mathbb{P}^1$. This surface was constructed as \mathbb{Z}_2^4 -cover of $\mathbb{P}^1 \times \mathbb{P}^1$ (see [7]). It is worth pointing out that C. Gleissner, R. Pignatelli and C. Rito constructed a family of surfaces with $K_X^2 = 24, p_g(X) = 3, q(X) = 1$ and $d = 24$ ([3]). Their example has a very similar construction as \mathbb{Z}_2^3 -cover of $\mathbb{P}^1 \times C$ branched on “fibers” of the obvious trivial fibrations.

Throughout this note all surfaces are projective algebraic over the complex numbers. The linear equivalence of divisors is denoted by \equiv . A character χ of the group G is a homomorphism from G to \mathbb{C}^* , the multiplicative group of the non-zero complex numbers. The rest of the notation is standard in algebraic geometry.

2 \mathbb{Z}_2^3 -coverings

The construction of abelian covers was studied by R. Pardini in [9]. For details about the building data of abelian covers and their notations, we refer the reader to

Section 1 and Section 2 of R. Pardini's work ([9]). For the sake of completeness, we recall some facts on \mathbb{Z}_2^3 -covers, in a form which is convenient for our later constructions.

We denote by $\chi_{j_1 j_2 j_3}$ the character of \mathbb{Z}_2^3 defined by

$$\chi_{j_1 j_2 j_3}(a_1, a_2, a_3) := e^{(\pi a_1 j_1)\sqrt{-1}} e^{(\pi a_2 j_2)\sqrt{-1}} e^{(\pi a_3 j_3)\sqrt{-1}}$$

for all $j_1, j_2, j_3, a_1, a_2, a_3 \in \mathbb{Z}_2$. A \mathbb{Z}_2^3 -cover $X \longrightarrow Y$ can be determined by a collection of non-trivial divisors L_χ labelled by characters of \mathbb{Z}_2^3 and effective divisors D_σ labelled by non-trivial elements of \mathbb{Z}_2^3 of the surface Y . More precisely, from [9, Theorem 2.1] we can define \mathbb{Z}_2^3 -covers as follows:

Proposition 1. *Given Y a smooth projective surface, let L_χ be divisors of Y such that $L_\chi \not\equiv \mathcal{O}_Y$ for all non-trivial characters χ of \mathbb{Z}_2^3 and let D_σ be effective divisors of Y for all $\sigma \in \mathbb{Z}_2^3 \setminus \{(0, 0, 0)\}$ such that the total branch divisor $B := \sum_{\sigma \neq 0} D_\sigma$ is reduced. Then $\{L_\chi, D_\sigma\}_{\chi, \sigma}$ is the building data of a \mathbb{Z}_2^3 -cover $f : X \longrightarrow Y$ if and only if*

$$L_\chi + L_{\chi'} \equiv L_{\chi\chi'} + \sum_{\chi(\sigma)=\chi'(\sigma)=-1} D_\sigma \quad (1)$$

for all non-trivial characters χ, χ' of \mathbb{Z}_2^3 .

For the reader's convenience, we leave here the relations (1) of the reduced building data of \mathbb{Z}_2^3 -covers:

$$\begin{array}{llllll} L_{100} + L_{100} & \equiv & & D_{100} & +D_{101} & +D_{110} & +D_{111} \\ L_{100} + L_{010} & \equiv & & & & D_{110} & +D_{111} & +L_{110} \\ L_{100} + L_{001} & \equiv & & & D_{101} & & +D_{111} & +L_{101} \\ L_{010} + L_{010} & \equiv & D_{010} & +D_{011} & & D_{110} & +D_{111} & \\ L_{010} + L_{001} & \equiv & & D_{011} & & & +D_{111} & +L_{011} \\ L_{001} + L_{001} & \equiv & D_{001} & +D_{011} & +D_{101} & & +D_{111}. & \end{array}$$

By [9, Theorem 3.1] if each branch component D_σ is smooth and the total branch locus B is a simple normal crossings divisor, the surface X is smooth.

Also from [9, Lemma 4.2, Proposition 4.2] we have:

Proposition 2. *If Y is a smooth surface and $f : X \longrightarrow Y$ is a smooth \mathbb{Z}_2^3 -cover with the building data $\{L_\chi, D_\sigma\}_{\chi, \sigma}$, the surface X satisfies the following:*

$$\begin{aligned} 2K_X &\equiv f^* \left(2K_Y + \sum_{\sigma \neq 0} D_\sigma \right); \\ f_* \mathcal{O}_X &= \mathcal{O}_Y \oplus \bigoplus_{\chi \neq \chi_{000}} L_\chi^{-1}. \end{aligned}$$

This implies that

$$\begin{aligned}
H^0(X, K_X) &= H^0(Y, K_Y) \oplus \bigoplus_{\chi \neq \chi_{000}} H^0(Y, K_Y + L_\chi); \\
K_X^2 &= 2 \left(2K_Y + \sum_{\sigma \neq 0} D_\sigma \right)^2; \\
p_g(X) &= p_g(Y) + \sum_{\chi \neq \chi_{000}} h^0(L_\chi + K_Y); \\
\chi(\mathcal{O}_X) &= 8\chi(\mathcal{O}_Y) + \sum_{\chi \neq \chi_{000}} \frac{1}{2} L_\chi (L_\chi + K_Y).
\end{aligned}$$

Moreover, the canonical linear system $|K_X|$ is generated by

$$f^*(|K_Y + L_\chi|) + \sum_{\chi(\sigma)=1} R_\sigma, \quad \forall \chi \in J$$

where $J := \{\chi' : |K_Y + L_{\chi'}| \neq \emptyset\}$ and R_σ is the reduced divisor supported on $f^*(D_\sigma)$.

For the proof of the last statement of Proposition 2, we refer the reader to [3, Page 3].

3 Construction

Throughout this section, we denote by $Y := \mathbb{P}^1 \times C$ the product surface of the rational curve \mathbb{P}^1 and a smooth elliptic curve C . Let $p_1 : \mathbb{P}^1 \times C \longrightarrow \mathbb{P}^1$ be the projection of the product surface $\mathbb{P}^1 \times C$ on \mathbb{P}^1 . We denote by E a general fiber of p_1 . The canonical class of Y is $K_Y \equiv -2E$.

Let E_1, E_2, \dots, E_6 be distinct elliptic fibres and let $F_1, F_2, \dots, F_n, F'_1, F'_2, \dots, F'_n, F''_1, F''_2, F''_3$ be distinct rational fibres (with $n \geq 3$) such that $2F'_1 \equiv 2F''_2 \equiv 2F''_3$. Because the sum of two points in an elliptic curve is divisible by 2 in the Picard group, there are fibres F_{ii} such that $2F_{ii} \equiv F_i + F'_i$, for all $i \in \{1, 2, \dots, n\}$. We consider the following divisors

$$\begin{aligned}
D_{100} &:= E_1 + E_2, & D_{101} &:= E_3 + E_4, & D_{110} &:= E_5 + E_6, & D_{111} &:= \sum_{i=1}^n (F_i + F'_i) \\
L_{100} &:= 3E + \sum_{i=1}^n F_{ii} & L_{010} &:= E + \sum_{i=1}^n F_{ii} + \eta_1 & L_{001} &:= E + \sum_{i=1}^n F_{ii} + \eta_2 & L_{110} &:= 2E + \eta_1 \\
L_{101} &:= 2E + \eta_2 & L_{011} &:= 2E + \eta_3 & L_{111} &:= E + \sum_{i=1}^n F_{ii} + \eta_3,
\end{aligned}$$

where $\eta_1 := F_1'' - F_2''$, $\eta_2 := F_2'' - F_3''$, $\eta_3 := F_1'' - F_3'' = \eta_1 + \eta_2$ are non-trivial 2-torsions. These divisors D_σ, L_χ satisfy the following relations

$$\begin{aligned}
L_{100} + L_{100} &\equiv D_{100} + D_{101} + D_{110} + D_{111} &&\equiv 6E + \sum_{i=1}^n 2F_{ii} \\
L_{100} + L_{010} &\equiv &&D_{110} + D_{111} + L_{110} &&\equiv 4E + \sum_{i=1}^n 2F_{ii} + \eta_1 \\
L_{100} + L_{001} &\equiv &&D_{101} + D_{111} + L_{101} &&\equiv 4E + \sum_{i=1}^n 2F_{ii} + \eta_2 \\
L_{010} + L_{010} &\equiv &&D_{110} + D_{111} &&\equiv 2E + \sum_{i=1}^n 2F_{ii} \\
L_{010} + L_{001} &\equiv &&+ D_{111} + L_{011} &&\equiv 2E + \sum_{i=1}^n 2F_{ii} + \eta_3 \\
L_{001} + L_{001} &\equiv &&D_{101} + D_{111} &&\equiv 2E + \sum_{i=1}^n 2F_{ii}.
\end{aligned}$$

Thus by Proposition 1, the divisors D_σ, L_χ define a \mathbb{Z}_2^3 -cover $f : X \longrightarrow Y$. Because each branch component D_σ is smooth and the total branch locus B is a normal crossings divisor, the surface X is smooth. Moreover, by Proposition 2, the surface X satisfies the following

$$2K_X \equiv f^* \left(2E + \sum_{i=1}^n (F_i + F_i') \right).$$

We notice that a surface is of general type and minimal if the canonical divisor is big and nef (see e.g. [5, Section 2]). Since the divisor $2K_X$ is the pull-back of a nef and big divisor, the canonical divisor K_X is nef and big. Thus, the surface X is of general type and minimal. Moreover, by Proposition 2 the invariants of X are as follows

$$K_X^2 = 16n, p_g(X) = 2n, \chi(\mathcal{O}_X) = 2n, q(X) = 1.$$

We show that the canonical map is of degree 8. By Proposition 2, we have the following decomposition

$$H^0(X, K_X) = H^0(Y, K_Y) \oplus \bigoplus_{\chi \neq \chi_{000}} H^0(Y, K_Y + L_\chi).$$

Moreover, the choice of L_χ 's yields that

$$h^0(L_\chi + K_Y) = 0$$

for all $\chi \neq \chi_{100}$. By Proposition 2, the linear system $|K_X|$ is generated by

$$f^*(K_Y + L_{100}) = f^* \left(E + \sum_{i=1}^n F_{ii} \right).$$

This implies that the canonical map of the surface X factors through the \mathbb{Z}_2^3 -cover $f : X \longrightarrow Y$. Thus the following diagram commutes

$$\begin{array}{ccc}
 X & \xrightarrow[\quad f \quad]{\mathbb{Z}_2^3} & Y \\
 & \searrow \varphi|_{K_X} & \downarrow \varphi|_{E + \sum_{i=1}^n F_{ii}} \\
 & & \mathbb{P}^{2n-1}.
 \end{array}$$

Since the map $\varphi|_{E + \sum_{i=1}^n F_{ii}}$ is of degree one for all $n \geq 3$, the canonical map of X is of degree 8.

Remark 1. Let $Z := X/\Gamma$ be the quotient surface of X , where $\Gamma := \langle (0, 0, 1), (0, 1, 0) \rangle$ is the subgroup of \mathbb{Z}_2^3 . The surface Z is a surface of general type whose only singularities are $12n$ nodes. Moreover, the canonical map $\varphi|_{K_Z}$ is a map of degree 2 (see [4, Theorem 5.1] and [2, Theorem 1.1]). The canonical map $\varphi|_{K_X}$ is the composition of the degree 4 quotient map $X \longrightarrow Z := X/\Gamma$ with the canonical map $\varphi|_{K_Z}$ of Z (see e.g. [10, Example 2.1]).

Remark 2. In the above construction, there are four different possible choices for each F_{ii} . A different choice produces a different surface X .

The previous observation leads to the following interesting question:

Question 1. *Are the surfaces in Remark 2 deformation equivalent?*

Acknowledgments

The author is deeply indebted to Margarida Mendes Lopes for all her help. Thanks are also due to Jungkai Alfred Chen for many interesting conversations. The author would like to express his gratitude to the anonymous referee for his/her thorough reading of the paper and valuable suggestion. This paper was finished during the author's postdoctoral fellowship at the National Center for Theoretical Sciences (NCTS), Taiwan, under the grant MOST 110-2123-M-002-005. The author would like to thank NCTS for the financial support and kind hospitality.

References

- [1] BEAUVILLE, A. L'application canonique pour les surfaces de type général. *Invent. Math.* 55, 2 (1979), 121–140.
- [2] CILIBERTO, C., MENDES LOPES, M., AND PARDINI, R. The classification of minimal irregular surfaces of general type with $K^2 = 2p_g$. *Algebr. Geom.* 1, 4 (2014), 479–488.

- [3] GLEISSNER, C., PIGNATELLI, R., AND RITO, C. New surfaces with canonical map of high degree. *arXiv e-prints* (July 2018), arXiv:1807.11854.
- [4] HORIKAWA, E. Algebraic surfaces of general type with small c_1^2 . V. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* 28, 3 (1981), 745–755 (1982).
- [5] LOPES, M. M., AND PARDINI, R. The geography of irregular surfaces. In *Current developments in algebraic geometry*, vol. 59 of *Math. Sci. Res. Inst. Publ.* Cambridge Univ. Press, Cambridge, 2012, pp. 349–378.
- [6] LOPES, M. M., AND PARDINI, R. On the degree of the canonical map of a surface of general type, 2021. *arXiv:2103.01912*.
- [7] NGUYEN BIN A new example of an algebraic surface with canonical map of degree 16. *Arch. Math. (Basel)* 113, 4 (2019), 385–390.
- [8] NGUYEN BIN Some unlimited families of minimal surfaces of general type with the canonical map of degree 8. *Manuscripta Math.* 163, 1-2 (2020), 13–25.
- [9] PARDINI, R. Abelian covers of algebraic varieties. *J. Reine Angew. Math.* 417 (1991), 191–213.
- [10] PARDINI, R. Canonical images of surfaces. *J. Reine Angew. Math.* 417 (1991), 215–219.
- [11] XIAO, G. Algebraic surfaces with high canonical degree. *Math. Ann.* 274, 3 (1986), 473–483.

Mathematics Division,
National Center for Theoretical Sciences,
National Taiwan University,
Taiwan.
E-mail address: `nguyenbin@ncts.ntu.edu.tw`