SEPARATING MANY LOCALISATION CARDINALS ON THE GENERALISED BAIRE SPACE

Tristan van der Vlugt^{*}

March 5, 2022

Abstract

Given a cofinal cardinal function $h \in {}^{\kappa}\kappa$ for κ inaccessible, we consider the dominating *h*-localisation number, that is, the least cardinality of a dominating set of *h*-slaloms such that every κ -real is localised by a slalom in the dominating set. It was proved in [BBTFM18] that the dominating localisation numbers can be consistently different for two functions *h* (the identity function and the power function). We will construct a κ -sized family of functions *h* and their corresponding localisation numbers, and use a $\leq \kappa$ -supported product of a cofinality-preserving forcing to prove that any simultaneous assignment of these localisation numbers to cardinals above κ is consistent. This answers an open question from [BBTFM18].

In an effort to generalise the cardinal characteristics related to the null ideal from the context of the continuum ${}^{\omega}\omega$ to the generalised Baire space ${}^{\kappa}\kappa$, the authors of [BBTFM18] considered localisation cardinals. These cardinals were first described in the context of ${}^{\omega}\omega$ by Tomek Bartoszyński [Bar87], and are defined using slaloms. Let κ be a regular strong limit cardinal (hence κ is inaccessible or equal to ω) and let $h \in {}^{\kappa}\kappa$. An *h*-slalom is any function $\varphi : \kappa \to [\kappa]^{<\kappa}$ such that $|\varphi(\alpha)| \leq |h(\alpha)|$ for all $\alpha \in \kappa$.

For $f \in {}^{\kappa}\kappa$, we say $f \in {}^{*}\varphi$, or f is **localised** by φ , if there exists some $\xi < \kappa$ such that $f(\alpha) \in \varphi(\alpha)$ for all $\alpha \in [\xi, \kappa)$. We denote the set of all h-slaloms by Loc_h .

Using these concepts, we can define the following two cardinal characteristics, sometimes called **localisation cardinals**:

 $\mathfrak{b}_{\kappa}^{h}(\in^{*}) = \text{the least cardinality of a family } \mathcal{F} \subseteq {}^{\kappa}\kappa \text{ such that } \forall \varphi \in \operatorname{Loc}_{h} \exists f \in \mathcal{F}(f \notin^{*} \varphi),$

 $\mathfrak{d}^h_{\kappa}(\in^*)$ = the least cardinality of a family $\Phi \subseteq \operatorname{Loc}_h$ such that $\forall f \in {}^{\kappa}\kappa \exists \varphi \in \Phi(f \in^* \varphi)$.

In the case that $\kappa = \omega$ these cardinals give a combinatorial definition of two of the cardinal invariants of the Lebesgue null ideal \mathcal{N} :

 $\operatorname{add}(\mathcal{N}) = \operatorname{the} \operatorname{least} \operatorname{cardinality} \operatorname{of} \operatorname{a} \operatorname{family} \mathcal{A} \subseteq \mathcal{N} \operatorname{such} \operatorname{that} \bigcup \mathcal{A} \notin \mathcal{N},$

 $\operatorname{cof}(\mathcal{N}) = \operatorname{the} \operatorname{least} \operatorname{cardinality} \operatorname{of} \operatorname{a} \operatorname{family} \mathcal{C} \subseteq \mathcal{N} \operatorname{such} \operatorname{that} \forall N \in \mathcal{N} \exists C \in \mathcal{C}(N \subseteq C)$

Bartoszyński introduced slaloms in [Bar87] to give the following combinatorial definition to $\operatorname{add}(\mathcal{N})$ and $\operatorname{cof}(\mathcal{N})$.

Fact 1

$$\operatorname{add}(\mathcal{N}) = \mathfrak{b}^h_{\omega}(\in^*) \text{ and } \operatorname{cof}(\mathcal{N}) = \mathfrak{d}^h_{\omega}(\in^*).$$

^{*}UNIVERSITÄT HAMBURG, FACHBEREICH MATHEMATIK – tristan.van.der.vlugt@uni-hamburg.de – The author wishes to thank Jörg Brendle for valuable discussions, suggestions and extensive comments.

The choice of $h \in {}^{\omega}\omega$ is irrelevant here, as it does not influence the cardinality of $\mathfrak{b}^{h}_{\omega}(\in^{*})$ and $\mathfrak{d}^{h}_{\omega}(\in^{*})$. This turns out to be different in the generalised case. Recently, it was proved in [BBTFM18] that $\mathfrak{d}^{h}_{\kappa}(\in^{*})$ can consistently have different values for different $h \in {}^{\kappa}\kappa$. In particular, it was shown that $\mathfrak{d}^{pow}_{\kappa}(\in^{*}) < \mathfrak{d}^{id}_{\kappa}(\in^{*})$ is consistent, where pow : $\alpha \mapsto 2^{\alpha}$ and id : $\alpha \mapsto \alpha$.

This consistency was proved using a generalisation of Sacks forcing on κ^2 that has the generalised pow-Sacks property, but not the generalised id-Sacks property.

Definition 2

Let $h \in {}^{\kappa}\kappa$. A forcing \mathbb{P} has the **generalised** h-Sacks property if for every \mathbb{P} -name \dot{f} and $p \in \mathbb{P}$ such that $p \Vdash ``\dot{f} : \check{\kappa} \to \check{\kappa}$ " there exists an h-slalom φ and $q \leq p$ such that $q \Vdash ``\dot{f}(\check{\alpha}) \in \check{\varphi}(\check{\alpha})$ " for all $\alpha < \kappa$. We will from now on simply say h-Sacks property and omit "generalised".

Hence, if $\Phi \subseteq \operatorname{Loc}_h$ is a family as in the definition of $\mathfrak{d}^h_{\kappa}(\in^*)$ in the ground model, then Φ will still witness the size of $\mathfrak{d}^h_{\kappa}(\in^*)$ in the extension by a forcing with the *h*-Sacks property. Therefore, the generalised Sacks forcing is unable to increase the size of $\mathfrak{d}^{\mathsf{pow}}_{\kappa}(\in^*)$, but it is shown to be possible to increase the size of $\mathfrak{d}^{\mathsf{pd}}_{\kappa}(\in^*)$ by using an iteration or a product of the forcing.

In this text we will answer an open question from [BBTFM18] and prove that there exist κ many functions $h_{\xi} \in {}^{\kappa}\kappa$ and cardinals $\lambda_{\xi} > \kappa$ such that it is simultaneously consistent that $\mathfrak{d}_{\kappa}^{h_{\xi}}(\in^*) = \lambda_{\xi}$ for all $\xi < \kappa$. The strategy will be the same, in that we consider a product of Sacks-like forcings that have the h_{ξ} -Sacks property, but not the h_{η} -Sacks property for different ξ and η . In the first section, we introduce the Sacks-like forcing \mathbb{S}_{κ}^{h} and prove that it preserves cardinals and cofinalities and use fusion to show that it satisfies certain Sacks properties. In the second section we consider products of such forcings, we show that properties such as the preservation of cardinals and cofinalities and the relevant Sacks properties are preserved under $\leq \kappa$ -support products and we use this to prove the consistency of our result.

For the sake of brevity, from now on we will assume that κ denotes a strongly inaccessible cardinal. We will also fix the convention that $h, H, F \in {}^{\kappa}\kappa$ denote increasing cofinal cardinal functions (i.e. ran(h) is cofinal in κ and $h(\alpha)$ is a cardinal for each $\alpha < \kappa$). This convention extends to h_{ξ} , F_0 , and other subscripts.

The Forcing

Let us establish some notation to discuss trees on $\kappa \kappa$ before we define our forcing notion.

Let $T \subseteq {}^{<\kappa}\kappa$ be a tree, then we let [T] denote the set of branches of T. Usually $[T] \subseteq {}^{\kappa}\kappa$. We call $u \in T$ a **non-splitting** node (of T) if there is a unique $\beta < \kappa$ such that $u^{\frown}\beta \in T$ and otherwise we call u an α -splitting node (of T), where $\alpha \leq |\{\beta < \kappa \mid u^{\frown}\beta \in T\}|$. We say u is a **splitting** node if it is a 2-splitting node. If u is an α -splitting node, but not a ξ splitting node for any ξ with $|\xi| > |\alpha|$, then we say that u is a **sharp** α -splitting node. We let $u \in \text{Split}_{\alpha}(T)$ iff u is splitting and $\operatorname{ot}(\{\beta < \operatorname{ot}(u) \mid u \restriction \beta \text{ is splitting}\}) = \alpha$. If $u \in T$ is splitting, let $\operatorname{suc}(u, T) = \{v \in T \mid \exists \beta < \kappa(v = u^{\frown}\beta)\}$.

Definition 3

The conditions of the forcing \mathbb{S}^h_{κ} are trees $T \subseteq {}^{<\kappa}\kappa$ that satisfy the following properties:

- (i) for any $u \in T$ there exists $v \in T$ such that $u \subseteq v$ and v is splitting,
- (ii) if $u \in \text{Split}_{\alpha}(T)$, then u is an $h(\alpha)$ -splitting node in T,
- (iii) if $\langle u_{\alpha} \mid \alpha < \gamma \rangle$ is an increasing sequence of splitting nodes, then $\bigcup_{\alpha < \gamma} u_{\alpha}$ is splitting.

The order is defined as $T \leq_{\mathbb{S}_{\kappa}^{h}} S$ (here T is stronger than S) iff $T \subseteq S$ and for every $\alpha < \kappa$ and $u \in T$ a splitting node, if $\operatorname{suc}(u,T) \neq \operatorname{suc}(u,S)$, then $|\operatorname{suc}(u,T)| < |\operatorname{suc}(u,S)|$. If the forcing notion is clear from context, we will write $T \leq S$ in place of $T \leq_{\mathbb{S}_{\kappa}^{h}} S$

Note that conditions (i) and (iii) imply that $[T] \subseteq {}^{\kappa}\kappa$ and that the set of splitting nodes is club in T. We naturally want \mathbb{S}^h_{κ} to preserve cardinalities. If we assume that $\mathbf{V} \models "2^{\kappa} = \kappa^+$ ", then it is clear that \mathbb{S}^h_{κ} has the $\langle \kappa^{++}$ -chain condition, since $|\mathbb{S}^h_{\kappa}| \leq |\mathcal{P}(\langle \kappa)| = 2^{\kappa} = \kappa^+$, where the former equality is implied by $\kappa^{\langle \kappa} = \kappa$, which in turn follows from κ being inaccessible. It is therefore clear that cardinalities above κ^+ are preserved under assumption of $\mathbf{V} \models "2^{\kappa} = \kappa^+$ ".

To preserve cardinalities less than or equal to κ , we show that \mathbb{S}^h_{κ} is $<\kappa$ -closed.

Lemma 4

 \mathbb{S}^h_{κ} is $\langle \kappa$ -closed. That is, for any $\lambda < \kappa$, if $\langle T_{\xi} | \xi < \lambda \rangle$ is a descending chain of conditions, then there exists a condition T such that $T \leq T_{\xi}$ for all $\xi < \lambda$.

Proof. The condition T that is below all T_{ξ} will be simply $T = \bigcap_{\xi} T_{\xi}$. Clearly T is a tree as well. We have to show that $T \in \mathbb{S}_{\kappa}^{h}$, which we do by veryfing points (i), (ii) and (iii) from Definition 3. We will make use of the following claim.

Claim. If $u \in T = \bigcap T_{\xi}$, then there is $\eta < \lambda$ such that $suc(u, T) = suc(u, T_{\xi})$ for all $\xi \in [\eta, \lambda)$.

Proof of claim. Suppose that $u \in T$, and let $\lambda_{\xi} = |\operatorname{suc}(u, T_{\xi})|$, then the ordering on \mathbb{S}^{h}_{κ} dictates that $\langle \lambda_{\xi} | \xi < \lambda \rangle$ is a descending sequence of cardinals, hence there is $\eta < \lambda$ such that $\lambda_{\xi} = \lambda_{\eta}$ for all $\xi \in [\eta, \lambda)$. But then $\operatorname{suc}(u, T_{\xi}) = \operatorname{suc}(u, T_{\eta})$ for all $\xi \in [\eta, \lambda)$ by the ordering of \mathbb{S}^{h}_{κ} .

(i) Let $u \in T$, and let $f \in [T]$ be a branch for which $u \subseteq f$. If $\operatorname{ot}(f) < \kappa$, then $f \in T_{\xi}$ for each ξ , thus by the claim there is some $\xi < \lambda$ for which $\operatorname{suc}(f, T_{\xi}) = \operatorname{suc}(f, T) = \emptyset$. Then clearly $T_{\xi} \notin \mathbb{S}_{\kappa}^{h}$, which is a contradiction, hence $\operatorname{ot}(f) = \kappa$. Let $S_{\xi} = \{\alpha \in [\operatorname{ot}(u), \kappa) \mid f \restriction \alpha \text{ is splitting in } T_{\xi}\}$, then since $T_{\xi} \in \mathbb{S}_{\kappa}^{h}$ satisfies (i) and (iii), we see that S_{ξ} is a club set. But then $\bigcap_{\xi < \lambda} S_{\xi}$ is club. Any $v \in \bigcap_{\xi < \lambda} S_{\xi}$ is splitting in all T_{ξ} , thus by the claim it is splitting in T, and by definition of S_{ξ} it follows that $u \subseteq v$.

(ii) If $u \in \text{Split}_{\alpha}(T)$, then by the claim there is $\xi < \lambda$ such that $\text{suc}(u, T_{\xi}) = \text{suc}(u, T)$. If $v \subseteq u$ is splitting in T, then v is splitting in T_{ξ} as well, therefore $u \in \text{Split}_{\beta}(T_{\xi})$ for some $\beta \geq \alpha$. This shows that u is $h(\beta)$ -splitting in T_{ξ} , hence it is $h(\beta)$ -splitting in T and by $h(\alpha) \leq h(\beta)$ we see that u is also $h(\alpha)$ -splitting in T.

(iii) Let $\langle u_{\alpha} \mid \alpha < \gamma \rangle$ be an increasing sequence of splitting nodes in T, then for every $\xi < \lambda$ we also see that $\langle u_{\alpha} \mid \alpha < \gamma \rangle$ is a sequence of splitting nodes in T_{ξ} , and thus $u = \bigcup_{\alpha < \gamma} u_{\alpha}$ is a splitting node in all T_{ξ} , hence by the claim u is splitting in T.

Corollary 5

 \mathbb{S}^h_{κ} preserves all cardinalities and cofinalities $\leq \kappa$.

What is left, is to show that κ^+ is also preserved. This will be a consequence of showing that \mathbb{S}^h_{κ} has the *F*-Sacks property for some suitably large $F \in {}^{\kappa}\kappa$: if \dot{f} is a \mathbb{S}^h_{κ} -name and $T \Vdash {}^{"}\dot{f} : \check{\kappa} \to \check{\kappa}^+{}^{"}$ for some $T \in \mathbb{S}^h_{\kappa}$, then using (the proof of) the *F*-Sacks property we may produce sets A_{ξ} with $|A_{\xi}| = F(\xi)$ for each $\xi < \kappa$ such that $T' \Vdash {}^{"}\dot{f}(\check{\xi}) \in \check{A}_{\xi}{}^{"}$ for some stronger $T' \leq T$, and thus \dot{f} is forced to have a range contained in $\bigcup_{\xi} A_{\xi}$ and cannot be cofinal in κ^+ .

Before we can prove that \mathbb{S}^h_{κ} indeed has the *F*-Sacks property for some suitably large *F*, we will need to show that \mathbb{S}^h_{κ} is closed under fusion. It will be helpful to establish some notation for subtrees. If $u \in T$, let $T_u = \{v \in T \mid u \subseteq v \lor v \subseteq u\}$. It is clear that $T_u \leq T$.

Let a tree $T \in \mathbb{S}^h_{\kappa}$ be called **sharp** if every $u \in \operatorname{Split}_{\alpha}(T)$ is a sharp $h(\alpha)$ -splitting node. It is clear that by pruning we may find a sharp T^* below any condition $T \in \mathbb{S}^h_{\kappa}$ such that $\operatorname{Split}_{\alpha}(T^*) \subseteq \operatorname{Split}_{\alpha}(T)$ for every $\alpha < \kappa$, and we may assume that we can canonically do so, thus we will hereby fix the notation T^* to denote a sharp tree below condition T. We will write $(\mathbb{S}^h_{\kappa})^* = \{T \in \mathbb{S}^h_{\kappa} \mid T \text{ is sharp}\}$, which embeds densely into \mathbb{S}^h_{κ} .

For $T, S \in \mathbb{S}^h_{\kappa}$, we let $T \leq_{\alpha} S$ iff $T \leq S$ and $\operatorname{Split}_{\alpha}(T) = \operatorname{Split}_{\alpha}(S)$. A fusion sequence is a sequence $\langle T_{\alpha} \in \mathbb{S}^h_{\kappa} \mid \alpha < \kappa \rangle$ such that $T_{\beta} \leq_{\alpha} T_{\alpha}$ for all $\alpha \leq \beta < \kappa$.

 \triangleleft

Lemma 6

If $\langle T_{\alpha} \in \mathbb{S}_{\kappa}^{h} \mid \alpha < \kappa \rangle$ is a fusion sequence, then $T = \bigcap T_{\alpha} \in \mathbb{S}_{\kappa}^{h}$.

Proof. Clearly T is a tree on $<\kappa$. We check conditions (i), (ii) and (iii) of Definition 3.

(i) Let $u \in T$ and $\alpha = \operatorname{ot}(u)$, then for any $\beta, \beta' \geq \alpha$ we see that $\operatorname{Split}_{\alpha}(T_{\beta}) = \operatorname{Split}_{\alpha}(T_{\beta'})$, thus pick $\beta > \alpha$ arbitrarily. Since $\alpha = \operatorname{ot}(u)$, necessarily there exists some $v \in \operatorname{Split}_{\alpha}(T_{\beta})$ such that $u \subseteq v$, and since $\beta > \alpha$ we see that $v \in \operatorname{Split}_{\alpha}(T)$.

(ii) Let $s \in \text{Split}_{\alpha}(T)$, then s is $h(\alpha)$ -splitting in $T_{\alpha+1}$. Let $\lambda_s = |\operatorname{suc}(s, T_{\alpha+1})| \ge h(\alpha)$ and let $\langle t_{\xi} | \xi < \lambda_s \rangle$ enumerate those $t \supseteq s$ such that $t \in \text{Split}_{\alpha+1}(T_{\alpha+1})$. For all $\beta, \beta' \ge \alpha + 1$ we have $\operatorname{Split}_{\alpha+1}(T_{\beta}) = \operatorname{Split}_{\alpha+1}(T_{\beta'})$, therefore for each $\xi < \lambda_s$ we see that $t_{\xi} \in T_{\beta}$ for all $\beta > \alpha$, thus $t_{\xi} \in T$. Therefore s is $h(\alpha)$ -splitting in T.

(iii) Let γ be limit and let $\langle u_{\alpha} \in \text{Split}_{\alpha}(T) \mid \alpha < \gamma \rangle$ such that $u_{\alpha} \subseteq u_{\alpha'}$ for each $\alpha < \alpha'$, then $u_{\alpha} \in \text{Split}_{\alpha}(T_{\gamma})$ and thus $u = \bigcup u_{\alpha}$ is an $h(\gamma)$ -splitting node in T_{γ} , hence $u \in \text{Split}_{\gamma}(T_{\gamma})$. It follows that $u \in \text{Split}_{\gamma}(T_{\beta})$ for all $\beta \geq \gamma$, thus $u \in \text{Split}_{\gamma}(T)$.

We are now ready to prove the two main ingredients necessary for separating the localisation cardinals. We will show that for any h there is some faster growing F such that \mathbb{S}^h_{κ} has the F-Sacks property, and reversely for any F there exists some faster growing h such that \mathbb{S}^h_{κ} does not have the F-Sacks property. In other words, for any F_0 we may find h and F_1 such that \mathbb{S}^h_{κ} does not have the F_0 -Sacks property, but does have the F_1 -Sacks property.

Theorem 7

For any h there exists F such that
$$h \leq F$$
 and \mathbb{S}^h_{κ} has the F-Sacks property.

Proof. We will let $F: \alpha \mapsto h(\alpha)^{|\alpha|}$ and show that \mathbb{S}^h_{κ} has the F-Sacks property.

Let $T_0 \in \mathbb{S}^h_{\kappa}$ and let \dot{f} be a name such that $T_0 \Vdash ``\dot{f} : \check{\kappa} \to \check{\kappa}$ ". If $T_0 \Vdash ``\dot{f} \in \check{\mathbf{V}}$ ", then the existence of an appropriate F-slalom is obvious, so we assume that $T_0 \Vdash ``\dot{f} \notin \check{\mathbf{V}}$ ". We will construct a fusion sequence $\langle T_{\xi} \mid \xi < \kappa \rangle$ and a family of sets $\{B_{\xi} \subseteq \kappa \mid \xi < \kappa\}$ with $|B_{\xi}| \leq F(\xi)$ such that $\bigcap T_{\xi} = T \Vdash ``\dot{f}(\check{\xi}) \in \check{B}_{\xi}$ " for each $\xi < \kappa$. Consequently, we can define the F-slalom $\varphi : \xi \mapsto B_{\xi}$ to see that $T \Vdash ``\dot{f}(\check{\xi}) \in \check{\varphi}(\check{\xi})$ " for all $\xi < \kappa$.

In general, we will assume each T_{ξ} has the following property:

(*) for every $u \in \text{Split}_{\alpha}(T_{\xi})$ with $\alpha < \xi$ we have $|\operatorname{suc}(u, T_{\xi})| = h(\alpha)$.

This is vacuously true for T_0 , and by using sharp trees at successor stages of our construction, (*) will follow by induction. If γ is limit, we will let $T_{\gamma} = \bigcap_{\xi < \gamma} T_{\xi}$, which will be a condition by the proof of Lemma 4. T_{γ} need not necessarily be a sharp tree, but it is at least sharp for all splitting levels less than γ , which is enough for (*).

Suppose T_{ξ} has been defined, then we will define $T_{\xi+1}$ that limits the possible values of $f(\xi)$ and such that $T_{\xi+1} \leq_{\xi} T_{\xi}$. First note that if T_{ξ} has property (*), then $T_{\xi}^* \leq_{\xi} T_{\xi}$: If u is splitting

in T_{ξ} and $u \notin T_{\xi}^*$, then u was removed because there is some $v \subseteq u$ such that $suc(v, T_{\xi})$ is too large for sharpness. But then by (*) it follows that $v \in \text{Split}_{\alpha}(T_{\xi})$ for some $\alpha \geq \xi$, hence $u \in \text{Split}_{\beta}(T_{\xi}) \text{ for some } \beta > \xi.$

Let $V_{\xi} = \bigcup \{ \operatorname{suc}(u, T_{\xi}^*) \mid u \in \operatorname{Split}_{\xi}(T_{\xi}^*) \}$, and for each $v \in V_{\xi}$ find a condition $T^v \leq (T_{\xi}^*)_v$ such that $T^v \Vdash \check{f}(\check{\xi}) = \check{\beta}^v_{\xi}$ for some $\beta^v_{\xi} < \kappa$. Let $u \in \text{Split}_{\xi}(T^v)$ and $w \in \text{suc}(u, T^v)$ be arbitrary and consider the subtree T_w^v of T^v generated by the initial segment w. We let $G_{\xi}: V_{\xi} \to \mathcal{P}(T_{\xi})$ send $v \mapsto T_w^v$. Note that the α -th splitting level of $G_{\xi}(v) = T_w^v$ corresponds to the $(\xi + 1 + \alpha)$ -th splitting level of T^v .

Now we define:

$$T_{\xi+1} = \bigcup G_{\xi}[V_{\xi}] = \bigcup \left\{ G_{\xi}(v) \mid v \in V_{\xi} \right\},$$
$$B_{\xi} = \left\{ \beta_{\xi}^{v} \mid v \in V_{\xi} \right\}.$$

For each $v \in V_{\xi}$ we have $v \in G_{\xi}(v)$, thus each successor of a splitting node in $\text{Split}_{\xi}(T_{\xi})$ is in $T_{\xi+1}$, therefore we see that $\operatorname{Split}_{\xi}(T_{\xi+1}) = \operatorname{Split}_{\xi}(T_{\xi})$. If $u \in \operatorname{Split}_{\xi+1+\alpha}(T_{\xi+1})$ for some $\alpha < \kappa$, then $u \in \text{Split}_{\alpha}(G_{\xi}(v))$, thus $u \in \text{Split}_{\xi+1+\alpha}(T^v)$, and since $T^v \in \mathbb{S}^h_{\kappa}$, we see that u is $h(\xi + 1 + \alpha)$ -splitting. Therefore $T_{\xi+1}$ satisfies (ii) of Definition 3. It is easy to check (i) and (iii), thus we can conclude that $T_{\xi+1} \in \mathbb{S}^h_{\kappa}$ and that $T_{\xi+1} \leq_{\xi} T_{\xi}$.

Note that $|B_{\xi}| \leq |V_{\xi}| = |\operatorname{Split}_{\xi}(T_{\xi}^*)| \cdot h(\xi) \leq h(\xi)^{|\xi|} = F(\xi)$, as intended.

Now $T^v \Vdash ``\dot{f}(\check{\xi}) \in \check{B}_{\xi}$, and $\{T^v \mid v \in V_{\xi}\}$ is predense below $T_{\xi+1}$, thus $T_{\xi+1} \Vdash ``\dot{f}(\check{\xi}) \in \check{B}_{\xi}$. Let $T = \bigcap_{\xi \leq \kappa} T_{\xi}$, then by the fusion lemma $T \in \mathbb{S}^h_{\kappa}$, and $T \Vdash ``f(\xi) \in \check{B}_{\xi}$ " for all $\xi < \kappa$.

Corollary 8

 \mathbb{S}^h_{κ} preserves κ^+ .

Before we prove the existence of F such that \mathbb{S}^h_{κ} does not have the F-Sacks property, we will state a lemma that will be useful later on as well.

Lemma 9

Let $T \in \mathbb{S}^h_{\kappa}$ and let $C_T = \{ \alpha < \kappa \mid \text{Split}_{\alpha}(T) = T \cap {}^{\alpha}\kappa \}$, then C_T is a club set. \triangleleft

Proof. For $\alpha_0 \in \kappa$ we can recursively define α_{n+1} large enough such that $\text{Split}_{\alpha_n}(T) \subseteq \leq \alpha_{n+1}\kappa$ for each $n \in \omega$. Let $\alpha = \bigcup_{n \in \omega} \alpha_n$, then $\alpha \in C_T$, hence C_T is unbounded. It is easy to see that C_T is continuous.

Theorem 10

Let $F \in {}^{\kappa}\kappa$, then there exists h such that \mathbb{S}^{h}_{κ} does not have the F-Sacks property.

Proof. Let h be such that $F(\alpha) < h(\alpha)$ for all $\alpha \in S$, where S is a stationary subset of κ . We will show that \mathbb{S}^h_{κ} does not have the *F*-Sacks property.

Let \dot{f} be a name for the generic \mathbb{S}^h_{κ} -real in ${}^{\kappa}\kappa$, let φ be an F-slalom, let $T \in \mathbb{S}^h_{\kappa}$ and let $\alpha_0 < \kappa$. We want to find some $\alpha \geq \alpha_0$ and $S \leq T$ such that $S \Vdash ``\dot{f}(\check{\alpha}) \notin \check{\varphi}(\check{\alpha})$ ''. If we can find $u \in T \cap {}^{\alpha+1}\kappa$ such that $u(\alpha) \notin \varphi(\alpha)$, then T_u will be sufficient.

Let C_T be as defined in Lemma 9 and $\alpha \in C_T \cap S$ such that $\alpha_0 \leq \alpha$, then $\text{Split}_{\alpha}(T) = T \cap {}^{\alpha}\kappa$, thus each $t \in T \cap {}^{\alpha}\kappa$ is an $h(\alpha)$ -splitting node. Hence, there is a set $X \subseteq \kappa$ with $|X| = h(\alpha)$ such that $t^{\gamma} \in T$ for all $\gamma \in X$. Since $|\varphi(\alpha)| = F(\alpha) < h(\alpha)$, there is some $\gamma \in X$ such that $\gamma \notin \varphi(\alpha)$, and thus $u = t^{\gamma} \gamma$ is as desired.



 \triangleleft

PRODUCTS

We see that for any F_0 , we can find a faster growing F_1 and some suitable h such that the forcing \mathbb{S}^h_{κ} has the F_1 -Sacks property and not the F_0 -Sacks property, thus forcing with \mathbb{S}^h_{κ} will not increase $\mathfrak{d}_{\kappa}^{F_1}(\in^*)$, but has the potential to increase $\mathfrak{d}_{\kappa}^{F_0}(\in^*)$.

In order to increase $\mathfrak{d}_{\kappa}^{F_0}(\in^*)$ we will need to add many \mathbb{S}_{κ}^h -generic κ -reals to the ground model. This can be either done with an iteration, or with a product. Iteration has the drawback that once we have forced 2^{κ} to be of size κ^{++} , the forcing \mathbb{S}^{h}_{κ} no longer has the $<\kappa^{++}$ -cc, and thus we cannot sufficiently control the iteration past this point. We will therefore focus on the product instead.

Let A be a set of ordinals and h_{ξ} for each $\xi \in A$. Let C be the set of (choice) functions $p: A \to \bigcup_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ such that $p(\xi) \in \mathbb{S}_{\kappa}^{h_{\xi}}$ for each $\xi \in A$. For any $p \in C$, we define the **support** of $p \text{ as supp}(p) = \left\{ \xi \in A \mid p(\xi) \neq \mathbb{1}_{\mathbb{S}^{h_{\xi}}} \right\}.$ We define the $\leq \kappa$ -support product as follows:

$$\prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}} = \{ p \in \mathcal{C} \mid |\operatorname{supp}(p)| \le \kappa \}.$$

We will fix A, $\langle h_{\xi} | \xi \in A \rangle$ and C and use the shorthand $\overline{\mathbb{S}} = \prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$, unless stated otherwise. If $p, q \in \overline{\mathbb{S}}$, then $q \leq_{\overline{\mathbb{S}}} p$ iff $q(\xi) \leq_{\mathbb{S}_{\kappa}^{h_{\xi}}} p(\xi)$ for all $\xi \in A$. We will again write $q \leq p$ instead of $q \leq_{\overline{\mathbb{S}}} q$ if the context is clear.

If $\langle p_{\alpha} \mid \alpha < \gamma \rangle$ is a sequence of conditions in $\overline{\mathbb{S}}$ such that $p_{\alpha'} \leq_{\overline{\mathbb{S}}} p_{\alpha}$ for all $\alpha \leq \alpha'$, then we let $\bigwedge_{\alpha < \gamma} p_{\alpha} : A \to \bigcup_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ be the function sending $\xi \mapsto \bigcap_{\alpha < \gamma} p_{\alpha}(\xi)$. If we assume that $\gamma < \kappa$, then by the $\langle \kappa$ -closure of each $\mathbb{S}^{h_{\xi}}_{\kappa}$ (Lemma 4) it follows that $\bigcap_{\alpha < \gamma} p_{\alpha}(\xi) \in \mathbb{S}^{h_{\xi}}_{\kappa}$, and thus $\bigwedge_{\alpha < \gamma} p_{\alpha} \in \mathcal{C}$. It is easy to see that $|\operatorname{supp}(\bigwedge_{\alpha < \gamma} p_{\alpha})| \le \kappa$ as well, thus we see that $\overline{\mathbb{S}}$ is $<\kappa$ -closed:

Lemma 11

 $\overline{\mathbb{S}}$ is $<\kappa$ -closed.

We will also need a generalisation of the fusion lemma to work on product forcings. The generalisation of fusion described here is analogous to what is described in [Kan80] or [BBTFM18].

Given $p, q \in \overline{\mathbb{S}}$, $\alpha < \kappa$, and $Z \subseteq A$ with $|Z| < \kappa$, let $q \leq_{Z,\alpha} p$ iff $q \leq p$ and for each $\xi \in Z$ we have $q(\xi) \leq_{\alpha} p(\xi)$.

A generalised fusion sequence is a sequence $\langle (p_{\alpha}, Z_{\alpha}) \mid \alpha < \kappa \rangle$ such that:

- $p_{\alpha} \in \overline{\mathbb{S}}$ and $Z_{\alpha} \in [A]^{<\kappa}$ for each $\alpha < \kappa$,
- p_β ≤_{Zα,α} p_α and Z_α ⊆ Z_β for all α ≤ β < κ,
 for limit δ we have Z_δ = U_{α<δ} Z_α,
- $\bigcup_{\alpha < \kappa} Z_{\alpha} = \bigcup_{\alpha < \kappa} \operatorname{supp}(p_{\alpha}).$

Lemma 12

If $\langle (p_{\alpha}, Z_{\alpha}) \mid \alpha < \kappa \rangle$ is a generalised fusion sequence, then $\bigwedge_{\alpha < \kappa} p_{\alpha} \in \overline{\mathbb{S}}$.

Proof. Suppose that $\langle (p_{\alpha}, Z_{\alpha}) \mid \alpha < \kappa \rangle$ is a generalised fusion sequence, and let $p = \bigwedge_{\alpha < \kappa} p_{\alpha}$. The last condition of the definition above implies that for any $\xi \in \text{supp}(p)$ there is some $\eta_{\xi} \in \kappa$ such that $\xi \in Z_{\eta_{\xi}}$. This means that if $\beta \geq \alpha \geq \eta_{\xi}$, then $p_{\beta}(\xi) \leq_{\alpha} p_{\alpha}(\xi)$, and thus $\langle p_{\alpha}(\xi) \mid \alpha > \eta_{\xi} \rangle$ is a fusion sequence in $\mathbb{S}_{\kappa}^{h_{\xi}}$. Since $\mathbb{S}_{\kappa}^{h_{\xi}}$ is closed under fusion sequences, we therefore conclude that $p(\xi) = \bigcap_{\alpha < \kappa} p_{\alpha}(\xi) \in \mathbb{S}_{\kappa}^{h_{\xi}}$. Since $\operatorname{supp}(p) = \bigcup_{\alpha < \kappa} Z_{\alpha}$, we see that $|\operatorname{supp}(p)| \leq \kappa$, thus we can conclude that $p \in \overline{\mathbb{S}}$.

 \triangleleft

By Lemma 11, $\overline{\mathbb{S}}$ preserves all cardinalities up to and including κ . Suppose that each $\mathbb{S}_{\kappa}^{h_{\xi}}$ has the *F*-Sacks property for some suitably large *F*. We will show in the next lemma that this implies that $\overline{\mathbb{S}}$ has the *F*-Sacks property and therefore preserves κ^+ . Finally, if we assume that $\mathbf{V} \models 2^{\kappa} = \kappa^+$ ", then a standard Δ -system argument (see e.g. [Jec03] Lemma 15.4) shows that $\overline{\mathbb{S}}$ is $<\kappa^{++}$ -cc as well. Thus, $\overline{\mathbb{S}}$ preserves all cardinals and cofinalities assuming that each $\mathbb{S}_{\kappa}^{h_{\xi}}$ has the *F*-Sacks property for some fixed $F \in {}^{\kappa}\kappa$.

Before we prove the lemma, let us introduce some notation related to the product of forcings. Suppose $\overline{\mathbb{P}} = \prod_{\xi \in A} \mathbb{P}_{\xi}$ is a product with $\leq \kappa$ -support and G is $\overline{\mathbb{P}}$ -generic. If $B \subseteq A$, we define

$$\overline{\mathbb{P}} \upharpoonright B = \left\{ p \upharpoonright B \mid p \in \overline{\mathbb{P}} \right\} \qquad \qquad G \upharpoonright B = \left\{ p \upharpoonright B \mid p \in G \right\}$$

Let $B^c = A \setminus B$, then clearly $\overline{\mathbb{P}}$ and $(\overline{\mathbb{P}} \upharpoonright B) \times (\overline{\mathbb{P}} \upharpoonright B^c)$ are forcing equivalent, $(G \upharpoonright B) \times (G \upharpoonright B^c)$ is $(\overline{\mathbb{P}} \upharpoonright B) \times (\overline{\mathbb{P}} \upharpoonright B^c)$ -generic and $\mathbf{V}[G] = \mathbf{V}[(G \upharpoonright B) \times (G \upharpoonright B^c)] = \mathbf{V}[G \upharpoonright B][G \upharpoonright B^c].$

Lemma 13

Let $B \subseteq A$ be sets of ordinals and $B^c = A \setminus B$, and consider a sequence of functions $\langle h_{\xi} | \xi \in A \rangle$. We define the $\leq \kappa$ -support product $\overline{\mathbb{S}} = \prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ and we assume G is an $\overline{\mathbb{S}}$ -generic filter. If $F : \alpha \mapsto (\sup_{\xi \in B^c} h_{\xi}(\alpha))^{|\alpha|}$ is a well-defined function in κ_{κ} , then for each $f \in (\kappa_{\kappa})^{\mathbf{V}[G]}$ there is $\varphi \in (\operatorname{Loc}_F)^{\mathbf{V}[G \upharpoonright B]}$ such that $f \in \varphi$.

Proof. Let \dot{f} be a name such that $\Vdash_{\overline{\mathbb{S}}}$ " $\dot{f}: \check{\kappa} \to \check{\kappa}$ " and let $p \in \overline{\mathbb{S}}$, then we will construct a name $\dot{\varphi}$ and a condition $p' \leq p$ such that $p' \Vdash$ " $\dot{\varphi} \in (\operatorname{Loc}_{\check{F}})^{\mathbf{V}[\dot{G} \upharpoonright \check{B}]}$ ".

The proof is essentially the same as the proof of Theorem 7, except that we work with generalised fusion sequences and have to construct a name $\dot{\varphi}$ for the appropriate *F*-slalom, since such slalom is not generally present in the ground model. That is, we will construct a sequence $\langle (p_{\xi}, Z_{\xi}) | \xi < \kappa \rangle$ with each $p_{\xi} \in \overline{\mathbb{S}}$ that is a generalised fusion sequence in $\overline{\mathbb{S}}$ and names \dot{D}_{ξ} for sets of ordinals $D_{\xi} \in \mathbf{V}[G \upharpoonright B]$ with $|D_{\xi}| \leq F(\xi)$, such that $p_{\xi+1} \Vdash "\dot{f}(\check{\xi}) \in \dot{D}_{\xi}$ ". Furthermore, for each $\xi < \kappa$ and $\beta \in Z_{\xi}$ we will make sure that $p_{\xi}(\beta) \in (\mathbb{S}_{\kappa}^{h_{\beta}})^*$ is sharp. To start, we let $p_0 = p$ and we let $Z_0 = \emptyset$. At limit stages δ we can define $p'_{\delta} = \bigwedge_{\xi < \delta} p_{\xi}$ and let $p_{\delta} \leq p'_{\delta}$ be defined elementwise such that $p_{\delta}(\beta) = (p'_{\delta}(\beta))^*$ is sharp for each $\beta \in Z_{\delta}$.

Suppose we have defined $p_{\xi} \in \overline{\mathbb{S}}$ and Z_{ξ} and that $|Z_{\xi}| \leq |\xi|$, then for each $\beta \in Z_{\xi}$ we define

$$V_{\xi}^{\beta} = \bigcup \left\{ \mathrm{suc}(u, p_{\xi}(\beta)) \mid u \in \mathrm{Split}_{\xi}(p_{\xi}(\beta)) \right\}$$

Let \mathcal{V}_{ξ} be the set of choice functions on $\{V_{\xi}^{\beta} \mid \beta \in Z_{\xi}\}$ and \mathcal{V}'_{ξ} the set of choice functions on $\{V_{\xi}^{\beta} \mid \beta \in Z_{\xi} \setminus B\}$. By induction hypothesis $p_{\xi}(\beta) \in (\mathbb{S}_{\kappa}^{h_{\beta}})^{*}$ for each $\beta \in Z_{\xi} \setminus B$, hence we know that $|\text{Split}_{\xi}(p_{\xi}(\beta))| \leq h_{\beta}(\xi)^{|\xi|} \leq F(\xi)$, and thus $|V_{\xi}^{\beta}| \leq F(\xi)$ for all $\beta \in Z_{\xi} \setminus B$. Since we assume that $|Z_{\xi}| \leq |\xi|$, we therefore have $|\mathcal{V}'_{\xi}| \leq F(\xi)^{|\xi|} = F(\xi)$ (assuming without loss of generality that $F(\xi)$ is infinite). For any choice function $g \in \mathcal{V}_{\xi}$, let $(p_{\xi})_{g}$ be the condition with $(p_{\xi})_{g}(\beta) = p_{\xi}(\beta)$ if $\beta \notin Z_{\xi}$ and $(p_{\xi})_{g}(\beta) = (p_{\xi}(\beta))_{g(\beta)}$ if $\beta \in Z_{\xi}$. Here $(p_{\xi}(\beta))_{g(\beta)}$ is the subtree of $p_{\xi}(\beta)$ generated by the initial segment $g(\beta)$.

Let $\zeta = |\mathcal{V}_{\xi}|$ then $\zeta < \kappa$ by inaccessibility of κ . Fix some enumeration $\langle g_{\eta} | \eta < \zeta \rangle$ of \mathcal{V}_{ξ} , which we will use to define a decreasing sequence of conditions r_{η} with $r_{\eta} \leq_{Z_{\xi},\xi} p_{\xi}$ for each $\eta < \zeta$. Let $r_0 = p_{\xi}$. For limit $\delta < \zeta$ let $r_{\delta} = \bigwedge_{\eta < \delta} r_{\eta}$, which is a condition by $<\kappa$ -closure (Lemma 11). Assuming that $r_{\eta} \leq_{Z_{\xi},\xi} p_{\xi}$ for each $\eta < \delta$, it is easy to see that $r_{\delta} \leq_{Z_{\xi},\xi} p_{\xi}$ as well. Suppose r_{η} is defined and $r_{\eta} \leq_{Z_{\xi},\xi} p_{\xi}$, then in particular $r_{\eta}(\beta) \leq_{\xi} p_{\xi}(\beta)$ for all $\beta \in Z_{\xi}$, and thus $\operatorname{Split}_{\xi}(r_{\eta}(\beta)) = \operatorname{Split}_{\xi}(p_{\xi}(\beta))$ for all $\beta \in Z_{\xi}$, and therefore by definition of the ordering on $\mathbb{S}_{\kappa}^{h_{\beta}}$ and the fact that $p_{\xi}(\beta)$ is sharp, we see that V_{ξ}^{β} is exactly the set of successors of nodes at the ξ -th splitting level of $r_{\eta}(\beta)$. Take the η -th choice function $g_{\eta} \in \mathcal{V}_{\xi}$, and let $r'_{\eta} \leq (r_{\eta})_{g_{\eta}}$ be such that $r'_{\eta} \Vdash "f(\xi) = \check{\beta}_{\xi}^{\eta}$ " for some ordinal β_{ξ}^{η} . We define $r_{\eta+1}$ elementwise.

If $\beta \notin Z_{\xi}$, then we simply take $r_{\eta+1}(\beta) = r'_{\eta}(\beta)$.

If $\beta \in Z_{\xi}$, fix some $w \in \operatorname{suc}(u, r'_{\eta}(\beta))$ for some $u \in \operatorname{Split}_{\xi}(r'_{\eta}(\beta))$ and consider the subtree $(r'_{\eta}(\beta))_w$ generated by the initial segment w. Now we are ready to define $r_{\eta+1}(\beta)$ as

$$r_{\eta+1}(\beta) = (r'_{\eta}(\beta))_{w} \cup \left\{ u \in r_{\eta}(\beta) \mid \exists v \in V_{\xi}^{\beta} \setminus \{g_{\eta}(\beta)\} \ (u \subseteq v \text{ or } v \subseteq u) \right\}.$$

In words, $r_{\eta+1}(\beta)$ is the result of replacing the extensions of $g_{\eta}(\beta) \in r_{\eta}(\beta)$ by $(r'_{\eta}(\beta))_w$ that decides \dot{f} at ξ , where the function of taking the subtree of $r'_{\eta}(\beta)$ generated by w is to make sure $r_{\eta+1}(\beta)$ has enough successors at each splitting level to be in $\mathbb{S}^{h_{\beta}}_{\kappa}$.

To finish the construction of the next condition in the fusion sequence, we use $<\kappa$ -closure to define $p'_{\xi+1} = \bigwedge_{\eta < \zeta} r_{\eta}$ and let $p_{\xi+1} = (p'_{\xi+1})^*$ be sharp. To see that $p_{\xi+1} \leq_{Z_{\xi},\xi} p_{\xi}$, note that for every $\beta \in Z_{\xi}$ and $v \in V_{\xi}^{\beta}$ we have $v \in r_{\eta}(\beta)$ for all $\eta < \zeta$, hence $v \in p_{\xi+1}(\beta)$. This implies by definition of V_{ξ}^{β} that $p_{\xi+1}(\beta) \leq_{\xi} p_{\xi}(\beta)$ for all $\beta \in Z_{\xi}$. Finally, we can let $Z_{\xi+1} = Z_{\xi} \cup \{\delta\}$ for some ordinal δ , using bookkeeping to make sure that $\bigcup_{\xi < \kappa} Z_{\xi} = \bigcup_{\xi < \kappa} \operatorname{supp}(p_{\xi})$.

Note that the set of conditions $r \leq p_{\xi+1}$ such that $|r(\beta) \cap V_{\xi}^{\beta}| = 1$ for all $\beta \in Z_{\xi}$, is dense below $p_{\xi+1}$. For any such r, let g map β to the unique element of $r(\beta) \cap V_{\xi}^{\beta}$ for each $\beta \in Z_{\xi}$, then $g \in \mathcal{V}_{\xi}$, so we see that there exists $\eta < \zeta$ such that $g = g_{\eta}$. We will show that $r \leq r'_{\eta}$, which implies that $r \Vdash ``\dot{f}(\check{\xi}) = \check{\beta}_{\xi}^{\eta}$.

For any β we have $r(\beta) \leq p_{\xi+1}(\beta) \leq r_{\eta+1}(\beta)$. If $\beta \notin Z_{\xi}$, then we simply have $r_{\eta+1}(\beta) = r'_{\eta}(\beta)$, thus we are done. Otherwise $\beta \in Z_{\xi}$, and we know that $g(\beta)$ is an initial segment of the stem of $r(\beta)$, hence $r(\beta) = (r(\beta))_{g(\beta)} \subseteq (r_{\eta+1}(\beta))_{g(\beta)} = (r'_{\eta}(\beta))_w$, where w is as in the definition of $r_{\eta+1}(\beta)$ above. Since $r(\beta) \leq r_{\eta+1}(\beta)$, we also have $r(\beta) = (r(\beta))_w \leq (r_{\eta+1}(\beta))_w = (r'_{\eta}(\beta))_w$, and since $(r'_{\eta}(\beta))_w \leq r'_{\eta}(\beta)$, we see that $r(\beta) \leq r'_{\eta}(\beta)$.

We now construct the names D_{ξ} such that:

$$p_{\xi+1} \Vdash ``\dot{f}(\check{\xi}) \in \dot{D}_{\xi} \text{ and } \dot{D}_{\xi} \in \mathbf{V}[\dot{G} \upharpoonright \check{B}] \text{ and } |\dot{D}_{\xi}| \leq \check{F}(\check{\xi})``.$$

For any $g \in \mathcal{V}_{\xi}$, let $g'' = g \upharpoonright (Z_{\xi} \cap B)$, and let $E_g = \left\{ \eta < \zeta \mid \exists g' \in \mathcal{V}'_{\xi}(g' \cup g'' = g_{\eta}) \right\}$ and $D_{\xi}^g = \left\{ \beta_{\xi}^{\eta} \mid \eta \in E_g \right\}$. Since $|\mathcal{V}'_{\xi}| \leq F(\xi)$, we see that $|E_g| \leq F(\xi)$, hence $|D_{\xi}^g| \leq F(\xi)$. Clearly, if $g, \tilde{g} \in \mathcal{V}_{\xi}$ and $g \upharpoonright (Z_{\xi} \cap B) = \tilde{g} \upharpoonright (Z_{\xi} \cap B)$, then $D_{\xi}^g = D_{\xi}^{\tilde{g}}$.

Let \mathcal{A}_{ξ} be an antichain below $p_{\xi+1}$ such that $r \in \mathcal{A}_{\xi}$ implies $|r(\beta) \cap V_{\xi}^{\beta}| = 1$ for all $\beta \in Z_{\xi}$, and let $g_r \in \mathcal{V}_{\xi}$ be such that $g_r(\beta)$ is the single element of $r(\beta) \cap V_{\xi}^{\beta}$ for each $\beta \in Z_{\xi}$. We define the name $\dot{D}_{\xi} = \left\{ (r, \check{D}_{\xi}^{g_r}) \mid r \in \mathcal{A}_{\xi} \right\}$. It is clear by the above that for each $r \in \mathcal{A}$ and η such that $g_r = g_{\eta}$ we have $r \Vdash ``\dot{f}(\check{\xi}) = \check{\beta}_{\xi}^{\eta} \in \check{D}_{\xi}^{g_r}$ and $|\check{D}_{\xi}^{g_r}| \leq \check{F}(\check{\xi})$, so by denseness $p_{\xi+1} \Vdash ``\dot{f}(\check{\xi}) \in \dot{D}_{\xi}$ and $|\dot{D}_{\xi}| \leq \check{F}(\check{\xi})$. To see that $p_{\xi+1} \Vdash ``\dot{D}_{\xi} \in \mathbf{V}[\dot{G} \upharpoonright B]$ ", we argue within $\mathbf{V}[G \upharpoonright B]$. For every $r, \tilde{r} \in \mathcal{A}_{\xi}$ such that $r \upharpoonright B \in G \upharpoonright B$ and $\tilde{r} \upharpoonright B \in G \upharpoonright B$ we see that the corresponding g_r and $g_{\tilde{r}}$ have the property that $g_r \upharpoonright (Z_{\xi} \cap B) = g_{\tilde{r}} \upharpoonright (Z_{\xi} \cap B)$, and therefore $D_{\xi}^{g_r} = D_{\xi}^{g_{\tilde{r}}}$. Thus, we can fix any arbitrary such $r \in \mathcal{A}_{\xi}$ for which $r \upharpoonright B \in G \upharpoonright B$ holds, and see that $\mathbf{V}[G \upharpoonright B] \models ``p_{\xi+1} \upharpoonright B^c \Vdash \dot{D}_{\xi} = \check{D}_{\xi}^{g_r}$ ".

Let $p' = \bigwedge_{\xi < \kappa} p_{\xi}$ be the limit of the generalised fusion sequence, and let $\dot{\varphi}$ be a name such that $p' \Vdash "\dot{\varphi} : \check{\xi} \mapsto \dot{D}_{\xi}"$, then $\dot{\varphi}$ names an *F*-slalom in $\mathbf{V}[G \upharpoonright B]$ and $p' \Vdash "\dot{f} \in "\dot{\varphi}"$.

If we let $B = \emptyset$ in the definition of the lemma, then we can simplify this lemma to the following corollary, providing us with the preservation of the Sacks property.

Corollary 14

If $\overline{\mathbb{S}} = \prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ and each h_{ξ} is bounded by $h \in {}^{\kappa}\kappa$ and $F : \alpha \mapsto h(\alpha)^{|\alpha|}$, then $\overline{\mathbb{S}}$ has the *F*-Sacks property.

Finally the following lemma is based on Theorem 10 and shows how we can use products of forcings $\mathbb{S}_{\kappa}^{h_{\xi}}$ to increase the cardinality of $\mathfrak{d}_{\kappa}^{F}(\in^{*})$.

Lemma 15

Let $B \subseteq A$ be sets of ordinals, and consider a sequence of functions $\langle h_{\xi} | \xi \in A \rangle$. We define the $\leq \kappa$ -support product $\overline{\mathbb{S}} = \prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ and we assume G is a $\overline{\mathbb{S}}$ -generic filter. Let $\langle S_{\xi} | \xi \in B \rangle$ be a sequence of stationary sets. If F is such that for each $\xi \in B$ we have $F(\alpha) < h_{\xi}(\alpha)$ for all $\alpha \in S_{\xi}$, then $\mathbf{V}[G] \models "|B| \leq \mathfrak{d}_{\kappa}^{F}(\in^{*})$ ".

Proof. The lemma is trivial if $|B| \leq \kappa^+$, so we will assume that $|B| \geq \kappa^{++}$.

We work in $\mathbf{V}[G]$. Let $\mu < |B|$ and let $\{\varphi_{\xi} \mid \xi < \mu\}$ be a family of *F*-slaloms, then we want to describe some $f \in {}^{\kappa}\kappa$ such that $f \notin {}^{*}\varphi_{\xi}$ for each $\xi < \mu$. Since $\overline{\mathbb{S}}$ is $<\kappa^{++}$ -cc, we could find $A_{\xi} \subseteq A$ with $|A_{\xi}| \le \kappa^{+}$ for each $\xi < \mu$ such that $\varphi_{\xi} \in \mathbf{V}[G \upharpoonright A_{\xi}]$. Since $|B| > \mu \cdot \kappa^{+}$, we may fix some $\beta \in B \setminus \bigcup_{\xi < \mu} A_{\xi}$ for the remainder of this proof. Let $f = \bigcap_{p \in G} p(\beta)$, then $f \in {}^{\kappa}\kappa$ is the generic κ -real added by the β -th term of the product $\overline{\mathbb{S}}$.

Continuing the proof in the ground model, let \dot{f} be an $\overline{\mathbb{S}}$ -name for f and $\dot{\varphi}_{\xi}$ be an $\overline{\mathbb{S}}$ -name for φ_{ξ} , let $p \in \overline{\mathbb{S}}$ and $\alpha_0 < \kappa$. We want to find some $\alpha \geq \alpha_0$ and $q \leq p$ such that $q \Vdash ``\dot{f}(\check{\alpha}) \notin \dot{\varphi}_{\xi}(\check{\alpha})"$.

Let $C = \{\alpha < \kappa \mid p(\beta) \cap {}^{\alpha}\kappa = \text{Split}_{\alpha}(p(\beta))\}$, which is a club set by Lemma 9. Since S_{β} is stationary, there exists some $\alpha \ge \alpha_0$ such that $\alpha \in C \cap S_{\beta}$. Choose some $p_0 \le p$ such that $p_0(\beta) = p(\beta)$ and such that there is a $Y \in [\kappa]^{\le F(\alpha)}$ for which $p_0 \Vdash "\dot{\varphi}_{\xi}(\check{\alpha}) = \check{Y}"$. This is possible, since $\varphi_{\xi} \in \mathbf{V}[G \upharpoonright A_{\xi}]$ and $\beta \notin A_{\xi}$, therefore we could find $p'_0 \in \overline{\mathbb{S}} \upharpoonright A_{\xi}$ with $p'_0 \le p \upharpoonright A_{\xi}$ and Y with the aforementioned property, and then let $p_0(\eta) = p'_0(\eta)$ if $\eta \in A_{\xi}$ and $p_0(\eta) = p(\eta)$ otherwise.

Each $t \in p_0(\beta) \cap {}^{\alpha}\kappa$ is a $h_{\beta}(\alpha)$ -splitting node, hence the set $X = \{\chi < \kappa \mid t^{\frown}\chi \in p_0(\beta)\}$ has cardinality $|X| \ge h_{\beta}(\alpha)$. Because $\alpha \in S_{\beta}$ and $\beta \in B$, we have by our assumptions on F that $|Y| \le F(\alpha) < h_{\xi}(\alpha) = |X|$. We can therefore find some $\chi \in X$ such that $\chi \notin Y$. Let $q \le p_0$ be defined as $q(\beta) = (p_0(\beta))_{t^{\frown}\chi}$, which is the subtree of $p_0(\beta)$ generated by the initial segment $t^{\frown}\chi$, and $q(\eta) = p_0(\eta)$ for all $\eta \ne \beta$. Then $q \le p_0 \le p$ and $q \Vdash ``\dot{f}(\check{\alpha}) \notin \dot{\varphi}_{\xi}(\check{\alpha})"$

Lemma 16

Let A be a set of ordinals, let $\langle h_{\xi} | \xi \in A \rangle$ be a sequence of functions, let $\overline{\mathbb{S}} = \prod_{\xi \in A} \mathbb{S}_{\kappa}^{h_{\xi}}$ with $\overline{\mathbb{S}}$ -generic G, and let $F \in {}^{\kappa}\kappa$. Then $\mathbf{V} \models {}^{\circ}2^{\kappa} = \kappa^{+}$ implies $\mathbf{V}[G] \models {}^{\circ}2^{\kappa} = |\mathrm{Loc}_{F}| = \kappa^{+} \cdot |A|$.

Proof. This is a standard argument.

We're now ready to use our product forcing to separate many cardinals of the form $\mathfrak{d}^h_{\kappa}(\in^*)$.

Theorem 17

There exists a family of functions $\{g_{\eta} \mid \eta < \kappa\} \subseteq {}^{\kappa}\kappa$ such that for any $\gamma < \kappa^{+}$ and any increasing sequence $\langle \lambda_{\xi} \mid \xi < \gamma \rangle$ of cardinals with $\lambda_{0} \geq 2^{\kappa}$ and any $\sigma : \kappa \to \gamma$, there exists a forcing extension in which $\mathfrak{d}_{\kappa}^{g_{\eta}}(\in^{*}) = \lambda_{\sigma(\eta)}$ for all $\eta < \kappa$.

Proof. We assume that $\mathbf{V} \models 2^{\kappa} = \kappa^+$, or otherwise we first use a forcing to collapse 2^{κ} to become κ^+ . By a result of Solovay (theorem 8.10 in [Jec03]) there exists a family of κ many disjoint stationary subsets of κ , thus let $\{S_{\eta} \mid \eta < \kappa\}$ be such a family. Let $\kappa \leq \gamma < \kappa^+$ and $\sigma : \kappa \to \gamma$ be given. We will assume without loss of generality that σ is bijective, and hence that $\sigma^{-1} : \gamma \to \kappa$ is a well-defined bijection. Let $\langle \lambda_{\xi} \mid \xi < \gamma \rangle$ be an increasing sequence of cardinals with $\lambda_0 \geq \kappa^+$.

Fix some F such that $F(\alpha)^{|\alpha|} = F(\alpha)$. For each $\eta < \kappa$ we define a function g_{η} as follows:

$$g_{\eta}(\alpha) = \begin{cases} F(\alpha) & \text{if } \alpha \in S_{\eta} \\ 2^{F(\alpha)} & \text{otherwise} \end{cases}$$

For each $\xi < \gamma$ we define $H_{\xi} \in {}^{\kappa}\kappa$ as follows:

$$H_{\xi}(\alpha) = \begin{cases} F(\alpha) & \text{if } \alpha \in \bigcup_{\xi' < \xi} S_{\sigma^{-1}(\xi')} \\ 2^{F(\alpha)} & \text{otherwise} \end{cases}$$

Note that $g_{\eta} \geq H_{\sigma(\eta)+1}$: if $\alpha \in S_{\eta}$, then $\alpha \in S_{\sigma^{-1}(\sigma(\eta))}$, so $\alpha \in \bigcup_{\xi < \sigma(\eta)+1} S_{\sigma^{-1}(\xi)}$, and thus $H_{\sigma(\eta)+1}(\alpha) = F(\alpha)$. Therefore $\mathsf{ZFC} \vdash ``\mathfrak{d}_{\kappa}^{g_{\eta}}(\in^*) \leq \mathfrak{d}_{\kappa}^{H_{\sigma(\eta)+1}}(\in^*)$ ".

For each $\xi < \gamma$ let A_{ξ} be a set of ordinals with $|A_{\xi}| = \lambda_{\xi}$, such that $\langle A_{\xi} | \xi < \gamma \rangle$ is a sequence of mutually disjoint sets, and let $A = \bigcup_{\xi < \gamma} A_{\xi}$. For each $\xi < \gamma$ and $\beta \in A_{\xi}$, we define $h_{\beta} \in {}^{\kappa}\kappa$ as $h_{\beta} = H_{\xi}$.

The forcing that we will consider is $\overline{\mathbb{S}} = \prod_{\beta \in A} \mathbb{S}_{\kappa}^{h_{\beta}}$ with $\leq \kappa$ -support. Let G be $\overline{\mathbb{S}}$ -generic. We will fix some $\xi < \gamma$ for the remainder, and let $B = \bigcup_{\xi' \leq \xi} A_{\xi'}$ and $B^c = A \setminus B = \bigcup_{\xi < \xi' < \gamma} A_{\xi'}$. Since we assume $F(\alpha)^{|\alpha|} = F(\alpha)$, we see $\sup_{\beta \in B^c} h_{\beta}(\alpha) = \sup_{\xi < \xi' < \gamma} H_{\xi'}(\alpha) = H_{\xi+1}(\alpha)$ and therefore $H_{\xi+1} : \alpha \mapsto (\sup_{\beta \in B^c} h_{\beta}(\alpha))^{|\alpha|}$.

By Lemma 16 we see that $(\operatorname{Loc}_{H_{\xi+1}})^{\mathbf{V}[G \upharpoonright B]}$ has cardinality $\kappa^+ \cdot |B| = \kappa^+ \cdot |\sup_{\xi' \leq \xi} A_{\xi'}| = \lambda_{\xi}$. Combining this with Lemma 13, we see that there exists a family in $\mathbf{V}[G]$ of size λ_{ξ} that forms a witness for $\mathbf{V}[G] \models "\mathfrak{d}_{\kappa}^{H_{\xi+1}}(\in^*) \leq \lambda_{\xi}$ ".

Finally, for each $\eta < \kappa$ we see that if $\alpha \in S_{\eta}$ and $\beta \in A_{\sigma(\eta)}$, then $h_{\beta}(\alpha) = H_{\sigma(\eta)}(\alpha) = 2^{F(\alpha)}$, while $g_{\eta}(\alpha) = F(\alpha)$, thus using Lemma 15 we see that $\mathbf{V}[G] \models \ \ \lambda_{\sigma(\eta)} = |A_{\sigma(\eta)}| \leq \mathfrak{d}_{\kappa}^{g_{\eta}}(\in^{*})$.

In conclusion,
$$\mathbf{V}[G] \models ``\lambda_{\sigma(\eta)} \leq \mathfrak{d}_{\kappa}^{g_{\eta}}(\in^*) \leq \mathfrak{d}_{\kappa}^{H_{\sigma(\eta)+1}}(\in^*) \leq \lambda_{\sigma(\eta)}$$
 "for each $\eta < \kappa$

Corollary 18

There exists functions h_{ξ} for each $\xi < \kappa$ such that for any cardinals $\lambda_{\xi} > \kappa$ it is consistent that simultaneously $\mathfrak{d}_{\kappa}^{h_{\xi}}(\in^*) = \lambda_{\xi}$ for all $\xi < \kappa$.

References

- [Bar87] Tomek Bartoszyński. Combinatorial aspects of measure and category. Fundamenta Mathematicae, 127(3):225–239, 1987.
- [BBTFM18] Jörg Brendle, Andrew Brooke-Taylor, Sy-David Friedman, and Diana Carolina Montoya. Cichoń's diagram for uncountable cardinals. *Israel Journal of Mathematics*, 225(2):959–1010, 2018.
- [BGS20] Thomas Baumhauer, Martin Goldstern, and Saharon Shelah. The Higher Cichoń Diagram. Fundamenta Mathematicae, 252(3):241–314, 2020.
- [BJ95] Tomek Bartoszyński and Haim Judah. Set Theory: On the Structure of the Real Line. A.K. Peters, Wellesley, MA, 1995.
- [BL79] James E Baumgartner and Richard Laver. Iterated perfect-set forcing. Ann. Math. Logic, 17(3):271–288, 1979.
- [Jec03] Thomas Jech. Set Theory: Third Millennium Edition. Springer Monographs in Mathematics, 2003.
- [Kan80] Akihiro Kanamori. Perfect-set forcing for uncountable cardinals. Annals of Mathematical Logic, 19(1-2):97–114, 1980.
- [Sac71] Gerald E Sacks. Forcing with perfect closed sets. In *Axiomatic set theory*, volume 1, pages 331–355. Amer. Math. Soc Providence, RI, 1971.