New companions to the generations of the Göllnitz-Gordon identities

Thomas Y. He^1 and Alice X.H. Zhao²

¹ School of Mathematical Sciences, Laurent Mathematics Center and V.C. & V.R. Key Lab, Sichuan Normal University, Chengdu 610066, P.R. China

² College of Science, Tianjin University of Technology, Tianjin 300384, P.R. China

¹heyao@sicnu.edu.cn, ²zhaoxh@email.tjut.edu.cn

Abstract. The Göllnitz-Gordon identities were found by Göllnitz and Gordon independently. In 1967, Andrews obtained a combinatorial generalization of the Göllnitz-Gordon identities, called the Andrews-Göllnitz-Gordon theorem. In 1980, Bressoud extended the Andrews-Göllnitz-Gordon theorem to even moduli, called the Bressoud-Göllnitz-Gordon theorem. Furthermore, Bressoud gave the generating functions for the generalizations of the Göllnitz-Gordon identities. In this article, we will give new companions to the generalizations of the Göllnitz-Gordon identities.

Keywords: Göllnitz-Gordon identities, Bailey pairs, Göllnitz-Gordon marking

AMS Classifications: 05A17, 11P81, 11P84

1 Introduction

A partition of a positive integer n is a finite non-increasing sequence of positive integers $\lambda_1 \geq \lambda_2 \geq \cdots \geq \lambda_\ell > 0$ such that $\lambda_1 + \lambda_2 + \cdots + \lambda_\ell = n$. The λ_s are called the parts of the partition. We sometimes write $\lambda = (1^{f_1} 2^{f_2} 3^{f_3} \cdots)$, where $f_t(\lambda)$ denotes the number of parts equal to t in λ . Let $\ell(\lambda)$ be the number of parts of λ and $|\lambda|$ be the sum of parts of λ . The Göllnitz-Gordon identities were found by Göllnitz [10, 11] and Gordon [12, 13] independently.

Theorem 1.1 (First Göllnitz-Gordon identity). The number of partitions of n into distinct nonconsecutive parts with no even parts differing by exactly 2 equals the number of partitions of n into parts $\equiv 1, 4, 7 \pmod{8}$.

Theorem 1.2 (Second Göllnitz-Gordon identity). The number of partitions of n into distinct nonconsecutive parts with no even parts differing by exactly 2 and no parts equal to 1 or 2 equals the number of partitions of n into parts $\equiv 3, 4, 5 \pmod{8}$.

The generating function version of the Göllnitz-Gordon identities was given as follows.

$$\sum_{n\geq 0} \frac{q^{n^2}(-q;q^2)_n}{(q^2;q^2)_n} = \frac{1}{(q,q^4,q^7;q^8)_{\infty}},$$
(1.1)

and

$$\sum_{n\geq 0} \frac{q^{n^2+2n}(-q;q^2)_n}{(q^2;q^2)_n} = \frac{1}{(q^3,q^4,q^5;q^8)_\infty}.$$
(1.2)

Here and in the sequel, we assume that |q| < 1 and adopt the standard notation [3]:

$$(a;q)_{\infty} = \prod_{i=0}^{\infty} (1 - aq^i), \quad (a;q)_n = \frac{(a;q)_{\infty}}{(aq^n;q)_{\infty}},$$

and

$$(a_1, a_2, \ldots, a_m; q)_{\infty} = (a_1; q)_{\infty} (a_2; q)_{\infty} \cdots (a_m; q)_{\infty}$$

In 1967, Andrews [2] found the following combinatorial generalization of the Göllnitz-Gordon identities, which can be called the Andrews-Göllnitz-Gordon theorem.

Theorem 1.3 (Andrews-Göllnitz-Gordon identities). For $k \ge i \ge 1$, let $C_1(k, i; n)$ denote the number of partitions λ of n of the form $(1^{f_1} 2^{f_2} 3^{f_3} \cdots)$, such that

- (1) $f_1(\lambda) + f_2(\lambda) \le i 1;$
- (2) $f_{2t+1}(\lambda) \le 1;$

(3)
$$f_{2t}(\lambda) + f_{2t+1}(\lambda) + f_{2t+2}(\lambda) \le k - 1$$

For $k \ge i \ge 1$, let $D_1(k, i; n)$ denote the number of partitions of n into parts $\not\equiv 2 \pmod{4}$ and $\not\equiv 0, \pm (2i-1) \pmod{4k}$. Then for $k \ge i \ge 1$ and $n \ge 0$,

$$C_1(k, i; n) = D_1(k, i; n).$$

Setting k = 2 and i = 2 or 1, Theorem 1.3 reduces to Theorem 1.1 and Theorem 1.2, respectively. In 1980, Bressoud [7] extended the Andrews-Göllnitz-Gordon identities to even moduli, which has been called the Bressoud-Göllnitz-Gordon theorem.

Theorem 1.4 (Bressoud-Göllnitz-Gordon identities). For $k > i \ge 1$, let $C_0(k, i; n)$ denote the number of partitions of n of the form $\lambda = (1^{f_1} 2^{f_2} 3^{f_3} \cdots)$ such that

- (1) $f_1(\lambda) + f_2(\lambda) \le i 1;$
- (2) $f_{2t+1}(\lambda) \le 1;$
- (3) $f_{2t}(\lambda) + f_{2t+1}(\lambda) + f_{2t+2}(\lambda) \le k 1;$
- (4) if $f_{2t}(\lambda) + f_{2t+1}(\lambda) + f_{2t+2}(\lambda) = k 1$, then $tf_{2t}(\lambda) + tf_{2t+1}(\lambda) + (t+1)f_{2t+2}(\lambda) \equiv O_{\lambda}(2t+1) + i 1 \pmod{2}$, where $O_{\lambda}(N)$ denotes the number of odd parts not exceeding N in λ .

For $k > i \ge 1$, let $D_0(k, i; n)$ denote the number of partitions of n into parts $\not\equiv 2 \pmod{4}$ and $\not\equiv 0, \pm (2i-1) \pmod{4k-2}$. Then, for $k > i \ge 1$ and $n \ge 0$,

$$C_0(k,i;n) = D_0(k,i;n).$$

Bressoud [7] also gave the generating function versions of Theorems 1.3 and 1.4. **Theorem 1.5** (Bressoud, 1980). For j = 0 or 1 and $(2k + j)/2 > i \ge 1$,

$$\sum_{\substack{N_1 \ge \dots \ge N_{k-1} \ge 0}} \frac{(-q^{1-2N_1}; q^2)_{N_1} q^{2(N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1})}}{(q^2; q^2)_{N_1 - N_2} \cdots (q^2; q^2)_{N_{k-2} - N_{k-1}} (q^{4-2j}; q^{4-2j})_{N_{k-1}}} = \frac{(q^2; q^4)_{\infty} (q^{2i-1}, q^{4k-2i-1+2j}, q^{4k-2+2j}; q^{4k-2+2j})_{\infty}}{(q; q)_{\infty}}.$$
(1.3)

In this article, we will give a new generating function version of Theorems 1.3 and 1.4 with the aid of Bailey pairs, which is stated as follows.

Theorem 1.6. For j = 0 or 1, $k \ge i \ge 2$ and (2k + j)/2 > 2,

$$\sum_{\substack{N_1 \ge \dots \ge N_{k-1} \ge 0}} \frac{q^{2(N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1})} (-q^{1-2N_2}; q^2)_{N_2} (-q^{1+2N_1}; q^2)_{\infty}}{(q^2; q^2)_{N_1 - N_2} \cdots (q^2; q^2)_{N_{k-2} - N_{k-1}} (q^{4-2j}; q^{4-2j})_{N_{k-1}}} = \frac{(q^2; q^4)_{\infty} (q^{4k-2+2j}, q^{2i-1}, q^{4k-2i-1+2j}; q^{4k-2+2j})_{\infty}}{(q; q)_{\infty}}.$$
(1.4)

We will show that the left-hand side of (1.4) can be interpreted combinatorially as the generating function of $C_j(k, i; n)$ in Theorems 1.3 and 1.4. More precisely, We will give the following formula for the generating function of $C_j(k, i; m, n)$, where $C_j(k, i; m, n)$ denotes the number of partitions enumerated by $C_j(k, i; n)$ with exactly m parts.

Theorem 1.7. For j = 0 or 1, $k \ge i \ge 2$ and (2k + j)/2 > 2,

$$\sum_{m,n\geq 0} C_j(k,i;m,n) x^m q^n = \sum_{N_1\geq \cdots\geq N_{k-1}\geq 0} \frac{q^{2(N_1^2+\cdots+N_{k-1}^2+N_i+\cdots+N_{k-1})}(-q^{1-2N_2};q^2)_{N_2}(-xq^{1+2N_1};q^2)_{\infty} x^{N_1+\cdots+N_{k-1}}}{(q^2;q^2)_{N_1-N_2}\cdots(q^2;q^2)_{N_{k-2}-N_{k-1}}(q^{4-2j};q^{4-2j})_{N_{k-1}}}.$$
(1.5)

This paper is organized as follows. In Section 2, we will give a proof of Theorem 1.6 with the aid of Bailey pairs. Section 3 is a preliminary for the proof of Theorem 1.7, in which we introduce the definition of the Göllnitz-Gordon marking of a partition, and give equivalent statements of Theorem 1.7. Section 4 is devoted to introducing the insertion operation and the separation operation. These operations allow us to provide a proof of Theorem 3.4. In order to show Theorem 3.5, we introduce the reduction operation and the dilation operation in Section 5. In light of these operations, we finally give a proof of Theorem 3.5 in Section 6.

2 Proof of Theorem 1.6

In this section, we will give a proof of Theorem 1.6 by using some related results on Bailey pairs. To this end, we first give a brief review of Bailey pairs. For more information on Bailey pairs, see, for example, [1, 5, 6, 8, 16, 17, 19]. Recall that a pair of sequences $(\alpha_n(a, q), \beta_n(a, q))$ is called a Bailey pair relative to (a, q) if for $n \ge 0$,

$$\beta_n(a,q) = \sum_{r=0}^n \frac{\alpha_r(a,q)}{(q;q)_{n-r}(aq;q)_{n+r}}$$

The following formulation of Bailey's lemma was given by Andrews [4, 5].

Theorem 2.1 (Bailey's lemma). If $(\alpha_n(a,q), \beta_n(a,q))$ is a Bailey pair relative to (a,q), then $(\alpha'_n(a,q), \beta'_n(a,q))$ is also a Bailey pair relative to (a,q), where

$$\alpha_n'(a,q) = \frac{(\rho_1;q)_n(\rho_2;q)_n}{(aq/\rho_1;q)_n(aq/\rho_2;q)_n} \left(\frac{aq}{\rho_1\rho_2}\right)^n \alpha_n(a,q),$$

$$\beta_n'(a,q) = \sum_{r=0}^n \frac{(\rho_1;q)_r(\rho_2;q)_r(aq/\rho_1\rho_2;q)_{n-r}}{(aq/\rho_1;q)_n(aq/\rho_2;q)_n(q;q)_{n-r}} \left(\frac{aq}{\rho_1\rho_2}\right)^r \beta_r(a,q).$$

When $\rho_1, \rho_2 \to \infty$ and $\rho_1 \to \infty, \rho_2 = -\sqrt{aq}$, Bailey's lemma reduces to the following two forms, which play an important role in proof of Theorem 1.5.

Lemma 2.2. If $(\alpha_n(a,q), \beta_n(a,q))$ is a Bailey pair relative to (a,q), then $(\alpha'_n(a,q), \beta'_n(a,q))$ is also a Bailey pair relative to (a,q), where

$$\alpha'_{n}(a,q) = a^{n}q^{n^{2}}\alpha_{n}(a,q),$$

$$\beta'_{n}(a,q) = \sum_{r=0}^{n} \frac{a^{r}q^{r^{2}}}{(q;q)_{n-r}}\beta_{r}(a,q)$$

Lemma 2.3. If $(\alpha_n(a,q), \beta_n(a,q))$ is a Bailey pair relative to (a,q), then $(\alpha'_n(a,q), \beta'_n(a,q))$ is also a Bailey pair relative to (a,q), where

$$\alpha'_n(a,q) = a^{n/2} q^{n^2/2} \alpha_n(a,q),$$

$$\beta'_n(a,q) = \sum_{r=0}^n \frac{(-\sqrt{aq};q)_r}{(-\sqrt{aq};q)_n(q;q)_{n-r}} a^{r/2} q^{r^2/2} \beta_r(a,q).$$

By successively using Lemma 2.2 and the following proposition, together with Lemma 2.3, Bressoud, Ismail and Stanton found a proof of Theorem 1.5.

Proposition 2.4. [8, Proposition 4.1] If $(\alpha_n(1,q), \beta_n(1,q))$ is a Bailey pair relative to (1,q), where

$$\alpha_n(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{An^2} (q^{(A-1)n} + q^{-(A-1)n}), & \text{if } n \ge 1, \end{cases}$$

then $(\alpha'_n(1,q), \beta'_n(1,q))$ is also a Bailey pair relative to (1,q), where

$$\alpha'_n(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{An^2} (q^{An} + q^{-An}), & \text{if } n \ge 1, \end{cases}$$
$$\beta'_n(1,q) = q^n \beta_n(1,q).$$

Using Lemma 2.2 followed by Proposition 2.4, we derive the following corollary.

Corollary 2.5. If $(\alpha_n(1,q), \beta_n(1,q))$ is a Bailey pair relative to (1,q), where

$$\alpha_n(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{An^2} (q^{An} + q^{-An}), & \text{if } n \ge 1, \end{cases}$$

then $(\alpha'_n(1,q),\beta'_n(1,q))$ is also a Bailey pair relative to (1,q), where

$$\alpha'_{n}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^{n} q^{(A+1)n^{2}} (q^{(A+1)n} + q^{-(A+1)n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta'_{n}(1,q) = q^{n} \sum_{r=0}^{n} \frac{q^{r^{2}}}{(q;q)_{n-r}} \beta_{r}(1,q).$$

Now we are ready to show Theorem 1.6.

Proof of Theorem 1.6. We begin with the Bailey pair $(\alpha_n^{(0)}(1,q), \beta_n^{(0)}(1,q))$ relative to (1,q), where

$$\alpha_n^{(0)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{jn^2/2} (q^{-jn/2} + q^{jn/2}), & \text{if } n \ge 1. \end{cases}$$

It is easy to see that when j = 1,

$$\beta_n^{(0)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ 0, & \text{if } n \ge 1. \end{cases}$$

The Bailey pair $(\alpha_n^{(0)}(1,q), \beta_n^{(0)}(1,q))$ corresponds to the unit Bailey pair [18, H(17)]. And when j = 0,

$$\beta_n^{(0)}(1,q) = \frac{(-1)^n}{(q^2;q^2)_n}.$$

It corresponds to the Bailey pair [18, E(2)].

Applying Corollary 2.5 to $(\alpha_n^{(0)}(1,q), \beta_n^{(0)}(1,q))$, we get

$$\begin{aligned} \alpha_n^{(1)}(1,q) &= \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{(j+2)n^2/2} (q^{-(j+2)n/2} + q^{(j+2)n/2}), & \text{if } n \ge 1, \end{cases} \\ \beta_n^{(1)}(1,q) &= q^n \sum_{N_k=0}^n \frac{q^{N_k^2}}{(q;q)_{n-N_k}} \beta_{N_k}^{(0)}(1,q). \end{aligned}$$

It is easy to check that

$$\beta_n^{(1)}(1,q) = \frac{q^n}{(q^{2-j};q^{2-j})_n},$$

in which the j = 0 case follows from the q-Chu-Vandermonde formula [9, Eq. (II.7)] with c = -q and $a \to \infty$,

$$\sum_{r=0}^{n} \frac{(a;q)_r (q^{-n};q)_r}{(c;q)_r (q;q)_r} \left(\frac{cq^n}{a}\right)^r = \frac{(c/a;q)_n}{(c;q)_n}.$$
(2.1)

Applying Corollary 2.5 k - i - 1 times to $(\alpha_n^{(1)}(1,q), \beta_n^{(1)}(1,q))$ yields the Bailey pair $(\alpha_n^{(k-i)}(1,q), \beta_n^{(k-i)}(1,q))$, where

$$\alpha_n^{(k-i)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{\frac{2k-2i+j}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta_n^{(k-i)}(1,q) = q^n \sum_{n \ge N_{i+1} \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_{i+1}^2 + \dots + N_{k-1}^2 + N_{i+1} + \dots + N_{k-1}}}{(q;q)_{n-N_{i+1}}(q;q)_{N_{i+1}-N_{i+2}} \cdots (q^{2-j};q^{2-j})_{N_{k-1}}}.$$

Next we substitute the Bailey pair $(\alpha_n^{(k-i)}(1,q), \beta_n^{(k-i)}(1,q))$ into Lemma 2.2 to get the following Bailey pair

$$\alpha_n^{(k-i+1)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{\frac{2k-2i+j+2}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta_n^{(k-i+1)}(1,q) = \sum_{n \ge N_i \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_i^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}}}{(q;q)_{n-N_i}(q;q)_{N_i - N_{i+1}} \cdots (q^{2-j};q^{2-j})_{N_{k-1}}}$$

Applying Lemma 2.2 to the Bailey pair $(\alpha_n^{(k-i+1)}(1,q), \beta_n^{(k-i+1)}(1,q))$ i-3 times, we will obtain the following Bailey pair

$$\alpha_n^{(k-2)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{\frac{2k+j-4}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta_n^{(k-2)}(1,q) = \sum_{n \ge N_3 \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_3^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}}}{(q;q)_{n-N_3}(q;q)_{N_3 - N_4} \cdots (q^{2-j};q^{2-j})_{N_{k-1}}}.$$

Plugging $(\alpha_n^{(k-2)}(1,q), \beta_n^{(k-2)}(1,q))$ into Lemma 2.3, we get the Bailey pair $(\alpha_n^{(k-1)}(1,q), \beta_n^{(k-1)}(1,q))$, where

$$\alpha_n^{(k-1)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{\frac{2k+j-3}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta_n^{(k-1)}(1,q) = \sum_{n \ge N_2 \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_2^2/2 + N_3^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}} (-q^{1/2};q)_{N_2}}{(-q^{1/2};q)_n (q;q)_{n-N_2} (q;q)_{N_2 - N_3} \cdots (q^{2-j};q^{2-j})_{N_{k-1}}}.$$

Then we apply Lemma 2.2 to the Bailey pair $(\alpha_n^{(k-1)}(1,q), \beta_n^{(k-1)}(1,q))$. This gives

$$\alpha_n^{(k)}(1,q) = \begin{cases} 1, & \text{if } n = 0, \\ (-1)^n q^{\frac{2k+j-1}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}), & \text{if } n \ge 1, \end{cases}$$
$$\beta_n^{(k)}(1,q) = \sum_{n \ge N_1 \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_1^2 + N_2^2/2 + N_3^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}} (-q^{1/2};q)_{N_2}}{(q;q)_{n-N_1}(q;q)_{N_1-N_2} \cdots (q^{2-j};q^{2-j})_{N_{k-1}} (-q^{1/2};q)_{N_1}}.$$

According to the definition of a Bailey pair, we have

$$\beta_n^{(k)}(1,q) = \sum_{r=0}^n \frac{\alpha_r^{(k)}(1,q)}{(q;q)_{n-r}(q;q)_{n+r}}.$$

Letting $n \to \infty$ and multiplying both sides by $(-q^{1/2}; q)_{\infty}(q; q)_{\infty}$, we obtain

$$\sum_{N_1 \ge \dots \ge N_{k-1} \ge 0} \frac{q^{N_1^2 + N_2^2/2 + N_3^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}} (-q^{1/2}; q)_{N_2} (-q^{1/2}; q)_{\infty}}{(q; q)_{N_1 - N_2} \cdots (q^{2-j}; q^{2-j})_{N_{k-1}} (-q^{1/2}; q)_{N_1}} \\ = \frac{(-q^{1/2}; q)_{\infty}}{(q; q)_{\infty}} \left(1 + \sum_{n=1}^{\infty} (-1)^n q^{\frac{2k+j-1}{2}n^2} (q^{-\frac{2k-2i+j}{2}n} + q^{\frac{2k-2i+j}{2}n}) \right) \\ = \frac{(-q^{1/2}; q)_{\infty} (q^{i-1/2}, q^{2k+j-i-1/2}, q^{2k+j-1}; q^{2k+j-1})_{\infty}}{(q; q)_{\infty}},$$
(2.2)

where the last equality follows from Jacobi's triple product identity.

Observing that

$$\frac{(-q^{1/2};q)_{\infty}}{(q;q)_{\infty}} = \frac{(q;q^2)_{\infty}}{(q^{1/2};q^{1/2})_{\infty}}, \quad \frac{(-q^{1/2};q)_{\infty}}{(-q^{1/2};q)_{N_1}} = (-q^{(2N_1+1)/2};q)_{\infty},$$

and

$$(-q^{1/2};q)_{N_2} = q^{N_2^2/2} (-q^{(1-2N_2)/2};q)_{N_2}$$

by replacing q by q^2 we derive (1.4). This completes the proof.

Remark 2.6. We point out some special cases of (1.4). For k = i, the Bailey pair $(\alpha_n^{(k-i)}(1,q), \beta_n^{(k-i)}(1,q))$ is actually $(\alpha_n^{(0)}(1,q), \beta_n^{(0)}(1,q))$. Thus following the remainder process in the above proof, we find that (1.4) reduces to the following forms.

Case 1. k = i = 2 and j = 1

$$\sum_{N_1 \ge 0} \frac{q^{2N_1^2}(-q^{1+2N_1};q^2)_{\infty}}{(q^2;q^2)_{N_1}} = \frac{(q^2;q^4)_{\infty}(q^3,q^5,q^8;q^8)_{\infty}}{(q;q)_{\infty}}$$

Case 2. $k = i \ge 3$

$$\sum_{N_1 \ge \dots \ge N_{k-1} \ge 0} \frac{q^{2(N_1^2 + \dots + N_{k-1}^2)} (-q^{1-2N_2}; q^2)_{N_2} (-q^{1+2N_1}; q^2)_{\infty}}{(q^2; q^2)_{N_1 - N_2} \cdots (q^2; q^2)_{N_{k-2} - N_{k-1}} (q^{4-2j}; q^{4-2j})_{N_{k-1}}}$$
$$= \frac{(q^2; q^4)_{\infty} (q^{2k-1}, q^{2k-1+2j}, q^{4k-2+2j}; q^{4k-2+2j})_{\infty}}{(q; q)_{\infty}}.$$

In addition, for k = i = 2 and j = 0, the case excluded by Theorem 1.6, using the above process we can also derive the following identity :

$$\sum_{N_1>0} \frac{q^{2N_1^2}(q;q^2)_{N_1}(-q^{1+2N_1};q^2)_{\infty}}{(q^4;q^4)_{N_1}} = \frac{(q^2;q^4)_{\infty}(q^3,q^3,q^6;q^6)_{\infty}}{(q;q)_{\infty}}$$

3 The Göllnitz-Gordon marking

In this section, we introduce the notion of the Göllnitz-Gordon marking of a partition. Furthermore, we investigate some attributes of a partition. And at the end of this section we give an explanation of Theorem 1.7 on the combinatorial side, which is stated in Theorems 3.4 and 3.5.

Definition 3.1 (Göllnitz-Gordon marking). The Göllnitz-Gordon marking $GG(\lambda)$ of a partition $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_{\ell})$ is an assignment of positive integers (marks) to the parts of λ from smallest to largest such that the marks are as small as possible subject to the condition that for $1 \leq s \leq \ell$, the integer assigned to λ_s is different from the integers assigned to the parts λ_g such that g > s and $\lambda_s - \lambda_g \leq 2$ with strict inequality if λ_s is odd.

For example, the Göllnitz-Gordon marking of

 $\lambda = (18, 15, 14, 14, 12, 11, 10, 8, 8, 6, 4, 3, 2, 1)$

is

$$GG(\lambda) = (18_1, 15_3, 14_2, 14_1, 12_3, 11_2, 10_1, 8_3, 8_2, 6_1, 4_3, 3_1, 2_2, 1_1).$$

The Göllnitz-Gordon marking of a partition can be represented by an array, where the column indicates the size of a part and the row (counted from bottom to top) indicates the mark, so the Göllnitz-Gordon marking of λ above would be:

$$GG(\lambda) = \begin{bmatrix} 4 & 8 & 12 & 15 \\ 2 & 8 & 11 & 14 \\ 1 & 3 & 6 & 10 & 14 & 18 \end{bmatrix}.$$
 (3.1)

Let $N_r(\lambda)$ (or N_r for short) denote the number of parts in the *r*-th row of $GG(\lambda)$. From the definition of the Göllnitz-Gordon marking, it is not hard to show that $N_1 \ge N_2 \ge \cdots$. By definition, we have $N_1(\lambda) = 6$, $N_2(\lambda) = 4$ and $N_3(\lambda) = 4$.

For $k \ge i \ge 1$, a partition λ is counted by $C_1(k, i; n)$ if and only if the marks of 1 and 2 are not exceed to i-1, the marks of the odd parts are at most 1, and there are at most k-1 rows in $GG(\lambda)$.

For $k \ge i \ge 1$ and j = 0 or 1, let $\mathbb{C}_j(k, i; n)$ denote the set of partitions counted by $C_j(k, i; n)$. Define

$$\mathbb{C}_j(k,i) = \bigcup_{n \ge 0} \mathbb{C}_j(k,i;n).$$

We wish to give a criterion to determine whether a partition in $\mathbb{C}_1(k, i)$ also belongs to $\mathbb{C}_0(k, i)$. Let $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_\ell)$ be a partition in $\mathbb{C}_1(k, i)$. From the definitions of $\mathbb{C}_1(k, i)$ and $\mathbb{C}_0(k, i)$, we see that λ belongs to $\mathbb{C}_0(k, i)$ if and only if λ satisfies the item (4) in Theorem 1.4, that is, for $1 \leq s \leq \ell - k + 2$, if

$$\lambda_s \le \lambda_{s+k-2} + 2$$
 with strict inequality if λ_s is odd, (3.2)

then

$$[\lambda_s/2] + \dots + [\lambda_{s+k-2}/2] \equiv i - 1 + O_\lambda(\lambda_s) \pmod{2}. \tag{3.3}$$

For the consecutive parts $\lambda_s, \lambda_{s+1}, \ldots, \lambda_{s+k-2}$ of λ , if such k-1 parts satisfy the difference condition (3.2), then we say that $\{\lambda_{s+l}\}_{0 \le l \le k-2}$ is a (k-1)-band of λ .

Let $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ be a (k-1)-band of λ . We say that the (k-1)-band $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ is good if the k-1 parts in $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ also satisfy the congruence condition (3.3), otherwise, we say that the (k-1)-band $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ is bad.

For a partition $\lambda \in \mathbb{C}_1(k, i)$, we see that λ is also a partition in $\mathbb{C}_0(k, i)$ if and only if all the (k-1)-bands of λ are good. Next we restrict our attention to certain special (k-1)-bands of λ to determine whether λ is a partition in $\mathbb{C}_0(k, i)$. Assume that there are N_{k-1} (k-1)-marked parts in $GG(\lambda)$, denoted $\lambda_1^{(k-1)} > \lambda_2^{(k-1)} > \cdots > \lambda_{N_{k-1}}^{(k-1)}$. For $1 \leq p \leq N_{k-1}$, assume that $\lambda_p^{(k-1)}$ is the s_p -th part λ_{s_p} of λ , there must exist k-2 parts λ_g in λ such that $g > s_p$ and $\lambda_g \geq \lambda_{s_p} - 2$ with strict inequality if λ_{s_p} is odd. Then $\{\lambda_{s_p+l}\}_{0 \leq l \leq k-2}$ is a (k-1)-band of λ . We say that $\{\lambda_{s_p+l}\}_{0 \leq l \leq k-2}$ by $\{\lambda_p^{(k-1)}\}_{k-1}$:

$$\{\lambda_{p,1}^{(k-1)}, \lambda_{p,2}^{(k-1)}, \dots, \lambda_{p,k-1}^{(k-1)}\},\$$

where $\lambda_{p,1}^{(k-1)} \geq \lambda_{p,2}^{(k-1)} \geq \cdots \geq \lambda_{p,k-1}^{(k-1)}$, that is, $\lambda_{p,1}^{(k-1)} = \lambda_p^{(k-1)}$ and $\lambda_p^{(k-1)} - \lambda_{p,k-1}^{(k-1)} \leq 2$ with strict inequality if $\lambda_p^{(k-1)}$ is odd.

The following theorem is the main result of this section, which gives a criterion to determine whether a partition λ in $\mathbb{C}_1(k, i)$ also belongs to $\mathbb{C}_0(k, i)$.

Theorem 3.2. For $k \geq i \geq 1$, let λ be a partition in $\mathbb{C}_1(k, i)$ with N_{k-1} (k-1)-marked parts in $GG(\lambda)$, denoted $\lambda_1^{(k-1)} > \lambda_2^{(k-1)} > \cdots > \lambda_{N_{k-1}}^{(k-1)}$. Then λ is a partition in $\mathbb{C}_0(k, i)$ if and only if $\{\lambda_p^{(k-1)}\}_{k-1}$ is good for $1 \leq p \leq N_{k-1}$.

To prove Theorem 3.2, we need the following lemma.

Lemma 3.3. For $k \geq i \geq 1$, let $\lambda = (\lambda_1, \lambda_2, ..., \lambda_\ell)$ be a partition in $\mathbb{C}_1(k, i)$, and let $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ and $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ be two (k-1)-bands of λ . If $\lambda_s < \lambda_g$ and $\lambda_{s+k-2} \geq \lambda_g - 4$ with strict inequality if λ_g is odd, then $\{\lambda_{s+l}\}_{0\leq l\leq k-2}$ and $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ are both good or bad.

Proof. Since $\lambda_s < \lambda_g$, then we have s > g. We assume that s = g + t where $t \ge 1$. Note that λ is a partition in $\mathbb{C}_1(k, i)$, then for $1 \le s \le \ell - k + 1$, we have

$$\lambda_s \ge \lambda_{s+k-1} + 2$$
 with strict inequality if λ_s is even. (3.4)

It follows that there are at most 2k - 2 parts of λ in $[\lambda_g - 4, \lambda_g]$ (resp. $(\lambda_g - 4, \lambda_g]$) if λ_g is even (resp. odd). So $1 \le t \le k - 1$. Since (s + k - 1 - t) - g = k - 1, using (3.4), we have

$$\lambda_g \ge \lambda_{s+k-1-t} + 2$$
 with strict inequality if λ_g is even. (3.5)

Note that $\{\lambda_{g+l}\}_{0 \le l \le k-2}$ is a (k-1)-band of λ , we have

$$\lambda_g \le \lambda_{g+k-2} + 2$$
 with strict inequality if λ_g is odd. (3.6)

Combining (3.5) and (3.6), we arrive at $\lambda_{s+k-1-t} < \lambda_{s+k-2-t}$. With the similar argument, we derive that $\lambda_{g+t} < \lambda_{g+t-1}$. To summarize, the overlapping structure of $\{\lambda_{s+l}\}_{0 \le l \le k-2}$ and $\{\lambda_{g+l}\}_{0 \le l \le k-2}$ can be described as follows. There are two cases.

If $1 \leq t < k-1$, then $\{\lambda_{s+l}\}_{0 \leq l \leq k-2}$ and $\{\lambda_{g+l}\}_{0 \leq l \leq k-2}$ have overlapping parts.

If t = k - 1, then $\{\lambda_{s+l}\}_{0 \le l \le k-2}$ and $\{\lambda_{g+l}\}_{0 \le l \le k-2}$ have the following relation:

$$\lambda_{s+k-2} \leq \cdots \leq \lambda_s < \lambda_{g+k-2} \leq \cdots \leq \lambda_g.$$

Set

$$[\lambda_{s+k-2}/2] + \dots + [\lambda_s/2] \equiv a + O_{\lambda}(\lambda_s) \pmod{2}, \tag{3.7}$$

where a = i - 1 or *i*. We aim to prove that for the same *a* in (3.7), we have

$$[\lambda_{g+k-2}/2] + \dots + [\lambda_g/2] \equiv a + O_{\lambda}(\lambda_g) \pmod{2}. \tag{3.8}$$

Note that $\lambda_{g+l} = \lambda_{s+l-t}$ for $t \leq l \leq k-2$, so

$$\begin{split} [\lambda_{g+k-2}/2] + \cdots + [\lambda_{g+t}/2] + [\lambda_{g+t-1}/2] + \cdots + [\lambda_g/2] \\ &= [\lambda_{s+k-2-t}/2] + \cdots + [\lambda_s/2] + [\lambda_{g+t-1}/2] + \cdots + [\lambda_g/2] \\ &= [\lambda_{s+k-2}/2] + \cdots + [\lambda_s/2] \\ &+ [\lambda_{g+t-1}/2] + \cdots + [\lambda_g/2] - ([\lambda_{s+k-2}/2] + \cdots + [\lambda_{s+k-1-t}/2]) \,. \end{split}$$

Combining with (3.7), we find that in order to prove (3.8), it suffices to show that

$$\begin{aligned} [\lambda_{g+t-1}/2] + \cdots + [\lambda_g/2] - ([\lambda_{s+k-2}/2] + \cdots + [\lambda_{s+k-1-t}/2]) \\ \equiv O_\lambda(\lambda_g) - O_\lambda(\lambda_s) \pmod{2}. \end{aligned}$$
(3.9)

We consider the following two cases.

Case 1: λ_g is even. We may write $\lambda_g = 2b + 4$. Under the condition that $\lambda_{s+k-2} \ge \lambda_g - 4$ with strict inequality if λ_g is odd, we have $\lambda_{s+k-2} \ge 2b$. Using (3.6), we have $\lambda_{g+k-2} \ge \lambda_g - 2 \ge 2b + 2$. Furthermore, $\lambda_{s+k-1-t} < 2b + 2$ and $\lambda_{g+t-1} > 2b + 2$. Hence we conclude that

$$2b \le \lambda_{s+k-2} \le \dots \le \lambda_{s+k-1-t} < 2b+2,$$

and

$$2b+2 < \lambda_{g+t-1} \le \dots \le \lambda_g = 2b+4.$$

It yields that for $k - 1 - t \leq l \leq k - 2$, $[\lambda_{s+l}/2] = b$, and for $0 \leq l \leq t - 1$,

$$[\lambda_{g+l}/2] = \begin{cases} b+1, & \text{if } \lambda_{g+l} \text{ is odd,} \\ b+2, & \text{otherwise.} \end{cases}$$

It follows that

$$[\lambda_{g+t-1}/2] + \dots + [\lambda_g/2] - ([\lambda_{s+k-2}/2] + \dots + [\lambda_{s+k-1-t}/2])$$

 $\equiv \text{the number of } \lambda_{g+l} \ (0 \le l \le t-1) \text{ which are odd} \pmod{2}.$

By the definitions of $O_{\lambda}(\lambda_q)$ and $O_{\lambda}(\lambda_s)$, we arrive at (3.9).

Case 2: λ_g is odd. We may write $\lambda_g = 2b + 3$. In such case, we have $\lambda_{g+k-2} = \cdots = \lambda_{g-1} = 2b + 2$. Under the condition that $\lambda_{s+k-2} \ge \lambda_g - 4$ with strict inequality if λ_g is odd, we have $\lambda_{s+k-2} \ge 2b$. Furthermore, we see that $\lambda_{s+k-3} = \cdots = \lambda_s = 2b + 2$. It follows that t = 1, $\lambda_{s+k-2} = 2b$ or 2b + 1, and $O_{\lambda}(\lambda_g) - O_{\lambda}(\lambda_s) = 1$. Thus we have

$$[\lambda_{g+t-1}/2] + \dots + [\lambda_g/2] - ([\lambda_{s+k-2}/2] + \dots + [\lambda_{s+k-1-t}/2])$$

= $[\lambda_g/2] - [\lambda_{s+k-2}/2] = 1 = O_{\lambda}(\lambda_g) - O_{\lambda}(\lambda_s) \pmod{2}.$

We arrive at (3.9). This completes the proof.

We are now in a position to give a proof of Theorem 3.2.

Proof of Theorem 3.2. If λ is a partition in $\mathbb{C}_0(k, i)$, then all the (k-1)-bands of λ are good. Thus the (k-1)-band $\{\lambda_p^{(k-1)}\}_{k-1}$ induced by $\lambda_p^{(k-1)}$ is good for $1 \leq p \leq N_{k-1}$.

On the other hand, we assume that $\{\lambda_p^{(k-1)}\}_{k-1}$ is good for $1 \leq p \leq N_{k-1}$. To prove λ is a partition in $\mathbb{C}_0(k, i)$, it suffices to show all the (k-1)-bands of λ are good. Given a (k-1)-band $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ of λ , that is, $\lambda_g \leq \lambda_{g+k-2} + 2$ with strict inequality if λ_g is odd. We aim to show $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ is good.

By the definition of the Göllnitz-Gordon marking, we see that the parts in $\{\lambda_{g+l}\}_{0 \le l \le k-2}$ have different marks in $GG(\lambda)$. Thus there is a part, say λ_{g+l_g} $(0 \le l_g \le k-2)$, marked with k-1 in $GG(\lambda)$. Assume that λ_{g+l_g} is the p_g -th (k-1)-marked part $\lambda_{p_g}^{(k-1)}$ in $GG(\lambda)$, then we have $\lambda_g - \lambda_{p_g}^{(k-1)} \le 2$ with strict inequality if λ_g is odd. Let

$$\{\lambda_{p_g,1}^{(k-1)}, \lambda_{p_g,2}^{(k-1)}, \dots, \lambda_{p_g,k-1}^{(k-1)}\}$$

be the (k-1)-band $\{\lambda_{p_g}^{(k-1)}\}_{k-1}$ induced by $\lambda_{p_g}^{(k-1)}$, that is, $\lambda_{p_g,1}^{(k-1)} = \lambda_{p_g}^{(k-1)}$, and $\lambda_{p_g}^{(k-1)} - \lambda_{p_g,k-1}^{(k-1)} \leq 2$ with strict inequality if $\lambda_{p_g}^{(k-1)}$ is odd. Hence

$$\lambda_g - \lambda_{p_g,k-1}^{(k-1)} = (\lambda_g - \lambda_{p_g}^{(k-1)}) + (\lambda_{p_g}^{(k-1)} - \lambda_{p_g,k-1}^{(k-1)}) \le 4$$

with strict inequality if λ_g is odd. By Lemma 3.3, we deduce that $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ and $\{\lambda_{p_g}^{(k-1)}\}_{k-1}$ are both good or bad in λ . Under the assumption that $\{\lambda_p^{(k-1)}\}_{k-1}$ is good for $1 \leq p \leq N_{k-1}$, we derive that $\{\lambda_{g+l}\}_{0\leq l\leq k-2}$ is good. It follows that λ is a partition in $\mathbb{C}_0(k, i)$. This completes the proof.

Now, we turn to give equivalent statements of Theorem 1.7.

For $k \geq i \geq 1$ and j = 0 or 1, let $\mathbb{E}_j(k, i)$ denote the number of partitions in $\mathbb{C}_j(k, i)$ without odd parts. For $N_1 \geq N_2 \geq \cdots \geq N_{k-1} \geq 0$, let $\mathbb{E}_j(N_1, \ldots, N_{k-1}; i)$ denote the set of partitions λ in $\mathbb{E}_j(k, i)$ with N_r parts in the *r*-th row of $GG(\lambda)$ for $1 \leq r \leq k-1$. Kurşungöz [15] established the following identity based on the Gordon marking of a partition.

$$\sum_{\lambda \in \mathbb{E}_j(N_1, \dots, N_{k-1}; i)} q^{\frac{|\lambda|}{2}} = \frac{q^{N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1}}}{(q; q)_{N_1 - N_2} \cdots (q; q)_{N_{k-2} - N_{k-1}} (q^{2-j}; q^{2-j})_{N_{k-1}}}$$

Then, we can see that

$$\sum_{\lambda \in \mathbb{E}_j(N_1, \dots, N_{k-1}; i)} q^{|\lambda|} = \frac{q^{2(N_1^2 + \dots + N_{k-1}^2 + N_i + \dots + N_{k-1})}}{(q^2; q^2)_{N_1 - N_2} \cdots (q^2; q^2)_{N_{k-2} - N_{k-1}} (q^{4-2j}; q^{4-2j})_{N_{k-1}}}.$$

It follows that

$$\sum_{\lambda \in \mathbb{E}_{j}(k,i)} x^{\ell(\lambda)} q^{|\lambda|} = \sum_{N_{1} \ge \dots \ge N_{k-1} \ge 0} x^{N_{1} + \dots + N_{k-1}} \sum_{\lambda \in \mathbb{E}_{j}(N_{1},\dots,N_{k-1};i)} q^{|\lambda|}$$
$$= \sum_{N_{1} \ge \dots \ge N_{k-1} \ge 0} \frac{q^{2(N_{1}^{2} + \dots + N_{k-1}^{2} + N_{i} + \dots + N_{k-1})} x^{N_{1} + \dots + N_{k-1}}}{(q^{2};q^{2})_{N_{1} - N_{2}} \cdots (q^{2};q^{2})_{N_{k-2} - N_{k-1}} (q^{4-2j};q^{4-2j})_{N_{k-1}}}.$$

For $N \ge 0$, let \mathbb{I}_N denote the set of partitions $\eta = (\eta_1, \eta_2, \ldots, \eta_\ell)$ with distinct odd parts greater than or equal to 2N + 1, that is, $\eta_1 > \eta_2 > \cdots > \eta_\ell \ge 2N + 1$. The generating functions for partitions in \mathbb{I}_N is

$$\sum_{\eta \in \mathbb{I}_N} x^{\ell(\eta)} q^{|\eta|} = (1 + xq^{2N+1})(1 + xq^{2N+3}) \dots = (-xq^{2N+1}; q^2)_{\infty}$$

We define $\mathbb{F}_1(2,2)$ to be set of pairs $(\lambda, -, \eta)$ of partitions such that

$$\lambda \in \mathbb{E}_1(2,2)$$
 and $\eta \in \mathbb{I}_{N_1(\lambda)}$.

Then for k = i = 2 and j = 1, Theorem 1.7 is equivalent to the following combinatorial statement.

Theorem 3.4. There is a bijection $\Phi_{2,2}$ between $\mathbb{F}_1(2,2)$ and $\mathbb{C}_1(2,2)$. Moreover, for a pair $(\lambda, -, \eta) \in \mathbb{F}_1(2,2)$, we have $\pi = \Phi_{2,2}(\lambda, -, \eta) \in \mathbb{C}_1(2,2)$ such that

$$|\pi| = |\lambda| + |\eta|$$
 and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

For $N \ge 0$, let \mathbb{O}_N denote the set of partitions $\tau = (\tau_1, \tau_2, \ldots, \tau_\ell)$ with distinct negative odd parts which lie in [1 - 2N, -1], that is, $-1 \ge \tau_1 > \tau_2 > \cdots > \tau_\ell \ge 1 - 2N$. Clearly, the generating functions for partitions in \mathbb{O}_N is

$$\sum_{\tau \in \mathbb{O}_N} q^{|\tau|} = (1+q^{1-2N})(1+q^{3-2N})\cdots(1+q^{-1}) = (-q^{1-2N};q^2)_N$$

Then for $k \geq 3$, $k \geq i \geq 2$ and j = 0 or 1, we define $\mathbb{F}_j(k, i)$ to be set of triplets (λ, τ, η) of partitions such that

$$\lambda \in \mathbb{E}_i(k, i), \tau \in \mathbb{O}_{N_2(\lambda)} \text{ and } \eta \in \mathbb{I}_{N_1(\lambda)}.$$

In this case, Theorem 1.7 is equivalent to the following combinatorial statement.

Theorem 3.5. For $k \geq 3$, $k \geq i \geq 2$ and j = 0 or 1, there is a bijection $\Phi_{k,i}$ between $\mathbb{F}_j(k,i)$ and $\mathbb{C}_j(k,i)$, namely, for a triplet $(\lambda, \tau, \eta) \in \mathbb{F}_j(k,i)$, we have $\pi = \Phi_{k,i}(\lambda, \tau, \eta) \in \mathbb{C}_j(k,i)$ such that

$$|\pi| = |\lambda| + |\tau| + |\eta| \text{ and } \ell(\pi) = \ell(\lambda) + \ell(\eta)$$

4 Proof of Theorem 3.4

The main objective of this section is to give a proof of Theorem 3.4. To this end, we construct the bijection $\theta_{(p)}$ and the combination operation, where $\theta_{(p)}$ is essentially the same as that defined in Section 5 in [14]. Composing the combination operation with $\theta_{(p)}$ gives the insertion operation which is the main ingredient for the proof of Theorem 3.4.

4.1 The bijection $\theta_{(p)}$

In this subsection, we modify the results in Section 5 in [14], which have been proved for j = 1. We just show that the results in this subsection also hold for j = 0. Throughout this subsection, we assume that $k \ge i \ge 1$, j = 0 or 1, and $N_1 \ge \cdots \ge N_{k-1} \ge 0$.

Let $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$ denote the set of partitions λ in $\mathbb{C}_j(k, i)$ such that there are N_r (k-1)-marked parts in $GG(\lambda)$ for $1 \leq r \leq k-1$. Let λ be a partition in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$. We use $\lambda^{(1)} = (\lambda_1^{(1)}, \lambda_2^{(1)}, \ldots, \lambda_{N_1}^{(1)})$ to denote the 1-marked parts in $GG(\lambda)$, where $\lambda_1^{(1)} > \lambda_2^{(1)} > \cdots > \lambda_{N_1}^{(1)}$. We now divide the parts of $\lambda^{(1)}$ into the following two types. For $1 \leq s \leq N_1$, we say $\lambda_s^{(1)}$ is an odd type if $\lambda_s^{(1)}$ is an odd part, or there is an odd part $\lambda_s^{(1)} + 1$ in λ , otherwise, we say $\lambda_s^{(1)}$ is an even type.

For $1 \leq p \leq N_1$, we prepare three subsets $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$, $\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ of $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$, which are described by using the subpartition $\lambda^{(1)} = (\lambda_1^{(1)}, \lambda_2^{(1)}, \ldots, \lambda_{N_1}^{(1)})$ of a partition λ in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$.

 \diamond Let $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ be the set of partitions λ in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$, where $\lambda_p^{(1)}$ is odd type and $\lambda_s^{(1)}$ is even type for $1 \le p \le N_1$ and $1 \le s \le p-1$.

 \Diamond Let $\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ be the set of partitions λ in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$, where $\lambda_p^{(1)}$ is even type, $\lambda_{p-1}^{(1)}$ is odd type, and $\lambda_s^{(1)}$ is even type for $1 and <math>1 \le s \le p-2$.

 \diamond Let $\overrightarrow{\mathbb{C}}_{j}(N_{1},\ldots,N_{k-1};i|p)$ be the set of partitions λ in $\mathbb{C}_{j}(N_{1},\ldots,N_{k-1};i)$, where $\lambda_{s}^{(1)}$ is even type for $1 \leq p \leq N_{1}$ and $1 \leq s \leq p$.

We give a bijection Θ_p between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ for $1 , a bijection <math>\Theta_1$ between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|1)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|1)$, and a bijection $\Theta_{(p)}$ between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ for $1 \leq p \leq N_1$.

Lemma 4.1. For $1 , there is a bijection <math>\Theta_p$ between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$. Furthermore, for $\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$, we have $\mu = \Theta_p(\lambda) \in \overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ and $|\mu| = |\lambda| + 2$.

To show Lemma 4.1, for $1 , we divide <math>\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ into two disjoint subsets $\mathbb{C}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\mathbb{C}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$, and divide $\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ into two disjoint subsets $\overline{\mathbb{C}}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$. We then build the bijection Θ_p consisting of two bijections $\Theta_{l,p}$ between $\mathbb{C}_j^{(l)}(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j^{(l)}(N_1, \ldots, N_{k-1}; i|p)$, where $1 \leq l \leq 2$.

For $1 , we divide <math>\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ into the following two disjoint subsets $\mathbb{C}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\mathbb{C}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$. For $\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$, we set r to be the mark of the odd part $\lambda_p^{(1)} + 1$ in $GG(\lambda)$ when $\lambda_p^{(1)}$ is an even part. It is easy to see that $r \geq 2$.

- (1) $\mathbb{C}_{j}^{(1)}(N_{1},\ldots,N_{k-1};i|p)$ is the set of partitions λ in $\mathbb{C}_{j}(N_{1},\ldots,N_{k-1};i|p)$ such that there exists a 1-marked even part $\lambda_{p}^{(1)} + 3$ in $GG(\lambda)$ when $\lambda_{p}^{(1)}$ is an odd part or there exists a *r*-marked even part $\lambda_{p}^{(1)} + 4$ in $GG(\lambda)$ when $\lambda_{p}^{(1)}$ is an even part.
- (2) $\mathbb{C}_{j}^{(2)}(N_{1},\ldots,N_{k-1};i|p)$ is the set of partitions λ in $\mathbb{C}_{j}(N_{1},\ldots,N_{k-1};i|p)$ such that there does not exist a 1-marked even part $\lambda_{p}^{(1)} + 3$ in $GG(\lambda)$ when $\lambda_{p}^{(1)}$ is an odd part or there does not exist a *r*-marked even part $\lambda_{p}^{(1)} + 4$ in $GG(\lambda)$ when $\lambda_{p}^{(1)}$ is an even part.

For $1 , we divide <math>\overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ into the following two disjoint subsets $\overline{\mathbb{C}}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$. For $\mu \in \overline{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i, p)$, if $\mu_{p-1}^{(1)}$ is an even part, then set r' to be the mark of the odd part $\mu_{p-1}^{(1)} + 1$ in $GG(\mu)$. It is easy to see that $\mu_{p-1}^{(1)} \ge 5$ and $r' \ge 2$.

- (1) $\overline{\mathbb{C}}_{j}^{(1)}(N_{1},\ldots,N_{k-1};i|p)$ is the set of partitions μ in $\overline{\mathbb{C}}_{j}(N_{1},\ldots,N_{k-1};i|p)$ such that there exists a 1-marked even part $\mu_{p-1}^{(1)} - 3$ in $GG(\mu)$ if $\mu_{p-1}^{(1)}$ is an odd part or there exists a r'-marked even part $\mu_{p-1}^{(1)} - 2$ in $GG(\mu)$ if $\mu_{p-1}^{(1)}$ is an even part.
- (2) $\overline{\mathbb{C}}_{j}^{(2)}(N_{1},\ldots,N_{k-1};i|p)$ is the set of partitions μ in $\overline{\mathbb{C}}_{j}(N_{1},\ldots,N_{k-1};i|p)$ such that there does not exist a 1-marked even part $\mu_{p-1}^{(1)} - 3$ in $GG(\mu)$ if $\mu_{p-1}^{(1)}$ is an odd part or there does not exist a r'-marked even part $\mu_{p-1}^{(1)} - 2$ in $GG(\mu)$ if $\mu_{p-1}^{(1)}$ is an even part.

Lemma 4.2. For $1 , there is a bijection <math>\Theta_{1,p}$ between $\mathbb{C}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$. Moreover, for $\lambda \in \mathbb{C}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$ and $\mu = \Theta_{1,p}(\lambda) \in \overline{\mathbb{C}}_j^{(1)}(N_1, \ldots, N_{k-1}; i|p)$, we have $|\mu| = |\lambda| + 2$.

Proof. If $\lambda_p^{(1)}$ is an odd part, set $\lambda_p^{(1)} = 2t + 1$, then there is a 1-marked 2t + 4 in $GG(\lambda)$, that is, $\lambda_{p-1}^{(1)} = 2t + 4$; if $\lambda_p^{(1)}$ is an even part, set $\lambda_p^{(1)} = 2t$, then there is a *r*-marked 2t + 1 and a *r*-marked 2t + 4 in $GG(\lambda)$. For $1 , define <math>\mu = \Theta_{1,p}(\lambda)$ as follows.

- If $\lambda_p^{(1)} = 2t + 1$, then replace $\lambda_p^{(1)} = 2t + 1$ by 2t + 2 and replace $\lambda_{p-1}^{(1)} = 2t + 4$ by 2t + 5.
- If $\lambda_p^{(1)} = 2t$, then replace the *r*-marked 2t + 1 by 2t + 2 and replace the *r*-marked 2t + 4 by 2t + 5.

Let $\lambda^{(k-1)} = (\lambda_1^{(k-1)}, \lambda_2^{(k-1)}, \dots, \lambda_{N_{k-1}}^{(k-1)})$ and $\mu^{(k-1)} = (\mu_1^{(k-1)}, \mu_2^{(k-1)}, \dots, \mu_{N_{k-1}}^{(k-1)})$ be the (k-1)-marked parts in $GG(\lambda)$ and $GG(\mu)$, respectively. We just need to show that if λ is a partition in $\mathbb{C}_0^{(1)}(N_1, \dots, N_{k-1}; i|p)$, then $\{\mu_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$. Assume that λ is a partition in $\mathbb{C}_0^{(1)}(N_1, \dots, N_{k-1}; i, p)$, then by Theorem 3.2 we have $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$.

For each $s \in [1, N_{k-1}]$, by the construction of μ , we see that 2t + 1 and 2t + 3 do not occur in μ , and so $\mu_s^{(k-1)} \neq 2t + 1$, 2t + 3. To show that $\{\mu_s^{(k-1)}\}_{k-1}$ is good, we consider the following five cases.

Case 1: $\mu_s^{(k-1)} \leq 2t$ or $\mu_s^{(k-1)} \geq 2t + 7$. By the construction of μ , we see that $\{\mu_s^{(k-1)}\}_{k-1} = \{\lambda_s^{(k-1)}\}_{k-1}$ and $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)})$. Note that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good, so we get that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 2: $\mu_s^{(k-1)} = 2t + 2$. By the construction of μ , we see that $f_{2t}(\mu) = f_{2t}(\lambda)$, $f_{2t+1}(\mu) = f_{2t+1}(\lambda) - 1$, $f_{2t+2}(\mu) = f_{2t+2}(\lambda) + 1$ and $O_{\mu}(2t+2) = O_{\lambda}(2t+2) - 1$. We have

$$\begin{aligned} [\mu_{s,1}^{(k-1)}/2] + [\mu_{s,2}^{(k-1)}/2] + \cdots + [\mu_{s,k-1}^{(k-1)}/2] \\ &= tf_{2t}(\mu) + tf_{2t+1}(\mu) + (t+1)f_{2t+2}(\mu) \\ &= tf_{2t}(\lambda) + t(f_{2t+1}(\lambda) - 1) + (t+1)(f_{2t+2}(\lambda) + 1) \\ &= tf_{2t}(\lambda) + tf_{2t+1}(\lambda) + (t+1)f_{2t+2}(\lambda) + 1 \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] + 1 \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) + 1 = i - 1 + O_{\lambda}(2t+2) + 1 \\ &\equiv i - 1 + O_{\mu}(2t+2) = i - 1 + O_{\mu}(\mu_{s,1}^{(k-1)}) \pmod{2}. \end{aligned}$$

So, $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 3: $\mu_s^{(k-1)} = 2t + 4$. In this case, we have $\mu_s^{(k-1)} = \lambda_s^{(k-1)} = 2t + 4$, $f_{2t+2}(\mu) = f_{2t+2}(\lambda) + 1$, $f_{2t+3}(\mu) = f_{2t+3}(\lambda) = 0$, $f_{2t+4}(\mu) = f_{2t+4}(\lambda) - 1$ and $O_{\mu}(2t+4) = O_{\lambda}(2t+4) - 1$.

So, we obtain that

$$\begin{aligned} [\mu_{s,1}^{(k-1)}/2] + [\mu_{s,2}^{(k-1)}/2] + \cdots + [\mu_{s,k-1}^{(k-1)}/2] \\ &= (t+1)f_{2t+2}(\mu) + (t+2)f_{2t+4}(\mu) \\ &= (t+1)(f_{2t+2}(\lambda)+1) + (t+2)(f_{2t+4}(\lambda)-1) \\ &= (t+1)f_{2t+2}(\lambda) + (t+2)f_{2t+4}(\lambda) - 1 \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] - 1 \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) - 1 = i - 1 + O_{\lambda}(2t+4) - 1 \\ &= i - 1 + O_{\mu}(2t+4) = i - 1 + O_{\mu}(\mu_{s,1}^{(k-1)}) \pmod{2}. \end{aligned}$$

It follows that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 4: $\mu_s^{(k-1)} = 2t + 5$. By the definition of Göllnitz-Gordon marking, we see that $\mu_{s,2}^{(k-1)} = \cdots = \mu_{s,k-1}^{(k-1)} = 2t + 4$. Moreover, by the construction of μ , we have $\lambda_{s,1}^{(k-1)} = \lambda_{s,2}^{(k-1)} = \cdots = \lambda_{s,k-1}^{(k-1)} = 2t + 4$ and $O_{\mu}(2t+5) = O_{\lambda}(2t+4)$. So

$$\begin{aligned} [\mu_{s,1}^{(k-1)}/2] + [\mu_{s,2}^{(k-1)}/2] + \cdots + [\mu_{s,k-1}^{(k-1)}/2] \\ &= (t+2)f_{2t+4}(\mu) + (t+2)f_{2t+5}(\mu) = (t+2)(k-2) + (t+2) \\ &= (t+2)(k-1) = (t+2)f_{2t+4}(\lambda) \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) = i - 1 + O_{\lambda}(2t+4) \\ &= i - 1 + O_{\mu}(2t+5) = i - 1 + O_{\mu}(\mu_{s,1}^{(k-1)}) \pmod{2}, \end{aligned}$$

which implies that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 5: $\mu_s^{(k-1)} = 2t + 6$. By the construction of μ , we see that $\mu_s^{(k-1)} = \lambda_s^{(k-1)} = 2t + 6$, $f_{2t+4}(\mu) = f_{2t+4}(\lambda) - 1$, $f_{2t+5}(\mu) = f_{2t+5}(\lambda) + 1$, $f_{2t+6}(\mu) = f_{2t+6}(\lambda)$ and $O_{\mu}(2t+6) = O_{\lambda}(2t+6)$. We have

$$\begin{aligned} [\mu_{s,1}^{(k-1)}/2] + [\mu_{s,2}^{(k-1)}/2] + \cdots + [\mu_{s,k-1}^{(k-1)}/2] \\ &= (t+2)f_{2t+4}(\mu) + (t+2)f_{2t+5}(\mu) + (t+3)f_{2t+6}(\mu) \\ &= (t+2)(f_{2t+4}(\lambda) - 1) + (t+2)(f_{2t+5}(\lambda) + 1) + (t+3)f_{2t+6}(\lambda) \\ &= (t+2)f_{2t+4}(\lambda) + (t+2)f_{2t+5}(\lambda) + (t+3)f_{2t+6}(\lambda) \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) = i - 1 + O_{\lambda}(2t+6) \\ &= i - 1 + O_{\mu}(2t+6) = i - 1 + O_{\mu}(\mu_{s,1}^{(k-1)}) \pmod{2}. \end{aligned}$$

Therefore, $\{\mu_s^{(k-1)}\}_{k-1}$ is good. Hence, we conclude that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

To prove that $\Theta_{1,p}$ is a bijection, we construct the inverse map $\Lambda_{1,p}$ of $\Theta_{1,p}$. If $\mu_{p-1}^{(1)}$ is an odd part, set $\mu_{p-1}^{(1)} = 2t + 5$, then there is a 1-marked 2t + 2 in $GG(\mu)$. If $\mu_{p-1}^{(1)}$ is an even part, set $\mu_{p-1}^{(1)} = 2t + 4$, then there are r'-marked 2t + 2 and r'-marked 2t + 5 in $GG(\mu)$. For $1 , define <math>\lambda = \Lambda_{1,p}(\mu)$ as follows.

- If $\mu_{p-1}^{(1)} = 2t + 5$, then replace $\mu_p^{(1)} = 2t + 2$ by 2t + 1 and replace $\mu_{p-1}^{(1)} = 2t + 5$ by 2t + 4.
- If $\mu_{p-1}^{(1)} = 2t+4$, then replace the r'-marked 2t+2 by 2t+1 and replace the r'-marked 2t+5 by 2t+4.

It can be verified that $\Lambda_{1,p}$ is the inverse map of $\Theta_{1,p}$. This completes the proof.

Lemma 4.3. For $1 , there is a bijection <math>\Theta_{2,p}$ between $\mathbb{C}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$ and $\overline{\mathbb{C}}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$. Moreover, for $\lambda \in \mathbb{C}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$ and $\mu = \Theta_{2,p}(\lambda) \in \overline{\mathbb{C}}_j^{(2)}(N_1, \ldots, N_{k-1}; i|p)$, we have $|\mu| = |\lambda| + 2$.

Proof. If $\lambda_p^{(1)}$ is an odd part, set r = 1 and $\lambda_p^{(1)} = 2t + 1$, then there does not exist a 1-marked 2t + 4 in $GG(\lambda)$, that is $\lambda_{p-1}^{(1)} \ge 2t + 6$. If $\lambda_p^{(1)}$ is an even part, set $\lambda_p^{(1)} = 2t$, then there is a *r*-marked 2t + 1 and there does not exist a *r*-marked 2t + 4 in $GG(\lambda)$, where $r \ge 2$. Let $\lambda_{p-1}^{(1)} = 2b + 2$ and let *l* be the largest mark of the parts 2b + 2 in $GG(\lambda)$. If $\lambda_p^{(1)} = 2t + 1$, then $b \ge t + 2$. If $\lambda_p^{(1)} = 2t$, then $b \ge t + 1$. For $1 , define <math>\mu = \Theta_{2,p}(\lambda)$ as follows.

- If r = 1, then replace $\lambda_p^{(1)} = 2t + 1$ by 2t + 2 and replace the *l*-marked 2b + 2 by 2b + 3.
- If $r \ge 2$, then replace the *r*-marked 2t + 1 by 2t + 2 and replace the *l*-marked 2b + 2 by 2b + 3.

Let $\lambda^{(k-1)} = (\lambda_1^{(k-1)}, \lambda_2^{(k-1)}, \dots, \lambda_{N_{k-1}}^{(k-1)})$ and $\mu^{(k-1)} = (\mu_1^{(k-1)}, \mu_2^{(k-1)}, \dots, \mu_{N_{k-1}}^{(k-1)})$ be the (k-1)-marked parts in $GG(\lambda)$ and $GG(\mu)$, respectively. We just need to show that if λ is a partition in $\mathbb{C}_0^{(2)}(N_1, \dots, N_{k-1}; i|p)$, then $\{\mu_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$. Assume that λ is a partition in $\mathbb{C}_0^{(2)}(N_1, \dots, N_{k-1}; i, p)$, then by Theorem 3.2 it follows that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$.

For each $s \in [1, N_{k-1}]$, by the construction of μ , we see that 2t + 1 and 2t + 3 do not occur in μ , and so $\mu_s^{(k-1)} \neq 2t+1$, 2t+3. Then, we proceed to show that $\mu_s^{(k-1)} \neq 2t+4$. If $\lambda_p^{(1)} = 2t + 1$, then there is no 1-marked 2t+2, 2t+3 and 2t+4 in $GG(\lambda)$. If $\lambda_p^{(1)} = 2t$, then there is no *r*-marked 2t+2, 2t+3 and 2t+4 in $GG(\lambda)$. By definition of Göllnitz-Gordon marking, we see that there is no (k-1)-marked 2t+4 in $GG(\lambda)$. By the construction of μ , we see that there is no (k-1)-marked 2t+4 in $GG(\mu)$, and so $\mu_s^{(k-1)} \neq 2t+4$. Now, we turn to show that if $\mu_s^{(k-1)} \in [2t+2, 2b+2]$, then $\mu_s^{(k-1)} = 2t+2$. If b = t+1, then recall that $\mu_s^{(k-1)} \neq 2t+3$, 2t+4, we have $\mu_s^{(k-1)} = 2t+2$. If $b \ge t+2$, then by the construction of μ , we have $\mu_p^{(1)} \le 2t+2$, $\mu_{p-1}^{(1)} \ge 2b+2$ and $\mu_s^{(k-1)} \neq 2b+2$, which implies that $\mu_s^{(k-1)} \notin [2t+5, 2b+2]$. Again by $\mu_s^{(k-1)} \neq 2t+3$ and 2t+4, we have $\mu_s^{(k-1)} = 2t+2$. Hence we obtain that $\mu_s^{(k-1)} = 2t+2$ if $\mu_s^{(k-1)} \in [2t+2, 2b+2]$. Then, with the similar argument in the proof of Lemma 4.2, we can obtain that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

To prove that $\Theta_{2,p}$ is a bijection, we construct the inverse map $\Lambda_{2,p}$ of $\Theta_{2,p}$. If $\mu_{p-1}^{(1)}$ is an odd part, then set l' = 1, $\mu_{p-1}^{(1)} = 2b + 3$ and $\mu_p^{(1)} = 2t$, where $b \ge t + 1$. If $\mu_{p-1}^{(1)}$ is an even part, set $\mu_{p-1}^{(1)} = 2b + 2$ and $\mu_p^{(1)} = 2t$ where $b \ge t + 1$, then there is a l'-marked 2b + 3 and there is no l'-marked 2b in $GG(\mu)$, where $l' \ge 2$. If there exist parts 2t + 2 in $GG(\mu)$, where $r' \ge 2$, otherwise we set r' = 1. For $1 , define <math>\lambda = \Lambda_{2,p}(\mu)$ as follows.

- If r' = 1, then replace $\mu_p^{(1)} = 2t$ by 2t 1 and replace the *l'*-marked 2b + 3 by 2b + 2.
- If $r' \ge 2$, then replace the r'-marked 2t + 2 by 2t + 1 and replace the l'-marked 2b + 3 by 2b + 2.

It can be verified that $\Lambda_{2,p}$ is the inverse map of $\Theta_{2,p}$. This completes the proof.

Proof of Lemma 4.1. For $1 and <math>\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$, define

$$\mu = \Theta_p(\lambda) = \begin{cases} \Theta_{1,p}(\lambda), & \text{if } \lambda \in \mathbb{C}_j^{(1)}(N_1, \dots, N_{k-1}; i|p); \\ \Theta_{2,p}(\lambda), & \text{if } \lambda \in \mathbb{C}_j^{(2)}(N_1, \dots, N_{k-1}; i|p). \end{cases}$$

Combining Lemmas 4.2 and 4.3, we complete the proof of Lemma 4.1.

Lemma 4.4. There is a bijection Θ_1 between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|1)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|1)$. [1]. Moreover, for $\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|1)$ and $\mu = \Theta_1(\lambda) \in \overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|1)$, we have $|\mu| = |\lambda| + 1$.

Proof. By definition, we see that $\lambda_1^{(1)}$ is odd type. Let r be the mark of the odd part $\lambda_1^{(1)}$ (resp. $\lambda_1^{(1)}+1$) if $\lambda_1^{(1)}$ is odd (resp. even) in $GG(\lambda)$. More precisely, if $\lambda_1^{(1)}$ is an odd part, set $\lambda_1^{(1)} = 2t + 1$, then r = 1. If $\lambda_1^{(1)}$ is an even part, set $\lambda_1^{(1)} = 2t$, then there is r-marked 2t + 1 in $GG(\lambda)$. Define $\mu = \Theta_1(\lambda)$ by replacing the r-marked 2t + 1 by 2t + 2.

Let $\lambda^{(k-1)} = (\lambda_1^{(k-1)}, \lambda_2^{(k-1)}, \dots, \lambda_{N_{k-1}}^{(k-1)})$ and $\mu^{(k-1)} = (\mu_1^{(k-1)}, \mu_2^{(k-1)}, \dots, \mu_{N_{k-1}}^{(k-1)})$ be the (k-1)-marked parts in $GG(\lambda)$ and $GG(\mu)$, respectively. We just need to show that if λ is a partition in $\mathbb{C}_0(N_1, \dots, N_{k-1}; i|1)$, then $\{\mu_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$. Assume that λ is a partition in $\mathbb{C}_0(N_1, \dots, N_{k-1}; i|1)$, then $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$.

For each $s \in [1, N_{k-1}]$, by the construction of μ , we have $\mu_s^{(k-1)} \leq 2t+2$ and $\mu_s^{(k-1)} \neq 2t+1$. To show that $\{\mu_s^{(k-1)}\}_{k-1}$ is good, we consider the following two cases.

Case 1: $\mu_s^{(k-1)} \leq 2t$. By the construction of μ , we have $\{\mu_s^{(k-1)}\}_{k-1} = \{\lambda_s^{(k-1)}\}_{k-1}$ and $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)})$. Note that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good, so $\{\mu_s^{(k-1)}\}_{k-1}$ is also good.

Case 2: $\mu_s^{(k-1)} = 2t + 2$. With the similar argument of Case 2 in the proof of Lemma 4.2, we can obtain that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

In either case, we obtain that $\{\mu_s^{(k-1)}\}_{k-1}$ is good. To prove that Θ_1 is a bijection, we construct the inverse map Λ_1 of Θ_1 . By definition, $\mu_1^{(1)}$ is even type, set $\mu_1^{(1)} = 2t$. Let r' be the smallest mark of the largest even parts in $GG(\mu)$, that is, if there exist even parts 2t + 2 in μ , then $r' \geq 2$; otherwise, r' = 1. Define $\lambda = \Lambda_1(\mu)$ as follows.

- If r' = 1, then replace $\mu_1^{(1)} = 2t$ by 2t 1.
- If $r' \ge 2$, then replace the r'-marked 2t + 2 by 2t + 1.

It can be verified that Λ_1 is the inverse map of Θ_1 . This completes the proof.

Iterating the bijections in Lemma 4.1 p-1 times as well as the bijection Θ_1 in Lemma 4.4, we get a bijection $\Theta_{(p)}$ between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$.

Lemma 4.5. For $1 \leq p \leq N_1$, there is a bijection $\Theta_{(p)}$ between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$. Moreover, for $\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\mu = \Theta_{(p)}(\lambda) \in \overrightarrow{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, we have $|\mu| = |\lambda| + 2p - 1$.

Proof. Define $\Theta_{(p)} = \Theta_1 \Theta_2 \cdots \Theta_p$, then by Lemmas 4.1 and 4.4, we complete the proof.

4.2 The combination $C_{p,t}$ and the division $D_{p,t}$

In this subsection, we assume that j = 0 or 1, (2k + j)/2 > 2, $k \ge i \ge 2$, $N_1 \ge \cdots \ge N_{k-1} \ge 0$, $t \ge 0$ and $0 \le p \le N_1$. Let λ be a partition in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$. For convenience, we set $\lambda_0^{(1)} = +\infty$ and $\lambda_{N_1+1}^{(1)} = -\infty$. We will prepare two sets and then establish a bijection between these two subsets.

Let $\mathbb{C}_{j}^{<}(N_{1},\ldots,N_{k-1};i|p,t)$ denote the set of partitions λ in $\mathbb{C}_{j}(N_{1},\ldots,N_{k-1};i)$ satisfying the following conditions:

- (1) $\lambda_{p+1}^{(1)} + 2 \le 2t + 1 < \lambda_p^{(1)};$
- (2) if $p \ge 1$, then $\lambda \in \overrightarrow{\mathbb{C}}_j(N_1, \dots, N_{k-1}; i|p)$.

It is worth mentioning that if $\mathbb{C}_{j}^{<}(N_{1},\ldots,N_{k-1};i|p,t)$ is non-empty, then $p+t \geq N_{1}$. If $p = N_{1}$, then it is obviously right. If $p < N_{1}$, then by the definition of Göllnitz-Gordon marking, we see that

$$2t+1 \ge \lambda_{p+1}^{(1)}+2 \ge \dots \ge \lambda_{N_1}^{(1)}+2(N_1-p) \ge 2(N_1-p)+1,$$

which leads to $p + t \ge N_1$. Hence, $p + t \ge N_1$.

Let $\mathbb{C}_{j}^{\triangleleft}(N_{1}+1,\ldots,N_{k-1};i|p,t)$ denote the set of partitions μ in $\mathbb{C}_{j}(N_{1}+1,\ldots,N_{k-1};i)$ satisfying the following conditions:

- (1) $\mu_{p+1}^{(1)} = 2t + 1;$
- (2) if $p \ge 1$, then $\mu \in \mathbb{C}_j(N_1 + 1, \dots, N_{k-1}; i|p);$
- (3) if there exist parts 2t + 2 in μ , then $p \ge 1$ and $\mu_p^{(1)} = 2t + 3$.

We proceed to show that there is a bijection between $\mathbb{C}_{j}^{\leq}(N_{1},\ldots,N_{k-1};i|p,t)$ and $\mathbb{C}_{j}^{\leq}(N_{1}+1,\ldots,N_{k-1};i|p,t)$, from which we find that if $\mathbb{C}_{j}^{\leq}(N_{1}+1,\ldots,N_{k-1};i|p,t)$ is non-empty, then we have $p+t \geq N_{1}$.

Lemma 4.6. For $1 \leq p \leq N_1$, there is a bijection, the combination $C_{p,t}$, between $\mathbb{C}_j^<(N_1,\ldots,N_{k-1};i|p,t)$ and $\mathbb{C}_j^\lhd(N_1+1,\ldots,N_{k-1};i|p,t)$. Moreover, for $\lambda \in \mathbb{C}_j^\lt(N_1,\ldots,N_{k-1};i|p,t)$ and $\mu = C_{p,t}(\lambda) \in \mathbb{C}_j^\lhd(N_1+1,\ldots,N_{k-1};i|p,t)$, we have $|\mu| = |\lambda| + 2t + 2$.

Proof. For $k \geq i \geq 2$, let λ be a partition in $\mathbb{C}_1^<(N_1, \ldots, N_{k-1}; i|p, t)$. Note that $p \geq 1$, then we have $\lambda \in \mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$, which implies that $\lambda_p^{(1)}$ is even type, and so $\lambda_p^{(1)}$ is an even part. We may write $\lambda_p^{(1)} = 2b + 2$.

If b = t, namely $\lambda_p^{(1)} = 2t + 2$, then set r = 1. If $b \ge t + 1$, namely $\lambda_p^{(1)} \ge 2t + 4$, then set r be the largest mark of parts 2b + 2 in $GG(\lambda)$. Define the combination $\mu = C_{p,t}(\lambda)$ by adding 2t + 1 as a new part to λ and replacing the r-marked 2b + 2 in $GG(\lambda)$ by 2b + 3to obtain μ .

Obviously, $|\mu| = |\lambda| + 2t + 2$. To show that $\mu \in \mathbb{C}_1^{\triangleleft}(N_1 + 1, \dots, N_{k-1}; i|p, t)$, we first prove that $\mu \in \mathbb{C}_1(N_1 + 1, \dots, N_{k-1}; i)$. It is enough to verify that μ satisfies the following three conditions:

- (A) There is only one 2t + 1 in μ and 2t + 1 is marked with 1 in $GG(\mu)$;
- (B) The new generated part 2b + 3 is also marked with r in $GG(\mu)$.
- (C) The marks of the unchanged parts in $GG(\mu)$ are the same as those in $GG(\lambda)$.

Condition (A). Under the condition that $\lambda_{p+1}^{(1)} + 2 \leq 2t + 1 < \lambda_p^{(1)}$, we deduce that $\lambda_{p+1}^{(1)} \leq 2t - 1$ and $\lambda_p^{(1)} \geq 2t + 2$. So, there is no part 2t + 1 in λ . Otherwise, we must

have $\lambda_p^{(1)} = 2t + 1$, which contradicts the fact that $\lambda_p^{(1)} \ge 2t + 2$. Hence, there is only one 2t + 1 in μ , which is marked with 1 in $GG(\mu)$.

Condition (B). It suffices to show that there are 1-marked, ..., (r-1)-marked 2b+2in $GG(\lambda)$ when $r \geq 2$. If $r \geq 2$, then $b \geq t+1$, namely, $\lambda_p^{(1)} \geq 2t+4$. Note that $\lambda_{p+1}^{(1)} \leq 2t-1$, so 2t+1, ..., 2b and 2b+1 do not occur in λ . Under the condition that there is a *r*-marked 2b+2 in $GG(\lambda)$, by the definition of Göllnitz-Gordon marking, we see that there are 1-marked, ..., (r-1)-marked 2b+2 in $GG(\lambda)$.

Condition (C). From the proofs of Conditions (A) and (B), we find that the marks of the unchanged parts less than 2b + 3 in $GG(\mu)$ are the same as those in $GG(\lambda)$. Moreover, by the definition of Göllnitz-Gordon marking, we derive that the marks of the unchanged parts greater than 2b + 3 in $GG(\mu)$ are the same as those in $GG(\lambda)$. Hence, we conclude that μ is a partition in $\mathbb{C}_1(N_1 + 1, \ldots, N_{k-1}; i)$.

Now, we turn to show that $\mu \in \mathbb{C}_1^{\triangleleft}(N_1+1,\ldots,N_{k-1};i|p,t)$. From the proof above, we have $\mu_s^{(1)} = \lambda_s^{(1)}$ for $1 \leq s \leq p-1$, $\mu_{p+1}^{(1)} = 2t+1$, $\mu_s^{(1)} = \lambda_{s-1}^{(1)}$ for $p+2 \leq s \leq N_1+1$. If r = 1, then we have $\mu_p^{(1)} = \lambda_p^{(1)} + 1 = 2b+3$. If $r \geq 2$, then we have $\mu_p^{(1)} = \lambda_p^{(1)} = 2b+2$ and there is a *r*-marked $\lambda_p^{(1)} + 1 = 2b+3$ in $GG(\mu)$. So, $\mu_p^{(1)}$ is odd type. Moreover, by the construction of μ , we find that $\mu_s^{(1)}$ is even type for $1 \leq s \leq p-1$, and so we deduce that $\mu \in \mathbb{C}_1(N_1+1,\ldots,N_{k-1};i|p)$.

Assume that there exist parts 2t+2 in μ , then the marks of parts 2t+2 are greater than 1 in $GG(\mu)$ since $\mu_{p+1}^{(1)} = 2t + 1$. From the proof above, we find that the marks of parts 2t+2 in $GG(\mu)$ are the same as those in $GG(\lambda)$. By the definition of Göllnitz-Gordon marking, we derive that there is a 1-marked 2t or 2t+1 or 2t+2 in $GG(\lambda)$. Note that $\lambda_{p+1}^{(1)} \leq 2t-1$ and $\lambda_p^{(1)} \geq 2t+2$, so we get $\lambda_p^{(1)} = 2t+2$. Therefore, we deduce that r=1and $\mu_p^{(1)} = 2t+3$. Hence, we conclude that μ is a partition in $\mathbb{C}_1^{\triangleleft}(N_1+1,\ldots,N_{k-1};i|p,t)$.

We proceed to prove that for $k \geq 3$ and $k \geq i \geq 2$, if $\lambda \in \mathbb{C}_0^<(N_1, \ldots, N_{k-1}; i|p, t)$, then $\mu \in \mathbb{C}_0^{\triangleleft}(N_1+1, \ldots, N_{k-1}; i|p, t)$. Let $\lambda^{(k-1)} = (\lambda_1^{(k-1)}, \lambda_2^{(k-1)}, \ldots, \lambda_{N_{k-1}}^{(k-1)})$ and $\mu^{(k-1)} = (\mu_1^{(k-1)}, \mu_2^{(k-1)}, \ldots, \mu_{N_{k-1}}^{(k-1)})$ be the (k-1)-marked parts in $GG(\lambda)$ and $GG(\mu)$, respectively. Assume that λ is a partition in $\mathbb{C}_0^<(N_1, \ldots, N_{k-1}; i|p, t)$, then $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$.

For each $s \in [1, N_{k-1}]$, we have $\lambda_{p+1}^{(1)} \leq 2t - 1$ and $\lambda_p^{(1)} = 2b + 2 \geq 2t + 2$, and so $\lambda_s^{(k-1)} \notin [2t+1, 2b+1]$. By the construction of μ , we derive that $\mu_s^{(k-1)} \notin [2t+1, 2b+1]$. To show that $\{\mu_s^{(k-1)}\}_{k-1}$ is good, we consider the following four cases.

Case 1: $\mu_s^{(k-1)} \leq 2t$ or $\mu_s^{(k-1)} \geq 2b+5$. By the construction of μ , we have $\{\mu_s^{(k-1)}\}_{k-1} = \{\lambda_s^{(k-1)}\}_{k-1}$. Moreover, if $\mu_s^{(k-1)} \leq 2t$, then $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)})$; if $\mu_s^{(k-1)} \geq 2b+5$, then $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)}) + 2$. This implies that $O_{\mu}(\mu_s^{(k-1)}) \equiv O_{\lambda}(\lambda_s^{(k-1)}) \pmod{2}$. Note that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good, so $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 2: $\mu_s^{(k-1)} = 2b + 2$. By the construction of μ , we find that b = t and r = 1.

Moreover, we have $\mu_s^{(k-1)} = \lambda_s^{(k-1)} = 2b + 2 = 2t + 2$, $f_{2t}(\mu) = f_{2t}(\lambda)$, $f_{2t+1}(\mu) = f_{2t+1}(\lambda) + 1$, $f_{2t+2}(\mu) = f_{2t+2}(\lambda) - 1$ and $O_{\mu}(2t+2) = O_{\lambda}(2t+2) + 1$. So we get

$$\begin{split} [\mu_{s,1}^{(k-1)}/2] + [\mu_{s,2}^{(k-1)}/2] + \cdots + [\mu_{s,k-1}^{(k-1)}/2] \\ &= tf_{2t}(\mu) + tf_{2t+1}(\mu) + (t+1)f_{2t+2}(\mu) \\ &= tf_{2t}(\lambda) + t(f_{2t+1}(\lambda) + 1) + (t+1)(f_{2t+2}(\lambda) - 1) \\ &= tf_{2t}(\lambda) + tf_{2t+1}(\lambda) + (t+1)f_{2t+2}(\lambda) - 1 \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] - 1 \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) - 1 = i - 1 + O_{\lambda}(2t+2) - 1 \\ &\equiv i - 1 + O_{\mu}(2t+2) = i - 1 + O_{\mu}(\mu_{s,1}^{(k-1)}) \pmod{2} \end{split}$$

which implies that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 3: $\mu_s^{(k-1)} = 2b + 3$. By the construction of μ , we find that $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)}) + 2 \equiv O_{\lambda}(\lambda_s^{(k-1)}) \pmod{2}$. With the similar argument of Case 4 in the proof of Lemma 4.2, we derive that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Case 4: $\mu_s^{(k-1)} = 2b + 4$. By the construction of μ , we observe that $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)}) + 2 \equiv O_{\lambda}(\lambda_s^{(k-1)}) \pmod{2}$. With the similar argument of Case 5 in the proof of Lemma 4.2, we obtain that $\{\mu_s^{(k-1)}\}_{k-1}$ is good.

Therefore, we conclude that $\{\mu_s^{(k-1)}\}_{k-1}$ is good. Hence, by Theorem 3.2, we see that μ is a partition in $\mathbb{C}_0^{\triangleleft}(N_1+1,\ldots,N_{k-1};i|p,t)$.

To prove that the combination $C_{p,t}$ is a bijection, we construct the inverse map of $C_{p,t}$, the division $D_{p,t}$. Assume that μ is a partition in $\mathbb{C}_{j}^{\triangleleft}(N_{1} + 1, \ldots, N_{k-1}; i|p, t)$, then we see that $\mu_{p+1}^{(1)} = 2t + 1$ and $\mu_{p}^{(1)}$ is odd type. If $\mu_{p}^{(1)}$ is an odd part, then we may write $\mu_{p}^{(1)} = 2b + 3$ and set r = 1, where $b \geq t$. If $\mu_{p}^{(1)}$ is an even part, then we may write $\mu_{p}^{(1)} = 2b + 2$ and set r be the mark of 2b + 3 in $GG(\mu)$, where $b \geq t + 1$ and $r \geq 2$. Define $\lambda = D_{p,t}(\mu)$ by removing the part $\mu_{p+1}^{(1)} = 2t + 1$ from μ and replacing the r-marked 2b + 3 in $GG(\mu)$ by 2b + 2 to obtain λ . It can be verified that the division $D_{p,t}$ is the inverse map of the combination $C_{p,t}$. This completes the proof.

Lemma 4.7. There is a bijection, the combination $C_{0,t}$, between $\mathbb{C}_j^{\leq}(N_1, \ldots, N_{k-1}; i|0, t)$ and $\mathbb{C}_j^{\triangleleft}(N_1 + 1, \ldots, N_{k-1}; i|0, t)$. Moreover, for $\lambda \in \mathbb{C}_j^{\leq}(N_1, \ldots, N_{k-1}; i|0, t)$ and $\mu = C_{0,t}(\lambda) \in \mathbb{C}_j^{\triangleleft}(N_1 + 1, \ldots, N_{k-1}; i|0, t)$, we have $|\mu| = |\lambda| + 2t + 1$.

Proof. For $k \geq i \geq 2$, let λ be a partition in $\mathbb{C}_1^<(N_1, \ldots, N_{k-1}; i|0, t)$. Define $\mu = C_{0,t}(\lambda)$ by adding 2t + 1 as a new part of λ to obtain μ . Obviously, we have $|\mu| = |\lambda| + 2t + 1$. Note that $\lambda_1^{(1)} = \lambda_{p+1}^{(1)} \leq 2t - 1$, we find that the parts of λ are not exceed to 2t. So, 2t + 1 is marked with 1 in $GG(\mu)$ and the marks of the remaining parts in $GG(\mu)$ are the same as those in $GG(\lambda)$. Therefore, μ is a partition in $\mathbb{C}_1(N_1 + 1, \ldots, N_{k-1}; i)$ and $\mu_1^{(1)} = 2t + 1$, which implies that $\mu \in \mathbb{C}_1^{\triangleleft}(N_1 + 1, \ldots, N_{k-1}; i|0, t)$. We turn to show that for $k \geq 3$ and $k \geq i \geq 2$, if $\lambda \in \mathbb{C}_0^<(N_1, \ldots, N_{k-1}; i|0, t)$, then $\mu \in \mathbb{C}_0^<(N_1 + 1, \ldots, N_{k-1}; i|0, t)$. Let $\lambda^{(k-1)} = (\lambda_1^{(k-1)}, \lambda_2^{(k-1)}, \ldots, \lambda_{N_{k-1}}^{(k-1)})$ and $\mu^{(k-1)} = (\mu_1^{(k-1)}, \mu_2^{(k-1)}, \ldots, \mu_{N_{k-1}}^{(k-1)})$ be the (k-1)-marked parts in $GG(\lambda)$ and $GG(\mu)$, respectively. Assume that λ is a partition in $\mathbb{C}_0^<(N_1, \ldots, N_{k-1}; i|0, t)$, then $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$.

For each $s \in [1, N_{k-1}]$, by the construction of μ , we have $\mu_s^{(k-1)} = \lambda_s^{(k-1)}$, $\{\mu_s^{(k-1)}\}_{k-1} = \{\lambda_s^{(k-1)}\}_{k-1}$ and $O_{\mu}(\mu_s^{(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)})$. Note that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good, so we get that $\{\mu_s^{(k-1)}\}_{k-1}$ is good. Hence, by Theorem 3.2, we see that μ is a partition in $\mathbb{C}_0^{\triangleleft}(N_1 + 1, \ldots, N_{k-1}; i | 0, t)$.

To prove that the combination $C_{0,t}$ is a bijection, we construct the inverse map of $C_{0,t}$, the division $D_{0,t}$. Assume that μ is a partition in $\mathbb{C}_{j}^{\triangleleft}(N_{1}+1,\ldots,N_{k-1};i|0,t)$, then we see that $\mu_{1}^{(1)} = 2t + 1$. Define $\lambda = D_{0,t}(\mu)$ by removing the part $\mu_{1}^{(1)} = 2t + 1$ from μ to obtain λ . It can be verified that the division $D_{0,t}$ is the inverse map of the combination $C_{0,t}$. This completes the proof.

4.3 The insertion I_m and the separation S_m

In this subsection, we assume that j = 0 or 1, (2k + j)/2 > 2, $k \ge i \ge 2$ and $N_1 \ge \cdots \ge N_{k-1} \ge 0$. Let μ be a partition in $\mathbb{C}_j(N_1 + 1, \ldots, N_{k-1}; i)$ such that there exist odd parts in μ . We say that 2t + 1 is a insertion part of μ if 2t + 1 is the largest odd part 2t + 1 of μ , $\mu_{p+1}^{(1)} = 2t + 1$, where $0 \le p \le N_1$, and μ satisfies either of the following two conditions:

- (1) There do not exist parts 2t + 2 in μ .
- (2) There exist parts 2t + 2 in μ , and there exists $s \in [1, p]$ such that $\mu_{p+1-s}^{(1)} = 2t + 4s$ and there do not exist parts 2t + 4s + 2 in μ .

For $t \ge 0$ and $0 \le p \le N_1$, let $\mathbb{C}_j^=(N_1+1,\ldots,N_{k-1};i|p,t)$ denote the set of partitions μ in $\mathbb{C}_j(N_1+1,\ldots,N_{k-1};i)$ such that $\mu_{p+1}^{(1)}=2t+1$ is a insertion part of μ .

Theorem 4.8. For $t \ge 0$ and $0 \le p \le N_1$, there exists a bijection $I_{p,t}$ between $\mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|p, t)$ and $\mathbb{C}_j^=(N_1 + 1, \ldots, N_{k-1}; i|p, t)$. Moreover, for $\lambda \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|p, t)$ and $\mu = I_{p,t}(\lambda) \in \mathbb{C}_j^=(N_1 + 1, \ldots, N_{k-1}; i|p, t)$, we have $|\mu| = |\lambda| + 2p + 2t + 1$.

Proof. For convenience, we define $\Theta_{(0)}$ as the identity map. Then we define $\mu = I_{p,t}(\lambda) = \Theta_{(p)}(C_{p,t}(\lambda))$. By Lemmas 4.5, 4.6 and 4.7, we find that $I_{p,t}$ is well-defined, $\mu \in \mathbb{C}_j(N_1 + 1, \ldots, N_{k-1}; i)$, $\mu_{p+1}^{(1)} = 2t + 1$ is the largest odd part of μ and $|\mu| = |\lambda| + 2p + 2t + 1$. We proceed to show that $\mu \in \mathbb{C}_j^= (N_1 + 1, \ldots, N_{k-1}; i|p, t)$.

If p = 0, then by Lemma 4.7, we have $\mu \in \mathbb{C}_1^=(N_1 + 1, ..., N_{k-1}; i|0, t)$, which implies that $\mu \in \mathbb{C}_j^=(N_1 + 1, ..., N_{k-1}; i|0, t)$.

If $p \ge 1$, then it suffices to show that μ satisfies either condition (1) or condition (2) in the definition of an insertion part. If there do not exist parts 2t + 2 in μ , then it is clear that μ satisfies the condition (1) in the definition of an insertion part.

Assume that there exist parts 2t + 2 in μ , recall that $\mu_{p+1}^{(1)} = 2t + 1$, then we see that the marks of parts 2t + 2 are greater than 1 in $GG(\mu)$. By the construction of μ , we find that $\lambda_p^{(1)} = 2t + 2$ and there exist parts 2t + 2 with marks greater than 1 in $GG(\lambda)$. Set sbe the largest integer such that $\lambda_{p+1-s}^{(1)} = 2t + 4s - 2$ and there exist parts 2t + 4s - 2 with marks greater than 1 in $GG(\lambda)$. Using $\lambda_p^{(1)} = 2t + 2$, we deduce that $1 \le s \le p$. It follows from the construction of μ that $\lambda_{p+1-s}^{(1)} = 2t + 4s - 2$ is replaced by 2t + 4s in μ , and so $\mu_{p+1-s}^{(1)} = 2t + 4s$. If there do not exist parts 2t + 4s + 2 in λ , then by the construction of μ , we see that there also do not exist parts 2t + 4s + 2 in μ . If there exist parts 2t + 4s + 2in λ , then by the choice of s, we find that $\lambda_{p-s}^{(1)} = 2t + 4s + 2$ and there do not parts exist 2t + 4s + 2 with marks greater than 1 in $GG(\lambda)$. Again by the construction of μ , we see that $\lambda_{p-s}^{(1)} = 2t + 4s + 2$ is replaced by 2t + 4s + 4 in μ , which implies that there do not exist parts 2t + 4s + 2 in μ . So, μ satisfies the condition (2) in the definition of insertion part. Hence, μ is a partition in $\mathbb{C}_j^= (N_1 + 1, \dots, N_{k-1}; i | p, t)$.

To prove that $I_{p,t}$ is a bijection, we construct the inverse map of the insertion $I_{p,t}$, the separation $S_{p,t}$. Define $\Lambda_{(0)}$ as the identity map. Then we define $\lambda = S_{p,t}(\mu) = D_{p,t}(\Lambda_{(p)}(\mu))$. By Lemmas 4.5, 4.6 and 4.7, it can be verified that the separation $S_{p,t}$ is the inverse map of the insertion $I_{p,t}$. This completes the proof.

For $m \geq N_1$, define

$$\mathbb{C}_j^<(N_1,\ldots,N_{k-1};i|m) = \bigcup_{p+t=m} \mathbb{C}_j^<(N_1,\ldots,N_{k-1};i|p,t),$$

and

$$\mathbb{C}_{j}^{=}(N_{1}+1,\ldots,N_{k-1};i|m) = \bigcup_{p+t=m} \mathbb{C}_{j}^{=}(N_{1}+1,\ldots,N_{k-1};i|p,t)$$

We establish a bijection between $\mathbb{C}_{j}^{<}(N_{1},\ldots,N_{k-1};i|m)$ and $\mathbb{C}_{j}^{=}(N_{1}+1,\ldots,N_{k-1};i|m)$.

Theorem 4.9. For $m \ge N_1$, there is a bijection, the insertion I_m , between $\mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|m)$ and $\mathbb{C}_j^=(N_1 + 1, \ldots, N_{k-1}; i|m)$. Moreover, for $\lambda \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|m)$ and $\mu = I_m(\lambda) \in \mathbb{C}_j^=(N_1 + 1, \ldots, N_{k-1}; i|m)$, we have $|\mu| = |\lambda| + 2m + 1$.

Proof. Let λ be a partition in $\mathbb{C}_{j}^{<}(N_{1},\ldots,N_{k-1};i|m)$ and let p be the smallest integer such that $2(m-p)+1 \geq \lambda_{p+1}^{(1)}+2$, where $0 \leq p \leq N_{1}$. Set t=m-p, then we have $\lambda \in \mathbb{C}_{j}^{<}(N_{1},\ldots,N_{k-1};i|p,t)$. Define $\mu = I_{m}(\lambda) = I_{p,t}(\lambda)$. By Theorem 4.8, we see that $\mu \in \mathbb{C}_{j}^{=}(N_{1}+1,\ldots,N_{k-1};i|p,t) \subset \mathbb{C}_{j}^{=}(N_{1}+1,\ldots,N_{k-1};i|m)$ and $|\mu| = |\lambda|+2t+2p+1 = |\lambda|+2m+1$.

To prove that I_m is a bijection, we construct the inverse map of the insertion I_m , the separation S_m . Let μ be a partition in $\mathbb{C}_i^=(N_1+1,\ldots,N_{k-1};i|m)$. Assume that $\mu_{p+1}^{(1)} = 2t + 1$ is the insertion part of μ , where m = p + t. Define $\lambda = S_m(\mu) = S_{p,t}(\mu)$. It is clear that the separation S_m is the inverse map of I_m . This completes the proof.

The following theorem give a criterion to determine whether a partition in $\mathbb{C}_{j}^{=}(N_{1} + 1, \ldots, N_{k-1}; i|m)$ also belongs to $\mathbb{C}_{j}^{<}(N_{1} + 1, \ldots, N_{k-1}; i|m')$, which involves the successive applications of the insertion.

Theorem 4.10. For $m \ge N_1$, let μ be a partition in $\mathbb{C}_j^=(N_1+1,\ldots,N_{k-1};i|m)$. Then μ is a partition in $\mathbb{C}_i^<(N_1+1,\ldots,N_{k-1};i|m')$ if and only if m' > m.

Proof. Assume that $\mu_{p+1}^{(1)} = 2t + 1$ is the insertion part of μ , where m = p + t.

If m' > m, then we have $2m' + 1 - 2p \ge 2(m+1) + 1 - 2p = (2t+1) + 2 = \mu_{p+1}^{(1)} + 2$. Assume that p' is the smallest integer such that $2m' + 1 - 2p' \ge \mu_{p'+1}^{(1)} + 2$, we see that $p' \le p$. Set t' = m' - p', then we have $\mu \in \mathbb{C}_{j}^{<}(N_{1} + 1, \dots, N_{k-1}; i|p', t')$, which implies that $\mu \in \mathbb{C}_{j}^{<}(N_{1} + 1, \dots, N_{k-1}; i|m')$.

On the other hand, assume that μ is a partition in $\mathbb{C}_{j}^{<}(N_{1}+1,\ldots,N_{k-1};i|m')$. Let p' be the smallest integer such that $2(m'-p')+1 \geq \mu_{p'+1}^{(1)}+2$, where $0 \leq p' \leq N_{1}+1$. Set t'=m'-p', then we have $\mu \in \mathbb{C}_{j}^{<}(N_{1}+1,\ldots,N_{k-1};i|p',t')$, and so t'>t and $p' \leq p$. Hence

$$2t' + 1 \ge \mu_{p'+1}^{(1)} + 2 \ge \dots \ge \mu_{p+1}^{(1)} + 2(p - p' + 1) = 2t + 1 + 2(p - p' + 1),$$

which leads to $p' + t' \ge p + t + 1$. This implies m' = p' + t' > p + t = m.

Thus, we complete the proof.

4.4 Proof of Theorem 3.4

We are now in a position to give a proof of Theorem 3.4.

Proof of Theorem 3.4. Let $(\lambda, -, \eta)$ be a pair in $\mathbb{F}_1(2, 2)$. Then, we have $\lambda \in \mathbb{E}_1(2, 2)$. We define $\pi = \Phi_{2,2}(\lambda, -, \eta)$ as follows. There are two cases.

Case 1: $\eta = \emptyset$. Then we set $\pi = \lambda$. It is clear that $\pi \in \mathbb{E}_1(2,2) \subset C_1(2,2)$. Moreover, $|\pi| = |\lambda| + |\eta|$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

Case 2: $\eta \neq \emptyset$. Note that λ is a partition in $\mathbb{E}_1(2,2)$, so the parts of λ are even and the marks of parts in $GG(\lambda)$ are not exceed to 1. Assume that there are N_1 1-marked parts in $GG(\lambda)$, denoted $\lambda = \lambda^{(1)} = (\lambda_1^{(1)}, \lambda_2^{(1)}, \ldots, \lambda_{N_1}^{(1)}) \in \mathbb{E}_1(N_1; 2)$. Then we have $\eta \in \mathbb{I}_{N_1}$, denoted $\eta = (2m_1 + 1, 2m_2 + 1, \ldots, 2m_\ell + 1)$, where $m_1 > m_2 > \cdots > m_\ell \ge N_1$.

Starting with λ , apply the insertion repeatedly to get π . Denote the intermediate partitions by $\lambda^0, \lambda^1, \ldots, \lambda^\ell$ with $\lambda^0 = \lambda$ and $\lambda^\ell = \pi$. Since $m_\ell \ge N_1$, there exists p such that p is the smallest integer p satisfying $2m_\ell + 1 - 2p \ge \lambda_{p+1}^{(1)} + 2$. Note that there do not

exist odd parts in λ , so $\lambda_s^{(1)}$ is even type for $1 \leq s \leq N_1$. Set $t = m_\ell - p$, then we have $\lambda \in \mathbb{C}_1^<(N_1; 2|p, t) \subset \mathbb{C}_1^<(N_1; 2|m_\ell)$.

Set b = 0 and repeat following process until $b = \ell$.

(A) Note that $\lambda^b \in \mathbb{C}_1^{<}(N_1 + b; 2|m_{\ell-b})$, we apply the insertion $I_{m_{\ell-b}}$ to λ^b to get λ^{b+1} , that is,

$$\lambda^{b+1} = I_{m_{\ell-b}}(\lambda^b).$$

In view of Theorem 4.9, we deduce that

$$\lambda^{b+1} \in \mathbb{C}_1^{=}(N_1 + b + 1; 2|m_{\ell-b})$$

and

$$|\lambda^{b+1}| = |\lambda^b| + 2m_{\ell-b} + 1.$$

(B) Replace b by b + 1. If $b = \ell$, then we are done. If $b < \ell$, then by Theorem 4.10, we have

 $\lambda^b \in \mathbb{C}_1^<(N_1+b;2|m_{\ell-b}),$

since $m_{\ell-b} > m_{\ell-b+1}$. Go back to (A).

Eventually, the above process yields $\pi = \lambda^{\ell} \in \mathbb{C}_1^=(N_1 + \ell; 2|m_1)$ such that $|\pi| = |\lambda| + |\eta|$. Moreover, we have $\pi \in \mathbb{C}_1(2; 2)$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

To show that $\Phi_{2,2}$ is a bijection, we give the inverse map $\Psi_{2,2}$ of $\Phi_{2,2}$ by successively applying the separation defined in the proof of Theorem 4.9. Let π be a partition in $\mathbb{C}_1(2;2)$. We shall construct a pair $(\lambda, -, \eta)$, that is, $(\lambda, -, \eta) = \Psi_{2,2}(\pi)$, such that $(\lambda, -, \eta) \in \mathbb{F}_1(2;2), |\pi| = |\lambda| + |\eta|$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

Assume that there are $\ell \geq 0$ odd parts and N_1 even parts in π . We eliminate all odd parts of π . There are two cases.

Case 1: $\ell = 0$. Then set $\lambda = \pi$ and $\eta = \emptyset$. Clearly, $(\lambda, -, \eta) \in \mathbb{F}_1(2; 2)$, $|\pi| = |\lambda| + |\eta|$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

Case 2: $\ell \geq 1$. Assume that $2t_0 + 1 > 2t_1 + 1 > \cdots > 2t_{\ell-1} + 1$ are the odd parts of π . We eliminate the ℓ odd parts of π by successively applying the separation defined in the proof of Theorem 4.9. Denote the intermediate pairs by $(\pi^0, -, \eta^0), (\pi^1, -, \eta^1), \ldots, (\pi^\ell, -, \eta^\ell)$ with $(\pi^0, -, \eta^0) = (\pi, -, \emptyset)$.

Set b = 0 and carry out the following procedure.

(A) Note that π^b is a partition in $\mathbb{C}_1(N_1 + \ell - b; 2)$ and $2t_b + 1$ is the largest odd part of π^b , we see that there are no parts $2t_b + 2$ in π^b , otherwise the largest mark in $GG(\pi^b)$ are greater than 1, which leads to a contradiction. So, $2t_b + 1$ is a insertion part of π^b . Assume that $2t_b + 1$ is the $(p_b + 1)$ -th part of π^b . Let $m_b = p_b + t_b$. Then,

$$\pi^b \in \mathbb{C}_1^=(N_1 + \ell - b; 2|m_b)$$

Apply the separation S_{m_b} to π^b to get π^{b+1} , that is,

$$\pi^{b+1} = S_{m_b}(\pi^b).$$

Employing Theorem 4.9, we find that

$$\pi^{b+1} \in \mathbb{C}_1^<(N_1 + \ell - b - 1; 2|m_b)$$

and

$$|\pi^{b+1}| = |\pi^b| - 2m_b - 1.$$

Then insert $2m_b + 1$ into η^b to obtain η^{b+1} .

(B) Replace b by b + 1. If $b = \ell$, then we are done. Otherwise, go back to (A).

Observe that for $0 \le b \le \ell$, there are $\ell - b$ odd parts in π^b . Theorem 4.10 reveals that for $0 \le b < \ell - 1$,

$$m_b > m_{b+1} \ge N_1 + \ell - b - 2.$$

Therefore, there are no odd parts in π^{ℓ} and $\eta^{\ell} = (2m_0 + 1, 2m_1 + 1, \dots, 2m_{\ell-1} + 1)$ is a partition in \mathbb{I}_{N_1} . Set $(\lambda, -, \eta) = (\pi^{\ell}, -, \eta^{\ell})$. It is clear that $\lambda = \pi^{\ell} \in \mathbb{C}_1^< (N_1; 2|m_{\ell-1}) \subset \mathbb{C}_1(N_1; 2), \ \eta = \eta^{\ell} \in \mathbb{I}_{N_1}, \ |\pi| = |\lambda| + |\eta| \text{ and } \ell(\pi) = \ell(\lambda) + \ell(\eta)$. Recall that there do not exist odd parts in λ , so $\lambda \in \mathbb{E}_1(2; 2)$. Hence, $(\lambda, -, \eta)$ is a pair in $\mathbb{F}_1(2, 2)$.

It can be verified that $\Psi_{2,2}$ is the inverse map of $\Phi_{2,2}$. Thus, we complete the proof.

5 The reduction R_p and the dilation H_p

Assume that $k \geq 3$, $k \geq i \geq 1$, j = 0 or 1, $N_1 \geq \cdots \geq N_{k-1} \geq 0$ and $1 \leq p \leq N_2$. In this section, we introduce two subsets of $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$, $\hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, show the follow theorem which is proved by the reduction operation and the dilation operation.

Theorem 5.1. There is a bijection, the reduction R_p , between $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i|p)$ and the set $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$. Moreover, for $\lambda \in \hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\mu = R_p(\lambda) \in \check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, we have $|\mu| = |\lambda| - 2p + 1$.

5.1 The set $\hat{\mathbb{C}}_{j}(N_{1}, \ldots, N_{k-1}; i|p)$

Let λ be a partition in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$ and let $\lambda^{(r)} = (\lambda_1^{(r)}, \lambda_2^{(r)}, \ldots, \lambda_{N_r}^{(r)})$ denote *r*-marked parts in $GG(\lambda)$, where $\lambda_1^{(r)} > \lambda_2^{(r)} > \cdots > \lambda_{N_r}^{(r)}$ and $1 \le r \le k-1$. We will focus on the 2-marked parts in $GG(\lambda)$.

The types of $\lambda_1^{(2)}, \lambda_2^{(2)}, \ldots, \lambda_p^{(2)}$. Assume that $\lambda_p^{(2)}$ is even and there do not exist odd parts greater than $\lambda_p^{(2)}$ in λ , then we define the types of $\lambda_1^{(2)}, \lambda_2^{(2)}, \ldots, \lambda_p^{(2)}$ from largest to smallest as follows. We first define the type of $\lambda_1^{(2)}$. There are two cases.

- (1) If there exist parts $\lambda_1^{(2)} + 2$ with marks greater than 2 in $GG(\lambda)$, then by the definition of Göllnitz-Gordon marking, there is a 1-marked part $\lambda_{s_1}^{(1)} = \lambda_1^{(2)}$ or $\lambda_1^{(2)} + 2$ in $GG(\lambda)$. We set $\hat{R}_1(\lambda) = s_1$. There are two subcases.
 - (1.1) If $\lambda_{s_1}^{(1)} = \lambda_1^{(2)}$, then we say that $\lambda_1^{(2)}$ is of type \hat{A}_1 . (1.2) If $\lambda_{s_1}^{(1)} = \lambda_1^{(2)} + 2$, then we say that $\lambda_1^{(2)}$ is of type \hat{B} .
- (2) If there do not exist parts $\lambda_1^{(2)} + 2$ with marks greater than 2 in $GG(\lambda)$, then by the definition of Göllnitz-Gordon marking, there is a 1-marked part $\lambda_{s_1}^{(1)} = \lambda_1^{(2)} 2$ or $\lambda_1^{(2)} 1$ or $\lambda_1^{(2)}$ in $GG(\lambda)$. We set $\hat{R}_1(\lambda) = s_1$. There are three subcases.
 - (2.1) If $\lambda_{s_1}^{(1)} = \lambda_1^{(2)} 2$, then we say that $\lambda_1^{(2)}$ is of type \hat{C} .
 - (2.2) If $\lambda_{s_1}^{(1)} = \lambda_1^{(2)} 1$, then we say that $\lambda_1^{(2)}$ is of type \hat{O} .
 - (2.3) If $\lambda_{s_1}^{(1)} = \lambda_1^{(2)}$, then we say that $\lambda_1^{(2)}$ is of type \hat{A}_2 .

For $1 < b \leq p$, assume that $\hat{R}_{b-1}(\lambda)$ and the type of $\lambda_{b-1}^{(2)}$ have been defined, then we define the type of $\lambda_{b}^{(2)}$ as follows. There are two cases.

- (1) If there exist parts $\lambda_b^{(2)} + 2$ with marks greater than 2 in $GG(\lambda)$, then by the definition of Göllnitz-Gordon marking, there is a 1-marked part $\lambda_{s_b}^{(1)} = \lambda_b^{(2)}$ or $\lambda_b^{(2)} + 2$ in $GG(\lambda)$. There are four subcases.
 - (1.1) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)}$, then we say that $\lambda_b^{(2)}$ is of type \hat{A}_1 and set $\hat{R}_b(\lambda) = s_b$.
 - (1.2) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} + 2$ and $\hat{R}_{b-1}(\lambda) \neq s_b$, then we say that $\lambda_b^{(2)}$ is of type \hat{B} and set $\hat{R}_b(\lambda) = s_b$.
 - (1.3) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} + 2$, $\hat{R}_{b-1}(\lambda) = s_b$ and $\lambda_{s_b+1}^{(1)} = \lambda_b^{(2)} 2$, then we say that $\lambda_b^{(2)}$ is of type \hat{C} and set $\hat{R}_b(\lambda) = s_b + 1$.
 - (1.4) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} + 2$, $\hat{R}_{b-1}(\lambda) = s_b$ and $\lambda_{s_b+1}^{(1)} = \lambda_b^{(2)} 1$, then we say that $\lambda_b^{(2)}$ is of type \hat{O} and set $\hat{R}_b(\lambda) = s_b + 1$.
- (2) If there do not exist parts $\lambda_b^{(2)} + 2$ with marks greater than 2 in $GG(\lambda)$, then by the definition of Göllnitz-Gordon marking, there is a 1-marked part $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} 2$ or $\lambda_b^{(2)} 1$ or $\lambda_b^{(2)}$ in $GG(\lambda)$. We set $\hat{R}_l(\lambda) = s_b$. There are three subcases.
 - (2.1) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} 2$, then we say that $\lambda_b^{(2)}$ is of type \hat{C} .
 - (2.2) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)} 1$, then we say that $\lambda_b^{(2)}$ is of type \hat{O} .
 - (2.3) If $\lambda_{s_b}^{(1)} = \lambda_b^{(2)}$, then we say that $\lambda_b^{(2)}$ is of type \hat{A}_2 .

For each $1 \leq b \leq p$, we say that $\lambda_b^{(2)}$ is of type \hat{A} if $\lambda_b^{(2)}$ it is of type \hat{A}_1 or type \hat{A}_2 .

For example, let λ be a partition in $\mathbb{C}_1(10, 9, 6; 3)$, whose Göllnitz-Gordon marking is

$$GG(\lambda) = \begin{bmatrix} 4 & 12 & 16 & 22 & 30 & 36 \\ 2 & 6 & 10 & 14 & 20 & 24 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 20 & 24 & 28 & 34 & 38 \end{bmatrix}.$$
 (5.1)

For p = 9, it is clear that $\lambda_1^{(2)} = 38$, $\lambda_2^{(2)} = 34$, $\lambda_3^{(2)} = 28$, $\lambda_4^{(2)} = 24$ and $\lambda_5^{(2)} = 20$ are of types \hat{A}_2 , \hat{A}_1 , \hat{A}_1 , \hat{A}_2 and \hat{A}_1 , respectively, and so these 2-marked parts are of type \hat{A} . Moreover, we have $\hat{R}_l(\lambda) = 1$, $\hat{R}_2(\lambda) = 2$, $\hat{R}_3(\lambda) = 3$, $\hat{R}_4(\lambda) = 4$ and $\hat{R}_5(\lambda) = 5$.

There are 1-marked and 3-marked parts 16 in $GG(\lambda)$. Moreover, note that $\lambda_6^{(1)} = 16$ and $\hat{R}_5(\lambda) = 5 \neq 6$, we see that $\hat{R}_6(\lambda) = 6$ and $\lambda_6^{(2)} = 14$ is of type \hat{B} . Similarly, we get that $\hat{R}_7(\lambda) = 7$ and $\lambda_7^{(2)} = 10$ is of type \hat{B} .

Note that there do not exist parts 8 with marks greater than 2 and $\lambda_9^{(1)} = 4$, then we see that $\hat{R}_8(\lambda) = 9$ and $\lambda_8^{(2)} = 6$ is of type \hat{C} . There are 1-marked and 3-marked parts 4 in $GG(\lambda)$. Moreover, note that $\lambda_9^{(1)} = 4$, $\lambda_{10}^{(1)} = 1$ and $\hat{R}_8(\lambda) = 9$, we see that $\hat{R}_9(\lambda) = 10$ and $\lambda_9^{(2)} = 2$ is of type \hat{O} .

Now, we define $\hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ to be the set of partitions λ in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$ such that there do not exist odd parts greater than or equal to $\lambda_p^{(2)}$, and $\lambda_p^{(2)}$ is of type \hat{A} or type \hat{B} or type \hat{C} . For example, the partition λ in (5.1) belongs to $\hat{\mathbb{C}}_1(10, 9, 6; 3|8)$.

Let λ be a partition in $\hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, we define the clusters based on the types of $\lambda_1^{(2)}, \lambda_2^{(2)}, \ldots, \lambda_p^{(2)}$. For convenience, we assume that $\lambda_0^{(2)} = +\infty$. We start from the part $\lambda_p^{(2)}$, assume that p_1 is the smallest integer such that $p_1 \leq p$, $\lambda_{p_1}^{(2)} - \lambda_p^{(2)} = 4(p - p_1)$ and satisfying one of the following four conditions:

- (1) $\lambda_p^{(2)}, \ldots, \lambda_{p_1}^{(2)}$ are of type \hat{A}_1 and $\lambda_{p_1-1}^{(2)} > \lambda_{p_1}^{(2)} + 4$, we say that the set $\{p_1, \ldots, p\}$ is of cluster \hat{A}_1 .
- (2) $\lambda_p^{(2)}, \ldots, \lambda_{p_1}^{(2)}$ are of type \hat{A} and $\lambda_{p_1}^{(2)}$ is of type \hat{A}_2 , we say that the set $\{p_1, \ldots, p\}$ is of cluster \hat{A}_2 .
- (3) $\lambda_p^{(2)}, \ldots, \lambda_{p_1}^{(2)}$ are of type \hat{B} , we say that the set $\{p_1, \ldots, p\}$ is of cluster \hat{B} .
- (4) $\lambda_p^{(2)}, \ldots, \lambda_{p_1}^{(2)}$ are of type \hat{C} , we say that the set $\{p_1, \ldots, p\}$ is of cluster \hat{C} .

Define $\hat{\alpha}_p(\lambda) = \cdots = \hat{\alpha}_{p_1}(\lambda) = \{p_1, \ldots, p\}$. If $p_1 > 1$, then set b = 1 and execute the following procedure:

(E) Assume that p_{b+1} is the smallest integer such that $p_{b+1} \leq p_b - 1$, $\lambda_{p_{b+1}}^{(2)} - \lambda_{p_b-1}^{(2)} = 4(p_b - 1 - p_{b+1})$ and satisfying one of the following five conditions:

- (1) $\lambda_{p_{b-1}}^{(2)}, \ldots, \lambda_{p_{b+1}}^{(2)}$ are of type \hat{A}_1 and $\lambda_{p_{b+1}-1}^{(2)} > \lambda_{p_{b+1}}^{(2)} + 4$, we say that the set $\{p_{b+1}, \ldots, p_b 1\}$ is of cluster \hat{A}_1 .
- (2) $\lambda_{p_{b-1}}^{(2)}, \ldots, \lambda_{p_{b+1}}^{(2)}$ are of type \hat{A} , $\lambda_{p_{b+1}}^{(2)}$ is of type \hat{A}_2 , and there is no 1-marked part $\lambda_{p_{b-1}}^{(2)} 4$ in $GG(\lambda)$, then we say that the set $\{p_{b+1}, \ldots, p_b 1\}$ is of cluster \hat{A}_2 .
- (3) $\lambda_{p_{b-1}}^{(2)}, \ldots, \lambda_{p_{b+1}}^{(2)}$ are of type $\hat{A}, \lambda_{p_{b+1}}^{(2)}$ is of type \hat{A}_2 , and there is a 1-marked part $\lambda_{p_{b-1}}^{(2)} 4$ in $GG(\lambda)$, then we say that the set $\{p_{b+1}, \ldots, p_b 1\}$ is of cluster \hat{A}_3 .
- (4) $\lambda_{p_b-1}^{(2)}, \ldots, \lambda_{p_{b+1}}^{(2)}$ are of type \hat{B} , we say that the set $\{p_{b+1}, \ldots, p_b-1\}$ is of cluster \hat{B} .
- (5) $\lambda_{p_{b-1}}^{(2)}, \ldots, \lambda_{p_{b+1}}^{(2)}$ are of type \hat{C} , we say that the set $\{p_{b+1}, \ldots, p_b-1\}$ is of cluster \hat{C} .

Define $\hat{\alpha}_{p_b-1}(\lambda) = \dots = \hat{\alpha}_{p_{b+1}}(\lambda) = \{p_{b+1}, \dots, p_b-1\}.$

(F) Replace b by b + 1. If $p_b = 1$, then we are done. If $p_b > 1$, then go back to (E).

For example, let λ be the partition in (5.1). For p = 8, we find that the intervals $\{8\}$, $\{6,7\}$, $\{4,5\}$, $\{3\}$ and $\{1,2\}$ are of clusters \hat{C} , \hat{B} , \hat{A}_3 , \hat{A}_1 and \hat{A}_2 , respectively. For p = 5, we derive that the intervals $\{4,5\}$, $\{3\}$ and $\{1,2\}$ are of clusters \hat{A}_2 , \hat{A}_1 and \hat{A}_2 , respectively. respectively.

5.2 The set $\check{\mathbb{C}}_{j}(N_{1}, \ldots, N_{k-1}; i|p)$

Let μ be a partition in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$ such that there exist odd parts in μ . Let $\mu^{(r)} = (\mu_1^{(r)}, \mu_2^{(r)}, \ldots, \mu_{N_r}^{(r)})$ denote *r*-marked parts in $GG(\mu)$, where $\mu_1^{(r)} > \mu_2^{(r)} > \cdots > \mu_{N_r}^{(r)}$ and $1 \leq r \leq k-1$. We consider the largest odd part 2t+1 in μ . We say that 2t+1 is a reduction part of μ if 2t+1 is not a insertion part of μ , namely, μ satisfies either of the following two conditions:

- (1) The mark of 2t + 1 is greater than 1 in $GG(\mu)$.
- (2) $\mu_s^{(1)} = 2t + 1$, 2t + 2 occurs in μ , and 2t + 4b + 2 occurs in μ if $\mu_{s-b}^{(1)} = 2t + 4b$, where $1 \le b < s \le N_1$.

By definition, we see that if 2t+1 is a reduction part in μ , then there exists a 2-marked part $\mu_p^{(2)} = 2t$ or 2t+1 or 2t+2 in $GG(\mu)$. We say that the reduction index of 2t+1 is p.

We define $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ to be the set of partitions μ in $\mathbb{C}_j(N_1, \ldots, N_{k-1}; i)$ such that there exist odd parts in μ and the largest odd part is a reduction part with the reduction index p.

For example, let μ be a partition in $\mathbb{C}_1(10, 9, 6; 3)$, whose Göllnitz-Gordon marking is

$$GG(\mu) = \begin{bmatrix} 4 & 10 & 14 & 22 & 28 & 36 \\ 2 & 5 & 10 & 14 & 18 & 22 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 20 & 24 & 28 & 32 & 36 \end{bmatrix}.$$
 (5.2)

We see that $\lambda_8^{(2)} = 5$ is the largest odd part in μ . So we have $\mu \in \check{\mathbb{C}}_1(10, 9, 6; 3|8)$.

For another example, let μ be a partition in $\mathbb{C}_1(10, 9, 6; 3)$, whose Göllnitz-Gordon marking is

$$GG(\mu) = \begin{bmatrix} 4 & 12 & 16 & 22 & 28 & 36 \\ 2 & 6 & 10 & 14 & 20 & 24 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 19 & 22 & 28 & 32 & 36 \end{bmatrix}.$$
 (5.3)

We see that $\mu_5^{(1)} = 19$ is the largest odd part in μ . Observe that $\mu_4^{(1)} = 22$, $\mu_3^{(1)} = 28 > 26$, and 20 and 24 occur in μ , so 19 is a reduction of μ . Moreover, note that $\mu_5^{(2)} = 20$, we have $\mu \in \check{\mathbb{C}}_1(10, 9, 6; 3|5)$.

It is clear that for 1 ,

$$\check{\mathbb{C}}_j(N_1,\ldots,N_{k-1};i|p)\subseteq \hat{\mathbb{C}}_j(N_1,\ldots,N_{k-1};i|p-1).$$

Let μ be a partition in $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$. We will define the types of $\mu_1^{(2)}, \mu_2^{(2)}, \ldots, \mu_p^{(2)}$ and the clusters related to $\mu_1^{(2)}, \mu_2^{(2)}, \ldots, \mu_p^{(2)}$. We first define the type of the part $\mu_p^{(2)}$. There are four cases.

- (1) $\mu_p^{(2)}$ is even and there is a part $\mu_p^{(2)} + 1$ with mark greater than 2 in $GG(\mu)$. Then, $\mu_p^{(2)} + 1$ is the reduction part of μ . We say that $\mu_p^{(2)}$ is of type \check{A}_1 .
- (2) $\mu_p^{(2)}$ is even and there is no $\mu_p^{(2)} + 1$ in μ . Then, there is a 1-marked part $\mu_p^{(2)} 1$ in $GG(\mu)$, which is the reduction part of μ . We say that $\mu_p^{(2)}$ is of type \check{A}_2 .
- (3) $\mu_p^{(2)}$ is even and there is a 1-marked part $\mu_p^{(2)} + 1$ in $GG(\mu)$. Then, $\mu_p^{(2)} + 1$ is the reduction part of μ . We say that $\mu_p^{(2)}$ is of type \check{B} .
- (4) $\mu_p^{(2)}$ is odd. Then, $\mu_p^{(2)}$ is the reduction part of μ . We say that $\mu_p^{(2)}$ is of type \check{C} .

Assume that p_1 is the smallest integer such that $p_1 \leq p$, $\mu_s^{(2)} = \mu_p^{(2)} + 4(p-s)$ (resp. $\mu_p^{(2)} + 4(p-s) - 1$) if $\mu_p^{(2)}$ is an even part (resp. an odd part) for $p_1 \leq s \leq p$, and satisfying one of the following conditions:

- (1) $\mu_p^{(2)}$ is of type \check{A}_1 and there is a 3-marked part $\mu_s^{(2)}$ or $\mu_s^{(2)} + 1$ in $GG(\mu)$ for $p_1 \leq s \leq p$. We say that the set $\{p_1, \ldots, p\}$ is of cluster \check{A}_1 and $\mu_{p-1}^{(2)}, \ldots, \mu_{p_1}^{(2)}$ are of type \check{A}_1 .
- (2) $\mu_p^{(2)}$ is of type \check{A}_2 and there is a 1-marked part $\mu_{p_1}^{(2)} 2$ or $\mu_{p_1}^{(2)} 1$ in $GG(\mu)$. We say that the set $\{p_1, \ldots, p\}$ is of cluster \check{A}_2 and $\mu_{p-1}^{(2)}, \ldots, \mu_{p_1}^{(2)}$ are of type \check{A}_2 .

- (3) $\mu_p^{(2)}$ is of type \check{B} and there exist parts $\mu_l^{(2)} + 2$ in μ for $p_1 \leq l \leq p$. We say that the set $\{p_1, \ldots, p\}$ is of cluster \check{B} and $\mu_{p-1}^{(2)}, \ldots, \mu_{p_1}^{(2)}$ are of type \check{B} .
- (4) $\mu_p^{(2)}$ is of type \check{C} . We say that the set $\{p_1, \ldots, p\}$ is of cluster \check{C} and $\mu_{p-1}^{(2)}, \ldots, \mu_{p_1}^{(2)}$ are of type \check{C} .

Define $\check{\beta}_p(\mu) = \cdots = \check{\beta}_{p_1}(\mu) = \{p_1, \ldots, p\}$. If $p_1 > 1$, then set b = 1 and carry out the following procedure:

- (E) Assume that p_{b+1} is the smallest integer such that $p_{b+1} \leq p_b 1$, $\mu_{p_{b+1}}^{(2)} \mu_{p_b-1}^{(2)} = 4(p_b 1 p_{b+1})$, and satisfying one of the following conditions:
 - (1) There are 1-marked and 3-marked parts $\mu_s^{(2)}$ in $GG(\mu)$ for $p_{b+1} \leq s \leq p_b 1$. We say that the set $\{p_{b+1}, \ldots, p_b - 1\}$ is of cluster \check{A}_1 and $\mu_{p_b-1}^{(2)}, \ldots, \mu_{p_{b+1}}^{(2)}$ are of type \check{A}_1 .
 - (2) There is a 1-marked part $\mu_s^{(2)} 2$ in $GG(\mu)$ for $p_{b+1} \leq s \leq p_b 1$, and $\mu_{p_{b+1}}^{(2)} + 2$ does not occur in μ . We say that the set $\{p_{b+1}, \ldots, p_b 1\}$ is of cluster \check{A}_2 and $\mu_{p_b-1}^{(2)}, \ldots, \mu_{p_{b+1}}^{(2)}$ are of type \check{A}_2 .
 - (3) There is a 1-marked part $\mu_s^{(2)} 2$ in $GG(\mu)$ for $p_{b+1} \leq s \leq p_b 1$, $\mu_{p_{b+1}}^{(2)} + 2$ occurs in μ , and $\mu_{p_b-1}^{(2)}$ appears exactly once in μ . We say that the set $\{p_{b+1}, \ldots, p_b-1\}$ is of cluster \check{A}_3 and $\mu_{p_b-1}^{(2)}, \ldots, \mu_{p_{b+1}}^{(2)}$ are of type \check{A}_3 .
 - (4) There is a 1-marked part $\mu_s^{(2)} 2$ in $GG(\mu)$ and $\mu_s^{(2)}$ appears at least twice in μ for $p_{b+1} \leq s \leq p_b - 1$, and $\mu_{p_{b+1}}^{(2)} + 2$ occurs in μ . We say that the set $\{p_{b+1}, \ldots, p_b - 1\}$ is of cluster \check{B} and $\mu_{p_b-1}^{(2)}, \ldots, \mu_{p_{b+1}}^{(2)}$ are of type \check{B} .
 - (5) There is a 1-marked part $\mu_s^{(2)}$ in $GG(\mu)$ for $p_{b+1} \leq s \leq p_b 1$, and $\mu_{p_b-1}^{(2)}$ appears exactly twice in μ . We say that the set $\{p_{b+1}, \ldots, p_b - 1\}$ is of cluster \check{C} and $\mu_{p_b-1}^{(2)}, \ldots, \mu_{p_{b+1}}^{(2)}$ are of type \check{C} .

Define $\check{\beta}_{p_b-1}(\mu) = \cdots = \check{\beta}_{p_{b+1}}(\mu) = \{p_{b+1}, \dots, p_b - 1\}.$

(F) Replace b by b + 1. If $p_b = 1$, then we are done. If $p_b > 1$, then go back to (E).

For example, let μ be the partition in (5.2). We find that the intervals $\{8\}$, $\{6,7\}$, $\{4,5\}$, $\{3\}$ and $\{1,2\}$ are of clusters \check{C} , \check{B} , \check{A}_3 , \check{A}_1 and \check{A}_2 , respectively.

For another example, let μ be the partition in (5.3), we derive that the intervals $\{4, 5\}$, $\{3\}$ and $\{1, 2\}$ are of clusters \check{A}_2 , \check{A}_1 and \check{A}_2 , respectively.

5.3 Special partitions

In order to give a description of the reduction R_p between $\hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ and $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, we need to introduce the special partitions. A special partition λ is an ordinary partition in which the largest odd part in λ may be overlined. We generalize the definition of the Göllnitz-Gordon marking of an ordinary partition to a special partition as follows.

Definition 5.2. The Göllnitz-Gordon marking of a special partition λ , denoted by $GG(\lambda)$, is an assignment of positive integers (marks) to the parts of $\lambda = (\lambda_1, \lambda_2, \ldots, \lambda_\ell)$ from smallest to largest such that the marks are as small as possible subject to the conditions that for $1 \leq s \leq \ell$,

- (1) the integer assigned to λ_s is different from the integers assigned to the parts λ_g such that $\lambda_s \lambda_g \leq 2$ with strict inequality if λ_s is an odd part for g > s;
- (2) λ_s can not be assigned with 1 if λ_s is a special odd part.

Definition 5.3. We define $\overleftarrow{C}_i(k;i)$ to be the set of special partitions λ such that

(1)
$$f_1(\lambda) + f_2(\lambda) \le i - 1;$$

(2)
$$f_{2t+1}(\lambda) + f_{\overline{2t+1}}(\lambda) \le 1;$$

- (3) $f_{2t}(\lambda) + f_{2t+1}(\lambda) + f_{\overline{2t+1}}(\lambda) + f_{2t+2}(\lambda) \le k 1;$
- (4) if $f_{2t}(\lambda) + f_{2t+1}(\lambda) + f_{2t+2}(\lambda) = k 1$, then $tf_{2t}(\lambda) + tf_{2t+1}(\lambda) + (t+1)f_{2t+2}(\lambda) \equiv O_{\lambda}(2t+2) + i 1 \pmod{2-j}$, where $O_{\lambda}(N)$ denotes the number of odd parts not exceeding N in λ ;
- (5) if there is a $\overline{2s+1}$ in λ , then there are 1-marked 2s-2 and 2s+2 in $\overline{GG}(\lambda)$.

Let $N_r(\lambda)$ (or N_r for short) denote the number of r-marked parts in $\overline{GG}(\lambda)$. If λ is a partition in $\overline{C}_j(k;i)$, then it is clear that $N_1 \geq N_2 \geq \cdots \geq N_{k-1} \geq 0$. Let $\overline{C}_j(N_1,\ldots,N_{k-1};i)$ denote the set of special partitions λ counted by $\overline{C}_j(k,i)$ such that there are N_r r-marked parts in $\overline{GG}(\lambda)$ for $1 \leq r \leq k-1$.

We also generalize the definition of (k-1)-bands of an ordinary partition in $\mathbb{C}_1(k, i)$ to a special partition in $\overleftarrow{C}_1(k; i)$. For the consecutive k-1 parts $\lambda_s, \lambda_{s+1}, \ldots, \lambda_{s+k-2}$ of a special partition $\lambda \in \overleftarrow{C}_1(k; i)$, if such k-1 parts satisfy the difference condition

$$\lambda_s \le \lambda_{s+k-2} + 2$$
 with strict inequality if λ_s is odd, (5.4)

then we say that $\{\lambda_{s+l}\}_{0 \le l \le k-2}$ is a (k-1)-band of λ . If the k-1 parts in the (k-1)-band $\{\lambda_{s+l}\}_{0 \le l \le k-2}$ also satisfy the congruence

$$[\lambda_s/2] + \dots + [\lambda_{s+k-2}/2] \equiv i - 1 + O_\lambda(\lambda_s) \pmod{2}, \tag{5.5}$$

then we say that the (k-1)-band $\{\lambda_{s+l}\}_{0 \leq l \leq k-2}$ is good. Otherwise, we say that the (k-1)-band $\{\lambda_{s+l}\}_{0 \leq l \leq k-2}$ is bad. Then λ is a special partition in $\overleftarrow{C}_0(k;i)$ if and only if all the (k-1)-bands of λ are good. With the similar argument in the proof of Lemma 3.3, we can obtain the following lemma.

Lemma 5.4. Let $\{\lambda_{s+l}\}_{0 \leq l \leq k-2}$ and $\{\lambda_{g+l}\}_{0 \leq l \leq k-2}$ be two (k-1)-bands of a special partition $\lambda \in \overleftarrow{C}_1(k; i)$. If $\lambda_g > \lambda_s$ and $\lambda_{s+k-2} \geq \lambda_g - 4$ with strict inequality if λ_g is odd, then $\{\lambda_{s+l}\}_{0 \leq l \leq k-2}$ and $\{\lambda_{g+l}\}_{0 \leq l \leq k-2}$ are both good or bad.

Let λ be a special partition in $\overleftarrow{C}_1(k; i)$, assume that there are N_{k-1} (k-1)-marked parts in $\overline{GG}(\lambda)$, denoted $\lambda_1^{(k-1)} > \lambda_2^{(k-1)} > \cdots > \lambda_{N_{k-1}}^{(k-1)}$. For each $p \in [1, N_{k-1}]$, assume that $\lambda_p^{(k-1)}$ is the s_p -th part λ_{s_p} of λ . If $\lambda_p^{(k-1)}$ is not a special odd part, then there exist k-2 parts λ_g in λ such that g > s and $\lambda_g \ge \lambda_s - 2$ with strict inequality if λ_s is an odd part. Then, $\{\lambda_{s+l}\}_{0\le l\le k-2}$ is a (k-1)-band of λ . Such a (k-1)-band is called the (k-1)-band induced by $\lambda_p^{(k-1)}$, denoted $\{\lambda_p^{(k-1)}\}_{k-1} = \{\lambda_{p,1}^{(k-1)}, \lambda_{p,2}^{(k-1)}, \dots, \lambda_{p,k-1}^{(k-1)}\}$, where $\lambda_{p,1}^{(k-1)} \ge \lambda_{p,2}^{(k-1)} \ge \cdots \ge \lambda_{p,k-1}^{(k-1)}$.

If $\lambda_p^{(k-1)}$ is a special odd part, set $\lambda_p^{(k-1)} = \overline{2t+1}$, then there exist 2-marked, ..., (k-2)-marked parts 2t and a 1-marked part 2t+2 in $\overline{GG}(\lambda)$. We define the (k-1)-band induced by $\lambda_p^{(k-1)}$ as $\{\lambda_p^{(k-1)}\}_{k-1} = \{\lambda_{p,1}^{(k-1)}, \lambda_{p,2}^{(k-1)}, \dots, \lambda_{p,k-1}^{(k-1)}\}$, where $\lambda_{p,1}^{(k-1)} = 2t+2$, $\lambda_{p,2}^{(k-1)} = \overline{2t+1}$ and $\lambda_{p,3}^{(k-1)} = \cdots = \lambda_{p,k-1}^{(k-1)} = 2t$.

Clearly, the k-1 parts in the (k-1)-band $\{\lambda_p^{(k-1)}\}_{k-1}$ satisfy the difference condition (5.4). With the similar argument in the proof of Theorem 3.2, we obtain the following theorem, which gives a criterion to determine whether a special partition λ in $\overleftarrow{C}_1(k;i)$ also belongs to $\overleftarrow{C}_0(k;i)$.

Theorem 5.5. Let λ be a special partition in $\overleftarrow{C}_1(k;i)$ with N_{k-1} (k-1)-marked parts in $\overline{GG}(\lambda)$, denoted $\lambda_1^{(k-1)} > \lambda_2^{(k-1)} > \cdots > \lambda_{N_{k-1}}^{(k-1)}$. Then λ is a special partition in $\overleftarrow{C}_0(k;i)$ if and only if $\{\lambda_p^{(k-1)}\}_{k-1}$ is good for $1 \leq p \leq N_{k-1}$.

5.4 Proof of Theorem 5.1

We are now in a position to give a proof a Theorem 5.1.

Proof of Theorem 5.1. Let λ be a partition in $\hat{\mathbb{C}}_1(N_1, \ldots, N_{k-1}; i|p)$. Let $\lambda^{(r)} = (\lambda_1^{(r)}, \lambda_2^{(r)}, \ldots, \lambda_{N_r}^{(r)})$ denote the *r*-marked parts in $GG(\lambda)$, where $1 \leq r \leq k-1$. We will define $\mu = R_p(\lambda)$ related to $\lambda_1^{(2)}, \lambda_2^{(2)}, \ldots, \lambda_p^{(2)}$, and denote the intermediate special partitions by $\lambda^1, \lambda^2, \ldots, \lambda^p$.

Step 1: We do the following operation related to $\lambda_1^{(2)}$, called the basic operation of $\lambda_1^{(2)}$. There are five cases.

Case 1: $\hat{\alpha}_1(\lambda)$ is of cluster \hat{A}_1 . We may write $\lambda_1^{(2)} = 2t_1$. In this case, $\lambda_1^{(2)}$ is of type \hat{A}_1 . So, there is a 1-marked part $2t_1$ in $GG(\lambda)$ and there exist parts $2t_1 + 2$ in λ . Let r_1 be the smallest mark of parts $2t_1 + 2$ in $GG(\lambda)$. We have $r_1 > 2$. Then we replace the r_1 -marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$ to get λ^1 .

Case 2: $\hat{\alpha}_1(\lambda)$ is of cluster \hat{A}_2 . We may write $\lambda_1^{(2)} = 2t_1 + 2$. In this case, $\lambda_1^{(2)}$ is of type \hat{A}_2 . So, there is a 1-marked part $2t_1 + 2$ in $GG(\lambda)$ and there do not exist parts $2t_1 + 4$ in λ . Then we set $r_1 = 1$ and replace the 1-marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$ to obtain λ^1 .

Case 3: $\hat{\alpha}_1(\lambda)$ is of cluster \hat{A}_3 . We find that $\hat{\alpha}_1(\lambda) \neq \hat{\alpha}_p(\lambda)$ and $\lambda_1^{(2)}$ is of type \hat{A}_2 . Set $\lambda_1^{(2)} = 2t_1 + 2$. We see that there is a 1-marked part $2t_1 + 2$ in $GG(\lambda)$ and there do not exist parts $2t_1 + 4$ in λ . Then we set $r_1 = 2$ and replace the 2-marked $2t_1 + 2$ in $GG(\lambda)$ by $\overline{2t_1 + 1}$ to obtain λ^1 .

Case 4: $\hat{\alpha}_1(\lambda)$ is of cluster \hat{B} . Then, $\lambda_1^{(2)}$ is of type \hat{B} . Set $\lambda_1^{(2)} = 2t_1$. We see that there exist parts $2t_1 + 2$ with mark 1 and marks greater than 2 in $GG(\lambda)$. There are two subcases.

Case 4.1: $\hat{\alpha}_1(\lambda) = \hat{\alpha}_p(\lambda)$. Then we set $r_1 = 1$ and replace the 1-marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$ to get λ^1 .

Case 4.2: $\hat{\alpha}_1(\lambda) \neq \hat{\alpha}_p(\lambda)$. Let r_1 be the smallest mark except for 1 of parts $2t_1 + 2$ in $GG(\lambda)$. We have $r_1 > 2$. Then, the λ^1 is obtained by replacing the r_1 -marked $2t_1 + 2$ in $GG(\lambda)$ by $\overline{2t_1 + 1}$.

Case 5: $\hat{\alpha}_1(\lambda)$ is of cluster \hat{C} . We may write $\lambda_1^{(2)} = 2t_1 + 2$. In this case, $\lambda_1^{(2)}$ is of type \hat{C} . So, there is a 1-marked part $2t_1$ and there do not exist parts $2t_1 + 4$ with marks greater than 2 in $GG(\lambda)$. Then we set $r_1 = 2$ and replace the 2-marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$ to get λ^1 .

Obviously, $|\lambda^1| = |\lambda| - 1$. We see that λ^1 is obtained by replacing the r_1 -marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$ (resp. $\overline{2t_1 + 1}$). We proceed to prove that the new generated part $2t_1 + 1$ (resp. $\overline{2t_1 + 1}$) is marked with r_1 in $\overline{GG}(\lambda^1)$. It suffice to show that there are 2-marked, ..., $(r_1 - 1)$ -marked parts $2t_1$ and there is a 1-marked $2t_1$ (resp. $2t_1 + 2$) in $GG(\lambda)$.

If λ^1 is obtained by replacing the r_1 -marked $2t_1 + 2$ in $GG(\lambda)$ by $2t_1 + 1$, then r_1 is the smallest mark of parts $2t_1 + 2$ in $GG(\lambda)$, and so there are 1-marked, ..., $(r_1 - 1)$ -marked parts $2t_1$ in $GG(\lambda)$.

If λ^1 is obtained by replacing the r_1 -marked $2t_1 + 2$ in $GG(\lambda)$ by $\overline{2t_1 + 1}$, then there is a 1-marked $2t_1 + 2$ and r_1 is the smallest mark except for 1 of parts $2t_1 + 2$ in $GG(\lambda)$, and so there are 2-marked, ..., $(r_1 - 1)$ -marked $2t_1$ in $GG(\lambda)$.

We conclude that there are 2-marked, ..., $(r_1 - 1)$ -marked $2t_1$ and there is a 1marked $2t_1$ (resp. $2t_1 + 2$) in $GG(\lambda)$. So, the new generated part $2t_1 + 1$ (resp. $\overline{2t_1 + 1}$) is marked with r_1 in $\overline{GG}(\lambda^1)$. Moreover, the marks of the unchanged parts in $\overline{GG}(\lambda^1)$ are the same as those in $GG(\lambda)$. Hence, λ^1 is a special partition in $\overleftarrow{C}_1(N_1,\ldots,N_{k-1};i)$.

We turn to show that if $\lambda \in \mathbb{C}_0(N_1, \ldots, N_{k-1}; i)$, then $\lambda^1 \in \overleftarrow{C}_0(N_1, \ldots, N_{k-1}; i)$. Assume that $\lambda \in \mathbb{C}_0(N_1, \ldots, N_{k-1}; i)$, then $\{\lambda_s^{(k-1)}\}_{k-1}$ is good for $1 \leq s \leq N_{k-1}$. Let $\lambda^{1,(k-1)} = (\lambda_1^{1,(k-1)}, \lambda_2^{1,(k-1)}, \ldots, \lambda_{N_{k-1}}^{1,(k-1)})$ be the (k-1)-marked parts in $\overrightarrow{GG}(\lambda^1)$. From the proof above, we have $\lambda_{r_1}^{1,(2)} = 2t_1 + 1$ (resp. $\overrightarrow{2t_1 + 1}$). For each $s \in [1, N_{k-1}]$, we find that $\lambda_s^{1,(k-1)} \leq 2t_1 + 2$. To show that $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good, we consider the following four cases.

- (1) $\lambda_s^{1,(k-1)} \leq 2t_1$. By the construction of λ^1 , we have $\{\lambda_s^{1,(k-1)}\}_{k-1} = \{\lambda_s^{(k-1)}\}_{k-1}$ and $O_{\lambda^1}(\lambda_s^{1,(k-1)}) = O_{\lambda}(\lambda_s^{(k-1)})$. Note that $\{\lambda_s^{(k-1)}\}_{k-1}$ is good, so $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good.
- (2) $\lambda_s^{1,(k-1)} = 2t_1 + 1$. In this case, we have $r_1 = k 1$ and $\lambda_s^{(k-1)} = 2t_1 + 2$. Moreover, we derive that $\lambda_{s,2}^{1,(k-1)} = \cdots = \lambda_{s,k-1}^{1,(k-1)} = \lambda_{s,2}^{(k-1)} = \cdots = \lambda_{s,k-1}^{(k-1)} = 2t_1$ and $O_{\lambda^1}(2t_1 + 1) = O_{\lambda}(2t + 2) + 1$. So, we get

$$\begin{split} [\lambda_{s,1}^{1,(k-1)}/2] &+ [\lambda_{s,2}^{1,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{1,(k-1)}/2] \\ &= t_1 f_{2t_1}(\lambda^1) + t_1 f_{2t_1+1}(\lambda^1) = t_1(k-2) + t_1 \\ &= t_1(k-2) + (t_1+1) - 1 = t_1 f_{2t_1}(\lambda) + (t_1+1) f_{2t_1+2}(\lambda) - 1 \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \dots + [\lambda_{s,k-1}^{(k-1)}/2] - 1 \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) - 1 = i - 1 + O_{\lambda}(2t_1+2) - 1 \\ &\equiv i - 1 + O_{\lambda^1}(2t_1+1) = i - 1 + O_{\lambda^1}(\lambda_{s,1}^{1,(k-1)}) \pmod{2}, \end{split}$$

which implies that $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good.

(3) $\lambda_{s}^{1,(k-1)} = \overline{2t_1 + 1}$. In this case, we have $r_1 = k - 1$ and $\lambda_{s}^{(k-1)} = 2t_1 + 2$. Moreover, we find that $\lambda_{s,1}^{1,(k-1)} = \lambda_{s,2}^{1,(k-1)} = 2t_1 + 2$, $\lambda_{s,3}^{1,(k-1)} = \cdots = \lambda_{s,k-1}^{1,(k-1)} = \lambda_{s,3}^{1,(k-1)} = \cdots = \lambda_{s,k-1}^{1,(k-1)} = 2t_1$ and $O_{\lambda^1}(2t_1 + 2) = O_{\lambda}(2t + 2) + 1$. We obtain that $[\lambda_{s,1}^{1,(k-1)}/2] + [\lambda_{s,2}^{1,(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{1,(k-1)}/2]$ $= t_1 f_{2t_1}(\lambda^1) + t_1 f_{\overline{2t_1+1}}(\lambda^1) + (t_1 + 1) f_{2t_1+2}(\lambda^1) = t_1(k - 3) + t_1 + (t_1 + 1)$ $= t_1(k - 3) + 2(t_1 + 1) - 1 = t_1 f_{2t_1}(\lambda) + (t_1 + 1) f_{2t_1+2}(\lambda) - 1$ $= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{(k-1)}/2] - 1$ $\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) - 1 = i - 1 + O_{\lambda}(2t_1 + 2) - 1$ $\equiv i - 1 + O_{\lambda^1}(2t_1 + 2) = i - 1 + O_{\lambda^1}(\lambda_{s,1}^{1,(k-1)}) \pmod{2}.$

So, $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good.

(4)
$$\lambda_s^{1,(k-1)} = 2t_1 + 2$$
. By the construction of λ^1 , we find that $\lambda_s^{(k-1)} = 2t_1 + 2$,
 $f_{2t_1}(\lambda^1) = f_{2t_1}(\lambda), f_{2t_1+1}(\lambda^1) + f_{\overline{2t_1+1}}(\lambda^1) = f_{2t_1+1}(\lambda) + f_{\overline{2t_1+1}}(\lambda) + 1, f_{2t_1+2}(\lambda^1) = f_{2t_1+2}(\lambda) - 1$ and $O_{\lambda^1}(2t_1 + 2) = O_{\lambda}(2t + 2) + 1$, and so

$$\begin{split} [\lambda_{s,1}^{1,(k-1)}/2] &+ [\lambda_{s,2}^{1,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{1,(k-1)}/2] \\ &= t_1 f_{2t_1}(\lambda^1) + t_1 (f_{2t_1+1}(\lambda^1) + f_{\overline{2t_1+1}}(\lambda^1)) + (t_1+1) f_{2t_1+2}(\lambda^1) \\ &= t_1 f_{2t_1}(\lambda) + t_1 (f_{2t_1+1}(\lambda) + f_{\overline{2t_1+1}}(\lambda) + 1) + (t_1+1) (f_{2t_1+2}(\lambda) - 1) \\ &= t_1 f_{2t_1}(\lambda) + t_1 (f_{2t_1+1}(\lambda) + f_{\overline{2t_1+1}}(\lambda)) + (t_1+1) f_{2t_1+2}(\lambda) - 1 \\ &= [\lambda_{s,1}^{(k-1)}/2] + [\lambda_{s,2}^{(k-1)}/2] + \dots + [\lambda_{s,k-1}^{(k-1)}/2] - 1 \\ &\equiv i - 1 + O_{\lambda}(\lambda_{s,1}^{(k-1)}) - 1 = i - 1 + O_{\lambda}(2t_1+2) - 1 \\ &\equiv i - 1 + O_{\lambda^1}(2t_1+2) = i - 1 + O_{\lambda^1}(\lambda_{s,1}^{1,(k-1)}) \pmod{2}. \end{split}$$

Therefore, $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good.

Hence, the (k-1)-band $\{\lambda_s^{1,(k-1)}\}_{k-1}$ is good for $s \in [1, N_{k-1}]$. It follows from Theorem 5.5 that λ^1 is a special partition in $C_0(N_1, \ldots, N_{k-1}; i)$.

Step 2: If p = 1, then we shall do nothing. If $p \ge 2$, then set b = 1 and repeat the following process.

- (E) Replace the r_b -marked $2t_b + 1$ (or $\overline{2t_b + 1}$) in $\overline{GG}(\lambda^b)$ by $2t_b$ and apply the basic operation defined in Step 1 to $\lambda_{b+1}^{(2)}$ to get λ^{b+1} .
- (F) replace b by b + 1. If b = p, then we are done. If b < p, then go back to (F).

Clearly, for $1 \leq b < p$, we have $|\lambda^{b+1}| = |\lambda^b| - 2$. Moreover, λ^{b+1} is obtained by replacing the r_b -marked $2t_b + 1$ (resp. $\overline{2t_b + 1}$) in $\overline{GG}(\lambda^b)$ by $2t_b$ and replacing the r_{b+1} -marked $2t_{b+1} + 2$ in $\overline{GG}(\lambda^b)$ by $2t_{b+1} + 1$ or $\overline{2t_{b+1} + 1}$. With the similar argument in Step 1, we see that the new generated part $2t_{b+1} + 1$ (resp. $\overline{2t_{b+1} + 1}$) is marked with r_{b+1} in $\overline{GG}(\lambda^{b+1})$.

Now, we proceed to show that the new generated part $2t_b$ is marked with r_b in $\overline{GG}(\lambda^{b+1})$. If there does not exist a r_b -marked $2t_b - 2$ in $\overline{GG}(\lambda^b)$, then it is obviously right. If there is a r_b -marked $2t_b - 2$ in $\overline{GG}(\lambda^b)$, then we have $\hat{\alpha}_{b+1}(\lambda) = \hat{\alpha}_b(\lambda)$ and $r_{b+1} = r_b$. By the construction of λ^{b+1} , we see that we will replace the r_b -marked $2t_b - 2$ in $\overline{GG}(\lambda^b)$ by $2t_b - 3$ (resp. $\overline{2t_b - 3}$). So, the new generated part $2t_b$ is marked with r_b in $\overline{GG}(\lambda^{b+1})$.

Furthermore, the marks of the unchanged parts in $\overline{GG}(\lambda^{b+1})$ are the same as those in $\overline{GG}(\lambda^b)$. Hence, λ^{b+1} is a special partition in $C_1(N_1, \ldots, N_{k-1}; i)$.

We turn to show that if $\lambda^b \in \overleftarrow{C}_0(N_1, \dots, N_{k-1}; i)$, then $\lambda^{b+1} \in \overleftarrow{C}_0(N_1, \dots, N_{k-1}; i)$. Assume that $\lambda^b \in \overleftarrow{C}_0(N_1, \dots, N_{k-1}; i)$, let $\lambda^{b,(k-1)} = (\lambda_1^{b,(k-1)}, \dots, \lambda_{N_{k-1}}^{b,(k-1)})$ and $\lambda^{b+1,(k-1)} = (\lambda_1^{b+1,(k-1)}, \dots, \lambda_{N_{k-1}}^{b+1,(k-1)}) \text{ be the } (k-1)\text{-marked parts in } \overline{GG}(\lambda^b) \text{ and } \overline{GG}(\lambda^{b+1}), \text{ respectively. Then, } \{\lambda_s^{b,(k-1)}\}_{k-1} \text{ is good for } 1 \leq s \leq N_{k-1}.$

For each $s \in [1, N_{k-1}]$, by the construction of λ^{b+1} , we find that $\lambda_s^{b+1,(k-1)} \notin (2t_{b+1}+2, 2t_b)$ and there are no odd parts greater than $2t_b$ in λ^{b+1} . To show that $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good, we consider the following six cases.

- (1) $\lambda_s^{b+1,(k-1)} \leq 2t_{b+1} + 2$. With the similar argument in Step 1, we derive that $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.
- (2) $\lambda_s^{b+1,(k-1)} = 2t_b, r_b = k-1$ and there is a (k-1)-marked $2t_b + 1$ in $\overline{GG}(\lambda^b)$. In this case, we have $\lambda_s^{b,(k-1)} = 2t_b + 1, \lambda_{s,2}^{b,(k-1)} = \cdots = \lambda_{s,k-1}^{b,(k-1)} = \lambda_{s,2}^{b+1,(k-1)} = \cdots = \lambda_{s,k-1}^{b+1,(k-1)} = 2t_b$ and $O_{\lambda^{b+1}}(2t_b) = O_{\lambda^b}(2t_b + 1)$, which implies that

$$\begin{aligned} [\lambda_{s,1}^{b+1,(k-1)}/2] + [\lambda_{s,2}^{b+1,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b+1,(k-1)}/2] \\ &= t_b f_{2t_b}(\lambda^{b+1}) = t_b(k-1) \\ &= t_b(k-2) + t_b = t_b f_{2t_b}(\lambda^b) + t_b f_{2t_b+1}(\lambda^b) \\ &= [\lambda_{s,1}^{b,(k-1)}/2] + [\lambda_{s,2}^{b,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b,(k-1)}/2] \\ &\equiv i - 1 + O_{\lambda^b}(\lambda_{s,1}^{b,(k-1)}) = i - 1 + O_{\lambda^b}(2t_b + 1) \\ &= i - 1 + O_{\lambda^{b+1}}(2t_b) = i - 1 + O_{\lambda^{b+1}}(\lambda_{s,1}^{b+1,(k-1)}) \pmod{2}. \end{aligned}$$

So, $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.

(3) $\lambda_s^{b+1,(k-1)} = 2t_b, r_b = k-1$ and there is a (k-1)-marked $\overline{2t_b+1}$ in $\overline{GG}(\lambda^b)$. In this case, we have $\lambda_{s,2}^{b,(k-1)} = \overline{2t_b+1}, \lambda_{s,1}^{b,(k-1)} = 2t_b+2, \lambda_{s,3}^{b,(k-1)} = \cdots = \lambda_{s,k-1}^{b,(k-1)} = \lambda_{s,2}^{b+1,(k-1)} = \cdots = \lambda_{s,k-2}^{b+1,(k-1)} = 2t_b, \lambda_{s,k-1}^{b+1,(k-1)} = 2t_b - 2$ and $O_{\lambda^{b+1}}(2t_b) = O_{\lambda^b}(2t_b+2)$. So, we get

$$\begin{split} [\lambda_{s,1}^{b+1,(k-1)}/2] + [\lambda_{s,2}^{b+1,(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{b+1,(k-1)}/2] \\ &= (t_b - 1)f_{2t_b - 2}(\lambda^{b+1}) + t_bf_{2t_b}(\lambda^{b+1}) = (t_b - 1) + t_b(k - 2) \\ &\equiv t_b(k - 3) + t_b + (t_b + 1) \\ &= t_bf_{2t_b}(\lambda^b) + t_bf_{\overline{2t_b + 1}}(\lambda^b) + (t_b + 1)f_{2t_b + 2}(\lambda^b) \\ &= [\lambda_{s,1}^{b,(k-1)}/2] + [\lambda_{s,2}^{b,(k-1)}/2] + \cdots + [\lambda_{s,k-1}^{b,(k-1)}/2] \\ &\equiv i - 1 + O_{\lambda^b}(\lambda_{s,1}^{b,(k-1)}) = i - 1 + O_{\lambda^b}(2t_b + 2) \\ &= i - 1 + O_{\lambda^{b+1}}(2t_b) = i - 1 + O_{\lambda^{b+1}}(\lambda_{s,1}^{b+1,(k-1)}) \pmod{2}, \end{split}$$

which implies that $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.

(4) $\lambda_s^{b+1,(k-1)} = 2t_b$ and $r_b < k-1$. By the construction of λ^{b+1} , we find that there is a r_b -marked $2t_b - 2$ in $\overline{GG}(\lambda^b)$, which is replaced by $2t_b - 3$ or $\overline{2t_b - 3}$ in λ^{b+1} .

Moreover, we have $\lambda_s^{b,(k-1)} = 2t_b$, $f_{2t_b-2}(\lambda^{b+1}) = f_{2t_b-2}(\lambda^b) - 1$, $f_{2t_b}(\lambda^{b+1}) = f_{2t_b}(\lambda^b) + 1$ and $O_{\lambda^{b+1}}(2t_b) = O_{\lambda^b}(2t_b) + 1$. So, we get

$$\begin{split} [\lambda_{s,1}^{b+1,(k-1)}/2] &+ [\lambda_{s,2}^{b+1,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b+1,(k-1)}/2] \\ &= (t_b - 1)f_{2t_b - 2}(\lambda^{b+1}) + t_b f_{2t_b}(\lambda^{b+1}) \\ &= (t_b - 1)(f_{2t_b - 2}(\lambda^b) - 1) + t_b(f_{2t_b}(\lambda^b) + 1) \\ &= (t_b - 1)f_{2t_b - 2}(\lambda^b) + t_b f_{2t_b}(\lambda^b) + 1 \\ &= [\lambda_{s,1}^{b,(k-1)}/2] + [\lambda_{s,2}^{b,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b,(k-1)}/2] + 1 \\ &\equiv i - 1 + O_{\lambda^b}(\lambda_{s,1}^{b,(k-1)}) + 1 = i - 1 + O_{\lambda^b}(2t_b) + 1 \\ &= i - 1 + O_{\lambda^{b+1}}(2t_b) = i - 1 + O_{\lambda^{b+1}}(\lambda_{s,1}^{b+1,(k-1)}) \pmod{2}. \end{split}$$

Therefore, $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.

(5) $\lambda_s^{b+1,(k-1)} = 2t_b + 2$. In this case, we have $\lambda_s^{b,(k-1)} = 2t_b + 2$. Moreover, we find that $f_{2t_b}(\lambda^{b+1}) = f_{2t_b}(\lambda^b) + 1$, $f_{2t_b+1}(\lambda^{b+1}) + f_{\overline{2t_b+1}}(\lambda^{b+1}) = f_{2t_b+1}(\lambda^b) + f_{\overline{2t_b+1}}(\lambda^b) - 1$, $f_{2t_b+2}(\lambda^{b+1}) = f_{2t_b+2}(\lambda^b)$ and $O_{\lambda^{b+1}}(2t_b+2) = O_{\lambda^b}(2t_b+2)$. So

$$\begin{split} [\lambda_{s,1}^{b+1,(k-1)}/2] &+ [\lambda_{s,2}^{b+1,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b+1,(k-1)}/2] \\ &= t_b f_{2t_b}(\lambda^{b+1}) + t_b (f_{2t_b+1}(\lambda^{b+1}) + f_{\overline{2t_b+1}}(\lambda^{b+1})) + (t_b+1) f_{2t_b+2}(\lambda^{b+1}) \\ &= t_b (f_{2t_b}(\lambda^b) + 1) + t_b (f_{2t_b+1}(\lambda^b) + f_{\overline{2t_b+1}}(\lambda^b) - 1) + (t_b+1) f_{2t_b+2}(\lambda^b) \\ &= t_b f_{2t_b}(\lambda^b) + t_b (f_{2t_b+1}(\lambda^b) + f_{\overline{2t_b+1}}(\lambda^b)) + (t_b+1) f_{2t_b+2}(\lambda^b) \\ &= [\lambda_{s,1}^{b,(k-1)}/2] + [\lambda_{s,2}^{b,(k-1)}/2] + \dots + [\lambda_{s,k-1}^{b,(k-1)}/2] \\ &\equiv i - 1 + O_{\lambda^b}(\lambda_{s,1}^{b,(k-1)}) + 1 = i - 1 + O_{\lambda^b}(2t_b+2) \\ &= i - 1 + O_{\lambda^{b+1}}(2t_b+2) = i - 1 + O_{\lambda^{b+1}}(\lambda_{s,1}^{b+1,(k-1)}) \pmod{2}, \end{split}$$

which yields that $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.

(6) $\lambda_s^{b+1,(k-1)} \ge 2t_b + 4$. By the construction of λ^{b+1} , we derive that $\{\lambda_s^{b+1,(k-1)}\}_{k-1} = \{\lambda_s^{b,(k-1)}\}_{k-1}$ and $O_{\lambda^{b+1}}(\lambda_s^{b+1,(k-1)}) = O_{\lambda^b}(\lambda_s^{b,(k-1)})$. Note that $\{\lambda_s^{b,(k-1)}\}_{k-1}$ is good, So $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good.

Hence, the (k-1)-band $\{\lambda_s^{b+1,(k-1)}\}_{k-1}$ is good for $s \in [1, N_{k-1}]$. By Theorem 5.5, we see that λ^{b+1} is a special partition in $C_0(N_1, \ldots, N_{k-1}; i)$.

Finally, we set $\mu = \lambda^p$. Clearly, $|\mu| = |\lambda| - 2p + 1$, μ is a special partition in $\mathbb{C}_1(N_1, \ldots, N_{k-1}; i)$ and the largest odd part in μ is $2t_p + 1$, which is marked with r_p in $GG(\mu)$. Furthermore, if λ is a partition in $\mathbb{C}_0(N_1, \ldots, N_{k-1}; i)$, then μ belongs to $\mathbb{C}_0(N_1, \ldots, N_{k-1}; i)$. Now, we proceed to show that $2t_p + 1$ is a reduction part of μ and the reduction index of $2t_p + 1$ is p. Assume that $\hat{\alpha}_p(\lambda) = \{p_1, \ldots, p\}$, we consider the following four cases.

- (1) $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_1 . Then, we have $r_p > 2$ and $\mu_p^{(2)} = 2t_p$.
- (2) $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_2 . Then, we find that $r_p = 1$, $\mu_p^{(2)} = 2t_p + 2$ and $\check{\beta}_p(\mu) = \{p_1, \ldots, p\}$ is of cluster \check{A}_2 .
- (3) $\hat{\alpha}_p(\lambda)$ is of cluster \hat{B} . Then, we derive that $r_p = 1$, $\mu_p^{(2)} = 2t_p$ and $\check{\beta}_p(\mu) = \{p_1, \ldots, p\}$ is of cluster \check{B} .
- (4) $\hat{\alpha}_p(\lambda)$ is of cluster \hat{C} . Then, we have $r_p = 2$ and $\mu_p^{(2)} = 2t_p + 1$.

So, we conclude that $2t_p + 1$ is a reduction part in μ with reduction index p. Hence μ is a partition in $\check{\mathbb{C}}_1(N_1, \ldots, N_{k-1}; i|p)$. Moreover, if λ is a partition in $\hat{\mathbb{C}}_0(N_1, \ldots, N_{k-1}; i|p)$, then μ belongs to $\check{\mathbb{C}}_0(N_1, \ldots, N_{k-1}; i|p)$.

To prove that the reduction R_p is a bijection, we construct the inverse map of the reduction R_p , the dilation H_p . Let μ be a partition in $\check{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$. Let $\mu^{(r)} = (\mu_1^{(r)}, \mu_2^{(r)}, \ldots, \mu_{N_r}^{(r)})$ denote r-marked parts in $GG(\mu)$, where $1 \leq r \leq k-1$. Assume that the r_p -marked odd part $2t_p + 1$ in $GG(\mu)$ is the reduction part of μ . We will define $\lambda = H_p(\mu)$ related to $\mu_p^{(2)}, \mu_{p-1}^{(2)}, \ldots, \mu_1^{(2)}$ successively, and denote the intermediate special partitions by $\mu^p, \mu^{p-1}, \ldots, \mu^1, \lambda$ with $\mu^p = \mu$.

Step 1: If p = 1, then go to Step 2 directly. If p > 1, then set b = p and repeat the following process.

(E) There is a r_b -marked $2t_b + 1$ (resp. $\overline{2t_b + 1}$) in $\overline{GG}(\mu^b)$. There are the following two cases.

Case 1: $\check{\beta}_b(\mu) = \check{\beta}_{b-1}(\mu)$. In this case, we find that $2t_b + 4$ occurs in μ , which also appears μ^b . Set r_{b-1} to the largest integer such that $r_{b-1} \leq r_b$ and there is a r_{b-1} -marked $2t_b + 4$ in $\overline{GG}(\mu^b)$. Set $t_{b-1} = t_b + 2$, we see that there is a r_{b-1} -marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$. Then μ^{b-1} is obtained by replacing the r_b -marked $2t_b + 1$ (resp. $\overline{2t_b + 1}$) by $2t_b + 2$ and replacing the r_{b-1} in $\overline{GG}(\mu^b)$ by $2t_{b-1} + 1$ (resp. $\overline{2t_{b-1} + 1}$).

Case 2: $\check{\beta}_b(\mu) \neq \check{\beta}_{b-1}(\mu)$. We first replace the r_b -marked $2t_b + 1$ (resp. $\overline{2t_b + 1}$) in $\overline{GG}(\mu^b)$ by $2t_b + 2$. For $\mu_{b-1}^{(2)}$, there are five subcases.

Case 2.1: $\mu_{b-1}^{(2)}$ is of type \check{A}_1 . We may write $\mu_{b-1}^{(2)} = 2t_{b-1}$. Let r_{b-1} be the largest mark of parts $2t_{b-1}$ in $\overline{GG}(\mu^b)$. We have $r_{b-1} > 2$. Then replace the r_{b-1} -marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$ by $2t_{b-1} + 1$ to get μ^{b-1} .

Case 2.2: $\mu_{b-1}^{(2)}$ is of type \check{A}_2 . We may write $\mu_{b-1}^{(2)} = 2t_{b-1} + 2$. Then set $r_{b-1} = 1$ and replace the 1-marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$ by $2t_{b-1} + 1$ to obtain μ^{b-1} .

Case 2.3: $\mu_{b-1}^{(2)}$ is of type \check{A}_3 . We may write $\mu_{b-1}^{(2)} = 2t_{b-1}$. Then set $r_{b-1} = 2$ and replace the 2-marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$ by $\overline{2t_{b-1}+1}$ to get μ^{b-1} .

Case 2.4: $\mu_{b-1}^{(2)}$ is of type \check{B} . We may write $\mu_{b-1}^{(2)} = 2t_{b-1}$. Let r_{b-1} be the largest mark of parts $2t_{b-1}$ in $\overline{GG}(\mu^b)$. We have $r_{b-1} > 2$. Then replace the r_{b-1} -marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$ by $\overline{2t_{b-1} + 1}$ to obtain μ^{b-1} . · Case 2.5: $\mu_{b-1}^{(2)}$ is of type \check{C} . We may write $\mu_{b-1}^{(2)} = 2t_{b-1}$. Then set $r_{b-1} = 2$ and replace the 2-marked $2t_{b-1}$ in $\overline{GG}(\mu^b)$ by $2t_{b-1} + 1$ to get μ^{b-1} .

(F) Replace b by b - 1. If b = 1, then go the Step 2. If b > 1, then go back to (E).

Step 2: There is a r_1 -marked $2t_1 + 1$ (resp. $\overline{2t_1 + 1}$) in $\overline{GG}(\mu^1)$. λ is obtained by replacing the r_1 -marked $2t_1 + 1$ (resp. $\overline{2t_1 + 1}$) in $\overline{GG}(\mu^1)$ by $2t_1 + 2$.

It can be verified that the dilation H_p is the inverse map of R_p . This completes the proof.

We conclude this section with an example illustrating the reduction R_p . Let λ be the partition in (5.1). We apply the reduction R_8 to λ to get the partition μ in (5.2), that is, $\mu = R_8(\lambda)$. Here we just give the intermediate special partitions $\lambda^1, \lambda^2, \ldots, \lambda^7$, where the parts in boldface are changed parts.

$\overline{GG}(\lambda^1) = \left[\begin{array}{c}\\1\end{array}\right]$	$\begin{array}{c} 4\\2\\4\end{array}$	6	10 3	12 12	14	16 16	20 20	22	24 24	28 28	30	$\frac{34}{34}$	36	37	38].
$ \downarrow $ $\overline{GG}(\lambda^2) = \begin{bmatrix} 4 & 12 & 16 & 22 & 30 & 36 \\ 2 & 6 & 10 & 14 & 20 & 24 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 20 & 24 & 28 & 33 & 36 \end{bmatrix}. $															
$\overline{GG}(\lambda^2) = \begin{bmatrix} \\ 1 \end{bmatrix}$	$\begin{array}{c} 4\\2\\4\end{array}$	6	10 8	12 12	14	16 16	20 20	22	24 24	28 28	30	33	34	36 36	38 .
r				10		10	\downarrow	00						0.0	7
$\overline{GG}(\lambda^3) = \left[\begin{array}{c} \\ 1 \end{array}\right]$	$\begin{array}{c} 4\\2\\4\end{array}$	6	10 8	12 12	14	16 16	20 20	22	24 24	28 28	29	32	34	36 36	38].
Г	4			10		16	\downarrow	00			20			96	٦
$\overline{GG}(\lambda^4) = \left[\begin{array}{c} \\ 1 \end{array}\right]$	$\begin{array}{c} 4\\2\\4\end{array}$	6 8	10	12 12	14	16	20 20	22	23	24	28 28 28	32	34	36	38].
							\downarrow								
$\overline{GG}(\lambda^5) = \left[\begin{array}{c}\\1\end{array}\right]$	4 2 4	6	10 8	12 12	14	16 16	19	20	22 22	24	28 28 28	32	34	36 36	38].

 \downarrow

$$\overline{GG}(\lambda^6) = \begin{bmatrix} 4 & 12 & \overline{\mathbf{15}} & 22 & 28 & 36 \\ 2 & 6 & 10 & 14 & \mathbf{18} & 22 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 20 & 24 & 28 & 32 & 36 \end{bmatrix}.$$

$$\overline{GG}(\lambda^7) = \begin{bmatrix} 4 & \overline{\mathbf{11}} & \mathbf{14} & 22 & 28 & 36 \\ 2 & 6 & 10 & 14 & 18 & 22 & 28 & 34 & 38 \\ 1 & 4 & 8 & 12 & 16 & 20 & 24 & 28 & 32 & 36 \end{bmatrix}.$$

Finally, we get $\mu = \lambda^8$ in (5.2). On the contrary, the same process could be run in reverse, which illustrates the dilation $\lambda = H_8(\mu)$.

6 Proof of Theorem 3.5

Assume that $k \ge 3$, $k \ge i \ge 2$ and j = 0 or 1. The goal of this section is to give a proof of Theorem 3.5. To this end, we first establish the following theorem.

Theorem 6.1. For $N_1 \geq \cdots \geq N_{k-1} \geq 0$, $1 \leq p \leq N_2$ and $0 \leq p_1 \leq N_1$, assume that $\lambda \in \hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$, $\hat{R}_p(\lambda) = s$ and $\lambda \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|p_1, t_1)$. Let $\lambda^{(r)} = (\lambda_1^{(r)}, \lambda_2^{(r)}, \ldots, \lambda_{N_r}^{(r)})$ denote r-marked parts in $GG(\lambda)$, where $1 \leq r \leq k-1$. Set $\mu^1 = I_{p_1,t_1}(\lambda)$ and $\mu^2 = R_p(\lambda)$. Then we have

- (1) If $2t_1 + 1 < \lambda_s^{(1)} 2$, then $\mu^1 \in \hat{\mathbb{C}}_j(N_1 + 1, \dots, N_{k-1}; i|p)$.
- (2) If $2t_1+1 = \lambda_s^{(1)}-1$ and $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_1 or \hat{C} , then $\mu^1 \in \hat{\mathbb{C}}_j(N_1+1,\ldots,N_{k-1};i|p)$.
- (3) If $2t_1+1 = \lambda_s^{(1)}-1$ and $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_2 or \hat{B} , then $\mu^2 \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|p_1-1, t_1+1)$.
- (4) If $2t_1+1 > \lambda_s^{(1)}+2$, then $\mu^2 \in \mathbb{C}_j^<(N_1, \dots, N_{k-1}; i|p_1, t_1)$ or $\mu^2 \in \mathbb{C}_j^<(N_1, \dots, N_{k-1}; i|p_1, t_1)$.

Proof. Let $\mu^{1,(1)} = (\mu_1^{1,(1)}, \dots, \mu_{N_1}^{1,(1)}, \mu_{N_1+1}^{1,(1)})$ denote 1-marked parts in $GG(\mu^1)$ and let $\mu^{1,(r)} = (\mu_1^{1,(r)}, \dots, \mu_{N_r}^{1,(r)})$ denote r-marked parts in $GG(\mu^1)$, where $2 \le r \le k-1$. Let $\mu^{2,(r)} = (\mu_1^{2,(r)}, \dots, \mu_{N_r}^{2,(r)})$ denote r-marked parts in $GG(\mu^2)$ where $1 \le r \le k-1$.

(1) If $2t_1 + 1 < \lambda_s^{(1)} - 2$, then by the construction of μ^1 , we find that $p_1 \ge s$, $2t_1 + 1$ is the largest odd part in μ^1 , and $\mu_p^{1,(2)} = \lambda_p^{(2)}$ or $\lambda_p^{(2)} + 2$, and so $2t_1 + 1 < \mu_p^{1,(2)} - 2$. It implies that there does not exist a 1-marked $\mu_p^{1,(2)} - 1$ in $GG(\mu^1)$ and there do not exist odd parts greater than or equal to $\mu_p^{1,(2)}$ in μ^1 . Moreover, $\mu_p^{1,(2)}$ is of type \hat{A} or \hat{B} or \hat{C} in μ^1 . Hence, $\mu^1 \in \hat{\mathbb{C}}_j(N_1 + 1, \dots, N_{k-1}; i|p)$.

(2) If $2t_1 + 1 = \lambda_s^{(1)} - 1$ and $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_1 or \hat{C} , then by the construction of μ^1 , we derive that $p_1 = s$, $2t_1 + 1$ is the largest odd part in μ^1 and $\mu_p^{1,(2)} = \lambda_p^{(2)}$. Note that

 $2t_1 + 1 = \lambda_s^{(1)} - 1 \leq \lambda_p^{(2)} - 1 = \mu_p^{1,(2)} - 1$, we see that there do not exist odd parts greater than or equal to $\mu_p^{1,(2)}$ in μ^1 . Moreover, $\mu_p^{1,(2)}$ is of type \hat{B} (resp. \hat{A}) if $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_1 (resp. \hat{C}). Hence, $\mu^1 \in \hat{\mathbb{C}}_j(N_1 + 1, \dots, N_{k-1}; i|p)$.

(3) If $2t_1 + 1 = \lambda_s^{(1)} - 1$ and $\hat{\alpha}_p(\lambda)$ is of cluster \hat{A}_2 or \hat{B} , then we have $p_1 = s$ and $\lambda_{p_1}^{(1)} = \lambda_s^{(1)} = 2t_1 + 2$. By the construction of μ^2 , we see that the 1-marked $2t_1 + 2$ in $GG(\lambda)$ is replaced by $\mu_{p_1}^{2,(1)} = \lambda_{p_1}^{(1)} - 1 = 2t_1 + 1$, which is the largest odd part in μ^2 . Moreover, we derive that $\mu_{p_1}^{2,(1)} + 2 = 2t_1 + 3$ and $\mu_{p_1-1}^{2,(1)} \ge 2t_1 + 4 > 2t_1 + 3$. This implies that $\mu^2 \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|p_1 - 1, t_1 + 1)$.

(4) If $2t_1 + 1 > \lambda_s^{(1)} + 2$, then we have $p_1 < s$. By the construction of μ^2 , we replace the 1-marked $\lambda_s^{(1)}$ (resp. a *r*-marked $\lambda_s^{(1)} + 2$ $(r \ge 2)$) in $GG(\lambda)$ by $\lambda_s^{(1)} - 1$ (resp. $\lambda_s^{(1)} + 1$) in μ^2 . We see that the largest odd part of μ^2 is $\lambda_s^{(1)} - 1$ (resp. $\lambda_s^{(1)} + 1$), and so the largest odd part in μ^2 is less than $2t_1 + 1$. Moreover, we have $2t_1 + 1 \ge \lambda_{p_1+1}^{(1)} + 2 \ge \mu_{p_1+1}^{2,(1)} + 2$ and $\mu_{p_1}^{2,(1)} = \lambda_{p_1}^{(1)}$ or $\lambda_{p_1}^{(1)} - 2$. Note that $2t_1 + 1 < \lambda_{p_1}^{(1)}$, then we consider the following two cases.

Case 1: If $2t_1 + 1 < \mu_{p_1}^{2,(1)}$, then it is clear that $\mu^2 \in \mathbb{C}_j^<(N_1, \dots, N_{k-1}; i|p_1, t_1)$.

Case 2: If $2t_1+1 > \mu_{p_1}^{2,(1)}$, then we have $\mu_{p_1}^{2,(1)} = \lambda_{p_1}^{(1)} - 2$, and so $\lambda_{p_1}^{(1)} - 2 < 2t_1 + 1 < \lambda_{p_1}^{(1)}$. This implies that $\lambda_{p_1}^{(1)} = 2t_1 + 2$ and $\mu_{p_1}^{2,(1)} = 2t_1$. Moreover, we have $2t_1 + 3 > \mu_{p_1}^{2,(1)} + 2$ and $\mu_{p_1-1}^{2,(1)} \ge 2t_1 + 4 > 2t_1 + 3$. So, $\mu^2 \in \mathbb{C}_j^< (N_1, \dots, N_{k-1}; i | p_1 - 1, t_1 + 1)$.

Thus, we complete the proof.

Proof of Theorem 3.5. Let (λ, τ, η) be a triplet in $\mathbb{F}_j(k, i)$. We define $\pi = \Phi_{k,i}(\lambda, \tau, \eta)$ as follows. Assume that there are N_r *r*-marked parts in $GG(\lambda)$, where $1 \leq r \leq k-1$, then we have $\tau \in \mathbb{O}_{N_2}$ and $\eta \in \mathbb{I}_{N_1}$. Set $\ell_1 = \ell(\tau)$, $\ell_2 = \ell(\eta)$ and $\ell = \ell_1 + \ell_2$.

Note that there do not exist odd parts in λ , then we see that $\lambda \in \hat{\mathbb{C}}_j(N_1, \ldots, N_{k-1}; i|p)$ for $1 \leq p \leq N_2$ and $\lambda \in \mathbb{C}_j^<(N_1, \ldots, N_{k-1}; i|m)$ for $m \geq N_1$. Let $\lambda^0 = \lambda$, $\tau^0 = \tau$ and $\eta^0 = \eta$. If $\ell \geq 1$, then set b = 0 and repeat the following process until $b = \ell$.

(E) If $\tau^b \neq \emptyset$, then assume that $1 - 2p_b$ is the smallest part in τ^b , where $1 \leq p_b \leq N_2$. Otherwise, if $\tau^b = \emptyset$, then set $p_b = 0$. If $\eta^b \neq \emptyset$, then assume that $2m_b + 1$ is the smallest part in η^b , where $m_b \geq N_1 + \ell_2 - \ell(\eta^b)$. Otherwise, if $\eta^b = \emptyset$, then set $m_b = 0$. If $p_b = 0$, then apply the insertion I_{m_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = I_{m_b}(\lambda^b)$. If $m_b = 0$, then apply the reduction R_{p_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = R_{p_b}(\lambda^b)$.

If $p_b \neq 0$ and $m_b \neq 0$, then we have $\lambda^b \in \hat{\mathbb{C}}_j(N_{1_b}, \ldots, N_{k-1}; i|p_b)$ and $\lambda^b \in \mathbb{C}_j(N_{1_b}, \ldots, N_{k-1}; i|p_b)$, where $N_{1_b} = N_1 + \ell_2 - \ell(\eta^b)$. Let $\lambda^{b,(1)} = (\lambda_1^{b,(1)}, \ldots, \lambda_{N_{1_b}}^{b,(1)})$ denote 1-marked parts in $GG(\lambda^b)$ and let $\lambda^{b,(r)} = (\lambda_1^{b,(r)}, \ldots, \lambda_{N_r}^{b,(r)})$ denote r-marked parts in $GG(\lambda^b)$, where $2 \leq r \leq k-1$. Assume that p'_b is the smallest integer such that $2(m_b - p'_b) + 1 \geq \lambda^{b,(1)}_{p'_b+1} + 2$. Set $t'_b = m_b - p'_b$, then we have $\lambda^b \in \mathbb{C}^{<}_j(N_1, \ldots, N_{k-1}; i | p'_b, t'_b)$. Assume that $\hat{R}_{p_b}(\lambda^b) = s_b$, there are four cases.

- (1) If $2t'_b + 1 \leq \lambda_{s_b}^{b,(1)} 2$, then apply the insertion I_{m_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = I_{m_b}(\lambda^b)$.
- (2) If $2t'_b + 1 = \lambda_{s_b}^{b,(1)} 1$ and $\hat{\alpha}_{p_b}(\lambda^b)$ is of cluster \hat{A}_1 or \hat{C} , then apply the insertion I_{m_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = I_{m_b}(\lambda^b)$.
- (3) If $2t'_b + 1 = \lambda_{s_b}^{b,(1)} 1$ and $\hat{\alpha}_{p_b}(\lambda^b)$ is of cluster \hat{A}_2 or \hat{B} , then apply the reduction R_{p_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = R_{p_b}(\lambda^b)$.
- (4) If $2t'_b + 1 > \lambda_{s_b}^{b,(1)} + 2$, then we apply the reduction R_{p_b} to λ^b to get λ^{b+1} , that is, $\lambda^{b+1} = R_{p_b}(\lambda^b)$.

If $\lambda^{b+1} = R_{p_b}(\lambda^b)$, then set $\eta^{b+1} = \eta^b$ and remove $1 - 2p_b$ from τ^b to obtain τ^{b+1} . If $\lambda^{b+1} = I_{m_b}(\lambda^b)$, then set $\tau^{b+1} = \tau^b$ and remove $2m_b + 1$ from η^b to obtain η^{b+1} .

(F) Replace b by b + 1. If $b = \ell$, then we are done. If $b < \ell$, then go back to (E).

By Theorem 4.10 and Theorem 6.1, we see that the process above is well-defined. Finally, we set $\pi = \lambda^{\ell}$. Clearly, $\pi \in \mathbb{C}_j(N_1 + \ell_2, \ldots, N_{k-1}; i)$ and $|\pi| = |\lambda| + |\tau| + |\eta|$. It implies that $\pi \in \mathbb{C}_j(k; i)$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$.

To show that $\Phi_{k,i}$ is a bijection, we need to give the inverse map $\Psi_{k,i}$ of $\Phi_{k,i}$. Let π be a partition in $\mathbb{C}_j(k;i)$. Assume that there are $\ell \geq 0$ odd parts in π . Let $\pi^{\ell} = \pi$, $\tau^{\ell} = \emptyset$ and $\eta^{\ell} = \emptyset$. If $\ell \geq 1$, then set $b = \ell$ and repeat the following process until b = 0.

- (E) Let $\pi^{b,(r)} = (\pi_1^{b,(r)}, \ldots, \pi_{N_{r_b}}^{b,(r)})$ denote the *r*-marked parts in $GG(\pi^b)$, where $1 \le r \le k-1$. Assume that $2t'_b + 1$ is the largest odd part in π^b . There are two cases.
 - (1) If $2t'_b + 1$ is a reduction part of π^b , then assume that the reduction index of $2t'_b + 1$ is p_b . We apply the dilation H_{p_b} to π^b to get π^{b-1} , that is, $\pi^{b-1} = H_{p_b}(\pi^b)$. Then set $\eta^{b-1} = \eta^b$, and add $1 - 2p_b$ as a new part to τ^b to obtain τ^{b-1} .
 - (2) If $2t'_b + 1$ is a insertion part of π^b , then assume that $\pi^{b,(1)}_{p'_b+1} = 2t'_b + 1$. Let $m_b = p'_b + t'_b$. We apply the separation S_{m_b} to π^b to get π^{b-1} , that is, $\pi^{b-1} = S_{m_b}(\pi^b)$. Then set $\tau^{b-1} = \tau^b$ and add $2m_b + 1$ as a new part to η^b to obtain η^{b-1} .
- (F) Replace b by b 1. If b = 0, then we are done. If b > 0, then go back to (E).

Finally, we set $\lambda = \pi^0$, $\tau = \tau^0$ and $\eta = \eta^0$. Clearly, $\lambda \in \mathbb{E}_j(k, i)$, $|\pi| = |\lambda| + |\tau| + |\eta|$ and $\ell(\pi) = \ell(\lambda) + \ell(\eta)$. Moreover, we find that for $0 \le b \le \ell$, $N_{1_b} = N_{1_\ell} - \ell(\eta^b)$ and $N_{r_b} = N_{r_\ell}$, where $2 \le r \le k - 1$. Setting $N_r = N_{r_0}$, where $1 \le r \le k - 1$, then we have $\tau \in \mathbb{O}_{N_2}$. Assume that $\eta = (\eta_1, \ldots, \eta_{\ell_2})$, where $\eta_s = 1 + 2m_{b_s}$, where $1 \leq s \leq \ell_2$. For $1 \leq s < \ell_2$, we see that

$$\eta_s = 1 + 2m_{b_s} = 1 + 2p'_{b_s} + 2t'_{b_s} \ge 1 + 2p'_{b_s} + 2(p'_{b_{s+1}} + 1 - p'_{b_s}) + 2t'_{b_{s+1}} = 1 + 2(p'_{b_{s+1}} + 1) + 2t'_{b_{s+1}} > 1 + 2p'_{b_{s+1}} + 2t'_{b_{s+1}} = 1 + 2m_{b_{s+1}} = \eta_{s+1}.$$

Moreover, we have $\eta_{\ell_2} = 1 + 2m_{b_{\ell_2}} \ge 1 + 2N_{1_{b_{\ell_2}-1}} \ge 1 + 2N_{1_0} = 1 + 2N_1$. So, $\eta_1 > \eta_2 > \cdots > \eta_{\ell_2} \ge 1 + 2N_1$, which implies $\eta \in \mathbb{I}_{N_1}$. Hence, (λ, τ, η) is a triplet in $\mathbb{F}_j(k, i)$. It follows from Theorem 4.9 and Theorem 5.1 that $\Psi_{k,i}$ is the inverse map of $\Phi_{k,i}$. Thus, we complete the proof.

Acknowledgments. This work was supported by the National Science Foundation of China (Nos. 12101437 and 11901430).

References

- A.K. Agarwal, G.E. Andrews and D.M. Bressoud, The Bailey lattice, J. Indian. Math. Soc. 51 (1987) 57–73.
- G.E. Andrews, A generalization of the Göllnitz-Gordon partition theorem, Proc. Amer. Math. Soc. 18 (1967) 945–952.
- [3] G.E. Andrews, The Theory of partitions, Addison-Wesley Publishing Co., 1976.
- [4] G.E. Andrews, Multiple series Rogers-Ramanujan type identities, Pacific J. Math. 114 (1984) 267–283.
- [5] G.E. Andrews, q-Series: Their Development and Application in Analysis, Number Theory, Combinatorics, Physics, and Computer Algebra, American Mathematical Soc, 1986.
- [6] G.E. Andrews, Bailey's transform, lemma, chains and tree, In: Special Functions 2000, Kluwer Acad. Publ., Dordrecht (2001), pp. 1–22.
- [7] D.M. Bressoud, Analytic and combinatorial generalizations of Rogers-Ramanujan identities, Mem. Amer. Math. Soc. 24(227) (1980) 54pp.
- [8] D.M. Bressoud, M. Ismail and D. Stanton, Change of base in Bailey pairs, Ramanujan J. 4 (2000) 435–453.
- [9] G. Gasper and M. Rahman, Basic Hypergeometric Series, Encyclopedia of Mathematics and its Applications, Vol. 35, (Cambridge University Press, Cambridge, 1990).
- [10] H. Göllnitz, Einfache Partionen, Diplomarbeit W. S. Gottingen, 65, 1960.

- [11] H. Göllnitz, Partitionen mit differenzenbedingungen, J. reine angew. Math. 225, 1967, 154–190
- [12] B. Gordon, Some Ramanujan-like continued fractions. Abstracts of Short Communications. Int. 1962.
- [13] B. Gordon, Some continued fractions of the Rogers-Ramanujan type, Duke Math. J. 31, 1965, 741–748.
- [14] Thomas Y. He, Kathy Q. Ji, Allison Y.F. and Alice X.H. Zhao, An overpartition analogue of the Andrews-Göllnitz-Gordon theorem, J. Combin. Theory, Ser. A 166 (2019) 254–296.
- [15] K. Kurşungöz, Parity considerations in Andrews-Gordon identities, European J. Combin. 31 (2010) 976–1000.
- [16] J. Lovejoy, A Bailey lattice, Proc. Amer. Math. Soc. 132 (2004) 1507–1516.
- [17] P. Paule, A note on Bailey's lemma, J. Combin. Theory Ser. A 44 (1987) 164–167.
- [18] L.J. Slater, Further identities of Rogers-Ramanujan type, Proc. London Math. Soc. 54 (2) (1952) 147–167.
- [19] S.O. Warnaar, 50 years of Bailey's lemma, In: Algebraic combinatorics and applications, Springer, Berlin (2001) 333–347.