

THE STRUCTURE OF THE LINEARIZER OF A CONNECTED COMPLEX LIE GROUP

O. YU. ARISTOV

ABSTRACT. The Morimoto theorem states that each connected abelian complex Lie group A can be decomposed into the direct product of a group on which all holomorphic functions are constant, finitely many copies of \mathbb{C}^\times and a vector group. We prove that if A is the complex linearizer of a connected complex Lie group then the last factor of the product is trivial.

Recall that the *linearizer* of a complex Lie group G is the intersection of the kernels of all holomorphic finite-dimensional representations. We denote the linearizer of G by $\text{Lin}_{\mathbb{C}}(G)$. This notion has been known circa the 1950s, but the main attention of researchers was paid to the conditions for the *linearity* of G , i.e., to the triviality of $\text{Lin}_{\mathbb{C}}(G)$. At the same time the structure of the linearizer has been not considered earlier, to the author's knowledge. The aim of the note is to fill this gap. Addressing the problem, the author pursues his goals, since the description of the linearizer is used in questions about the homological properties of the algebras of analytical functionals [1]. However, it seems that the result of this article is of interest in its own right.

The kernel of the adjoint representation of a connected Lie group G coincides with the center of G . Therefore, the subgroup $\text{Lin}_{\mathbb{C}}(G)$ is central and, in particular, abelian. Furthermore, $\text{Lin}_{\mathbb{C}}(G)$ is connected (see Proposition 6 below). The structure of connected abelian complex Lie groups was studied by Matsushima and Morimoto in [2, 3, 4]. Namely, such a group has the following form (here \mathbb{C}^\times is the group of the invertible elements of \mathbb{C}):

$$M \times (\mathbb{C}^\times)^k \times \mathbb{C}^m \quad (k, m \in \mathbb{Z}_+), \quad (1)$$

where M is an abelian complex Lie group such that all holomorphic functions on M are constant [4, Theorem 3.2] (see [5], Lemma 15.2.12(ii), Proposition 15.3.4 and Corollary 15.3.5 as well). Note that M need not be compact (see [3] or [5, Example 15.3.10]).

By [3, Theorem 1], if G is a connected complex Lie group, then there is a smallest closed normal subgroup M such that G/M is a Stein group. In this case all holomorphic functions on M are constant. We refer to M as the *Morimoto subgroup* of G . In particular, if G is abelian then M in (1) is the Morimoto subgroup of G [4, Theorem 3.2].

Our main result states that the last factor is trivial in the decomposition (1) of $\text{Lin}_{\mathbb{C}}(G)$. More accurately, the following theorem holds.

Theorem 1. *Let G be a connected complex Lie group, and let M be the Morimoto subgroup of G . Then $\text{Lin}_{\mathbb{C}}(G)$ is isomorphic to $M \times (\mathbb{C}^\times)^k$ for some $k \in \mathbb{Z}_+$.*

From this we immediately obtain the linearizer of a Stein group (by the triviality of the Morimoto subgroup in this case).

Corollary 2. *Let G be a connected Stein group. Then $\text{Lin}_{\mathbb{C}}(G)$ is isomorphic to $(\mathbb{C}^\times)^k$ for some $k \in \mathbb{Z}_+$.*

Recall that the subgroup generated by $\exp \mathfrak{h}$ for a Lie subalgebra \mathfrak{h} of the Lie algebra \mathfrak{g} corresponding to G is said to be *integral*; and, by definition, the *solvable radical* is the integral subgroup corresponding to the solvable radical of \mathfrak{g} [5, p. 591, Definition 16.2.1]. The following assertion is used in the proof of Theorem 1.

Theorem 3. [6, Theorem 2.1] *A connected complex Lie group is linear if and only if its solvable radical is linear.*

Note that in 1943 Malcev published his version of Theorem 3 for real Lie groups (the linearity of the Levi subgroup is required additionally, but the analogous condition holds automatically in the complex case) in [7] (for example, see a proof in [8, p. 221, Chapter XVIII, Theorem 4.2]). However, to the author's knowledge, the complex case has not attracted the attention of specialists over the past decades and was only studied in [6].

First, we prove some consequence of Theorem 3.

Theorem 4. *Let G be a connected complex Lie group, and let R be the solvable radical of G . Then $\text{Lin}_{\mathbb{C}}(R) = \text{Lin}_{\mathbb{C}}(G)$.*

Proof. The inclusion $\text{Lin}_{\mathbb{C}}(R) \subset \text{Lin}_{\mathbb{C}}(G)$ is obvious. To prove the inverse inclusion we recall that $\text{Lin}_{\mathbb{C}}(G)$ is central. Hence, $\text{Lin}_{\mathbb{C}}(R)$ is central too, and so it is normal in G . We claim that $G/\text{Lin}_{\mathbb{C}}(R)$ is linear.

Indeed, denote by \mathfrak{g} , \mathfrak{r} and \mathfrak{l} the Lie algebras associated with G , R , and $\text{Lin}_{\mathbb{C}}(R)$. Since \mathfrak{l} is solvable, the radical of $\mathfrak{g}/\mathfrak{l}$ coincides with $\mathfrak{r}/\mathfrak{l}$ (for example, see [9, Lemma 4.10]). Since the radical of a Lie group is the integral subgroup corresponding to the radical of the Lie algebra; therefore, as the exponential mapping is functorial, $R/\text{Lin}_{\mathbb{C}}(R)$ is the integral subgroup generated by $\mathfrak{r}/\mathfrak{l}$, whence $R/\text{Lin}_{\mathbb{C}}(R)$ coincides with the radical of $G/\text{Lin}_{\mathbb{C}}(R)$.

The quotient group by the linearizer of a group is always linear [5, p. 578, Lemma 15.2.14]; in particular, $R/\text{Lin}_{\mathbb{C}}(R)$ is linear. Since the linearity of the radical of a connected complex Lie group implies the linearity of the initial group by Theorem 3, we infer that $G/\text{Lin}_{\mathbb{C}}(R)$ is linear. Hence, $\text{Lin}_{\mathbb{C}}(G) \subset \text{Lin}_{\mathbb{C}}(R)$. \square

Furthermore, we need two auxiliary assertions.

Lemma 5. *Let G be a complex Lie group, and let H be a closed subgroup of $\text{Lin}_{\mathbb{C}}(G)$. Then $\text{Lin}_{\mathbb{C}}(G/H) = \text{Lin}_{\mathbb{C}}(G)/H$.*

Proof. Note that $\text{Lin}_{\mathbb{C}}(G/H) \subset \text{Lin}_{\mathbb{C}}(G)/H$ because

$$(G/H)/(\text{Lin}_{\mathbb{C}}(G)/H) \cong G/\text{Lin}_{\mathbb{C}}(G)$$

and the latter group is linear.

On the other hand, assume that $g \in \text{Lin}_{\mathbb{C}}(G)$. If α is a finite-dimensional holomorphic representation of G/H , then the composition of α with $G \rightarrow G/H$ is a finite-dimensional holomorphic representation of G which sends g to 1. Thus, the kernel of α contains gH . Since $\text{Lin}_{\mathbb{C}}(G/H)$ is the intersection of the kernels of all such representations, $gH \in \text{Lin}_{\mathbb{C}}(G/H)$. Hence, $\text{Lin}_{\mathbb{C}}(G)/H \subset \text{Lin}_{\mathbb{C}}(G/H)$. \square

Proposition 6. *If a complex Lie group G is connected then so is $\text{Lin}_{\mathbb{C}}(G)$.*

Proof. Let Λ_0 be a connected component of $\text{Lin}_{\mathbb{C}}(G)$. Putting $H = \Lambda_0$ in Lemma 5, we see that $\text{Lin}_{\mathbb{C}}(G/\Lambda_0) \cong \text{Lin}_{\mathbb{C}}(G)/\Lambda_0$. Since $\text{Lin}_{\mathbb{C}}(G)/\Lambda_0$ is discrete, it suffices to prove that the linearizer of a connected Lie group is discrete if and only if it is trivial.

So, let $\text{Lin}_{\mathbb{C}}(G)$ be discrete. Recall that every solvable subgroup of a connected locally compact group is compactly generated (see [10]). In particular, this holds for abelian subgroups, and so $\text{Lin}_{\mathbb{C}}(G)$ is finitely generated since it is discrete. As it is easy to see, every finitely generated abelian group is residually finite. In particular, given a nonunit $g \in \text{Lin}_{\mathbb{C}}(G)$, there is a subgroup Λ_1 of $\text{Lin}_{\mathbb{C}}(G)$ such that $\text{Lin}_{\mathbb{C}}(G)/\Lambda_1$ is finite and $g \notin \Lambda_1$. Moreover, we may assume that $\text{Lin}_{\mathbb{C}}(G)/\Lambda_1$ is a finite cyclic group. Passing to G/Λ_1 and applying Lemma 5, we reduce the claim to the case when $\text{Lin}_{\mathbb{C}}(G)$ is finite.

Assume now that $\text{Lin}_{\mathbb{C}}(G)$ is finite. In this case, since $G/\text{Lin}_{\mathbb{C}}(G)$ is connected and linear, and $\text{Lin}_{\mathbb{C}}(G)$ is central, G is linear [5, Corollary 16.3.9]; i.e., $\text{Lin}_{\mathbb{C}}(G)$ is trivial. \square

Remark 7. The linearizer of a connected real Lie group (in the sense of real Lie groups) need not be connected. For example, it is well known that the linearizer of the universal covering group of $\text{SL}(2, \mathbb{R})$ is isomorphic to \mathbb{Z} .

Proof of Theorem 1. Notice first that the Morimoto subgroups of G and $\text{Lin}_{\mathbb{C}}(G)$ coincide. Indeed, denote them by M and M_1 , respectively. Since every holomorphic function on M is constant, so are all holomorphic representations. Hence, $M \subset \text{Lin}_{\mathbb{C}}(G)$. Since G/M is a Stein group, so is its every Lie subgroup; in particular, $\text{Lin}_{\mathbb{C}}(G/M)$. Putting $H = M$ in Lemma 5, we get $\text{Lin}_{\mathbb{C}}(G/M) = \text{Lin}_{\mathbb{C}}(G)/M$. Hence, $\text{Lin}_{\mathbb{C}}(G)/M$ is a Stein group. Using the definition of M_1 we infer that $M_1 \subset M$. On the other hand, it follows from the explicit construction of the Morimoto subgroup [3, §2] that M_1 is maximal among those Lie subgroups of $\text{Lin}_{\mathbb{C}}(G)$ that have only constant holomorphic functions on $\text{Lin}_{\mathbb{C}}(G)$. Thus, $M_1 = M$.

By Theorem 4, we may assume that G is solvable. By [4, Theorem 3.2] $\text{Lin}_{\mathbb{C}}(G)$ is of the form (1), and M coincides with the Morimoto subgroup of G , as was just shown. It remains to demonstrate that $m = 0$. Assume the contrary. Then $\text{Lin}_{\mathbb{C}}(G) \cong N \times \mathbb{C}$, where $N \cong M \times (\mathbb{C}^{\times})^k \times \mathbb{C}^{m-1}$. Since N is central, we may consider the quotient group G/N . Putting $H = N$ in Lemma 5, we get $\text{Lin}_{\mathbb{C}}(G/N) \cong \text{Lin}_{\mathbb{C}}(G)/N$. So, it suffices to consider the case when $\text{Lin}_{\mathbb{C}}(G)$ is isomorphic to \mathbb{C} .

Assume that $\text{Lin}_{\mathbb{C}}(G) \cong \mathbb{C}$. Denote $G/\text{Lin}_{\mathbb{C}}(G)$ by G_1 and the quotient homomorphism $G \rightarrow G_1$ by σ . Let K be a maximal compact subgroup of G . Put $K_1 = \sigma(K)$. Then by [5, p. 545, Theorem 14.3.13] K_1 is a maximal compact subgroup of G_1 . Furthermore, it follows from [5, p. 540, Lemma 14.3.3(4)] that a maximal compact subgroup of a solvable Lie group is abelian and, consequently, it is a torus. Thus, we have a homomorphism $\sigma': K \rightarrow K_1$ of tori.

From the exactness of the homotopy sequence of pointed bundle (for example, see [11, Lecture 5, pp. 219–220, Formula (15) and Proposition 2]) and from the contractibility of $\text{Lin}_{\mathbb{C}}(G)$ we infer that σ induces an isomorphism $\pi_1(G) \cong \pi_1(G_1)$ of the fundamental groups. It follows from the second splitting theorem [5, p. 544, Theorem 14.3.11] that G and G_1 are homotopically equivalent to K and K_1 . Hence, $\pi_1(K) \cong \pi_1(K_1)$. Since $K \cong \mathbb{T}^n$ and $K_1 \cong \mathbb{T}^{n_1}$ for some $n, n_1 \in \mathbb{Z}$, $\pi_1(K) \cong \mathbb{Z}^n$ and $\pi_1(K_1) \cong \mathbb{Z}^{n_1}$. It is easy to verify that every homomorphism $\mathbb{T}^n \rightarrow \mathbb{T}^{n_1}$ is uniquely defined by the corresponding homomorphism $\mathbb{Z}^n \rightarrow \mathbb{Z}^{n_1}$. In particular, this implies that σ' is an isomorphism.

Note that G and G_1 are Stein groups. Indeed, the Morimoto subgroup lies in the linearizer. In our case, it is either trivial or isomorphic to \mathbb{C} . By the connectedness of Morimoto subgroups [3, Lemma 4], we infer that they are trivial for both groups, as required.

Denote by K^* and K_1^* the smallest complex integral subgroups containing K and K_1 , respectively. These subgroups are closed [5, Corollary 14.5.6]; and since G and G_1 are Stein groups, K^* and K_1^* have no compact factors. Consequently, these are the universal complexifications of K and K_1 [5, Proposition 15.3.4 (i)]. Since universal complexification is a functor, σ maps K^* onto K_1^* isomorphically.

Since $G/\text{Lin}_{\mathbb{C}}(G)$ is connected and linear, there is a decomposition into a semidirect product: $G/\text{Lin}_{\mathbb{C}}(G) = B_1 \rtimes K_1^*$, where B_1 is simply connected and solvable [5, p.601, Theorem 16.3.7]. Put $B := \sigma^{-1}(B_1)$. Obviously, $\text{Lin}_{\mathbb{C}}(G) \subset B$, and $\text{Lin}_{\mathbb{C}}(G)$ coincides with the kernel of the restriction of σ to B . Thus, $B/\text{Lin}_{\mathbb{C}}(G) \cong B_1$; and, consequently, B is also simply connected and solvable (since $\text{Lin}_{\mathbb{C}}(G) \cong \mathbb{C}$).

We claim that $G = B \rtimes K^*$. Indeed, firstly $\sigma(g) = b_1 l_1$ for every $g \in G$, where $b_1 \in B_1$ and $l_1 \in K_1^*$. Let $l \in K^*$ be the preimage of l_1 . Then $g = (gl^{-1})l$, and $gl^{-1} \in B$ since $\sigma(gl^{-1}) = b_1$. Thus $G = BK^*$. Secondly, if $g \in B \cap K^*$, then $\sigma(g) \in B_1 \cap K_1^* = \{1\}$, whence $g = 1$ since $K^* \rightarrow K_1^*$ is an isomorphism. Thus, $B \cap K^* = \{1\}$. So $G = B \rtimes K^*$ with B simply connected and solvable, and K is a maximal compact subgroup. Applying in the inverse direction Theorem 16.3.7 from [5], we infer that G is linear. The so-obtained contradiction with the nontriviality of the linearizer completes the proof of Theorem 1. \square

Acknowledgment. The author thanks the referee for the useful comments that improved the presentation.

REFERENCES

- [1] Aristov O.Yu., *Analytic functionals on complex Lie groups, smash products, and resolutions*, Preprint, 2022 (in Russian).
- [2] Matsushima Y., Morimoto A., *Sur certains espaces fibrés holomorphes sur une variété de Stein*, Bull. Soc. Math. France 88 (1960), 137–155.
- [3] Morimoto A., *Non-compact complex Lie groups without non-constant holomorphic functions*, Proc. of the Conf. on Complex Analysis, Minneapolis 1964, Springer, 1965, 256–272.
- [4] Morimoto A., *On the classification of non-compact abelian Lie groups*, Trans. Amer. Math. Soc. 123 (1966), 200–228.
- [5] Hilgert J., Neeb K.-H., *Structure and geometry of Lie groups*, Springer, 2011.
- [6] Aristov O.Yu., *Holomorphic functions of exponential type on connected complex Lie groups*, J. Lie Theory 29:4 (2019), 1045–1070, arXiv:1903.08080.
- [7] Malcev A.I., *On arcwise connected locally closed groups*, Dokl. Akad. Nauk SSSR, 40:3 (1943), 108–110.
- [8] Hochschild G., *The structure of Lie groups*, Holden Day, San Francisco, 1965.
- [9] Aristov O.Yu., *Functions of class C^∞ in non-commuting variables in the context of triangular Lie algebras*, Izv. RAN. Ser. Mat., 86:6 (2022), 5–46, arXiv:2103.06143.
- [10] Hofmann K. H., Neeb K.-H., *The compact generation of closed subgroups of locally compact groups*, J. Group Theory 12 (2009), 555–559.
- [11] Postnikov M. M., *Lectures on algebraic topology. Homotopy theory basics*, Nauka, Moscow, 1984 (in Russian).

Email address: aristovoyu@inbox.ru