

On large partial ovoids of symplectic and Hermitian polar spaces

Michela Ceria Jan De Beule Francesco Pavese Valentino Smaldore

Abstract

In this paper we provide constructive lower bounds on the sizes of the largest partial ovoids of the symplectic polar spaces $\mathcal{W}(3, q)$, q odd square, $q \not\equiv 0 \pmod{3}$, $\mathcal{W}(5, q)$ and of the Hermitian polar spaces $\mathcal{H}(4, q^2)$, q even or q odd square, $q \not\equiv 0 \pmod{3}$, $\mathcal{H}(6, q^2)$, $\mathcal{H}(8, q^2)$.

Keywords: Partial ovoid, symplectic polar space; Hermitian polar space.

1 Introduction

Let \mathcal{P} be a finite classical polar space, i.e., \mathcal{P} arises from a vector space of finite dimension over a finite field equipped with a non-degenerate reflexive sesquilinear or quadratic form. A projective subspace of maximal dimension contained in \mathcal{P} is called a *generator* of \mathcal{P} . A (*partial*) *ovoid* \mathcal{O} of a polar space \mathcal{P} is a set of points of \mathcal{P} such that every generator of \mathcal{P} contains (at most) one point of \mathcal{O} . A partial ovoid is said to be *maximal* if it is maximal with respect to set-theoretic inclusion.

Here we focus on the symplectic polar space $\mathcal{W}(2n - 1, q)$ and the Hermitian polar space $\mathcal{H}(2n, q^2)$, consisting of the absolute projective subspaces with respect to a non-degenerate symplectic polarity of $\text{PG}(2n - 1, q)$ and a non-degenerate unitary polarity of $\text{PG}(2n, q^2)$, respectively. It is well known that $\mathcal{W}(2n - 1, q)$ has ovoids if and only if $n = 2$ and q is even, whereas $\mathcal{H}(2n, q^2)$ does not possess ovoids. Hence the question of the largest (maximal) partial ovoids of $\mathcal{W}(2n - 1, q)$, $(n, q) \neq (2, 2^h)$, and $\mathcal{H}(2n, q^2)$ naturally arises. Constructions of maximal partial ovoids of $\mathcal{W}(3, q)$, q even, of size an integer between about $q^2/10$ and $9q^2/10$ or of size $q^2 - hq + 1$, $1 \leq h \leq q/2$, have been provided in [12, 13]. The situation is somewhat different for q odd where the lack of examples is transparent. In [13] Tallini proved that a partial ovoid of $\mathcal{W}(3, q)$, q odd, has size at most $q^2 - q + 1$ and constructed a maximal partial ovoid of $\mathcal{W}(3, q)$ of size $2q + 1$. Regarding symplectic polar spaces in higher dimensions an upper bound on the size of

M. Ceria, F. Pavese: Department of Mechanics, Mathematics and Management, Polytechnic University of Bari, Via Orabona 4, 70125 Bari, Italy; *e-mail*: {michela.ceria, francesco.pavese}@poliba.it

J. De Beule: Department of Mathematics and Data Science, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussel, Belgium and Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Krijgslaan 281 (S8), 9000 Gent, Belgium; *e-mail*: jan@debeule.eu, <https://orcid.org/0000-0001-5333-5224>

V. Smaldore: Department of Mathematics, Computer Science and Economics, University of Basilicata, Contrada Macchia Romana, 85100, Potenza, Italy; *e-mail*: valentino.smaldore@unibas.it

Mathematics Subject Classification (2020): Primary 51E20. Secondary 05B25.

the largest partial ovoid has been provided in [8] and, if q is even, a partial ovoid of an elliptic or hyperbolic quadric is also a partial ovoid of a symplectic polar space.

As for $\mathcal{H}(2n, q^2)$, an upper bound on the size of the largest partial ovoid can be found in [7]. In particular, a partial ovoid has at most $q^5 - q^4 + q^3 + 1$ points if $n = 2$. The largest known example of a maximal partial ovoid of $\mathcal{H}(2n, q^2)$, $n = 2, 3$, occurs when $q = 3^h$ and has size $q^4 + 1$ [11]. A straightforward check shows that a non-degenerate plane section of $\mathcal{H}(2n, q^2)$ is an example of maximal partial ovoid of $\mathcal{H}(2n, q^2)$ of size $q^3 + 1$. Other examples of maximal partial ovoids of $\mathcal{H}(4, q^2)$ of size $2q^3 + q^2 + 1$ have been constructed in [4] and of size $q^3 + 1$ in [5, 11].

In Section 3, partial ovoids of symplectic polar spaces are studied. We show the existence of a partial ovoid of $\mathcal{W}(3, q)$, q an odd square, $q \not\equiv 0 \pmod{3}$, of size $(q^{3/2} + 3q - q^{1/2} + 3)/3$. It is obtained by glueing together a twisted cubic \mathcal{C} of $\text{PG}(3, q)$ and an orbit of a subgroup of $\text{PSp}(4, q)$ isomorphic to $\text{PGL}(2, q^{1/2})$ stabilizing \mathcal{C} . Next maximal partial ovoids of $\mathcal{W}(5, q)$ of size $q^2 + q + 1$ and of size $2q^2 - q + 1$, if q is even, are exhibited.

In Section 4 we introduce the notion of *tangent-set* of a Hermitian variety $\mathcal{H}(2n - 1, q^2)$, that is a set \mathcal{T} of points of $\text{PG}(2n - 1, q^2)$ such that every line that is either tangent or contained in $\mathcal{H}(2n - 1, q^2)$ has at most one point in common with \mathcal{T} . In particular we show that starting from a partial ovoid of $\mathcal{W}(2n - 1, q)$ of size x , it is possible to obtain a tangent-set of $\mathcal{H}(2n - 1, q^2)$ of size roughly xq , which in turn gives rise to a partial ovoid of $\mathcal{H}(2n, q^2)$ of size roughly xq^2 . Applying this construction technique to the known (partial) ovoids of $\mathcal{W}(2n - 1, q)$, $n \in \{2, 3, 4\}$, the following are obtained: maximal partial ovoids of $\mathcal{H}(4, q^2)$, q even, of size $q^4 + 1$ and of size $q^4 - q^3 + q + 1$, a partial ovoid of $\mathcal{H}(4, q^2)$, q an odd square, $q \not\equiv 0 \pmod{3}$, of size $(q^{7/2} + 3q^3 - q^{5/2} + 3q^2)/3$, partial ovoids of $\mathcal{H}(6, q^2)$ of size $q^4 + q^3 + 1$ and of size $2q^4 - q^3 + 1$, if q is even, and a partial ovoid of $\mathcal{H}(8, q^2)$ of size $q^5 + 1$. Table 1 and Table 2 summarize old and new results regarding large partial ovoids of symplectic and Hermitian polar spaces in small dimensions.

	lower bound	upper bound
$\mathcal{W}(3, q)$, q even	$q^2 + 1$	$q^2 + 1$
$\mathcal{W}(3, q)$, q odd	$(q^{3/2} + 3q - q^{1/2} + 3)/3$, $q = p^{2h}$, $p \neq 3$, $2q + 1$, $q = p^{2h+1}$ or $q = 3^h$ [13]	$q^2 - q + 1$ [13]
$\mathcal{W}(5, q)$	$2q^2 - q + 1$, q , even $q^2 + q + 1$, q odd 7 , $q = 2$	$\frac{q\sqrt{5q^4+6q^3+7q^2+6q+1}-q^3-q^2-q+2}{2}$ [8] 7 , $q = 2$
$\mathcal{W}(7, q)$	$q^3 + 1$ [2, 3]	$q^4 - q^3 - q(\sqrt{q} - 1)(q - \sqrt{q} + 1) + 3$, $q > 2$ [8] 9 , $q = 2$

Table 1: Large partial ovoids of $\mathcal{W}(2n - 1, q)$, $n \in \{2, 3, 4\}$.

	lower bound	upper bound
	$q^4 + 1, q = 2^h \text{ or } q = 3^h$ [11]	
$\mathcal{H}(4, q^2)$	$(q^{7/2} + 3q^3 - q^{5/2} + 3q^2)/3,$ $q = p^{2h}, p \text{ odd}, p \neq 3$	$q^5 - q^4 + q^3 + 1$ [7]
	$2q^3 + q^2 + 1,$ $q = p^{2h+1}, p \neq 2, 3$ [4]	
$\mathcal{H}(6, q^2)$	$2q^4 - q^3 + 1, q, \text{ even}$	$q^7 - q^6 + q^5 - q^3 + 2$ [7]
	$q^4 + q^3 + 1, q \text{ odd}$	
$\mathcal{H}(8, q^2)$	$q^5 + 1$	$q^9 - q^8 + q^7 - q^5 - q^3 + q^2 + 1$ [7]

Table 2: Large partial ovoids of $\mathcal{H}(2n, q^2)$, $n \in \{2, 3, 4\}$.

2 Preliminaries

Let q be a prime power and let \mathbb{F}_q be the finite field of order q . Let $\text{PG}(n, q)$ be the n -dimensional projective space over \mathbb{F}_q equipped with homogeneous projective coordinates $(X_1, X_2, \dots, X_{n+1})$. Denote by U_i the point having 1 in the i -th position and 0 elsewhere. A non-degenerate symplectic or unitary polarity of $\text{PG}(n, q^2)$ is induced by a non-degenerate alternating or Hermitian form on the underlying vector space.

The symplectic polar space $\mathcal{W}(2n - 1, q)$ is formed by the projective subspaces that are absolute with respect to a non-degenerate symplectic polarity \mathfrak{s} of $\text{PG}(2n - 1, q)$. It is left invariant by the group of projectivities $\text{PSp}(4, q)$. It consists of all the points of $\text{PG}(2n - 1, q)$ and of $(q + 1) \dots (q^n + 1)$ generators. Through every point P of $\text{PG}(2n - 1, q)$ there pass $(q + 1) \dots (q^{n-1} + 1)$ generators and these generators all lie in a hyperplane. The hyperplane containing these generators is the polar hyperplane $P^{\mathfrak{s}}$ of P with respect to the symplectic polarity \mathfrak{s} defining $\mathcal{W}(2n - 1, q)$.

A (non-degenerate) Hermitian variety $\mathcal{H}(n, q^2)$ consists of the absolute points of a unitary polarity of $\text{PG}(n, q^2)$. It has $(q^{n+1} + (-1)^n)(q^n - (-1)^n) / (q^2 - 1)$ points. Subspaces of maximal dimension contained in $\mathcal{H}(n, q^2)$ are $\lfloor \frac{n-1}{2} \rfloor$ -dimensional projective spaces and are called *generators*. There are either $(q + 1)(q^3 + 1) \dots (q^n + 1)$ or $(q^3 + 1)(q^5 + 1) \dots (q^{n+1} + 1)$ generators in $\mathcal{H}(n, q^2)$, depending on whether n is odd or even, respectively. A line of $\text{PG}(n, q^2)$ meets $\mathcal{H}(n, q^2)$ in 1, $q + 1$ or, if $n \geq 3$, in $q^2 + 1$ points. The latter lines are the generators of $\mathcal{H}(n, q^2)$ if $n \in \{3, 4\}$; lines meeting $\mathcal{H}(n, q^2)$ in one or $q + 1$ points are called *tangent lines* or *secant lines*, respectively. Through a point P of $\mathcal{H}(n, q^2)$ there pass $(q^{n-1} + (-1)^{n-2})(q^{n-2} - (-1)^{n-2}) / (q^2 - 1)$ generators and these generators are contained in a hyperplane. The hyperplane containing these generators is the polar hyperplane of P with respect to the unitary polarity of $\text{PG}(n, q^2)$ defining $\mathcal{H}(n, q^2)$ and it is also called the *tangent hyperplane* to $\mathcal{H}(n, q^2)$ at P . The tangent lines through P are precisely the remaining lines of its polar hyperplane that are incident with P . If $P \notin \mathcal{H}(n, q^2)$ then the polar hyperplane of P is a hyperplane of $\text{PG}(n, q^2)$ meeting $\mathcal{H}(n, q^2)$ in a non-degenerate Hermitian variety $\mathcal{H}(n - 1, q^2)$ and it is said to be *secant* to $\mathcal{H}(n, q^2)$. The

stabilizer of $\mathcal{H}(n, q^2)$ in $\text{PGL}(n+1, q^2)$ is the group $\text{PGU}(n+1, q^2)$. Recall that the symplectic polar space $\mathcal{W}(2n-1, q)$ can be embedded in $\mathcal{H}(2n-1, q^2)$.

3 Partial ovoids of symplectic polar spaces

3.1 $\mathcal{W}(3, q)$, q odd square, $q \not\equiv 0 \pmod{3}$

Let $\mathcal{W}(3, q)$, $q \not\equiv 0 \pmod{3}$, be the symplectic polar space consisting of the subspaces of $\text{PG}(3, q)$ induced by the totally isotropic subspaces of \mathbb{F}_q^4 with respect to the non-degenerate alternating form β given by

$$x_1y_4 + x_2y_3 - x_3y_2 - x_4y_1.$$

Denote by \mathfrak{s} the symplectic polarity of $\text{PG}(3, q)$ defining $\mathcal{W}(3, q)$. Let \mathcal{C} be the twisted cubic of $\text{PG}(3, q)$ consisting of the $q+1$ points $\{P_t \mid t \in \mathbb{F}_q\} \cup \{(0, 0, 0, 1)\}$, where $P_t = (1, -3t, t^2, t^3)$. It is well known that a line of $\text{PG}(3, q)$ meets \mathcal{C} in at most 2 points and a plane shares with \mathcal{C} at most 3 points (i.e., \mathcal{C} is a so called $(q+1)$ -arc). A line of $\text{PG}(3, q)$ joining two distinct points of \mathcal{C} is called a *real chord* and there are $q(q+1)/2$ of them. Let $\bar{\mathcal{C}} = \{P_t \mid t \in \mathbb{F}_{q^2}\} \cup \{(0, 0, 0, 1)\}$ be the twisted cubic of $\text{PG}(3, q^2)$ which extends \mathcal{C} over \mathbb{F}_{q^2} . The line of $\text{PG}(3, q^2)$ obtained by joining P_t and P_{t^q} , with $t \notin \mathbb{F}_q$, meets the canonical Baer subgeometry $\text{PG}(3, q)$ in the $q+1$ points of a line skew to \mathcal{C} . Such a line is called an *imaginary chord* and they are $q(q-1)/2$ in number. If r is a (real or imaginary) chord, then the line r^s is called (*real or imaginary*) *axis*. Also, for each point P of \mathcal{C} , the line $\ell_P = \langle P, P' \rangle$, where P' equals $(0, -3, 2t, 3t^2)$ or U_3 if $P = P_t$ or $P = U_4$, respectively, is called the *tangent* line to \mathcal{C} at P . With each point P_t (resp. U_4) of \mathcal{C} there corresponds the *osculating plane* P_t^s (resp. U_4^s) with equation $t^3X_1 + t^2X_2 + 3tX_3 - X_4 = 0$ (resp. $X_1 = 0$), meeting \mathcal{C} only at P_t (resp. U_4) and containing the tangent line. Hence the $q+1$ lines tangent to \mathcal{C} are generators of $\mathcal{W}(3, q)$ and they form a regulus \mathcal{R} if q even. Every point of $\text{PG}(3, q) \setminus \mathcal{C}$ lies on exactly one chord or a tangent of \mathcal{C} . For more properties and results on \mathcal{C} the reader is referred to [9, Chapter 21]. Let G be the group of projectivities of $\text{PG}(3, q)$ stabilizing \mathcal{C} . Then $G \simeq \text{PGL}(2, q)$ whenever $q \geq 5$, and elements of G are induced by the matrices

$$M_{a,b,c,d} = \begin{pmatrix} a^3 & -a^2b & 3ab^2 & b^3 \\ -3a^2c & a^2d + 2abc & -3b^2c - 6abd & -3b^2d \\ ac^2 & \frac{-bc^2 - 2acd}{3} & ad^2 + 2bcd & bd^2 \\ c^3 & -c^2d & 3cd^2 & d^3 \end{pmatrix},$$

where $a, b, c, d \in \mathbb{F}_q$, $ad - bc \neq 0$. The group G leaves invariant $\mathcal{W}(3, q)$ since $M_{a,b,c,d}^t J M_{a,b,c,d} = (ad - bc)^3 J$, where

$$J = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ -1 & 0 & 0 & 0 \end{pmatrix}$$

is the Gram matrix of β .

Assume q to be an odd square. Here we show the existence of a partial ovoid of $\mathcal{W}(3, q)$ obtained by glueing together the twisted cubic \mathcal{C} of $\text{PG}(3, q)$ and an orbit of size $\sqrt{q}(q-1)/3$ of a subgroup G_ϵ of $\text{PSp}(4, q) \cap G$ isomorphic to $\text{PGL}(2, \sqrt{q})$ stabilizing \mathcal{C} . In particular such a subgroup fixes a twisted cubic $\mathcal{C}_\epsilon = \mathcal{C} \cap \Sigma_\epsilon$ of a Baer subgeometry Λ_ϵ of $\text{PG}(3, q)$.

$$\boxed{\sqrt{q} \equiv 1 \pmod{3}}$$

Assume $\sqrt{q} \equiv 1 \pmod{3}$. Let $\mathcal{C}_1 = \{P_t \mid t \in \mathbb{F}_{\sqrt{q}}\} \cup \{U_4\}$. Thus $\mathcal{C}_1 \subset \mathcal{C}$ is a twisted cubic of the canonical Baer subgeometry $\Lambda_1 = \text{PG}(3, \sqrt{q})$ of $\text{PG}(3, q)$. The group G_1 of projectivities stabilizing \mathcal{C}_1 is isomorphic to $\text{PGL}(2, \sqrt{q})$ and it is induced by the matrices

$$M_{a,b,c,d}, a, b, c, d \in \mathbb{F}_{\sqrt{q}}, ad - bc \neq 0.$$

Let $R = U_1 + xU_4$, for a fixed $x \in \mathbb{F}_q \setminus \mathbb{F}_{\sqrt{q}}$ where x is not a cube in \mathbb{F}_q . Set

$$\mathcal{O}_1 = R^{G_1} = \{(a^3 + xb^3, -3a^2c - 3xb^2d, ac^2 + xbd^2, c^3 + xd^3) \mid a, b, c, d \in \mathbb{F}_{\sqrt{q}}, ad - bc \neq 0\}.$$

Lemma 3.1. *The set \mathcal{O}_1 is a partial ovoid of $\mathcal{W}(3, q)$ of size $\sqrt{q}(q-1)/3$.*

Proof. Since the stabilizer of R in G_1 has order 3 and $|G_1| = \sqrt{q}(q-1)$, by applying the Orbit-Stabilizer Theorem it follows that \mathcal{O}_1 has the required size. In order to show that \mathcal{O}_1 is a partial ovoid of $\mathcal{W}(3, q)$, it is enough to see that the line joining R and a further point R^g of \mathcal{O}_1 is not a generator of $\mathcal{W}(3, q)$. Here $g \in G_1$ is induced by $M_{a,b,c,d}$. Assume by contradiction that this is not the case, then there are $a, b, c, d \in \mathbb{F}_{\sqrt{q}}$, with $ad - bc \neq 0$, such that $A(x) = 0$, where

$$A(x) = c^3 + xd^3 - xa^3 - x^2b^3.$$

Hence $A(x) + A(x)\sqrt{q} = A(x) - A(x)\sqrt{q} = 0$, that is

$$\begin{aligned} 2c^3 + (d^3 - a^3)(x + x\sqrt{q}) - (x^2 + x^2\sqrt{q})b^3 &= 0, \\ d^3 - a^3 - (x + x\sqrt{q})b^3 &= 0. \end{aligned}$$

If $b = 0$, then $a^3 = d^3$ and $c = 0$. Thus g fixes R , i.e., $R = R^g$. If $b \neq 0$, then the previous equations imply

$$x\sqrt{q}+1 = -\frac{c^3}{b^3},$$

that is a contradiction since $x\sqrt{q}+1$ is not a cube in $\mathbb{F}_{\sqrt{q}}$. Indeed x is not a cube in \mathbb{F}_q and $(\sqrt{q}+1, 3) = 1$. \square

$$\boxed{\sqrt{q} \equiv -1 \pmod{3}}$$

Assume $\sqrt{q} \equiv -1 \pmod{3}$. Let $\mathcal{C}_{-1} = \{P_t \mid t \in \mathbb{F}_q, t\sqrt{q}+1 = 1\}$. Thus $\mathcal{C}_{-1} \subset \mathcal{C}$ is a twisted cubic of the Baer subgeometry

$$\Lambda_{-1} = \{(\alpha, -3\beta, \beta^q, \alpha^q) \mid \alpha, \beta \in \mathbb{F}_{q^2}, (\alpha, \beta) \neq (0, 0)\} \simeq \text{PG}(3, \sqrt{q})$$

of $\text{PG}(3, q)$. In this case the group G_{-1} of projectivities stabilizing \mathcal{C}_{-1} is isomorphic to $\text{PGL}(2, \sqrt{q})$ and it is induced by the matrices

$$M_{a,b,c,d}, a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0, ab\sqrt{q} - cd\sqrt{q} = 0, a\sqrt{q}+1 + b\sqrt{q}+1 - c\sqrt{q}+1 - d\sqrt{q}+1 = 0.$$

Let $S = U_1 + xU_4$, for a fixed $x \in \mathbb{F}_q \setminus \mathbb{F}_{\sqrt{q}}$ where x is not a cube in \mathbb{F}_q and $x^{\sqrt{q}+1} \neq 1$. Set

$$\mathcal{O}_{-1} = S^{G_{-1}} = \{(a^3 + xb^3, -3a^2c - 3xb^2d, ac^2 + xbd^2, c^3 + xd^3) \mid a, b, c, d \in \mathbb{F}_q, ad - bc \neq 0, \\ ab\sqrt{q} - cd\sqrt{q} = 0, a\sqrt{q}+1 + b\sqrt{q}+1 - c\sqrt{q}+1 - d\sqrt{q}+1 = 0\}.$$

Lemma 3.2. *The set \mathcal{O}_{-1} is a partial ovoid of $\mathcal{W}(3, q)$ of size $\sqrt{q}(q-1)/3$.*

Proof. Since the stabilizer of S in G_{-1} has order 3 and $|G_{-1}| = \sqrt{q}(q-1)$, by applying the Orbit-Stabilizer Theorem it follows that \mathcal{O}_{-1} has the required size. In order to show that \mathcal{O}_{-1} is a partial ovoid of $\mathcal{W}(3, q)$, it is enough to see that the line joining R and a further point S^g of \mathcal{O}_{-1} is not a generator of $\mathcal{W}(3, q)$. Here $g \in G_{-1}$ is induced by $M_{a,b,c,d}$. Assume by contradiction that this is not the case, then there are $a, b, c, d \in \mathbb{F}_q$, with $ad - bc \neq 0$, $ab\sqrt{q} - cd\sqrt{q} = 0$, $a\sqrt{q}+1 + b\sqrt{q}+1 - c\sqrt{q}+1 - d\sqrt{q}+1 = 0$, such that $A(x) = 0$, where

$$A(x) = c^3 + xd^3 - xa^3 - x^2b^3.$$

If $b = 0$, then $c = 0$, since $cd\sqrt{q} = 0$ and $ad \neq 0$. Moreover $a\sqrt{q}+1 = d\sqrt{q}+1$ and $a^3 = d^3$, since $A(x) = 0$. Thus g fixes S , i.e., $S = S^g$. If $b \neq 0$, we may assume w. l. o. g. that $b = 1$. Then $a = cd\sqrt{q}$, $c(d\sqrt{q}+1 - 1) \neq 0$ and $(1 - c\sqrt{q}+1)(1 - d\sqrt{q}+1) = 0$. Therefore $c\sqrt{q}+1 = 1$ and $d\sqrt{q}+1 \neq 1$. Moreover $A(x) = 0$ gives

$$d^3\sqrt{q} - \frac{d^3}{c^3} - \frac{c^3 - x^2}{c^3x} = 0.$$

By considering the last equation in the unknown d^3 , by [10, Theorem 1.9.3], it admits solutions in \mathbb{F}_q if and only

$$0 = \frac{c^3\sqrt{q} - x^2\sqrt{q}}{c^3\sqrt{q}x\sqrt{q}} + \frac{c^3 - x^2}{c^3x} \frac{1}{c^3\sqrt{q}} = \frac{x(1 - x\sqrt{q}+1)(c^3\sqrt{q} + x\sqrt{q}-1)}{c^3\sqrt{q}x\sqrt{q}+1}.$$

Hence either $x = 0$ or $x\sqrt{q}+1 = 1$ or $x\sqrt{q}-1$ is a cube in \mathbb{F}_q . If the last possibility occurs then x is a cube in \mathbb{F}_q since $\sqrt{q} \equiv -1 \pmod{3}$. We infer that none of the three cases arises. \square

Theorem 3.3. *Let q be an odd square with $\sqrt{q} \equiv \epsilon \pmod{3}$, $\epsilon \in \{\pm 1\}$. Then the set $\mathcal{O}_\epsilon \cup \mathcal{C}$ is a partial ovoid of $\mathcal{W}(3, q)$, of size $(q^{3/2} + 3q - q^{1/2} + 3)/3$.*

Proof. Since in both cases x is not a cube in \mathbb{F}_q , it follows that there is no line of $\mathcal{W}(3, q)$ through R or S meeting \mathcal{C} . \square

Remark 3.4. The partial ovoid $\mathcal{O}_\epsilon \cup \mathcal{C}$ of $\mathcal{W}(3, q)$ is not maximal. Indeed if N is a point of Λ_ϵ not lying on an osculating plane of \mathcal{C}_ϵ , then $\mathcal{O}_\epsilon \cup \mathcal{C} \cup \{N\}$ is a partial ovoid of $\mathcal{W}(3, q)$. In order to see this fact let ℓ be a line of $\mathcal{W}(3, q)$. If ℓ passes through a point of \mathcal{C}_ϵ then it lies in an osculating plane of \mathcal{C}_ϵ , whereas if ℓ contains a point of $\mathcal{C} \setminus \Lambda_\epsilon$, then $\ell \cap \Lambda_\epsilon$ is a point lying on an imaginary axis of \mathcal{C}_ϵ . Similarly, since the points of \mathcal{O}_ϵ are on extended chords of \mathcal{C}_ϵ , if ℓ contains a point of \mathcal{O}_ϵ , then $\ell \cap \Lambda_\epsilon$ is a point lying on an axis of \mathcal{C}_ϵ . On the other hand every axis of \mathcal{C}_ϵ is contained in an osculating plane of \mathcal{C}_ϵ .

Problem 3.5. Find a maximal partial ovoid of $\mathcal{W}(3, q)$ containing $\mathcal{O}_\epsilon \cup \mathcal{C}$.

3.2 Maximal partial ovoids of $\mathcal{W}(5, q)$ of size $q^2 + q + 1$

Here we consider a symplectic polar space in $\text{PG}(5, q)$, where $\text{PG}(5, q)$ is embedded as a subgeometry in $\text{PG}(5, q^3)$. We denote by $N : x \in \mathbb{F}_{q^3} \mapsto x^{q^2+q+1} \in \mathbb{F}_q$ the *norm function* of \mathbb{F}_{q^3} over \mathbb{F}_q and by $T : x \mapsto x + x^q + x^{q^2}$ the *trace function* of \mathbb{F}_{q^3} over \mathbb{F}_q . Let V be the 6-dimensional \mathbb{F}_q -vector subspace of $\mathbb{F}_{q^3}^6$ given by

$$\{P_{a,b} = (a, a^q, a^{q^2}, b^{q^2}, b^q, b) \mid a, b \in \mathbb{F}_{q^3}\}.$$

Then $\text{PG}(V)$ is a q -order subgeometry of $\text{PG}(5, q^3)$. If $\mathcal{W}(5, q^3)$ is the symplectic polar space consisting of the subspaces of $\text{PG}(5, q^3)$ induced by the totally isotropic subspaces of $\mathbb{F}_{q^3}^6$ with respect to the non-degenerate alternating form given by

$$x_1y_6 + x_2y_5 + x_3y_4 - x_4y_3 - x_5y_2 - x_6y_1,$$

then $\mathcal{W}(5, q^3)$ induces in $\text{PG}(V)$ a symplectic polar space, say $\mathcal{W}(5, q)$. It is straightforward to check that the cyclic group of order $q^2 + q + 1$ formed by the projectivities of $\text{PG}(V)$ induced by the matrices

$$D_x = \begin{pmatrix} x & 0 & 0 & 0 & 0 & 0 \\ 0 & x^q & 0 & 0 & 0 & 0 \\ 0 & 0 & x^{q^2} & 0 & 0 & 0 \\ 0 & 0 & 0 & x^{-q^2} & 0 & 0 \\ 0 & 0 & 0 & 0 & x^{-q} & 0 \\ 0 & 0 & 0 & 0 & 0 & x^{-1} \end{pmatrix}, x \in \mathbb{F}_{q^3}, N(x) = 1,$$

preserves $\mathcal{W}(5, q)$. The group K fixes the two planes

$$\pi_1 = \{P_{a,0} \mid a \in \mathbb{F}_{q^3} \setminus \{0\}\}, \quad \pi_2 = \{P_{0,b} \mid b \in \mathbb{F}_{q^3} \setminus \{0\}\}.$$

Lemma 3.6. *There are one or three K -orbits on points of π_i , $i = 1, 2$, according as $q \not\equiv 1 \pmod{3}$ or $q \equiv 1 \pmod{3}$. In the latter case their representatives are as follows*

$$\pi_1 : P_{z^i,0}, \pi_2 : P_{0,z^i}, i = 1, 2, 3,$$

for some $z \in \mathbb{F}_{q^3}$ such that $z^{q-1} = \xi$, where ξ is a fixed element in \mathbb{F}_q such that $\xi^2 + \xi + 1 = 0$.

The group K has $q^3 - 1$ orbits on points of $\text{PG}(V) \setminus (\pi_1 \cup \pi_2)$. Each of them has size $q^2 + q + 1$ and their representatives are the points

$$P_{1,b}, b \in \mathbb{F}_{q^3} \setminus \{0\}, \quad q \not\equiv 1 \pmod{3}, \\ P_{z^i,b^3}, i = 1, 2, 3, b \in \mathbb{F}_{q^3} \setminus \{0\}, \quad q \equiv 1 \pmod{3}.$$

Proof. The element of K induced by D_x fixes a point of π_1 (or of π_2) if and only if $x^{q-1} = x^{q^2+q+1} = 1$, that is $x^3 = 1$, with $x \in \mathbb{F}_q$. Therefore if $q \not\equiv 1 \pmod{3}$, then K permutes in a single orbit the points of π_1 (or of π_2). If $q \equiv 1 \pmod{3}$, then the kernel of the action of K on both π_1 and π_2 consists of the subgroup of order three induced by $\langle D_\xi \rangle$. Let $z \in \mathbb{F}_{q^3}$ be such

that $z^{q-1} = \xi$, where ξ is a fixed element in \mathbb{F}_q such that $\xi^2 + \xi + 1 = 0$. In order to see that the representatives of the K -orbits on points of π_1 (or of π_2) are $P_{z^i,0}$ (or P_{0,z^i}), $i = 1, 2, 3$, observe that

$$\mathbb{F}_{q^3} \setminus \{0\} = \bigcup_{i=1}^3 \{xz^i \mid x \in \mathbb{F}_{q^3}, x^{q^2+q+1} = 1\}.$$

Let R be a point of $\text{PG}(V) \setminus (\pi_1 \cup \pi_2)$. It is straightforward to check that no non-trivial element of K leaves R invariant, that is $|R^K| = q^2 + q + 1$. Since the planes π_1, π_2 are disjoint, there exists a unique line ℓ of $\text{PG}(V)$ intersecting both π_1 and π_2 and containing R . Hence if $|\ell \cap R^K| \geq 2$, then there exists a non-trivial element in K fixing ℓ and hence stabilizing both $\ell \cap \pi_1$ and $\ell \cap \pi_2$. From the discussion above such a non-trivial element exists if and only if $q \equiv 1 \pmod{3}$. If $q \not\equiv 1 \pmod{3}$, it follows that the representatives of the K -orbits on points of $\text{PG}(V) \setminus (\pi_1 \cup \pi_2)$ can be taken as the points on the $q^2 + q + 1$ lines through $P_{1,0}$ and meeting π_2 in a point. Similarly, if $q \equiv 1 \pmod{3}$, then these representatives can be chosen among the points of $\text{PG}(V) \setminus (\pi_1 \cup \pi_2)$ lying on the $3(q^2 + q + 1)$ lines through $P_{z^i,0}$, $i = 1, 2, 3$, and meeting π_2 in a point. In particular, in this case, $P_{z^i,b}$ and $P_{z^i,b'}$ lie in the same K -orbit if $b' = \xi^i b$, $i = 1, 2, 3$. The result now follows. \square

Theorem 3.7. *If $c \in \mathbb{F}_q \setminus \{0\}$, the orbit $\mathcal{O} = P_{1,c}^K$ is a partial ovoid of $\mathcal{W}(5, q)$ of size $q^2 + q + 1$.*

Proof. Let P' be a point of $P_{1,c}^K \setminus \{P_{1,c}\}$. Then there exists $x \in \mathbb{F}_{q^3}$, with $N(x) = 1$, $x \neq 1$, such that

$$P' = (x, x^q, x^{q^2}, cx^{-q^2}, cx^{-q}, cx^{-1}).$$

If the line joining $P_{1,c}$ with P' is a line of $\mathcal{W}(5, q)$, then

$$0 = cT(x^{-1} - x) = cT(x^{q+1} - x) = cN(1 - x)$$

Therefore $x = 1$ and $P_{1,c} = P'$, a contradiction. \square

In order to prove that \mathcal{O} is maximal we need to recall some results obtained by Culbert and Ebert in [6] in terms of Sherk surfaces. A *Sherk surface* $S(\alpha, \beta, \gamma, \delta)$, where $\alpha, \delta \in \mathbb{F}_q$, $\beta, \gamma \in \mathbb{F}_{q^3}$, can be seen as a hypersurface of the projective line $\text{PG}(1, q^3)$. More precisely

$$S(\alpha, \beta, \gamma, \delta) = \{x \in \mathbb{F}_{q^3} \cup \{\infty\} \mid \alpha N(x) + T(\beta^{q^2} x^{q+1}) + T(\gamma x) + \delta = 0\}.$$

Lemma 3.8 ([6]). *Let $\alpha, \delta \in \mathbb{F}_q$, $\beta, \gamma \in \mathbb{F}_{q^3}$.*

- 1) $|S(\alpha, \beta, \gamma, \delta)| \in \{1, q^2 - q + 1, q^2 + 1, q^2 + q + 1\}$.
- 2) $\infty \in S(\alpha, \beta, \gamma, \delta)$ if and only if $\alpha = 0$.
- 3) $|S(\alpha, \beta, \gamma, \delta)| = 1$ if and only if $(\alpha, \beta, \gamma, \delta) \in \{(1, \beta, \beta^{q^2+q}, N(\beta)), (0, 0, 0, 1)\}$.

Note that $\lambda S(\alpha_1, \beta_1, \gamma_1, \delta_1) + \mu S(\alpha_2, \beta_2, \gamma_2, \delta_2) = S(\lambda\alpha_1 + \mu\alpha_2, \lambda\beta_1 + \mu\beta_2, \lambda\gamma_1 + \mu\gamma_2, \lambda\delta_1 + \mu\delta_2)$ for any $\lambda, \mu \in \mathbb{F}_q$, $(\lambda, \mu) \neq (0, 0)$. Therefore any two distinct Sherk surfaces give rise to a pencil containing $q + 1$ distinct Sherk surfaces which together contain all the points of $\text{PG}(1, q^3)$. Moreover, any two distinct Sherk surfaces in a given pencil intersect in the same set of points, called the *base locus* of the pencil, which is the intersection of all the Sherk surfaces in that pencil.

Lemma 3.9. *Let $\beta, \gamma \in \mathbb{F}_{q^3}$ not both zero. If one of the following is satisfied*

- $\beta\gamma = 0$,
- $N(\beta) \neq c^3$, for all $c \in \mathbb{F}_q \setminus \{0\}$,
- $N(\beta) = c^3$, for some $c \in \mathbb{F}_q \setminus \{0\}$, and $\gamma\beta \neq -c^2$,

then the base locus of the pencil generated by $S(1, 0, 0, -1)$ and $S(0, \beta, \gamma, 0)$ is not empty.

Proof. The pencil consists of $S(0, \beta, \gamma, 0)$ and $S(1, 0, 0, -1) + \lambda S(0, \beta, \gamma, 0) = S(1, \lambda\beta, \lambda\gamma, -1)$, where $\lambda \in \mathbb{F}_q$. Let y_i be the number of Sherk surfaces in the pencil having i points. Hence y_i are non-negative integer and $y_{q^2+q+1} \geq 1$ since $|S(1, 0, 0, -1)| = q^2 + q + 1$. By Lemma 3.8, it follows that $|S(0, \beta, \gamma, 0)| \neq 1$ and $|S(1, \lambda\beta, \lambda\gamma, -1)| \neq 1$, for $\lambda \in \mathbb{F}_q$, i.e., $y_1 = 0$. Assume by contradiction that the base locus of the pencil is empty, then the following equations hold

$$\begin{aligned} y_{q^2-q+1} + y_{q^2+1} + y_{q^2+q+1} &= q + 1, \\ (q^2 - q + 1)y_{q^2-q+1} + (q^2 + 1)y_{q^2+1} + (q^2 + q + 1)y_{q^2+q+1} &= q^3 + 1. \end{aligned}$$

By substituting $y_{q^2-q+1} = q + 1 - y_{q^2+1} - y_{q^2+q+1}$ in the second equation, we have that

$$y_{q^2+q+1} = -\frac{y_{q^2+1}}{2}.$$

Therefore necessarily

$$y_{q^2+q+1} = y_{q^2+1} = 0, y_{q^2-q+1} = q + 1,$$

which is a contradiction. □

Theorem 3.10. *The partial ovoid \mathcal{O} of $\mathcal{W}(5, q)$ is maximal.*

Proof. It is enough to show that for each of the representatives R of the K -orbits on points of $\text{PG}(V) \setminus \mathcal{O}$, there exists a point $P = (x, x^q, x^{q^2}, cx^{-q^2}, cx^{-q}, cx^{-1}) \in \mathcal{O}$ such that the line joining R and P is a line of $\mathcal{W}(5, q)$. If R is a point of π_1 , then $R = P_{1,0}$ if $q \not\equiv 1 \pmod{3}$ or $R \in \{P_{z^i,0} \mid i = 1, 2, 3\}$, if $q \equiv 1 \pmod{3}$. The line RP belongs to $\mathcal{W}(5, q)$ if and only if there exists $x \in \mathbb{F}_{q^3}$, with $N(x) = 1$ such that

$$cT(ux^{-1}) = 0, \tag{3.1}$$

where $u = 1$ or $u \in \{z^i \mid i = 1, 2, 3\}$ according as $q \not\equiv 1 \pmod{3}$ or $q \equiv 1 \pmod{3}$. Since $c \neq 0$ and $N(x) = 1$, (3.1) is equivalent to $T(u^{q^2}x^{q+1}) = 0$, i.e., $x \in S(0, u, 0, 0)$. Hence the existence of a line RP of $\mathcal{W}(5, q)$ is equivalent to the following

$$|S(1, 0, 0, -1) \cap S(0, u, 0, 0)| \geq 1. \quad (3.2)$$

As before let $u = 1$ or $u \in \{z^i \mid i = 1, 2, 3\}$ according as $q \not\equiv 1 \pmod{3}$ or $q \equiv 1 \pmod{3}$. Let R be the point $P_{0,u} \in \pi_2$ or $P_{u,b} \in \text{PG}(V) \setminus (\pi_1 \cup \pi_2 \cup \mathcal{O})$, where $b \in \mathbb{F}_{q^3} \setminus \{0\}$, with $b^3 \neq c^3$, if $u = 1$, otherwise $P_{1,b} \in \mathcal{O}$. Arguing in a similar way we get that the existence of a line RP of $\mathcal{W}(5, q)$ is equivalent to

$$|S(1, 0, 0, -1) \cap S(0, 0, u, 0)| \geq 1 \text{ or } |S(1, 0, 0, -1) \cap S(0, cu, -b, 0)| \geq 1. \quad (3.3)$$

Since $N(u)$ is a cube in \mathbb{F}_q if and only if $u = 1$ and if $u = 1$, then $b \neq c$, the inequalities in (3.2) and (3.3) are satisfied by Lemma 3.9. \square

3.3 Maximal partial ovoids of $\mathcal{W}(5, q)$, q even, of size $2q^2 - q + 1$

Assume q to be even and let $\mathcal{W}(5, q)$ be the symplectic polar space of $\text{PG}(5, q)$ induced by the totally isotropic subspaces of \mathbb{F}_q^6 with respect to the non-degenerate alternating form given by

$$x_1y_2 + x_2y_1 + x_3y_4 + x_4y_3 + x_5y_6 + x_6y_5.$$

Recall that \mathfrak{s} is the symplectic polarity of $\text{PG}(5, q)$ associated with $\mathcal{W}(5, q)$. Fix $\delta_i \in \mathbb{F}_q$ such that the polynomial $X^2 + X + \delta_i$ is irreducible over \mathbb{F}_q , $i = 1, 2$. Let Δ_1 and Δ_2 be the three-spaces given by $X_5 = X_6 = 0$ and $X_3 = X_4 = 0$, respectively; consider the three-dimensional elliptic quadrics $\mathcal{E}_i \subset \Delta_i$ defined as follows

$$\begin{aligned} \mathcal{E}_1 : X_1X_2 + X_3^2 + X_3X_4 + \delta_1X_4^2 &= 0, \\ \mathcal{E}_2 : X_1X_2 + X_5^2 + X_5X_6 + \delta_2X_6^2 &= 0. \end{aligned}$$

Denote by σ the plane of Δ_1 spanned by U_1, U_2, U_3 ; hence $\mathcal{E}_1 \cap \sigma$ is a conic and $U_3^{\mathfrak{s}} \cap \Delta_1 = \sigma$. In the hyperplane $U_3^{\mathfrak{s}} : X_4 = 0$, let \mathcal{P}_1 be the cone having as vertex the point U_3 and as base \mathcal{E}_2 and let \mathcal{P}_2 be the cone having as vertex the line $\Delta_1^{\mathfrak{s}}$ and as base the conic $\mathcal{E}_1 \cap \sigma$. Set

$$\mathcal{A} = \mathcal{P}_1 \cap \mathcal{P}_2 = \{(1, c^2 + cd + \delta_2d^2, \sqrt{c^2 + cd + \delta_2d^2}, 0, c, d) \mid c, d \in \mathbb{F}_q\} \cup \{U_2\}.$$

Lemma 3.11. *The point sets \mathcal{E}_1 and \mathcal{A} are partial ovoids of $\mathcal{W}(5, q)$ of size $q^2 + 1$.*

Proof. The elliptic quadric \mathcal{E}_1 is an ovoid of $\Delta_1 \cap \mathcal{W}(5, q)$. Hence it is a partial ovoid of $\mathcal{W}(5, q)$. The set \mathcal{A} is contained in $U_3^{\mathfrak{s}}$ and each of the lines obtained by joining U_3 with a point of \mathcal{A} intersects Δ_2 in a point of \mathcal{E}_2 . In particular this gives a bijective correspondence between the points of \mathcal{A} and those of \mathcal{E}_2 . In order to see that \mathcal{A} is a partial ovoid of $\mathcal{W}(5, q)$ note that the elliptic quadric \mathcal{E}_2 is an ovoid of $\Delta_2 \cap \mathcal{W}(5, q)$. \square

Theorem 3.12. *The point set $\mathcal{A} \cup (\mathcal{E}_1 \setminus \sigma)$ is a maximal partial ovoids of $\mathcal{W}(5, q)$ of size $2q^2 - q + 1$.*

Proof. Taking into account the previous lemma, it is enough to show that if P is a point of $\mathcal{E}_1 \setminus \sigma$, then $|P^5 \cap \mathcal{A}| = 0$. By contradiction let $P = (1, a^2 + ab + \delta_1 b^2, a, b, 0, 0) \in \mathcal{E}_1 \setminus \sigma$ and $R = (1, c^2 + cd + \delta_2 d^2, \sqrt{c^2 + cd + \delta_2 d^2}, 0, c, d) \in \mathcal{A}$, for some $a, b, c, d \in \mathbb{F}_q$, $b \neq 0$, with $R \in P^5$. Then $\sqrt{c^2 + cd + \delta_2 d^2} \in \mathbb{F}_q$ is a root of $X^2 + bX + a^2 + ab + \delta_1 b^2$, which is irreducible over \mathbb{F}_q , a contradiction.

In order to prove the maximality, let Q be a point not in $\mathcal{A} \cup (\mathcal{E}_1 \setminus \sigma)$ and assume that $|Q^5 \cap (\mathcal{E}_1 \setminus \sigma)| = 0$. There are two possibilities: either $Q^5 \cap \Delta_1$ coincides with σ or $Q^5 \cap \Delta_1$ is a plane meeting \mathcal{E}_1 in exactly one point of the conic $\mathcal{E}_1 \cap \sigma$. In the former case Q^5 meets \mathcal{A} in the points $\{U_1, U_2\}$. If the latter possibility occurs, let R be the point of the conic $\mathcal{E}_1 \cap \sigma$ contained in $Q^5 \cap \Delta_1$; since $R^5 \cap \Delta_1 \subset Q^5$, the point Q lies in the plane spanned by R and Δ_1^5 . Hence the line joining R and Q is a line of $\mathcal{W}(5, q)$ and contains a unique point of \mathcal{A} . \square

Remark 3.13. Some computations performed with Magma [1] show that the largest partial ovoid of $\mathcal{W}(5, q)$, has size 7 for $q = 2$ and 13 for $q = 3$. In particular, $\mathcal{W}(5, 2)$ has a unique partial ovoid of size 7, up to projectivities, which coincides with the partial ovoids described above.

4 Partial ovoids of Hermitian polar spaces

Let $\mathcal{H}(2n, q^2)$, $n \geq 2$, be the Hermitian polar space of $\text{PG}(2n, q^2)$ and let \perp be its associated unitary polarity. Fix a point $P \notin \mathcal{H}(2n, q^2)$ and a pointset $\mathcal{T} \subset P^\perp$ suitably chosen. In this section we explore the partial ovoids of $\mathcal{H}(2n, q^2)$ that are constructed by taking the points of $\mathcal{H}(2n, q^2)$ on the lines obtained by joining P with the points of \mathcal{T} . It turns out that in order to get a partial ovoid, the set \mathcal{T} has to intersect the lines that are tangent or contained in $P^\perp \cap \mathcal{H}(2n, q^2)$ in at most one point.

4.1 Tangent-sets of $\mathcal{H}(2n - 1, q^2)$, $n \in \{2, 3, 4\}$

Let $\mathcal{H}(2n - 1, q^2)$ be a non-degenerate Hermitian variety of $\text{PG}(2n - 1, q^2)$. We introduce the notion of *tangent set* of $\mathcal{H}(2n - 1, q^2)$.

Definition 4.1. A *tangent-set* of a Hermitian variety $\mathcal{H}(2n - 1, q^2)$ is a set \mathcal{T} of points of $\text{PG}(2n - 1, q^2)$ such that every line that is either tangent or contained in $\mathcal{H}(2n - 1, q^2)$ has at most one point in common with \mathcal{T} . A tangent-set is said to be maximal if it is not contained in a larger tangent-set.

Here the main aim is to show that starting from a (partial) ovoid of $\mathcal{W}(2n - 1, q)$ it is possible to obtain a tangent set of $\mathcal{H}(2n - 1, q^2)$. In order to do that some preliminary results are needed.

Lemma 4.2. An ovoid of $\mathcal{W}(3, q) \subset \mathcal{H}(3, q^2)$ is a maximal partial ovoid of $\mathcal{H}(3, q^2)$.

Proof. Let \mathcal{X} be an ovoid of $\mathcal{W}(3, q) \subset \mathcal{H}(3, q^2)$. Then obviously \mathcal{X} is a partial ovoid of $\mathcal{H}(3, q^2)$. In order to see that \mathcal{X} is maximal as a partial ovoid of $\mathcal{H}(3, q^2)$, observe that every point of $\mathcal{H}(3, q^2) \setminus \mathcal{W}(3, q)$ lies on a unique extended line of $\mathcal{W}(3, q)$. \square

Let us fix an element $\iota \in \mathbb{F}_{q^2}$ such that $\iota + \iota^q = 0$ and let ξ_1, \dots, ξ_q be q elements of \mathbb{F}_{q^2} such that

$$\{\iota(\xi_i^q - \xi_i) \mid i = 1, \dots, q\} = \mathbb{F}_q. \quad (4.1)$$

In particular we may assume $\xi_1 = 0$. Let \mathcal{H}_i be the Hermitian variety given by

$$X_1 X_{n+1}^q - X_1^q X_{n+1} + \dots + X_n X_{2n}^q - X_n^q X_{2n} + (\xi_i^q - \xi_i) X_{2n}^{q+1} = 0, \quad i = 1, \dots, q.$$

Then, if Π denotes the hyperplane $X_{2n} = 0$, the set $\{\mathcal{H}_i \mid i = 1, \dots, q\} \cup \{\Pi\}$ is a pencil of Hermitian varieties whose base locus consists of $\Pi \cap \mathcal{H}_i$. Let Σ_i denote the Baer subgeometry of $\text{PG}(2n-1, q^2)$ consisting of the points

$$\{(a_1, \dots, a_{n-1}, a_n + \xi_i, a_{n+1}, \dots, a_{2n-1}, 1) \mid a_1, \dots, a_{2n-1} \in \mathbb{F}_q, (a_1, \dots, a_{2n-1}) \neq (0, \dots, 0)\} \\ \cup \{(a_1, \dots, a_{2n-1}, 0) \mid a_1, \dots, a_{2n-1} \in \mathbb{F}_q, (a_1, \dots, a_{2n-1}) \neq (0, \dots, 0)\},$$

for $i = 1, \dots, q$. It is easily seen that $\Sigma_i \subset \mathcal{H}_i$, for $i = 1, \dots, q$. Hence $|\mathcal{H}_1 \cap (\Sigma_i \setminus \Pi)| = 0$, if $i \neq 1$. Moreover, the lines of \mathcal{H}_i having $q+1$ points in common with Σ_i are those of a symplectic polar space \mathcal{W}_i of Σ_i . Let K be the group of projectivities of $\text{PG}(2n-1, q^2)$ of order q^{2n-1} induced by the matrices

$$\begin{pmatrix} 1 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & a_1 \\ 0 & 1 & \dots & 0 & 0 & 0 & \dots & 0 & a_2 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 1 & 0 & 0 & \dots & 0 & a_{n-1} \\ -a_{n+1} & -a_{n+2} & \dots & -a_{2n-1} & 1 & a_1 & \dots & a_{n-1} & a_n \\ 0 & 0 & \dots & 0 & 0 & 1 & \dots & 0 & a_{n+1} \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & a_{2n-1} \\ 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & 1 \end{pmatrix}, \quad a_1, \dots, a_{2n-1} \in \mathbb{F}_q.$$

Then K fixes \mathcal{W}_i and \mathcal{H}_i ; furthermore it permutes in a single orbit the q^{2n-1} points of $\Sigma_i \setminus \Pi$, $i = 1, \dots, q$. The polar hyperplane of U_n with respect to \mathcal{H}_i is Π ; if Γ is a $(2n-3)$ -space of Π such that $U_n \notin \Gamma$ and $\Gamma \cap \Sigma_i$ is a $(2n-3)$ -space, then $|\Gamma^\perp \cap \Sigma_i| = q+1$. Hence the q^{2n-1} points of $\bigcup_i \Sigma_i \setminus \Pi$ lie on q^{2n-2} lines of $\text{PG}(2n-1, q^2)$ through U_n . Denote by \mathcal{L} this set of q^{2n-2} lines, that is

$$\mathcal{L} = \left\{ \Gamma^\perp \mid \Gamma \text{ is a } (2n-3)\text{-space of } \Pi, U_n \notin \Gamma, |\Gamma \cap \Sigma_i| = \frac{q^{2n-2} - 1}{q-1} \right\}.$$

Note that K acts transitively on \mathcal{L} . In the next lemma we restrict attention on the case $n = 2$.

Lemma 4.3. *Let $\ell \in \mathcal{L}$. The intersection of Π with the tangent lines to \mathcal{H}_1 through a point of $(\ell \cap \Sigma_i) \setminus \Pi$, $i = 2, \dots, q$, is a non-degenerate Hermitian curve of Π belonging to the pencil determined by ℓ^\perp and $\Pi \cap \mathcal{H}_1$. Viceversa, if r is a line of Π with $|r \cap \Sigma_i| = |r \cap \mathcal{H}_1| = q+1$ and \mathcal{U}_r is a non-degenerate Hermitian curve of Π belonging to the pencil determined by r and $\Pi \cap \mathcal{H}_1$, then there exists k such that the lines joining a point of $(r^\perp \cap \Sigma_k) \setminus \Pi$ with a point of \mathcal{U}_r are tangent to \mathcal{H}_1 .*

Proof. If $P \in (\ell \cap \Sigma_i) \setminus \Pi$, $i = 2, \dots, q$, we may assume $P = (0, a_2 + \xi_i, 0, 1)$, $\xi_i \neq 0$, $a_2 \in \mathbb{F}_q$. Then the points of $(x, y, z, 0)$ of Π lying on a line through P and tangent to \mathcal{H}_1 satisfy

$$\iota^2 y^{q+1} + \iota \alpha (x^q z - x z^q) = 0, \quad (4.2)$$

where $\alpha = \iota(\xi_i^q - \xi_i)$. Hence they form a non-degenerate Hermitian curve of Π belonging to the pencil determined by r and $\Pi \cap \mathcal{H}_1$. Viceversa, if r is a line of Π with $|r \cap \Sigma_i| = |r \cap \mathcal{H}_1| = q + 1$, we may assume that r is the line joining U_1 and U_3 . It follows that \mathcal{U}_r is given by (4.2), for some $\alpha \in \mathbb{F}_q \setminus \{0\}$. Thus if $\alpha = \iota(\xi_k^q - \xi_k)$, the lines joining a point of $(r^\perp \cap \Sigma_k) \setminus \Pi$ with a point of \mathcal{U}_r are tangent to \mathcal{H}_1 . \square

Lemma 4.4. *Let ℓ be a line that is either tangent or contained in \mathcal{H}_1 . Then $|\ell \cap (\bigcup_i^q \Sigma_i)| \in \{0, 1, q + 1\}$. In particular, if $|\ell \cap (\bigcup_i^q \Sigma_i)| = q + 1$ then there exists k such that $\ell \cap \Sigma_k = q + 1$ and $\ell \subset \mathcal{H}_k$.*

Proof. If ℓ is contained in \mathcal{H}_1 , then there is nothing to prove. If ℓ is tangent, let P_i, P_j be two points belonging to $\ell \cap \Sigma_i$ and $\ell \cap \Sigma_j$, respectively, where $i \neq 1$. Since K is transitive on points of $\Sigma_i \setminus \Pi$, we may assume $P_i = (0, \dots, 0, \xi_i, 0, \dots, 0, 1)$. If $P_j = (a_1, \dots, a_{n-1}, a_n + \xi_j, a_{n+1}, \dots, a_{2n-1}, 1)$ for some $a_1, \dots, a_{2n-1} \in \mathbb{F}_q$ not all zero, and $\lambda \in \mathbb{F}_{q^2}$, then the point $P_j + \lambda P_i$ belongs to \mathcal{H}_1 if and only if λ is a root of

$$\iota(\xi_i - \xi_j)X^{q+1} + \iota(\xi_j - \xi_i^q + a_n)X^q + \iota^q(\xi_j^q - \xi_i + a_n)X + \iota(\xi_j - \xi_j^q) = 0.$$

Some straightforward calculations show that ℓ is tangent if and only if $\xi_j = \xi_i - a_n$. If $i \neq j$, then $\iota(\xi_j^q - \xi_j) = \iota(\xi_i^q - \xi_i)$, contradicting (4.1). Therefore $i = j$ and $a_n = 0$. This means that $|\ell \cap \Sigma_i| = q + 1$. Moreover $\ell \subset \mathcal{H}_i$. If $P_j = (a_1, \dots, a_{2n-1}, 0)$ for some $a_1, \dots, a_{2n-1} \in \mathbb{F}_q$, then $P_j \in \Sigma_i \cap \Sigma_j$. Hence $|\ell \cap \Sigma_i| = q + 1$ and we may repeat the previous argument. \square

Theorem 4.5. *Let \mathcal{O}_i be a (partial) ovoid of \mathcal{W}_i , $i = 1, \dots, q$, such that $\mathcal{O}_i \cap \Pi = \mathcal{O}_j \cap \Pi$, if $i \neq j$. Then $\mathcal{T} = \bigcup_{i=1}^q \mathcal{O}_i$ is a tangent-set of \mathcal{H}_1 . Moreover if $n = 2$ and \mathcal{O}_i is an ovoid of \mathcal{W}_i , for every $i = 1, \dots, q$, then \mathcal{T} is a maximal tangent-set.*

Proof. Assume by contradiction that \mathcal{T} is not a tangent-set. If ℓ is a line of \mathcal{H}_1 and $|\ell \cap \mathcal{T}| \geq 2$, then $\ell \cap \mathcal{T} \subset \mathcal{O}_1$, since $\mathcal{T} \cap \mathcal{H}_1 = \mathcal{O}_1$. Hence ℓ is an extended line of \mathcal{W}_1 containing two points of \mathcal{O}_1 , contradicting the assumption that \mathcal{O}_1 is a (partial) ovoid of \mathcal{W}_1 . Similarly, if ℓ is tangent to \mathcal{H}_1 and $|\ell \cap \mathcal{T}| \geq 2$, then, by Lemma 4.4, there exists a k such that ℓ is an extended line of \mathcal{W}_k and $\ell \cap \mathcal{T} \subset \mathcal{O}_k$. Therefore \mathcal{O}_k is not a (partial) ovoid of \mathcal{W}_k , since $|\ell \cap \mathcal{O}_k| \geq 2$, contradicting the hypotheses.

Assume now that $n = 2$ and \mathcal{O}_i is an ovoid of \mathcal{W}_i , for every $i = 1, \dots, q$. We claim that every point R of $\text{PG}(3, q^2) \setminus \mathcal{T}$ lies on at least one line that is either tangent or contained in \mathcal{H}_1 and contains one point of \mathcal{T} . If $R \notin \Pi$, then there exactly one \mathcal{H}_k with $R \in \mathcal{H}_k$; hence there is at least one extended line, say r_k of \mathcal{W}_k containing R . Such a line r_k has to contain exactly one point of \mathcal{O}_k (see Lemma 4.2) and by Lemma 4.4 it is contained or tangent to \mathcal{H}_1 according as $k = 1$ or $k \neq 1$. Similarly if $R \in \mathcal{H}_i \cap \Pi$. Let R be a point of $\Pi \setminus \mathcal{H}_i$. There are two possibilities:

either $U_2 \in \mathcal{O}_i$ for every $i = 1, \dots, q$, and in this case the line joining R and U_2 is tangent to \mathcal{H}_1 or $U_2 \notin \mathcal{O}_i$, for $i = 1, \dots, q$. If the latter possibility occurs, then there exists a line r of Π such that $R \notin r$, $|r \cap \mathcal{O}_i| = 0$ and $|r \cap \Sigma_i| = |r \cap \mathcal{H}_1| = q + 1$. Let \mathcal{U}_r be the (unique) non-degenerate Hermitian curve of Π belonging to the pencil determined by r and $\Pi \cap \mathcal{H}_1$ such that $R \in \mathcal{U}_r$. By Lemma 4.3, there exists a k such that the lines joining a point of $(r^\perp \cap \Sigma_k) \setminus \Pi$ with R are tangent to \mathcal{H}_1 . The Baer subline $r^\perp \cap \Sigma_k$ contains two points of \mathcal{O}_k being the polar line of $r \cap \Sigma_k$ with respect to the symplectic polarity of Σ_k associated with \mathcal{W}_k . Therefore the line joining a point of $r^\perp \cap \Sigma_k \cap \mathcal{O}_k$ with R is tangent to \mathcal{H}_1 . \square

Proposition 4.6. *If there is a (partial) ovoid of $\mathcal{W}(2n - 1, q)$ of size $x + y$, with y points on a hyperplane, then there exists a tangent-set of $\mathcal{H}(2n - 1, q^2)$ of size $xq + y$.*

The next result is reached by applying Theorem 4.5, where \mathcal{O}_i is either an ovoid of $\mathcal{W}(3, q)$, q even, or the partial ovoid of $\mathcal{W}(3, q)$, obtained in Theorem 3.3, or the partial ovoids of $\mathcal{W}(5, q)$ described in Theorem 3.7 and in Theorem 3.12, or the partial ovoid of $\mathcal{W}(7, q)$ of size $q^3 + 1$ given in [2, 3].

Corollary 4.7. *The following hold true.*

- *If q is even, there exist maximal tangent-sets of $\mathcal{H}(3, q^2)$ of size $q^3 + 1$ and of size $q^3 - q^2 + q + 1$.*
- *If q is an odd square and $q \not\equiv 0 \pmod{3}$, there exists a tangent-set of $\mathcal{H}(3, q^2)$ of size $(q^{5/2} + 3q^2 - q^{3/2} + 3q)/3$.*
- *If q is even, there exists a tangent-set of $\mathcal{H}(5, q^2)$ of size $2q^3 - q^2 + 1$.*
- *There exists a tangent-set of $\mathcal{H}(5, q^2)$ of size $q^3 + q^2 + 1$.*
- *There exists a tangent-set of $\mathcal{H}(7, q^2)$ of size $q^4 + 1$.*

4.2 Partial ovoids of $\mathcal{H}(2n, q^2)$ from tangent-sets of $\mathcal{H}(2n - 1, q^2)$, $n \in \{2, 3, 4\}$

The next result shows that starting from a (maximal) tangent-set of $\mathcal{H}(2n - 1, q^2)$ it is possible to obtain a (maximal) partial ovoid of $\mathcal{H}(2n, q^2)$.

Lemma 4.8. *Let P be a point of $\text{PG}(2n, q^2) \setminus \mathcal{H}(2n, q^2)$ and let \mathcal{T} be a (maximal) tangent-set of $\mathcal{H}(2n - 1, q^2) = P^\perp \cap \mathcal{H}(2n, q^2)$. Then*

$$\mathcal{O} = \left(\bigcup_{R \in \mathcal{T}} \langle P, R \rangle \right) \cap \mathcal{H}(2n, q^2)$$

is a (maximal) partial ovoid of $\mathcal{H}(2n, q^2)$ of size $(q + 1)|\mathcal{T} \setminus \mathcal{H}(2n - 1, q^2)| + |\mathcal{T} \cap \mathcal{H}(2n - 1, q^2)|$.

Proof. Assume by contradiction that \mathcal{O} is not a partial ovoid, then there would exist a line ℓ of $\mathcal{H}(2n, q^2)$ containing two distinct points P_1 and P_2 of \mathcal{O} . If at least one among P_1, P_2 is not in P^\perp , then the line $\bar{\ell} \subset P^\perp$ obtained by projecting ℓ from P to P^\perp is tangent to

$\mathcal{H}(2n-1, q^2) = P^\perp \cap \mathcal{H}(2n, q^2)$ and contains two points of \mathcal{T} , a contradiction. If both P_1 and P_2 are in P^\perp , then ℓ is a line of $\mathcal{H}(2n-1, q^2)$ and contains two points of \mathcal{T} , a contradiction.

As for maximality, note that if T is a point of $\mathcal{H}(2n, q^2)$ such that $\mathcal{O} \cup \{T\}$ is a partial ovoid of $\mathcal{H}(2n, q^2)$ and $\bar{T} = \langle P, T \rangle \cap P^\perp$, then $\mathcal{T} \cup \{\bar{T}\}$ is a tangent-set of $\mathcal{H}(2n-1, q^2)$. \square

Combining Theorem 4.5 and Lemma 4.8, the following result arises.

Theorem 4.9. *If there is a (partial) ovoid of $\mathcal{W}(2n-1, q)$ of size $x+y$, with y points on a hyperplane, then there exists a partial ovoid of $\mathcal{H}(2n, q^2)$ of size xq^2+y .*

Hence starting from (partial) ovoids of $\mathcal{W}(2n-1, q)$, we get partial ovoids of $\mathcal{H}(2n, q^2)$, $n \in \{2, 3, 4\}$, as described below.

Corollary 4.10. *The following hold true.*

- *If q is even, there exist maximal partial ovoids of $\mathcal{H}(4, q^2)$ of size q^4+1 and of size q^4-q^3+q+1 .*
- *If q is an odd square and $q \not\equiv 0 \pmod{3}$, there exists a partial ovoid of $\mathcal{H}(4, q^2)$ of size $(q^{7/2}+3q^3-q^{5/2}+3q^2)/3$.*
- *If q is even, there exists a partial ovoid of $\mathcal{H}(6, q^2)$ of size $2q^4-q^3+1$.*
- *There exists a partial ovoid of $\mathcal{H}(6, q^2)$ of size q^4+q^3+1 .*
- *There exists a partial ovoid of $\mathcal{H}(8, q^2)$ of size q^5+1 .*

Acknowledgments. This work was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA–INdAM).

References

- [1] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.*, 24, (1997), 235-265.
- [2] A. Cossidente, On twisted tensor product group embeddings and the spin representation of symplectic groups: the case q odd, *ISRN Geom.*, 2011 (2011), 694605, 6 pp.
- [3] A. Cossidente, O.H. King, Twisted tensor product group embeddings and complete partial ovoids on quadrics in $\text{PG}(2^t-1, q)$, *J. Algebra* 273 (2004), 854-868.
- [4] A. Cossidente, F. Pavese, On $(0, \alpha)$ -sets of generalized quadrangles, *Finite Fields Appl.*, 30 (2014), 139-152.
- [5] A. Cossidente, A. Siciliano, Commuting polarities and maximal partial ovoids of $\mathcal{H}(4, q^2)$, *J. Combin. Designs*, 17 (2009), no. 4, 307-313.

- [6] C. Culbert, G.L. Ebert, Circle geometry and three-dimensional subregular translation planes, *Innov. Incidence Geom.*, 1 (2005), 3-18.
- [7] J. De Beule, A. Klein, K. Metsch, L. Storme, Partial ovoids and partial spread in Hermitian polar space, *Des. Codes Cryptogr.*, 47 (2008), 21-34.
- [8] J. De Beule, A. Klein, K. Metsch, L. Storme, Partial ovoids and partial spreads in symplectic and orthogonal polar spaces, *European J. Combin.*, 29 (2008), 1280-1297.
- [9] J.W.P. Hirschfeld, *Finite Projective Spaces of Three Dimensions*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1985.
- [10] J.W.P. Hirschfeld, *Projective Geometries over Finite Fields*, Oxford Mathematical Monographs, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1998.
- [11] F. Mazzocca, O. Polverino, L. Storme, Blocking sets in $PG(r, q^n)$, *Des. Codes Cryptogr.*, 44 (2007), 97-113.
- [12] C. Röbbing, L. Storme, A spectrum result on maximal partial ovoids of the generalized quadrangle $Q(4, q)$, q even, *European J. Combin.*, 31 (2010), no. 1, 349-361
- [13] G. Tallini, Fibrations through lines in a nonsingular quadric $Q_{4,q}$ of $PG(4, q)$, *Atti Accad. Peloritana Pericolanti. Cl. Sci. Fis. Mat. Natur.*, 66 (1988), 127-146 (1989).