

Interplay of charge and spin fluctuations in a Hund's coupled impurity

Victor Drouin-Touchette,^{1,*} Elio J. König,² Yashar Komijani,³ and Piers Coleman^{1,4}

¹Center for Materials Theory, Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854 USA

²Max-Planck Institute for Solid State Research, 70569 Stuttgart, Germany

³Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221-0011, USA

⁴Department of Physics, Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK

(Dated: March 11, 2022)

In Hund's metals, the local ferromagnetic interaction between orbitals leads to an emergence of complex electronic states with large and slowly fluctuating magnetic moments. Introducing a new analysis of a Hund's coupled mixed valence quantum impurity, we gain analytic insight into recent numerical renormalization group studies. We show that valence fluctuations drastically impedes the development of a large fluctuating moment over a wide swath of temperatures and energy, characterized by quenched orbital degrees of freedom and a singular logarithmic behavior of the spin susceptibility $\chi''_{\text{sp}}(\omega) \propto [\omega \log(\omega/T_K^{\text{eff}})]^{-1}$, closely resembling power-law scaling $\chi''_{\text{sp}}(\omega) \sim \omega^{-\gamma}$. These singular spin fluctuations may play an important role in future models of Hund's driven Cooper pairing.

Introduction - The concept of Hund's metals was first introduced in the context of iron-based superconductors [1–3], with the nomenclature now being extended to include the ruthenates [4–6]. In both cases, the local physics is characterized by electronic shells that are one filling away from half-filling [7, 8], such that the ferromagnetic inter-orbital Hund interaction overshadows the Coulomb interaction [9]. This class of materials presents an intermediate paramagnetic regime dominated by slowly fluctuating high-spin configurations [10–12] and largely suppressed Fermi liquid coherence scales, leading to anomalous transport properties [13–15]. It has been speculated that Cooper pairing emerges out of this intermediate state in iron-based compounds [16–22].

The link between large moments generated by Hund's coupling and the exponential reduction of Fermi liquid scales in Kondo impurity models has been studied extensively [23–28]. However, the physical valence of Fe or Ru atoms in Hund's metals deviates from half-filled shells. This has led to a vigorous interest in doped multi-orbital models, where an intermediate coupling non-Fermi-Liquid fixed point was pointed out through analytical [29] and NRG studies [30–32] for the $S = 1$, three orbital system. Although these studies are performed in the physical subspace of two electrons among three orbitals, they did not explicitly treat charge fluctuations out of this state.

Motivated by the potentially new physics in the mixed valence regime, here we extend our previous work on the Hund-Kondo impurity [28]. This, coupled with the unique property of the large- N self-consistent equations which enable a direct access to real frequency correlation functions [33], has led us to study in detail the dynamical properties of the intermediate regime generated by Hund's coupling.

In this Letter, we show that the treatment of charge and spin dynamics on equal footing within the dynamical large- N approach [28, 33–41] leads to new insight into




FIG. 1. (a-b) Schematic mixed valence states and of the Hund-coupled three-orbital Anderson model, referring to Eq. (2). (c) The obtained phase diagram as a function of total impurity filling n_{imp} (left side is holon dominated). T_{orb} and T_K^{eff} are crossover temperatures where the specific heat c_v has a local maxima, as in Fig. 2. Generically, screening occurs in two steps. The formation of a large emergent fluctuating moment, while orbital degrees of freedom are quenched, occurs at T_{orb} . In this spin-orbital separated (S.O.S.) regime, we see a Curie-like spin susceptibility $\chi_{\text{sp}} \sim \mu^2/T$ while the orbital susceptibility reaches a plateau. Then, at $T_K^{\text{eff}} \ll T_K^0$, with T_K^0 the bare Kondo temperature for $J_H = 0$, the large moment is screened and forms a local Fermi Liquid. $\Gamma = \pi\rho V^2$ is the bare hybridization width.

the dynamical properties of hole doped multi-orbital impurities. Based on computed thermodynamic quantities, we unveil the complete phase diagram [see Fig. 1] as a function of impurity occupancy. Furthermore, we characterize the large emergent moment regime [28] as one with spin-orbital separation [42]. The spin susceptibility in this regime shows logarithmic corrections due to the extremely slow approach to Kondo screening, reminiscent of the low-energy properties of the underscreened Kondo model [43, 44]. At a strongly renormalized Kondo tem-

perature $T_K^{\text{eff}} \ll T_K^0$, the holons eventually gap out at low energy, resulting in a local Fermi liquid. The scaling of the local spin susceptibility in the spin-orbital separated regime is found to exhibit quasi-power-law scaling due to the unusually strong logarithmic corrections.

Model - We consider the paradigmatic example of a three-orbital impurity model, resembling the t_{2g} orbital subset generated due to the tetrahedral environment around the iron atoms in Fe-based Hund metals. Mixed-valence states are included via a Hund-Anderson model. We take the limit of $U \rightarrow \infty$, such that the Coulomb interaction enforces a “no double occupancy” rule for each of the m orbitals. This is enforced through the use of Hubbard operators [45], which transform the K empty states $|d^0 : a\rangle$ into the N magnetic states $|d^1 : \alpha\rangle$ filled with local d electrons for each orbital m . In the large- N formalism, local moments transform as spin- S representations of $SU(N)$, and there are $K = 2S$ electronic channels present to maintain perfect screening. Schwinger bosons $X_{\alpha,\beta}^{(m)} = b_{m\alpha}^\dagger b_{m\beta}$ are used to express the spin degrees of freedom through spinons b^\dagger which form a symmetric representation of the spins. Together with the use of slave fermions (holons χ^\dagger), Hubbard operators can faithfully be represented as

$$\begin{aligned} X_{\alpha,a}^{(m)} &\equiv |d^1 : \alpha, m\rangle \langle d^0 : a, m| = b_{m\alpha}^\dagger \chi_{ma}, \\ X_{a,\alpha}^{(m)} &\equiv |d^0 : a, m\rangle \langle d^1 : \alpha, m| = \chi_{ma}^\dagger b_{m\alpha}, \\ X_{a,b}^{(m)} &\equiv |d^0 : a, m\rangle \langle d^0 : b, m| = \chi_{ma}^\dagger \chi_{mb}, \\ X_{\alpha,\beta}^{(m)} &\equiv |d^1 : \alpha, m\rangle \langle d^1 : \beta, m| = b_{m\alpha}^\dagger b_{m\beta}, \end{aligned} \quad (1)$$

while the Hamiltonian is itself expressed as $H = \sum_m H_c^{(m)} + \sum_m H_K^{(m)} + H_H$, with individual terms

$$H_c^{(m)} = \sum_{\mathbf{k}} \epsilon_{\mathbf{k}} c_{\mathbf{k}m\alpha a}^\dagger c_{\mathbf{k}m\alpha a}, \quad (2a)$$

$$H_K^{(m)} = V(c_{0m\alpha a}^\dagger X_{a,\alpha}^{(m)} + X_{\alpha,a}^{(m)} c_{0m\alpha a}) + \epsilon_f X_{a,a}^{(m)}, \quad (2b)$$

$$H_H = -\frac{J_H}{N} \sum_m X_{\alpha\beta}^{(m)} X_{\beta\alpha}^{(m+1)}. \quad (2c)$$

A visual representation of the mixed valence states and the Hund-Anderson model is presented at Fig. 1 (a). Here, V is the hybridization between conduction electrons and the Hubbard operators corresponding to adding or removing an impurity electron ($\Gamma = \pi\rho V^2$ is the bare hybridization width). We denote the energy of a hole as ϵ_f , which, when tuned, leads to different n_{imp} as the average valence of each state is changed. The Hund’s term H_H can be treated through a Hubbard-Stratonovich decoupling in the hopping channel:

$$H_H \rightarrow \sum_m [\bar{\Delta}_m (b_{m+1,\alpha}^\dagger b_{m\alpha}) + h.c.] + \frac{N|\Delta_m|^2}{J_H}. \quad (3)$$

A mean-field equation relates J_H to the spinon gap Δ . If J_H is large enough to generate a finite Δ , their relationship will be $N\Delta = 2S n_{\text{imp}} J_H$. Furthermore, the total charge

$$Q_m = \sum_{\alpha} b_{m\alpha}^\dagger b_{m\alpha} + \sum_a \chi_{ma}^\dagger \chi_{ma} = 2S, \quad (4)$$

at each orbital commutes with the Hubbard operators and is a conserved quantity, setting the size of the local moments. This is a constraint on the spinons/holons, enforced through a common Lagrange multiplier λ . In the large- N limit, the dynamics of holons and spinons is dominated by the non-crossing Feynman diagrams, which leads to the self-energy equations

$$\Sigma_\chi(\tau) = g_c(-\tau)G_B(\tau), \quad \Sigma_B(\tau) = -kg_c(\tau)G_\chi(\tau), \quad (5)$$

where $k = K/N = 2S/N$ and $g_{c,0}(z) = \sum_{\mathbf{k}} [z - \epsilon_{\mathbf{k}}]^{-1}$ corresponds to the conduction electron’s bare propagator for imaginary frequencies z . Eqs. (5) are solved self-consistently together with the Dyson equations $G_b(z) = \sum_p [z - \epsilon_p - V^2 \Sigma_b(z)]^{-1}$ and $G_\chi(z) = [z - \epsilon_f^* - V^2 \Sigma_\chi(z)]^{-1}$, where $\epsilon_p = (\lambda - 2\Delta \cos p)$ and $\epsilon_f^* = \lambda + \epsilon_f$. Here, we have defined the chirality $p = 0, \pm 2\pi/3$ which denotes the chosen Hund energy levels such that $p = 0$ is the aligned state. λ and Δ are adjusted to fit the constraints of Eqs. (3) and (4).

Thermodynamic and dynamical observables are obtained from the Green’s functions [28, 33, 34]. Notably, the impurity entropy $S_{\text{imp}}(T)$ can be extracted exactly in the large- N limit [46, 47] and is given by

$$S_{\text{imp}} = -\text{Tr}_\xi \int d\omega \frac{\partial n_\xi}{\partial T} [\text{Im} \ln(-G_\xi^{-1}) + G'_\xi \Sigma''_\xi], \quad (6)$$

where the trace is over all α and a spin and channel indices, chiralities p and the various fields χ (where $\xi = F$) and b (where $\xi = B$). G'_ξ is the real part of the retarded Green’s function while Σ''_ξ is the imaginary part of the self-energy. From this, we can extract the specific heat as $c_{v,\text{imp}} = T \partial S_{\text{imp}} / \partial T$.

Summary of results - In the absence of Hund’s coupling, the presence of holons increases the bare Kondo temperature $T_K^0 = D e^{-1/J_K^0 \rho} \sim D \exp(-|\epsilon_f|/\Gamma)$ for a fixed hybridization width Γ . This expected trend is seen in black in Fig. 1 (b), and we expand further on it in the supplementary materials [48].

For a finite Hund’s coupling, the situation changes drastically. The emergence of an intermediate large moment phase in the Kondo limit is consistent with our previous work on the Hund-Kondo model [28]. We can now connect this phase continuously throughout the hole-doped regime. At some critical holon doping, there is no longer enough local moments to lock together - the




FIG. 2. **(a)** The impurity's local moment μ_{imp}^2 obtained from the spin susceptibility, for varied impurity occupations n_{imp} . The large emergent moment, seen as a low-temperature higher plateau, is destroyed as more holons are added, following the trend of Fig. 1. **(b)** The specific heat $c_{v,\text{imp}}$, obtained from the closed form of the entropy for the multiorbital Anderson model, see Eq. (6). Both T_{orb} and T_K^{eff} are crossovers associated with c_v peaks. Dashed lines correspond to the high and intermediate temperature limits for the impurity moment obtained in Ref. [28].

two-step Kondo screening reverts to a single step Kondo crossover. The obtained phase diagram of Fig. 1 is consistent with other NRG+DMFT studies [9] of hole doped multiorbital impurity models.

In Fig. 2, we present the thermodynamic measurements of the total impurity's magnetic moment $\mu \sim T\chi_{\text{sp}}$ as well as the impurity specific heat for select impurity occupations n_{imp} throughout the entire temperature range. For Kondo-like systems ($n_{\text{imp}} = 2.7$) there is a clear non-monotonicity in the local moment, signaling the intermediate formation of an emergent large moment due to Hund's coupling. As the holon occupancy increases, there are less spinons in the system and the emergent moment can no longer form - the shoulder disappears at $n_{\text{imp}} \approx 0.7$. In the specific heat, this disappearance of the intermediate phase is seen as the low and high temperature crossover features merge to become one as $n_{\text{imp}} = 0.7$, where single-step Kondo screening occurs.

Previous works [9, 31, 32, 42, 49] have characterized the intermediate large moment phase as a regime with spin-orbital separation. We can take advantage of the Hubbard operators' description in terms of spinons and holons to write composite orbital operators. This procedure would not be possible in the absence of physical holons [28].

We start with the total impurity spin operators at imaginary time τ , described as $S_{\alpha\beta}(\tau) = \sum_m X_{\alpha\beta}^{(m)}(\tau) = \sum_m b_{m\alpha}^\dagger(\tau)b_{m\beta}(\tau)$ (α, β are SU(N) spin states). Then, we harness the SO(3) orbital symmetry and describe impurity orbital operators \hat{L}_γ ($\gamma = 1, 2, 3$ corresponding to the three degenerate orbitals) as $\hat{L}_\gamma = (1/NK) \sum_{mm'\alpha\alpha'} X_{\alpha\alpha}^{(m)}(L_\gamma)_{mm'} X_{\alpha\alpha'}^{(m')}$. The L_γ are generators of the SO(3) group and linear combinations of

the $(\lambda_2, \lambda_5, \lambda_7)$ Gell-Mann matrices [50]. Following these definitions, spin and orbital susceptibilities in the large-N limit can be expressed as

$$\chi_{\text{sp}}(\tau) = \frac{1}{N^2} \sum_\gamma \sum_{\alpha\beta} \langle S_{\alpha\beta}^{(\gamma)}(\tau) S_{\beta\alpha}^{(0)}(0) \rangle, \quad (7a)$$

$$\chi_{\text{orb}}(\tau) = \frac{1}{3} \sum_\gamma \langle \hat{L}_\gamma(\tau) \hat{L}_\gamma(0) \rangle. \quad (7b)$$

Thus, the inclusion of holons as bona fide particles (as opposed to Grassmannian variables in the Hund-Kondo model [28]) allows us to define a clear orbital degree of freedom and subsequently, to study its dynamical and static properties. The full expression for the dynamical susceptibilities in real frequency are presented in the supplementary materials [48], as well as further details on this derivation. Note that the orbital susceptibility has a $1/K^2$ factor reduction compared to χ_{sp} ; we nevertheless can plot $K^2\chi_{\text{orb}}$ and obtain valuable insight. The static components of these susceptibilities, $\chi_{\text{orb}}(\tau = 0)$ and $\chi_{\text{sp}}(\tau = 0)$, are shown in panel (a) of Fig. 3. The clear splitting of both susceptibilities at T_{orb} , and the subsequent plateau in χ_{orb} , heralds the formation of the large moment and the separation of spin and orbital scales (S.O.S). Throughout this regime, orbital and charge fluctuations are nearly frozen while the spin susceptibility remains Curie-like.

Dynamical susceptibilities - The emergent moment regime has clear thermodynamic attributes, as described above and in particular in Fig. 2. Further insight into this phase is provided by the dynamical spin and orbital susceptibilities, as defined in Eqs. (7). We show these in Fig. 3 for two different total impurity valences.

Firstly, it can be clearly seen that at high frequencies, both spin and orbital degrees of freedom fluctuate freely. Then, for $\omega < T_{\text{orb}}$, the lower-energy high-spin configurations split off, which is associated with the separation of spin and orbital fluctuation. The charge fluctuations are nearly frozen and the valence stabilizes below T_{orb} ; this quenching of orbital degrees of freedom leads to the decrease in χ_{orb}'' .

From Fig. 3, we see that, for many decades in frequency between T_K^{eff} and T_{orb} , the spin susceptibility seems to grow in a power-law $\chi_{\text{sp}} \sim \omega^{-\gamma}$ (dot-dashed green line). In Kondo impurity problems, such behavior is often indicative of non-Fermi-liquid fixed points [31, 32], for example in the 2-channel spin-1/2 Kondo model [51–54]. Closer examination reveals that this is not the case in this system, having maintained perfect screening throughout. Instead, we find a good agreement with the following scaling at low temperatures and intermediate frequencies:

$$\chi_{\text{sp}}''(\omega) = \frac{(J_K^{\text{eff}} \rho)^2}{\omega} \propto \left(\omega \left[\log \left(\frac{\omega}{T_K^{\text{eff}}} \right) \right]^2 \right)^{-1}. \quad (8)$$




FIG. 3. **(a)** Static spin and orbital susceptibilities for $n_{\text{imp}} = 2.7$, showing a clear separation at T_{orb} below which the Curie-like $\chi_{\text{sp}} \sim \mu^2/T$ spin susceptibility is in stark contrast to the constant χ_{orb} . **(b-c)** Imaginary part of the dynamical spin and orbital susceptibilities, Eqs. (7), presented for different occupations of the three-orbital impurity: $n_{\text{imp}} = 2$ and $n_{\text{imp}} = 2.7$ from left to right. The value of the Hund coupling $J_H/D = 0.37$ is fixed. The dashed black line corresponds to the derived second-order perturbation scaling of the spinon susceptibility, Eqs. (8), while the dot-dashed green line is the quasi-power-law form of Eq. (10) (with small offset for readability).

Here we cover the basic steps of this derivation and leave the details to [48]. A careful consideration of a single iteration [28] of the self-energy equations of Eq. (5) for $T_K^{\text{eff}} \leq \max(\omega, T) \leq T_{\text{orb}}$, along with a mapping of the mixed valence problem onto a Kondo problem, leads to an effective holon propagator $\tilde{G}_\chi(\omega) = -J_K^{\text{eff}}(\omega)$, with

$$\frac{1}{\rho J_K^{\text{eff}}(\omega)} \simeq \log\left(\frac{\max(\omega, T)}{T_K^{\text{eff}}}\right). \quad (9)$$

After having obtained the effective running Kondo coupling, we proceed in a second-order perturbation in J_K^{eff} of the spinon bubble of the spin susceptibility [55]. This leads to the scaling of Eq. (8). One can see in Fig 3 that it agrees perfectly within the intermediate regime $T_K^{\text{eff}} < \omega < T_{\text{orb}}$ with only T_K^{eff} as an input parameter. A downturn is observed at lower frequencies consistent with $\text{Im} \chi_{\text{sp}} \propto \omega$ in the Fermi liquid regime. This scaling holds for all n_{imp} of the phase diagram where the SOS phase is present.

In the SOS regime, the large separation of scales between T_K^{eff} and T_{orb} allows us to manipulate the logarithm in Eq. (8) [48]. For intermediate frequencies, we find that a quasi-power-law spin susceptibility

$$\chi_{\text{sp}}'' \sim \omega^{-\gamma} \quad \text{and} \quad \gamma = 1 - 2/\log\left(\frac{T_K^{\text{eff}}}{T_{\text{orb}}}\right). \quad (10)$$

For very small $T_K^{\text{eff}}/T_{\text{orb}}$, due to strong Hund's coupling and the resulting nearly frozen charge fluctuations, the slow logarithmic scaling presents itself as a quasi-power-law such that $\gamma \simeq 1.2$ for $n_{\text{imp}} = 2.7$ and $\gamma \simeq 1.4$ for $n_{\text{imp}} = 2.0$. This exponent γ changes continuously as n_{imp} is varied.

We note that $\chi_{\text{sp}}'' \sim \omega^{-1.2}$ was seen in a different but related model [31, 32], and was related to spin-fluctuation-induced Cooper pairing in the iron-based superconductors [56, 57]. Future work should be devoted to the inclusion of the dynamical spin susceptibility from Eq. (8) into an Eliashberg treatment of the formation of a superconducting condensate from local spin fluctuations [56]. Since the quasi-power-law is present for many decades in frequency, we expect the same conclusion as of Ref. [56] of a universal superconducting gap to T_c ratio at a given n_{imp} , e.g. $\Delta_{\text{max}}/k_B T_c \sim 7.2$, in the case of the iron-based superconductors.

Conclusion - We have shown that the dynamical large-N approach can capture the destruction of the Hund's coupled emergent large moment due to hole doping. Furthermore, the inclusion of bona fide holons leads to a description of orbital degrees of freedom and the conclusion that the intermediate regime is well described through the concept of spin-orbital separation (SOS) [9, 42]. In this phase, the dynamical spin susceptibility has a logarithmic component due to the extremely slow approach to the Kondo fixed point and to the nearly frozen charge fluctuations. Because $T_K^{\text{eff}} \ll T_{\text{orb}}$, this presents itself as a quasi-power-law for an extended intermediate frequency regime. The non-Fermi-liquid-like features in the emergent moment regime can therefore be continuously connected to the integer valence limit. Such dynamical local spin susceptibility could act as a glue for Cooper pairs in the iron-based superconductors, a subject of a future work.

Acknowledgments - VDT would like to thank Elias Walter, Jan von Delft, Fabian Kugler and Abhishek Kumar for illuminating discussions. We also acknowledge useful discussions with Thomas Schäfer. This

work was supported by DOE Basic Energy Sciences grant DE-FG02-99ER45790 (VDT, PC) and the Fonds de Recherche Québécois en Nature et Technologie (VDT).

* vdrouin@physics.rutgers.edu

- [1] K. Haule and G. Kotliar, *New Journal of Physics* **11**, 025021 (2009).
- [2] Z. Yin, K. Haule, and G. Kotliar, *Physical Review B* **86**, 195141 (2012).
- [3] L. Fanfarillo and E. Bascones, *Physical Review B* **92**, 075136 (2015).
- [4] H. T. Dang, J. Mravlje, A. Georges, and A. J. Millis, *Physical Review B* **91**, 195149 (2015).
- [5] X. Deng, K. Haule, and G. Kotliar, *Physical review letters* **116**, 256401 (2016).
- [6] Y. Wang, C.-J. Kang, H. Miao, and G. Kotliar, *Physical Review B* **102**, 161118 (2020).
- [7] L. de Medici, *Physical Review B* **83**, 205112 (2011).
- [8] A. Georges, L. de Medici, and J. Mravlje, *Annu. Rev. Condens. Matter Phys.* **4**, 137 (2013).
- [9] K. M. Stadler, G. Kotliar, A. Weichselbaum, and J. von Delft, *Annals of Physics* **405**, 365 (2019).
- [10] P. Werner, E. Gull, M. Troyer, and A. J. Millis, *Physical Review Letters* **101**, 166405 (2008).
- [11] C. Watzenböck, M. Edelmann, D. Springer, G. Sangiovanni, and A. Toschi, *Physical Review Letters* **125**, 086402 (2020).
- [12] P. Hansmann, R. Arita, A. Toschi, S. Sakai, G. Sangiovanni, and K. Held, *Phys. Rev. Lett.* **104**, 197002 (2010).
- [13] F. Hardy, A. Böhmer, D. Aoki, P. Burger, T. Wolf, P. Schweiss, R. Heid, P. Adelmann, Y. Yao, G. Kotliar, *et al.*, *Physical review letters* **111**, 027002 (2013).
- [14] F. Hardy, A. Böhmer, L. de'Medici, M. Capone, G. Giovannetti, R. Eder, L. Wang, M. He, T. Wolf, P. Schweiss, *et al.*, *Physical Review B* **94**, 205113 (2016).
- [15] R. Yang, Z. Yin, Y. Wang, Y. Dai, H. Miao, B. Xu, X. Qiu, and C. C. Homes, *Physical Review B* **96**, 201108 (2017).
- [16] G. R. Stewart, *Rev. Mod. Phys.* **83**, 1589 (2011).
- [17] H. Hosono, A. Yamamoto, H. Hiramatsu, and Y. Ma, *Materials today* **21**, 278 (2018).
- [18] C. M. Puetter and H.-Y. Kee, *EPL (Europhysics Letters)* **98**, 27010 (2012).
- [19] S. Hoshino and P. Werner, *Physical Review Letters* **115**, 247001 (2015).
- [20] O. Vafek and A. V. Chubukov, *Physical review letters* **118**, 087003 (2017).
- [21] A. K. Cheung and D. Agterberg, *Physical Review B* **99**, 024516 (2019).
- [22] P. Coleman, Y. Komijani, and E. J. König, *Physical Review Letters* **125**, 077001 (2020).
- [23] J. Schrieffer, *Journal of Applied Physics* **38**, 1143 (1967).
- [24] I. Okada and K. Yosida, *Progress of Theoretical Physics* **49**, 1483 (1973).
- [25] P. Nozières and A. Blandin, *Journal de Physique* **41**, 193 (1980).
- [26] C. Jayaprakash, H. Krishna-Murthy, and J. Wilkins, *Physical Review Letters* **47**, 737 (1981).
- [27] A. H. Nevidomskyy and P. Coleman, *Physical Review Letters* **103**, 147205 (2009).
- [28] V. Drouin-Touchette, E. J. König, Y. Komijani, and P. Coleman, *Physical Review B* **103**, 205147 (2021).
- [29] C. Aron and G. Kotliar, *Physical Review B* **91**, 041110 (R) (2015).
- [30] A. Horvat, R. Zitko, and J. Mravlje, *arXiv preprint arXiv:1907.07100* (2019).
- [31] Y. Wang, E. Walter, S. Lee, K. Stadler, J. von Delft, A. Weichselbaum, and G. Kotliar, *Physical Review Letters* **124**, 136406 (2020).
- [32] E. Walter, K. Stadler, S. Lee, Y. Wang, G. Kotliar, A. Weichselbaum, and J. von Delft, *Physical Review X* **10**, 031052 (2020).
- [33] Y. Komijani and P. Coleman, *Physical Review Letters* **120**, 157206 (2018).
- [34] J. Rech, P. Coleman, G. Zarand, and O. Parcollet, *Physical Review Letters* **96**, 016601 (2006).
- [35] Y. Komijani and P. Coleman, *Physical Review Letters* **122**, 217001 (2019).
- [36] J. Wang, Y.-Y. Chang, C.-Y. Mou, S. Kirchner, and C.-H. Chung, *Physical Review B* **102**, 115133 (2020).
- [37] B. Shen, Y. Zhang, Y. Komijani, M. Nicklas, R. Borth, A. Wang, Y. Chen, Z. Nie, R. Li, X. Lu, *et al.*, *Nature* **579**, 51 (2020).
- [38] Y. Komijani, *Phys. Rev. B* **101**, 235131 (2020).
- [39] J. Wang and Y.-f. Yang, *Physical Review B* **104**, 165120 (2021).
- [40] R. Han, D. Hu, J. Wang, and Y.-f. Yang, *Physical Review B* **104**, 245132 (2021).
- [41] Y. Ge and Y. Komijani, *arXiv preprint arXiv:2201.06681* (2022).
- [42] K. Stadler, Z. Yin, J. Von Delft, G. Kotliar, and A. Weichselbaum, *Physical review letters* **115**, 136401 (2015).
- [43] P. Coleman and C. Pépin, *Physical Review B* **68**, 220405 (2003).
- [44] P. Coleman and I. Paul, *Physical Review B* **71**, 035111 (2005).
- [45] P. Coleman, *Introduction to many-body physics* (Cambridge University Press, 2015).
- [46] P. Coleman, I. Paul, and J. Rech, *Physical Review B* **72**, 094430 (2005).
- [47] E. Lebanon, J. Rech, P. Coleman, and O. Parcollet, *Physical Review Letters* **97**, 106604 (2006).
- [48] See Supplemental Material at [link] for further information on the bare mixed valence problem, the derivation of spin and orbital susceptibilities, details on the single iteration approach, the scaling of the spin susceptibility and the quasi-power-law exponent.
- [49] K. M. Stadler, *A model study of strong correlations in Hund metals*, *Ph.D. thesis*, lmu (2019).
- [50] Another definition of $\hat{L}_\gamma = \sum L_\gamma^{mm'} X_{ab}^m X_{ba}^{m'}$ purely in terms of orbital charge can generically be written. For the antisymmetric L_γ matrices, this leads to an identically zero operator.
- [51] I. Affleck, A. W. W. Ludwig, H.-B. Pang, and D. L. Cox, *Phys. Rev. B* **45**, 7918 (1992).
- [52] I. Affleck and A. W. W. Ludwig, *Phys. Rev. B* **48**, 7297 (1993).
- [53] P. Coleman, L. Ioffe, and A. Tsvelik, *Physical Review B* **52**, 6611 (1995).
- [54] A. Georges and A. M. Sengupta, *Physical review letters* **74**, 2808 (1995).
- [55] W. Koller, A. C. Hewson, and D. Meyer, *Physical Review B* **72**, 045117 (2005).

- [56] T.-H. Lee, A. Chubukov, H. Miao, and G. Kotliar, *Physical review letters* **121**, 187003 (2018).
- [57] Y.-M. Wu, A. Abanov, and A. V. Chubukov, *Physical Review B* **99**, 014502 (2019).
- [58] O. Parcollet and A. Georges, *Physical Review Letters* **79**, 4665 (1997).
- [59] O. Parcollet, A. Georges, G. Kotliar, and A. Sengupta, *Physical Review B* **58**, 3794 (1998).

Supplementary materials on

”Interplay of charge and spin fluctuations in a Hund’s coupled impurity”

Victor Drouin-Touchette¹, Elio J. König², Yashar Komijani³, Piers Coleman^{1,4}

¹*Center for Materials Theory, Department of Physics and Astronomy, Rutgers University, Piscataway, NJ 08854 USA*

²*Max-Planck Institute for Solid State Research, 70569 Stuttgart, Germany*

³*Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221-0011, USA*

⁴*Department of Physics, Royal Holloway, University of London, Egham, Surrey TW20 0EX, UK*

These supplements contain: (I) details on the bare mixed valence problem in the absence of Hund’s coupling; (II) the derivation of the spin and orbital susceptibilities; (III) details on the single iteration approach to logarithmic corrections of the Kondo coupling; (IV) the consequence of the logarithmic approach to the Fermi liquid for the dynamical spin susceptibility and the obtained scaling; (V) the derivation of the quasi-power-law exponent γ .

I. INFINITE-U ANDERSON MODEL IN THE LARGE-N APPROACH

We here present further details on the black curve from Fig. 1 (c), since only once [S47] has the infinite-U Anderson model been studied in the large-N approach. In this context, the interorbital Hund’s coupling J_H is set to 0. While the self-energy equations are the same, the Dyson equations for the holons and spinons are

$$G_b(z)^{-1} = [z - \lambda - V^2 \Sigma_b(z)], \quad (\text{S1})$$

$$G_\chi(z)^{-1} = [z - \lambda - \epsilon_f - V^2 \Sigma_\chi(z)]. \quad (\text{S2})$$

We numerically solve these Dyson equations together with the self-energy self-consistent equations of Eq. (5), for a fixed bare hybridization width $\Gamma = \pi\rho V^2$ with $\rho = 1/(2D)$ the conduction electron’s density of states, and a fixed electronic bandwidth D .

For such an infinite-U Anderson-model, it is known [S47] that the characteristic Kondo temperature is given by

$$T_K^0 = D \left(\frac{\Gamma}{\pi D} \right)^k \exp \left\{ - \frac{\pi |\epsilon_f|}{\Gamma} \right\}, \quad (\text{S3})$$

where $k = q = 2S/N$ for this perfectly screened case. We show in Fig. (S1) the obtained Kondo temperature as a function of the holon energy level from numerically solving the large-N equations. We can see that, having defined the holon energy in the action as $\epsilon_f \chi_i^\dagger \chi_i$, $\epsilon_f > 0$ will result in a suppression of the holon occupation (red curve) and an enhancement of the spinon occupation (blue curve). That is the Kondo limit. In this limit, the Kondo temperature is very small because of the small holon occupation. This can be understood through as a Gutzwiller renormalization of the hybridization V due to finite holon occupancy $\tilde{V} \sim V \sqrt{\langle n_\chi \rangle}$ [S45]. Therefore, as $n_\chi \rightarrow 0$, the exponential in Eq. (S3) tends to 0 exponentially.

On the other side, as $\epsilon_f < 0$, the holons dominate over the spinons and the impurity is in the strongly mixed valence regime. There, the Fermi-liquid ground state is still reached, but it is one filled with Kondo singlets, which are easy to form with the large presence of holons, hence the large Kondo temperature. Since Finally, on Fig. (S1) (b), we see the reproduced trend of Eq. (S3), and the exponentially smaller Kondo temperature in the bare Kondo limit.

II. SUSCEPTIBILITIES AND HUBBARD OPERATORS

Well defined susceptibilities need to be expressed in terms of the Hubbard operators. These operators, which we use to capture the constraint of no double occupancy, are defined in terms of the Schwinger bosons b and holons χ :

$$\begin{aligned} X_{\alpha,a}^{(m)} &= b_{m\alpha}^\dagger \chi_{ma}, & X_{a,\alpha}^{(m)} &= \chi_{ma}^\dagger b_{m\alpha}, \\ X_{a,b}^{(m)} &= \chi_{ma}^\dagger \chi_{mb}, & X_{\alpha,\beta}^{(m)} &= b_{m\alpha}^\dagger b_{m\beta}. \end{aligned} \quad (\text{S4})$$

In [S9, S31, S32, S42, S49], the spin and orbital susceptibilities for the impurity are studied, and their behavior brings clarity to the concept of spin-orbital separation. We therefore wish to connect these observables with the large-N Schwinger boson method presented in this paper. These papers define the spin and orbital susceptibility as




FIG. S1. (a) Display of the variation in spinon and holon proportional occupation versus the holon energy level (q_B/q and q_χ/q respectively), with $q = 2S/N$ and $q_\xi = -(1/\pi) \int d\omega n_\xi(\omega) \text{Im} G_\xi(\omega + i\delta)$. We also plot the effective Kondo temperature, taken as the temperature at which the holon phase shift reaches $\delta_\chi > 0.98\pi$. (b) The same data for the Kondo temperature T_K , but in a logarithmic vertical axis to show the exponential spread of T_K (following Eq. (S3)).

$$\begin{aligned}\chi_{\text{sp}} &= \frac{1}{3} \sum_{\alpha} \langle \hat{S}^{\alpha} | | \hat{S}^{\alpha} \rangle_{\omega}, \\ \chi_{\text{orb}} &= \frac{1}{8} \sum_a \langle \hat{T}^a | | \hat{T}^a \rangle_{\omega}.\end{aligned}\tag{S5}$$

with $\hat{T}^a = \frac{1}{2} \sum_{nn'\sigma} d_{n\sigma}^\dagger \tau_{nn'}^a d_{n'\sigma}$, τ^a the 8 Gell-Mann matrices such that $\text{Tr}[\tau^a \tau^b] = 2\delta_{ab}$. Similarly, the spin operators are taken as $\hat{S}^\alpha = \frac{1}{2} \sum_{n\sigma\sigma'} d_{n\sigma}^\dagger \sigma_{\sigma\sigma'}^\alpha d_{n\sigma'}$, with $\text{Tr}[\sigma^\alpha \sigma^\beta] = 2\delta_{\alpha\beta}$. Their system has SU(2) spin symmetry and SU(3) orbital symmetry, hence the definition for the \hat{S}^α and \hat{T}^a operators.

The expression $\langle X | | X \rangle_{\omega}$ refers to the Fourier-transformed retarded correlation functions $-i\Theta(t)\langle [X(t), X(0)] \rangle$, with real frequency ω . In order to connect with these definitions, we need to define spin and orbital operators, and then obtain the susceptibility bubbles in the large-N limit.

Spin Susceptibility

Consider the total spin operator on the impurity to be

$$S_{\alpha\beta}(\tau) = \sum_m X_{\alpha\beta}^{(m)}(\tau) = \sum_m b_{m,\alpha}^\dagger(\tau) b_{m,\beta}(\tau),\tag{S6}$$

with m the orbital index, and $\alpha, \beta \in [1, \dots, N]$ are indices of SU(N). The total spin-spin susceptibility in imaginary time τ is then defined as

$$\begin{aligned}\chi_{\text{sp}}(\tau) &= \langle \vec{S}(\tau) \cdot \vec{S}(0) \rangle = \frac{1}{N^2} \frac{1}{3} \sum_{\alpha\beta} \langle S_{\alpha\beta}(\tau) S_{\beta\alpha}(0) \rangle \\ &= \frac{1}{N^2} \frac{1}{3} \sum_{\alpha\beta} \langle \left[\sum_m X_{\alpha\beta}^{(m)}(\tau) \right] \left[\sum_{m'} X_{\beta\alpha}^{(m')}(0) \right] \rangle = \frac{1}{N^2} \sum_m \sum_{\alpha\beta} \langle X_{\alpha\beta}^{(m)}(\tau) X_{\beta\alpha}^{(0)}(0) \rangle\end{aligned}\tag{S7}$$

where we used orbital invariance on the m index. We obtain

$$\chi_{\text{sp}}(\tau) = \frac{1}{N^2} \sum_m \sum_{\alpha\beta} \langle b_{m,\alpha}^\dagger(\tau) b_{m\beta}(\tau) b_{0\beta}^\dagger(0) b_{0,\alpha}(0) \rangle \quad (\text{S8})$$

$$= \frac{1}{N^2} \sum_m \sum_{\alpha\beta} \left[\langle b_{m,\alpha}^\dagger(\tau) b_{m\beta}(\tau) b_{0\beta}^\dagger(0) b_{0,\alpha}(0) \rangle + \langle b_{m,\alpha}^\dagger(\tau) b_{m\beta}(\tau) b_{0\beta}^\dagger(0) b_{0,\alpha}(0) \rangle \right] \quad (\text{S9})$$

$$= \frac{1}{N^2} \sum_m \sum_{\alpha\beta} \langle b_{m,\alpha}^\dagger(\tau) b_{0,\alpha}(0) \rangle \langle b_{m\beta}(\tau) b_{0\beta}^\dagger(0) \rangle \quad (\text{S10})$$

$$= \sum_m G_B(m, \tau) G_B(-m, -\tau) \quad (\text{S11})$$

with the definition that $G_B(m, \tau) = -\sum_{\alpha\beta} \langle T b_{m\alpha}(\tau) b_{0\beta}^\dagger(0) \rangle$. The Linked-Cluster theorem allows us to reject disconnected parts (first term that is crossed), which anyway lead to a $O(1/N)$ contribution to the susceptibility.

After converting to Matsubara frequencies and doing the analytic continuation $i\nu_p \rightarrow \omega + i\eta$, we obtain the finite frequency result

$$\chi_{\text{sp}}(\omega) = \sum_k \int \frac{d\omega'}{2\pi} n_B(\omega') G_B''(k, \omega' + i\eta) [G_B(k, \omega' - \omega - i\eta) + G_B(k, \omega' + \omega + i\eta)] \quad (\text{S12})$$

with the bosonic Green's functions $G_B(k, z) = [z - \lambda + 2\Delta \cos(k) - \Sigma_B(z)]^{-1}$ as defined in the main text. The dynamical spin susceptibility can therefore be computed as $\chi_{\text{sp}}''(\omega) = -\frac{1}{\pi} \text{Im} \chi_{\text{sp}}(\omega)$.

Orbital Susceptibility

The model we consider has $SO(3)$ symmetry, representing the angular momentum subspace of the t_{2g} ($L = 1$) orbitals. The $SO(3)$ group is represented using the iL_γ operators (which are themselves linear combinations of the $(\lambda_2, \lambda_5, \lambda_7)$ Gell-Mann matrices, as $SO(3)$ is a subgroup of $SU(3)$). The matrices acting in the 3×3 orbital site space representing the angular momenta are

$$L_x = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad L_y = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad L_z = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}. \quad (\text{S13})$$

so that the angular momentum operator in the γ direction, defined through the Hubbard operators, is

$$\hat{L}_\gamma = \frac{1}{NK} \sum_{mm'\alpha\alpha} X_{\alpha\alpha}^{(m)}(L_\gamma)_{mm'} X_{\alpha\alpha}^{(m')} = \frac{1}{NK} \sum_{mm'\alpha\alpha} b_{m,\alpha}^\dagger \chi_{m,a}(L_\gamma)_{mm'} \chi_{m',a}^\dagger b_{m',\alpha} \quad (\text{S14})$$

with a the channel index $a \in [1, \dots, K]$, and $\alpha \in [1, \dots, K]$ the $SU(N)$ spin index, and $X_{\alpha\alpha}^{(m)}$ ($X_{\alpha\alpha}^{(m)}$) being the Hubbard operator associated with creation (annihilation) of an impurity electron at site m , respectively. It is clear that acting \hat{L}_z on a filled state such as $\prod_{\alpha\alpha} b_{1,\alpha}^\dagger \chi_{1,a} |0\rangle$ leads to its m_z , in this case $m_z = 1$ (one impurity electronic state is filled at $m = 1$). We then define the orbital susceptibility as

$$\chi_{\text{orb}}(\tau) = \frac{1}{3} \sum_\gamma \langle \hat{L}_\gamma(\tau) \hat{L}_\gamma(0) \rangle \quad (\text{S15})$$

$$= \frac{1}{3} \frac{1}{N^2 K^2} \sum_{mm'nn'} \sum_{\alpha\beta ab} \langle X_{\alpha\alpha}^{(m)}(\tau) X_{\alpha\alpha}^{(m')}(\tau) X_{\beta\beta}^{(n)}(0) X_{\beta\beta}^{(n')}(0) \rangle \left[\sum_\gamma (L_\gamma)_{mm'} (L_\gamma)_{nn'} \right] \quad (\text{S16})$$

$$= \frac{1}{3} \sum_{mm'nn'} \left[\sum_\gamma (L_\gamma)_{mm'} (L_\gamma)_{nn'} \right] \mathcal{A}_{mm'}^{nn'}(\tau) \quad (\text{S17})$$

along the following definition:




FIG. S2. (a) The vertex for the Hubbard operators $X_{\alpha a}^{(m)}$ for sites m and m' . Dashed lines are holons propagators G_χ and wavy lines are spinons propagators G_b . (b-c) The four different Wick contractions that can be created from the vertex presented in (a). Each closed dashed loop carries a summation over a indices ($O(K)$ sum) while each closed wavy loop carries a summation over α indices ($O(N)$ sum). (b) is the disconnected part, and does not correspond to any dynamical behavior. Diagrams (c-d) have disconnected parts to them, but cannot be made fully disconnected without cutting through a propagator. Whereas (c) and (d) scale like $O(N^2K)$ and $O(K^2N)$ respectively, (d) scales like $O(NK)$, and we discard it in the large- N limit.

$$\mathcal{A}_{mm'}^{nn'}(\tau) = \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} X_{\alpha a}^{(m)}(\tau) X_{\alpha a}^{(m')}(\tau) X_{\beta b}^{(n)}(0) X_{\beta b}^{(n')}(0) \rangle \quad (\text{S18})$$

Vertices at time τ have one Hubbard operator entering and one leaving; this creates a complicated vertex for bosons and holons (see Fig. S2 (a)). There are four general diagrams one can create after Wick contracting the boson and holon lines (see Fig. S2 (b-e)), with (b) the truly disconnected one which can be disregarded through the Linked-Cluster theorem.

Thus, the calculation of the susceptibilities can include the 3 remaining objects (c-e) from figure S2. (c-d) are $O(1/K)$ and $O(1/N)$ contributions respectively, and (e) is $O(1/NK)$, seen by summing repeated indices for each bosonic and fermionic loop. The expression for the sum of these three remaining connected graphs is therefore, in the most general case,

$$\mathcal{A}_{mm'}^{nn'}(\tau) = \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} X_{\alpha a}^{(m)}(\tau) X_{\alpha a}^{(m')}(\tau) X_{\beta b}^{(n)}(0) X_{\beta b}^{(n')}(0) \rangle \quad (\text{S19})$$

$$= \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} b_{m,\alpha}^\dagger(\tau) \chi_{m,a}(\tau) \chi_{m'a}^\dagger(\tau) b_{m'\alpha}(\tau) b_{n\beta}^\dagger(0) \chi_{nb}(0) \chi_{n',b}^\dagger(0) b_{n',\beta}(0) \rangle \quad (\text{S20})$$

$$= \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} \overbrace{b_{m,\alpha}^\dagger(\tau) b_{m'\alpha}(\tau)} \overbrace{b_{n\beta}^\dagger(0) b_{n',\beta}(0)} \overbrace{\chi_{m,a}(\tau) \chi_{m'a}^\dagger(\tau)} \overbrace{\chi_{nb}(0) \chi_{n',b}^\dagger(0)} \rangle \quad (\text{S21})$$

$$+ \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} \overbrace{b_{m,\alpha}^\dagger(\tau) b_{m'\alpha}(\tau)} \overbrace{b_{n\beta}^\dagger(0) b_{n',\beta}(0)} \overbrace{\chi_{m,a}(\tau) \chi_{m'a}^\dagger(\tau)} \overbrace{\chi_{nb}(0) \chi_{n',b}^\dagger(0)} \rangle \quad (\text{S21})$$

$$+ \frac{1}{N^2K^2} \sum_{\alpha\beta ab} \langle \mathcal{T} \overbrace{b_{m,\alpha}^\dagger(\tau) b_{m'\alpha}(\tau)} \overbrace{b_{n\beta}^\dagger(0) b_{n',\beta}(0)} \overbrace{\chi_{m,a}(\tau) \chi_{m'a}^\dagger(\tau)} \overbrace{\chi_{nb}(0) \chi_{n',b}^\dagger(0)} \rangle \quad (\text{S21})$$

$$= -\frac{1}{K} G_B(m', m, 0^-) G_B(n', n, 0^-) G_h(m, n', \tau) G_h(n, m', -\tau) \quad (\text{S22})$$

$$+ \frac{1}{N} G_B(n', m, -\tau) G_B(m', n, \tau) G_h(m, m', 0^+) G_h(n, n', 0^+)$$

$$- \frac{1}{NK} G_B(n', m, -\tau) G_B(m', n, \tau) G_h(m, n', \tau) G_h(n, m', -\tau)$$

where the $1/N$ and $1/K$ factors come from $G_B = \frac{1}{N} \sum_\alpha G_{B,\alpha}(\tau) = -\frac{1}{N} \sum_\alpha \langle \mathcal{T} b_\alpha(\tau) b_\alpha^\dagger(0) \rangle$ and $G_{\alpha\beta} = \delta_{\alpha\beta} G_\alpha$. For

Eqs. (S21) and (S22), the three expressions correspond respectively to the diagrams (c) (d) and (e) from figure S2. Furthermore, we are using the notation that

$$G(m, n, \tau) = -\langle T \mathcal{O}_m(\tau) \mathcal{O}_n^\dagger(0) \rangle \quad (\text{S23})$$

with \mathcal{O} a bosonic or fermionic operator. Returning to the expression for the orbital susceptibility, we have that

$$\chi_{\text{orb}}^x(\tau) = \frac{1}{3} \left[-\mathcal{A}_{3,2}^{3,2}(\tau) - \mathcal{A}_{2,3}^{2,3}(\tau) + \mathcal{A}_{3,2}^{2,3}(\tau) + \mathcal{A}_{2,3}^{3,2}(\tau) \right] \quad (\text{S24})$$

$$\chi_{\text{orb}}^y(\tau) = \frac{1}{3} \left[-\mathcal{A}_{3,1}^{3,1}(\tau) - \mathcal{A}_{1,3}^{1,3}(\tau) + \mathcal{A}_{3,1}^{1,3}(\tau) + \mathcal{A}_{1,3}^{3,1}(\tau) \right] \quad (\text{S25})$$

$$\chi_{\text{orb}}^z(\tau) = \frac{1}{3} \left[-\mathcal{A}_{1,2}^{1,2}(\tau) - \mathcal{A}_{2,1}^{2,1}(\tau) + \mathcal{A}_{1,2}^{2,1}(\tau) + \mathcal{A}_{2,1}^{1,2}(\tau) \right] \quad (\text{S26})$$

In the case of $\chi_{\text{orb}}^x(\tau)$, we therefore need to evaluate $\mathcal{A}_{3,2}^{3,2}(\tau)$ and $\mathcal{A}_{2,3}^{2,3}(\tau)$. To do so, we first remark that since the term in the action for the holons is diagonal in orbital space ($S_\chi \propto \epsilon_f \sum_m \chi_{ma}^\dagger \chi_{ma}$), we have the holon's Green's function is diagonal in orbital space. Hence,

$$G_h(x, y, \tau) = -\frac{1}{K} \sum_a \langle T \chi_{x,a}(\tau) \chi_{y,a}^\dagger(0) \rangle = \delta_{xy} G_h(\tau). \quad (\text{S27})$$

Once we feed those results into the expression for $\mathcal{A}_{mm'}^{nn'}(\tau)$, we obtain

$$\begin{aligned} \mathcal{A}_{3,2}^{3,2}(\tau) &= -\frac{1}{K} G_B(2, 3, 0^-) G_B(2, 3, 0^-) \cancel{G_h(3, 2, \tau) G_h(2, 3, -\tau)} \\ &\quad + \frac{1}{N} G_B(2, 3, -\tau) G_B(2, 3, \tau) \cancel{G_h(3 - 2, 0^+) G_h(3, 2, 0^+)} \\ &\quad - \frac{1}{NK} G_B(2, 3, -\tau) G_B(2, 3, \tau) \cancel{G_h(3, 2, \tau) G_h(3, 2, -\tau)} \\ &= 0 = \mathcal{A}_{2,3}^{2,3}(\tau) \end{aligned} \quad (\text{S28})$$

and

$$\begin{aligned} \mathcal{A}_{3,2}^{2,3}(\tau) &= -\frac{1}{K} G_B(2, 3, 0^-) G_B(3, 2, 0^-) G_h(3, 3, \tau) G_h(2, 2, -\tau) \\ &\quad + \frac{1}{N} G_B(3, 3, -\tau) G_B(2, 2, \tau) \cancel{G_h(3, 2, 0^+) G_h(2, 3, 0^+)} \\ &\quad - \frac{1}{NK} G_B(3, 3, -\tau) G_B(2, 2, \tau) G_h(3, 3, \tau) G_h(2, 2, -\tau) \\ &= -\frac{1}{K} G_h(\tau) G_h(-\tau) \left[G_B(2, 3, 0^-) G_B(3, 2, 0^-) + \frac{1}{N} G_B(3, 3, -\tau) G_B(2, 2, \tau) \right] \end{aligned} \quad (\text{S29})$$

In the limit of large-N, only the first contribution will be relevant. The other contributions from the y and z sectors will lead to essentially the same results, with a change of index. Therefore, we identify

$$\chi_{\text{orb}}(\tau) = \sum_\gamma \chi_{\text{orb}}^\gamma = \frac{1}{K} \left(\frac{\Delta}{J_H} \right)^2 G_h(\tau) G_h(-\tau) \quad (\text{S30})$$

The constant factor of $\left(\frac{\Delta}{J_H} \right)^2$ in this expression can be obtained after Fourier transforming the bosonic Green's function:

$$\frac{1}{3} \sum_m G_B(m, m+1, 0^-) G_B(m+1, m, 0^-) = [G_B(k=0, 0^-) - G_B(k=\pm 2\pi/3, 0^-)]^2 \quad (\text{S31})$$

where $G_B(k, z) = [z - \lambda - 2\Delta \cos(k) - \Sigma_B(z)]^{-1}$. We have that

$$G_B(k=0, 0^-) - G_B(k=\pm 2\pi/3, 0^-) = \sum_k \int_{-\infty}^{\infty} \frac{d\omega}{\pi} n_B(\omega) \text{Im}[\cos(k) G_B(k, \omega + i\eta)] = \frac{\Delta}{J_H} \quad (\text{S32})$$

This is the same definition as the constraint on the Schwinger bosons, which leads to the Δ/J_H factor, and hence completes the expression.

After converting to Matsubara frequencies and doing the analytic continuation $i\omega_p \rightarrow \omega + i\eta$, we obtain the finite frequency result

$$\chi_{\text{orb}}(\omega) = \frac{1}{K} \left(\frac{\Delta}{J_H} \right)^2 \int \frac{d\omega'}{2\pi} n_F(\omega') G_h''(\omega' + i\eta) [G_h(\omega' - \omega - i\eta) + G_h(\omega' + \omega + i\eta)] \quad (\text{S33})$$

with the holonic Green's functions $G_h(z) = [z - \lambda - \epsilon_f - \Sigma_\chi(z)]^{-1}$ as defined in the main text. The dynamical orbital susceptibility can therefore be computed as $\chi_{\text{orb}}''(\omega) = -\frac{1}{\pi} \text{Im} \chi_{\text{orb}}(\omega)$. This concludes the derivation of the expressions presented in Eq. (7) and Fig. 3.

III. DETAILS ON SINGLE ITERATION APPROACH

In this section, we analytically evaluate a single self-energy correction to the holon's propagator, therefore obtaining the logarithmically dependent effective Kondo coupling, as mentioned in Eq. (9) in the main text.

We consider the single loop correction to the holon propagator,

$$G_{\chi,0}^{-1}(z) = z - \epsilon_f - \lambda, \quad (\text{S34})$$

$$G_{\chi,1}^{-1}(z) = G_{\chi,0}^{-1}(z) - V^2 \Sigma_{\chi,0}(z). \quad (\text{S35})$$

In a previous approach [S28], the evaluation of the first order self-energy at $z = 0 + i\delta$ led to an analytical expression for the effective Kondo temperature in presence of Hund's coupling. The argument there was that the holons have a pole that crosses from $\omega > 0$ to $\omega < 0$ at $T = T_K^{\text{eff}}$. Here, we expand this work to include the frequency dependence of the self-energy, based on other works on underscreened Kondo models [S43, S44].

A single iteration is conducted by estimating $\Sigma_{\chi,0}$ in a single loop, i.e. through the bare propagators for the spinons and conduction electrons. In imaginary time, this is written as

$$\Sigma_{\chi,0}(\tau) = g_{c,0}(-\tau) G_{B,0}(\tau) \quad (\text{S36})$$

where we use the following bare propagators in the presence of Hund's coupling.

$$g_{c,0}(z) = \int d\epsilon \frac{\rho(\epsilon)}{z - \epsilon}, \quad G_{B,0}(z) = \frac{1}{3} \left[(z - \lambda')^{-1} + 2(z - \lambda' - 3\Delta)^{-1} \right] \quad (\text{S37})$$

Passing from imaginary time to matsubara frequencies leads to

$$\Sigma_{\chi,0}(i\omega_m) = \frac{1}{3} \int d\epsilon \rho(\epsilon) \sum_n \frac{1}{i\omega_n - \epsilon} \left[\frac{1}{i\omega_n + i\omega_m - \lambda'} + \frac{2}{i\omega_n + i\omega_m - \lambda' - 3\Delta} \right] \quad (\text{S38})$$

which, after doing the Matsubara sum over fermionic frequencies, leads to

$$\Sigma_{\chi,0}(i\omega_m) = \frac{1}{3} \int d\epsilon \rho(\epsilon) \left[\frac{n_F(\epsilon_k) + n_B(\lambda')}{\epsilon - (\lambda' - i\omega_n)} + 2 \frac{n_F(\epsilon_k) + n_B(\lambda' + 3\Delta)}{\epsilon - (\lambda' + 3\Delta - i\omega_n)} \right] \quad (\text{S39})$$

We can then analytically continue this expression $i\omega_n \rightarrow \omega + i\delta$. Finally, progress can be made to evaluate analytically this integral. Using $\rho(\epsilon) = \rho\Phi(\epsilon)$ with $\rho = 1/2D$, such that $\Phi(\epsilon)D^2/(\omega^2 + D^2)$ for the conduction electrons' density of states (Φ acts as a soft cutoff at $\omega = D$, the bandwidth) simplifies the problem. Integrals of this sort are regularly found in Kondo problems [S45], and can be evaluated through the use of the digamma function $\Psi(z) = d \ln \Gamma(z) / dz$. The general result we use is that

$$\int d\omega' \Phi(\omega') \frac{n_F(\omega')}{\omega' - A} = \left[\Psi \left(\frac{1}{2} - \frac{A}{2\pi iT} \right) - \ln \left(\frac{D}{2\pi T} \right) \right] - \frac{\pi}{2} \frac{D}{iD - A} \quad (\text{S40})$$

Therefore, the full result is obtained:

$$\begin{aligned} \Sigma_{\chi,0}(\omega) = & -\frac{\rho}{3} \left(1 + \frac{2}{1 + \frac{\Delta^2}{D^2}}\right) \log\left(\frac{D}{2\pi T}\right) + \frac{\rho}{3} \left[\Psi\left(\frac{1}{2} + \frac{\lambda' - \omega}{2\pi iT}\right) + \frac{2}{1 + \frac{\Delta^2}{D^2}} \Psi\left(\frac{1}{2} + \frac{\lambda' + 3\Delta - \omega}{2\pi iT}\right)\right] \\ & + \frac{\pi\rho}{6} [\coth(-\beta\lambda'/2) + 2 \coth(-\beta(\lambda' + 3\Delta)/2)], \end{aligned} \quad (\text{S41})$$

and the single iteration result for the holon's Green's function is then

$$\begin{aligned} G_{\chi,1}^{-1}(\omega) = & \omega - \epsilon_f - \lambda' + 2\Delta \\ & + \frac{\rho V^2}{3} \left(1 + \frac{2}{1 + \frac{\Delta^2}{D^2}}\right) \log\left(\frac{D}{2\pi T}\right) - \frac{\rho V^2}{3} \left[\Psi\left(\frac{1}{2} + \frac{\lambda' - \omega}{2\pi iT}\right) + \frac{2}{1 + \frac{\Delta^2}{D^2}} \Psi\left(\frac{1}{2} + \frac{\lambda' + 3\Delta - \omega}{2\pi iT}\right)\right] \\ & - \frac{\pi\rho V^2}{6} [\coth(-\beta\lambda'/2) + 2 \coth(-\beta(\lambda' + 3\Delta)/2)], \end{aligned} \quad (\text{S42})$$

where $\lambda' = \lambda + 2\Delta$ is the chemical potential for the Schwinger bosons in the aligned state ($k = 0$). The unaligned state is gapped. The value of λ' is obtained by satisfying the constraint concerning the number of bosons. Quite generally, we will have that $n_B(\lambda) = q/\alpha$ with $\alpha > 1$ where $q = 2S/N$. $\alpha = 1$ corresponds to no Hund's coupling and no holons ($n_\chi = 0$). The presence of holons reduces the $T = 0$ spinon occupation and increases α . Therefore, as $T \rightarrow 0$, $\lambda' \rightarrow T \log(1 + \alpha/q)$.

Before exploring the explicit frequency dependence of the holon's Green's function, it is very useful to extract the universal temperature scale T_K^{eff} . The Kondo temperature, signaling the advent of the Nozières Fermi liquid at $T < T_K^{\text{eff}}$, is intimately related to a shift of the pole G_χ . As it was used before [S28], a useful tool to estimate T_K^{eff} is that

$$\text{Re } G_\chi^{-1}(z)|_{z=0+i\delta} = 0 \quad \text{at} \quad T = T_K^{\text{eff}} \quad (\text{S43})$$

In extracting the effective Kondo temperature, we are mainly concerned with two limits. Firstly, in the absence of Hund's coupling, we can set $\Delta = 0$. This is a conventional Anderson impurity problem, and an effective Kondo coupling $J_K \sim \frac{V^2}{\epsilon_f + \lambda}$ can be obtained by a Schrieffer Wolff transformation. The same J_K scale emerges out of this method. In this case, the holon's Green's function is simply

$$G_{\chi,1}^{-1}(\omega) = \omega - \epsilon_f - \lambda' + \rho V^2 \log\left(\frac{D}{2\pi T}\right) - \rho V^2 \Psi\left(\frac{1}{2} + \frac{\lambda' - \omega}{2\pi iT}\right) - \frac{\pi\rho V^2}{2} \coth(-\beta\lambda'/2), \quad (\text{S44})$$

Therefore, the Kondo temperature criteria of Eq. (S43) leads to the following equation

$$\frac{\epsilon_f + \lambda'}{\rho V^2} = \log\left(\frac{D}{2\pi T_K^{\text{eff}}}\right) - \text{Re } \Psi\left(\frac{1}{2} + \frac{\lambda'}{2\pi iT_K^{\text{eff}}}\right) - \frac{\pi}{2} \coth(-\beta\lambda'/2), \quad (\text{S45})$$

Quite generally, we will have that $n_B(\lambda) = q/\alpha$ with $\alpha > 1$ where $q = 2S/N$. $\alpha = 1$ corresponds to no Hund's coupling and no holons ($n_\chi = 0$). The presence of holons reduces the $T = 0$ spinon occupation and increases α . Therefore, we have $\lambda \rightarrow T \log(1 + \alpha/q)$. This simplifies the digamma term as $\log[\log(1 + \alpha/q)/2\pi]$. Furthermore, we approximate the digamma function such that $\text{Re } \Psi(z) \simeq \log(|z|)$ and simplifying Eq. (S45), we get

$$\frac{T_K^{\text{eff}}}{D} = \mathcal{F}(q) \exp\left[-\frac{\epsilon_f + \lambda}{\rho V^2}\right] \quad (\text{S46})$$

$$\mathcal{F}(q) = \frac{\exp\left[-\frac{\pi}{2} \coth(-\log(1 + \alpha/q)/2)\right]}{\log(1 + \alpha/q)} \quad (\text{S47})$$

where we recover an analogous form to the typical Kondo temperature $T_K/D \simeq \exp[-1/\rho J_K]$ with $J_K \sim \frac{V^2}{\epsilon_f + \lambda}$ as expected from a Schrieffer Wolff transformation.

In the case of large Hund's coupling, the spinon gap can be taken to be larger than the electronic bandwidth ($\Delta \gg D$), and all spinons will be in the aligned ground state. Furthermore, the chemical potential adjusts itself so that $\lambda' = \lambda - 2\Delta \simeq T \log(1 + \alpha/q)$. In this case, the holon's Green's function is

$$G_{\chi,1}^{-1}(\omega) = \omega - \epsilon_f - \lambda + \frac{\rho V^2}{3} \log\left(\frac{D}{2\pi T}\right) - \frac{\rho V^2}{3} \Psi\left(\frac{1}{2} + \frac{\lambda' - \omega}{2\pi iT}\right) - \frac{\pi\rho V^2}{6} [\coth(-\beta\lambda'/2) - 2], \quad (\text{S48})$$

Therefore, the Kondo temperature criteria of Eq. (S43) leads to

$$3 \frac{\epsilon_f + \lambda}{\rho V^2} = \log \left(\frac{D}{2\pi T_K^{\text{eff}}} \right) - \text{Re} \Psi \left(\frac{1}{2} + \frac{\log(1 + \alpha/q)}{2\pi i} \right) - \frac{\pi}{2} [\coth(-\log(1 + \alpha/q)/2) - 2], \quad (\text{S49})$$

Solving for T_K^{eff} , we obtain

$$\frac{T_K^{\text{eff}}}{D} = \mathcal{K}(q) \exp \left[-3 \frac{\epsilon_f + \lambda}{\rho V^2} \right] \quad (\text{S50})$$

$$\mathcal{K}(q) = \frac{\exp \left[-\frac{\pi}{2} \coth(-\log(1 + \alpha/q)/2) + \pi \right]}{\log(1 + \alpha/q)} \quad (\text{S51})$$

which is equivalent, generically, to the following form

$$\frac{T_K^{\text{eff,H}}}{D} \propto \left(\frac{T_K^{\text{eff,0}}}{D} \right)^3 \quad (\text{S52})$$

$$\frac{T_K^{\text{eff,0}}}{D} \propto \exp \left[-\frac{\epsilon_f + \lambda}{\rho V^2} \right] \quad (\text{S53})$$

with the proportionality factor being a function of q , comprising the functions $\mathcal{F}(q)$ and $\mathcal{K}(q)$. This is a simple recovery of both the effective Kondo temperature for an Anderson model [S45], as well as the Schrieffer effect for strong Hund's coupling [S23]. Therefore, we will generically associate

$$\frac{\epsilon_f + \lambda}{\rho V^2} = \frac{\mathcal{C}}{n} \log \left(\frac{D}{2\pi T_K^{\text{eff}}} \right) \quad (\text{S54})$$

with $n = 1$ for $J_H = 0$ (no Hund's coupling) and $n = 3$ for $J_H \gg D$ (strong Hund's coupling). \mathcal{C} is a $O(1)$ number depending on α/q .

We are now fully able to study the holon Green's function's full frequency dependence of Eq. S48. We consider frequencies $\omega > \max\{T, T_K^{\text{eff}}\}$. In that case, the digamma functions in Eq. (S48) can be approximated as $\Psi(a + \omega) \simeq \log(\omega)$ for large arguments ω . Furthermore, we take advantage of the relation of Eq. (S54), and are left with

$$G_{\chi,1}^{-1}(\omega) = \omega - \rho V^2 \left[\frac{\mathcal{C}}{n} \log \left(\frac{D}{2T_K^{\text{eff}}} \right) - \frac{1}{n} \log \left(\frac{D}{2\pi T} \right) + \frac{1}{n} \log \left(\frac{\omega}{2\pi T} \right) - \mathcal{D} \right], \quad (\text{S55})$$

for the general holonic Green's function at any temperature. Simplifying the logarithms, one finds that the holon's Green's function for the mixed valence problem, once the self-energy has been taken into account, is

$$G_{\chi,1}^{-1}(\omega) = \omega - \frac{\rho V^2}{n} \log \left(\frac{\omega}{T_K^{\text{eff}}} \right) + \mathcal{B}. \quad (\text{S56})$$

In the cases studied in this paper, the effective Kondo temperature is very small due to both the presence of holons in the system (T_K^0 is small), and because of strong Hund's coupling ($n = 3$).

Mapping to a Kondo problem

When studying small frequencies but nevertheless such that $\omega/T_K^{\text{eff}} > 1$, the logarithm in the above expression dominates the ω component. In this regime, we find that the physical holons χ^\dagger can be replaced by simple Grassmann fields $\bar{\chi} = V\chi^\dagger$, where we have absorbed the hybridization V into their definition. This is to connect to the Parcollet-Georges decoupling [S34, S46, S58, S59] of the Kondo interaction through the use of Grassmann variables, as we have used in Ref. [S28, S33]. In the infinite-U Anderson model, we have that the holon part of the action is [see Eq. (2)]

$$H_{\text{mv}} = \sum_m V (c_{0m\alpha\alpha}^\dagger \chi_{m,a}^\dagger b_{m\alpha} + h.c.) + (\epsilon_f + \lambda) \chi_{m,a}^\dagger \chi_{m,a} \quad (\text{S57})$$

where the effect of Eq. (S56) is to renormalize the holon energy level from $\epsilon_f + \lambda$ to $\epsilon^*(\omega) = \frac{\rho V^2}{n} \log\left(\frac{\omega}{T_K^{\text{eff}}}\right)$. In the Kondo model [S28], we had

$$H_{\text{kondo}} = \sum_m (c_{0m\alpha\alpha}^\dagger \bar{\chi}_{m,a} b_{m\alpha} + h.c.) + \frac{\bar{\chi}_{m,a} \chi_{m,a}}{J_K} \quad (\text{S58})$$

We can see that one can connect the two Hamiltonians with the association $\bar{\chi} = V\chi^\dagger$. This leads to an effective Kondo coupling

$$J_K^{\text{eff}} \simeq \frac{V^2}{\epsilon^*(\omega)} = \frac{n}{\rho \log\left(\frac{\omega}{T_K^{\text{eff}}}\right)}, \quad (\text{S59})$$

The new Kondo Green's function for our holons will be noted as $\tilde{G}_\chi(\omega) = -\langle T\chi(\tau)\bar{\chi}(0) \rangle$, in terms of the newly introduced Grassmann quantities. Because of the absorption of a factor of V into the definition of $\bar{\chi}$, we get $V^2\tilde{G}_\chi(\omega) = G_{\chi,1}(\omega)$. There is no dynamics (no ∂_τ term) in the action for these Grassmannian variables. Therefore, the ω in Eq. (S56) can be removed in the expression for \tilde{G}_χ . The final Kondo Green's function is

$$\tilde{G}_\chi(\omega) \simeq \left[-\frac{1}{J_K^{\text{eff}}} \right]^{-1} \quad (\text{S60})$$

This replacement is only valid in the limit where $\omega/T_K^{\text{eff}} > 1$ while $\max(T, \omega) < J_H$. In this regime, the nearly frozen charge degrees of freedom means that we can neglect the holon's dynamics and get a very simple instantaneous effective imaginary time Green's function

$$\tilde{G}_\chi(\tau) = -J_K^{\text{eff}} \delta(\tau), \quad (\text{S61})$$

This concludes the derivation of Eq. (9) in the main text. This result is further used in the next section for second order contribution to the dynamical spin susceptibility. For large frequencies $\omega = D$ one recovers the bare J_K^{eff} value of Eq. (S46).

IV. LOGARITHMIC CORRECTION TO THE SPIN SUSCEPTIBILITY

In this section, we consider whether the combination of the single iteration holon propagator, as derived above in Eq. (S59), with the spinon susceptibility bubble, is enough to obtain the correct scaling form for the dynamical spin susceptibility, as seen in Fig. 3. Former insight on this is provided by Ref. [S55], in which a running Kondo coupling is introduced in the context of underscreened Kondo models such that $\tilde{J}(\omega) = 1/\ln(\omega/T_0)$ with $T_0 \sim T_K$ the onset on underscreened features. Through a second order correction to the impurity spin dynamics as electrons are scattered on the impurity through the Kondo coupling J_K , this running coupling is then used to predict the spin susceptibility behavior of $\chi''(\omega) \sim [\omega \ln^2(\omega/T_K)]^{-1}$.

The question then becomes whether we obtain such a correction to χ_{sp} in this Schwinger boson formalism too. The previous section showed that a single iteration of the self-energy equations leads to a running effective Kondo coupling as the holon propagator. This section focuses on the use of this propagator to calculate the second order correction to spinon dynamics as they scatter into holons and conduction electrons.

We first consider the zeroth order contribution to the spin susceptibility. We use the result of Eq. (S12), and replace $G_B(k, z) \rightarrow G_{B,0}(k, z)$. Furthermore, we can consider both cases of strong Hund's coupling and no Hund's coupling in the same breath. In the case $J_H = 0$, $G_{B,0}(k, z) = G_{B,0}(z) = [z - \lambda]^{-1}$. The chemical potential λ is found by satisfying the constraint on boson number, i.e. we have that $n_B(\lambda) \simeq q = 2S/N$.

In the case of finite J_K , the bare spinon propagators are $G_{B,0}(k = 0, z) = [z - \lambda + 2\Delta]^{-1}$ and $G_{B,0}(k = \pm 2\pi/3, z) = [z - \lambda - \Delta]^{-1}$. The chemical potential adjusts itself to the gap in the spinon dispersion, and we have that $n_B(\lambda - 2\Delta) \simeq 3q$. Noting $\lambda' = \lambda - 2\Delta$, we therefore have $G_{B,0}(k = 0, z) = [z - \lambda']^{-1}$ and $G_{B,0}(k = \pm 2\pi/3, z) = [z - \lambda' - 3\Delta]^{-1}$. In the limit of large J_H and therefore large Δ , $G_{B,0}(k = \pm 2\pi/3, z) \rightarrow 0$. Therefore, the bare propagator to be used for the case of strong Hund's coupling and absent Hund's coupling has the same form: $G_{B,0}(z) = [z - \lambda]^{-1}$, where λ has to be adjusted for the context.




FIG. S3. (a) Full spinon susceptibility with the dressed propagators. (b) Zeroth order contribution to the spin susceptibility where only the bare spinon propagators are taken into account. (b) First order scattering of a spinon into the conduction sea through the Kondo effect. In the limit of $T_K^{\text{eff}} \ll \omega \ll T_{\text{orb}}$, due to strong Hund's coupling, the holon propagator can be replaced by the Kondo-like instantaneous propagator. This diagram then has a bubble correction from the conduction electrons. (c) Second order correction to the spinon susceptibility. Taking the same limit of an instantaneous holon propagator, we recover the rightmost diagram of (c), which is calculated explicitly in Eq. (S79) and is responsible for the logarithmic features in the dynamical spin susceptibility.

Zeroth order

The zeroth order is readily obtained.

$$\chi_{\text{sp},0}(i\nu_p) = T \sum_n G_{B,0}(i\nu_n) G_{B,0}(i\nu_n - i\nu_p) = T \sum_n \frac{1}{i\nu_n - \lambda} \frac{1}{i\nu_n - i\nu_p - \lambda} \quad (\text{S62})$$

$$= T \sum_n \frac{1}{(i\nu_n - \lambda)^2} \delta_{p,0} = \beta n_B(\lambda) (1 + n_B(\lambda)) \delta_{p,0} \quad (\text{S63})$$

Therefore there is no finite frequency impurity spin susceptibility absent hybridization with the electronic degrees of freedom. The $p = 0$ (zero frequency contribution) leads to the Curie-like behavior of the static susceptibility: $\chi_{\text{sp}} \simeq \beta n_B(\lambda) [1 + n_B(\lambda)]$.

First order

We then consider the first order correction to this through the addition of a self-energy bubble for the spinons, see Fig. S3 (c). In this section, we concern ourselves with the intermediate energy window where $T_K^{\text{eff}} < \omega < T_{\text{orb}}$, in which the results of the preceding section are applicable. In essence, instead of considering a full G_χ holon propagator, we will be replacing it such that $V^2 G_\chi = \tilde{G}_\chi$, i.e. a Kondo holon's Green's function. We use the instantaneous Green's function in imaginary time obtained in Eq. (S61).

Therefore, the first order correction to the spinon bubble, as shown in Fig. S3 (b), is

$$\chi_{\text{sp},1}(\tau) = G_{B,0}(-\tau) \int d\tau_1 d\tau_2 G_{B,0}(\tau_1) g_{c,0}(\tau_2 - \tau_1) \tilde{G}_\chi(\tau_2 - \tau_1) G_{B,0}(\tau - \tau_2) \quad (\text{S64})$$

$$= -J_K^{\text{eff}} G_{B,0}(-\tau) g_{c,0}(0^+) \int d\tau_1 d\tau_2 G_{B,0}(\tau_1) G_{B,0} g_{c,0}(\tau_2 - \tau_1) \delta(\tau_2 - \tau_1) (\tau - \tau_2) \quad (\text{S65})$$

$$(\text{S66})$$

In the next step, we need to use our form for a flat electronic density of states. We have that and the resultant propagator for the conduction electrons.

$$g_{c,0}(i\omega_n) = \rho \int_{-D}^D d\epsilon \frac{1}{i\omega_n - \epsilon} \quad (\text{S67})$$

where $\rho = \frac{1}{2D}$, and D the electron bandwidth. This leads to $\rho_{c,0}(\omega) = \rho$. Because of particle-hole symmetry and the fermionic commutation relation, we then get that $g_{c,0}(\tau = 0^+) = -\frac{1}{2}$. Therefore, after transforming to Matsubara frequencies, we obtain

$$\chi_{\text{sp},1}(i\nu_p) = (J_K^{\text{eff}})^2 T \sum_n [G_{B,0}(i\nu_n)]^2 G_{B,0}(i\nu_n - i\nu_p) \quad (\text{S68})$$

$$= (J_K^{\text{eff}})^2 T \sum_n \frac{1}{(i\nu_n - \lambda)^2} \frac{1}{i\nu_n - i\nu_p - \lambda} = \frac{(J_K^{\text{eff}})^2 n'_B(\lambda)}{i\nu_p} = \frac{\beta (J_K^{\text{eff}})^2 n_B(\lambda) [1 + n_B(\lambda)]}{i\nu_p} \quad (\text{S69})$$

After doing the analytic continuation $i\nu_p \rightarrow \omega + i\delta$, we see that taking the imaginary part leaves $\chi''_{\text{sp},1}(\omega) \propto \delta(\omega)$ - there is no dissipative feature at the first order.

Second order

Finally, we can proceed to the second order correction to the spin susceptibility, which is outlined in Fig. S3 (c). Again, we use the simplification of an instantaneous holon propagator after mapping the mixed valence problem onto an effective Kondo one: $\tilde{G}_\chi(\tau) \simeq -J_K^{\text{eff}} \delta(\tau)$. The second order susceptibility becomes

$$\chi_{\text{sp},2}(\tau) = (J_K^{\text{eff}})^2 G_{B,0}(-\tau) \int d\tau_1 d\tau_2 G_{B,0}(\tau_1) g_{c,0}(\tau_2 - \tau_1) g_{c,0}(\tau_1 - \tau_2) G_{B,0}(\tau_2 - \tau_1) G_{B,0}(\tau - \tau_2) \quad (\text{S70})$$

We then Fourier transform these expressions to Matsubara frequencies. We obtain

$$\begin{aligned} \chi_{\text{sp},2}(i\nu_p) &= (J_K^{\text{eff}})^2 T^3 \sum_{n_1, \dots, n_6} \left[T \int d\tau e^{(i\nu_p + i\nu_{n_1} - i\nu_{n_6})\tau} \right] \\ &\quad \times \left[T \int d\tau_1 e^{(-i\nu_{n_2} + i\omega_{n_3} - i\omega_{n_4} + i\nu_{n_5})\tau_1} \right] \left[T \int d\tau_2 e^{(-i\omega_{n_3} + i\omega_{n_4} - i\nu_{n_5} + i\nu_{n_6})\tau_2} \right] \\ &\quad \times G_{B,0}(i\nu_{n_1}) G_{B,0}(i\nu_{n_2}) g_{c,0}(i\omega_{n_3}) g_{c,0}(i\omega_{n_4}) G_{B,0}(i\nu_{n_5}) G_{B,0}(i\nu_{n_6}) \\ &= (\rho J_K^{\text{eff}})^2 \int d\epsilon d\epsilon' T^3 \sum_{n,m,q} \frac{1}{i\omega_n + i\omega_m + i\nu_p - i\nu_q - \lambda} \frac{1}{(i\omega_n + i\omega_m + i\nu_p - \lambda)^2} \frac{1}{i\nu_p - \lambda} \frac{1}{i\omega_n - \epsilon} \frac{1}{i\omega_m - \epsilon'} \end{aligned} \quad (\text{S71})$$

$$\quad (\text{S72})$$

where the integral form for the conduction electron's Green's function (Eq. (S67)) was included in the last line. The Matsubara sums were performed analytically on Mathematica. The result is

$$\chi_{\text{sp},2}(i\nu_p) = (\rho J_K^{\text{eff}})^2 \int d\epsilon d\epsilon' \left(\frac{\mathcal{F}_1(\epsilon, \epsilon', \lambda, i\nu_p)}{i\nu_p} + \frac{\mathcal{F}_2(\epsilon, \epsilon', \lambda, i\nu_p)}{\epsilon - \epsilon' - i\nu_p} \right) \quad (\text{S73})$$

We are only interested in this section in a Mathematica expression for the dynamical susceptibility, i.e. $\text{Im } \chi_{\text{sp}}(\omega)$. Therefore, we first analytically the bosonic Matsubara frequencies from the imaginary axis to the real axis: $i\nu_p \rightarrow \omega + i\delta$, and then take the imaginary part of that. We then obtain

$$\chi''_{\text{sp},2}(\omega) = (\rho J_K^{\text{eff}})^2 \int d\epsilon d\epsilon' \left(\mathcal{F}_1(\epsilon, \epsilon', \lambda) \delta(\omega) + \mathcal{F}_2(\epsilon, \epsilon', \lambda) \delta(\epsilon - \epsilon' - \omega) \right) \quad (\text{S74})$$

The factors \mathcal{F}_1 and \mathcal{F}_2 are obtained from the Matsubara sums. We simply state the result here

$$\begin{aligned} \mathcal{F}_1(\epsilon, \epsilon', \lambda) &= \frac{1}{\epsilon - \epsilon'} \left(n_B(\lambda)^2 n'_F(\epsilon) - [n_F(\epsilon') - n_F(\epsilon - \lambda)] [n_F(\epsilon) n'_B(\lambda - \epsilon + \epsilon') - n_B(\lambda - \epsilon + \epsilon') n'_F(\epsilon)] \right. \\ &\quad \left. - n_B(\lambda - \epsilon + \epsilon') n_F(\epsilon) n'_F(\epsilon') + n_B(\lambda) [n_F(\epsilon) n'_F(\epsilon) - n_B(\lambda - \epsilon + \epsilon') n'_F(\epsilon')] \right. \\ &\quad \left. + n_B(\lambda) [n_F(\epsilon - \lambda) (n'_B(\lambda - \epsilon + \epsilon') + n'_F(\epsilon)) - n_F(\epsilon) (n'_B(\lambda - \epsilon + \epsilon') + n'_F(\epsilon))] \right) \end{aligned} \quad (\text{S75})$$

$$\mathcal{F}_2(\epsilon, \epsilon', \lambda) = \frac{n_B(\lambda) - n_B(\lambda - \epsilon + \epsilon')}{(\epsilon - \epsilon')^2} [n_B(\lambda) + n_F(\epsilon)] [n_F(\epsilon') - n_F(\epsilon - \lambda)] \quad (\text{S76})$$

Clearly, the first term in Eq. (S74) leads to a contribution only at $\omega = 0$, and therefore does not contribute to the finite frequency behavior of the spin susceptibility. We then proceed to do the integral over ϵ , which picks up the Kronecker δ -function. We are therefore left with

$$\chi''_{\text{sp},2}(\omega) = (\rho J_K^{\text{eff}})^2 \frac{n_B(\lambda) - n_B(\lambda - \omega)}{\omega^2} \int d\epsilon' [n_B(\lambda) + n_F(\epsilon' + \omega)][n_F(\epsilon') - n_F(\epsilon' + \omega - \lambda)] \quad (\text{S77})$$

Taking the low temperature limit ($\omega/T \gg 1$), we find that $n_B(\lambda - \omega) \rightarrow -1$. Furthermore, the low-temperature substitution for the Fermi-Dirac distribution leads to

$$\begin{aligned} \chi''_{\text{sp},2}(\omega) &= (\rho J_K^{\text{eff}})^2 \frac{(1 + n_B(\lambda))}{\omega^2} \int d\epsilon' [n_B(\lambda) + \Theta(-\epsilon' - \omega)][\Theta(-\epsilon') - \Theta(-\epsilon' - \omega + \lambda)] \\ &\simeq \frac{(\rho J_K^{\text{eff}})^2 n_B(\lambda) [1 + n_B(\lambda)]}{\omega} \end{aligned} \quad (\text{S78})$$

since the integrand is only nonzero in the interval $[-\omega, 0]$, where its value is $n_B(\lambda)$. We can complete the analogy to the result of Eq. C7 in Ref. [S55] by noting that $n_B(\lambda) \simeq q = 2S/N$.

Furthermore, connecting the formula with the one obtained in Eq. (S59), we get that

$$\chi''_{\text{sp},2}(\omega) \simeq \frac{n_B(\lambda)(1 + n_B(\lambda))}{\omega \left[\log \left(\frac{\omega}{T_K^{\text{eff}}} \right) \right]^2} \quad (\text{S79})$$

This completes the derivation of Eq. (8) in the main text.

V. QUASI-POWER-LAW DERIVATION

In this section, we derive the argument given in the main text where the logarithmic corrections give rise to a quasi-power-law for a large frequency window of the intermediate regime. We start with $1/\chi''_{\text{sp}}(\omega)$ and then

$$1/\chi''_{\text{sp}}(\omega) = \omega \left[\log \left(\frac{\omega}{T_K^{\text{eff}}} \right) \right]^2 = \omega \left[\log \left(\frac{\omega}{T_{\text{orb}}} \right) - \log \left(\frac{T_K^{\text{eff}}}{T_{\text{orb}}} \right) \right]^2 \quad (\text{S80})$$

$$= \omega \left[-\log \left(\frac{T_K^{\text{eff}}}{T_{\text{orb}}} \right) \right]^2 \left[1 - \frac{\log \omega / T_{\text{orb}}}{\log T_K^{\text{eff}} / T_{\text{orb}}} \right]^2 \quad (\text{S81})$$

T_{orb} is the upper energy cutoff of the S.O.S. regime where this form applies to. It acts as a reference scale in the logarithm so that both $\omega/T_{\text{orb}}, T_K^{\text{eff}}/T_{\text{orb}} \in (0, 1)$. Therefore, in the final line, the quotient of logarithms is well defined, and for $\omega > T_K^{\text{eff}}$, $|\log \omega / T_{\text{orb}}| < |\log T_K^{\text{eff}} / T_{\text{orb}}|$. Since the quotient of logarithms in Eq. (S81) is $\ll 1$, we replace $(1 - x)^2 \sim \exp(-2x)$, and obtain

$$1/\chi''_{\text{sp}}(\omega) = \omega \left[-\log(T_K^{\text{eff}}/T_{\text{orb}}) \right]^2 \exp \left[-2 \frac{\log \omega / T_{\text{orb}}}{\log T_K^{\text{eff}} / T_{\text{orb}}} \right] \quad (\text{S82})$$

$$= \frac{\omega}{T_{\text{orb}}} \left[-\log(T_K^{\text{eff}}/T_{\text{orb}}) \right]^2 \omega^{-2/\log \left(\frac{T_K^{\text{eff}}}{T_{\text{orb}}} \right)} \quad (\text{S83})$$

By rearranging terms and only keeping ω dependent terms, we get

$$\chi''_{\text{sp}}(\omega) \sim \omega^{-\gamma} \quad (\text{S84})$$

$$\gamma = 1 - \frac{2}{\log \frac{T_K^{\text{eff}}}{T_{\text{orb}}}}, \quad (\text{S85})$$




FIG. S4. Value of the quasi-power-law exponent γ from Eq. (S85), extracted for the various impurity doping values n_{imp} presented in the main phase diagram and their corresponding ratios $T_K^{\text{eff}}/T_{\text{orb}}$.

which completes the derivation of Eq.(10) in the main text. We plot this relation in Fig. S4 for the different extracted T_K^{eff} and T_{orb} as a function of n_{imp} from our phase diagram [Fig. 1].

As one approaches $n_{\text{imp}} \sim 1.0$, the extent of the S.O.S. phase shrinks so dramatically that the assumption that $\omega \gg T_K^{\text{eff}}$ is no longer valid, leading to an ever increasing γ . As $n_{\text{imp}} \rightarrow 2.7$, γ shrinks and approaches 1.2.
