

Complete theory of radiative corrections to $K_{\ell 3}$ decays and the V_{us} update

Chien-Yeah Seng¹, Daniel Galviz^{1,2}, Mikhail Gorchtein^{3,4}, and Ulf-G. Meißner^{1,5,6}

¹*Helmholtz-Institut für Strahlen- und Kernphysik and Bethe Center for Theoretical Physics,
Universität Bonn, 53115 Bonn, Germany*

²*Yau Mathematical Sciences Center,
Tsinghua University, Beijing 100084, China*

³*Institut für Kernphysik, Johannes Gutenberg-Universität,
J.J. Becher-Weg 45, 55128 Mainz, Germany*

⁴*PRISMA Cluster of Excellence, Johannes Gutenberg-Universität, Mainz, Germany*

⁵*Institute for Advanced Simulation,
Institut für Kernphysik and Jülich Center for Hadron Physics,
Forschungszentrum Jülich, 52425 Jülich, Germany and*

⁶*Tbilisi State University, 0186 Tbilisi, Georgia*

(Dated: July 13, 2022)

Abstract

We fill up the missing piece in our own re-analysis of the long-distance electromagnetic radiative corrections to semileptonic kaon decays by performing a rigorous study in the $K \rightarrow \pi \mu^+ \nu_\mu(\gamma)$ channels. With appropriate experimental and lattice inputs, we achieve a precision level of 10^{-4} in these channels. This is comparable to our previous analysis in the $K \rightarrow \pi e^+ \nu_e(\gamma)$ channels. With this new result, we present an updated global analysis to extract the Standard Model parameter $|V_{us}|$ from semileptonic kaon decays. We obtain $|V_{us}| = 0.22308(55)$ and $0.22356(73)$, using the lattice average of the $K^0 \rightarrow \pi^-$ transition form factor at $N_f = 2+1+1$ and $N_f = 2+1$, respectively.

I. INTRODUCTION

A series of new applications of the classical Sirlin representation of the Standard Model (SM) radiative corrections (RCs) [1, 2] has recently brought our understanding of semileptonic decays of mesons, nucleons and nuclei to the next level. For instance, the implementation of a dispersion relation analysis on top of this framework has significantly improved the accuracy of the SM predictions of the free neutron [3–7] and nuclear beta decays [4, 8], leading to a more precise extraction of the top-row Cabibbo-Kobayashi-Maskawa (CKM) matrix element V_{ud} . These new developments unveiled an apparent violation of the top-row CKM unitarity [9], which provides one of the most promising evidences of the breakdown of precise SM theory predictions at low-energy precision experiments, alongside with the anomalies observed in the muon anomalous magnetic moment [10–14] and decays of B-mesons [15–18].

It was first thought that Sirlin’s representation is only useful for decay processes where the parent and daughter particles are nearly degenerate, because some of the more complicated hadronic matrix elements involving products of three electroweak currents vanish in this limit. However, some of the authors in this paper pointed out that this framework is equally applicable to semileptonic decay with large mass gaps, e.g. the kaon semileptonic decays ($K_{\ell 3}$), upon combining it with more modern techniques such as Chiral Perturbation Theory (ChPT) [19]. In a follow-up work [20] we outlined an efficient approach for lattice Quantum Chromodynamics (QCD) to help fixing the large theory uncertainties in the $K_{\ell 3}$ RC that originate from non-perturbative QCD, which was then partially put into practice [21]. With these new theory basis and lattice inputs, we improved the existing theory precision of the K_{e3} RC [22] by almost an order of magnitude [23, 24], and presented an updated global analysis to the value of the top-row CKM matrix element $|V_{us}|$ extracted from semileptonic kaon decays [25]. This new analysis supports the current top-row CKM unitarity violation, and further sharpens a previously-observed discrepancy between the values of $|V_{us}|$ obtained from $K_{\ell 3}$ and the leptonic kaon decay ($K_{\mu 2}$) [9].

This paper is a generalization of the aforementioned works to cover all channels in semileptonic kaon decays, in particular the $K_{\mu 3}$ channels. A reason why only the K_{e3} decays were previously focused on is that in these channels the effect of the (more) poorly-constrained RCs to the intrinsically non-forward charged weak form factor $f_-(t)$ to the decay rate is suppressed by $m_e^2/M_K^2 \sim 10^{-6}$ (on top of α/π), making them theoretically clean. In this work

we show that, with the precision goal of 10^{-4} , this can even be done for the $K_{\mu 3}$ channels by first separating the infrared (IR)-divergent and IR-finite corrections to $f_-(t)$, calculating the former precisely and matching the latter to a fixed-order ChPT expression. This simple prescription enables us to calculate the $K_{\mu 3}$ RCs to a same level of precision with their $K_{e 3}$ counterparts.

The content of this work is arranged as follows. In Section II we introduce the basic notations and formula for the $K_{\ell 3}$ decay rate. In Section III we outline the theory framework of the $\mathcal{O}(\alpha)$ virtual corrections, most of them are similar to Refs. [23, 24] except a novel and more rigorous treatment of the RCs to $f_-(t)$. In Section IV we discuss the bremsstrahlung contributions. The numerical results and error analysis are presented in Section V together with a new global analysis of $|V_{us}|$ from semileptonic kaon decays. In Section VI we state our final conclusions.

II. BASIC FORMALISM

This work is a straightforward extension of our previous $K_{e 3}$ calculation. It is rather meaningless to copy everything from the existing literature, so most of the time we simply refer the reader to the notations, formulas and descriptions in specific sections of Ref. [24]. Some trivial generalizations of notations, e.g. $e \rightarrow \ell$, are automatically implied if not stated otherwise.

First of all, we define $K_{\ell 3}$ (where $\ell = e, \mu$) as the fully-inclusive semileptonic decay process $K(p) \rightarrow \pi(p') + \ell^+(p_\ell) + \nu_\ell(p_\nu) + n\gamma$, with $n \geq 0$. At tree-level, the decay amplitude is given by:

$$M_0 = -\frac{G_F}{\sqrt{2}} \bar{u}_\nu \gamma^\mu (1 - \gamma_5) v_\ell F_\mu(p', p), \quad (1)$$

where

$$F_\mu(p', p) = \langle \pi(p') | (J_\lambda^W)^\dagger | K(p) \rangle = V_{us}^* [f_+^{K\pi}(t)(p + p')_\mu + f_-^{K\pi}(t)(p - p')_\mu] \quad (2)$$

defines the (real) charged weak form factors $f_\pm^{K\pi}(t)$ (for notational simplicity, from now on we will suppress the superscript $K\pi$ whenever it does not cause any confusion). The standard Mandelstam variables are defined as $s = (p' + p_\ell)^2$, $t = (p - p')^2$ and $u = (p - p_\ell)^2$, but for $n = 0$ only two kinematic variables are independent, which are often chosen as

$y = 2p \cdot p_\ell / M_K^2$ and $z = 2p \cdot p' / M_K^2$. The total tree-level decay rate is then given by:

$$(\Gamma_{K_{\ell 3}})_{\text{tree}} = \frac{M_K}{256\pi^3} \int_{\mathcal{D}_3} dydz G_F^2 |V_{us}|^2 \{ f_+^2 H(+1, +1) + 2f_+ f_- H(+1, -1) + f_-^2 H(-1, -1) \} , \quad (3)$$

where the integration region \mathcal{D}_3 is defined in Appendix A of Ref. [24], while the function $H(a, b)$ is defined in Eq. (2.7) of the same reference.

The full $\mathcal{O}(\alpha)$ electroweak RC to $K_{\ell 3}$ include one-loop virtual corrections at $n = 0$ and tree-level real (i.e. bremsstrahlung) corrections at $n = 1$. It is most conveniently expressed as a fractional correction to the tree-level decay rate:

$$\delta_{K_{\ell 3}} = \frac{\delta\Gamma_{K_{\ell 3}}}{(\Gamma_{K_{\ell 3}})_{\text{tree}}} . \quad (4)$$

However, one must be aware that in the existing ChPT treatment some of the short-distance electromagnetic corrections are not included in the definition of $\delta_{K_{\ell 3}}$, but rather redistributed into the charged weak form factors as well as the isospin-breaking corrections (see Sec. 8 of Ref. [24] for a discussion). After removing those terms, our result can be directly compared to the ‘‘long-distance’’ electromagnetic RC $\delta_{\text{EM}}^{K\ell}$ studied in the standard ChPT literature by further removing a short-distance, channel-independent electroweak correction factor:

$$\delta_{\text{EM}}^{K\ell} = \delta_{K_{\ell 3}} - (S_{\text{EW}} - 1) , \quad (5)$$

where $S_{\text{EW}} = 1.0232(3)_{\text{HO}}$ [26].

Below we discuss the virtual and real corrections separately.

III. VIRTUAL CORRECTIONS

Using on-shell relations, the virtual corrections to the amplitude M_0 can always be expressed in terms of a shift to f_\pm , i.e.

$$M_0 \rightarrow M_0 + \delta M_{\text{vir}}, \quad \text{with} \quad f_\pm(t) \rightarrow f_\pm(t) + \delta f_\pm(y, z) . \quad (6)$$

The corrections δf_\pm are generally complex functions of two variables y, z instead of a single variable t . The corresponding change in the decay rate is:

$$\begin{aligned} (\delta\Gamma_{K_{\ell 3}})_{\text{vir}} = & \frac{M_K}{256\pi^3} \int_{\mathcal{D}_3} dydz 2G_F^2 |V_{us}|^2 \{ H(+1, +1) f_+ \Re \delta f_+ \\ & + H(+1, -1) [f_+ \Re \delta f_- + f_- \Re \delta f_+] + H(-1, -1) f_- \Re \delta f_- \} . \end{aligned} \quad (7)$$

An important feature of the equation above is that both $H(+1, -1)$ and $H(-1, -1)$ contain an explicit factor $r_\ell \equiv m_\ell^2/M_K^2$ which suppresses the contribution of δf_- to the decay rate. In the K_{e3} channels, $r_e \sim 10^{-6}$ and δf_- is completely negligible, which is one of the basic assumptions of our previous work. However, even in the $K_{\mu 3}$ channels, the suppression $r_\mu \sim 0.04$ is still quite significant. Therefore, apart from the IR-singular pieces in δf_- that need to be rigorously calculated for the exact cancellation of IR-divergences, one is allowed to adopt an approximate representation for the remaining, IR-finite pieces without introducing a large error budget to the decay rate. This is the main spirit of this work.

The general structure of the virtual corrections to the decay amplitude can be summarized following Section 3 of Ref. [24] as

$$\delta M_{\text{vir}} = \delta M_{\text{I}} + (\delta M_2 + \delta M_{\gamma W}^a)_{\text{int}} + \delta M_{\gamma W}^b + \delta M_3. \quad (8)$$

The first term collects all universal, model-independent analytical pieces that originate from corrections sensitive to IR and UV physics. The second and third terms combine the remnants of the two-point function and the γW -box contributions upon accounting for the respective analytic contributions in the first term. The last term represents the contribution of the three-point function. As explained in Ref. [24], the full γW -box correction stems from the generalized Compton tensor $T^{\mu\nu}$ which contains a symmetric and an antisymmetric pieces. Correspondingly, the two contributions $\delta M_{\gamma W}^a$ and $\delta M_{\gamma W}^b$ are distinguished. Below, we discuss all different contributions to δf_\pm in Sirlin's representation in necessary detail.

A. Analytic pieces

Firstly, part of the loop integrals in the $\mathcal{O}(\alpha)$ electroweak RC can be calculated analytically (bearing errors of the order $\mathcal{O}(m_\ell^2/M_W^2)$) and the outcome is proportional to the tree-level amplitude,

$$(\delta f_\pm)_{\text{I}} = \left\{ \frac{\alpha}{2\pi} \left[\ln \frac{M_Z^2}{m_\ell^2} - \frac{1}{4} \ln \frac{M_W^2}{m_\ell^2} + \frac{1}{2} \ln \frac{m_\ell^2}{M_\gamma^2} - \frac{3}{8} + \frac{1}{2} \tilde{a}_g \right] + \frac{1}{2} \delta_{\text{HO}}^{\text{QED}} \right\} f_\pm(t), \quad (9)$$

see Eq. (3.11) and below of Ref. [24] for explanations of the notations. Throughout this work we introduce a fictitious photon mass M_γ to regularize the IR-divergence.

B. From $(\delta M_2 + \delta M_{\gamma W}^a)_{\text{int}} + \delta M_{\gamma W}^{b,V}$

Next we have the integrals $(\delta M_2 + \delta M_{\gamma W}^a)_{\text{int}} + \delta M_{\gamma W}^{b,V}$ defined in Eqs. (3.9) and (3.10) of Ref. [24]. Two relevant comments are in order:

- They give IR-divergent contributions to both δf_{\pm} .
- One can show that their contributions to δf_+ cannot depend on physics at large loop momentum; the same argument, however, does not apply to δf_- .

Therefore, we adopt the following strategy:

1. To study δf_+ from these integrals, we apply the same technique as in Sec. 4 of Ref. [24], to write

$$(\delta f_+)_{\text{this subsection}} = (\delta f_+)_{\text{II}} + (\delta f_+)_{\text{conv}}^{\text{fin}} + (\delta f_+)_{\text{Born-conv}} + (\delta f_+)_{\text{inel}} . \quad (10)$$

The calculable piece is the so-called ‘‘Born contribution’’ to the integral, and within it is the ‘‘convection term’’ contribution that is analytically calculable and contains the full IR-divergence. We split the latter into the IR divergent piece $(\delta f_+)_{\text{II}}$ and the IR-finite piece $(\delta f_+)_{\text{conv}}^{\text{fin}}$. The remaining ‘‘Born minus convection’’ contribution can be computed numerically making use of the π^- and K^+ electromagnetic form factors, for which we adopt a monopole parameterization [27, 28]. Finally, the systematic uncertainty due to the incalculable inelastic (or ‘‘non-Born’’) pieces is estimated using the resonance chiral theory [29–31], see Appendix B of Ref. [24] for details. We find that their corrections to $\delta_{K\ell 3}$ in all channels are at most 1×10^{-4} , so we simply assign a conservative uncertainty of 2×10^{-4} to them in each channel.

2. For δf_- , we do not attempt to calculate the full Born contribution because the convergence of the loop integrals depends on the ultraviolet (UV)-behavior of the form factors, which introduces extra model-dependence to the outcome. Instead, we compute only the convection term contribution (both the IR-divergent and IR-finite pieces, of which loop integrals are explicitly UV-finite) and leave the remaining, non-convection contribution as an incalculable piece at the moment:

$$(\delta f_-)_{\text{this subsection}} = (\delta f_-)_{\text{II}} + (\delta f_-)_{\text{conv}}^{\text{fin}} + (\delta f_-)_{\text{non-conv}} . \quad (11)$$

Below we provide the analytic expressions of the convection term contribution to δf_{\pm} from $(\delta M_2 + \delta M_{\gamma W}^a)_{\text{int}} + \delta M_{\gamma W}^{b,V}$:

$$\begin{aligned}
(\delta f_{\pm}^{K^0\pi^-})_{\text{II}} &= -\frac{\alpha}{4\pi} \left\{ -\frac{4p' \cdot p_{\ell} x_s}{M_{\pi} m_{\ell} (1-x_s^2)} \ln x_s \ln \left(\frac{M_{\gamma}^2}{M_{\pi} m_{\ell}} \right) f_{\pm}^{K^0\pi^-} \right. \\
&\quad \left. \pm \left(\frac{5}{2} - \ln \frac{M_{\pi}^2}{M_{\gamma}^2} \right) \left(\frac{p' \cdot (p+p')}{2M_{\pi}^2} f_{+}^{K^0\pi^-} + \frac{p' \cdot (p-p')}{2M_{\pi}^2} f_{-}^{K^0\pi^-} \right) \right\} \\
(\delta f_{\pm}^{K^+\pi^0})_{\text{II}} &= -\frac{\alpha}{4\pi} \left\{ \frac{4p \cdot p_{\ell} x_u}{M_K m_{\ell} (1-x_u^2)} \ln x_u \ln \left(\frac{M_{\gamma}^2}{M_K m_{\ell}} \right) f_{\pm}^{K^+\pi^0} \right. \\
&\quad \left. + \left(\frac{5}{2} - \ln \frac{M_K^2}{M_{\gamma}^2} \right) \left(\frac{p \cdot (p+p')}{2M_K^2} f_{+}^{K^+\pi^0} + \frac{p \cdot (p-p')}{2M_K^2} f_{-}^{K^+\pi^0} \right) \right\} \quad (12)
\end{aligned}$$

$$\begin{aligned}
(\delta f_{+}^{K^0\pi^-})_{\text{conv}}^{\text{fin}} &= -\frac{\alpha}{4\pi} \left\{ (C_{00}^{\text{fin}} + 4p' \cdot p_{\ell} C_0^{\text{fin}} + 2p' \cdot p_{\ell} C_1 - 2m_{\ell}^2 C_2) f_{+}^{K^0\pi^-} \right. \\
&\quad + \left(p' \cdot (p+p') f_{+}^{K^0\pi^-} + p' \cdot (p-p') f_{-}^{K^0\pi^-} \right) (C_1 + C_{11}/2) \\
&\quad - \left(p_{\ell} \cdot (p+p') f_{+}^{K^0\pi^-} + p_{\ell} \cdot (p-p') f_{-}^{K^0\pi^-} \right) C_{12}/2 \\
&\quad \left. + (p_{\ell} \cdot (p'-p) + m_{\ell}^2) \left(f_{+}^{K^0\pi^-} + f_{-}^{K^0\pi^-} \right) C_2 \right\} \\
(\delta f_{-}^{K^0\pi^-})_{\text{conv}}^{\text{fin}} &= -\frac{\alpha}{4\pi} \left\{ (C_{00}^{\text{fin}} + 4p' \cdot p_{\ell} C_0^{\text{fin}} + 2p' \cdot p_{\ell} C_1 - 2m_{\ell}^2 C_2) f_{-}^{K^0\pi^-} \right. \\
&\quad - \left(p' \cdot (p+p') f_{+}^{K^0\pi^-} + p' \cdot (p-p') f_{-}^{K^0\pi^-} \right) (C_1 + C_{11}/2 + 2C_2 + C_{12}) \\
&\quad + \left(p_{\ell} \cdot (p+p') f_{+}^{K^0\pi^-} + p_{\ell} \cdot (p-p') f_{-}^{K^0\pi^-} \right) (C_{12}/2 + C_{22}) \\
&\quad \left. + (p_{\ell} \cdot (p-3p') + 2p \cdot p' - 2M_{\pi}^2 - m_{\ell}^2) \left(f_{+}^{K^0\pi^-} + f_{-}^{K^0\pi^-} \right) C_2 \right\} \quad (13)
\end{aligned}$$

$$\begin{aligned}
(\delta f_{+}^{K^+\pi^0})_{\text{conv}}^{\text{fin}} &= -\frac{\alpha}{4\pi} \left\{ (C_{00}^{\text{fin}} - 4p \cdot p_{\ell} C_0^{\text{fin}} - 2p \cdot p_{\ell} C_1 - 2m_{\ell}^2 C_2) f_{+}^{K^+\pi^0} \right. \\
&\quad + \left(p \cdot (p+p') f_{+}^{K^+\pi^0} + p \cdot (p-p') f_{-}^{K^+\pi^0} \right) (C_1 + C_{11}/2) \\
&\quad + \left(p_{\ell} \cdot (p+p') f_{+}^{K^+\pi^0} + p_{\ell} \cdot (p-p') f_{-}^{K^+\pi^0} \right) C_{12}/2 \\
&\quad \left. + (p_{\ell} \cdot (p'-p) + m_{\ell}^2) \left(f_{+}^{K^+\pi^0} - f_{-}^{K^+\pi^0} \right) C_2 \right\} \\
(\delta f_{-}^{K^+\pi^0})_{\text{conv}}^{\text{fin}} &= -\frac{\alpha}{4\pi} \left\{ (C_{00}^{\text{fin}} - 4p \cdot p_{\ell} C_0^{\text{fin}} - 2p \cdot p_{\ell} C_1 - 2m_{\ell}^2 C_2) f_{-}^{K^+\pi^0} \right. \\
&\quad + \left(p \cdot (p+p') f_{+}^{K^+\pi^0} + p \cdot (p-p') f_{-}^{K^+\pi^0} \right) (C_1 + C_{11}/2 + 2C_2 + C_{12}) \\
&\quad + \left(p_{\ell} \cdot (p+p') f_{+}^{K^+\pi^0} + p_{\ell} \cdot (p-p') f_{-}^{K^+\pi^0} \right) (C_{12}/2 + C_{22}) \\
&\quad \left. + (p_{\ell} \cdot (p-3p') + 2p \cdot p' - 2M_{\pi}^2 - m_{\ell}^2) \left(f_{+}^{K^+\pi^0} - f_{-}^{K^+\pi^0} \right) C_2 \right\} \quad (14)
\end{aligned}$$

We refer the reader to Appendix C in Ref. [24] for the definitions of x_v and the C -functions; in $K_{\ell 3}^0$ their arguments are $m_1 = M_\pi$, $m_2 = m_\ell$, $v = s$, whereas in $K_{\ell 3}^+$ their arguments are $m_1 = M_K$, $m_2 = m_\ell$, $v = u$. We defer the discussion of their numerical impact on $\delta_{K_{\ell 3}}$ (as well as the ‘‘Born minus convection’’ contribution to δf_+) to Sections IV and V.

C. From the three-point function

Next we have the so-called ‘‘three-point function’’ correction to the form factors which we denote as $\delta f_{\pm,3}$, see Eq. (5.14) in Ref. [19]. The same reference provides an approximate expression for these quantities to $\mathcal{O}(e^2 p^2)$ in the chiral power counting (see Sec. 8 of that paper), but naïvely applying those expressions here will cause an incomplete cancellation of IR-divergences. Therefore, adopting the strategy in Sec. 5 of Ref. [24], we split the full $\delta f_{\pm,3}$ into the IR-divergent and the IR-finite pieces:

$$\delta f_{\pm,3} = (\delta f_{\pm})_{\text{III}} + (\delta f_{\pm,3})_{\text{fin}} . \quad (15)$$

Starting from its $\mathcal{O}(e^2 p^2)$ expression, one could resum the IR-divergent piece $(\delta f_{\pm})_{\text{III}}$ to all orders in the chiral expansion by simply adding back the full form factors and to ensure the exact cancellation of IR-divergences between the real and virtual corrections, as we pointed out in Sec. 5 of Ref. [24]. The resummed version reads:

$$\begin{aligned} (\delta f_{\pm}^{K^0\pi^-})_{\text{III}} &= \mp \frac{\alpha}{4\pi} \left(\ln \frac{M_\pi^2}{M_\gamma^2} - \frac{5}{2} \right) \frac{p' \cdot (p \mp p')}{2M_\pi^2} \left(f_+^{K^0\pi^-} + f_-^{K^0\pi^-} \right) \\ (\delta f_{\pm}^{K^+\pi^0})_{\text{III}} &= \pm \frac{\alpha}{4\pi} \left(\ln \frac{M_K^2}{M_\gamma^2} - \frac{5}{2} \right) \frac{p \cdot (p \mp p')}{2M_K^2} \left(f_+^{K^+\pi^0} - f_-^{K^+\pi^0} \right) . \end{aligned} \quad (16)$$

In the following, it is sufficient to adopt an $\mathcal{O}(e^2 p^2)$ approximation to $(\delta f_{\pm,3})_{\text{fin}}$, which could be inferred from Ref. [19]. However, there is one further complication as discussed in Sec. 8 of Ref. [24], namely: Following the standard ChPT treatment, some of the terms in $(\delta f_{\pm,3})_{\text{fin}}$ are in fact not counted as a part of the long-distance electromagnetic corrections, but rather redistributed into the extra t -dependence of the form factors (Eqs. (5.3), (5.7), (5.11), (5.12) in Ref. [32]) as well as the isospin-breaking correction $\delta_{\text{SU}(2)}^{K\pi}$ through the electromagnetically-induced $\pi^0 - \eta$ mixing (Eq. (5.5) in Ref. [32]). Therefore, in order to appropriately compare with the existing literature, we have to remove these terms from our

definitions of $(\delta f_{\pm,3})_{\text{fin}}$ as well. After doing so, we obtain the following $\mathcal{O}(e^2 p^2)$ expressions:

$$\begin{aligned}
(\delta f_{+,3}^{K^0\pi^-})_{\text{fin}}^{e^2 p^2} &= 0 \\
(\delta f_{+,3}^{K^+\pi^0})_{\text{fin}}^{e^2 p^2} &= 0 \\
(\delta f_{-,3}^{K^0\pi^-})_{\text{fin}}^{e^2 p^2} &= \frac{\alpha}{4\pi} \left[\frac{3}{4} \ln \frac{M_\pi^2}{\mu^2} - 1 \right] - \frac{Z\alpha}{8\pi} \ln \frac{M_\pi^2}{\mu^2} + \frac{8\pi\alpha}{3} (K_5^r + K_6^r) \\
(\delta f_{-,3}^{K^+\pi^0})_{\text{fin}}^{e^2 p^2} &= -\frac{\alpha}{4\sqrt{2}\pi} \left[\frac{3}{4} \ln \frac{M_K^2}{\mu^2} - 1 \right] - \frac{5Z\alpha}{8\sqrt{2}\pi} \ln \frac{M_K^2}{\mu^2} \\
&\quad + \frac{8\pi\alpha}{\sqrt{2}} \left(-2K_3^r + K_4^r + \frac{1}{3}K_5^r + \frac{1}{3}K_6^r \right), \tag{17}
\end{aligned}$$

where $Z \approx 0.8$ and K_i^r are low-energy constants (LECs) in ChPT, with μ the scale of dimensional regularization, which will be taken as $M_\rho = 770$ MeV in the numerical analysis.

D. From $\delta M_{\gamma W}^{b,A}$

Finally, we have the contribution from the axial γW -box amplitude $\delta M_{\gamma W}^{b,A}$, namely the axial charged weak current contribution to the loop integral $\delta M_{\gamma W}^b$ defined in Eq. (3.10) of Ref. [24]. In Sec. 6 of that paper, we outlined the strategy to fix its contribution to δf_+ , namely to make use of the existing lattice QCD calculations of the forward axial γW -box diagrams in the meson sector. In short, we can write:

$$(\delta f_+)_{\gamma W}^{b,A} = \left\{ \square_{\gamma W}^{V,A>} + \left[\square_{\gamma W}^{V,A<}(K, \pi, M_\pi) + \mathcal{O}\left(\frac{M_K^2}{\Lambda_\chi^2}\right) \right] \right\} f_+(t). \tag{18}$$

First, $\square_{\gamma W}^{V,A>} \approx 2.16 \times 10^{-3}$ is the channel-independent component originating from the integral of the forward box diagram at $Q^2 > 2 \text{ GeV}^2$, which was calculated to high precision from perturbative QCD. In the meantime, $\square_{\gamma W}^{V,A<}(K, \pi, M_\pi)$ is the remaining, channel-dependent part of the forward $K\pi$ box diagram calculated in the flavor SU(3) limit $M_K = M_\pi$. Existing lattice calculations are for $\square_{\gamma W}^{V,A<}(\pi^+, \pi^0, M_\pi)$ [33] and $\square_{\gamma W}^{V,A<}(K^0, \pi^-, M_\pi)$ [21], from which we can also obtain $\square_{\gamma W}^{V,A<}(K^+, \pi^0, M_\pi) = 2\square_{\gamma W}^{V,A<}(\pi^+, \pi^0, M_\pi) - \square_{\gamma W}^{V,A<}(K^0, \pi^-, M_\pi)$ through a ChPT matching [20]. The uncertainties due to the non-forward (NF) corrections are conservatively estimated by multiplying the central values of $\square_{\gamma W}^{V,A<}$ by the factor M_K^2/Λ_χ^2 , where $\Lambda = 4\pi F_\pi \approx 1.2 \text{ GeV}$ is the chiral symmetry breaking scale.

Unfortunately, these lattice calculations of the forward axial γW -box diagrams do not provide any useful information for $(\delta f_-)_{\gamma W}^{b,A}$ because $p - p' = 0$ by definition. So at this point we just leave it as another incalculable piece.

E. Matching the incalculable pieces in δf_- to ChPT

So far we have discussed the full $\mathcal{O}(\alpha)$ electroweak RC to δf_{\pm} . For δf_+ we have:

$$\delta f_+ = (\delta f_+)_{\text{I+II+III}} + (\delta f_+)_{\text{conv}}^{\text{fin}} + (\delta f_+)_{\text{Born-conv}} + (\delta f_+)_{\text{inel}} + (\delta f_{+,3})_{\text{fin}} + (\delta f_+)_{\gamma W}^{b,A} . \quad (19)$$

From the discussions above, we see that each term at the right-hand side can either be determined to high precision or is small enough to assign a controllable theory uncertainty.

Meanwhile, for δf_- we have:

$$\delta f_- = (\delta f_-)_{\text{I+II+III}} + (\delta f_-)_{\text{conv}}^{\text{fin}} + (\delta f_-)_{\text{non-conv}} + (\delta f_{-,3})_{\text{fin}} + (\delta f_-)_{\gamma W}^{b,A} , \quad (20)$$

among which $(\delta f_-)_{\text{I+II+III}}$ and $(\delta f_-)_{\text{conv}}^{\text{fin}}$ are exactly known, $(\delta f_{-,3})_{\text{fin}}$ is known to $\mathcal{O}(e^2 p^2)$, while $(\delta f_-)_{\text{non-conv}}$ and $(\delta f_-)_{\gamma W}^{b,A}$ are so far completely unknown. Fortunately, since they are IR-regular terms whose contributions to $\delta_{K\ell_3}$ are naturally suppressed by r_ℓ , there is a simple strategy to deal with them: Existing ChPT studies provided the $\mathcal{O}(e^2 p^2)$ expression for the full δf_- (i.e. the left-hand side of Eq. (20)) [32], and we can further take the $\mathcal{O}(e^2 p^2)$ approximation for $(\delta f_-)_{\text{I+II+III}}$ and $(\delta f_-)_{\text{conv}}^{\text{fin}}$ by replacing:

$$f_+^{K^0\pi^-}(t) \rightarrow -1 , \quad f_+^{K^+\pi^0}(t) \rightarrow -\frac{1}{\sqrt{2}} , \quad f_-(t) \rightarrow 0 . \quad (21)$$

By doing so, one could then equate both sides in Eq. (20) to extract the $\mathcal{O}(e^2 p^2)$ expression of $(\delta f_-)_{\text{non-conv}} + (\delta f_-)_{\gamma W}^{b,A}$. As a consequence, we arrive the following representation:

$$\delta f_- = (\delta f_-)_{\text{I+II+III}} + (\delta f_-)_{\text{conv}}^{\text{fin}} + (\delta f_-)_{\text{rem}} \quad (22)$$

where the first two terms at the right-hand side are exactly known, while the remaining piece $(\delta f_-)_{\text{rem}}$ adopts an $\mathcal{O}(e^2 p^2)$ approximation:

$$\begin{aligned} (\delta f_-^{K^0\pi^-})_{\text{rem}}^{e^2 p^2} &= -\frac{\alpha}{4\pi} \left[r_{0,1} \Lambda(s, M_\pi, m_\ell) + r_{0,2} \ln \frac{M_\pi^2}{m_\ell^2} + r_{0,3} \ln \frac{M_\pi^2}{\mu^2} + r_{0,4} \right] \\ (\delta f_-^{K^+\pi^0})_{\text{rem}}^{e^2 p^2} &= -\frac{\alpha}{4\sqrt{2}\pi} \left[r_{+,1} \Lambda(u, M_K, m_\ell) + r_{+,2} \ln \frac{M_K^2}{m_\ell^2} + r_{+,3} \ln \frac{M_K^2}{\mu^2} + r_{+,4} \right] , \quad (23) \end{aligned}$$

where the function $\Lambda(v, m_1, m_2)$ is defined in Eq. (C.5) of Ref. [24]. The coefficients $r_{i,j}$ are

given by:

$$\begin{aligned}
r_{0,1} &= -\frac{1}{8\lambda s} \left\{ M_K^2 [2s(m_\ell^2 - 2M_\pi^2) + (m_\ell^2 - M_\pi^2)^2 + 3s^2] - m_\ell^6 + m_\ell^4 [3M_\pi^2 + s] \right. \\
&\quad \left. - m_\ell^2 [3M_\pi^4 + 13M_\pi^2 s + s(t - 20s)] + M_\pi^6 + 12M_\pi^4 s + 7M_\pi^2 s^2 + st[M_\pi^2 - 5s] \right. \\
&\quad \left. - 20s^3 \right\} \\
r_{0,2} &= \frac{1}{16s^2} [M_K^2 (-m_\ell^2 + M_\pi^2 - 3s) + m_\ell^4 - 2m_\ell^2 M_\pi^2 + M_\pi^4 + 13M_\pi^2 s + 28s^2 + st] \\
r_{0,3} &= \frac{Z - 6}{2} \\
r_{0,4} &= 32\pi^2 [X_1 + X_2^r - X_3^r - (K_5^r + K_6^r)/3] + \frac{m_\ell^2 - M_K^2 - M_\pi^2}{8s} + \frac{3}{4} \tag{24}
\end{aligned}$$

and

$$\begin{aligned}
r_{+,1} &= \frac{1}{8\lambda u} \left\{ u^2 [7M_K^2 + 20m_\ell^2 + 3M_\pi^2 - 5t] + [M_K^2 - m_\ell^2]^2 [M_K^2 - m_\ell^2 + M_\pi^2] \right. \\
&\quad \left. + u [12M_K^4 + M_K^2 (-13m_\ell^2 - 4M_\pi^2 + t) + m_\ell^2 (m_\ell^2 + 2M_\pi^2 - t)] - 20u^3 \right\} \\
r_{+,2} &= -\frac{1}{16u^2} [(M_K^2 - m_\ell^2)(M_K^2 - m_\ell^2 + M_\pi^2) + u(13M_K^2 - 3M_\pi^2 + t) + 28u^2] \\
r_{+,3} &= \frac{5Z + 6}{2} \\
r_{+,4} &= 32\pi^2 [X_1 - X_2^r + X_3^r + 2K_3^r - K_4^r - (K_5^r + K_6^r)/3] + \frac{M_K^2 + M_\pi^2 - m_\ell^2}{8u} - \frac{3}{4}, \tag{25}
\end{aligned}$$

where $\lambda = \lambda(m_1^2, m_2^2, v)$ is the Källén function.

IV. REAL CORRECTIONS

Our previous treatment of the K_{e3} bremsstrahlung correction detailed in Sec. 7 of Ref. [24] can be directly applied to all $K_{\ell 3}$ channels without any further modification. During the bremsstrahlung, the real photon can be emitted either by the charged lepton or by the meson; in the latter case, the decay amplitude involves the generalized, non-forward Compton tensor $T_{\mu\nu}(q'; p', p)$ (see Eq. (3.5) in Ref. [24]). Our strategy is to split $T_{\mu\nu}$ into the convection term contribution and a remainder:

$$T^{\mu\nu} = T_{\text{conv}}^{\mu\nu} + [T^{\mu\nu} - T_{\text{conv}}^{\mu\nu}]. \tag{26}$$

In practice, we retain the full $T_{\text{conv}}^{\mu\nu}$ but take the $\mathcal{O}(p^2)$ approximation for $T^{\mu\nu} - T_{\text{conv}}^{\mu\nu}$; this simple implementation ensures exact electromagnetic gauge invariance. Consequently, we split the full tree-level $K \rightarrow \pi \ell^+ \nu_\ell \gamma$ amplitudes into two pieces:

$$M_{K \rightarrow \pi \ell^+ \nu_\ell \gamma} = M_A + M_B, \tag{27}$$

$\delta_{K_{\ell 3}}$	$(\delta f_+)_{\text{conv}}^{\text{fin}}$	$(\delta f_+)_{\text{Born-conv}}$	$(\delta f_+)_{\gamma W}^{b,A}$	(*) $(\delta f_-)_{\text{rem}}$
K_{e3}^0	-5.0	4.1	4.9(1) _{lat} (1) _{NF}	0.0
K_{e3}^+	9.6	0.1	6.4(1) _{lat} (4) _{NF}	0.0
$K_{\mu 3}^0$	18.6	0.4	4.9(1) _{lat} (1) _{NF}	0.3(2) _{LEC}
$K_{\mu 3}^+$	-0.1	1.0	6.5(1) _{lat} (4) _{NF}	-1.1(2) _{LEC}

Table I: Contribution to $\delta_{K_{\ell 3}}$ from $(\delta f_+)_{\text{conv}}^{\text{fin}}$, $(\delta f_+)_{\text{Born-conv}}$, $(\delta f_+)_{\gamma W}^{b,A}$ and $(\delta f_-)_{\text{rem}}$, in units of 10^{-3} .

$(\delta_{K_{\ell 3}})_{\text{I+II+III+brem}(\mathcal{D}_3)}$	(*) From $2\Re\{M_A^* M_B\} + M_B ^2$	Remainder
K_{e3}^0	1.0	24.1
K_{e3}^+	-0.3	4.4
$K_{\mu 3}^0$	0.6	13.4
$K_{\mu 3}^+$	-0.1	17.4

Table II: Sum of the IR-divergent one-loop contributions I, II, III and the bremsstrahlung contribution in the \mathcal{D}_3 region, in units of 10^{-3} .

where $T^{\mu\nu} - T_{\text{conv}}^{\mu\nu}$ is fully contained in M_B . Therefore, in calculating the squared amplitude, the result of $|M_A|^2$ is exact while $2\Re\{M_A^* M_B\} + |M_B|^2$ acquires a chiral expansion uncertainty.

The bremsstrahlung process resides at a larger allowed phase space area $\mathcal{D}_4 = \mathcal{D}_3 \oplus \mathcal{D}_{4-3}$ (see Appendix A of Ref. [24]). The IR-divergence is fully contained in $|M_A|^2$ integrating over the \mathcal{D}_3 region, and is exactly canceled by those in $(\delta f_{\pm})_{\text{I+II+III}}$ from the virtual corrections.

V. NUMERICAL RESULTS, ERRORS AND CORRELATIONS

Now we are ready to present the numerical results of $\delta_{K_{\ell 3}}$ in all channels. The IR-finite contribution from the virtual corrections is summarized in Table I, the combination of the IR-divergent parts in the virtual corrections and the bremsstrahlung corrections in the \mathcal{D}_3 region is given in Table II, and the bremsstrahlung contribution in the \mathcal{D}_{4-3} region is given in Table III. Columns in these three tables with an asterisk (*) denote quantities that have taken an $\mathcal{O}(e^2 p^2)$ approximation and are subject to chiral uncertainties at $\mathcal{O}(e^2 p^4)$.

$(\delta_{K\ell 3})_{\text{brem}(\mathcal{D}_{4-3})}$	(*) From $2\Re\{M_A^* M_B\} + M_B ^2$	From $ M_A ^2$
K_{e3}^0	0.2	5.6
K_{e3}^+	-0.1	5.3
$K_{\mu 3}^0$	0.1	0.2
$K_{\mu 3}^+$	0.0	0.1

Table III: The bremsstrahlung contribution in the \mathcal{D}_{4-3} region, in units of 10^{-3} .

For the error analysis we include only theory uncertainties that after a single rounding-up are of the size 1×10^{-4} or larger. After subtracting out $S_{\text{EW}} - 1$ and its associated short-distance error, there are five major sources of theory uncertainties:¹

1. The lattice uncertainty in $(\delta f_+)_{\gamma W}^{b,A}$,
2. The LEC uncertainties in $(\delta f_-)_{\text{rem}}$,
3. The NF uncertainty in $(\delta f_+)_{\gamma W}^{b,A}$,
4. The $\mathcal{O}(e^2 p^4)$ chiral uncertainty, and
5. The unknown $(\delta f_+)_{\text{inel}}$ in Eq.(10).

The first three are already displayed in Table I, but the last two not yet. In any case, in order to discuss later the average of $|V_{us}|$ over different channels, it is not sufficient to know just the uncertainties in each separate channel, but their correlations must also be understood. To do so one needs to first identify the independent theory inputs that carry the uncertainties, and then the correlation matrix can be calculated using the formula outlined in the Appendix of Ref. [25]. This can be rigorously done for the first two types, which we describe below:

- For $(\delta f_+)_{\gamma W}^{b,A}$, the independent lattice QCD inputs are [21, 33]:

$$\square_{\gamma W}^{VA<}(\pi^+, \pi^0, M_\pi) = 0.671(28)_{\text{lat}} \times 10^{-3} \quad , \quad \square_{\gamma W}^{VA<}(K^0, \pi^-, M_\pi) = 0.278(44)_{\text{lat}} \times 10^{-3} \quad . \quad (28)$$

¹ The uncertainty due to the kaon charge radius in K_{e3}^+ channel is around 4.8×10^{-5} . In Ref. [24], we somewhat exaggerated this uncertainty by first rounding it up to 5×10^{-5} and then displaying it as 1×10^{-4} . In this work we do not perform such two-time rounding up so this uncertainty is not taken into account anymore.

	$\delta_{\text{EM}}^{K\ell}$	ChPT
$K^0 e$	11.6(2) _{inel(1)lat(1)NF(2)$e^2 p^4$}	9.9(1.9) $e^2 p^4$ (1.1) _{LEC}
$K^+ e$	2.1(2) _{inel(1)lat(4)NF(1)$e^2 p^4$}	1.0(1.9) $e^2 p^4$ (1.6) _{LEC}
$K^0 \mu$	15.4(2) _{inel(1)lat(1)NF(2)LEC(2)$e^2 p^4$}	14.0(1.9) $e^2 p^4$ (1.1) _{LEC}
$K^+ \mu$	0.5(2) _{inel(1)lat(4)NF(2)LEC(2)$e^2 p^4$}	0.2(1.9) $e^2 p^4$ (1.6) _{LEC}

Table IV: Final result for $\delta_{\text{EM}}^{K\ell}$, in units of 10^{-3} . The ChPT result from Ref.[22] is given in the last column for comparison.

- For $(\delta f_-)_{\text{rem}}$, the independent combinations of LECs are X_1 , $C_1 \equiv X_2^r - X_3^r$, $C_2 \equiv 2K_3^r - K_4^r$ and $C_3 \equiv K_5^r + K_6^r$. Among them, $X_1 = -2.2(4) \times 10^{-3}$ was fixed to good precision with the recent lattice calculations [21], and its resulting uncertainty to $\delta_{K\ell 3}$ is negligible. Similar calculations are not yet done for C_{1-3} , so we infer their values at $\mu = M_\rho$ from resonance models [34–36], and assign a 100% uncertainty to each of them:

$$C_1 = -1.4(1.4)_{\text{LEC}} \times 10^{-3}, \quad C_2 = 4.0(4.0)_{\text{LEC}} \times 10^{-3}, \quad C_3 = 14.4(14.4)_{\text{LEC}} \times 10^{-3}. \quad (29)$$

Meanwhile, the next three uncertainties are estimated as follows:

- The NF uncertainty in $(\delta f_+)_{\gamma W}^{b,A}$ is estimated by multiplying $\square_{\gamma W}^{VA<}$ in each channel by M_K^2/Λ_χ^2 ;
- The $\mathcal{O}(e^2 p^4)$ chiral uncertainty is obtained by first adding all the columns with asterisks in Table I–III, and then multiply the sum by M_K^2/Λ_χ^2 ;
- Finally, a conservative uncertainty of 2×10^{-4} is assign to each channel to account for the poorly-constrained contribution from $(\delta f_+)_{\text{inel}}$ (see discussions in Sec.III B).

Unlike the first two, these three errors are deduced using naïve power counting and order-of-magnitude estimations, and it is difficult to identify independent sources of uncertainties within each type. In fact, we consider it as arbitrary to take these uncertainties to be uncorrelated as to assume any correlation. Therefore, we simply take them to be uncorrelated, following the same strategy adopted by some of us in Ref. [25].

	$ V_{us}f_+^{K^0\pi^-}(0) $	Correlation Matrix
$K_L e$	$0.21617(46)_{\text{exp}}(10)_{I_K}(4)_{\delta_{\text{EM}}}$	1 0.021 0.025 0.567 0.004 0.017
$K_S e$	$0.21530(122)_{\text{exp}}(10)_{I_K}(4)_{\delta_{\text{EM}}}$	1 0.009 0.014 0.000 0.006
$K^+ e$	$0.21714(88)_{\text{exp}}(10)_{I_K}(21)_{\delta_{\text{SU}(2)}}(5)_{\delta_{\text{EM}}}$	1 0.018 0.002 0.894
$K_L \mu$	$0.21649(50)_{\text{exp}}(16)_{I_K}(4)_{\delta_{\text{EM}}}$	1 0.011 0.044
$K_S \mu$	$0.21251(466)_{\text{exp}}(16)_{I_K}(4)_{\delta_{\text{EM}}}$	1 0.005
$K^+ \mu$	$0.21699(108)_{\text{exp}}(16)_{I_K}(21)_{\delta_{\text{SU}(2)}}(6)_{\delta_{\text{EM}}}$	1
Average: Ke	$0.21626(40)_K(3)_{\text{HO}}$	
Average: $K\mu$	$0.21654(48)_K(3)_{\text{HO}}$	
Average: tot	$0.21634(38)_K(3)_{\text{HO}}$	

Table V: Updated determination of $|V_{us}f_+^{K^0\pi^-}(0)|$, which should be compared to Table I in Ref.[25].

We present our final result of $\delta_{\text{EM}}^{K\ell}$ in Table IV, with the correlation matrix given by:

$$\text{Corr}(\delta_{\text{EM}}) = \begin{pmatrix} 1 & -0.050 & 0.069 & -0.043 \\ & 1 & -0.049 & 0.079 \\ & & 1 & 0.088 \\ & & & 1 \end{pmatrix}, \quad (30)$$

where $\delta_{\text{EM}} = \left(\delta_{\text{EM}}^{K^0e} \delta_{\text{EM}}^{K^+e} \delta_{\text{EM}}^{K^0\mu} \delta_{\text{EM}}^{K^+\mu} \right)^T$. The results in the two Ke channels are obviously the same as in Ref. [24]. As a comparison, we also quote the ChPT result from Ref. [22]. Our new determinations agree with them within error bars in all four channels, but with significant improvements in precision by almost an order of magnitude.

With these new theory inputs of $\delta_{\text{EM}}^{K\ell}$, we also present here an updated global analysis of $|V_{us}f_+^{K^0\pi^-}(0)|$ from semileptonic kaon decays. Essentially, we take all the theory and experimental inputs quoted in Ref. [25], and only replace the values of δ_{EM} and $\text{Corr}(\delta_{\text{EM}})$ by our new results. The outcome is summarized in Table V, which has only very minor changes comparing to Table I in Ref. [25]. Supplementing the result with the most recent FLAG averages of $|f_+^{K^0\pi^-}(0)|$ with different numbers of active quark flavors [37]:

$$|f_+^{K^0\pi^-}(0)| = \begin{cases} 0.9698(17) & N_f = 2 + 1 + 1 & \text{Refs. [38, 39]} \\ 0.9677(27) & N_f = 2 + 1 & \text{Refs. [40, 41]} \end{cases}, \quad (31)$$

we obtain:

$$|V_{us}|_{K_{\ell 3}} = \begin{cases} 0.22308(39)_{\text{lat}}(39)_K(3)_{\text{HO}} & N_f = 2 + 1 + 1 \\ 0.22356(62)_{\text{lat}}(39)_K(3)_{\text{HO}} & N_f = 2 + 1 \end{cases} \quad (32)$$

respectively.

The value of $|V_{us}|$ can also be extracted by first obtaining $|V_{us}/V_{ud}|$ from the ratio between the leptonic kaon ($K_{\mu 2}$) and pion ($\pi_{\mu 2}$) decay rates, and then substituting $|V_{ud}|$ by its most precise determination from superallowed nuclear decays [42]. With this method, we obtain [9]:

$$|V_{us}|_{K_{\mu 2}} = \begin{cases} 0.2252(5) & N_f = 2 + 1 + 1 \\ 0.2255(8) & N_f = 2 + 1 \end{cases} . \quad (33)$$

Comparing Eqs. (32) and (33), we see that the so-called $K_{\ell 3} - K_{\mu 2}$ discrepancy in the $|V_{us}|$ determination is still very much alive.

VI. CONCLUSION

In this work we present for the first time a unified analysis of the long-distance electromagnetic RC to all channels in $K_{\ell 3}$ decays based on the novel hybrid method combining Sirlin's representation with ChPT, that we developed a few years ago. With an appropriate use of the electromagnetic/charged weak form factors, lattice QCD inputs of the forward axial γW -box as well as the existing knowledge of the remaining, poorly-constrained LECs, we are able to control the theory uncertainties in all four independent channels in $K_{\ell 3}$ at the level of 10^{-4} .

Several important implications of this new calculation are as follows. First, no significant change of $\delta_{\text{EM}}^{K\ell}$ from the previous ChPT result is observed; this allows us to state with more confidence that the outstanding discrepancy between the $|V_{us}|$ extracted from $K_{\mu 2}/\pi_{\mu 2}$ and $K_{\ell 3}$ does not originate from the long-distance electromagnetic corrections. Second, separate global analysis of $|V_{us}f_+^{K^0\pi^-}(0)|$ from Ke and $K\mu$ channels show no significant difference, which supports the SM prediction of lepton flavor universality. Third, the further reduction of theory uncertainties provides an even stronger motivation for next-generation experiments to measure the kaon lifetimes and $K_{\ell 3}$ branching ratios with higher precision. From the theory side, the dominant sources of uncertainty are now the phase space factor I_K and the isospin-breaking correction $\delta_{\text{SU}(2)}$, which await new breakthroughs from phenomenology and

lattice QCD.

Finally, we stress that every important theory input should always be cross-checked with several independent methods. In this sense, the existing plan to compute the full (virtual and real) $K_{\ell 3}$ RC directly on the lattice [43] is very much anticipated.

Acknowledgements

This work is supported in part by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) and the NSFC through the funds provided to the Sino-German Collaborative Research Center TRR110 “Symmetries and the Emergence of Structure in QCD” (DFG Project-ID 196253076 - TRR 110, NSFC Grant No. 12070131001) (U-G.M and C.Y.S), by the Chinese Academy of Sciences (CAS) through a President’s International Fellowship Initiative (PIFI) (Grant No. 2018DM0034) and by the VolkswagenStiftung (Grant No. 93562) (U-G.M), by EU Horizon 2020 research and innovation programme, STRONG-2020 project under grant agreement No 824093, by the German-Mexican research collaboration Grant No. 278017 (CONACyT) and No. SP 778/4-1 (DFG), and partially by the DFG personal grant No GO 2604/3-1 (M.G).

-
- [1] A. Sirlin, *Current Algebra Formulation of Radiative Corrections in Gauge Theories and the Universality of the Weak Interactions*, *Rev. Mod. Phys.* **50** (1978) 573. [Erratum: *Rev. Mod. Phys.* 50,905(1978)].
 - [2] C.-Y. Seng, *Radiative Corrections to Semileptonic Beta Decays: Progress and Challenges*, *Particles* **4** (2021), no. 4 397–467 [2108.03279].
 - [3] C.-Y. Seng, M. Gorchtein, H. H. Patel and M. J. Ramsey-Musolf, *Reduced Hadronic Uncertainty in the Determination of V_{ud}* , *Phys. Rev. Lett.* **121** (2018), no. 24 241804 [1807.10197].
 - [4] C. Y. Seng, M. Gorchtein and M. J. Ramsey-Musolf, *Dispersive evaluation of the inner radiative correction in neutron and nuclear β decay*, *Phys. Rev.* **D100** (2019), no. 1 013001 [1812.03352].
 - [5] C.-Y. Seng, X. Feng, M. Gorchtein and L.-C. Jin, *Joint lattice QCD–dispersion theory*

- analysis confirms the quark-mixing top-row unitarity deficit, *Phys. Rev. D* **101** (2020), no. 11 111301 [2003.11264].
- [6] K. Shiells, P. G. Blunden and W. Melnitchouk, *Electroweak axial structure functions and improved extraction of the V_{ud} CKM matrix element*, *Phys. Rev. D* **104** (2021), no. 3 033003 [2012.01580].
- [7] M. Gorchtein and C.-Y. Seng, *Dispersion relation analysis of the radiative corrections to g_A in the neutron β -decay*, *JHEP* **10** (2021) 053 [2106.09185].
- [8] M. Gorchtein, *γW Box Inside Out: Nuclear Polarizabilities Distort the Beta Decay Spectrum*, *Phys. Rev. Lett.* **123** (2019), no. 4 042503 [1812.04229].
- [9] **Particle Data Group** Collaboration, P. Zyla *et. al.*, *Review of Particle Physics*, *PTEP* **2020** (2020), no. 8 083C01.
- [10] **Muon $g-2$** Collaboration, B. Abi *et. al.*, *Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm*, *Phys. Rev. Lett.* **126** (2021) 141801.
- [11] T. Aoyama *et. al.*, *The anomalous magnetic moment of the muon in the Standard Model*, *Phys. Rept.* **887** (2020) 1–166 [2006.04822].
- [12] J. P. Miller, E. de Rafael and B. L. Roberts, *Muon ($g-2$): Experiment and theory*, *Rept. Prog. Phys.* **70** (2007) 795 [hep-ph/0703049].
- [13] J. P. Miller, E. de Rafael, B. L. Roberts and D. Stöckinger, *Muon ($g-2$): Experiment and Theory*, *Ann. Rev. Nucl. Part. Sci.* **62** (2012) 237–264.
- [14] F. Jegerlehner and A. Nyffeler, *The Muon $g-2$* , *Phys. Rept.* **477** (2009) 1–110 [0902.3360].
- [15] **LHCb** Collaboration, R. Aaij *et. al.*, *Search for lepton-universality violation in $B^+ \rightarrow K^+ \ell^+ \ell^-$ decays*, *Phys. Rev. Lett.* **122** (2019), no. 19 191801 [1903.09252].
- [16] **LHCb** Collaboration, R. Aaij *et. al.*, *Test of lepton universality using $B^+ \rightarrow K^+ \ell^+ \ell^-$ decays*, *Phys. Rev. Lett.* **113** (2014) 151601 [1406.6482].
- [17] **LHCb** Collaboration, R. Aaij *et. al.*, *Measurement of the ratio of branching fractions $\mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \tau^- \bar{\nu}_\tau) / \mathcal{B}(\bar{B}^0 \rightarrow D^{*+} \mu^- \bar{\nu}_\mu)$* , *Phys. Rev. Lett.* **115** (2015), no. 11 111803 [1506.08614]. [Erratum: *Phys.Rev.Lett.* 115, 159901 (2015)].
- [18] **LHCb** Collaboration, R. Aaij *et. al.*, *Angular analysis of the $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ decay using 3 fb^{-1} of integrated luminosity*, *JHEP* **02** (2016) 104 [1512.04442].
- [19] C.-Y. Seng, D. Galviz and U.-G. Meißner, *A New Theory Framework for the Electroweak Radiative Corrections in K_{l3} Decays*, *JHEP* **02** (2020) 069 [1910.13208].

- [20] C.-Y. Seng, X. Feng, M. Gorchtein, L.-C. Jin and U.-G. Meißner, *New method for calculating electromagnetic effects in semileptonic beta-decays of mesons*, *JHEP* **10** (2020) 179 [2009.00459].
- [21] P.-X. Ma, X. Feng, M. Gorchtein, L.-C. Jin and C.-Y. Seng, *Lattice QCD calculation of the electroweak box diagrams for the kaon semileptonic decays*, *Phys. Rev. D* **103** (2021) 114503 [2102.12048].
- [22] V. Cirigliano, M. Giannotti and H. Neufeld, *Electromagnetic effects in $K(l3)$ decays*, *JHEP* **11** (2008) 006 [0807.4507].
- [23] C.-Y. Seng, D. Galviz, M. Gorchtein and U. G. Meißner, *High-precision determination of the $Ke3$ radiative corrections*, *Phys. Lett. B* **820** (2021) 136522 [2103.00975].
- [24] C.-Y. Seng, D. Galviz, M. Gorchtein and U.-G. Meißner, *Improved $Ke3$ radiative corrections sharpen the $K_{\mu 2}$ - K_{l3} discrepancy*, *JHEP* **11** (2021) 172 [2103.04843].
- [25] C.-Y. Seng, D. Galviz, W. J. Marciano and U.-G. Meißner, *Update on $|V_{us}|$ and $|V_{us}/V_{ud}|$ from semileptonic kaon and pion decays*, *Phys. Rev. D* **105** (2022), no. 1 013005 [2107.14708].
- [26] W. J. Marciano and A. Sirlin, *Radiative corrections to $\pi(\text{lepton } 2)$ decays*, *Phys. Rev. Lett.* **71** (1993) 3629–3632.
- [27] **NA7** Collaboration, S. Amendolia *et. al.*, *A Measurement of the Space - Like Pion Electromagnetic Form-Factor*, *Nucl. Phys. B* **277** (1986) 168.
- [28] S. Amendolia *et. al.*, *A Measurement of the Kaon Charge Radius*, *Phys. Lett. B* **178** (1986) 435–440.
- [29] G. Ecker, J. Gasser, A. Pich and E. de Rafael, *The Role of Resonances in Chiral Perturbation Theory*, *Nucl. Phys. B* **321** (1989) 311–342.
- [30] G. Ecker, J. Gasser, H. Leutwyler, A. Pich and E. de Rafael, *Chiral Lagrangians for Massive Spin 1 Fields*, *Phys. Lett. B* **223** (1989) 425–432.
- [31] V. Cirigliano, G. Ecker, M. Eidemuller, R. Kaiser, A. Pich and J. Portoles, *Towards a consistent estimate of the chiral low-energy constants*, *Nucl. Phys. B* **753** (2006) 139–177 [hep-ph/0603205].
- [32] V. Cirigliano, M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, *Radiative corrections to $K(l3)$ decays*, *Eur. Phys. J.* **C23** (2002) 121–133 [hep-ph/0110153].
- [33] X. Feng, M. Gorchtein, L.-C. Jin, P.-X. Ma and C.-Y. Seng, *First-principles calculation of*

- electroweak box diagrams from lattice QCD*, *Phys. Rev. Lett.* **124** (2020), no. 19 192002 [2003.09798].
- [34] S. Descotes-Genon and B. Moussallam, *Radiative corrections in weak semi-leptonic processes at low energy: A Two-step matching determination*, *Eur. Phys. J.* **C42** (2005) 403–417 [hep-ph/0505077].
- [35] B. Ananthanarayan and B. Moussallam, *Four-point correlator constraints on electromagnetic chiral parameters and resonance effective Lagrangians*, *JHEP* **06** (2004) 047 [hep-ph/0405206].
- [36] J. Bijnens and G. Ecker, *Mesonic low-energy constants*, *Ann. Rev. Nucl. Part. Sci.* **64** (2014) 149–174 [1405.6488].
- [37] Y. Aoki *et. al.*, *FLAG Review 2021*, 2111.09849.
- [38] N. Carrasco, P. Lami, V. Lubicz, L. Riggio, S. Simula and C. Tarantino, *$K \rightarrow \pi$ semileptonic form factors with $N_f = 2 + 1 + 1$ twisted mass fermions*, *Phys. Rev. D* **93** (2016), no. 11 114512 [1602.04113].
- [39] **Fermilab Lattice, MILC** Collaboration, A. Bazavov *et. al.*, *$|V_{us}|$ from $K_{\ell 3}$ decay and four-flavor lattice QCD*, *Phys. Rev.* **D99** (2019), no. 11 114509 [1809.02827].
- [40] A. Bazavov *et. al.*, *Kaon semileptonic vector form factor and determination of $|V_{us}|$ using staggered fermions*, *Phys. Rev. D* **87** (2013) 073012 [1212.4993].
- [41] **RBC/UKQCD** Collaboration, P. A. Boyle *et. al.*, *The kaon semileptonic form factor in $N_f = 2 + 1$ domain wall lattice QCD with physical light quark masses*, *JHEP* **06** (2015) 164 [1504.01692].
- [42] W. J. Marciano, *Precise determination of $-V(us)$ — from lattice calculations of pseudoscalar decay constants*, *Phys. Rev. Lett.* **93** (2004) 231803 [hep-ph/0402299].
- [43] P. Boyle *et. al.* *High-precision determination of V_{us} and V_{ud} from lattice QCD*. [Link].