

Likelihood ratio tests under model misspecification in high dimensions

Nina Dörnemann^{a,*}

^a*Fakultät für Mathematik, Ruhr-Universität Bochum, 44801 Bochum, Deutschland*

Abstract

We investigate the likelihood ratio test for a large block-diagonal covariance matrix with an increasing number of blocks under the null hypothesis. While so far the likelihood ratio statistic has only been studied for normal populations, we establish that its asymptotic behavior is invariant under a much larger class of distributions. This implies robustness against model misspecification, which is common in high-dimensional regimes. Demonstrating the flexibility of our approach, we additionally establish asymptotic normality of the log-likelihood ratio test statistic for the equality of many large sample covariance matrices under model uncertainty. For this statistic, a subtle adjustment to the centering term is needed compared to normal case. A simulation study and an analysis of a data set from psychology emphasize the usefulness of our findings.

Keywords: Block-diagonal covariance matrix, Central limit theorem, Equality of covariance matrices, High-dimensional inference, Likelihood ratio test, Model misspecification, Non-normal population.

2020 MSC: Primary 62H15, Secondary 62H10

1. Introduction

Over the last decades, the availability of high-dimensional data sets across diverse disciplines such as biostatistics, wireless communications and finance has transformed statistical practice (see, e.g., [15, 30] and references therein). Traditional multivariate analysis, as outlined in the text book [2, 40], is developed under the paradigm that the dimension is negligible compared to the sample size and breaks down seriously if this assumption is violated. Such problems have spurred the development of new analysis tools, that work for dimensions of the same order as and even larger than the sample size. The literature on these topics is so large, that we can only cite a few illustrative examples, related to the present work: The works [7, 51] address whether a large covariance matrix admits a block-diagonal structure. Tests for independence in various setting are discussed in [21, 36]. In the work [24], Hu et al. concentrate on tests for the equality of high-dimensional mean vectors, while [22] take a broader perspective on high-dimensional testing by investigating a class of U -statistics.

Turning closer to the scope of this work, the likelihood ratio method has received much attention in the literature on high-dimensional statistical inference since the past decade. The starting point for the investigating of various classical testing problems transferred to a high-dimensional setting can be seen in the work of [29] establishing CLTs for the corresponding log-likelihood ratio tests, including the two main testing problems investigated in this work. The authors of [28] tried to relax the assumptions on the parameters, while other authors extended these results in various directions. For example, [27] proved a moderate deviation principle for these likelihood ratio tests. More recent generalizations include the works of [14, 19, 44]. All of these works rely on normally distributed data, and the asymptotic behavior of these test statistics under model misspecification has received little attention in the literature on high-dimensional statistics so far. A few works investigating likelihood ratio tests in different settings under model uncertainty include [25, 32, 33, 37, 48]. We add to this line of literature by dropping the restrictive distributional assumption of normality. In particular, we find that the CLTs for the log-likelihood of two specific testing problems remain still valid when only assuming moments of order $(4 + \delta)$ for some $\delta > 0$. Besides the theoretical importance

*Corresponding author. Email address: nina.doernemann@rub.de

of our findings, these results ensure more robust statistical guarantees for practitioners, as the validity of the normal assumption is not a priori clear for high-dimensional data sets.

Interestingly, our results reveal that the limiting null distribution of the log-likelihood for testing for a block-diagonal structure does not depend on specific characteristics of the underlying data generating distribution, such as the fourth moment. This observation is illustrated in Fig. 1 where we consider the problem of testing whether the covariance matrix of a p -dimensional random vector admits a block-diagonal structure with q blocks. Here, we display three histograms for the corresponding log-likelihood ratio test statistic under the null hypothesis based on three different distributions for the samples. The components of all vectors are independent identically distributed with respect to a standard normal distribution (left column), standardized t-distribution (middle column) and centered exponential distribution (right column), respectively. Thus, the null hypothesis of a block-diagonal covariance matrix is obviously satisfied (in this case, the covariance matrix equals the identity matrix). We observe that the histograms look very similar. This testing problem will be examined in detail in Section 2 of this work.




Fig. 1: Histograms for the log-likelihood ratio test statistic (4) under the hypothesis (3) of a block diagonal structure of $q = 30$ blocks of equal size 2 in a $p = 60$ -dimensional random vector (sample size 100, simulation runs 10,000). Left column: standard normally distributed data, middle column: standardized t-distributed data with 15 degrees of freedom, right column: centered exponentially distributed data with parameter 1. The gray curve indicates the density of the standard normal distribution.

Before concluding this introduction, we would like to discuss the main ideas for our proofs. Under the normal assumption, the exact distribution of the test statistic is available under the null hypothesis, on which proofs of previous works crucially depend on. Equipped with such a knowledge, the moment-generating of the log-likelihood test statistic is investigated [see, e.g., 19, 29, 44] or a general central limit theorem is applied [see 14]. Obviously, we cannot hope for an analogue exact representation without knowing the underlying distribution of the data.

In order to tackle the difficulties arising in the proof for non-normal populations, we derive a novel representation of the log-likelihood test statistic involving random quadratic forms without imposing restrictive distributional assumptions. For this purpose, we perform a QR-decomposition for the (sub)data matrices. Such QR-decompositions are useful in a broader context: The authors of [49] used this tool in order to derive the logarithmic law of the sample covariance matrix for the case $p/n \rightarrow 1$ near singularity, while [23] investigated the log-determinant of the sample correlation matrix under infinite fourth moment. These papers were partially inspired by works of [5, 41], in which the authors proved Girko’s logarithmic law for a general random matrix with independent entries and brought his “method of perpendiculars” [see 17] to a mathematically rigorous level. Via our representation, we are in the position to decompose the test statistic into three parts: we will prove that the dominating linear term satisfies a central limit theorem for martingale difference schemes, while the quadratic term converges to constant and the remainders are asymptotically negligible. Heuristically, this decomposition can be motivated by Taylor’s expansion $\log(1 + x) = x - x^2/2 + \mathcal{O}(x^3)$, though one needs more delicate arguments in order to justify this step mathematically correct.

This work is structured as follows. In Section 2, we present a CLT for the log-likelihood ratio test of a block-diagonal covariance matrix under the null hypothesis. Here, the number of blocks may increase together with the dimension of the data and sample size while we do not assume that the data is generated by a normal distribution. As a corollary, the distribution of a test for a diagonal covariance matrix is derived. In Section 3, we apply our method

to another classical likelihood ratio test and provide the asymptotic distribution of the log-likelihood ratio test on the equality of many large covariance matrices. The main results of Section 2 and 3 are proven in Section 5. We illustrate our findings with a simulation study, including a comparison to other criteria, and a real data analysis in Section 4.

2. Testing for a block-diagonal covariance matrix

In the main part of this work, we revisit a very prominent problem in high-dimensional data analysis, namely a test for uncorrelation of sub-vectors of a multivariate distribution. For normally distributed data, this coincides with a test for independence of these sub-vectors. To be precise, let $\mathbf{y} = \Sigma^{\frac{1}{2}} \mathbf{x}$ denote a p -dimensional random vector with mean $\boldsymbol{\mu} = \mathbf{0} \in \mathbb{R}^p$ and covariance matrix $\Sigma \in \mathbb{R}^{p \times p}$, where $\Sigma^{\frac{1}{2}}$ denotes the symmetric square root of Σ . We assume that \mathbf{y} is decomposed as

$$\mathbf{y} = (\mathbf{y}^{(1)\top}, \dots, \mathbf{y}^{(q)\top})^\top, \quad (1)$$

where $\mathbf{y}^{(i)}$ are vectors of dimension $p_i \in \mathbb{N}$, $1 \leq i \leq q$, such that $\sum_{i=1}^q p_i = p$ for some integer $q \geq 2$. Moreover, we assume that the components of \mathbf{x} are i.i.d. with respect to some centered and standardized distribution. Let

$$\Sigma = \begin{pmatrix} \Sigma_{11} & \Sigma_{12} & \dots & \Sigma_{1q} \\ \Sigma_{21} & \Sigma_{22} & \dots & \Sigma_{2q} \\ \vdots & \vdots & \ddots & \vdots \\ \Sigma_{q1} & \Sigma_{q2} & \dots & \Sigma_{qq} \end{pmatrix} \quad (2)$$

denote the corresponding decomposition of the covariance matrix, where $\Sigma_{ij} := \text{Cov}(\mathbf{y}^{(i)}, \mathbf{y}^{(j)})$. The hypothesis of uncorrelated sub-vectors is formulated as

$$H_0 : \Sigma_{ij} = \mathbf{0} \text{ for all } i \neq j. \quad (3)$$

Let $\mathbf{x}_1, \dots, \mathbf{x}_n \stackrel{\text{i.i.d.}}{\sim} \mathbf{x}$ be a sample of independent identically distributed random variables according to \mathbf{x} and denote $\mathbf{y}_k = \Sigma^{1/2} \mathbf{x}_k$ for $1 \leq k \leq n$. Under the normal assumption $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, the likelihood ratio test is given by

$$\Lambda_n = \frac{|\hat{\Sigma}|^{n/2}}{\prod_{i=1}^q |\hat{\Sigma}_{ii}|^{n/2}} = V_n^{\frac{n}{2}}, \quad \hat{\Sigma} = \frac{1}{n} \sum_{k=1}^n \mathbf{y}_k \mathbf{y}_k^\top, \quad (4)$$

where $\hat{\Sigma}$ denotes the sample covariance matrix of $\mathbf{y}_1, \dots, \mathbf{y}_n$ and $\hat{\Sigma}_{ij}$ denotes the block in the i th row and j th column of the estimate $\hat{\Sigma}$ corresponding to the decomposition (2).

The authors of [28, 29] derived a central limit theorem for the corresponding log-likelihood ratio test statistic assuming that $\mathbf{y} \sim \mathcal{N}(\boldsymbol{\mu}, \Sigma)$ and that the number q of blocks is fixed. Several authors such as [14, 44] demonstrated that such a CLT still holds true if the parameter $q = q_n$ is allowed to increase with sample size and dimension. All of these works rely on normally distributed data. Dropping the normal assumption, the following theorem provides the asymptotic distribution of $\log \Lambda_n$ under the null hypothesis of uncorrelation without assuming a normal distribution for $\mathbf{x}_1, \dots, \mathbf{x}_n \sim \mathbf{x}$. The proof is deferred to Section 5.

Theorem 1. *Let the components of \mathbf{x} be i.i.d. centered random variables following a continuous distribution with finite $(4 + \delta)$ th moment for some $\delta > 0$. Assume that $q = q_n \geq 2$ is a possibly increasing integer, $2 \leq p = p_n < n$ with $0 < \inf_{n \in \mathbb{N}} \min_{1 \leq i \leq q} (p_i q) / n \leq \sup_{n \in \mathbb{N}} p / n < 1$ and $\max_{1 \leq i \leq q} p_i \leq \eta p$ for each $n \in \mathbb{N}$ and some fixed $\eta \in (0, 1)$. Then, it holds under the null hypothesis in (3) that*

$$\frac{\log V_n - \mu_n}{\sigma_n} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1),$$

where

$$\mu_n = \sum_{i=1}^q \left(n - p_i - \frac{1}{2} \right) \log \left(1 - \frac{p_i}{n} \right) - \left(n - p - \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right), \quad \sigma_n^2 = 2 \left\{ \sum_{i=1}^q \log \left(1 - \frac{p_i}{n} \right) - \log \left(1 - \frac{p}{n} \right) \right\}.$$

Remark 1. (i) Choose $\alpha \in (0, 1)$. We propose to reject the null hypothesis whenever

$$\log V_n \leq \sigma_n u_\alpha + \mu_n, \quad (5)$$

where u_α denotes the α -quantile of the standard normal distribution. Thus, we have by Theorem 1

$$\lim_{n \rightarrow \infty} \Pr_{H_0}(\log V_n \leq \sigma_n u_\alpha + \mu_n) = \Pr(\mathcal{N}(0, 1) \leq u_\alpha) = \alpha,$$

which means that the test keeps asymptotically its nominal level α .

- (ii) We would like to comment on the continuity assumption on the distribution of the random vector \mathbf{x} , which is needed in the proof of Theorem 1 in order to ensure formal correctness of the QR-decomposition. In general, one could replace the entries of \mathbf{x} by some random vector $\bar{\mathbf{x}}$ following a continuous distribution without changing the asymptotic distribution of $\log V_n$. One choice for $\bar{\mathbf{x}}$ could be a mixture of the original distribution and a uniform distribution (see, e.g., Proposition 2.1 in [49] for the log-determinant of the sample covariance matrix). If $q_n = q$ denotes a fixed integer, the continuity assumption is not necessary for proving Theorem 1. For an increasing parameter q_n , however, this would require a more restrictive regime for $q = q_n$, since the log-likelihood test statistics involves $q_n + 1$ log-determinants of sample covariance matrices. We do not pursue in this direction here.

As a noteworthy-by-product of Theorem 1, we are able to construct a test for a diagonal covariance matrix based on the sample correlation matrix. For this purpose, we consider the special case of testing for a diagonal covariance matrix which coincides with complete independence of the p components of \mathbf{x} in the normal case. In this case, the test in (3) is equivalent to

$$H_0 : \mathbf{R} = \mathbf{I}_p, \quad (6)$$

where $\mathbf{R} = \text{diag}(\boldsymbol{\Sigma})^{-\frac{1}{2}} \boldsymbol{\Sigma} \text{diag}(\boldsymbol{\Sigma})^{-\frac{1}{2}}$ denotes the population correlation matrix of \mathbf{y} . Then, the statistic V_n defined in (4) can be written as the determinant of the sample correlation matrix, that is,

$$V_n = |\hat{\mathbf{R}}|,$$

where $\hat{\mathbf{R}} = \text{diag}(\hat{\boldsymbol{\Sigma}})^{-\frac{1}{2}} \hat{\boldsymbol{\Sigma}} \text{diag}(\hat{\boldsymbol{\Sigma}})^{-\frac{1}{2}}$ denotes the sample correlation matrix of $\mathbf{y}_1, \dots, \mathbf{y}_n$. Several authors investigated tests for the hypothesis given in (6) in different frameworks (e.g., see [16, 23, 28, 29, 39, 43, 44]). We observe that testing for (6) is a special case of testing for (3) by letting $q = p$ and $p_1 = \dots = p_q = 1$. Then, Theorem 1 gives us the following result.

Corollary 1. *Let the components of \mathbf{x} be i.i.d. centered random variables following a continuous distribution with finite $(4 + \delta)$ th moment for some $\delta > 0$. Assume that $2 \leq p = p_n < n$ and $0 < \inf_{n \in \mathbb{N}} p/n \leq \sup_{n \in \mathbb{N}} p/n < 1$. Then, it holds under the null hypothesis in (6) that,*

$$\frac{\log |\mathbf{R}| - \bar{\mu}_n}{\bar{\sigma}_n} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1),$$

where

$$\bar{\mu}_n = p \left(n - \frac{3}{2} \right) \log \left(1 - \frac{1}{n} \right) - \left(n - p - \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right), \quad \bar{\sigma}_n^2 = 2 \left\{ p \log \left(1 - \frac{1}{n} \right) - \log \left(1 - \frac{p}{n} \right) \right\}.$$

Note that [43] investigated the log-determinant of the sample correlation matrix in a more general context. In fact, they cover the case $\mathbf{R} \neq \mathbf{I}$, while Corollary 1 is formulated under the null hypothesis $\mathbf{R} = \mathbf{I}$. If we assume that $p/n \rightarrow \gamma \in (0, 1)$, then Corollary 1 yields a special version of their Theorem 2.1, since

$$\bar{\mu}_n = - \left(n - p - \frac{1}{2} \right) \log \left(1 - \frac{p-1}{n} \right) - (p-1) + \frac{p}{n} + o(1), \quad \bar{\sigma}_n^2 = -2 \left\{ \frac{p}{n} - \log \left(1 - \frac{p-1}{n} \right) \right\} + o(1),$$

which coincides with mean and variance given in their Theorem 2.1 in the case $\mathbf{R} = \mathbf{I}$. In a follow-up work, [23] showed that the CLT for the sample correlation matrix in the case $\mathbf{R} = \mathbf{I}$ still holds true under infinite fourth moment. Other authors were interested in the study of the spatial-sign covariance matrix, which includes the sample correlation matrix as a special case. For example, [35, 52] proved CLTs for linear spectral statistics of this general class of random matrices.

3. Testing for equality of covariance matrices

We expect that our method for proving a CLT as given in Theorem 1 can be adapted to the investigation of other classical likelihood ratio tests in a non-normal setting. In order to demonstrate this adaption, we consider in this section the comparison of q centered distributions with covariance matrices $\Sigma_1, \dots, \Sigma_q \in \mathbb{R}^{p \times p}$ and generic elements $\mathbf{y}_1 = \Sigma_1^{1/2} \mathbf{x}, \dots, \mathbf{y}_q = \Sigma_q^{1/2} \mathbf{x}$. We assume that for each group j a sample of size n_j is available, $j \in \{1, \dots, q\}$. When considering asymptotics, the dimension p and the number q of groups increase with the (sub)sample sizes. As before, we assume that the components of \mathbf{x} are i.i.d. with respect to some centered and standardized distribution.

An important assumption for multivariate analysis of variance (MANOVA) is that of equal covariances in the different groups, which motivates our interest in testing the hypothesis

$$H_0 : \Sigma_1 = \dots = \Sigma_q. \quad (7)$$

This problem has been considered by several authors in the context of high-dimensional inference (see [14, 19, 28, 29, 42, 46, 47] among others). In this section, we add to this line of literature and investigate the asymptotic distribution of the likelihood ratio test based on samples of independent distributed observations $\mathbf{y}_{jk} \stackrel{\text{i.i.d.}}{\sim} \mathbf{y}_j$, $1 \leq k \leq n_j$, $1 \leq j \leq q$. To be precise, let $n = \sum_{j=1}^q n_j$ be the total sample size, then the likelihood ratio test for the hypothesis (7) under the normal assumption $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ is given by

$$\Lambda_{n,2} = \frac{\prod_{j=1}^q |\mathbf{A}_j/n_j|^{\frac{1}{2}n_j}}{|\mathbf{A}/n|^{\frac{1}{2}n}}, \quad (8)$$

where the $p \times p$ matrices \mathbf{A}_j and \mathbf{A} are defined as

$$\mathbf{A}_j = \sum_{k=1}^{n_j} \mathbf{y}_{jk} \mathbf{y}_{jk}^\top, \quad \mathbf{A} = \sum_{j=1}^q \mathbf{A}_j.$$

In the case $\mathbf{y}_j \sim \mathcal{N}(\boldsymbol{\mu}_j, \Sigma_j)$, $1 \leq j \leq q$, [28, 29] proved asymptotic normality of the corresponding log-likelihood ratio test statistic under the null hypothesis if the number q of groups is fixed. These results were generalized by [14, 19] for the case of an increasing number $q = q_n$ of groups. All of these works dealt only with normally distributed data, while the following result shows that a subtle adjustment to the centering term is necessary for non-normal data. In the following theorem, we provide the limiting distribution of $\log \Lambda_{n,2}$ under the null hypothesis without imposing a normal assumption on $\mathbf{y}_1, \dots, \mathbf{y}_q$ in a high-dimensional setting, where the number of groups is allowed to increase.

Theorem 2. *Let the components of \mathbf{x} be i.i.d. centered random variables following a continuous distribution with finite $(4 + \delta)$ th moment for some $\delta > 0$. Assume that $q = q_n \geq 2$ is a possibly increasing integer, $n_j = n_j(n) > p = p_n$ for every $n \in \mathbb{N}$ and $0 < \inf_{n \in \mathbb{N}} p/n \leq \sup_{n \in \mathbb{N}} \max_{1 \leq j \leq q} p/n_j < 1$. Then it holds under the null hypothesis (7)*

$$\frac{2(\log \Lambda_{n,2} - \mu_n)}{n\sigma_n} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1),$$

where

$$\begin{aligned} \mu_n &= n \left(n - p - \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \left(n_j - p - \frac{1}{2} \right) \log \left(1 - \frac{p}{n_j} \right) + \frac{\mathbb{E}[x_{11}^4] - 3}{2} p(1 - q), \\ \sigma_n^2 &= \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q \left(\frac{n_j}{n} \right)^2 \log \left(1 - \frac{p}{n_j} \right). \end{aligned} \quad (9)$$

The proof is provided in Section 5. Similarly to Remark 1, an asymptotic level α test for the hypothesis (7) can be constructed using Theorem 2. For the sake of brevity, we omit the details.




Fig. 2: Rejection rates for the log-likelihood ratio test (5) for Scenario (i) (first row) and Scenario (ii) (second row) under (10) based on 2,000 simulation runs. First column: standard normally distributed data, second column: standardized t-distributed data with 15 degrees of freedom, third column: centered exponentially distributed data with parameter 1. The triangle indicates $n = 100, p = 60$, the square $n = 120, p = 90$ and the circle $n = 180, p = 120$. The vertical grey line in each figure defines the nominal level $\alpha = 5\%$.

4. Finite-sample properties

In this section, we investigate the finite-sample properties of the test (5) under both the hypothesis and the alternative. Following [44], we consider the following alternative

$$\Sigma = (1 - \delta)\mathbf{I} + \delta\mathbf{1}, \quad (10)$$

where $\mathbf{1}$ denotes the $p \times p$ matrix filled with ones and \mathbf{I} denotes the $p \times p$ identity matrix. Here, the parameter $\delta \geq 0$ determines the “distance” to the null hypothesis (3) (note that the choice $\delta = 0$ corresponds to the null hypothesis (3)). In Fig. 2, we display the empirical rejection rates of the test (5) for different choices of δ, n, p, q, p_i and different distributions for the random vector \mathbf{x} . All results are based on 2,000 simulation runs and the components of \mathbf{x} are independent identically distributed with respect to a standard normal distribution (first column), standardized t-distribution (second column) and centered exponential distribution (third column), respectively. The vertical grey line in each figure defines the nominal level $\alpha = 5\%$. For the choice of the different groups, we consider the following two scenarios:

- (i) $q = 3, p_1 = p_2 = p_3 = p/3$,
- (ii) $q = p/2, p_1 = \dots = p_{q-1} = 1, p_q = q + 1$.

We observe a good approximation of the nominal level in all cases under consideration. Moreover, the power increases reasonably as δ increases. It should be noted that the increase in power is a bit stronger for Scenario (ii) than for Scenario (i). The finite-sample properties do not significantly differ for the three underlying data generating distributions, as indicated by the asymptotic result provided in Theorem 1. Overall, the test admits a desirable performance for

finite-sample sizes, both for a large number and relatively small number of groups as covered by Scenarios (i) and (ii), and the accuracy improves for large sample size and dimension.

4.1. Comparison to other tests

In this section, we compare the test given in (5) to two trace criteria for the null hypothesis (3) of blockdiagonality. Both procedures rely on the normal assumption for the data. In this case, substitution principles are widely available, and the tests are based on the centered sample covariance matrix

$$\hat{\Sigma}^{\text{cen}} = \frac{1}{n} \sum_{k=1}^n (\mathbf{y}_k - \bar{\mathbf{y}})(\mathbf{y}_k - \bar{\mathbf{y}})^\top$$

and its blocks Σ_{ij}^{cen} corresponding to the decomposition in (2), where

$$\bar{\mathbf{y}} = \frac{1}{n} \sum_{k=1}^n \mathbf{y}_k$$

denotes the sample mean. On the one hand, we consider the tests proposed by [4, 26] who assume that $q \geq 2$ is fixed. On the other hand, we investigate a criterion proposed by [34] for $q = 2$ groups. To make a meaningful comparison, we restrict our analysis to the case $q = 2$ with normally distributed data. In this case, the test statistics proposed by [4, 26] both are equal to

$$L_n = \text{tr} \left(\hat{\Sigma}_{21}^{\text{cen}} (\hat{\Sigma}_{11}^{\text{cen}})^{-1} \hat{\Sigma}_{12}^{\text{cen}} (\hat{\Sigma}_{22}^{\text{cen}})^{-1} \right),$$

which is shown to satisfy

$$T_1 = \frac{L_n - m_n}{\sqrt{v}} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1)$$

under H_0 given in (3) [see Theorem 3.2 in the work of 26]. Here, we set

$$m_n = \frac{p_2 r_{n2}}{r_{n1} + r_{n2}}, v = \frac{2h^2 r_1^2 r_2^2}{(r_1 + r_2)^4}, h = \sqrt{r_1 + r_2 - r_1 r_2}$$

and assume that $r_{n1} = p_2/p_1 \rightarrow r_1 \in (0, \infty)$ and $r_{n2} = p_2/(n-1-p_1) \rightarrow r_2 \in (0, \infty)$, $p_2 < n$. Coming back to the test investigated by [34], we define

$$\hat{\gamma}_{ij} = \frac{1}{(n-2)(n+1)} \left\{ \text{tr} \left(\hat{\Sigma}_{ij}^{\text{cen}} \hat{\Sigma}_{ji}^{\text{cen}} \right) - \frac{1}{n-1} \text{tr} \left(\hat{\Sigma}_{ii}^{\text{cen}} \right) \text{tr} \left(\hat{\Sigma}_{jj}^{\text{cen}} \right) \right\}, i, j \in \{1, 2\}.$$

If $p \rightarrow \infty$ as $n \rightarrow \infty$, the authors show that

$$T_2 = \sqrt{\frac{(n-2)(n+1)}{2}} \frac{\hat{\gamma}_{12}}{\sqrt{\hat{\gamma}_{11} \hat{\gamma}_{22}}} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1)$$

under H_0 in (3) assuming that

$$0 < \lim_{n \rightarrow \infty} \text{tr} \Sigma^k < \infty, k \in \{1, 2, 4\}.$$

For a prescribed level $\alpha \in (0, 1)$, we reject H_0 if $T_2 > u_{1-\alpha}$ (or $T_1 > u_{1-\alpha}$ when using the test based on T_1). For simplicity of presentation, we refer to the test decisions based upon the statistics T_1 and T_2 also by T_1 and T_2 . We expect the test T_2 to admit a strong rejection rate under the alternative (10) since the core part $\hat{\gamma}_{12}$ of the statistic T_2 compares the trace of the product of the two off-diagonal blocks to the product of traces of the two diagonal blocks. This difference is expected to be large under the alternative (10). In Table 1, the empirical rejection rates of T_1, T_2

and the log-likelihood ratio test LLRT (see (5)) under the null hypothesis (3) and the alternative (10) are displayed. Additionally, we consider the alternative model

$$\mathbf{y}^{(1)} = (1 + \delta)\mathbf{z}^{(1)}, \mathbf{y}^{(2)} = \mathbf{z}^{(2)} + \delta\mathbf{z}^{(1)} \quad (11)$$

(recall the notation in (1)), where $\mathbf{z}^{(i)} \sim \mathcal{N}_{p_i}(\mathbf{0}, \mathbf{I})$ are independent for $i \in \{1, 2\}$. Note that the choice $\delta = 0$ in (11) corresponds to the null hypothesis (3). This model has been considered in previous works [see, e.g., 44]. The corresponding empirical rejection rates can be found in Table 2. In most cases, the nominal level $\alpha = 5\%$ is approximated well by all tests under the null hypothesis. Under the alternative (10), the test T_2 outperforms T_1 as well as the LLRT in terms of empirical power. The LLRT still performs reasonably well under this alternative, while the performance of T_1 is not satisfying in most cases. Under model (11), we observe for all three tests a strong rejection rate under the alternative with T_2 leading the field.

In summary, we observe that T_2 is slightly preferable in the cases under consideration, which is not surprising as T_2 is just tailored to the case of two groups with normally distributed data, while the user may apply the LLRT in a broader context. However, even in this restrictive situation, the LLRT still forms a reliable criterion with a reasonable rejection rate. We emphasize that the theoretical statistical guarantees for T_1 and T_2 rely on normally distributed data and a fixed number q of groups, while the LLRT is shown to satisfy a CLT beyond the normal assumption for a possibly increasing number of groups (Theorem 1).

4.2. The almost-singular case




Fig. 3: Rejection rates for the log-likelihood ratio test (5) for Scenario (i) (first row, defined at the beginning of Section 4) and Scenario (ii) (second row) under (10) based on 2,000 simulation runs. First column: standard normally distributed data, second column: standardized t-distributed data with 15 degrees of freedom, third column: centered exponentially distributed data with parameter 1. The triangle indicates $n = 100, p = 98$, the square $n = 120, p = 118$ and the circle $n = 180, p = 178$ for Scenario (i). For Scenario (ii), the corresponding parameters are $n = 100, p = 99$ (triangle), $n = 120, p = 117$ (square) and $n = 180, p = 177$ (circle), respectively. The vertical gray line in each figure defines the nominal level $\alpha = 5\%$.

Table 1: Empirical rejection rates of the tests T_1, T_2 and the LLRT under the null hypothesis (3) and the alternative (10) for different values of $n, p_1, p_2, q = 2, \mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ and 2,000 simulation runs.

(n, p, p_1)	$\delta = 0$			$\delta = 0.15$			$\delta = 0.3$		
	T_1	T_2	LLRT	T_1	T_2	LLRT	T_1	T_2	LLRT
(150, 80, 70)	0.048	0.041	0.051	0.529	1.000	0.787	0.717	1.000	0.995
(150, 80, 40)	0.054	0.054	0.059	0.418	1.000	0.838	0.518	1.000	0.995
(150, 100, 80)	0.057	0.060	0.049	0.331	1.000	0.674	0.423	1.000	0.949
(200, 50, 25)	0.048	0.051	0.050	0.949	1.000	1.000	0.997	1.000	1.000
(200, 120, 60)	0.048	0.045	0.044	0.395	1.000	0.889	0.454	1.000	0.994
(200, 120, 90)	0.059	0.053	0.056	0.420	1.000	0.857	0.497	1.000	0.997
(250, 100, 75)	0.048	0.061	0.050	0.810	1.000	0.997	0.907	1.000	1.000
(250, 150, 100)	0.053	0.052	0.053	0.410	1.000	0.936	0.482	1.000	0.999
(250, 200, 100)	0.040	0.046	0.045	0.251	1.000	0.665	0.270	1.000	0.890

Table 2: Empirical rejection rates of the tests T_1, T_2 and the LLRT under the null hypothesis (3) and the alternative model (11) for different values of $n, p_1, p_2, q = 2, \mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ and 2,000 simulation runs.

(n, p, p_1)	$\delta = 0$			$\delta = 0.1$			$\delta = 0.2$		
	T_1	T_2	LLRT	T_1	T_2	LLRT	T_1	T_2	LLRT
(150, 80, 70)	0.048	0.042	0.051	0.086	0.103	0.092	0.277	0.457	0.290
(150, 80, 40)	0.054	0.054	0.059	0.188	0.287	0.183	0.915	0.990	0.895
(150, 100, 80)	0.057	0.060	0.049	0.090	0.122	0.093	0.356	0.642	0.349
(200, 50, 25)	0.048	0.051	0.050	0.334	0.407	0.340	0.998	0.100	0.999
(200, 120, 60)	0.048	0.045	0.044	0.251	0.397	0.238	0.981	0.100	0.976
(200, 120, 90)	0.059	0.053	0.056	0.140	0.204	0.147	0.682	0.923	0.671
(250, 100, 75)	0.048	0.061	0.050	0.181	0.259	0.180	0.921	0.981	0.915
(250, 150, 100)	0.053	0.052	0.053	0.212	0.350	0.203	0.947	0.100	0.937
(250, 200, 100)	0.040	0.046	0.045	0.281	0.543	0.243	0.993	1.000	0.969

In Theorem 1, we assume that the dimension-to-sample-size ratio p/n is uniformly bounded away from 1, which ensures that the variance σ_n^2 remains bounded. In this subsection, we will demonstrate that the test proposed in (5) can admit a desirable performance, especially under the null hypothesis (3), even if the variance explodes. For this purpose, we consider Scenarios (i) and (ii) in cases where the dimension p is close to the sample size n . The two scenarios determining the group structure are defined at the beginning of Section 4. In Fig. 3, we display the empirical rejection rates of the test (5) for different choices of δ , n , p , q , p_i and different distributions for the random vector \mathbf{x} . The nominal level $\alpha = 5\%$ is approximated well under the null hypothesis ($\delta = 0$) in all cases under consideration. Comparing Fig. 3 to Fig. 2, the increase of the power curve is a little flatter if the dimension p is close to the sample size n for Scenario (ii) and significantly flatter for Scenario (i). This indicates that a central limit theorem for the log-likelihood test statistic under the null hypothesis might still hold true if $p/n \rightarrow 1$ as $n \rightarrow \infty$ but, unsurprisingly, at the cost of a lower speed of convergence as the variance is no longer bounded. Under the alternative for Scenario (i), the test does not yield a useful decision criterion. However, we still observe a reasonable behavior of the test under H_0 for both Scenarios (i) and (ii) as well as under the alternative for Scenario (ii).

4.3. Analysis of the 16 Personality Factor Questionnaire

To further illustrate the usefulness of our method, we analyze the 16 Personality Factor Questionnaire (16PF) data set. The questionnaire goes back to [10, 11] and provides a comprehensive measure of normal-range personality characteristics [see also 9]. The 16 proposed factors include warmth, reasoning, emotional stability, dominance, liveliness, rule-consciousness, social boldness, sensitivity, vigilance, abstractedness, privateness, apprehension, openness to change, self-reliance, perfectionism, tension. In psychology, there is a controversy about the appropriate number of personality factors in connection with 16PF [e.g., see 12, 38]. Nowadays, most psychologists can agree on a number of 5 factors, the so-called Big Five: openness, conscientiousness, extraversion, agreeableness, neuroticism [see, e.g. 6, 13, 18, 45]. Nevertheless, the 16PF is widely applied in various contexts where an in-depth personality analysis is needed. For example, it can be used in order to construct effective self-development goals and career plans adjusted to a worker's individual strengths and limitations [see, among many others, 8, 31, 50]. The relevance of 16PF is emphasized by the fact that it had been adapted to over 35 languages worldwide.

The data set "PF16" under consideration (available at https://openpsychometrics.org/_rawdata/) includes the ratings of 49159 persons concerning 163 statements about their personality on the following scale of accuracy: (1) disagree, (2) slightly disagree, (3) neither agree nor disagree, (4) slightly agree, (5) agree.

For the following analysis, we centered the data with the sample mean. Since we will perform the test only on subsamples of size $n \ll 49159$, this will not cause a mean shift in the CLT given in Theorem 1 as discussed in detail at the end of this subsection. The number of groups is $q = 16$ with sizes $p_i = 10$ for $i \in \{1, \dots, 16\} \setminus \{2\}$ and $p_2 = 13$, so that the total dimension indeed sums up to $p = 163$. In order to quantify how far the sample covariance matrix $\hat{\Sigma}$ deviates from a block diagonal structure, we propose the following quantity

$$\tau(\hat{\Sigma}) = \frac{\|\hat{\Sigma} - \hat{\Sigma}_{\text{block}}\|}{\|\hat{\Sigma}\|} \geq 0.$$

Here,

$$\hat{\Sigma}_{\text{block}} = \begin{pmatrix} \hat{\Sigma}_{11} & \mathbf{0} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & \hat{\Sigma}_{22} & \mathbf{0} & \dots & \mathbf{0} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \dots & \hat{\Sigma}_{16,16} \end{pmatrix}$$

is the block-diagonal version of $\hat{\Sigma}$ and $\|\cdot\|$ denotes an appropriate matrix norm. Note that $\tau(\hat{\Sigma})$ equals 0 if and only if $\hat{\Sigma} = \hat{\Sigma}_{\text{block}}$ and $\hat{\Sigma} \neq \mathbf{0}$. For the 1-norm or the Frobenius norm, we have $\tau(\hat{\Sigma}) \in [0, 1]$ and $\tau(\hat{\Sigma})$ equals 1 if and only if $\hat{\Sigma}_{ii} = \mathbf{0}$ for $1 \leq i \leq 16$. ($\mathbf{0}$ denotes always a matrix of appropriate dimension filled with zeros.) In Table 3, we display the values of $\tau(\hat{\Sigma})$ for different matrix norms. Since the values of $\tau(\hat{\Sigma})$ are not close to 0, assuming a block diagonal structure seems not reasonable and the null hypothesis of blockdiagonality should be rejected. In order to demonstrate the usefulness of our approach for high-dimensional data sets, we proceed with a data reduction. We

Table 3: Values of $\tau(\hat{\Sigma})$ for different matrix norms.

Norm	spectral	Frobenius	1-norm
$\tau(\hat{\Sigma})$	0.7962393	0.8005626	0.825113

interpret the answers to the questionnaire as i.i.d. observations from a random vector of dimension $p = 163$. Then we choose a sample of persons of size $n \ll 49159$ uniformly at random and iterate this procedure. The test in (5) rejects the null hypothesis (3) of blockdiagonality for p, q, p_i as given above and all values of the size n of the subsample under consideration (e.g., $n \in \{170, 200, 250\}$), that is, the empirical rejection rate equals 1. This means that the LLRT detects the dependency between the 16 factors indicated by the values of $\tau(\hat{\Sigma})$.

Motivated by the Big Five, we choose five subgroups corresponding to the factors openness to change, perfectionism, privateness, warmth and apprehension, and analyze the corresponding sample covariance matrix $\hat{\Sigma}_{\text{sub}}$. The “measure” $\tau(\hat{\Sigma}_{\text{sub}})$ for blockdiagonality of $\hat{\Sigma}_{\text{sub}}$ can be defined similarly to $\tau(\hat{\Sigma})$ and is displayed in Table 4. The

Table 4: Values of $\tau(\hat{\Sigma}_{\text{sub}})$ for different matrix norms.

Norm	spectral	Frobenius	1-norm
$\tau(\hat{\Sigma}_{\text{sub}})$	0.5270513	0.5003968	0.4909513

values of $\tau(\hat{\Sigma}_{\text{sub}})$ displayed above differ significantly from 0 but less extremely than those of $\tau(\hat{\Sigma})$. Still, we expect that a reasonable test rejects the null hypothesis at a high rate when iterated over randomly chosen subsamples. In the Table 5, the empirical rejection rate is displayed for the test given in (5) applied to 500 subsamples of size $n \in \{51, 55, 60, 65, 70, 75\}$ chosen at random and dimension $p = 50, q = 5, p_i = 10$ for $1 \leq i \leq 5$. The choice of the subgroups is motivated by the Big Five, as explained above. Desirably, as the values of $\tau(\hat{\Sigma})$ and $\tau(\hat{\Sigma}_{\text{sub}})$ are not close

Table 5: Empirical rejection rate for the test (5) applied to 500 subsamples of size $n \in \{51, 55, 60, 65, 70, 75\}$ chosen at random and dimension $p = 50, q = 5, p_i = 10$ for $1 \leq i \leq 5$.

n	51	55	60	65	70	75
rejection rate	0.774	0.92	0.976	0.992	0.966	1

to 0, we observe that the null hypothesis is rejected for an overwhelming number of cases under consideration. In this section, we have seen that the LLRT does not only perform well for simulated data in various scenarios, but it can also be used in order to detect dependent factors appearing in real data sets.

Moreover, our statistical analysis reflects the controversy in psychology about the appropriate number of factors describing a human’s personality. It confirms that the factors given in 16PF are not independent, even when choosing a smaller group of factors which can be associated with the Big Five. This observation sheds light on a fundamental difference between these two approaches: Although personality traits are thought to be dependent, the Big Five in contrast to 16PF are modelled as independent factors received through the factor analysis using orthogonal rotations, which simplifies the statistical analysis and the psychological interpretation. This may explain why the Big Five are very popular as a universal model in personality research, while the 16PF is widely applied in various everyday scenarios, e.g., by mental health professionals and career coaches.

Concluding this data example, we would like to comment on two technical issues relating to Theorem 1. First, the assumption that the data follows a continuous distribution is obviously violated. However, as the number q of groups is relatively small in comparison to the considered sample sizes n , we can interpret q asymptotically as a fixed (or negligible) quantity in comparison to n . Then, the continuity assumption in Theorem 1 is not necessary (see

Remark 1). Second, the data are assumed to be centered, and in general, we expect a mean shift in Theorem 1 when centering with the sample mean. In our application, we center with the full sample mean, while the statistical analysis is restricted to subsamples of much smaller size $n \ll 49159$. Thus, the full sample mean can be interpreted as the true population mean and does not cause a shift in the mean structure.

5. Proofs

In this section, we provide the proof of Theorem 1 and present the necessary auxiliary results. We conclude with the proof of Theorem 2. In the following, we make use of the notation $a \lesssim b$ which means that a is less than or equal to b up to a positive constant, that is, there exists some $C > 0$ independent of $n \in \mathbb{N}$ such that $a \leq Cb$.

Proof of Theorem 1. Note that, under the null hypothesis (3), we have

$$V_n = \frac{|\Sigma| |\hat{\mathbf{I}}|}{\prod_{i=1}^q (|\Sigma_{ii}| |\hat{\mathbf{I}}_{ii}|)} = \frac{|\hat{\mathbf{I}}|}{\prod_{i=1}^q |\hat{\mathbf{I}}_{ii}|}, \quad (12)$$

where

$$\hat{\mathbf{I}} = \frac{1}{n} \mathbf{X}_n \mathbf{X}_n^\top, \hat{\mathbf{I}}_{ii} = \frac{1}{n} \mathbf{X}_{n,i} \mathbf{X}_{n,i}^\top, \mathbf{X}_n = (\mathbf{x}_1, \dots, \mathbf{x}_n) = (\mathbf{b}_1, \dots, \mathbf{b}_p)^\top,$$

$$\mathbf{X}_{n,i} = (\mathbf{b}_{p_{i-1}^*+1}, \dots, \mathbf{b}_{p_i^*})^\top, p_i^* = \sum_{j=1}^i p_j,$$

for $1 \leq i \leq q$, where we set $p_0^* = 0$. In order to establish a more handy representation for determinants of the sample covariance matrix, we proceed with a QR decomposition of \mathbf{X}_n^\top and $\mathbf{X}_{n,i}^\top$ as explained in detail in Section 2 of [49] and get

$$\hat{\mathbf{I}} = \frac{1}{n} \prod_{i=1}^p \mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i, \hat{\mathbf{I}}_{ii} = \frac{1}{n} \prod_{j=p_{i-1}^*+1}^{p_i^*} \mathbf{b}_j^\top \mathbf{P}(p_{i-1}^*+1; j-1) \mathbf{b}_j, \quad 1 \leq i \leq q,$$

where

$$\mathbf{P}(p_{i-1}^*+1; j-1) = \mathbf{I} - \mathbf{X}_n(p_{i-1}^*+1; j-1)^\top \left(\mathbf{X}_n(p_{i-1}^*+1; j-1) \mathbf{X}_n(p_{i-1}^*+1; j-1)^\top \right)^{-1} \mathbf{X}_n(p_{i-1}^*+1; j-1), \quad (13)$$

and

$$\mathbf{P}(j) = \mathbf{P}(1; j),$$

denote the projection matrices on the orthogonal complements of $\text{span}(\mathbf{b}_{p_{i-1}^*+1}, \dots, \mathbf{b}_{j-1})$ and $\text{span}(\mathbf{b}_1, \dots, \mathbf{b}_j)$, respectively. Here, we denote

$$\mathbf{X}_n(i; j) = (\mathbf{b}_i, \dots, \mathbf{b}_j)^\top, \quad 1 \leq i \leq j \leq p$$

and $\mathbf{P}(0) = \mathbf{I} = \mathbf{P}(i; j)$ for $j < i$. This implies

$$\begin{aligned} \log V_n &= \sum_{i=1}^p \log \left(\mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i \right) - \sum_{i=1}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \log \left(\mathbf{b}_j^\top \mathbf{P}(p_{i-1}^*+1; j-1) \mathbf{b}_j \right) \\ &= \sum_{i=p_1+1}^p \log \left(\mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i \right) - \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \log \left(\mathbf{b}_j^\top \mathbf{P}(p_{i-1}^*+1; j-1) \mathbf{b}_j \right), \end{aligned}$$

where we used $\mathbf{P}(i-1) = \mathbf{P}(1; i-1)$ for $1 \leq i \leq p_1$. In the following, we will make use of Stirling's formula

$$\log n! = n \log n - n + \frac{1}{2} \log(2\pi n) + \frac{1}{12n} + O(n^{-3}), \quad n \rightarrow \infty.$$

As a preparation, we note that

$$\begin{aligned} & \sum_{i=p_1+1}^p \log(n-i+1) - \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \log(n-j+1+p_{i-1}^*) = \log \frac{(n-p_1)!}{(n-p)!} - \sum_{i=2}^q \log \frac{n!}{(n-p_i)!} \\ &= \sum_{i=2}^q \left(n-p_i + \frac{1}{2} \right) \log \left(1 - \frac{p_i}{n} \right) - \left(n-p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right) - \sum_{i=2}^q \left(\frac{1}{12n} - \frac{1}{12(n-p_i)} \right) + o(1) \\ &= \mu_n + \frac{\sigma_n^2}{2} - \frac{1}{12} \sum_{i=2}^q \frac{p_i}{n(n-p_i)} + o(1) = \mu_n + \frac{\sigma_n^2}{2} + o(1), \end{aligned} \quad (14)$$

where we used the fact

$$\min_{1 \leq i \leq q} (n-p_i) \rightarrow \infty, \quad (15)$$

which is a consequence of our assumptions. Defining for $p_1+1 \leq i \leq p, 2 \leq j \leq q$

$$\begin{aligned} X_i &= \frac{\mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i - (n-i+1)}{n-i+1}, \quad X_{j,i} = \frac{\mathbf{b}_i^\top \mathbf{P}(p_{j-1}^*+1; i-1) \mathbf{b}_i - (n-i+1+p_{j-1}^*)}{n-i+1+p_{j-1}^*}, \\ Y_i &= \log(1+X_i) - \left(X_i - \frac{X_i^2}{2} \right), \quad Y_{j,i} = \log(1+X_{j,i}) - \left(X_{j,i} - \frac{X_{j,i}^2}{2} \right), \end{aligned}$$

we decompose, using (14),

$$\log V_n - \mu_n = \sum_{i=p_1+1}^p X_i - \sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} X_{j,i} - \left(\sum_{i=p_1+1}^p \frac{X_i^2}{2} - \sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} \frac{X_{j,i}^2}{2} - \frac{\sigma_n^2}{2} \right) + \sum_{i=p_1+1}^p Y_i - \sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} Y_{j,i} + o_{\text{Pr}}(1).$$

The assertion of Theorem 1 is then implied by Lemmas 1, 2 and 3. \square

Auxiliary results for the proof of Theorem 1

This section contains the auxiliary results needed for the proof of Theorem 1 and its proofs.

Lemma 1. *Under the assumptions of Theorem 1, it holds*

$$\frac{\sum_{i=p_1+1}^p X_i - \sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} X_{j,i}}{\sigma_n} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1), \quad n \rightarrow \infty.$$

Proof of Lemma 1. Let $\mathcal{F}_i = \sigma(\{\mathbf{b}_1, \dots, \mathbf{b}_i\})$ denote the σ -field generated by $\mathbf{b}_1, \dots, \mathbf{b}_i$ for $1 \leq i \leq p$. We write

$$\sum_{j=p_1+1}^p X_j - \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} X_{i,j} = \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} (X_j - X_{i,j}) = \sum_{i=p_1+1}^p Z_i,$$

where

$$Z_i = X_i - X_{g(i),i}, \quad p_1+1 \leq i \leq p,$$

where $g(i) = k$ if $\mathbf{x}^{(i)}$ belongs to the k th group, that is, if $p_{k-1} + 1 \leq i \leq p_k$. We observe that $\mathbb{E}[Z_i | \mathcal{F}_{i-1}] = 0$ and that Z_i is measurable with respect to \mathcal{F}_i for $p_1 + 1 \leq i \leq p$. Thus, we conclude that $(Z_i)_{p_1+1 \leq i \leq p}$ forms a martingale difference scheme with respect to the filtration $(\mathcal{F}_i)_{p_1+1 \leq i \leq p}$ scheme for every $n \in \mathbb{N}$. We aim to apply a central limit theorem for this dependency structure. In order to calculate the limiting variance, we write

$$Z_i = \mathbf{b}_i^\top \frac{(n-i+1 + p_{g(i-1)}^*)\mathbf{P}(i-1) - (n-i+1)\mathbf{P}(p_{g(i-1)}^* + 1; i-1)}{(n-i+1)(n-i+1 + p_{g(i-1)}^*)} \mathbf{b}_i \\ - \text{tr} \left(\frac{(n-i+1 + p_{g(i-1)}^*)\mathbf{P}(i-1) - (n-i+1)\mathbf{P}(p_{g(i-1)}^* + 1; i-1)}{(n-i+1)(n-i+1 + p_{g(i-1)}^*)} \right), \quad p_1 + 1 \leq i \leq p,$$

and use the fact

$$\mathbb{E} \left(\mathbf{b}_i^\top \mathbf{A} \mathbf{b}_i - \text{tr} \mathbf{A} \right)^2 = 2 \text{tr} \mathbf{A}^2 + (v_4 - 3) \text{tr} (\mathbf{A}^{\odot 2}), \quad 1 \leq i \leq p, \quad (16)$$

for any non-random matrix symmetric $\mathbf{A} \in \mathbb{R}^{n \times n}$, where $v_4 = \mathbb{E}[b_{11}^4]$. Here, \odot denotes the Hadamard product of matrices (entry-wise multiplication) and we use the notation

$$\mathbf{A} \odot \mathbf{A} = \mathbf{A}^{\odot 2}.$$

Consequently, we observe

$$\mathbb{E}[Z_i^2 | \mathcal{F}_{i-1}] = \mathbb{E}[(X_i - X_{g(i),i})^2 | \mathcal{F}_{i-1}] = 2 \left(\frac{\text{tr} \mathbf{P}(i-1)^2}{(n-i+1)^2} + \frac{\text{tr} \mathbf{P}(p_{g(i-1)}^* + 1; i-1)^2}{(n-i+1 + p_{g(i-1)}^*)^2} - 2 \frac{\text{tr} (\mathbf{P}(p_{g(i-1)}^* + 1; i-1) \mathbf{P}(i-1))}{(n-i+1 + p_{g(i-1)}^*)(n-i+1)} \right) \\ + (v_4 - 3) \left(\frac{\text{tr} (\mathbf{P}(i-1)^{\odot 2})}{(n-i+1)^2} + \frac{\text{tr} (\mathbf{P}(p_{g(i-1)}^* + 1; i-1)^{\odot 2})}{(n-i+1 + p_{g(i-1)}^*)^2} - 2 \frac{\text{tr} (\mathbf{P}(p_{g(i-1)}^* + 1; i-1) \odot \mathbf{P}(i-1))}{(n-i+1 + p_{g(i-1)}^*)(n-i+1)} \right) \\ = \sigma_{n,1,i}^2 + (v_4 - 3) \sigma_{n,2,i}^2,$$

where

$$\sigma_{n,1,i}^2 = 2 \left(\frac{1}{n-i+1} - \frac{1}{n-i+1 + p_{g(i-1)}^*} \right), \\ \sigma_{n,2,i}^2 = \frac{\text{tr} (\mathbf{P}(i-1)^{\odot 2})}{(n-i+1)^2} + \frac{\text{tr} (\mathbf{P}(p_{g(i-1)}^* + 1; i-1)^{\odot 2})}{(n-i+1 + p_{g(i-1)}^*)^2} - 2 \frac{\text{tr} (\mathbf{P}(p_{g(i-1)}^* + 1; i-1) \odot \mathbf{P}(i-1))}{(n-i+1 + p_{g(i-1)}^*)(n-i+1)}. \quad (17)$$

Note that $\sum_{i=p_1+1}^p \sigma_{n,2,i}^2 = o_{\text{Pr}}(1)$ by Lemma 5. For the term $\sigma_{n,1,i}^2$, we used that $\mathbf{P}(i-1)\mathbf{P}(p_{g(i-1)}^* + 1; i-1) = \mathbf{P}(i-1)$. Thus, we have for this term

$$\sum_{i=p_1+1}^p \sigma_{n,1,i}^2 = 2 \left(\log(n-p_1) - \log(n-p) - \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \frac{1}{n-j+1 + p_{i-1}^*} \right) + o(1) \\ = 2 \left\{ \log(n-p_1) - \log(n-p) - \sum_{i=2}^q \left\{ \log(n) - \log(n-p_i^* + p_{i-1}^*) \right\} - \sum_{i=2}^q \left(\frac{1}{2n} - \frac{1}{2(n-p_i)} \right) \right\} + o(1) \\ = 2 \left(\log \left(1 - \frac{p_1}{n} \right) - \log \left(1 - \frac{p}{n} \right) + \sum_{i=2}^q \log \left(1 - \frac{p_i}{n} \right) + \frac{1}{2} \sum_{i=2}^q \frac{p_i}{n(n-p_i)} \right) + o(1) \\ = 2 \left(\sum_{i=1}^q \log \left(1 - \frac{p_i}{n} \right) - \log \left(1 - \frac{p}{n} \right) \right) + o(1) = \sigma_n^2 + o(1), \quad (18)$$

where we used (15), $\sum_{i=2}^q p_i \leq p$ and the expansion for the partial sums of the harmonic series

$$\sum_{k=1}^n \frac{1}{k} = \log n + \gamma + \frac{1}{2n} + \mathcal{O}(n^{-2}), \quad n \rightarrow \infty. \quad (19)$$

Here, γ denotes the Euler-Mascheroni constant. Note that the term in (18) is bounded away from zero for all $n \in \mathbb{N}$. More precisely, we have applying inequality (33) of [44]

$$\sigma_n^2 \geq 2 \left\{ \sum_{i=1}^q \log \left(1 - \frac{p_i}{n} \right) - \log \left(1 - \frac{p}{n} \right) \right\} \geq -2 \left(\log \left(1 - \frac{p}{n} \right) + \frac{p}{n} \right) (1 - \eta) > 0 \quad (20)$$

uniformly over $n \in \mathbb{N}$ (recall that $\inf_{n \in \mathbb{N}} p/n > 0$ and $\max_{1 \leq i \leq q} p_i \leq \eta p$). These considerations imply

$$\sum_{i=p_1+1}^p \mathbb{E} \left[\frac{Z_i^2}{\sigma_n^2} \middle| \mathcal{F}_{i-1} \right] \xrightarrow{\text{Pr}} 1, \quad n \rightarrow \infty. \quad (21)$$

Let $\varepsilon > 0$. Then, using Lemma B.26 in [3] and recalling that $\mathbb{E}|x_{11}|^{4+\delta} < \infty$, we get

$$\begin{aligned} \sum_{i=p_1+1}^p \mathbb{E}[Z_i^2 I\{|Z_i| > \varepsilon\}] &\leq \frac{1}{\varepsilon^{\frac{\delta}{2}}} \sum_{i=p_1+1}^p \mathbb{E}[|Z_i|^{2+\frac{\delta}{2}}] \lesssim \sum_{i=p_1+1}^p \frac{1}{(n-i+1)^{2+\frac{\delta}{2}}} \mathbb{E} |\mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i - (n-i+1)|^{2+\frac{\delta}{2}} \\ &+ \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \frac{1}{(n-j+1+p_{i-1}^*)^{2+\frac{\delta}{2}}} \mathbb{E} |\mathbf{b}_j^\top \mathbf{P}(p_{i-1}^*+1; i-1) \mathbf{b}_j - (n-j+1+p_{i-1}^*)|^{2+\frac{\delta}{2}} \\ &\lesssim \sum_{i=p_1+1}^p \frac{1}{(n-i+1)^{1+\frac{\delta}{4}}} + \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \frac{1}{(n-j+1+p_{i-1}^*)^{1+\frac{\delta}{4}}} = o(1), \quad n \rightarrow \infty. \end{aligned}$$

Using (20) and the fact that $(\sigma_n^2)_{n \in \mathbb{N}}$ is bounded, we see that $(Z_i/\sigma_n)_{p_1+1 \leq i \leq p}$ satisfies the following Lindeberg condition for all $\varepsilon > 0$:

$$\sum_{i=p_1+1}^p \mathbb{E} \left[\frac{Z_i^2}{\sigma_n^2} I\{|Z_i/\sigma_n| > \varepsilon\} \right] = o(1), \quad n \rightarrow \infty. \quad (22)$$

Since (21) and (22) hold true, we may apply a CLT for martingale difference schemes (e.g., see Corollary 3.1 in [20]) and the proof of Lemma 1 concludes. \square

Lemma 2. *Under the assumptions of Theorem 1, it holds*

$$\frac{\sum_{i=p_1+1}^p \frac{X_i^2}{2} - \sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} \frac{X_{i,j}^2}{2} - \frac{\sigma_n^2}{2}}{\sigma_n} \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty.$$

Proof of Lemma 2. Note that for each $n \in \mathbb{N}$, both $(X_i)_{p_1+1 \leq i \leq p}$ and $(X_{g(i),i})_{p_1+1 \leq i \leq p}$ form a martingale difference scheme with respect to the filtration $(\mathcal{F}_i)_{p_1+1 \leq i \leq p}$ defined previously. We obtain from the proof of Lemma 1

$$\sum_{i=p_1+1}^p \mathbb{E}[X_i^2 | \mathcal{F}_{i-1}] = \check{\sigma}_{n,1}^2 + o_{\text{Pr}}(1), \quad n \rightarrow \infty, \quad (23)$$

and

$$\sum_{i=2}^q \mathbb{E} \left[X_{g(i),i}^2 \middle| \mathcal{F}_{i-1} \right] = \sum_{i=2}^q \sum_{j=p_{i-1}^*+1}^{p_i^*} \mathbb{E}[X_{i,j}^2 | \mathcal{F}_{i-1}] = \check{\sigma}_{n,2}^2 + o_{\text{Pr}}(1), \quad n \rightarrow \infty, \quad (24)$$

where we define

$$\begin{aligned}\check{\sigma}_{n,1}^2 &= -2 \log\left(1 - \frac{p}{n}\right) + (v_4 - 3) \sum_{i=p_1+1}^p \frac{\text{tr}(\mathbf{P}(i-1)^{\otimes 2})}{(n-i+1)^2}, \\ \check{\sigma}_{n,2}^2 &= -2 \sum_{i=1}^q \log\left(1 - \frac{p_i}{n}\right) + (v_4 - 3) \sum_{i=p_1+1}^p \frac{\text{tr}(\mathbf{P}(p_{g(i)-1}^* + 1; i-1)^{\otimes 2})}{(n-i+1 + p_{g(i)-1}^*)^2}.\end{aligned}$$

Recalling that $0 < \inf_{n \in \mathbb{N}} \min_{1 \leq i \leq q} (p_i q) / n \leq \sup_{n \in \mathbb{N}} p/n < 1$ and using the inequality $\log(1+x) \leq x$ for $x > -1$, we note that

$$0 < \inf_{n \in \mathbb{N}} \check{\sigma}_{n,1}^2 \leq \sup_{n \in \mathbb{N}} \check{\sigma}_{n,1}^2 < \infty, \quad 0 < \inf_{n \in \mathbb{N}} \check{\sigma}_{n,2}^2 \leq \sup_{n \in \mathbb{N}} \check{\sigma}_{n,2}^2 < \infty. \quad (25)$$

Taking a closer look at the proof of (22), we observe that both schemes satisfy the Lindeberg condition, that is, we have for $\varepsilon > 0$

$$\sum_{i=p_1+1}^p \mathbb{E} \left[\frac{X_i^2}{\check{\sigma}_{n,1}^2} I\{|X_i|/\check{\sigma}_{n,1} > \varepsilon\} \right] = o(1), \quad (26)$$

$$\sum_{i=p_1+1}^p \mathbb{E} \left[\frac{X_{g(i),i}^2}{\check{\sigma}_{n,2}^2} I\{|X_{g(i),i}|/\check{\sigma}_{n,2} > \varepsilon\} \right] = o(1). \quad (27)$$

By Theorem 2.23 in [20], we see that (23), (24), (26) and (27) imply that the conditional variance can be approximated by the sum of squares, that is,

$$\sum_{i=p_1+1}^p \frac{X_i^2 - \mathbb{E}[X_i^2 | \mathcal{F}_{i-1}]}{\check{\sigma}_{n,1}^2} \xrightarrow{\text{Pr}} 0, \quad \sum_{i=p_1+1}^p \frac{X_{g(i),i}^2 - \mathbb{E}[X_{g(i),i}^2 | \mathcal{F}_{i-1}]}{\check{\sigma}_{n,2}^2} \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty,$$

Combining these observations with (23), (24) and (25), we get

$$\sum_{i=p_1+1}^p X_i^2 - \check{\sigma}_{n,1}^2 \xrightarrow{\text{Pr}} 0, \quad \sum_{i=p_1+1}^p X_{g(i),i}^2 - \check{\sigma}_{n,2}^2 \xrightarrow{\text{Pr}} 0.$$

Using (20) and $\check{\sigma}_{n,1}^2 - \check{\sigma}_{n,2}^2 = \sigma_n^2 + o_{\text{Pr}}(1)$ by Lemma 4, the proof of Lemma 2 concludes. \square

Lemma 3. *Under the assumptions of Theorem 1, it holds*

$$\sum_{i=p_1+1}^p \frac{Y_i}{\sigma_n} \xrightarrow{\text{Pr}} 0, \quad (28)$$

$$\sum_{j=2}^q \sum_{i=p_{j-1}^*+1}^{p_j^*} \frac{Y_{ji}}{\sigma_n} \xrightarrow{\text{Pr}} 0, \quad (29)$$

as $n \rightarrow \infty$.

Proof of Lemma 3. In the following, we will show that the convergence in (28) holds true. Then, the assertion (29) can be shown similarly.

Let $0 < \varepsilon < 1$. Then, we estimate for $1 + p_1 \leq i \leq p$ using Taylor's expansion

$$\mathbb{E} [Y_i I\{|X_i| \leq 1 - \varepsilon\}] \lesssim \mathbb{E} [X_i^3 I\{|X_i| \leq 1 - \varepsilon\}] \lesssim \mathbb{E} |X_i|^{2+\frac{\delta}{2}}.$$

We also have

$$\begin{aligned} \mathbb{E} [|Y_i|I\{|X_i| > 1 - \varepsilon\}] &\leq \mathbb{E} [|\log(1 + X_i)|I\{|X_i| > 1 - \varepsilon\}] + \mathbb{E} [|X_i|I\{|X_i| > 1 - \varepsilon\}] + \mathbb{E} [X_i^2 I\{|X_i| > 1 - \varepsilon\}] \\ &\lesssim \mathbb{E} [|X_i|I\{|X_i| > 1 - \varepsilon\}] + \mathbb{E} [X_i^2 I\{|X_i| > 1 - \varepsilon\}] \lesssim \mathbb{E}|X_i|^{2+\frac{\varepsilon}{2}}, \end{aligned}$$

where we used the inequality $\log(1 + x) \leq x$ for all $x > -1$. These two estimates imply

$$\sum_{i=p_1+1}^p \mathbb{E}|Y_i| \lesssim \sum_{i=p_1+1}^p \mathbb{E}|X_i|^{2+\frac{\varepsilon}{2}} = o(1), \quad n \rightarrow \infty,$$

where we used Lemma B.26 in [3] as in the proof of (22). Thus, we obtain

$$\sum_{i=p_1+1}^p Y_i \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty,$$

which implies the assertion of Lemma 3 recalling $\inf_{n \in \mathbb{N}} \sigma_n^2 > 0$. □

Lemma 4. *It holds, as $n \rightarrow \infty$,*

$$\sum_{i=p_1+1}^p \left(\frac{\text{tr}(\mathbf{P}(i-1)^{\otimes 2})}{(n-i+1)^2} - \frac{1}{n} \right) \xrightarrow{\text{Pr}} 0, \quad (30)$$

$$\sum_{i=p_1+1}^p \left(\frac{\text{tr}(\mathbf{P}(p_{g(i)-1}^* + 1; i-1)^{\otimes 2})}{(n-i+1+p_{g(i)-1}^*)^2} - \frac{1}{n} \right) \xrightarrow{\text{Pr}} 0, \quad (31)$$

where the projection matrices are defined in (13).

Proof of Lemma 4. As a preparation, we will first show that for any sequence $(i_n)_{n \in \mathbb{N}}$ such that $2 \leq i_n \leq p_n$ for all $n \in \mathbb{N}$ and the limit $\lim_{n \rightarrow \infty} i_n/n \in [0, 1)$ exists, it holds

$$a_{i_n, n} - b_{i_n, n} \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty, \quad (32)$$

where we define for $2 \leq i \leq p$

$$a_{i, n} = \frac{\text{tr}(\mathbf{P}(i-1)^{\otimes 2})}{n-i+1}, \quad b_{i, n} = \left(1 - \frac{i}{n}\right), \quad c_n = \sum_{i=p_1+1}^p \frac{a_{i, n} - b_{i, n}}{n-i+1}.$$

In the following, we denote the diagonal entries of $\mathbf{P}(i_n-1)$ by p_{ii} ($1 \leq i \leq n$). First, we consider the case $\lim_{n \rightarrow \infty} i_n/n = 0$. For this case, we note that

$$\begin{aligned} \frac{\text{tr}(\mathbf{P}(i_n-1)^{\otimes 2})}{n} &= \frac{1}{n} \sum_{i=1}^n (1-p_{ii})^2 - 1 + \frac{2}{n} \sum_{i=1}^n p_{ii} = \frac{2(n-i_n+1)}{n} - 1 + o_{\text{Pr}}(1) \\ &= 1 + o_{\text{Pr}}(1), \quad n \rightarrow \infty, \end{aligned}$$

where we used

$$\frac{1}{n} \sum_{i=1}^n \mathbb{E}(1-p_{ii})^2 \leq \frac{1}{n} \sum_{i=1}^n \mathbb{E}[1-p_{ii}] = \frac{1}{n} \text{tr}(\mathbf{I} - \mathbf{P}) = \frac{i_n-1}{n} = o(1), \quad n \rightarrow \infty.$$

In this case, we conclude

$$a_{i_n, n} - b_{i_n, n} = \frac{\text{tr}(\mathbf{P}(i_n-1)^{\otimes 2})}{n} - 1 + o_{\text{Pr}}(1) = o_{\text{Pr}}(1), \quad n \rightarrow \infty.$$

Now consider the case $\lim_{n \rightarrow \infty} i_n/n = \gamma \in (0, 1)$. Then we have from Theorem 3.2 in [1]

$$\frac{1}{n} \sum_{i=1}^n (1 - p_{ii} - \gamma)^2 \xrightarrow{\Pr} 0, \quad n \rightarrow \infty,$$

which implies

$$\begin{aligned} \frac{\text{tr}(\mathbf{P}(i_n - 1)^{\odot 2})}{n} &= \frac{1}{n} \sum_{i=1}^n (1 - p_{ii} - \gamma)^2 - (1 - \gamma)^2 + \frac{2(1 - \gamma)}{n} \sum_{i=1}^n p_{ii} \\ &= \frac{2(1 - \gamma)(n - i_n + 1)}{n} - (1 - \gamma)^2 + o_{\Pr}(1) = (1 - \gamma)^2 + o_{\Pr}(1), \quad n \rightarrow \infty, \end{aligned}$$

which implies that (32) holds also true in this case. We continue with a proof of (30) by showing that any subsequence of $(c_n)_{n \in \mathbb{N}}$ admits a further subsequence converging in probability to 0. Let $(c_{n_j})_{j \in \mathbb{N}}$ be an arbitrary subsequence of $(c_n)_{n \in \mathbb{N}}$. We choose

$$i_{n_j} \in \arg \max_{p_1+1 \leq i \leq p} (a_{i, n_j} - b_{i, n_j}).$$

Not that there exists a subsequence $(i_{n_{j_k}})_{k \in \mathbb{N}}$ of $(i_{n_j})_{j \in \mathbb{N}}$ which admits a limit $\lim_{k \rightarrow \infty} i_{n_{j_k}}/k \in [0, 1)$ (that is, this subsequence satisfies the assumption for (32)). Then, it holds using (32)

$$c_{n_{j_k}} \lesssim \max_{p_1+1 \leq i \leq p} (a_{i, n_{j_k}} - b_{i, n_{j_k}}) = a_{i_{n_{j_k}}, n_{j_k}} - b_{i_{n_{j_k}}, n_{j_k}} \xrightarrow{\Pr} 0, \quad n \rightarrow \infty.$$

This implies the convergence $c_n \xrightarrow{\Pr} 0$ of the whole sequence $(c_n)_{n \in \mathbb{N}}$ for $n \rightarrow \infty$ and thus, the convergence in (35) holds true. The second assertion (31) of Lemma 4 can be shown similarly. \square

Lemma 5. *It holds*

$$\sum_{i=p_1+1}^p \sigma_{n,2,i}^2 \xrightarrow{\Pr} 0, \quad n \rightarrow \infty,$$

where the term $\sigma_{n,2,i}^2$ is defined in (17).

Proof of Lemma 5. Recalling the definition of $\sigma_{n,2,i}^2$ and using Lemma 4, it suffices to show

$$\sum_{i=p_1+1}^p \left(\text{tr} \mathbf{A}_i \odot \mathbf{B}_i - \frac{1}{n} \right) \xrightarrow{\Pr} 0, \quad n \rightarrow \infty,$$

where we denote

$$\mathbf{A}_i = \frac{\mathbf{P}(i-1)}{n-i+1}, \quad \mathbf{B}_i = \frac{\mathbf{P}(p_{g(i)-1}^* + 1; i-1)}{n-i+1 + p_{g(i)-1}^*}, \quad p_1 + 1 \leq i \leq p.$$

Note that $\text{tr} \mathbf{A}_i = \text{tr} \mathbf{B}_i = 1$ for all $p_1 + 1 \leq i \leq p$. This gives

$$\begin{aligned} \sum_{i=p_1+1}^p \left(\text{tr} \mathbf{A}_i \odot \mathbf{B}_i - \frac{1}{n} \right) &= \sum_{i=p_1+1}^p \text{tr} \left\{ \left(\mathbf{A}_i - \frac{1}{n} \mathbf{I} \right) \odot \left(\mathbf{B}_i - \frac{1}{n} \mathbf{I} \right) \right\} \leq \sum_{i=p_1+1}^p \left\{ \text{tr} \left(\mathbf{A}_i - \frac{1}{n} \mathbf{I} \right)^{\odot 2} \text{tr} \left(\mathbf{B}_i - \frac{1}{n} \mathbf{I} \right)^{\odot 2} \right\}^{\frac{1}{2}} \\ &\leq \left\{ \sum_{i=p_1+1}^p \text{tr} \left(\mathbf{A}_i - \frac{1}{n} \mathbf{I} \right)^{\odot 2} \sum_{i=p_1+1}^p \text{tr} \left(\mathbf{B}_i - \frac{1}{n} \mathbf{I} \right)^{\odot 2} \right\}^{\frac{1}{2}} = \left\{ \sum_{i=p_1+1}^p \left(\text{tr} \mathbf{A}_i^{\odot 2} - \frac{1}{n} \right) \sum_{i=p_1+1}^p \left(\text{tr} \mathbf{B}_i^{\odot 2} - \frac{1}{n} \right) \right\}^{\frac{1}{2}} \\ &\xrightarrow{\Pr} 0, \end{aligned}$$

as $n \rightarrow \infty$, where we applied the Cauchy-Schwarz inequality twice and Lemma 4. \square

Proof of Theorem 2. For proving Theorem 2, we need the following properties of the variance.

Lemma 6. *Under the assumptions of Theorem 2, we have*

$$0 < \inf_{n \in \mathbb{N}} \sigma_n^2 \leq \sup_{n \in \mathbb{N}} \sigma_n^2 < \infty,$$

where σ_n^2 denotes the variance defined in (9).

Proof of Lemma 6. Define the functions

$$\xi(x) = -(\log(1-x) + x), \quad \eta(x) = \frac{\xi(x)}{x^2},$$

where $x \in (0, 1)$. Note that η is a monotone increasing function with $\eta(x) \geq 1/2$. Using $\sum_{j=1}^q n_j = n$ and the definition $n_{\max} = \max_{1 \leq j \leq q} n_j$, we obtain the estimate

$$\begin{aligned} n^2 \sigma_n^2 &= \sum_{j=1}^q n_j^2 \xi\left(\frac{p}{n_j}\right) - n^2 \xi\left(\frac{p}{n}\right) = p^2 \left\{ \sum_{j=1}^q \eta\left(\frac{p}{n_j}\right) - \eta\left(\frac{p}{n}\right) \right\} \\ &\geq p^2 \left\{ q \eta\left(\frac{p}{n_{\max}}\right) - \eta\left(\frac{p}{n_{\max}}\right) \right\} = p^2 (q-1) \eta\left(\frac{p}{n_{\max}}\right). \end{aligned}$$

Using $\inf_{n \in \mathbb{N}} p/n > 0$, we conclude $\inf_{n \in \mathbb{N}} \sigma_n^2 > 0$. Moreover, from our assumption on the dimension-to-subsample-size ratios p/n_j , we obtain

$$\sup_{n \in \mathbb{N}} \sigma_n^2 \leq \sup_{n \in \mathbb{N}} \max_{1 \leq j \leq q} \left(-\log\left(1 - \frac{p}{n_j}\right) \right) < \infty.$$

This concludes the proof of Lemma 6. □

Continuing with the proof of Theorem 2, we follow the same strategy as in the proof of Theorem 1 and concentrate on discussing the main steps. Recalling the definition (8) of the likelihood ratio test statistic, we note that under the null hypothesis (7)

$$\begin{aligned} 2 \log \Lambda_{n,1} &= \sum_{j=1}^q n_j \log |\mathbf{A}_j| - n \log |\mathbf{A}| + pn \log n - \sum_{j=1}^q n_j p \log n_j \\ &= \sum_{j=1}^q n_j \log |n_j \hat{\mathbf{\Gamma}}_j| - n \log |n \hat{\mathbf{\Gamma}}| + pn \log n - \sum_{j=1}^q n_j p \log n_j, \end{aligned}$$

where $\hat{\mathbf{\Gamma}}$ is defined in the proof of Theorem 1 and

$$\hat{\mathbf{\Gamma}}_j = \frac{1}{n_j} \sum_{k=1}^{n_j} \mathbf{x}_{jk} \mathbf{x}_{jk}^\top.$$

Applying the QR-procedure to the matrices $\hat{\mathbf{\Gamma}}$ and $\hat{\mathbf{\Gamma}}_j$ ($1 \leq j \leq q$), we obtain for their determinants

$$|n \hat{\mathbf{\Gamma}}| = \prod_{i=1}^p \mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i, \quad |n_j \hat{\mathbf{\Gamma}}_j| = \prod_{i=1}^p \mathbf{b}_{ji}^\top \mathbf{P}(j; i-1) \mathbf{b}_{ji},$$

where

$$\mathbf{b}_i = (\mathbf{b}_{1i}^\top, \dots, \mathbf{b}_{qi}^\top)^\top \in \mathbb{R}^n, \quad \mathbf{b}_{ji} \in \mathbb{R}^{n_j}, \quad 1 \leq j \leq q,$$

and throughout this proof, $\mathbf{P}(j; i-1) \in \mathbb{R}^{n_j \times n_j}$ denotes the projection matrix on the orthogonal complement of

$$\text{span}\{\mathbf{b}_{j1}, \dots, \mathbf{b}_{j,i-1}\}$$

for $1 \leq i \leq p$, $1 \leq j \leq q$ (note that we have a different definition than in the proof of Theorem 1). We set $\mathbf{P}(j; 0) = \mathbf{I} \in \mathbb{R}^{n_j \times n_j}$. The remaining quantities are defined as in the proof of Theorem 1. With a slight abuse of notation, we define for $1 \leq i \leq p$, $1 \leq j \leq q$

$$X_i = \frac{\mathbf{b}_i^\top \mathbf{P}(i-1) \mathbf{b}_i - (n-i+1)}{n-i+1}, \quad X_{j,i} = \frac{\mathbf{b}_{j,i}^\top \mathbf{P}(j; i-1) \mathbf{b}_{j,i} - (n_j-i+1)}{n_j-i+1},$$

$$Y_i = \log(1 + X_i) - \left(X_i - \frac{X_i^2}{2} \right), \quad Y_{j,i} = \log(1 + X_{j,i}) - \left(X_{j,i} - \frac{X_{j,i}^2}{2} \right).$$

Similarly to (14), we obtain using Stirling's formula

$$\begin{aligned} & \sum_{i=1}^p \sum_{j=1}^q n_j \log(n_j - i + 1) - n \sum_{i=1}^p \log(n - i + 1) + pn \log n - \sum_{j=1}^q n_j p \log n_j \\ &= \sum_{j=1}^q n_j \log \left(\frac{n_j!}{(n_j - p)!} \right) - n \log \left(\frac{n!}{(n - p)!} \right) + pn \log n - \sum_{j=1}^q n_j p \log n_j \\ &= n \left(n - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \left(n_j - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n_j} \right) + \sum_{j=1}^q \frac{n_j}{12} \left(\frac{1}{n_j} - \frac{1}{n_j - p} \right) + o(n) \\ &= n \left(n - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \left(n_j - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n_j} \right) - \sum_{j=1}^q \frac{p}{12(n_j - p)} + o(n) \\ &= n \left(n - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \left(n_j - p + \frac{1}{2} \right) \log \left(1 - \frac{p}{n_j} \right) + o(n) \\ &= \mu_n + n \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \log \left(1 - \frac{p}{n_j} \right) - \frac{1}{2}(\nu_4 - 3)p(1 - q) + o(n), \quad n \rightarrow \infty. \end{aligned} \quad (33)$$

For (33), we used that under the assumptions of Theorem 2

$$\sum_{j=1}^q \frac{p}{12n(n_j - p)} \lesssim \frac{q}{n} \leq \frac{1}{\min_j n_j} = o(1), \quad n \rightarrow \infty. \quad (34)$$

Consequently, we may decompose

$$\begin{aligned} & 2(\log \Lambda_{n,1} - \mu_n) \\ &= \sum_{j=1}^q \sum_{i=1}^p n_j X_{j,i} - n \sum_{i=1}^p X_i - \left(\sum_{j=1}^q \sum_{i=1}^p n_j \frac{X_{j,i}^2}{2} - n \sum_{i=1}^p \frac{X_i^2}{2} - \tau_n \right) + \sum_{j=1}^q \sum_{i=1}^p n_j Y_{j,i} - n \sum_{i=1}^p Y_i + o(n), \quad n \rightarrow \infty, \end{aligned}$$

where

$$\tau_n = 2 \left\{ n \log \left(1 - \frac{p}{n} \right) - \sum_{j=1}^q n_j \log \left(1 - \frac{p}{n_j} \right) \right\} + (\nu_4 - 3)p(q - 1).$$

Note that $(W_i)_{1 \leq i \leq p}$ with

$$W_i = \sum_{j=1}^q n_j X_{j,i} - n X_i, \quad 1 \leq i \leq p,$$

forms a martingale difference scheme with respect to the filtration $(\mathcal{A}_i)_{1 \leq i \leq p}$, where the σ -field \mathcal{A}_i is generated by the random variables $\mathbf{b}_1, \dots, \mathbf{b}_i$ for $1 \leq i \leq p$. One can show that

$$\frac{\sum_{j=1}^q \sum_{i=1}^p n_j \frac{X_{j,i}^2}{2} - n \sum_{i=1}^p \frac{X_i^2}{2} - \tau_n}{n\sigma_n} \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty, \quad (35)$$

and

$$\frac{\sum_{j=1}^q \sum_{i=1}^p n_j Y_{j,i} - n \sum_{i=1}^p Y_i}{n\sigma_n} \xrightarrow{\text{Pr}} 0, \quad n \rightarrow \infty. \quad (36)$$

For the sake of brevity, we omit the proofs of (35) and (36) as they are very similar to the proofs of Lemma 2 and Lemma 3. For (35), we additionally note that (similarly to Lemma 5)

$$n \sum_{i=1}^p \frac{\text{tr} \mathbf{P}(i-1)^{\otimes 2}}{(n-i+1)^2} - \sum_{j=1}^q n_j \sum_{i=1}^p \frac{\text{tr} \mathbf{P}(j; i-1)^{\otimes 2}}{(n_j-i+1)^2} = \frac{np}{n} - \sum_{j=1}^q \frac{n_j p}{n_j} + o_{\text{Pr}}(n) = p - qp + o_{\text{Pr}}(n) = p(1-q) + o_{\text{Pr}}(n).$$

We continue with a proof of the asymptotic normality of the scheme $(W_i/(n\sigma_n))_{1 \leq i \leq p}$. To begin with, we show that

$$\sum_{i=1}^p \mathbb{E} \left[\left(\frac{W_i}{n\sigma_n} \right)^2 \middle| \mathcal{A}_{i-1} \right] = 1 + o_{\text{Pr}}(1), \quad n \rightarrow \infty. \quad (37)$$

As a preparation for (37), note that

$$\begin{aligned} \sum_{i=1}^p \sum_{j=1}^q n_j n \mathbb{E}[X_i X_{j,i} | \mathcal{A}_{i-1}] &= \sum_{i=1}^p \sum_{j=1}^q n n_j \left\{ \sum_{k=n_{j-1}^*+1}^{n_j^*} (\nu_4 - 3) \frac{(\mathbf{P}(j; i-1))_{kk} (\mathbf{P}(i-1))_{kk}}{(n-i+1)(n_j-i+1)} + 2 \frac{\text{tr}(\mathbf{P}(j-1) \tilde{\mathbf{P}}(j; i-1))}{(n-i+1)(n_j-i+1)} \right\} \\ &= \sum_{i=1}^p \sum_{j=1}^q n n_j \left\{ (\nu_4 - 3) \frac{\text{tr}(\tilde{\mathbf{P}}(j; i-1) \odot \mathbf{P}(i-1))}{(n-i+1)(n_j-i+1)} + 2 \frac{\text{tr} \tilde{\mathbf{P}}(j; i-1)}{(n_j-i+1)(n-i+1)} \right\} \\ &= (\nu_4 - 3) \sum_{i=1}^p \sum_{j=1}^q n n_j \frac{\text{tr}(\tilde{\mathbf{P}}(j; i-1) \odot \mathbf{P}(i-1))}{(n-i+1)(n_j-i+1)} + \sum_{i=1}^p \frac{2n^2}{(n-i+1)}, \end{aligned}$$

where $\tilde{\mathbf{P}}(j; i-1)$ denotes the $(n \times n)$ dimensional embedded matrix of $\mathbf{P}(j; i-1) \in \mathbb{R}^{n_j \times n_j}$, that is,

$$(\tilde{\mathbf{P}}(j; i-1))_{kl} = \begin{cases} (\mathbf{P}(j; i-1))_{kl}, & n_{j-1}^* + 1 \leq k, l \leq n_j^*, \\ 0, & \text{else,} \end{cases}$$

for $1 \leq k, l \leq n$, $1 \leq j \leq q$, $1 \leq i \leq p$. In order to prove (37), we calculate

$$\begin{aligned} \sum_{i=1}^p \mathbb{E}[W_i^2 | \mathcal{A}_{i-1}] &= n^2 \sum_{i=1}^p \mathbb{E}[X_i^2 | \mathcal{A}_{i-1}] + \sum_{i=1}^p \sum_{j=1}^q n_j^2 \mathbb{E}[X_{j,i}^2 | \mathcal{A}_{i-1}] - 2 \sum_{i=1}^p \sum_{j=1}^q n_j n \mathbb{E}[X_i X_{j,i} | \mathcal{A}_{i-1}] \\ &= 2 \sum_{j=1}^q \sum_{i=1}^p n_j^2 \frac{1}{n_j - i + 1} - 2n^2 \sum_{i=1}^p \frac{1}{n - i + 1} + \tilde{\sigma}_{n,2}^2, \end{aligned}$$

where we used the fact $\mathbb{E}[X_{j,i} X_{j',i} | \mathcal{A}_{i-1}] = 0$ for different groups $j, j' \in \{1, \dots, q\}$, $j \neq j'$ and (16) and we define

$$\tilde{\sigma}_{n,2}^2 = (\nu_4 - 3) \left\{ n^2 \sum_{i=1}^p \frac{\text{tr} \mathbf{P}(i-1)^{\otimes 2}}{(n-i+1)^2} + \sum_{i=1}^p \sum_{j=1}^q n_j^2 \frac{\text{tr} \mathbf{P}(j; i-1)^{\otimes 2}}{(n_j-i+1)^2} - 2 \sum_{i=1}^p \sum_{j=1}^q n n_j \frac{\text{tr}(\tilde{\mathbf{P}}(j; i-1) \odot \mathbf{P}(i-1))}{(n-i+1)(n_j-i+1)} \right\}.$$

Note that $\tilde{\sigma}_{n,2}^2/n^2 = o_{\text{Pr}}(1)$ as $n \rightarrow \infty$ (similarly to Lemma 4 and Lemma 5). Using (19) and (34), we obtain

$$\begin{aligned} \sum_{i=1}^p \mathbb{E} \left[\left(\frac{W_i}{n} \right)^2 \mid \mathcal{A}_{i-1} \right] &= \sum_{i=1}^p \log \left(1 - \frac{p}{n} \right) - \sum_{i=1}^p \sum_{j=1}^q \left(\frac{n_j}{n} \right)^2 \log \left(1 - \frac{p}{n_j} \right) + \sum_{j=1}^q \left(\frac{n_j}{n} \right)^2 \left\{ \frac{1}{2n_j} - \frac{1}{2(n_j - p)} \right\} + \frac{\tilde{\sigma}_{n,2}^2}{n^2} + o(1) \\ &= \sigma_n^2 + o_{\text{Pr}}(1), \quad n \rightarrow \infty, \end{aligned}$$

which implies (37) by an application of Lemma 6. Note that the Lindeberg condition for the scheme $(W_i/(n\sigma_n))_{1 \leq i \leq p}$ can be shown similarly to (22) using Lemma 6. Combining (35) and (36), we conclude

$$\frac{2(\log \Lambda_{n,1} - \mu_n)}{n\sigma_n} = \frac{\sum_{j=1}^q \sum_{i=1}^p n_j X_{j,i} - n \sum_{i=1}^p X_i}{n\sigma_n} + o_{\text{Pr}}(1), \quad n \rightarrow \infty.$$

By an application of Corollary 3.1 of [20], we obtain

$$\frac{\sum_{j=1}^q \sum_{i=1}^p n_j X_{j,i} - n \sum_{i=1}^p X_i}{n\sigma_n} \xrightarrow{\mathcal{D}} \mathcal{N}(0, 1), \quad n \rightarrow \infty.$$

The proof Theorem 2 concludes. \square

Acknowledgments

This work was supported by the German Research Foundation (DFG Research Unit 1735, DE 502/26-2, RTG 2131, *High-dimensional Phenomena in Probability - Fluctuations and Discontinuity* and project number 460867398, DFG Research unit 5381, *Mathematical Statistics in the Information Age*). The author would like to thank Tim Kutta for his helpful comments on an earlier version of this manuscript. The very constructive comments of the Editor, Associate Editor and referees are kindly acknowledged.

References

- [1] S. Anatolyev, P. Yaskov, Asymptotics of diagonal elements of projection matrices under many instruments/regressors, *Econometric Theory* 33 (2017) 717–738.
- [2] T. W. Anderson, *An introduction to multivariate statistical analysis*, Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics, John Wiley & Sons, Inc., New York, second edition, 1984.
- [3] Z. Bai, J. W. Silverstein, *Spectral analysis of large dimensional random matrices*, volume 20, Springer, 2010.
- [4] Z. Bao, J. Hu, G. Pan, W. Zhou, Test of independence for high-dimensional random vectors based on freeness in block correlation matrices, *Electronic Journal of Statistics* 11 (2017) 1527–1548.
- [5] Z. Bao, G. Pan, W. Zhou, The logarithmic law of random determinant, *Bernoulli* 21 (2015) 1600–1628.
- [6] M. R. Barrick, M. K. Mount, The big five personality dimensions and job performance: a meta-analysis, *Personnel psychology* 44 (1991) 1–26.
- [7] T. Bodnar, H. Dette, N. Parolya, Testing for independence of large dimensional vectors, *The Annals of Statistics* 47 (2019) 2977–3008.
- [8] A. D. Carson, The integration of interests, aptitudes, and personality traits: A test of lowman’s matrix, *Journal of Career Assessment* 6 (1998) 83–105.
- [9] H. E. Cattell, A. D. Mead, The sixteen personality factor questionnaire (16pf), *The SAGE handbook of personality theory and assessment* 2 (2008) 135–159.
- [10] R. B. Cattell, *Personality and motivation structure and measurement*. (1957).
- [11] R. B. Cattell, *Personality and mood by questionnaire*, Jossey-Bass, 1973.
- [12] R. B. Cattell, S. E. Krug, The number of factors in the 16pf: A review of the evidence with special emphasis on methodological problems, *Educational and Psychological Measurement* 46 (1986) 509–522.
- [13] B. De Raad, *The big five personality factors: the psycholexical approach to personality*, Hogrefe & Huber Publishers, 2000.
- [14] H. Dette, N. Dörnemann, Likelihood ratio tests for many groups in high dimensions, *Journal of Multivariate Analysis* 178 (2020) 104605.
- [15] J. Fan, R. Li, Statistical challenges with high dimensionality: Feature selection in knowledge discovery, *Proceedings of the International Congress of Mathematicians, Madrid* 3 (2006).
- [16] J. Gao, X. Han, G. Pan, Y. Yang, High dimensional correlation matrices: The central limit theorem and its applications, *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* 79 (2017) 677–693.
- [17] V. L. Girko, A refinement of the central limit theorem for random determinants, *Theory of Probability & Its Applications* 42 (1998) 121–129.

- [18] S. D. Gosling, P. J. Rentfrow, W. B. Swann Jr, A very brief measure of the big-five personality domains, *Journal of Research in personality* 37 (2003) 504–528.
- [19] W. Guo, Y. Qi, Asymptotic distributions for likelihood ratio tests for the equality of covariance matrices, arXiv preprint arXiv:2110.02384 (2021).
- [20] P. Hall, C. C. Heyde, *Martingale limit theory and its application*, Academic press, 1980.
- [21] F. Han, S. Chen, H. Liu, Distribution-free tests of independence in high dimensions, *Biometrika* 104 (2017) 813–828.
- [22] Y. He, G. Xu, C. Wu, W. Pan, Asymptotically independent u-statistics in high-dimensional testing, *The Annals of Statistics* 49 (2021) 154–181.
- [23] J. Heiny, N. Parolya, Log determinant of large correlation matrices under infinite fourth moment, arXiv preprint arXiv:2112.15388 (2021).
- [24] J. Hu, Z. Bai, C. Wang, W. Wang, On testing the equality of high dimensional mean vectors with unequal covariance matrices, *Annals of the Institute of Statistical Mathematics* 69 (2017) 365–387.
- [25] R. Ishii, K. Maruo, H. Noma, M. Goshio, Statistical inference based on accelerated failure time models under model misspecification and small samples, *Statistics in Biopharmaceutical Research* 13 (2021) 384–394.
- [26] D. Jiang, Z. Bai, S. Zheng, Testing the independence of sets of large-dimensional variables, *Science China Mathematics* 56 (2013) 135–147.
- [27] H. Jiang, S. Wang, Moderate deviation principles for classical likelihood ratio tests of high-dimensional normal distributions, *Journal of Multivariate Analysis* 156 (2017) 57–69.
- [28] T. Jiang, Y. Qi, Likelihood ratio tests for high-dimensional normal distributions, *Scandinavian Journal of Statistics* 42 (2015) 988–1009.
- [29] T. Jiang, F. Yang, Central limit theorems for classical likelihood ratio tests for high-dimensional normal distributions, *The Annals of Statistics* 41 (2013) 2029–2074.
- [30] I. M. Johnstone, High dimensional statistical inference and random matrices, *Proceedings of the International Congress of Mathematicians, Madrid* (2006).
- [31] S. Krug, E. Johns, *The 16 personality factor questionnaire*, Testing and Counseling Practice. Mahwah, NJ: Lawrence Erlbaum Associates (1990) 63–90.
- [32] A. Lemonte, *The gradient test: another likelihood-based test*, Academic Press, 2016.
- [33] A. J. Lemonte, On the gradient statistic under model misspecification. *Statistics & Probability Letters* 83 (2013) 390–398.
- [34] W. Li, J. Chen, J. Yao, Testing the independence of two random vectors where only one dimension is large, *Statistics* 51 (2017) 141–153.
- [35] W. Li, Q. Wang, J. Yao, W. Zhou, On eigenvalues of a high-dimensional spatial-sign covariance matrix, *Bernoulli* 28 (2022) 606–637.
- [36] P. Loubaton, A. Rosuel, Properties of linear spectral statistics of frequency-smoothed estimated spectral coherence matrix of high-dimensional gaussian time series, *Electronic Journal of Statistics* 15 (2021) 5380–5454.
- [37] X. Luo, W. Y. Tsai, A proportional likelihood ratio model, *Biometrika* 99 (2012) 211–222.
- [38] B. Mershon, R. L. Gorsuch, Number of factors in the personality sphere: Does increase in factors increase predictability of real-life criteria?, *Journal of Personality and Social Psychology* 55 (1988) 675.
- [39] X. Mestre, P. Vallet, Correlation tests and linear spectral statistics of the sample correlation matrix, *IEEE Transactions on Information Theory* 63 (2017) 4585–4618.
- [40] R. J. Muirhead, *Aspects of Multivariate Statistical Theory*, John Wiley & Sons, New York, 1982.
- [41] H. H. Nguyen, V. Vu, Random matrices: Law of the determinant, *The Annals of Probability* 42 (2014) 146–167.
- [42] P. O’Brien, Robust procedures for testing equality of covariance matrices, *Biometrics* 48 (1992) 819–827.
- [43] N. Parolya, J. Heiny, D. Kurowicka, Logarithmic law of large random correlation matrix, arXiv preprint arXiv:2103.13900 (2021).
- [44] Y. Qi, F. Wang, L. Zhang, Limiting distributions of likelihood ratio test for independence of components for high-dimensional normal vectors, *Annals of the Institute of Statistical Mathematics* 71 (2019) 911–946.
- [45] S. Roccas, L. Sagiv, S. H. Schwartz, A. Knafo, The big five personality factors and personal values, *Personality and social psychology bulletin* 28 (2002) 789–801.
- [46] J. R. Schott, A test for the equality of covariance matrices when the dimension is large relative to the sample sizes, *Computational Statistics and Data Analysis* 51 (2007) 6535–6542.
- [47] M. S. Srivastava, H. Yanagihara, Testing the equality of several covariance matrices with fewer observations than the dimension, *Journal of Multivariate Analysis* 101 (2010) 1319–1329.
- [48] L. J. Strug, The evidential statistical paradigm in genetics, *Genetic Epidemiology* 42 (2018) 590–607.
- [49] X. Wang, X. Han, G. Pan, The logarithmic law of sample covariance matrices near singularity, *Bernoulli* 24 (2018) 80–114.
- [50] D. Watterson, *The 16pf leadership coaching report manual*, 2002.
- [51] Y. Yamada, M. Hyodo, T. Nishiyama, Testing block-diagonal covariance structure for high-dimensional data under non-normality, *Journal of Multivariate Analysis* 155 (2017) 305–316.
- [52] X. Yang, X. Zheng, J. Chen, Testing high-dimensional covariance matrices under the elliptical distribution and beyond, *Journal of Econometrics* 221 (2021) 409–423.