A CHARACTERIZATION OF THE INDIVIDUAL MAXIMUM AND ANTI-MAXIMUM PRINCIPLE

SAHIBA ARORA AND JOCHEN GLÜCK

ABSTRACT. Abstract approaches to maximum and anti-maximum principles for differential operators typically rely on the condition that all vectors in the domain of the operator are dominated by the leading eigenfunction of the operator. We study the necessity of this condition. In particular, we show that under a number of natural assumptions, so-called individual versions of both the maximum and the anti-maximum principle simultaneously hold if and only if the aforementioned domination condition is satisfied.

Consequently, we are able to show that a variety of concrete differential operators do not satisfy an anti-maximum principle.

1. INTRODUCTION

Maximum and anti-maximum principles have been long used to obtain information about PDEs without explicitly knowing their solutions. While (anti-)maximum principles have been proved/disproved, for various concrete differential operators – using, for instance, kernel estimates – only a few endeavours have been made to integrate these arguments in an abstract setting. One attempt in this direction was made by Takáč in [36]. Inspired by his techniques, the present authors recently proved necessary and sufficient conditions for the so-called *uniform* (anti-)maximum principles in [8]. On the other hand, a characterization for the *individual* maximum and anti-maximum principle was given in [19, Theorem 4.4].

More precisely, let Ω be a bounded domain in \mathbb{R}^d and let $A : E \supseteq \operatorname{dom} A \to E$ be a differential operator on a function space E over Ω . Consider the equation

$$(\lambda - A)u = f$$

for real numbers λ in the resolvent set of A. If $\lambda_0 \in \mathbb{R}$ is an isolated spectral value of A, then we say that A satisfies the *individual* maximum principle if the inequality $f \geq 0$ implies $u \geq 0$ for all λ in a f-dependent right neighbourhood of λ_0 . If the right neighbourhood can be chosen independently of f, then we call the maximum principle *uniform*. Analogously, if $f \geq 0$ implies $u \leq 0$ in a left neighbourhood of λ_0 , then we speak of (individual or uniform) anti-maximum principles.

Eventual positivity and a domination assumption. Recall that A is said to be resolvent positive if the resolvent $\mathcal{R}(\lambda, A)$ exists and is positive for all sufficiently large real numbers λ . Resolvent positivity was studied in detail in [3] and is also related to the theory of positive C_0 -semigroups (see [28, Chapters B-II and C-II] and [10, Corollary 11.4]). This notion also appears in [36].

As a generalization of the above notion, eventual positivity and eventual negativity of the resolvent was studied, as a complement to eventually positive C_0 semigroups, in [16–20]. The notions of individual and uniform eventual positivity

Date: 30th August 2023.

²⁰¹⁰ Mathematics Subject Classification. 35B09; 47B65; 46B42.

Key words and phrases. Maximum principle; individual anti-maximum principle; eventual positivity; eventually positive resolvents.

of the resolvent, actually coincide with the individual and uniform maximum principles discussed above. The same is true for the anti-maximum principle and the eventual negativity of the resolvent.

In the present article, we deal with the individual (anti-)maximum principles and, in particular, with a domination condition that occurred in the characterization theorem [19, Theorem 4.4]. To recall this condition, let E be a complex Banach lattice whose real part and positive cone will be denoted by $E_{\mathbb{R}}$ and E_+ respectively. For $0 \leq u \in E$, the set

$$E_u := \{ f \in E : \text{ there exists } c > 0 \text{ such that } |f| \le cu \}$$

is called the *principal ideal* generated by u. Equipped with the gauge norm

$$||f||_u := \inf\{c > 0 : |f| \le cu\}$$
 for $f \in E_u$,

it is also a complex Banach lattice. The canonical embedding $E_u \to E$ is continuous and a lattice homomorphism. If E_u is dense in E, then we say that u is a quasiinterior point of E (or more precisely of E_+). As an example, let (Ω, μ) be a finite measure space and $p \in [1, \infty]$. Then $\mathbb{1}$ – which stands for the constant function taking the value 1 – is a quasi-interior point of the Banach lattice $L^p(\Omega, \mu)$ and the corresponding principal ideal is given by $L^p(\Omega, \mu)_1 = L^{\infty}(\Omega, \mu)$ (here the gauge norm is simply the $\|\cdot\|_{\infty}$ -norm). We refer to the standard monographs [34] and [27] for the general theory of Banach lattices.

For a linear operator $A: E \supseteq \operatorname{dom} A \to E$, the condition

dom $A \subseteq E_u$

is referred to as the domination condition in [16,19] and plays a significant role in the characterization of the individual (anti-)maximum principle in [19, Theorem 4.4]. In [16, Theorem 4.1], the condition was analysed in detail and it was shown that the necessary conditions in [19, Theorem 4.4] remain true without it. In this paper, we show that under a spectral assumption and a much weaker domination assumption (see Setting 1.1 below), the individual maximum and anti-maximum principles are concomitantly satisfied if and only if dom $A \subseteq E_u$. This helps us in disproving anti-maximum principles for various differential operators for which the maximum principle is already known.

Main result. We confine ourselves to the study of real linear operators. This is not too restrictive, for instance, differential operators with real coefficients are typically real. Concretely, we call a linear operator $A : E \supseteq \operatorname{dom} A \to E$ real if $\operatorname{dom} A = \operatorname{dom} A \cap E_{\mathbb{R}} + i \operatorname{dom} A \cap E_{\mathbb{R}}$ and $A(\operatorname{dom} A \cap E_{\mathbb{R}}) \subseteq E_{\mathbb{R}}$. Observe that if A is real, then so are the operators $\mathcal{R}(\lambda, A)$ for all real numbers λ in the resolvent set $\rho(A)$. For two vectors $u, v \in E_{\mathbb{R}}$, we write $u \succeq v$ (equivalently, $v \preceq u$) if there exists a real number c > 0 such that $u \ge cv$. We use the same notation not only for vectors but also for operators.

Most of our results are formulated in the following setting:

Setting 1.1. Let $A : E \supseteq \text{dom } A \to E$ be a closed, densely defined, and real linear operator on a complex Banach lattice E and let $\lambda_0 \in \mathbb{R}$ be an isolated spectral value of A and a pole of the resolvent $\mathcal{R}(\cdot, A)$. Moreover, fix a quasi-interior point $u \in E$.

• We say that the spectral assumption is satisfied if and only if the following holds: the eigenvalue λ_0 of A is geometrically simple and the corresponding eigenspace ker $(\lambda_0 - A)$ is spanned by a vector v which satisfies $v \succeq u$; moreover, the dual eigenspace ker $(\lambda_0 - A')$ contains a strictly positive functional φ .

• We say that the *domination assumption* is satisfied if and only if there exists an integer $n \ge 0$ such that dom $A^n \subseteq E_u$.

Observe that if the domination assumption above holds, then the condition "u is a quasi-interior point of E" is redundant because A (and hence A^n) is a densely defined operator.

The following is our main result.

Theorem 1.2. In Setting 1.1, let the domination assumption be satisfied. Then the following assertions are equivalent.

(i) Strong individual (anti-)maximum principle: For every $0 \leq f \in E$, we have

$$\mathcal{R}(\mu, A)f \preceq -u \quad and \quad \mathcal{R}(\lambda, A)f \succeq u$$

for all μ in a f-dependent left neighbourhood of λ_0 and for all λ in a f-dependent right neighbourhood of λ_0 .

(ii) Individual (anti-)maximum principle plus spectral assumption: The spectral assumption is satisfied and for every $0 \le f \in E$, we have

 $\mathcal{R}(\mu, A)f \leq 0 \qquad and \qquad \mathcal{R}(\lambda, A)f \geq 0$

for all μ in a f-dependent left neighbourhood of λ_0 and for all λ in a f-dependent right neighbourhood of λ_0 .

(iii) Improved domination assumption plus spectral assumption: The spectral assumption is satisfied and dom $A \subseteq E_u$.

Let us briefly discuss the particular case where E = C(K), the space of continuous functions on a compact Hausdorff set K and $u \in E$ is the constant one function. Since $E_u = E$, the domination condition is always satisfied in Setting 1.1 for n = 1. Thus, according to Theorem 1.2, both strong individual maximum and anti-maximum principles hold at λ_0 if and only if the spectral assumption is satisfied.

Next, let $E = L^p(\Omega, \mu)$, where (Ω, μ) is a finite measure space and $p \in [1, \infty)$. If u is the constant one function, then as mentioned before, $E_u = L^{\infty}(\Omega, \mu)$. For numerous differential operators A, one can show using Sobolev embedding theorems that the domination condition holds for sufficiently large $n \in \mathbb{N}$ but not for n = 1. As a result, if one knows a priori that the maximum principle holds at λ_0 for those operators that in addition satisfy the spectral assumption, then the anti-maximum principle cannot hold at λ_0 . Analogously, if the anti-maximum principle holds at λ_0 , then the maximum principle cannot. We will illustrate this in Section 5.

It is natural to ask whether the assertions in Theorem 1.2 are equivalent without the domination assumption in Setting 1.1. However, one can immediately check that this is not the case. Indeed, on $L^2[0, 1]$ consider the bounded operator defined by $Af = (\mathbb{1} \otimes \mathbb{1})f - f$ for all $f \in L^2[0, 1]$, and let $\lambda_0 = 0$. The resolvent can be computed directly and is given by

$$\mathcal{R}(\lambda, A)f = \frac{1}{\lambda(\lambda+1)} \big((\mathbb{1} \otimes \mathbb{1})f + \lambda f \big), \qquad (\lambda \in \rho(A)).$$

Thus, one has $\mathcal{R}(\lambda, A) \succeq \mathbb{1} \otimes \mathbb{1}$ and $\mathcal{R}(\mu, A) \preceq -\mathbb{1} \otimes \mathbb{1}$ for all $\lambda > 0$ and for all $\mu \in (-1, 0)$. However, dom $(A^n) = L^2[0, 1] \not\subseteq L^{\infty}[0, 1] = L^2[0, 1]_1$ for any $n \in \mathbb{N}_0$.

Related literature. Individual maximum and anti-maximum principles for concrete differential operators have been intensively studied for several decades. For instance, we refer the reader to the articles [12,13,24,31,32] and to the references in Section 5 for various examples. Abstract approaches to (anti-)maximum principles with several applications have been taken [8, 19, 20, 36]. Recently, the first author proved *local* versions of the (anti-)maximum principles in [6], see also [7, Chapter 9], and gave a number of applications. **Organization of the article.** In Section 2, we recall the notion of operator ranges, a concept which is instrumental in proving Theorem 1.2 in Section 3. In Section 4, we discuss a characterization of the domination assumption in Setting 1.1 in the important special case where A generates a C_0 -semigroup. Applications of our result to various concrete differential operators are presented in Section 5.

2. Eventual conditions on operator ranges

The improved domination condition dom $A \subseteq E_u$ from Theorem 1.2 can equivalently be written in the form $\mathcal{R}(\lambda, A)E \subseteq E_u$ for one number λ in the resolvent set of A or, equivalently, for all λ in the resolvent set. This raises the following question: what happens if we only know that $\mathcal{R}(\lambda_f, A)f \in E_u$ for all $f \in E$ and certain f-dependent numbers λ_f ? The purpose of this section is to analyse this situation; see Corollary 2.2(a) below. In fact, this will be a consequence of the following more general result:

Theorem 2.1. Let E, F, and V be Banach spaces such that V is continuously embedded in F. Let $\mathcal{T} \subseteq \mathcal{L}(E; F)$ be a countable set of bounded linear operators and assume that for each $f \in E$ there exists $T \in \mathcal{T}$ such that $Tf \in V$. Then there even exists an operator $T_0 \in \mathcal{T}$ such that $T_0 E \subseteq V$.

We will show below that this result is a consequence of Baire's category theorem. But first, let us list a few consequences:

Corollary 2.2. Let E and V be two complex Banach spaces such that V is continuously embedded in E, and let $A : E \supseteq \text{dom } A \to E$ be a linear operator.

- (a) Let $C \subseteq \mathbb{C}$ be a countable subset of the resolvent set of A and assume that for each $f \in E$ there exists a number $\lambda \in C$ such that $\mathcal{R}(\lambda, A)f \in V$. Then dom $A \subseteq V$.
- (b) Assume that A generates a C_0 -semigroup $(e^{tA})_{t \in [0,\infty)}$ on E and assume that there exists a countable set $S \subseteq [0,\infty)$ such that for each $f \in E$ there exists a time $t \in S$ for which we have $e^{tA}f \in V$. Then there exists a time $t_0 \in [0,\infty)$ such that $e^{tA}E \subseteq V$ for each $t \ge t_0$.

Proof. (a) From the assumption and Theorem 2.1 we conclude that, for some $\lambda \in C$, dom $A = \mathcal{R}(\lambda, A)E \subseteq V$.

(b) It follows from the assumption and from Theorem 2.1 that, for some $t_0 \in S$, we have $e^{t_0A}E \subseteq V$. By the semigroup law, the same is also true for all times $t \geq t_0$.

We will use assertion (a) of the corollary in the next section in the proof of our main theorem.

Remarks 2.3. (a) Corollary 2.2(b) is related to the theory of eventually positive semigroups: If $(e^{tA})_{t\in[0,\infty)}$ is a C_0 -semigroup on a Banach lattice E, and $u \in E_+$ is a quasi-interior point, an assumption that occurs frequently in the context of eventually positive semigroups is the existence of a time t_0 such that $e^{t_0A}E \subseteq E_u$; see, for instance, [20, Section 5]. Corollary 2.2(b) shows that this is equivalent to the a priori weaker assumption that each orbit of the semigroup is eventually in E_u ; here we have tacitly used the semigroup law.

(b) On the other hand, the theory of eventually positive semigroups also shows that a similar result as in Corollary 2.2(b) does not hold if we replace subspaces with cones: In [20, Examples 5.7 and 5.8], one can find an example of a semigroup $(e^{tA})_{t\in[0,\infty)}$ on a Banach lattice E such that, for each $f \in E_+$, the orbit of f is eventually in E_+ (thus, we can take the countable subset S to be \mathbb{N}), but one does not have $e^{tA}E_+ \subseteq E_+$ for any $t \in (0,\infty)$.

(c) In a similar way and again with the aid of [20, Example 5.7], one can see that Corollary 2.2(a) need not hold if V is merely a cone but not a subspace.

(d) By applying Corollary 2.2(b) to V = dom A, we can see that a C_0 -semigroup is eventually differentiable (in the sense of [22, Definition II.4.13]) if each of its individual orbits is eventually differentiable. It turns out, though, that the latter result even holds for much more general classes of operator-valued functions than only C_0 -semigroups; this was recently proved by Peruzzetto in [30, Proposition 3.11] (also based on Baire's category theorem, but with a different technical set-up).

To derive Theorem 2.1 from Baire's theorem, we need the concept of *operator* ranges. A subspace V of a Banach space F is called an *operator* range if there exists a Banach space E and a bounded linear operator $T: E \to F$ such that T(E) = V. For more details about operator ranges, we refer the reader to [14, 15]. Recent results about the eventual invariance of operator ranges were given by the present authors in [9, Section 2]. Here, we only recall some of their basic properties. The proof of the following proposition can be found in [14, Proposition 2.1]:

Proposition 2.4. Let F be a Banach space and let $V \subseteq F$ be a vector subspace of F. The following are equivalent.

- (i) The subspace V is an operator range.
- (ii) There exists a closed operator on F whose domain is V.
- (iii) There exists a complete norm $\|\cdot\|_V$ on V which makes the embedding $(V, \|\cdot\|_V) \hookrightarrow (F, \|\cdot\|_F)$ is continuous.

Obviously, the image of an operator range under a bounded linear operator is again an operator range. A bit more interestingly, the same holds for pre-images:

Proposition 2.5. Let $S : E \to F$ be a bounded linear operator between Banach spaces E and F. If a vector subspace $V \subseteq F$ is an operator range in F, then the pre-image $U := S^{-1}(V)$ is an operator range in E.

Proof. Let $\|\cdot\|_V$ denote a complete norm on V which makes the embedding into F continuous (such a norm exists by Proposition 2.4). We define a norm $\|\cdot\|_U$ on $U = S^{-1}(V)$ by the formula

$$||x||_U := ||x||_E + ||Sx||_V$$

for all $x \in U$. It is not difficult to check that $(U, \|\cdot\|_U)$ is a Banach space, and obviously the inclusion $(U, \|\cdot\|_U) \hookrightarrow (E, \|\cdot\|_E)$ is continuous; thus, U is an operator range.

It is a classical observation in functional analysis that, as a consequence of Baire's theorem, a countable union of closed proper subspaces of a Banach space F cannot be equal to F. For the proof of Theorem 2.1, we recall that the same is true not only for closed subspaces but even for operator ranges:

Proposition 2.6. Let F be a Banach space and let $U \subseteq F$ be an operator range. If U is a proper subspace (i.e., $U \neq F$), then it is a meagre set in F.

Consequently, if $F = \bigcup_{k \in \mathbb{N}} U_k$, where each $U_k \subseteq F$ is an operator range, then there exists $k \in \mathbb{N}$ such that $U_k = F$.

Proof. Since U is an operator range, there exists a Banach space E and $T \in \mathcal{L}(E, F)$ such that T(E) = U. If U is not a meagre set, then by a version of open mapping theorem [33, Theorem 2.11], we have U = F. The second assertion follows now by Baire's theorem.

We conclude this section with the proof of Theorem 2.1.

Proof of Theorem 2.1. Enumerate the elements of \mathcal{T} as T_1, T_2, \ldots (where some operators might occur infinitely often if \mathcal{T} is finite). For each $k \in \mathbb{N}$, consider the subspace

$$U_k := T_k^{-1}(V)$$

of E. According to Propositions 2.4 and 2.5, each of the spaces U_k is an operator range in E, and the assumption of the theorem implies that $\bigcup_{k \in \mathbb{N}} U_k = E$. Hence, Proposition 2.6 shows that there exists an index k_0 for which we have $U_{k_0} = E$; thus, $T_{k_0}E \subseteq V$.

3. Proof of the main result

The proof of Theorem 1.2 is based on a couple of auxiliary results. We start with the following finite series expansion of resolvents.

Lemma 3.1. Let $A : E \supseteq \operatorname{dom} A \to E$ be a closed linear operator on a complex Banach space E, let $m \ge 0$ be an integer and let $\lambda, \mu_1, \ldots, \mu_m \in \mathbb{C}$ be points in the resolvent set of A. Then the expansion formula

$$\mathcal{R}(\lambda, A) = \sum_{k=1}^{m} \Big(\prod_{j=1}^{k-1} (\mu_j - \lambda) \Big) \Big(\prod_{j=1}^{k} \mathcal{R}(\mu_j, A) \Big) \\ + \mathcal{R}(\lambda, A) \Big(\prod_{j=1}^{m} (\mu_j - \lambda) \Big) \Big(\prod_{j=1}^{m} \mathcal{R}(\mu_j, A) \Big)$$

holds.

Proof. The formula follows from an iterative application of the resolvent identity: first apply it to the numbers λ, μ_1 , then to the numbers λ, μ_2 , and proceed inductively.

Remark 3.2. Suppose the domination assumption in Setting 1.1 is satisfied and $\mu_1, \mu_2, \ldots, \mu_n$ are real numbers in the resolvent set of A. Then by the closed graph theorem, $\prod_{j=1}^n \mathcal{R}(\mu_j, A)$ is a bounded operator from E to E_u . Hence for every $f \in E_{\mathbb{R}}$, we have $-u \preceq \prod_{j=1}^n \mathcal{R}(\mu_j, A)f \preceq u$.

By using the previous lemma, we now show how upper/lower estimates for the resolvent at a point can be transferred to other points in the resolvent set. Related results can be found in [8, Section 4], but the point of the following theorem is that the number μ is allowed to depend on f, which makes the situation more subtle.

Theorem 3.3. In Setting 1.1, let both the domination and spectral assumption be satisfied.

(a) Assume that for each $0 \leq f \in E$ there exists a number $\mu > \lambda_0$ in the resolvent set of A such that $\mathcal{R}(\mu, A)f \succeq -u$.

Then we have $\mathcal{R}(\lambda, A) f \succeq -u$ for all $0 \leq f \in E$ and all $\lambda < \lambda_0$ in the resolvent set of A.

(b) Assume that for each $0 \leq f \in E$ there exists a number $\mu < \lambda_0$ in the resolvent set of A such that $\mathcal{R}(\mu, A)f \preceq u$.

Then we have $\mathcal{R}(\lambda, A)f \leq u$ for all $0 \leq f \in E$ and all $\lambda > \lambda_0$ in the resolvent set of A.

Proof. It is sufficient to prove (a); assertion (b) will then follow by replacing A with -A. To this end, let $0 \leq f \in E$ and let $n \geq 0$ be the integer from the domination assumption (for which we have dom $A^n \subseteq E_u$). We will recursively construct numbers μ_1, \ldots, μ_n larger than λ_0 and in the resolvent set of A such that

$$\mathcal{R}(\mu_k, A) \cdots \mathcal{R}(\mu_1, A) f \succeq -u$$
 (3.1)

for all $k \in \{0, ..., n\}$. Then Lemma 3.1 (applied to m = n) along with Remark 3.2 yields the required estimate.

For the construction, first note that since $f \ge 0 \ge -u$, the case k = 0 is trivial. Now, assume that μ_1, \ldots, μ_k have already been constructed for some k < n such that the estimate (3.1) holds. Then, because the eigenvector v from the spectral assumption satisfies $v \succeq u$, there exists a constant c > 0 such that

$$g := \mathcal{R}(\mu_k, A) \dots \mathcal{R}(\mu_1, A)f + cv \ge 0.$$

By assumption, this implies that there exists $\mu_{k+1} > \lambda_0$ and a constant $d_1 > 0$ such that $\mathcal{R}(\mu_{k+1}, A)g + d_1u \ge 0$. On the other hand, combining the domination and spectral assumptions, we have that $0 \le v \in \ker(\lambda_0 - A) \subseteq \operatorname{dom} A^n \subseteq E_u$. We can thus choose a constant $d_2 > 0$ such that $(\mu_{k+1} - \lambda_0)\mathcal{R}(\mu_{k+1}, A)v = v \le d_2u$. Therefore,

$$0 \leq \mathcal{R}(\mu_{k+1}, A) \mathcal{R}(\mu_k, A) \dots \mathcal{R}(\mu_1, A) f + c \ \mathcal{R}(\mu_{k+1}, A) v + d_1 u$$
$$\leq \mathcal{R}(\mu_{k+1}, A) \mathcal{R}(\mu_k, A) \dots \mathcal{R}(\mu_1, A) f + \left(\frac{cd_2}{\mu_{k+1} - \lambda_0} + d_1\right) u;$$

which readily yields

$$\mathcal{R}(\mu_{k+1}, A)\mathcal{R}(\mu_k, A)\dots\mathcal{R}(\mu_1, A)f \succeq -u.$$

Now we can prove our main result.

Proof of Theorem 1.2. "(iii) \Rightarrow (i)": This implication is a direct consequence of [19, Theorem 4.4 and Corollary 3.3].

"(i) \Rightarrow (ii)": If (i) holds, then so does the spectral assumption – due to [16, Theorem 4.1] and [19, Corollary 3.3] – and in turn, (ii).

"(ii) \Rightarrow (iii)": We assume (ii), and we only have to show that dom $A \subseteq E_u$. Due to the individual anti-maximum principle in assertion (ii), Theorem 3.3(b) tells us that $\mathcal{R}(\lambda, A)f \preceq u$ for all $\lambda > \lambda_0$ and all $f \in E_+$. By combining this with the individual maximum principal in assertion (ii), we see that, for all $f \in E_+$, the estimate $0 \leq \mathcal{R}(\lambda, A)f \preceq u$ holds for all λ in a f-dependent right neighbourhood of λ_0 .

Because the positive cone of a Banach lattice is generating, we've shown that, each $f \in E$ satisfies $\mathcal{R}(\lambda, A) f \in E_u$ for all λ in a *f*-dependent right neighbourhood of λ_0 . We can now employ Corollary 2.2(a) to conclude that dom $A \subseteq E_u$.

Remark 3.4. Since both [16, Theorem 4.1] and [19, Corollary 3.3] do not impose any domination assumptions, such assumptions are not needed for the implication (i) \Rightarrow (ii) in Theorem 1.2 to be true.

It is possible to give a proof of the implication "(ii) \Rightarrow (iii)" which does not rely on the concept of operator ranges and Baire's theorem. A major drawback of the alternative proof is that it is technically more involved and, thus, less transparent. Moreover, it uses consequences of the resolvent equation in more than one place, due to which it is less likely to be adaptable to different situations. However, for the benefit of the interested reader, we include the proof here.

Alternate proof of "(ii) \Rightarrow (iii)" in Theorem 1.2. Assume that (ii) is true. We want to prove that dom $A \subseteq E_u$, for which it suffices to prove that $\mathcal{R}(\mu, A)E \subseteq E_u$ for some $\mu \in \rho(A)$.

Recall again that $n \ge 0$ denotes the integer that occurs in the domination assumption. If n = 0, the conclusion is, of course, trivial. So, let $n \ge 1$ and fix $f \in E_+$. We assert that there exist $\mu_1, \ldots, \mu_n \in \rho(A)$ such that

$$0 \le \mathcal{R}(\mu_k, A) \dots \mathcal{R}(\mu_1, A) f \preceq u \tag{3.2}$$

for all $k \in \{1, ..., n\}$.

To see this, first note that since $f \geq 0$, condition (ii) implies that for some $\lambda < \lambda_0$, we have $\mathcal{R}(\lambda, A)f \leq 0 \leq u$. We can hence apply Theorem 3.3(b), and doing so in conjunction with (ii), gives the existence of $\mu_1 > \lambda_0$ such that $0 \leq \mathcal{R}(\mu_1, A)f \leq u$. This establishes the assertion for the case k = 1.

To proceed inductively, we now assume that for some k < n, there exist real numbers $\mu_1, \ldots, \mu_k \in \rho(A)$ satisfying the estimate (3.2). Then

$$0 \le g := \mathcal{R}(\mu_k, A) \dots \mathcal{R}(\mu_1, A) f \in E.$$

Repeating the above argument for g instead of f, we find $\mu_{k+1} > \lambda_0$ such that $0 \leq \mathcal{R}(\mu_{k+1}, A)g \leq u$, which completes the induction step.

In particular, it follows that $\prod_{j=1}^{k} \mathcal{R}(\mu_j, A) f \in E_u$ for all $k \in \{1, \ldots, n\}$. Now, appealing to Lemma 3.1 together with Remark 3.2, we obtain that $\mathcal{R}(\mu, A) f \in E_u$ for each $\mu \in \rho(A)$. Since E_+ spans E, ergo, for each $\mu \in \rho(A)$ one has the inclusion $\mathcal{R}(\mu, A) E \subseteq E_u$, as required.

For eventually differentiable semigroups, [19, Corollary 5.3] gives a characterization for individual eventual positivity of a semigroup whose generator satisfies the domination assumption from Setting 1.1. In this case, Theorem 1.2 says that if the domination assumption does not hold for n = 1, then the individual maximum and anti-maximum principle cannot both be true. This is also illustrated in Section 5.

4. A sufficient condition for the domination assumption

For sufficiently well-behaved concrete differential operators A, one way to check the domination assumption in Setting 1.1 will typically be as follows: By employing an elliptic regularity result one gets that dom A^n is contained in a Sobolev space of an order that scales with n. So if n is sufficiently large, one can employ a Sobolev embedding theorem (and combine it, if necessary, with the boundary conditions encoded in the domain of A) to obtain the embedding dom $(A^n) \subseteq E_u$. One problem with this approach is that it fails for operators for which elliptic regularity results are not available.

In fact, there are various differential operators for which elliptic regularity does not hold or is not known, but for which one has a parabolic estimate that yields $e^{tA}E \subseteq E_u$ for all t > 0, where $(e^{tA})_{t \in [0,\infty)}$ denotes the C_0 -semigroup generated by A. The following theorem (when applied to $V = E_u$) shows that a specific quantitative form of the property $e^{tA}E \subseteq E_u$ is equivalent to our domination assumption dom $(A^n) \subseteq E_u$.

Theorem 4.1. Let E and V be Banach spaces and assume that V is continuously embedded in E. For a C_0 -semigroup $(e^{tA})_{t \in [0,\infty)}$ on E, consider the following assertions:

(i) There exist real numbers $t_0, q, c > 0$ such that, for all $t \in (0, t_0]$, we have

 $e^{tA}E \subseteq V$ and $\|e^{tA}\|_{E \to V} \leq c t^{-q}$.

(ii) There exists an integer $n \ge 0$ such that dom $(A^n) \subseteq V$.

One always has (i) \Rightarrow (ii). If the semigroup is analytic, then also (ii) \Rightarrow (i).

For the case $E = L^2$ and $V = L^{\infty}$, and under additional assumptions on the semigroup, implication "(i) \Rightarrow (ii)" in Theorem 4.1 can essentially be found in [21, Theorem 2.4.1]. The same argument can be adapted to the general case; though for the convenience of the reader, we prefer to include the details.

8

Proof of Theorem 4.1. "(i) \Rightarrow (ii)": Choose a real number λ which is strictly larger than the growth bound of $(e^{tA})_{t \in [0,\infty)}$, and an integer $n \ge 1$ which satisfies n > q. Fix $f \in E$. Using

$$\|e^{tA}f\|_{V} \le \|e^{t_{0}A}\|_{E \to V} \|e^{(t-t_{0})}f\|_{E}$$
 for $t \in (t_{0}, \infty)$

along with the estimate in (i), we get that

$$\int_0^\infty t^{n-1} e^{-\lambda t} \left\| e^{tA} f \right\|_V \, \mathrm{d}t < \infty;$$

the essential point is that the singularity of $t \mapsto \|e^{tA}f\|_V$ at t = 0 gets suppressed since n - 1 - q > -1. (Observe that the mapping $(0, \infty) \ni t \mapsto e^{tA}f \in V$ is continuous, so the integrand is also continuous, and in turn, measurable.) Therefore,

$$\int_0^\infty t^{n-1} e^{-\lambda t} e^{tA} f \, \mathrm{d}t \in V,$$

because the integral above exists as a Bochner integral in V. At the same time, the integral also exists as a Bochner integral in E, and this integral is equal to $(n-1)!\mathcal{R}(\lambda, A)^n f$; see for instance [22, Corollary II.1.1].

The continuity of the embedding V into E implies that both Bochner integrals coincide, so $(n-1)!\mathcal{R}(\lambda, A)^n f \in V$. In particular, $\mathcal{R}(\lambda, A)^n f \in V$. This shows that $\operatorname{dom}(A^n) \subseteq V$.

"(ii) \Rightarrow (i)": We now assume that the semigroup is analytic. By replacing A with $A - \lambda$ for a sufficiently large number $\lambda > 0$ if necessary, we may assume that 0 is in the resolvent set of A. Now we endow the spaces dom (A^k) for $k = 0, 1, \ldots, n$ with the graph norms

$$\|f\|_{\operatorname{dom}(A^k)} := \|A^k f\|_E,$$

which renders them Banach spaces.

If n = 0, then V = E, so there is nothing to prove; hence, let $n \ge 1$ for the rest of the proof. We show that

$$\left\| e^{tA} \right\|_{E \to \operatorname{dom}(A^n)} \le \tilde{c} t^{-n}$$

for a constant $\tilde{c} > 0$ and all $t \in (0, n]$. Since the inclusion dom $(A^n) \hookrightarrow V$ is, by the closed graph theorem, continuous, assertion (i) then follows using the analyticity of the semigroup with q = n and $t_0 = n$.

Since the semigroup is analytic, the number

$$M := \sup_{t \in (0,1]} \left\| tAe^{tA} \right\|_{E \to E}$$

is finite [22, Theorem II.4.6(a) and (c)] (this reference is formulated for semigroups that are bounded on a sector, and thus has to be applied to a rescaled version of our semigroup).

Consider an integer $k \in \{0, ..., n-1\}$ and a vector $f \in \text{dom}(A^k)$. Then, for $t \in (0, 1]$, we have $e^{tA}f \in \text{dom}(A^{k+1})$ and

$$\|e^{tA}f\|_{\operatorname{dom}(A^{k+1})} = \|Ae^{tA}A^{k}f\|_{E} \le \frac{M}{t} \|f\|_{\operatorname{dom}(A^{k})},$$

so $\|e^{tA}\|_{\operatorname{dom}(A^k)\to\operatorname{dom}(A^{k+1})} \leq Mt^{-1}$ for $t \in (0,1]$. For $t \in (0,n]$ this implies

$$|e^{tA}||_{E \to \operatorname{dom}(A^n)} \le \prod_{k=0}^{n-1} \left\| e^{\frac{t}{n}A} \right\|_{\operatorname{dom}(A^k) \to \operatorname{dom}(A^{k+1})} \le (nM)^n t^{-n}$$

which concludes the proof.

Remarks 4.2. (a) Observe that the implication "(ii) \Rightarrow (i)" in Theorem 4.1 is, in general, false without the assumption of analyticity. Just consider a C_0 -group with unbounded generator A and define V := dom A to obtain a counterexample where one does not even have $e^{tA}E \subseteq V$ for any time t.

(b) The proof of Theorem 4.1 shows that, in the implication "(i) \Rightarrow (ii)", one can choose n to be the smallest integer which satisfies n > q (and hence, one can choose n = q + 1 if q is an integer). For the converse implication, "(ii) \Rightarrow (i)", one can choose q to be equal to n.

(c) Since the condition dom $(A^n) \subseteq V$ is equivalent to $\mathcal{R}(\lambda, A)^n E \subseteq V$ for one (equivalently: all) λ in the resolvent set of A, it follows from Theorem 2.1 that condition (ii) in Theorem 4.1 is equivalent to the condition that, for some λ in the resolvent set, the following holds: For each $f \in E$, there exists an (*f*-dependent) integer $n \geq 0$ such that $\mathcal{R}(\lambda, A)^n f \in V$.

5. Examples and applications

In this section, we consider a couple of concrete differential operators to demonstrate how Theorem 1.2 can be applied to characterize when they satisfy the individual anti-maximum principle.

5.1. The Laplacian with Robin boundary conditions. Let Ω be a bounded domain in \mathbb{R}^d ; for the sake of simplicity we assume that Ω has C^{∞} -boundary, and we let $\beta : \partial \Omega \to \mathbb{R}$ be a C^1 -function. Fix $p \in (1, \infty)$. We consider the Robin Laplace operator $\Delta_{\beta} : L^p(\Omega) \supseteq \operatorname{dom}(\Delta_{\beta}) \to L^p(\Omega)$ given by

$$\operatorname{dom}(\Delta_{\beta}) = \left\{ u \in W^{2,p}(\Omega) : \frac{\partial}{\partial \nu} u = \beta u \text{ on } \partial\Omega \right\},\$$
$$\Delta_{\beta} u = \Delta u;$$

where $\frac{\partial}{\partial u} u$ denotes the outer normal derivative.

The operator Δ_{β} generates an analytic C_0 -semigroup on $L^p(\Omega)$ [37, Theorem 5.6 on p. 189] and it can be shown – for instance, by applying form methods to the case p = 2 and then extrapolating those properties to other L^p -spaces – that this semigroup is both positive and irreducible.

Let v denote the eigenvector of Δ_{β} associated to the eigenvalue s(A). Then v is an element of $C(\overline{\Omega})$ and it satisfies $v \succeq 1$, where 1 denotes the constant one function. This is true even for more general elliptic operators and much rougher domains and can, for instance, be found in [5, Theorem 4.5(b)]. Also by the positivity and the irreducibility, the eigenspace of the dual operator A' corresponding to the eigenvalue s(A) contains a strictly positive functional [28, Proposition C-III-3.5].

We also note that, for p = 2, the associated sesquilinear form has form domain $H^1(\Omega)$, which embeds continuously into $L^q(\Omega)$ for some q > 2, so using an ultracontractivity argument (see [4, Theorem in Section 7.3.2]), we obtain that $e^{t\Delta_{\beta}}L^p(\Omega) \subseteq L^{\infty}(\Omega) = L^p(\Omega)_1$ and, even more, that the growth condition in Theorem 4.1(i) is satisfied. As a result, the aforementioned theorem implies that there exists $n \in \mathbb{N}$ such that dom $((\Delta_{\beta})^n) \subseteq L^p(\Omega)_1$. We have thus shown that both the domination and the spectral assumptions in Setting 1.1 are satisfied for $\lambda_0 = s(\Delta_{\beta})$ and $u = \mathbb{1}$.

We also note that the resolvent of Δ_{β} is positive on the right of $s(\Delta_{\beta})$ since the semigroup $(e^{t\Delta_{\beta}})_{t\in[0,\infty)}$ is positive. In particular, in the terminology of Theorem 1.2, the individual (and in fact, even the uniform) maximum principle is satisfied at $\lambda_0 = s(\Delta_{\beta})$. Let us now discuss under which conditions the individual anti-maximum principle is also satisfied. **Theorem 5.1.** The Robin Laplace operator Δ_{β} satisfies the individual anti-maximum principle at $\lambda_0 = s(\Delta_{\beta})$ if and only if p > d/2.

Proof. Since the domination and the spectral assumption are satisfied for u = 1 and since Δ_{β} satisfies the individual maximum principle, it follows from Theorem 1.2 that the individual anti-maximum principle is satisfied if and only if

$$\operatorname{dom}(\Delta_{\beta}) \subseteq L^2(\Omega)_{\mathbb{1}} = L^{\infty}(\Omega).$$

If p > d/2, then by the Sobolev embedding theorem, $W^{2,p}(\Omega)$ embeds into $L^{\infty}(\Omega)$ and hence, indeed, dom $(\Delta_{\beta}) \subseteq L^{\infty}(\Omega)$.

If, on the other hand, $p \leq d/2$, then there exists a function in $W^{2,p}(\Omega)$ which has compact support within Ω (and is thus an element of dom (Δ_{β})), but is not contained in $L^{\infty}(\Omega)$ [2, Examples 5.25 and 5.26]. Wherefore, dom $(\Delta_{\beta}) \not\subseteq L^{\infty}(\Omega)$ in this case.

A generalisation of this result will (under slightly more restrictive assumptions on β) be discussed in Section 5.4.

5.2. On the Laplacian with Dirichlet boundary conditions. For the Laplace operator with Dirichlet boundary conditions Δ_{Dir} , it is also possible to characterize whether an anti-maximum principle at $s(\Delta_{\text{Dir}})$ holds. This was done by Sweers [35] who proved that, on a smooth bounded domain in \mathbb{R}^d , the individual anti-maximum principle for the Dirichlet Laplacian on L^p holds if and only if p > d. The crucial point here is the boundary behaviour of functions lying in the domain of the operator, which explains that p needs to be larger (when compared to the Robin Laplacian) for the individual anti-maximum principle to hold.

Sweers' argument can also be rephrased to fit into the framework of our Theorem 1.2 – however, our theorem does not appear to significantly simplify the argument, so we refrain from discussing the details here.

We point out that for $d \ge 2$, the individual anti-maximum principle is not uniform; see [8, Proposition 6.1(a)] and the introductions of [13, 35].

An important observation here is that while the individual anti-maximum principle holds only for some p, its uniform counterpart is not true for any p. In fact, this characteristic is not unique to the Dirichlet Laplacian. The uniform (anti-)maximum principle is an operator inequality; hence, for a differential operator which acts on the entire L^{p} -scale, if the principle holds for some p, then by a simple density argument, it automatically holds for each p. The individual (anti-)maximum principle, on the other hand, is more subtle and its validity may rely on the choice of p.

5.3. A system of Neumann Laplacians coupled by a matrix-valued potential. We consider a situation similar to Subsection 5.1 – however, we couple multiple Laplace operators by a matrix-valued potential now. For the sake of simplicity, we consider Neumann boundary conditions only.

Let $\Omega \subseteq \mathbb{R}^d$ be a bounded domain with C^{∞} -boundary, let $N \geq 1$ be an integer and let $V : \Omega \to \mathbb{R}^{N \times N}$ be a measurable and bounded function. We assume that, for each $x \in \Omega$, all off-diagonal entries of the matrix V(x) are non-negative and that the matrix that is obtained from V(x) by setting all diagonal entries to 0 is irreducible. On the space $L^p(\Omega; \mathbb{C}^N)$, we consider the operator A given by

$$D(A) = D(\Delta_{\text{Neu}}) \times \ldots \times D(\Delta_{\text{Neu}})$$
$$Aw = \begin{pmatrix} \Delta w_1 \\ \vdots \\ \Delta w_N \end{pmatrix} + Vw;$$

where $D(\Delta_{\text{Neu}}) = \{h \in W^{2,p}(\Omega; \mathbb{C}) \ \frac{\partial}{\partial \nu}h = 0 \text{ on } \partial\Omega\}$ denotes the domain of the Neumann Laplace operator on $L^p(\Omega)$. Since V acts as a bounded linear operator on $L^p(\Omega; \mathbb{C})$ and the semigroup generated by this operator is positive, it follows that the semigroup generated by A is also positive; in fact, it is even irreducible, as follows, for example, from [28, Proposition C-III-3.3].

Because A has compact resolvent, we can conclude that s(A) is an algebraically simple eigenvalue of A and that the corresponding eigenspace is spanned by a vector v which is a quasi-interior point of the positive cone of $L^p(\Omega; \mathbb{C}^N)$; moreover, the dual eigenspace is also spanned by a strictly positive functional. This follows from [28, Proposition C-III-3.5].

That fact that v is a quasi-interior point means that each of the components $v_1, \ldots, v_N \in L^p(\Omega; \mathbb{C})$ is > 0 almost everywhere on Ω . In fact, we even have $v_1, \ldots, v_n \succeq \mathbb{1}$. To see this, choose a number $c \in [0, \infty)$ sufficiently large such that $V(x) + c \operatorname{id}_{\mathbb{C}^N} \ge 0$ for each $x \in \Omega$. Then $(e^{tc}e^{tA})_{t \in [0,\infty)}$ dominates the semigroup generated by



Hence, we obtain for the k-th component of the vector v, that

$$e^{tc}e^{ts(A)}v_k = (e^{tc}e^{tA}v)_k \ge e^{t\Delta_{\text{Neu}}}v_k \succeq 1,$$

and thus $v_k \succeq 1$. So the spectral assumption in Setting 1.1 is satisfied for $\lambda_0 = s(A)$ and u := (1, ..., 1). On the other hand, the domination assumption in Setting 1.1 is satisfied for this u, as well. This follows from $(L^p(\Omega; \mathbb{C}^N))_u = L^{\infty}(\Omega; \mathbb{C}^N)$ along with Theorem 4.1 and an ultracontractivity argument (see for instance [4, Theorem in Section 7.3.2]).

Since the semigroup generated by A is positive, the resolvent of A is positive on the right of A, i.e., the individual (and in fact, even uniform) maximum principle holds at $\lambda_0 = s(A)$. For the individual anti-maximum principle, we now get the same result as in Theorem 5.1:

Theorem 5.2. The coupled Neumann Laplace operator A satisfies the individual anti-maximum principle at $\lambda_0 = s(A)$ if and only if p > d/2.

Proof. Given the facts discussed before the theorem, the proof is now the same as for Theorem 5.1. $\hfill \Box$

5.4. Powers of the Robin Laplacian. We consider the same situation as in Section 5.1, but now we assume in addition that $\beta \succeq 1$ and, for the sake of simplicity, that β is a C^{∞} -function. Since $\beta \succeq 1$, the spectral bound $s(\Delta_{\beta})$ of the Robin Laplacian Δ_{β} is strictly negative.

Now, we fix an integer $k \ge 1$ and consider the operator $B := -(-\Delta_{\beta})^k$ on $L^p(\Omega)$. The semigroup generated by B is not positive unless k = 1 (however, we note that eventual positivity of the semigroup generated by the square of the Robin Laplacian on the space $C(\overline{\Omega})$ was discussed in [20, Section 6.4]).

By the spectral mapping theorem, the spectral bound of B also satisfies s(B) < 0, and for the resolvent at the point 0 we have

$$\mathcal{R}(0,B) = \mathcal{R}(0,\Delta_{\beta})^k \ge 0.$$

Hence, it follows from the Taylor series expansion of the resolvent that $\mathcal{R}(\lambda, B) \geq 0$ for all $\lambda \in (s(B), 0]$ (see [18, Proposition 3.2(i)] and [20, Proposition 4.2(i)]).

In particular, the operator B satisfies the individual (even the uniform) maximum principle at $\lambda_0 = s(B)$. As pointed out in Subsection 5.1 the operator Δ_β satisfies the domination assumption in Setting 1.1 for u = 1 and hence, so does B. Moreover, since $s(\Delta_{\beta}) < 0$, the eigenspace of B for the eigenvalue s(B) coincides with the eigenspace of Δ_{β} for the eigenvalue $s(\Delta_{\beta})$, and the same is true for the dual operators. Thus, B also satisfies the spectral assumption from Setting 1.1 (since the same is true for Δ_{β}).

Let us now analyse whether B satisfies the individual anti-maximum principle at s(B):

Theorem 5.3. The operator B satisfies the individual anti-maximum principle at $\lambda_0 = s(B)$ if and only if kp > d/2.

Proof. Since both $\partial\Omega$ and β are smooth, it follows from elliptic regularity that $\operatorname{dom}(B) \subseteq W^{2k,p}(\Omega)$. So if kp > d/2, then by the Sobolev embedding theorem, $\operatorname{dom}(B) \subseteq L^{\infty}(\Omega) = (L^p(\Omega))_1$, and consequently Theorem 1.2 implies that B satisfies the individual anti-maximum principle at s(B).

Conversely, if $kp \leq d/2$, then we can find a function $w : \Omega \to \mathbb{R}$ which has compact support within Ω and is in $W^{2k,p}(\Omega)$, but not in $L^{\infty}(\Omega)$. So, this function w is an element of dom(B), but not of $L^{\infty}(\Omega) = (L^p(\Omega))_1$. Whence, B does not satisfy the anti-maximum principle at s(B) according to Theorem 1.2.

5.5. The Dirichlet-to-Neumann operator. Let $\Omega \subseteq \mathbb{R}^d$ be a bounded domain with C^{∞} -boundary. Let Δ_{Dir} denote the Dirichlet Laplace operator on $L^2(\Omega)$ and consider a function $V \in L^{\infty}(\Omega, \mathbb{R})$ such that the infinity norm of the negative part of V, $||V^-||_{\infty}$, is strictly smaller than the smallest eigenvalue of $-\Delta_{\text{Dir}}$. In this situation, the operator $-\Delta_{\text{Dir}} + V$ is positive definite; in particular, 0 is not in its spectrum. This enables us, for every $\phi \in H^{1/2}(\partial\Omega)$, to uniquely solve

$$(-\Delta_{\text{Dir}} + V)u = 0$$
 weakly on Ω ,
Tr $u = \phi$;

where $u \in W^{1,2}(\Omega)$. Finally, denote the weak outer normal derivative of u on the boundary $\partial\Omega$ by $\partial_{\nu}u$ whenever it exists (see [40] or [39] for a definition of weak normal derivative). Then the operator $\phi \mapsto \partial_{\nu}u$ is called the *Dirichlet-to-Neumann* operator. We denote this operator by D_V and its domain by dom D_V . For an exact definition, and more details about the Dirichlet-to-Neumann operator, we refer the reader to [40] or [39]. This operator is self-adjoint on $L^2(\partial\Omega)$ with compact resolvent and $-D_V$ generates a holomorphic C_0 -semigroup on $L^2(\partial\Omega)$ of angle $\frac{\pi}{2}$. Moreover, the semigroup generated by $-D_V$ is positive (see [38, Theorem 2.3] or [1, Theorem 1.1(a)]) and even irreducible, as shown in [1, Theorem 1.1(c)] (note that there is a small inaccuracy in the statement of this result: the estimate that is assumed there should actually be the stronger estimate that is also assumed in [1, Theorem 1.1(a)]).

Due to the positivity of the semigroup, the resolvent satisfies $\mathcal{R}(\mu, -D_V) \geq 0$ for all $\mu > s(-D_V)$. In particular, we have an individual maximum principle at $\lambda_0 = s(-D_V)$. We now show that for $d \geq 3$, there is no anti-maximum principle.

Theorem 5.4. The individual anti-maximum principle holds at $\lambda_0 = s(-D_V)$ if and only if $d \leq 2$.

Proof. According to [40, Theorem 2.3], there exist c, q > 0 such that

 $e^{-tD_V}L^2(\partial\Omega) \subseteq L^\infty(\partial\Omega)$ and $\left\|e^{-tD_V}\right\|_{L^2(\partial\Omega) \to L^\infty(\partial\Omega)} \le c t^{-q}.$

for all $t \in (0, 1]$. Therefore, by virtue of Theorem 4.1, there exists an integer $n \ge 0$ such that dom $(D_V^n) \subseteq L^{\infty}(\partial \Omega)$. Combining this with $L^{\infty}(\partial \Omega) = L^2(\partial \Omega)_1$ – where 1 denotes the constant function with value one – we obtain that the domination assumption in Setting 1.1 is fulfilled for u = 1.

Moreover, since the semigroup generated by $-D_V$ is both positive and irreducible, the spectral bound $s(-D_V)$ is an eigenvector of $-D_V$ with corresponding eigenspace spanned by a vector v which is strictly positive almost everywhere; see [28, Proposition C-III-3.5]. Let us show that even $v \succeq 1$, so that also the spectral assumption in Setting 1.1 is satisfied for u = 1 and $\lambda_0 = s(-D_V)$. To this end, we note that it follows from $e^{-tD_V}L^2(\partial\Omega) \subseteq L^\infty(\partial\Omega)$ and from [39, Lemma 2.2] that e^{-tD_V} maps $L^2(\partial\Omega)$ into $C(\partial\Omega)$ for each t > 0; moreover, this mapping is continuous by the closed graph theorem. Now, we use an argument inspired by [5, Section 3]: It was shown in [39, Theorem 1.1] that $(e^{-tD_V}|_{C(\partial\Omega)})_{t\geq 0}$ is a C_0 -semigroup on $C(\partial \Omega)$. Hence, there exists a time t > 0 such that $e^{-tD_V} \mathbb{1} \succeq \mathbb{1}$, i.e., such that e^{-tD_V} maps a positive element of $L^2(\partial\Omega)$ to a quasi-interior point of $C(\partial \Omega)$. Such an operator automatically maps all quasi-interior points of $L^2(\partial \Omega)$ to quasi-interior points of $C(\partial \Omega)$ (see [23, Proposition 2.21] and note that the notion almost interior point that occurs there is, within Banach lattices, equivalent to quasi-interior point [23, pp. 242–243]). As a result, $v = e^{-tD_V}v$ is a quasi-interior point of $C(\partial \Omega)$ and thus satisfies $v \succeq \mathbb{1}$.

As mentioned above, the positivity of the semigroup implies that the (uniform and hence, in particular, individual) maximum principle holds at $\lambda_0 = s(-D_V)$. Lastly, we know that dom $(-D_V) = H^1(\partial\Omega)$; this follows from [11, Theorem 5.2] since $0 \in \rho(-\Delta_{\text{Dir}} + V)$. Using Sobolev embedding theorems on manifolds (see, for instance, [25, Theorem 6.2]), we obtain

$$\operatorname{dom}(-D_V) \subseteq C(\partial\Omega) \subseteq L^{\infty}(\partial\Omega) = L^2(\partial\Omega)_{\mathbb{1}}$$

if d < 3. On the other hand, if $d \ge 3$, then $\partial\Omega$ is a manifold of dimension at least two and hence, $H^1(\partial\Omega) \not\subseteq L^{\infty}(\partial\Omega)$ (see [2, Examples 5.25 and 5.26], [25, Proposition 4.1], or [26, Example 2.5.1]). The result thus follows from Theorem 1.2.

Acknowledgements. The authors would like to express their gratitude to Mikhail Ostrovskii [29] and Jochen Wengenroth [41] for two answers on MathOverflow that contain useful references about operator ranges. The question of whether assertion (b) in Corollary 2.2 holds was kindly brought to the authors' attention by Bálint Farkas.

Part of the work on this article was done when the first author visited the second author at Universität Passau, Germany.

The first named author was supported by Deutscher Akademischer Austauschdienst (Forschungsstipendium-Promotion in Deutschland).

References

- Jamil Abreu and Érika Capelato. Dirichlet-to-Neumann semigroup with respect to a general second order eigenvalue problem. Semigroup Forum, 97(2):183-202, 2018. doi:10.1007/s00233-018-9913-x.
- [2] Robert A. Adams. Sobolev spaces. New York-San Francisco-London: Academic Press, Inc., 1975.
- Wolfgang Arendt. Resolvent positive operators. Proc. Lond. Math. Soc. (3), 54:321-349, 1987. doi:10.1112/plms/s3-54.2.321.
- [4] Wolfgang Arendt. Semigroups and evolution equations: functional calculus, regularity and kernel estimates. In *Handbook of differential equations: Evolutionary equations. Vol. I*, pages 1–85. Amsterdam: Elsevier/North-Holland, 2004. doi:10.1016/S1874-5717(04)80003-3.
- [5] Wolfgang Arendt, A. F. M. ter Elst, and Jochen Glück. Strict positivity for the principal eigenfunction of elliptic operators with various boundary conditions. Adv. Nonlinear Stud., 20(3):633-650, 2020. doi:10.1515/ans-2020-2091.
- [6] Sahiba Arora. Locally eventually positive operator semigroups. J. Oper. Theory, 88(1):203–242, 2022. doi:10.7900/jot.2021jan26.2316.

- [7] Sahiba behavior ofArora. Long-term operator semigroups and (anti-)maximum principles. PhDthesis, Technische Universität Dresden. 2023.https://nbn-resolving.org/urn:nbn:de:bsz:14-qucosa2-834573.
- [8] Sahiba Arora and Jochen Glück. An operator theoretic approach to uniform (anti-)maximum principles. J. Differ. Equations, 310:164–197, 2022. doi:10.1016/j.jde.2021.11.037.
- [9] Sahiba Arora and Jochen Glück. Irreducibility of eventually positive semigroups, 2023. Preprint. arXiv:2307.04627.
- [10] András Bátkai, Marjeta K. Fijavž, and Abdelaziz Rhandi. Positive operator semigroups: From finite to infinite dimensions, volume 257. Basel: Birkhäuser/Springer, 2017. doi:10.1007/978-3-319-42813-0.
- [11] Jussi Behrndt and Antonius F.M. ter Elst. Dirichlet-to-Neumann maps on bounded Lipschitz domains. J. Differ. Equations, 259(11):5903-5926, 2015. doi:10.1016/j.jde.2015.07.012.
- [12] Isabeau Birindelli. Hopf's lemma and anti-maximum principle in general domains. J. Differ. Equations, 119(2):450–472, 1995. doi:10.1006/jdeq.1995.1098.
- [13] Philippe Clément and Lambertus A. Peletier. An anti-maximum principle for second order elliptic operators. J. Differ. Equations, 34:218–229, 1979. doi:10.1016/0022-0396(79)90006-8.
- [14] R. W. Cross. On the continuous linear image of a Banach space. J. Aust. Math. Soc., 29:219– 234, 1980. doi:10.1017/S1446788700021200.
- [15] Ronald W. Cross, Mikhail I. Ostrovskii, and Vitalii V. Shevchik. Operator ranges in Banach spaces, I. Math. Nachr., 173:91–114, 1995. doi:10.1002/mana.19951730107.
- [16] Daniel Daners and Jochen Glück. The role of domination and smoothing conditions in the theory of eventually positive semigroups. Bull. Aust. Math. Soc., 96(2):286–298, 2017. doi:10.1017/S0004972717000260.
- [17] Daniel Daners and Jochen Glück. A criterion for the uniform eventual positivity of operator semigroups. *Integral Equations Oper. Theory*, 90(4):19, 2018. Id/No 46. doi:10.1007/s00020-018-2478-y.
- [18] Daniel Daners and Jochen Glück. Towards a perturbation theory for eventually positive semigroups. J. Oper. Theory, 79(2):345–372, 2018. doi:10.7900/jot.2017mar29.2148.
- [19] Daniel Daners, Jochen Glück, and James B. Kennedy. Eventually and asymptotically positive semigroups on Banach lattices. J. Differ. Equations, 261(5):2607-2649, 2016. doi:10.1016/j.jde.2016.05.007.
- [20] Daniel Daners, Jochen Glück, and James B. Kennedy. Eventually positive semigroups of linear operators. J. Math. Anal. Appl., 433(2):1561–1593, 2016. doi:10.1016/j.jmaa.2015.08.050.
- [21] E. Brian Davies. Heat kernels and spectral theory, volume 92 of Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1989. doi:10.1017/CB09780511566158.
- [22] Klaus-Jochen Engel and Rainer Nagel. One-parameter semigroups for linear evolution equations, volume 194. Berlin: Springer, 2000.
- [23] Jochen Glück and Martin R. Weber. Almost interior points in ordered Banach spaces and the long-term behaviour of strongly positive operator semigroups. *Stud. Math.*, 254(3):237–263, 2020. doi:10.4064/sm190111-18-10.
- [24] Hans-Christoph Grunau and Guido Sweers. Optimal conditions for anti-maximum principles. Ann. Sc. Norm. Super. Pisa, Cl. Sci., IV. Ser., 30(3-4):499-513, 2001. URL: http://www.numdam.org/item/ASNSP_2001_4_30_3-4_499_0/.
- [25] Emmanuel Hebey and Frédéric Robert. Sobolev spaces on manifolds. In Handbook of global analysis, pages 375–415. Amsterdam: Elsevier, 2008. doi:10.1016/B978-044452833-9.50008-5.
- [26] Srinivasan Kesavan. Topics in functional analysis and applications. New York: John Wiley & Sons, Inc.; New Delhi: Wiley Eastern Limited, 1989.
- [27] Peter Meyer-Nieberg. Banach lattices. Berlin etc.: Springer-Verlag, 1991.
- [28] Rainer Nagel, editor. One-parameter semigroups of positive operators, volume 1184. Springer, Cham, 1986.
- [29] Mikhail Ostrovskii, 2019. MathOverflow post. URL: https://mathoverflow.net/a/332155/102946.
- [30] Marco Peruzzetto. On eventual regularity properties of operator-valued functions. Stud. Math., 265(2):141–176, 2022. doi:10.4064/sm201202-21-10.
- [31] Yehuda Pinchover. Maximum and anti-maximum principles and eigenfunctions estimates via perturbation theory of positive solutions of elliptic equations. *Math. Ann.*, 314(3):555–590, 1999. doi:10.1007/s002080050307.
- [32] Murray H. Protter and Hans F. Weinberger. Maximum Principles in Differential Equations. Partial differential equations. Springer New York, 1999. doi:10.1007/978-1-4612-5282-5.
- [33] Walter Rudin. Functional analysis. International series in pure and applied mathematics. New York, NY: McGraw-Hill, 1991.

- [34] Helmut H. Schaefer. Banach lattices and positive operators, volume 215. Springer, Berlin, 1974.
- [35] Guido Sweers. Lⁿ is sharp for the anti-maximum principle. J. Differ. Equations, 134(1):148– 153, 1997. doi:10.1006/jdeq.1996.3211.
- [36] Peter Takáč. An abstract form of maximum and anti-maximum principles of Hopf's type. J. Math. Anal. Appl., 201(2):339–364, 1996. doi:10.1006/jmaa.1996.0259.
- [37] Hiroki Tanabe. Functional analytic methods for partial differential equations, volume 204. New York, NY: Marcel Dekker, 1997. doi:10.1201/9780203755419.
- [38] Antonius F.M. ter Elst and El M. Ouhabaz. Analysis of the heat kernel of the Dirichlet-to-Neumann operator. J. Funct. Anal., 267(11):4066-4109, 2014. doi:10.1016/j.jfa.2014.09.001.
- [39] Antonius F.M. ter Elst and El M. Ouhabaz. Analyticity of the Dirichlet-to-Neumann semigroup on continuous functions. J. Evol. Equ., 19(1):21–31, 2019. doi:10.1007/s00028-018-0467-x.
- [40] Antonius F.M. ter Elst and El M. Ouhabaz. Dirichlet-to-Neumann and elliptic operators on C^{1+κ}-domains: Poisson and Gaussian bounds. J. Differ. Equations, 267(7):4224–4273, 2019. doi:10.1016/j.jde.2019.04.034.
- [41] Jochen Wengenroth, 2021. MathOverflow post. URL: https://mathoverflow.net/a/393345/102946.

Sahiba Arora, Technische Universität Dresden, Institut für Analysis, Fakultät für Mathematik , 01062 Dresden, Germany

 $Email \ address: \verb"sahiba.arora@mailbox.tu-dresden.de"$

Jochen Glück, Bergische Universität Wuppertal, Fakultät für Mathematik und Naturwissenschaften, 42119 Wuppertal, Germany

Email address: glueck@uni-wuppertal.de