
FORMATION OF COLLOIDAL CHAINS AND DRIVEN CLUSTERS WITH OPTICAL BINDING.

A PREPRINT

✉ **Dominique Davenport***
School of Natural Sciences
University of California - Merced
Merced, CA 95340
ddavenport3@ucmerced.edu

✉ **Dustin Kleckner**
School of Natural Sciences
University of California - Merced
Merced, CA 95340
dkleckner@ucmerced.edu

May 28, 2022

ABSTRACT

We study the effects of the optical binding force on wavelength sized colloidal particles free to move in a counter-propagating beam. This work is motivated by the idea of using optical binding to direct the assembly of large numbers of colloidal particles; previous work has used small numbers of particles and/or 1D or 2D restricted geometries. In this study, we utilize a unique geometry which allows us to image at high light intensities—maintaining strong optical binding interactions for many particles in a 3-dimensional space. Utilizing this novel experimental scheme, we describe the general static and dynamic self-organization behaviors for 20–100 particles. Among the observations, we describe the self-organization into large optically bound structures along with the formation of driven particle clusters. Furthermore we show that the appearance and behavior of these optically bound structures can be tuned using different particle properties. We find a relationship between scattering strength, by altering relative refractive index, and the presence of driven particle clusters. We propose that N -body mechanisms, expected to be prevalent in stronger scattering systems drive the emergent clustering behaviors.

Keywords optical binding · self-assembly · non-equilibrium · N -body · scattering · optical force · driven · colloid

1 Introduction

Self-organization is the spontaneous formation of structure via the interactions of their constituent particles (1). The resulting structures from a self-organized system can be found in equilibrium and non-equilibrium states; the latter provides a pathway towards active and adaptive matter. While adaptive materials are common in living systems such as cells, it is possible to artificially introduce dynamic forces which can externally deliver energy to a system while mediating self-organization (2). Dynamic self-organization is relatively new but active field of self-assembly with the goal of extending our knowledge of equilibrium thermodynamics onto living systems. Some examples include the dynamic assembly of non-biological components including active colloids (3; 4; 5), rotating discs (6; 7), and magnetic swimmers (8; 9; 10).

Optical binding is a long-range light-induced force which causes dielectric particles to interact mutually through light-scattering (11). In theory, optical binding can be felt between polarizable molecules (~ 1 nm) (12) up to biological cells (13) (~ 5 μ m). In practice, because the force is relatively weak and requires a strong scattering response, the effects of optical binding are observed most strongly for objects that are on the order of the wavelength of light in size. Moreover, optical binding can give rise to non-equilibrium forces because the optical field applied to the system is also a constant supply of external momentum which can contribute to particle motion (14), although this effect has been largely unexplored in previous experimental work. In principle optical binding is well understood – forces can be

*Use footnote for providing further information about author (webpage, alternative address)—*not* for acknowledging funding agencies.




Figure 1: (a) Experimental setup. The laser beam is split into two paths that are aligned to counter propagate through the colloidal sample. Both beams are adjusted so that they are totally internally reflected at the coverslip. The polarization and incident angle of each beam can be adjusted individually. The camera, electronically tunable lens, and illumination source are all synchronized to a wave-function generator for the acquisition of 3-dimensional videos. (b) Depth profile of sample cell, we define z -axis as direction along the sample depth and \hat{k} as the axis corresponding to the beam propagation axis. (c) image still of HIPs in the optical field. (d) image still of LIPs in the optical field.

exactly predicted using a multiple scattering approach – but nonetheless it remains difficult to predict and explain the behavior of systems composed of many particles (14).

Numerous past studies have illustrated the complexity of the optical binding force. Two-particle studies have shown that the force can contain multiple stable points defined by multiples of the light wavelength (11; 15; 16). By increasing the number of particles, previous experimental and computational studies have shown the self-organization of 1-dimensional chains exhibiting breathing modes (17), static and drifting 2-dimensional lattices exhibiting stable and quasi-stable behaviors (18), and observations of bistability (19) and multistability (20) in the equilibrium positions of optically bound particles (21). The complexity that arises from optical binding in multi-particle systems can largely be attributed to feedback in the form of interference with the incoming field through multiple scattering effects (21). One often neglected fact is that optical binding forces are sensitive to N -body effects, and can not be treated as strictly pair-wise interactions (22). These N -body interactions can lead to highly correlated and emergent behaviors but can consequently be difficult to model.

In this manuscript we study optical binding of large numbers of particles ($N > 20$) free to move in three-dimensional space. We assess which features of the assembly can be described by pairwise interactions versus features which emerge from higher-order effects. By comparing strong and weak scatterers, we share new insights on the relationship between refractive index, scattering strength, non-conservative forces, and non-linear optical binding. We present observations of unexpected emergent driven behaviors which appear to manifest from the deviation from pairwise forces. We also use the coupled dipole method (CDM) (23; 24), also known as the discrete dipole approximation (DDA), to explain the origin of these effects.

2 Experimental Design

2.1 Experimental Parameters

Generalizations about the structures resulting from optical binding are difficult to make. There are a number of relevant parameters which can alter how an object scatters light and consequentially alter the properties of the interactions. In practice, the most important quantities are the size parameter, ka , and the relative refractive index $m = n_p/n_0$ (where k is the wavenumber of the incident field in the background medium, a is the particle radius, and n_p/n_0 is the particle/background medium index of refraction).

Tuning the parameters will not necessarily yield linear responses, which is why we must be careful about generalizing optical binding behavior. For instance, the size parameter alone can span three different regimes: the Rayleigh limit, $ka \ll 1$, the ray-optics limit, $ka \gg 1$, and the Mie scattering regime, $ka \sim 1$, each of which are separated by drastically different assumptions including the relevance of multiple scattering, scattering modes, and how the strength of the forces scale. Our current study is focused in the Mie regime.

The relative refractive index, m , affects the scattering strength of a single particle, and so modulates the importance of multiple-scattering events (25). We use this study as an opportunity to contrast the behaviors of strong and weak scatterers. We perform a side-by-side analysis of two commonly available colloidal microspheres which differ in refractive index. We refer to these as high-index particles (HIPs, $m=1.2$, made of polystyrene) and low-index particles (LIPs, $m=1.1$, made of SiO₂).

Unique to field-driven self-organization, direction and geometry become particularly important. Because optical scattering is directional, optical binding is an interaction which breaks symmetry along the axis of the field direction. As a result, experimental and numerical studies have often focused on either two distinct optical binding geometries: lateral (11; 15; 16; 26; 18; 14) and longitudinal(27; 20; 28; 29; 17) optical binding. Lateral optical binding describes binding that occurs between particles with a displacement perpendicular to the direction of propagation, \hat{k} , while longitudinal optical binding describes binding of particles displaced along \hat{k} . In the current study, we implement a novel experimental approach to present for the first time the self-organization of many particles free to interact in all directions. We believe this configuration is particularly illuminating for understanding potential bulk behaviors.

Finally, previous studies have focused on how the stability of optically bound matter is affected by damping conditions (18; 14) suggesting that the low damping can lead to instabilities due to the systems inability to remove energy. Because we use sub-micron particles suspended in a fluid, our study is in an over-damped regime. Our estimated Reynold's number is on the order of $Re = 10^{-6}$ (assuming 500 nm diameter particles in water with a characteristic velocity of 1 $\mu\text{m/s}$).

2.2 Optical setup

The experiments are performed on a custom designed inverted microscope (**Fig. 1** (a)). Two Gaussian beams ($\lambda = 532$ nm in vacuum, or 400 nm in solution) are aligned and focused through the sample tube to generate a counter-propagating crossed polarized optical field ($w_0 = 27.0 \pm 0.3 \mu\text{m}$). The orientation of the beam through the sample is at an angle of approximately $\phi_g = 45^\circ$ through the glass layers and $\phi = 36^\circ$ through the water. We achieve total-internal reflection at the glass-air interface at the bottom surface of the coverslip depicted in **Fig. 1** (b). The sample tube depth is $10 \pm 1 \mu\text{m}$. We collect 3-dimensional video microscopy data using a combination of a microscope objective (40X, NA = 0.75) and an electronically tunable lens (ETL). The scanning of the ETL is synchronized with the imaging camera acquisition to capture 10 volumes per second with 50 frames per volume. The signal output to the ETL is in the form of a periodic sawtooth pattern (driven at 10 Hz) which increases linearly for 70 % of the total period and quickly decreases linearly for the remaining 30 %. As such, the first 70 % of the period is used to create the full volume. The 3D scanning method gave us a spatial resolution of 0.8 $\mu\text{m}/\text{pixel}$ in the z-direction and a time resolution of 10 Hz which was sufficient to reliably locate particles within the volume.

The colloidal particles in the sample exhibit only Brownian motion until they enter the beam region. Particles that diffuse into the region remain confined within the Gaussian beam but can move thermally in all 3-dimensions within the area of confinement. Radiation pressure forces, which can be a strong optical force in the direction of the field propagation (25), are carefully balanced by the two counter-propagating beams. As nearby colloids drift into the beam, the density of the particles increase slowly. While there is no direct control to stop diffusive colloids from drifting into the beam, the variability in the number of particles over the entire 2 mins of observation are between 1 and 5 particles ($\sim 3-7$ % of total particles).

The light intensity is a parameter which plays a linear role in the strength of the optical binding force and can be tuned quite easily. For the current study, we adjust the peak light intensity of the Gaussian beam up to $\sim 2.5 \text{ mW}/\mu\text{m}^2$. For




Figure 2: Strength of optical forces for a range of size-parameters (ka). The optical gradient force (blue) was numerically simulated by placing a single particle in a node of a standing wave, the optical scattering force (orange) was numerically simulated by placing a single particle in a propagating plane-wave. The optical binding force was obtained by placing to particles laterally in a counter-propagating field at a separation of 2λ . The reference force, F_0 , is also plotted. In all simulations we assume a light intensity of $1 \text{ mW}/\mu\text{m}^2$.

the numerical simulations we use a reference intensity of $I_0 = 1 \text{ mW}/\mu\text{m}^2$. Subsequently we define a reference force, F_0 , which is the total optical momentum impinging on a single particle in a plane wave of intensity I_0 :

$$F_0 = \frac{I_0 \pi r^2}{c}, \quad (1)$$

where r is the radius of the particle and c is the speed of light. The value approximately 650 fN for 500 nm diameter particles. It is also useful to compare this to thermal excitation. We can convert the thermal energy, kT , to an equivalent force by including a length scale. Given that the acoustic binding force has spatial oscillations with the wavelength of the incoming light, an effective thermal force is given by $F_T \sim kT/\lambda \sim 10 \text{ fN}$.

We performed a numerical analysis to predict suitable light intensity and particle parameters which should allow us to observe optical binding effects (**Fig. 2**). By comparing the scaling of optical forces – including binding – with an estimated force of thermal excitation, we show that the size parameter is important in determining the strength of optical binding effects relative to thermal motion. Working in the $ka = 4$ regime guarantees that all the optical forces are stronger than the thermal background at a power density of $1 \text{ mW}/\mu\text{m}^2$.

Potentially more important than the strength of the force is the nature of the interaction. We note that the optical binding force is a combination of second-order gradient and scattering forces. In **Fig. 2**, we plot the relative strength of first-order gradient and scattering forces for HIPs and LIPs to approximate their relative contributions to the binding force. Note that the optical binding force is computed only for a pair of particles spaced by 2λ ; for larger numbers of closely spaced particles it can be 1–2 orders of magnitude stronger, while other forces. We observe for $ka = 4$ that the HIPs have a strong scattering force response, relative to the gradient force. Because scattering forces are generally non-conservative, we expect that it is the stronger presence of scattering forces in optical binding for the HIPs that give them a more dynamic self-organization behavior as opposed to the LIPs.

2.3 Sample preparation

Colloidal samples are diluted in water ($\sim 0.01 \text{ w/v } \%$) and placed into thin rectangular Borosilicate tubes ($100 \mu\text{m}$ wide \times $10 \mu\text{m}$ thick \times 50 mm long) purchased from *VitroCom*. We performed experiments with polystyrene (‘HIPs’, diameter $d = 518 \pm 10 \text{ nm}$, relative index $m = 1.20$ (30; 31)) and silicon dioxide (‘LIPs’, diameter $d = 518 \pm 20 \text{ nm}$,




Figure 3: **(a,b)** Two-body force map of optical binding forces for HIPs and LIPs ($ka = 4$) placed in counter-propagating crossed-polarized plane wave ($\lambda = 0.4\mu\text{m}$ in solution) propagating in the $+\hat{k} = +z'$ and $-\hat{k} = -z'$ direction. Maps are obtained by placing two particles in a CDM simulation and calculating the force applied on each particle at various displacements which fill the map space. The forces are provided in a unit-less scale normalized by the reference force, F_0 , described by the single scattering force on a single particle of the same size and material. The direction of the arrows represent the direction of the force, while the size represents the relative strength. **(c-f)** 2D pair-correlation functions for HIPs in the x - y plane at various light intensities. **(g-j)** 2D pair-correlation functions for LIPs in the x - y plane at various light intensities. The color bar in the pair-correlation functions gives the 2D probability density of the particles, and has units of μm^{-2} .

relative index $m = 1.10$ (32; 31)) purchased from *microParticles GmbH*. In the experiments, the number of colloidal particles are in the range of 30-50 for HIP experiments and 40-100 for the LIP experiments. We found that LIPs were more readily collected into the beam and comparable initial densities to the HIPs. The tubes allow us to keep a precise sample depth of $10 \pm 1\mu\text{m}$. The tubes are coated with index matching fluid and placed in-between a coverslip and microscope slide. Finally, the coverslip is glued to the microscope slide, using UV curable Epoxy (*Norland Optical Adhesive NOA61*), which seals the tube and index matching fluid. The samples are fixed onto the *Mad City Labs - RM21* microscopy base allowing for micro-precision movement of the sample in the x - y plane.

3 Results

3.1 Observations of self-organization behavior

Optical binding effects are immediately apparent between particles within the beam region. For instance, spatial ordering is immediately observable within the beam area – effects which can be enhanced by increasing the light intensity. The most apparent feature is the generation of multiple chains of particles aligned along the beam propagation

direction (**Fig. 1 (c)**). Once collected into a chain, particles within the chain can be observed to move collectively together within the beam region. Being a larger region which can collect many-particles, we have the opportunity to observe the formation of multiple chains, and subsequent inter-chain behaviors.

Other features of the formed structures arise on further examination. For instance, the HIPs are found to self-organize into extended chains which extend along the sample depth—or–small clusters of typically (3-5) particles long. Both HIP structures can be observed in **Fig. 1 (c)** and †ESI Movie S1. Unlike the HIPs, the LIPs tend to form extremely long and close packed chains of particles (rods) (†ESI Movie S2). Two LIP rods can be observed in **Fig. 1 (d)**.

3.2 Comparing observations to two-body forces

To understand the dominating interactions which are leading to the general behavior of both systems, we numerically generated two-body force maps for the two particle types (**Fig. 3(a,b)**). The force maps were created using a CDM based simulation (23; 24) (see supplementary material for details). Comparing force maps for HIP ($m = 1.2$) and LIP ($m = 1.1$) particles reveals surprisingly little difference, apart from the overall strength which scales approximately like the reflectivity of a dielectric plane, which scales like $\sim (m - 1)^2$.

Despite this, the two-body force maps can be used to describe some of the overall structures observed in the system. The long range alignment along the beam propagation for the HIPs and LIPs are qualitatively consistent with the two-body force maps (**Fig. 3**), which suggest strong forces can tend to pull particles onto the axis of propagation (\hat{k}). The \hat{k} axis, as shown in **Fig. 1 (b)** is rotated 54° from z . For comparison, we approximate the energy $F_0\lambda \sim 65kT$, thus the HIPs can experience $\sim 20kT$ of energy keeping them aligned in the \hat{k} axis in the beam at $1 \text{ mW}/\mu\text{m}^2$. The LIPs experience $\sim 5kT$ at $1 \text{ mW}/\mu\text{m}^2$.

The HIPs force-map shows a repulsive force between particles that are aligned along the field propagation (**Fig. 3 (a)**). This fits with the observation of HIPs forming extended particle chains (**Fig. 1 (c)**), rather than tightly packing. What the force-map fails to predict for the HIPs are the presence of the smaller tightly packed HIP clusters also found in the experiment. While it is expected that we observe structures that cannot be predicted from two-body optical binding interactions, here we observe an emergence of a force which is not only strongly contributing, but can act in the opposite direction of the two-body force. For the HIPs, the two-body force is clearly represented but we observe experimentally an unexpected close-range attractive force suddenly drives particles together into small clusters.

3.3 Pair-correlation function

We use a pair-correlation function to map the time-average structure created by optical binding forces. The form of the pair-correlation function also gives an approximation for an effective potential through which the particles interact. We collected the data for the pair-correlation function by recording the positions of colloidal particles all located within the same weakly focused beam of light. The beam is fixed at a constant light intensity for a duration of 2 mins during which we collected 3-dimensional imaging data. The particles were located using the *trackpy* implementation of Crocker and Grier (33; 34). While the particle speeds and dense particle clustering made it challenging to track particles through time, we found that the algorithms were quite efficient at locating particles within a frame. We found relatively low fluctuations in the total number of particles from frame to frame suggesting that the particle locating algorithms were performing consistently. Inter-particle displacements were determined by particle locations given for each time-step, thus the frequencies of each displacement over the entire run were available. The pair-correlation functions are projected against the x - y plane.

The 2D pair-correlation function (PCF) for the HIPs suggest an increase in spatial order with the increase in light intensity **Fig. 3 (c-f)**. The form of the PCF shows the dominating feature that the HIPs tend to align along the beam propagation. At higher light intensities, $I > 1 \text{ mW}/\mu\text{m}^2$, multiple lines off-axis begin to appear. The HIP two-body force map (**Fig. 3 (a)**) can be used to help explain the multiple off-axis lines that appear in the PCF at higher light intensities, as we can observe multiple off-axis lines in which the force arrows converge.

The 2D PCF for the LIPs are very consistent with the observation that the particles collapse into tightly bound rods (**Fig. 3 (g-j)**). One feature that becomes prominent are secondary off-axis lines at higher light intensities. The secondary lines are evidence of multiple rods interacting to form a long-range regular spacing over time.

3.4 Frame Mean Squared Difference (FMSD)

We use the mean squared displacement of particles to determine their dynamic behavior. Due to high-density particle clustering and the high velocities of clusters evidenced in **Fig. 5**, it is difficult to reliably track individual particles through time. This is especially true for close-packed clusters, which – as we shall show later – appear to drive motion in




Figure 4: **(a)** Distribution of relative FMSD values over 2 min duration for various intensities. For the high light intensity case ($2.3 \text{ mW}/\mu\text{m}^2$) we share a comparative distribution of FMSD values for the same system in a counter-propagating standing wave. **(b)** The average relative FMSD value over 2 minutes for various light intensities. The shaded regions represent the variability in the relative FMSD values over the 2 minutes. Values over the dotted line represent the average motion of the particles greater than what would be observed in a purely diffusive system. **(g)** Distribution of relative FMSD values over 2 min duration for various light intensities. For the higher light intensity case ($2.3 \text{ mW}/\mu\text{m}^2$) we share a comparative distribution of FMSD values for the same system in a counter-propagating standing wave. **(h)** The average relative FMSD value over 2 minutes for various light intensities. The shaded regions represent the variability in the relative FMSD values over the 2 minutes. Values over the dotted line represent the average motion of the particles greater than what would be observed in a purely diffusive system.

the HIPs. Instead of attempting to directly track the particles in time, we use a proxy for the mean squared displacement by obtaining the mean squared difference between two subsequent image volumes. We found that the mean squared difference between frames (FMSD) gives us more consistent measures while remaining highly correlated with the true mean squared displacement (details of the method are described in the supplementary material). The FMSD, Δ , is given by:

$$\Delta(t) = \frac{1}{N_p(t)} \left[\sum_{i,j,k}^{volume} (p_{i,j,k,t} - p_{i,j,k,t-dt})^2 - \Delta_0 \right] \quad (2)$$

where $p_{i,j,k,t}$ is the pixel value at a given location and time, $dt = 0.1$ is the time between frames, and Δ_0 is the background FMSD caused primarily by camera noise, and is computed for when there are no moving particles in the frame.

As shown in the supplemental materials, Δ is proportional to the squared particle displacement provided the particles average motion between frames is smaller to or comparable the particle size. In practice, we normalize this value

relative to one obtained by turning the optical binding laser off, in which case the particles experience only Brownian motion.

The distribution average particle motion for the HIPs is provided in **Fig. 4** (a). At lower light intensities, we find narrow distributions of the average particle motion and average values lower than what is found in a Brownian system, reflecting the fact that the particles are being confined by the optical binding forces. At higher light intensities, we find that the distributions shift to higher average values and the size of the fluctuations are greater, indicating the presence of a new driving force.

Surprisingly, comparing the pair-correlation function to the FMSD values suggests that the average motion is increasing even as the particles are becoming more ordered. This would not be expected for a conservative pair-wise force; in this case ordering will result in lower fluctuations. We do indeed observe this for lower power levels ($<0.4 \text{ mW}/\mu\text{m}^2$). Above these power levels, the increasing motion suggests that we are forming collections of particles which experience additional non-conservative forces from the optical field.

We found that the behavior can be altered by aligning the polarizations of the counter-propagating beams to generate a standing wave pattern. In this configuration, the optical binding area includes multiple planes of high light intensity perpendicular to the propagation axis and separated by $\lambda/2$. While the overall average motion is suppressed in the standing wave, the system is still observed to fluctuate strongly between low average motion ($\sim 0.5 \times$ Brownian motion) and high average motion ($\sim 2.5 \times$ Brownian motion). At the same light intensities to the previous configuration, the standing wave had the effect of dramatically suppressing the overall motion (\dagger ESI Movie 3).

Using the same range of intensities, the same analysis was done for the LIP system (**Fig. 4** (b,d)). At lower light intensities, we find narrow distributions of the average particle motion and mean values lower than what is found in a Brownian system. At higher light intensities, the average values do not exceed what is found in the Brownian system; however the distributions are far less Gaussian. The distribution suggests that the average motion is low with occasional rare events that lead to high motion. The LIPs in the standing wave do not show significant differences than particles in the cross-polarized counter-propagating beam.

Comparing the dynamic behavior between the HIPs and LIPs, there are clear differences between the two systems. The HIPs are move faster due to the optical binding force. We believe that non-conservative second-order scattering forces, expected to be stronger for the HIPs, contribute to the higher overall motion. The LIPS are likely dominated by second-order gradient forces thus pack in the rod structure until density limitations require the formation of new rods. Where the standing-wave generates first-order gradient forces which compete with the second-order scattering forces to suppress the overall motion, the same standing-wave does not show much of an effect on the Lower-index particles which are already dominated by gradient forces.

The two-body force maps cannot describe how a cluster or rod forms; in fact, the two body force maps suggest a weak repulsive force among particles that are aligned along the beam propagation. As a result, we believe the cluster formation can only be explained by considering a complete N -body force which we compute using the CDM.

We find numerically that when the separation between aligned particles is sufficiently large ($> 0.7 \mu\text{m}$) the total force is repulsive for HIP chains and increases in strength with the number of particles (**Fig. 6** (a)). This suggests that a chain of particles aligned along the propagation axis should remain spread apart. This is consistent with some of the structures found in the experiment: the lower density HIP chains are often found extending from the lower to upper boundary of the sample. The strong repulsion at long range could also act as a barrier to cluster formation at very high light intensity and low particle density. The trend is similar for LIP chains at separations $> 0.4 \mu\text{m}$ (**Fig. 6** (b)).

At smaller distances, we observe a decrease in repulsive strength for all HIP chains. HIP chains of more than 3 particles, we also observe a change in sign (**Fig. 6** (a)). This result suggests that the two-body repulsive force (**Fig. 3** (a)) dominates as long as the particles are far away; however, when many HIPs become in close range, N -body forces are strong enough to overcome two-body repulsion and switch the sign of the force entirely.

The same analysis of the LIP chains of particles shows a similar reduction of the repulsive force at shorter distances; however, an attractive force does not appear until the number of particles exceeds $N = 14$ (**Fig. 6** (b)). The initial reduction of the force, even at small N , is indicative of the presence of non-pairwise forces; however the effects are clearly weaker in comparison to the HIPs.

The numerical results provide a potential pathway to the previously unexpected formation of clusters. The results also highlight the unique nature of the N -body optical binding forces. These effects are shown to be comparatively much stronger for the HIPs. This supports the hypothesis that a major factor driving the differences in behavior between the HIP and LIP systems is a stronger presence of N -body forces due to the higher scattering strength of HIPs.




Figure 5: **(a-c)** Three snapshots of the optically bound HIPs over a 0.2 s duration. Two clusters are circled: (red) cluster of 5 particles ballistically moving ($>80 \mu\text{m/s}$) through the sample and (blue) isolated cluster of 4 particles remaining relatively still.

3.5 Non-conservative forces on clusters

We simulated the effects of a particle cluster as it grows in size and interacts with neighboring particles. This simulation was simplified by focusing on 1-dimensional interactions that occur along the propagation of the beam. The clusters are in the form of a line of N touching particles aligned along the beam propagation direction. One extra particle is added to the simulation, along the same center-line but not in contact with the cluster. We then compute the total force on the clustered particles, F_c , as well as the force on the lone particle, F_p . To compare the relative presence of conservative and non-conservative forces, we use the following quantity:

$$\gamma = \frac{|F_c + F_p|}{|F_c - F_p|} \quad (3)$$

If the forces are equal and opposite – indicating a conservative interaction – we will obtain $\gamma = 0$. In contrast, if the system is equal but in the same direction, this would be indicative of a system which would be driven ($\gamma \rightarrow \infty$).

Observing γ for the interaction between the HIP cluster and particle, we find that the value increases exponentially as the cluster size increases from 2-6 particles (**Fig. 6 (c)**). At $\gamma > 1$ the force between the cluster and the particle are in the same direction, while $\gamma \gg 1$ the forces are highly non-conservative.

For HIPs, there are configurations which could lead to uni-directional interactions first appearing for clusters of size 4. For clusters of sizes > 5 particles, $\gamma > 1$ at long ranges. This result is in good agreement with our observations of driven HIP clusters. At certain cluster sizes, non-conservative forces appear to allow clusters of particles to propel in an otherwise symmetric initial field. Here we study a limited set of configurations due to their common observable presence in the experimental data; however, we expect there are a number of alternative configurations that can also exhibit highly propelled motion. We expect that these strongly non-conservative interactions are cause for the increased average motion in the high light intensity binding experiments (**Fig. 4 (a,b)**).

For a similar set of cluster sizes in the LIP case, we find $\gamma < 0.1$ regardless of cluster size (**Fig. 6 (d)**). Indeed, we observed in the experiments that LIPs can build themselves into extremely long rods without exhibiting strongly non-conservative responses at small cluster sizes, as we found for the HIPs. It is likely that non-conservative forces won't have a strong presence for LIPs until they've assembled into elongated rods for which we do observe fluctuations in the FMSD values.




Figure 6: (a,b) 1-dimensional simulation of forces on particle chains. Particles are placed into an equidistant configuration aligned with the beam axis of propagation ($\hat{k} = \hat{z}'$) with inter-particle distance, z'_s . At each inter-particle spacing, the total outward force is calculated. A positive force represents a net-force pushing particles away from each other while a negative force represents a net-force in which particles are attracted. We performed the simulation for (a) HIPs ($m=1.2$, $ka=4$) and (b) LIPs ($m=1.1$, $ka=4$) for various cluster sizes (N) (c,d) 1-dimensional simulation of forces between clusters and neighboring particles. A cluster of N touching spheres is aligned along the axis of propagation with a single particle and the absolute sum over the absolute difference of force on the cluster and the force on the particle are plotted for various particle-cluster spacings, z'_p . Simulations were performed for (c) HIP ($m=1.2$, $ka=4$) and (d) LIP ($m=1.1$, $ka=4$) particles.

4 Conclusion

Optical binding of many particles in 3D gives rise to complex behavior that can not be predicted from two-body interactions alone. A simplified view of the interaction does, however, predict some features of the resulting structures: in particular the formation of extended chains of particles along the axis of light propagation. However, comparing low to high index particles – whose two-body interaction is nearly identical apart from overall strength – already demonstrates differences that must arise from N -body interactions. In particular, the lower index particles form close packed (or nearly close packed) chains, while the higher index particles are aligned with the beam, but usually spaced out by several particle diameters. More strikingly, the dynamic behavior of these particles demonstrates effects that can only be explained by considering non-conservative N -body interactions. This is most dramatically demonstrated by the temporary formation of close-packed clusters of 4-5 high index particles, which appear to drive an instability of the system as a whole.

Our results suggest that although optical binding can be used to guide the self-organization of colloidal particles, the effects of N -body forces are critical for the assembly of many-particle systems. Potentially, these forces offer new possibilities: with proper system design these forces could be used to promote or evade specific configurations or produce driven arrangements with dynamic behavior. Unfortunately, predicting the behavior of large scale optically bound systems remains a difficult task without a quick and accurate method of computing the forces for a fully coupled N -body system. Efforts on this front can be aided by further experimentation which can validate assumptions regarding N -body and non-conservative effects over a larger parameter space than the current study.

References

- [1] George M. Whitesides and Bartosz Grzybowski. Self-Assembly at All Scales. *Science*, 295(5564):2418–2421, March 2002. ISSN 0036-8075, 1095-9203. doi:10.1126/science.1070821. URL <https://www.science.org/>

- doi/10.1126/science.1070821.
- [2] Bartosz A. Grzybowski, Christopher E. Wilmer, Jiwon Kim, Kevin P. Browne, and Kyle J. M. Bishop. Self-assembly: from crystals to cells. *Soft Matter*, 5(6):1110, 2009. ISSN 1744-683X, 1744-6848. doi:10.1039/b819321p. URL <http://xlink.rsc.org/?DOI=b819321p>.
- [3] F. Ginot, I. Theurkauff, F. Detcheverry, C. Ybert, and C. Cottin-Bizonne. Aggregation-fragmentation and individual dynamics of active clusters. *Nature Communications*, 9(1):696, February 2018. ISSN 2041-1723. doi:10.1038/s41467-017-02625-7. URL <https://www.nature.com/articles/s41467-017-02625-7>.
- [4] Andong Wang, Wenyue Shi, Jianbin Huang, and Yun Yan. Adaptive soft molecular self-assemblies. *Soft Matter*, 12(2):337–357, December 2015. ISSN 1744-6848. doi:10.1039/C5SM02397A. URL <https://pubs.rsc.org/en/content/articlelanding/2016/sm/c5sm02397a>.
- [5] Jonathan R. Howse, Richard A. L. Jones, Anthony J. Ryan, Tim Gough, Reza Vafabakhsh, and Ramin Golestanian. Self-Motile Colloidal Particles: From Directed Propulsion to Random Walk. *Physical Review Letters*, 99(4):048102, July 2007. doi:10.1103/PhysRevLett.99.048102. URL <https://link.aps.org/doi/10.1103/PhysRevLett.99.048102>.
- [6] Bartosz A. Grzybowski, Howard A. Stone, and George M. Whitesides. Dynamic self-assembly of magnetized, millimetre-sized objects rotating at a liquid–air interface. *Nature*, 405(6790):1033–1036, June 2000. ISSN 1476-4687. doi:10.1038/35016528. URL <https://www.nature.com/articles/35016528>.
- [7] Bartosz A. Grzybowski, Howard A. Stone, and George M. Whitesides. Dynamics of self assembly of magnetized disks rotating at the liquid–air interface. *Proceedings of the National Academy of Sciences*, 99(7):4147–4151, April 2002. ISSN 0027-8424, 1091-6490. doi:10.1073/pnas.062036699. URL <https://www.pnas.org/content/99/7/4147>.
- [8] Gašper Kokot, German V. Kolmakov, Igor S. Aranson, and Alexey Snezhko. Dynamic self-assembly and self-organized transport of magnetic micro-swimmers. *Scientific Reports*, 7(1):14726, November 2017. ISSN 2045-2322. doi:10.1038/s41598-017-15193-z. URL <https://www.nature.com/articles/s41598-017-15193-z>.
- [9] Pietro Tierno, Ramin Golestanian, Ignacio Pagonabarraga, and Francesc Sagués. Controlled Swimming in Confined Fluids of Magnetically Actuated Colloidal Rotors. *Physical Review Letters*, 101(21):218304, November 2008. doi:10.1103/PhysRevLett.101.218304. URL <https://link.aps.org/doi/10.1103/PhysRevLett.101.218304>.
- [10] Alexey Snezhko and Igor S. Aranson. Magnetic manipulation of self-assembled colloidal asters. *Nature Materials*, 10(9):698–703, September 2011. ISSN 1476-4660. doi:10.1038/nmat3083. URL <https://www.nature.com/articles/nmat3083>.
- [11] Michael M. Burns, Jean-Marc Fournier, and Jene A. Golovchenko. Optical binding. *Physical Review Letters*, 63(12):1233–1236, September 1989. doi:10.1103/PhysRevLett.63.1233. URL <https://link.aps.org/doi/10.1103/PhysRevLett.63.1233>.
- [12] T. Thirunamachandran. Intermolecular interactions in the presence of an intense radiation field. *Molecular Physics*, 40(2):393–399, June 1980. ISSN 0026-8976. doi:10.1080/00268978000101561. URL <https://doi.org/10.1080/00268978000101561>.
- [13] Jing Bai, Zhen-Sen Wu, and Cheng-Xian Ge. Lateral Optical Binding of Stratified Biological Cells Induced by High-Order Bessel Beams. In *2018 12th International Symposium on Antennas, Propagation and EM Theory (ISAPE)*, pages 1–3, December 2018. doi:10.1109/ISAPE.2018.8634151.
- [14] Xiao Li, Yineng Liu, Zhifang Lin, Jack Ng, and C. T. Chan. Non-Hermitian physics for optical manipulation uncovers inherent instability of large clusters. *Nature Communications*, 12(1):6597, November 2021. ISSN 2041-1723. doi:10.1038/s41467-021-26732-8. URL <https://www.nature.com/articles/s41467-021-26732-8>.
- [15] F Dapasse and J M Vigoureux. Optical binding force between two Rayleigh particles. *Journal of Physics D: Applied Physics*, 27(5):914–919, May 1994. ISSN 0022-3727, 1361-6463. doi:10.1088/0022-3727/27/5/006. URL <https://iopscience.iop.org/article/10.1088/0022-3727/27/5/006>.
- [16] Samarendra Kumar Mohanty, Joseph Thomas Andrews, and Pradeep Kumar Gupta. Optical binding between dielectric particles. *Optics Express*, 12(12):2746–2753, June 2004. ISSN 1094-4087. doi:10.1364/OPEX.12.002746. URL <https://www.osapublishing.org/oe/abstract.cfm?uri=oe-12-12-2746>.
- [17] S. A. Tatarkova, A. E. Carruthers, and K. Dholakia. One-Dimensional Optically Bound Arrays of Microscopic Particles. *Physical Review Letters*, 89(28):283901, December 2002. doi:10.1103/PhysRevLett.89.283901. URL <https://link.aps.org/doi/10.1103/PhysRevLett.89.283901>.

- [18] Jack Ng, Z. F. Lin, C. T. Chan, and Ping Sheng. Photonic clusters formed by dielectric microspheres: Numerical simulations. *Physical Review B*, 72(8):085130, August 2005. doi:10.1103/PhysRevB.72.085130. URL <https://link.aps.org/doi/10.1103/PhysRevB.72.085130>.
- [19] N. K. Metzger, K. Dholakia, and E. M. Wright. Observation of Bistability and Hysteresis in Optical Binding of Two Dielectric Spheres. *Physical Review Letters*, 96(6):068102, February 2006. doi:10.1103/PhysRevLett.96.068102. URL <https://link.aps.org/doi/10.1103/PhysRevLett.96.068102>.
- [20] V. Karásek, T. Čižmár, O. Brzobohatý, P. Zemánek, V. Garcés-Chávez, and K. Dholakia. Long-Range One-Dimensional Longitudinal Optical Binding. *Physical Review Letters*, 101(14):143601, October 2008. doi:10.1103/PhysRevLett.101.143601. URL <https://link.aps.org/doi/10.1103/PhysRevLett.101.143601>.
- [21] Kishan Dholakia and Pavel Zemánek. Colloquium: Grippled by light: Optical binding. *Reviews of Modern Physics*, 82(2):1767–1791, June 2010. doi:10.1103/RevModPhys.82.1767. URL <https://link.aps.org/doi/10.1103/RevModPhys.82.1767>.
- [22] Justo Rodríguez, Luciana C. Dávila Romero, and David L. Andrews. Optical binding in nanoparticle assembly: Potential energy landscapes. *Physical Review A*, 78(4):043805, October 2008. ISSN 1050-2947, 1094-1622. doi:10.1103/PhysRevA.78.043805. URL <https://link.aps.org/doi/10.1103/PhysRevA.78.043805>.
- [23] Bruce T. Draine and Piotr J. Flatau. Discrete-Dipole Approximation For Scattering Calculations. *JOSA A*, 11(4):1491–1499, April 1994. ISSN 1520-8532. doi:10.1364/JOSAA.11.001491. URL <https://www.osapublishing.org/josaa/abstract.cfm?uri=josaa-11-4-1491>.
- [24] A. G. Hoekstra, M. Frijlink, L. B. F. M. Waters, and P. M. A. Sloot. Radiation forces in the discrete-dipole approximation. *JOSA A*, 18(8):1944–1953, August 2001. ISSN 1520-8532. doi:10.1364/JOSAA.18.001944. URL <https://www.osapublishing.org/josaa/abstract.cfm?uri=josaa-18-8-1944>.
- [25] A. Ashkin, J. M. Dziedzic, J. E. Bjorkholm, and Steven Chu. Observation of a single-beam gradient force optical trap for dielectric particles. *Optics Letters*, 11(5):288–290, May 1986. ISSN 1539-4794. doi:10.1364/OL.11.000288. URL <https://www.osapublishing.org/ol/abstract.cfm?uri=ol-11-5-288>.
- [26] Ming-Tzo Wei, Jack Ng, C. T. Chan, and H. Daniel Ou-Yang. Lateral optical binding between two colloidal particles. *Scientific Reports*, 6(1):38883, December 2016. ISSN 2045-2322. doi:10.1038/srep38883. URL <https://www.nature.com/articles/srep38883>.
- [27] V. Karásek, K. Dholakia, and P. Zemánek. Analysis of optical binding in one dimension. *Applied Physics B*, 84(1):149–156, July 2006. ISSN 1432-0649. doi:10.1007/s00340-006-2297-8. URL <https://doi.org/10.1007/s00340-006-2297-8>.
- [28] V. Karásek, O. Brzobohatý, and P. Zemánek. Longitudinal optical binding of several spherical particles studied by the coupled dipole method. *Journal of Optics A: Pure and Applied Optics*, 11(3):034009, March 2009. ISSN 1464-4258, 1741-3567. doi:10.1088/1464-4258/11/3/034009. URL <https://iopscience.iop.org/article/10.1088/1464-4258/11/3/034009>.
- [29] Wolfgang Singer, Manfred Frick, Stefan Bernet, and Monika Ritsch-Marte. Self-organized array of regularly spaced microbeads in a fiber-optical trap. *JOSA B*, 20(7):1568–1574, July 2003. ISSN 1520-8540. doi:10.1364/JOSAB.20.001568. URL <https://www.osapublishing.org/josab/abstract.cfm?uri=josab-20-7-1568>.
- [30] Xiaoning Zhang, Jun Qiu, Jun Qiu, Jun Qiu, Xingcan Li, Xingcan Li, Junming Zhao, Junming Zhao, Linhua Liu, and Linhua Liu. Complex refractive indices measurements of polymers in visible and near-infrared bands. *Applied Optics*, 59(8):2337–2344, March 2020. ISSN 2155-3165. doi:10.1364/AO.383831. URL <https://www.osapublishing.org/ao/abstract.cfm?uri=ao-59-8-2337>.
- [31] G. M. Hale and M. R. Querry. Optical Constants of Water in the 200-nm to 200-microm Wavelength Region. *Applied Optics*, 12(3):555–563, March 1973. ISSN 1559-128X. doi:10.1364/AO.12.000555.
- [32] I. H. Malitson. Interspecimen Comparison of the Refractive Index of Fused Silica*,†. *JOSA*, 55(10):1205–1209, October 1965. doi:10.1364/JOSA.55.001205. URL <https://www.osapublishing.org/josa/abstract.cfm?uri=josa-55-10-1205>.
- [33] John C. Crocker and David G. Grier. Methods of Digital Video Microscopy for Colloidal Studies. *Journal of Colloid and Interface Science*, 179(1):298–310, April 1996. ISSN 0021-9797. doi:10.1006/jcis.1996.0217. URL <https://www.sciencedirect.com/science/article/pii/S0021979796902179>.
- [34] Dan Allan, Thomas Caswell, Nathan Keim, and Casper van der Wel. trackpy: Trackpy v0.3.0, November 2015. URL <https://zenodo.org/record/34028>.

FORMATION OF COLLOIDAL CHAINS AND DRIVEN CLUSTERS WITH OPTICAL BINDING: SUPPLEMENTAL

A PREPRINT

May 28, 2022

1 Optical scattering, gradient, and binding forces

The analysis in the manuscript was guided by comparing two types of scattering bodies, Polystyrene (PS) and Silica (SiO_2) particles, denoted as HIP and LIP. Initial calculations were done comparing the relative strengths of three optical forces: the scattering force, gradient force, and binding force.

The scattering force is a first-order optical force which arises from the back-reflection of incident photons (1). The scattering force is non-conservative and is dependent on the direction of light propagation. We determine the strength of the force through simulating a single particle in single plane wave.

The gradient force is a first-order optical force due to the induced dipole moments generated in the scattering object by the incident light (2). This force is completely conservative and is dependent on the gradient of the electric field. We determine the strength of the force through simulating a single particle in a standing wave generated by two counter-propagating plane waves. The two counter-propagating waves negate scattering forces while the standing wave pattern generates gradients in the optical field thus applying optical gradient forces onto the single object.

The optical binding force is a second-order optical force due to the modification of an incident field on an object by the scattered fields of other objects (3). Optical binding is a combination of conservative and non-conservative forces. We simulate the force by placing two objects separated in cross-polarized counter-propagating plane waves. The counter-propagation negates the first-order scattering forces while cross-polarizing removes the standing wave suppressing the first-order gradient force. Remaining forces must be due to optical binding effects.

At the size range of interest we find that PS particles have relatively stronger scattering forces while LIPs have relatively stronger gradient forces. The conclusion made is that HIPs are stronger scatterers.

2 Numerical Simulations

The numerical simulations were achieved using a Coupled Dipole Method (CDM) based scattering code. In each case, we ensure 10 dipoles per wavelength is used. We use an iterative solver in which the number of iterations are set until the root mean square of the dipole moments converge to $< 1\%$.




Figure 1: A simulated PS particle displaced along the x, y, and z axis plotted against a corresponding squared frame difference.

The 2-dimensional maps were generated by running a stand-alone simulation at each particle displacement. A single particle was fixed at an origin while a second particle was moved along a 2-dimensional space on the plane in the immediate vicinity of the fixed particle.

3 Force map generation

The force maps were generated using CDM based simulation (4; 5). By placing two particles in a cross-polarized counter-propagating field, we calculate the radiation forces placed on each particle. Filling out the 2-dimensional space is done by fixing one object at an origin and repeating the simulation for various displacements.

4 Correlating frame mean squared difference (FMSD) with mean squared displacement (MSD)

3-Dimensional videos were taken of particles undergoing Brownian motion. 19 individual particles located in a single volume were averaged to generate a reference particle. We used the reference particle to simulate particle motion by shifting it a known amount in a generated noisy background. The results of the simulated displacements along different axes are provided in **Fig. 1**. We can correlate the size of the simulated step to the value of taking the sum squared frame difference (FSD) of the array before and after the step. We found that larger displacements correspond to a larger




Figure 2: FMSD values compared to 2-dimensional mean squared displacement (MSD) in the xy -plane for (a) HIP (PS) and (b) LIP (SiO_2) particles.

FSD in general. One limitation is that the motion in the z -direction is underrepresented compared to motion in the x - y plane due to the inherent asymmetry of the particles imaging profile along the z -axis. This limitation does not allow us to place an absolute estimate of how fast the objects are moving in physical units; however we can still measure the relative motion under the assumption that the motion in the compared systems share similar axial preferences. Observations of the data support this assumption. The second limitation is that the FSD value is capped at displacements that are slightly larger than the particle diameter. At the volume rate of 10 volumes/s, particles undergoing Brownian motion are moving slow enough; however we have observed that particles undergoing optical binding forces can be propelled much faster. One way to alleviate the limitation is to record at a higher volume rate; however, it is currently unfeasible to cover the full range of possible particle speeds. This limitation is also present in traditional tracking methods.

Although the motion of the fastest particles are underrepresented in the FSD values, the FMSD method still allows a maximum value to be assigned. Traditional tracking methods would not be able to assign a velocity value at all. We also note that underrepresenting the motion of the faster moving particles only acts to underrepresent the strength of the findings.

The FMSD at any given time is determined by subtracting two subsequent volumes from each other, removing the contribution from the background, and dividing by the total number of particles in the frame:

$$\Delta(t) = \frac{1}{N_p(t)} \left[\sum_{i,j,k}^{volume} (p_{i,j,k,t} - p_{i,j,k,t-dt})^2 - \Delta_0 \right] \quad (1)$$

The background contribution, Δ_0 , is determined in each video by monitoring the squared difference of sections that have zero particles. We find a non-zero constant value which can be subtracted to remove contributions that are due to noise. The number of particles, N_p , is estimated by the

following:

$$N_p = c_0 \sum_{i,j,k}^{volume} |p_{i,j,k,t} - b_{i,j,k}| + c_1 \quad (2)$$

Where $b_{i,j,k}$ is belongs to a volume representing the sample devoid of particles. c_0 and c_1 are fitting parameters determined by linearly fitting the number of particles found by traditional tracking methods to $\sum_{i,j,k}^{volume} |p_{i,j,k,t} - b_{i,j,k}|$ for a system of particles undergoing Brownian motion. Some results in **Fig. 2**. For the HIPs, we show that for diffusive particles and particles interacting in a weak field, there is a clear correlation between the FMSD and the true MSD (**Fig. 2**). For the LIPs, we also find correlation between the FMSD and true MSD for the diffusive particles; however in a weak field, the correlation is reduced. We found that in a weak field, the Lower-index particles tended to pack into tight formations which effected the individual tracking capabilities. Thus we utilize the purely diffusive correlation to justify the use of FMSD in a difficult tracking atmosphere.

We represent the motion in terms of the FMSD values relative to the Brownian Motion case. Doing so allows us to compare the HIP and LIP systems in how their motion changes relative to diffusive motion.

5 Electronic Supplementary Information

All movies are 2 mins played in real-time.

5.1 Movie S1

HIP (Polystyrene) particles in cross polarized counter-propagating beam (peak power density - 1.5 mW/ μm^2).

5.2 Movie S2

LIP (SiO₂) particles in cross polarized counter-propagating beam (peak power density - 1.5 mW/ μm^2).

5.3 Movie S2

HIP (Polystyrene) particles in standing wave (peak power density - 1.5 mW/ μm^2).

References

- [1] A. Ashkin. Acceleration and trapping of particles by radiation pressure. *Physical Review Letters*, 24(4):156–159. doi:10.1103/PhysRevLett.24.156. URL <https://link.aps.org/doi/10.1103/PhysRevLett.24.156>.
- [2] A. Ashkin, J. M. Dziedzic, J. E. Bjorkholm, and Steven Chu. Observation of a single-beam gradient force optical trap for dielectric particles. *Optics Letters*, 11(5):288–290. ISSN 1539-4794. doi:10.1364/OL.11.000288. URL <https://www.osapublishing.org/ol/abstract.cfm?uri=ol-11-5-288>.
- [3] Michael M. Burns, Jean-Marc Fournier, and Jene A. Golovchenko. Optical binding. *Physical Review Letters*, 63(12):1233–1236. doi:10.1103/PhysRevLett.63.1233. URL <https://link.aps.org/doi/10.1103/PhysRevLett.63.1233>.

- [4] A. G. Hoekstra, M. Frijlink, L. B. F. M. Waters, and P. M. A. Sloot. Radiation forces in the discrete-dipole approximation. *JOSA A*, 18(8):1944–1953. ISSN 1520-8532. doi:10.1364/JOSAA.18.001944. URL <https://www.osapublishing.org/josaa/abstract.cfm?uri=josaa-18-8-1944>.
- [5] Bruce T. Draine and Piotr J. Flatau. Discrete-dipole approximation for scattering calculations. *JOSA A*, 11(4):1491–1499. ISSN 1520-8532. doi:10.1364/JOSAA.11.001491. URL <https://www.osapublishing.org/josaa/abstract.cfm?uri=josaa-11-4-1491>.