ODD RANK VECTOR BUNDLES IN ETA-PERIODIC MOTIVIC HOMOTOPY THEORY

OLIVIER HAUTION

ABSTRACT. We observe that, in the eta-periodic motivic stable homotopy category, odd rank vector bundles behave to some extent as if they had a nowhere vanishing section. We discuss some consequences concerning SL^c -orientations of motivic ring spectra, and the étale classifying spaces of certain algebraic groups. In particular, we compute the classifying spaces of diagonalisable groups in the eta-periodic motivic stable homotopy category.

INTRODUCTION

Around forty years ago, Arason computed the Witt groups of projective spaces [Ara80]. This computation was later revisited by Gille [Gil01], Walter [Wal03], and Nenashev [Nen09]. It exhibited Witt groups as a somewhat exotic cohomology theory, whose value on projective spaces differs quite drastically from what is obtained in more classical cohomology theories such as Chow groups or K-theory. It is now understood that this behaviour reflects the lack of GL-orientation in Witt theory.

Ananyevskiy observed in [Ana16a] that the key property of Witt groups permitting to perform these computations turns out to be the fact that the Hopf map

$$\eta \colon \mathbb{A}^2 \smallsetminus \{0\} \to \mathbb{P}^1, \quad (x:y) \mapsto [x:y]$$

induces by pullback an isomorphism of Witt groups $W(\mathbb{P}^1) \xrightarrow{\sim} W(\mathbb{A}^2 \setminus \{0\})$. He thus extended in [Ana16a] the above-mentioned computations to arbitrary cohomology theories in which η induces an isomorphism.

Inverting the Hopf map η in the motivic stable homotopy category $\operatorname{SH}(S)$ over a base scheme S yields its η -periodic version $\operatorname{SH}(S)[\eta^{-1}]$, a category which has been studied in details by Bachmann-Hopkins [BH20]. In this paper, we lift Ananyevskiy's computations of the cohomology of projective spaces to the η -periodic stable homotopy category: we obtain for instance that a projective bundle of even relative dimension becomes an isomorphism in $\operatorname{SH}(S)[\eta^{-1}]$.

Another familiar feature of Witt groups is that twisting these groups by squares of line bundles has no effect, which may be viewed as a manifestation of the SL^c -orientability

Date: July 12, 2023.

²⁰²⁰ Mathematics Subject Classification. 14F42.

Key words and phrases. motivic homotopy theory, orientations, motivic Hopf map, étale classifying spaces of linear groups.

This work was supported by the DFG research grant HA 7702/5-1 and Heisenberg grant HA 7702/4-1.

of Witt groups (see below). We show that that property of Witt groups in fact follows from their η -periodicity alone (see (4.3.2) for a more general statement):

Proposition. Let $V \to X$ be a vector bundle and $L \to X$ a line bundle. Then we have an isomorphism of Thom spaces

$$\operatorname{Th}_X(V) \simeq \operatorname{Th}_X(V \otimes L^{\otimes 2}) \in \operatorname{SH}(S)[\eta^{-1}].$$

Panin and Walter introduced [PW18, §3] the notion of SL^c -orientability for algebraic cohomology theories, which consists in the data of Thom classes for vector bundles equipped with a square root of their determinant, and proved that Hermitian K-theory is SL^c -oriented. Ananyevskiy later showed [Ana20, Theorem 1.2] that a cohomology theory is SL^c -oriented as soon as it is a Zariski sheaf in bidegree (0, 0), and pointed out [Ana20, Theorem 1.1] the close relations between SL^c -orientations and SL-orientations (the latter consisting in the data of Thom classes for vector bundles with trivialised determinant). We show in this paper that the two notions actually coincide in the η -periodic context:

Theorem. Every SL-orientation of an η -periodic motivic commutative ring spectrum is induced by a unique SL^c-orientation.

These results are obtained as consequences of the following observation:

Proposition. Let E be a vector bundle of odd rank over a smooth S-scheme X, and E° the complement of the zero-section in E.

- (i) The projection $E^{\circ} \to X$ admits a section in $SH(S)[\eta^{-1}]$.
- (ii) The diagram $E^{\circ} \times_X E^{\circ} \Rightarrow E^{\circ} \to X$ becomes a split coequaliser diagram in $SH(S)[\eta^{-1}]$.

The first assertion may be viewed as a splitting principle, while the second permits to perform a form of descent. To some extent, this proposition allows us to assume that odd rank vector bundle admit a nowhere-vanishing section (once η is inverted); in particular that line bundles are trivial.

Finally, we provide applications to the computation in $SH(S)[\eta^{-1}]$ of the étale classifying spaces of certain algebraic groups.

Theorem. For $r \in \mathbb{N} \setminus \{0\}$, there exist natural maps

 $S \to \mathbb{B}\mathbb{G}_m$; $S \to \mathbb{B}\mu_{2r+1}$; $\mathbb{G}_m \to \mathbb{B}\mu_{2r}$

which become isomorphisms in $SH(S)[\eta^{-1}]$.

From this theorem, we deduce a computation in $SH(S)[\eta^{-1}]$ of the classifying space of an arbitrary diagonalisable group. We obtain that all invariants of torsors under a diagonalisable group with values in an η -periodic cohomology theory arise from a single invariant of μ_2 -torsors. In the appendix, we present an explicit construction of that invariant, exploiting the identification of the group μ_2 with the orthogonal group O_1 .

Next, we obtain "relative" computations in $SH(S)[\eta^{-1}]$ of certain étale classifying spaces in terms of others:

Theorem. For $n \in \mathbb{N} \setminus \{0\}$ and $r \in \mathbb{N}$, the natural morphisms

 $BSL_n \to BSL_n^c$; $BGL_{2r} \to BGL_{2r+1}$; $BSL_{2r+1} \to BGL_{2r+1}$

become isomorphisms in $SH(S)[\eta^{-1}]$.

The first result can be viewed as a companion of the theorem on orientations stated above, and cements the idea that the groups SL_n^c and SL_n are the same in the eyes of η -periodic stable homotopy theory. The second (resp. third) result expresses the fact that odd-dimensional vector bundles behave as if they had a nowhere-vanishing section (resp. trivial determinant) from the point of view of η -periodic stable homotopy theory.

The morphism $BSL_{2r} \to BGL_{2r}$ is not an isomorphism in $SH(S)[\eta^{-1}]$, but we show that it admits a section, expressing the fact that every invariant (with values in an η -periodic cohomology theory) of even-dimensional vector bundles is determined by its value on those bundles having trivial determinant.

Finally, let us mention that the results of this paper serve as a starting point for the paper [Hau22] on Pontryagin classes.

Acknowledgments. I am grateful to the referee for his/her suggestions, and in particular for noticing that the coequaliser diagram of (2.2.7) is split.

1. NOTATION AND BASIC FACTS

1.1. Throughout the paper, we work over a noetherian base scheme S of finite dimension. The category of smooth separated S-schemes of finite type will be denoted by Sm_S . All schemes will be implicitly assumed to belong to Sm_S , and the notation $\mathbb{A}^n, \mathbb{P}^n, \mathbb{G}_m$ will refer to the corresponding S-schemes. We will denote by 1 the trivial line bundle over a given scheme in Sm_S .

1.2. We will use the \mathbb{A}^1 -homotopy theory introduced by Morel–Voevodsky [MV99]. We will denote by $\operatorname{Spc}(S)$ the category of motivic spaces (i.e. simplicial presheaves on Sm_S), by $\operatorname{Spc}_{\bullet}(S)$ its pointed version, and by $\operatorname{Spt}(S)$ the category of *T*-spectra, where $T = \mathbb{A}^1/\mathbb{G}_m$. We endow these with the motivic equivalences, resp. stable motivic equivalences, and denote by $\operatorname{H}(S)$, $\operatorname{H}_{\bullet}(S)$, $\operatorname{SH}(S)$ the respective homotopy categories. We refer to e.g. [PPR09, Appendix A] for more details.

We have an infinite suspension functor Σ^{∞} : $\operatorname{Spc}_{\bullet}(S) \to \operatorname{Spt}(S)$. Composing with the functor $\operatorname{Spc}(S) \to \operatorname{Spc}_{\bullet}(S)$ adding an external base-point, we obtain a functor Σ^{∞}_{+} : $\operatorname{Spc}(S) \to \operatorname{Spt}(S)$.

The spheres are denoted as usual by $S^{p,q} \in \operatorname{Spc}_{\bullet}(S)$ for $p, q \in \mathbb{N}$ with $p \ge q$ (where $T \simeq S^{2,1}$). The motivic sphere spectrum $\Sigma^{\infty}_{+}S$ will be denoted by $\mathbf{1}_{S} \in \operatorname{Spt}(S)$. When A is a motivic spectrum, we denote its (p,q)-th suspension by $\Sigma^{p,q}A = S^{p,q} \wedge A$. This yields functors $\Sigma^{p,q} \colon \operatorname{SH}(S) \to \operatorname{SH}(S)$ for $p, q \in \mathbb{Z}$.

1.3. When $E \to X$ is a vector bundle with $X \in \operatorname{Sm}_S$, we denote by $E^\circ = E \setminus X$ the complement of the zero-section. The Thom space of E is the pointed motivic space $\operatorname{Th}_X(E) = E/E^\circ$. We will write $\operatorname{Th}_X(E) \in \operatorname{Spt}(S)$ instead of $\Sigma^\infty \operatorname{Th}_X(E)$, in order to lighten the notation. When $g: Y \to X$ is a morphism in Sm_S , we will usually write

 $\mathbf{3}$

 $\operatorname{Th}_Y(E)$ instead of $\operatorname{Th}_Y(g^*E)$. Since $E \to X$ is a weak equivalence, we have a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$, where $p \colon E^{\circ} \to X$ is the projection,

(1.3.a)
$$(E^{\circ})_+ \xrightarrow{p_+} X_+ \to \operatorname{Th}_X(E).$$

If $F \to S$ is a vector bundle and $f: X \to S$ the structural morphism, we have by [MV99, Proposition 3.2.17 (1)] a natural identification in $\text{Spc}_{\bullet}(S)$

(1.3.b)
$$\operatorname{Th}_X(E \oplus f^*F) = \operatorname{Th}_X(E) \wedge \operatorname{Th}_S(F).$$

When $V \to S$ is a vector bundle, we denote by $\Sigma^V \colon \operatorname{SH}(S) \to \operatorname{SH}(S)$ the derived functor induced by $A \mapsto A \wedge \operatorname{Th}_S(V)$. It is an equivalence of categories, with inverse denoted by Σ^{-V} .

1.4. Let $i: Y \to X$ be a closed immersion in Sm_S , with normal bundle $N \to Y$, and open complement $u: U \to X$. The purity equivalence $X/U \simeq \operatorname{Th}_Y(N)$ (see e.g. [MV99, Theorem 3.2.23]) yields a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$

(1.4.a)
$$U_+ \xrightarrow{u_+} X_+ \to \operatorname{Th}_Y(N).$$

More generally, if $V \to X$ is a vector bundle, we have a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$

 $\operatorname{Th}_U(V) \to \operatorname{Th}_X(V) \to \operatorname{Th}_Y(N \oplus i^*V).$

This may be deduced from (1.4.a) by first reducing to the case X = S using the functor f_{\sharp} of (1.9) below, and then applying the functor $-\wedge \operatorname{Th}_X(V)$ (both of which preserve homotopy colimits), in view of (1.3.b).

1.5. Let $\varphi \colon E \xrightarrow{\sim} F$ be an isomorphism of vector bundles over $X \in \mathrm{Sm}_S$. Then φ induces a weak equivalence in $\mathrm{Spc}_{\bullet}(S)$ (and $\mathrm{Spt}(S)$)

$$\operatorname{Th}(\varphi) \colon \operatorname{Th}_X(E) \to \operatorname{Th}_X(F).$$

If $\psi \colon F \xrightarrow{\sim} G$ is an isomorphism of vector bundles over X, we have

(1.5.a)
$$\operatorname{Th}(\psi \circ \varphi) = \operatorname{Th}(\psi) \circ \operatorname{Th}(\varphi)$$

If X = S and $f \in \operatorname{End}_{\operatorname{SH}(S)}(\mathbf{1}_S)$, then we have in $\operatorname{SH}(S)$

(1.5.b)
$$(\Sigma^F f) \circ \operatorname{Th}(\varphi) = \operatorname{Th}(\varphi) \circ (\Sigma^E f) \colon \operatorname{Th}_S(E) \to \operatorname{Th}_S(F).$$

(This follows from the fact that, as morphisms $\mathbf{1}_S \wedge \operatorname{Th}_S(E) \to \mathbf{1}_S \wedge \operatorname{Th}_S(F)$

$$(f \wedge \mathrm{id}_{\mathrm{Th}_{S}(F)}) \circ (\mathrm{id}_{\mathbf{1}_{S}} \wedge \mathrm{Th}(\varphi)) = f \wedge \mathrm{Th}(\varphi) = (\mathrm{id}_{\mathbf{1}_{S}} \wedge \mathrm{Th}(\varphi)) \circ (f \wedge \mathrm{id}_{\mathrm{Th}_{S}(E)}).)$$

1.6. (See [Mor04, Lemma 6.3.4].) Let $X \in \text{Sm}_S$ and $u \in H^0(X, \mathbb{G}_m)$. Consider the automorphism $u \operatorname{id}_1 \colon 1 \to 1$ of the trivial line bundle over X, and set

 $\langle u \rangle = \Sigma^{-2,-1} \operatorname{Th}(u \operatorname{id}_1) \in \operatorname{Aut}_{\operatorname{SH}(S)}(\Sigma^{\infty}_+ X).$

It follows from (1.5.a) that

(1.6.a)
$$\langle uv \rangle = \langle u \rangle \circ \langle v \rangle \quad \text{for } u, v \in H^0(X, \mathbb{G}_m).$$

When X = S and $A \in SH(S)$, we will denote again by $\langle u \rangle \in Aut_{SH(S)}(A)$ the morphism

$$A = \mathbf{1}_S \land A \xrightarrow{\langle u \rangle \land \mathrm{id}_A} \mathbf{1}_S \land A = A.$$

If $f: A \to B$ is a morphism in SH(S), then

(1.6.b)
$$f \circ \langle u \rangle = \langle u \rangle \circ f.$$

1.7. We denote by $\eta: \mathbb{A}^2 \setminus \{0\} \to \mathbb{P}^1$ in $\operatorname{Spc}_{\bullet}(S)$ the map $(x, y) \mapsto [x: y]$, where $\mathbb{A}^2 \setminus \{0\}$ is pointed by (1, 1) and \mathbb{P}^1 by [1: 1].

1.8. We will consider the categories $\operatorname{Spc}_{\bullet}(S)[\eta^{-1}]$ and $\operatorname{Spt}(S)[\eta^{-1}]$ obtained by monoidally inverting the map η of (1.7), which can be constructed as left Bousfield localisations, as discussed in [Bac18, §6]. Their respective homotopy categories will be denoted by $\operatorname{H}_{\bullet}(S)[\eta^{-1}]$ and $\operatorname{SH}(S)[\eta^{-1}]$, and we will usually omit the mention of the localisation functors.

A spectrum $A \in \operatorname{Spt}(S)$ is called η -periodic if the map

(1.8.a)
$$A \wedge \Sigma^{\infty}(\mathbb{A}^2 \setminus \{0\}) \xrightarrow{\mathrm{id} \wedge \Sigma^{\infty} \eta} A \wedge \Sigma^{\infty} \mathbb{P}^1$$

is an isomorphism in SH(S). The full subcategory of such objects in Spt(S) can be identified $Spt(S)[\eta^{-1}]$.

1.9. Let $X \in \text{Sm}_S$ with structural morphism $f: X \to S$. Then there are Quillen adjunctions

$$f_{\sharp} \colon \operatorname{Spc}_{\bullet}(X) \leftrightarrows \operatorname{Spc}_{\bullet}(S) \colon f^* \quad ; \quad f_{\sharp} \colon \operatorname{Spt}(X) \leftrightarrows \operatorname{Spt}(S) \colon f^*.$$

The functor f^* is induced by base-change, while f_{\sharp} arises from viewing a smooth X-scheme as a smooth S-scheme by composing with f. These induce Quillen adjunctions

$$f_{\sharp} \colon \operatorname{Spc}_{\bullet}(S)[\eta^{-1}] \leftrightarrows \operatorname{Spc}_{\bullet}(S)[\eta^{-1}] \colon f^* \quad ; \quad f_{\sharp} \colon \operatorname{Spt}(X)[\eta^{-1}] \leftrightarrows \operatorname{Spt}(S)[\eta^{-1}] \colon f^*.$$

We will also use the notation f^* , f_{\sharp} for the derived functors on the respective homotopy categories.

1.10. (See e.g. [DHI04].) Let $V \to X$ be a vector bundle with $X \in \text{Sm}_S$, and U_{α} an open covering of X. Then the map

hocolim
$$\left(\cdots \rightrightarrows \prod_{\alpha,\beta} \operatorname{Th}_{U_{\alpha} \cap U_{\beta}}(V|_{U_{\alpha} \cap U_{\beta}}) \rightrightarrows \prod_{\alpha} \operatorname{Th}_{U_{\alpha}}(V|_{U_{\alpha}}) \right) \to \operatorname{Th}_{X}(V)$$

is a weak equivalence.

2. Splitting \mathbb{G}_m -torsors

2.1. Local splitting. In this section we consider the schemes $\mathbb{A}^1, \mathbb{G}_m, \mathbb{P}^1, \mathbb{A}^2 \setminus \{0\}$ as pointed motivic spaces, respectively via 1, 1, [1:1], (1,1). We recall that $T = \mathbb{A}^1/\mathbb{G}_m$. We have a chain of weak equivalences

(2.1.0.a)
$$T = \mathbb{A}^1 / \mathbb{G}_m \xrightarrow{\sim} \mathbb{P}^1 / \mathbb{A}^1 \xleftarrow{\sim} \mathbb{P}^1,$$

where the first arrow is induced by the immersion $\mathbb{A}^1 \to \mathbb{P}^1, x \mapsto [x:1]$, and the quotient $\mathbb{P}^1/\mathbb{A}^1$ is taken with respect to the immersion $\mathbb{A}^1 \to \mathbb{P}^1, y \mapsto [1:y]$.

We first recall a well-known fact (see e.g. [Ana20, Lemma 6.2] for a stable version):

2.1.1. Lemma. Let $u \in H^0(S, \mathbb{G}_m)$. Then the morphism $\operatorname{Th}(u^2 \operatorname{id}_1) \colon T \to T$ (see (1.5)) coincides with the identity in $\operatorname{H}_{\bullet}(S)$.

5

Proof. The endomorphism $\varphi \colon \mathbb{P}^1 \to \mathbb{P}^1$ given by $[x : y] \mapsto [u^2x : y] = [ux : u^{-1}y]$ is induced by the matrix

$$A = \begin{pmatrix} u & 0 \\ 0 & u^{-1} \end{pmatrix}.$$

Since

$$A = \begin{pmatrix} 1 & u \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -u^{-1} & 1 \end{pmatrix} \begin{pmatrix} 1 & u - 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}$$

is a product of transvections, the endomorphism φ induces the identity endomorphism of \mathbb{P}^1_+ in $\mathrm{H}_{\bullet}(S)$ (see e.g. [Ana16a, Lemma 1]). The map φ stabilises the copies of \mathbb{A}^1 given by $x \mapsto [x:1]$ and $y \mapsto [1:y]$, and restricts to $u^2 \operatorname{id}_1$ on the former. Thus the statement follows from the isomorphism (2.1.0.a). \Box

2.1.2. Excision yields isomorphisms in $H_{\bullet}(S)$

$$(\mathbb{A}^2 \setminus \{0\})/(\mathbb{G}_m \times \mathbb{A}^1) \stackrel{\sim}{\leftarrow} (\mathbb{A}^1 \times \mathbb{G}_m)/(\mathbb{G}_m \times \mathbb{G}_m) \stackrel{\sim}{\to} (\mathbb{A}^1/\mathbb{G}_m) \wedge (\mathbb{G}_m)_+ = T \wedge (\mathbb{G}_m)_+$$

Composing with the quotient $\mathbb{A}^2 \setminus \{0\} \to (\mathbb{A}^2 \setminus \{0\})/(\mathbb{A}^1 \times \mathbb{G}_m)$, this yields a map

(2.1.2.a) $\mathbb{A}^2 \setminus \{0\} \to T \land (\mathbb{G}_m)_+.$

2.1.3. Lemma. The projection $p: \mathbb{G}_m \to S$ induces a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$

$$\mathbb{A}^2 \smallsetminus \{0\} \xrightarrow{(2.1.2.a)} T \land (\mathbb{G}_m)_+ \xrightarrow{\operatorname{id} \land p_+} T$$

Proof. This follows from the consideration of the following commutative diagram in $\operatorname{Spc}_{\bullet}(S)$, whose rows are cofiber sequences

$$\begin{array}{c} \mathbb{A}^{2} \smallsetminus \{0\} \longrightarrow (\mathbb{A}^{2} \smallsetminus \{0\}) / (\mathbb{G}_{m} \times \mathbb{A}^{1}) \longrightarrow S^{1,0} \land (\mathbb{G}_{m} \times \mathbb{A}^{1}) \\ \uparrow & \uparrow^{\sim} & \uparrow \\ \mathbb{A}^{1} \times \mathbb{G}_{m} \longrightarrow (\mathbb{A}^{1} \times \mathbb{G}_{m}) / (\mathbb{G}_{m} \times \mathbb{G}_{m}) \longrightarrow S^{1,0} \land (\mathbb{G}_{m} \times \mathbb{G}_{m}) \\ \downarrow & \downarrow^{\sim} & \downarrow \\ \mathbb{A}^{1} \land (\mathbb{G}_{m})_{+} \longrightarrow (\mathbb{A}^{1} / \mathbb{G}_{m}) \land (\mathbb{G}_{m})_{+} \longrightarrow S^{1,0} \land \mathbb{G}_{m} \land (\mathbb{G}_{m})_{+} \\ \stackrel{\mathrm{id} \land p_{+}}{\longrightarrow} & \stackrel{\mathrm{id} \land p_{+}}{\longrightarrow} & \stackrel{\mathrm{id} \land \mathrm{id} \land p_{+}}{\longrightarrow} \\ \mathbb{A}^{1} \longrightarrow \mathbb{A}^{1} / \mathbb{G}_{m} \xrightarrow{\sim} & S^{1,0} \land \mathbb{G}_{m} \end{array}$$

and where the curved arrow is the weak equivalence induced by the projection $\mathbb{G}_m \times \mathbb{A}^1 \to \mathbb{G}_m$.

The next lemma is reminiscent of [Ana16b, Theorem 3.8]:

2.1.4. Lemma. The morphism η of (1.7) factors in $H_{\bullet}(S)$ as

$$\mathbb{A}^2 \smallsetminus \{0\} \xrightarrow{(2.1.2.\mathrm{a})} T \land (\mathbb{G}_m)_+ \xrightarrow{\mathrm{Th}(t^{-1} \operatorname{id}_1)} T \land (\mathbb{G}_m)_+ \xrightarrow{\mathrm{id} \land p_+} T \xrightarrow{(2.1.0.\mathrm{a})} \mathbb{P}^1,$$

where $t \in H^0(\mathbb{G}_m, \mathbb{G}_m)$ is the tautological section and $p: \mathbb{G}_m \to S$ the projection (and $T \land (\mathbb{G}_m)_+$ is identified with $\mathrm{Th}_{\mathbb{G}_m}(1)$).

Proof. Consider the commutative diagram in Sm_S



where the upper horizontal arrow is the natural open immersion, the lower horizontal arrow is given by $x \mapsto [x : 1]$, and μ is given by $(x, y) \mapsto xy^{-1}$. Excision yields the isomorphisms in the commutative diagram in $H_{\bullet}(S)$

$$\begin{array}{c|c} (\mathbb{A}^1 \times \mathbb{G}_m) / (\mathbb{G}_m \times \mathbb{G}_m) & \xrightarrow{\sim} & (\mathbb{A}^2 \smallsetminus \{0\}) / (\mathbb{G}_m \times \mathbb{A}^1) \longleftarrow \mathbb{A}^2 \smallsetminus \{0\} \\ & & \downarrow & & \downarrow \eta \\ & & \mathbb{A}^1 / \mathbb{G}_m & \xrightarrow{\sim} & \mathbb{P}^1 / \mathbb{A}^1 \longleftarrow \mathbb{P}^1 \end{array}$$

where $\tilde{\mu}$ is induced by μ , and the lower horizontal arrows are the morphisms of (2.1.0.a). To conclude, observe that the morphism $\tilde{\mu}$ factors as the upper horizontal composite in the following commutative diagram in $H_{\bullet}(S)$:

2.1.5. **Proposition.** Let $p: \mathbb{G}_m \to S$ be the projection, and $T = \mathbb{A}^1/\mathbb{G}_m = \mathrm{Th}_S(1)$. Consider the composite (see (1.5))

$$\pi \colon T \land (\mathbb{G}_m)_+ \xrightarrow{\mathrm{Th}(t \operatorname{id}_1)} T \land (\mathbb{G}_m)_+ \xrightarrow{\mathrm{id} \land p_+} T$$

Then the following square is homotopy cocartesian in $\operatorname{Spc}_{\bullet}(S)[\eta^{-1}]$

$$\begin{array}{ccc} T \land (\mathbb{G}_m)_+ & \stackrel{\pi}{\longrightarrow} T \\ & & \downarrow \\ & & \downarrow \\ & & T & \stackrel{\pi}{\longrightarrow} * \end{array}$$

Proof. From (2.1.3) we deduce a commutative diagram

where the left inner square is homotopy cocartesian. Applying (2.1.1) over the base \mathbb{G}_m and using the functor p_{\sharp} of (1.9), we have in $\mathrm{H}_{\bullet}(S)$

$$\operatorname{Th}(t \operatorname{id}_1) = \operatorname{Th}(t^{-1} \operatorname{id}_1) \colon T \land (\mathbb{G}_m)_+ \to T \land (\mathbb{G}_m)_+.$$

7

It thus follows from (2.1.4) that the upper composite in the diagram (2.1.5.a) is an isomorphism in $H_{\bullet}(S)[\eta^{-1}]$, hence the exterior square in (2.1.5.a) is homotopy cocartesian. We conclude that the right inner square is homotopy cocartesian (by [Hir03, Proposition 13.3.15, Remark 7.1.10]).

2.2. Global splitting.

2.2.1. **Definition.** Let $L \to S$ be a line bundle. Denote by L° the complement of the zero-section in L, and by $p: L^{\circ} \to S$ the projection. Then the graph $L^{\circ} \to L^{\circ} \times_{S} L$ of the open immersion $L^{\circ} \to L$ may be viewed as a nowhere vanishing section of the line bundle $p^{*}L$ over L° , which induces the *tautological trivialisation* $\tau: 1 \xrightarrow{\sim} p^{*}L$. We define a morphism in $H_{\bullet}(S)$ (recall that $T = Th_{S}(1)$, and see (1.5))

$$\pi_L \colon T \land (L^\circ)_+ = \operatorname{Th}_{L^\circ}(1) \xrightarrow{\operatorname{Th}(\tau)} \operatorname{Th}_{L^\circ}(p^*L) \xrightarrow{p} \operatorname{Th}_S(L).$$

2.2.2. **Example.** Assume that L = 1. Then the tautological trivialisation $\tau: 1 \to 1$ of the trivial line bundle over $L^{\circ} = \mathbb{G}_m$ is given by multiplication by the tautological section $t \in H^0(\mathbb{G}_m, \mathbb{G}_m)$, hence π_1 coincides with morphism π of (2.1.5).

2.2.3. Let $L \to S$ be a line bundle. If $f: R \to S$ is a scheme morphism, then the functor $f^*: \operatorname{Spc}_{\bullet}(S) \to \operatorname{Spc}_{\bullet}(R)$ maps π_L to π_{f^*L} .

2.2.4. If $\varphi \colon L \xrightarrow{\sim} M$ is an isomorphism of line bundles over S, then the following diagram commutes in $H_{\bullet}(S)$

2.2.5. **Proposition.** Let $L \to S$ be a line bundle. Then the following square is homotopy cocartesian in $\operatorname{Spc}_{\bullet}(S)[\eta^{-1}]$

$$\begin{array}{c} T \land (L^{\circ})_{+} \xrightarrow{\pi_{L}} \operatorname{Th}_{S}(L) \\ \stackrel{\operatorname{id} \land p_{+}}{\downarrow} & \downarrow \\ T \xrightarrow{} & * \end{array}$$

Proof. Let F be the homotopy colimit of the diagram $\operatorname{Th}_S(L) \xleftarrow{}{}{}^{T} \wedge (L^\circ)_+ \xrightarrow{\mathrm{id} \wedge p_+} T$. By (1.10) and (1.9) (and in view of (2.2.3)), the fact that $F \simeq *$ may be verified Zariskilocally on S. We may thus assume that L is trivial. By (2.2.4) we may further assume that L = 1, so that $L^\circ = \mathbb{G}_m$. Then, in view of (2.2.2) the result follows from (2.1.5)

The next statement was initially inspired by [Lev19, Proof of Theorem 4.1]:

2.2.6. Corollary. Let $L \to S$ be a line bundle. Then in the notation of (2.2.1), we have an isomorphism in $SH(S)[\eta^{-1}]$

$$(\Sigma^{\infty}_{+}p, \Sigma^{-2,-1}\Sigma^{\infty}\pi_L): \Sigma^{\infty}_{+}L^{\circ} \xrightarrow{\sim} \mathbf{1}_S \oplus \Sigma^{-2,-1} \operatorname{Th}_S(L).$$

Proof. The square induced in $\text{Spt}(S)[\eta^{-1}]$ by the square of (2.2.5) is homotopy cocartesian, hence also homotopy cartesian (see e.g. [Hov99, Remark 7.1.12]). This yields an isomorphism

$$(\Sigma^{\infty}(\mathrm{id} \wedge p_+), \Sigma^{\infty}\pi_L) \colon \Sigma^{\infty}(T \wedge (L^{\circ})_+) \xrightarrow{\sim} \Sigma^{\infty}T \oplus \mathrm{Th}_S(L),$$

from which the result follows by applying the functor $\Sigma^{-2,-1}$.

2.2.7. Corollary. Let $L \to S$ be a line bundle, and $V \to S$ a vector bundle.

(i) The natural map $\operatorname{Th}_{L^{\circ}}(V) \to \operatorname{Th}_{S}(V)$ extends to a natural isomorphism in $\operatorname{SH}(S)[\eta^{-1}]$

$$\operatorname{Th}_{L^{\circ}}(V) \simeq \operatorname{Th}_{S}(V) \oplus \Sigma^{-2,-1} \operatorname{Th}_{S}(V \oplus L).$$

(ii) Denote by $p: L^{\circ} \to S$ and $p_1, p_2: L^{\circ} \times_S L^{\circ} \to L^{\circ}$ the projections. Then

$$\operatorname{Th}_{L^{\circ}\times_{S}L^{\circ}}(V) \xrightarrow{p_{1}}{\longrightarrow} \operatorname{Th}_{L^{\circ}}(V) \xrightarrow{p} \operatorname{Th}_{S}(V)$$

is a split coequaliser diagram in $SH(S)[\eta^{-1}]$.

Proof. Statement (i) follows by applying the auto-equivalence $\Sigma^V \colon \operatorname{SH}(S)[\eta^{-1}] \to \operatorname{SH}(S)[\eta^{-1}]$ to the decomposition of (2.2.6), in view of (1.3.b).

Certainly in the diagram of (ii) we have $p \circ p_1 = p \circ p_2$. The isomorphism (i) yields a section $s: \operatorname{Th}_S(V) \to \operatorname{Th}_{L^\circ}(V)$ of p in $\operatorname{SH}(S)[\eta^{-1}]$. Then, in $\operatorname{SH}(S)[\eta^{-1}]$, the composite

$$t: \operatorname{Th}_{L^{\circ}}(V) = (\Sigma^{\infty}_{+}L^{\circ}) \wedge \operatorname{Th}_{S}(V) \xrightarrow{\operatorname{id} \wedge s} (\Sigma^{\infty}_{+}L^{\circ}) \wedge \operatorname{Th}_{L^{\circ}}(V) = \operatorname{Th}_{L^{\circ} \times_{S}L^{\circ}}(V)$$

is a section of

$$p_1: \operatorname{Th}_{L^{\circ} \times_S L^{\circ}}(V) = (\Sigma^{\infty}_+ L^{\circ}) \wedge \operatorname{Th}_{L^{\circ}}(V) \xrightarrow{\operatorname{id} \wedge p} (\Sigma^{\infty}_+ L^{\circ}) \wedge \operatorname{Th}_S(V) = \operatorname{Th}_{L^{\circ}}(V).$$

On the other hand, in the commutative diagram in $SH(S)[\eta^{-1}]$

$$\begin{aligned} \operatorname{Th}_{L^{\circ}}(V) &= (\Sigma^{\infty}_{+}L^{\circ}) \wedge \operatorname{Th}_{S}(V) \xrightarrow{\operatorname{id} \wedge s} (\Sigma^{\infty}_{+}L^{\circ}) \wedge \operatorname{Th}_{L^{\circ}}(V) = \operatorname{Th}_{L^{\circ} \times_{S}L^{\circ}}(V) \\ p & \downarrow p_{\wedge \operatorname{id}} & \downarrow p_{\wedge} \operatorname{id} & \downarrow p_{2} \\ \operatorname{Th}_{S}(V) &= \mathbf{1}_{S} \wedge \operatorname{Th}_{S}(V) \xrightarrow{\operatorname{id} \wedge s} \mathbf{1}_{S} \wedge \operatorname{Th}_{L^{\circ}}(V) = \operatorname{Th}_{L^{\circ}}(V) \end{aligned}$$

the upper composite is t, while the lower one is s. Therefore $p_2 \circ t = s \circ p$ as endomorphisms of $\operatorname{Th}_{L^{\circ}}(V)$ in $\operatorname{SH}(S)[\eta^{-1}]$, proving (ii).

2.2.8. Corollary. Let $L \to S$ be a line bundle, and denote by $p: L^{\circ} \to S$ the projection. Then the functor $p^*: \operatorname{SH}(S)[\eta^{-1}] \to \operatorname{SH}(L^{\circ})[\eta^{-1}]$ is faithful and conservative.

Proof. By the smooth projection formula and (2.2.6), the composite $p_{\sharp} \circ p^* \colon \operatorname{SH}(S)[\eta^{-1}] \to \operatorname{SH}(S)[\eta^{-1}]$ decomposes as

$$p_{\sharp} \circ p^* = \operatorname{id} \wedge (\Sigma^{\infty}_{+} L^{\circ}) = \operatorname{id} \wedge (\mathbf{1}_S \oplus \Sigma^{-2,-1} \operatorname{Th}_S(L)) = \operatorname{id} \oplus (\Sigma^{-2,-1} \circ \Sigma^L),$$

which is faithful, hence so is p^* . The above formula also shows that $p_{\sharp} \circ p^*$ reflects zero-objects, hence so does p^* . Since p^* is triangulated, it is conservative.

2.2.9. **Remark.** The results of this section on line bundles will be generalised to odd rank vector bundles in §4.2.

9

3. Applications to twisted cohomology

3.1. Cohomology theories represented by ring spectra.

3.1.1. Let $A \in \operatorname{Spt}(S)$ be a motivic spectrum. For a pointed motivic space \mathcal{X} we write

$$A^{p,q}(\mathcal{X}) = \operatorname{Hom}_{\operatorname{SH}(S)}(\Sigma^{\infty}\mathcal{X}, \Sigma^{p,q}A),$$

and $A^{*,*}(\mathcal{X}) = \bigoplus_{p,q \in \mathbb{Z}} A^{p,q}(\mathcal{X})$. When X is a smooth S-scheme, we will write $A^{*,*}(X)$ instead of $A^{*,*}(X_+)$. If $E \to X$ is a vector bundle of constant rank r, we write

$$A^{p,q}(X;E) = A^{p+2r,q+r}(\operatorname{Th}_X(E)),$$

and extend this notation to arbitrary vector bundles in an obvious way. A morphism $f: \mathcal{Y} \to \mathcal{X}$ of pointed motivic spaces (resp. of smooth *S*-schemes) induces a pullback $f^*: A^{*,*}(\mathcal{X}) \to A^{*,*}(\mathcal{Y})$.

3.1.2. A commutative ring spectrum will mean a commutative monoid in $(SH(S), \land, \mathbf{1}_S)$. When $A \in SH(S)$ is a commutative ring spectrum and $X \in Sm_S$, then $A^{*,*}(X)$ is naturally a ring, and $A^{*,*}(X; E)$ an $A^{*,*}(X)$ -module. When $u \in H^0(X, \mathbb{G}_m)$, we will write $\langle u \rangle \in A^{0,0}(X)$ instead of $\langle u \rangle^*(1)$ (see (1.6)).

3.1.3. If A is an η -periodic motivic spectrum, for any pointed motivic space \mathcal{X} , we have natural isomorphisms for $p, q \in \mathbb{Z}$

$$A^{p,q}(\mathcal{X}) = \operatorname{Hom}_{\operatorname{SH}(S)[\eta^{-1}]}(\Sigma^{\infty}\mathcal{X}, \Sigma^{p,q}A).$$

3.1.4. **Proposition.** Let A be an η -periodic motivic spectrum. Let $X \in \text{Sm}_S$. Let $L \to X$ be a line bundle, and $V \to X$ a vector bundle.

(i) Denoting by $p: L^{\circ} \to X$ the projection, we have a split short exact sequence

$$0 \to A^{*,*}(X;V) \xrightarrow{p^*} A^{*,*}(L^{\circ};V) \to A^{*,*}(X;V \oplus L) \to 0.$$

(ii) Denoting by $p_1, p_2: L^{\circ} \times_X L^{\circ} \to L^{\circ}$ the projections, we have an exact sequence

$$0 \to A^{*,*}(X;V) \xrightarrow{p^*} A^{*,*}(L^\circ;V) \xrightarrow{p_1^* - p_2^*} A^{*,*}(L^\circ \times_X L^\circ;V).$$

Proof. This follows by applying (2.2.7) over the base X to the image of A under the pullback $\operatorname{Spt}(S) \to \operatorname{Spt}(X)$.

3.2. SL- and SL^c -orientations.

3.2.1. **Definition.** An SL-oriented vector bundle over a scheme X is a pair (E, δ) , where $E \to X$ is a vector bundle and $\delta: 1 \xrightarrow{\sim} \det E$ is an isomorphism of line bundles. We will also say that δ is an SL-orientation of the vector bundle $E \to X$. An isomorphism of SL-oriented vector bundles $(E, \delta) \xrightarrow{\sim} (F, \epsilon)$ is an isomorphism of vector bundles $\varphi: E \xrightarrow{\sim} F$ such that $(\det \varphi) \circ \delta = \epsilon$.

3.2.2. **Definition** (See [PW18, §3]). An SL^c-oriented vector bundle over a scheme X is a triple (E, L, λ) , where $E \to X$ is a vector bundle and $L \to X$ a line bundle, and $\lambda: L^{\otimes 2} \xrightarrow{\sim} \det E$ is an isomorphism. We will also say that (L, λ) is an SL^c-orientation of the vector bundle $E \to X$. An isomorphism of SL^c-oriented vector bundles $(E, L, \lambda) \xrightarrow{\sim} (F, M, \mu)$ is an isomorphism of vector bundles $\varphi: E \xrightarrow{\sim} F$ and an isomorphism of line bundles $\psi: L \xrightarrow{\sim} M$ such that $(\det \varphi) \circ \lambda = \mu \circ \psi^{\otimes 2}$.

3.2.3. Observe that each SL-orientation δ of a vector bundle E induces an SL^c-orientation (L, λ) of E, where L = 1 and λ is the composite $1^{\otimes 2} \simeq 1 \xrightarrow{\delta} \det E$.

3.2.4. Let (E, L, λ) be an SL^c-oriented vector bundle, and assume that the line bundle L is trivial. Then every trivialisation $\alpha \colon 1 \xrightarrow{\sim} L$ induces an SL-orientation of E given by

$$\delta_{\alpha} \colon 1 \simeq 1^{\otimes 2} \xrightarrow{\alpha^{\otimes 2}} L^{\otimes 2} \xrightarrow{\lambda} \det E.$$

Observe that the SL^c-oriented vector bundle induced (in the sense of (3.2.3)) by δ_{α} is isomorphic to (E, L, λ) .

3.2.5. Consider a commutative ring spectrum $A \in SH(S)$. By a SL-, resp. SL^c -, orientation of A, we will mean a normalised orientation in the sense of [Ana20, Definition 3.3]. Such data consists in Thom classes $th_{(E,\delta)} \in A^{*,*}(X; E)$ for each SL-oriented vector bundle (E, δ) over $X \in Sm_S$, resp. $th_{(E,L,\lambda)} \in A^{*,*}(X; E)$ for each SL^c-oriented vector bundle (E, L, λ) over $X \in Sm_S$, satisfying a series of axioms.

3.2.6. Let $A \in SH(S)$ be a commutative ring spectrum. Then each SL^c -orientation of A induces an SL-orientation of A, by letting the Thom class of an SL-oriented vector bundle be the Thom class of the induced SL^c -oriented vector bundle, in the sense of (3.2.3).

3.2.7. Lemma. Let $A \in SH(S)$ be an SL-oriented commutative ring spectrum. Let (E, L, λ) be an SL^c-oriented vector bundle over $X \in Sm_S$, and assume that the line bundle L is trivial. Then, in the notation of (3.2.4), the Thom class $th_{(E,\delta_{\alpha})} \in A^{*,*}(X; E)$ does not depend on the choice of the trivialisation α of L.

Proof. If $\alpha \colon 1 \xrightarrow{\sim} L$ is a trivialisation, then every trivialisation is of the form $u\alpha$ for some $u \in H^0(X, \mathbb{G}_m)$. In the notation of (3.2.4) we then have $\delta_{u\alpha} = u^2 \delta_{\alpha}$. By [Ana20, Lemma 7.3], we have

$$\operatorname{th}_{(E,\delta_{u\alpha})} = \operatorname{th}_{(E,u^2\delta_{\alpha})} = \langle u^2 \rangle \operatorname{th}_{(E,\delta_{\alpha})} \in A^{*,*}(X;E).$$

Since $\langle u^2 \rangle = 1 \in A^{0,0}(X)$ by (2.1.1) (or [Ana20, Lemma 6.2]), the statement follows. \Box

3.2.8. **Proposition.** Let $A \in SH(S)$ be an η -periodic commutative ring spectrum. Then every SL-orientation of A is induced (in the sense of (3.2.6)) by a unique SL^corientation.

Proof. We assume given an SL-orientation of A. Let (E, L, λ) be an SL^c-oriented vector bundle. Denoting by $p: L^{\circ} \to X$ the projection, the line bundle p^*L over L° admits a tautological trivialisation $\tau: 1 \xrightarrow{\sim} p^*L$. In view of (3.2.4), this yields an SL-orientation δ_{τ} of p^*E .

First assume given an SL^c-orientation of A compatible with its SL-orientation. As observed in (3.2.4), the SL^c-oriented vector bundle $(p^*E, p^*L, p^*\lambda)$ is isomorphic to the one induced by the SL-oriented vector bundle (p^*E, δ_{τ}) . Thus we must have

$$p^* \operatorname{th}_{(E,L,\lambda)} = \operatorname{th}_{(p^*E,p^*L,p^*\lambda)} = \operatorname{th}_{(p^*E,\delta_\tau)} \in A^{*,*}(L^\circ; p^*E).$$

Since $p^* \colon A^{*,*}(X; E) \to A^{*,*}(L^{\circ}; p^*E)$ is injective by (3.1.4), we obtain the uniqueness part of the statement.

We now construct an SL^c-orientation of A from its SL-orientation. In the situation considered at the beginning of the proof, let $p_1, p_2: L^{\circ} \times_X L^{\circ} \to L^{\circ}$ be the projections, and set $q = p \circ p_1 = p \circ p_2$. The tautological trivialisation τ of p^*L yields two trivialisations $p_1^*\tau$ and $p_2^*\tau$ of q^*L , and thus two SL-orientations $\alpha_1 = \delta_{p_1^*\tau}$ and $\alpha_2 = \delta_{p_2^*\tau}$ of E. However it follows from (3.2.7) that their Thom classes coincide, so that (observe that $\alpha_i = p_i^*(\delta_{\tau})$ for i = 1, 2)

$$p_1^* \operatorname{th}_{(p^*E,\delta_{\tau})} = \operatorname{th}_{(E,\alpha_1)} = \operatorname{th}_{(E,\alpha_2)} = p_2^* \operatorname{th}_{(p^*E,\delta_{\tau})} \in A^{*,*}(L^{\circ} \times_X L^{\circ}; q^*E).$$

Therefore it follows from (3.1.4.ii) that the element $\operatorname{th}_{(p^*E,\delta_\tau)} \in A^{*,*}(L^\circ; E)$ is the image of a unique element $\theta_{(E,L,\lambda)} \in A^{*,*}(X; E)$.

From the fact that $(E, \delta) \mapsto \operatorname{th}_{(E,\delta)}$ defines an SL-orientation of A, we deduce at once that $(E, L, \lambda) \mapsto \theta_{(E,L,\lambda)}$ defines an SL^c-orientation of A: indeed, each axiom of [Ana20, Definition 3.3] can be verified after pulling back along $p: L^{\circ} \to X$, since $p^*: A^{*,*}(X; E) \to A^{*,*}(L^{\circ}; p^*E)$ is injective by (3.1.4).

To conclude the proof, it remains to show the SL^c-orientation $(E, L, \lambda) \mapsto \theta_{(E,L,\lambda)}$ induces the original SL-orientation of A. So let us assume that the SL^c-oriented vector bundle (E, L, λ) is induced by an SL-oriented vector bundle (E, δ) , in the sense of (3.2.3). In particular L = 1. Then the tautological trivialisation $\tau \colon 1 \xrightarrow{\sim} p^*L$ and the trivialisation $1 = p^*1 = p^*L$ yield two SL-orientations of p^*E . Their Thom classes in $A^{*,*}(L^\circ; p^*E)$ coincide by (3.2.7), and they are respectively $p^*\theta_{(E,L,\lambda)}$ and $p^* \operatorname{th}_{(E,\delta)}$. Since $p^* \colon A^{*,*}(X; E) \to A^{*,*}(L^\circ; p^*E)$ is injective by (3.1.4), we have $\theta_{(E,L,\lambda)} = \operatorname{th}_{(E,\delta)} \in$ $A^{*,*}(X; E)$, as required.

3.2.9. **Remark.** Ananyevskiy constructed "Thom isomorphisms" associated with SL^c bundles in [Ana20, §4] when A is an arbitrary SL-oriented theory, but as explained in [Ana20, Remark 4.4] it is not clear whether this yields an SL^c -orientation, the problem being the multiplicativity axiom. When A is η -periodic, our construction leads to the same Thom isomorphisms for SL^c -bundles (in fact the proof of (3.2.8) shows that there is at most one way to construct such functorial isomorphisms compatibly with the SLorientation). Thus the Thom isomorphisms constructed by Ananyevskiy do give rise to an SL^c -orientation when A is η -periodic.

3.3. Twisting by doubles and squares of line bundles.

3.3.1. **Proposition.** Let L be a line bundle over S.

- (i) There exist an isomorphism $\Sigma^{4,2}\mathbf{1}_S \simeq \mathrm{Th}_S(L^{\oplus 2})$ in $\mathrm{SH}(S)[\eta^{-1}]$.
- (ii) If $s_1, s_2 \in \mathbb{Z}$ are of the same parity, there exist an isomorphism

$$\operatorname{Th}_{S}(L^{\otimes s_{1}}) \simeq \operatorname{Th}_{S}(L^{\otimes s_{2}}) \text{ in } \operatorname{SH}(S)[\eta^{-1}].$$

Proof. Let us first assume that the line bundle $L \to S$ admits a trivialisation $\alpha \colon 1 \xrightarrow{\sim} L$. Then we have an isomorphism in SH(S) (see (1.5))

(3.3.1.a)
$$\operatorname{Th}(\alpha^{\oplus 2}) \colon \operatorname{Th}_{S}(1^{\oplus 2}) \xrightarrow{\sim} \operatorname{Th}_{S}(L^{\oplus 2}).$$

Every trivialisation of L is of the form $u\alpha$ with $u \in H^0(S, \mathbb{G}_m)$. As automorphisms of $\operatorname{Th}_S(1^{\oplus 2})$ in $\operatorname{SH}(S)$ we have,

$$\operatorname{Th}((u \operatorname{id}_1)^{\oplus 2}) = \operatorname{Th}((u^2 \operatorname{id}_1) \oplus \operatorname{id}_1) \circ \operatorname{Th}((u^{-1} \operatorname{id}_1) \oplus (u \operatorname{id}_1)) = \operatorname{Th}((u^2 \operatorname{id}_1) \oplus \operatorname{id}_1)$$

because $\operatorname{Th}((u^{-1}\operatorname{id}_1) \oplus (u\operatorname{id}_1))$ is the identity of $\operatorname{Th}_S(1^{\oplus 2})$, being given by a product transvections (see e.g. [Ana16a, Lemma 1]). Now by (2.1.1), under the identification $\operatorname{Th}_S(1^{\oplus 2}) = \operatorname{Th}_S(1) \wedge \operatorname{Th}_S(1)$ (see (1.3.b)), we have

$$\operatorname{Th}((u^2 \operatorname{id}_1) \oplus \operatorname{id}_1) = \operatorname{Th}(u^2 \operatorname{id}_1) \wedge \operatorname{id}_{\operatorname{Th}_S(1)} = \operatorname{id}_{\operatorname{Th}_S(1)} \wedge \operatorname{id}_{\operatorname{Th}_S(1)} = \operatorname{id}_{\operatorname{Th}_S(1\oplus 2)}.$$

Therefore $\operatorname{Th}((u \operatorname{id}_1)^{\oplus 2})$ is the identity of $\operatorname{Th}_S(1^{\oplus 2})$ in $\operatorname{SH}(S)$, hence

$$\operatorname{Th}((u\alpha)^{\oplus 2}) = \operatorname{Th}(\alpha^{\oplus 2}) \circ \operatorname{Th}((u\operatorname{id}_1)^{\oplus 2}) = \operatorname{Th}(\alpha^{\oplus 2}),$$

so that the isomorphism $\operatorname{Th}(\alpha^{\oplus 2})$ in $\operatorname{SH}(S)$ considered in (3.3.1.a) is independent of the choice of the trivialisation α .

Next let us consider the case (ii). If $\alpha \colon 1 \xrightarrow{\sim} L$ is a trivialisation, we have an isomorphism in SH(S)

(3.3.1.b)
$$\operatorname{Th}(\operatorname{id}_{L^{\otimes s_1}} \otimes \alpha^{\otimes s_2 - s_1}) \colon \operatorname{Th}_S(L^{\otimes s_1}) \xrightarrow{\sim} \operatorname{Th}_S(L^{\otimes s_2})$$

(Here and below, for $r \in \mathbb{N}$, the notation $\alpha^{\otimes -r}$ refers to the morphism $((\alpha^{\vee})^{-1})^{\otimes r}$.) Now for $u \in H^0(S, \mathbb{G}_m)$, the composite in SH(S)

$$\operatorname{Th}_{S}(1) \xrightarrow{\operatorname{Th}(\alpha^{\otimes s_{1}})} \operatorname{Th}_{S}(L^{\otimes s_{1}}) \xrightarrow{\operatorname{Th}(\operatorname{id}_{L^{\otimes s_{1}}} \otimes (u\alpha)^{\otimes s_{2}-s_{1}})} \operatorname{Th}_{S}(L^{\otimes s_{2}}) \xrightarrow{\operatorname{Th}(\alpha^{\otimes s_{2}})^{-1}} \operatorname{Th}_{S}(1)$$

coincides with $\operatorname{Th}(u^{s_2-s_1}\operatorname{id}_1)$, which is the identity by (2.1.1) (recall that $s_2 - s_1$ is even), and in particular does not depend on $u \in H^0(S, \mathbb{G}_m)$. Since the left and right arrows in the above composite are isomorphisms, we deduce that the middle arrow does not depend on $u \in H^0(S, \mathbb{G}_m)$, which shows as above that the isomorphism (3.3.1.b) is independent of the choice of the trivialisation α .

Let us come back to the general case, where $L \to S$ is a possibly nontrivial line bundle. Let $p: L^{\circ} \to S$ be the projection, and consider the tautological trivialisation τ of the line bundle p^*L over L° . Let us consider the isomorphism $\varphi \colon \operatorname{Th}_{L^{\circ}}(B) \xrightarrow{\sim} \operatorname{Th}_{L^{\circ}}(C)$ in $\operatorname{SH}(S)[\eta^{-1}]$, where

•
$$B = 1^{\oplus 2}, C = L^{\oplus 2}, \varphi = \operatorname{Th}(\tau^{\oplus 2})$$
 in case (i).

• $B = L^{\otimes s_1}, C = L^{\otimes s_2}, \varphi = \text{Th}(\text{id}_{L^{\otimes s_1}} \otimes \alpha^{\otimes s_2 - s_1})$ in case (ii),

Let $p_1, p_2: L^{\circ} \times_S L^{\circ} \to L^{\circ}$ be the projections, and set $q = p \circ p_1 = p \circ p_2$. For $i \in \{1, 2\}$, the isomorphism $p_i^* \varphi: \operatorname{Th}_{L^{\circ} \times_S L^{\circ}}(B) \to \operatorname{Th}_{L^{\circ} \times_S L^{\circ}}(C)$ in $\operatorname{SH}(S)[\eta^{-1}]$ is induced by the trivialisation $p_i^* \tau$ of the line bundle q^*L over $L^{\circ} \times_S L^{\circ}$, hence does not depend on i, by the special case considered at the beginning of the proof. Thus, by (2.2.7.ii) there exists a unique morphism f fitting into the commutative diagram in $\operatorname{SH}(S)[\eta^{-1}]$

$$\begin{array}{c|c} \operatorname{Th}_{L^{\circ}\times_{S}L^{\circ}}(B) \xrightarrow{p_{1}} \operatorname{Th}_{L^{\circ}}(B) \xrightarrow{p} \operatorname{Th}_{S}(B) \\ p_{1}^{*}\varphi = p_{2}^{*}\varphi & & \downarrow^{\varphi} & & \downarrow^{f} \\ \operatorname{Th}_{L^{\circ}\times_{S}L^{\circ}}(C) \xrightarrow{p_{1}} \operatorname{Th}_{L^{\circ}}(C) \xrightarrow{p} \operatorname{Th}_{S}(C) \end{array}$$

as well as a unique morphism g into the commutative diagram in $SH(S)[\eta^{-1}]$

$$\begin{array}{ccc} \operatorname{Th}_{L^{\circ}\times_{S}L^{\circ}}(C) \xrightarrow{p_{1}} \operatorname{Th}_{L^{\circ}}(C) \xrightarrow{p} \operatorname{Th}_{S}(C) \\ p_{1}^{*}\varphi^{-1} = p_{2}^{*}\varphi^{-1} & & \downarrow \varphi^{-1} & & \downarrow g \\ \operatorname{Th}_{L^{\circ}\times_{S}L^{\circ}}(B) \xrightarrow{p_{1}} \operatorname{Th}_{L^{\circ}}(B) \xrightarrow{p} \operatorname{Th}_{S}(B) \end{array}$$

As $p: \operatorname{Th}_{L^{\circ}}(B) \to \operatorname{Th}_{S}(B)$ and $p: \operatorname{Th}_{L^{\circ}}(C) \to \operatorname{Th}_{S}(C)$ are epimorphisms in $\operatorname{SH}(S)[\eta^{-1}]$ (see (2.2.7)), it follows that f and g are mutually inverse isomorphisms in $\operatorname{SH}(S)[\eta^{-1}]$.

3.3.2. Remark. Proposition (3.3.1) will be improved in (4.3.2).

4. Nowhere vanishing sections of odd rank bundles

4.1. **Projective bundles.** The results of this section are slight generalisations of those of [Ana16a, §4].

4.1.1. Let us consider the linear embeddings $i_k \colon \mathbb{P}^k \to \mathbb{P}^{k+1}$ given by the vanishing of the last coordinate. Denote by $\iota_k \colon S \to \mathbb{P}^k$ the S-point given by the composite $S = \mathbb{P}^0 \xrightarrow{i_0} \mathbb{P}^1 \xrightarrow{i_1} \cdots \xrightarrow{i_{k-1}} \mathbb{P}^k$.

4.1.2. Assume given a collection $\mathcal{D} = (d_1, \ldots, d_r) \in \mathbb{Z}^r$ for some $r \in \mathbb{N}$. We will denote by $\mathcal{O}(\mathcal{D})$ the vector bundle $\mathcal{O}(d_1) \oplus \cdots \oplus \mathcal{O}(d_r)$ over \mathbb{P}^k , for each $k \in \mathbb{N}$. When k = 0, we have a canonical isomorphism $\mathcal{O}(\mathcal{D}) \simeq 1^{\oplus r}$ over $\mathbb{P}^0 = S$. This yields, for any $k \in \mathbb{N}$, a canonical map in $\operatorname{Spt}(S)$

(4.1.2.a)
$$\Sigma^{2r,r} \mathbf{1}_S = \operatorname{Th}_S(1^{\oplus r}) \simeq \operatorname{Th}_{\mathbb{P}^0}(\mathcal{O}(\mathcal{D})) \xrightarrow{\iota_k} \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})).$$

4.1.3. **Proposition.** Let $k, r \in \mathbb{N}$, and $d_1, \ldots, d_r \in \mathbb{Z}$. Set $\mathcal{D} = (d_1, \ldots, d_r)$, and let $d = d_1 + \cdots + d_r$. We use the notation $\mathcal{O}(\mathcal{D})$ described in (4.1.2).

- (i) If k and d are odd, then $\operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})) = 0$ in $\operatorname{SH}(S)[\eta^{-1}]$.
- (ii) If k and d are even, then (4.1.2.a) induces an isomorphism $\Sigma^{2r,r} \mathbf{1}_S \simeq \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D}))$ in $\operatorname{SH}(S)[\eta^{-1}]$.
- (iii) If k is even and d is odd, then $\operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})) \simeq \Sigma^{2(k+r),k+r} \mathbf{1}_S$.
- (iv) If k is odd and d is even, then $\operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})) \simeq \Sigma^{2(k+r),k+r} \mathbf{1}_S \oplus \Sigma^{2r,r} \mathbf{1}_S$.

Proof. Let us first prove (i). Assume that k and d are odd. Consider a linear embedding $\mathbb{P}^1 \to \mathbb{P}^k$. Its normal bundle is $\mathcal{O}(1)^{\oplus k-1}$, and its open complement is a vector bundle over \mathbb{P}^{k-2} . The corresponding zero-section $\mathbb{P}^{k-2} \to \mathbb{P}^k \setminus \mathbb{P}^1$ induces an isomorphism in $\mathrm{SH}(S)$, and is the restriction of a linear embedding $\mathbb{P}^{k-2} \to \mathbb{P}^k$. Thus (1.4) yields a distinguished triangle in $\mathrm{SH}(S)$

$$\mathrm{Th}_{\mathbb{P}^{k-2}}(\mathcal{O}(\mathcal{D})) \to \mathrm{Th}_{\mathbb{P}^{k}}(\mathcal{O}(\mathcal{D})) \to \mathrm{Th}_{\mathbb{P}^{1}}(\mathcal{O}(\mathcal{D}) \oplus \mathcal{O}(1)^{\oplus k-1}) \to \Sigma^{1,0} \mathrm{Th}_{\mathbb{P}^{k-2}}(\mathcal{O}(\mathcal{D})).$$

Using induction on the odd integer k, we are reduced to assuming that k = 1. Now by (3.3.1.ii) we have in $SH(S)[\eta^{-1}]$

$$\operatorname{Th}_{\mathbb{P}^1}(\mathcal{O}(\mathcal{D})) \simeq \Sigma^{2s,s}(\operatorname{Th}_{\mathbb{P}^1}(\mathcal{O}(-1))^{\wedge r-s}),$$

where s is the number of indices $i \in \{1, ..., r\}$ such that d_i is even. Since d is odd, so is r - s, and using (3.3.1.i) we deduce that

$$\operatorname{Th}_{\mathbb{P}^1}(\mathcal{O}(\mathcal{D})) \simeq \Sigma^{2(r-1),r-1} \operatorname{Th}_{\mathbb{P}^1}(\mathcal{O}(-1)) \quad \text{in } \operatorname{SH}(S)[\eta^{-1}].$$

But $\operatorname{Th}_{\mathbb{P}^1}(\mathcal{O}(-1))$ vanishes in $\operatorname{SH}(S)[\eta^{-1}]$, because of the distinguished triangle (see (1.3.a))

$$\Sigma^{\infty}_{+}\mathcal{O}(-1)^{\circ} \to \Sigma^{\infty}_{+}\mathbb{P}^{1} \to \operatorname{Th}_{\mathbb{P}^{1}}(\mathcal{O}(-1)) \to \Sigma^{1,0}\Sigma^{\infty}_{+}\mathcal{O}(-1)^{\circ}$$

and the definition of the map η (recall that $\mathcal{O}(-1)^{\circ} = \mathbb{A}^2 \setminus \{0\}$). We have proved (i).

Let us come back to the situation when k and d are arbitrary. Consider a linear embedding $\mathbb{P}^{k-1} \to \mathbb{P}^k$ avoiding the S-point $\iota_k \colon S \to \mathbb{P}^k$ (we write $\mathbb{P}^{-1} = \emptyset$). It is a closed immersion defined by the vanishing of a regular section of $\mathcal{O}(1)$. Its open complement is isomorphic to \mathbb{A}^k , and the morphism $j_k \colon S \to \mathbb{A}^k$ induced by ι_k induces an isomorphism in $\mathrm{SH}(S)$. The canonical trivialisation of $\mathcal{O}(\mathcal{D})$ over $\mathbb{P}^0 = S$ is the restriction along j_k of a trivialisation of $\mathcal{O}(\mathcal{D})|_{\mathbb{A}^k}$ (induced by the trivialisation of $\mathcal{O}(1)|_{\mathbb{A}^k}$ corresponding to the regular section of $\mathcal{O}(1)$ mentioned above). It follows that the map $\mathrm{Th}_{\mathbb{P}^0}(\mathcal{O}(\mathcal{D})) \to \mathrm{Th}_{\mathbb{A}^k}(\mathcal{O}(\mathcal{D}))$ induced by j_k induces an isomorphism in $\mathrm{SH}(S)$. Thus (1.4) yields a distinguished triangle in $\mathrm{SH}(S)$

$$\Sigma^{2r,r}\mathbf{1}_S \xrightarrow{(4.1.2.a)} \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})) \to \operatorname{Th}_{\mathbb{P}^{k-1}}(\mathcal{O}(\mathcal{D}) \oplus \mathcal{O}(1)) \to \Sigma^{2r+1,r}\mathbf{1}_S,$$

so that (ii) follows from (i).

Consider now a linear embedding $s: S = \mathbb{P}^0 \to \mathbb{P}^k$ avoiding $i_{k-1}(\mathbb{P}^{k-1})$. Its open complement U is a line bundle over \mathbb{P}^{k-1} . The corresponding zero-section $\mathbb{P}^{k-1} \to U$ induces an isomorphism in $\mathrm{SH}(S)$, and is the restriction of the linear embedding $i_{k-1}: \mathbb{P}^{k-1} \to \mathbb{P}^k$. Since the vector bundle $s^*\mathcal{O}(\mathcal{D})$ and the normal bundle $s^*\mathcal{O}(1)^{\oplus k}$ to s are both trivial, we have by (1.4) a distinguished triangle in $\mathrm{SH}(S)$

(4.1.3.a)
$$\operatorname{Th}_{\mathbb{P}^{k-1}}(\mathcal{O}(\mathcal{D})) \to \operatorname{Th}_{\mathbb{P}^{k}}(\mathcal{O}(\mathcal{D})) \to \Sigma^{2(k+r),k+r}\mathbf{1}_{S} \to \Sigma^{1,0} \operatorname{Th}_{\mathbb{P}^{k-1}}(\mathcal{O}(\mathcal{D})).$$

Therefore (iii) follows from (i).

Finally, assume that k is odd and d is even. It follows from (ii) that the composite $\operatorname{Th}_{\mathbb{P}^{k-1}}(\mathcal{O}(\mathcal{D})) \to \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D})) \to \operatorname{Th}_{\mathbb{P}^{k+1}}(\mathcal{O}(\mathcal{D}))$ is an isomorphism in $\operatorname{SH}(S)[\eta^{-1}]$, hence $\operatorname{Th}_{\mathbb{P}^{k-1}}(\mathcal{O}(\mathcal{D})) \to \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(\mathcal{D}))$ admits a retraction, giving a splitting of the triangle (4.1.3.a). In view of (ii), this proves (iv).

4.1.4. Corollary. If $k \in \mathbb{N}$ is even, the structural morphism $\mathbb{P}^k \to S$ induces an isomorphism $\Sigma^{\infty}_{+}\mathbb{P}^k \xrightarrow{\sim} \mathbf{1}_S$ in $\mathrm{SH}(S)[\eta^{-1}]$.

Proof. The structural morphism is retraction of ι_k , so the corollary follows from (4.1.3.ii) applied with r = 0.

4.1.5. **Proposition.** Let E, V_1, \ldots, V_n be vector bundles of constant rank over S, and $d_1, \ldots, d_n \in \mathbb{Z}$. Assume that rank E is even and that $d_1 \operatorname{rank} V_1 + \cdots + d_n \operatorname{rank} V_n$ is odd. Then

 $\operatorname{Th}_{\mathbb{P}(E)}((\mathcal{O}(d_1)\otimes q^*V_1)\oplus\cdots\oplus(\mathcal{O}(d_n)\otimes q^*V_n))=0\in \operatorname{SH}(S)[\eta^{-1}],$

where $q: \mathbb{P}(E) \to S$ is the projective bundle.

Proof. By (1.10) and (1.9), this may be verified Zariski-locally on S, so we may assume that E, V_1, \ldots, V_n are all trivial. Then the statement follows from (4.1.3.i).

4.1.6. **Proposition.** Let E, V be vector bundles over S. Assume that E has constant odd rank. Then $\operatorname{Th}_{\mathbb{P}(E)}(V) \to \operatorname{Th}_{S}(V)$ is an isomorphism in $\operatorname{SH}(S)[\eta^{-1}]$. In particular $\Sigma^{\infty}_{+}\mathbb{P}(E) \xrightarrow{\sim} \mathbf{1}_{S}$ in $\operatorname{SH}(S)[\eta^{-1}]$.

Proof. By (1.10) and (1.9), this may be verified Zariski-locally on S, so we may assume that E and V are both trivial. Then the statement follows after suspending (4.1.4). \Box

4.2. Odd rank vector bundles.

4.2.1. Let $E \to S$ be a vector bundle. The composite $\mathcal{O}(-1) \subset E \times_S \mathbb{P}(E) \to E$ restricts to an isomorphism $\mathcal{O}(-1)^{\circ} \xrightarrow{\sim} E^{\circ}$, which is \mathbb{G}_m -equivariant. We thus obtain a commutative diagram in $\operatorname{Spt}(S)$

which induces a morphism of the homotopy cofibers of the horizontal morphisms:

(4.2.1.b) $\operatorname{Th}_{\mathbb{P}(E)}(\mathcal{O}(-1)) \to \operatorname{Th}_{S}(E) \in \operatorname{SH}(S).$

4.2.2. Lemma. Let $E \to S$ be a vector bundle of constant odd rank. Then (4.2.1.b) is an isomorphism $\operatorname{Th}_{\mathbb{P}(E)}(\mathcal{O}(-1)) \xrightarrow{\sim} \operatorname{Th}_{S}(E)$ in $\operatorname{SH}(S)[\eta^{-1}]$.

Proof. It follows from (4.1.6) that both vertical arrows in the diagram (4.2.1.a) are isomorphisms in $SH(S)[\eta^{-1}]$, hence so the induced map on homotopy cofibers.

4.2.3. **Proposition.** Let $E \to S$ be a vector bundle of constant odd rank. Then the projection $E^{\circ} \to S$ admits a section in $SH(S)[\eta^{-1}]$, inducing a decomposition in $SH(S)[\eta^{-1}]$

$$\Sigma^{\infty}_{+}E^{\circ} \simeq \mathbf{1}_{S} \oplus \Sigma^{-2,-1} \operatorname{Th}_{S}(E).$$

Proof. By (2.2.6), we have a splitting in $SH(S)[\eta^{-1}]$

$$\Sigma^{\infty}_{+}\mathcal{O}(-1)^{\circ} \simeq \Sigma^{\infty}_{+}\mathbb{P}(E) \oplus \Sigma^{-2,-1} \operatorname{Th}_{\mathbb{P}(E)}(\mathcal{O}(-1))$$

and the statement follows from (4.2.1), (4.1.6) and (4.2.2).

4.2.4. **Remark.** One may deduce that (2.2.7) and (2.2.8) remain valid when L is an odd rank vector bundle instead of a line bundle, using exactly the same arguments, but substituting (4.2.3) for (2.2.6).

We deduce the following splitting principle:

4.2.5. Corollary. Let $X \in \text{Sm}_S$, and $E \to X$ be a vector bundle of constant odd rank. Then there exists a morphism $f: Y \to X$ in Sm_S whose image in $\text{SH}(S)[\eta^{-1}]$ admits a section, and a vector bundle $F \to Y$ such that $f^*E \simeq F \oplus 1$. *Proof.* Applying the functor $\operatorname{SH}(X)[\eta^{-1}] \to \operatorname{SH}(S)[\eta^{-1}]$ of (1.9) we may assume that X = S. Let us denote by $p: (E^{\vee})^{\circ} \to S$ the projection. Then p^*E^{\vee} admits a nowhere vanishing section s. Its dual $s^{\vee}: p^*E \to 1$ is surjective. Letting $Q = \ker s^{\vee}$, we have an exact sequence of vector bundles over $(E^{\vee})^{\circ}$

$$(4.2.5.a) 0 \to Q \to p^*E \to 1 \to 0.$$

Then we may find an affine bundle $g: Y \to (E^{\vee})^{\circ}$ along which the pullback of the sequence (4.2.5.a) splits (we may take for Y the scheme parametrising the sections of $p^*E^{\vee} \to Q^{\vee}$, see e.g. [Rio10, p.243]). Then $\Sigma^{\infty}_+g: \Sigma^{\infty}_+Y \to \Sigma^{\infty}_+(E^{\vee})^{\circ}$ is an isomorphism in SH(S), and $\Sigma^{\infty}_+p: \Sigma^{\infty}_+(E^{\vee})^{\circ} \to \mathbf{1}_S$ admits a section in SH(S)[η^{-1}] by (4.2.3). So we may set $f = p \circ g$.

4.3. Thom spaces of tensor products by line bundles. We are now in position to slightly improve the result obtained in §3.3.

4.3.1. Lemma. Let $E \to S$ be a vector bundle and $u \in H^0(S, \mathbb{G}_m)$. Assume that E has constant rank r. Then in the notation of (1.5) and (1.6), we have in $SH(S)[\eta^{-1}]$

$$\operatorname{Th}(u \operatorname{id}_E) = \langle u^r \rangle \colon \operatorname{Th}_S(E) \to \operatorname{Th}_S(E)$$

Proof. Since, under the identification $\operatorname{Th}_S(E \oplus 1) = \operatorname{Th}_S(E) \wedge \operatorname{Th}_S(1)$ (see (1.3.b)) we have

$$\operatorname{Th}(u \operatorname{id}_{E \oplus 1}) = \operatorname{Th}(u \operatorname{id}_E) \wedge \operatorname{Th}(u \operatorname{id}_1) = \operatorname{Th}(u \operatorname{id}_E) \wedge \langle u \rangle,$$

we may replace E with $E \oplus 1$ if necessary, and thus assume that r is odd. The \mathbb{G}_m equivariant isomorphism $E^{\circ} \simeq \mathcal{O}(-1)^{\circ}$ (see (4.2.1)) yields a commutative square in SH(S)

where the vertical arrows coincide, and are isomorphisms in $\operatorname{SH}(S)[\eta^{-1}]$ by (4.2.2). In view of (1.6.b) we may replace $E \to S$ with $\mathcal{O}(-1) \to \mathbb{P}(E)$, and thus assume that Eis a line bundle. By (2.2.8), we may replace S with E° , and thus assume that the line bundle $E \to S$ admits a trivialisation $\alpha \colon 1 \xrightarrow{\sim} E$. Then we have a commutative square of isomorphisms in $\operatorname{SH}(S)$

$$\begin{array}{c} \operatorname{Th}_{S}(E) \xrightarrow{\operatorname{Th}(u \operatorname{id}_{E})} \operatorname{Th}_{S}(E) \\ \xrightarrow{\operatorname{Th}(\alpha)} & & \uparrow^{\operatorname{Th}(\alpha)} \\ \operatorname{Th}_{S}(1) \xrightarrow{\operatorname{Th}(u \operatorname{id}_{1})} & \operatorname{Th}_{S}(1) \end{array}$$

By definition $\operatorname{Th}(u \operatorname{id}_1) = \langle u \rangle$, and we deduce using (1.6.b) that $\operatorname{Th}(u \operatorname{id}_E) = \langle u \rangle \in \operatorname{Aut}_{\operatorname{SH}(S)}(\operatorname{Th}_S(E))$. Since $\langle u^2 \rangle = \operatorname{id}$ by (2.1.1) and r is odd, it follows that $\langle u \rangle = \langle u^r \rangle$, concluding the proof.

4.3.2. **Proposition.** Let $L \to S$ be a line bundle, and $V \to S$ a vector bundle of constant rank r. If $s \in \mathbb{Z}$ is such that rs is even, then there exists an isomorphism in $SH(S)[\eta^{-1}]$

$$\operatorname{Th}_{S}(V) \simeq \operatorname{Th}_{S}(V \otimes L^{\otimes s}).$$

Proof. Upon replacing V with $V \otimes L^{\otimes s}$, we may assume that $s \ge 0$. When $\alpha: 1 \xrightarrow{\sim} L$ is a trivialisation of the line bundle L over S we have an isomorphism in SH(S)

(4.3.2.a)
$$\operatorname{Th}(\operatorname{id}_V \otimes \alpha^{\otimes s}) \colon \operatorname{Th}_S(V) \to \operatorname{Th}_S(V \otimes L^{\otimes s}).$$

Any trivialisation of L is of the form $u\alpha$ for some $u \in H^0(S, \mathbb{G}_m)$, and we have in $SH(S)[\eta^{-1}]$, by (4.3.1) and (2.1.1)

 $\operatorname{Th}(\operatorname{id}_V \otimes (u\alpha)^{\otimes s}) = \operatorname{Th}(\operatorname{id}_V \otimes \alpha^{\otimes s}) \circ \operatorname{Th}(u^s \operatorname{id}_V) = \operatorname{Th}(\operatorname{id}_V \otimes \alpha^{\otimes s}) \circ \langle u^{rs} \rangle = \operatorname{Th}(\operatorname{id}_V \otimes \alpha^{\otimes s}).$ It follows that the image of the isomorphism (4.3.2.a) in $SH(S)[\eta^{-1}]$ is independent of the choice of the trivialisation α , and we conclude as in the proof of (3.3.1).

5. Classifying spaces and characters

5.1. Models for étale classifying spaces. Here we recall some facts concerning the geometric models of the étale classifying space of a linear algebraic group given in [MV99, §4.2].

5.1.1. Let G be a linear algebraic group over S. Let (V_m, U_m, f_m) , for $m \in \mathbb{N} \setminus \{0\}$, be an admissible gadget with a nice (right) G-action, in the sense of [MV99, Definition 4.2.1]. Here $V_m \to S$ are G-equivariant vector bundles, and $U_m \subset V_m$ are G-invariant open subschemes where the G-action is free. Set $E_m G = U_m$ and $B_m G = (E_m G)/G$. Let us define $BG \in \operatorname{Spc}(S)$ as the colimit of the motivic spaces B_mG as m runs over $\mathbb{N} \setminus \{0\}$. It is proved in [MV99, Proposition 4.2.6] that the weak-equivalence class of BG does not depend on the choice of (V_m, U_m, f_m) . More precisely if $(V_m, U_m, f_m), (V'_m, U'_m, f'_m)$ are admissible gadgets with a nice G-action, and $U_m \to U'_m$ are G-equivariant morphisms commuting with the morphisms f_m, f'_m , then the induced morphism of motivic spaces $\operatorname{colim}_m(U_m/G) \to \operatorname{colim}_m(U'_m/G)$ is a weak equivalence. In the sequel we will refer to a system (V_m, U_m, f_m) as above as a model for BG, and use the notation E_mG, B_mG .

5.1.2. In the situation of (5.1.1), since B_1G is cofibrant and each $B_mG \to B_{m+1}G$ is a cofibration (for the model structure of [MV99]), it follows that the colimit BG is canonically weakly equivalent to the homotopy colimit of the motivic spaces $B_m G$ in $\operatorname{Spc}(S)$ (see e.g. [Hir03, Theorem 19.9.1]).

5.1.3. Let G be a linear algebraic group over S, and choose a model for BG. Since the map $\operatorname{colim}_m E_m G \to S$ is a weak-equivalence of motivic spaces [MV99, Proposition 4.2.3], we obtain a canonical morphism $S \to BG$ in H(S). We say that the model is pointed if we are given an S-point of E_1G . This yields map $S \to BG$ in Spc(S), whose image in H(S) is the canonical morphism described just above.

5.1.4. (See also [MV99, p.133].) Let us fix an integer $n \in \mathbb{N}$ and describe an explicit model for BGL_n. Fix an integer $p \ge n$ (we will typically take p = n). For $s \in \mathbb{N}$, we denote by Gr(n,s) the grassmannian of rank n subbundles $U \subset 1^{\oplus s}$ over S (for us a subbundle is locally split, so $1^{\oplus s}/U$ is a vector bundle). For each $m \in \mathbb{N} \setminus \{0\}$,

consider the S-scheme $V_{m,p}$ parametrising the vector bundles maps $1^{\oplus n} \to 1^{\oplus pm}$; then $V_{m,p} \to S$ is a vector bundle. Let $U_{m,p}$ the open subscheme of $V_{m,p}$ parametrising those vector bundle maps admitting Zariski-locally a retraction (i.e. making $1^{\oplus n}$ a subbundle of $1^{\oplus pm}$). Then the natural left GL_n -action on $1^{\oplus n}$ induces a right GL_n -action on $U_{m,p}$, which is free, and the quotient $U_{m,p}/\operatorname{GL}_n$ can be identified with the grassmannian $\operatorname{Gr}(n, pm)$. The inclusion $1^{\oplus m} \subset 1^{\oplus m+1}$ given by the vanishing of the last coordinate induces an inclusion

$$1^{\oplus pm} = (1^{\oplus m})^{\oplus p} \subset (1^{\oplus m+1})^{\oplus p} = 1^{\oplus p(m+1)},$$

which yields a GL_n -equivariant morphism $f_{m,p}: U_{m,p} \to U_{m+1,p}$.

Then the family $(V_{m,p}, U_{m,p}, f_{m,p})$ is an admissible gadget with a nice GL_n -action. Indeed the first condition of [MV99, Definition 4.2.1] is satisfied because $U_{1,p}$ possesses an S-point, and the second condition is satisfied with j = 2i. The fact that the group GL_n is special implies the validity of condition (3) of [MV99, Definition 4.2.4]. We thus obtained a model for BGL_n . We have just seen that this model is pointed (in the sense of (5.1.3)); a canonical pointing when p = n is induced by the identity of $1^{\oplus n}$.

5.1.5. Let $H \subset G$ be an inclusion of linear algebraic groups over S. Then any admissible gadget with a nice G-action is also one with a nice H-action (where the H-action is given by restricting the G-action). Indeed, the only non-immediate point is condition (3) of [MV99, Definition 4.2.4]. So let F be a smooth S-scheme with a free right Haction. Consider the quotient $E = (F \times G)/H$, where the right H-action on G is given by letting $h \in H$ act via $g \mapsto h^{-1}g$. Right multiplication in G induces a free right G-action on E. For any $U \in \text{Sm}_S$ with a right G-action we have isomorphisms

$$(E \times U)/G \simeq ((F \times G)/H \times U)/G \simeq (F \times (G \times U)/G)/H \simeq (F \times U)/H,$$

which are functorial in U, and thus permit to identify the morphisms $(E \times U)/G \rightarrow E/G$ and $(F \times U)/H \rightarrow F/H$. Since the former is an epimorphism in the Nisnevich topology (as the group G is nice), so is the latter.

Thus given a model for BG, we obtain a model for BH, where $E_m H = E_m G$ with the induced H-action. This yields morphisms

$$B_m H = (E_m H)/H = (E_m G)/H \rightarrow (E_m G)/G = B_m G$$

which are compatible with the transition maps as m varies, and thus a map $BH \rightarrow BG$.

5.1.6. (See also [MV99, Remark 4.2.7].) Assume that G is a linear algebraic group over S, and fix an embedding $G \subset GL_n$ as a closed subgroup. By (5.1.5) every (pointed) model for BGL_n induces a (pointed) model for BG. Since BGL_n admits a pointed model by (5.1.4), so does BG.

5.2. Products.

5.2.1. Let G, G' be linear algebraic groups over S. Choose admissible gadgets (V_m, U_m, f_m) with a nice G-action, and (V'_m, U'_m, f'_m) with a nice G'-action (in the sense of [MV99, Definition 4.2.1], recall from (5.1.6) that such exist). Then the family $(V_m \times V'_m, U_m \times U'_m, f_m \times f'_m)$ constitutes an admissible gadget with a nice $G \times G'$ -action. Indeed, to

check the last condition of [MV99, Definition 4.2.4], let $T \to X$ be a $G \times G'$ -torsor in Sm_S. Then the projection $(T \times U \times U')/(G \times G') \to T/(G \times G')$ factors as

$$(T_1 \times U')/G' \rightarrow T_1/G'$$

where $T_1 = (T \times U)/G$ followed by

$$(T_2 \times U)/G \to T_2/G$$

where $T_2 = T/G'$. Each morphism is an epimorphism in the Nisnevich topology by assumption, hence so is their composite.

Under this choice of a model for BG, we have

(5.2.1.a)
$$B_m(G \times G') = B_m G \times B_m G'.$$

5.2.2. Lemma. If G, G' are linear algebraic groups over S, we have an isomorphism

$$\mathcal{B}(G \times G') \simeq \mathcal{B}G \times \mathcal{B}G' \in \mathcal{H}(S).$$

Proof. Since the product with a given motivic space commutes with homotopy colimits, we have isomorphisms in H(S)

$$B(G \times G') \simeq \operatorname{hocolim}_m(B_m G \times B_m G') \qquad \text{by (5.2.1.a) and (5.1.2)}$$

$$\simeq \operatorname{hocolim}_m \operatorname{hocolim}_d(B_m G \times B_d G') \qquad \text{by a cofinality argument}$$

$$\simeq \operatorname{hocolim}_m(B_m G \times \operatorname{hocolim}_d B_d G')$$

$$\simeq (\operatorname{hocolim}_m B_m G) \times (\operatorname{hocolim}_d B_d G')$$

$$\simeq BG \times BG' \qquad \text{by (5.1.2).} \qquad \Box$$

5.3. Characters. In this section, we discuss general facts relating the classifying space of a linear algebraic group G to that of the kernel H of a character of G, which will be applied to explicit situations in §6.

5.3.1. Let G be a linear algebraic group over S, and fix a model for BG (see (5.1.1)). Assume given a character of G, that is a morphism of algebraic groups $\chi: G \to \mathbb{G}_m$. Considering the right G-action on \mathbb{A}^1 given by letting $g \in G$ act via $\lambda \mapsto \chi(g)^{-1}\lambda$, we define for each $m \in \mathbb{N} \setminus \{0\}$ a line bundle over $B_m G$:

(5.3.1.a)
$$C_m(\chi) = (E_m G \times \mathbb{A}^1)/G.$$

The assignment $\chi \mapsto C_m(\chi)$ satisfies

(5.3.1.b)
$$C_m(\chi\chi') = C_m(\chi) \otimes C_m(\chi'),$$

yielding group morphisms

 $\operatorname{Hom}_{\operatorname{alg. groups}}(G, \mathbb{G}_m) \to \operatorname{Pic}(B_m G) \quad ; \quad \chi \mapsto C_m(\chi).$

Let now $H \subset G$ be a closed subgroup, and consider the morphisms $B_m H \to B_m G$ defined in (5.1.5). If $\chi|_H$ denotes the restriction of the character χ to H, then

(5.3.1.c)
$$C_m(\chi) \times_{B_m G} B_m H \simeq C_m(\chi|_H).$$

If G' is a linear algebraic group over S, letting $\tilde{\chi} \colon G \times G' \to G \xrightarrow{\chi} \mathbb{G}_m$ be the induced character of $G \times G'$, we have

(5.3.1.d)
$$C_m(\tilde{\chi}) = C_m(\chi) \times B_m G'.$$

5.3.2. Let G be a linear algebraic group over S, and χ a surjective character of G. Letting $H = \ker \chi$, we thus have an exact sequence of algebraic groups over S

(5.3.2.a)
$$1 \to H \to G \xrightarrow{\chi} \mathbb{G}_m \to 1.$$

Let us fix a model for BG. As explained in (5.1.5), this yields a model for BH, and morphisms $p_m: B_m H \to B_m G$ for $m \in \mathbb{N} \setminus \{0\}$. By (5.3.1), we also have a line bundle $C_m(\chi)$ over $B_m G$, such that

$$C_m(\chi)^\circ = ((E_m G \times \mathbb{A}^1)/G)^\circ = (E_m G \times \mathbb{G}_m)/G = ((E_m G)/H \times \mathbb{G}_m)/\mathbb{G}_m = (E_m G)/H = B_m H.$$

In view of (1.3.a), this yields a cofiber sequence in $\text{Spc}_{\bullet}(S)$, for each $m \in \mathbb{N} \setminus \{0\}$

(5.3.2.b)
$$(B_m H)_+ \xrightarrow{p_{m+}} (B_m G)_+ \to \operatorname{Th}_{B_m G}(C_m(\chi)).$$

More generally (as in (1.4)), if $V \to B_m G$ is a vector bundle, we have a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$,

(5.3.2.c)
$$\operatorname{Th}_{B_m H}(V) \to \operatorname{Th}_{B_m G}(V) \to \operatorname{Th}_{B_m G}(C_m(\chi) \oplus V).$$

5.3.3. In the situation of (5.3.2), let us define

$$\operatorname{Th}_{BG}(C(\chi)) = \operatorname{colim}_m \operatorname{Th}_{B_m G}(C_m(\chi)) \in \operatorname{Spc}_{\bullet}(S).$$

As in (5.1.2), this coincides with the homotopy colimit (the transition morphisms are again monomorphisms, being directed colimits of such). We will also write $\operatorname{Th}_{BG}(C(\chi)) \in$ $\operatorname{Spt}(S)$ instead of $\Sigma^{\infty} \operatorname{Th}_{BG}(C(\chi))$. Taking the (homotopy) colimit of (5.3.2.b) yields a cofiber sequence in $\operatorname{Spc}_{\bullet}(S)$

(5.3.3.a)
$$(BH)_+ \to (BG)_+ \to \operatorname{Th}_{BG}(C(\chi))$$

5.3.4. In the situation of (5.3.2), assume that the model for BG is pointed. Then we have a commutative diagram of S-schemes with cartesian squares

where e_1 is induced by the S-point and the G-action on E_1G , and j_1 , resp. i_1 , is obtained by taking the H-quotient, resp. G-quotient of e_1 . Composing i_1 and j_1 with the natural maps $B_1G \to BG$ and $B_1H \to BH$ respectively, we obtain maps $i: S \to BG$ and $j: \mathbb{G}_m \to BH$ in Spc(S). Note that i is the map described in (5.1.3), and that the left-hand cartesian square in (5.3.4.a) shows that the map $j: \mathbb{G}_m \to BH$ classifies the H-torsor $\chi: G \to \mathbb{G}_m$.

The right-hand cartesian square in (5.3.4.a) shows that the \mathbb{G}_m -torsor $B_1H \to B_1G$ pulls back to the trivial torsor along i_1 , which yields a trivialisation of the line bundle $i_1^*C_1(\chi)$ over S, and thus a morphism in Spc_•(S)

$$t: T = \operatorname{Th}_{S}(1) \to \operatorname{Th}_{B_{1}G}(C_{1}(\chi)) \to \operatorname{Th}_{BG}(C(\chi)).$$

We thus obtain a commutative diagram in $\operatorname{Spc}_{\bullet}(S)$, whose rows are cofiber sequences

(5.3.4.b)
$$(BH)_{+} \longrightarrow (BG)_{+} \longrightarrow Th_{BG}(C(\chi))$$
$$i_{+} \uparrow \qquad i_{+} \uparrow \qquad t \uparrow$$
$$(\mathbb{G}_{m})_{+} \longrightarrow S_{+} \longrightarrow T$$

5.3.5. In the situation of (5.3.2), using (2.2.5) for the line bundle $C_m(\chi) \to B_m G$, and applying the functor $\operatorname{Spc}_{\bullet}(B_m G)[\eta^{-1}] \to \operatorname{Spc}_{\bullet}(S)[\eta^{-1}]$ of (1.9), we have homotopy cocartesian squares in $\operatorname{Spc}_{\bullet}(S)[\eta^{-1}]$

which are compatible with the transition maps as m varies by (2.2.3). Taking the homotopy colimit, and proceeding as in the proof of (2.2.6), we obtain an isomorphism in $SH(S)[\eta^{-1}]$

(5.3.5.a)
$$\Sigma^{\infty}_{+} \mathbf{B}H \simeq \Sigma^{\infty}_{+} \mathbf{B}G \oplus \Sigma^{-2,-1} \operatorname{Th}_{\mathbf{B}G}(C(\chi)) \in \mathrm{SH}(S)[\eta^{-1}].$$

6. Computations of classifying spaces

6.1. **Diagonalisable groups.** Using the embeddings $\mathbb{P}^k \subset \mathbb{P}^{k+1}$ of (4.1.1) for $k \in \mathbb{N}$, we define, for $n \in \mathbb{Z}$

 $\mathbb{P}^{\infty} = \operatorname{hocolim}_{k} \mathbb{P}^{k} \in \operatorname{Spc}(S) \quad \text{and} \quad \operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n)) = \operatorname{hocolim}_{k} \operatorname{Th}_{\mathbb{P}^{k}}(\mathcal{O}(n)) \in \operatorname{Spc}_{\bullet}(S),$

and as usual write $\operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n)) \in \operatorname{Spt}(S)$ instead of $\Sigma^{\infty} \operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n))$. We have a natural map $\iota_{\infty} \colon S = \mathbb{P}^{0} \to \mathbb{P}^{\infty}$ in $\operatorname{Spc}(S)$. For each $n \in \mathbb{Z}$, the line bundle $\mathcal{O}(n)$ over \mathbb{P}^{0} admits a canonical trivialisation, so that (4.1.2.a) yields a canonical map in $\operatorname{Spt}(S)$

(6.1.0.a)
$$\Sigma^{2,1}\mathbf{1}_S \to \operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n)).$$

6.1.1. **Proposition.** Let $n \in \mathbb{Z}$. The following hold in $SH(S)[\eta^{-1}]$:

- (i) The morphism ι_{∞} induces an isomorphism $\mathbf{1}_{S} \simeq \Sigma^{\infty}_{+} \mathbb{P}^{\infty}$.
- (ii) If n is odd, then $\operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n)) = 0$.

(iii) If n is even, then (6.1.0.a) induces an isomorphism $\Sigma^{2,1}\mathbf{1}_S \simeq \mathrm{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(n)).$

Proof. We apply (4.1.3) with $\mathcal{D} = (n)$, and so $\mathcal{O}(\mathcal{D}) = \mathcal{O}(n)$. We obtain that $\Sigma^{2,1}\mathbf{1}_S \to \operatorname{Th}_{\mathbb{P}^k}(\mathcal{O}(n))$ is a weak equivalence in $\operatorname{Spt}(S)[\eta^{-1}]$ when n, k are even. Taking the homotopy colimit over k yields a weak equivalence $\Sigma^{2,1}\mathbf{1}_S \to \operatorname{Th}_{\mathbb{P}^\infty}(\mathcal{O}(n))$ in $\operatorname{Spt}(S)[\eta^{-1}]$ when n is even. This proves (iii). The other statements are deduced in a similar way from (4.1.3).

6.1.2. Consider the model for $\mathbb{B}\mathbb{G}_m$ described in (5.1.4) with p = n = 1, under the identification $\mathbb{G}_m = \mathrm{GL}_1$. Then $B_m \mathbb{G}_m = \mathbb{P}^{m-1}$, and thus $\mathbb{B}\mathbb{G}_m = \mathbb{P}^{\infty}$. Furthermore, the line bundle $C_m(\mathrm{id}_{\mathbb{G}_m})$ over $B_m G$ defined in (5.3.1.a) may be identified with the tautological bundle $\mathcal{O}(-1)$ over \mathbb{P}^{m-1} .

6.1.3. Theorem. Let $n \in \mathbb{N} \setminus \{0\}$. The following hold in $SH(S)[\eta^{-1}]$:

- (i) The natural morphism $\mathbf{1}_S \to \Sigma^{\infty}_+ \mathbb{B}\mathbb{G}_m$ is an isomorphism.
- (ii) If n is odd, the natural morphism $\mathbf{1}_S \to \Sigma^{\infty}_+ B\mu_n$ is an isomorphism.
- (iii) If n is even, the morphism $\mathbb{G}_m \to \mathbb{B}\mu_n$ classifying the μ_n -torsor $\mathbb{G}_m \to \mathbb{G}_m$ given by taking n-th powers induces an isomorphism $\Sigma^{\infty}_+ \mathbb{G}_m \simeq \Sigma^{\infty}_+ \mathbb{B}\mu_n$.

Proof. Let us consider the model for $\mathbb{B}\mathbb{G}_m$ described in (6.1.2), where $B_m\mathbb{G}_m = \mathbb{P}^{m-1}$ and $\mathbb{B}\mathbb{G}_m = \mathbb{P}^{\infty}$. Then the first statement follows from (6.1.1.i).

Next consider the character $n: \mathbb{G}_m \to \mathbb{G}_m$ given by taking *n*-th powers. Its kernel is μ_n , and the line bundle $C_m(n)$ over $B_m\mathbb{G}_m$ (defined in (5.3.1.a)) corresponds to the line bundle $\mathcal{O}(-n)$ over \mathbb{P}^{m-1} (this may be seen for instance by combining (6.1.2) with (5.3.1.b)). So we are in the situation of (5.3.2) with $G = \mathbb{G}_m, \chi = n, H = \mu_n$. Thus (5.3.3.a) yields a distinguished triangle in SH(S)

$$\Sigma^{\infty}_{+} \mathrm{B}\mu_n \to \Sigma^{\infty}_{+} \mathrm{B}\mathbb{G}_m \to \mathrm{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(-n)) \to \Sigma^{1,0}\Sigma^{\infty}_{+} \mathrm{B}\mu_n.$$

If n is odd, then $\operatorname{Th}_{\mathbb{P}^{\infty}}(\mathcal{O}(-n)) = 0$ in $\operatorname{SH}(S)[\eta^{-1}]$ by (6.1.1.ii), and the above distinguished triangle shows that the morphism $\Sigma^{\infty}_{+} \mathrm{B}\mu_{n} \to \Sigma^{\infty}_{+} \mathrm{B}\mathbb{G}_{m}$ is an isomorphism in $\operatorname{SH}(S)[\eta^{-1}]$. Thus the second statement follows from the first.

Assume that n is even. Then in the diagram (5.3.4.b) the maps i_+ and t become isomorphisms in $SH(S)[\eta^{-1}]$ by (6.1.1.i) and (6.1.1.iii), hence $\Sigma^{\infty}_+ j \colon \Sigma^{\infty}_+ \mathbb{G}_m \to \Sigma^{\infty}_+ \mathbb{B}\mu_n$ is also one. As observed in (5.3.2), the latter is induced by the μ_n -torsor $n \colon \mathbb{G}_m \to \mathbb{G}_m$. \Box

6.1.4. **Remark.** Combining (6.1.3) with (5.2.2), we have thus obtained a "computation" in $SH(S)[\eta^{-1}]$ of the classifying space BG of every finitely generated diagonalisable group G.

6.2. SL versus SL^c. Let $n \in \mathbb{N} \setminus \{0\}$. Consider the character

$$\nu_n \colon \operatorname{GL}_n \times \mathbb{G}_m \to \mathbb{G}_m \quad ; \quad (M, t) \mapsto t^{-2} \det M.$$

By definition (see [PW18, §3]), we have $SL_n^c = \ker \nu_n$. We view SL_n as a subgroup of SL_n^c via the mapping $M \mapsto (M, 1)$.

6.2.1. **Proposition.** For $n \in \mathbb{N} \setminus \{0\}$ the inclusion $SL_n \subset SL_n^c$ induces an isomorphism

$$\Sigma^{\infty}_{+} \operatorname{BSL}_{n} \xrightarrow{\sim} \Sigma^{\infty}_{+} \operatorname{BSL}_{n}^{c} \text{ in } \operatorname{SH}(S)[\eta^{-1}].$$

Proof. The character

$$\delta_n \colon \operatorname{SL}_n^c \to \mathbb{G}_m \quad ; \quad (M,t) \mapsto t$$

is surjective (recall that $n \ge 1$), and satisfies $SL_n = \ker \delta_n$.

Set $P_n = \operatorname{GL}_n \times \mathbb{G}_m$, and denote by $q_n \colon P_n \to \mathbb{G}_m$ the second projection. Let us fix an arbitrary model for BGL_n , but choose the model for BG_m described in (6.1.2), so that $B_m \mathbb{G}_m = \mathbb{P}^{m-1}$. Recall from (5.2.1.a) that this yields a model for BP_n , such that

$$B_m P_n = (B_m \operatorname{GL}_n) \times (B_m \mathbb{G}_m) = (B_m \operatorname{GL}_n) \times \mathbb{P}^{m-1}.$$

Letting $g_n: P_n \to \operatorname{GL}_n$ be the first projection, we have, as characters $P_n \to \mathbb{G}_m$

$$\nu_n = g_n^*(\det_n) \cdot q_n^*(\operatorname{id}_{\mathbb{G}_m})^{-2},$$

where det_n: GL_n $\rightarrow \mathbb{G}_m$ denotes the determinant morphism. It follows from (5.3.1.b) and (5.3.1.d) that, as line bundles over $B_m P_n$, we have in the notation of (5.3.1.a)

$$C_m(\nu_n) \simeq C_m(\det_n) \boxtimes C_m(\operatorname{id}_{\mathbb{G}_m})^{\otimes -2} \quad ; \quad C_m(q_n) \simeq 1 \boxtimes C_m(\operatorname{id}_{\mathbb{G}_m}).$$

Recall from (6.1.2) that the line bundle $C_m(\mathrm{id}_{\mathbb{G}_m}) \to B_m \mathbb{G}_m$ corresponds to $\mathcal{O}(-1) \to \mathbb{P}^{m-1}$. Applying (4.1.5) to the projective bundle $B_m P_n \to B_m \operatorname{GL}_n$, for m even we have

(6.2.1.a)
$$\operatorname{Th}_{B_m P_n}(C_m(\nu_n) \oplus C_m(q_n)) = 0 = \operatorname{Th}_{B_m P_n}(C_m(q_n)) \in \operatorname{SH}(S)[\eta^{-1}].$$

Since the character δ_n is the restriction of $q_n \colon P_n \to \mathbb{G}_m$, it follows from (5.3.1.c) that the line bundle $C_m(\delta_n)$ over $B_m \operatorname{SL}_n^c$ is the pullback of $C_m(q_n)$ over $B_m P_n$. By (5.3.2.c), we have a distinguished triangle in $\operatorname{SH}(S)$

$$\operatorname{Th}_{B_m \operatorname{SL}_n^c}(C_m(\delta_n)) \to \operatorname{Th}_{B_m P_n}(C_m(q_n)) \to \operatorname{Th}_{B_m P_n}(C_m(\nu_n) \oplus C_m(q_n))$$
$$\to \Sigma^{1,0} \operatorname{Th}_{B_m \operatorname{SL}_n^c}(C_m(\delta_n)).$$

so that, in view of (6.2.1.a)

$$\operatorname{Th}_{B_m \operatorname{SL}_n^c}(C_m(\delta_n)) = 0 \in \operatorname{SH}(S)[\eta^{-1}] \text{ for } m \text{ even}$$

Now the distinguished triangle in SH(S) (see (5.3.2.b))

$$\Sigma^{\infty}_{+} B_m \operatorname{SL}_n \to \Sigma^{\infty}_{+} B_m \operatorname{SL}_n^c \to \operatorname{Th}_{B_m \operatorname{SL}_n^c}(C_m(\delta_n)) \to \Sigma^{1,0} \Sigma^{\infty}_{+} B_m \operatorname{SL}_n^c$$

implies that, for m even, the natural map induces an isomorphism

$$\Sigma^{\infty}_{+} B_m \operatorname{SL}_n \xrightarrow{\sim} \Sigma^{\infty}_{+} B_m \operatorname{SL}_n^c \quad \text{in } \operatorname{SH}(S)[\eta^{-1}].$$

The statement follows by taking the homotopy colimit.

6.2.2. **Remark.** Let $A \in SH(S)$ be an η -periodic commutative ring spectrum, and consider the corresponding cohomology theory $A^{*,*}(-)$ (see (3.1.1)). Then by (6.2.1), we have a natural isomorphism

$$A^{*,*}(\mathrm{BSL}_n^c) \simeq A^{*,*}(\mathrm{BSL}_n).$$

If A is SL-oriented (see (3.2.5)) and $S = \operatorname{Spec} k$ with k a field of characteristic not two, Ananyevskiy computed in [Ana15, Theorem 10] that

$$A^{*,*}(BSL_n) = \begin{cases} A^{*,*}(S)[[p_1, \dots, p_{r-1}, e]]_h & \text{if } n = 2r \text{ with } r \in \mathbb{N} \setminus \{0\} \\ A^{*,*}(S)[[p_1, \dots, p_r]]_h & \text{if } n = 2r+1 \text{ with } r \in \mathbb{N}, \end{cases}$$

where p_i has degree (4i, 2i) and e has degree (2r, r) (here the notation $R[[x_1, \ldots, x_m]]_h$ refers to the homogeneous power series ring in m variables over the graded ring R, see [Ana15, Definition 27]). This computation remains valid (with exactly the same arguments) when S is an arbitrary noetherian scheme of finite dimension, under the assumption that 2 is invertible in S. Removing that last assumption seems to require a modification of the arguments of [Ana15], the problem being with [Ana15, Lemma 6] (which is used to prove [Ana15, Theorem 9]).

6.3. GL and SL. In this section, we compare the classifying spaces BGL_{2r} , BGL_{2r+1} , BSL_{2r} , BSL_{2r+1} .

6.3.1. Recall that under the model for BGL_n described in (5.1.4) for p = n, the scheme $B_m \operatorname{GL}_n$ is identified with the grassmannian $\operatorname{Gr}(n, nm)$. The closed immersion $\operatorname{Gr}(n, nm) \to \operatorname{Gr}(n + 1, (n + 1)m)$ mapping a subbundle $E \subset (1^{\oplus m})^{\oplus n}$ to $E \oplus 1 \subset (1^{\oplus m})^{\oplus n} \oplus 1^{\oplus m} = (1^{\oplus m})^{\oplus n+1}$, where the inclusion $1 \subset 1^{\oplus m}$ is given by the vanishing of the last m - 1 coordinates, induces a morphism $f_m \colon B_m \operatorname{GL}_n \to B_m \operatorname{GL}_{n+1}$ which is compatible with the transition maps as m varies. This yields a morphism in $\operatorname{Spc}(S)$

$$(6.3.1.a) \qquad \qquad BGL_n \to BGL_{n+1}.$$

6.3.2. For integers $u, v, w \in \mathbb{N}$, we denote by $\operatorname{Gr}(u \subset v, w)$ the flag variety of subbundles $P \subset Q \subset 1^{\oplus w}$ with rank P = u and rank Q = v. Let $r, s \in \mathbb{N}$, and consider the morphisms

$$\operatorname{Gr}(2r,s) \xleftarrow{p} \operatorname{Gr}(2r \subset 2r+1,s) \xrightarrow{q} \operatorname{Gr}(2r+1,s)$$

given by mapping a flag $P \subset Q$ to P, resp. Q.

For $n \in \{2r, 2r+1\}$, let us denote by $\mathcal{U}_n \subset 1^{\oplus s}$ the tautological rank n subbundle over $\operatorname{Gr}(n, s)$, and write $\mathcal{Q}_n = 1^{\oplus s}/\mathcal{U}_n$. Then the morphism p may be identified with the projective bundle $\mathbb{P}(\mathcal{Q}_{2r})$, and the morphism q is the projective bundle $\mathbb{P}(\mathcal{U}_{2r+1}^{\vee})$.

6.3.3. **Proposition.** The map $BGL_{2r} \to BGL_{2r+1}$ of (6.3.1.a) becomes an isomorphism in $SH(S)[\eta^{-1}]$.

Proof. Let n = 2r. For $m \in \mathbb{N} \setminus \{0\}$, consider the commutative diagram in Sm_S

$$\operatorname{Gr}(n, (n+1)m) \xrightarrow{g_m} \operatorname{Gr}(n \subset n+1, (n+1)m) \xrightarrow{f_m} \operatorname{Gr}(n+1, (n+1)m)$$

where the morphism j_m is given by mapping $E \subset (1^{\oplus m})^{\oplus n}$ to

$$E \subset E \oplus 1 \subset (1^{\oplus m})^{\oplus n} \oplus 1^{\oplus m} = (1^{\oplus m})^{\oplus n+1},$$

with the inclusion $1 \subset 1^{\oplus m}$ given by the vanishing of the m-1 last coordinates. Here the morphisms p_m, q_m are the morphisms p, q described in (6.3.2) when s = (n+1)m. The morphism f_m is the one described in (6.3.1), and the morphism g_m is induced by the inclusion

(6.3.3.a)
$$(1^{\oplus m})^{\oplus n} = (1^{\oplus m})^{\oplus n} \oplus 0 \subset (1^{\oplus m})^{\oplus n} \oplus 1^{\oplus m} = (1^{\oplus m})^{\oplus n+1}.$$

The morphisms of this diagram are compatible with the transition maps as m varies, induced by the inclusions $1^{\oplus m} \subset 1^{\oplus m+1}$ given by the vanishing of the last coordinate.

The morphism q_m is a \mathbb{P}^n -bundle, hence is an isomorphism in $\mathrm{SH}(S)[\eta^{-1}]$ by (4.1.6) (recall that n = 2r is even). The morphism p_m is a $\mathbb{P}^{(n+1)m-n-1}$ -bundle, hence is also an isomorphism in $\mathrm{SH}(S)[\eta^{-1}]$ when m is odd by (4.1.6).

In the notation of (5.1.4) the morphism g_m is the GL_n -quotient of the morphism $U_{m,n} \to U_{m,n+1}$ induced by (6.3.3.a). Therefore it follows from (5.1.1) that the map $\operatorname{colim}_m g_m$ is a weak equivalence of motivic spaces.

Applying the functor Σ^{∞}_+ : $\operatorname{Sm}_S \to \operatorname{Spt}(S)[\eta^{-1}]$ to the above diagram, and taking the homotopy colimit over m, we thus obtain a commutative diagram in $\operatorname{Spt}(S)[\eta^{-1}]$, where all maps are weak equivalences. Since the map (6.3.1.a) is obtained as $\operatorname{colim}_m f_m$, the proposition follows.

6.3.4. Let $n \in \mathbb{N} \setminus \{0\}$. The group SL_n is the kernel of the determinant morphism $\det_n: \mathrm{GL}_n \to \mathbb{G}_m$, which is surjective (as $n \ge 1$). We are thus in the situation of (5.3.2), so that we have by (5.3.5.a), a splitting

(6.3.4.a)
$$\Sigma_{+}^{\infty} \operatorname{BSL}_{n} = \Sigma_{+}^{\infty} \operatorname{BGL}_{n} \oplus \Sigma^{-2,-1} \operatorname{Th}_{\operatorname{BGL}_{n}}(C(\operatorname{det}_{n})) \in \operatorname{SH}(S)[\eta^{-1}]$$

6.3.5. Using the model for BGL_n described in (5.1.4) with p = n, the variety $B_m \operatorname{GL}_n$ coincides with the grassmannian $\operatorname{Gr}(n, nm)$. Observe that the tautological bundle \mathcal{U}_n over this variety is isomorphic to the quotient $(E_m \operatorname{GL}_n \times \mathbb{A}^n)/\operatorname{GL}_n$, where the right GL_n -action on \mathbb{A}^n is given by letting $\varphi \in \operatorname{GL}_n$ act via $v \mapsto \varphi^{-1}(v)$. The GL_n -equivariant isomorphism $\det(E_m \operatorname{GL}_n \times \mathbb{A}^n) \simeq E_m \operatorname{GL}_n \times \mathbb{A}^1$, where the right GL_n -action on \mathbb{A}^1 is given by letting $\varphi \in \operatorname{GL}_n$ act via $\lambda \mapsto \det_n(\varphi^{-1})\lambda$, yields an isomorphism of line bundles over $B_m \operatorname{GL}_n$

(6.3.5.a)
$$\det \mathcal{U}_n \simeq C_m(\det_n).$$

6.3.6. **Proposition.** Let $r \in \mathbb{N}$. Then in $SH(S)[\eta^{-1}]$, the natural morphism $BSL_{2r} \rightarrow BGL_{2r}$ acquires a section, and the natural morphism $BSL_{2r+1} \rightarrow BGL_{2r+1}$ becomes an isomorphism.

Proof. The first statement follows from (6.3.4.a). Let us prove the second. We use the model for BGL_{2r+1} described in (5.1.4) with n = p = 2r + 1, so that $B_m GL_{2r+1} = Gr(2r + 1, s)$ where s = (2r + 1)m. We consider the situation of (6.3.2), and use the notation thereof. We have an exact sequence of vector bundles over $Y_m = Gr(2r \subset 2r + 1, s) = \mathbb{P}(\mathcal{Q}_{2r})$

$$0 \to p^* \mathcal{U}_{2r} \to q^* \mathcal{U}_{2r+1} \to \mathcal{O}_{\mathbb{P}(\mathcal{Q}_{2r})}(-1) \to 0,$$

Taking determinants and using (6.3.5.a) we obtain an isomorphism of line bundles

(6.3.6.a)
$$p^*(\det \mathcal{U}_{2r}) \otimes \mathcal{O}_{\mathbb{P}(\mathcal{Q}_{2r})}(-1) \simeq q^*(\det \mathcal{U}_{2r+1}) \simeq q^*C_m(\det_{2r+1}).$$

When *m* is even, the vector bundle $Q_{2r} = 1^{\oplus s}/\mathcal{U}_{2r}$ has even rank, hence it follows from (4.1.5) and (6.3.6.a) that $\operatorname{Th}_{Y_m}(q^*C_m(\det_{2r+1})) = 0$ in $\operatorname{SH}(S)[\eta^{-1}]$. Since $q: Y_m \to \operatorname{Gr}(2r+1,s)$ is a \mathbb{P}^{2r} -bundle, it then follows from (4.1.6) (applied with $V = C_m(\det_{2r+1}))$ that $\operatorname{Th}_{\operatorname{Gr}(2r+1,s)}(C_m(\det_{2r+1})) = 0$ in $\operatorname{SH}(S)[\eta^{-1}]$ for *m* even. Taking the homotopy colimit over *m*, we deduce that $\operatorname{Th}_{\operatorname{BGL}_{2r+1}}(C(\det_{2r+1})) = 0$ in $\operatorname{SH}(S)[\eta^{-1}]$, and the second statement of the proposition follows from (6.3.4.a). \Box

6.3.7. **Remark.** Let $A \in SH(S)$ be an η -periodic SL-oriented commutative ring spectrum (see (3.2.5)), and consider the corresponding cohomology theory $A^{*,*}(-)$ (see (3.1.1)). Assume that 2 is invertible in S. Combining (6.3.6) and (6.3.3) with Ananyevskiy's computation of $A^{*,*}(BSL_{2r+1})$ (see (6.2.2)), we recover Levine's computation [Lev19, Theorem 4.1]

$$A^{*,*}(BGL_{2r}) = A^{*,*}(BGL_{2r+1}) = A^{*,*}(S)[[p_1, \dots, p_r]]_h$$

(Note that this permits to remove some of the technical assumptions present in the statement of [Lev19, Theorem 4.1].)

Appendix A. An invariant of μ_2 -torsors

We have proved in (6.1.3) that in $SH(S)[\eta^{-1}]$, for $n \in \mathbb{N}$

$$\Sigma^{\infty}_{+} \mathbf{B} \mu_{n} = \begin{cases} \mathbf{1}_{S} & \text{if } n = 0 \text{ (i.e. } \mu_{n} = \mathbb{G}_{m} \text{) or } n \text{ is odd,} \\ \Sigma^{\infty}_{+} \mathbb{G}_{m} = \mathbf{1}_{S} \oplus \mathbf{1}_{S} & \text{if } n > 0 \text{ is even.} \end{cases}$$

Thus, when n > 0 is even, there is essentially one nontrivial invariant of μ_n -torsors over S in $\mathrm{SH}(S)[\eta^{-1}]$, in the form of an element of $\mathrm{End}_{\mathrm{SH}(S)[\eta^{-1}]}(\mathbf{1}_S)$. Moreover, it also follows from (6.1.3) that the morphism $\mathrm{B}\mu_2 \to \mathrm{B}\mu_{2r}$ becomes an isomorphism in $\mathrm{SH}(S)[\eta^{-1}]$ for r > 0. Therefore the above-mentioned invariant of μ_n -torsors is induced by an invariant of μ_2 -torsors, which is however not really explicit from this description. In this section we provide an explicit construction of this invariant (the connection with the above discussion is made in (A.9) below).

A.1. Let $L \to S$ be a line bundle. As observed by Ananyevskiy [Ana20, Lemma 4.1], the isomorphism of S-schemes $L^{\circ} \xrightarrow{\sim} (L^{\vee})^{\circ}$, given locally by $l \mapsto l^{\vee}$ where $l^{\vee}(l) = 1$, induces an isomorphism in $H_{\bullet}(S)$

$$\sigma_L \colon \operatorname{Th}_S(L) \xrightarrow{\sim} \operatorname{Th}_S(L^{\vee}).$$

A.2. If $\varphi: L \to M$ is an isomorphism of line bundles over X, we have (see (1.5))

(A.2.a) $\sigma_L = \operatorname{Th}(\varphi^{\vee}) \circ \sigma_M \circ \operatorname{Th}(\varphi).$

A.3. **Definition.** It will be convenient to think of a μ_2 -torsor over S as a pair (L, λ) , where $L \to S$ is a line bundle, and $\lambda \colon L \xrightarrow{\sim} L^{\vee}$ is an isomorphism of line bundles over S. Isomorphisms $(L, \lambda) \to (L', \lambda')$ are given by isomorphisms of line bundles $\varphi \colon L \xrightarrow{\sim} L'$ such that $\lambda = \varphi^{\vee} \circ \lambda' \circ \varphi$. The set of isomorphism classes of μ_2 -torsors is denoted $H^1_{et}(S, \mu_2)$; it is endowed with a group structure induced by the tensor product of line bundles.

A.4. **Definition.** Consider a μ_2 -torsor, given by a line bundle $L \to S$ and an isomorphism $\lambda \colon L \xrightarrow{\sim} L^{\vee}$. Let us consider the composite isomorphism in SH(S) (see (A.1) for the definition of σ_L , and (1.5) for that of $Th(\lambda)$)

$$\operatorname{Th}_{S}(L) \xrightarrow{\operatorname{Th}(\lambda)} \operatorname{Th}_{S}(L^{\vee}) \xrightarrow{\sigma_{L}^{-1}} \operatorname{Th}_{S}(L).$$

This yields an element $a_{(L,\lambda)} = \Sigma^{-L}(\sigma_L^{-1} \circ \operatorname{Th}(\lambda)) \in \operatorname{Aut}_{\operatorname{SH}(S)}(\mathbf{1}_S)$. We define

$$\alpha(L,\lambda) = (a_{(1,\operatorname{can})})^{-1} \circ a_{(L,\lambda)} \in \operatorname{Aut}_{\operatorname{SH}(S)}(\mathbf{1}_S).$$

where can: $1 \to 1^{\vee}$ is the canonical isomorphism of line bundles over S. (The element $a_{(1,\text{can})}$ corresponds to the element ϵ of [Mor04, §6.1].)

A.5. This construction is compatible with pullbacks, in the sense that if $f: R \to S$ is a morphism of noetherian schemes of finite dimension, and (L, λ) a μ_2 -torsor, then the composite in SH(R)

$$\mathbf{1}_R = f^* \mathbf{1}_S \xrightarrow{f^* \alpha(L,\lambda)} f^* \mathbf{1}_S = \mathbf{1}_R$$

is $\alpha(f^*L, f^*\lambda)$.

A.6. **Example.** Consider a μ_2 -torsor (L, λ) where L = 1 is the trivial line bundle. Then $\lambda = u$ can for some $u \in H^0(S, \mathbb{G}_m)$, hence $\operatorname{Th}(\lambda) = \operatorname{Th}(\operatorname{can}) \circ \operatorname{Th}(u \operatorname{id}_1)$. Therefore (see (1.6))

$$\begin{aligned} \alpha(1,\lambda) &= \Sigma^{-2,-1}((\sigma_1^{-1} \circ \operatorname{Th}(\operatorname{can}))^{-1} \circ (\sigma_1^{-1} \circ \operatorname{Th}(\lambda)) \\ &= \Sigma^{-2,-1}(\operatorname{Th}(\operatorname{can})^{-1} \circ \operatorname{Th}(\lambda)) \\ &= \Sigma^{-2,-1}\operatorname{Th}(u \operatorname{id}_1) \\ &= \langle u \rangle. \end{aligned}$$

A.7. **Proposition.** The assignment $(L, \lambda) \mapsto \alpha(L, \lambda)$ induces a morphism of pointed sets

$$\alpha \colon H^1_{et}(S, \mu_2) \to \operatorname{Aut}_{\operatorname{SH}(S)}(\mathbf{1}_S).$$

Proof. By construction we have $\alpha(1, \operatorname{can}) = \operatorname{id}$. Consider an isomorphism of μ_2 -torsors $(L, \lambda) \xrightarrow{\sim} (M, \mu)$ (see (A.3)), given by an isomorphism $\varphi \colon L \xrightarrow{\sim} M$. Let us set $a_L = a_{(L,\lambda)}$ and $a_M = a_{(M,\mu)}$ (see (A.4)). Then, in $\operatorname{Aut}_{\operatorname{SH}(S)}(\operatorname{Th}_S(L))$,

$$\begin{split} \Sigma^{L} a_{L} &= \sigma_{L}^{-1} \circ \operatorname{Th}(\lambda) = \sigma_{L}^{-1} \circ \operatorname{Th}(\varphi^{\vee}) \circ \operatorname{Th}(\mu) \circ \operatorname{Th}(\varphi) & \text{(as } \lambda = \varphi^{\vee} \circ \mu \circ \varphi) \\ &= \operatorname{Th}(\varphi)^{-1} \circ \sigma_{M}^{-1} \circ \operatorname{Th}(\mu) \circ \operatorname{Th}(\varphi) & \text{by (A.2.a)} \\ &= \operatorname{Th}(\varphi)^{-1} \circ (\Sigma^{M} a_{M}) \circ \operatorname{Th}(\varphi) \\ &= \operatorname{Th}(\varphi)^{-1} \circ \operatorname{Th}(\varphi) \circ (\Sigma^{L} a_{M}) & \text{(by (1.5.b))} \\ &= \Sigma^{L} a_{M}, \end{split}$$

whence $a_L = a_M$, and $\alpha(L, \lambda) = \alpha(M, \mu)$.

A.8. **Proposition.** The assignment $(L, \lambda) \mapsto \alpha(L, \lambda)$ induces a group morphism

$$\alpha \colon H^1_{et}(S, \mu_2) \to \operatorname{Aut}_{\operatorname{SH}(S)[\eta^{-1}]}(\mathbf{1}_S).$$

Proof. Consider μ_2 -torsors given by line bundles L, M over S, and isomorphisms $L \xrightarrow{\sim} L^{\vee}$ and $M \xrightarrow{\sim} M^{\vee}$. As the functor $f^* \colon \operatorname{SH}(S)[\eta^{-1}] \to \operatorname{SH}(L^{\circ})[\eta^{-1}]$ is faithful (2.2.8), by functoriality (A.5) we may assume that the line bundle L is trivial. Similarly, we may assume that M is also trivial. By (A.7), we may assume that L = M = 1. Then, in view of (A.6), the statement follows from (1.6.a).

A.9. **Remark.** Let $\rho: \mathbb{G}_m \to \mathbb{B}\mu_2$ be the map classifying the μ_2 -torsor $(1, t \operatorname{id})$ over \mathbb{G}_m , where $t \in H^0(\mathbb{G}_m, \mathbb{G}_m)$ is the tautological section. Recall from (6.1.3.iii) that $\Sigma^{\infty}_{+}\rho: \Sigma^{\infty}_{+}\mathbb{G}_m \to \Sigma^{\infty}_{+}\mathbb{B}\mu_2$ is an isomorphism in $\mathrm{SH}(S)[\eta^{-1}]$. If a μ_2 -torsor (L, λ) is classified by the map $f: S \to \mathbb{B}\mu_2$, we claim that $\alpha(L, \lambda)$ is the composite in $\mathrm{SH}(S)[\eta^{-1}]$ (the map π_1 was defined in (2.2.1))

(A.9.a)
$$\mathbf{1}_S \xrightarrow{\Sigma_+^{\infty} f} \Sigma_+^{\infty} B\mu_2 \xrightarrow{(\Sigma_+^{\infty} \rho)^{-1}} \Sigma_+^{\infty} \mathbb{G}_m \xrightarrow{\Sigma^{-2,-1} \Sigma_+^{\infty} \pi_1} \mathbf{1}_S.$$

Indeed, applying the functor $SH(S)[\eta^{-1}] \rightarrow SH(L^{\circ})[\eta^{-1}]$ which is faithful by (2.2.8), and using (A.7), we may assume that L = 1. Then λ is given by multiplication by

an element $u \in H^0(S, \mathbb{G}_m)$, and the map f factors as $S \xrightarrow{u} \mathbb{G}_m \xrightarrow{\rho} B\mu_2$. Denoting by $p: \mathbb{G}_m \to S$ is the projection, the composite (A.9.a) is given by

$$\mathbf{1}_S \xrightarrow{\Sigma_+^{\infty} u} \Sigma_+^{\infty} \mathbb{G}_m \xrightarrow{\langle t \rangle} \Sigma_+^{\infty} \mathbb{G}_m \xrightarrow{\Sigma_+^{\infty} p} \mathbf{1}_S,$$

which coincides with $\langle u \rangle \in \operatorname{Aut}_{\operatorname{SH}(S)[n^{-1}]}(S)$. Thus the claim follows from (A.6).

References

- [Ana15] Alexey Ananyevskiy. The special linear version of the projective bundle theorem. Compos. Math., 151(3):461–501, 2015.
- [Ana16a] Alexey Ananyevskiy. On the push-forwards for motivic cohomology theories with invertible stable Hopf element. *Manuscripta Math.*, 150(1-2):21–44, 2016.
- [Ana16b] Alexey Ananyevskiy. On the relation of special linear algebraic cobordism to Witt groups. Homology Homotopy Appl., 18(1):204–230, 2016.
- [Ana20] Alexey Ananyevskiy. SL-oriented cohomology theories. In Motivic homotopy theory and refined enumerative geometry, volume 745 of Contemp. Math., pages 1–19. Amer. Math. Soc., Providence, RI, 2020.
- [Ara80] Jón Kr. Arason. Der Wittring projektiver Räume. Math. Ann., 253(3):205–212, 1980.
- [Bac18] Tom Bachmann. Motivic and real étale stable homotopy theory. *Compos. Math.*, 154(5):883–917, 2018.
- [BH20] Tom Bachmann and Michael J. Hopkins. η-periodic motivic stable homotopy theory over fields. Preprint, 2020. arXiv:2005.06778.
- [DHI04] Daniel Dugger, Sharon Hollander, and Daniel C. Isaksen. Hypercovers and simplicial presheaves. *Math. Proc. Cambridge Philos. Soc.*, 136(1):9–51, 2004.
- [Gil01] Stefan Gille. A note on the Witt group of \mathbb{P}^n . Math. Z., 237(3):601–619, 2001.
- [Hau22] Olivier Haution. Motivic Pontryagin classes and hyperbolic orientations. *Preprint*, 2022. arXiv:2203.06017.
- [Hir03] Philip S. Hirschhorn. *Model categories and their localizations*, volume 99 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2003.
- [Hov99] Mark Hovey. *Model categories*, volume 63 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1999.
- [Lev19] Marc Levine. Motivic Euler characteristics and Witt-valued characteristic classes. Nagoya Math. J., 236:251–310, 2019.
- [Mor04] Fabien Morel. An introduction to \mathbb{A}^1 -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441. Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.
- [MV99] Fabien Morel and Vladimir Voevodsky. A¹-homotopy theory of schemes. Inst. Hautes Études Sci. Publ. Math., (90):45–143 (2001), 1999.
- [Nen09] Alexander Nenashev. On the Witt groups of projective bundles and split quadrics: geometric reasoning. J. K-Theory, 3(3):533–546, 2009.
- [PPR09] Ivan Panin, Konstantin Pimenov, and Oliver Röndigs. On Voevodsky's algebraic K-theory spectrum. In Algebraic topology, volume 4 of Abel Symp., pages 279–330. Springer, Berlin, 2009.
- [PW18] Ivan Panin and Charles Walter. On the motivic commutative ring spectrum **BO**. Algebra i Analiz, 30(6):43–96, 2018.
- [Rio10] Joël Riou. Algebraic K-theory, \mathbf{A}^1 -homotopy and Riemann-Roch theorems. J. Topol., 3(2):229-264, 2010.
- [Wal03] Charles Walter. Grothendieck-Witt groups of projective bundles. Preprint available at http://www.math.uiuc.edu/K-theory/0644/, 2003.

Email address: olivier.haution at gmail.com

Mathematisches Institut, Ludwig-Maximilians-Universität München, Theresienstr. 39, D-80333 München, Germany