

Low c -Differential Uniformity of the Swapped Inverse Function in Odd Characteristic

Jaeseong Jeong¹, Namhun Koo², Soonhak Kwon¹

Email: wotjd012321@naver.com, nhkoo@ewha.ac.kr, shkwon@skku.edu

¹Applied Algebra and Optimization Research Center, Sungkyunkwan University, Suwon, Korea

²Institute of Mathematical Sciences, Ewha Womans University, Seoul, Korea

Abstract

The study of Boolean functions with low c -differential uniformity has become recently an important topic of research. However, in odd characteristic case, there are not many results on the (c -)differential uniformity of functions that are not power functions. In this paper, we investigate the c -differential uniformity of the swapped inverse functions in odd characteristic, and show that their c -differential uniformities are at most 6 except for some special case.

Keywords. c -Differential Uniformity, Differential Uniformity, Permutation,

Mathematics Subject Classification(2020) 94A60, 06E30

1 Introduction

In this paper, we mainly focus on the finite fields in odd characteristic. Throughout this paper, let

- \mathbb{F}_{p^n} be the finite field of p^n elements and $\mathbb{F}_{p^n}^*$ and $\mathbb{F}_{p^n}^* = \mathbb{F}_{p^n} \setminus \{0\}$ be the multiplicative group, where p is an odd prime.
- Inv be the multiplicative inverse function on \mathbb{F}_{p^n} such that $Inv(x) = x^{-1}$ for all $x \in \mathbb{F}_{p^n}^*$ and $Inv(0) = 0^{-1} = 0$.

Recently, Ellinsen et. al. [6] introduced a new concept c -differential uniformity, which is useful to estimate resistance against some variants of differential attack [3].

Definition 1.1. (*[6]*) Let $F : \mathbb{F}_{p^n} \rightarrow \mathbb{F}_{p^n}$ be a function and $c \in \mathbb{F}_{p^n}$.

(i) We denote the c -differential of F by ${}_cD_aF(x) = F(x+a) - cF(x)$.

(ii) Let $a, b \in \mathbb{F}_{p^n}$. We denote ${}_c\Delta_F(a, b)$ by the number of solutions in \mathbb{F}_{p^n} of ${}_cD_aF(x) = b$.

(iii) The c -differential uniformity of F is defined by ${}_c\Delta_F = \max\{{}_c\Delta_F(a, b) : a, b \in \mathbb{F}_{p^n} \text{ and } a \neq 0 \text{ if } c = 1\}$.

(iv) F is called a perfect c -nonlinear(PcN) function if ${}_c\Delta_F = 1$.

(v) F is called a almost perfect c -nonlinear(APcN) function if ${}_c\Delta_F = 2$.

Functions with low c -differential uniformity provide good resistance to differential attacks of this type. Therefore, finding functions with low c -differential uniformity has been a good research topic, and many new classes of functions with low c -differential uniformity were proposed. Please see [1, 2, 7, 11, 16, 19–24, 26] for some new progress.

In even characteristic, it is known that $Inv \circ (0, 1)$ has good cryptographic properties, for example, low differential uniformity [15], high nonlinearity [15], low boomerang uniformity [14], and low differential-linear uniformity [10]. Very recently, the c -differential uniformity and the c -boomerang uniformity are investigated in [19].

However, the c -differential uniformity of $Inv \circ (0, 1)$ in odd characteristic has not been investigated, and the author of [19] suggested that it would be a good research topic. In this paper, we investigate the c -differential uniformity of the permutation obtained by swapping two points from Inv , which can be expressed as $Inv \circ (\alpha, \beta)$ with $\alpha \neq \beta \in \mathbb{F}_{p^n}$. In [11], we show that the c -differential uniformity is not preserved under affine equivalence but is preserved under some specific affine equivalence in even characteristic. The next proposition shows that the c -differential uniformity is also preserved under this specific affine equivalence in odd characteristic.

Proposition 1.2. *We call an affine permutation of the form $A(x) = a_1x + a_0$ where $a_0 \in \mathbb{F}_{p^n}$ and $a_1 \in \mathbb{F}_{p^n}^*$ an affine permutation of degree one. We call two functions F and F' are affine equivalent of degree one if $F = A_1 \circ F' \circ A_2$ where A_1 and A_2 are affine permutations of degree one. If two functions F and F' defined on \mathbb{F}_{p^n} are affine equivalent of degree one, then ${}_c\Delta_F = {}_c\Delta_{F'}$.*

Proof. We write $F = A_1 \circ F' \circ A_2$ and $F'' = F' \circ A_2$, where A_1 and A_2 are affine permutations of degree one. F' and F'' have the same c -differential uniformity in even characteristic [7], and the proof in [7] can be naturally extended for odd characteristic case. It remains to show that $F = A_1 \circ F''$ and F'' has the same c -differential uniformity. We denote $A_1(x) = a_1x + a_0$ where $a_1 \in \mathbb{F}_{p^n}^*$ and $a_0 \in \mathbb{F}_{p^n}$. Let $a, b \in \mathbb{F}_{p^n}$. Then ${}_cD_aF(x) = b$ implies that

$$\begin{aligned} b &= F(x+a) - cF(x) = (A_1 \circ F'')(x+a) - c(A_1 \circ F'')(x) = A_1(F''(x+a)) - cA_1(F''(x)) \\ &= a_1F''(x+a) + a_0 - c(a_1F''(x) + a_0) = a_1(F''(x+a) - cF''(x)) + a_0(1-c). \end{aligned}$$

So we have ${}_cD_aF''(x) = a_1^{-1}(b + a_0(c-1))$ and hence ${}_c\Delta_F(a, b) = {}_c\Delta_{F''}(a, a_1^{-1}(b + a_0(c-1)))$. Since the map from b to $a_1^{-1}(b + a_0(c-1))$ is bijective for all $c \in \mathbb{F}_{p^n}$, we obtain ${}_c\Delta_F = {}_c\Delta_{F''}$. \square

If $F' = Inv \circ (\alpha, \beta)$, $A(x) = \alpha x$ where $\alpha \in \mathbb{F}_{p^n}^*$ and $\beta \in \mathbb{F}_{p^n}$, then $F = Inv \circ (1, \alpha^{-1}\beta) = A \circ F' \circ A$ and F' have the same c -differential uniformity for all $c \in \mathbb{F}_{p^n}$ by Proposition 1.2. Therefore, in this paper, we investigate the c -differential uniformity of $Inv \circ (1, \gamma)$ where $\gamma \in \mathbb{F}_{p^n} \setminus \{1\}$.

For even characteristic case, there are many non-power functions with low differential uniformity (see [11] and references therein). The situation for odd characteristic case is quite different. That is, though there are many functions with low differential uniformity (see [4, 5, 8, 9, 12, 13, 18, 26–28] and the references therein), most of them are power functions. Also, to the best of our knowledge, there is no published article on the differential uniformity of $Inv \circ (1, \gamma)$ in odd characteristic. Therefore, we will also study the special case $c = 1$, i.e., the differential uniformity of $Inv \circ (1, \gamma)$.

The rest of this paper is organized as follows. In section 2, we introduce some preliminaries which are necessary in our subsequent sections. In section 3, we investigate the c -differential uniformity of $Inv \circ (0, 1)$ in odd characteristic, i.e., the case $\gamma = 0$. In section 4, we investigate the c -differential uniformity of $Inv \circ (1, \gamma)$ with $\gamma \neq 0, 1$. In section 5, we give a concluding remark.

2 Some Previous Results

In this section, we recall some previous results. First we briefly mention the c -differential uniformity of Inv and $Inv \circ (0,1)$ in even characteristic. It is known [6] that Inv is APcN if and only if $\text{Tr}(c) = \text{Tr}(\frac{1}{c}) = 1$, otherwise ${}_c\Delta_{Inv} = 3$, where $p = 2$ with $c \neq 0, 1$ and Tr is the field trace. It is also known [19] that ${}_c\Delta_{Inv \circ (0,1)} = 4$ when $n \geq 4$. Hence we can see that the swapped inverse function has slightly higher c -differential uniformity than the inverse function when $p = 2$.

For odd characteristic case, if $c \neq 0, 1$, then Inv is APcN if and only if $c \in \{4, 4^{-1}\}$ or both $c^2 - 4c$ and $1 - 4c$ are not squares in \mathbb{F}_{p^n} , otherwise, ${}_c\Delta_{Inv} = 3$ (see [6]). The differential uniformity of Inv is as follows [5, 8] (as in usual notation, we use the notation $D_a F(x) = {}_1D_a F(x)$ and $\Delta_F(a, b) = {}_1\Delta_F(a, b)$, throughout this paper).

$$\Delta_{Inv} = \begin{cases} 2 & \text{if } p^n \equiv 2 \pmod{3}, \\ 3 & \text{if } p = 3, \\ 4 & \text{if } p^n \equiv 1 \pmod{3}. \end{cases}$$

To summarize, in odd characteristic, one has $2 \leq \Delta_{Inv} \leq 4$ and $2 \leq {}_c\Delta_{Inv} \leq 3$ where $c \neq 0, 1$. We will later show that the swapped inverse function has slightly higher (c -)differential uniformities than the inverse function in odd characteristic. For the details, see section 3 and section 4.

The quadratic character in \mathbb{F}_{p^n} is defined as follows

$$\chi(x) = \begin{cases} 1 & \text{if } x \in \mathbb{F}_{p^n}^* \text{ is a square,} \\ -1 & \text{if } x \in \mathbb{F}_{p^n}^* \text{ is a nonsquare,} \\ 0 & \text{if } x = 0. \end{cases}$$

It is known that $\chi(x) = x^{\frac{p^n-1}{2}}$. The followings are also well known and useful for our results.

$$\chi(-1) = \begin{cases} 1 & \text{if } p^n \equiv 1 \pmod{4}, \\ -1 & \text{if } p^n \equiv 3 \pmod{4}, \end{cases} \quad \chi(5) = \begin{cases} 1 & \text{if } p^n \equiv \pm 1 \pmod{5}, \\ -1 & \text{if } p^n \equiv \pm 2 \pmod{5}, \end{cases}$$

$$\chi(-3) = \begin{cases} 1 & \text{if } p^n \equiv 1 \pmod{3}, \\ -1 & \text{if } p^n \equiv 2 \pmod{3}, \end{cases} \quad \chi(2) = \begin{cases} 1 & \text{if } p^n \equiv \pm 1 \pmod{8}, \\ -1 & \text{if } p^n \equiv \pm 3 \pmod{8}. \end{cases}$$

We also denote \sqrt{x} by any square root of x in \mathbb{F}_{p^n} if $\chi(x) = 1$ throughout this paper. Next we introduce a well-known lemma about the number of solutions of quadratic equations.

Lemma 2.1. *Let $a_2 \in \mathbb{F}_{p^n}^*$ and $a_1, a_0 \in \mathbb{F}_{p^n}$ and $D = a_1^2 - 4a_0a_2$. Then,*

$$\#\{x \in \mathbb{F}_{p^n} : a_2x^2 + a_1x + a_0 = 0\} = \begin{cases} 2 & \text{if } \chi(D) = 1, \\ 1 & \text{if } \chi(D) = 0, \\ 0 & \text{if } \chi(D) = -1. \end{cases}$$

The following lemma is a generalization of Lemma 2.6 in [11] to the odd characteristic case. This lemma is very simple and useful to our results.

Lemma 2.2. *Let $c \neq 0$. Then ${}_c\Delta_F(a, b) = {}_{c^{-1}}\Delta_F(-a, -bc^{-1})$ for all $a \in \mathbb{F}_{p^n}$ and $b \in \mathbb{F}_{p^n}$, and hence ${}_c\Delta_F = {}_{c^{-1}}\Delta_F$.*

Proof. Let $a \in \mathbb{F}_{p^n}$ and $b \in \mathbb{F}_{p^n}$. Then, it is clear that x is a solution of ${}_cD_aF(x) = b$ if and only if $x + a$ is a solution of ${}_{c^{-1}}D_{-a}(x) = -bc^{-1}$. Hence we obtain ${}_c\Delta_F(a, b) = {}_{c^{-1}}\Delta_F(-a, -bc^{-1})$. Therefore, we get ${}_c\Delta_F = {}_{c^{-1}}\Delta_F$. \square

3 The c -differential uniformity of $Inv \circ (0, 1)$

Let $F = Inv \circ (0, 1)$. Since F is permutation, we have ${}_c\Delta_F(0, b) = 1$ for all $b \in \mathbb{F}_{p^n}$ and $c \in \mathbb{F}_{p^n} \setminus \{1\}$, and we also have ${}_0\Delta_F(a, b) = 1$ for all $a, b \in \mathbb{F}_{p^n}$. Hence, throughout this section we assume that

$$F = Inv \circ (0, 1), \quad a \neq 0, \quad c \neq 0. \quad (1)$$

We also denote throughout this section that

$$P = \{x \in \mathbb{F}_{p^n} : F(x) \neq Inv(x)\} = \{0, 1\}, \quad P_a = P \cup \{x - a : x \in P\} = \{0, 1, -a, 1 - a\}.$$

It is easy to see that

$${}_c\Delta_F(a, b) = \#\{x \in P_a : {}_cD_aF(x) = b\} + \#\{x \in \mathbb{F}_{p^n} \setminus P_a : {}_cD_aF(x) = b\}.$$

The following lemma characterizes the case $\#\{x \in \mathbb{F}_{p^n} \setminus P_a : D_aF(x) = b\} = 2$.

Lemma 3.1. *Under the same assumption as in (1) let $a \in \mathbb{F}_{p^n}^*$ and $b \in \mathbb{F}_{p^n}$. Then ${}_cD_aF(x) = b$ has at most two solutions in $\mathbb{F}_{p^n} \setminus P_a$. Furthermore, ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ if and only if $(ab + c - 1)^2 - 4abc$ is a nonzero square and $b \neq 0$, $(b + c)a + b + c - 1 \neq 0$, $(1 - b)a + b + c - 1 \neq 0$.*

Proof. If $x \in \mathbb{F}_{p^n} \setminus P_a$ then $b = {}_cD_aF(x) = {}_cD_aInv(x)$ implies that $bx^2 + (ab + c - 1)x + ac = 0$ which has at most two solutions. Hence ${}_cD_aF(x) = b$ has at most two solutions in $\mathbb{F}_{p^n} \setminus P_a$.

${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ implies that $bx^2 + (ab + c - 1)x + ac = 0$ has two solutions which is equivalent to $(ab + c - 1)^2 - 4abc$ is a nonzero square and $b \neq 0$ by Lemma 2.1. If there is a solution $x_0 \in P_a$ of $bx^2 + (ab + c - 1)x + ac = 0$ then x_0 cannot be a solution of ${}_cD_aF(x) = b$. So we require that $bx^2 + (ab + c - 1)x + ac = 0$ has no solutions in P_a . We check this by substituting every element in P_a to $bx^2 + (ab + c - 1)x + ac = 0$.

- If $x = 0$ or $x = -a$ then we have $a = 0$ which is a contradiction to $a \neq 0$.
- If $x = 1$ then we have $(b + c)a + b + c - 1 = 0$.
- If $x = 1 - a$ then we have $(1 - b)a + b + c - 1 = 0$.

Thus $bx^2 + (ab + c - 1)x + ac = 0$ has no solutions in P_a if and only if $(b + c)a + b + c - 1 \neq 0$ and $(1 - b)a + b + c - 1 \neq 0$.

If all conditions in this theorem are satisfied, then $bx^2 + (ab + c - 1)x + ac = 0$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$. Since all solutions are not contained in P_a , they are also solutions of ${}_cD_aF(x) = b$. \square

In the following lemma, we provide a lower bound on ${}_c\Delta_F$ where $c \neq 0$. Unfortunately, the following lemma shows that F is not APcN for all $c \neq 0$, for every odd prime p and $n > 0$.

Lemma 3.2. *Under the same assumption as in (1), if $p^n > 5$ then ${}_c\Delta_F \geq 3$.*

Proof. Let $b = a^{-1} - c = {}_cD_aF(0)$. Then ${}_cD_aF(x) = b$ has at least one solution $x = 0$ in P_a . We investigate required conditions that ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ using Lemma 3.1. We require that $(ab + c - 1)^2 - 4abc = (c - ca)^2 - 4c(1 - ca) = c^2a^2 + 2c^2a + c^2 - 4c = c^2(a + 1)^2 - 4c$ is a nonzero square. And we have $(b + c)a + b + c - 1 = a^{-1} \cdot a + a^{-1} - 1 = a^{-1}$ and $(1 - b)a + b + c - 1 = (c + 1 - a^{-1}) \cdot a + a^{-1} - 1 = a^{-1}((c + 1)a^2 - 2a + 1)$. So we require $a \neq 0$ and $(c + 1)a^2 - 2a + 1 \neq 0$. Since $(c + 1)a^2 - 2a + 1 = 0$ has no solutions if $(-2)^2 - 4(c + 1) = -4c$ is not a square by Lemma 2.1, it is enough to check $-c$ is not a square.

(i) We assume that $-c$ is a square.

- If $c \neq -4$, then we set $a = -1$. Then $c^2(a + 1)^2 - 4c = -4c$ is a nonzero square and $(c + 1)a^2 - 2a + 1 = c + 4 \neq 0$.
- If $c = -4$ and $p \neq 3, 5, 7$, then we choose $a = -\frac{1}{4} = c^{-1}$. Then $c^2(a + 1)^2 - 4c = 25$ is a nonzero square and $(c + 1)a^2 - 2a + 1 = -\frac{3}{16} - \frac{1}{2} + 1 = \frac{21}{16} \neq 0$.
- If $c = -4$ and $p = 3, 5, 7$, then $c^2(a + 1)^2 - 4c = 16(a + 1)^2 + 16 = (a + 1)^2 + 1$ when $p = 3, 5$ and $c^2(a + 1)^2 - 4c = 2((a + 1)^2 + 1)$ when $p = 7$. Since 2 is a square for every n when $p = 7$, it is enough to show that there is $\alpha_0 \in \mathbb{F}_{p^n} \setminus \mathbb{F}_p$ such that both α_0 and $\alpha_0 + 1$ are squares. We denote $\mathbb{F}_{p^n} = \mathbb{F}_p[X]/\langle f(X) \rangle$ for some $f(X)$ and $f(g) = 0$ for some $g \in \mathbb{F}_{p^n}$. Then \mathbb{F}_{p^n} is partitioned by subsets of the form $\{x + \alpha : \alpha \in \mathbb{F}_p\}$ where $x \in \mathbb{F}_{p^n} \cdot g$. If all such subsets have at most $\frac{p-1}{2}$ squares then there are at most $\frac{p-1}{2p}(p^n - p)$ which is a contradiction to the fact that there are $\frac{1}{2}(p^n - p)$ squares in $\mathbb{F}_{p^n} \setminus \mathbb{F}_p$. Hence there is $\{x + \alpha : \alpha \in \mathbb{F}_p\} \neq \mathbb{F}_p$ containing $\frac{p+1}{2}$ squares. If there are at least $\frac{p+1}{2}$ squares in $\{x_0 + \alpha : \alpha \in \mathbb{F}_p\}$ for some $x_0 \in \mathbb{F}_{p^n} \cdot g$, then there is $\alpha_0 \in \{x_0 + \alpha : \alpha \in \mathbb{F}_p\}$ such that both α_0 and $\alpha_0 + 1$ are squares. Then there is $a \in \mathbb{F}_{p^n}$ such that $\alpha_0 = (a + 1)^2$, and then $(a + 1)^2 + 1 = \alpha_0 + 1$ is also a square. Since $\alpha_0 \notin \mathbb{F}_p$ and hence $a \notin \mathbb{F}_p$. When $p = 3$ we have $(c + 1)a^2 - 2a + 1 = a + 1 \neq 0$. When $p = 5$ we have $(c + 1)a^2 - 2a + 1 = 2(a + 1)(a - 2) \neq 0$. When $p = 7$ we have $(c + 1)a^2 - 2a + 1 = 4(a + 2)(a + 1) \neq 0$.

(ii) We assume that $-c$ is not a square. As mentioned above, we only require to check that there is $a \in \mathbb{F}_{p^n}$ such that $c^2(a + 1)^2 - 4c$ is a nonzero square.

- If -1 is a square, then one of $c - 4$ and $-c^2 + 4c = -c(c - 4)$ is a square.
 - If $c - 4$ is a square then $4(c - 4) = 4c - 16$ is also a square. If we choose $a \in \mathbb{F}_{p^n}$ with $c^2(a + 1)^2 = 4c - 16$ then $c^2(a + 1)^2 - 4c = -16$ is a nonzero square.
 - If $-c^2 + 4c$ is a square then we choose $a \in \mathbb{F}_{p^n}$ with $c^2(a + 1)^2 = -c^2 + 4c$. Then $c^2(a + 1)^2 - 4c = -c^2$ is a nonzero square.
- If -1 is not a square, then c is a square. Then one of $2c$ and $-2c$ is a square.
 - If $2c$ is a square then $8c$ is a square. If we choose $a \in \mathbb{F}_{p^n}$ with $c^2(a + 1)^2 = 8c$ then $c^2(a + 1)^2 - 4c = 4c$ is a nonzero square.
 - If $-2c$ is a square then $-8c$ is a square. In this case, $p = 3$ or one of 3 and -3 is a nonzero square.

- * If $p = 3$ and $n > 1$ then \mathbb{F}_{p^n} is partitioned by p^{n-1} subsets of the form $\{x + \alpha c : \alpha \in \mathbb{F}_p\}$. By similar argument with the case $c = -4$ and $p \in \{3, 5, 7\}$ above, there is a subset $\{x + \alpha c : \alpha \in \mathbb{F}_p\} \neq \{\alpha c : \alpha \in \mathbb{F}_p\}$ containing 2 squares. Then there is $\alpha_0 \in \{x_0 + \alpha : \alpha \in \mathbb{F}_p\}$ such that both α_0 and $\alpha_0 + c$ are nonzero squares. We choose $a \in \mathbb{F}_{p^n}$ such that $c^2(a+1)^2 = \alpha_0 + c$, then $c^2(a+1)^2 - 4c = \alpha_0$ is also a nonzero square.
- * If 3 is a nonzero square then we choose $a \in \mathbb{F}_{p^n}$ satisfying $c^2(a+1)^2 = -2c$. Then $c^2(a+1)^2 - 4c = -6c$ is also a nonzero square.
- * If -3 is a nonzero square then -12 is also a square. If we choose a satisfying $c^2(a+1)^2 = -8c$ then $c^2(a+1)^2 - 4c = -12c$ is also a nonzero square.

Therefore, there is $a \in \mathbb{F}_{p^n}^*$ such that ${}_c D_a(x) = a^{-1} - c$ has two solutions in $\mathbb{F}_{2^n} \setminus P_a$, and hence ${}_c \Delta_F(a, a^{-1} - c) \geq 3$ for all $p^n > 7$. When $p^n = 7$ we already show that there is $a \in \mathbb{F}_7$ such that ${}_c \Delta(a, a^{-1} - c) \geq 3$ for all $c \neq -4$. Since $-4^{-1} = 5 \neq -4$, we have ${}_5 \Delta(a, a^{-1} - c) \geq 3$ and hence ${}_{-4} \Delta(-a, c^{-1}(c - a^{-1})) = {}_5 \Delta(a, a^{-1} - c) \geq 3$, by Lemma 2.2. Therefore, we have ${}_c \Delta_F \geq 3$ for all $p^n > 5$, which completes the proof. \square

We obtain $\#P_a < 4$ only when $a = \pm 1$. In the following lemma we investigate ${}_c \Delta_F(a, b)$ when $a = \pm 1$.

Lemma 3.3. *Under the same assumption as in (1) let $p^n > 5$. Then ${}_c \Delta_F(1, b) \leq 3$ for all $b \in \mathbb{F}_{p^n} \setminus \{2^{-1}\}$ and $c \in \mathbb{F}_{p^n}^* \setminus \{-2^{-1}\}$, and ${}_c \Delta_F(-1, b) \leq 3$ for all $b \in \mathbb{F}_{p^n} \setminus \{1\}$ and $c \in \mathbb{F}_{p^n}^* \setminus \{-2\}$. Furthermore,*

$${}_{-2^{-1}} \Delta_F(1, 2^{-1}) = {}_{-2} \Delta_F(-1, 1) = \begin{cases} 3 & \text{if } p^n \equiv \pm 3 \pmod{8}, \\ 5 & \text{if } p^n \equiv \pm 1 \pmod{8}. \end{cases}$$

Proof. If $a = 1$ then $P_a = \{0, 1, -1\}$ and then we have ${}_c D_1 F(0) = F(1) - cF(0) = -c$, ${}_c D_1 F(1) = F(2) - cF(1) = 2^{-1}$ and ${}_c D_1(-1) = F(0) - cF(-1) = 1 + c$.

- ${}_c D_1 F(0) = {}_c D_1 F(1) \Leftrightarrow -c = 2^{-1}$: Then we have $c = -2^{-1}$.
- ${}_c D_1 F(0) = {}_c D_1 F(-1) \Leftrightarrow -c = 1 + c$: Then we have $c = -2^{-1}$.
- ${}_c D_1 F(1) = {}_c D_1 F(-1) \Leftrightarrow 2^{-1} = 1 + c$: Then $2c + 2 = 1$ and hence we have $c = -2^{-1}$.

So if $c = -2^{-1}$ and $b = 2^{-1}$ then ${}_c D_1 F(x) = b$ has three solutions in P_a . If $c \neq -2^{-1}$ or $b \neq 2^{-1}$ then ${}_c D_1 F(x) = b$ has at most one solution, and hence we have ${}_c \Delta_F(1, b) \leq 3$ for all $b \in \mathbb{F}_{p^n}$, by Lemma 3.1. Now assume that $c = -2^{-1}$ and $b = 2^{-1}$. Then ${}_{-2^{-1}} D_1 F(x) = 2^{-1}$ has three solutions $x = 0, \pm 1$ in P_a . If $a = 1$, $b = 2^{-1}$ and $c = -2^{-1}$ in Lemma 3.1 then

- $(ab + c - 1)^2 - 4abc = (-1)^2 - 4 \cdot (-\frac{1}{4}) = 2$ is a square if and only if $p^n \equiv \pm 1 \pmod{8}$.
- $(b + c)a + b + c - 1 = -1 \neq 0$ and $(1 - b)a + b + c - 1 = -2^{-1} \neq 0$

Hence ${}_{-2^{-1}} D_1 F(x) = 2^{-1}$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ if and only if $p^n \equiv \pm 1 \pmod{8}$, by Lemma 3.1.

By Lemma 2.2, ${}_{c^{-1}} D_{-1} F(x) = -bc^{-1}$ and ${}_c D_1 F(x) = b$ has the same number of solutions. Hence ${}_{-2} \Delta_F(-1, 1) = {}_{-2^{-1}} \Delta_F(1, 2^{-1})$, and ${}_c D_{-1} F(x) = b$ has at most three solutions for all $b \in \mathbb{F}_{p^n} \setminus \{1\}$ and $c \in \mathbb{F}_{p^n}^* \setminus \{-2\}$. \square

Theorem 3.4. *Under the same assumption as in (1) if $p^n > 5$ then $3 \leq {}_c\Delta_F \leq 5$. Moreover, ${}_c\Delta_F = 5$ if and only if at least one of the followings are satisfied :*

(i) $p^n \equiv \pm 1 \pmod{8}$ and $c \in \{-2, -2^{-1}\}$

(ii) $p^n \equiv 1 \pmod{3}$ and one of the followings holds :

- $c \in \left\{ \sqrt{-3}, \frac{1}{\sqrt{-3}} \right\}$, $\sqrt{-3} \neq 5$, and $\frac{-3+5\sqrt{-3}}{6}$ is a nonzero square.
- $c \in \left\{ -\sqrt{-3}, -\frac{1}{\sqrt{-3}} \right\}$, $\sqrt{-3} \neq -5$, and $\frac{-3-5\sqrt{-3}}{6}$ is a nonzero square.

(iii) $p^n \equiv \pm 1 \pmod{5}$ and one of the followings holds :

- $c^2 + 4c - 1 = 0$, and $7 - 30c$ is a nonzero square.
- $c^2 - 4c - 1 = 0$, and $7 - 30c^{-1}$ is a nonzero square.

Proof. The case when $a = \pm 1$ is covered by Lemma 3.3 and we have (i). Hence it remains to show (ii) and (iii) when $a \neq \pm 1$. We can see that

$$\begin{aligned} {}_cD_aF(0) &= F(a) - cF(0) = a^{-1} - c, & {}_cD_aF(1) &= F(1+a) - cF(1) = (1+a)^{-1}, \\ {}_cD_aF(-a) &= F(0) - cF(-a) = 1 + ca^{-1}, & {}_cD_aF(1-a) &= F(1) - cF(1-a) = c(a-1)^{-1}. \end{aligned} \quad (2)$$

We investigate all possible cases that ${}_cD_aF(x) = b$ has at least two solutions in P_a .

$$\begin{aligned} {}_cD_aF(0) = {}_cD_aF(1) &\Leftrightarrow ca^2 + ca - 1 = 0, & {}_cD_aF(0) = {}_cD_aF(-a) &\Leftrightarrow a = \frac{1-c}{c+1} \text{ when } c \neq \pm 1, \\ {}_cD_aF(0) = {}_cD_aF(1-a) &\Leftrightarrow ca^2 - a + 1 = 0, & {}_cD_aF(1) = {}_cD_aF(-a) &\Leftrightarrow a^2 + ca + c = 0, \\ {}_cD_aF(1) = {}_cD_aF(1-a) &\Leftrightarrow a = \frac{c+1}{1-c} \text{ when } c \neq \pm 1, & {}_cD_aF(-a) = {}_cD_aF(1-a) &\Leftrightarrow a^2 - a - c = 0. \end{aligned}$$

If ${}_cD_aF(0) = {}_cD_aF(-a) = {}_cD_aF(1) = {}_cD_aF(1-a)$ then $a = \frac{1-c}{c+1} = \frac{c+1}{1-c}$ and hence we have $c = 0$, a contradiction. Hence ${}_cD_aF(x) = b$ has at most three solutions in P_a . We investigate all possible cases that ${}_cD_aF(x) = b$ has three solutions in P_a .

- If ${}_cD_aF(0) = {}_cD_aF(1) = {}_cD_aF(-a)$ then $a = \frac{1-c}{c+1}$ is a solution of two equations $ca^2 + ca - 1 = 0$ and $a^2 + ca + c = 0$. Then we have $0 = ca^2 + ca - 1 = c \left(\frac{1-c}{c+1} \right)^2 + c \cdot \frac{1-c}{c+1} - 1 = -\frac{3c^2+1}{(c+1)^2}$ and $0 = a^2 + ca + c = \left(\frac{1-c}{c+1} \right)^2 + c \cdot \frac{1-c}{c+1} + c = \frac{3c^2+1}{(c+1)^2}$. Such c exists when -3 is a nonzero square, if and only if $p^n \equiv 1 \pmod{3}$, and then $c = \pm \frac{1}{\sqrt{-3}}$. Then $a = \frac{\sqrt{-3} \mp 1}{\sqrt{-3} \pm 1} = \frac{1 \pm \sqrt{-3}}{2}$ and $b = {}_cD_aF(0) = a^{-1} - c = \frac{2}{1 \pm \sqrt{-3}} \mp \frac{1}{\sqrt{-3}} = \frac{3 \mp \sqrt{-3}}{6}$. Since $(ab+c-1)^2 - 4abc = \left(\frac{1 \pm \sqrt{-3}}{2} \cdot \frac{3 \mp \sqrt{-3}}{6} \pm \frac{1}{\sqrt{-3}} - 1 \right)^2 \mp 4 \cdot \frac{1 \pm \sqrt{-3}}{2} \cdot \frac{3 \mp \sqrt{-3}}{6} \cdot \frac{1}{\sqrt{-3}} = \left(\frac{-3 \mp \sqrt{-3}}{6} \right)^2 + \frac{-2 \pm 2\sqrt{-3}}{3} = \frac{-3 \pm 5\sqrt{-3}}{6}$. Using $b+c = \frac{3 \mp \sqrt{-3}}{6} \mp \frac{\sqrt{-3}}{3} = \frac{1 \mp \sqrt{-3}}{2}$, we have $a(b+c) + b + c + 1 = \frac{1 \pm \sqrt{-3}}{2} \cdot \frac{1 \mp \sqrt{-3}}{2} + \frac{1 \mp \sqrt{-3}}{2} + 1 = \frac{5 \mp \sqrt{-3}}{2}$ and $(1-b)a + b + c - 1 = \frac{3 \pm \sqrt{-3}}{6} \cdot \frac{1 \pm \sqrt{-3}}{2} + \frac{1 \mp \sqrt{-3}}{2} - 1 = \frac{-3 \mp \sqrt{-3}}{6} \neq 0$. So, we have ${}_c\Delta_F(a, b) = 5$ if and only if $p^n \equiv 1 \pmod{3}$, $\frac{5 \mp \sqrt{-3}}{2} \neq 0$, and $\frac{-3 \pm 5\sqrt{-3}}{6}$ is a square. Note that all double signs are in same order in above computations this case.
- If ${}_cD_aF(0) = {}_cD_aF(1) = {}_cD_aF(1-a)$ then $a = \frac{c+1}{1-c}$ is a solution of two equations $ca^2 + ca - 1 = 0$ and $ca^2 - a + 1 = 0$. Then we have $0 = c \cdot \left(\frac{c+1}{1-c} \right)^2 + c \cdot \frac{c+1}{1-c} - 1 = \frac{c^2+4c-1}{(1-c)^2}$ and $0 = c \cdot \left(\frac{c+1}{1-c} \right)^2 - \frac{c+1}{1-c} +$

$1 = \frac{c(c^2+4c-1)}{(1-c)^2}$, and hence $c^2+4c-1 = 0$. Such c exists if $(-4)^2 - 4 \cdot (-1) = 20 = 2^2 \cdot 5$ is a square, if and only if $p = 5$ or $p^n \equiv \pm 1 \pmod{5}$. We set $b = {}_cD_aF(0) = a^{-1} - c = \frac{1-c}{c+1} - c = \frac{1-2c-c^2}{c+1}$. Using the proof of Lemma 3.2, we have $(ab+c-1)^2 - 4abc = c^2(a+1)^2 - 4c = c^2 \cdot \left(\frac{2}{1-c}\right)^2 - 4c = -\frac{4(c^3-3c^2+c)}{(c-1)^2} = -\frac{4((c-7)(c^2+4c-1)+30c-7)}{(c-1)^2} = -\frac{4(30c-7)}{(c-1)^2}$ and $(b+c)a + b + c + 1 = a^{-1} \neq 0$ and $(1-b)a+b+c-1 = a^{-1}((c+1)a^2-2a+1) = a^{-1} \left((c+1) \cdot \left(\frac{c+1}{1-c}\right)^2 - 2 \cdot \frac{c+1}{1-c} + 1 \right) = \frac{c(c^2+6c+1)}{a(c-1)^2} = \frac{2c(c+1)}{a(c-1)^2} \neq 0$ since $c \neq -1$. So, we have ${}_c\Delta_F(a, b) = 5$ if and only if $p^n \equiv 0, \pm 1 \pmod{5}$ and $7 - 30c$ is a nonzero square. If $p = 5$ then we have $c = -2$, and $7 - 30c = 2$ is a square if and only if $p^n \equiv \pm 1 \pmod{8}$ and hence this case is already covered by (i).

- If ${}_cD_aF(0) = {}_cD_aF(-a) = {}_cD_aF(1-a)$ then we have ${}_{c^{-1}}D_{-a}F(0) = {}_{c^{-1}}D_{-a}F(1) = {}_{c^{-1}}D_{-a}F(-(-a))$ by Lemma 2.2. We set $c^{-1} = \pm \frac{1}{\sqrt{-3}}$ and $-a = \frac{1-c^{-1}}{c^{-1}+1} = \frac{1 \pm \sqrt{-3}}{2}$ or equivalently $c = \pm \sqrt{-3}$ and $a = -\frac{1 \pm \sqrt{-3}}{2}$, and $b = (-a)^{-1} - c^{-1} = \frac{3 \mp \sqrt{-3}}{6}$. By the above observation and Lemma 2.2, $5 = {}_{c^{-1}}\Delta_F(-a, -bc^{-1}) = {}_c\Delta_F(a, b)$ if and only if $p^n \equiv 1 \pmod{3}$, $\frac{5 \mp \sqrt{-3}}{2} \neq 0$, and $\frac{-3 \pm 5\sqrt{-3}}{6}$ is a square.
- If ${}_cD_aF(1) = {}_cD_aF(-a) = {}_cD_aF(1-a)$ then we have ${}_{c^{-1}}D_{-a}F(0) = {}_{c^{-1}}D_{-a}F(1) = {}_{c^{-1}}D_{-a}F(1-(-a))$ by Lemma 2.2. We assume that $(c^{-1})^2 + 4c^{-1} - 1 = 0$ and $-a = \frac{c^{-1}+1}{1-c^{-1}}$ or equivalently $c^2 - 4c - 1 = 0$ and $a = \frac{1+c}{1-c}$, and $b = -a^{-1} - c^{-1} = \frac{1-2c^{-1}-(c^{-1})^2}{1+c^{-1}}$. By the above observation and Lemma 2.2, $5 = {}_{c^{-1}}\Delta_F(-a, b) = {}_c\Delta_F(a, b)$ if and only if $p \equiv 0, \pm 1 \pmod{5}$ and $7 - 30c^{-1}$ is a nonzero square. Similarly with the above case, the case $p = 5$ can be covered by (i).

Conversely, suppose each condition in (ii) and (iii) is satisfied. Using the same a and b in previous analysis, we can see that ${}_cD_aF(x) = b$ has three solutions in P_a . By Lemma 3.1, the given element is a nonzero square, then ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$. Hence we have ${}_c\Delta_F(a, b) = 5$ which completes the proof. \square

Next we further investigate the differential uniformity of F , the special case $c = 1$.

Theorem 3.5. *Let $F = \text{Inv} \circ (0, 1)$ and $p^n > 5$. Then $3 \leq \Delta_F \leq 5$. Furthermore,*

- (i) $\Delta_F = 5$ if and only if $p = 3$ and n is even.
(ii) $\Delta_F = 4$ if and only if at least one of the followings is satisfied :

- $p = 5$ and $n \equiv 0 \pmod{4}$.
- $p^n \equiv \pm 1 \pmod{5}$ with $p \neq 3$, and at least one of $\frac{-5+\sqrt{5}}{2}$ and $\frac{-5-\sqrt{5}}{2}$ is a square.
- $p^n \equiv 1 \pmod{3}$ with $p \neq 5$, and at least one of $\frac{-5+3\sqrt{-3}}{2}$ and $\frac{-5-3\sqrt{-3}}{2}$ is a nonzero square.

Proof. By Theorem 3.4, $3 \leq \Delta_F \leq 5$ is straightforward.

- (i) We first show that $D_aF(x) = b$ has at most two solutions in $\mathbb{F}_{p^n} \setminus P_a$ and hence $\Delta_F(a, b) \leq 4$, when $a \in \mathbb{F}_{p^n} \setminus \{0, \pm 1\}$ and $b \in \mathbb{F}_{p^n}$. We have the following to substitute $c = 1$ in (2)

$$\begin{aligned} D_aF(0) &= F(a) - F(0) = a^{-1} - 1, & D_aF(1) &= F(1+a) - F(1) = (1+a)^{-1}, \\ D_aF(-a) &= F(0) - F(-a) = 1 + a^{-1}, & D_aF(1-a) &= F(1) - F(1-a) = (a-1)^{-1}. \end{aligned}$$

We can see that $D_a F(x-a) = F(x) - F(x-a) = -D_{-a}(x)$. If $D_a F(0) = D_a F(-a)$ we have $1 = -1$, a contradiction, and hence we have $D_a F(0) \neq D_a F(-a)$. Similarly we have $D_a F(1) \neq D_a F(1-a)$. Hence $D_a F(u) \neq D_a F(u-a)$ for $u \in P$. If there is $b \in \mathbb{F}_{p^n}$ such that $D_a F(x) = b$ has at least three solutions in P_a , then there is $u \in P_a$ such that both u and $u-a$ are solutions of $D_a F(x) = b$, which is a contradiction. Hence $D_a F(x) = b$ has at most two solutions in P_a .

So we have $\Delta_F = 5$ if and only if $\Delta_F(1, b) = 5$ for some $b \in \mathbb{F}_{p^n}$. We have $c = 1 = -2^{-1}$ which is equivalent to $1 = (-2) \cdot (-2)^{-1} = (-2) \cdot 1 = -2$ if and only if $p = 3$. Then $3^n \equiv \pm 1 \pmod{8}$ if and only if n is even. Hence by Lemma 3.3, $\Delta_F = 5$ if and only if $p = 3$ and n is even.

(ii) We investigate all possible cases that $D_a F(x) = b$ has at least two solutions in P_a applying $D_a F(u) \neq D_a F(u-a)$ for $u \in P$. Since $D_a F(x-a) = F(x) - F(x-a) = -D_{-a}(x)$, we have $D_a F(-a) = D_a F(u-a)$ if and only if $D_{-a} F(0) = D_{-a} F(u)$ for $u \in \{1, 1-a\}$. Hence it is enough to investigate the following cases.

- $D_a F(0) = D_a F(1)$ implies $a^2 + a - 1 = 0$. $D = (-1)^2 - 4 \cdot (-1) = 5$ is a nonzero square if and only if $p \equiv \pm 1 \pmod{5}$.
- $D_a F(0) = D_a F(1-a)$ implies $a^2 - a + 1 = 0$. $D = (-1)^2 - 4 \cdot 1 = -3$ is a nonzero square if and only if $p \equiv 1 \pmod{3}$.

We set $b = D_a F(0) = a^{-1} - 1$. Applying the proof of Lemma 3.2 we have $(b+c)a + b + c - 1 = a^{-1} \neq 0$ and $(1-b)a + b + c - 1 = a^{-1}(2a^2 - 2a + 1)$.

- If $D_a F(0) = D_a F(1)$ then $a^2 = -a + 1$. Hence $0 = 2a^2 - 2a + 1 = 2(-a + 1) - 2a + 1 = -4a + 3$ implies $a = \frac{3}{4}$. Then $0 = a^2 + a - 1 = (\frac{3}{4})^2 + \frac{3}{4} - 1 = \frac{5}{16}$ requires $p = 5$.
- If $D_a F(0) = D_a F(1-a)$ then $a^2 - a = -1$. Hence we have $2a^2 - 2a + 1 = 2(a^2 - a) + 1 = -1 \neq 0$.

1. If $p = 3$ then it remains to show that $\Delta_F(a, b) \leq 3$ for all $a \in \mathbb{F}_{p^n} \setminus \{0, \pm 1\}$ and $b \in \mathbb{F}_{p^n}$ when n is odd, by Lemma 3.3. Then $a^2 - a + 1 = 0$ has one solution $a = -1$, contradicts to the assumption. Furthermore we have $3^n \equiv \pm 2 \pmod{5}$ if n is odd. So $D_a F(x) = b$ has at most one solution in P_a when $a \in \mathbb{F}_{p^n} \setminus \{0, 1\}$, and hence we have $\Delta_F(a, b) \leq 3$ for all $a \in \mathbb{F}_{p^n} \setminus \{0, 1\}$ and $b \in \mathbb{F}_{p^n}$.

2. If $p = 5$ then $a^2 + a - 1 = 0$ has one solution. It is easy to see that $a = 2$ is the only solution of $a^2 + a - 1 = 0$ and then $b = D_2 F(0) = 2^{-1} - 1 = 2$. As mentioned above, $D_2 F(x) = 2$ has two solutions $x = 0, 1$ in P_a . And then we have $(ab + c - 1)^2 - 4abc = 0$ and hence $bx^2 + (ab + c - 1)x + ac = 0$ has exactly one solution, but by the above observation $x = 1 - a = -1 \in P_a$ is the solution of $bx^2 + (ab + c - 1)x + ac = 0$ and hence it cannot be a solution of $D_2 F(x) = 2$. Hence $D_2 F(x) = 2$ has no solutions in $\mathbb{F}_{p^n} \setminus P_a$. So we have $\Delta_F(2, 2) = 2$.

If n is odd then $5^n \equiv 2 \pmod{3}$ and hence $D_a F(x) = b$ has at most one solution in P_a . By Lemma 3.3 we have $\Delta_F = 3$. If n is even then $5^n \equiv 1 \pmod{3}$ so $a^2 + a + 1 = 0$ has two solutions. Then $a^3 = -1$ and hence $a^{-1} = -a^2$. So we have $b = D_a F(0) = a^{-1} - 1 = -a^2 - 1 = -a$ then $(ab + c - 1)^2 - 4abc = a^4 + 4a^2 = -a + 4(a - 1) = 3a + 1$. Note that $(3a + 1)^2 = 9a^2 + 6a + 1 = -a^2 + a + 1 = 2$ and hence $3a + 1$ is a square root of 2. Since the multiplicative order of 2 in modulo 5 is 4, we have $1 = \chi(3a + 1) = (3a + 1)^{\frac{5^n - 1}{2}} = 2^{\frac{5^n - 1}{4}}$ if and only if $4 \mid \frac{5^n - 1}{4}$ if and only if $5^n \equiv 1 \pmod{16}$. Since the multiplicative order of 5 in modulo 16 is 4, we have $5^n \equiv 1 \pmod{16}$ if and only if $n \equiv 0 \pmod{4}$. Hence, we have $\Delta_F = 4$ if and only if $n \equiv 0 \pmod{4}$.

3. Now we investigate the case $p \neq 3, 5$.

- If $p^n \equiv \pm 1 \pmod{5}$, then 5 is a square and hence $D_a F(0) = D_a F(1)$ where $a^2 + a - 1 = 0$. Then $a = \frac{-1 \pm \sqrt{5}}{2}$ and $b = a^{-1} - 1$ and $(ab + c - 1)^2 - 4abc = a^2 + 2a - 3 = a^2 + a - 1 + a - 2 = a - 2 = \frac{-5 \pm \sqrt{5}}{2}$. Hence by Lemma 3.1 we have $\Delta_F(a, b) = 4$ if and only if at least one of $\frac{-5 \pm \sqrt{5}}{2}$ and $\frac{-5 - \sqrt{5}}{2}$ is a square. Note that $\frac{-5 \pm \sqrt{5}}{2} \neq 0$, since if $\frac{-5 \pm \sqrt{5}}{2} = 0$ then $x = \pm\sqrt{5}$ is a solution of $x^2 - x = 0$ which implies $1 = 5$ or $0 = 5$, which is a contradiction to $p \neq 5$.
- If $p^n \equiv 1 \pmod{3}$ then -3 is a square and hence $D_a F(0) = D_a F(1 - a)$ where $a^2 - a + 1 = 0$. Then $a = \frac{1 \pm \sqrt{-3}}{2}$ and $b = a^{-1} - 1$ and $(ab + c - 1)^2 - 4abc = a^2 + 2a - 3 = a^2 - a + 1 + 3a - 4 = 3a - 4 = \frac{-5 \pm 3\sqrt{-3}}{2}$. Hence by Lemma 3.1 we have $\Delta_F(a, b) = 4$ if and only if at least one of $\frac{-5 + 3\sqrt{-3}}{2}$ and $\frac{-5 - 3\sqrt{-3}}{2}$ is a nonzero square.

Conversely, suppose each condition in (ii) is satisfied. Using the same a and b in previous analysis, we can see that ${}_c D_a F(x) = b$ has two solutions in P_a . By Lemma 3.1, the given element in each case is a nonzero square, then $D_a F(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$. Hence we have $\Delta_F(a, b) = 4$ which completes the proof. \square

4 The c -differential uniformity of $Inv \circ (1, \gamma)$

Let $F = Inv \circ (1, \gamma)$ where $\gamma \in \mathbb{F}_{p^n} \setminus \{0, 1\}$. Similarly with section 3 since F is permutation, we have ${}_c \Delta_F(0, b) = 1$ for all $b \in \mathbb{F}_{p^n}$ and $c \in \mathbb{F}_{p^n} \setminus \{1\}$, and we also have ${}_0 \Delta_F(a, b) = 1$ for all $a, b \in \mathbb{F}_{p^n}$. Hence, throughout this section we assume that

$$F = Inv \circ (1, \gamma), \quad \gamma \in \mathbb{F}_{p^n} \setminus \{0, 1\}, \quad a \neq 0, \quad c \neq 0 \quad (3)$$

We also denote throughout this section that

$$P = \{x \in \mathbb{F}_{p^n} : F(x) \neq Inv(x)\} \cup \{0\} = \{0, 1, \gamma\}, \quad P_a = P \cup \{x - a : x \in P\} = \{0, 1, \gamma, -a, 1 - a, \gamma - a\}.$$

It is easy to see that

$${}_c \Delta_F(a, b) = \#\{x \in P_a : {}_c D_a F(x) = b\} + \#\{x \in \mathbb{F}_{p^n} \setminus P_a : {}_c D_a F(x) = b\}.$$

The following lemma characterizes the case $\#\{x \in \mathbb{F}_{p^n} \setminus P_a : D_a F(x) = b\} = 2$.

Lemma 4.1. *Under the same assumption as in (3) let $a \in \mathbb{F}_{p^n}^*$ and $b \in \mathbb{F}_{p^n}$. Then ${}_c D_a F(x) = b$ has at most two solutions in $\mathbb{F}_{p^n} \setminus P_a$. Furthermore, ${}_c D_a F(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ if and only if $(ab + c - 1)^2 - 4abc$ is a nonzero square and $b \neq 0$, $(b + c)a + b + c - 1 \neq 0$, $(1 - b)a + b + c - 1 \neq 0$, $(b\gamma + c)a + \gamma(b\gamma + c - 1) \neq 0$ and $(1 - b\gamma)a + \gamma(b\gamma + c - 1) \neq 0$.*

Proof. Similarly with Lemma 3.1, since $F(x) = Inv(x)$ for all $x \in \mathbb{F}_{2^n} \setminus P_a$, ${}_c D_a F(x) = b$ has at most two solutions in $\mathbb{F}_{2^n} \setminus P_a$, and has two solutions if and only if $(ab + c - 1)^2 - 4abc$ is a nonzero square and $b \neq 0$. We require that $bx^2 + (ab + c - 1)x + ac = 0$ has no solutions in P_a . We check this by substituting every element in P_a to $bx^2 + (ab + c - 1)x + ac = 0$.

- If $x = 0$ or $x = -a$ then we have $a = 0$ which is a contradiction to $a \neq 0$.
- If $x = 1$ then we have $(b + c)a + b + c - 1 = 0$.

- If $x = 1 - a$ then we have $(1 - b)a + b + c - 1 = 0$.
- If $x = \gamma$ then we have $(b\gamma + c)a + \gamma(b\gamma + c - 1) = 0$.
- If $x = \gamma - a$ then we have $(1 - b\gamma)a + \gamma(b\gamma + c - 1) = 0$.

Thus $bx^2 + (ab + c - 1)x + ac = 0$ has no solutions in P_a if and only if $(b + c)a + b + c - 1 \neq 0$, $(1 - b)a + b + c - 1 \neq 0$, $(b\gamma + c)a + \gamma(b\gamma + c - 1) \neq 0$ and $(1 - b\gamma)a + \gamma(b\gamma + c - 1) \neq 0$. Similarly with Lemma 3.1, if all conditions in this theorem are satisfied, then ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$. \square

The following lemma shows symmetry of the c -differential uniformity with respect to γ , and it is useful for our results.

Lemma 4.2. *Let $\gamma \neq 0, 1$ and $F = \text{Inv} \circ (1, \gamma)$ and $F' = \text{Inv} \circ (1, \gamma^{-1})$. Then ${}_c\Delta_{F'}(a, b) = {}_c\Delta_F(\gamma a, \gamma^{-1}b)$ for all $a, b \in \mathbb{F}_{p^n}$ and hence ${}_c\Delta_F = {}_c\Delta_{F'}$ for all $c \in \mathbb{F}_{p^n}$.*

Proof. Let $A(x) = \gamma x$. Then it is not difficult to see that $F' = A \circ F \circ A$. Then ${}_cD_aF'(x) = b$ implies

$$\begin{aligned} b &= {}_cD_aF'(x) = (A \circ F \circ A)(x + a) - c(A \circ F \circ A)(x) = (A \circ F)(\gamma x + \gamma a) - c(A \circ F)(\gamma x) \\ &= \gamma(F(\gamma x + \gamma a) - F(\gamma x)) = \gamma(F(X + \gamma a) - cF(X)) = \gamma \cdot {}_cD_{\gamma a}F(X) \end{aligned}$$

if and only if $\gamma^{-1}b = {}_cD_{\gamma a}F(X)$, where $X = \gamma x$ in above. \square

We can see that $\#P_a < 6$ if and only if $a \in \{\pm 1, \pm \gamma, \pm(\gamma - 1)\}$. We investigate the number of solutions of ${}_cD_aF(x) = b$ in P_a for this case.

Lemma 4.3. *Let $a \in \{\pm 1, \pm \gamma, \pm(\gamma - 1)\}$. Then*

- (i) ${}_cD_aF(x) = b$ has five solutions in P_a if and only if $p = 5$, $\gamma = -1$, $c = 1$ and $(a, b) \in \{(2, 3), (3, 2)\}$.
- (ii) ${}_cD_aF(x) = b$ has four solutions in P_a if and only if $p = 5$, $\gamma = -1$, $c = 1$ and $(a, b) \in \{(1, -1), (-1, 1)\}$.

Otherwise, ${}_cD_aF(x) = b$ has at most three solutions in P_a .

Proof. We investigate all possible cases, and details are in Appendix A. \square

In Lemma 3.2 we show that the c -differential uniformity of F is lower bounded by 3 for all $c \in \mathbb{F}_{p^n}^*$ when $\gamma = 0$. But the method used in the proof of Lemma 3.2 is not applicable when $\gamma \neq 0$, and it is not easy to show a lower bound of the c -differential uniformity of F . But we confirm that ${}_c\Delta_F \geq 3$ when $7 < p^n < 500$ by using SageMath.

4.1 The differential uniformity

In this subsection we investigate the differential uniformity of F , that is, the case $c = 1$. We first investigate the number of solutions of $D_aF(x) = b$ in P_a when $\#P_a = 6$.

Lemma 4.4. *Let $a \in \mathbb{F}_{p^n} \setminus \{0, \pm 1, \pm \gamma, \pm(\gamma - 1)\}$ and $b \in \mathbb{F}_{p^n}$. Then, $D_aF(x) = b$ has four solutions in P_a if and only if $\gamma = -1$ and at least one of the followings is satisfied:*

- (i) $p^n \equiv \pm 1 \pmod{5}$, $a^4 - 3a^2 + 1 = 0$ and $b = a^{-1}$,
- (ii) $p^n \equiv \pm 1 \pmod{8}$, $a^2 = 2$ and $b = a$.

Otherwise, $D_aF(x) = b$ has at most three solutions.

Proof. $D_a F(x)$ for each $x \in P_a$ is as follows:

$$\begin{aligned} D_a F(0) &= F(a) - F(0) = a^{-1}, & D_a F(-a) &= F(0) - F(-a) = a^{-1}, \\ D_a F(1) &= F(1+a) - F(1) = (a+1)^{-1} - \gamma^{-1}, & D_a F(1-a) &= F(1) - F(1-a) = \gamma^{-1} - (1-a)^{-1}, \\ D_a F(\gamma) &= F(\gamma+a) - F(\gamma) = (a+\gamma)^{-1} - 1, & D_a F(\gamma-a) &= F(\gamma) - F(\gamma-a) = 1 + (a-\gamma)^{-1}. \end{aligned}$$

We first assume that $b = D_a F(0) = D_a F(-a) = a^{-1}$. We investigate $D_a F(x) = a^{-1}$ has at least three solutions in P_a . First we investigate equivalent condition for each case that $D_a F(x) = a^{-1}$ has at least two solutions in P_a .

$$\begin{aligned} a^{-1} = D_a F(1) &\Leftrightarrow a^2 + a + \gamma = 0, & a^{-1} = D_a F(1-a) &\Leftrightarrow a^2 - a + \gamma = 0, \\ a^{-1} = D_a F(\gamma) &\Leftrightarrow a^2 + \gamma a + \gamma = 0, & a^{-1} = D_a F(\gamma-a) &\Leftrightarrow a^2 - \gamma a + \gamma = 0. \end{aligned}$$

We can easily see that if $a^{-1} = D_a F(1) = D_a F(1-a)$ or $a^{-1} = D_a F(\gamma) = D_a F(\gamma-a)$ then we have $a = 0$, a contradiction. Hence $D_a F(x) = a^{-1}$ has at most four solutions. If $a^{-1} = D_a F(1) = D_a F(\gamma-a)$ then we have $\gamma = -1$ and hence $a^2 + a - 1 = 0$, such a exists if and only if $1^2 - 4 \cdot (-1) = 5$ is a square which is equivalent to $p^n \equiv 0, \pm 1 \pmod{5}$. But if $p = 5$ then we have $a = 2 = 1 - \gamma$ which contradicts to assumption, and hence we have $p \neq 5$. Similarly, if $a^{-1} = D_a F(\gamma) = D_a F(1-a)$ we also have $\gamma = -1$ and hence $a^2 - a - 1 = 0$, such a exists if and only if $(-1)^2 - 4 \cdot (-1) = 5$ is a square, and we have $p^n \equiv \pm 1 \pmod{5}$. Hence, $D_a F(x) = a^{-1}$ has four solutions if and only if $p^n \equiv \pm 1 \pmod{5}$, $\gamma = -1$ and $0 = (a^2 + a - 1)(a^2 - a - 1) = a^4 - 3a^2 + 1$. Otherwise, $D_a F(x) = a^{-1}$ has at most three solutions. Conversely, if $p \equiv \pm 1 \pmod{5}$ then there exists $a \in \mathbb{F}_{p^n}$ with $a^2 + a - 1 = 0$ or $a^2 - a - 1 = 0$ and then by the above analysis we can see that $D_a F(x) = a^{-1}$ has four solutions in P_a when $\gamma = -1$.

Next we investigate the case $b \neq a^{-1}$. In this case, $D_a F(x) = b$ has four solutions in P_a if and only if $b = D_a F(1) = D_a F(\gamma) = D_a F(1-a) = D_a F(\gamma-a)$. First we investigate equivalent condition for each case that $D_a F(x) = b$ has at least two solutions in $\{1, \gamma, 1-a, \gamma-a\}$.

$$\begin{aligned} D_a F(1) = D_a F(\gamma) &\Leftrightarrow a^2 + (\gamma+1)a + 2\gamma = 0, & D_a F(1) = D_a F(\gamma-a) &\Leftrightarrow a(\gamma+1)(a-\gamma+1)a = 0, \\ D_a F(1) = D_a F(1-a) &\Leftrightarrow a^2 + \gamma - 1 = 0, & D_a F(\gamma) = D_a F(1-a) &\Leftrightarrow a(\gamma+1)(a+\gamma-1)a = 0, \\ D_a F(\gamma) = D_a F(\gamma-a) &\Leftrightarrow a^2 + \gamma(1-\gamma) = 0, & D_a F(1-a) = D_a F(\gamma-a) &\Leftrightarrow a^2 - (\gamma+1)a + 2\gamma = 0. \end{aligned}$$

From $a^2 + (\gamma+1)a + 2\gamma = 0$ and $a^2 - (\gamma+1)a + 2\gamma = 0$ we have $\gamma = -1$. Then we have $a^2 = 2$, such a exists if and only if $p^n \equiv \pm 1 \pmod{8}$. Then we have $b = D_a F(1) = \frac{1}{a+1} + 1 = \frac{a+2}{a+1} = \frac{a^2+a-2}{a^2-1} = a$. Conversely, if $p^n \equiv 1 \pmod{8}$ then there exists $a \in \mathbb{F}_{p^n}$ with $a^2 = 2$ and then by the above analysis we can see that $D_a F(x) = a$ has four solutions in P_a , when $\gamma = -1$. \square

Next we show the main theorem of this subsection.

Theorem 4.5. *Let $F = \text{Inv} \circ (1, \gamma)$ with $\gamma \neq 0, 1$. Then,*

- (i) $\Delta_F = 7$ if and only if $\gamma = -1$, $p = 5$ and n is even.
 - (ii) $\Delta_F = 6$ if and only if $\gamma = -1$, and $p^n \equiv 1 \pmod{8}$ or $p^n \equiv 1, 4 \pmod{15}$.
- Otherwise, $\Delta_F \leq 5$.

Proof. Using Lemma 4.3 and Lemma 4.4 we investigate all the cases that $D_a F(x) = b$ has at least four solutions, which requires $\gamma = -1$. By Lemma 4.1 with $c = 1$ and $\gamma = -1$, $D_a F(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ if and only if $(ab)^2 - 4ab$ is a nonzero square, $(b+1)a + b = ab + a + b \neq 0$ and $(1-b)a + b = a + b - ab \neq 0$.

- If $p = 5$ and $(a, b) \in \{(2, 3), (3, 2)\}$ then $D_a F(x) = b$ has five solutions in P_a . Then $(ab)^2 - 4ab = 1^2 - 4 \cdot 1 = -3$ is a square if and only if $5^n \equiv 1 \pmod{3}$ which is equivalent to n is even. And, we have $ab + a + b = 1 \neq 0$ and $a + b - ab = -1 \neq 0$. Hence we have $\Delta_F(a, b) = 7$ if n is even, $\Delta_F(a, b) = 5$ if n is odd.
- If $p = 5$ and $(a, b) \in \{(1, -1), (-1, 1)\}$ then $D_a F(x) = b$ has four solutions in P_a . Then we have $(ab)^2 - 4ab = 1^2 + 4 \cdot 1 = 5 = 0$ and hence we have $\Delta_F(a, b) \leq 5$ in this case.
- If $p^n \equiv \pm 1 \pmod{5}$ and $a^4 - 3a^2 + 1 = (a^2 + a - 1)(a^2 - a - 1) = 0$ then $D_a(x) = a^{-1}$ has four solutions in P_a . Then we require $(ab)^2 - 4ab = 1^2 - 4 \cdot 1 = -3$ is a nonzero square if and only if $p^n \equiv 1 \pmod{3}$. If $a^2 + a - 1 = 0$ then $ab + a + b = a^{-1}(a^2 + a + 1) = 2a^{-1} \neq 0$ and $a + b - ab = a^{-1}(a^2 - a + 1) = 2a^{-1}(1 - a) \neq 0$ since $a \neq 1$. Similarly, if $a^2 - a - 1 = 0$ then $ab + a + b \neq 0$ and $a + b - ab \neq 0$. Hence we have $\Delta_F(a, a^{-1}) = 6$ where $a^4 - 3a^2 + 1 = 0$, when $p^n \equiv \pm 1 \pmod{5}$ and $p^n \equiv 1 \pmod{3}$, or equivalently $p^n \equiv 1, 4 \pmod{15}$.
- If $p^n \equiv \pm 1 \pmod{8}$ and $a^2 = 2$ then $D_a(x) = a$ has four solutions in P_a . Then we have $(ab)^2 - 4ab = a^4 - 4a^2 = 4 - 8 = -4$ is a square if and only if $p^n \equiv 1 \pmod{4}$. Since $a \neq 1$ from $(\pm 1)^2 = 1 \neq 2$, we have $ab + a + b = a^2 + 2a = 2(a + 1) \neq 0$ and $a + b - ab = 2a - 2 = 2(a - 1) \neq 0$. Hence we have $\Delta_F(a, a) = 6$ where $a^2 = 2$, when $p^n \equiv \pm 1 \pmod{8}$ and $p^n \equiv 1 \pmod{4}$, or equivalently $p^n \equiv 1 \pmod{8}$.

Otherwise, $D_a F(x) = b$ at most three solutions in P_a by Lemma 4.3 and Lemma 4.4, and hence we have $\Delta_F \leq 5$ by Lemma 4.1. \square

4.2 The c -differential uniformity with $c \neq 0, 1$

In the above subsection we investigate the differential uniformity of $Inv \circ (1, \gamma)$. In this subsection we investigate the c -differential uniformity of $Inv \circ (1, \gamma)$ where $c \neq 0, 1$. First we investigate the number of solutions of ${}_c D_a F(x) = b$ in P_a when $\#P_a = 6$.

Lemma 4.6. *Let $F = Inv \circ (1, \gamma)$ with $\gamma \neq 0, 1$ and $c \neq 0, 1$. Let $a \in \mathbb{F}_{p^n} \setminus \{0, \pm 1, \pm \gamma, \pm(\gamma - 1)\}$ and $b \in \mathbb{F}_{p^n}$. Then, ${}_c D_a F(x) = b$ has four solutions in P_a if and only if $\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 0$, $c \neq -1$, $a = \frac{\gamma^2 - c}{\gamma + c}$ and $b = \frac{1 - c}{1 + \gamma}$. Otherwise, ${}_c D_a F(x) = b$ has at most three solutions in P_a .*

Proof. ${}_c D_a F(x)$ for each $x \in P_a$ is as follows :

$$\begin{aligned} {}_c D_a F(0) &= F(a) - cF(0) = a^{-1}, & {}_c D_a F(-a) &= F(0) - cF(-a) = ca^{-1}, \\ {}_c D_a F(1) &= F(1 + a) - cF(1) = (a + 1)^{-1} - c\gamma^{-1}, & {}_c D_a F(1 - a) &= F(1) - cF(1 - a) = \gamma^{-1} - c(1 - a)^{-1}, \\ {}_c D_a F(\gamma) &= F(\gamma + a) - cF(\gamma) = (a + \gamma)^{-1} - c, & {}_c D_a F(\gamma - a) &= F(\gamma) - cF(\gamma - a) = 1 + c(a - \gamma)^{-1}. \end{aligned}$$

We investigate some possible cases to be used in this proof such that ${}_c D_a F(x) = b$ has at least two solutions in P_a and we obtain an equation on a in each case.

$$\begin{aligned} {}_c D_a F(0) = {}_c D_a F(1) &\Leftrightarrow ca^2 + ca + \gamma = 0, & {}_c D_a F(0) = {}_c D_a F(1 - a) &\Leftrightarrow a^2 + (c\gamma - \gamma - 1)a + \gamma = 0, \\ {}_c D_a F(0) = {}_c D_a F(\gamma) &\Leftrightarrow ca^2 + c\gamma a + \gamma = 0, & {}_c D_a F(0) = {}_c D_a F(\gamma - a) &\Leftrightarrow a^2 + (c - \gamma - 1)a + \gamma = 0, \\ {}_c D_a F(-a) = {}_c D_a F(1 - a) &\Leftrightarrow a^2 - a + c\gamma = 0, & {}_c D_a F(1) = {}_c D_a F(-a) &\Leftrightarrow ca^2 + (c\gamma + c - \gamma)a + c\gamma = 0, \\ {}_c D_a F(-a) = {}_c D_a F(\gamma - a) &\Leftrightarrow a^2 - \gamma a + c\gamma = 0, & {}_c D_a F(\gamma) = {}_c D_a F(-a) &\Leftrightarrow ca^2 + (c\gamma + c - 1)a + c\gamma = 0. \end{aligned}$$

We investigate possible cases such that ${}_c D_a F(x) = b$ has at least four solutions in P_a . Since $c \neq 1$ we have ${}_c D_a F(0) \neq {}_c D_a F(-a)$, and hence ${}_c D_a F(x) = b$ has at most one solution in $\{0, -a\}$. We first investigate the cases that ${}_c D_a F(x) = b$ has one solution in $\{0, -a\}$.

- If ${}_cD_aF(0) = {}_cD_aF(1) = {}_cD_aF(\gamma)$ then ${}_cD_aF(0) = {}_cD_aF(1)$ and ${}_cD_aF(0) = {}_cD_aF(\gamma)$, and hence we have $ca^2 + ca + \gamma = 0$ and $ca^2 + c\gamma a + \gamma = 0$. Then we have $\gamma = 1$, a contradiction, so this case cannot happen.
- If ${}_cD_aF(0) = {}_cD_aF(1-a) = {}_cD_aF(\gamma-a)$ then ${}_cD_aF(0) = {}_cD_aF(1-a)$ and ${}_cD_aF(0) = {}_cD_aF(\gamma-a)$, and hence we have $a^2 + (c\gamma - \gamma - 1)a + \gamma = 0$ and $a^2 + (c - \gamma - 1)a + \gamma = 0$. Then we have $\gamma = 1$, a contradiction, so this case cannot happen.
- ${}_cD_aF(1) = {}_cD_aF(\gamma) = {}_cD_aF(-a)$ then ${}_cD_aF(1) = {}_cD_aF(-a)$ and ${}_cD_aF(\gamma) = {}_cD_aF(-a)$, and hence $ca^2 + (c\gamma + c - \gamma)a + c\gamma = 0$ and $ca^2 + (c\gamma + c - 1)a + c\gamma = 0$. Then we have $\gamma = 1$, a contradiction, so this case cannot happen.
- If ${}_cD_aF(-a) = {}_cD_aF(1-a) = {}_cD_aF(\gamma-a)$ then ${}_cD_aF(-a) = {}_cD_aF(1-a)$ and ${}_cD_aF(-a) = {}_cD_aF(\gamma-a)$, and hence we have $a^2 - a + c\gamma = 0$ and $a^2 - \gamma a + c\gamma = 0$. Then we have $\gamma = 1$, a contradiction, so this case cannot happen.

In this case, ${}_cD_aF(x) = b$ has at least four solutions in P_a requires it has at least three solutions in $\{1, \gamma, 1-a, \gamma-a\}$. Then at least one case that is considered above should be satisfied, which is a contradiction. Hence the only possible case (that ${}_cD_aF(x) = b$ has four solutions) is $b = {}_cD_aF(1) = {}_cD_aF(\gamma) = {}_cD_aF(1-a) = {}_cD_aF(\gamma-a)$. We first investigate all possible cases that ${}_cD_aF(x) = b$ has at least two solutions in $\{1, \gamma, 1-a, \gamma-a\}$.

- ${}_cD_aF(1) = {}_cD_aF(\gamma) \Leftrightarrow ca^2 + c(\gamma+1)a + \gamma(c+1) = 0$.
- ${}_cD_aF(1) = {}_cD_aF(1-a) \Leftrightarrow (c+1)a^2 + \gamma(c-1)a + (\gamma-1)(c+1) = 0$.
- ${}_cD_aF(1) = {}_cD_aF(\gamma-a) \Leftrightarrow (c+\gamma)a^2 + (c-\gamma^2)a = 0$ and hence $a = \frac{\gamma^2-c}{\gamma+c}$ when $c \neq -\gamma$.
- ${}_cD_aF(\gamma) = {}_cD_aF(1-a) \Leftrightarrow (c\gamma+1)a^2 + (c\gamma^2-1)a = 0$ and hence $a = \frac{1-c\gamma^2}{1+c\gamma}$ when $c \neq -\gamma^{-1}$.
- ${}_cD_aF(\gamma) = {}_cD_aF(\gamma-a) \Leftrightarrow (c+1)a^2 + (c-1)a + \gamma(1-\gamma)(c+1) = 0$

If $c = -1$, then ${}_cD_aF(1) = {}_cD_aF(1-a)$ and ${}_cD_aF(\gamma) = {}_cD_aF(\gamma-a)$ imply $a = 0$, a contradiction. Hence we require $c \neq -1$. Next we investigate all possible cases that ${}_cD_aF(x) = b$ has at least three solutions in $\{1, \gamma, 1-a, \gamma-a\}$.

- If ${}_cD_aF(1) = {}_cD_aF(\gamma) = {}_cD_aF(1-a)$ then we have $ca^2 + c(\gamma+1)a + \gamma(c+1) = 0$, $(c+1)a^2 + \gamma(c-1)a + (\gamma-1)(c+1) = 0$ and $(c\gamma+1)a^2 + (c\gamma^2-1)a = 0$. We substitute $ca^2 = -c(\gamma+1)a - \gamma(c+1)$ which is obtained from the first equation to the second equation to have $a^2 - (c+\gamma)a - (c+1) = 0$. We substitute $a = \frac{1-c\gamma^2}{c\gamma+1}$ to $a^2 - (c+\gamma)a - (c+1) = 0$ to have $(\gamma^3 - \gamma^2)c^3 + (2\gamma^4 - 3\gamma)c^2 + (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c - \gamma = 0$ when $c \neq -\gamma^{-1}$. But if $c = -\gamma^{-1}$, then we have $0 = (\gamma^3 - \gamma^2)c^3 + (2\gamma^4 - 3\gamma)c^2 + (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c - \gamma = (\gamma+1)^2$ and then $\gamma = -1$ and $c = 1$, a contradiction.
- If ${}_cD_aF(1) = {}_cD_aF(\gamma) = {}_cD_aF(\gamma-a)$ then we have $ca^2 + c(\gamma+1)a + \gamma(c+1) = 0$, $(c+\gamma)a^2 + (c-\gamma^2)a = 0$ and $(c+1)a^2 + (c-1)a + \gamma(1-\gamma)(c+1) = 0$. We substitute $ca^2 = -c(\gamma+1)a - \gamma(c+1)$ which is obtained from the first equation to the third equation to have $a^2 - (c\gamma+1)a - \gamma^2(c+1) = 0$. We substitute $a = \frac{\gamma^2-c}{\gamma+c}$ from the second equation to $a^2 - (c\gamma+1)a - \gamma^2(c+1) = 0$ to have

$(\gamma^2 - \gamma)c^3 + (3\gamma^3 - 2)c^2 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c + \gamma^3 = 0$ when $c \neq -\gamma$. But if $c = -\gamma$ then $0 = (\gamma^2 - \gamma)c^3 + (3\gamma^3 - 2)c^2 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c + \gamma^3 = -\gamma^2(\gamma + 1)^2$ which implies $\gamma = -1$ and $c = 1$, a contradiction.

- If ${}_cD_aF(1) = {}_cD_aF(1-a) = {}_cD_aF(\gamma-a)$ then we have ${}_{c^{-1}}D_{-a}F(1-(-a)) = {}_{c^{-1}}D_{-a}F(1-a-(-a)) = {}_{c^{-1}}D_{-a}F(\gamma-a-(-a))$ by Lemma 2.2, or equivalently ${}_{c^{-1}}D_{a'}F(1) = {}_{c^{-1}}D_{a'}F(\gamma) = {}_{c^{-1}}D_{a'}F(1-a')$ where $a' = -a$. By the above analysis, we have $(\gamma^3 - \gamma^2)(c^{-1})^3 + (2\gamma^4 - 3\gamma)(c^{-1})^2 + (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c^{-1} - \gamma = 0$ or equivalently $\gamma c^3 - (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c^2 - (2\gamma^4 - 3\gamma)c - \gamma^3 + \gamma^2 = 0$.
- If ${}_cD_aF(\gamma) = {}_cD_aF(1-a) = {}_cD_aF(\gamma-a)$ then we have ${}_{c^{-1}}D_{a'}F(1) = {}_{c^{-1}}D_{a'}F(\gamma) = {}_{c^{-1}}D_{a'}F(\gamma-a')$ where $a' = -a$, similarly with the above case. Then we have $(\gamma^2 - \gamma)(c^{-1})^3 + (3\gamma^3 - 2)(c^{-1})^2 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c^{-1} + \gamma^3 = 0$ or equivalently $\gamma^3 c^3 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c^2 + (3\gamma^3 - 2)c + \gamma^2 - \gamma = 0$.

So we have the following four cubic equations on c .

$$(\gamma^3 - \gamma^2)c^3 + (2\gamma^4 - 3\gamma)c^2 + (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c - \gamma = 0, \quad (4)$$

$$(\gamma^2 - \gamma)c^3 + (3\gamma^3 - 2)c^2 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c + \gamma^3 = 0, \quad (5)$$

$$\gamma c^3 - (\gamma^3 - 3\gamma^2 - 2\gamma - 2)c^2 - (2\gamma^4 - 3\gamma)c - \gamma^3 + \gamma^2 = 0, \quad (6)$$

$$\gamma^3 c^3 + (2\gamma^4 + 2\gamma^3 + 3\gamma^2 - \gamma)c^2 + (3\gamma^3 - 2)c + \gamma^2 - \gamma = 0. \quad (7)$$

Then (5) - $(\gamma - 1) \times$ (6) is the following quadratic equation.

$$\begin{aligned} 0 &= (\gamma^4 - \gamma^3 + \gamma^2)c^2 + 2(\gamma^5 + \gamma^3 + \gamma)c + \gamma^4 - \gamma^3 + \gamma^2 \\ &= \gamma(\gamma^2 - \gamma + 1)(\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma) \end{aligned} \quad (8)$$

We also confirm that $-(4) + \gamma \times (5)$ and $(-6) \times \gamma^2 + (7) / (\gamma - 1)$ are same with $(8) \times (\gamma + 1) / \gamma$. If $\gamma^2 - \gamma + 1 = 0$ then equations (4), (5), (6) and (7) imply $0 = c^3 + 5c^2 + 5c + 1 = (c + 1)(c^2 + 4c + 1)$. Since $c \neq -1$, we have $c^2 + 4c + 1 = 0$ and then $\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 2c(\gamma^2 - \gamma + 1) = 0$ when $\gamma^2 - \gamma + 1 = 0$. Hence we only require that $\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 0$ whether $\gamma^2 - \gamma + 1 = 0$ or not. In this case we set $a = \frac{\gamma^2 - c}{\gamma + c}$ and $b = {}_cD_aF(1) = (a + 1)^{-1} - c\gamma^{-1} = \frac{1-c}{1+\gamma}$.

Conversely, we assume that $\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 0$ with $c = -1$. If $c = -\gamma$ then we have $0 = -\gamma^3 - 2\gamma^2 - \gamma = -\gamma(\gamma + 1)^2$ so $\gamma = -1$ and hence $c = -\gamma = 1$ which contradicts to the assumption, so we have $c \neq -\gamma$. Similarly, we have $c \neq -\gamma^{-1}$. Since $(\gamma^2 - c)(1 + c\gamma) - (\gamma + c)(1 - c\gamma^2) = (\gamma - 1)(\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma) = 0$ we set $a = \frac{\gamma^2 - c}{\gamma + c} = \frac{1 - c\gamma^2}{1 + c\gamma}$. By the above analysis, we have ${}_cD_aF(1) = {}_cD_aF(\gamma - a)$ and ${}_cD_aF(\gamma) = {}_cD_aF(1 - a)$. Since $a = \frac{\gamma^2 - c}{\gamma + c}$, we have $(c + 1)a^2 + \gamma(c - 1)a + (\gamma - 1)(c + 1) = \frac{\gamma(\gamma - 1)(\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma)}{(c + \gamma)^2} = 0$ and hence ${}_cD_aF(1) = {}_cD_aF(1 - a)$. Hence ${}_cD_aF(x) = b$ has four solutions in P_a , which completes the proof. \square

Next we show the main theorem of this subsection.

Theorem 4.7. *Let $F = \text{Inv} \circ (1, \gamma)$ with $\gamma \neq 0, 1$ and $c \neq 0, 1$. Then ${}_c\Delta_F = 6$ if and only if $\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 0$, $c \neq -1$, and $-c$ is a square. Otherwise, ${}_c\Delta_F \leq 5$.*

Proof. By Lemma 4.3 and Lemma 4.6, ${}_cD_aF(x) = b$ has four solutions in P_a if and only if

$$\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = 0, \quad c \neq -1, \quad a = \frac{\gamma^2 - c}{\gamma + c}, \quad b = \frac{1 - c}{1 + \gamma}, \quad (9)$$

otherwise ${}_cD_aF(x) = b$ has at most three solutions in P_a . Hence it remains to investigate required conditions that ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ using Lemma 4.1, under the assumption as in (9). We require that

$$\begin{aligned} (ab + c - 1)^2 - 4abc &= \frac{\gamma^2 c^4 + 4(\gamma^4 - \gamma - 1)c^3 + 2(2\gamma^2 - \gamma^2 + 2)c^2 - 4(\gamma^4 + \gamma^3 - \gamma)c + \gamma^2}{(\gamma + 1)^2(c + \gamma)^2} \\ &= \frac{(\gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma)^2 - 4c(\gamma + 1)^2(c + \gamma)^2}{(\gamma + 1)^2(c + \gamma)^2} = -4c \end{aligned}$$

is a square. We can see that $c \neq -\gamma$ in the proof of Lemma 4.6. If $\gamma = -1$ then we have $0 = \gamma c^2 + 2(\gamma^2 + \gamma + 1)c + \gamma = -(c - 1)^2$, and then $c = 1$ which is a contradiction, so we have $\gamma \neq -1$. Hence we get $(b + c)a + b + c - 1 = \frac{c(\gamma^2 - 1)}{c + \gamma} \neq 0$, $(1 - b)a + b + c - 1 = \frac{(\gamma - 1)(c + \gamma)}{\gamma + 1} \neq 0$, $(b\gamma + c)a + \gamma(b\gamma + c - 1) \frac{(\gamma - 1)(c + \gamma)}{\gamma + 1} \neq 0$ and $(1 - b\gamma)a + \gamma(b\gamma + c - 1) = \frac{c(\gamma^2 - 1)}{c + \gamma} \neq 0$. Therefore, under the assumption (9) ${}_cD_aF(x) = b$ has two solutions in $\mathbb{F}_{p^n} \setminus P_a$ if and only if $-c$ is a square, by Lemma 4.1. \square

5 Concluding Remark

In this paper, we studied the c -differential uniformity of the swapped inverse function $Inv \circ (1, \gamma)$. In section 3, we showed that the c -differential uniformity of $Inv \circ (0, 1)$ is at most 5. In section 4, we showed that the c -differential uniformity of $Inv \circ (1, \gamma)$ is at most 6 when $\gamma \neq 1$. For each case (i.e., $c = 1$ or $c \neq 1$), we gave a detailed classification of the (c -)differential uniformity of $Inv \circ (1, \gamma)$, which has been unknown as far as we know. Unfortunately we could not find (A)PcN permutations except when p^n is very small. In future studies, we will continue to find permutations $Inv \circ h$ with cyclic permutation h having low c -differential uniformities.

Acknowledgements : This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2021R1C1C2003888). Soonhak Kwon was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (No. 2016R1A5A1008055, No. 2019R1F1A1058920 and 2021R1F1A1050721).

References

- [1] D. Bartoli, and M. Calderini, *On construction and (non)existence of c -(almost) perfect nonlinear functions*, Finite Fields Appl., Vol. 72, 101835, 2021. DOI : 10.1016/j.ffa.2021.101835
- [2] D. Bartoli, M. Calderini, C. Riera, and P. Stănică, *Low c -differential uniformity for functions modified on subfields*, Cryptogr. Commun., in press, 2022. DOI : 10.1007/s12095-022-00554-x
- [3] N. Borisov, M. Chew, R. Johnson, and D. Wagner, *Multiplicative Differentials*, FSE 2002, Lect. Notes Comput. Sci. Vol. 2365, pp.17-33, 2002. DOI : 10.1007/3-540-45661-9_2

- [4] R. S. Coulter, R. W. Matthews, *Planar Functions and Planes of Lenz-Barlotti Class II*, Des. Codes Cryptogr., Vol. 10, issue 2, pp.167-184, 1997. DOI : 10.1023/A:1008292303803
- [5] H. Dobbertin, D. Mills, E. N. Müller, A. Pott, and W. Willems, *APN functions in odd characteristic*, Discret. Math., Vol. 267, pp.95-112, 2003. DOI : 10.1016/S0012-365X(02)00606-4
- [6] P. Ellingsen, P. Felke, C. Riera, P. Stănică, and A. Tkachenko, *c-Differentials, Multiplicative Uniformity, and (Almost) Perfect c-Nonlinearity*, IEEE Trans. Inf. Theory, Vol. 66, No. 9, pp.5781-5789, 2020. DOI : 10.1109/TIT.2020.2971988
- [7] S. U. Hasan, M. Pal, C. Riera, and P. Stănică, *On the c-differential uniformity of certain maps over finite fields*, Des. Codes Cryptogr., Vol. 89, issue 2, pp.221-239, 2021. DOI : 10.1007/s10623-020-00812-0
- [8] T. Helleseth, C. Rong, and D. Sandberg, *New Families of Almost Perfect Nonlinear Power Mappings*, IEEE Trans. Inf. Theory, Vol. 45, No. 2, pp. 475-485, 1999. DOI : 10.1109/18.748997
- [9] T. Helleseth, and D. Sandberg, *Some Power Mappings with Low Differential Uniformity*, Appl. Algebr. Eng. Commun. Comput., Vol. 8, issue 5, pp.363-370, 1997. DOI : 10.1007/s002000050073
- [10] J. Jeong, N. Koo, and S. Kwon, *New differentially 4-uniform permutations from modifications of the inverse function*, Finite Fields Appl., Vol. 77, 101931, 2022. DOI : 10.1016/j.ffa.2021.101931
- [11] J. Jeong, N. Koo, and S. Kwon, *Investigations of c-Differential Uniformity of Permutations with Carlitz Rank 3*, a preprint, available at <https://arxiv.org/abs/2202.02185>
- [12] W. Jia, X. Zeng, C. Li, T. Helleseth, and L. Hu, *Permutation polynomials with low differential uniformity over finite fields of odd characteristic*, Sci. China-Math., Vol. 56, No. 7, pp.1429-1440, 2013.
- [13] L. Lei, W. Ren, and C. Fan, *The Differential Spectrum of a Class of Power Functions over Finite Fields*, Adv. Math. Commun., Vol. 15, No. 3, pp. 525-537, 2021. DOI : 10.3934/amc.2020080
- [14] K. Li, L. Qu, B. Sun, and C. Li, *New Results about the Boomerang Uniformity of Permutation Polynomials*, IEEE Trans. Inf. Theory, Vol. 65 (2019) 7542-7553, 2019. DOI : 10.1109/TIT.2019.2918531
- [15] Y. Li, M. Wang and Y. Yu, *Constructing Differentially 4-uniform Permutations over $GF(2^{2k})$ from the Inverse Function Revisited*, IACR ePrint Archive 2013/731.
- [16] S. Mesnager, C. Riera, P. Stănică, H. Yan, and Z. Zhou, *Investigations on c-(Almost) Perfect Nonlinear Functions*, IEEE Trans. Inf. Theory, Vol. 67, No. 10, pp. 6916-6925, 2021. DOI : 10.1109/TIT.2021.3081348
- [17] K. Nyberg, *Differentially uniform mappings for cryptography*, EUROCRYPT '93, Lect. Notes Comput. Sci. Vol. 765, pp. 55-64, 1994. DOI : 10.1007/3-540-48285-7_6
- [18] A. Pott, *Almost perfect and planar functions*, Des. Codes Cryptogr., Vol. 78, issue 1, pp.141-195, 2016. DOI : 10.1007/s10623-015-0151-x

- [19] P. Stănică, *Low c -differential and c -boomerang uniformity of the swapped inverse function*, Discret. Math., Vol. 344, Vol. 344, 112543, 2021. DOI : 10.1016/j.disc.2021.112543
- [20] P. Stănică, *Low c -Differential Uniformity for the Gold Function Modified on a Subfield*, Security and Privacy, Lecture Notes in Electrical Engineering, Vol. 744, pp.131-137, 2021. DOI : 10.1007/978-981-33-6781-4_11
- [21] Z. Tu, X. Zeng, Y. Jiang, and X. Tang, *A class of APcN power functions over finite fields of even characteristic*, a preprint, available at <https://arxiv.org/abs/2107.06464v1>.
- [22] X. Wang, and D. Zheng, *Several classes of PcN power functions over finite fields*, a preprint, available at <https://arxiv.org/abs/2104.12942v1>.
- [23] Y. Wu, N. Li, and X. Zeng, *New PcN and APcN functions over finite fields*, Des. Codes Cryptogr., Vol. 89, issue 11, pp. 2637-2651, 2021. DOI : 10.1007/s10623-021-00946-9
- [24] H. Yan, *On (-1) -differential uniformity of ternary APN power functions*, Cryptogr. Commun., in press, 2021. DOI : 10.1007/s12095-021-00526-7
- [25] H. Yan, and D. Han, *New Ternary Power Mapping with Differential Uniformity $\Delta_f \leq 3$ and Related Optimal Cyclic Codes*, IEICE Trans. Fundam. Electron. Commun. Comput. Sci., Vol. E102-A, No. 6, pp. 849-853, 2019. DOI: 10.1587/transfun.E102.A.849
- [26] Z. Zha, and L. Hu, *Some classes of power functions with low c -differential uniformity over finite fields*, Des. Codes Cryptogr., Vol. 89, Issue 6, pp.1193-1210, 2021. DOI : 10.1007/s10623-021-00866-8
- [27] Z. Zha, and X. Wang, *Power functions with low uniformity on odd characteristic finite fields*, Sci. China-Math., Vol. 53, No. 8, pp.1931-1940, 2010. DOI : 10.1007/s11425-010-3149-x
- [28] Z. Zha, and X. Wang, *Almost Perfect Nonlinear Power Functions in Odd Characteristic*, IEEE Trans. Inf. Theory, Vol. 57, No. 7, pp.4826-4832, 2011. DOI : 10.1109/TIT.2011.2145130
- [29] K. Zhang, and H. Yan, *On the c -differential spectrum of power functions over finite fields*, a preprint, available at : <https://arxiv.org/abs/2201.06172v1>

A Proof of Lemma 4.3

By Lemma 2.2, we have ${}_c\Delta_F(-a, b) = {}_{c^{-1}}\Delta_F(a, -bc^{-1})$. Hence it is enough to investigate the case $a \in \{1, \gamma, \gamma - 1\}$.

If $a = 1$ with $\gamma = -1$ then $P_a = \{0, \pm 1, -2\}$ and we have ${}_cD_1F(0) = -1$, ${}_cD_1F(1) = 2^{-1} + c$, ${}_cD_1F(-1) = -c$ and ${}_cD_1F(-2) = 1 + 2^{-1}c$.

$$\begin{aligned} {}_cD_1F(0) = {}_cD_1F(1) &\Leftrightarrow c = -\frac{3}{2}, & {}_cD_1F(0) = {}_cD_1F(-1) &\Leftrightarrow c = 1, \\ {}_cD_1F(0) = {}_cD_1F(-2) &\Leftrightarrow c = -4, & {}_cD_1F(1) = {}_cD_1F(-1) &\Leftrightarrow c = -\frac{1}{4}, \\ {}_cD_1F(1) = {}_cD_1F(-2) &\Leftrightarrow c = 1, & {}_cD_1F(-1) = {}_cD_1F(-2) &\Leftrightarrow c = -\frac{2}{3}. \end{aligned}$$

We investigate all possible cases such that ${}_cD_1F(x) = b$ has at least three solutions in P_a .

$$\begin{aligned} {}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-1) &\Leftrightarrow c = -\frac{3}{2} = 1 = -\frac{1}{4} \Rightarrow 2 = -3, 6 = 1 \text{ and } 1 = -4 \Rightarrow p = 5. \\ {}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-2) &\Leftrightarrow c = -\frac{3}{2} = -4 = 1 \Rightarrow -8 = -3, 2 = -3 \text{ and } 1 = -4 \Rightarrow p = 5. \\ {}_cD_1F(0) = {}_cD_1F(-1) = {}_cD_1F(-2) &\Leftrightarrow c = 1 = -4 = -\frac{2}{3} \Rightarrow 1 = -4, 2 = -3 \text{ and } -12 = -2 \Rightarrow p = 5. \\ {}_cD_1F(1) = {}_cD_1F(-1) = {}_cD_1F(-2) &\Leftrightarrow c = -\frac{1}{4} = 1 = -4 = -\frac{2}{3} \Rightarrow 1 = -4 = 16 \Rightarrow p = 5. \end{aligned}$$

So we have ${}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-1) = {}_cD_1F(-2) = -1$ when $c = 1$ and $p = 5$.

If $a = 1$ with $\gamma = 2$ then $P_a = \{0, \pm 1, 2\}$. If $p = 3$ then we have $P_a = \mathbb{F}_3$ and hence ${}_cD_aF(x) = b$ has at most three solutions in P_a . If $p \neq 3$, then we have ${}_cD_1F(0) = 2^{-1}$, ${}_cD_1F(1) = 1 - 2^{-1}c$, ${}_cD_1F(-1) = c$ and ${}_cD_1F(2) = 3^{-1} - c$. We investigate equivalent condition for each case that $D_aF(x) = b$ has at least two solutions in P_a .

$$\begin{aligned} {}_cD_1F(0) = {}_cD_1F(1) &\Leftrightarrow c = 1, & {}_cD_1F(0) = {}_cD_1F(-1) &\Leftrightarrow c = 2^{-1}, \\ {}_cD_1F(0) = {}_cD_1F(2) &\Leftrightarrow c = -\frac{1}{6}, & {}_cD_1F(1) = {}_cD_1F(-1) &\Leftrightarrow c = \frac{2}{3}, \\ {}_cD_1F(1) = {}_cD_1F(2) &\Leftrightarrow c = -\frac{4}{3}, & {}_cD_1F(-1) = {}_cD_1F(2) &\Leftrightarrow c = \frac{1}{6}. \end{aligned}$$

We investigate all possible cases such that ${}_cD_1F(x) = b$ has at least three solutions in P_a .

$$\begin{aligned} {}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-1) &\Leftrightarrow c = 1 = 2^{-1} \Rightarrow 2 = 1, \text{ a contradiction.} \\ {}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(2) &\Leftrightarrow c = 1 = -\frac{1}{6} = -\frac{4}{3} \Rightarrow 1 = -6, 3 = -4 \text{ and } 3 = 24 \Rightarrow p = 7. \\ {}_cD_1F(0) = {}_cD_1F(-1) = {}_cD_1F(2) &\Leftrightarrow c = \frac{1}{6} = -\frac{1}{6}, \text{ a contradiction.} \\ {}_cD_1F(1) = {}_cD_1F(-1) = {}_cD_1F(2) &\Leftrightarrow c = \frac{2}{3} = -\frac{4}{3} \Rightarrow 2 = 0, \text{ a contradiction.} \end{aligned}$$

So we can see that ${}_cD_1F(x) = b$ has at most three solutions.

If $a = 1$ with $\gamma \neq -1, 2$ then $P_a = \{0, 1, \gamma, -1, \gamma - 1\}$ and we have ${}_cD_1F(0) = \gamma^{-1}$, ${}_cD_1F(1) = 2^{-1} - c\gamma^{-1}$, ${}_cD_1F(\gamma) = (\gamma + 1)^{-1} - c$, ${}_cD_1F(-1) = c$ and ${}_cD_1F(\gamma - 1) = 1 - c(\gamma - 1)^{-1}$. We investigate equivalent condition for each case that $D_aF(x) = b$ has at least two solutions in P_a .

$$\begin{aligned} {}_cD_1F(0) = {}_cD_1F(1) &\Leftrightarrow c = \frac{\gamma-2}{2}, & {}_cD_1F(0) = {}_cD_1F(\gamma) &\Leftrightarrow c = -\frac{1}{\gamma(\gamma+1)}, \\ {}_cD_1F(0) = {}_cD_1F(-1) &\Leftrightarrow c = \gamma^{-1}, & {}_cD_1F(0) = {}_cD_1F(\gamma-1) &\Leftrightarrow c = \frac{(\gamma-1)^2}{\gamma}, \\ {}_cD_1F(1) = {}_cD_1F(\gamma) &\Leftrightarrow c = -\frac{\gamma}{2(\gamma+1)}, & {}_cD_1F(1) = {}_cD_1F(-1) &\Leftrightarrow c = \frac{2\gamma}{\gamma+1}, \\ {}_cD_1F(1) = {}_cD_1F(\gamma-1) &\Leftrightarrow c = \frac{\gamma(\gamma-1)}{2}, & {}_cD_1F(\gamma) = {}_cD_1F(-1) &\Leftrightarrow c = \frac{1}{2(\gamma+1)}, \\ {}_cD_1F(\gamma) = {}_cD_1F(\gamma-1) &\Leftrightarrow c = -\frac{\gamma(\gamma-1)}{(\gamma+1)(\gamma-2)}, & {}_cD_1F(-1) = {}_cD_1F(\gamma-1) &\Leftrightarrow c = \frac{\gamma-1}{\gamma}. \end{aligned}$$

Next, we can see that ${}_cD_1F(x) = b$ cannot have four solutions in P_a by showing all possible cases cannot happen.

- ${}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-1) \Leftrightarrow c = \frac{\gamma-2}{2} = \gamma^{-1} = \frac{2\gamma}{\gamma+1}$. Then we have $\gamma^2 - 2\gamma - 2 = 2\gamma^2 - \gamma - 1 = \gamma^2 - 5\gamma - 2 = 0$ and then $\gamma = 0$, a contradiction. Hence ${}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(\gamma) = {}_cD_1F(-1)$ and ${}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(-1) = {}_cD_1F(\gamma - 1)$ cannot happen.
- ${}_cD_1F(1) = {}_cD_1F(\gamma) = {}_cD_1F(-1) \Leftrightarrow c = -\frac{\gamma}{2(\gamma+1)} = \frac{2\gamma}{\gamma+1} = \frac{1}{2(\gamma+1)}$. This case cannot happen, since $-\frac{\gamma}{2(\gamma+1)} = \frac{1}{2(\gamma+1)}$ implies $\frac{1}{2} = 0$, a contradiction. So ${}_cD_1F(0) = {}_cD_1F(\gamma) = {}_cD_1F(-1) = {}_cD_1F(\gamma - 1)$ is also impossible.

- ${}_cD_1F(1) = {}_cD_1F(\gamma) = {}_cD_1F(\gamma-1) \Leftrightarrow c = -\frac{\gamma}{2(\gamma+1)} = \frac{\gamma(\gamma-1)}{2} = -\frac{\gamma(\gamma-1)}{(\gamma+1)(\gamma-2)}$. This case cannot happen, since $-\frac{\gamma}{2(\gamma+1)} = \frac{\gamma(\gamma-1)}{2}$ implies $\gamma = 0$, a contradiction. Hence ${}_cD_1F(0) = {}_cD_1F(1) = {}_cD_1F(\gamma) = {}_cD_1F(\gamma-1)$ and ${}_cD_1F(1) = {}_cD_1F(\gamma) = {}_cD_1F(-1) = {}_cD_1F(\gamma-1)$ cannot happen, neither.

If $a = \gamma$ with $\gamma = -1$, then we have ${}_cD_{-1}F(1) = {}_cD_{-1}F(2) = {}_cD_{-1}F(0) = {}_cD_{-1}F(-1) = 1$ when $c = 1$ and $p = 5$ by applying Lemma 2.2 for the already investigated case $a = 1$ and $\gamma = -1$. If $a = \gamma$ with $\gamma = 2^{-1}$ then ${}_cD_{2^{-1}}F_1(x) = b$ and ${}_cD_1F_2(x) = b$ have the same number of solutions in P_a where $F_1 = Inv \circ (1, 2^{-1})$ and $F_2 = Inv \circ (1, 2)$, by applying $\gamma = 2$ in Lemma 4.2. Hence ${}_cD_{2^{-1}}F_1(x) = b$ has at most three solutions in P_a . If $a = \gamma$ with $\gamma \neq -1, 2^{-1}$ then $P_a = \{0, 1, \gamma, -\gamma, 1 - \gamma\}$, and we have ${}_cD_\gamma F(0) = 1$, ${}_cD_\gamma F(1) = (\gamma + 1)^{-1} - c\gamma^{-1}$, ${}_cD_\gamma F(\gamma) = (2\gamma)^{-1} - c$, ${}_cD_\gamma F(-\gamma) = c\gamma^{-1}$ and ${}_cD_\gamma F(1 - \gamma) = \gamma^{-1} + c(\gamma - 1)^{-1}$. We investigate equivalent condition for each case that $D_a F(x) = b$ has at least two solutions in P_a .

$$\begin{aligned}
{}_cD_\gamma F(0) = {}_cD_\gamma F(1) &\Leftrightarrow c = -\frac{\gamma^2}{\gamma+1}, & {}_cD_\gamma F(0) = {}_cD_\gamma F(\gamma) &\Leftrightarrow c = -\frac{2\gamma-1}{2\gamma}, \\
{}_cD_\gamma F(0) = {}_cD_\gamma F(-\gamma) &\Leftrightarrow c = \gamma, & {}_cD_\gamma F(0) = {}_cD_\gamma F(1-\gamma) &\Leftrightarrow c = \frac{(\gamma-1)^2}{\gamma}, \\
{}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) &\Leftrightarrow c = -\frac{1}{2(\gamma+1)}, & {}_cD_\gamma F(1) = {}_cD_\gamma F(-\gamma) &\Leftrightarrow c = \frac{\gamma}{2(\gamma+1)}, \\
{}_cD_\gamma F(1) = {}_cD_\gamma F(1-\gamma) &\Leftrightarrow c = -\frac{\gamma-1}{(\gamma+1)(2\gamma-1)}, & {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(-\gamma) &\Leftrightarrow c = \frac{1}{2(\gamma+1)}, \\
{}_cD_\gamma F(\gamma) = {}_cD_\gamma F(1-\gamma) &\Leftrightarrow c = -\frac{\gamma-1}{2\gamma^2}, & {}_cD_\gamma F(-\gamma) = {}_cD_\gamma F(1-\gamma) &\Leftrightarrow c = -\gamma + 1.
\end{aligned}$$

Next, we can see that ${}_cD_1F(x) = b$ cannot have four solutions in P_a by showing all possible cases cannot happen.

- ${}_cD_\gamma F(0) = {}_cD_\gamma F(-\gamma) = {}_cD_\gamma F(1 - \gamma) \Leftrightarrow c = \gamma = \frac{(\gamma-1)^2}{\gamma} = -\gamma + 1$. Then we have $c = \gamma = 2^{-1}$, a contradiction. Hence ${}_cD_\gamma F(0) = {}_cD_\gamma F(1) = {}_cD_\gamma F(-\gamma) = {}_cD_\gamma F(1 - \gamma)$ and ${}_cD_\gamma F(0) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(-\gamma) = {}_cD_\gamma F(1 - \gamma)$ are also impossible.
- ${}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(-\gamma) \Leftrightarrow c = -\frac{1}{2(\gamma+1)} = \frac{\gamma}{2(\gamma+1)} = \frac{1}{2(\gamma+1)}$. This case cannot happen, because we have $\frac{1}{2} = 0$ from $-\frac{1}{2(\gamma+1)} = \frac{\gamma}{2(\gamma+1)}$ a contradiction. Hence ${}_cD_\gamma F(0) = {}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(-\gamma)$ and ${}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(-\gamma) = {}_cD_\gamma F(1 - \gamma)$ are also impossible.
- ${}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(1 - \gamma) \Leftrightarrow c = -\frac{1}{2(\gamma+1)} = -\frac{\gamma-1}{(\gamma+1)(2\gamma-1)} = -\frac{\gamma-1}{2\gamma^2}$. This case cannot happen, because we have $-1 = 0$ from $-\frac{1}{2(\gamma+1)} = -\frac{\gamma-1}{2\gamma^2}$, a contradiction. So, ${}_cD_\gamma F(0) = {}_cD_\gamma F(1) = {}_cD_\gamma F(\gamma) = {}_cD_\gamma F(1 - \gamma)$ is also impossible.

If $a = \gamma - 1$ with $\gamma = 2$ then we have $a = 1$ which is already investigated. If $a = \gamma - 1$ with $\gamma = 2^{-1}$ then we have $a = -2^{-1}$. By Lemma 2.2, ${}_cD_{2^{-1}}F(x) = b$ and ${}_{c^{-1}}D_{2^{-1}}F(x) = -b$ have the same number of solutions in P_a . By Lemma 4.2, ${}_{c^{-1}}D_{2^{-1}}F(x) = -b$ and ${}_{c^{-1}}D_1F_2(x) = -2^{-1}b$ have the same number of solutions in P_a where $F_2 = Inv \circ (1, 2)$. Hence ${}_cD_{2^{-1}}F(x) = b$ has at most three solutions in P_a . If $a = \gamma - 1$ with $\gamma \neq 2, 2^{-1}$ then $P_a = \{0, 1, \gamma, 1 - \gamma, 2 - \gamma\}$ and we have ${}_cD_{\gamma-1}F(0) = (\gamma - 1)^{-1}$, ${}_cD_{\gamma-1}F(1) = 1 - c\gamma^{-1}$, ${}_cD_{\gamma-1}F(\gamma) = (2\gamma - 1)^{-1} - c$, ${}_cD_{\gamma-1}F(1 - \gamma) = c(\gamma - 1)^{-1}$ and ${}_cD_{\gamma-1}F(2 - \gamma) = \gamma^{-1} + c(\gamma - 2)^{-1}$. We investigate equivalent condition for each case

that $D_a F(x) = b$ has at least two solutions in P_a .

$$\begin{aligned}
{}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1) &\Leftrightarrow c = \frac{\gamma(\gamma-2)}{\gamma-1}, & {}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(\gamma) &\Leftrightarrow c = \frac{-\gamma}{(2\gamma-1)(\gamma-1)}, \\
{}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1-\gamma) &\Leftrightarrow c = 1, & {}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(2-\gamma) &\Leftrightarrow c = \frac{\gamma-2}{\gamma(\gamma-1)}, \\
{}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(\gamma) &\Leftrightarrow c = -\frac{2\gamma}{2\gamma-1}, & {}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(1-\gamma) &\Leftrightarrow c = \frac{\gamma(\gamma-1)}{2\gamma-1}, \\
{}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(2-\gamma) &\Leftrightarrow c = \frac{\gamma-2}{2}, & {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(1-\gamma) &\Leftrightarrow c = \frac{\gamma-1}{\gamma(2\gamma-1)}, \\
{}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(2-\gamma) &\Leftrightarrow c = \frac{2-\gamma}{\gamma(2\gamma-1)}, & {}_c D_{\gamma-1} F(1-\gamma) = {}_c D_{\gamma-1} F(2-\gamma) &\Leftrightarrow c = -\frac{(\gamma-1)(\gamma-2)}{\gamma}.
\end{aligned}$$

- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(\gamma) \Leftrightarrow c = \frac{\gamma(\gamma-2)}{\gamma-1} = \frac{-\gamma}{(2\gamma-1)(\gamma-1)} = -\frac{2\gamma}{2\gamma-1}$. Then we have $\gamma = \frac{3}{2}$ and then $c = -\frac{3}{2}$.
- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(1-\gamma) \Leftrightarrow c = \frac{\gamma(\gamma-2)}{\gamma-1} = \frac{\gamma(\gamma-1)}{2\gamma-1} = 1$. Then we have $\gamma^2 - 3\gamma + 1 = 0$, such γ exists if and only if $(-3)^2 - 4 = 5$ is a square if and only if $p^n \equiv 0, \pm 1 \pmod{5}$.
- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = \frac{\gamma(\gamma-2)}{\gamma-1} = \frac{\gamma-2}{\gamma(\gamma-1)} = \frac{\gamma-2}{2}$. Then we have $\gamma = -1$ and then $c = -\frac{3}{2}$.
- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(1-\gamma) \Leftrightarrow c = \frac{-\gamma}{(2\gamma-1)(\gamma-1)} = \frac{\gamma-1}{\gamma(2\gamma-1)} = 1$. Then we have $2\gamma^2 - 2\gamma + 1 = 0$, such γ exists if and only if $(-2)^2 - 8 = -4$ is a square if and only if $p^n \equiv 1 \pmod{4}$.
- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = \frac{-\gamma}{(2\gamma-1)(\gamma-1)} = \frac{\gamma-2}{\gamma(\gamma-1)} = \frac{2-\gamma}{\gamma(2\gamma-1)}$. Then we have $\gamma = \frac{2}{3}$ and then $c = 6$.
- ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1-\gamma) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = 1 = \frac{\gamma-2}{\gamma(\gamma-1)} = -\frac{(\gamma-1)(\gamma-2)}{\gamma}$. Then we have $\gamma^2 - 2\gamma + 2 = 0$, such γ exists if and only if $(-2)^2 - 8 = -4$ is a square if and only if $p^n \equiv 1 \pmod{4}$.
- ${}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(1-\gamma) \Leftrightarrow c = -\frac{2\gamma}{2\gamma-1} = \frac{\gamma(\gamma-1)}{2\gamma-1} = \frac{\gamma-1}{\gamma(2\gamma-1)}$. Then we have $\gamma = -1$ and then $c = -\frac{2}{3}$.
- ${}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = -\frac{2\gamma}{2\gamma-1} = \frac{\gamma-2}{2} = \frac{2-\gamma}{\gamma(2\gamma-1)}$. Then we have $2\gamma^2 - \gamma + 2 = 0$, such γ exists if and only if $(-1)^2 - 4 \cdot 2 \cdot 2 = -15$ is a square if and only if $p^n \equiv 0, 1, 2, 4, 6, 8, 9, 10 \pmod{15}$.
- ${}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(1-\gamma) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = \frac{\gamma(\gamma-1)}{2\gamma-1} = \frac{\gamma-2}{2} = -\frac{(\gamma-1)(\gamma-2)}{\gamma}$. Then we have $\gamma = \frac{2}{3}$ and then $c = -\frac{2}{3}$.
- ${}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(1-\gamma) = {}_c D_{\gamma-1} F(2-\gamma) \Leftrightarrow c = \frac{\gamma-1}{\gamma(2\gamma-1)} = \frac{2-\gamma}{\gamma(2\gamma-1)} = -\frac{(\gamma-1)(\gamma-2)}{\gamma}$. Then we have $\gamma = \frac{3}{2}$ and then $c = \frac{1}{6}$.

From above we can see that ${}_c D_{\gamma-1} F(x) = b$ has at least four solutions only when $p = 5$, $\gamma = -1$ and $c = 1$. For example, if ${}_c D_{\gamma-1} F(0) = {}_c D_{\gamma-1} F(1) = {}_c D_{\gamma-1} F(\gamma) = {}_c D_{\gamma-1} F(1-\gamma)$ then $c = -\frac{3}{2} = 1 = -\frac{2}{3}$ and $\gamma = \frac{3}{2} = -1$ and $\gamma^2 - 3\gamma + 1 = 2\gamma^2 - 2\gamma + 1 = 0$ can happen only when $p = 5$. Hence we can see that all element in P_a is a solution of ${}_c D_{\gamma-1} F(x) = b$ if and only if $p = 5$, $\gamma = -1$ and $c = 1$, and note that $a = 3$ and $b = 2$ in this case. By Lemma 2.2, ${}_1 D_2 F(x) = 3$ also have five solutions in P_a .