

# APERIODIC AND LINEARLY REPETITIVE LORENTZ GASES OF FINITE HORIZON ARE NOT EXPONENTIALLY MIXING

RODRIGO TREVIÑO AND AGNIESZKA ZELEROWICZ

**ABSTRACT.** We prove that aperiodic and linearly repetitive Lorentz gases with finite horizon are not mixing with exponential or stretched exponential speed in any dimension for any class of Hölder observables. We also bound the polynomial speed of mixing for observables in the Hölder space  $H_\alpha$  depending on  $\alpha$ .

The Lorentz gas is one of the oldest and best-studied models from statistical mechanics. It consists of a massless point particle moving through Euclidean space bouncing off a given set of scatterers  $\mathcal{S}$  with elastic collisions at the boundaries  $\partial\mathcal{S}$ . If the set of scatterers is periodic in space, then the quotient system, which is compact, is known as the Sinai billiard. The statistical properties of Sinai billiards in two dimensions are very well understood (see [Szá08, Gol12, Det14, CM06]) and one of the fundamental tools developed for their study are Young towers [You98]. Through the use of towers, Young proved that the Sinai billiard in two dimensions is mixing with exponential speed. This was an improvement of the result of Bunimovich, Sinai and Chernov [BSC90], who proved that the Sinai billiard in two dimensions was mixing with stretched exponential speed. By exponential and stretched exponential speed, we mean the following.

**Definition 1.** Let  $F : X \rightarrow X$  be continuous map on a compact metric space  $X$  which preserves a probability measure  $\mu$ , and let  $H_\alpha$  be the set of Hölder functions on  $X$  with exponent  $\alpha > 0$ . The system  $(X, F, \mu)$  is **exponentially mixing** in  $H_\alpha$  if there exists a  $C > 0$  and  $\tau \in (0, 1)$  such that for any two functions  $\psi_1, \psi_2 \in H_\alpha$  we have

$$\left| \int_X \psi_1 \circ F^n \cdot \psi_2 d\mu - \mu(\psi_1)\mu(\psi_2) \right| \leq C \|\psi_1\|_\alpha \|\psi_2\|_\alpha \tau^n$$

for all  $n \geq 0$ . It has **stretched exponential mixing** in  $H_\alpha$  if there exist  $\gamma, \tau \in (0, 1)$  and  $C > 0$  such that for any two functions  $\psi_1, \psi_2 \in H_\alpha$  we have

$$\left| \int_X \psi_1 \circ F^n \cdot \psi_2 d\mu - \mu(\psi_1)\mu(\psi_2) \right| \leq C \|\psi_1\|_\alpha \|\psi_2\|_\alpha \tau^{n^\gamma}$$

for all  $n \geq 0$ .

In contrast to the Lorentz gas with periodic configurations of scatterers, very little is known about the behavior of the Lorentz gas with aperiodic configurations of

---

*Date:* March 15, 2022.

This work was partially supported by the Simons Collaboration Grant 712227.




FIGURE 1. A trajectory in an aperiodic Lorentz gas.

scatterers which model quasicrystals and other low-complexity aperiodic sets. The work of Marklof-Strömbergsson [MS14, Mar14] studied the Boltzmann-Grad limit, which is the limit as the size of the scatterers goes to zero, for special types of aperiodic scatterer configurations in any dimension. In [TZ21] we proved a particular type of mixing for a large class of aperiodic Lorentz gases in two dimensions, but the question of speed of mixing was left open. Here we prove that exponential mixing never happens for the billiard map of aperiodic Lorentz gases with linearly repetitive scatterer configurations in any dimension.

Let us be more specific: just as there is a compactification of a Lorentz gas with periodic scatterer configuration (which defines the Sinai billiard), there is a compactification of Lorentz gases when the scatterer configuration  $\mathcal{S}$  is aperiodic and of finite local complexity. There are a couple of equivalent ways to do this (see [TZ21]), but the point is that the resulting collision space  $\mathcal{D}$  parametrizes not only a particular aperiodic Lorentz gas but a continuum of Lorentz gases which are locally indistinguishable to the original one. The space  $\mathcal{D}$  is not a manifold but has a nice product structure (see §2 below). If the billiard map  $F : \mathcal{D} \rightarrow \mathcal{D}$  is mixing with respect to some measure  $\mu$ , and the measure  $\mu$  is nice enough, then this implies the **leafwise mixing** of the billiard map of the Lorentz gas (called *global-global mixing* by Lenci [Len10]):

$$\lim_{n \rightarrow \infty} \lim_{R \rightarrow \infty} \frac{1}{\text{Vol}(B_R(0))} \int_{S_+^{d-1}} \int_{B_R(0) \cap \mathcal{S}} g_1 \circ f^n \cdot g_2 d\Xi = \beta(g_1)\beta(g_2),$$

for  $g_i : \partial\mathcal{S} \times S_+^{d-1} \rightarrow \mathbb{R}$ , where  $S_+^{d-1}$  is the upper  $(d-1)$ -sphere, parametrizing directions in the unit tangent bundle, and  $\beta(g_i)$  is the average of  $g_i$  in  $\mathbb{R}^d$ . Here  $g_i$  belong to a special class of functions which contains functions whose values depend on the local patterns of the aperiodic set underlying the aperiodic scatterer configuration  $\mathcal{S}$ , and  $\Xi$  is a smooth invariant measure on the Poincaré section  $\partial\mathcal{S} \times S_+^{d-1}$  of the Lorentz

gas. Exponential mixing of the billiard map  $F : \mathcal{D} \rightarrow \mathcal{D}$  would imply that the rate of convergence in  $n$  of the outer limit above would also be exponential.

In [TZ21] leafwise mixing was proved for the billiard map of repetitive, aperiodic Lorentz gases of finite local complexity with bounded horizon in two dimensions (in fact  $K$ -mixing was proved). The mixing rate was left as an open question. The main result in this paper shows that this cannot be exponentially fast, and in fact it is significantly restricted.

**Theorem 1.** The billiard map of aperiodic Lorentz gases in  $\mathbb{R}^d$  with finite horizon and linearly repetitive scatterer configurations is not mixing with exponential or stretched exponential speed in any  $H_\alpha$ ,  $\alpha > 0$ . Moreover, if for some  $\alpha > 0$  and  $C > 0$  we have that

$$\left| \int_{\mathcal{D}} \psi_1 \circ F^n \cdot \psi_2 d\mu - \mu(\psi_1)\mu(\psi_2) \right| \leq C \|\psi_1\|_\alpha \|\psi_2\|_\alpha n^{-\gamma}$$

for all  $n \in \mathbb{N}$ ,  $\psi_1, \psi_2 \in H_\alpha$ , then  $\gamma \leq 2(1 + \alpha)$ .

**Remark 1.** Some remarks:

- (i) This should be contrasted with the exponential mixing of two-dimensional Sinai billiards, revealing a strong qualitative difference between the periodic and the aperiodic cases in two dimensions.
- (ii) The class of aperiodic scatterer configurations for which we prove this, the so-called linearly repetitive sets, is a very broad class and covers all scatterer configurations that can be built from quasicrystals (cut-and-project sets), substitution tilings, and mixed/S-adic/globally-random substitution tilings.
- (iii) The collision spaces for which these results hold are not locally Euclidean (manifolds). As such, non-constant functions with Hölder exponents  $\alpha > 1$  exist, and this exponent in spirit describes the regularity in the non-Euclidean direction of the space.

This paper is organized as follows. In §1 we review aperiodic Delone sets and the structure of pattern spaces. In §2 we recall the setup of aperiodic Lorentz gases from [TZ21] and their relation to aperiodic Delone sets. In §3 we recall relevant facts about the structure of linearly repetitive Delone sets, following mostly the results of [APC11, APCC+15]. Sections 4 and 5 show how the structure of pattern spaces of linearly repetitive aperiodic sets prohibit fast mixing, that is, they give the proof of the main result.

## 1. BACKGROUND: DELONE MULTISSETS AND REPETITIVITY

A **Delone set** in  $\mathbb{R}^d$  is a countable subset  $\Lambda \subset \mathbb{R}^d$  that is

**uniformly discrete:** there exists an  $r_\Lambda > 0$  such that  $B_{r_\Lambda}(x) \cap \Lambda$  contains at most one point, and

**relatively dense:** there exists a  $R_\Lambda > 0$  such that  $B_{R_\Lambda}(x) \cap \Lambda$  contains at least one point.

These numbers are respectively called the **packing** and **covering** radii, respectively. A **Delone multiset** is a Delone set  $\Lambda$  of the form of a finite union

$$\Lambda = \bigsqcup_i \Lambda_i$$

where each  $\Lambda_i$  is a Delone set and such that the Delone sets are pairwise disjoint. In this case the index  $i$  of a set  $\Lambda$  can be thought of as a label or color. The translation of a Delone set  $\Lambda$  by  $t \in \mathbb{R}^d$  is denoted by  $\varphi_t(\Lambda) := \Lambda - t$ . The set  $\Lambda$  is **aperiodic** if  $\varphi_t(\Lambda) = \Lambda$  implies that  $t = 0$ .

A **cluster** of  $\Lambda$  is a finite subset  $C \subset \Lambda$ . An  **$R$ -cluster** is a cluster of the form  $C = \Lambda \cap B_R(p)$  for some  $p \in \Lambda$ . A Delone set  $\Lambda$  has **finite local complexity** (FLC) if for any  $R > 0$  the collection of all  $R$ -clusters, up to translation equivalence, is finite. A Delone set  $\Lambda$  is **repetitive** if for any  $R$ -cluster  $C \subset \Lambda$  there exists a  $T > 0$  such that for any  $p \in \Lambda$  the  $T$ -cluster around  $p$  contains point  $p'$  such that  $B_R(p') \cap \Lambda$  is translation equivalent to  $C$ .

For a Delone set  $\Lambda$  and some translates  $\varphi_s(\Lambda), \varphi_t(\Lambda)$  consider the quantity

$$d^*(\varphi_s(\Lambda), \varphi_t(\Lambda)) := \inf_{\varepsilon > 0} \{ \text{there exist } x, y \in B_\varepsilon(0) \text{ such that } B_{1/\varepsilon}(0) \cap \varphi_{s+x}(\Lambda) = B_{1/\varepsilon}(0) \cap \varphi_{t+y}(\Lambda) \},$$

and introduce the function  $d$  on the set of all translates

$$d(\varphi_s(\Lambda), \varphi_t(\Lambda)) = \min\{d^*(\varphi_s(\Lambda), \varphi_t(\Lambda)), 2^{-1/2}\}.$$

This function is a metric on this set [LMS02], and we consider the completion, with respect to this metric, of the set of all translates of  $\Lambda$ :

$$\Omega_\Lambda := \overline{\{\varphi_t(\Lambda) : t \in \mathbb{R}^d\}}.$$

This set is called the **pattern space** of  $\Lambda$ . If  $\Lambda$  is repetitive and FLC, which will be assumed throughout the paper, then  $\Omega_\Lambda$  is a compact metric space with an  $\mathbb{R}^d$  action defined by the translation  $\varphi_t$ . The pattern space has a **canonical transversal** defined by the set

$$\mathcal{U}_\Lambda := \{\Lambda' \in \Omega_\Lambda : 0 \in \Lambda'\}.$$

For any  $R > 0$ , the canonical transversal is partitioned into finitely many subsets  $\mathcal{C}_P$ , where  $P$  is an  $R$ -cluster. That is,  $\mathcal{C}_P$  is the set of all  $\Lambda' \in \Omega_\Lambda$  with a point in the origin such that the  $R$ -cluster around the origin is  $P$ . It follows that the space  $\Omega_\Lambda$  has a local product structure of the form  $B_\varepsilon(0) \times \mathcal{C}$ , where  $B_\varepsilon(0)$  is the Euclidean ball of radius  $\varepsilon$  and  $\mathcal{C}$  is a Cantor set.

An invariant measure  $\mu$  for the  $\mathbb{R}^d$  action on  $\Omega_\Lambda$  has a local product structure  $\text{Leb} \times \nu$ , where  $\text{Leb}$  is the Lebesgue measure and  $\nu$  is a transverse invariant measure. The

measure  $\nu$  is defined as follows. Let  $\Lambda' \in \Omega_\Lambda$  and  $A \subset \mathbb{R}^d$  a bounded subset. For a cluster  $P$ , define

$$L_P^{\Lambda'}(A) := |\{t \in \mathbb{R}^d : \varphi_t(P) \subset A \cap \Lambda'\}|$$

to be the number of translates of  $P$  contained in  $A$ . Taking a limit we get the frequency of a cluster

$$\text{freq}_{\Lambda'}(P) = \lim_{R \rightarrow \infty} \frac{L_P^{\Lambda'}(B_R(0))}{\text{Vol}(B_R(0))}.$$

When the frequencies are independent of the set  $\Lambda'$  used to compute them, the system has **uniform patch frequencies**. The frequency map induces a transverse  $\mathbb{R}^d$ -invariant measure  $\nu_{\Lambda'}$  on  $\mathcal{U}_\Lambda$ , defined for a cylinder set  $\mathcal{C}_P$  of an  $R$ -patch  $P$ ,

$$\nu_{\Lambda'}(\mathcal{C}_P) := \text{freq}_{\Lambda'}(P).$$

In the case of uniform patch frequencies this measure is unique and the system is uniquely ergodic. This will be the only relevant case in this paper.

## 2. APERIODIC LORENTZ GASES

Let  $\mathcal{S} \subset \mathbb{R}^d$  be a countable collection of disjoint convex topological balls with  $C^{d-1}$  boundaries, called a set of **scatterers**. The **Lorentz gas** on  $\mathbb{R}^d$  with scatterers  $\mathcal{S}$  is the system describing the free motion of a point particle in  $\mathbb{R}^d$  having elastic collisions at  $\partial\mathcal{S}$ . That is, it is the flow on  $\mathbb{R}^d \times T^1\mathbb{R}^d$  defined by geodesics on  $(\mathbb{R}^d \setminus \mathcal{S}) \times T^1\mathbb{R}^d$  and changing directions at  $\partial\mathcal{S}$ . The system has **finite horizon** if there is a constant  $M$  such that the time between any two collisions is bounded by  $M$ . In this case, the set  $\partial\mathcal{S}$  serves as a Poincaré section for this flow and the Poincaré map for this section is called the **billiard map**  $f : \partial\mathcal{S} \rightarrow \partial\mathcal{S}$ .

To any collection of scatterers we will assign a countable set  $\Lambda_{\mathcal{S}}$  as follows. Define an equivalence relation on the connected components of  $\mathcal{S}$  by  $S_i \sim S_j$  if they are translation equivalent, that is, if  $\varphi_s(S_i) = S_j$  for some  $s \in \mathbb{R}^d$ . Let  $\mathfrak{C}$  be the set of classes of translation-equivalent scatterers. For each  $i \in \mathfrak{C}$ , let  $\Lambda_i^{\mathcal{S}}$  be the union of centers of masses of components in the class  $i$ , and define

$$\Lambda_{\mathcal{S}} := \bigsqcup_{i \in \mathfrak{C}} \Lambda_i^{\mathcal{S}}.$$

**Definition 2.** The scatterer configuration  $\mathcal{S}$  is aperiodic, linearly repetitive, has finite local complexity, etc, if the associated set  $\Lambda_{\mathcal{S}}$  is a Delone multiset with the same properties.

Suppose that  $\mathcal{S}$  is aperiodic, repetitive and of finite local complexity. Then we can identify it with a subset  $\Sigma_{\mathcal{S}} \subset \Omega_{\Lambda_{\mathcal{S}}} := \Omega_{\Lambda_{\mathcal{S}}}$  as follows. Define  $R_{\mathcal{S}} = \{t \in \mathbb{R}^d : \bar{0} \in \varphi_t(\mathcal{S})\}$  to be the set of translates of  $\mathcal{S}$  which cover the origin and note that this contains  $\Lambda_{\mathcal{S}}$ . This defines a subset  $\Sigma' = \{\varphi_t(\Lambda_{\mathcal{S}}) : t \in R_{\mathcal{S}}\} \subset \Omega_{\Lambda_{\mathcal{S}}}$ . This set is defined along the orbit of  $\Lambda_{\mathcal{S}}$ , so taking the closure in  $\Omega_{\Lambda_{\mathcal{S}}}$  we obtain the set  $\Sigma_{\mathcal{S}} = \bar{\Sigma}'$ . As such, the Lorentz gas can be represented as a flow in  $(\Omega_{\Lambda_{\mathcal{S}}} \setminus \Sigma_{\mathcal{S}}) \times T^1\mathbb{R}^d$ . In this space, which involves the translation closure of  $\Lambda_{\mathcal{S}}$ , not only do we find the Lorentz gas defined by  $\mathcal{S}$  but

a continuum of Lorentz gases defined on scatterer configurations which are locally indistinguishable from  $\mathcal{S}$ .

The local product structure of  $\Omega_{\mathcal{S}}$  gives us a nice system of coordinates for the Poincaré map. Denoting by  $S_+^{d-1}$  the (closed) upper  $d - 1$  sphere, let

$$(1) \quad \mathcal{D} := \left( \bigsqcup_{i \in \mathfrak{C}} S_i \times \mathcal{C}_i \right) \times S_+^{d-1},$$

where  $S_i$  is a representative of the scatterer class  $i$  and  $\mathcal{C}_i \subset \mathcal{U}_{\mathcal{S}}$  is the subset of the canonical transversal having a point of  $\Lambda_i^{\mathcal{S}}$  at the origin. Thus a point in  $\mathcal{D}$  records what type of scatterer it is on (the index  $i$ ), where the center of mass of this scatterer is located in  $\Lambda_{\mathcal{S}}$  (the coordinate  $\mathcal{C}_i$ ), where on the scatterer it is (the coordinate in  $S_i$ ), and in which direction it is going (the direction defined by the upper tangent space  $S_+^{d-1}$  at the point of collision). The set  $\mathcal{D}$  is then the **collision space** and the Poincaré map  $F : \mathcal{D} \rightarrow \mathcal{D}$  is defined on this set. An invariant measure  $\mu$  for  $F$  is **absolutely continuous** if it is locally of the form  $L_1 \times \nu \times L_2$ , where  $L_1$  is an absolutely continuous measure on  $S_i$ ,  $\nu$  is a frequency measure, and  $L_2$  is an absolutely continuous measure on  $S_+^{d-1}$ .

### 3. THE STRUCTURE OF LINEARLY REPETITIVE SETS

In this section we recall some properties of linearly repetitive Delone sets which will be crucial to the proof of the main result. Our main reference for this section is [APC11], see also [APCC+15].

The **repetitivity function**  $M_{\Lambda}$  assigns to each  $R > 0$  the least  $T$  such that every ball of radius  $T$  contains a copy of every  $R$ -cluster that is found in  $\Lambda$ . A Delone set  $\Lambda$  is **linearly repetitive** if there exists a  $L > 1$  such that the repetitivity function is bounded as  $M_{\Lambda}(R) \leq LR$ . It is known that linear repetitivity implies unique ergodicity [APCC+15] but this can fail for very mild superlinear repetitivity [LP03, CN16].

For any Delone set  $\Lambda$  and  $R > 0$ , let  $C_{\Lambda,R}$  be the **cylinder** set defined by the  $R$ -neighborhood of the origin in  $\Lambda$ , that is,

$$C_{\Lambda,R} := \{\Lambda' \in \Omega_{\Lambda} : \Lambda' \cap B_R(0) = \Lambda \cap B_R(0)\}.$$

Any clopen subset of a cylinder in  $\Omega_{\Lambda}$  is called a **local transversal**. For any local transversal  $C$ , its **recognition radius** is defined as

$$\text{rec}(C) := \inf\{R > 0 : C_{\Lambda',R} \subset C \text{ for all } \Lambda' \in C\}$$

which is finite for any repetitive Delone set of finite local complexity. If  $C$  is a local transversal and  $D \subset \mathbb{R}^d$ , set

$$C[D] := \{\varphi_t(\Lambda') : t \in D, \Lambda' \in C\}.$$

This type of set is called a **box** if the map from  $D \times C$  to  $B = C[D]$  given by  $(t, \Lambda') \mapsto \varphi_t(\Lambda')$  is a homeomorphism. If  $C$  is a local transversal, then there exists  $r(C) > 0$  such that  $C[D]$  is a box provided that  $D \subset B_{r(C)}(0)$ . A **box decomposition** of  $\Omega_{\Lambda}$  is a finite

collection of pairwise disjoint boxes  $\mathcal{B} = \{B_1, \dots, B_k\}$  such that the closures of the boxes form a cover of  $\Omega_\Lambda$ . In this case, the boxes will be denoted by  $C_i[D_i]$ , where  $C_i$ , called the **base**, is contained in  $B_i$  and  $D_i$  contains the origin.

**Definition 3.** A box decomposition  $\mathcal{B}' = \{C'_i[D'_i]\}_{i=1}^{k'}$  is **zoomed out** of another box decomposition  $\mathcal{B} = \{C_i[D_i]\}_{i=1}^k$  if the following properties hold between them:

- (i) If  $\Lambda' \in C'_i$  satisfies  $\varphi_{x'}(\Lambda') \in \varphi_y(C_j)$  for some  $x' \in \bar{D}'_i$  and  $y \in \bar{D}_j$ , then  $\varphi_{x'}(C'_i) \subset \varphi_y(C_j)$ ;
- (ii) if  $x' \in \partial D'_i$ , there exists  $j$  and  $y \in \partial D_j$  such that  $\varphi_{x'}(C'_i) \subset \varphi_y(C_j)$ ;
- (iii) For every box  $B' \in \mathcal{B}'$  there is  $B \in \mathcal{B}$  such that  $B \cap B' \neq \emptyset$  and  $\partial B \cap \partial B' = \emptyset$ ;
- (iv) If for each  $1 \leq i \leq k'$  and  $1 \leq j \leq k$  we define

$$O_{i,j} = \{x' \in D'_i : \varphi_{x'}(C'_i) \subset C_j\},$$

then for each  $1 \leq i \leq k'$  and  $1 \leq j \leq k$  we have that

$$\bar{D}'_i = \bigcup_{j=1}^k \bigcup_{x' \in O_{i,j}} \varphi_{x'}(\bar{D}_j)$$

where all the sets on the right have pairwise disjoint interiors;

- (v) The base of  $\mathcal{B}'$  is included in the base of  $\mathcal{B}$ , that is,  $\bigcup_i C'_i \subset \bigcup_j C_j$ .

It follows from these properties that we must also have that

$$(2) \quad C_j = \bigcup_{i=1}^{k'} \bigcup_{x' \in O_{i,j}} \varphi_{x'}(C'_i).$$

A **tower system** is a sequence of box decompositions  $\{\mathcal{B}_n\}$  such that  $\mathcal{B}_{n+1}$  is zoomed out of  $\mathcal{B}_n$ .

Tower systems give Delone sets a hierarchical structure and they always exist for pattern spaces of aperiodic, repetitive Delone sets. In the linearly repetitive case, they have a particular nice structure which we will use.

If  $\{C_n\}$  is a decreasing sequence of local transversals with diameter going to zero and  $\{\mathcal{B}_n\}$  is a tower system, then  $\{\mathcal{B}_n\}$  is **adapted to**  $\{C_n\}$  if we have that  $\mathcal{B}_n = \{C_{n,i}[D_{n,i}]\}_{i=1}^{k_n}$  such that  $C_n = \bigcup_i C_{n,i}$ . In such case define

$$O_{i,j}^{(n)} = \{x \in D_{n,i} : \varphi_x(C_{n,i}) \subset C_{n-1,j}\} \quad \text{and} \quad m_{i,j}^{(n)} = |O_{i,j}^{(n)}|.$$

The numbers  $m_{i,j}^{(n)}$  are the entries of a  $k_{n-1} \times k_n$  matrix, called the **transition matrix**. Since each local transversal  $C_{n,i}$  is defined by the translation-equivalence class of some cluster  $P_{n,i}$ , these numbers record the number of clusters of type  $P_{n-1,j}$  which are found in the cluster of type  $P_{n,i}$ .

Given a box decomposition  $\mathcal{B} = \{C_i[D_i]\}_{i=1}^k$  its **internal and external radii** are defined respectively as

$$\begin{aligned} r_{int}(\mathcal{B}) &= \min_{i \in \{1, \dots, k\}} \sup\{R > 0 : B_R(0) \subset D_i\}, \\ R_{ext}(\mathcal{B}) &= \max_{i \in \{1, \dots, k\}} \inf\{R > 0 : D_i \subset B_R(0)\}, \\ \text{rec}(\mathcal{B}) &= \max_{i \in \{1, \dots, k\}} \text{rec}(C_i). \end{aligned}$$

We can now state the main theorem on the structure of linearly repetitive Delone sets.

**Theorem** ([APC11]). Let  $\Lambda$  be an aperiodic linearly repetitive Delone set with repetitivity constant  $L > 1$  and  $0 \in \Lambda$ . Define  $\lambda = 6L(L + 1)^2$  and  $C_n := C_{\Lambda, \lambda^n}$  for all  $n > 0$ . Then there exists a tower system  $\{\mathcal{B}_n\}$  of  $\Omega_\Lambda$  adapted to  $\{C_n\}$  such that

- (i)  $C_{n+1} \subset C_{n,1}$
- (ii) setting

$$K_1 := \frac{1}{2(L+1)} - \frac{L}{\lambda-1} \text{ and } K_2 := \frac{\lambda L}{\lambda-1}$$

we have that  $0 < K_1 < 1 < K_2$  and moreover

$$K_1 \lambda^n \leq r_{int}(\mathcal{B}_n) < R_{ext}(\mathcal{B}_n) \leq K_2 \lambda^n,$$

- (iii)  $\text{rec}(\mathcal{B}_n) \leq (2L + 1)\lambda^n$

for all  $n$ . In addition, the transition matrices have strictly positive entries and are uniformly bounded in size and norm.

If  $\mathcal{B} = \{C_{n,i}[D_{n,i}]\}$  is a tower system adapted to some  $\{C_n\} = \{C_{\Lambda, \lambda^n}\}$  as in the theorem above, we will refer to it as a **nice tower system**. Observe that the bounds on the covering and packing radii  $r_{int}$  and  $R_{ext}$  for clusters which give the  $n^{\text{th}}$  box decomposition imply that for all  $T > \lambda^n$

$$K_3' \frac{\text{Vol}(B_T(0))}{K_2 \lambda^n} \leq L_{B_{\lambda^n}(0) \cap \Lambda}^\Lambda(B_T(0)) \leq K_4' \frac{\text{Vol}(B_T(0))}{K_1 \lambda^n},$$

where the constants  $K_3', K_4'$  depend only on the dimension. From these observations we obtain the following corollary.

**Corollary 1.** Let  $\{\mathcal{B}_n\}$  be a tower system adapted to  $\{C_n\}$  for an aperiodic, linearly-repetitive Delone set of the type given by the theorem above. There exists  $K_3 > 1$  such that

$$K_3^{-1} \lambda^{-n} \leq \nu(C_{n,i}) \leq K_3 \lambda^{-n}$$

for all  $1 \leq i \leq k_n$ , where  $\nu$  is the unique transverse  $\mathbb{R}^d$ -invariant measure on  $\Omega_\Lambda$ .

Given a tower system  $\mathcal{B} = \{C_{n,i}[D_{n,i}]\}$  as in the theorem above adapted to  $\{C_n\} = \{C_{\Lambda, \lambda^n}\}$ , let  $B_{n,i}$  be a largest circumscribed ball inside  $D_{n,i}$  centered at a point  $p_{n,i} \in$



$D_{n,i}$  of the cluster inside  $D_{n,i}$ . Then  $B_{n,i}$  has radius  $R \geq K_1\lambda^n - 2R_\Lambda$ . We can change coordinates for our tower system and write it as

$$(3) \quad \mathcal{B}' = \{C'_{n,i}[D'_{n,i}]\} = \{\varphi_{p_{n,i}}(C_{n,i}[D_{n,i}])\}$$

so that  $D'_{n,i}$  contains a ball of radius  $K_1\lambda^n - 2R_\Lambda$  centered at the origin for every  $i$ . Defining

$$C'_n = \bigcup_i C'_{n,i},$$

the set of return vectors to  $C'_n$

$$\mathcal{R}_n := \{t \in \mathbb{R}^d : \varphi_t(\Lambda') \in C'_n \text{ for some } \Lambda' \in C'_n\}$$

is a Delone set with packing radius  $r_{\mathcal{R}_n}$  greater than  $2(K_1\lambda^n - 2R_\Lambda)$ , since it was obtained by modifying the box decomposition given by the tower system of the theorem above. That is

$$(4) \quad 2(K_1\lambda^n - 2R_\Lambda) \leq r_{\mathcal{R}_n}.$$

#### 4. FAR AWAY FUNCTIONS

Consider an aperiodic, linearly repetitive collection of scatterers  $\mathcal{S} \subset \mathbb{R}^d$ , their corresponding aperiodic, linearly repetitive Delone set  $\Lambda_{\mathcal{S}}$ , its nice tower system  $\mathcal{B}_{\mathcal{S}}$ , and its modified tower system  $\mathcal{B}'_{\mathcal{S}}$  as in (3). Using the coordinates (1), and the modified tower define the sets

$$A_{n,i} := S_{j_{n,i}} \times C'_{n,i} \times S_+^{d-1} \subset \mathcal{D},$$

where  $j_{n,i} \in \mathfrak{C}$  is the index associated to the point defining the base of the transversal in  $C'_{n,i}[D'_{n,i}]$ . Define the functions

$$\psi_i^{(n)} := \chi_{A_{n,i}}.$$

We want to estimate the Hölder norm of these functions, that is, we are interested in bounding

$$\|\psi_i^{(n)}\|_\alpha := \sup_{(p,q \neq p) \in \mathcal{D}^2} \frac{|\psi_i^{(n)}(p) - \psi_i^{(n)}(q)|}{\text{dist}(p,q)^\alpha}.$$

for  $\alpha > 0$ . By definition, the quantity  $|\psi_i^{(n)}(p) - \psi_i^{(n)}(q)|/\text{dist}(p,q)^\alpha$  is nonzero only when  $p$  is in  $A_{n,i}$  and  $q$  is not (or vice-versa). In that case, if  $p \in A_{n,i}$  and  $q \notin A_{n,i}$ , they are closest whenever they have the same smooth coordinates, and the coordinates in the transversal direction are as close as possible.

By the estimates on the box decompositions of the tower system, they cannot have the same  $K_2\lambda^n$  neighborhood around the origin as otherwise they would have to be in the same box. This means that  $\text{dist}(p,q) \geq (K_2\lambda^n)^{-1}$  in this case, and so for any  $i$  we have

$$(5) \quad \|\psi_i^{(n)}\|_\alpha \leq K_2\lambda^{\alpha n}.$$

## 5. NON-FAST MIXING

Here we prove the main result. Let  $\psi_i^{(n)}$  and  $\psi_j^{(n)}$  be two functions constructed from a nice box decomposition as in the previous section and consider the correlation

$$\int_{\mathcal{D}} \psi_i^{(n)} \circ F^k \cdot \psi_j^{(n)} d\mu,$$

where  $\mu$  is any absolutely continuous  $F$ -invariant finite measure on  $\mathcal{D}$ . Let  $P_{n,i}$  be the clusters defined by the modified tower system, that is,

$$P_{n,i} := \{t \in D'_{n,i} : \varphi_t(\Lambda') \in \mathcal{U}_\Lambda \text{ for every } \Lambda' \in C'_{n,i}\}.$$

Since the function  $\psi_i^{(n)}$  is supported on scatterers whose  $(K_1\lambda^n - 2R_\Lambda)$ -neighborhood is determined by the cluster  $P_{n,i}$ , and the distance between two points at the centers of different clusters is at least  $2(K_1\lambda^n - 2R_\Lambda)$  by (4),  $\psi_i^{(n)} \circ F^k(x) \cdot \psi_j^{(n)}(x) = 0$  for all the  $k$  so that no orbit starting at the center of a cluster of the form  $P_{n,j}$  reaches a scatterer at the center of the cluster of the form  $P_{n,i}$  after  $k$  collisions. By finite horizon, a trajectory can travel at most a distance of  $Mk$  after  $k$  collisions. Thus,

$$\psi_i^{(n)} \circ F^k(x) \cdot \psi_j^{(n)}(x) = 0 \quad \text{for all} \quad |k| < \frac{2}{M}(K_1\lambda^n - 2R_\Lambda).$$

So we have that

$$(6) \quad \left| \int_{\mathcal{D}} \psi_i^{(n)} \circ F^k \cdot \psi_j^{(n)} d\mu - \mu(\psi_i^{(n)})\mu(\psi_j^{(n)}) \right| = \left| \mu(\psi_i^{(n)})\mu(\psi_j^{(n)}) \right| \geq K_3^{-2}\lambda^{-2n}$$

for  $|k| < \frac{2}{M}(K_1\lambda^n - 2R_\Lambda)$  by Corollary 1.

We now show how exponential mixing is incompatible with the nice tower system which linearly repetitive systems have. Suppose that for some  $\alpha > 0$  and  $\tau \in (0, 1)$  there is a  $C > 0$  such that

$$\left| \int_{\mathcal{D}} h_1 \circ F^k \cdot h_2 d\mu - \mu(h_1)\mu(h_2) \right| \leq C\|h_1\|_\alpha\|h_2\|_\alpha\tau^k$$

for all  $k \in \mathbb{N}$  and  $h_i \in H_\alpha$ . Using the functions  $\psi_i^{(n)}$  we constructed above, by (5) and (6), we would have

$$K_3^{-2}\lambda^{-2n} \leq C\|\psi_i^{(n)}\|_\alpha\|\psi_j^{(n)}\|_\alpha\tau^k \leq CK_2^2\lambda^{2\alpha n}\tau^k$$

for all  $|k| < \frac{2}{M}(K_1\lambda^n - 2R_\Lambda)$ . In particular this should hold for  $k = \lfloor \frac{2}{M}(K_1\lambda^n - 2R_\Lambda) \rfloor$ , so

$$K_3^{-2}\lambda^{-2n} \leq CK_2^2\lambda^{2\alpha n}\tau^{\lfloor \frac{2}{M}(K_1\lambda^n - 2R_\Lambda) \rfloor}.$$

This is equivalent to

$$\log(C^*) - 2n(1 + \alpha) \log \lambda \leq \left\lfloor \frac{2}{M}(K_1\lambda^n - 2R_\Lambda) \right\rfloor \log \tau.$$

Now, the right hand side is negative and decreasing at an exponential rate with  $n$ , while the left hand side is negative and decreasing at a linear rate. This of course

cannot happen for all  $n$ , and so this obstructs the system from mixing at an exponential rate. The same argument prevents the system from mixing with any stretched exponential rate.

Polynomial mixing is equally restricted by the nice tower system which linearly repetitive systems have. Suppose that for some  $\alpha > 0$  and  $\gamma > 0$  there is a  $C > 0$  such that

$$\left| \int_{\mathcal{D}} h_1 \circ F^k \cdot h_2 d\mu - \mu(h_1)\mu(h_2) \right| \leq C \|h_1\|_{\alpha} \|h_2\|_{\alpha} |k|^{-\gamma}$$

for all  $k \in \mathbb{Z}$  and  $h_i \in H_{\alpha}$ . Using the functions  $\psi_i^{(n)}$  we constructed above, by (5) and (6), we would have

$$K_3^{-2} \lambda^{-2n} \leq C \|\psi_i^{(n)}\|_{\alpha} \|\psi_j^{(n)}\|_{\alpha} k^{-\gamma} \leq CK_2^2 \lambda^{2\alpha n} k^{-\gamma}$$

for all  $|k| < \frac{2}{M}(K_1 \lambda^n - 2R_{\Lambda})$ . In particular this should hold for  $k = \lfloor \frac{2}{M}(K_1 \lambda^n - 2R_{\Lambda}) \rfloor$ , so

$$K_3^{-2} \lambda^{-2n} \leq CK_2^2 \lambda^{2\alpha n} \left[ \frac{2}{M}(K_1 \lambda^n - 2R_{\Lambda}) \right]^{-\gamma}.$$

For sufficiently large  $n$  this implies the inequality

$$C^{**} \lambda^{-2(1+\alpha)n} \leq \lambda^{-\gamma n}.$$

This cannot hold for all  $n$  unless  $\gamma \leq 2(1 + \alpha)$ .

## REFERENCES

- [APC11] José Aliste-Prieto and Daniel Coronel, *Tower systems for linearly repetitive Delone sets*, Ergodic Theory Dynam. Systems **31** (2011), no. 6, 1595–1618. MR 2851668
- [APCC+15] José Aliste-Prieto, Daniel Coronel, María Isabel Cortez, Fabien Durand, and Samuel Petite, *Linearly repetitive Delone sets*, Mathematics of aperiodic order, Progr. Math., vol. 309, Birkhäuser/Springer, Basel, 2015, pp. 195–222. MR 3381482
- [BSC90] L. A. Bunimovich, Ya. G. Sinai, and N. I. Chernov, *Markov partitions for two-dimensional hyperbolic billiards*, Uspekhi Mat. Nauk **45** (1990), no. 3(273), 97–134, 221. MR 1071936
- [CM06] Nikolai Chernov and Roberto Markarian, *Chaotic billiards*, Mathematical Surveys and Monographs, vol. 127, American Mathematical Society, Providence, RI, 2006. MR 2229799
- [CN16] María Isabel Cortez and Andrés Navas, *Some examples of repetitive, nonrectifiable Delone sets*, Geom. Topol. **20** (2016), no. 4, 1909–1939. MR 3548461
- [Det14] Carl P. Dettmann, *Diffusion in the Lorentz gas*, Commun. Theor. Phys. (Beijing) **62** (2014), no. 4, 521–540. MR 3363162
- [Gol12] François Golse, *Recent results on the periodic Lorentz gas*, Nonlinear partial differential equations, Adv. Courses Math. CRM Barcelona, Birkhäuser/Springer Basel AG, Basel, 2012, pp. 39–99. MR 3059762
- [Len10] Marco Lenci, *On infinite-volume mixing*, Comm. Math. Phys. **298** (2010), no. 2, 485–514. MR 2669446
- [LMS02] J.-Y. Lee, R. V. Moody, and B. Solomyak, *Pure point dynamical and diffraction spectra*, Ann. Henri Poincaré **3** (2002), no. 5, 1003–1018. MR 1937612
- [LP03] Jeffrey C. Lagarias and Peter A. B. Pleasants, *Repetitive Delone sets and quasicrystals*, Ergodic Theory Dynam. Systems **23** (2003), no. 3, 831–867. MR 1992666

- [Mar14] Jens Marklof, *The low-density limit of the Lorentz gas: periodic, aperiodic and random*, Proceedings of the International Congress of Mathematicians—Seoul 2014. Vol. III, Kyung Moon Sa, Seoul, 2014, pp. 623–646. MR 3729044
- [MS14] Jens Marklof and Andreas Strömbergsson, *Free path lengths in quasicrystals*, Comm. Math. Phys. **330** (2014), no. 2, 723–755. MR 3223485
- [Sz08] Domokos Szász, *Some challenges in the theory of (semi)-dispersing billiards*, Nonlinearity **21** (2008), no. 10, T187–T193. MR 2439473
- [TZ21] Rodrigo Treviño and Agnieszka Zelerowicz, *Statistical properties of Lorentz gases on aperiodic tilings, part 1*, 2021.
- [You98] Lai-Sang Young, *Statistical properties of dynamical systems with some hyperbolicity*, Ann. of Math. (2) **147** (1998), no. 3, 585–650. MR 1637655

UNIVERSITY OF MARYLAND, COLLEGE PARK  
Email address: [rodrigo@umd.edu](mailto:rodrigo@umd.edu)

UNIVERSITY OF MARYLAND, COLLEGE PARK  
Email address: [azelerow@umd.edu](mailto:azelerow@umd.edu)