Rank Two Approximations of $2 \times 2 \times 2$ Tensors over \mathbb{R}

David Warren Katz

We provide a coordinate-free proof that real $2 \times 2 \times 2$ rank three tensors do not have optimal rank two approximations with respect to the Frobenius norm. This result was first proved in [1, Thm. 8.1] by considering the $GL(V^1) \times GL(V^2) \times GL(V^3)$ orbit classes of $V^1 \otimes V^2 \otimes V^3$ and the $2 \times 2 \times 2$ hyperdeterminant. Our coordinate-free proof expands on the result in [1] by developing a proof method that can be generalized more readily to higher dimensional $n_1 \times n_2 \times n_3$ tensor spaces.

1 Introduction

Let V^1 , V^2 , and V^3 be two-dimensional real vector spaces, respectively, and let $V^1 \otimes V^2 \otimes V^3$ denote the tensor product of these spaces. The tensors in the form $v^1 \otimes v^2 \otimes v^3$ for some vectors v^1 , v^2 , and v^3 are called simple tensors. Every tensor can be written as the sum of finitely many simple tensors. The rank of a tensor τ is the minimum number n such that τ is the sum of n simple tensors. That is,

$$\mathsf{rk}(\tau) \ = \ \min\{n \in \mathbb{N} \ | \ \tau \ = \sum_{i=1}^n \ v_i^1 \otimes v_i^2 \otimes v_i^3 \ \text{ for some } v_i^s \in V^s\}.$$

Once bases $\{e_t^s\}_{t=1}^2$ of V^s are chosen for s = 1, 2, 3, a tensor in $V^1 \otimes V^2 \otimes V^3$ can be coordinitized as a hypermatrix in $\mathbb{R}^{2 \times 2 \times 2}$ by the isomorphism from $V^1 \otimes V^2 \otimes V^3$ to $\mathbb{R}^{2 \times 2 \times 2}$ defined on simple tensors as

for some real constants a_t^s . A lot of information about the rank of the $2 \times 2 \times 2$ real hypermatrix

$$(a_{ijk}) = \underbrace{\begin{vmatrix} & & & \\ & &$$

can be inferred from sign of the polynomial

L

$$\Delta(a_{ijk}) = (a_{111}^2 a_{222}^2 + a_{112}^2 a_{221}^2 + a_{121}^2 a_{212}^2 + a_{122}^2 a_{211}^2) - 2(a_{111}a_{112}a_{221}a_{222} + a_{111}a_{121}a_{212}a_{222} + a_{111}a_{122}a_{211}a_{222}) - 2(a_{112}a_{121}a_{212}a_{221} + a_{112}a_{122}a_{221}a_{211} + a_{121}a_{122}a_{212}a_{211}) + 4(a_{111}a_{122}a_{212}a_{221} + a_{112}a_{121}a_{211}a_{222}).$$

The polynomial Δ is called the $2 \times 2 \times 2$ hyperdeterminant. $\Delta(a_{ijk}) > 0$ implies hypermatrix (a_{ijk}) is rank two, and $\Delta(a_{ijk}) < 0$ implies (a_{ijk}) is rank three. The existence of such a polynomial is not particular to the $2 \times 2 \times 2$ case. The sets of constant rank real hypermatrices are semialgebraic for any dimensions $n_1 \times n_2 \times \cdots \times n_d$. Hence, the rank of a real hypermatrix can always be computed by evaluating a finite number of polynomial equalities and inequalities, where the variables of the polynomials are the entries of the hypermatrix. Because of this, real tensors are usually studied in coordinates by representing tensors as hypermatrices and then exploiting properties of these polynomials. The polynomials that define the sets of constant rank $2 \times 2 \times 2$ tensors are different, however, from the polynomials that define the sets of constant rank $n_1 \times n_2 \times n_3$ tensors for any other dimensions $n_1 \times n_2 \times n_3$. As a result, coordinate proofs about $2 \times 2 \times 2$ tensors that rely on these polynomials cannot be readily generalized to the arbitrary $n_1 \times n_2 \times n_3$ case. In this paper, we take an alternative coordinate-free approach in our study of $2 \times 2 \times 2$ tensors that lends itself more readily to generalization. We prove that real $2 \times 2 \times 2$ rank three tensors do not have optimal rank two approximations with respect to the Frobenius norm without relying on hypermatrices or the hyperdeterminant.

Matrix rank is lower semi-continuous for matrices of real numbers. That is, if a sequence of rank r real matrices converges to a matrix A in the norm topology, then the rank of A is less than or equal to r. This guarantees the existence of optimal low-rank approximations of real matrices. A sequence of real hypermatrices, however, can converge to a hypermatrix of greater rank. As a result, optimal low rank approximations may not exist for real hypermatrices. In fact, the set of real hypermatrices with no optimal low rank approximation often has positive Lebesgue measure [2], so characterizing such real hypermatrices is an important step in implementing algorithms with real hypermatrices.

Let $\tau \in V^1 \otimes V^2 \otimes V^3$ be a tensor of rank r, and let $s \leq r$. An optimal rank s approximation of τ with respect to norm $\|\cdot\|$ is a tensor v such that

$$\|\tau - v\| = \inf_{\substack{v' \in V^1 \otimes V^2 \otimes V^3 \\ \mathrm{rk}(v') \le s}} \|\tau - v'\|$$

It is a classical observation that tensors in the form

$$\beta = x_2^1 \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes x_2^2 \otimes x_1^3 + x_1^1 \otimes x_2^1 \otimes x_2^3$$
(1)

are limit points of the sequence of rank at most two tensors

$$\tau_n = n \left(x_1^1 + \frac{1}{n} x_2^1 \right) \otimes \left(x_1^2 + \frac{1}{n} x_2^2 \right) \otimes \left(x_1^3 + \frac{1}{n} x_2^3 \right) - n x_1^1 \otimes x_1^2 \otimes x_1^3, \ n \in \mathbb{N}$$

$$\tag{2}$$

in the norm topology. It follows from the triangle inequality and the multilinearity of the tensor product that

$$\|\tau_n - \beta\| \le \frac{1}{n} \|x_2^1 \otimes x_2^2 \otimes x_1^3 + x_2^1 \otimes x_1^2 \otimes x_2^3 + x_1^1 \otimes x_2^2 \otimes x_2^3\| + \frac{1}{n^2} \|x_2^1 \otimes x_2^2 \otimes x_2^3\|.$$

Hence, $\lim_{n\to\infty} ||\tau_n - \beta|| = 0$, so $\lim_{n\to\infty} \tau_n$ indeed equals β . In Section 2, we give a coordinate-free proof that tensors in the form of (2) are in fact rank three when the sets $\{x_1^1, x_2^1\}$, $\{x_1^2, x_2^2\}$, and $\{x_1^3, x_2^3\}$ are independent. Thus, when $\{x_1^1, x_2^1\}$, $\{x_1^2, x_2^2\}$, and $\{x_1^3, x_2^3\}$ are independent, β is an example of a rank three tensor with no an optimal rank two approximation. The main result of this paper is a coordinate-free proof that in fact *every* rank three $2 \times 2 \times 2$ tensor has no optimal rank two approximation with respect to the Frobenius norm, not just the tensors in the form of (1).

This result was first proved in coordinates in [1, Thm. 8.1]. The argument in [1] can be summarized as follows. Suppose for contradiction that the $2 \times 2 \times 2$ real hypermatrix B is an optimal rank two approximation of rank three $2 \times 2 \times 2$ real hypermatrix A with respect to the Frobenius norm. By considering properties of the polynomial Δ , it follows that $\Delta(B) = 0$. The polynomial Δ is invariant on the $GL(V^1) \times GL(V^2) \times GL(V^3)$ orbit classes of $V^1 \otimes V^2 \otimes V^3$, and only three of the eight orbit classes are zero on Δ . Hypermatrices in these three orbit classes are equivalent up to an orthogonal change of coordinates to hypermatrices in the form

for some real λ and μ . Since B is rank two, we may thus assume B is in form (3) with both λ and μ nonzero. Finally, it is shown that if H is a $2 \times 2 \times 2$ hypermatrix such that

$$\Delta(B + \epsilon H) = 0 \tag{4}$$

for all real ϵ , then A - B is orthogonal to H. By considering various hypermatrices H that satisfy (4), the authors of [1] then conclude that

$$A - B = \begin{pmatrix} & & & \\ & & & & \\ & & & &$$

for some constant a. This implies that

$$A = \begin{pmatrix} & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & & \\ & & & &$$

However, this hypermatrix is rank two, contradicting that A is rank three.

Our coordinate-free proof expands on the result in [1] by developing a proof method that can be generalized more readily to higher dimensional tensor spaces. Our proof is also a proof by contradiction. We suppose for contradiction that rank two tensor v in $V^1 \otimes V^2 \otimes V^3$ is an optimal rank two approximation of rank three tensor τ . By considering the relationship between the mode-1 contraction maps of τ and v, we also derive the contradiction that τ is rank two. Our proof has the interesting geometric corollary that the nearest point of a rank three tensor to the second secant variety of the Segre variety is a rank three tensor in the tangent space of the Segre variety.

2 Contraction Maps of Tensors

Let V^1 , V^2 , and V^3 be finite n_1 , n_2 , and n_3 -dimensional real vector spaces, and let V^{1*} , V^{2*} , and V^{3*} denote their dual spaces. The tensor $\tau = \sum_{i=1}^r v_i^1 \otimes v_i^2 \otimes v_i^3 \in V^1 \otimes V^2 \otimes V^3$ induces the three linear maps

$$\begin{array}{cccc} V^{1*} \xrightarrow{\Pi_1(\tau)} V^2 \otimes V^3 & V^{2*} \xrightarrow{\Pi_2(\tau)} V^1 \otimes V^3 & V^{3*} \xrightarrow{\Pi_3(\tau)} V^1 \otimes V^2 \\ v^{1*} \mapsto \sum_{i=1}^r v^{1*}(v_i^1) v_i^2 \otimes v_i^3, & v^{2*} \mapsto \sum_{i=1}^r v^{2*}(v_i^2) v_i^1 \otimes v_i^3, & \text{and} & v^{3*} \mapsto \sum_{i=1}^r v^{3*}(v_i^3) v_i^1 \otimes v_i^2. \end{array}$$

These maps are called the mode-1, mode-2, and mode-3 contraction maps of τ , respectively.

Theorem 1. The mode-*i* contraction map Π_i is well-defined.

Proof. Suppose that $\sum_{t=1}^{r} v_t^1 \otimes v_t^2 \otimes v_t^3 = \sum_{t=1}^{s} w_t^1 \otimes w_t^2 \otimes w_t^3$ for vectors $v_t^i, w_t^i \in V^i$. We need to show that

$$\Pi_i \left(\sum_{t=1}^r v_t^1 \otimes v_t^2 \otimes v_t^3 \right) (v^{i*}) = \Pi_i \left(\sum_{t=1}^s w_t^1 \otimes w_t^2 \otimes w_t^3 \right) (v^{i*})$$

for every $v^{i*} \in V^{i*}$ for i = 1, 2 and 3. Choose bases $\{e_t^i\}_{t=1}^{n_i}$ of V^i with corresponding dual bases $\{e_t^{i*}\}_{t=1}^{n_i}$ for each i. It is sufficient to show that

$$\Pi_1\left(\sum_{t=1}^r v_t^1 \otimes v_t^2 \otimes v_t^3\right)(e_u^{1*}) = \Pi_1\left(\sum_{t=1}^s w_t^1 \otimes w_t^2 \otimes w_t^3\right)(e_u^{1*})$$

for each e_u^{1*} in our dual basis. Let $a_{t,u}^i$ and $b_{t,u}^i$ be scalars such that $v_t^i = \sum_{u=1}^{n_i} a_{t,u}^i e_u^i$ and $w_t^i = \sum_{u=1}^{n_i} b_{t,u}^i e_u^i$ for all i and t. It follows that $\sum_{t=1}^r v_t^1 \otimes v_t^2 \otimes v_t^3 = \sum_{t,i,j,k=1}^{r,n_1,n_2,n_3} a_{ti}^1 a_{tj}^2 a_{tk}^3 e_i^1 \otimes e_j^2 \otimes e_k^3$, which must equal $\sum_{t=1}^s w_t^1 \otimes w_t^2 \otimes w_t^3 = \sum_{t,i,j,k=1}^{r,n_1,n_2,n_3} a_{ti}^1 a_{tj}^2 a_{tk}^3 e_i^1 \otimes e_j^2 \otimes e_k^3$.

$$\begin{split} \sum_{t,i,j,k=1}^{s,n_1,n_2,n_3} b_{ti}^1 b_{tj}^2 b_{tk}^3 e_i^1 \otimes e_j^2 \otimes e_k^3, & \text{so} \sum_{t=1}^r a_{ti}^1 a_{tj}^2 a_{tk}^3 = \sum_{t=1}^s b_{ti}^1 b_{tj}^2 b_{tk}^3 \text{ for all } i, j, k. \text{ Hence,} \\ \Pi_1 \left(\sum_{t=1}^r v_t^1 \otimes v_t^2 \otimes v_t^3 \right) (e_u^{1*}) &= \sum_{t,j,k=1}^{r,n_2,n_3} a_{tu}^1 a_{tj}^2 a_{tk}^3 e_j^2 \otimes e_k^3 \\ &= \sum_{j,k=1}^{n_2,n_3} \left(\sum_{t=1}^r a_{tu}^1 a_{tj}^2 a_{tk}^3 \right) e_j^2 \otimes e_k^3 \\ &= \sum_{j,k=1}^{n_2,n_3} \left(\sum_{t=1}^s b_{tu}^1 b_{tj}^2 b_{tk}^3 \right) e_j^2 \otimes e_k^3 \\ &= \sum_{t,j,k=1}^{s,n_2,n_3} b_{tu}^1 b_{tj}^2 b_{tk}^3 e_j^2 \otimes e_k^3 \\ &= \prod_1 \left(\sum_{t=1}^s w_t^1 \otimes w_t^2 \otimes w_t^3 \right) (e_u^{1*}). \end{split}$$

The relationship between the contraction maps of a tensor generalizes, in a coordinate-free way, the fundamental relationship between the rows and columns of a matrix to hypermatrices. The rank of a two-fold tensor $\tau \in V^1 \otimes V^2$ is equal to the dimension of the image of $\Pi_1(\tau)$, which is also equal to the dimension of the image of $\Pi_2(\tau)$. We now use contraction maps to show when tensors in the form (1) are rank three. First, we need the following lemma.

Lemma 2. Let V^1 and V^2 be finite dimensional real vector spaces, let $\{x_i^1\}_{i=1}^r$ and $\{y_i^1\}_{i=1}^s$ be linearly independent subsets of V^1 , and let $\{x_i^2\}_{i=1}^r$ and $\{y_i^2\}_{i=1}^s$ be linearly independent subsets of V^2 . If $\sum_{i=1}^r x_i^1 \otimes x_i^2 = \sum_{i=1}^s y_i^1 \otimes y_i^2$, then r = s.

Proof. Since $\{x_i^1\}_{i=1}^r$ is independent, we can choose $\{x_j^{1*}\}_{j=1}^r \subseteq V^{1*}$ such that $x_j^{1*}(x_i^1) = \delta_{ij}$. By taking the mode-1 contraction of the tensor with respect to each of the two representations, it follows that

$$\sum_{i=1}^{r} x_{j}^{1*}(x_{i}^{1}) x_{i}^{2} = \sum_{i=1}^{s} x_{j}^{1*}(y_{i}^{1}) y_{i}^{2} \text{ for } j = 1, 2, \dots, r.$$

This implies $x_j^2 = \sum_{i=1}^s x_j^{1*}(y_i^1) y_i^2$ for j = 1, 2, ..., r, so each $x_j^2 \in \langle y_i^2 \rangle_{i=1}^s$. Thus, we conclude that the linear space $\langle x_j^2 \rangle_{j=1}^r$ is a subset of $\langle y_i^2 \rangle_{i=1}^s$. Similarly, $\langle y_i^2 \rangle_{i=1}^s \subseteq \langle x_j^2 \rangle_{j=1}^r$, so by the independence of each set, r = s.

Theorem 3. Let V^1 , V^2 , and V^3 be finite dimensional real vector spaces. If the sets $\{x_1^1, x_2^1\}$, $\{x_1^2, x_2^2\}$, and $\{x_1^3, x_2^3\}$ in V^1 , V^2 , and V^3 , respectively, are linearly independent, then the tensor

$$\beta = x_2^1 \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes x_2^2 \otimes x_1^3 + x_1^1 \otimes x_1^2 \otimes x_2^3$$

is rank three.

Proof. Suppose for contradiction that β were rank less than three. That is, suppose that

$$eta \ = \ y_1^1 \otimes y_1^2 \otimes y_1^3 \ + \ y_2^1 \otimes y_2^2 \otimes y_2^3$$

for some vectors y_j^i . Since im $\Pi_i(\beta) = \langle x_1^i, x_2^i \rangle = \langle y_1^i, y_2^i \rangle$ for each *i*, it follows that the sets $\{y_1^i, y_2^i\}$ are independent for each *i*. Let $\{x_1^{1*}, x_2^{1*}\}$ be the dual basis of the basis $\{x_1^1, x_2^1\}$ of the subspace $\langle x_1^1, x_2^1 \rangle$ of V^1 . By considering the mode-1 contraction of both representations of β , it follows that

$$\Pi_1(\beta)(x_2^{1*}) = x_2^{1*}(x_2^1)x_1^2 \otimes x_1^3 + x_2^{1*}(x_1^1)x_2^2 \otimes x_1^3 + x_2^{1*}(x_1^1)x_1^2 \otimes x_2^3 = x_2^{1*}(y_1^1)y_1^2 \otimes y_1^3 + x_2^{1*}(y_2^1)y_2^2 \otimes y_2^3.$$

This implies that

This implies that

$$x_1^2 \otimes x_1^3 = x_2^{1*}(y_1^1)y_1^2 \otimes y_1^3 + x_2^{1*}(y_2^1)y_2^2 \otimes y_2^3.$$

Since the sets $\{y_1^i, y_2^i\}$ are independent, Lemma 2 implies that either $x_2^{1*}(y_1^1) = 0$ or $x_2^{1*}(y_2^1) = 0$. We consider the case when $x_2^{1*}(y_2^1) = 0$, and leave the remaining similar case to the reader. It follows that

$$\operatorname{\mathsf{im}} \Pi_2(\Pi_1(\beta)(x_2^{1*})) = \operatorname{\mathsf{im}} \Pi_2(x_1^2 \otimes x_1^3) = \operatorname{\mathsf{im}} \Pi_2(x_2^{1*}(y_1^1)y_1^2 \otimes y_1^3),$$

which implies that $\langle x_1^2 \rangle = \langle y_1^2 \rangle$. Let k be a scalar such that $y_1^2 = kx_1^2$. Similarly,

$$\operatorname{im}\Pi_1(\Pi_1(\beta)(x_2^{1*})) = \operatorname{im}\Pi_1(x_1^2 \otimes x_1^3) = \operatorname{im}\Pi_1(x_2^{1*}(y_1^1)y_1^2 \otimes y_1^3),$$

so $\langle x_1^3 \rangle = \langle y_1^3 \rangle$. Finally, we derive a contradiction by considering the mode-2 contraction of both representations of β .

$$\begin{aligned} \Pi_2(\beta)(x_2^{2^*}) &= x_1^1 \otimes x_1^3 = x_2^{2^*}(y_1^2)y_1^1 \otimes y_1^3 + x_2^{2^*}(y_2^2)y_1^2 \otimes y_2^3 \\ &= x_2^{2^*}(kx_1^2)y_1^1 \otimes y_1^3 + x_2^{2^*}(y_2^2)y_1^2 \otimes y_2^3 \\ &= x_2^{2^*}(y_2^2)y_1^2 \otimes y_2^3. \end{aligned}$$

However, this implies that $y_2^3 \in \langle x_1^3 \rangle$. This is a contradiction, since we have already shown that $y_1^3 \in \langle x_1^3 \rangle$ and the set $\{y_1^3, y_2^3\}$ is independent. We leave it to the reader to check the similar case of $x_2^{1*}(y_1^1) = 0$.

We have now shown that when V^1 , V^2 , and V^3 are finite dimensional real vector spaces and the sets $\{x_1^i, x_2^i\} \subseteq V^i$ are linearly independent, then the tensor

$$\beta = x_2^1 \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes x_2^2 \otimes x_1^3 + x_1^1 \otimes x_1^2 \otimes x_2^3$$

is rank three. However, it is also the limit of the sequence of rank at most two tensors

$$\tau_n = n \left(x_1^1 + \frac{1}{n} x_2^1 \right) \otimes \left(x_1^2 + \frac{1}{n} x_2^2 \right) \otimes \left(x_1^3 + \frac{1}{n} x_2^3 \right) - n x_1^1 \otimes x_1^2 \otimes x_1^3, \ n \in \mathbb{N}$$

in the norm topology. Hence, such tensors do not have optimal rank two approximations. We next characterize optimal rank two $2 \times 2 \times 2$ approximations geometrically, and show that, in fact, *all* rank three $2 \times 2 \times 2$ tensors do not have optimal rank two approximations with respect to the Frobenius norm.

3 A Characterization of Optimal Rank Two Approximations

The set of simple tensors is a variety, and it is called the Segre variety. It is the image of the map

$$V^1 \times V^2 \times V^3 \to V^1 \otimes V^2 \otimes V^3$$
$$v^1 \times v^2 \times v^3 \mapsto v^1 \otimes v^2 \otimes v^3.$$


The Segre variety of the $n_1 \times n_2 \times n_3$ tensor space $V^1 \otimes V^2 \otimes V^3$ is denoted $X^{n_1 \times n_2 \times n_3}$ or simply X when the dimensions are clear from context. The tangent space of the Segre variety at a point v is denoted as $T_v(X)$, and is characterized in [3] as in the following theorem.

Theorem 4. Let V^1 , V^2 , and V^3 be finite n_1 , n_2 , n_3 -dimensional real vector spaces, respectively, and let $x_1^1 \otimes x_1^2 \otimes x_1^3$ be a rank one tensor in the Segre variety X. The tangent space of X at $x_1^1 \otimes x_1^2 \otimes x_1^3$ is the space of all tensors in the form

$$x_{2}^{1} \otimes x_{1}^{2} \otimes x_{1}^{3} + x_{1}^{1} \otimes x_{2}^{2} \otimes x_{1}^{3} + x_{1}^{1} \otimes x_{1}^{2} \otimes x_{2}^{3}$$
(5)

for some $x_2^1 \in V^1$, $x_2^2 \in V^2$, and $x_2^3 \in V^3$.

Tensors in the form of (5) are precisely the tensors that we previously showed were limit points of the sequence (2) of tensors of rank at most two. This gives us geometric insight into the tensor rank-jumping phenomenon. Secant lines of the Segre variety contain rank at most two tensors, and tangent lines are limits of secant lines. Thus, rank three tensors on lines tangent to the Segre variety are limit points of rank at most two tensors.



We have seen that the set of rank at most two tensors is not closed with respect to the norm topology. This implies that it is also not closed with respect to the Zariski topology. This motivates the following definition of the 2^{nd} secant variety of the Segre variety.

Definition 5. The 2^{nd} secant variety of the Segre variety, denoted $\sigma_2(X)$, is the Zariski closure of all the secant lines of the Segre variety X.

If v is an optimal rank two approximation of τ with respect to an inner product norm, then $v - \tau$ must be orthogonal to the tangent space of $\sigma_2(X)$ at v, which we characterize in the following theorem using Theorem 4 and Terracini's Lemma [4].

Theorem 6. Let V^1 , V^2 , and V^3 be finite n_1 , n_2 , n_3 -dimensional real vector spaces, respectively, and let $v = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_2^3$ be a rank two tensor in $V^1 \otimes V^2 \otimes V^3$. The tangent space of $\sigma_2(X)$ at v is the space of all tensors in the form

$$x_{3}^{1} \otimes x_{1}^{2} \otimes x_{1}^{3} + x_{1}^{1} \otimes x_{3}^{2} \otimes x_{1}^{3} + x_{1}^{1} \otimes x_{1}^{2} \otimes x_{3}^{3} + x_{4}^{1} \otimes x_{2}^{2} \otimes x_{2}^{3} + x_{2}^{1} \otimes x_{4}^{2} \otimes x_{2}^{3} + x_{2}^{1} \otimes x_{2}^{2} \otimes x_{4}^{3}$$
(6)

for some $x_3^1, x_4^1 \in V^1$, $x_3^2, x_4^2 \in V^2$, and $x_3^3, x_4^3 \in V^3$.

Until now, we have worked with $n_1 \times n_2 \times n_3$ tensors. The next theorem, however, is our first statement that must be restricted to $2 \times 2 \times 2$ tensors.

Theorem 7. Let V^1 , V^2 , and V^3 be 2-dimensional real vector spaces, and let $\tau \in V^1 \otimes V^2 \otimes V^3$ be of rank greater than two. If $v \in V^1 \otimes V^2 \otimes V^3$ is an optimal rank two approximation of τ with respect to an inner product norm, then im $\Pi_i(v)$ is not dimension two for some i = 1, 2, or 3.

Proof. Suppose for contradiction that im $\Pi_i(v)$ were dimension two for i = 1, 2, and 3. Then, there would exist three linearly independent sets $\{x_1^i, x_2^i\}$ in V^i for i = 1, 2, and 3, such that

$$\upsilon = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_2^3$$

It follows that the set $\{x_i^1 \otimes x_j^2 \otimes x_k^3\}_{i,j,k=1}^{2,2,2}$ is a basis of $V^1 \otimes V^2 \otimes V^3$, so there must exist constants c_{ijk} such that $\tau = \sum_{i,j,k=1}^{2,2,2} c_{ijk} x_i^1 \otimes x_j^2 \otimes x_k^3$. By multilinearity,

$$\begin{aligned} \tau &= (c_{111}x_1^1 + c_{211}x_2^1) \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes c_{121}x_2^2 \otimes x_1^3 + x_1^1 \otimes x_1^2 \otimes c_{112}x_2^3 \\ &+ c_{122}x_1^1 \otimes x_2^2 \otimes x_2^3 + x_2^1 \otimes c_{212}x_1^2 \otimes x_2^3 + x_2^1 \otimes x_2^2 \otimes (c_{221}x_1^3 + c_{222}x_2^3) \end{aligned}$$

Hence, τ is in the form of (6), and, thus, it is in the tangent space of $\sigma_2(X)$ at v. This implies that $v - \tau$ is both in $T_v \sigma_2(X)$ and orthogonal to $T_v \sigma_2(X)$, which implies that $\tau = v$. However, this is a contradiction, since the rank of τ is not equal to the rank of v by hypothesis.

4 P-Norms and Optimal Rank Two Approximations

Once bases of vector spaces V^1 , V^2 , and V^3 are chosen, we can define explicit norms on the tensor space $V^1 \otimes V^2 \otimes V^3$. Choose basis $\{e_j^i\}_{j=1}^{n_i}$ of V^i for i = 1, 2, and 3, and denote the corresponding dual basis as $\{e_j^{i*}\}_{j=1}^{n_i}$ also for i = 1, 2, and 3. Let B denote the collection of these bases. For $\tau \in V^1 \otimes V^2 \otimes V^3$, we define the following class of norms for any positive integer p.

$$\|\tau\|_{B,p} = \left(\sum_{i,j,k=1,1,1}^{n_1,n_2,n_3} |a_{ijk}|^p\right)^{\frac{1}{p}},$$

where $\tau = \sum_{i,j,k} a_{ijk} e_i^1 \otimes e_j^2 \otimes e_k^3$. Similarly, for $\kappa \in V^r \otimes V^s$ and $v^r \in V^r$ for $1 \le r, s \le 3$, we define

$$\|\kappa\|_{B,p} = \left(\sum_{i,j=1,1}^{n_r,n_s} |a_{ij}|^p\right)^{\frac{1}{p}}, \text{ and } \|v^r\|_{B,p} = \left(\sum_{i=1}^{n_r} |a_i|^p\right)^{\frac{1}{p}},$$

where $\kappa = \sum_{i,j} a_{ij} e_i^r \otimes e_j^s$ and $v^r = \sum_i a_i e_i^r$. These norms are a convenient choice for working in tensor spaces as they work well with contraction maps.

Theorem 8. Let V^1 , V^2 , and V^3 be finite n_1 , n_2 , and n_3 -dimensional real vector spaces, and let B be the collection of bases of V^1 , V^2 , and V^3 defined above. For $\tau \in V^1 \otimes V^2 \otimes V^3$,

$$\|\tau\|_{B,p}^{p} = \sum_{t=1}^{n_{i}} \|\Pi_{i}(\tau)(e_{t}^{i*})\|_{B,p}^{p}$$

for any mode-*i* contraction Π_i for i = 1, 2, 3.

Proof. Without loss of generality, we prove the theorem for the the mode-1 contraction. Suppose $\tau = \sum_{s=1}^{r} v_s^1 \otimes v_s^2 \otimes v_s^3$ for some vectors v_s^i , and let $a_{s,t}^i$ be scalars such that $v_s^i = \sum_{t=1}^{n_i} a_{s,t}^i e_t^i$ for all i and s. It follows that

$$\sum_{s=1}^{r} v_s^1 \otimes v_s^2 \otimes v_s^3 = \sum_{i,j,k=1}^{n_1,n_2,n_3} \left(\sum_{s=1}^{r} a_{si}^1 a_{sj}^2 a_{sk}^3 \right) e_i^1 \otimes e_j^2 \otimes e_k^3.$$

Hence,

$$\begin{aligned} \|\tau\|_{B,p}^{p} &= \sum_{i,j,k=1}^{n_{1},n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \sum_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{si}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{n_{2},n_{3}} \left| \sum_{s=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{1} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{n_{1}} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{r} \left\| \prod_{j,k=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{3} \right\|^{p} \\ &= \sum_{i=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} \right\|^{p} \\ &= \sum_{i=1}^{r} \left\| \prod_{j,k=1}^{r} a_{sj}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^{2} a_{sk}^$$

Furthermore, when p = 2, the norm $\|\cdot\|_{B,2}$ is induced by the inner product

$$\langle \tau, \upsilon \rangle_{B,2} = \sum_{i,j,k} a_{ijk} b_{ijk},$$

where $\tau = \sum_{ijk} a_{ijk} e_i^1 \otimes e_j^2 \otimes e_k^3$ and $v = \sum_{ijk} b_{ijk} e_i^1 \otimes e_j^2 \otimes e_k^3$. The norm $\|\cdot\|_{B,2}$ is called the Frobenius norm with respect to bases B.

Theorem 9. The Frobenius inner product has the following property on rank one tensors.

$$\langle v_1^1 \otimes v_1^2 \otimes v_1^3 \mid v_2^1 \otimes v_2^2 \otimes v_2^3 \rangle_{B,2} = \langle v_1^1 \mid v_2^1 \rangle_{B,2} \langle v_1^2 \mid v_2^2 \rangle_{B,2} \langle v_1^3 \mid v_2^3 \rangle_{B,2}$$

for any rank one tensors $v_1^1 \otimes v_1^2 \otimes v_1^3$, $v_2^1 \otimes v_2^2 \otimes v_2^3 \in V^1 \otimes V^2 \otimes V^3$.

Proof. Let $a_{t,s}^i$ be scalars such that $v_t^i = \sum_{t=1}^{n_i} a_{t,s}^i e_s^i$ for i = 1, 2, 3 and t = 1, 2. It follows that

$$\begin{split} v_1^1 \otimes v_1^2 \otimes v_1^3 \ = \ \sum_{i,j,k=1}^{n_1,n_2,n_3} a_{1i}^1 a_{1j}^2 a_{1k}^3 \ e_i^1 \otimes e_j^2 \otimes e_k^3, \text{ and} \\ v_2^1 \otimes v_2^2 \otimes v_2^3 \ = \ \sum_{i,j,k=1}^{n_1,n_2,n_3} a_{2i}^1 a_{2j}^2 a_{2k}^3 \ e_i^1 \otimes e_j^2 \otimes e_k^3. \end{split}$$

Hence,

$$\langle v_1^1 \otimes v_1^2 \otimes v_1^3 \mid v_2^1 \otimes v_2^2 \otimes v_2^3 \rangle_{B,2} = \sum_{i,j,k=1}^{n_1,n_2,n_3} a_{1i}^1 a_{1j}^2 a_{1k}^3 a_{2i}^1 a_{2j}^2 a_{2k}^3$$

$$= \left(\sum_{i=1}^{n_1} a_{1i}^1 a_{2i}^1 \right) \left(\sum_{j=1}^{n_2} a_{1j}^2 a_{2j}^2 \right) \left(\sum_{k=1}^{n_3} a_{1k}^3 a_{2k}^3 \right)$$

$$= \langle v_1^1 \mid v_2^1 \rangle_{B,2} \langle v_1^2 \mid v_2^2 \rangle_{B,2} \langle v_1^3 \mid v_2^3 \rangle_{B,2}.$$

An optimal rank r approximation could be rank strictly less than r by our definition. We now show that an optimal rank r approximations with respect to p-norms must be rank r. This theorem is modified from [1, Lemma 8.2].

Theorem 10. Let V^i be finite n_i -dimensional real vector spaces, and let $\tau \in V^1 \otimes V^2 \otimes V^3$ have rank greater than r. If v is an optimal rank r approximation of τ with respect to $\|\cdot\|_{B,p}$, then v must be rank r.

Proof. Suppose for contradiction that there existed an optimal rank r approximation v with rank strictly less than r. Let a_{ijk} and b_{ijk} be scalars such that $\tau = \sum_{ijk} a_{ijk} e_i^1 \otimes e_j^2 \otimes e_k^3$ and $v = \sum_{ijk} b_{ijk} e_i^1 \otimes e_j^2 \otimes e_k^3$. Since τ and v have different ranks, there must exists some triple (α, β, γ) such that $a_{\alpha\beta\gamma} \neq b_{\alpha\beta\gamma}$. The tensor $v' = v + (a_{\alpha\beta\gamma} - b_{\alpha\beta\gamma})e_{\alpha}^1 \otimes e_{\beta}^2 \otimes e_{\gamma}^3$ is rank less than or equal to r by construction. It follows that

$$\begin{aligned} |\tau - \upsilon'||_{B,p} &= \|\tau - \left(\upsilon + (a_{\alpha\beta\gamma} - b_{\alpha\beta\gamma})e_{\alpha}^{1} \otimes e_{\beta}^{2} \otimes e_{\gamma}^{3}\right)\|_{B,p} \\ &= \left(\sum_{ijk \neq \alpha\beta\gamma} |a_{ijk} - b_{ijk}|^{p}\right)^{\frac{1}{p}} \\ &< \left(\sum_{ijk} |a_{ijk} - b_{ijk}|^{p}\right)^{\frac{1}{p}} = \|\tau - \upsilon\|_{B,p}, \end{aligned}$$

which contradicts that v is an optimal rank r approximation of τ .

If v is an optimal rank two approximation of τ with respect to the Frobenius norm, then the contractions of v must be related to the contractions of τ in the following way.

Theorem 11. Let V^1 , V^2 , and V^3 be n_1 , n_2 , and n_3 -dimensional real vector spaces, respectively. Let B denote the collection of bases $\{e_j^i\}_{j=1}^{n_i}$ of V^i for i = 1, 2, and 3. Furthermore, denote the corresponding dual bases as $\{e_j^{i*}\}_{j=1}^{n_i}$ for i = 1, 2, and 3. Let $\tau \in V^1 \otimes V^2 \otimes V^3$ be of rank greater than or equal to two, and let $v \in V^1 \otimes V^2 \otimes V^3$ be an optimal rank two approximation of τ with respect to the Frobenius norm $\|\cdot\|_{B,2}$. Let $P_{im \prod_i(v)}$ denote the projection onto the image of the mode-*i* contraction of v. It follows that

$$P_{im \prod_i(v)}(\prod_i(\tau)(e_j^{i*})) = \prod_i(v)(e_j^{i*})$$
 for any i, j .

Proof. As v is rank two, there must exist vectors x_i^i such that

$$v = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_2^3$$

Without loss of generality, we prove the theorem for i and j both equal to 1. First, note that the image of $\Pi_1(v)$ contains vectors in the form $ax_1^2 \otimes x_1^3 + bx_2^2 \otimes x_2^3$ for some constants a and b. If the set $\{x_1^1, x_2^1\}$ is independent, it

can be extended to a basis $\{x_j^1\}_{j=1}^{n_1}$ with dual basis $\{x_j^{1*}\}_{j=1}^{n_1}$. It then follows that the image of $\Pi_1(v)$ is the span of $\Pi_1(v)(x_1^{1*}) = x_1^2 \otimes x_1^3$ and $\Pi_1(v)(x_2^{1*}) = x_2^2 \otimes x_2^3$, so every element in the image of $\Pi_1(v)$ can indeed be written as $ax_1^2 \otimes x_1^3 + bx_2^2 \otimes x_2^3$ for some constants a and b. On the other hand, if $x_2^1 = kx_1^1$ for some constant k, then the set $\{x_1^1\}$ can similarly be extended to a basis $\{x_1^1, y_j^1\}_{j=2}^{n_1}$ of V^1 with the corresponding dual basis $\{x_1^{1*}, y_j^{1*}\}_{j=2}^{n_2}$. It follows that the image of $\Pi_1(v)$ is the span of $\Pi_1(v)(x_1^{1*}) = x_1^2 \otimes x_1^3 + kx_2^2 \otimes x_2^3$. Hence, in this case, it is also true that every element in the image of $\Pi_1(v)$ can be written as $ax_1^2 \otimes x_1^3 + bx_2^2 \otimes x_2^3$ for some constants a and b.

Suppose for contradiction that $P_{\text{im}\,\Pi_1(v)}(\Pi_1(\tau)(e_1^{1*})) \neq \Pi_1(v)(e_1^{1*})$. Let α and β be scalars such that

$$P_{\operatorname{im}\Pi_{1}(\upsilon)}(\Pi_{1}(\tau)(e_{1}^{1*})) = \alpha x_{1}^{2} \otimes x_{1}^{3} + \beta x_{2}^{2} \otimes x_{2}^{3}.$$

Furthermore, let a_i and b_i be scalars such that

$$\Pi_1(v)(e_i^{1*}) = a_i x_1^2 \otimes x_1^3 + b_i x_2^2 \otimes x_2^3$$

for $i = 2, ..., n_1$. Define $v' \in V^1 \otimes V^2 \otimes V^3$ as the unique tensor with the following mode-1 contraction:

$$\Pi_{1}(v')(e_{1}^{1*}) = \alpha x_{1}^{2} \otimes x_{1}^{3} + \beta x_{2}^{2} \otimes x_{2}^{3} = P_{\operatorname{im}\Pi_{1}(v)}(\Pi_{1}(\tau)(e_{1}^{1*})),$$

$$\Pi_{1}(v')(e_{i}^{1*}) = a_{i} x_{1}^{2} \otimes x_{1}^{3} + b_{i} x_{2}^{2} \otimes x_{2}^{3} = \Pi_{1}(v)(e_{i}^{1*}) \text{ for } i = 2, \dots, n_{1}.$$

It follows that

$$\nu' = (\alpha e_1^1 + \sum_{j=2}^{n_1} a_j e_j^1) \otimes x_1^2 \otimes x_1^3 + (\beta e_1^1 + \sum_{j=2}^{n_1} b_j e_j^1) \otimes x_2^2 + \otimes x_2^3$$

so v' is rank ≤ 2 . Furthermore,

$$\|\tau - v'\|_{B,2}^2 = \sum_{i=1}^{n_1} \|\Pi_1(\tau)(e_i^{1*}) - \Pi_1(v')(e_i^{1*})\|_{B,2}^2$$
(7)

$$= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(\upsilon')(e_1^{1*})\|_{B,2}^2 + \sum_{i=2}^{n_1} \|\Pi_1(\tau)(e_i^{1*}) - \Pi_1(\upsilon)(e_i^{1*})\|_{B,2}^2$$
(8)

$$< \|\Pi_{1}(\tau)(e_{1}^{1*}) - \Pi_{1}(\upsilon)(e_{1}^{1*})\|_{B,2}^{2} + \sum_{i=2}^{n_{1}} \|\Pi_{1}(\tau)(e_{i}^{1*}) - \Pi_{1}(\upsilon)(e_{i}^{1*})\|_{B,2}^{2}$$

$$(9)$$

$$= \|\tau - v\|_{B,2}^2.$$

Equation (7) follows from Theorem 8, and equation (8) follows from the fact that $\Pi_1(v')(e_i^{1*}) = \Pi_1(v)(e_i^{1*})$ for $i = 2, 3, \ldots, n_1$. Equation (9) follows from our hypothesis that $\alpha x_1^2 \otimes x_1^3 + \beta x_2^2 \otimes x_2^3$ is a better approximation of $\Pi_1(\tau)(e_1^{1*})$ with respect to $\|\cdot\|_{B,2}$ than $\Pi_1(v)(e_1^{1*})$. Hence, $\|\tau - v'\|_{B,2} < \|\tau - v\|_{B,2}$, contradicting that v is a best rank two approximation of τ .

5 Optimal Rank Two Approximations of Rank Three 2 × 2 × 2 Tensors Do Not Exist With Respect to the Frobenius Norm

In this section, we show that rank three $2 \times 2 \times 2$ tensors over \mathbb{R} do not have optimal rank two approximations with respect to the Frobenius norm. Given any rank two tensor v and rank three tensor τ , we construct a rank two tensor v' that is a better approximation of τ than v with respect to the Frobenius norm. Let V^i be two-dimensional real vector spaces for i = 1, 2, 3. As in the previous section, let B denote a collection of bases $\{e_j^i\}_{j=1}^2$ of V^i for i = 1, 2, 3. Furthermore, denote the corresponding dual bases as $\{e_i^{i*}\}_{i=1}^2$ for i = 1, 2, 3.

Theorem 12. If $\tau \in V^1 \otimes V^2 \otimes V^3$ is rank three, then there does not exist an optimal rank two approximation of τ with respect to $\|\cdot\|_{B,2}$.

Proof. Suppose for contradiction that there existed a tensor v that was an optimal rank two approximation of τ . By Theorem 7, we may assume

$$v = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_1^3$$




Figure 2: In this figure, we consider the $2 \times 2 \times 2$ case. Let $v = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_1^3$ be an optimal rank two approximation of rank three tensor $\tau \in V^1 \otimes V^2 \otimes V^3$. In this case, im $\Pi_1(v)$ is a plane in $V^2 \otimes V^3$ that contains all the rank one tensors in the span of $x_1^2 \otimes x_1^3$ and $x_2^2 \otimes x_1^3$. This plane is denoted as M in the figure. Theorem 11 shows that the projection of $\Pi_1(\tau)(e_i^{1*})$ onto the plane M must be $\Pi_1(v)(e_i^{1*})$ for i = 1, 2.

for some vectors x_i^i . Since $\{e_1^1, e_2^1\}$ is a basis of V^1 , there exists scalars a, b, c, and d such that

$$x_1^1 = ae_1^1 + ce_2^1$$
 and $x_2^1 = be_1^1 + de_2^1$

It follows that

$$v = e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 + e_2^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3$$

If the set $\{x_1^1, x_2^1\}$ were linearly dependent, then v would be rank one by multilinearity, which would contradict Theorem 10. Hence, $\{x_1^1, x_2^1\}$ is independent, and is thus a basis of V^1 . Let $\{x_1^{1*}, x_2^{1*}\}$ denote its dual basis. The image of $\Pi_1(v)$ is the span of $\Pi_1(v)(x_1^{1*}) = x_1^2 \otimes x_1^3$ and $\Pi_1(v)(x_2^{1*}) = x_2^2 \otimes x_1^3$. These two tensors are linearly independent in $V^2 \otimes V^3$ since the set $\{x_1^2, x_2^2\}$ is independent, which also follows from the fact that v is rank two. Hence, the image of $\Pi_1(v)$ is the plane spanned by the tensors $x_1^2 \otimes x_1^3$ and $x_2^2 \otimes x_1^3$. Let M denote this plane and let P_M denote the projection onto this plane. By Theorem 11,

$$P_M(\Pi_1(\tau)(e_1^{1*})) = \Pi_1(\upsilon)(e_1^{1*}) = (ax_1^2 + bx_2^2) \otimes x_1^3, \text{ and} P_M(\Pi_1(\tau)(e_2^{1*})) = \Pi_1(\upsilon)(e_2^{1*}) = (cx_1^2 + dx_2^2) \otimes x_1^3.$$

Let $x_2^3 \in V^3$ be a vector such that the set $\{x_1^3, x_2^3\}$ is orthogonal with respect to $\langle \cdot | \cdot \rangle_{B,2}$. The set $\{x_1^2 \otimes x_2^3, x_2^2 \otimes x_2^3\}$ spans the orthogonal complement of the plane M in $V^2 \otimes V^3$ since

$$\langle x_i^2 \otimes x_2^3 \mid x_j^2 \otimes x_1^3 \rangle_{B,2} = \langle x_i^2 \mid x_j^2 \rangle_{B,2} \ \langle x_2^3 \mid x_1^3 \rangle_{B,2} = 0 \text{ for } i, j \in \{1,2\},$$

by property (9). Thus, there must exist some constants r, s, p, and q such that

$$\Pi_1(\tau)(e_1^{1*}) = (ax_1^2 + bx_2^2) \otimes x_1^3 + (rx_1^2 + sx_2^2) \otimes x_2^3, \text{ and} \\ \Pi_1(\tau)(e_2^{1*}) = (cx_1^2 + dx_2^2) \otimes x_1^3 + (px_1^2 + qx_2^2) \otimes x_2^3.$$

It follows that τ can be written in the form

$$\begin{array}{rcl} \tau \ = \ e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 \ + \ e_1^1 \otimes (rx_1^2 + sx_2^2) \otimes x_2^3 \\ & + \ e_2^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3 \ + \ e_2^1 \otimes (px_1^2 + qx_2^2) \otimes x_2^3. \end{array}$$

If we can show that

$$\langle (cx_1^2 + dx_2^2) | (rx_1^2 + sx_2^2) \rangle_{B,2} = 0,$$
 (10)

$$\langle (ax_1^2 + bx_2^2) | (px_1^2 + qx_2^2) \rangle_{B,2} = 0$$
, and (11)

$$\langle (ax_1^2 + bx_2^2) | (rx_1^2 + sx_2^2) \rangle_{B,2} = 0,$$
(12)

then this would imply that

$$(px_1^2 + qx_2^2) = k_1(rx_1^2 + sx_2^2)$$
 and $(cx_1^2 + dx_2^2) = k_2(ax_1^2 + bx_2^2)$

for some constants k_1 , k_2 , since these vectors are in a two-dimensional space. It then follows by the multilinearity of the tensor product that

$$\begin{aligned} \tau \ &= \ e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 \ + \ e_1^1 \otimes (rx_1^2 + sx_2^2) \otimes x_2^3 \\ &+ \ e_2^1 \otimes k_2 (ax_1^2 + bx_2^2) \otimes x_1^3 \ + \ e_2^1 \otimes k_1 (rx_1^2 + sx_2^2) \otimes x_2^3 \\ &= \ (e_1^1 + k_2 e_2^1) \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 \ + \ (e_1^1 + k_1 e_2^1) \otimes (rx_1^2 + sx_2^2) \otimes x_2^3, \end{aligned}$$

contradicting that τ is rank three.

We first prove equality (10) by considering the tensor

$$v_1(\epsilon) = e_2^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3 + e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 + e_1^1 \otimes (cx_1^2 + dx_2^2) \otimes \epsilon x_2^3$$

Suppose for contradiction that (10) were not true. It would then follow that

$$\langle (rx_1^2 + sx_2^2) \otimes x_2^3 | (cx_1^2 + dx_2^2) \otimes x_2^3 \rangle_{B,2} = \langle (rx_1^2 + sx_2^2) | (cx_1^2 + dx_2^2) \rangle_{B,2} ||x_2^3||_{B,2}^2 \neq 0.$$

We can always choose a real ϵ small enough in absolute value such that

$$B_1(\epsilon) = -2\epsilon \left\langle \left(rx_1^2 + sx_2^2 \right) \otimes x_2^3 \right| \left(cx_1^2 + dx_2^2 \right) \otimes x_2^3 \right\rangle_{B,2} + \epsilon^2 \left\| \left(cx_1^2 + dx_2^2 \right) \otimes x_2^3 \right\|_{B,2}^2$$

is negative. For example, if $\langle (rx_1^2 + sx_2^2) \otimes x_2^3 | (cx_1^2 + dx_2^2) \otimes x_2^3 \rangle_{B,2} < 0$, an $\epsilon < 0$ small enough in absolute value would result in a $B_1(\epsilon)$ negative. Suppose such an ϵ is chosen. Observe that

$$\Pi_1(\upsilon_1(\epsilon))(e_1^{1*}) = (ax_1^2 + bx_2^2) \otimes x_1^3 + \epsilon(cx_1^2 + dx_2^2) \otimes x_2^3, \text{ and} \\ \Pi_1(\upsilon_1(\epsilon))(e_2^{1*}) = (cx_1^2 + dx_2^2) \otimes x_1^3 = \Pi_1(\upsilon)(e_2^{1*}).$$

It thus follows that

$$\|\tau - v_1(\epsilon)\|_{B,2}^2 = \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(v_1(\epsilon))(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(v_1(\epsilon))(e_2^{1*})\|_{B,2}^2$$
(13)

$$= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(v_1(\epsilon))(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(v)(e_2^{1*})\|_{B,2}^2.$$
(14)

Equation (13) follows from Theorem 8, and equation (14) follows from the fact that $\Pi_1(\upsilon)(e_2^{1*}) = \Pi_1(\upsilon_1(\epsilon))(e_2^{1*})$. By the multilinearity of the tensor product and the multilinearity of the inner product, we conclude that

$$\begin{split} \|\tau - v_{1}(\epsilon)\|_{B,2}^{2} &= \langle \Pi_{1}(\tau)(e_{1}^{1*}) - \Pi_{1}(v_{1}(\epsilon))(e_{1}^{1*}) | \Pi_{1}(\tau)(e_{1}^{1*}) - \Pi_{1}(v_{1}(\epsilon))(e_{1}^{1*}) \rangle_{B,2} \\ &+ \|\Pi_{1}(\tau)(e_{2}^{1*}) - \Pi_{1}(v)(e_{2}^{1*})\|_{B,2}^{2} \\ &= \langle (rx_{1}^{2} + sx_{2}^{2}) \otimes x_{2}^{3} - \epsilon(cx_{1}^{2} + dx_{2}^{2}) \otimes x_{2}^{3} | (rx_{1}^{2} + sx_{2}^{2}) \otimes x_{2}^{3} - \epsilon(cx_{1}^{2} + dx_{2}^{2}) \otimes x_{2}^{3} \rangle_{B,2} \\ &+ \|\Pi_{1}(\tau)(e_{2}^{1*}) - \Pi_{1}(v)(e_{2}^{1*})\|_{B,2}^{2} \\ &= \|(rx_{1}^{2} + sx_{2}^{2}) \otimes x_{2}^{3}\|_{B,2}^{2} - 2\epsilon \langle (rx_{1}^{2} + sx_{2}^{2}) \otimes x_{2}^{3} | (cx_{1}^{2} + dx_{2}^{2}) \otimes x_{2}^{3} \rangle_{B,2} \\ &+ \epsilon^{2} \|(cx_{1}^{2} + dx_{2}^{2}) \otimes x_{2}^{3}\|_{B,2}^{2} + \|\Pi_{1}(\tau)(e_{2}^{1*}) - \Pi_{1}(v)(e_{2}^{1*})\|_{B,2}^{2} \\ &= \|(rx_{1}^{2} + sx_{2}^{2}) \otimes x_{2}^{3}\|_{B,2}^{2} + B_{1}(\epsilon) + \|\Pi_{1}(\tau)(e_{2}^{1*}) - \Pi_{1}(v)(e_{2}^{1*})\|_{B,2}^{2}. \end{split}$$

Since ϵ was chosen specifically to make $B_1(\epsilon)$ negative, we can conclude that

$$\begin{aligned} \|\tau - \upsilon_1(\epsilon)\|_{B,2}^2 &= \|(rx_1^2 + sx_2^2) \otimes x_2^3\|_{B,2}^2 + B_1(\epsilon) + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &< \|(rx_1^2 + sx_2^2) \otimes x_2^3\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(\upsilon)(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &= \|\tau - \upsilon\|_{B,2}^2. \end{aligned}$$

Hence, $v_1(\epsilon)$ is a better approximation of τ than v. The tensor $v_1(\epsilon)$ is also in the tangent space of the Segre variety at $e_1^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3$. Hence, there exists a sequence of rank two tensors that converges to $v_1(\epsilon)$. Thus, there must be some rank two tensor in the sequence that is better approximation to τ than v, contradicting that v is an optimal rank two approximation.

Equality (11) can be proven in the same way by considering the tensor

$$v_2(\epsilon) = e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3 + e_2^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3 + e_2^1 \otimes (ax_1^2 + bx_2^2) \otimes \epsilon x_2^3$$

which is in the tangent space of the Segre variety at $e_2^1 \otimes (ax_1^2 + bx_2^2) \otimes x_1^3$.

It thus remains to show equality (12), which we prove by considering the rank two tensor

$$\upsilon_3(\epsilon) = e_1^1 \otimes (ax_1^2 + bx_2^2) \otimes (x_1^3 + \epsilon(x_1^3 + x_2^3)) + e_2^1 \otimes (cx_1^2 + dx_2^2) \otimes x_1^3.$$

It follows that the mode-1 contractions of $v_3(\epsilon)$ are as follows.

$$\Pi_1(\upsilon_3(\epsilon))(e_1^{1*}) = (ax_1^2 + bx_2^2) \otimes x_1^3 + \epsilon(ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3), \text{ and} \\ \Pi_1(\upsilon_3(\epsilon))(e_2^{1*}) = (cx_1^2 + dx_2^2) \otimes x_1^3 = \Pi_1(\upsilon)(e_2^{1*}).$$

Suppose for contradiction that (12) were nonzero. It would then follow that

$$\langle (rx_1^2 + sx_2^2) \otimes x_2^3 | (ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3) \rangle_{B,2} = \langle (rx_1^2 + sx_2^2) | (ax_1^2 + bx_2^2) \rangle_{B,2} \|x_2^3\|_{B,2}^2 \neq 0.$$

We could then choose an ϵ small enough in absolute value such that $D(\epsilon) < 0$, where

$$D(\epsilon) = -2\epsilon \left\langle (rx_1^2 + sx_2^2) \otimes x_2^3 \mid (ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3) \right\rangle_{B,2} + \epsilon^2 \left\| (ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3) \right\|_{B,2}^2.$$

For example, if $\langle (rx_1^2 + sx_2^2) \otimes x_2^3 | (ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3) \rangle_{B,2} < 0$, an $\epsilon < 0$ small enough in absolute value will yield a negative $D(\epsilon)$. Suppose such an ϵ is chosen. Then,

$$\begin{aligned} |\tau - \upsilon_3(\epsilon)||_{B,2}^2 &= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(\upsilon_3(\epsilon))(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon_3(\epsilon))(e_2^{1*})\|_{B,2}^2 \\ &= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(\upsilon_3(\epsilon))(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2. \end{aligned}$$

Observe that

$$\|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(v_3(\epsilon))(e_1^{1*})\|_{B,2}^2 = \|(rx_1^2 + sx_2^2) \otimes x_2^3 - \epsilon(ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3)\|_{B,2}^2.$$

Writing this as an inner product, we see that

$$\begin{aligned} \|\tau - \upsilon_3(\epsilon)\|_{B,2}^2 &= \|(rx_1^2 + sx_2^2) \otimes x_2^3\|_{B,2}^2 - 2\epsilon \left\langle (rx_1^2 + sx_2^2) \otimes x_2^3 \mid (ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3) \right\rangle_{B,2} \\ &+ \epsilon^2 \|(ax_1^2 + bx_2^2) \otimes (x_1^3 + x_2^3)\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &= \|(rx_1^2 + sx_2^2) \otimes x_2^3\|_{B,2}^2 + D(\epsilon) + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2. \end{aligned}$$

Finally, since ϵ was chosen so that $D(\epsilon)$ would be negative, it follows that

$$\begin{aligned} \|\tau - \upsilon_3(\epsilon)\|_{B,2}^2 &< \|(rx_1^2 + sx_2^2) \otimes x_2^3\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &= \|\Pi_1(\tau)(e_1^{1*}) - \Pi_1(\upsilon)(e_1^{1*})\|_{B,2}^2 + \|\Pi_1(\tau)(e_2^{1*}) - \Pi_1(\upsilon)(e_2^{1*})\|_{B,2}^2 \\ &= \|\tau - \upsilon\|_{B,2}^2. \end{aligned}$$

Hence, the rank two tensor $v_3(\epsilon)$ is a better approximation of τ than v, contradicting that v is an optimal rank two approximation.

We have thus shown that rank three $2 \times 2 \times 2$ real tensors have no optimal rank two approximations with respect to the Frobenius norm. This implies that the nearest point of a rank three $2 \times 2 \times 2$ tensor τ to the second secant variety $\sigma_2(X)$ of the Segre variety X with respect to the Frobenius norm must be rank three. In fact, our proof above demonstrates that the nearest point of τ to $\sigma_2(X)$ is in fact on the tangential variety of the Segre variety, which is the set of all tensors contained in the tangent space of the Segre variety at some point $u \in X$. This is a variety as is it the daul projective variety of X, and is denoted TX. That is,

$$TX = \{ \tau \in V^1 \otimes V^2 \otimes V^3 \mid \tau \in T_u(X) \text{ for some } u \in X \}$$

From Theorem (4), we know that

$$TX = \{ \tau \in V^1 \otimes V^2 \otimes V^3 \mid \tau = x_2^1 \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes x_2^2 \otimes x_1^3 + x_1^1 \otimes x_1^2 \otimes x_2^3 \\ \text{for some } x_j^i \in V^i \}.$$

There is an open, dense subset of TX of rank three tensors. However, there are rank two elements in TX, and the fact that these rank two tensors can never be the the nearest point on TX of any rank three tensor implies there is an interesting curvature of TX at these points. We now consider an example of such a rank two tensor in TX. Let $\{x_1^i, x_2^i\}$ be independent vectors in V^i for i = 1, 2, 3. The tensor

$$\nu = x_1^1 \otimes x_1^2 \otimes x_1^3 + x_2^1 \otimes x_2^2 \otimes x_1^3$$

is a rank two tensor in TX. For every $\epsilon \neq 0$, the tensor

$$\tau(\epsilon) \ = \ x_1^1 \otimes x_1^2 \otimes x_1^3 \ + \ x_2^1 \otimes x_2^2 \otimes x_1^3 \ + \ \epsilon(x_1^1 + x_2^1) \otimes (x_1^2 + x_2^2) \otimes x_2^3$$

is rank three. Clearly, $\lim_{\epsilon \to 0} \tau(\epsilon) = \nu$. However, the nearest point to $\tau(\epsilon)$ on TX with respect to the Frobenius norm is never ν , even when ϵ is infinitesimally small, since the nearest point to $\tau(\epsilon)$ on TX must be rank three. The tangential variety of the $2 \times 2 \times 2$ Segre variety must thus have significant curvature at its rank two points, which is already suggested by the fact that ν is tangent to the Segre variety at any tensors in the form $x_2^1 \otimes x_1^2 \otimes y_1^3$ and $x_1^1 \otimes x_2^2 \otimes y_1^3$ for some $y_1^3 \in V^3$. In contrast, the rank three tensors in TX are tangent to distinct point on the Segre variety up to a multiplicative constant.

Theorem 13. Let V^1 , V^2 , and V^3 be two-dimensional real vector spaces. Let X denote the Segre variety of simple tensors in $V^1 \otimes V^2 \otimes V^3$, and let TX denote the tangential variety of X. If $\tau \in TX$ is rank three, then it is tangent to a unique point of X up to a multiplicative constant.

Proof. Suppose τ is tangent to both the tensors $x_1^1 \otimes x_1^2 \otimes x_1^3$ and $x_2^1 \otimes x_2^2 \otimes x_2^3$. It then follows from Theorem 4 that

$$\tau = y_1^1 \otimes x_1^2 \otimes x_1^3 + x_1^1 \otimes y_1^2 \otimes x_1^3 + x_1^1 \otimes x_1^2 \otimes y_1^3, \text{ and}$$
(15)

$$= y_2^1 \otimes x_2^2 \otimes x_2^3 + x_2^1 \otimes y_2^2 \otimes x_2^3 + x_2^1 \otimes x_2^2 \otimes y_2^3,$$
 (16)

for some vectors y_j^i . Since τ is rank three, the sets $\{y_j^i, x_j^i\}$ must be independent for all i, j. Let $\{y_j^{i*}, x_j^{i*}\}$ be the corresponding dual bases. Considering the contraction maps of τ with respect to both of these representation (15) and (16), we conclude that

$$\Pi_1(\tau)(y_1^{1*}) = x_1^2 \otimes x_1^3 = x_2^2 \otimes \left(y_1^{1*}(y_2^1)x_2^3 + y_1^{1*}(x_2^1)y_2^3\right) + y_1^{1*}(x_2^1)y_2^2 \otimes x_2^3.$$

By Lemma 2, it follows that $y_1^{1*}(x_2^1) = 0$, so $x_1^2 \otimes x_1^3 = y_1^{1*}(y_2^1) x_2^2 \otimes x_2^3$. The fact that $y_1^{1*}(x_2^1) = 0$ implies that $x_2^1 = k_1 x_1^1$ for some constant k_1 . Furthermore, im $\Pi_2(x_1^2 \otimes x_1^3) = \text{im } \Pi_2(y_1^{1*}(y_2^1)x_2^2 \otimes x_2^3)$ implies that $x_1^2 = k_2 x_2^2$ for some constant k_2 , and im $\Pi_1(x_1^2 \otimes x_1^3) = \text{im } \Pi_1(y_1^{1*}(y_2^1)x_2^2 \otimes x_2^3)$ implies that $x_2^3 = k_3 x_1^3$ for some constant k_3 . Hence, $x_2^1 \otimes x_2^2 \otimes x_2^3 = k_1 k_2 k_3 x_1^1 \otimes x_1^2 \otimes x_1^3$.

Acknowledgments

The author would like to thank Saugata Basu and Ryan Vitale for their valuable discussions about this paper.

References

- [1] Vin De Silva and Lek-Heng Lim. Tensor rank and the ill-posedness of the best low-rank approximation problem. *SIAM Journal on Matrix Analysis and Applications*, 30(3):1084–1127, 2008.
- [2] Alwin Stegeman. Nonexistence of best low-rank approximations for real-valued three-way arrays and what to do about it. In 21st International Symposium on Mathematical Theory of Networks and Systems (MTNS2014), pages 954–957, 2014.
- [3] Giorgio Ottaviani. Introduction to the hyperdeterminant and to the rank of multidimensional matrices. In *Commutative algebra*, pages 609–638. Springer, 2013.
- [4] Fedor Lazarevič Zak. *Tangents and secants of algebraic varieties*, volume 127. American Mathematical Soc., 1993.