ANALOGIES OF JACOBI'S FORMULA

KEIJI MATSUMOTO

ABSTRACT. By considering Schwarz's map for the hypergeometric differential equation with parameters (a, b, c) = (1/6, 1/2, 1) or (1/12, 5/12, 1), we give some analogies of Jacobi's formula $\vartheta_{00}(\tau)^2 = F(1/2, 1/2, 1; \lambda(\tau))$, where $\vartheta_{00}(\tau)$ and $\lambda(\tau)$ are the theta constant and the lambda function defined on the upper-half plane, and F(a, b, c; z) is the hypergeometric series defined on the unit disk. As applications of our formulas, we give several functional equations for F(a, b, c; z).

1. INTRODUCTION

Schwarz's map is defined by the analytic continuation of the ratio of linearly independent local solutions to the hypergeometric differential equation E(a, b, c) given in (2.3). It is a multi-valued map from $Z = \mathbb{C} - \{0, 1\}$ to the complex projective line \mathbb{P}^1 in general. Schwarz's map φ_0 for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$ is given by the map

$$z \mapsto \varphi_0(z) = \mathbf{i} \cdot \frac{F(\frac{1}{2}, \frac{1}{2}, 1; 1-z)}{F(\frac{1}{2}, \frac{1}{2}, 1; z)},\tag{1.1}$$

around $z = \frac{1}{2}$, where $\mathbf{i} = \sqrt{-1}$, and F(a, b, c; z) is the hypergeometric series given in (2.1). It is known that the image $\varphi_0(Z)$ is the upper-half plane \mathbb{H} , and that the projective monodromy of φ_0 is $\Pr(2) = \Gamma(2)/\{\pm I_2\}$, where I_2 is the unit matrix of size 2, and $\Gamma(2)$ is the principally congruence subgroup of $\mathrm{SL}_2(\mathbb{Z})$ of level 2. The inverse of φ_0 is expressed by the lambda function

$$\tau \mapsto \lambda(\tau) = \frac{\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4},\tag{1.2}$$

where $\vartheta_{pq}(\tau)$ is the theta constant with characteristics p, q given in (3.1). Jacobi's formula is

$$\vartheta_{00}(\tau)^2 = F(\frac{1}{2}, \frac{1}{2}, 1; \lambda(\tau))$$

where τ is a complex number with large imaginary part. Here note that this formula is an equality between the denominators in (1.1) and (1.2) under the relation $z = \lambda(\tau)$ in (1.2). Some analogies of Jacobi's formula are given in [BB2, Theorem 2.3, Theorem 2.6], which relate $F(\frac{1}{3}, \frac{2}{3}, 1; z)$ and $F(\frac{1}{4}, \frac{3}{4}, 1; z)$ to modular forms with respect to their projective monodromy group isomorphic to the triangle groups $(3, \infty, \infty)$ and $(2, \infty, \infty)$, respectively.

In this paper, we give some analogies of Jacobi's formula by considering Schwarz's maps φ_1 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$ and φ_2 for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$. There exists a basis of the space of local solutions to $E(\frac{1}{6}, \frac{1}{2}, 1)$ (or $E(\frac{1}{12}, \frac{5}{12}, 1)$) around $z = \frac{1}{2}$ such that the image of φ_1 (or φ_2) is an open dense subset in \mathbb{H} and that the projective monodromy of φ_1 (or φ_2)

Date: today.

²⁰¹⁰ Mathematics Subject Classification. Primary 14K25; Secondary 33C05, 11F03.

Key words and phrases. Jacobi's formula, Theta function, Hypergeometric function.

is a subgroup of $\text{PSL}_2(\mathbb{Z}) = \text{SL}_2(\mathbb{Z})/\{\pm I_2\}$ isomorphic to the triangle group $(3, 3, \infty)$ (or $(2, 3, \infty)$). We give each basis as linear combinations of F(a, b, a + b - c + 1; 1 - z) and F(a, b, c; z) for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1), (\frac{1}{12}, \frac{5}{12}, 1)$, and the circuit matrices along ρ_0 and ρ_1 with respect to them as

$$\begin{array}{ll} T^2, \ \omega^2 W & \text{for} \quad (a,b,c) = (\frac{1}{6}, \frac{1}{2}, 1), \\ T, \quad \mathbf{i}J & \text{for} \quad (a,b,c) = (\frac{1}{12}, \frac{5}{12}, 1), \end{array}$$

where ρ_k (k = 0, 1) is a positively oriented circle with terminal $z = \frac{1}{2}$, radius $\frac{1}{2}$ and center z = k, and

$$\omega = \frac{-1 + \sqrt{3}\mathbf{i}}{2}, \quad T = \begin{pmatrix} 1 & 1\\ 0 & 1 \end{pmatrix}, \quad W = \begin{pmatrix} -1 & -1\\ 1 & 0 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 1\\ -1 & 0 \end{pmatrix}.$$

The group generated by T and J is $SL_2(\mathbb{Z})$, and that by T^2 and W is

$$\Gamma(2)^{1/3} = \{g \in \mathrm{SL}_2(\mathbb{Z}) \mid g^3 \in \Gamma(2)\},\$$

which is an intermediate group of $\mathrm{SL}_2(\mathbb{Z})$ and $\Gamma(2)$. Note that the quotients $\mathrm{SL}_2(\mathbb{Z})/\Gamma(2)$ and $\Gamma(2)^{1/3}/\Gamma(2)$ are isomorphic to the symmetric group S_3 of degree 3 and the alternating group A_3 of degree 3, respectively. We express the inverses of φ_1 and φ_2 by the lambda function; in fact, φ_1^{-1} is given in Theorem 5.5 by an invariant

$$\nu(\tau) = \frac{3\sqrt{3i\lambda(\tau)(1-\lambda(\tau))}}{(\lambda(\tau)+\omega^2)^3}$$

under the action of $\Gamma(2)^{1/3}/\Gamma(2) \simeq A_3$, and φ_2^{-1} is given in Theorem 6.5 by the reciprocal of the *j*-invariant

$$j(\tau) = \frac{4}{27} \frac{(\lambda(\tau)^2 - \lambda(\tau) + 1)^3}{\lambda(\tau)^2 (1 - \lambda(\tau))^2}.$$

By using the denominators of $\varphi_1, \varphi_1^{-1}$ and $\varphi_2, \varphi_2^{-1}$, we have analogies of Jacobi' formula:

$$\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4 = F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2,$$
$$E_4(\tau) = F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{i(\tau)})^4,$$

in Theorems 5.6, 6.7, where $E_4(\tau)$ is the normalized Eisenstein series of weight 4. We remark that the factor ω^2 in the circuit matrix $\omega^2 W$ along ρ_1 plays an important role in our proof of the former equality.

By eliminating $\vartheta_{pq}(\tau)$ from Jacobi's formula and its analogies, we have functional equations for F(a, b, c; z) such as

$$F(\frac{1}{12}, \frac{5}{12}, 1; \frac{z^2}{4(z-1)})^2 = F(\frac{1}{6}, \frac{1}{2}, 1; z)F(\frac{1}{6}, \frac{1}{2}, 1; \frac{z}{z-1});$$

see Corollary 7.1 for other formulas.

We summarize the contents of this paper. In §2, we collect facts for the hypergeometric series and the hypergeometric differential equations. We cannot select a well-used pair of $z^{1-c}F(a-c+1, b-c+1, 2-c; z)$ and F(a, b, c; z) as a basis of the space of local solutions to E(a, b, c) around $\dot{z} = \frac{1}{2}$, since we set c = 1 in our study. We show the linear independence of F(a, b, a + b - c + 1; 1 - z) and F(a, b, c; z) under some conditions on a, b, c, and give a simple way to obtain circuit matrices of ρ_0 and ρ_1 with respect to constant multiples of them. In §3, we review properties of the theta constant by referring to [Mu]. We give actions of $\operatorname{SL}_2(\mathbb{Z})/\Gamma(2) \simeq S_3$ on $\vartheta_{pq}(\tau)^2$. In §4, we give facts about Schwarz's map φ_0 for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$ by referring to [Yo]. We study Schwarz's map φ_1 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$ in §5 and Schwarz's map φ_2 for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$ in §6, and give functional equations for F(a, b, c; z) in §7.

2. Fundamental properties of F(a, b, c; z)

We begin with defining the hypergeometric series.

DEFINITION 2.1. The hypergeometric series F(a, b, c; z) is defined by

$$F(a, b, c; z) = \sum_{n=0}^{\infty} \frac{(a, n)(b, n)}{(c, n)(1, n)} z^n,$$
(2.1)

where z is a complex variable, a, b, c are complex parameters with $c \neq 0, -1, -2, \cdots$, and $(a, n) = \Gamma(a+n)/\Gamma(a) = a(a+1)\cdots(a+n-1)$. It converges absolutely and uniformly on any compact set in the unit disk $\mathbb{B}_0 = \{z \in \mathbb{C} \mid |z| < 1\}$ for any fixed a, b, c.

The radius of the convergence of F(a, b, c; z) is 1 for generic a, b, c. Gauss's identity gives the limit of F(a, b, c; z) as $z \to 1$ with |z| < 1.

FACT 2.2 (Gauss's identity [Er, (14) in p.61]). If $\operatorname{Re}(c-a-b) > 0$ then

$$\lim_{z \to 1, |z| < 1} F(a, b, c; z) = \frac{\Gamma(c)\Gamma(c - a - b)}{\Gamma(c - a)\Gamma(c - b)}.$$
(2.2)

The series F(a, b, c; z) satisfies the hypergeometric differential equation

$$E(a,b,c): \left[z(1-z)\frac{d^2}{dz^2} + \{c - (a+b+1)z\}\frac{d}{dz} - ab\right] \cdot f(z) = 0,$$
(2.3)

which is a second order linear ordinary differential equation with regular singular points $z = 0, 1, \infty$. The space $\mathcal{F}_{\dot{z}}(a, b, c)$ of (single-valued) local solutions to E(a, b, c) around a point $\dot{z} \in Z = \mathbb{C} - \{0, 1\}$ is a 2-dimensional complex vector space. We take $\dot{z} = \frac{1}{2} \in Z$, and fix it as a base point of Z. We give a basis of $\mathcal{F}_{\dot{z}}(a, b, c)$ for $\dot{z} = \frac{1}{2}$.

LEMMA 2.3 (A fundamental system of solutions to E(a, b, c)). Suppose that a, b, c satisfy $a, b \notin \mathbb{Z}$ and $\operatorname{Re}(a), \operatorname{Re}(b - c + 1), \operatorname{Re}(c - a) > 0$.

(1) The integrals

$$f_1(z) = \int_{-\infty}^0 e^{-\pi \mathbf{i}(a-1)} (-t)^{b-c} (z-t)^{-b} (1-t)^{c-a-1} dt,$$

$$f_2(z) = \int_{1}^{\infty} t^{b-c} (t-z)^{-b} (t-1)^{c-a-1} dt,$$
(2.4)

converge and satisfy the hypergeometric differential equation E(a, b, c), where we assign each branch of the integrands in (2.4) over the integral intervals to positive real values for real z near to $\dot{z} = \frac{1}{2}$. They admit expressions in terms of the hypergeometric series as

$$f_1(z) = e^{-\pi \mathbf{i}(a-1)} B(a, b-c+1) F(a, b, a+b-c+1; 1-z),$$

$$f_2(z) = B(a, c-a) F(a, b, c; z),$$
(2.5)

where B denotes the beta function.

(2) The functions $f_1(z)$ and $f_2(z)$ span the space $\mathcal{F}_{\dot{z}}(a,b,c)$. In particular, we have

$$\det \begin{pmatrix} F(a,b,a+b-c+1;1-z) & \frac{d}{dz}F(a,b,a+b-c+1;1-z) \\ F(a,b,c;z) & \frac{d}{dz}F(a,b,c;z) \end{pmatrix}$$
(2.6)
= $\frac{\Gamma(c)\Gamma(a+b-c+1)}{\Gamma(a)\Gamma(b)}z^{-c}(1-z)^{c-a-b-1}.$

PROOF. (1) It is elementary to show the convergence of these integrals, which are given by a variable change s = 1/t to usual Euler type integrals

$$\int_{\ell} s^{a-1} (1-s)^{c-a-1} (1-sz)^{-b} ds$$

satisfying E(a, b, c), where ℓ is the open interval (0, 1) or $(1, \infty)$. We remark that the integrand $e^{-\pi i (a-1)} (-t)^{b-c} (z-t)^{-b} (1-t)^{c-a-1}$ in $f_1(z)$ is the analytic continuation of $u(t) = t^{b-c} (t-z)^{-b} (t-1)^{c-a-1}$ via the upper-half plane of the t-space. The expression of $f_2(z)$ in (2.5) is a direct consequence of the above, and we obtain that of $f_1(z)$ in (2.5) by a further variable change t' = 1 - t to the integral expressing $f_1(z)$. Note that neither the parameter c nor a + b - c + 1 in (2.5) becomes a non-positive integer under the conditions $\operatorname{Re}(a)$, $\operatorname{Re}(b-c+1)$, $\operatorname{Re}(c-a) > 0$.

(2) By [Va, Theorem 1.1], we have

$$\det \begin{pmatrix} f_1(z) & \frac{d}{dz} f_1(z) \\ f_2(z) & \frac{d}{dz} f_2(z) \end{pmatrix} = e^{-\pi \mathbf{i}(a-1)} \frac{\Gamma(a)\Gamma(c-a)\Gamma(b-c+1)}{\Gamma(b)} z^{-c} (1-z)^{c-a-b-1}$$

Since the last term never vanishes under the assumption, $f_1(z)$ and $f_2(z)$ are linearly independent, and span the space $\mathcal{F}_{\dot{z}}(a, b, c)$. We can rewrite this equality to (2.6) by (2.5). \Box

On a small disk with a point hole in the center z_0 ($z_0 = 0, 1, \infty$), there are (multi-valued) solutions to E(a, b, c) of the forms

$$z^{e_{0,i}} f_{0,i}(z), \quad (1-z)^{e_{1,i}} f_{1,i}(z), \quad (1/z)^{e_{\infty,i}} f_{\infty,i}(z),$$

where $f_{z_0,i}(z)$ are holomorphic functions around $z = z_0$ with $f_{z_0,i}(z_0) = 1$ $(z_0 = 0, 1, \infty)$, and $e_{z_0,i}$ are given in Table 1 called Riemann's scheme.

z_0	0	1	∞
$e_{z_{0},1}$	0	0	a
$e_{z_{0},2}$	1 - c	c-a-b	b
$e_{z_0,2} - e_{z_0,1}$	1 - c	c-a-b	b-a

TABLE 1. Riemann's scheme

Any element in $\mathcal{F}_{\dot{z}}(a,b,c)$ admits the analytic continuation along any path in Z. In particular, a loop ρ in Z with the base point \dot{z} leads to a linear transformation of $\mathcal{F}_{\dot{z}}(a,b,c)$ called the circuit transformation along ρ . We have a homomorphism from the fundamental group $\pi_1(Z, \dot{z})$ to the general linear group $\mathrm{GL}(\mathcal{F}_{\dot{z}}(a,b,c))$ of $\mathcal{F}_{\dot{z}}(a,b,c)$, which is called the monodromy representation of E(a, b, c). We take loops ρ_0 and ρ_1 with terminal \dot{z} as positively oriented circles with radius $\frac{1}{2}$ and center 0 and 1, respectively. Since $\pi_1(Z, \dot{z})$ is a free group generated by ρ_0 and ρ_1 , the monodromy representation of E(a, b, c) is uniquely characterized by the images of ρ_0 and ρ_1 . We give their explicit forms with respect to the basis of $\mathcal{F}_{\dot{z}}(a, b, c)$ in Lemma 2.3.

LEMMA 2.4. The loops ρ_0 and ρ_1 lead to linear transformations sending the basis $\mathbf{F}(z) = {}^t(f_1(z), f_2(z))$ of $\mathcal{F}_{\dot{z}}(a, b, c)$ to $M_0\mathbf{F}(z)$ and $M_1\mathbf{F}(z)$, where the circuit matrices M_0 and M_1 are

$$M_0 = \begin{pmatrix} e^{-2\pi \mathbf{i}c} & e^{-2\pi \mathbf{i}c} - e^{-2\pi \mathbf{i}a} \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & 0 \\ e^{2\pi \mathbf{i}(c-b)} - 1 & e^{2\pi \mathbf{i}(c-a-b)} \end{pmatrix}.$$
(2.7)

PROOF. Since $f_2(z)$ and $f_1(z)$ are single valued on $\mathbb{B}_0 = \{z \in \mathbb{C} \mid |z| < 1\}$ and $\mathbb{B}_1 = \{z \in \mathbb{C} \mid |1 - z| < 1\}$, respectively, the circuit matrices M_0 and M_1 take forms of

$$M_0 = \begin{pmatrix} y'_0 & y_0 \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & 0 \\ y_1 & y'_1 \end{pmatrix}.$$

Since the eigenvalues of M_0 are 1 and $e^{-2\pi i c}$ and those of M_1 are 1 and $e^{2\pi i (c-a-b)}$ by Riemann's scheme in Table 1, we have $y'_0 = e^{-2\pi i c}$ and $y'_1 = e^{2\pi i (c-a-b)}$. We set

$$M'_{\infty} = (M_1 M_0)^{-1} = \begin{pmatrix} y_0 y_1 e^{2\pi \mathbf{i}(a+b)} + e^{2\pi \mathbf{i}c} & -y_0 e^{2\pi \mathbf{i}(a+b)} \\ -y_1 e^{2\pi \mathbf{i}(a+b-c)} & e^{2\pi \mathbf{i}(a+b-c)} \end{pmatrix},$$

which is the circuit matrix of the loop $(\rho_1\rho_0)^{-1} = \rho_0^{-1}\rho_1^{-1}$ with respect to $\mathbf{F}(z)$. Since $\rho_0^{-1}\rho_1^{-1}$ is a loop turning $z = \infty$ once around positively, the eigenvalues of M'_{∞} are $e^{2\pi i a}$, $e^{2\pi i b}$ by Riemann's scheme in Table 1. Thus we have

$$\operatorname{tr}(M'_{\infty}) = y_0 y_1 e^{2\pi \mathbf{i}(a+b)} + e^{2\pi \mathbf{i}c} + e^{2\pi \mathbf{i}(a+b-c)} = e^{2\pi \mathbf{i}a} + e^{2\pi \mathbf{i}b},$$

which relates y_0 and y_1 as

$$y_1 = -\frac{(e^{2\pi i a} - e^{2\pi i c})(e^{2\pi i b} - e^{2\pi i c})}{y_0 e^{2\pi i (a+b+c)}},$$

where $\operatorname{tr}(M'_{\infty})$ denotes the trace of M'_{∞} . By eliminating y_1 in the expression of M'_{∞} , we obtain an $e^{2\pi i b}$ -eigenvector

$$(e^{2\pi i a} - e^{2\pi i c}, e^{2\pi i (a+c)}y_0)$$

of M'_{∞} .

On the other hand, since $\rho_0^{-1}\rho_1^{-1}$ is the loop connecting the end of ρ_0^{-1} to the start of ρ_1^{-1} , it is homotopic to a loop ρ'_{∞} starting from \dot{z} approaching ∞ via the lower-half plane of Z, turning $z = \infty$ once around positively, and tracing back to \dot{z} , see Figure 1. Note the path ℓ_{01} from t = 0 to t = 1 via the upper-half plane of the t-space is not involved in the movement of z along ρ'_{∞} , and that $\arg(t-z)$ on ℓ_{01} decreases 2π by the continuation along ρ'_{∞} . Thus $\int_{\ell_{01}} u(t)dt$ is an $e^{2\pi i b}$ -eigen function under the analytic continuation along ρ'_{∞} , where the branch of u(t) on ℓ_{01} is assigned by the continuation of that on the open interval $(1, \infty)$ defining $f_2(z)$ via the upper-half plane of the t-space. By Cauchy's integral theorem, this integral is expressed as a linear combination

$$-f_1(z) - f_2(z) = -(1,1)\mathbf{F}(z).$$

Since each of $(e^{2\pi i a} - e^{2\pi i c}, e^{2\pi i (a+c)}y_0)$ and (1, 1) is an $e^{2\pi i b}$ -eigenvector of M'_{∞} , they are parallel. Hence we have

$$e^{2\pi \mathbf{i}(a+c)}y_0 = e^{2\pi \mathbf{i}a} - e^{2\pi \mathbf{i}c} \Leftrightarrow y_0 = e^{-2\pi \mathbf{i}c} - e^{-2\pi \mathbf{i}a}$$

This expression of y_0 and the relation between y_0 and y_1 yield (2.7).



FIGURE 1. Loops in Z and the path ℓ_{01}

Remark 2.5. The integrals

$$\int_0^z u(t)dt, \quad \int_z^1 u(t)dt$$

are an eigen function of the circuit transformation along ρ_0 with eigenvalue $e^{-2\pi i c}$ and that along ρ_1 with eigenvalue $e^{2\pi i (c-a-b)}$, respectively, where branches of u(t) on (0, z) and (z, 1) are assigned as the analytic continuations of that on (0, 1) via the upper-half space of the *t*-space. These are expressed by linear combinations of $f_1(z)$ and $f_2(z)$ as

$$\int_{0}^{z} u(t)dt = -\frac{e^{2\pi i c} - 1}{e^{2\pi i b} - 1}f_{1}(z) + \frac{e^{2\pi i a} - e^{2\pi i c}}{e^{2\pi i (a+b)} - e^{2\pi i a}}f_{2}(z),$$
$$\int_{z}^{1} u(t)dt = \frac{e^{2\pi i c} - e^{2\pi i b}}{e^{2\pi i b} - 1}f_{1}(z) - \frac{e^{2\pi i (a+b)} - e^{2\pi i c}}{e^{2\pi i (a+b)} - e^{2\pi i a}}f_{2}(z).$$

REMARK 2.6. In Proof of Lemma 2.4, we consider the loop ρ'_{∞} homotopic to $(\rho_1 \rho_0)^{-1}$ and its circuit matrix

$$M'_{\infty} = (M_1 M_0)^{-1} = \begin{pmatrix} e^{2\pi \mathbf{i}a} + e^{2\pi \mathbf{i}b} - e^{2\pi \mathbf{i}(a+b-c)} & e^{2\pi \mathbf{i}b} - e^{2\pi \mathbf{i}(a+b-c)} \\ e^{2\pi \mathbf{i}(a+b-c)} - e^{2\pi \mathbf{i}a} & e^{2\pi \mathbf{i}(a+b-c)} \end{pmatrix}$$

with respect to $\mathbf{F}(z)$. Its eigen row vectors of eigenvalues $e^{2\pi i a}$ and $e^{2\pi i b}$ are

$$(e^{2\pi \mathbf{i}(a+b)} - e^{2\pi \mathbf{i}(a+c)}, e^{2\pi \mathbf{i}(a+b)} - e^{2\pi \mathbf{i}(b+c)}), \quad (1,1),$$

respectively. We set $\rho_{\infty} = (\rho_0 \rho_1)^{-1}$, which is homotopic to a loop starting from \dot{z} approaching ∞ via the upper-half plane of Z, turning $z = \infty$ once around positively, and

tracing back to \dot{z} . Its circuit matrix M_{∞} with respect to $\mathbf{F}(z)$ is

$$M_{\infty} = (M_0 M_1)^{-1} = \begin{pmatrix} e^{2\pi \mathbf{i}c} & e^{2\pi \mathbf{i}(c-a)} - 1\\ e^{2\pi \mathbf{i}(a+b)} - e^{2\pi \mathbf{i}(a+c)} & e^{2\pi \mathbf{i}a} + e^{2\pi \mathbf{i}b} - e^{2\pi \mathbf{i}c} \end{pmatrix}.$$

Its eigen row vectors of eigenvalues $e^{2\pi i a}$ and $e^{2\pi i b}$ are

$$(e^{2\pi \mathbf{i}(a+b)} - e^{2\pi \mathbf{i}(a+c)}, e^{2\pi \mathbf{i}a} - e^{2\pi \mathbf{i}c}), \quad (e^{2\pi \mathbf{i}a}, 1),$$

respectively.

REMARK 2.7. We can make the analytic continuation of the hypergeometric series F(a, b, c; z) to the simply connected domain $\mathbb{C} - [1, \infty)$ as a solution to E(a, b, c). We use the same symbol F(a, b, c; z) for this continuation, which is a single-valued holomorphic function on $\mathbb{C} - [1, \infty)$.

REMARK 2.8. We set $B(a, c - a)F(a, b, c; z) = \int_{1}^{\infty} u(t)dt$ in the second entry of the column vector $\mathbf{F}(z)$ for the compatibility of the monodromy representation with the action of linear fractional transformations on the upper-half plane \mathbb{H} .

DEFINITION 2.9 (Schwarz's map). For a basis ${}^{t}(\varphi_{1}(z), \varphi_{2}(z))$ of $\mathcal{F}_{\dot{z}}(a, b, c)$, we have a map from a neighborhood of $\dot{z} = \frac{1}{2}$ to the complex projective line \mathbb{P}^{1} by

$$\varphi: z \mapsto \varphi(z) = \frac{\varphi_1(z)}{\varphi_2(z)} \tag{2.8}$$

Schwarz's map is define by its analytic continuation to $Z = \mathbb{C} - \{0, 1\}$, which is also denoted by φ . Though Schwarz's map φ is regarded as multi-valued on Z, its restriction to a simply connected domain $\mathbb{C} - (-\infty, 0] - [1, \infty)$ is single valued.

FACT 2.10. If the parameters a, b, c are real and satisfy

$$|1 - c| + |c - a - b| + |a - b| < 1,$$

then there exists a basis $\varphi_1(z), \varphi_2(z)$ of $\mathcal{F}_{\dot{z}}(a, b, c)$ such that the image of Schwarz's map φ is in the upper-half plane $\mathbb{H} = \{\tau \in \mathbb{C} \mid \operatorname{Im}(\tau) > 0\}$. For this basis, the image of the upper-half plane in $Z = \mathbb{C} - \{0, 1\}$ under Schwarz's map φ becomes a triangle with angles $|1 - c|\pi, |c - a - b|\pi, |a - b|\pi$ with respect to the hyperbolic metric on \mathbb{H} , which is called Schwarz's triangle and denoted by $\varphi(\mathbb{H})$. Moreover, a, b, c satisfy

$$\frac{1}{|1-c|}, \frac{1}{|c-a-b|}, \frac{1}{|a-b|} \in \{2, 3, 4, \dots\} \cup \{\infty\}$$

then the image of Schwarz's map is an open dense subset in \mathbb{H} , and its inverse is single valued. The projectivization of the monodromy representation of E(a, b, c) is represented as a discrete subgroup of $\mathrm{PSL}_2(\mathbb{R}) = \mathrm{SL}_2(\mathbb{R})/\{\pm I_2\}$ generated by two elements g_0 and g_1 with

$$\operatorname{ord}(g_0) = \frac{1}{|1-c|}, \quad \operatorname{ord}(g_1) = \frac{1}{|c-a-b|}, \quad \operatorname{ord}(g_\infty) = \frac{1}{|a-b|},$$

where I_2 is the unit matrix of size 2, $g_{\infty} = (g_0 g_1)^{-1}$, and $\operatorname{ord}(g_0)$ denotes the order of g_0 .

3. Fundamental properties of $\vartheta_{pq}(\tau)$

We next introduce theta constants and their properties by referring to [Mu].

DEFINITION 3.1. The theta constant $\vartheta_{pq}(\tau)$ with characteristics p, q is defined by

$$\vartheta_{pq}(\tau) = \sum_{n=-\infty}^{\infty} \exp\left(\pi \mathbf{i}(n+\frac{p}{2})^2 \tau + 2\pi \mathbf{i}(n+\frac{p}{2})\frac{q}{2}\right),\tag{3.1}$$

where τ is a variable in the upper-half plane \mathbb{H} , and p, q are parameters taking the value 0 or 1. It converges absolutely and uniformly on any compact set in \mathbb{H} for any fixed p, q.

There are four functions $\vartheta_{00}(\tau)$, $\vartheta_{01}(\tau)$, $\vartheta_{10}(\tau)$, $\vartheta_{11}(\tau)$; the last one vanishes identically on \mathbb{H} , and the rests satisfy Jacobi's identity

$$\vartheta_{00}(\tau)^4 = \vartheta_{01}(\tau)^4 + \vartheta_{10}(\tau)^4 \tag{3.2}$$

for any $\tau \in \mathbb{H}$.

The group $SL_2(\mathbb{Z})$ acts on \mathbb{H} by the linear fractional transformation

$$g \cdot \tau = \frac{g_{11}\tau + g_{12}}{g_{21}\tau + g_{22}}$$

where $g = (g_{ij}) \in \mathrm{SL}_2(\mathbb{Z})$ and $\tau \in \mathbb{H}$. Note that $(-I_2) \cdot \tau = \tau$ for any $\tau \in \mathbb{H}$ and that the projectivization $\mathrm{PSL}_2(\mathbb{Z}) = \mathrm{SL}_2(\mathbb{Z})/\{\pm I_2\}$ acts effectively on \mathbb{H} .

The group $SL_2(\mathbb{Z})$ is generated by two elements

$$T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$
 (3.3)

We give transformation formulas of $\vartheta_{pq}(\tau)^2$ under the actions T and J on τ :

$$T \cdot \tau = \tau + 1, \quad J \cdot \tau = \frac{-1}{\tau}.$$

FACT 3.2 ([Mu, Table V in p.36]). We have

$$\vartheta_{00}(\tau+1)^2 = \vartheta_{01}(\tau)^2, \quad \vartheta_{01}(\tau+1)^2 = \vartheta_{00}(\tau)^2, \quad \vartheta_{10}(\tau+1)^2 = \mathbf{i}\vartheta_{10}(\tau)^2, \\ \vartheta_{00}(\frac{-1}{\tau})^2 = (-\mathbf{i}\tau)\vartheta_{00}(\tau)^2, \quad \vartheta_{01}(\frac{-1}{\tau})^2 = (-\mathbf{i}\tau)\vartheta_{10}(\tau)^2, \quad \vartheta_{10}(\frac{-1}{\tau})^2 = (-\mathbf{i}\tau)\vartheta_{01}(\tau)^2.$$

The theta constant $\vartheta_{00}(\tau)$ is quasi invariant under the action of the subgroup

$$\Gamma_{12} = \{ g = (g_{ij}) \in \mathrm{SL}_2(\mathbb{Z}) \mid g_{11}g_{12} \equiv g_{21}g_{22} \equiv 0 \bmod 2 \}$$

of $SL_2(\mathbb{Z})$.

FACT 3.3 ([Mu, Theorem 7.1, Table V in p.36]). Let $g = (g_{ij})$ be an element in the projectivization $P\Gamma_{12}$ of Γ_{12} . By multiplying -1 to g if necessary, we may assume that g satisfies either $g_{21} > 0$, or $g_{21} = 0$ and $g_{22} > 0$. Then we have

$$\vartheta_{00}(g \cdot \tau)^2 = \chi(g) \cdot (g_{21}\tau + g_{22})\vartheta_{00}(\tau)^2$$

for any $\tau \in \mathbb{D}_{12}$, where

$$\chi(g) = \begin{cases} \mathbf{i}^{g_{22}-1} & \text{if } g_{21} \in 2\mathbb{Z}, g_{22} \notin 2\mathbb{Z}, \\ \mathbf{i}^{-g_{21}} & \text{if } g_{21} \notin 2\mathbb{Z}, g_{22} \in 2\mathbb{Z}, \end{cases}$$

Let $\Gamma(2)$ be the principal congruence subgroup of $SL_2(\mathbb{Z})$ of level 2, i.e.,

$$\Gamma(2) = \{ g = (g_{ij}) \in \mathrm{SL}_2(\mathbb{Z}) \mid g_{11} - 1, g_{22} - 1, g_{12}, g_{21} \equiv 0 \bmod 2 \}.$$

The group $\Gamma(2)$ is generated by $T^2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$, $J^{-1}T^2J = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}$ and $-I_2$. Facts 3.2 and 3.3 imply the following.

FACT 3.4. The theta constants $\vartheta_{pq}(\tau)$ ((p,q) = (0,0), (0,1), (1,0)) satisfy

$$\vartheta_{pq}(g \cdot \tau)^4 = (g_{21}\tau + g_{22})^2 \vartheta_{pq}(\tau)^4$$

for any $g = (g_{ij}) \in \Gamma(2)$.

The group $\Gamma(2)$ is normal in $SL_2(\mathbb{Z})$, and the quotient

$$\operatorname{SL}_2(\mathbb{Z})/\Gamma(2) = \{I_2, W, W^2, J, T, J^{-1}TJ\}, \quad W = (JT)^{-1} = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$$

is isomorphic to the symmetric group S_3 of degree 3. There exists a subgroup G of $SL_2(\mathbb{Z})$ such that the quotient $G/\Gamma(2)$ is isomorphic to the alternating group A_3 of degree 3. This group is generated by $\Gamma(2)$ and W, and characterized by

$$\{g \in \mathrm{SL}_2(\mathbb{Z}) \mid g^3 \in \Gamma(2)\},\$$

which is denoted by $\Gamma(2)^{1/3}$. Note that

$$[\operatorname{SL}_2(\mathbb{Z}): \Gamma(2)^{1/3}] = 2, \quad [\Gamma(2)^{1/3}: \Gamma(2)] = 3.$$

We give a fundamental region of $P\Gamma(2)^{1/3} = \Gamma(2)^{1/3}/\{\pm I_2\}$ and that of $PSL_2(\mathbb{Z})$ as

$$\widetilde{\mathbb{D}} = \{ \tau \in \mathbb{H} \mid -\frac{3}{2} < \operatorname{Re}(\tau) \le \frac{1}{2}, \ |\tau| \ge 1, \ |\tau+1| > 1 \},$$
(3.4)

$$\mathbb{D} = \{\tau \in \mathbb{H} \mid -\frac{1}{2} < \operatorname{Re}(\tau) \le \frac{1}{2}, \ |\tau| > 1\} \cup \{\tau \in \mathbb{H} \mid |\tau| = 1, 0 \le \operatorname{Re}(\tau) \le \frac{1}{2}\}, \quad (3.5)$$

see Figure 2.

LEMMA 3.5. By the actions of $W = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$ and $W^2 = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$ on $\tau \in \mathbb{H}$, $\vartheta_{00}(\tau)^2$, $\vartheta_{01}(\tau)^2$ and $\vartheta_{10}(\tau)^2$ are transformed into

$$\vartheta_{00}(W \cdot \tau)^{2} = -\mathbf{i}\tau \cdot \vartheta_{10}(\tau)^{2}, \quad \vartheta_{00}(W^{2} \cdot \tau)^{2} = -\mathbf{i}(\tau+1) \cdot \vartheta_{01}(\tau)^{2}, \\
\vartheta_{01}(W \cdot \tau)^{2} = -\mathbf{i}\tau \cdot \vartheta_{00}(\tau)^{2}, \quad \vartheta_{01}(W^{2} \cdot \tau)^{2} = (\tau+1) \cdot \vartheta_{10}(\tau)^{2}, \\
\vartheta_{10}(W \cdot \tau)^{2} = -\tau \cdot \vartheta_{01}(\tau)^{2}, \quad \vartheta_{10}(W^{2} \cdot \tau)^{2} = -\mathbf{i}(\tau+1) \cdot \vartheta_{00}(\tau)^{2}.$$
(3.6)

PROOF. By Fact 3.2 together with $W = T^{-1}J^{-1}$ and $J^{-1} = -J$, we have

$$\vartheta_{00}(W\cdot\tau)^2 = \vartheta_{00}(T^{-1}\cdot(J\cdot\tau))^2 = \vartheta_{01}(J\cdot\tau)^2 = (-\mathbf{i}\tau)\cdot\vartheta_{10}(\tau)^2,$$

$$\vartheta_{01}(W\cdot\tau)^2 = \vartheta_{01}(T^{-1}\cdot(J\cdot\tau))^2 = \vartheta_{00}(J\cdot\tau)^2 = (-\mathbf{i}\tau)\cdot\vartheta_{00}(\tau)^2,$$

$$\vartheta_{10}(W\cdot\tau)^2 = \vartheta_{10}(T^{-1}\cdot(J\cdot\tau))^2 = -\mathbf{i}\vartheta_{10}(J\tau)^2 = -\mathbf{i}\cdot(-\mathbf{i}\tau)\cdot\vartheta_{01}(\tau)^2.$$

Similarly, we can show the transformations of $\vartheta_{pq}(\tau)^2$ under the action of $W^2 = JT$. \Box



FIGURE 2. Fundamental regions $\widetilde{\mathbb{D}}$ and \mathbb{D} for $\mathrm{P}\Gamma(2)^{1/3}$ and $\mathrm{PSL}_2(\mathbb{Z})$

DEFINITION 3.6. We define the (normalized) Eisenstein series $E_{2k}(\tau)$ of weight 2k $(k \in \mathbb{N})$ by

$$E_{2k}(\tau) = \frac{1}{2\zeta(2k)} \sum_{n_1, n_2 \in \mathbb{Z}^2 - \{(0,0)\}} \frac{1}{(n_1\tau + n_2)^{2k}},$$

where $\tau \in \mathbb{H}$ and ζ denotes Riemann's zeta function.

If $k \geq 2$ then this series converges absolutely, and it satisfies

$$E_{2k}(g \cdot \tau) = (g_{21}\tau + g_{22})^{2k} E_{2k}(\tau)$$

for any $g \in \mathrm{SL}_2(\mathbb{Z})$.

FACT 3.7 ([Mu, §15]). The Eisenstein series $E_4(\tau)$ is expressed by a Fourier expansion and by the theta constants as

$$E_4(\tau) = 1 + 240 \sum_{n=1}^{\infty} \left(\sum_{d|n} d^3\right) e^{2\pi i n \tau}$$

= 1 + 240 e^{2\pi i \tau} + 2160 (e^{2\pi i \tau})^2 + 6720 (e^{2\pi i \tau})^3 + \cdots (3.7)

$$=\frac{\vartheta_{00}(\tau)^8 + \vartheta_{01}(\tau)^8 + \vartheta_{10}(\tau)^8}{2}.$$
(3.8)

4. Schwarz's map for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$

To study Schwarz's map φ_0 for parameters $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$, we select $\mathbf{F}(z) = {}^t(f_1, f_2)$ as a basis of $\mathcal{F}_{\dot{z}}(\frac{1}{2}, \frac{1}{2}, 1)$. Schwarz's map φ_0 is the analytic continuation of

$$\varphi_0: z \mapsto \tau = \varphi_0(z) = \frac{f_1(z)}{f_2(z)} = \mathbf{i} \cdot \frac{F(\frac{1}{2}, \frac{1}{2}, 1, 1 - z)}{F(\frac{1}{2}, \frac{1}{2}, 1, z)}$$
(4.1)

defined on $\mathbb{B}_0 \cap \mathbb{B}_1 = \{z \in \mathbb{C} \mid |z| < 1, |1-z| < 1\}$ to $Z = \mathbb{C} - \{0,1\}$. We have $\left(\frac{1}{|1-c|}, \frac{1}{|c-a-b|}, \frac{1}{|a-b|}\right) = (\infty, \infty, \infty)$, and the circuit matrices M_0, M_1 and $(M_0M_1)^{-1}$ are

$$M_0 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix}, \quad (M_0 M_1)^{-1} = \begin{pmatrix} 1 & -2 \\ 2 & -3 \end{pmatrix}.$$

The group generated by these matrices is not the principal congruence subgroup $\Gamma(2)$ but

$$\Gamma(2,4) = \{g = (g_{ij}) \in \mathrm{SL}_2(\mathbb{Z}) \mid g_{11}, g_{22} \equiv 1 \mod 4, g_{12}, g_{21} \equiv 0 \mod 2\}.$$

Note that $\Gamma(2,4)$ is a subgroup in $\Gamma(2)$ of index 2, and that $P\Gamma(2) = \Gamma(2)/\{\pm I_2\}$ is isomorphic to $\Gamma(2,4)$.

Since $F(\frac{1}{2}, \frac{1}{2}, 1, 1-z)$ and $F(\frac{1}{2}, \frac{1}{2}, 1, z)$ take positive real values for z in the open interval $(0, 1), \varphi_0(z)$ for $z \in (0, 1)$ is a pure imaginary number in \mathbb{H} . Since the image of the monodromy representation is in $\mathrm{PSL}_2(\mathbb{R})$ and $f_1(z)/f_2(z)$ is in \mathbb{H} for $z \in (0, 1)$, the image of the analytic continuation of $f_1(z)/f_2(z)$ to $Z = \mathbb{C} - \{0, 1\}$ is an open dense subset in \mathbb{H} .

FACT 4.1 (Schwarz's map for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$). [Yo, §5.6,5.7,5.8 in Chap. II, Proposition 8.1 in Chap. III]

(1) Schwarz's triangle $\varphi_0(\mathbb{H})$ for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$ is

$$\{\tau \in \mathbb{H} \mid 0 < \operatorname{Re}(\tau) < 1, \ |\tau - \frac{1}{2}| > \frac{1}{2}\}$$

and its vertices $\varphi_0(z_0) = \lim_{z \to z_0, z \in \mathbb{H}} \varphi_0(z) \ (z_0 = 0, 1, \infty)$ are

$$\varphi_0(0) = \mathbf{i}\infty, \quad \varphi_0(1) = 0, \quad \varphi_0(\infty) = 1.$$

The images of the intervals (0, 1), $(-\infty, 0)$, $(1, \infty)$ consisting of the boundary of $\mathbb{H}(\subset Z = \mathbb{C} - \{0, 1\})$ under φ_0 are $\{\tau \in \mathbb{H} \mid \operatorname{Re}(\tau) = 0\}$, $\{\tau \in \mathbb{H} \mid \operatorname{Re}(\tau) = 1\}$, $\{\tau \in \mathbb{H} \mid |\tau - \frac{1}{2}| = \frac{1}{2}\}$, respectively.



FIGURE 3. Schwarz's triangle for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$

- (2) The image $\varphi_0(\overline{\mathbb{H}})$ of the lower-half plane $\overline{\mathbb{H}}$ under φ_0 is the reflection of Schwarz's triangle $\varphi_0(\mathbb{H})$ with respect to the imaginary axis. The union $\varphi_0(\mathbb{H})^c \cup \varphi_0(\overline{\mathbb{H}})$ forms a fundamental region $\mathbb{D}(2)$ of $\mathrm{P}\Gamma(2)$, where $\varphi_0(\mathbb{H})^c$ denotes the closure of $\varphi_0(\mathbb{H})$ in \mathbb{H} .
- (3) The inverse of Schwarz's map for $(a, b, c) = (\frac{1}{2}, \frac{1}{2}, 1)$ is

$$\mathbb{H} \ni \tau \mapsto \lambda(\tau) = \frac{\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4} \in Z.$$
(4.2)

The function $\lambda(\tau)$ is invariant under the action $\tau \mapsto (g_{11}\tau + g_{12})/(g_{21}\tau + g_{22})$ of $g = (g_{ij}) \in P\Gamma(2)$ on $\tau \in \mathbb{H}$.

(4) Schwarz's map φ_0 and its inverse λ are extended to isomorphisms between the complex projective line $\mathbb{P}^1(\supset Z = \mathbb{C} - \{0, 1\})$ and the compactification $\widehat{\mathbb{H}}/\mathrm{P\Gamma}(2)$ of $\mathbb{H}/\mathrm{P\Gamma}(2)$, where $\widehat{\mathbb{H}} = \mathbb{H} \cup \{\mathbf{i}\infty\} \cup \mathbb{Q}$ and $\mathrm{P\Gamma}(2)$ acts naturally on $\{\mathbf{i}\infty\} \cup \mathbb{Q}$ and there are three orbits represented by the vertices $\mathbf{i}\infty = \varphi_0(0), 0 = \varphi_0(1), 1 = \varphi_0(\infty)$ of Schwarz's triangle. These extended isomorphisms are also denoted by the same symbols φ_0 and λ .

LEMMA 4.2. (1) The function λ is transformed into $\lambda(T \cdot \tau) = \frac{\lambda(\tau)}{\lambda(\tau) - 1}$ under the action of $T \in SL_2(\mathbb{Z})$.

(2) The function λ sends the mid points \mathbf{i} , $\frac{1+\mathbf{i}}{2}$, $1+\mathbf{i}$ of the edges of Schwarz's triangle $\varphi_0(\mathbb{H})$ to

$$\lambda(\mathbf{i}) = \frac{1}{2}, \quad \lambda(\frac{1+\mathbf{i}}{2}) = 2, \quad \lambda(1+\mathbf{i}) = -1.$$

(3) The function λ sends the centers ω^2 and ω of gravity of Schwarz's triangles $\varphi_0(\mathbb{H})$ and $\varphi_0(\overline{\mathbb{H}})$ to

$$\lambda(-\omega^2) = -\omega^2, \quad \lambda(\omega) = -\omega.$$

PROOF. (1) By Fact 3.2 and (3.2), we have

$$\lambda(T \cdot \tau) = \frac{\vartheta_{10}(T \cdot \tau)^4}{\vartheta_{00}(T \cdot \tau)^4} = \frac{-\vartheta_{10}(\tau)^4}{\vartheta_{01}(\tau)^4} = \frac{-\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4 - \vartheta_{10}(\tau)^4} = \frac{\lambda(\tau)}{\lambda(\tau) - 1}.$$

(2) Note that $\mathbf{i} \in \mathbb{H}$ is a fix point under the action of $J \in SL_2(\mathbb{Z})$. By Fact 3.2 and (3.2), the function $\lambda(\tau)$ is transformed into

$$\lambda(J \cdot \tau) = \frac{\vartheta_{10}(J \cdot \tau)^4}{\vartheta_{00}(J \cdot \tau)^4} = \frac{\vartheta_{01}(\tau)^4}{\vartheta_{00}(\tau)^4} = \frac{\vartheta_{00}(\tau)^4 - \vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4} = 1 - \lambda(\tau)$$

under the action of J. Thus we have

 $\lambda(\mathbf{i}) = \lambda(J \cdot \mathbf{i}) = 1 - \lambda(\mathbf{i}),$

which yields $\lambda(\mathbf{i}) = \frac{1}{2}$. The points $\frac{1+\mathbf{i}}{2}$ and $1+\mathbf{i}$ in \mathbb{H} are fixed by $J_1 = \begin{pmatrix} -1 & 1 \\ -2 & 1 \end{pmatrix}$ and $J_2 = \begin{pmatrix} -1 & 2 \\ -1 & 1 \end{pmatrix}$, which transform $\lambda(\tau)$ into $\frac{\lambda(\tau)}{\lambda(\tau)-1}$ and $\frac{1}{\lambda(\tau)}$, respectively by Fact 3.2 and (3.2) together with $J_1 = J^{-1}T^{-1}JTJ$, $J_2 = TJT^{-1}$. Thus we have $\lambda(\frac{1+\mathbf{i}}{2}) = 2$ and $\lambda(1+\mathbf{i}) = -1$ since $\lambda(\tau) \neq 0, 1$ for any $\tau \in \mathbb{H}$.

(3) The point $\omega \in \mathbb{H}$ is fixed by $W = (JT)^{-1}$, which transform $\lambda(\tau)$ into $1 - \frac{1}{\lambda(\tau)}$ by Lemma 3.5. Thus we have $\lambda(\omega) = 1 - \frac{1}{\lambda(\omega)}$, which yields that $\lambda(\omega)$ is $-\omega$ or $-\omega^2$. Since ω is in the left-half plane of \mathbb{C} , $\lambda(\omega)$ should be $-\omega$ in the lower-half plane of \mathbb{C} by Fact 4.1 (2). Note that $\lambda(\omega) = -\omega$ is equivalent to $\varphi_0(-\omega) = \omega$. Since $-\omega = -\omega^2$ and the reflection image of ω with respect to the imaginary axis is $-\omega^2$, Fact 4.1 (2) also implies $\varphi_0(-\omega^2) = -\omega^2$, which is equivalent to $\lambda(-\omega^2) = -\omega^2$.

As given in [BB1, Theorem 2.1], the hypergeometric series $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ and the theta constants are related by Jacobi's formula:

FACT 4.3 (Jacobi's formula). Jacobi's formula is

$$\vartheta_{00}(\tau)^2 = F(\frac{1}{2}, \frac{1}{2}, 1; \lambda(\tau)) \tag{4.3}$$

for any τ in the interior $\mathbb{D}(2)^{\circ}$ of the fundamental region $\mathbb{D}(2)$ of $\mathrm{PF}(2)$, where $F(\frac{1}{2}, \frac{1}{2}, 1; z)$ denotes the single-valued function on $\mathbb{C} - [1, \infty)$ given in Remark 2.7. By acting T on τ in (4.3), we have

$$\vartheta_{01}(\tau)^2 = F(\frac{1}{2}, \frac{1}{2}, 1; \frac{\lambda(\tau)}{\lambda(\tau) - 1})$$
(4.4)

for any $\tau \in \mathbb{D}(2)^{\circ}$.

REMARK 4.4. Since \mathbb{H} is simply connected, we can extend (4.3) and (4.4) to equalities holding for any τ in the whole space \mathbb{H} . For these extensions, we should carefully trace the continuations of the right hand sides of (4.3) and (4.4), for details refer to [CM, Corollary 4.7]. By Schwarz's map

$$\varphi_0: Z \ni z \mapsto \tau = \mathbf{i} \frac{F(\frac{1}{2}, \frac{1}{2}, 1; 1 - z)}{F(\frac{1}{2}, \frac{1}{2}, 1; z)} \in \mathbb{H},$$

the equalities (4.3) and (4.4) are pulled back to

$$F(\frac{1}{2}, \frac{1}{2}, 1; z) = \vartheta_{00}(\varphi_0(z))^2, \quad F(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1}) = \vartheta_{01}(\varphi_0(z))^2, \tag{4.5}$$

which are initially equalities on $\mathbb{B}_0 \cap \mathbb{B}_1$ and extended to those on the whole space Z by the analytic continuation.

5. Schwarz's map for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$

We consider the hypergeometric differential equation $E(\frac{1}{6}, \frac{1}{2}, 1)$. Its Riemann's scheme becomes as in Table 2. The basis of the space of its local solutions around $\dot{z} = \frac{1}{2}$ in Lemma 2.3 becomes

$$\begin{pmatrix} \exp(\frac{5\pi \mathbf{i}}{6}) \int_{-\infty}^{0} (-t)^{-1/2} (z-t)^{-1/2} (1-t)^{-1/6} dt \\ \int_{1}^{\infty} (t)^{-1/2} (t-z)^{-1/2} (t-1)^{-1/6} dt \end{pmatrix} = \begin{pmatrix} \exp(\frac{5\pi \mathbf{i}}{6}) B(\frac{1}{6}, \frac{1}{2}) F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}, 1-z) \\ B(\frac{1}{6}, \frac{5}{6}) F(\frac{1}{6}, \frac{1}{2}, 1, z) \end{pmatrix}$$

z_0	0	1	∞
$e_{z_0,1}$	0	0	1/6
$e_{z_0,2}$	0	1/3	1/2
$e_{z_0,2} - e_{z_0,1}$	$1/\infty$	1/3	1/3

TABLE 2. Riemann's scheme for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$

The circuit matrices with respect to this basis are

$$M_0 = \begin{pmatrix} 1 & -\omega^2 \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & 0 \\ -2 & \omega \end{pmatrix}.$$

Since these matrices are not in $\text{PSL}_2(\mathbb{R})$, the analytic continuation of $f_1(z)/f_2(z)$ does not stay in \mathbb{H} . We find a matrix $R \in \text{GL}_2(\mathbb{C})$ such that

$$RM_0R^{-1}, RM_1R^{-1} \in PSL_2(\mathbb{Z}).$$

We can diagonalize M_1 by $P_x = \begin{pmatrix} \frac{-1+\omega}{2} & 0\\ 1 & x \end{pmatrix}$ with a parameter x, i.e.,

$$P_x^{-1}M_1P_x = \begin{pmatrix} 1 & 0\\ 0 & \omega \end{pmatrix}$$

holds for any $x \in \mathbb{C}$. On the other hand, since

$$\begin{pmatrix} \omega & \omega^2 \\ 1 & 1 \end{pmatrix}^{-1} \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \omega & \omega^2 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} \omega & 0 \\ 0 & \omega^2 \end{pmatrix},$$

we have

$$R_x M_1 R_x^{-1} = \omega^2 \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix} = N_1$$

by setting

$$R_x = \begin{pmatrix} \omega & \omega^2 \\ 1 & 1 \end{pmatrix} P_x^{-1}.$$

The matrix $R_x M_0 R_x^{-1}$ becomes

$$\frac{1}{3x} \left(\begin{array}{c} \left(-1+\sqrt{3}\mathbf{i}\right)x^2+5x-1-\sqrt{3}\mathbf{i} & \left(1+\sqrt{3}\mathbf{i}\right)x^2+4x+1-\sqrt{3}\mathbf{i} \\ 2x^2-4x+2 & \left(1-\sqrt{3}\mathbf{i}\right)x^2+x+1+\sqrt{3}\mathbf{i} \end{array} \right).$$

Since this matrix is not diagonalizable and has an eigenvalue 1, if the (2, 1)-entry of this matrix vanishes then the action of this matrix becomes a translation on \mathbb{C} . By solving $2x^2 - 4x + 2 = 0$ equivalent to the vanishing of (2, 1)-entry of $R_x M_0 R_x^{-1}$, we have x = 1 and the above matrix becomes $N_0 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}$ and R_x for x = 1 is

$$R = R_1 = \begin{pmatrix} \omega & \omega^2 \\ 1 & 1 \end{pmatrix} P_1^{-1} = \begin{pmatrix} -2\omega & \omega^2 \\ 0 & 1 \end{pmatrix}.$$

Thus the circuit matrices M_0 and M_1 are transformed into

$$RM_0R^{-1} = N_0 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \quad RM_1R^{-1} = N_1 = \omega^2 \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}.$$

The circuit matrix N_{∞} around $z = \infty$ is

$$N_{\infty} = (N_0 N_1)^{-1} = (-\omega) \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix},$$

which satisfy $N_{\infty}^3 = -I_2$. Thus N_{∞} is of order 6, and of order 3 as a projective element. Since

$$N_1^{-1}N_0N_1 = \begin{pmatrix} 1 & 0 \\ -2 & 1 \end{pmatrix},$$

the group generated by N_0 and N_1 are

$$\Gamma(2) \cup (\omega^2 W \cdot \Gamma(2)) \cup (\omega W^2 \cdot \Gamma(2)),$$

where
$$W = \begin{pmatrix} -1 & -1 \\ 1 & 0 \end{pmatrix}$$
, $W^2 = \begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$, $W^3 = I_2$ and
 $\omega^2 W \cdot \Gamma(2) = \{\omega^2 W g \in \operatorname{GL}_2(\mathbb{Z}[\omega]) \mid g \in \Gamma(2)\}$
 $\omega W^2 \cdot \Gamma(2) = \{\omega W^2 g \in \operatorname{GL}_2(\mathbb{Z}[\omega]) \mid g \in \Gamma(2)\}$

The group $\Gamma(2)$ is normal in this group, and we have

$$[\Gamma(2) \cup \omega^2 W \cdot \Gamma(2) \cup \omega W^2 \cdot \Gamma(2)] / \Gamma(2) \simeq \{I_2, W, W^2\},\$$

$$[\Gamma(2) \cup (\omega^2 W \cdot \Gamma(2)) \cup (\omega W^2 \cdot \Gamma(2))] / \{\pm I_2, \pm \omega I_2, \pm \omega^2 I_2\} \simeq \Pr(2)^{1/3}.$$

We change the basis $\mathbf{F}(z)$ into $R \cdot \mathbf{F}(z)$, then $R \cdot \mathbf{F}(z)$ takes the form

$$\begin{pmatrix} -2\omega f_1(z) + \omega^2 f_2(z) \\ f_2(z) \end{pmatrix} = \begin{pmatrix} 2\mathbf{i}B(\frac{1}{6}, \frac{1}{2})F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z) + 2\omega^2\pi F(\frac{1}{6}, \frac{1}{2}, 1; z) \\ 2\pi F(\frac{1}{6}, \frac{1}{2}, 1; z) \end{pmatrix}.$$

DEFINITION 5.1 (Schwarz's map φ_1 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$). We define Schwarz's map φ_1 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$ by the analytic continuation of the map

$$\varphi_{1}: z \mapsto \tau = \frac{2\mathbf{i}B(\frac{1}{6}, \frac{1}{2})F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z) + 2\omega^{2}\pi F(\frac{1}{6}, \frac{1}{2}, 1; z)}{2\pi F(\frac{1}{6}, \frac{1}{2}, 1; z)}$$
$$= \omega^{2} + \mathbf{i} \cdot \frac{B(\frac{1}{6}, \frac{1}{2})}{\pi} \cdot \frac{F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z)}{F(\frac{1}{6}, \frac{1}{2}, 1; z)}$$

defined on $\mathbb{B}_0 \cap \mathbb{B}_1$ to $\mathbb{C} - \{0, 1\}$. Its projective monodromy representation is isomorphic to $\mathrm{P}\Gamma(2)^{1/3}$.

PROPOSITION 5.2. Schwarz's triangle $\varphi_1(\mathbb{H})$ for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$ is

$$\{\tau \in \mathbb{H} \mid -\frac{1}{2} < \operatorname{Re}(\tau) < \frac{1}{2}, \ |\tau| > 1\},\$$

which is the interior of the fundamental domain \mathbb{D} for $PSL_2(\mathbb{Z})$, and its vertices $\varphi_1(z_0) = \lim_{z \to z_0, z \in \mathbb{H}} \varphi_1(z)$ $(z_0 = 0, 1, \infty)$ are

$$\varphi_1(0) = \mathbf{i}\infty, \quad \varphi_1(1) = \omega, \quad \varphi_1(\infty) = -\omega^2.$$

The images of the intervals (0,1), $(-\infty,0)$, $(1,\infty)$ consisting of the boundary of $\mathbb{H}(\subset Z = \mathbb{C} - \{0,1\})$ under φ_1 are

$$\begin{aligned} \{\tau \in \mathbb{H} \mid \operatorname{Re}(\tau) &= -\frac{1}{2}, \ \operatorname{Im}(\tau) > \frac{\sqrt{3}}{2} \}, \\ \{\tau \in \mathbb{H} \mid \operatorname{Re}(\tau) &= \frac{1}{2}, \ \operatorname{Im}(\tau) > \frac{\sqrt{3}}{2} \}, \\ \{\tau \in \mathbb{H} \mid |\tau| = 1, \ |\operatorname{Re}(\tau)| < \frac{1}{2} \}, \end{aligned}$$

respectively. See Figure 2 for their shape.

PROOF. We firstly consider the image of (0, 1) under φ_1 . Note that $F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z)$ and $F(\frac{1}{6}, \frac{1}{2}, 1; z)$ take positive real values. Since

$$\lim_{z \to 0, z \in (0,1)} F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z) = \infty, \quad \lim_{z \to 0, z \in (0,1)} F(\frac{1}{6}, \frac{1}{2}, 1; z) = 1$$

by Fact 2.2 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, \frac{2}{3})$ together with c - a - b = 0, we have $\varphi_1(0) = \mathbf{i}\infty$. Since

$$\lim_{z \to 1, z \in (0,1)} F(\frac{1}{6}, \frac{1}{2}, \frac{2}{3}; 1-z) = 1, \quad \lim_{z \to 1, z \in (0,1)} F(\frac{1}{6}, \frac{1}{2}, 1; z) = \frac{\Gamma(1)\Gamma(1 - \frac{1}{6} - \frac{1}{2})}{\Gamma(1 - \frac{1}{6})\Gamma(1 - \frac{1}{2})} = \frac{\Gamma(1)\Gamma(\frac{1}{3})}{\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})}$$

by Fact 2.2 for $(a, b, c) = (\frac{1}{6}, \frac{1}{2}, 1)$ together with $c - a - b = \frac{1}{3} > 0$, we have

$$\varphi_1(1) = \omega^2 + \mathbf{i} \cdot \frac{B(\frac{1}{6}, \frac{1}{2})}{\pi} \cdot \frac{\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})}{\Gamma(1)\Gamma(\frac{1}{3})} = \omega^2 + \mathbf{i} \cdot \frac{\Gamma(\frac{1}{6})\Gamma(\frac{1}{2})\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})}{\pi\Gamma(\frac{2}{3})\Gamma(\frac{1}{3})}$$
$$= \omega^2 + \mathbf{i} \cdot \frac{\sin\frac{\pi}{3}}{\sin\frac{\pi}{6}} = \omega^2 + \sqrt{3}\mathbf{i} = \omega,$$

where we use the reflection formula $\Gamma(a)\Gamma(1-a) = \frac{\pi}{\sin(\pi a)}$ for the Gamma function.

We secondly consider the image $\varphi_1(\infty)$. Note that the eigenvalues of N_{∞} are $-\omega^2$ and -1 and that their eigen row vectors are $(\omega^2, 1)$ and $(\omega, 1)$, respectively. Thus $(\omega^2, 1) \cdot R \cdot \mathbf{F}(z)$ and $(\omega, 1) \cdot R \cdot \mathbf{F}(z)$ take forms of

$$\alpha(1/z)^{1/6} f_{\infty,1}(z), \quad \beta(1/z)^{1/2} f_{\infty,2}(z)$$

by Table 2, where α and β are non-zero complex numbers, and $f_{\infty,i}(z)$ (i = 1, 2) are holomorphic functions around $z = \infty$. Thus $R \cdot \mathbf{F}(z)$ admits an expression

$$R \cdot \mathbf{F}(z) = \begin{pmatrix} \omega^2 & 1 \\ \omega & 1 \end{pmatrix}^{-1} \begin{pmatrix} \alpha(1/z)^{1/6} f_{\infty,1}(z) \\ \beta(1/z)^{1/2} f_{\infty,2}(z) \end{pmatrix}$$

$$= \frac{1}{\omega^2 - \omega} \begin{pmatrix} \alpha(1/z)^{1/6} f_{\infty,1}(z) - \beta(1/z)^{1/2} f_{\infty,2}(z) \\ -\omega\alpha(1/z)^{1/6} f_{\infty,1}(z) + \omega^2\beta(1/z)^{1/2} f_{\infty,2}(z) \end{pmatrix},$$
(5.1)

which implies

$$\varphi_1(\infty) = \lim_{z \to \infty} \frac{\alpha (1/z)^{1/6} f_{\infty,1}(z) - \beta (1/z)^{1/2} f_{\infty,2}(z)}{-\omega \alpha (1/z)^{1/6} f_{\infty,1}(z) + \omega^2 \beta (1/z)^{1/2} f_{\infty,2}(z)}$$
$$= \lim_{z \to \infty} \frac{\alpha (1/z)^{1/6} f_{\infty,1}(z)}{-\omega \alpha (1/z)^{1/6} f_{\infty,1}(z)} = -\omega^2.$$

We finally determine the images of $(-\infty, 0)$ and $(1, \infty)$ under φ_1 . Since each of them is a part of a circle or line perpendicular to the real axis, $\varphi_1(-\infty, 0)$ is the line from $-\omega^2$ to $\mathbf{i}\infty$ parallel to the imaginary axis, and $\varphi_1(1, \infty)$ is the arc from ω to $-\omega^2$ with radius 1 and center at the origin.

REMARK 5.3. The image of $Z = \mathbb{C} - \{0, 1\}$ under φ_1 is

$$\mathbb{H}_1 = \mathbb{H} - \{ g \cdot \omega, \ g \cdot (-\omega^2) \in \mathbb{H} \mid g \in \mathrm{P}\Gamma(2)^{1/3} \}.$$

We can extend Schwarz's map φ_1 to a map from $\mathbb{P}^1 - \{0\}$ to \mathbb{H} by corresponding 1 and ∞ to the vertices $\omega = \varphi_1(1)$ and $-\omega^2 = \varphi_1(\infty)$ of Schwarz's triangles or their equivalent points under the action of $\mathrm{P}\Gamma(2)^{1/3}$. Moreover, we can extend it to a map from \mathbb{P}^1 to $\widehat{\mathbb{H}} = \mathbb{H} \cup \{\mathbf{i}\infty\} \cup \mathbb{Q}$. Note that any point in \mathbb{Q} is equivalent to $\mathbf{i}\infty = \varphi_1(0)$ under the action of $\mathrm{P}\Gamma(2)^{1/3}$. In fact, for any irreducible fraction r/s, there exists $r', s' \in \mathbb{Z}$ such that rr' + ss' = 1. Thus $g = \begin{pmatrix} r & -s' \\ s & r' \end{pmatrix}$ belongs to $\mathrm{SL}_2(\mathbb{Z})$ and sends $\mathbf{i}\infty$ to r/s. If this matrix does not belong to $\Gamma(2)^{1/3}$ then $g \cdot T$ belongs to $\Gamma(2)^{1/3}$ and sends $\mathbf{i}\infty$ to r/s, since $T \notin \Gamma(2)^{1/3}$ fixes $\mathbf{i}\infty$ and $[\mathrm{SL}_2(\mathbb{Z}) : \Gamma(2)^{1/3}] = 2$. These extensions of Schwarz's map φ_1 are denoted by the same symbol φ_1 .

REMARK 5.4. To obtain $\varphi_1(\infty)$ in Proof of Proposition 5.2, we cannot use $N'_{\infty} = (N_1 N_0)^{-1} = N_0^{-1} N_1^{-1}$ which is a circuit matrix with respect to $R \cdot \mathbf{F}(z)$ along the loop connecting the end of ρ_0^{-1} and the start of ρ_1^{-1} . This matrix is useful to study the reflection of Schwarz's triangle with respect to the image of (0, 1) under φ_1 .

THEOREM 5.5. The inverse of Schwarz's map φ_1 is given by

$$\nu(\tau) = 3\sqrt{3}\mathbf{i}\frac{\vartheta_{00}(\tau)^4\vartheta_{01}(\tau)^4\vartheta_{10}(\tau)^4}{(\vartheta_{00}(\tau)^4 + \omega\vartheta_{10}(\tau)^4)^3}.$$
(5.2)

PROOF. Schwarz's map φ_1 leads to an isomorphism between Z and $\mathbb{H}_1/\mathrm{P}\Gamma(2)^{1/3}$. Since $\mathrm{P}\Gamma(2)^{1/3}/\mathrm{P}\Gamma(2) \simeq \{I_2, W, W^2\} \simeq A_3$, the space $\mathbb{H}_1/\mathrm{P}\Gamma(2)^{1/3}$ is regarded as the quotient of $\mathbb{H}/\mathrm{P}\Gamma(2)$ by the cyclic group $\{I_2, W, W^2\}$. Here $\mathbb{H}/\mathrm{P}\Gamma(2)$ is isomorphic to Z by

$$\lambda : \mathbb{H}/\mathrm{P}\Gamma(2) \ni \tau \mapsto \lambda(\tau) = \frac{\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4} \in Z.$$

To distinguish the space $\mathbb{C} - \{0, 1\}$ isomorphic to $\mathbb{H}_1/\mathrm{P}\Gamma(2)^{1/3}$ and that to $\mathbb{H}/\mathrm{P}\Gamma(2)$, the former is denoted by Z_{ν} and the latter by Z_{λ} . Note that Z_{λ} is a cyclic triple covering of Z_{ν} . By tracing the action of W via the map λ , we characterize the covering transformation group for the projection $\mathrm{pr}_{\lambda\nu}: Z_{\lambda} \to Z_{\nu}$. The function $\lambda(\tau)$ is transformed into

$$\lambda(W \cdot \tau) = 1 - \frac{1}{\lambda(\tau)}, \quad \lambda(W^2 \cdot \tau) = 1 - \frac{1}{\lambda(W \cdot \tau)} = \frac{1}{1 - \lambda(\tau)}$$
(5.3)

by the actions of W and W^2 on τ as shown in Proof of Lemma 4.2 (3). Thus the covering transformation group is generated by a map

$$\sigma: Z_{\lambda} \ni \lambda \mapsto 1 - \frac{1}{\lambda} \in Z_{\lambda}$$

satisfying

$$\sigma^2(\lambda) = \frac{1}{1-\lambda}, \quad \sigma^3(\lambda) = \lambda.$$

By an invariant function

$$\nu^{\sigma}(\lambda) = \frac{\lambda + \sigma(\lambda) + \sigma^2(\lambda)}{3} = \frac{1}{3}\left(\lambda + (1 - \frac{1}{\lambda}) + \frac{1}{1 - \lambda}\right) = \frac{\lambda^3 - 3\lambda + 1}{3\lambda(\lambda - 1)}$$

under σ , we have a projection

$$\mathrm{pr}_{\nu}^{\sigma}: Z_{\lambda} \ni \lambda \mapsto \nu^{\sigma}(\lambda) \in Z_{\lambda}/\langle \sigma \rangle,$$

where $Z_{\lambda}/\langle \sigma \rangle$ is isomorphic to Z_{ν} . We find an isomorphism $\psi : Z_{\lambda}/\langle \sigma \rangle \to Z_{\nu}$ such that $\psi(\nu^{\sigma}(\lambda(\tau)))$ becomes the inverse of φ_1 . The fixed points in Z_{λ} of σ are $-\omega^2$ and $-\omega$, which are solutions to $\sigma(\lambda) = \lambda$ equivalent to $\lambda^2 - \lambda + 1 = 0$. Since these points are fixed by σ^2 , these points are ramification points of index 3 of $\mathrm{pr}_{\nu} : Z_{\lambda} \to Z_{\nu}$, and the function ν^{σ} sends these points to

$$\nu^{\sigma}(-\omega^2) = -\omega^2, \quad \nu^{\sigma}(-\omega) = -\omega.$$

On the other hand, the preimages of these points under the map $\lambda(\tau) : \mathbb{H}/\mathrm{P}\Gamma(2) \to Z_{\lambda}$ are $-\omega^2$ and ω by Fact 4.1, i.e.

$$\lambda(-\omega^2) = -\omega^2, \quad \lambda(\omega) = -\omega.$$

Hence the map $\nu^{\sigma}(\lambda(\tau))$ sends $\tau = \mathbf{i}\infty, \omega, -\omega^2$ as

$$\nu^{\sigma}(\lambda(\mathbf{i}\infty)) = \infty, \quad \nu^{\sigma}(\lambda(\omega)) = -\omega, \quad \nu^{\sigma}(\lambda(-\omega^2)) = -\omega^2.$$

Here recall that

$$\varphi_1(0) = \mathbf{i}\infty, \quad \varphi_1(1) = \omega, \quad \varphi_1(\infty) = -\omega^2.$$

Since $Z_{\lambda}/\langle \sigma \rangle$ and Z_{ν} are the complex projective line \mathbb{P}^1 minus 3 points, the inverse of φ_1 is give as $\psi(\nu(\lambda(\tau)))$ by the linear fractional transformation ψ sending ∞ , $-\omega$, $-\omega^2$ to 0, 1, ∞ . In fact, ψ is given by

$$\psi(z) = \frac{\omega^2 - \omega}{z + \omega^2}$$

and

$$\psi(\nu^{\sigma}(\lambda)) = -\sqrt{3}\mathbf{i}\frac{3\lambda(\lambda-1)}{\lambda^3 - 3\lambda + 1 + 3\omega^2\lambda(\lambda-1)} = \frac{3\sqrt{3}\mathbf{i}\lambda(1-\lambda)}{(\lambda+\omega^2)^3}$$

By substituting $\lambda = \frac{\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4}$ into the last term and using Jacobi's identity (3.2), we have the expression $\nu(\tau)$ of φ_1^{-1} .

THEOREM 5.6. We have

$$\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4 = F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$$
(5.4)

for any τ in the interior $\widetilde{\mathbb{D}}^{\circ}$ of $\widetilde{\mathbb{D}}$, where $\nu(\tau)$ is given in (5.2). It is extended to an equality on the whole space \mathbb{H} .

18

PROOF. We first show that the both sides of (5.4) behave in the same manner as holomorphic functions on \mathbb{H} under the action of N_0 and N_1 . It is easy to see that the left hand side $\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4$ of (5.4) is invariant under the action of N_0 by Fact 3.2 together with $N_0 = T^2$. Since $N_1 = \omega^2 W$ acts on it projectively, we may use W instead of N_1 . By (3.6) and Jacobi's identity (3.2), we have

$$\vartheta_{00}(W\cdot\tau)^4 + \omega\vartheta_{10}(W\cdot\tau)^4 = -\tau^2\vartheta_{10}(\tau)^4 + \omega\tau^2\vartheta_{01}(\tau)^4 = \omega\tau^2(\vartheta_{00}(\tau)^4 + \omega\vartheta_{10}(\tau)^4).$$

We study the right hand side $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$ of (5.4) under the actions of N_0 and N_1 . Though the function $\nu(\tau)$ is invariant under the actions of N_0 and N_1 , we should study the analytic continuation of $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$ along paths γ_0 and γ_1 from τ to $N_0 \cdot \tau = \tau + 2$ and to $N_1 \cdot \tau = 1 - \frac{1}{\tau}$ since $F(\frac{1}{6}, \frac{1}{2}, 1; z)$ is not single valued. Note that the function ν sends these paths to loops homotopic to ρ_0 and ρ_1 in $Z = \mathbb{C} - \{0, 1\}$. Thus $2\pi F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))$ is invariant under the analytic continuation along γ_0 , and transformed into ω^2 times the first entry of $R \cdot \mathbf{F}(\nu(\tau))$ by the analytic continuation along γ_1 , since $2\pi F(\frac{1}{6}, \frac{1}{2}, 1; z)$ is the second entry of $R \cdot \mathbf{F}(\nu(\tau))$, and the second rows of N_0 and N_1 are (0, 1) and $(\omega^2, 0)$. Recall that τ is the quotient of the first entry of $R \cdot \mathbf{F}(\nu(\tau))$ by the second. Thus $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$ is transformed into

$$\left(\omega^{2}\tau \cdot F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))\right)^{2} = \omega\tau^{2} \cdot F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^{2}$$

by the analytic continuation along γ_1 .

We next study the zeros of the both sides of (5.4). The function $\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4$ appears in the denominator of $\nu(\tau)$. Since $\nu(\tau)$ has a simple zero at $\tau = \mathbf{i}\infty$ and a simple pole at $\tau = -\omega^2$ as a function on $\widehat{\mathbb{H}}/\mathrm{P\Gamma}(2)^{1/3}$, and the factor $\vartheta_{00}(\tau)\vartheta_{01}(\tau)\vartheta_{10}(\tau)$ in its numerator never vanishes on \mathbb{H} , $\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4$ has a simple zero at $\tau = -\omega^2$ as a function on \mathbb{H} , and never vanishes at any point not in the $\mathrm{P\Gamma}(2)^{1/3}$ -orbit of $-\omega^2$. Since $\tau \in \mathbb{H}$ is the quotient of the first entry of $R \cdot \mathbf{F}(z)$ by the second, if its denominator $2\pi F(\frac{1}{6}, \frac{1}{2}, 1; z)$ vanishes at $z_0 \in \mathbb{Z}$, then its numerator also vanishes at this point. By the multi-linearity of the determinant, the Wronskian of $\mathbf{F}(z)$ should has a factor $z - z_0$. However, this contradicts (2.6), which admits the analytic continuation along any path in \mathbb{Z} . Thus if $\tau \in \mathbb{H}$ does not belong to the $\mathrm{P\Gamma}(2)^{1/3}$ -orbit of $-\omega^2$ or the $\mathrm{P\Gamma}(2)^{1/3}$ -orbit of ω , then $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$ does not vanish. Since $\nu(\omega) = 1$ and

$$F(\frac{1}{6}, \frac{1}{2}, 1; 1) = \frac{\Gamma(\frac{1}{3})}{\Gamma(\frac{5}{6})\Gamma(\frac{1}{2})} \neq 0$$

by (2.2), $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\omega))^2$ does not vanish at $\tau = \omega$. Since $\nu(\tau)$ has a pole at $\tau = -\omega^2$ with order 3 as a function on \mathbb{H} , and the behavior of $F(\frac{1}{6}, \frac{1}{2}, 1; z)$ around $z = \infty$ is expressed by Landau's symbol $O((1/z)^{1/6})$ $(z = \infty)$ by (5.1), $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\omega))^2$ has a simple zero at $\tau = -\omega^2$ as a function on \mathbb{H} .

We finally show (5.4). As shown until now, the ratio

$$\frac{F(\frac{1}{6},\frac{1}{2},1;\nu(\tau))^2}{\vartheta_{00}(\tau)^4 + \omega\vartheta_{10}(\tau)^4}$$

is regarded as a holomorphic function on the quotient space $\mathbb{H}/\mathrm{P}\Gamma(2)^{1/3}$, whose one-point compactification by $\mathbf{i}\infty$ is isomorphic to \mathbb{P}^1 . Since

$$\lim_{\tau \to i\infty} (\vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4) = 1, \quad \lim_{\tau \to i\infty} F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2 = F(\frac{1}{6}, \frac{1}{2}, 1; 0)^2 = 1,$$

the ratio becomes a holomorphic function on \mathbb{P}^1 . It should be a constant, and this constant is 1.

REMARK 5.7. Though $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2$ is a single-valued holomorphic function on \mathbb{H} , $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))$ is not. As shown in Proof of Theorem 5.6, any element of the $P\Gamma(2)^{1/3}$ -orbit

$$\{g \cdot (-\omega^2) \in \mathbb{H} \mid g \in \mathrm{P}\Gamma(2)^{1/3}\}$$

of $-\omega^2$ is a simple zero of $F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2 = \vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4$. Note also that the monodromy representation for $E(\frac{1}{6}, \frac{1}{2}, 1)$ includes the scalar multiple of -1.

COROLLARY 5.8. We have

$$\vartheta_{01}(\tau)^{4} - \omega\vartheta_{10}(\tau)^{4} = F(\frac{1}{6}, \frac{1}{2}, 1; -3\sqrt{3}\mathbf{i}\frac{\vartheta_{00}(\tau)^{4}\vartheta_{01}(\tau)^{4}\vartheta_{10}(\tau)^{4}}{(\vartheta_{01}(\tau)^{4} - \omega\vartheta_{10}(\tau)^{4})^{3}})^{2}
= F(\frac{1}{6}, \frac{1}{2}, 1; \frac{\nu(\tau)}{\nu(\tau) - 1})^{2}$$
(5.5)

for any $\tau \in \widetilde{\mathbb{D}}^{\circ}$, where $\nu(\tau)$ is given in (5.2). They are extended to equalities on the whole space \mathbb{H} .

PROOF. We have only to act T on the equality (5.4). By Fact 3.2, the left hand side of (5.4) is transformed into

$$\vartheta_{00}(T\cdot\tau)^4 + \omega\vartheta_{10}(T\cdot\tau)^4 = \vartheta_{01}(\tau)^4 - \omega\vartheta_{10}(T\cdot\tau)^4.$$

The function $\nu(\tau)$ in the right hand side of (5.4) is transformed into

$$\begin{split} \nu(T \cdot \tau) &= 3\sqrt{3} \mathbf{i} \frac{\vartheta_{00}(T \cdot \tau)^4 \vartheta_{01}(T \cdot \tau)^4 \vartheta_{10}(T \cdot \tau)^4}{(\vartheta_{00}(T \cdot \tau)^4 + \omega \vartheta_{10}(T \cdot \tau)^4)^3} = 3\sqrt{3} \mathbf{i} \frac{-\vartheta_{00}(\tau)^4 \vartheta_{01}(\tau)^4 \vartheta_{10}(\tau)^4}{(\vartheta_{01}(\tau)^4 - \omega \vartheta_{10}(\tau)^4)^3}.\\ \text{Since } \lambda(T \cdot \tau) &= \frac{\lambda(\tau)}{\lambda(\tau) - 1},\\ \nu(T \cdot \tau) &= \frac{3\sqrt{3}\mathbf{i}\lambda(T \cdot \tau)(1 - \lambda(T \cdot \tau))}{(\lambda(T \cdot \tau) + \omega^2)^3} = \frac{3\sqrt{3}\mathbf{i}\frac{\lambda(\tau)}{\lambda(\tau) - 1}(1 - \frac{\lambda(\tau)}{\lambda(\tau) - 1})}{(\frac{\lambda(\tau)}{\lambda(\tau) - 1} + \omega^2)^3} = \frac{3\sqrt{3}\mathbf{i}\lambda(\tau)(1 - \lambda(\tau))}{(-\omega\lambda(\tau) - 1)^3},\\ \frac{\nu(\tau)}{\nu(\tau) - 1} &= \frac{3\sqrt{3}\mathbf{i}\lambda(\tau)(1 - \lambda(\tau))}{(\lambda(\tau) + \omega^2)^3} \cdot \frac{(\lambda(\tau) + \omega^2)^3}{3\sqrt{3}\mathbf{i}\lambda(\tau)(1 - \lambda(\tau)) - (\lambda(\tau) + \omega^2)^3} \\ &= \frac{3\sqrt{3}\mathbf{i}\lambda(\tau)(1 - \lambda(\tau))}{-\lambda(\tau)^3 - 3(\omega^2 + \sqrt{3}\mathbf{i})\lambda(\tau)^2 - 3(\omega - \sqrt{3}\mathbf{i})\lambda(\tau) - 1} = \frac{3\sqrt{3}\mathbf{i}\lambda(\tau)(1 - \lambda(\tau))}{-(\lambda(\tau) + \omega)^3}, \end{split}$$

we have $\nu(T \cdot \tau) = \frac{\nu(\tau)}{\nu(\tau) - 1}$.

20

REMARK 5.9. By Jacobi's identity (3.2), we have

$$\begin{split} \vartheta_{00}(\tau)^4 + \omega \vartheta_{10}(\tau)^4 &= \vartheta_{01}(\tau)^4 - \omega^2 \vartheta_{10}(\tau)^4 = -\omega^2 \vartheta_{00}(\tau)^4 - \omega \vartheta_{01}(\tau)^4 \\ &= \frac{1 - \omega^2}{3} \left(\vartheta_{00}(\tau)^4 - \omega \vartheta_{01}(\tau)^4 - \omega^2 \vartheta_{10}(\tau)^4 \right), \\ \vartheta_{00}(\tau)^4 + \omega^2 \vartheta_{10}(\tau)^4 &= \vartheta_{01}(\tau)^4 - \omega \vartheta_{10}(\tau)^4 = -\omega \vartheta_{00}(\tau)^4 - \omega^2 \vartheta_{01}(\tau)^4 \\ &= \frac{1 - \omega}{3} \left(\vartheta_{00}(\tau)^4 - \omega^2 \vartheta_{01}(\tau)^4 - \omega \vartheta_{10}(\tau)^4 \right), \end{split}$$

$$E_4(\tau) = \frac{\vartheta_{00}(\tau)^8 + \vartheta_{10}(\tau)^8 + \vartheta_{01}(\tau)^8}{2} = \vartheta_{00}(\tau)^8 - \vartheta_{00}(\tau)^4 \vartheta_{10}(\tau)^4 + \vartheta_{10}(\tau)^8$$
$$= \vartheta_{00}(\tau)^8 - \vartheta_{00}(\tau)^4 \vartheta_{01}(\tau)^4 + \vartheta_{01}(\tau)^8 = \vartheta_{01}(\tau)^8 + \vartheta_{01}(\tau)^4 \vartheta_{10}(\tau)^4 + \vartheta_{10}(\tau)^8.$$

COROLLARY 5.10. For any $\tau \in \widetilde{\mathbb{D}}^{\circ}$, we have

$$E_4(\tau) = \frac{\vartheta_{00}(\tau)^8 + \vartheta_{01}(\tau)^8 + \vartheta_{10}(\tau)^8}{2} = F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2 F(\frac{1}{6}, \frac{1}{2}, 1; \frac{\nu(\tau)}{\nu(\tau) - 1})^2, \quad (5.6)$$

which is extended to an equality on the whole space \mathbb{H} .

PROOF. We have

$$F(\frac{1}{6}, \frac{1}{2}, 1; \nu(\tau))^2 F(\frac{1}{6}, \frac{1}{2}, 1; \frac{\nu(\tau)}{\nu(\tau) - 1})^2 = (\vartheta_{00}(\tau)^4 + \omega\vartheta_{10}(\tau)^4)(\vartheta_{00}(\tau)^4 + \omega^2\vartheta_{10}(\tau)^4)$$
$$= \vartheta_{00}(\tau)^8 - \vartheta_{00}(\tau)^4 \vartheta_{10}(\tau)^4 + \vartheta_{10}(\tau)^8 = \frac{\vartheta_{00}(\tau)^8 + \vartheta_{10}(\tau)^8 + \vartheta_{01}(\tau)^8}{2} = E_4(\tau)$$

by Theorem 5.6, Corollary 5.8 and Remark 5.9.

By pulling back identities in Theorem 5.6 and Corollaries 5.8 and 5.10 under Schwarz's map $\varphi_1 : Z \to \mathbb{H}_1$, we have the following.

COROLLARY 5.11. We have

$$F(\frac{1}{6}, \frac{1}{2}, 1; z)^{2} = \vartheta_{00}(\varphi_{1}(z))^{4} + \omega\vartheta_{10}(\varphi_{1}(z))^{4},$$

$$F(\frac{1}{6}, \frac{1}{2}, 1; \frac{z}{z-1})^{2} = \vartheta_{00}(\varphi_{1}(z))^{4} + \omega^{2}\vartheta_{10}(\varphi_{1}(z))^{4},$$

$$F(\frac{1}{6}, \frac{1}{2}, 1; z)^{2}F(\frac{1}{6}, \frac{1}{2}, 1; \frac{z}{z-1})^{2} = \frac{\vartheta_{00}(\varphi_{1}(z))^{8} + \vartheta_{01}(\varphi_{1}(z))^{8} + \vartheta_{10}(\varphi_{1}(z))^{8}}{2} = E_{4}(\varphi_{1}(z)),$$
(5.7)

which are initially equalities on $\mathbb{B}_0 \cap \mathbb{B}_1$ and extended to those on the whole space Z by the analytic continuation.

6. Schwarz's map for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$

We consider the hypergeometric differential equation $E(\frac{1}{12}, \frac{5}{12}, 1)$. Its Riemann's scheme becomes as in Table 3. The basis of the space of its local solutions around $\dot{z} = \frac{1}{2}$ in Lemma

z_0	0	1	∞
$e_{z_0,1}$	0	0	1/12
$e_{z_{0},2}$	0	1/2	5/12
$e_{z_0,2} - e_{z_0,1}$	$1/\infty$	1/2	1/3

TABLE 3. Riemann's scheme for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$

2.3 becomes

$$\begin{pmatrix} \exp(\frac{11\pi \mathbf{i}}{12}) \int_{-\infty}^{0} (-t)^{-7/12} (z-t)^{-5/12} (1-t)^{-1/12} dt \\ \int_{1}^{\infty} (t)^{-7/12} (t-z)^{-5/12} (t-1)^{-1/12} dt \end{pmatrix}$$

$$= \begin{pmatrix} \exp(\frac{11\pi \mathbf{i}}{12}) B(\frac{1}{12}, \frac{5}{12}) F(\frac{1}{12}, \frac{5}{12}, \frac{1}{2}, 1-z) \\ B(\frac{1}{12}, \frac{11}{12}) F(\frac{1}{12}, \frac{5}{12}, 1, z) \end{pmatrix}.$$

The circuit matrices with respect to this basis are

$$M_0 = \begin{pmatrix} 1 & 1 - e^{-\pi \mathbf{i}/6} \\ 0 & 1 \end{pmatrix}, \quad M_1 = \begin{pmatrix} 1 & 0 \\ e^{7\pi \mathbf{i}/6} - 1 & -1 \end{pmatrix}.$$

As in the previous section, we can find a matrix $R = \begin{pmatrix} -\mathbf{i} - \omega & -\mathbf{i} \\ 0 & 1 \end{pmatrix}$, which satisfies

$$N_0 = RM_0R^{-1} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} = T, \quad N_1 = RM_0R^{-1} = \mathbf{i} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \mathbf{i}J$$
$$N_\infty = (N_0N_1)^{-1} = \mathbf{i} \begin{pmatrix} 0 & 1 \\ -1 & 1 \end{pmatrix}.$$

Since $N_{\infty}^3 = \mathbf{i}I_2$ and $\mathrm{SL}_2(\mathbb{Z})$ is generated by T and J, the group generated by N_0 and N_1 is

$$\operatorname{SL}_2(\mathbb{Z})\langle \mathbf{i}I_2 \rangle = \{ \mathbf{i}^k g \in \operatorname{GL}_2(\mathbb{Z}[\mathbf{i}]) \mid g \in \operatorname{SL}_2(\mathbb{Z}), k \in \mathbb{Z} \},\$$

whose projectivization is isomorphic to $PSL_2(\mathbb{Z})$.

DEFINITION 6.1 (Schwarz's map φ_2 for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$). We define Schwarz's map φ_2 for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$ by the analytic continuation of the map

$$\varphi_2 : z \mapsto \tau = \tau(z) = \mathbf{i} \cdot \frac{B\left(\frac{1}{12}, \frac{5}{12}\right)}{2\pi} \cdot \frac{F(\frac{1}{12}, \frac{5}{12}, \frac{1}{2}; 1-z)}{F(\frac{1}{12}, \frac{5}{12}, 1; z)} - \mathbf{i}.$$
(6.1)

defined on $\mathbb{B}_0 \cap \mathbb{B}_1$ to $\mathbb{C} - \{0, 1\}$. Its projective monodromy representation is isomorphic to $\mathrm{PSL}_2(\mathbb{Z})$.

REMARK 6.2. Schwarz's map for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$ is studied in [Sh, Proposition 6.4.1] by a different basis of the space of local solutions from ours.

PROPOSITION 6.3. Schwarz's triangle $\varphi_2(\mathbb{H})$ for $(a, b, c) = (\frac{1}{12}, \frac{5}{12}, 1)$ is

$$\{\tau \in \mathbb{H} \mid 0 < \operatorname{Re}(\tau) < \frac{1}{2}, \ |\tau| > 1\},\$$

which is the interior of the right half of the fundamental region \mathbb{D} for $\mathrm{PSL}_2(\mathbb{Z})$, and its vertices $\varphi_2(z_0) = \lim_{z \to z_0, z \in \mathbb{H}} \varphi_2(z)$ $(z_0 = 0, 1, \infty)$ are

$$\varphi_2(0) = \mathbf{i}\infty, \quad \varphi_2(1) = \mathbf{i}, \quad \varphi_2(\infty) = -\omega^2.$$
 (6.2)

See Figure 2 for their shape.

PROOF. Fact 2.2 together with c-a-b=0 for $(a,b,c) = (\frac{1}{12}, \frac{5}{12}, \frac{1}{2})$ yields $\varphi_2(0) = \mathbf{i}\infty$. It also yields that

$$\varphi_{2}(1) = \lim_{z \to 1, z \in (0,1)} \left(\mathbf{i} \cdot \frac{B\left(\frac{1}{12}, \frac{5}{12}\right)}{2\pi} \cdot \frac{F\left(\frac{1}{12}, \frac{5}{12}; \frac{1}{2}; 1-z\right)}{F\left(\frac{1}{12}, \frac{5}{12}; 1; z\right)} - \mathbf{i} \right)$$

= $\mathbf{i} \cdot \frac{\Gamma\left(\frac{1}{12}\right) \Gamma\left(\frac{5}{12}\right)}{2\pi\Gamma\left(\frac{1}{2}\right)} \cdot \frac{\Gamma\left(\frac{11}{12}\right) \Gamma\left(\frac{7}{12}\right)}{\Gamma(1)\Gamma\left(\frac{1}{2}\right)} - \mathbf{i} = \frac{\mathbf{i} \cdot \pi^{2}}{2\pi^{2} \cdot \sin\left(\frac{\pi}{12}\right) \sin\left(\frac{5\pi}{12}\right)} - \mathbf{i} = \mathbf{i}.$

Note that the eigenvalues of N_{∞} are $e^{\pi \mathbf{i}/6} = \omega/\mathbf{i}$ and $e^{5\pi \mathbf{i}/6} = \omega^2/\mathbf{i}$ and that their eigen row vectors are $(1, \omega)$ and $(1, \omega^2)$, respectively. We can show $\varphi_2(\infty) = -\omega^2$ quite similarly to $\varphi_1(\infty) = -\omega^2$ in Proof of Proposition 5.2. Since each edge of Schwarz's triangle $\varphi_2(\mathbb{H})$ is parallel to the imaginary axis or in a circle with center in the real axis, and bounded by two of $\varphi_2(0) = \mathbf{i}\infty$, $\varphi_2(1) = \mathbf{i}$, $\varphi_2(\infty) = -\omega^2$, the edges are contained in the lines $\operatorname{Re}(\tau) = 0$ or $\operatorname{Re}(\tau) = \frac{1}{2}$, or the circle $|\tau| = 1$.

REMARK 6.4. The image of
$$Z = \mathbb{C} - \{0, 1\}$$
 under φ_2 is

$$\mathbb{H}_2 = \mathbb{H} - \{g \cdot \mathbf{i}, \ g \cdot (-\omega^2) \in \mathbb{H} \mid g \in \mathrm{PSL}_2(\mathbb{Z})\}$$

We can extend Schwarz's map φ_2 to a map from $\mathbb{P}^1 - \{0\}$ to \mathbb{H} by corresponding 1 and ∞ to the vertices $\mathbf{i} = \varphi_2(1)$ and $-\omega^2 = \varphi_2(\infty)$ of Schwarz's triangles or their equivalent points under the action of $\mathrm{PSL}_2(\mathbb{Z})$. Moreover, we can extend it to a map from \mathbb{P}^1 to $\widehat{\mathbb{H}} = \mathbb{H} \cup \{\mathbf{i}\infty\} \cup \mathbb{Q}$. Note that any point in \mathbb{Q} is equivalent to $\mathbf{i}\infty = \varphi_2(0)$ under the action of $\mathrm{PSL}_2(\mathbb{Z})$. These extensions of Schwarz's map φ_2 are denoted by the same symbol φ_2 .

THEOREM 6.5. The inverse of Schwarz's map φ_2 is given by

$$\mathbb{H}_2 \ni \tau \mapsto z = \frac{1}{j(\tau)} \in \mathbb{C} - \{0, 1\},\tag{6.3}$$

where the *j*-invariant is defined by

$$j(\tau) = \frac{4}{27} \frac{(\lambda(\tau)^2 - \lambda(\tau) + 1)^3}{\lambda(\tau)^2 (1 - \lambda(\tau))^2} = \frac{1}{54} \frac{(\vartheta_{00}(\tau)^8 + \vartheta_{01}(\tau)^8 + \vartheta_{10}(\tau)^8)^3}{\vartheta_{00}(\tau)^8 \vartheta_{01}(\tau)^8 \vartheta_{10}(\tau)^8}.$$
 (6.4)

REMARK 6.6. We refer to [Yo, p.42] for our definition of $j(\tau)$. As in [Se, (23) in p. 90], $1728j(\tau)$ admits a Fourier expansion with integral coefficients:

$$1728j(\tau) = \frac{1}{q} + 744 + 196884q + \cdots,$$

where $q = \exp(2\pi \mathbf{i}\tau)$.

PROOF. Schwarz's map φ_2 leads to an isomorphism between Z and $\mathbb{H}_2/\mathrm{PSL}_2(\mathbb{Z})$, which is extended to that between \mathbb{P}^1 and $\widehat{\mathbb{H}}_2/\mathrm{PSL}_2(\mathbb{Z})$. Since $\mathrm{PSL}_2(\mathbb{Z})/\mathrm{P\Gamma}(2)$ is isomorphic to S_3 , the space $\mathbb{H}_2/\mathrm{PSL}_2(\mathbb{Z})$ is regarded as the quotient of $\mathbb{H}/\mathrm{P\Gamma}(2)$ by the symmetric group

 S_3 . To distinguish the space $\mathbb{C} - \{0, 1\}$ isomorphic to $\mathbb{H}_2/\mathrm{PSL}_2(\mathbb{Z})$ and that to $\mathbb{H}/\mathrm{P\Gamma}(2)$, the former is denoted by $Z_{1/j}$ and the latter by Z_{λ} . The space Z_{λ} is regarded as a 6-fold covering of $Z_{1/j}$. Thus we have a diagram

$$\mathbb{H}/\mathrm{P}\Gamma(2) \xrightarrow{\lambda} Z_{\lambda}$$

$$\downarrow \qquad \qquad \downarrow \frac{1}{\jmath}$$

$$\mathbb{H}/\mathrm{PSL}_{2}(\mathbb{Z}) \xrightarrow{\sim} Z_{1/\jmath}.$$

Since the right hand side of (6.4) is invariant under the action of $SL_2(\mathbb{Z})$ by Fact 3.2, the function $\frac{1}{j(\tau)}$ on $\mathbb{H}/\mathrm{P}\Gamma(2)$ descends to a function on $\mathbb{H}/\mathrm{P}\mathrm{SL}_2(\mathbb{Z})$. We decompose the function $\frac{1}{j(\tau)}$ into the λ -function and a rational function

$$\frac{1}{j}: Z_{\lambda} \ni \lambda \mapsto \frac{27}{4} \frac{\lambda^2 (1-\lambda)^2}{(\lambda^2 - \lambda + 1)^3} \in Z_{1/j}$$

of degree 6. This function is regarded as a branched covering map from \mathbb{P}^1 to \mathbb{P}^1 of degree 6. Since the natural projection from $\mathbb{H}/\mathrm{PF}(2)$ to $\mathbb{H}/\mathrm{PSL}_2(\mathbb{Z})$ is of degree 6, it turns out that the function $\frac{1}{j(\tau)}$ on $\mathbb{H}/\mathrm{PSL}_2(\mathbb{Z})$ is of degree 1. We can easily see that the function $\frac{1}{j}$ maps $\lambda = 0, 1, \infty$ to 0 and $\lambda = -\omega, -\omega^2$ to ∞ . By solving an equation

$$\frac{27}{4} \frac{\lambda^2 (1-\lambda)^2}{(\lambda^2 - \lambda + 1)^3} = 1,$$

we also see that $\frac{1}{j}$ maps $\lambda = -1, \frac{1}{2}, 2$ to 1. Recall that the function λ sends $\tau = \mathbf{i}\infty, \mathbf{i}, -\omega^2$ to $0, \frac{1}{2}, -\omega^2$ as in Fact 4.1(1) and Lemma 4.2 (2). Since $\frac{1}{j(\tau)}$ is extended to a map $\widehat{\mathbb{H}}/\mathrm{PSL}_2(\mathbb{Z}) \to \mathbb{P}^1$ of degree 1 with sending $\tau = \mathbf{i}\infty, \mathbf{i}, -\omega^2$ to $0, 1, \infty$, it coincides with the inverse of φ_2 by (6.2).

THEOREM 6.7. We have

$$E_4(\tau) = \frac{\vartheta_{00}(\tau)^8 + \vartheta_{01}(\tau)^8 + \vartheta_{10}(\tau)^8}{2} = F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})^4$$
(6.5)

for any τ in the interior \mathbb{D}° of \mathbb{D} , where the *j*-invariant $j(\tau)$ is given in (6.4). It is extended to an equality on the whole space \mathbb{H} .

PROOF. We can show this theorem quite similarly to Proof of Theorem 5.6. \Box

REMARK 6.8. Though $F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})^4$ is a single-valued holomorphic function on \mathbb{H} , neither $F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})$ nor $F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})^2$ is so. We can see that any element of the $PSL_2(\mathbb{Z})$ -orbit

$$\{g \cdot (-\omega^2) \in \mathbb{H} \mid g \in \mathrm{PSL}_2(\mathbb{Z})\}$$

of $-\omega^2$ is a simple zero of $F(\frac{1}{12}, \frac{5}{12}, 1; \nu(\tau))^4 = E_4(\tau)$. Note also that the monodromy representation for $E(\frac{1}{12}, \frac{5}{12}, 1)$ includes the scalar multiple of **i**. Since their restrictions to $\{\tau \in \mathbb{H} \mid \text{Im}(\tau) > 1\}$ are single valued and have a period 1, each of them admits a Fourier expansion:

$$F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)}) = 1 + 60e^{2\pi i\tau} - 4860(e^{2\pi i\tau})^2 + 660480(e^{2\pi i\tau})^3 - \cdots,$$

$$F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})^2 = 1 + 120e^{2\pi i\tau} - 6120(e^{2\pi i\tau})^2 + 737760(e^{2\pi i\tau})^3 - \cdots;$$

see (3.7) for the Fourier expansion of $F(\frac{1}{12}, \frac{5}{12}, 1; \frac{1}{j(\tau)})^4 = E_4(\tau)$.

By pulling back equalities in Theorem 6.7 under Schwarz's map φ_2 in Definition 6.1, we have the following.

COROLLARY 6.9. We have

$$F(\frac{1}{12}, \frac{5}{12}, 1; z)^4 = \frac{\vartheta_{00}(\varphi_2(z))^8 + \vartheta_{01}(\varphi_2(z))^8 + \vartheta_{10}(\varphi_2(z))^8}{2} = E_4(\varphi_2(z)), \quad (6.6)$$

which is initially an equality on $\mathbb{B}_0 \cap \mathbb{B}_1$ and extended to that on the whole space Z by the analytic continuation.

7. Functional equations for F(a, b, c; z)

COROLLARY 7.1. The hypergeometric series F(a, b, c; z) satisfies functional equations:

$$F\left(\frac{1}{6}, \frac{1}{2}, 1; \frac{3\sqrt{3}iz(1-z)}{(1+\omega z)^3}\right)^2 = -\omega^2 F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)^2 - \omega F\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1}\right)^2,$$

$$F\left(\frac{1}{6}, \frac{1}{2}, 1; -\frac{3\sqrt{3}iz(1-z)}{(z+\omega)^3}\right)^2 = -\omega F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)^2 - \omega^2 F\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1}\right)^2,$$

$$F\left(\frac{1}{12}, \frac{5}{12}, 1; \frac{z^2}{4(z-1)}\right)^2 = F\left(\frac{1}{6}, \frac{1}{2}, 1; z\right) F\left(\frac{1}{6}, \frac{1}{2}, 1; \frac{z}{z-1}\right),$$

$$F\left(\frac{1}{12}, \frac{5}{12}, 1; \frac{27z^2(1-z)^2}{4(z^2-z+1)^3}\right)^4 = F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)^4 + F\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1}\right)^4$$

$$-F\left(\frac{1}{2}, \frac{1}{2}, 1; z\right)^2 F\left(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1}\right)^2,$$

$$(7.1)$$

where z is in a small neighborhood of z = 0.

PROOF. We set $z = \lambda(\tau) = \frac{\vartheta_{10}(\tau)^4}{\vartheta_{00}(\tau)^4}$. Then Theorem 5.6 and Fact 4.3 yield that

$$F(\frac{1}{6}, \frac{1}{2}, 1; \frac{3\sqrt{3l^2(1-z)}}{(1+\omega z)^3})^2 = \vartheta_{00}(\tau)^4 + \omega\vartheta_{10}(\tau)^4 = -\omega^2\vartheta_{00}(\tau)^4 - \omega\vartheta_{01}(\tau)^4$$
$$= -\omega^2 F(\frac{1}{2}, \frac{1}{2}, 1; z)^2 - \omega F(\frac{1}{2}, \frac{1}{2}, 1; \frac{z}{z-1})^2.$$

We obtain the second equality by acting $T \in SL_2(\mathbb{Z})$ on τ in the first equality as in Proof of Corollary 5.8.

Theorem 6.7, Fact 4.3, Remark 5.9 and the equality $j(\tau) = \frac{4}{27} \cdot \frac{(\lambda(\tau)^2 - \lambda(\tau) + 1)^3}{\lambda(\tau)^2(1 - \lambda(\tau))^2}$ yield the last equality.

To obtain the third equality, we set $z = \nu(\tau) = \frac{3\sqrt{3}i\lambda(\tau)(1-\lambda(\tau))}{(1+\omega\lambda(\tau))^3}$. Then we have

$$F(\frac{1}{6}, \frac{1}{2}, 1; z)F(\frac{1}{6}, \frac{1}{2}, 1; \frac{z}{z-1}) = \frac{\vartheta_{00}(\tau)^8 + \vartheta_{01}(\tau)^8 + \vartheta_{10}(\tau)^8}{2} = F(\frac{1}{12}, \frac{5}{12}, 1; \frac{z^2}{4(z-1)})^4$$

by Corollary 5.10, Theorem 6.7 and the equality $j(\tau) = 4 \cdot \frac{\nu(\tau) - 1}{\nu(\tau)^2}$.

References

- [BB1] Borwein J.M. and Borwein P.B., Pi and the AGM, John Wiley & Sons, Inc., New York, 1998.
- [BB2] Borwein J.M. and Borwein P.B., A cubic counterpart of Jacobi's identity and the AGM, Trans. Amer. Math. Soc. 323 (1991), 691–701.
- [Er] Erdélyi A., Magnus W., Oberhettinger F., and Tricomi F.G., Higher transcendental functions, Vol. I, McGraw-Hill, 1953.
- [CM] Chiba J. and Matsumoto K., An analogy of Jacobi's formula and its applications, preprint, 2021.
- [Mu] Mumford D., Tata Lectures on Theta I, Birkhäuser, Boston, 2007.
- [Se] Serre, J.-P., A course in arithmetic, Graduate Texts in Mathematics 7, Springer-Verlag, New York-Heidelberg, 1973.
- [Sh] Shiga, H., Modular functions (in Japanese), Kyoritsu Shuppan, Tokyo, 2017.
- [Va] Varchenko, A. N., The Euler beta-function, the Vandermonde determinant, the Legendre equation, and critical values of linear functions on a configuration of hyperplanes, I, II. Math. USSR-Izv. 35 (1990), 543–571, 36 (1991), 155–167.
- [Yo] Yoshida, M., Hypergeometric functions, my love, Friedr. Vieweg & Sohn, Braunschweig, 1997.

(Matsumoto) DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE HOKKAIDO UNIVERSITY, SAP-PORO 060-0810, JAPAN