

THE GROUP INVERSE OF THE SUM IN A BANACH ALGEBRA

HUANYIN CHEN AND MARJAN SHEIBANI*

ABSTRACT. In this paper, we present new necessary and sufficient conditions under which the sum of two group invertible elements in a Banach algebra has group inverse. We then apply these results to block operator matrices over Banach spaces. The group inverses of certain operator block matrices are thereby obtained. Additionally, this paper extends the results obtained in [5].

1. INTRODUCTION

Let \mathcal{A} be a complex Banach algebra with an identity 1. An element a in a Banach algebra \mathcal{A} has group inverse provided that there exists $b \in \mathcal{A}$ such that $ab = ba, b = bab$ and $a = aba$. Such b is unique if exists, denoted by $a^\#$, and called the group inverse of a . As is well known, a square complex matrix A has group inverse if and only if $\text{rank}(A) = \text{rank}(A^2)$.

The group invertibility in a Banach algebra is attractive. Many authors have studied such problems from many different views, e.g., [1, 5, 6, 7, 10]. It was also extensively investigated under the concept "strongly regularity" in ring theory.

In Section 2, we present necessary and sufficient conditions under which the sum of two group invertible elements in a Banach algebra. Let $a, b \in \mathcal{A}^\#$. If $ab = \lambda ba$ for some $\lambda \in \mathbb{C}$, then $a + b \in \mathcal{A}^\#$ if and only if $abb^\# + baa^\# \in \mathcal{A}^\#$. In this case, $(a + b)^\# = (abb^\# + baa^\#)^\# + a^\#b^\# + b^\#a^\#$.

2010 *Mathematics Subject Classification.* 15A09, 47A11, 47A53.

Key words and phrases. group inverse; operator matrix; Banach algebra; Banach space.

*Corresponding author.

Let X and Y be Banach spaces. In Section 3, we investigate the group inverse of a 2×2 operator matrix

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \quad (*)$$

where $A \in \mathcal{L}(X)$, $D \in \mathcal{L}(Y)$. Here, M is a bounded linear operator on $X \oplus Y$. This problem is quite complicated and was extensively studied by many authors. Let A, D, BC have group inverses. If $AB = 0, BD = 0, B(CB)^\pi = 0, C(BC)^\pi = 0$ and $DC = \lambda CA$, we prove that M has group inverse. The explicit formula of the group inverse of M is also given.

If $a \in \mathcal{A}$ has the group inverse a^d , the element $a^\pi = 1 - aa^d$ is called the spectral idempotent of a . Let $p \in \mathcal{A}$ be an idempotent, and let $x \in \mathcal{A}$. Then we write

$$x = pxp + px(1-p) + (1-p)xp + (1-p)x(1-p),$$

and induce a Pierce representation given by the matrix

$$x = \begin{pmatrix} pxp & px(1-p) \\ (1-p)xp & (1-p)x(1-p) \end{pmatrix}_p.$$

Throughout the paper, all Banach algebras are complex with an identity. We use $\mathcal{A}^\#$ to denote the set of all group invertible elements in \mathcal{A} . λ stands for a nonzero complex numbers.

2. ADDITIVE PROPERTIES

The purpose of this section is to establish the new formulas for the group of the sum $a + b$ which will be used in the sequel. We begin with

Lemma 2.1. *Let \mathcal{A} be a Banach algebra, and let $a, b \in \mathcal{A}^\#$. If $ab = \lambda ba$, then the following hold:*

- (1) $abb^\# = bb^\#a$.
- (2) $baa^\# = aa^\#b$.

Proof. (1) As b has group inverse, $b^\#$ coincides with its Drazin inverse, and so it follows by [2, Theorem 2.3] that $ab^\# = \frac{1}{\lambda}b^\#a$.

Therefore

$$\begin{aligned} abb^\# &= \lambda(ba)b^\# \\ &= \lambda b\left(\frac{1}{\lambda}b^\#a\right) \\ &= bb^\#a. \end{aligned}$$

(2) This is similar to (1). □

Theorem 2.2. *Let $a, b \in \mathcal{A}^\#$. If $ab = \lambda ba$, then the following are equivalent:*

- (1) $a + b \in \mathcal{A}^\#$.
- (2) $a(1 + a^\#b) \in \mathcal{A}^\#$.
- (3) $abb^\# + baa^\# \in \mathcal{A}^\#$.

In this case,

$$\begin{aligned} (a + b)^\# &= a^\#b^\pi + [a(1 + a^\#b)]^\# \\ &= a^\#b^\pi + a^\pi b^\# + [abb^\# + baa^\#]^\#. \end{aligned}$$

Proof. (1) \Leftrightarrow (2) Let $p = aa^\#$. Then we have

$$a = \begin{pmatrix} a_1 & 0 \\ 0 & 0 \end{pmatrix}_p, b = \begin{pmatrix} b_1 & b_{12} \\ b_{21} & b_2 \end{pmatrix}_p.$$

Since $ab = \lambda ba$, we get $b_{12} = b_{21} = 0$. Thus,

$$a = \begin{pmatrix} a_1 & 0 \\ 0 & 0 \end{pmatrix}_p, b = \begin{pmatrix} b_1 & 0 \\ 0 & b_2 \end{pmatrix}_p,$$

Clearly, $a_1 = aa^\#aaa^\#, b_1 = aa^\#baa^\#$ and $b_2 = a^\pi ba^\pi$. Hence,

$$a + b = \begin{pmatrix} aa^\#(a + b)aa^\# & 0 \\ 0 & a^\pi ba^\pi \end{pmatrix}.$$

In view of Lemma 2.1, $a^\pi b = ba^\pi$. Then we easily check that $a^\pi ba^\pi$ has group inverse. Therefore $a + b$ has group inverse if and only if $aa^\#(a + b)aa^\#$ has group inverse. In this case,

$$(a + b)^\# = a^\#b^\pi + [a(1 + a^\#b)]^\#.$$

(2) \Leftrightarrow (3) Write $aa^\#(a + b)aa^\# = a + aa^\#b := a_1 + b_1$. Choose $q = b_1b_1^\#$. Then Then we have

$$a_1 = \begin{pmatrix} a'_1 & 0 \\ 0 & a'_2 \end{pmatrix}_p, b_1 = \begin{pmatrix} b'_1 & 0 \\ 0 & 0 \end{pmatrix}_q,$$

where $a_1 = b_1 b_1^\# a b_1 b_1^\#$, $a'_2 = b_1^\pi a b_1^\pi$. Hence,

$$a_1 + b_1 = \begin{pmatrix} a a^\# (a + b) b b^\# & 0 \\ 0 & b_1^\pi a b_1^\pi \end{pmatrix}.$$

In view of Lemma 2.1, $b_1^\pi a b_1^\pi$ has group inverse. Therefore $a_1 + b_1$ has group inverse if and only if $a a^\# (a + b) b b^\#$ has group inverse. In this case,

$$\begin{aligned} (a_1 + b_1)^\# &= a^\# b^\pi + [a a^\# (a + b) b b^\#]^\# \\ &= a^\# b^\pi + [a a^\# (a + b) b b^\#]^\#. \end{aligned}$$

Combing the preceding formulas, we derive that

$$(a + b)^\# = a^\# b^\pi + a^\pi b^\# + [a b b^\# + b a a^\#]^\#,$$

as asserted. \square

Corollary 2.3. *Let \mathcal{A} be a Banach algebra, and let $a, b \in \mathcal{A}^\#$. If $ab = \lambda ba$, then the following are equivalent:*

- (1) $a - b \in \mathcal{A}^\#$.
- (2) $a(1 - a^\# b) \in \mathcal{A}^\#$.

In this case,

$$(a - b)^\# = a^\# b^\pi - a^\pi b^\# + [a(1 - a^\# b)]^\#.$$

Proof. Clearly, $-b \in \mathcal{A}^\#$. Applying Theorem 2.2 to a and $-b$, we obtain the result. \square

Corollary 2.4. *Let $a, b \in \mathcal{A}^\#$. If $ab = ba$, then the following are equivalent:*

- (1) $a + b \in \mathcal{A}^\#$.
- (2) $1 + a^\# b \in \mathcal{A}^\#$.

In this case,

$$(a + b)^\# = a^\# (1 + a^\# b)^\# b b^\# + a^\# b^\pi + b^\# a^\pi.$$

Proof. (1) \Rightarrow (2) We check that

$$1 + a^\# b = a^\pi + a^\# (a + b).$$

Since $a^\pi a^\# (a + b) = 0 = a^\# (a + b) a^\pi$, it follows by [1, Theorem 2.1] that $1 + a^\# b \in \mathcal{A}^\#$.

(2) \Rightarrow (1) In view of Lemma 2.1, we see that

$$abb^\# + baa^\# = abb^\#(1 + a^\#b).$$

Clearly, $abb^\# \in \mathcal{A}^\#$. Since $abb^\#, 1 + a^\#b \in \mathcal{A}^\#$ and $abb^\#(1 + a^\#b) = (1 + a^\#b)abb^\#$, it follows that $abb^\#(1 + a^\#b) \in \mathcal{A}^\#$. Hence $abb^\# + baa^\# \in \mathcal{A}^\#$. Moreover,

$$(abb^\# + baa^\#)^\# = [abb^\#(1 + a^\#b)]^\# = a^\#(1 + a^\#b)^\#bb^\#.$$

According to Theorem 2.2, we complete the proof. \square

We come now to extend [5, Corollary 3.3] to the group inverse of an element in Banach algebra.

Corollary 2.5. *Let $a, b \in \mathcal{A}^\#$ and $\lambda, \mu \in \mathbb{C} \setminus \{0\}$. If $ab = ba$, then the following are equivalent:*

- (1) $\lambda a + \mu b \in \mathcal{A}^\#$.
- (2) $\lambda abb^\# + \mu baa^\# \in \mathcal{A}^\#$.

In this case,

$$(\lambda a + \mu b)^\# = (\lambda abb^\# + \mu baa^\#)^\# + \frac{1}{\lambda}ab^\# + \frac{1}{\mu}b^\#a^\#.$$

Proof. Since $ab = ba$, we have $(\lambda a)(\mu b) = (\mu b)(\lambda a)$. Clearly, $(\lambda a)^\# = \frac{1}{\lambda}a^\#, (\mu b)^\# = \frac{1}{\mu}b^\#$. Applying Theorem 2.2 to $\lambda a, \mu b$, we obtain the result. \square

Example 2.6. *Let $a = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, b = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \in \mathbb{C}^{2 \times 2}$. Then $ab = -ba$ and*

$$a^\# = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, b^\# = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

In view of Theorem 2.2, $a + b$ has group inverse. Evidently,

$$(a + b)^\# = \begin{pmatrix} -\frac{1}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}.$$

In this case, $ab \neq ba$.

3. BLOCK OPERATOR MATRICES

The aim of this section is to present certain simpler representations of the group inverse of the block matrix M by applying Theorem 2.2.

Lemma 3.1. *Let A and D have group inverses. If $D^\pi C = 0$, then $\begin{pmatrix} A & 0 \\ C & D \end{pmatrix}$ has group inverse and*

$$\begin{pmatrix} A & 0 \\ C & D \end{pmatrix}^\# = \begin{pmatrix} A^\# & 0 \\ -D^\#CA^\# + (D^\#)^2CA^\pi & D^\# \end{pmatrix}.$$

Proof. This is obvious by [1, Theorem 3.4]. \square

Recall that $a \in \mathcal{A}$ is called Drazin invertible provided that there is a common solution to the equations $ab = ba$, $b = bab$ and $a^k = a^kba$ for some $k \geq 1$. If such a solution exists, then it is unique and is called a Drazin inverse of a , denoted it by $b = a^D$ (see [3, 4]). We now prove the main result of this section.

Theorem 3.2. *Let A, D and BC have group inverses. If $AB = 0$, $BD = 0$, $B(CB)^\pi = 0$, $C(BC)^\pi = 0$ and $DC = \lambda CA$, then $M \in M_2(\mathcal{A})^\#$ and*

$$M^\# = \begin{pmatrix} 2A^\# & 0 \\ -D^\#CA^\# & D^\# + D^\#(CB)^\pi \end{pmatrix}.$$

Proof. Write $M = P + Q$, where

$$P = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix}, Q = \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix}.$$

Then

$$P^\# = \begin{pmatrix} A^\# & 0 \\ 0 & D^\# \end{pmatrix}, P^\pi = \begin{pmatrix} A^\pi & 0 \\ 0 & D^\pi \end{pmatrix}.$$

Since BC has group inverse, we have

$$(Q^2)^D = \begin{pmatrix} BC & 0 \\ 0 & CB \end{pmatrix}^D = \begin{pmatrix} (BC)^\# & 0 \\ 0 & (CB)^\# \end{pmatrix}.$$

In view of —cite[Theorem 2.1]J, Q has Drazin inverse and

$$Q^D = Q(Q^2)^D = \begin{pmatrix} 0 & B(CB)^\# \\ C(BC)^\# & 0 \end{pmatrix}.$$

Moreover, we have

$$Q^\pi = \begin{pmatrix} (BC)^\pi & 0 \\ 0 & (CB)^\pi \end{pmatrix}.$$

Hence,

$$QQ^D = Q^DQ, Q^D = Q^DQQ^D.$$

It is easy to verify that

$$\begin{aligned} QQ^\pi &= \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix} \begin{pmatrix} (BC)^\pi & 0 \\ 0 & (CB)^\pi \end{pmatrix} \\ &= \begin{pmatrix} 0 & B(CB)^\pi \\ C(BC)^\pi & 0 \end{pmatrix} \\ &= 0; \end{aligned}$$

hence, $Q = Q^2Q^D$. This implies that Q has group inverse and

$$Q^\# = \begin{pmatrix} 0 & B(CB)^\# \\ C(BC)^\# & 0 \end{pmatrix}.$$

Therefore we get

$$P^\#Q^\pi = \begin{pmatrix} A^\#(BC)^\pi & 0 \\ 0 & D^\#(CB)^\pi \end{pmatrix}.$$

We easily check that

$$PQ = \begin{pmatrix} 0 & 0 \\ DC & 0 \end{pmatrix} = \lambda \begin{pmatrix} 0 & 0 \\ CA & 0 \end{pmatrix} = \lambda QP.$$

It follows by Lemma 2.1 that $PQQ^\# = QQ^\#P$ and $QPP^\# = PP^\#Q$. Then we have

$$\begin{aligned} BCA &= 0, CAA^\# = DD^\#C, \\ D(CB)(CB)^\# &= (CB)(CB)^\#D. \end{aligned}$$

Obviously, we have

$$= \begin{pmatrix} A & AA^\#B \\ DD^\#C & D \end{pmatrix} = \begin{pmatrix} A & 0 \\ 0 & D \end{pmatrix} \left[\begin{pmatrix} I & 0 \\ 0 & I \end{pmatrix} + \begin{pmatrix} A^\# & 0 \\ 0 & D^\# \end{pmatrix} \begin{pmatrix} 0 & B \\ C & 0 \end{pmatrix} \right].$$

According to Theorem 2.2, M has group inverse if and only if $\begin{pmatrix} A & 0 \\ DD^\#C & D \end{pmatrix}$ has group inverse. Obviously, $D^\pi(DD^\#C) = 0$.

In view of Lemma 3.1, $\begin{pmatrix} A & 0 \\ DD^\#C & D \end{pmatrix}$ has group inverse. Moreover, we have

$$\begin{pmatrix} A & 0 \\ DD^\#C & D \end{pmatrix}^\# = \begin{pmatrix} A^\# & 0 \\ -D^\#CA^\# + (D^\#)^2CA^\pi & D^\# \end{pmatrix}.$$

Therefore

$$\begin{aligned} M^\# &= [P(I + P^\#Q)]^\# + P^\#Q^\pi \\ &= \begin{pmatrix} A & 0 \\ DD^\#C & D \end{pmatrix}^\# + P^\#Q^\pi \\ &= \begin{pmatrix} A^\# + A^\#(BC)^\pi & 0 \\ -D^\#CA^\# + (D^\#)^2CA^\pi & D^\# + D^\#(CB)^\pi \end{pmatrix}, \end{aligned}$$

as asserted. \square

Corollary 3.3. *Let A, D and BC have group inverses. If $CA = 0, DC = 0, B(CB)^\pi = 0, C(BC)^\pi = 0$ and $AB = \lambda BD$, then $M \in M_2(\mathcal{A})^\#$ and*

$$M^\# = \begin{pmatrix} A^\# + A^\#(BC)^\pi & -A^\#BD^\# \\ 0 & 2D^\# \end{pmatrix}.$$

Proof. Since BC has group inverse, it has Drazin inverse. According to Cline's formula ([4, Theorem 2.3]), CB has Drazin inverse. Hence, $(CB)(CB)^D = (CB)^D(CB), (CB)^D = (CB)^D(CB)(CB)^D$. Since $B(CB)^\pi = 0$, we have $CB = (CB)(CB)^D(CB)$. Therefore CB has group invrse. Applying Theorem 3.2, we prove that

$\begin{pmatrix} D & C \\ B & A \end{pmatrix}$ has group inverse. Clearly,

$$M = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \begin{pmatrix} D & C \\ B & A \end{pmatrix} \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix},$$

and so M has group inverse. Moreover, $\begin{pmatrix} D & C \\ B & A \end{pmatrix}$ has group inverse. Clearly,

$$M^\# = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix} \begin{pmatrix} D & C \\ B & A \end{pmatrix}^\# \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix},$$

as required. \square

Corollary 3.4. *Let $A \in \mathbb{C}^{m \times m}$, $D \in \mathbb{C}^{n \times n}$ have group inverses. If $AB = 0$, $BD = 0$, $B(BC)^\pi = 0$, $C(BC)^\pi = 0$ and $DC = \lambda CA$, then $M \in \mathbb{C}^{(m+n) \times (m+n)}$ has group inverse and*

$$M^\# = \begin{pmatrix} A^\# + A^\#(BC)^\pi & 0 \\ -D^\#CA^\# + (D^\#)^2CA^\pi & D^\# + D^\#(CB)^\pi \end{pmatrix}.$$

Proof. Since $C(BC)^\pi = 0$, we have $BC = (BC)^2(BC)^D$, hence, BC has group inverse. This completes the proof by Theorem 3.2. \square

We are now ready to prove:

Theorem 3.5. *Let A, D and BC have group inverses. If $CA = 0$, $DC = 0$, $(CB)^\pi C = 0$, $(BC)^\pi B = 0$ and $AB = \lambda BD$, then $M \in M_2(\mathcal{A})^\#$ and*

$$M^\# = \begin{pmatrix} A^\# + (BC)^\pi A^\# & -A^\#BD^\# + A^\pi B(D^\#)^2 \\ 0 & D^\# + (CB)^\pi D^\# \end{pmatrix}.$$

Proof. By hypothesis, A^T, D^T and $C^T B^T$ have group inverses. Moreover, we check that

$$\begin{aligned} D^T B^T &= \lambda B^T A^T, A^T C^T = 0, C^T D^T = 0, \\ C^T (B^T C^T)^\pi &= 0, B^T (C^T B^T)^\pi = 0. \end{aligned}$$

Applying Theorem 3.2 to the transpose M^T of M , where

$$M^T = \begin{pmatrix} A^T & C^T \\ B^T & D^T \end{pmatrix}.$$

Then M^T has group inverse, and so has M . Moreover, we derive

$$(M^T)^\# = \begin{pmatrix} (A^T)^\# + (A^T)^\#(C^T B^T)^\pi & -(D^T)^\# B^T (A^T)^\# + [(D^T)^\#]^2 B^T (A^T)^\pi \\ 0 & (D^T)^\# + (D^T)^\#(B^T C^T)^\pi \end{pmatrix}.$$

This completes the proof. \square

As a consequence of the above, we derive

Corollary 3.6. *Let A, D and BC have group inverses. If $AB = 0, BD = 0(BC)^\pi B = 0, (CB)^\pi C = 0$ and $DC = \lambda CA$, then $M \in M_2(\mathcal{A})^\#$ and*

$$M^\# = \begin{pmatrix} A^\# + (BC)^\pi A^\# & 0 \\ -D^\# C A^\# + D^\pi C (A^\#)^2 & D^\# + (CB)^\pi D^\# \end{pmatrix}.$$

Proof. Applying Theorem 3.5, $\begin{pmatrix} D & C \\ B & A \end{pmatrix}$ has group inverse. As in the proof of Corollary 3.3, we obtain the result. \square

Corollary 3.7. *Let $A \in \mathbb{C}^{m \times m}, D \in \mathbb{C}^{n \times n}$ have group inverses. If $CA = 0, DC = 0, (CB)^\pi C = 0, (BC)^\pi B = 0$ and $AB = \lambda BD$, then $M \in \mathbb{C}^{(m+n) \times (m+n)}$ has group inverse and*

$$M^\# = \begin{pmatrix} A^\# + (BC)^\pi A^\# & -A^\# B D^\# + A^\pi B (D^\#)^2 \\ 0 & D^\# + (CB)^\pi D^\# \end{pmatrix}.$$

REFERENCES

- [1] J. Benitez; X. Liu and T. Zhu, Additive results for the group inverse in an algebra with applications to block operators, *Linear Multilinear Algebra*, **59**(2011), 279–289.
- [2] D.S. Cvetkovic-Ilic, The generalized Drazin inverse with commutativity up to a factor in a Banach algebra, *Linear Algebra Appl.*, **431**(2009), 783–791.
- [3] D.S. Cvetkovic-Ilic, Some results on the $(2, 2, 0)$ Drazin inverse problem, *Linear Algebra Appl.*, **438**(2013), 4726–4741.
- [4] Y. Jiang; Y. Wen and Q. Zeng, Generalizations of Cline’s formula for three generalized inverses, *Revista de la UMA*, **58**(2017), 127–134.
- [5] X. Liu; L. Wu and J. Benitez, On the group inverse of linear combinatons of two group invertible matrices, *Electronic J. Linear Algebra*, **22**(2011), 490–503.

- [6] N. Mihajlovic and D.S. Djordjevic, On group invertibility in rings, *Filomat*, **33**(2019), 6141–6150.
- [7] D. Mosaic, Group inverse and generalized Drazin inverse of block matrices in a Banach algebra, *Bull. Korean Math. Soc.*, **51**(2014), 765–771.
- [8] D. Zhang, Representations for generalized Drazin inverse of operator matrices over a Banach space, *Turk. J. Math.*, **40**(2016), 428–437.
- [9] D. Zhang and D. Mosić, Explicit formulae for the generalized Drazin inverse of block matrices over a Banach algebra, *Filomat*, **32**(2018), 5907–5917.
- [10] M. Zhou; J. Chen and X. Zhu, The group inverse and core inverse of sum of two elements in a ring, *Comm. Algebra*, **48**(2020), 676–690.
- [11] H. Zhu and J. Chen, Additive property of pseudo Drazin inverse of elements in Banach algebras, *Filomat*, **28**(2014), 1773–1781.
- [12] H. Zhu and J. Chen, Additive and product properties of Drazin inverses of elements in a ring, *Bull. Malays. Math. Sci. Soc.*, **40**(2017), 259–278.

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANGZHOU, CHINA

Email address: <huanyinchen@aliyun.com>

WOMEN'S UNIVERSITY OF SEMNAN (FARZANEGAN), SEMNAN, IRAN

Email address: <sheibani@fgusem.ac.ir>