BOUNDARY POINTS, MINIMAL L² INTEGRALS AND CONCAVITY PROPERTY II: ON WEAKLY PSEUDOCONVEX KÄHLER MANIFOLDS

QI'AN GUAN, ZHITONG MI, AND ZHENG YUAN

ABSTRACT. In this article, we consider minimal L^2 integrals on the sublevel sets of plurisubharmonic functions on weakly pseudoconvex Kähler manifolds with Lebesgue measurable gain related to modules at boundary points of the sublevel sets, and establish a concavity property of the minimal L^2 integrals. As applications, we present a necessary condition for the concavity degenerating to linearity, a concavity property related to modules at inner points of the sublevel sets, an optimal support function related to the modules, a strong openness property of the modules and a twisted version, an effectiveness result of the strong openness property of the modules.

1. INTRODUCTION

Let φ be a plurisubharmonic function on a complex manifold M (see [12]). Recall that the multiplier ideal sheaf $\mathcal{I}(\varphi)$ is the sheaf of germs of holomorphic functions f such that $|f|^2 e^{-\varphi}$ is locally integrable, which plays an important role in several complex variables, complex algebraic geometry and complex differential geometry (see e.g. [50, 45, 46, 15, 16, 13, 17, 43, 48, 49, 14, 37]).

The strong openness property of multiplier ideal sheaves [33], i.e. $\mathcal{I}(\varphi) = \mathcal{I}_+(\varphi) := \bigcup_{\epsilon>0} \mathcal{I}((1+\epsilon)\varphi)$ (conjectured by Demailly [13]) is an important feature of multiplier ideal sheaves and has opened the door to new types of approximation technique (see e.g. [33, 44, 41, 4, 5, 19, 7, 51, 36, 52, 53, 20, 42, 8]). Guan-Zhou [33] proved the strong openness property (the 2-dimensional case was proved by Jonsson-Mustață [39]). After that, Guan-Zhou [34] established an effectiveness result of the strong openness property by considering the minimal L^2 integral on the pseudoconvex domain D.

Considering the minimal L^2 integrals on the sublevel sets of the weight φ , Guan [23] obtained a sharp version of Guan-Zhou's effectiveness result, and established a concavity property of the minimal L^2 integrals (see also [22],[25],[26],[28],[27]).

Recall that the minimal L^2 integrals on the sublevel sets of φ in [23] (see also [22],[25],[26],[28],[27]) are related to the modules (ideals) at the inner points of the sublevel sets. In [2], Bao-Guan-Yuan considered the minimal L^2 integrals related to modules at boundary points of the sublevel sets of plurisubharmonic functions on pseudoconvex domains, and gave a concavity property of the minimal L^2 integrals, which deduced a sharp effectiveness result related to a conjecture posed

Date: June 6, 2022.

²⁰²⁰ Mathematics Subject Classification. 32Q15, 32F10, 32U05, 32W05.

Key words and phrases. minimal L^2 integrals, plurisubharmonic functions, boundary points, weakly pseudoconvex Kähler manifold.

by Jonsson-Mustață [39], and completed the approach from the conjecture to the strong openness property.

In the present article, we consider minimal L^2 integrals on sublevel sets of plurisubharmonic functions on weakly pseudoconvex Kähler manifolds with Lebesgue measurable gain related to modules at boundary points of the sublevel sets, and establish a concavity property of the minimal L^2 integrals.

1.1. Main result. Let M be a complex manifold. Let X and Z be closed subsets of M. We call that a triple (M, X, Z) satisfies condition (A), if the following two statements hold:

I. X is a closed subset of M and X is locally negligible with respect to L^2 holomorphic functions; i.e., for any local coordinated neighborhood $U \subset M$ and for any L^2 holomorphic function f on $U \setminus X$, there exists an L^2 holomorphic function \tilde{f} on U such that $\tilde{f}|_{U \setminus X} = f$ with the same L^2 norm;

II. Z is an analytic subset of M and $M \setminus (X \cup Z)$ is a weakly pseudoconvex Kähler manifold.

Let M be an n-dimensional complex manifold, and let (M, X, Z) satisfy condition (A). Let K_M be the canonical line bundle on M. Let F be a holomorphic function on M. We assume that F is not identically zero. Let ψ be a plurisubharmonic function on M. Let φ_{α} be a Lebesgue measurable function on M such that $\varphi_{\alpha} + \psi$ is a plurisubharmonic function on M.

Let $T \in [-\infty, +\infty)$. Denote that

$$\Psi := \min\{\psi - 2\log|F|, -T\}.$$

For any $z \in M$ satisfying F(z) = 0, we set $\Psi(z) = -T$. Note that for any $t \geq T$, the holomorphic function F has no zero points on the set $\{\Psi < -t\}$. Hence $\Psi = \psi - 2 \log |F| = \psi + 2 \log |\frac{1}{F}|$ is a plurisubharmonic function on $\{\Psi < -t\}$.

Definition 1.1. We call that a positive measurable function c (so-called "gain") on $(T, +\infty)$ is in class $P_{T,M,\Psi}$ if the following two statements hold:

(1) $c(t)e^{-t}$ is decreasing with respect to t;

(2) There exist $T_1 > T$ and a closed subset E of M such that $E \subset Z \cap \{\Psi(z) = -\infty\}$ and for any compact subset $K \subset M \setminus E$, $e^{-\varphi_{\alpha}}c(-\Psi_1)$ has a positive lower bound on K, where $\Psi_1 := \min\{\psi - 2\log|F|, -T_1\}$.

Let z_0 be a point in M. Denote that $\tilde{J}(\Psi)_{z_0} := \{f \in \mathcal{O}(\{\Psi < -t\} \cap V) : t \in \mathbb{R}$ and V is a neighborhood of $z_0\}$. We define an equivalence relation \backsim on $\tilde{J}(\Psi)_{z_0}$ as follows: for any $f, g \in \tilde{J}(\Psi)_{z_0}$, we call $f \backsim g$ if f = g holds on $\{\Psi < -t\} \cap V$ for some $t \gg T$ and open neighborhood $V \ni o$. Denote $\tilde{J}(\Psi)_{z_0} / \backsim$ by $J(\Psi)_{z_0}$, and denote the equivalence class including $f \in \tilde{J}(\Psi)_{z_0}$ by f_{z_0} .

If $z_0 \in \bigcap_{t>T} \{\Psi < -t\}$, then $J(\Psi)_{z_0} = \mathcal{O}_{M,z_0}$ (the stalk of the sheaf \mathcal{O}_M at z_0), and f_{z_0} is the germ (f, z_0) of holomorphic function f.

Let $f_{z_0}, g_{z_0} \in J(\Psi)_{z_0}$ and $(h, z_0) \in \mathcal{O}_{M, z_0}$. We define $f_{z_0} + g_{z_0} := (f + g)_{z_0}$ and $(h, z_0) \cdot f_{z_0} := (hf)_{z_0}$. Note that $(f + g)_{z_0}$ and $(hf)_{z_0} \ (\in J(\Psi)_{z_0})$ are independent of the choices of the representatives of f, g and h. Hence $J(\Psi)_{z_0}$ is an \mathcal{O}_{M, z_0} -module.

For $f_{z_0} \in J(\Psi)_{z_0}$ and $a, b \ge 0$, we call $f_{z_0} \in I(a\Psi + b\varphi_{\alpha})_{z_0}$ if there exist $t \gg T$ and a neighborhood V of z_0 , such that $\int_{\{\Psi < -t\} \cap V} |f|^2 e^{-a\Psi - b\varphi_{\alpha}} < +\infty$. Note that $I(a\Psi + b\varphi_{\alpha})_{z_0}$ is an \mathcal{O}_{M,z_0} -submodule of $J(\Psi)_{z_0}$. If $z_0 \in \cap_{t>T} \{\Psi < -t\}$, then $I_{z_0} = \mathcal{O}_{M,z_0}$, where $I_{z_0} := I(0\Psi + 0\varphi_{\alpha})_{z_0}$.

 $\mathbf{2}$

Let Z_0 be a subset of M. Let f be a holomorphic (n, 0) form on $\{\Psi < -t_0\} \cap V$, where $V \supset Z_0$ is an open subset of M and $t_0 > T$ is a real number. Let J_{z_0} be an \mathcal{O}_{M,z_0} -submodule of $J(\Psi)_{z_0}$ such that $I(\Psi + \varphi_\alpha)_{z_0} \subset J_{z_0}$, where $z_0 \in Z_0$. Denote

$$\inf \left\{ \int_{\{\Psi < -t\}} |\tilde{f}|^2 e^{-\varphi_{\alpha}} c(-\Psi) : \tilde{f} \in H^0(\{\Psi < -t\}, \mathcal{O}(K_M)) \\ \& (\tilde{f} - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}, \text{for any } z_0 \in Z_0 \right\}$$
(1.1)

by $G(t; c, \Psi, \varphi_{\alpha}, J, f)$, where $t \in [T, +\infty)$, c is a nonnegative function on $(T, +\infty)$ and $|f|^2 := \sqrt{-1}^{n^2} f \wedge \bar{f}$ for any (n, 0) form f. Without misunderstanding, we simply denote $G(t; c, \Psi, \varphi_{\alpha}, J, f)$ by G(t). For various c, ψ and φ_{α} , we simply denote $G(t; c, \Psi, \varphi_{\alpha}, J, f)$ by $G(t; c), G(t; \Psi)$ and $G(t; \varphi_{\alpha})$ respectively.

In this article, we obtain the following concavity property of G(t).

Theorem 1.2. Let $c \in \mathcal{P}_{T,M,\Psi}$. If there exists $t \in [T, +\infty)$ satisfying that $G(t) < +\infty$, then $G(h^{-1}(r))$ is concave with respect to $r \in (\int_{T_1}^T c(t)e^{-t}dt, \int_{T_1}^{+\infty} c(t)e^{-t}dt),$ $\lim_{t\to T+0} G(t) = G(T)$ and $\lim_{t\to +\infty} G(t) = 0$, where $h(t) = \int_{T_1}^t c(t_1)e^{-t_1}dt_1$ and $T_1 \in (T, +\infty)$.

When M is a pseudoconvex domain D in \mathbb{C}^n , $\varphi_{\alpha} \equiv 0$, $c(t) \equiv 1$ and T = 0, Theorem 1.2 degenerates to the concavity property in [2].

Remark 1.3. Let $c \in \mathcal{P}_{T,M,\Psi}$. If $\int_{T_1}^{+\infty} c(t)e^{-t}dt = +\infty$ and $f_{z_0} \notin \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for some $z_0 \in Z_0$, then $G(t) = +\infty$ for any $t \geq T$. Thus, when there exists $t \in [T, +\infty)$ satisfying that $G(t) \in (0, +\infty)$, we have $\int_{T_1}^{+\infty} c(t)e^{-t}dt < +\infty$ and $G(\hat{h}^{-1}(r))$ is concave with respect to $r \in (0, \int_T^{+\infty} c(t)e^{-t}dt)$, where $\hat{h}(t) = \int_t^{+\infty} c(t)e^{-t}dt$.

Let c(t) be a nonnegative measurable function on $(T, +\infty)$. Set

$$\mathcal{H}^{2}(c,t) = \left\{ \tilde{f} : \int_{\{\Psi < -t\}} |\tilde{f}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) < +\infty, \ \tilde{f} \in H^{0}(\{\Psi < -t\}, \mathcal{O}(K_{M})) \\ \& (\tilde{f} - f)_{z_{0}} \in \mathcal{O}(K_{M})_{z_{0}} \otimes J_{z_{0}}, \text{for any } z_{0} \in Z_{0}) \right\},$$

where $t \in [T, +\infty)$.

As a corollary of Theorem 1.2, we give a necessary condition for the concavity property degenerating to linearity.

Corollary 1.4. Let $c \in \mathcal{P}_{T,M,\Psi}$. Assume that $G(t) \in (0, +\infty)$ for some $t \geq T$, and $G(\hat{h}^{-1}(r))$ is linear with respect to $r \in [0, \int_T^{+\infty} c(s)e^{-s}ds)$, where $\hat{h}(t) = \int_t^{+\infty} c(l)e^{-l}dl$.

Then there exists a unique holomorphic (n,0) form \tilde{F} on $\{\Psi < -T\}$ such that $(\tilde{F}-f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ holds for any $z_0 \in Z_0$, and $G(t) = \int_{\{\Psi < -t\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi)$ holds for any $t \geq T$.

Furthermore

$$\int_{\{-t_1 \le \Psi < -t_2\}} |\tilde{F}|^2 e^{-\varphi_\alpha} a(-\Psi) = \frac{G(T_1;c)}{\int_{T_1}^{+\infty} c(t)e^{-t}dt} \int_{t_2}^{t_1} a(t)e^{-t}dt \qquad (1.2)$$

holds for any nonnegative measurable function a on $(T, +\infty)$, where $T \leq t_2 < t_1 \leq$ $+\infty$ and $T_1 \in (T, +\infty)$.

Remark 1.5. If $\mathcal{H}^2(\tilde{c}, t_0) \subset \mathcal{H}^2(c, t_0)$ for some $t_0 \geq T$, we have

$$G(t_0; \tilde{c}) = \int_{\{\Psi < -t_0\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \tilde{c}(-\Psi) = \frac{G(T_1; c)}{\int_{T_1}^{+\infty} c(t) e^{-t} dt} \int_{t_0}^{+\infty} \tilde{c}(s) e^{-s} ds, \quad (1.3)$$

where \tilde{c} is a nonnegative measurable function on $(T, +\infty)$ and $T_1 \in (T, +\infty)$.

1.2. Applications. In this section, we give some applications of Theorem 1.2.

1.2.1. Concavity property at inner points. Let M be a complex manifold. Let Xand Z be closed subsets of M. Assume that the triple (M, X, Z) satisfies condition (A). Let ψ be a plurisubharmonic function on M. Let φ be a Lebesgue measurable function on M, such that $\psi + \varphi$ is a plurisubharmonic function on M. Denote $T = -\sup \psi.$ M

Definition 1.6. We call a positive measurable function c (so-called "gain") on $(T, +\infty)$ in class $P_{T,M}$ if the following two statements hold:

(1) $c(t)e^{-t}$ is decreasing with respect to t;

(2) there is a closed subset E of M such that $E \subset Z \cap \{\psi(z) = -\infty\}$ and for any compact subset $K \subset M \setminus E$, $e^{-\varphi}c(-\psi)$ has a positive lower bound on K.

Let Z_0 be a subset of $\{\psi = -\infty\}$ such that $Z_0 \cap Supp(\mathcal{O}/\mathcal{I}(\varphi + \psi)) \neq \emptyset$. Let $U \supset Z_0$ be an open subset of M, and let f be a holomorphic (n, 0) form on U. Let $\mathcal{F}_{z_0} \supset \mathcal{I}(\varphi + \psi)_{z_0}$ be an ideal of \mathcal{O}_{z_0} for any $z_0 \in Z_0$.

Denote

$$\inf\left\{\int_{\{\psi<-t\}} |\tilde{f}|^2 e^{-\varphi} c(-\psi) : \tilde{f} \in H^0(\{\psi<-t\}, \mathcal{O}(K_M)) \\ & \& (\tilde{f}-f) \in H^0(Z_0, (\mathcal{O}(K_M) \otimes \mathcal{F})|_{Z_0})\right\}$$
(1.4)

by $G(t; c, \psi, \varphi, f, \mathcal{F})$, where $t \in [T, +\infty)$, c is a nonnegative function on $(T, +\infty)$, $|f|^2 := \sqrt{-1}^{n^2} f \wedge \bar{f}$ for any (n, 0) form f, and $(\tilde{f} - f) \in H^0(Z_0, (\mathcal{O}(K_M) \otimes \mathcal{F})|_{Z_0})$ means $(f - f, z_0) \in (\mathcal{O}(K_M) \otimes \mathcal{F})_{z_0}$ for all $z_0 \in Z_0$.

Using Theorem 1.2, we obtain the following concavity property of $G(t; c, \psi, \varphi, f, \mathcal{F})$.

Corollary 1.7. Assume that $c \in \mathcal{P}_{T,M}$. If there exists $t_0 \geq T$ satisfying that $\lim_{t \to +\infty} G(t; c, \psi, \varphi, f, \mathcal{F}) = 0, \text{ where } h(t) = \int_{T_1}^t c(t_1) e^{-t_1} dt_1 \text{ and } T_1 \in (T, +\infty).$

It follows from Corollary 1.4 and Corollary 1.7 that we have the following remark.

Remark 1.8. Let $c \in \mathcal{P}_{T,M}$. Assume that $G(t; c, \psi, \varphi, f, \mathcal{F}) \in (0, +\infty)$ for some $t \geq T$, and $G(\hat{h}^{-1}(r); c, \psi, \varphi, f, \mathcal{F})$ is linear with respect to $r \in [0, \int_{T}^{+\infty} c(s)e^{-s}ds)$, where $\hat{h}(t) = \int_{t}^{+\infty} c(l)e^{-l}dl$.

Then there exists a unique holomorphic (n, 0) form \tilde{F} on M such that $(\tilde{F} - f) \in H^0(Z_0, (\mathcal{O}(K_M) \otimes \mathcal{F})|_{Z_0})$ and $G(t; c, \psi, \varphi, f, \mathcal{F}) = \int_{\{\psi < -t\}} |\tilde{F}|^2 e^{-\varphi} c(-\psi)$ holds for any $t \geq T$.

Furthermore

$$\int_{\{-t_1 \le \psi < -t_2\}} |\tilde{F}|^2 e^{-\varphi} a(-\psi) = \frac{G(T_1; c, \psi, \varphi, f, \mathcal{F})}{\int_{T_1}^{+\infty} c(t) e^{-t} dt} \int_{t_2}^{t_1} a(t) e^{-t} dt$$

holds for any nonnegative measurable function a on $(T, +\infty)$, where $T \le t_2 < t_1 \le +\infty$ and $T_1 \in (T, +\infty)$.

When ψ is a plurisubharmonic function on M, and $\{\psi < -t\} \setminus (X \cup Z)$ is a weakly pseudoconvex Kähler manifold for any $t \in \mathbb{R}$, Corollary 1.7 and Remark 1.8 can be referred to [27] (see also [26] and [1]).

1.2.2. An optimal support function related to $I(a\Psi)$. Let M be an n-dimensional complex manifold. Let X and Z be closed subsets of M, and let (M, X, Z) satisfy condition (A). Let K_M be the canonical line bundle on M. Let F be a holomorphic function on M. Assume that F is not identically zero. Let ψ be a plurisubharmonic function on M.

Denote that

$\Psi := \min\{\psi - 2\log|F|, 0\}.$

For any $z \in M$ satisfying F(z) = 0, we set $\Psi(z) = 0$. Denote that $M_t := \{z \in M : -t \leq \Psi(z) < 0\}$. Let Z_0 be a subset of M, and let f be a holomorphic (n, 0) form on $\{\Psi < 0\}$. Denote

$$\inf\left\{\int_{M_t} |\tilde{f}|^2 : f \in H^0(\{\Psi < 0\}, \mathcal{O}(K_M)) \\ \& (\tilde{f} - f)_{z_0} \in \mathcal{O}(K_M) \otimes I(\Psi)_{z_0} \text{ for any } z_0 \in Z_0\right\}$$

by $C_{f,\Psi,t}(Z_0)$ for any $t \ge 0$. When $C_{f,\Psi,t}(Z_0) = 0$ or $+\infty$, we set $\frac{\int_{M_t} |f|^2 e^{-\Psi}}{C_{f,\Psi,t}(Z_0)} = +\infty$. We obtain the following optimal support function of $\frac{\int_{M_t} |f|^2 e^{-\Psi}}{C_{f,\Psi,t}(Z_0)}$.

Proposition 1.9. Assume that $\int_{\{\Psi < -l\}} |f|^2 < +\infty$ holds for any l > 0. Then the inequality

$$\frac{\int_{M_t} |f|^2 e^{-\Psi}}{C_{f,\Psi,t}(Z_0)} \ge \frac{t}{1 - e^{-t}} \tag{1.5}$$

holds for any $t \ge 0$, where $\frac{t}{1-e^{-t}}$ is the optimal support function.

Proposition 1.9 can be referred to [29] when M is a pseudoconvex domain in \mathbb{C}^n , $F \equiv 1, Z_0 = \{z_0\} \subset M$ and $\psi(z_0) = -\infty$.

Take $M = \Delta \subset \mathbb{C}$, $Z_0 = o$ the origin of \mathbb{C} , $F \equiv 1$, $\psi = 2 \log |z|$ and $f \equiv dz$. It is clear that $\int_M |f|^2 < +\infty$. Note that $C_{f,\Psi,t}(Z_0) = 2\pi(1-e^{-t})$ and $\int_{M_t} |f|^2 e^{-\Psi} = 2t\pi$. Then $\frac{\int_{M_t} |f|^2 e^{-\Psi}}{C_{f,\Psi,t}(Z_0)} = \frac{t}{1-e^{-t}}$, which shows the optimality of the support function $\frac{t}{1-e^{-t}}$.

1.2.3. Strong openness property of $I(a\Psi + \varphi)_o$ and a twisted version. Let $D \subseteq \mathbb{C}^n$ be a pseudoconvex domain containing the origin o, and let ψ be a plurisubharmonic function on D. Let $F \neq 0$ be a holomorphic function on D, and let φ be a plurisubharmonic function on D. Denote that

$$\Psi := \min\{\psi - 2\log|F|, 0\}.$$

For any $z \in M$ satisfying F(z) = 0, we set $\Psi(z) = 0$.

Recall that $f_o \in I(a\Psi + b\varphi)_o$ if and only if there exist $t \gg 0$ and a neighborhood V of o such that $\int_{\{\Psi < -t\} \cap V} |h|^2 e^{-a\Psi - b\varphi} < +\infty$, where $a \ge 0$ and $b \ge 0$. Denote that

$$I_+(a\Psi + b\varphi)_o := \bigcup_{s>a} I(s\Psi + b\varphi)_o$$

for any $b \ge 0$. In [2], Bao-Guan-Yuan gave the strong openness property of $I(a\Psi)_o$.

Theorem 1.10 ([2]). $I(a\Psi)_o = I_+(a\Psi)_o$ holds for any $a \ge 0$.

When $F \equiv 1$ and $\psi(o) = -\infty$, Theorem 1.10 is the strong openness property of multiplier ideal sheaves [33], i.e. $\mathcal{I}(\psi)_o = \mathcal{I}_+(\psi)_o := \bigcup_{s>1} \mathcal{I}(s\psi)_o$. We present the following strong openness property of $I(a\Psi + \varphi)_o$.

Theorem 1.11. $I(a\Psi + \varphi)_o = I_+(a\Psi + \varphi)_o$ holds for any $a \ge 0$.

Let p be a positive real number. Denote that

 $c_{o,p}^{h}(\varphi) := \sup\{c \ge 0 : |h|^{p} e^{-2c\varphi} \text{ is } L^{1} \text{ on a neighborhood of } o\}.$

When p = 2, $c_{o,2}^{h}(\psi)$ is the jumping number $c_{o}^{h}(\psi)$ (see [40]). Note that the strong openness property of multiplier ideal sheaves is equivalent to $|h|^{2}e^{-2c_{o}^{h}(\psi)\psi}$ is not integrable near o for any holomorphic function h on a neighborhood of o.

Using the strong openness property of multiplier ideal sheaves, Fornæss [18] obtained the following strong openness property of multiplier ideal sheaves in L^p :

 $|h|^{p}e^{-2c_{o,p}^{h}(\psi)\psi}$ is not integrable near o for any holomorphic function h on a neighborhood of o.

In [31], Guan-Yuan gave the following twisted version of strong openness property of multiplier ideal sheaves in L^p (some related results can be referred to [35] and [6]).

Theorem 1.12 ([31]). Let $p \in (0, +\infty)$, and let a(t) be a positive measurable function on $(-\infty, +\infty)$. If one of the following conditions holds:

(1) a(t) is decreasing near $+\infty$;

(2) $a(t)e^t$ is increasing near $+\infty$,

then the following three statements are equivalent:

(A') a(t) is not integrable near $+\infty$;

(B') $a(-2c_{o,p}^{h}(\psi)\psi)\exp(-2c_{o,p}^{h}(\psi)\psi + p\log|h|)$ is not integrable near o for any ψ and h satisfying $c_{o,p}^{h}(\psi) < +\infty$;

 $(C') \ a(-2c^h_{o,p}(\psi)\psi + p\log|h|) \exp(-2c^h_{o,p}(\psi)\psi + p\log|h|) \ is \ not \ integrable \ near \ o for \ any \ \psi \ and \ h \ satisfying \ c^h_{o,p}(\psi) < +\infty.$

Denote that

$$a_o^J(\Psi;\varphi) := \sup\{a \ge 0 : f_o \in I(2a\Psi + \varphi)_o\}$$

for any $f_o \in I(\varphi)_o$. Especially, $a_o^f(\Psi; \varphi)$ is the jumping number $c_o^f(\psi)$ (see [40]) when $F \equiv 1, \psi(o) = -\infty$ and $\varphi \equiv 0$.

Note that the strong openness property of $I(a\Psi + \varphi)_o$ is equivalent to that $f_o \notin I(2a_o^f(\Psi)\Psi + \varphi)_o$ for any $f_o \in I(\varphi)_o$ satisfying $a_o^f(\Psi; \varphi) < +\infty$. We present a twisted version of the strong openness property of $I(a\Psi + \varphi)_o$.

Theorem 1.13. Let a(t) be a positive measurable function on $(-\infty, +\infty)$. If one of the following conditions holds:

- (1) a(t) is decreasing near $+\infty$;
- (2) $a(t)e^t$ is increasing near $+\infty$,
- then the following two statements are equivalent:
- (A) a(t) is not integrable near $+\infty$;
- (B) for any Ψ , φ and $f_o \in I(\varphi)_o$ satisfying $a_o^f(\Psi; \varphi) < +\infty$, we have

$$a(-2a_o^f(\Psi;\varphi)\Psi)\exp(-2a_o^f(\Psi;\varphi)\Psi-\varphi+2\log|f|)\notin L^1(U\cap\{\Psi<-t\}),$$

where U is any neighborhood of o and t > 0.

Remark 1.14. Theorem 1.13 is a generalization of Theorem 1.12. We prove this remark in Section 7.

1.2.4. Effectiveness of the strong openness property of $I(a\Psi + \varphi)_o$. Let D be a pseudoconvex domain in \mathbb{C}^n containing the origin o, and let ψ be a plurisubharmonic function on D. Let $F \neq 0$ be a holomorphic function on D, and let φ be a plurisubharmonic function on D. Denote that

$$\Psi := \min\{\psi - 2\log|F|, 0\}.$$

If F(z) = 0 for some $z \in M$, we set $\Psi(z) = 0$. Let f be a holomorphic function on $\{\Psi < 0\}$. Denote that

$$\frac{1}{K_{\Psi,f,a}(o)} := \inf \left\{ \int_{\{\Psi < 0\}} |\tilde{f}|^2 e^{-\varphi - (1-a)\Psi} : (\tilde{f} - f)_o \in I_+(\varphi + 2a_o^f(\Psi; \varphi)\Psi)_o \\ & \& \tilde{f} \in \mathcal{O}(\{\Psi < 0\}) \right\},$$

where $a \in (0, +\infty)$.

We present an effectiveness result of the strong openness property of $I(a\Psi + \varphi)_{z_0}$.

Theorem 1.15. Let C_1 and C_2 be two positive constants. If there exists a > 0, such that

(1) $\int_{\{\Psi < 0\}} |f|^2 e^{-\varphi - \Psi} \le C_1;$ (2) $\frac{1}{K_{\Psi,f,a}(o)} \ge C_2.$ Then for any q > 1 satisfying

$$\theta_a(q) > \frac{C_1}{C_2},$$

we have $f_o \in I(q\Psi + \varphi)_o$, where $\theta_a(q) = \frac{q+a-1}{q-1}$.

When $F \equiv 1$ and $\psi(o) = -\infty$, Theorem 1.15 degenerates to the effectiveness result of strong openness property of multiplier ideal sheaves in L^p [30] (some related results can be referred to [34, 23]).

2. Some preparations

In this section, we do some preparations.

2.1. L^2 method. Let M be an n-dimensional weakly pseudoconvex Kähler manifolds. Let ψ be a plurisubharmonic function on M. Let F be a holomorphic function on M. We assume that F is not identically zero. Let φ_{α} be a Lebesgue measurable function on M such that $\varphi_{\alpha} + \psi$ is a plurisubharmonic function on M.

Let δ be a positive integer. Let T be a real number. Denote

$$\varphi := \varphi_{\alpha} + (1+\delta) \max\{\psi + T, 2\log|F|\}$$

and

$$\Psi := \min\{\psi - 2\log|F|, -T\}.$$

If F(z) = 0 for some $z \in M$, we set $\Psi(z) = -T$.

Let c(t) be a positive measurable function on $[T, +\infty)$ such that $c(t)e^{-t}$ is decreasing with respect t. We have the following lemma.

Lemma 2.1. Let $B \in (0, +\infty)$ and $t_0 > T$ be arbitrarily given. Let f be a holomorphic (n, 0) form on $\{\Psi < -t_0\}$ such that

$$\int_{\{\Psi<-t_0\}\cap K} |f|^2 < +\infty,$$

for any compact subset $K \subset M$, and

$$\int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi} < +\infty$$

Then there exists a holomorphic (n, 0) form \tilde{F} on M such that

$$\int_{M} |\tilde{F} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq (\frac{1}{\delta}c(T)e^{-T} + \int_{T}^{t_0+B} c(s)e^{-s}ds) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0-B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi}$$

where $b_{t_0,B}(t) = \int_{-\infty}^t \frac{1}{B} \mathbb{I}_{\{-t_0 - B < s < -t_0\}} ds, \ v_{t_0,B}(t) = \int_{-t_0}^t b_{t_0,B}(s) ds - t_0.$

We introduce the following property of gain function c(t). Let $T_0 \in [-\infty, +\infty)$. Denote

$$\Psi_0 := \min\{\psi - 2\log|F|, -T_0\}.$$

If F(z) = 0 for some $z \in M$, we set $\Psi(z) = -T_0$. Let $c(t) \in \mathcal{P}_{T_0,M,\Psi_0}$, i.e.,

(1) $c(t)e^{-t}$ is decreasing with respect to t;

(2) There exist $T_1 > T_0$ and a closed subset E of M such that $E \subset Z \cap \{\Psi(z) = -\infty\}$ and for any compact subset $K \subset M \setminus E$, $e^{-\varphi_{\alpha}}c(-\Psi_1)$ has a positive lower bound on K, where $\Psi_1 := \min\{\psi - 2\log|F|, -T_1\}$.

Proposition 2.2. Let $c(t) \in \mathcal{P}_{T_0,M,\Psi_0}$. Let T be a real number such that $T > T_0$. Then for any compact subset $K \subset M \setminus E$, we have $e^{-\varphi_{\alpha}}c(-\Psi)$ has a positive lower bound on K, where $\Psi := \min\{\psi - 2\log|F|, -T\}$.

Proof. We note that c(t) is positive on $[T_1, T]$ (or $[T, T_1]$) and has positive lower bound and upper bound on $[T_1, T]$ (or $[T, T_1]$). Hence $\frac{c(-\Psi)}{c(-\Psi_1)}$ has a positive lower bound on M. Then we know that for any compact subset $K \subset M \setminus E$, $e^{-\varphi_{\alpha}}c(-\Psi)$ has a positive lower bound on K.

Let $T_0 \in [-\infty, +\infty)$. Let $c(t) \in \mathcal{P}_{T_0, M, \Psi_0}$. It follows from Lemma 2.1 that we have the following lemma.

Lemma 2.3. Let (M, X, Z) satisfies condition (A). Let $B \in (0, +\infty)$ and $t_0 > T > T_0$ be arbitrarily given. Let f be a holomorphic (n, 0) form on $\{\Psi < -t_0\}$ such that

$$\int_{\{\Psi<-t_0\}} |f|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty, \qquad (2.1)$$

9

Then there exists a holomorphic (n, 0) form \tilde{F} on M such that

$$\int_{M} |\tilde{F} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi - \Psi + v_{t_0,B}(\Psi)} c(-v_{t_0,B}(\Psi))$$

$$\leq \left(\frac{1}{\delta}c(T)e^{-T} + \int_{T}^{t_0+B} c(t)e^{-t}dt\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0-B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi},$$
(2.2)

where $b_{t_0,B}(t) = \int_{-\infty}^t \frac{1}{B} \mathbb{I}_{\{-t_0 - B < s < -t_0\}} ds$ and $v_{t_0,B}(t) = \int_{-t_0}^t b_{t_0,B}(s) ds - t_0.$

Proof. It follows from inequality (2.1) and $c(t)e^{-t}$ is decreasing with respect to t that

$$\int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi} < +\infty.$$

Note that $c(t) \in \mathcal{P}_{T_0,M,\Psi_0}$ and there exists a closed subset $E \subset Z \cap \{\Psi = -\infty\}$ such that $e^{-\varphi}c(-\Psi)$ has locally positive lower bound on $M \setminus E$. For any compact subset $K \subset M \setminus E$, we obtain from inequality (2.1) that

$$\int_{K \cap \{\Psi < -t_0\}} |f|^2 < +\infty.$$

As (M, X, Z) satisfies condition (A), $M \setminus (Z \cup X)$ is a weakly pseudoconvex Kähler manifold. It follows from Lemma 2.1 that there exists a holomorphic (n, 0) form \tilde{F}_Z on $M \setminus (Z \cup X)$ such that

$$\int_{M \setminus (Z \cup X)} |\tilde{F}_Z - (1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq (\frac{1}{\delta} c(T) e^{-T} + \int_T^{t_0+B} c(s) e^{-s} ds) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0-B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}.$$

For any $z \in ((Z \cup X) \setminus E)$, there exists an open neighborhood V_z of z such that $V_z \in M \setminus E$.

Note that $c(t)e^{-t}$ is decreasing with respect to t and $v_{t_0,B}(\Psi) \geq \Psi$, we have

$$\begin{split} &\int_{M \setminus (Z \cup X)} |\tilde{F}_Z - (1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi} c(-\Psi) \\ &\leq \int_{M \setminus (Z \cup X)} |\tilde{F}_Z - (1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) < +\infty. \end{split}$$

Note that $\varphi = \varphi_{\alpha} + (1 + \delta)2 \log |F|$ on $\{\Psi < -t_0\}$. We have

$$\int_{\{\Psi < -t_0\}} |fF^{1+\delta}|^2 e^{-\varphi} c(-\Psi) = \int_{\{\Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty.$$

Note that there exists a positive number C > 0 such that $c(-\Psi)e^{-\varphi} > C$ on V_z . Then we have

$$\begin{split} &\int_{V_z \setminus (Z \cup X)} |\tilde{F}_Z|^2 \\ \leq &\frac{1}{C} \int_{V_z \setminus (Z \cup X)} |\tilde{F}_Z|^2 c(-\Psi) e^{-\varphi} \\ \leq &\frac{2}{C} \Big(\int_{V_z \setminus (Z \cup X)} |\tilde{F}_Z - (1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi} c(-\Psi) \\ &+ \int_{V_z \setminus (Z \cup X)} |(1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi} c(-\Psi) \Big) \\ \leq &\frac{2}{C} \Big(\int_{M \setminus (Z \cup X)} |\tilde{F}_Z - (1 - b_{t_0,B}(\Psi)) f F^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) \\ &+ \int_{\{\Psi < -t_0\}} |f F^{1+\delta}|^2 e^{-\varphi} c(-\Psi) \Big) < +\infty. \end{split}$$

As $Z \cup X$ is locally negligible with respect to L^2 holomorphic function, we can find a holomorphic extension \tilde{F}_E of \tilde{F}_Z from $M \setminus (Z \cup X)$ to $M \setminus E$ such that

$$\int_{M\setminus E} |\tilde{F}_E - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq (\frac{1}{\delta}c(T)e^{-T} + \int_T^{t_0+B} c(s)e^{-s}ds) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0-B<\Psi<-t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}.$$

Note that $E \subset \{\Psi = -\infty\} \subset \{\Psi < -t_0\}$ and $\{\Psi < -t_0\}$ is open, then for any

There exists an open neighborhood U_z of z such that $U_z \in {\Psi < -t_0}$. As $v_{t_0,B}(t) \ge -t_0 - \frac{B}{2}$, we have $c(-v_{t_0,B}(\Psi))e^{v_{t_0,B}(\Psi)} \ge c(t_0 + \frac{B}{2})e^{-t_0 - \frac{B}{2}} > 0$. Note that $\varphi + \Psi$ is plurisubharmonic on M. Thus we have

$$\int_{U_z \setminus E} |\tilde{F}_E - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2$$

$$\leq \frac{1}{C_1} \int_{U_z \setminus E} |\tilde{F}_E - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) < +\infty,$$

where C_1 is some positive number.

As $U_z \in \{\Psi < -t_0\}$, we have

$$\int_{U_z \setminus E} |(1 - b_{t_0, B}(\Psi)) f F^{1+\delta}|^2 \le \left(\sup_{U_z} |F^{1+\delta}|^2 \right) \int_{U_z} |f|^2 < +\infty$$

Hence we have

$$\int_{U_z \setminus E} |\tilde{F}_E|^2 < +\infty.$$

As E is contained in some analytic subset of M, we can find a holomorphic extension \tilde{F} of \tilde{F}_E from $M \setminus E$ to M such that

$$\int_{M} |\tilde{F} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi - \Psi + v_{t_0,B}(\Psi)} c(-v_{t_0,B}(\Psi)) \\
\leq \left(\frac{1}{\delta}c(T)e^{-T} + \int_{T}^{t_0+B} c(t)e^{-t}dt\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0-B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi}.$$
(2.3)

Lemma 2.3 is proved.

Let $T_0 \in [-\infty, +\infty)$. Let $c(t) \in \mathcal{P}_{T_0, M, \Psi_0}$. Using Lemma 2.3, we have the following lemma, which will be used to prove Theorem 1.2.

Lemma 2.4. Let (M, X, Z) satisfies condition (A). Let $B \in (0, +\infty)$ and $t_0 > t_1 \ge T > T_0$ be arbitrarily given. Let f be a holomorphic (n, 0) form on $\{\Psi < -t_0\}$ such that

$$\int_{\{\Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha}} c(-\Psi) < +\infty,$$
(2.4)

Then there exists a holomorphic (n,0) form \tilde{F} on $\{\Psi < -t_1\}$ such that

$$\int_{\{\Psi < -t_1\}} |\tilde{F} - (1 - b_{t_0,B}(\Psi))f|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq \left(\int_{t_1}^{t_0 + B} c(s)e^{-s}ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi},$$

where $b_{t_0,B}(t) = \int_{-\infty}^{t} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < s < -t_0\}} ds, \ v_{t_0,B}(t) = \int_{-t_0}^{t} b_{t_0,B}(s) ds - t_0.$

Proof of Lemma 2.4. Denote that

$$\tilde{\varphi} := \varphi_{\alpha} + (1+\delta) \max\{\psi + t_1, 2\log|F|\}$$

and

$$\tilde{\Psi} := \min\{\psi - 2\log|F|, -t_1\}.$$

As $t_0 > t_1 \ge T$, we have $\{\Psi < -t_0\} = \{\tilde{\Psi} < -t_0\}$. It follows from inequality (2.4) and Lemma 2.3 that there exists a holomorphic function \tilde{F}_{δ} on M such that

$$\begin{split} &\int_{M} |\tilde{F}_{\delta} - (1 - b_{t_0,B}(\tilde{\Psi})) f F^{1+\delta}|^2 e^{-\tilde{\varphi} + v_{t_0,B}(\tilde{\Psi}) - \tilde{\Psi}} c(-v_{t_0,B}(\tilde{\Psi})) \\ &\leq \left(\frac{1}{\delta} c(t_1) e^{-t_1} + \int_{t_1}^{t_0+B} c(s) e^{-s} ds\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0-B < \tilde{\Psi} < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \tilde{\Psi}}. \end{split}$$

Note that on $\{\Psi < -t_1\}$, we have $\Psi = \tilde{\Psi} = \psi - 2 \log |F|$ and $\tilde{\varphi} = \varphi = \varphi_{\alpha} + (1 + \delta)2 \log |F|$. Hence

$$\begin{split} &\int_{\{\Psi < -t_1\}} |\tilde{F}_{\delta} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) \\ &= \int_{\{\Psi < -t_1\}} |\tilde{F}_{\delta} - (1 - b_{t_0,B}(\tilde{\Psi}))fF^{1+\delta}|^2 e^{-\tilde{\varphi} + v_{t_0,B}(\tilde{\Psi}) - \tilde{\Psi}} c(-v_{t_0,B}(\tilde{\Psi})) \\ &\leq \int_M |\tilde{F}_{\delta} - (1 - b_{t_0,B}(\tilde{\Psi}))fF^{1+\delta}|^2 e^{-\tilde{\varphi} + v_{t_0,B}(\tilde{\Psi}) - \tilde{\Psi}} c(-v_{t_0,B}(\tilde{\Psi})) \\ &\leq \left(\frac{1}{\delta} c(t_1) e^{-t_1} + \int_{t_1}^{t_0 + B} c(s) e^{-s} ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \tilde{\Psi} < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \tilde{\Psi}} \\ &= \left(\frac{1}{\delta} c(t_1) e^{-t_1} + \int_{t_1}^{t_0 + B} c(s) e^{-s} ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi} < +\infty. \end{split}$$
(2.5)

Let $F_{\delta} := \frac{\tilde{F}_{\delta}}{F^{\delta}}$ be a holomorphic function on $\{\Psi < -t_1\}$. Then it follows from (2.5) that

$$\int_{\{\Psi < -t_1\}} |F_{\delta} - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_{\alpha} + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \\
\leq \left(\frac{1}{\delta}c(t_1)e^{-t_1} + \int_{t_1}^{t_0 + B} c(s)e^{-s}ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi}.$$
(2.6)

Let K be any compact subset of $M \setminus E$. Note that $\inf_K e^{-\varphi_{\alpha} + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \ge \left(c(t_0 + \frac{2}{B})e^{-t_0 - \frac{2}{B}}\right) \inf_K e^{-\varphi_{\alpha} - \psi} > 0$. It follows from (2.6) that we have

$$\sup_{\delta} \int_{\{\Psi < -t_1\} \cap K} |F_{\delta} - (1 - b_{t_0,B}(\Psi))fF|^2 < +\infty.$$

We also note that

$$\int_{\{\Psi < -t_1\} \cap K} |(1 - b_{t_0, B}(\Psi))fF|^2 \le \left(\sup_K |F|^2\right) \int_{\{\Psi < -t_0\} \cap K} |f|^2 < +\infty.$$

Then we know that

$$\sup_{\delta} \int_{\{\Psi < -t_1\} \cap K} |F_{\delta}|^2 < +\infty,$$

and there exists a subsequence of $\{F_{\delta}\}$ (also denoted by F_{δ}) compactly convergent to a holomorphic (n, 0) form \tilde{F}_1 on $\{\Psi < -t_1\}\setminus E$. It follows from Fatou's Lemma and inequality (2.6) that we have

$$\begin{split} &\int_{\{\Psi < -t_1\} \setminus E} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \\ &\leq \liminf_{\delta \to +\infty} \int_{\{\Psi < -t_1\} \setminus E} |F_\delta - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \\ &\leq \liminf_{\delta \to +\infty} \int_{\{\Psi < -t_1\}} |F_\delta - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \\ &\leq \liminf_{\delta \to +\infty} \left(\frac{1}{\delta} c(t_1) e^{-t_1} + \int_{t_1}^{t_0 + B} c(s) e^{-s} ds \right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi} \\ &\leq \left(\int_{t_1}^{t_0 + B} c(s) e^{-s} ds \right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}. \end{split}$$

$$(2.7)$$

Note that $E \subset \{\Psi = -\infty\} \subset \{\Psi < -t_1\}$ and $\{\Psi < -t_1\}$ is open, then for any $z \in E$, there exists an open neighborhood U_z of z such that $U_z \Subset \{\Psi < -t_1\}$. Note that $\varphi_{\alpha} + \psi$ is plurisubharmonic function on M. As $v_{t_0,B}(t) \ge -t_0 - \frac{B}{2}$, we have $c(-v_{t_0,B}(\Psi_1))e^{v_{t_0,B}(\Psi_1)} \ge c(t_0 + \frac{B}{2})e^{-t_0 - \frac{B}{2}} > 0$. Thus by (2.7), we have

$$\int_{U_z \setminus E} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))fF|^2$$

$$\leq \frac{1}{C_1} \int_{U_z \setminus E} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) < +\infty,$$

where $C_1 := c(t_0 + \frac{B}{2})e^{-t_0 - \frac{B}{2}} \inf_{U_z} e^{-\varphi_\alpha - \psi}$ is some positive number. As $U_z \subseteq \{\Psi < -t_1\}$, we have

$$\int_{U_z \setminus E} |(1 - b_{t_0, B}(\Psi))fF|^2 \le \left(\sup_{U_z} |F|^2\right) \int_{U_z} |f|^2 < +\infty.$$

Hence we have

$$\int_{U_z \setminus E} |\tilde{F}_1|^2 < +\infty.$$

As E is contained in some analytic subset of M, we can find a holomorphic extension \tilde{F}_0 of \tilde{F}_1 from $\{\Psi < -t_1\} \setminus E$ to $\{\Psi < -t_1\}$ such that

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_0 - (1 - b_{t_0,B}(\Psi))fF|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \psi} c(-v_{t_0,B}(\Psi)) \\
\leq \left(\int_{t_1}^{t_0 + B} c(s)e^{-s}ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}.$$
(2.8)

Denote $\tilde{F} := \frac{\tilde{F}_0}{F}$. Note that on $\{\Psi < -t_1\}$, we have $\Psi = \psi - 2 \log |F|$. It follows from (2.8) that we have

$$\int_{\{\Psi < -t_1\}} |\tilde{F} - (1 - b_{t_0,B}(\Psi))f|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq \left(\int_{t_1}^{t_0 + B} c(s)e^{-s}ds\right) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}.$$

Lemma 2.4 is proved.

2.2. Properties of \mathcal{O}_{M,z_0} -module J_{z_0} . In this section, we present some properties of \mathcal{O}_{M,z_0} -module J_{z_0} .

Since the case is local, we assume that F is a holomorphic function on a pseudoconvex domain $D \subset \mathbb{C}^n$ containing the origin $o \in \mathbb{C}^n$. Let ψ be a plurisubharmonic function on D. Let φ_{α} be a Lebesgue measurable function on D such that $\psi + \varphi_{\alpha}$ is plurisubharmonic. Let $T_0 \in [-\infty, +\infty)$ and $T > T_0$ be a real number. Denote

$$\varphi_1 := 2 \max\{\psi + T, 2 \log |F|\}$$

and

$$\Psi := \min\{\psi - 2\log|F|, -T\}.$$

If F(z) = 0 for some $z \in D$, we set $\Psi(z) = -T$.

Let c(t) be a positive measurable function on $(T_0, +\infty)$ such that (1) $c(t)e^{-t}$ is decreasing with respect t;

(2) $c(-\Psi)e^{-\varphi_{\alpha}}$ has a positive lower bound on D.

Denote that $H_o := \{f_o \in J(\Psi)_o : \int_{\{\Psi < -t\} \cap V_0} |f|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty \text{ for some } t > T_0 \text{ and } V_0 \text{ is an open neighborhood of o} \}$ and $\mathcal{H}_o := \{(F, o) \in \mathcal{O}_{\mathbb{C}^n, o} : \int_{U_0} |F|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty \text{ for some open neighborhood } U_0 \text{ of } o\}.$

As $c(-\Psi)e^{-\varphi_{\alpha}}$ has a positive lower bound on D and $c(t)e^{-t}$ is decreasing with respect to t, we have $I(\Psi + \varphi_{\alpha})_o \subset H_o \subset I_o$. We also note that \mathcal{H}_o is an ideal of $\mathcal{O}_{\mathbb{C}^n,o}$.

Lemma 2.5. For any $f_o \in H_o$, there exist a pseudoconvex domain $D_0 \subset D$ containing o and a holomorphic function \tilde{F} on D_0 such that $(\tilde{F}, o) \in \mathcal{H}_o$ and

$$\int_{\{\Psi < -t_1\} \cap D_0} |\tilde{F} - fF^2| e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty,$$

for some $t_1 > T$.

Proof. It follows from $f_o \in H_o$ that there exists $t_0 > T > T_0$ and a pseudoconvex domain $D_0 \subset D$ containing o such that

$$\int_{\{\Psi < -t_0\} \cap D_0} |f|^2 e^{-\varphi_{\alpha}} c(-\Psi) < +\infty.$$
(2.9)

As $c(t)e^{-t}$ is decreasing with respect to t, it follows from inequality (2.9) that we have $\int_{D_0} \mathbb{I}_{\{-t_0-1<\Psi<-t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi} < +\infty$. As $c(-\Psi)e^{-\varphi_\alpha}$ has a positive lower bound on D, we have $\int_{\{\Psi<-t_0\}\cap D_0} |f|^2 < +\infty$. Then it follows from Lemma 2.1 that there exists a holomorphic function \tilde{F} on D_0 such that

$$\int_{D_0} |\tilde{F} - (1 - b_{t_0}(\Psi))fF^2|^2 e^{-\varphi_\alpha - \varphi_1 + v_{t_0}(\Psi) - \Psi} c(-v_{t_0}(\Psi))$$

$$\leq \left(c(T)e^{-T} + \int_T^{t_0 + 1} c(s)e^{-s}ds \right) \int_{D_0} \mathbb{I}_{\{-t_0 - 1 < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi},$$

where $b_{t_0}(t) = \int_{-\infty}^t \mathbb{I}_{\{-t_0 - 1 < s < -t_0\}} ds$, $v_{t_0}(t) = \int_{-t_0}^t b_{t_0}(s) ds - t_0$. Denote $C := c(T)e^{-T} + \int_T^{t_0 + B} c(s)e^{-s} ds$, we note that C is a positive number.

Note that $v_{t_0}(t) > -t_0 - 1$. We have $e^{v_{t_0}(\Psi)}c(-v_{t_0}(\Psi)) \ge c(t_0 + 1)e^{-(t_0+1)} > 0$. As $b_{t_0}(t) \equiv 0$ on $(-\infty, -t_0 - 1)$, we have

$$\int_{D_{0} \cap \{\Psi < -t_{0}-1\}} |\tilde{F} - fF^{2}|^{2} e^{-\varphi_{\alpha} - \varphi_{1} - \Psi} \\
\leq \frac{1}{c(t_{0}+1)e^{-(t_{0}+1)}} \int_{D_{0}} |\tilde{F} - (1 - b_{t_{0}}(\Psi))fF^{2}|^{2} e^{-\varphi_{\alpha} - \varphi_{1} - \Psi + v_{t_{0}}(\Psi)} c(-v_{t_{0}}(\Psi)) \\
\leq \frac{C}{c(t_{0}+1)e^{-(t_{0}+1)}} \int_{D_{0}} \mathbb{I}_{\{-t_{0}-1 < \Psi < -t_{0}\}} |f|^{2} e^{-\varphi_{\alpha} - \Psi} < +\infty.$$
(2.10)

Note that on $\{\Psi < -t_0\}, |F|^2 e^{-\varphi_1} = 1$. As $v_{t_0}(\Psi) \ge \Psi$, we have $c(-v_{t_0}(\Psi))e^{v_{t_0}(\Psi)} \ge c(-\Psi)e^{-\Psi}$. Hence we have

$$\begin{split} &\int_{D_0} |\tilde{F}|^2 e^{-\varphi_{\alpha}-\varphi_1} c(-\Psi) \\ &\leq 2 \int_{D_0} |\tilde{F} - (1 - b_{t_0}(\Psi)) f F^2|^2 e^{-\varphi_{\alpha}-\varphi_1} c(-\Psi) \\ &+ 2 \int_{D_0} |(1 - b_{t_0}(\Psi)) f F^2|^2 e^{-\varphi_{\alpha}-\varphi_1} c(-\Psi) \\ &\leq 2 \int_{D_0} |\tilde{F} - (1 - b_{t_0}(\Psi)) f F^2|^2 e^{-\varphi_{\alpha}-\varphi_1-\Psi+v_{t_0}(\Psi)} c(-v_{t_0}(\Psi)) \\ &+ 2 \int_{D_0 \cap \{\Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &< +\infty. \end{split}$$

Hence we know that $(\tilde{F}, o) \in \mathcal{H}_o$.

For any $(\tilde{F}, o) \in \mathcal{H}_o$ and $(\tilde{F}_1, o) \in \mathcal{H}_o$ such that $\int_{D_1 \cap \{\Psi < -t_1\}} |\tilde{F} - fF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$ and $\int_{D_1 \cap \{\Psi < -t_1\}} |\tilde{F}_1 - fF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, for some open neighborhood D_1 of o and $t_1 \geq T$, we have

$$\int_{D_1 \cap \{\Psi < -t_1\}} |\tilde{F}_1 - \tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

As $(\tilde{F}, o) \in \mathcal{H}_o$ and $(\tilde{F}_1, o) \in \mathcal{H}_o$, there exists a neighborhood D_2 of o such that

$$\int_{D_2} |\tilde{F}_1 - \tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty.$$
(2.11)

Note that we have $c(-\Psi)e^{\Psi} \ge c(t_1)e^{-t_1}$ on $\{\Psi \ge -t_1\}$. It follows from inequality (2.11) that we have

$$\int_{D_2 \cap \{\Psi \ge -t_1\}} |\tilde{F}_1 - \tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

Hence we have $(\tilde{F} - \tilde{F}_1, o) \in \mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$.

Thus it follows from Lemma 2.5 that there exists a map $\tilde{P}: H_o \to \mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$ given by

$$\tilde{P}(f_o) = [(\tilde{F}, o)]$$

for any $f_o \in H_o$, where (\tilde{F}, o) satisfies $(\tilde{F}, o) \in \mathcal{H}_o$ and $\int_{D_1 \cap \{\Psi < -t\}} |\tilde{F} - fF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, for some t > T and some open neighborhood D_1 of o, and $[(\tilde{F}, o)]$ is the equivalence class of (\tilde{F}, o) in $\mathcal{H}_o / \mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o$.

Proposition 2.6. \tilde{P} is an $\mathcal{O}_{\mathbb{C}^n,o}$ -module homomorphism and $Ker(\tilde{P}) = I(\varphi_{\alpha} + \Psi)_o$.

Proof. For any $f_o, g_o \in H_o$. Denote that $\tilde{P}(f_o) = [(\tilde{F}, o)], \ \tilde{P}(g_o) = [(\tilde{G}, o)]$ and $\tilde{P}(f_o + g_o) = [(\tilde{H}, o)].$

Note that there exists an open neighborhood D_1 of o and $t \geq T$ such that $\int_{D_1 \cap \{\Psi < -t\}} |\tilde{F} - fF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, $\int_{D_1 \cap \{\Psi < -t\}} |\tilde{G} - gF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, and $\int_{D_1 \cap \{\Psi < -t\}} |\tilde{H} - (f+g)F^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$. Hence we have

$$\int_{D_1 \cap \{\Psi < -t\}} |\tilde{H} - (\tilde{F} + \tilde{G})|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

As (\tilde{F}, o) , (\tilde{G}, o) and (\tilde{H}, o) belong to \mathcal{H}_o , there exists an open neighborhood $\tilde{D}_1 \subset D_1$ of o such that $\int_{\tilde{D}_1} |\tilde{H} - (\tilde{F} + \tilde{G})|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty$. As $c(t)e^{-t}$ is decreasing with respect to t, we have $c(-\Psi)e^{\Psi} \geq c(t)e^{-t}$ on $\{\Psi \geq -t\}$. Hence we have

$$\int_{\tilde{D}_1 \cap \{\Psi \ge -t\}} |\tilde{H} - (\tilde{F} + \tilde{G})|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} \le \frac{1}{c(t)e^{-t}} \int_{\tilde{D}_1 \cap \{\Psi \ge -t\}} |\tilde{H} - (\tilde{F} + \tilde{G})|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty$$

Thus we have $\int_{\tilde{D}_1} |\tilde{H} - (\tilde{F} + \tilde{G})|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, which implies that $\tilde{P}(f_o + g_o) = \tilde{P}(f_o) + \tilde{P}(g_o)$.

For any $(h, o) \in \mathcal{O}_{\mathbb{C}^n, o}$. Denote $\tilde{P}((hf)_o) = [(\tilde{F}_h, o)]$. Note that there exists an open neighborhood D_2 of o and $t \geq T$ such that $\int_{D_2 \cap \{\Psi < -t\}} |\tilde{F}_h - (hf)F^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$. It follows from $\int_{D_2 \cap \{\Psi < -t\}} |\tilde{F} - fF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$ and h is holomorphic on $\overline{D_2}$ (shrink D_2 if necessary) that $\int_{D_2 \cap \{\Psi < -t\}} |h\tilde{F} - hfF^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$. Then we have

$$\int_{D_2 \cap \{\Psi < -t\}} |\tilde{F}_h - h\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

Note that $(h\tilde{F}, o)$ and (\tilde{F}_h, o) belong to \mathcal{H}_o , we have $\int_{D_2} |\tilde{F}_h - h\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty$. As $c(t)e^{-t}$ is decreasing with respect to t, we have $c(-\Psi)e^{\Psi} \ge c(t)e^{-t}$ on $\{\Psi \ge -t\}$. Hence we have

$$\int_{D_2 \cap \{\Psi \ge -t\}} |\tilde{F}_h - h\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} \le \frac{1}{c(t)e^{-t}} \int_{D_2 \cap \{\Psi \ge -t\}} |\tilde{F}_h - h\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty.$$

Thus we have $\int_{D_2} |\tilde{F}_h - h\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty$, which implies that $\tilde{P}(hf_o) = (h, o)\tilde{P}(f_o)$.

Now we have proved that \tilde{P} is an $\mathcal{O}_{\mathbb{C}^n,o}$ -module homomorphism.

Next, we prove $Ker(\tilde{P}) = I(\varphi_{\alpha} + \Psi)_o$.

If $f_o \in I(\varphi_{\alpha} + \Psi)_o$. Denote $\tilde{P}(f_o) = [(\tilde{F}, o)]$. It follows from Lemma 2.5 that $(\tilde{F}, o) \in \mathcal{H}_o$ and there exists an open neighborhood D_3 of o and a real number $t_1 > T$ such that

$$\int_{\{\Psi < -t_1\} \cap D_3} |\tilde{F} - fF^2| e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

As $f_o \in I(\varphi_{\alpha} + \Psi)_o$, shrink D_3 and t_1 if necessary, we have

$$\int_{\{\Psi < -t_1\} \cap D_3} |\tilde{F}|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} \\
\leq 2 \int_{\{\Psi < -t_1\} \cap D_3} |\tilde{F} - fF^2|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} + 2 \int_{\{\Psi < -t_1\} \cap D_3} |fF^2|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} \\
\leq 2 \int_{\{\Psi < -t_1\} \cap D_3} |\tilde{F} - fF^2|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} + 2 \int_{\{\Psi < -t_1\} \cap D_3} |f|^2 e^{-\varphi_{\alpha} - \Psi} \\
< + \infty.$$
(2.12)

As $c(t)e^{-t}$ is decreasing with respect to t, $c(-\Psi)e^{\Psi} \ge C_0 > 0$ for some positive number C_0 on $\{\Psi \ge -t_1\}$. Then we have

$$\int_{\{\Psi \ge -t_1\} \cap D_3} |\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} \le \frac{1}{C_0} \int_{\{\Psi \ge -t_1\} \cap D_3} |\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty.$$
(2.13)

Combining inequality (2.12) and inequality (2.13), we know that $\tilde{F} \in \mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$, which means $\tilde{P}(f_o) = 0$ in $\mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$. Hence we know $I(\varphi_{\alpha} + \Psi)_o \subset Ker(\tilde{P})$.

If $f_o \in Ker(\tilde{P})$, we know $\tilde{F} \in \mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$. We can assume that \tilde{F} satisfies $\int_{D_4} |\tilde{F}|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} < +\infty$ for some open neighborhood D_4 of o. Then we have

$$\int_{\{\Psi < -t_1\} \cap D_4} |f|^2 e^{-\varphi_{\alpha} - \Psi} \\
= \int_{\{\Psi < -t_1\} \cap D_4} |fF^2|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} \\
\leq \int_{\{\Psi < -t_1\} \cap D_4} |\tilde{F}|^2 e^{-\varphi_{\alpha} - \varphi_1 - \Psi} + \int_{\{\Psi < -t_1\} \cap D_4} |\tilde{F} - fF^2| e^{-\varphi_{\alpha} - \varphi_1 - \Psi} \\
\leq + \infty.$$
(2.14)

By definition, we know $f_o \in I(\varphi_{\alpha} + \Psi)_o$. Hence $Ker(\tilde{P}) \subset I(\varphi_{\alpha} + \Psi)_o$. $Ker(\tilde{P}) = I(\varphi_{\alpha} + \Psi)_o$ is proved.

Now we can define an $\mathcal{O}_{\mathbb{C}^n,o}$ -module homomorphism $P: H_o/I(\varphi_{\alpha} + \Psi)_o \to \mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$ as follows,

$$P([f_o]) = \tilde{P}(f_o)$$

for any $[f_o] \in H_o/I(\varphi_{\alpha} + \Psi)_o$, where $f_o \in H_o$ is any representative of $[f_o]$. It follows from Proposition 2.6 that $P([f_o])$ is independent of the choices of the representatives of $[f_o]$.

Let $(\tilde{F}, o) \in \mathcal{H}_o$, i.e. $\int_U |\tilde{F}|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty$ for some neighborhood U of o. Note that $|F|^4 e^{-\varphi_1} \equiv 1$ on $\{\Psi < -T\}$. Hence we have $\int_{U \cap \{\Psi < -t\}} |\frac{\tilde{F}}{F^2}|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty$ for some t > T, i.e. $(\frac{\tilde{F}}{F^2})_o \in \mathcal{H}_o$. And if $(\tilde{F}, o) \in \mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o$, it is easy to verify that $(\frac{\tilde{F}}{F^2})_o \in I(\varphi_\alpha + \Psi)_o$. Hence we have an $\mathcal{O}_{\mathbb{C}^n,o}$ -module homomorphism $Q : \mathcal{H}_o/\mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o \to \mathcal{H}_o/I(\varphi_\alpha + \Psi)_o$ defined as follows,

$$Q([(\tilde{F},o)]) = [(\frac{\tilde{F}}{F^2})_o].$$

The above discussion shows that Q is independent of the choices of the representatives of $[(\tilde{F}, o)]$ and hence Q is well defined.

Proposition 2.7. $P: H_o/I(\varphi_{\alpha} + \Psi)_o \to \mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$ is an $\mathcal{O}_{\mathbb{C}^n,o}$ -module isomorphism and $P^{-1} = Q$.

Proof. It follows from Proposition 2.6 that we know P is injective.

Now we prove P is surjective.

For any $[(\tilde{F}, o)]$ in $\mathcal{H}_o/\mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o$. Let (\tilde{F}, o) be any representatives of $[(\tilde{F}, o)]$ in \mathcal{H}_o . Denote that $[(f_1)_o] := [(\frac{\tilde{F}}{F^2})_o] = Q([(\tilde{F}, o)])$. Let $(f_1)_o := (\frac{\tilde{F}}{F^2})_o \in \mathcal{H}_o$ be the representative of $[(f_1)_o]$. Denote $[(\tilde{F}_1, o)] := \tilde{P}((f_1)_o) = P([(f_1)_o])$. By the construction of \tilde{P} , we know that $(\tilde{F}_1, o) \in \mathcal{H}_o$ and

$$\int_{D_1 \cap \{\Psi < -t\}} |\tilde{F}_1 - f_1 F^2| e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty,$$

where t > T and D_1 is some neighborhood of o. Note that $(f_1)_o := (\frac{F}{F^2})_o$. Hence we have

$$\int_{D_1 \cap \{\Psi < -t\}} |\tilde{F}_1 - \tilde{F}| e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

It follows from $(\tilde{F}, o) \in \mathcal{H}_o$ and $(\tilde{F}_1, o) \in \mathcal{H}_o$ that there exists a neighborhood $D_2 \subset D_1$ of o such that

$$\int_{D_2} |\tilde{F} - \tilde{F}_1|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty$$

Note that on $\{\Psi \ge -t\}$, we have $c(-\Psi)e^{\Psi} \ge c(t)e^{-t} > 0$. Hence we have

$$\int_{D_2 \cap \{\Psi \ge -t\}} |\tilde{F} - \tilde{F}_1|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$

Thus we know that $(\tilde{F}_1 - \tilde{F}, o) \in \mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o$, i.e. $[(\tilde{F}, o)] = [(\tilde{F}_1, o)]$ in $\mathcal{H}_o/\mathcal{I}(\varphi_\alpha + \varphi_1 + \Psi)_o$. Hence we have $P \circ Q([(\tilde{F}, o)]) = [(\tilde{F}, o)]$, which implies that P is surjective.

We have proved that $P: H_o/I(\varphi_{\alpha} + \Psi)_o \to \mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$ is an $\mathcal{O}_{\mathbb{C}^n,o^-}$ module isomorphism and $P^{-1} = Q$.

We recall the following property of closedness of holomorphic functions on a neighborhood of o.

Lemma 2.8 (see [21]). Let N be a submodule of $\mathcal{O}^q_{\mathbb{C}^n,o}$, $1 \leq q < +\infty$, let $f_j \in \mathcal{O}_{\mathbb{C}^n}(U)^q$ be a sequence of q-tuples holomorphic in an open neighborhood U of the origin o. Assume that the f_j converge uniformly in U towards a q-tuples $f \in \mathcal{O}_{\mathbb{C}^n}(U)^q$, assume furthermore that all germs (f_j, o) belong to N. Then $(f, o) \in N$.

The following lemma shows the closedness of submodules of H_o .

Lemma 2.9. Let J_o be an $\mathcal{O}_{\mathbb{C}^n,o}$ -submodule of H_o such that $I(\varphi_{\alpha} + \Psi)_o \subset J_o$. Assume that $f_o \in J(\Psi)_o$. Let U_0 be a Stein open neighborhood of o. Let $\{f_j\}_{j\geq 1}$ be a sequence of holomorphic functions on $U_0 \cap \{\Psi < -t_j\}$ for any $j \geq 1$, where $t_j > T > T_0$. Assume that $t_0 := \lim_{j \to +\infty} t_j \in [T, +\infty)$,

$$\limsup_{j \to +\infty} \int_{U_0 \cap \{\Psi < -t_j\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) \le C < +\infty,$$
(2.15)

19

and $(f_j - f)_o \in J_o$. Assume that $e^{-\varphi_\alpha}c(-\Psi)$ has a positive lower bound on U_o . Then there exists a subsequence of $\{f_j\}_{j\geq 1}$ compactly convergent to a holomorphic function f_0 on $\{\Psi < -t_0\} \cap U_0$ which satisfies

$$\int_{U_0 \cap \{\Psi < -t_0\}} |f_0|^2 e^{-\varphi_\alpha} c(-\Psi) \le C,$$

and $(f_0 - f)_o \in J_o$.

Proof. It follows from $\limsup_{j \to +\infty} \int_{U_0 \cap \{\Psi < -t_j\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) \leq C < +\infty$ and $e^{-\varphi_\alpha} c(-\Psi)$ has a positive lower bound on U_o that

$$\limsup_{j \to +\infty} \int_{U_0 \cap \{\Psi < -t_j\}} |f_j|^2 < +\infty.$$

As $t_0 := \lim_{j \to +\infty} t_j \in [T, +\infty)$, by Montel's theorem, there exists a subsequence of $\{f_j\}_{j \ge 1}$ (also denoted by $\{f_j\}_{j \ge 1}$) compactly convergent to a holomorphic function f_0 on $\{\Psi < -t_0\} \cap U_0$. It follows from Fatou's Lemma that

$$\int_{U_0 \cap \{\Psi < -t_0\}} |f_0|^2 e^{-\varphi_\alpha} c(-\Psi) \le \liminf_{j \to +\infty} \int_{U_0 \cap \{\Psi < -t_j\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) \le C.$$

Now we prove $(f_0 - f)_o \in J_o$. We firstly recall some constructions in Lemma 2.5.

As $c(t)e^{-t}$ is decreasing with respect to t, it follows from inequality (2.15) that we have $\sup_{j\geq 1} \left(\int_{U_0} \mathbb{I}_{\{-t_j-1<\Psi<-t_j\}} |f_j|^2 e^{-\varphi_\alpha - \Psi} \right) < +\infty$. As $c(-\Psi)e^{-\varphi_\alpha}$ has a positive lower bound on U_0 , we have $\sup_{j\geq 1} \left(\int_{\{\Psi<-t_j\}\cap U_0} |f_j|^2 \right) < +\infty$. Then it follows from Lemma 2.1 that there exists a holomorphic function \tilde{F}_j on U_0 such that

$$\int_{U_0} |\tilde{F}_j - (1 - b_{t_j}(\Psi)) f_j F^2|^2 e^{-\varphi_\alpha - \varphi_1 + v_{t_j}(\Psi) - \Psi} c(-v_{t_j}(\Psi))$$

$$\leq \left(c(T) e^{-T} + \int_T^{t_j + 1} c(s) e^{-s} ds \right) \int_{U_0} \mathbb{I}_{\{-t_0 - 1 < \Psi < -t_0\}} |f_j|^2 e^{-\varphi_\alpha - \Psi},$$
(2.16)

where $b_{t_j}(t) = \int_{-\infty}^t \mathbb{I}_{\{-t_j-1 < s < -t_j\}} ds$, $v_{t_j}(t) = \int_{-t_j}^t b_{t_j}(s) ds - t_j$. Denote $C_j := c(T)e^{-T} + \int_T^{t_j+1} c(s)e^{-s} ds$. As $t_0 := \lim_{j \to +\infty} t_j \in [T, +\infty)$, we can assume that there exists a positive number $C_0 < +\infty$ such that $C_j \leq C_0$ for all $j \geq 1$.

Note that $v_{t_j}(t) > -t_j - 1$. We have $e^{v_{t_j}(\Psi) - \Psi} c(-v_{t_j}(\Psi)) \ge c(t_j + 1)e^{-(t_j + 1)} > 0$. As $b_{t_j}(t) \equiv 0$ on $(-\infty, -t_j - 1)$, we have

$$\int_{U_{0} \cap \{\Psi < -t_{j}-1\}} |\tilde{F}_{j} - f_{j}F^{2}|^{2}e^{-\varphi_{\alpha}-\varphi_{1}-\Psi} \\
\leq \frac{1}{c(t_{j}+1)e^{-(t_{j}+1)}} \int_{U_{0}} |\tilde{F}_{j} - (1 - b_{t_{j}}(\Psi))f_{j}F^{2}|^{2}e^{-\varphi_{\alpha}-\varphi_{1}-\Psi+v_{t_{j}}(\Psi)}c(-v_{t_{j}}(\Psi)) \\
\leq \frac{C}{c(t_{j}+1)e^{-(t_{j}+1)}} \int_{U_{0}} \mathbb{I}_{\{-t_{j}-1<\Psi<-t_{j}\}} |f_{j}|^{2}e^{-\varphi_{\alpha}-\Psi} < +\infty.$$
(2.17)

Note that $|F^2|^2 e^{-\varphi_1} = 1$ on $\{\Psi < -t_j\}$. As $v_{t_j}(\Psi) \ge \Psi$, we have $c(-v_{t_j}(\Psi))e^{v_{t_j}(\Psi)} \ge c(-\Psi)e^{-\Psi}$. Hence we have

$$\int_{U_{0}} |\tilde{F}_{j}|^{2} e^{-\varphi_{\alpha}-\varphi_{1}} c(-\Psi)
\leq 2 \int_{U_{0}} |\tilde{F}_{j} - (1 - b_{t_{j}}(\Psi))f_{j}F^{2}|^{2} e^{-\varphi_{\alpha}-\varphi_{1}} c(-\Psi)
+ 2 \int_{U_{0}} |(1 - b_{t_{j}}(\Psi))f_{j}F^{2}|^{2} e^{-\varphi_{\alpha}-\varphi_{1}} c(-\Psi)
\leq 2 \int_{U_{0}} |\tilde{F}_{j} - (1 - b_{t_{j}}(\Psi))f_{j}F^{2}|^{2} e^{-\varphi_{\alpha}-\varphi_{1}-\Psi+v_{t_{j}}(\Psi)} c(-v_{t_{j}}(\Psi))
+ 2 \int_{U_{0} \cap \{\Psi < -t_{j}\}} |f_{j}|^{2} e^{-\varphi_{\alpha}} c(-\Psi)
< + \infty.$$
(2.18)

Hence we know that $(\tilde{F}_j, o) \in \mathcal{H}_o$.

It follows from inequality (2.15), $\sup_{j\geq 1} \left(\int_{U_0} \mathbb{I}_{\{-t_j-1<\Psi<-t_j\}} |f_j|^2 e^{-\varphi_\alpha - \Psi} \right) < +\infty$ and inequality (2.18) that we actually have $\sup_j \left(\int_{U_0} |\tilde{F}_j|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) \right) < +\infty$. Note that φ_1 is a plurisubharmonic function and $c(-\Psi)e^{-\varphi_\alpha}$ has a positive lower bound on U_0 . We have (shrink U_0 if necessary)

$$\sup_{j} \left(\int_{U_0} |\tilde{F}_j|^2 \right) < +\infty.$$

Hence we know there exists a subsequence of $\{\tilde{F}_j\}_{j\geq 1}$ (also denoted by $\{\tilde{F}_j\}_{j\geq 1}$) compactly convergent to a holomorphic function \tilde{F}_0 on U_0 . It follows from Fatou's Lemma and inequality (2.18) that

$$\int_{U_0} |\tilde{F}_0|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) \le \liminf_{j \to +\infty} \int_{U_0} |\tilde{F}_j|^2 e^{-\varphi_\alpha - \varphi_1} c(-\Psi) < +\infty.$$
(2.19)

As f_j converges to f_0 , it follows from Fatou's Lemma and inequality (2.16) that

$$\int_{U_0} |\tilde{F}_0 - (1 - b_{t_0}(\Psi)) f_0 F^2|^2 e^{-\varphi_\alpha - \varphi_1 + v_{t_0}(\Psi) - \Psi} c(-v_{t_0}(\Psi))$$

$$\leq \liminf_{j \to +\infty} \int_{U_0} |\tilde{F}_j - (1 - b_{t_j}(\Psi)) f_j F^2|^2 e^{-\varphi_\alpha - \varphi_1 + v_{t_j}(\Psi) - \Psi} c(-v_{t_j}(\Psi))$$

$$< +\infty,$$

which implies that

$$\int_{U_0 \cap \{\Psi < -t_0 - 1\}} |\tilde{F}_0 - f_0 F^2|^2 e^{-\varphi_\alpha - \varphi_1 - \Psi} < +\infty.$$
(2.20)

It follows from inequality (2.17), inequality (2.18), inequality (2.19), inequality (2.20) and definition of $P: H_o/I(\varphi_{\alpha} + \Psi)_o \to \mathcal{H}_o/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o$ that for any $j \geq 0$, we have

$$P([(f_j)_o]) = [(\tilde{F}_j, o)].$$

As $(f_j - f)_o \in J_o$ for any $j \ge 1$, we have $(f_j - f_1)_o \in J_o$ for any $j \ge 1$. It follows from Proposition 2.7 that there exists an ideal \tilde{J} of $\mathcal{O}_{\mathbb{C}^n,o}$ such that $\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o \subset \tilde{J} \subset \mathcal{H}_o \text{ and } \tilde{J}/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o = \operatorname{Im}(P|_{J_o/I(\varphi_{\alpha} + \Psi)_o}). \text{ It follows from } (f_j - f_1)_o \in J_o \text{ and } P([(f_j)_o]) = [(F_j, o)] \text{ for any } j \ge 1 \text{ that we have } (\tilde{F}_j - \tilde{F}_1) \in \tilde{J},$

for any $j \ge 1$.

As \tilde{F}_j compactly converges to \tilde{F}_0 , using Lemma 2.8, we obtain that $(\tilde{F}_0 - \tilde{F}_1, o) \in \tilde{J}$. Note that P is an $\mathcal{O}_{\mathbb{C}^n,o}$ -module isomorphism and $\tilde{J}/\mathcal{I}(\varphi_{\alpha} + \varphi_1 + \Psi)_o = \text{Im}(P|_{J_o/I(\varphi_{\alpha} + \Psi)_o})$. We have $(f_0 - f_1)_o \in J_o$, which implies that $(f_0 - f)_o \in J_o$. Lemma 2.9 is proved.

Let φ_{α} be a plurisubharmonic function on D, and let $c \equiv 1$. Note that $H_o = I(\varphi_{\alpha})_o$ and $\mathcal{H}_o = \mathcal{I}(\varphi_{\alpha} + \varphi_1)_o$. We know that $I(a\Psi + \varphi_{\alpha})_o \subset I(a'\Psi + \varphi_{\alpha})_o$ for any $0 \leq a' < a < +\infty$. Denote that $I_+(a\Psi + \varphi_{\alpha})_o := \bigcup_{p>a} I(p\Psi + \varphi_{\alpha})_o$ is an $\mathcal{O}_{\mathbb{C}^n,o}$ -submodule of H_o , where $a \geq 0$.

Lemma 2.10. There exists a' > a such that $I(a'\Psi + \varphi_{\alpha})_o = I_+(a\Psi + \varphi_{\alpha})_o$ for any $a \ge 0$.

Proof. By the definition of $I_+(a\Psi + \varphi_\alpha)_o$, we know $I(p\Psi + \varphi_\alpha)_o \subset I_+(a\Psi + \varphi_\alpha)_o$ for any p > a. It suffices to prove that there exists a' > a such that $I_+(a\Psi + \varphi_\alpha)_o \subset I(a'\Psi + \varphi_\alpha)_o$.

Let k > a be an integer. Denote that $\tilde{\varphi}_1 := k\varphi_1 = 2 \max\{k\psi + kT, 2\log|F^k|\}$ and $\tilde{\Psi} := k\Psi = \min\{k\psi - 2\log|F^k|, -kT\}$. Proposition 2.7 shows that there exists an $\mathcal{O}_{\mathbb{C}^n,o}$ -module isomorphism P from $I(\varphi_\alpha)_o/I(\varphi_\alpha + \tilde{\Psi})_o \to \mathcal{I}(\varphi_\alpha + \varphi_1)_o/\mathcal{I}(\varphi_\alpha + \tilde{\varphi}_1 + \tilde{\Psi})_o)$, which implies that there exists an ideal K_p of $\mathcal{O}_{\mathbb{C}^n,o}$ such that

$$P(I(\varphi_{\alpha} + p\Psi)_o / I(\varphi_{\alpha} + \Psi)_o) = K_p / \mathcal{I}(\varphi_{\alpha} + \tilde{\varphi}_1 + \Psi)_o,$$

where $p \in (0, k)$. Denote that

$$L := \bigcup_{a$$

be an ideal of $\mathcal{O}_{\mathbb{C}^n,o}$. Hence $P|_{I_+(a\Psi+\varphi_\alpha)_o/I(\varphi_\alpha+\tilde{\Psi})_o}$ is an $\mathcal{O}_{\mathbb{C}^n,o}$ -module isomorphism from $I_+(a\Psi+\varphi_\alpha)_o/I(\varphi_\alpha+\tilde{\Psi})_o$ to $L/\mathcal{I}(\varphi_\alpha+\tilde{\varphi}_1+\tilde{\Psi})_o$. As $\mathcal{O}_{\mathbb{C}^n,o}$ is a Noetherian ring (see [38]), we get that L is finitely generated. Hence there exists a finite set $\{(f_1)_o,\ldots,(f_m)_o\} \subset I_+(a\Psi+\varphi_\alpha)_o$, which satisfies that for any $f_o \in I_+(a\Psi+\varphi_\alpha)_o$ there exists $(h_j,o) \in \mathcal{O}_{\mathbb{C}^n,o}$ for any $1 \leq j \leq m$ such that

$$f_o - \sum_{j=1}^{m} (h_j, o) \cdot (f_j)_o \in I(\varphi_\alpha + \tilde{\Psi})_o.$$

By the definition of $I_+(a\Psi+\varphi_\alpha)_o$, there exists $a' \in (a,k)$ such that $\{(f_1)_o, \ldots, (f_m)_o\} \subset I(a'\Psi+\varphi_\alpha)_o$. As $(h_j, o) \cdot (f_j)_o \in I(a'\Psi+\varphi_\alpha)_o$ for any $1 \leq j \leq m$ and $I(\varphi_\alpha + \tilde{\Psi})_o = I(k\Psi+\varphi_\alpha)_o \subset I(a'\Psi+\varphi_\alpha)_o$, we obtain that $I_+(a\Psi+\varphi_\alpha)_o \subset I(a'\Psi+\varphi_\alpha)_o$. Thus, Lemma 2.10 holds.

3. Properties of G(t)

Following the notations in Section 1.1, we present some properties of the function G(t) in this section.

Lemma 3.1 (see [28]). Let M be a complex manifold. Let S be an analytic subset of M. Let $\{g_j\}_{j=1,2,...}$ be a sequence of nonnegative Lebesgue measurable functions on M, which satisfies that g_j are almost everywhere convergent to g on M when $j \to +\infty$, where g is a nonnegative Lebesgue measurable function on M. Assume that for any compact subset K of $M \setminus S$, there exist $s_K \in (0, +\infty)$ and $C_K \in (0, +\infty)$ such that

$$\int_{K} g_j^{-s_K} dV_M \le C_K$$

for any j, where dV_M is a continuous volume form on M.

Let $\{F_j\}_{j=1,2,...}$ be a sequence of holomorphic (n,0) form on M. Assume that $\liminf_{j\to+\infty} \int_M |F_j|^2 g_j \leq C$, where C is a positive constant. Then there exists a subsequence $\{F_{j_l}\}_{l=1,2,...}$, which satisfies that $\{F_{j_l}\}$ is uniformly convergent to a holomorphic (n,0) form F on M on any compact subset of M when $l \to +\infty$, such that

$$\int_M |F|^2 g \le C$$

Let $c(t) \in P_{T,M,\Psi}$. The following lemma will be used to discuss the convergence property of holomorphic forms on $\{\Psi < -t\}$.

Lemma 3.2. Let f be a holomorphic (n, 0) form on $\{\Psi < -\hat{t}_0\} \cap V$, where $V \supset Z_0$ is an open subset of M and $\hat{t}_0 > T$ is an real number. For any $z_0 \in Z_0$, let J_{z_0} be an \mathcal{O}_{M,z_0} -submodule of $J(\Psi)_{z_0}$ such that $I(\Psi + \varphi_\alpha)_{z_0} \subset J_{z_0}$.

Let $\{f_j\}_{j\geq 1}$ be a sequence of holomorphic (n,0) form on $\{\Psi < -t_j\}$. Assume that $t_0 := \lim_{j \to +\infty} t_j \in [T, +\infty)$,

$$\limsup_{j \to +\infty} \int_{\{\Psi < -t_j\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) \le C < +\infty, \tag{3.1}$$

and $(f_j - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Then there exists a subsequence of $\{f_j\}_{j \in \mathbb{N}^+}$ compactly convergent to a holomorphic (n, 0) form f_0 on $\{\Psi < -t_0\}$ which satisfies

$$\int_{\{\Psi < -t_0\}} |f_0|^2 e^{-\varphi_\alpha} c(-\Psi) \le C,$$

and $(f_0 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$.

Proof. As $e^{-\varphi_{\alpha}}c(-\Psi)$ has positive lower bound on any compact subset of $M \setminus Z$, where Z is some analytic subset of M, it follows from Lemma 3.1 that there exists a subsequence of $\{f_j\}_{j \in \mathbb{N}^+}$ (also denoted by $\{f_j\}_{j \in \mathbb{N}^+}$) that compactly convergent to a holomorphic (n, 0) form f_0 on $\{\Psi < -t_0\}$ which satisfies

$$\int_{\{\Psi < -t_0\}} |f_0|^2 e^{-\varphi} c(-\Psi) \le \liminf_{j \to +\infty} \int_{\{\Psi < -t_j\}} |f_j|^2 e^{-\varphi} c(-\Psi) \le C.$$

Next we prove $(f_0 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$.

For any $z_0 \in Z_0 \cap \{\Psi = -\infty\}$, we know that $\{f_j\}_{j\geq 0}$ and f are holomorphic (n,0) forms on some neighborhood U_{z_0} of z_0 . It is also easy to verify that $J(\Psi)_{z_0} = I_{z_0} = \mathcal{O}_{M,z_0}$ and J_{z_0} is an \mathcal{O}_{M,z_0} -submodule of \mathcal{O}_{M,z_0} . As $J_{z_0} \subset \mathcal{O}_{M,z_0}$ is an \mathcal{O}_{M,z_0} -submodule, it follows from Lemma 2.8, $(f_j - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ and $\{f_j\}_{j\in\mathbb{N}^+}$ compactly converges to f_0 that we know $(f_0 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0 \cap \{\Psi = -\infty\}$.

Let $z_0 \in Z_0 \setminus \{\Psi = -\infty\}$. As $\limsup_{j \to +\infty} \int_{\{\Psi < -t_j\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) \le C < +\infty$, we

know $(f_j - f_1) \in H_{z_0}$. The definition of H_{z_0} can be referred to Section 2.2. It follows from $(f_j - f)_{z_0} \in J_{z_0}$ that we know $(f_j - f_1)_{z_0} \in J_{z_0}$. Hence we have $(f_j - f_1) \in H_{z_0} \cap J_{z_0}$. We note that $e^{-\varphi_{\alpha}}c(-\Psi)$ has a positive lower bound on

some open neighborhood of z_0 . It follows from inequality (3.1), $(f_j - f_1) \in H_{z_0} \cap J_{z_0}$, the uniqueness of limit function and Lemma 2.9 that we know $(f_0 - f_1)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes (H_{z_0} \cap J_{z_0})$. Hence we know that $(f_0 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0 \setminus \{\Psi = -\infty\}$.

Now we have $(f_0 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Lemma 3.2 is proved.

Lemma 3.3. Let $t_0 > T$. The following two statements are equivalent, (1) $G(t_0) = 0$;

(2)
$$f_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$$
, for any $z_0 \in Z_0$

Proof. If $f_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$, then take $\tilde{f} \equiv 0$ in the definition of G(t) and we get $G(t_0) \equiv 0$.

If $G(t_0) = 0$, by definition, there exists a sequence of holomorphic (n, 0) forms $\{f_j\}_{j \in \mathbb{Z}^+}$ on $\{\Psi < -t_0\}$ such that

$$\lim_{j \to +\infty} \int_{\{\Psi < -t_0\}} |f_j|^2 e^{-\varphi} c(-\Psi) = 0, \qquad (3.2)$$

and $(f_j - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$ and $j \geq 1$. It follows from Lemma 3.2 that there exists a subsequence of $\{f_j\}_{j \in \mathbb{N}^+}$ compactly convergent to a holomorphic (n, 0) form f_0 on $\{\Psi < -t_0\}$ which satisfies

$$\int_{\{\Psi < -t_0\}} |f_0|^2 e^{-\varphi_\alpha} c(-\Psi) = 0$$

and $(f_0-f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. It follows from $\int_{\{\Psi < -t_0\}} |f_0|^2 e^{-\varphi_\alpha} c(-\Psi) = 0$ that we know $f_0 \equiv 0$. Hence we have $f_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Statement (2) is proved.

The following lemma shows the existence and uniqueness of the holomorphic (n, 0) form related to G(t).

Lemma 3.4. Assume that $G(t) < +\infty$ for some $t \in [T, +\infty)$. Then there exists a unique holomorphic (n, 0) form F_t on $\{\Psi < -t\}$ satisfying

$$\int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi_\alpha} c(-\Psi) = G(t)$$

and $(F_t - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$.

Furthermore, for any holomorphic (n,0) form \hat{F} on $\{\Psi < -t\}$ satisfying

$$\int_{\{\Psi<-t\}} |\hat{F}|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty$$

and $(\hat{F} - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$. We have the following equality

$$\int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{\Psi < -t\}} |\hat{F} - F_t|^2 e^{-\varphi_{\alpha}} c(-\Psi)$$

$$= \int_{\{\Psi < -t\}} |\hat{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi).$$
(3.3)

Proof. We firstly show the existence of F_t . As $G(t) < +\infty$, then there exists a sequence of holomorphic (n, 0) forms $\{f_i\}_{i \in \mathbb{N}^+}$ on $\{\Psi < -t\}$ such that

$$\lim_{j \to +\infty} \int_{\{\Psi < -t\}} |f_j|^2 e^{-\varphi_\alpha} c(-\Psi) = G(t)$$

and $(f_j - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$ and any $j \ge 1$. It follows from Lemma 3.2 that there exists a subsequence of $\{f_j\}_{j\in\mathbb{N}^+}$ compactly convergent to a holomorphic (n, 0) form F on $\{\Psi < -t\}$ which satisfies

$$\int_{\{\Psi < -t\}} |F|^2 e^{-\varphi_\alpha} c(-\Psi) \le G(t)$$

and $(F-f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. By the definition of G(t), we have $\int_{\{\Psi \le -t\}} |F|^2 e^{-\varphi_{\alpha}} c(-\Psi) = G(t)$. Then we obtain the existence of $F_t(=F)$.

We prove the uniqueness of F_t by contradiction: if not, there exist two different holomorphic (n,0) forms f_1 and f_2 on $\{\Psi < -t\}$ satisfying $\int_{\{\Psi < -t\}} |f_1|^2 e^{-\varphi_\alpha}$ $c(-\Psi) = \int_{\{\Psi < -t\}} |f_2|^2 e^{-\varphi_\alpha} c(-\Psi) = G(t), \ (f_1 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$ and $(f_2 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Note that

$$\begin{split} &\int_{\{\Psi<-t\}} |\frac{f_1+f_2}{2}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{\Psi<-t\}} |\frac{f_1-f_2}{2}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &= \frac{1}{2} (\int_{\{\Psi<-t\}} |f_1|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{\Psi<-t\}} |f_1|^2 e^{-\varphi_{\alpha}} c(-\Psi)) = G(t), \end{split}$$

then we obtain that

$$\int_{\{\Psi < -t\}} |\frac{f_1 + f_2}{2}|^2 e^{-\varphi_\alpha} c(-\Psi) < G(t)$$

and $(\frac{f_1+f_2}{2}-f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$, which contradicts to the definition of G(t).

Now we prove the equality (3.3). Let h be any holomorphic (n,0) form on $\{\Psi < -t\}$ such that $\int_{\{\Psi < -t\}} |h|^2 e^{-\varphi_{\alpha}} c(-\Psi) < +\infty$ and $h \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. It is clear that for any complex number α , $F_t + \alpha h$ satisfying $((F_t + \alpha h) - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$ and $\int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi_{\alpha}} c(-\Psi) \leq \int_{\{\Psi < -t\}} |F_t + \alpha h|^2 e^{-\varphi_{\alpha}} c(-\Psi)$. Note that

$$\int_{\{\Psi < -t\}} |F_t + \alpha h|^2 e^{-\varphi_{\alpha}} c(-\Psi) - \int_{\{\psi < -t\}} |F_t|^2 e^{-\varphi_{\alpha}} c(-\Psi) \ge 0$$

(By considering $\alpha \to 0$) implies

$$\Re \int_{\{\Psi < -t\}} F_t \bar{h} e^{-\varphi_\alpha} c(-\Psi) = 0,$$

then we have

$$\int_{\{\Psi < -t\}} |F_t + h|^2 e^{-\varphi_\alpha} c(-\Psi) = \int_{\{\Psi < -t\}} (|F_t|^2 + |h|^2) e^{-\varphi_\alpha} c(-\Psi).$$

Letting $h = \hat{F} - F_t$, we obtain equality (3.3).

The following lemma shows the lower semicontinuity property of G(t).

Lemma 3.5. G(t) is decreasing with respect to $t \in [T, +\infty)$, such that $\lim_{t \to t_0+0} G(t) = G(t_0)$ for any $t_0 \in [T, +\infty)$, and if $G(t) < +\infty$ for some t > T, then $\lim_{t \to +\infty} G(t) = 0$. Especially, G(t) is lower semicontinuous on $[T, +\infty)$.

Proof. By the definition of G(t), it is clear that G(t) is decreasing on $[T, +\infty)$. If $G(t) < +\infty$ for some t > T, by the dominated convergence theorem, we know $\lim_{t\to+\infty}G(t)=0.$ It suffices to prove $\lim_{t\to t_0+0}G(t)=G(t_0)$. We prove it by contradiction: if not, then $\lim_{t \to t_0+0} G(t) < G(t_0)$.

By using Lemma 3.4, for any $t > t_0$, there exists a unique holomorphic (n, 0)form F_t on $\{\Psi < -t\}$ satisfying $\int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi_\alpha} c(-\Psi) = G(t)$ and $(F_t - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Note that G(t) is decreasing with respect to t. We have $\int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi} c(-\psi) \leq \lim_{t \to t_0+0} G(t)$ for any $t > t_0$. If $\lim_{t \to t_0+0} G(t) = +\infty$, the equality $\lim_{t \to t_0+0} G(t) = G(t_0)$ obviously holds, thus it suffices to prove the case $\lim_{t \to t_0+0} G(t) < +\infty. \text{ It follows from } \int_{\{\Psi < -t\}} |F_t|^2 e^{-\varphi_\alpha} c(-\Psi) \leq \lim_{t \to t_0+0} G(t) < +\infty$ holds for any $t \in (t_0, t_1]$ (where $t_1 > t_0$ is a fixed number) and Lemma 3.2 that there exists a subsequence of $\{F_t\}$ (denoted by $\{F_{t_i}\}$) compactly convergent to a holomorphic (n, 0) form \hat{F}_{t_0} on $\{\Psi < -t_0\}$ satisfying

$$\int_{\{\Psi < -t_0\}} |\hat{F}_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \le \lim_{t \to t_0 + 0} G(t) < +\infty$$

and $(F_{t_0} - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$.

Then we obtain that $G(t_0) \leq \int_{\{\Psi < -t_0\}} |\hat{F}_{t_0}|^2 e^{-\varphi_\alpha} c(-\Psi) \leq \lim_{t \to t_0+0} G(t)$, which contradicts $\lim_{t \to t_0+0} G(t) < G(t_0)$. Thus we have $\lim_{t \to t_0+0} G(t) = G(t_0)$.

We consider the derivatives of G(t) in the following lemma.

Lemma 3.6. Assume that $G(t_1) < +\infty$, where $t_1 \in (T, +\infty)$. Then for any $t_0 > t_1$, we have

$$\frac{G(t_1) - G(t_0)}{\int_{t_1}^{t_0} c(t)e^{-t}dt} \le \liminf_{B \to 0+0} \frac{G(t_0) - G(t_0 + B)}{\int_{t_0}^{t_0 + B} c(t)e^{-t}dt},$$

i.e.

$$\frac{G(t_0) - G(t_1)}{\int_{T_1}^{t_0} c(t)e^{-t}dt - \int_{T_1}^{t_1} c(t)e^{-t}dt} \ge \limsup_{B \to 0+0} \frac{G(t_0 + B) - G(t_0)}{\int_{T_1}^{t_0+B} c(t)e^{-t}dt - \int_{T_1}^{t_0} c(t)e^{-t}dt}$$

Proof. It follows from Lemma 3.5 that $G(t) < +\infty$ for any $t > t_1$. By Lemma 3.4,

there exists a holomorphic (n,0) form F_{t_0} on $\{\Psi < -t_0\}$, such that $(F_{t_0} - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$ and $G(t_0) = \int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_\alpha} c(-\Psi)$. It suffices to consider that $\liminf_{B \to 0+0} \frac{G(t_0) - G(t_0 + B)}{\int_{t_0}^{t_0 + B} c(t) e^{-t} dt} \in [0, +\infty)$ because of the decreasing property of G(t). Then there exists $1 \geq B_j \to 0 + 0$ (as $j \to +\infty$) such that

$$\lim_{j \to +\infty} \frac{G(t_0) - G(t_0 + B_j)}{\int_{t_0}^{t_0 + B_j} c(t)e^{-t}dt} = \liminf_{B \to 0+0} \frac{G(t_0) - G(t_0 + B)}{\int_{t_0}^{t_0 + B} c(t)e^{-t}dt}$$
(3.4)

and $\{\frac{G(t_0)-G(t_0+B_j)}{\int_{t_0}^{t_0+B_j}c(t)e^{-t}dt}\}_{j\in\mathbb{N}^+}$ is bounded. As $c(t)e^{-t}$ is decreasing and positive on $(t, +\infty)$, then

$$\lim_{j \to +\infty} \frac{G(t_0) - G(t_0 + B_j)}{\int_{t_0}^{t_0 + B_j} c(t)e^{-t}dt} = \left(\lim_{j \to +\infty} \frac{G(t_0) - G(t_0 + B_j)}{B_j}\right) \left(\frac{1}{\lim_{t \to t_0 + 0} c(t)e^{-t}}\right) = \left(\lim_{j \to +\infty} \frac{G(t_0) - G(t_0 + B_j)}{B_j}\right) \left(\frac{e^{t_0}}{\lim_{t \to t_0 + 0} c(t)}\right).$$
(3.5)

Hence $\{\frac{G(t_0)-G(t_0+B_j)}{B_j}\}_{j\in\mathbb{N}^+}$ is uniformly bounded with respect to j. As $t \leq v_{t_0,j}(t)$, the decreasing property of $c(t)e^{-t}$ shows that

$$e^{-\Psi + v_{t_0, B_j}(\Psi)} c(-v_{t_0, B_j}(\Psi)) \ge c(-\Psi).$$

It follows from Lemma 2.4 that, for any B_j , there exists holomorphic (n, 0) form \tilde{F}_j on $\{\Psi < -t_1\}$ such that

$$\begin{split} &\int_{\{\Psi<-t_1\}} |\tilde{F}_j - (1 - b_{t_0,B_j}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &\leq \int_{\{\Psi<-t_1\}} |\tilde{F}_j - (1 - b_{t_0,B_j}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha}} e^{-\Psi+v_{t_0,B_j}(\Psi)} c(-v_{t_0,B_j}(\Psi)) \\ &\leq \int_{t_1}^{t_0+B_j} c(t) e^{-t} dt \int_{\{\Psi<-t_1\}} \frac{1}{B_j} \mathbb{I}_{\{-t_0-B_j<\Psi<-t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}-\Psi} \\ &\leq \frac{e^{t_0+B_j} \int_{t_1}^{t_0+B_j} c(t) e^{-t} dt}{i \epsilon_{(t_0,t_0+B_j)}} \int_{\{\Psi<-t_1\}} \frac{1}{B_j} \mathbb{I}_{\{-t_0-B_j<\Psi<-t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &= \frac{e^{t_0+B_j} \int_{t_1}^{t_0+B_j} c(t) e^{-t} dt}{i \epsilon_{(t_0,t_0+B_j)} c(t)} \times \left(\int_{\{\Psi<-t_1\}} \frac{1}{B_j} \mathbb{I}_{\{\Psi<-t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \right) \\ &- \int_{\{\Psi<-t_1\}} \frac{1}{B_j} \mathbb{I}_{\{\Psi<-t_0-B_j\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \right) \\ &\leq \frac{e^{t_0+B_j} \int_{t_1}^{t_0+B_j} c(t) e^{-t} dt}{i \epsilon_{(t_0,t_0+B_j)} c(t)} \times \frac{G(t_0) - G(t_0+B_j)}{B_j} < +\infty. \end{split}$$
(3.6)

Note that $b_{t_0,B_j}(t) = 0$ for $t \leq -t_0 - B_j$, $b_{t_0,B_j}(t) = 1$ for $t \geq t_0$, $v_{t_0,B_j}(t) > -t_0 - B_j$ and $c(t)e^{-t}$ is decreasing with respect to t. It follows from inequality (3.6) that $(F_j - F_{t_0})_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes I(\Psi + \varphi_\alpha)_{z_0} \subset \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Note that

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_j|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq 2 \int_{\{\Psi < -t_1\}} |\tilde{F}_j - (1 - b_{t_0, B_j}(\Psi)) F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + 2 \int_{\{\Psi < -t_1\}} |(1 - b_{t_0, B_j}(\Psi)) F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq 2 \frac{e^{t_0 + B_j} \int_{t_1}^{t_0 + B_j} c(t) e^{-t} dt}{\sum_{t \in (t_0, t_0 + B_j)} c(t)} \times \frac{G(t_0) - G(t_0 + B_j)}{B_j} + 2 \int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi). \tag{3.7}$$

We also note that $B_j \leq 1$, $\frac{G(t_0)-G(t_0+B_j)}{B_j}$ is uniformly bounded with respect to j and $G(t_0) = \int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi)$. It follows from inequality (3.7) that we know $\int_{\{\Psi < -t_1\}} |\tilde{F}_j|^2 e^{-\varphi_{\alpha}} c(-\Psi)$ is uniformly bounded with respect to j.

It follows from Lemma 3.2 that there exists a subsequence of $\{\tilde{F}_j\}_{j\in\mathbb{N}^+}$ compactly convergent to a holomorphic (n,0) form \tilde{F}_{t_1} on $\{\Psi < -t_1\}$ which satisfies

$$\int_{\{\Psi<-t_1\}} |\tilde{F}_{t_1}|^2 e^{-\varphi_\alpha} c(-\Psi) \le \liminf_{j \to +\infty} \int_{\{\Psi<-t_1\}} |\tilde{F}_j|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty,$$

and $(\tilde{F}_{t_1} - F_{t_0})_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. Note that $\lim_{j \to +\infty} b_{t_0,B_j}(t) = \mathbb{I}_{\{t \ge -t_0\}}$ and

$$v_{t_0}(t) := \lim_{j \to +\infty} v_{t_0, B_j}(t) = \begin{cases} -t_0 & \text{if } x < -t_0, \\ t & \text{if } x \ge t_0. \end{cases}$$

It follows from inequality (3.6) and Fatou's lemma that

$$\int_{\{\Psi < -t_0\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{-t_0 \le \Psi < -t_1\}} |\tilde{F}_{t_1}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq \int_{\{\Psi < -t_1\}} |\tilde{F}_{t_1} - \mathbb{I}_{\{\Psi < -t_0\}} F_{t_0}|^2 e^{-\varphi_{\alpha}} e^{-\Psi + v_{t_0}(\Psi)} c(-v_{t_0}(\Psi)) \\
\leq \liminf_{j \to +\infty} \int_{\{\Psi < -t_1\}} |\tilde{F}_j - (1 - b_{t_0, B_j}(\Psi)) F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq \liminf_{j \to +\infty} \left(\frac{e^{t_0 + B_j} \int_{t_1}^{t_0 + B_j} c(t) e^{-t} dt}{\inf_{t \in (t_0, t_0 + B_j)} c(t)} \times \frac{G(t_0) - G(t_0 + B_j)}{B_j} \right).$$
(3.8)

It follows from Lemma 3.4, equality (3.4), equality (3.5) and inequality (3.8) that we have

$$\begin{split} &\int_{\{\Psi<-t_1\}} |\tilde{F}_{t_1}|^2 e^{-\varphi_{\alpha}} c(-\Psi) - \int_{\{\Psi<-t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &\leq \int_{\{\Psi<-t_0\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{-t_0 \le \Psi<-t_1\}} |\tilde{F}_{t_1}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &\leq \int_{\{\Psi<-t_1\}} |\tilde{F}_{t_1} - \mathbb{I}_{\{\Psi<-t_0\}} F_{t_0}|^2 e^{-\varphi_{\alpha}} e^{-\Psi+v_{t_0}(\Psi)} c(-v_{t_0}(\Psi)) \\ &\leq \liminf_{j \to +\infty} \int_{\{\Psi<-t_1\}} |\tilde{F}_j - (1 - b_{t_0,B_j}(\Psi)) F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &\leq \liminf_{j \to +\infty} \left(\frac{e^{t_0+B_j} \int_{t_1}^{t_0+B_j} c(t) e^{-t} dt}{i \epsilon_{(t_0,t_0+B_j)}} \times \frac{G(t_0) - G(t_0+B_j)}{B_j} \right) \\ &\leq \left(\int_{t_1}^{t_0} c(t) e^{-t} dt \right) \liminf_{B \to 0+0} \frac{G(t_0) - G(t_0+B)}{\int_{t_0}^{t_0+B} c(t) e^{-t} dt}. \end{split}$$

Note that $(\tilde{F}_{t_1} - F_{t_0})_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$. It follows from the definition of G(t) and inequality (3.9) that we have

$$G(t_{1}) - G(t_{0}) \leq \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) - \int_{\{\Psi < -t_{0}\}} |F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) \leq \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}} - \mathbb{I}_{\{\Psi < -t_{0}\}} F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) \leq \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}} - \mathbb{I}_{\{\Psi < -t_{0}\}} F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} e^{-\Psi + v_{t_{0}}(\Psi)} c(-v_{t_{0}}(\Psi)) \leq \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}} - \mathbb{I}_{\{\Psi < -t_{0}\}} F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} e^{-\Psi + v_{t_{0}}(\Psi)} c(-v_{t_{0}}(\Psi)) \leq \int_{\{1}^{t_{0}} c(t) e^{-t} dt) \liminf_{B \to 0 + 0} \frac{G(t_{0}) - G(t_{0} + B)}{\int_{t_{0}}^{t_{0} + B} c(t) e^{-t} dt}.$$
(3.10)

Lemma 3.6 is proved.

The following property of concave functions will be used in the proof of Theorem 1.2.

Lemma 3.7 (see [23]). Let H(r) be a lower semicontinuous function on (0, R]. Then H(r) is concave if and only if

$$\frac{H(r_1) - H(r_2)}{r_1 - r_2} \le \liminf_{r_3 \to r_2 - 0} \frac{H(r_3) - H(r_2)}{r_3 - r_2}$$

holds for any $0 < r_2 < r_1 \leq R$.

4. Proofs of Theorem 1.2, Remark 1.3, Corollary 1.4 and Remark 1.5

We firstly prove Theorem 1.2.

Proof. We firstly show that if $G(t_0) < +\infty$ for some $t_0 > T$, then $G(t_1) < +\infty$ for any $T < t_1 < t_0$. As $G(t_0) < +\infty$, it follows from Lemma 3.4 that there exists a unique holomorphic (n, 0) form F_{t_0} on $\{\Psi < -t\}$ satisfying

$$\int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_\alpha} c(-\Psi) = G(t_0) < +\infty$$

and $(F_{t_0} - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$.

It follows from Lemma 2.4 that there exists a holomorphic (n,0) form \tilde{F}_1 on $\{\Psi < -t_1\}$ such that

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))F_{t_0}|^2 e^{-\varphi_\alpha + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) \\
\leq (\int_{t_1}^{t_0 + B} c(s)e^{-s}ds) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_\alpha - \Psi} < +\infty.$$
(4.1)

Note that $b_{t_0,B}(t) = 0$ on $\{\Psi < -t_0 - B\}$ and $v_{t_0,B}(\Psi) > -t_0 - B$. We have $e^{v_{t_0,B}(\Psi)}c(-v_{t_0,B}(\Psi))$ has a positive lower bound. It follows from inequality (4.1) that we have $(F_1 - F_{t_0})_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes I(\Psi + \varphi_\alpha)_{z_0} \subset \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$, which implies that $(F_1 - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$. As

_	_	

 $v_{t_0,B}(\Psi) \geq \Psi$ and $c(t)e^{-t}$ is decreasing with respect to t, it follows from inequality (4.1) that we have

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq \int_{\{\Psi < -t_1\}} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha} + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi)) \\
\leq (\int_{t_1}^{t_0 + B} c(s)e^{-s} ds) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha} - \Psi} < +\infty.$$
(4.2)

Then we have

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_1|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq 2 \int_{\{\Psi < -t_1\}} |\tilde{F}_1 - (1 - b_{t_0,B}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + 2 \int_{\{\Psi < -t_1\}} |(1 - b_{t_0,B}(\Psi))F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
\leq 2 (\int_{t_1}^{t_0 + B} c(s)e^{-s} ds) \int_M \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha} - \Psi} + 2 \int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\
< + \infty.$$
(4.3)

Hence we have $G(t_1) \leq \int_{\{\Psi < -t_1\}} |\tilde{F}_1|^2 e^{-\varphi_\alpha} c(-\Psi) < +\infty.$

Now, it follows from Lemma 3.5, Lemma 3.6 and Lemma 3.7 that we know $G(h^{-1}(r))$ is concave with respect to r. It follows from Lemma 3.5 that $\lim_{t \to T^+ 0} G(t) =$ G(T) and $\lim_{t \to +\infty} G(t) = 0.$

Theorem 1.2 is proved.

Now we prove Remark 1.3.

Proof. Note that if there exists a positive decreasing concave function q(t) on $(a,b) \subset \mathbb{R}$ and g(t) is not a constant function, then $b < +\infty$.

Assume that $G(t_0) < +\infty$ for some $t_0 \ge T$. As $f_{z_0} \notin \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for some $z_0 \in Z_0$, Lemma 3.3 shows that $G(t_0) \in (0, +\infty)$. Following from Theorem 1.2 we know $G(h^{-1}(r))$ is concave with respect to $r \in (\int_{T_1}^T c(t)e^{-t}dt, \int_{T_1}^{+\infty} c(t)e^{-t}dt)$ and $G(h^{-1}(r))$ is not a constant function, therefore we obtain $\int_{T_1}^{+\infty} c(t)e^{-t}dt < +\infty$, which contradicts to $\int_{T_1}^{+\infty} c(t)e^{-t}dt = +\infty$. Thus we have $G(t) \equiv +\infty$. When $G(t_2) \in (0, +\infty)$ for some $t_2 \in [T, +\infty)$, Lemma 3.3 shows that $f_{z_0} \notin C$.

 $\mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$. Combining the above discussion, we know $\int_{T_1}^{+\infty} c(t) e^{-t} dt < +\infty$. Using Theorem 1.2, we obtain that $G(\hat{h}^{-1}(r))$ is concave with respect to $r \in (0, \int_T^{+\infty} c(t)e^{-t}dt)$, where $\hat{h}(t) = \int_t^{+\infty} c(t)e^{-t}dt$.

Thus, Remark 1.3 holds.

Now we prove Corollary 1.4

Proof. As $G(h^{-1}(r))$ is linear with respect to $r \in [0, \int_T^{+\infty} c(s)e^{-s}ds)$, we have $G(t) = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s)e^{-s}ds} \int_t^{+\infty} c(s)e^{-s}ds$ for any $t \in [T, +\infty)$ and $T_1 \in (T, +\infty)$.

We follow the notation and the construction in Lemma 3.6. Let $t_0 > t_1 > T$ be given. It follows from $G(h^{-1}(r))$ is linear with respect to $r \in [0, \int_T^{+\infty} c(s)e^{-s}ds)$ that we know that all inequalities in (3.10) should be equalities, i.e., we have

$$G(t_{1}) - G(t_{0})$$

$$= \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) - \int_{\{\Psi < -t_{0}\}} |F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi)$$

$$= \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}} - \mathbb{I}_{\{\Psi < -t_{0}\}} F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi)$$

$$= \int_{\{\Psi < -t_{1}\}} |\tilde{F}_{t_{1}} - \mathbb{I}_{\{\Psi < -t_{0}\}} F_{t_{0}}|^{2} e^{-\varphi_{\alpha}} e^{-\Psi + v_{t_{0}}(\Psi)} c(-v_{t_{0}}(\Psi))$$

$$= \left(\int_{t_{1}}^{t_{0}} c(t) e^{-t} dt\right) \liminf_{B \to 0+0} \frac{G(t_{0}) - G(t_{0} + B)}{\int_{t_{0}}^{t_{0} + B} c(t) e^{-t} dt}.$$
(4.4)

Note that $G(t_0) = \int_{\{\Psi < -t_0\}} |F_{t_0}|^2 e^{-\varphi_{\alpha}} c(-\Psi)$. Equality (4.4) shows that $G(t_1) = \int_{\{\Psi < -t_1\}} |\tilde{F}_{t_1}|^2 e^{-\varphi_{\alpha}} c(-\Psi)$.

Note that on $\{\Psi \geq -t_0\}$, we have $e^{-\Psi + v_{t_0}(\Psi)}c(-v_{t_0}(\Psi)) = c(-\Psi)$. It follows from

$$\int_{\{\Psi < -t_1\}} |\tilde{F}_{t_1} - \mathbb{I}_{\{\Psi < -t_0\}} F_{t_0}|^2 e^{-\varphi_\alpha} c(-\Psi)$$

=
$$\int_{\{\Psi < -t_1\}} |\tilde{F}_{t_1} - \mathbb{I}_{\{\Psi < -t_0\}} F_{t_0}|^2 e^{-\varphi_\alpha} e^{-\Psi + v_{t_0}(\Psi)} c(-v_{t_0}(\Psi))$$

that we have (note that $v_{t_0}(\Psi) = -t_0$ on $\{\Psi < -t_0\}$)

$$\int_{\{\Psi < -t_0\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_\alpha} c(-\Psi)$$

$$= \int_{\{\Psi < -t_0\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_\alpha} e^{-\Psi - t_0} c(t_0).$$
(4.5)

As $\int_T^{+\infty} c(t)e^{-t}dt < +\infty$ and $c(t)e^{-t}$ is decreasing with respect to t, we know that there exists $t_2 > t_0$ such that $c(t)e^{-t} < c(t_0)e^{-t_0} - \epsilon$ for any $t \ge t_2$, where $\epsilon > 0$ is a constant. Then equality (4.5) implies that

$$\begin{aligned} &\epsilon \int_{\{\Psi < -t_2\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_\alpha - \Psi} \\ &\leq \int_{\{\Psi < -t_2\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_\alpha} (e^{-\Psi - t_0} c(t_0) - c(-\Psi)) \\ &\leq \int_{\{\Psi < -t_0\}} |\tilde{F}_{t_1} - F_{t_0}|^2 e^{-\varphi_\alpha} (e^{-\Psi - t_0} c(t_0) - c(-\Psi)) \\ &= 0. \end{aligned}$$

$$(4.6)$$

Note that $e^{-\varphi_{\alpha}-\Psi} \ge e^{-(\varphi_{\alpha}+\psi)}|F|^2$, $\varphi_{\alpha}+\psi$ is a plurisubharmonic function and the integrand in (4.6) is nonnegative, we must have $\tilde{F}_{t_1}|_{\{\Psi < -t_0\}} = F_{t_0}$.

It follows from Lemma 3.4 that for any t > T, there exists a unique holomorphic (n, 0) form F_t on $\{\Psi < -t\}$ satisfying

$$\int_{\{\Psi<-t\}}|F_t|^2e^{-\varphi_\alpha}c(-\Psi)=G(t)$$

and $(F_t - f) \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$. By the above discussion, we know $F_t = F_{t'}$ on $\{\Psi < -\max\{t, t'\}\}$ for any $t \in (T, +\infty)$ and $t' \in (T, +\infty)$. Hence combining $\lim_{t\to T+0} G(t) = G(T)$, we obtain that there exists a unique holomorphic (n, 0) form \tilde{F} on $\{\Psi < -T\}$ satisfying $(\tilde{F} - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$ for any $z_0 \in Z_0$ and $G(t) = \int_{\{\Psi < -t\}} |\tilde{F}|^2 e^{-\varphi_\alpha} c(-\Psi)$ for any $t \geq T$.

Secondly, we prove equality (1.2).

As a(t) is a nonnegative measurable function on $(T, +\infty)$, then there exists a sequence of functions $\{\sum_{j=1}^{n_i} a_{ij} \mathbb{I}_{E_{ij}}\}_{i \in \mathbb{N}^+}$ $(n_i < +\infty \text{ for any } i \in \mathbb{N}^+)$ satisfying that $\sum_{j=1}^{n_i} a_{ij} \mathbb{I}_{E_{ij}}$ is increasing with respect to i and $\lim_{i \to +\infty} \sum_{j=1}^{n_i} a_{ij} \mathbb{I}_{E_{ij}} = a(t)$ for any $t \in (T, +\infty)$, where E_{ij} is a Lebesgue measurable subset of $(T, +\infty)$ and $a_{ij} \ge 0$ is a constant for any i, j. It follows from Levi's Theorem that it suffices to prove the case that $a(t) = \mathbb{I}_E(t)$, where $E \subset (T, +\infty)$ is a Lebesgue measurable set.

case that $a(t) = \mathbb{I}_E(t)$, where $E \subset (T, +\infty)$ is a Lebesgue measurable set. Note that $G(t) = \int_{\{\Psi < -t\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi) = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s)e^{-s}ds} \int_t^{+\infty} c(s)e^{-s}ds$ where $T_1 \in (T, +\infty)$, then

$$\int_{\{-t_1 \le \Psi < -t_2\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi) = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2}^{t_1} c(s) e^{-s} ds \tag{4.7}$$

holds for any $T \leq t_2 < t_1 < +\infty$. It follows from the dominated convergence theorem and equality (4.7) that

$$\int_{\{z\in M:-\Psi(z)\in N\}} |\tilde{F}|^2 e^{-\varphi_\alpha} = 0 \tag{4.8}$$

holds for any $N \subset (T, +\infty)$ such that $\mu(N) = 0$, where μ is the Lebesgue measure on \mathbb{R} .

As $c(t)e^{-t}$ is decreasing on $(T, +\infty)$, there are at most countable points denoted by $\{s_j\}_{j\in\mathbb{N}^+}$ such that c(t) is not continuous at s_j . Then there is a decreasing sequence of open sets $\{U_k\}$, such that $\{s_j\}_{j\in\mathbb{N}^+} \subset U_k \subset (T, +\infty)$ for any k, and $\lim_{k\to+\infty} \mu(U_k) = 0$. Choosing any closed interval $[t'_2, t'_1] \subset (T, +\infty)$, then we have

$$\int_{\{-t_{1}'\leq\Psi<-t_{2}'\}} |\tilde{F}|^{2}e^{-\varphi_{\alpha}} \\
= \int_{\{z\in M:-\Psi(z)\in(t_{2}',t_{1}']\setminus U_{k}\}} |\tilde{F}|^{2}e^{-\varphi_{\alpha}} + \int_{\{z\in M:-\Psi(z)\in[t_{2}',t_{1}']\cap U_{k}\}} |\tilde{F}|^{2}e^{-\varphi_{\alpha}} \\
= \lim_{n\to+\infty} \sum_{i=0}^{n-1} \int_{\{z\in M:-\Psi(z)\in I_{i,n}\setminus U_{k}\}} |\tilde{F}|^{2}e^{-\varphi_{\alpha}} + \int_{\{z\in M:-\Psi(z)\in[t_{2}',t_{1}']\cap U_{k}\}} |\tilde{F}|^{2}e^{-\varphi_{\alpha}}, \tag{4.9}$$

where $I_{i,n} = (t'_1 - (i+1)\alpha_n, t'_1 - i\alpha_n]$ and $\alpha_n = \frac{t'_1 - t'_2}{n}$. Note that

$$\lim_{n \to +\infty} \sum_{i=0}^{n-1} \int_{\{z \in M: -\Psi(z) \in I_{i,n} \setminus U_k\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}}$$

$$\leq \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{I_{i,n} \setminus U_k} c(t)} \int_{\{z \in M: -\Psi(z) \in I_{i,n} \setminus U_k\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi).$$
(4.10)

It follows from equality (4.7) that inequality (4.10) becomes

$$\lim_{n \to +\infty} \sum_{i=0}^{n-1} \int_{\{z \in M: -\Psi(z) \in I_{i,n} \setminus U_k\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}}$$

$$\leq \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{I_{i,n} \setminus U_k} c(t)} \int_{I_{i,n} \setminus U_k} c(s) e^{-s} ds.$$

$$(4.11)$$

It is clear that c(t) is uniformly continuous and has positive lower bound and upper bound on $[t'_2, t'_1] \setminus U_k$. Then we have

$$\limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{I_{i,n} \setminus U_k} c(t)} \int_{I_{i,n} \setminus U_k} c(s) e^{-s} ds$$

$$\leq \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{\sup_{I_{i,n} \setminus U_k} c(t)}{\inf_{I_{i,n} \setminus U_k} c(t)} \int_{I_{i,n} \setminus U_k} e^{-s} ds$$

$$= \int_{(t'_2, t'_1] \setminus U_k} e^{-s} ds.$$
(4.12)

Combining inequality (4.9), (4.11) and (4.12), we have

$$\int_{\{-t_{1}' \leq \Psi < -t_{2}'\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} \\
= \int_{\{z \in M: -\Psi(z) \in (t_{2}', t_{1}'] \setminus U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in [t_{2}', t_{1}'] \cap U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} \\
\leq \frac{G(T_{1})}{\int_{T_{1}}^{+\infty} c(s) e^{-s} ds} \int_{(t_{2}', t_{1}'] \setminus U_{k}} e^{-s} ds + \int_{\{z \in M: -\Psi(z) \in [t_{2}', t_{1}'] \cap U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}}.$$
(4.13)

Let $k \to +\infty$, following from equality (4.8) and inequality (4.13), then we obtain that

$$\int_{\{-t_1' \le \psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \le \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2'}^{t_1'} e^{-s} ds.$$
(4.14)

Following from a similar discussion we can obtain that

$$\int_{\{-t_1' \leq \psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \geq \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2'}^{t_1'} e^{-s} ds.$$

Then combining inequality (4.14), we know

$$\int_{\{-t_1' \le \Psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2'}^{t_1'} e^{-s} ds.$$
(4.15)

Then it is clear that for any open set $U \subset (T, +\infty)$ and compact set $V \subset (T, +\infty)$,

$$\int_{\{z \in M; -\Psi(z) \in U\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_U e^{-s} ds,$$
$$\int_{\{z \in M; -\Psi(z) \in V\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_V e^{-s} ds.$$

and

As
$$E \subset (T, +\infty)$$
, then $E \cap (t_2, t_1]$ is a Lebesgue measurable subset of $(T + \frac{1}{n}, n)$ for some large n , where $T \leq t_2 < t_1 \leq +\infty$. Then there exists a sequence of compact sets $\{V_j\}$ and a sequence of open subsets $\{V'_j\}$ satisfying $V_1 \subset \ldots \subset V_j \subset V_{j+1} \subset \ldots \subset E \cap (t_2, t_1] \subset \ldots \subset V'_{j+1} \subset V'_j \subset \ldots \subset V'_1 \subset (T, +\infty)$ and $\lim_{j \to +\infty} \mu(V'_j - V_j) = 0$, where μ is the Lebesgue measure on \mathbb{R} . Then we have

$$\begin{split} \int_{\{-t_1' \leq \Psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \mathbb{I}_E(-\Psi) &= \int_{z \in M: -\Psi(z) \in E \cap (t_2, t_1]} |\tilde{F}|^2 e^{-\varphi_\alpha} \\ &\leq \liminf_{j \to +\infty} \int_{\{z \in M: -\Psi(z) \in V_j'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \\ &\leq \liminf_{j \to +\infty} \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{V_j'} e^{-s} ds \\ &\leq \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{E \cap (t_2, t_1]} e^{-s} ds \\ &= \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2}^{t_1} e^{-s} \mathbb{I}_E(s) ds, \end{split}$$

and

$$\int_{\{-t_1' \le \Psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} \mathbb{I}_E(-\Psi) \ge \liminf_{j \to +\infty} \int_{\{z \in M: -\Psi(z) \in V_j\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}}$$
$$\ge \liminf_{j \to +\infty} \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{V_j} e^{-s} ds$$
$$= \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2}^{t_1} e^{-s} \mathbb{I}_E(s) ds,$$

which implies that

$$\int_{\{-t_1' \le \Psi < -t_2'\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \mathbb{I}_E(-\Psi) = \frac{G(T_1)}{\int_{T_1}^{+\infty} c(s) e^{-s} ds} \int_{t_2}^{t_1} e^{-s} \mathbb{I}_E(s) ds.$$

Hence we know that equality (1.2) holds. Corollary 1.4 is proved.

Now we prove Remark 1.5.

Proof of Remark 1.5. By the definition of $G(t; \tilde{c})$, we have $G(t_0; \tilde{c}) \leq \int_{\{\Psi < -t_0\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \tilde{c}(-\Psi)$, where \tilde{F} is the holomorphic (n, 0) form on $\{\Psi < -T\}$ such that $G(t) = \int_{\{\Psi < -t\}} |\tilde{F}|^2 e^{-\varphi_\alpha} c(-\Psi)$ for any $t \geq T$. Hence we only consider the case $G(t_0; \tilde{c}) < +\infty$.

By the definition of $G(t; \tilde{c})$, we can choose a holomorphic (n, 0) form $F_{t_0, \tilde{c}}$ on $\{\Psi < -t_0\}$ satisfying $(F_{t_0, \tilde{c}} - f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes J_{z_0}$, for any $z_0 \in Z_0$ and

 $\int_{\{\Psi < -t_0\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} \tilde{c}(-\Psi) < +\infty. \text{ As } \mathcal{H}^2(\tilde{c},t_0) \subset \mathcal{H}^2(c,t_0), \text{ we have } \int_{\{\Psi < -t_0\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} c(-\Psi) < +\infty. \text{ Using Lemma 3.4, we obtain that}$

$$\begin{split} \int_{\{\Psi < -t\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} c(-\Psi) &= \int_{\{\Psi < -t\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \\ &+ \int_{\{\Psi < -t\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi) \end{split}$$

for any $t \geq t_0$, then

$$\int_{\{-t_3 \le \Psi < -t_4\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} c(-\Psi) = \int_{\{-t_3 \le \Psi < -t_4\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi) + \int_{\{-t_3 \le \Psi < -t_4\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_{\alpha}} c(-\Psi)$$
(4.16)

holds for any $t_3 > t_4 \ge t_0$. It follows from the dominant convergence theorem, equality (4.16), (4.8) and c(t) > 0 for any t > T, that

$$\int_{\{z \in M: -\Psi(z)=t\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha} = \int_{\{z \in M: -\Psi(z)=t\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_\alpha}$$
(4.17)

holds for any $t > t_0$.

Choosing any closed interval $[t'_4, t'_3] \subset (t_0, +\infty) \subset (T, +\infty)$. Note that c(t) is uniformly continuous and have positive lower bound and upper bound on $[t'_4, t'_3] \setminus U_k$, where $\{U_k\}$ is the decreasing sequence of open subsets of $(T, +\infty)$, such that c is continuous on $(T, +\infty) \setminus U_k$ and $\lim_{k \to +\infty} \mu(U_k) = 0$. Take $N = \bigcap_{k=1}^{+\infty} U_k$. Note that

$$\int_{\{-t'_{3} \leq \Psi < -t'_{4}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\
= \lim_{n \to +\infty} \sum_{i=0}^{n-1} \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4}, t'_{3}] \cap U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\
\leq \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{S_{i,n}} c(t)} \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) \\
+ \int_{\{z \in M: -\Psi(z) \in (t'_{4}, t'_{3}] \cap U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}}, \tag{4.18}$$

where $S_{i,n} = (t'_4 - (i+1)\alpha_n, t'_3 - i\alpha_n]$ and $\alpha_n = \frac{t'_3 - t'_4}{n}$. It follows from equality (4.16),(4.17), (4.8) and the dominated theorem that

$$\int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_k\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha} c(-\Psi) \\
= \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_k\}} |\tilde{F}|^2 e^{-\varphi_\alpha} c(-\Psi) + \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_k\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_\alpha} c(-\Psi). \tag{4.19}$$

As c(t) is uniformly continuous and have positive lower bound and upper bound on $[t'_3, t'_4] \setminus U_k$, combining equality (4.19), we have

$$\begin{split} &\lim_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{S_{i,n} \setminus U_{k}} c(t)} \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) \\ &= \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{1}{\inf_{S_{i,n} \setminus U_{k}} c(t)} (\int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} c(-\Psi) \\ &+ \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}} c(-\Psi)) \\ &\leq \limsup_{n \to +\infty} \sum_{i=0}^{n-1} \frac{\sup_{S_{i,n} \setminus U_{k}} c(t)}{\inf_{S_{i,n} \setminus U_{k}} c(t)} (\int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} \\ &+ \int_{\{z \in M: -\Psi(z) \in S_{i,n} \setminus U_{k}\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}}) \\ &= \int_{\{z \in M: -\Psi(z) \in (t'_{4}, t'_{3}] \setminus U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4}, t'_{3}] \setminus U_{k}\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}}. \end{split}$$

$$(4.20)$$

If follows from inequality (4.18) and (4.20) that

$$\int_{\{-t'_{3} \leq \Psi < -t'_{4}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\
\leq \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \setminus U_{k}\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \setminus U_{k}\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}} \\
+ \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \cap U_{k}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}}.$$
(4.21)

It follows from $F_{t_0,\tilde{c}} \in \mathcal{H}^2(c,t_0)$ that $\int_{\{-t'_3 \leq \Psi < -t'_4\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} < +\infty$. Let $k \to +\infty$, by equality (4.8), inequality (4.21) and the dominated theorem, we have

$$\int_{\{-t'_{3} \leq \Psi < -t'_{4}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\
\leq \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}]\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \setminus N\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}} \\
+ \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \cap N\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}}.$$
(4.22)

By similar discussion, we also have that

$$\begin{split} &\int_{\{-t'_{3} \leq \Psi < -t'_{4}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\ \geq &\int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}]\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \setminus N\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}} \\ &+ \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \cap N\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}}. \end{split}$$

then combining inequality (4.22), we have

$$\int_{\{-t'_{3} \leq \Psi < -t'_{4}\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}} \\
= \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}]\}} |\tilde{F}|^{2} e^{-\varphi_{\alpha}} + \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \setminus N\}} |F_{t_{0},\tilde{c}} - \tilde{F}|^{2} e^{-\varphi_{\alpha}} \\
+ \int_{\{z \in M: -\Psi(z) \in (t'_{4},t'_{3}] \cap N\}} |F_{t_{0},\tilde{c}}|^{2} e^{-\varphi_{\alpha}}.$$
(4.23)

Using equality (4.8), (4.17) and Levi's Theorem, we have

$$\int_{\{z \in M: -\Psi(z) \in U\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha}
= \int_{\{z \in M: -\Psi(z) \in U\}} |\tilde{F}|^2 e^{-\varphi} + \int_{\{z \in M: -\Psi(z) \in U \setminus N\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_\alpha}
+ \int_{\{z \in M: -\Psi(z) \in U \cap N\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha}$$
(4.24)

holds for any open set $U \subset \subset (t_0, +\infty)$, and

$$\int_{\{z\in M:-\Psi(z)\in V\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}}$$

$$= \int_{\{z\in M:-\Psi(z)\in V\}} |\tilde{F}|^2 e^{-\varphi} + \int_{\{z\in M:-\Psi(z)\in V\setminus N\}} |F_{t_0,\tilde{c}} - \tilde{F}|^2 e^{-\varphi_{\alpha}} \qquad (4.25)$$

$$+ \int_{\{z\in M:-\Psi(z)\in V\cap N\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}}$$

holds for any compact set $V \subset (t_0, +\infty)$. For any measurable set $E \subset (t_0, +\infty)$, there exists a sequence of compact set $\{V_l\}$, such that $V_l \subset V_{l+1} \subset E$ for any l and $\lim_{l \to +\infty} \mu(V_l) = \mu(E)$, hence by equality (4.25), we have

$$\int_{\{\Psi<-t_0\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha} \mathbb{I}_E(-\psi) \ge \lim_{l \to +\infty} \int_{\{\psi<-t_0\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_\alpha} \mathbb{I}_{V_j}(-\psi)
\ge \lim_{l \to +\infty} \int_{\{\psi<-t_0\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \mathbb{I}_{V_j}(-\psi)
= \int_{\{\psi<-t_0\}} |\tilde{F}|^2 e^{-\varphi} \mathbb{I}_{V_j}(-\psi).$$
(4.26)

It is clear that for any $t > t_0$, there exists a sequence of functions $\{\sum_{j=1}^{n_i} \mathbb{I}_{E_{i,j}}\}_{i=1}^{+\infty}$ defined on $(t, +\infty)$, satisfying $E_{i,j} \subset (t, +\infty)$, $\sum_{j=1}^{n_{i+1}} \mathbb{I}_{E_{i+1,j}}(s) \geq \sum_{j=1}^{n_i} \mathbb{I}_{E_{i,j}}(s)$ and $\lim_{i \to +\infty} \sum_{j=1}^{n_i} \mathbb{I}_{E_{i,j}}(s) = \tilde{c}(s)$ for any s > t. Combining Levi's Theorem and inequality (4.26), we have

$$\int_{\{\Psi < -t_0\}} |F_{t_0,\tilde{c}}|^2 e^{-\varphi_{\alpha}} \tilde{c}(-\Psi) \ge \int_{\{\Psi < -t_0\}} |\tilde{F}|^2 e^{-\varphi_{\alpha}} \tilde{c}(-\Psi).$$
(4.27)

By the definition of $G(t_0, \tilde{c})$, we have $G(t_0, \tilde{c}) = \int_{\{\Psi < -t_0\}} |\tilde{F}|^2 e^{-\varphi_\alpha} \tilde{c}(-\Psi)$. Equality (1.3) is proved.

c

5. Proof of Corollary 1.7

In this section, we prove Corollary 1.7.

Proof. Let $Z_0 \subset \{\psi = -\infty\}$. Let $F \equiv 1$ on M. Denote $\varphi_{\alpha} := \varphi$. As $\psi < -T$ on M, we have $\Psi := \min\{\psi - 2\log|F|, -T\} = \psi$. which implies that $H_{z_0} = \mathcal{H}_{z_0}$ for any $z_0 \in Z_0$. Denote $J_{z_0} := \mathcal{F}_{z_0}$ for any $z_0 \in Z_0$.

Let $c(t) \in P_{T,M}$. Now we prove that c(t) belongs to $P_{T,M,\Psi}$. Let $T_1 > T$ be a real number. Denote $\Psi_1 := \min\{\psi, -T_1\}$. Let K be any compact subset of $M \setminus E$. If $K \cap \{\psi < -T_1\} \neq \emptyset$, as $\psi = \Psi_1$ on $\{\psi < -T_1\}$, it follows from $e^{-\varphi}c(-\psi)$ has a positive lower bound on K that $e^{-\varphi}c(-\Psi_1)$ has a positive lower bound on $K \cap \{\psi < -T_1\}$. If $K \cap \{\psi \ge -T_1\} \neq \emptyset$, as $\varphi + \psi$ is a plurisubharmonic function on M, then the function $e^{-\varphi}c(-\Psi_1) = e^{-\varphi - \psi}c(-\Psi_1)e^{\psi} \ge e^{-(\varphi + \psi)}c(T_1)e^{-T_1}$ has a positive lower bound on $K \cap \{\psi \ge -T_1\}$. Hence $e^{-\varphi}c(-\Psi_1)$ has a positive lower bound on any compact set $K \subset M \setminus E$. We know $c(t) \in P_{T,M,\Psi}$.

By the definition (1.1) of $G(t; c, \Psi, \varphi_{\alpha}, J, f)$ and the definition (1.4) of $G(t; c, \psi, \varphi, \mathcal{F}, f)$, we know that $G(t; c, \Psi, \varphi_{\alpha}, J, f) = G(t; c, \psi, \varphi, \mathcal{F}, f)$ for any $t \in [T, +\infty)$. As there exists a $t_0 \geq T$ such that $G(t_0; c, \psi, \varphi, f, \mathcal{F}) < +\infty$, we know that $G(t_0; c, \Psi, \varphi_{\alpha}, J, f) < +\infty$.

Then it follows from Theorem 1.2 that Corollary 1.7 holds.

6. Proof of Proposition 1.9

In this section, we prove Proposition 1.9 by using Theorem 1.2.

 $\begin{array}{l} \textit{Proof. Let } h(x) = \left\{ \begin{array}{l} e^{-\frac{1}{1-(x-1)^2}} & \text{if } |x-1| < 1 \\ 0 & \text{if } |x-1| \geq 1 \end{array} \right. \text{ be a real function defined on } \mathbb{R}, \\ \text{and let } g_n(x) = \frac{n}{(n+1)d} \int_0^{nx} h(s) ds, \text{ where } d = \int_{\mathbb{R}} h(s) ds. \text{ Note that } h(x) \in C_0^{\infty}(\mathbb{R}) \\ \text{and } h(x) \geq 0 \text{ for any } x \in \mathbb{R}. \text{ Then we get that } g_n(x) \text{ is increasing with respect to } x, \\ g_n(x) \leq g_{n+1}(x) \text{ for any } n \in \mathbb{N} \text{ and } x \in \mathbb{R}, \text{ and } \lim_{n \to +\infty} g_n(x) = \mathbb{I}_{\{s \in \mathbb{R}: s > 0\}}(x) \text{ for } \\ \text{any } x \in \mathbb{R}. \text{ Setting } c_t^n(x) = 1 - g_n(x-t), \text{ where } t \text{ is the given positive number in } \\ \text{Proposition 1.9, it follows from the properties of } \{g_n(x)\}_{n \in \mathbb{N}} \text{ that } c_t^n(x) \text{ is decreasing } \\ \text{with respect to } x, c_t^n(x) \geq c_t^{n+1}(x) \text{ for any } n \in \mathbb{N} \text{ and } x \in \mathbb{R}, \text{ and } \lim_{n \to +\infty} c_t^n(x) = \\ \mathbb{I}_{\{s \in \mathbb{R}: s \leq t\}}(x) \text{ for any } x \in \mathbb{R}. \text{ Let } \varphi_\alpha \equiv 0. \text{ Note that } c_t^n(x) \in [\frac{1}{n+1}, 1] \text{ on } (0, +\infty), \\ \text{then } c_t^n(x) \in P_{0,M,\Psi} \text{ for any } n \in \mathbb{N}. \\ \end{array} \right.$

$$\inf\left\{\int_{\{\Psi<-t\}} |\tilde{f}|^2 c_t^n(-\Psi) : \tilde{f} \in H^0(\{\Psi<-t\}, \mathcal{O}(K_M)) \\ \& (\tilde{f}-f)_{z_0} \in \mathcal{O}(K_M)_{z_0} \otimes I(\Psi)_{z_0} \text{ for any } z_0 \in Z_0\right\}$$

by $G_{t,n}(s)$. It follows from Theorem 1.2 that

$$\int_{\{\Psi < -l\}} |f|^2 c_t^n(-\Psi) \ge G_{t,n}(l) \ge \frac{\int_l^{+\infty} c_t^n(s) e^{-s} ds}{\int_0^{+\infty} c_t^n(s) e^{-s} ds} G_{t,n}(0)$$
(6.1)

for any l > 0. Following from $\int_{\{\Psi < -l\}} |f|^2 < +\infty$ for any l > 0, the properties of $\{c_t^n\}_{n \in \mathbb{N}}$ and the dominated convergence theorem, we obtain that

$$\lim_{n \to +\infty} \int_{\{\Psi < -l\}} |f|^2 c_t^n(-\Psi) = \int_{\{-t \le \Psi < -l\}} |f|^2.$$
(6.2)

As $c_t^n(x) \geq \mathbb{I}_{\{s \in \mathbb{R}: s \leq t\}}(x)$ for any x > 0 and $n \in \mathbb{N}$, then it follows from the definitions of $G_{t,n}(0)$ and $C_{f,\Psi,t}(Z_0)$ that

$$G_{t,n}(0) \ge C_{f,\Psi,t}(Z_0).$$
 (6.3)

Combining inequality (6.1), equality (6.2), and inequality (6.3), we obtain that

$$\int_{\{-t \le \Psi < -l\}} |f|^2 = \lim_{n \to +\infty} \int_{\{\Psi < -l\}} |f|^2 c_t^n (-\Psi)$$

$$\geq \lim_{n \to +\infty} \frac{\int_l^{+\infty} c_t^n (s) e^{-s} ds}{\int_0^{+\infty} c_t^n (s) e^{-s} ds} C_{f,\Psi,t}(Z_0)$$

$$= \frac{e^{-l} - e^{-t}}{1 - e^{-t}} C_{f,\Psi,t}(Z_0)$$

for any $l \in (0, t)$. Following from the difinition of $C_{f,\Psi,t}(Z_0)$, we have $\int_{\{-t \leq \Psi < 0\}} |f|^2 \geq C_{f,\Psi,t}(Z_0)$. Thus, we have

$$\int_{\{-t \le \Psi < -l\}} |f|^2 \ge \frac{e^{-l} - e^{-t}}{1 - e^{-t}} C_{f,\Psi,t}(Z_0) \tag{6.4}$$

for any $l \in [0, t)$.

Following from Fubini's Theorem and inequality (6.4), we obtain that

$$\begin{split} \int_{M_t} |f|^2 e^{-\Psi} &= \int_{M_t} \left(|f|^2 \int_0^{e^{-\Psi}} dr \right) \\ &= \int_0^{+\infty} \left(\int_{M_t \cap \{r < e^{-\Psi}\}} |f|^2 \right) dr \\ &= \int_{-\infty}^t \left(\int_{\{-t \le \Psi < \min\{-l,0\}\}} |f|^2 \right) e^l dl \\ &= \int_{-\infty}^0 \left(\int_{\{-t \le \Psi < \min\{-l,0\}\}} |f|^2 \right) e^l dl + \int_0^t \left(\int_{\{-t \le \Psi < -l\}} |f|^2 \right) e^l dl \\ &\ge C_{f,\Psi,t}(Z_0) \left(\int_{-\infty}^0 e^l dl + \int_0^t \frac{1 - e^{l-t}}{1 - e^{-t}} dl \right) \\ &= \frac{t}{1 - e^{-t}} C_{f,\Psi,t}(Z_0). \end{split}$$

Then Proposition 1.9 has thus been proved.

7. Proofs of Theorem 1.11, Theorem 1.13 and Remark 1.14

The following estimate will be used in the proofs of Theorem 1.11 and Theorem 1.13.

Proposition 7.1. Let φ be a plurisubharmonic function D, and let f be a holomorphic function on $\{\Psi < -t_0\}$ such that $f_o \in I(\varphi)_o$. Assume that $a_o^f(\Psi; \varphi) < +\infty$, then we have $a_o^f(\Psi; \varphi) > 0$ and

$$\frac{1}{r^2} \int_{\{a_o^f(\Psi;\varphi)\Psi < \log r\}} |f|^2 e^{-\varphi} \ge G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o, f) > 0$$

holds for any $r \in (0, e^{-a_o^f(\Psi; \varphi)t_0}]$, where the definition of $G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f)$ can be found in Section 1.1.

Proof. Lemma 2.10 tells us that there exists $p_0 > 2a_o^f(\Psi;\varphi)$ such that $I(p_0\Psi+\varphi)_o = I_+(2a_o^f(\Psi;\varphi)\Psi+\varphi)_o$. Following from the definition of $a_o^f(\Psi;\varphi)$ and Lemma 3.3, we obtain that

$$G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f) > 0.$$

$$(7.1)$$

Without loss of generality, assume that there exists $t > t_0$ such that $\int_{\{\Psi < -t\}} |f|^2 e^{-\varphi} < +\infty$. Denote that $t_1 := \inf\{t \ge t_0 : \int_{\{\Psi < -t\}} |f|^2 e^{-\varphi} < +\infty\}$. Denote

$$\inf\left\{\int_{\{p\Psi<-t\}} |\tilde{f}|^2 e^{-\varphi} : \tilde{f} \in \mathcal{O}(\{p\Psi<-t\}) \& (\tilde{f}-f)_o \in I(p\Psi+\varphi)_o\right\}$$

by $G_p(t)$, where $t \in [0, +\infty)$ and $p > 2a_o^f(\Psi; \varphi)$. Then we know that $G_p(0) \ge G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f)$ for any $p > 2a_o^f(\Psi; \varphi)$. Note that

$$p\Psi = \min\{p\psi + (2\lceil p \rceil - 2p)\log|F| - 2\log|F^{\lceil p \rceil}|, 0\},\$$

where $[p] = \min\{n \in \mathbb{Z} : n \ge p\}$, and

$$G_p(pt) \le \int_{\{\Psi < -t\}} |f|^2 e^{-\varphi} < +\infty$$

for any $t > t_1$. Theorem 1.2 shows that $G_p(-\log r)$ is concave with respect to $r \in (0,1]$ and $\lim_{t\to+\infty} G_p(t) = 0$, which implies that

$$\frac{1}{r_1^2} \int_{\{p\Psi < 2\log r_1\}} |f|^2 e^{-\varphi} \ge \frac{1}{r_1^2} G_p(-2\log r_1)
\ge G_p(0)
\ge G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f),$$
(7.2)

where $0 < r_1 \le e^{-\frac{pt_0}{2}}$.

We prove $a_o^f(\Psi; \varphi) > 0$ by contradiction: if $a_o^f(\Psi; \varphi) = 0$, as $\int_{\{\Psi < -t_1 - 1\}} |f|^2 e^{-\varphi} < +\infty$, it follows from the dominated convergence theorem and inequality (7.2) that

$$\frac{1}{r_1^2} \int_{\{\Psi=-\infty\}} |f|^2 e^{-\varphi} = \lim_{p \to 0+0} \frac{1}{r_1^2} \int_{\{p\Psi<2\log r_1\}} |f|^2 e^{-\varphi} \\
\geq G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f).$$
(7.3)

Note that $\mu(\{\Psi = -\infty\}) = \mu(\{\psi = -\infty\}) = 0$, where μ is the Lebesgue measure on \mathbb{C}^n . Inequality (7.3) implies that $G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o, f) = 0$, which contradicts inequality (7.1). Thus, we get that $a_o^f(\Psi; \varphi) > 0$.

which contradicts inequality (7.1). Thus, we get that $a_o^f(\Psi;\varphi) > 0$. For any $r_2 \in (0, e^{-a_o^f(\Psi;\varphi)t_1})$, note that $\frac{2\log r_2}{p} < -t_1$ for any $p \in (2a_0^f(\Psi;\varphi), -\frac{2\log r_2}{t_1})$, which implies that $\int_{\{p\Psi < 2\log r_2\}} |f|^2 e^{-\varphi} < +\infty$ for any $p \in (2a_0^f(\Psi;\varphi), -\frac{2\log r_2}{t_1})$. Then it follows from the dominated convergence theorem and inequality (7.2) that

$$\frac{1}{r_2^2} \int_{\{2a_0^f(\Psi;\varphi)\Psi \le 2\log r_2\}} |f|^2 e^{-\varphi} = \lim_{p \to 2a_0^f(\Psi;\varphi)+0} \frac{1}{r_2^2} \int_{\{p\Psi < 2\log r_2\}} |f|^2 e^{-\varphi} \\
\ge G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o, f).$$
(7.4)

For any $r \in (0, e^{-a_o^f(\Psi;\varphi)t_0}]$, if $r > e^{-a_o^f(\Psi;\varphi)t_1}$, we have $\int_{\{a_0^f(\Psi;\varphi)\Psi < \log r\}} |f|^2 e^{-\varphi} = +\infty > G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o, f)$, and if $r \in (0, e^{-a_o^f(\Psi;\varphi)t_1}]$, it follows from $\{a_o^f(\Psi;\varphi)\Psi < \log r\} = \bigcup_{0 < r_2 < r}\{a_o^f(\Psi;\varphi)\Psi < \log r_2\}$ and inequality (7.4) that

$$\begin{split} \int_{\{a_0^f(\Psi;\varphi)\Psi < \log r\}} |f|^2 e^{-\varphi} &= \sup_{r_2 \in (0,r)} \int_{\{2a_0^f(\Psi;\varphi)\Psi \le 2\log r_2\}} |f|^2 e^{-\varphi} \\ &\geq \sup_{r_2 \in (0,r)} r_2^2 G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o, f) \\ &= r^2 G(0; c \equiv 1, \Psi, \varphi, I_+(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o, f). \end{split}$$

Thus, Proposition 7.1 holds.

Proof of Theorem 1.11. It is clear that $I_+(a\Psi+\varphi)_o \subset I(a\Psi+\varphi)_o$, hence it suffices to prove that $I(a\Psi+\varphi) \subset I_+(a\Psi+\varphi)_o$.

If there exists $f_o \in I(a\Psi + \varphi)_o$ such that $f_o \notin I_+(a\Psi + \varphi)_o$, then $a_o^f(\Psi; \varphi)_o = \frac{a}{2} < +\infty$. Proposition 7.1 shows that a > 0. Without loss of generality, assume that f is a holomorphic function on $\{\Psi < -t_0\} \cap D$, where $t_0 > 0$. For any neighborhood $U \subset D$ of o, it follows from Proposition 7.1 that there exists $C_U > 0$ such that

$$\frac{1}{r^2} \int_{\{a\Psi < 2\log r\} \cap U} |f|^2 e^{-\varphi} \ge C_U \tag{7.5}$$

for any $r \in (0, e^{-\frac{at_0}{2}}]$. For any $t > at_0$, it follows from Fubini's Theorem and inequality (7.5) that

$$\int_{\{a\Psi<-t\}\cap U} |f|^2 e^{-a\Psi-\varphi} = \int_{\{a\Psi<-t\}\cap U} \left(|f|^2 e^{-\varphi} \int_0^{e^{-a\Psi}} dl \right)$$
$$= \int_0^{+\infty} \left(\int_{\{l< e^{-a\Psi}\}\cap \{a\Psi<-t\}\cap U} |f|^2 e^{-\varphi} \right) dl$$
$$\geq \int_{e^t}^{+\infty} \left(\int_{\{a\Psi<-\log l\}\cap U} |f|^2 e^{-\varphi} \right) dl$$
$$\geq C_U \int_{e^t}^{+\infty} \frac{1}{l} dl$$
$$= +\infty.$$

which contradicts $f_o \in I(a\Psi + \varphi)_o$. Thus, there is no $f_o \in I(a\Psi + \varphi)_o$ such that $f_o \notin I_+(a\Psi + \varphi)_o$, which implies that $I(a\Psi + \varphi) = I_+(a\Psi + \varphi)_o$ for any $a \ge 0$. \Box

We recall two lemmas, which will be used in the proof of Theorem 1.13.

Lemma 7.2 (see [31]). Let a(t) be a positive measurable function on $(-\infty, +\infty)$, such that $a(t)e^t$ is increasing near $+\infty$, and a(t) is not integrable near $+\infty$. Then there exists a positive measurable function $\tilde{a}(t)$ on $(-\infty, +\infty)$ satisfying the following statements:

(1) there exists $T < +\infty$ such that $\tilde{a}(t) \leq a(t)$ for any t > T;

(2) $\tilde{a}(t)e^t$ is strictly increasing and continuous near $+\infty$;

(3) $\tilde{a}(t)$ is not integrable near $+\infty$.

Lemma 7.3 (see [35]). For any two measurable spaces (X_i, μ_i) and two measurable functions g_i on X_i respectively $(i \in \{1, 2\})$, if $\mu_1(\{g_1 \ge r^{-1}\}) \ge \mu_2(\{g_2 \ge r^{-1}\})$ for any $r \in (0, r_0]$, then $\int_{\{g_1 \ge r_0^{-1}\}} g_1 d\mu_1 \ge \int_{\{g_2 \ge r_0^{-1}\}} g_2 d\mu_2$.

Proof of Theorem 1.13. We prove Theorem 1.13 in two cases, that a(t) satisfies condition (1) or condition (2).

Case (1). a(t) is decreasing near $+\infty$.

Firstly, we prove $(B) \Rightarrow (A)$. Consider $F \equiv f \equiv 1$, $\varphi \equiv 0$ and $\psi = \log |z_1|$ on the unit polydisc $\Delta^n \subset \mathbb{C}^n$. Note that $a_o^1(\log |z_1|; 0) = 1$ and

$$\int_{\Delta_{r_0}^n} a(-2\log|z_1|) \frac{1}{|z_1|^2} = (\pi r_0^2)^{n-1} \int_{\Delta_{r_0}} a(-2\log|z_1|) \frac{1}{|z_1|^2}$$
$$= (\pi r_0^2)^{n-1} 2\pi \int_0^{r_0} a(-2\log r) r^{-1} dr$$
$$= (\pi r_0^2)^{n-1} \pi \int_{-2\log r_0}^{+\infty} a(t) dt,$$

hence we obtain $(B) \Rightarrow (A)$.

Then, we prove $(A) \Rightarrow (B)$. Theorem 1.11 shows that $f_o \notin I(2a_o^f(\Psi; \varphi)\Psi + \varphi)_o$ and $a_o^f(\Psi; \varphi) > 0$. Now we assume that there exist $t_0 > 0$ and a pseudoconvex domain $D_0 \subset D$ containing o such that $\int_{\{\Psi < -t_0\} \cap D_0} |f|^2 e^{-2a_o^f(\Psi; \varphi)\Psi - \varphi} a(-2a_o^f(\Psi; \varphi)\Psi) < +\infty$ to get a contradiction. As $f_o \in I(\varphi)_o$, there exist $t_1 > t_0$ and a pseudoconvex domain $D_1 \subset D_0$ containing o such that $\int_{D_1 \cap \{\Psi < -t_1\}} |f|^2 e^{-\varphi} < +\infty$. Set $c(t) = a(t)e^t + 1$, then we have

$$\int_{D_1 \cap \{\Psi < -t_1\}} |f|^2 e^{-\varphi} c(-2a_o^f(\Psi;\varphi)\Psi) < +\infty.$$
(7.6)

Without loss of generality, assume that a(t) is decreasing on $(2a_o^f(\Psi; \varphi)t_1, +\infty)$. Note that $c(t)e^{-t} = a(t)+e^{-t}$ is decreasing on $(2a_o^f(\Psi; \varphi)t_1, +\infty)$ and $\liminf_{t\to+\infty} c(t) > 0$. As a(t) is not integrable near $+\infty$, so is $c(t)e^{-t}$. Note that there exist a plurisub-harmonic function ψ_1 and a holomorphic function F_1 on D_1 such that

$$|\psi_1 - 2\log|F_1| = 2a_o^f(\Psi;\varphi)(\psi - 2\log|F|)$$

on D_1 . Denote that $\Psi_1 := \min\{\psi_1 - 2\log|F_1|, -2a_o^f(\Psi;\varphi)t_1\}$ on D_1 . Using Remark 1.3 (replacing M, Ψ and T by D_1 , Ψ_1 and $2a_o^f(\Psi;\varphi)t_1$ respectively), as $f_o \notin I(2a_o^f(\Psi;\varphi)\Psi + \varphi)_o = I(\Psi_1 + \varphi)_o$, then we have $G(2a_o^f(\Psi;\varphi)t_1;c,\Psi_1,\varphi,I(\Psi_1 + \varphi)_o,f) = +\infty$, which contradicts to inequality (7.6). Thus, we obtain $(A) \Rightarrow (B)$.

Case (2). $a(t)e^t$ is increasing near $+\infty$.

In this case, the proof of $(B) \Rightarrow (A)$ is the same as the case (1), therefore it suffices to prove $(A) \Rightarrow (B)$.

Assume that statement (A) holds. It follows from Lemma 7.2 that there exists a positive function $\tilde{a}(t)$ on $(-\infty, +\infty)$ satisfying that: $\tilde{a}(t) \leq a(t)$ near $+\infty$; $\tilde{a}(t)e^t$ is strictly increasing and continuous near $+\infty$; $\tilde{a}(t)$ is not integrable near $+\infty$. Thus, it suffices to prove that for any Ψ , φ and $f_o \in I(\varphi)_o$ satisfying $a_o^f(\Psi; \varphi) < +\infty$,

 $\tilde{a}(-2a_{\rho}^{f}(\Psi;\varphi)\Psi)\exp(-2a_{\rho}^{f}(\Psi;\varphi)\Psi-\varphi+2\log|f|)\notin L^{1}(U\cap\{\Psi<-t\}),$ where U is any neighborhood of o and t > 0.

Take any $t_0 \gg 0$ and any small pseudoconvex domain $D_0 \subset D$ containing the origin o such that $f \in \mathcal{O}(D_0 \cap \{\Psi < -t_0\})$. Let $\mu_1(X) = \int_X |f|^2 e^{-\varphi}$, where X is a Lebesgue measurable subset of $D_0 \cap \{\Psi < -t_0\}$, and let μ_2 be the Lebesgue measure on (0,1]. Denote that $Y_r = \{-2a_o^f(\Psi;\varphi)\Psi \ge -\log r\}$. Proposition 7.1 shows that there exists a positive constant C such that $\mu_1(Y_r) \geq Cr$ holds for any $r \in (0, e^{-2a_o^f(\Psi;\varphi)t_0}].$

Let $g_1 = \tilde{a}(-2a_o^f(\Psi;\varphi)\Psi) \exp(-2a_o^f(\Psi;\varphi)\Psi)$ and $g_2(x) = \tilde{a}(-\log x + \log C)Cx^{-1}$. As $\tilde{a}(t)e^{-t}$ is strictly increasing near $+\infty$, then $g_1 \ge \tilde{a}(-\log r)r^{-1}$ on Y_r implies that

$$\mu_1(\{g_1 \ge \tilde{a}(-\log r)r^{-1}\}) \ge \mu_1(Y_r) \ge Cr$$
(7.7)

holds for any r > 0 small enough. As $\tilde{a}(t)e^t$ is strictly increasing near $+\infty$, then there exists $r_0 \in (0, e^{-2a_o^f(\Psi)t_0})$ such that

$$\mu_2(\{x \in (0, r_0] : g_2(x) \ge \tilde{a}(-\log r)r^{-1}\}) = \mu_2(\{0 < x \le Cr\}) = Cr$$
(7.8)

for any $r \in (0, r_0]$. As $\tilde{a}(-\log r)r^{-1}$ converges to $+\infty$ (when $r \to 0+0$) and $\tilde{a}(t)$ is continuous near $+\infty$, we obtain that

$$\mu_1(\{g_1 \ge r^{-1}\}) \ge \mu_2(\{x \in (0, r_0] : g_2(x) \ge r^{-1}\})$$

holds for any r > 0 small enough. Following from Lemma 7.3 and $\tilde{a}(t)$ is not integrable near $+\infty$, we obtain $\tilde{a}(-2a_{\rho}^{2}(\Psi;\varphi)\Psi)\exp(-2a_{\rho}^{2}(\Psi;\varphi)\Psi-\varphi+2\log|f|) \notin$ $L^1(D_0 \cap \{\Psi < -t_0\}).$

Thus, Theorem 1.13 holds.

Lemma 7.4. Let h be a holomorphic function on a pseudoconvex domain $D \subset \mathbb{C}^n$ containing the origin o, and let ψ be a plurisubharmonic function on D such that $c^h_{o,p}(\psi) < +\infty$, where p > 0. Then we have $a^1_o(\Psi; 0) = \frac{1}{2}$, where $\Psi = 0$ $\min\{2c_{o,p}^{h}(\psi)\psi - p\log|h|, 0\}.$

Proof. As ψ is a plurisubharmonic function, there exist $\delta > 0$ and a neighborhood U of o such that $\int_U e^{-\delta\psi} < +\infty$. For any $r \in (0,1)$, there exist $r' \in (0,r)$ and a neighborhood $U' \Subset U$ of o such that $2\frac{r-r'}{1-r}c^h_{o,p}(\psi) < \delta$ and $\int_{U'} |h|^p e^{-2\frac{r'}{r}c^h_{o,p}(\psi)\psi} < \delta$ $+\infty$. Following from Hölder Inequality, we get that

$$\begin{split} \int_{U'} |h|^{pr} e^{-2rc_{o,p}^{h}(\psi)\psi} &= \int_{U'} |h|^{pr} e^{-2r'c_{o,p}^{h}(\psi)\psi} e^{-2(r-r')c_{o,p}^{h}(\psi)\psi} \\ &\leq \left(\int_{U'} |h|^{p} e^{-2\frac{r'}{r}c_{o,p}^{h}(\psi)\psi}\right)^{r} \times \left(\int_{U'} e^{-2\frac{r-r'}{1-r}c_{o,p}^{h}(\psi)\psi}\right)^{1-r} \\ &< +\infty, \end{split}$$

which implies that $a_o^1(\Psi; 0) \geq \frac{1}{2}$. We prove $a_o^1(\Psi; 0) = \frac{1}{2}$ by contradiction: if not, there exist $s > \frac{1}{2}$, t > 0 and a neighborhood $V \in U$ of o such that

$$\int_{\{\Psi < -t\} \cap V} |h|^{2sp} e^{-4sc_{o,p}^{h}(\psi)\psi} = \int_{\{\Psi < -t\} \cap V} e^{-2s\Psi} < +\infty.$$
(7.9)

There exists s' > 1 such that $\frac{4s(s'-1)}{2s-1}c_{o,p}^{h}(\psi) < \delta$. Following from inequality (7.9) and Hölder Inequality, we get that

$$\begin{split} &\int_{\{\Psi<-t\}\cap V} |h|^p e^{-2s'c_{o,p}^h(\psi)\psi} \\ &= \int_{\{\Psi<-t\}\cap V} |h|^p e^{-2c_{o,p}^h(\psi)\psi} e^{-2(s'-1)c_{o,p}^h(\psi)\psi} \\ &\leq \left(\int_{\{\Psi<-t\}\cap V} |h|^{2sp} e^{-4sc_{o,p}^h(\psi)\psi}\right)^{\frac{1}{2s}} \times \left(\int_{\{\Psi<-t\}\cap V} e^{-\frac{4s(s'-1)}{2s-1}c_{o,p}^h(\psi)\psi}\right)^{1-\frac{1}{2s}} \\ &< +\infty, \end{split}$$

which contradicts to the definition of $c_{o,p}^h(\psi)$. Thus, we obtain $a_o^1(\Psi; 0) = \frac{1}{2}$. \Box

Proof of Remark 1.14. We give a proof of Theorem 1.12 by using Theorem 1.13. The proofs of $(B') \Rightarrow (A')$ and $(C') \Rightarrow (A')$ are as the same as the proof of $(B) \Rightarrow (A)$ in Theorem 1.13. It suffices to prove $(A') \Rightarrow (B')$ and $(A') \Rightarrow (C')$.

Assume that statement (A') holds, then we have the statement (B) in Theorem 1.13 holds.

For any h and ψ satisfying $c_{o,p}^{h}(\psi) < +\infty$, let $F \equiv 1$, $f = h^{\lceil \frac{p}{2} \rceil}$ and $\varphi = (2\lceil \frac{p}{2} \rceil - p) \log |h|$, where $\lceil m \rceil := \min\{n \in \mathbb{Z} : n \ge m\}$. Note that $|h|^{p} = |f|^{2}e^{-\varphi}$, $\psi(o) = -\infty$ and $a_{o}^{f}(\Psi; \varphi) = c_{o,p}^{h}(\psi)$. The holding of statement (B) in Theorem 1.13 implies that $a(-2c_{o,p}^{h}(\psi)\psi) \exp(-2c_{o,p}^{h}(\psi)\psi + p \log |h|)$ is not integrable near o.

For any h and ψ satisfying $c_{o,p}^{h}(\psi) < +\infty$, let $\Psi = \min\{2c_{o,p}^{h}(\psi)\psi - p\log|h|, 0\}$, $\varphi \equiv 0$ and $f \equiv 1$. Lemma 7.4 shows that $a_{o}^{f}(\Psi;\varphi) = \frac{1}{2}$. The holding of statement (B) in Theorem 1.13 implies that $a(-\Psi)\exp(-\Psi) \notin L^{1}(U \cap \{\Psi < 0\})$ for any neighborhood U of o, which implies that $a(-2c_{o,p}^{h}(\psi)\psi + p\log|h|)\exp(-2c_{o,p}^{h}(\psi)\psi + p\log|h|)\exp(-2c_{o,p}^{h}(\psi)\psi + p\log|h|)$ is not integrable near o.

8. Proof of Theorem 1.15

In this section, we prove Theorem 1.15 by using Theorem 1.2.

Proof of Theorem 1.15. Following from Fubini's Theorem, we have

$$\begin{split} &\int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi} \\ &= \int_{\{\Psi<0\}} \left(|f|^2 e^{-\varphi-\Psi+a\Psi} \int_0^{e^{-a\Psi}} ds \right) \\ &= \int_0^{+\infty} \left(\int_{\{\Psi<0\}\cap\{s< e^{-a\Psi}\}} |f|^2 e^{-\varphi-\Psi+a\Psi} \right) ds \qquad (8.1) \\ &= \int_{-\infty}^{+\infty} \left(\int_{\{\Psi<-\frac{l}{a}\}\cap\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi+a\Psi} \right) e^l dl \\ &= \int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi+a\Psi} + \int_0^{+\infty} \left(\int_{\{\Psi<-\frac{l}{a}\}} |f|^2 e^{-\varphi-\Psi+a\Psi} \right) e^l dl. \end{split}$$

Denote

$$\inf\left\{\int_{\{q'\Psi<-t\}}|\tilde{f}|^2e^{-\varphi}c(-q'\Psi):(\tilde{f}-f)_o\in I(q'\Psi+\varphi)_o\,\&\,\tilde{f}\in\mathcal{O}(\{q'\Psi<-t\})\right\}$$

by $G_{c,q'}(t)$, where c is a Lebesgue measurable function on $(0, +\infty)$ and $q' > 2a_o^f(\Psi; \varphi) \ge 1$.

Next we prove inequality

$$\int_{\{\Psi < -\frac{l}{a}\}} |f|^2 e^{-\varphi - \Psi + a\Psi} \ge e^{-\frac{a-1+q'}{a}l} \frac{1}{K_{\Psi,f,a}(o)}$$
(8.2)

holds for $l \ge 0$, a > 0 and $q' > 2a_o^f(\Psi; \varphi)$.

We prove inequality (8.2) for the case $a \in (0,1]$. Let $c_1(t) = e^{\frac{1-a}{q'}t}$ on $(0, +\infty)$. Note that $c_1(t)e^{-t}$ is decreasing on $(0, +\infty)$ and $e^{-\varphi}c_1(-q'\Psi) = e^{-\varphi-(1-a)\Psi}$ has a positive lower bound on any compact subset of D. Theorem 1.2 shows that $G_{c_1,q'}(h^{-1}(r))$ is concave with respect to r, where $h(t) = \int_t^{+\infty} c_1(s)e^{-s}ds$. Note that $G_{c_1,q'}(0) \geq \frac{1}{K_{\Psi,f,a}(o)}$ for any $q' > 2a_o^f(\Psi; \varphi)$. Hence we have

$$\begin{split} \int_{\{\Psi < -\frac{l}{a}\}} |f|^2 e^{-\varphi - (1-a)\Psi} \geq &G_{c_1,q'}\left(\frac{q'l}{a}\right) \\ \geq &\frac{\int_{\frac{q'l}{a}}^{+\infty} c_1(s) e^{-s} ds}{\int_0^{+\infty} c_1(s) e^{-s} ds} G_{c_1,q'}(0) \\ \geq &e^{-\frac{a-1+q'}{a}l} \frac{1}{K_{\Psi,f,a}(o)}. \end{split}$$

We prove inequality (8.2) for the case a > 1. Let $\tilde{c}_m(t)$ be a continuous function on $(0, +\infty)$ such that $\tilde{c}_m(t) = e^{\frac{1-a}{q'}t}$ on (0, m) and $\tilde{c}_m(t) = e^{\frac{1-a}{q'}m}$ on $(m, +\infty)$ for any positive integer m. Note that $\tilde{c}_m(t)e^{-t}$ is decreasing on $(0, +\infty)$ and $e^{-\varphi}\tilde{c}_m(-q'\Psi)$ has a positive lower bound on any compact subset of D. Theorem 1.2 shows that $G_{\tilde{c}_m,q'}(h_m^{-1}(r))$ is concave with respect to r, where $h_m(t) = \int_t^{+\infty} \tilde{c}_m(s)e^{-s}ds$. Note that $G_{\tilde{c}_m,q'}(0) \geq \frac{1}{K_{\Psi,f,a}(o)}$ for any $q' > 2a_o^f(\Psi;\varphi)$. Hence we have

$$\int_{\{\Psi<-\frac{l}{a}\}} |f|^2 e^{-\varphi} \tilde{c}_m(-q'\Psi) \ge G_{\tilde{c}_m,q'}\left(\frac{q'l}{a}\right) \\
\ge \frac{\int_{\frac{q'l}{a}}^{+\infty} \tilde{c}_m(s)e^{-s}ds}{\int_0^{+\infty} \tilde{c}_m(s)e^{-s}ds} G_{\tilde{c}_m,q'}(0) \\
\ge \frac{\int_{\frac{q'l}{a}}^{+\infty} \tilde{c}_m(s)e^{-s}ds}{\int_0^{+\infty} \tilde{c}_m(s)e^{-s}ds} \frac{1}{K_{\Psi,f,a}(o)}.$$
(8.3)

As $\int_{\{\Psi < 0\}} |f|^2 e^{-\varphi - \Psi} \leq C_1 < +\infty$, it follows from $\tilde{c}_m(-q'\Psi) \leq e^{-\Psi}$, the dominated convergence theorem and inequality (8.3) that

$$\begin{split} \int_{\{\Psi<-\frac{l}{a}\}} |f|^2 e^{-\varphi-(1-a)\Psi} &= \lim_{m \to +\infty} \int_{\{\Psi<-\frac{l}{a}\}} |f|^2 e^{-\varphi} \tilde{c}_m(-q'\Psi) \\ &\geq \lim_{m \to +\infty} \frac{\int_{q'a}^{+\infty} \tilde{c}_m(s) e^{-s} ds}{\int_0^{+\infty} \tilde{c}_m(s) e^{-s} ds} \frac{1}{K_{\Psi,f,a}(o)} \\ &= e^{-\frac{a-1+q'}{a}} l \frac{1}{K_{\Psi,f,a}(o)}. \end{split}$$

Combining equality (8.1), inequality (8.2) and the definition of $K_{\Psi,f,a}(o)$, we obtain that

$$\int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi}
= \int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi+a\Psi} + \int_0^{+\infty} \left(\int_{\{\Psi<-\frac{l}{a}\}} |f|^2 e^{-\varphi-\Psi+a\Psi} \right) e^l dl
\ge \left(1 + \int_0^{+\infty} e^{-\frac{-1+q'}{a}l}\right) \frac{1}{K_{\Psi,f,a}(o)}
= \frac{a+q'-1}{q'-1} \cdot \frac{1}{K_{\Psi,f,a}(o)}$$
(8.4)

for any $q' > 2a_o^f(\Psi; \varphi)$. Let $q' \to 2a_o^f(\Psi; \varphi)$, we get that inequality (8.4) also holds when $q' = 2a_o^f(\Psi; \varphi)$. Thus, if q > 1 satisfies

$$\frac{q+a-1}{q-1} > \frac{C_1}{C_2} \ge K_{\Psi,f,a}(o) \int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi},$$

$$\int_{\{\Psi<0\}} |f|^2 e^{-\varphi-\Psi},$$

we have $p < 2a_o^f(\Psi; \varphi)$, i.e. $f_o \in I(p\Psi + \varphi)_o$

45

9. Appendix: Proof of Lemma 2.1

In this section, we prove Lemma 2.1.

9.1. Some results used in the proof of Lemma 2.1.

Lemma 9.1 (see [11]). Let Q be a Hermitian vector bundle on a Kähler manifold M of dimension n with a Kähler metric ω . Assume that $\eta, g > 0$ are smooth functions on M. Then for every form $v \in D(M, \wedge^{n,q}T^*M \otimes Q)$ with compact support we have

$$\int_{M} (\eta + g^{-1}) |D^{''*}v|_{Q}^{2} dV_{M} + \int_{M} \eta |D^{''}v|_{Q}^{2} dV_{M} \\
\geq \int_{M} \langle [\eta \sqrt{-1}\Theta_{Q} - \sqrt{-1}\partial \bar{\partial}\eta - \sqrt{-1}g \partial \eta \wedge \bar{\partial}\eta, \Lambda_{\omega}]v, v \rangle_{Q} dV_{M}.$$
(9.1)

Lemma 9.2 (Lemma 4.2 in [32]). Let M and Q be as in the above lemma and θ be a continuous (1,0) form on M. Then we have

$$[\sqrt{-1}\theta \wedge \bar{\theta}, \Lambda_{\omega}]\alpha = \bar{\theta} \wedge (\alpha_{\vdash}(\bar{\theta})^{\sharp}), \qquad (9.2)$$

for any (n,1) form α with value in Q. Moreover, for any positive (1,1) form β , we have $[\beta, \Lambda_{\omega}]$ is semipositive.

Lemma 9.3 (Remark 3.2 in [11]). Let (M, ω) be a complete Kähler manifold equipped with a (non-necessarily complete) Kähler metric ω , and let Q be a Hermitian vector bundle over M. Assume that η and g are smooth bounded positive functions on M and let $B := [\eta \sqrt{-1}\Theta_Q - \sqrt{-1}\partial\bar{\partial}\eta - \sqrt{-1}g\partial\eta \wedge \bar{\partial}\eta, \Lambda_\omega]$. Assume that $\delta \geq 0$ is a number such that $B + \delta I$ is semi-positive definite everywhere on $\wedge^{n,q}T^*M \otimes Q$ for some $q \geq 1$. Then given a form $v \in L^2(M, \wedge^{n,q}T^*M \otimes Q)$ such that D''v = 0 and $\int_M \langle (B+\delta I)^{-1}v, v \rangle_Q dV_M < +\infty$, there exists an approximate solution $u \in L^2(M, \wedge^{n,q-1}T^*M \otimes Q)$ and a correcting term $h \in L^2(M, \wedge^{n,q}T^*M \otimes Q)$ such that $D''u + \sqrt{\delta}h = v$ and

$$\int_{M} (\eta + g^{-1})^{-1} |u|_{Q}^{2} dV_{M} + \int_{M} |h|_{Q}^{2} dV_{M} \le \int_{M} \langle (B + \delta I)^{-1} v, v \rangle_{Q} dV_{M}.$$
(9.3)

Lemma 9.4 (Theorem 6.1 in [10], see also Theorem 2.2 in [52]). Let (M, ω) be a complex manifold equipped with a Hermitian metric ω , and $\Omega \subset \subset M$ be an open set. Assume that $T = \tilde{T} + \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \varphi$ is a closed (1,1)-current on M, where \tilde{T} is a smooth real (1,1)-form and φ is a quasi-plurisubharmonic function. Let γ be a continuous real (1,1)-form such that $T \geq \gamma$. Suppose that the Chern curvature tensor of TM satisfies

$$(\sqrt{-1}\Theta_{TM} + \varpi \otimes Id_{TM})(\kappa_1 \otimes \kappa_2, \kappa_1 \otimes \kappa_2) \ge 0$$

$$\forall \kappa_1, \kappa_2 \in TM \quad with \quad \langle \kappa_1, \kappa_2 \rangle = 0$$
(9.4)

for some continuous nonnegative (1,1)-form ϖ on M. Then there is a family of closed (1,1)-current $T_{\zeta,\rho} = \widetilde{T} + \frac{\sqrt{-1}}{\pi} \partial \overline{\partial} \varphi_{\zeta,\rho}$ on M ($\zeta \in (0, +\infty)$) and $\rho \in (0, \rho_1)$ for some positive number ρ_1) independent of γ , such that

(i) $\varphi_{\zeta,\rho}$ is quasi-plurisubharmonic on a neighborhood of Ω , smooth on $M \setminus E_{\zeta}(T)$, increasing with respect to ζ and ρ on Ω and converges to φ on Ω as $\rho \to 0$.

(*ii*) $T_{\zeta,\rho} \geq \gamma - \zeta \varpi - \delta_{\rho} \omega$ on Ω .

where $E_{\zeta}(T) := \{x \in M : v(T, x) \ge \zeta\}$ ($\zeta > 0$) is the ζ -upper level set of Lelong numbers and $\{\delta_{\rho}\}$ is an increasing family of positive numbers such that $\lim_{\rho \to 0} \delta_{\rho} = 0$.

Remark 9.5 (see Remark 2.1 in [52]). Lemma 9.4 is stated in [10] in the case M is a compact complex manifold. The similar proof as in [10] shows that Lemma 9.4 on noncompact complex manifold still holds where the uniform estimate (i) and (ii) are obtained only on a relatively compact subset Ω .

Lemma 9.6 (Theorem 1.5 in [9]). Let M be a Kähler manifold, and Z be an analytic subset of M. Assume that Ω is a relatively compact open subset of M possessing a complete Kähler metric. Then $\Omega \setminus Z$ carries a complete Kähler metric.

Lemma 9.7 (Lemma 6.9 in [9]). Let Ω be an open subset of \mathbb{C}^n and Z be a complex analytic subset of Ω . Assume that v is a (p,q-1)-form with L^2_{loc} coefficients and h is a (p,q)-form with L^1_{loc} coefficients such that $\overline{\partial}v = h$ on $\Omega \setminus Z$ (in the sense of distribution theory). Then $\overline{\partial}v = h$ on Ω .

Let M be a complex manifold. Let ω be a continuous Hermitian metric on M. Let dV_M be a continuous volume form on M. We denote by $L^2_{p,q}(M, \omega, dV_M)$ the spaces of L^2 integrable (p,q) forms over M with respect to ω and dV_M . It is known that $L^2_{p,q}(M, \omega, dV_M)$ is a Hilbert space.

Lemma 9.8. Let $\{u_n\}_{n=1}^{+\infty}$ be a sequence of (p,q) forms in $L^2_{p,q}(M,\omega,dV_M)$ which is weakly convergent to u. Let $\{v_n\}_{n=1}^{+\infty}$ be a sequence of Lebesgue measurable real functions on M which converges pointwise to v. We assume that there exists a constant C > 0 such that $|v_n| \leq C$ for any n. Then $\{v_n u_n\}_{n=1}^{+\infty}$ weakly converges to vu in $L^2_{p,q}(M, \omega, dV_M)$.

Proof. Let $g \in L^2_{p,q}(M, \omega, dV_M)$. Consider

$$\begin{split} I &= |\langle v_n u_n, g \rangle - \langle v u, g \rangle| \\ &= |\int_M (v_n u_n, g)_\omega dV_M - \int_M (v u, g)_\omega dV_M| \\ &\leq |\int_M (v_n u_n - v u_n, g)_\omega dV_M| + |\int_M (v u_n - v u, g)_\omega dV_M| \\ &= |\int_M (u_n, v_n g - v g)_\omega dV_M| + |\int_M (u_n - u, v g)_\omega dV_M| \\ &\leq ||u_n|| \cdot ||v_n g - v g|| + |\int_M (u_n - u, v g)_\omega dV_M|. \end{split}$$

Denote $I_1 := ||u_n|| \cdot ||v_n g - vg||$ and $I_2 := |\int_M (u_n - u, vg)_\omega dV_M|$. It follows from $\{u_n\}_{n=1}^{+\infty}$ weakly converges to u that $||u_n||$ is uniformly bounded with respect to n. Note that $|v_n|$ is uniformly bounded with respect to n. We know |v| < C and then $vg \in L^2_{p,q}(M, \omega, dV_M)$. Hence we have $I_2 \to 0$ as $n \to +\infty$. It follows from Lebesgue dominated convergence theorem that we have $\lim_{n \to +\infty} I_1 = 0$.

Hence $\lim_{n \to +\infty} I = 0$ and we know $\{v_n u_n\}_{n=1}^{+\infty}$ weakly converges to vu in $L^2_{p,q}(M, \omega, dV_M)$.

The following notations can be referred to [3].

Let X be a complex manifold. An upper semi-continuous function $u: X \to [-\infty, +\infty)$ is quasi-plurisubharmonic if it is locally of the form $u = \varphi + f$ where φ is plurisubharmonic and f is smooth. Let θ be a closed, real (1, 1) form on X. By Poincaré lemma, θ is locally of the form $\theta = dd^c f$ for a smooth real-valued function f which is called a local potential of θ . We call a quasi-plurisubharmonic function u is θ -plurisubharmonic if $\theta + dd^c u \geq 0$ in the sense of currents.

Lemma 9.9 (see [12], see also [3]). For arbitrary $\eta = (\eta_1, \ldots, \eta_p) \in (0, +\infty)^p$, the function

$$M_{\eta}(t_1,\ldots,t_p) = \int_{\mathbb{R}^p} \max\{t_1+h_1,\ldots,t_p+h_p\} \prod_{1 \le j \le p} \theta(\frac{h_j}{\eta_j}) dh_1 \ldots dh_p$$

possesses the following properties:

(1) $M_{\eta}(t_1, \ldots, t_p)$ is non decreasing in all variables, smooth and convex on \mathbb{R}^p ; (2) $\max\{t_1, \ldots, t_p\} \leq M_{\eta}(t_1, \ldots, t_p) \leq \max\{t_1 + \eta_1, \ldots, t_p + \eta_p\};$

(3)
$$M_{\eta}(t_1, \ldots, t_p) = M_{\eta_1, \ldots, \hat{\eta}_j, \ldots, \eta_p}(t_1, \ldots, \hat{t}_j, \ldots, t_p)$$
 if $t_j + \eta_j \le \max_{j \le j \le j} \{t_k - \eta_k\}$;

(4) $M_{\eta}(t_1 + a, \dots, t_p + a) = M_{\eta}(t_1, \dots, t_p) + a \text{ for any } a \in \mathbb{R};$

(5) if u_1, \ldots, u_p are plurisubharmonic functions, then $u = M_\eta(u_1, \ldots, u_p)$ is plurisubharmonic;

(6) if u_1, \ldots, u_p are θ -plurisubharmonic functions, then $u = M_\eta(u_1, \ldots, u_p)$ is θ -plurisubharmonic function.

Proof. The proof of (1)-(5) can be referred to [12] and the proof of (6) can be referred to [3]. For the convenience of the readers, we recall the proof of (6).

Let f be a local potential of θ . We know $f + u_i$ is plurisubharmonic function. It follows from (4) and (5) that $M_{\eta}(u_1 + f, \ldots, u_p + f) = M_{\eta}(u_1, \ldots, u_p) + f$ is plurisubharmonic. Hence $u = M_{\eta}(u_1, \ldots, u_p)$ is θ -plurisubharmonic function. \Box

9.2. Proof of Lemma 2.1. Now we begin to prove Lemma 2.1.

Note that $M \setminus \{F = 0\}$ is a weakly pseudoconvex Kähler manifold. The following remark shows that we can assume that F has no zero points on M.

Remark 9.10. Assume that there exists a holomorphic (n,0) form \hat{F} on $M \setminus \{F =$ 0 such that

$$\int_{M \setminus \{F=0\}} |\hat{F} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq \left(\frac{1}{\delta}c(T)e^{-T} + \int_T^{t_0+B} c(s)e^{-s}ds\right) \int_{M \setminus \{F=0\}} \frac{1}{B} \mathbb{I}_{\{-t_0-B < \Psi < -t_0\}} |f|^2 e^{-\varphi_\alpha - \Psi}.$$

As $v_{t_0,B}(\Psi) \geq \Psi$ and $c(t)e^{-t}$ is decreasing with respect to t, we have $e^{-\varphi+v_{t_0,B}(\Psi)-\Psi}c(-v_{t_0,B}(\Psi)) \ge e^{-\varphi}c(-\Psi) = e^{-\varphi_{\alpha}}c(-\Psi)e^{-(1+\delta)\max\{\psi+T, 2\log|F|\}}.$

Let K be any compact subset of M. Note that $\varphi + \Psi$ is plurisubharmonic function on M, $v_{t_0,B}(t) \geq -t_0 - B$ and $c(t)e^{-t}$ is decreasing with respect to t. Then we have

$$\begin{split} &\int_{(M\setminus\{F=0\})\cap K} |\hat{F}|^2 \\ \leq & 2\int_{(M\setminus\{F=0\})\cap K} |(1-b_{t_0,B}(\Psi))fF^{1+\delta}|^2 + 2\int_{(M\setminus\{F=0\})\cap K} |\hat{F}-(1-b_{t_0,B}(\Psi))fF^{1+\delta}|^2 \\ \leq & 2\left(\sup_K |F^{1+\delta}|^2\right) \int_{\{\Psi<-t_0\}\cap K} |f|^2 \\ &+ \frac{2}{M_K} \int_{(M\setminus\{F=0\})\cap K} |\hat{F}-(1-b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi+v_{t_0,B}(\Psi)-\Psi} c(-v_{t_0,B}(\Psi)) \\ < & +\infty, \end{split}$$

where M_K is a positive number. As K is arbitrarily chosen, we know that there exists a holomorphic (n,0) form \tilde{F} on M such that $\tilde{F} = \hat{F}$ on $M \setminus \{F = 0\}$. And we have

$$\int_{M} |\tilde{F} - (1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 e^{-\varphi + v_{t_0,B}(\Psi) - \Psi} c(-v_{t_0,B}(\Psi))$$

$$\leq \left(\frac{1}{\delta}c(T)e^{-T} + \int_{T}^{t_0+B} c(s)e^{-s}ds\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0-B<\Psi<-t_0\}} |f|^2 e^{-\varphi_{\alpha}-\Psi}$$

The following remark shows that we can assume that c(t) is a smooth function.

Remark 9.11. We firstly introduce the regularization process of c(t).

Let $f(x) = 2\mathbb{I}_{(-\frac{1}{2},\frac{1}{2})} * \rho(x)$ be a smooth function on \mathbb{R} , where ρ is the kernel of

convolution satisfying $supp(\rho) \subset (-\frac{1}{3}, \frac{1}{3})$ and $\rho > 0$. Let $g_i(x) = \begin{cases} if(ix) & \text{if } x \leq 0\\ if(i^2x) & \text{if } x > 0 \end{cases}$, then $\{g_i\}_{i \in \mathbb{N}^+}$ is a family of smooth function.

tions on \mathbb{R} satisfying: (1) $supp(g) \subset \left[-\frac{1}{i}, \frac{1}{i}\right], g_i(x) \ge 0$ for any $x \in \mathbb{R}$,

(2) $\int_{-\frac{1}{i}}^{0} g_i(x) dx = 1$, $\int_{0}^{\frac{1}{i}} g_i(x) dx \leq \frac{1}{i}$ for any $i \in \mathbb{N}^+$. Let $\tilde{h}(t)$ be an extension of the function $c(t)e^{-t}$ from $[T, +\infty)$ to \mathbb{R} such that (1) $\tilde{h}(t) = h(t) := c(t)e^{-t}$ on $[T, +\infty)$; (2) $\tilde{h}(t)$ is decreasing with respect to t; (3) $\lim_{t \to T-0} \tilde{h}(t) = c(T)e^{-T}$.

Denote $c_i(t) := e^t \int_{\mathbb{R}} \tilde{h}(t+y)g_i(y)dy$. By the construction of convolution, we know $c_i(t) \in C^{\infty}(-\infty, +\infty)$. For any $t \geq T$, we have

$$c_i(t) - c(t) \ge e^t \left(\int_{-\frac{1}{i}}^0 (\tilde{h}(t+y) - \tilde{h}(t))g_i(y)dy \right) \ge 0.$$

As $\tilde{h}(t)$ is decreasing with respect to t, we know that $c_i(t)e^{-t}$ is also decreasing with respect to t. Hence $c_i(t)e^{-t}$ is locally L^1 integrable on \mathbb{R} .

As $\tilde{h}(t)$ is decreasing with respect to t, then set $\tilde{h}^-(t) = \lim_{s \to t^-0} \tilde{h}(s) \ge h(t)$ for any $t \in \mathbb{R}$. Note that $c^-(t) := \lim_{s \to t^-0} \tilde{h}(s)e^t \ge c(t)$ for any $t \ge T$.

Now we prove $\lim_{i \to +\infty} c_i(t)e^{-t} = \tilde{h}^-(t)$. In fact, we have

$$|c_{i}(t)e^{-t} - \tilde{h}^{-}(t)| \leq \int_{-\frac{1}{i}}^{0} |\tilde{h}(t+y) - h^{-}(t)|g_{i}(y)dy + \int_{0}^{\frac{1}{i}} \tilde{h}(t+y)g_{i}(y)dy.$$
(9.5)

For any $\epsilon > 0$, there exists $\delta > 0$ such that $|h(t - \delta) - h^-(t)| < \epsilon$. Then $\exists N > 0$, such that for any n > N, $t \ge t + y > t - \delta$ for all $y \in [-\frac{1}{i}, 0)$ and $\frac{1}{i} < \epsilon$. It follows from (9.5) that

$$|c_i(t)e^{-t} - \tilde{h}^-(t)| \le \epsilon + \epsilon \tilde{h}(t),$$

hence $\lim_{i \to +\infty} c_i(t)e^{-t} = \tilde{h}^-(t)$ for any $t \in \mathbb{R}$. Especially, we have $\lim_{i \to +\infty} c_i(T)e^{-T} = \tilde{h}^-(T) = c(T)e^{-T}$.

Assume that for each i, we have a holomorphic (n,0) form \tilde{F}_i on M such that

$$\int_{M} |\tilde{F}_{i} - (1 - b_{t_{0},B}(\Psi))fF^{1+\delta}|^{2}e^{-\varphi + v_{t_{0},B}(\Psi) - \Psi}c_{i}(-v_{t_{0},B}(\Psi))$$

$$\leq \left(\frac{1}{\delta}c_{i}(T)e^{-T} + \int_{T}^{t_{0}+B}c_{i}(s)e^{-s}ds\right)\int_{M}\frac{1}{B}\mathbb{I}_{\{-t_{0}-B<\Psi<-t_{0}\}}|f|^{2}e^{-\varphi_{\alpha}-\Psi}.$$
(9.6)

By construction of $c_i(t)$, we have

$$\int_{T}^{t_{0}+B} c_{i}(t_{1})e^{-t_{1}}dt_{1}$$

$$= \int_{T}^{t_{0}+B} \int_{\mathbb{R}} \tilde{h}(t_{1}+y)g_{i}(y)dydt_{1}$$

$$= \int_{\mathbb{R}} g_{i}(y) \left(\int_{T}^{t_{0}+B} \tilde{h}(t_{1}+y)dt_{1}\right)dy$$

$$= \int_{\mathbb{R}} g_{i}(y) \left(\int_{T+y}^{t_{0}+B+y} \tilde{h}(s)ds\right)dy$$

$$= \int_{\mathbb{R}} g_{i}(y) \left(\int_{T}^{t_{0}+B} \tilde{h}(s)ds + \int_{t_{0}+B}^{t_{0}+B+y} \tilde{h}(s)ds - \int_{T}^{T+y} \tilde{h}(s)ds\right)dy,$$
(9.7)

then it follows from the construction of $g_i(t)$, $\tilde{h}(t)$ is decreasing with respect to t, inequality (9.7) and $\tilde{h}(t) = c(t)e^{-t}$ on $[T, +\infty)$ that we have

$$\lim_{i \to +\infty} \int_{T}^{t_0 + B} c_i(t_1) e^{-t_1} dt_1 = \int_{T}^{t_0 + B} c(t_1) e^{-t_1} dt_1.$$
(9.8)

For any compact subset K of M, we have $\inf_{i} \inf_{K} e^{v_{t_0,B}(\Psi) - \varphi - \Psi} c_i(-v_{t_0,B}(\Psi)) \geq \inf_{K} e^{v_{t_0,B}(\Psi) - \varphi - \Psi} c(-v_{t_0,B}(\Psi)), \text{ then }$

$$\sup_{i} \int_{K} |\tilde{F}_{i} - (1 - b_{t_{0},B}(\Psi))fF^{1+\delta}|^{2} < +\infty.$$

Note that

$$\int_{K} |(1 - b_{t_0,B}(\Psi))fF^{1+\delta}|^2 \le (\sup_{K} |F^{1+\delta}|^2) \int_{K \cap \{\psi < -t_0\}} |f|^2 < +\infty,$$

then $\sup_{i} \int_{K} |\tilde{F}_{i}|^{2} < +\infty$, which implies that there exists a subsequence of $\{\tilde{F}_{i}\}$ (also denoted by $\{\tilde{F}_{i}\}$), which is compactly convergent to a holomorphic (n, 0) form \tilde{F} on M. Then it follows from inequality (9.6) and Fatou's Lemma that

$$\begin{split} &\int_{M} |\tilde{F} - (1 - b_{t_{0},B}(\Psi))fF^{1+\delta}|^{2}e^{-\varphi + v_{t_{0},B}(\Psi) - \Psi}c(-v_{t_{0},B}(\Psi)) \\ &\leq \int_{M} |\tilde{F} - (1 - b_{t_{0},B}(\Psi))fF^{1+\delta}|^{2}e^{-\varphi + v_{t_{0},B}(\Psi) - \Psi}c^{-}(-v_{t_{0},B}(\Psi)) \\ &\leq \liminf_{i \to +\infty} \int_{M} |\tilde{F}_{i} - (1 - b_{t_{0},B}(\Psi))fF^{1+\delta}|^{2}e^{-\varphi + v_{t_{0},B}(\Psi) - \Psi}c_{i}(-v_{t_{0},B}(\Psi)) \\ &\leq \liminf_{i \to +\infty} \left(\frac{1}{\delta}c_{i}(T)e^{-T} + \int_{T}^{t_{0}+B}c_{i}(s)e^{-s}ds\right) \int_{M} \frac{1}{B}\mathbb{I}_{\{-t_{0}-B < \Psi < -t_{0}\}}|f|^{2}e^{-\varphi_{\alpha}-\Psi} \\ &= \left(\frac{1}{\delta}c(T)e^{-T} + \int_{T}^{t_{0}+B}c(s)e^{-s}ds\right) \int_{M} \frac{1}{B}\mathbb{I}_{\{-t_{0}-B < \Psi < -t_{0}\}}|f|^{2}e^{-\varphi_{\alpha}-\Psi}. \end{split}$$

In the following discussion, we assume that F has no zero points on M and c(t) is smooth.

As M is weakly pseudoconvex, there exists a smooth plurisubharmonic exhaustion function P on M. Let $M_j := \{P < j\}$ (k = 1, 2, ...,). We choose P such that $M_1 \neq \emptyset$.

Then M_j satisfies $M_1 \in M_2 \in ... \in M_j \in M_{j+1} \in ...$ and $\bigcup_{j=1}^n M_j = M$. Each M_j is weakly pseudoconvex Kähler manifold with exhaustion plurisubharmonic function $P_j = 1/(j-P)$.

We will fix j during our discussion until step 8.

Step 1: Regularization of Ψ and $\varphi_{\alpha} + \psi$.

We note that there must exists a continuous nonnegative (1, 1)-form ϖ on M_{j+1} satisfying

$$(\sqrt{-1}\Theta_{TM} + \varpi \otimes Id_{TM})(\kappa_1 \otimes \kappa_2, \kappa_1 \otimes \kappa_2) \ge 0,$$

for $\forall \kappa_1, \kappa_2 \in TM$ on M_{j+1} .

Let $M = M_{j+1}$, $\Omega = M_j$, $T = \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \psi$, $\gamma = 0$ in Lemma 9.4, then there exists a family of functions $\psi_{\zeta,\rho}$ ($\zeta \in (0, +\infty)$) and $\rho \in (0, \rho_1)$ for some positive ρ_1) on M_{j+1} such that

(1) $\psi_{\zeta,\rho}$ is a quasi-plurisubharmonic function on a neighborhood of $\overline{M_j}$, smooth on $M_{j+1} \setminus E_{\zeta}(\psi)$, increasing with respect to ζ and ρ on M_j and converges to ψ on M_j as $\rho \to 0$,

(2) $\frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \psi_{\zeta,\rho} \geq -\zeta \varpi - \delta \omega$ on M_j ,

where $E_{\zeta}(\psi) := \{x \in M : v(\psi, x) \ge \zeta\}$ is the upper-level set of Lelong number and $\{\delta_{\rho}\}$ is an increasing family of positive numbers such that $\lim_{\rho \to 0} \delta_{\rho} = 0$.

Let $\rho = \frac{1}{m}$. Let $\tilde{\delta}_m := \delta_{\frac{1}{m}}$ and $\zeta = \tilde{\delta}_m$. Denote $\psi_m := \psi_{\tilde{\delta}_m, \frac{1}{m}}$. Then we have a sequence of functions $\{\psi_m\}$ satisfying

(1') ψ_m is quasi-plurisubharmonic function on $\overline{M_j}$, smooth on $M_{j+1} \setminus E_m(\psi)$, decreasing with respect to m and converges to ψ on M_j as $m \to +\infty$,

(2') $\frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \psi_m \ge -\tilde{\delta}_m \varpi - \tilde{\delta}_m \omega$ on M_j ,

where $E_m(\psi) = \{x \in X : v(\psi, x) \ge \frac{1}{m}\}$ is the upper level set of Lelong number and $\{\tilde{\delta}_m\}$ is an decreasing family of positive numbers such that $\lim_{m \to +\infty} \tilde{\delta}_m = 0$.

As M_j is relatively compact in M, there exists a positive number $b \ge 1$ such that $b\omega \ge \varpi$ on M_j . Then condition (2') becomes

 $(2") \frac{\sqrt{-1}}{\pi} \partial \bar{\partial} \psi_m \ge -\tilde{\delta}_m \varpi - \tilde{\delta}_m \omega \ge -2b \tilde{\delta}_m \omega \text{ on } M_j.$

Denote $h := \varphi_{\alpha} + \psi$. Note that h is a plurisubharmonic function on M. Denote $h_l := \max\{h, -l\}$, where $l \in \mathbb{Z}^+$. Note that h_l is a plurisubharmonic function. As $h_l \geq -l$, we know $v(h_l, z) = 0$ for all $z \in M$. By using the similarly discussion as above, we have a sequence of functions $\{h_{m',l}\}$ on M_{j+1} such that

(i) $h_{m',l}$ is quasi-plurisubharmonic function on $\overline{M_j}$, smooth on M_{j+1} , decreasing with respect to m' and converges to h_l on M_j as $m' \to +\infty$,

(ii)
$$\frac{\sqrt{-1}}{\pi} \partial \bar{\partial} h_{m',l} \geq -2b\delta_{m',l}\omega$$
 on M_i ,

where $\{\delta_{m',l}\}$ is an decreasing family of positive numbers such that $\lim_{m'\to+\infty} \delta_{m',l} = 0$.

From now on, we will fix the positive integer l during our discussion until step 7.

For fixed l, we can assume that δ_n and $\delta_{n,l}$ are the same sequence of variable $n \in \mathbb{Z}^+$, since we can replace them by the term $\max\{\tilde{\delta}_n, \delta_{n,l}\}$. We denote both $\tilde{\delta}_n$ and $\delta_{n,l}$ by $\tilde{\delta}_n$ for simplicity.

Let $\eta_m = \{\frac{t_0 - T}{3m}, \frac{t_0 - T}{3m}\}$ and we have the function $M_{\eta_m}(\psi_m + T, 2\log|F|)$. Denote $M_{\eta_m} := M_{\eta_m}(\psi_m + T, 2\log|F|)$ for simplicity. Note that $\psi_m + T$ is a $2b\tilde{\delta}_m\omega$ plurisubharmonic function. As F is a holomorphic function, ω is a Kähler form and $b\delta_m > 0$, we know that $2\log|F|$ is a $2b\delta_m\omega$ -plurisubharmonic function. It follows from Lemma 9.9 that M_{η_m} is a $2b\delta_m\omega$ -plurisubharmonic function, i.e.,

$$\frac{\sqrt{-1}}{\pi}\partial\bar{\partial}M_{\eta_m} \ge -2\pi b\tilde{\delta}_m\omega.$$

Denote $\Psi_m := \psi_m - M_{\eta_m}(\psi_m + T, 2\log|F|)$. Then Ψ_m is smooth on $M_j \setminus E_m$. It is easy to verify that when $m \to +\infty$, $\Psi_m \to \Psi$. It follows from Lemma 9.9 that we know

we know (1) if $\psi_m + T \leq 2 \log |F| - \frac{2(t_0 - T)}{3m}$ holds, we have $\Psi_m = \psi_m - 2 \log |F|$; (2) if $\psi_m + T \geq 2 \log |F| + \frac{2(t_0 - T)}{3m}$ holds, we have $\Psi_m = -T$; (3) if $2 \log |F| - \frac{2(t_0 - T)}{3m} < \psi_m + T < 2 \log |F| + \frac{2(t_0 - T)}{3m}$ holds, we have $\max\{\psi_m + T, 2 \log |F|\} \leq M_{\eta_m} \leq (\psi_m + T + \frac{t_0 - T}{m})$ and hence $-T - \frac{t_0 - T}{m} \leq \Psi_m \leq -T$. Thus we have $\{\Psi_m < -t_0\} = \{\psi_m - 2 \log |F| < -t_0\} \subset \{\psi - 2 \log |F| < -t_0\} = \{\Psi < -t_0\}$. We also note that $\Psi_m \leq -T$ on M_{j+1} .

Step 2: Recall some constructions.

To simplify our notations, we denote $b_{t_0,B}(t)$ by b(t) and $v_{t_0,B}(t)$ by v(t).

Let $\epsilon \in (0, \frac{1}{8}B)$. Let $\{v_{\epsilon}\}_{\epsilon \in (0, \frac{1}{8}B)}$ be a family of smooth increasing convex functions on \mathbb{R} , such that:

(1) $v_{\epsilon}(t) = t$ for $t \ge -t_0 - \epsilon$, $v_{\epsilon}(t) = constant$ for $t < -t_0 - B + \epsilon$;

(2) $v_{\epsilon}''(t)$ are convergence pointwise to $\frac{1}{B}\mathbb{I}_{(-t_0-B,-t_0)}$, when $\epsilon \to 0$, and $0 \leq 1$ $v_{\epsilon}''(t) \leq \frac{2}{B} \mathbb{I}_{(-t_0 - B + \epsilon, -t_0 - \epsilon)}$ for ant $t \in \mathbb{R}$; (3) $v_{\epsilon}'(t)$ are convergence pointwise to b(t) which is a continuous function on R

when $\epsilon \to 0$ and $0 \le v_{\epsilon}'(t) \le 1$ for any $t \in \mathbb{R}$.

One can construct the family $\{v_{\epsilon}\}_{\epsilon \in (0,\frac{1}{8}B)}$ by setting

$$v_{\epsilon}(t) := \int_{-\infty}^{t} (\int_{-\infty}^{t_{1}} (\frac{1}{B - 4\epsilon} \mathbb{I}_{(-t_{0} - B + 2\epsilon, -t_{0} - 2\epsilon)} * \rho_{\frac{1}{4}\epsilon})(s)ds)dt_{1} \\ - \int_{-\infty}^{-t_{0}} (\int_{-\infty}^{t_{1}} (\frac{1}{B - 4\epsilon} \mathbb{I}_{(-t_{0} - B + 2\epsilon, -t_{0} - 2\epsilon)} * \rho_{\frac{1}{4}\epsilon})(s)ds)dt_{1} - t_{0}ds$$

where $\rho_{\frac{1}{4}\epsilon}$ is the kernel of convolution satisfying $\operatorname{supp}(\rho_{\frac{1}{4}\epsilon}) \subset (-\frac{1}{4}\epsilon, \frac{1}{4}\epsilon)$. Then it follows that

$$v_{\epsilon}''(t) = \frac{1}{B - 4\epsilon} \mathbb{I}_{(-t_0 - B + 2\epsilon, -t_0 - 2\epsilon)} * \rho_{\frac{1}{4}\epsilon}(t),$$

and

$$v_{\epsilon}'(t) = \int_{-\infty}^{t} \left(\frac{1}{B - 4\epsilon} \mathbb{I}_{(-t_0 - B + 2\epsilon, -t_0 - 2\epsilon)} * \rho_{\frac{1}{4}\epsilon}\right)(s) ds.$$

Let $\eta = s(-v_{\epsilon}(\Psi_m))$ and $\phi = u(-v_{\epsilon}(\Psi_m))$, where $s \in C^{\infty}([T, +\infty))$ satisfies $s \geq \frac{1}{\delta}$ and $u \in C^{\infty}([T, +\infty))$, such that $s'(t) \neq 0$ for any t, u''s - s'' > 0 and s' - u's = 1. Let $\Phi = \phi + h_{m',l} + \delta M_{\eta_m}$. Denote $\tilde{h} := e^{-\Phi}$.

Step 3: Solving $\bar{\partial}$ -equation with error term.

Set $B = [\eta \sqrt{-1}\Theta_{\tilde{h}} - \sqrt{-1}\partial \bar{\partial}\eta - \sqrt{-1}g\partial \eta \wedge \bar{\partial}\eta$, $\Lambda_{\omega}]$, where g is a positive function. We will determine g by calculations. On $M_j \setminus E_m$, direct calculation shows that

$$\begin{aligned} \partial \partial \eta &= -s'(-v_{\epsilon}(\Psi_{m}))\partial \bar{\partial}(v_{\epsilon}(\Psi_{m})) + s''(-v_{\epsilon}(\Psi_{m}))\partial(v_{\epsilon}(\Psi_{m})) \wedge \bar{\partial}(v_{\epsilon}(\Psi_{m})), \\ \eta \Theta_{\tilde{h}} &= \eta \partial \bar{\partial} \phi + \eta \partial \bar{\partial} h_{m',l} + \eta \partial \bar{\partial}(\delta M_{\eta_{m}}) \\ &= su''(-v_{\epsilon}(\Psi_{m}))\partial(v_{\epsilon}(\Psi_{m})) \wedge \bar{\partial}(v_{\epsilon}(\Psi_{m})) - su'(-v_{\epsilon}(\Psi_{m}))\partial \bar{\partial}(v_{\epsilon}(\Psi_{m})) \\ &+ s \partial \bar{\partial} h_{m',l} + s \partial \bar{\partial}(\delta M_{\eta_{m}}). \end{aligned}$$

Hence

$$\begin{split} \eta \sqrt{-1} \Theta_{\tilde{h}} &- \sqrt{-1} \partial \bar{\partial} \eta - \sqrt{-1} g \partial \eta \wedge \bar{\partial} \eta \\ = s \sqrt{-1} \partial \bar{\partial} h_{m',l} + s \sqrt{-1} \partial \bar{\partial} (\delta M_{\eta_m}) \\ &+ (s' - su') (v'_{\epsilon}(\Psi_m) \sqrt{-1} \partial \bar{\partial} (\Psi_m) + v''_{\epsilon}(\psi_m) \sqrt{-1} \partial (\Psi_m) \wedge \bar{\partial} (\Psi_m)) \\ &+ [(u''s - s'') - gs'^2] \sqrt{-1} \partial (v_{\epsilon}(\Psi_m)) \wedge \bar{\partial} (v_{\epsilon}(\Psi_m)), \end{split}$$

where we omit the term $-v_{\epsilon}(\Psi_m)$ in $(s'-su')(-v_{\epsilon}(\Psi_m))$ and $[(u''s-s'')-gs'^2](-v_{\epsilon}(\Psi_m))$ for simplicity.

Let
$$g = \frac{u''s-s''}{s'^2}(-v_{\epsilon}(\Psi_m))$$
 and note that $s' - su' = 1, 0 \le v'_{\epsilon}(\Psi_m) \le 0$. Then
 $\eta\sqrt{-1}\Theta_{\tilde{h}} - \sqrt{-1}\partial\bar{\partial}\eta - \sqrt{-1}g\partial\eta \wedge \bar{\partial}\eta$
 $=s\sqrt{-1}\partial\bar{\partial}h_{m',l} + s\sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_m}) + v'_{\epsilon}(\Psi_m)\sqrt{-1}\partial\bar{\partial}(\Psi_m) + v''_{\epsilon}(\psi_m)\sqrt{-1}\partial(\Psi_m) \wedge \bar{\partial}(\Psi_m)$
 $=v''_{\epsilon}(\psi_m)\sqrt{-1}\partial(\Psi_m) \wedge \bar{\partial}(\Psi_m) + v'_{\epsilon}(\Psi_m)\sqrt{-1}\partial\bar{\partial}(\Psi_m)$
 $+ s(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega) - 2\pi bs\tilde{\delta}_{m'}\omega + s(\sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_m}) + 2\pi b\delta\tilde{\delta}_m\omega) - 2\pi bs\delta\tilde{\delta}_m\omega$
 $\ge v''_{\epsilon}(\psi_m)\sqrt{-1}\partial(\Psi_m) \wedge \bar{\partial}(\Psi_m) + v'_{\epsilon}(\Psi_m)\sqrt{-1}\partial\bar{\partial}(\Psi_m)$
 $+ \frac{1}{\delta}(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega) + \frac{1}{\delta}(\sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_m}) + 2\pi b\delta\tilde{\delta}_m\omega) - 2\pi bs(\tilde{\delta}_{m'} + \delta\tilde{\delta}_m)\omega$
(9.9)

Note that

$$\delta v'_{\epsilon}(\Psi_{m})\sqrt{-1}\partial\bar{\partial}(\Psi_{m}) + (\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega) + (\sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_{m}}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$= (1 - v'_{\epsilon}(\Psi_{m}))(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega + \sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_{m}}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$+ v'_{\epsilon}(\Psi_{m})(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega + \sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_{m}}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$+ v'_{\epsilon}(\Psi_{m})(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega + \sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_{m}}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$+ v'_{\epsilon}(\Psi_{m})(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega + \sqrt{-1}\partial\bar{\partial}(\delta M_{\eta_{m}}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$+ v'_{\epsilon}(\Psi_{m})(\sqrt{-1}\partial\bar{\partial}h_{m',l} + 2\pi b\tilde{\delta}_{m'}\omega + \sqrt{-1}\partial\bar{\partial}(\delta\psi_{m}) + 2\pi b\delta\tilde{\delta}_{m}\omega)$$

$$\geq 0.$$

$$(0.10)$$

(9.10)

It follows from inequality (9.9) and inequality (9.10) that

$$\eta \sqrt{-1} \Theta_{\tilde{h}} - \sqrt{-1} \partial \bar{\partial} \eta - \sqrt{-1} g \partial \eta \wedge \bar{\partial} \eta$$

$$\geq v_{\epsilon}''(\Psi_m) \sqrt{-1} \partial (\Psi_m) \wedge \bar{\partial} (\Psi_m) - 2\pi b s(\tilde{\delta}_{m'} + \delta \tilde{\delta}_m) \omega.$$

By the constructions of s(t), $v_{\epsilon}(t)$ and $\sup_{m} \sup_{M_{j}} \Psi_{m} \leq -T$, we have $s(-v_{\epsilon}(\Psi_{m}))$ is uniformly bounded on M_{j} with respect to ϵ and m. Let \tilde{M} be the uniformly upper bound of $s(-v_{\epsilon}(\Psi_{m}))$ on M_{j} . Then on $M_{j} \setminus E_{m}$, we have

$$\eta \sqrt{-1} \Theta_{\tilde{h}} - \sqrt{-1} \partial \bar{\partial} \eta - \sqrt{-1} g \partial \eta \wedge \bar{\partial} \eta$$

$$\geq v_{\epsilon}''(\Psi_m) \sqrt{-1} \partial (\Psi_m) \wedge \bar{\partial} (\Psi_m) - 2\pi b \tilde{M} (\tilde{\delta}_{m'} + \delta \tilde{\delta}_m) \omega.$$

Hence, for any (n, 1) form α , we have

$$\langle (B + 2\pi b \tilde{M}(\tilde{\delta}_{m'} + \delta \tilde{\delta}_m)I)\alpha, \alpha \rangle_{\tilde{h}}$$

$$\geq \langle [v_{\epsilon}''(\Psi_m)\partial(\Psi_m) \wedge \bar{\partial}(\Psi_m), \Lambda_{\omega}]\alpha, \alpha \rangle_{\tilde{h}}$$

$$= \langle (v_{\epsilon}''(\Psi_m)\bar{\partial}(\Psi_m) \wedge (\alpha \llcorner (\bar{\partial}\Psi_m)^{\sharp})), \alpha \rangle_{\tilde{h}}.$$

$$(9.11)$$

It follows from Lemma 9.2 that $B + 2\pi b \tilde{M} (\tilde{\delta}_{m'} + \delta \tilde{\delta}_m) I$ is semi-positive. Using the definition of contraction, Cauchy-Schwarz inequality and inequality (9.11), we have

$$\begin{aligned} |\langle v_{\epsilon}''(\Psi_{m})\bar{\partial}\Psi_{m}\wedge\gamma,\widetilde{\alpha}\rangle_{\tilde{h}}|^{2} &= |\langle v_{\epsilon}''(\Psi_{m})\gamma,\widetilde{\alpha}\llcorner(\bar{\partial}\Psi_{m})^{\sharp}\rangle_{\tilde{h}}|^{2} \\ &\leq \langle (v_{\epsilon}''(\Psi_{m})\gamma,\gamma)\rangle_{\tilde{h}}(v_{\epsilon}''(\Psi_{m}))|\widetilde{\alpha}\llcorner(\bar{\partial}\Psi_{m})^{\sharp}|_{\tilde{h}}^{2} \\ &= \langle (v_{\epsilon}''(\Psi_{m})\gamma,\gamma)\rangle_{\tilde{h}}\langle (v_{\epsilon}''(\Psi_{m}))\bar{\partial}\Psi_{m}\wedge(\widetilde{\alpha}\llcorner(\bar{\partial}\Psi_{m})^{\sharp}),\widetilde{\alpha}\rangle_{\tilde{h}} \\ &\leq \langle (v_{\epsilon}''(\Psi_{m})\gamma,\gamma)\rangle_{\tilde{h}}\langle (B+2\pi b\tilde{M}(\tilde{\delta}_{m'}+\delta\tilde{\delta}_{m})I)\widetilde{\alpha},\widetilde{\alpha})\rangle_{\tilde{h}} \end{aligned}$$
(9.12)

for any (n,0) form γ and (n,1) form $\tilde{\alpha}$.

As $fF^{1+\delta}$ is holomorphic on $\{\Psi < -t_0\}$ and $\{\Psi_m < -t_0 - \epsilon\} \subset \{\Psi_m < -t_0\} \subset \{\Psi_m < -t_0\} \subset \{\Psi < -t_0\}$, then $\lambda := \bar{\partial}((1-v'_{\epsilon}(\Psi_m))fF^{1+\delta})$ is well defined and smooth on $M_j \setminus E_m$. Taking $\gamma = fF^{1+\delta}$, $\tilde{\alpha} = (B + 2\pi b\tilde{M}(\tilde{\delta}_{m'} + \delta\tilde{\delta}_m)I)^{-1}(\bar{\partial}v'_{\epsilon}(\Psi_m)) \wedge fF^{1+\delta}$. Then it follows from inequality (9.12) that

$$\langle (B+2\pi b\tilde{M}(\tilde{\delta}_{m'}+\delta\tilde{\delta}_m)I)^{-1}\lambda,\lambda\rangle_{\widetilde{h}}\leq v_\epsilon''(\Psi_m)|fF^{1+\delta}|^2e^{-\Phi}$$

Thus we have

$$\int_{M_j \setminus E_m} \langle (B + 2\pi b \tilde{M}(\tilde{\delta}_{m'} + \delta \tilde{\delta}_m) I)^{-1} \lambda, \lambda \rangle_{\tilde{h}} \leq \int_{M_j \setminus E_m} v_{\epsilon}''(\Psi_m) |fF^{1+\delta}|^2 e^{-\Phi}.$$

Recall that $e^{-\Phi} = e^{-\phi - h_{m',l} - \delta M_{\eta_m}}$. Note that $h_{m',l} \ge -l$ and $\delta M_{\eta_m} \ge \delta 2 \log |F|$. By the construction of ϕ , we know that $\sup_{\overline{M_j}} e^{-\phi} < +\infty$. Then

$$\int_{M_j \setminus E_m} v_{\epsilon}''(\Psi_m) |fF^{1+\delta}|^2 e^{-\Phi} \le \sup_{\overline{M_j}} \left(e^{-\phi+l} |F|^2 \right) \int_{\overline{M_j}} \frac{2}{B} \mathbb{I}_{\{\Psi \le -t_0\}} |f|^2 < +\infty.$$

By Lemma 9.6, $M_j \setminus E_m$ carries a complete Kähler metric. Then it follows from Lemma 9.3 that there exists

$$u_{m,m',l,\epsilon,j} \in L^2(M_j \backslash E_m, K_M),$$

$$h_{m,m',l,\epsilon,j} \in L^2(M_j \setminus E_m, \wedge^{n,1}T^*M)$$

such that $\bar{\partial} u_{m,m',l,\epsilon,j} + \sqrt{2\pi b \tilde{M}(\tilde{\delta}_{m'} + \delta \tilde{\delta}_m)} h_{m,m',l,\epsilon,j} = \lambda$ holds on $M_j \setminus E_m$, and

$$\begin{split} &\int_{M_j \setminus E_m} \frac{1}{\eta + g^{-1}} |u_{m,m',l,\epsilon,j}|^2 e^{-\Phi} + \int_{M_j \setminus E_m} |h_{m,m',l,\epsilon,j}|^2 e^{-\Phi} \\ &\leq \int_{M_j \setminus E_m} \langle (B + 2\pi b \tilde{M}(\tilde{\delta}_{m'} + \delta \tilde{\delta}_m) I)^{-1} \lambda, \lambda \rangle_{\tilde{h}} \\ &\leq \int_{M_j \setminus E_m} v_{\epsilon}''(\Psi_m) |fF^{1+\delta}|^2 e^{-\Phi} < +\infty. \end{split}$$

Assume that we can choose η and ϕ such that $(\eta + g^{-1})^{-1} = e^{v_{\epsilon}(\Psi_m)} e^{\phi} c(-v_{\epsilon}(\Psi_m)).$ Then we have

$$\int_{M_{j}\setminus E_{m}} |u_{m,m',l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi_{m}) - h_{m',l} - \delta M_{\eta_{m}}} c(-v_{\epsilon}(\Psi_{m})) + \int_{M_{j}\setminus E_{m}} |h_{m,m',l,\epsilon,j}|^{2} e^{-\phi - h_{m',l} - \delta M_{\eta_{m}}} \\
\leq \int_{M_{j}\setminus E_{m}} v_{\epsilon}''(\Psi_{m}) |fF^{1+\delta}|^{2} e^{-\phi - h_{m',l} - \delta M_{\eta_{m}}} < +\infty.$$
(9.13)

By the construction of $v_{\epsilon}(t)$ and $c(t)e^{-t}$ is decreasing with respect to t, we know $c(-v_{\epsilon}(\Psi_m))e^{v_{\epsilon}(\Psi_m)}$ has a positive lower bound on $M_j \subseteq M$. By the constructions of $v_{\epsilon}(t)$ and u, we know $e^{-\phi} = e^{-u(-v_{\epsilon}(\Psi_m))}$ has a positive lower bound on $M_j \subseteq M$. By the upper semi-continuity of M_{η_m} , we know $e^{-\delta M_{\eta_m}}$ has a positive lower bound on $M_j \in M$. Note that $h_{m',l}$ is smooth on $M_j \in M$. Hence it follows from inequality (9.13) that

$$u_{m,m',l,\epsilon,j} \in L^2(M_j, K_M),$$
$$h_{m,m',l,\epsilon,j} \in L^2(M_j, \wedge^{n,1}T^*M).$$

It follows from Lemma 9.7 that we know

$$\bar{\partial}u_{m,m',l,\epsilon,j} + \sqrt{2\pi b\tilde{M}(\tilde{\delta}_{m'} + \delta\tilde{\delta}_m)}h_{m,m',l,\epsilon,j} = \lambda$$
(9.14)

holds on M_i . And we have

$$\int_{M_{j}} |u_{m,m',l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi_{m}) - h_{m',l} - \delta M_{\eta_{m}}} c(-v_{\epsilon}(\Psi_{m})) + \int_{M_{j}} |h_{m,m',l,\epsilon,j}|^{2} e^{-\phi - h_{m',l} - \delta M_{\eta_{m}}} \\
\leq \int_{M_{j}} v_{\epsilon}''(\Psi_{m}) |fF^{1+\delta}|^{2} e^{-\phi - h_{m',l} - \delta M_{\eta_{m}}} < +\infty.$$
(9.15)

Step 4: Letting $m \to +\infty$. Note that $\sup_m \sup_{M_j} e^{-\phi} = \sup_m \sup_{M_j} e^{-u(-v_{\epsilon}(\Psi_m))} < +\infty, e^{-h_{m',l}} \le e^l$ and $e^{-\delta M_{\eta_m}} \le e^{-\delta 2\log|F|}$. As $\{\Psi_m < -t_0 - \epsilon\} \subset \{\Psi_m < -t_0\} \subset \{\Psi < -t_0\}$, we have

$$v_{\epsilon}''(\Psi_m)|fF^{1+\delta}|^2 e^{-\phi - h_{m',l} - \delta M_{\eta_m}} \le \frac{2}{B} \left(\sup_{m \ M_j} \sup_{M_j} e^{-\phi} \right) e^l \left(\sup_{M_j} |F|^2 \right) \mathbb{I}_{\{\Psi < -t_0\}} |f|^2$$

holds on M_j . It follows from $\int_{\{\Psi < -t_0\} \cap \overline{M_j}} |f|^2 < +\infty$ and dominated convergence theorem that

$$\begin{split} &\lim_{m \to +\infty} \int_{M_j} v_{\epsilon}''(\Psi_m) |fF^{1+\delta}|^2 e^{-\phi - h_{m',l} - \delta M_{\eta_m}} \\ &= \int_{M_j} v_{\epsilon}''(\Psi) |fF^{1+\delta}|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} \end{split}$$

Note that

$$v_{\epsilon}''(\Psi)|fF^{1+\delta}|^{2}e^{-u(-v_{\epsilon}(\Psi))-h_{m',l}-\delta\max\{\psi+T,2\log|F|\}} \leq \frac{2}{B}\left(\sup_{M_{j}}e^{-u(-v_{\epsilon}(\Psi))}|F|^{2}\right)e^{l}\mathbb{I}_{\{\Psi<-t_{0}\}}|f|^{2}$$

holds on M_j . We have

$$\int_{M_j} v_{\epsilon}''(\Psi) |fF^{1+\delta}|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max{\{\psi + T, 2\log{|F|}\}}} < +\infty.$$

Note that $\inf_{m} \inf_{M_{j}} c(-v_{\epsilon}(\Psi_{m}))e^{-v_{\epsilon}(\Psi_{m})-h_{m',l}} > 0$. It follows from Lemma 9.9 that $M_{\eta_{m}} \leq \max \{\psi_{m} + T, 2\log|F|\} + \frac{t_{0}-T}{3m} \leq \max \{\psi_{m} + T, 2\log|F|\} + t_{0} - T \leq \max \{\psi_{1} + T, 2\log|F|\} + t_{0} - T$. As ψ_{1} is a quasi-plurisubharmonic function on $\overline{M_{j}}$, we know $\max \{\psi_{1} + T, 2\log|F|\}$ is upper semi-continuous function on M_{j} . Hence

$$\inf_{m} \inf_{M_{j}} e^{-M_{\eta_{m}}} \ge \inf_{M_{j}} e^{-\max\left\{\psi_{1}+T, 2\log|F|\right\}-t_{0}} > 0.$$
(9.16)

Then it follows from inequality (9.15) that

$$\sup_m \int_{M_j} |u_{m,m',l,\epsilon,j}|^2 < +\infty.$$

Therefore the solutions $u_{m,m',l,\epsilon,j}$ are uniformly bounded in L^2 norm with respect to m on M_j . Since the closed unit ball of the Hilbert space is weakly compact, we can extract a subsequence $u_{m_1,m',l,\epsilon,j}$ weakly convergent to $u_{m',l,\epsilon,j}$ in $L^2(M_j, K_M)$ as $m_1 \to +\infty$.

Note that $\sup_m \sup_{M_j} e^{v_{\epsilon}(\Psi_m)} c(-v_{\epsilon}(\Psi_m)) e^{-h_{m',l}} < +\infty$. As $M_{\eta_m} \ge \max\{\psi_m + T, 2\log|F|\} \ge 2\log|F|$ and F has no zero points on M, we have $\sup_m \sup_{M_j} e^{-M_{\eta_m}} \le \sup_{M_j} \frac{1}{|F|^2} < +\infty$. Hence we know

$$\sup_{m} \sup_{M_{j}} e^{v_{\epsilon}(\Psi_{m})} c(-v_{\epsilon}(\Psi_{m})) e^{-h_{m',l} - \delta M_{\eta_{m}}} < +\infty.$$

It follows from Lemma 9.8 that we know $u_{m_1,m',l,\epsilon,j}\sqrt{e^{v_{\epsilon}(\Psi_{m_1})}c(-v_{\epsilon}(\Psi_{m_1}))e^{-h_{m',l}-\delta M_{\eta_{m_1}}}}$ weakly convergent to $u_{m',l,\epsilon,j}\sqrt{e^{v_{\epsilon}(\Psi)}c(-v_{\epsilon}(\Psi))e^{-h_{m',l}-\delta\max\{\psi+T,2\log|F|\}}}$. Hence we have

$$\int_{M_{j}} |u_{m',l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi) - h_{m',l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi))$$

$$\leq \liminf_{m_{1} \to +\infty} \int_{M_{j}} |u_{m_{1},m',l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi_{m_{1}}) - h_{m',l} - \delta M_{\eta_{m_{1}}}} c(-v_{\epsilon}(\Psi_{m_{1}})))$$

$$\leq \liminf_{m_{1} \to +\infty} \int_{M_{j}} v_{\epsilon}''(\Psi_{m_{1}}) |fF^{1+\delta}|^{2} e^{-u(-v_{\epsilon}(\Psi_{m_{1}})) - h_{m',l} - \delta M_{\eta_{m_{1}}}}$$

$$\leq \int_{M_{j}} v_{\epsilon}''(\Psi) |fF^{1+\delta}|^{2} e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\{\psi + T, 2\log|F|\}} < +\infty.$$
(9.17)

Note that $\inf_{M_1} \inf_{M_j} e^{-u(-v_{\epsilon}(\Psi_{m_1})) - h_{m',l}} > 0$. Combining inequality (9.15) and inequality (9.16) we know

$$\sup_{m_1} \int_{M_j} |h_{m_1,m',l,\epsilon,j}|^2 < +\infty.$$

Since the closed unit ball of the Hilbert space is weakly compact, we can extract a subsequence of $\{h_{m_1,m',l,\epsilon,j}\}$ (also denote by $h_{m_1,m',l,\epsilon,j}$) weakly convergent to $h_{m',l,\epsilon,j}$ in $L^2(M_j, \wedge^{n,1}T^*M)$ as $m_1 \to +\infty$.

 $\begin{array}{l} h_{m',l,\epsilon,j} \text{ in } L^2(M_j,\wedge^{n,1}T^*M) \text{ as } m_1 \to +\infty.\\ \text{ Note that } \sup_{M_j} \sup_{M_j} e^{-u(-v_\epsilon(\Psi_{m_1}))-h_{m',l}} < +\infty \text{ and } \sup_{M_j} \sup_{M_j} e^{-M_{\eta_{m_1}}} \leq \\ \sup_{M_j} \frac{1}{|F|^2} < +\infty. \text{ We know} \end{array}$

$$\sup_{m_1} \sup_{M_j} e^{-u(-v_{\epsilon}(\Psi_{m_1})) - h_{m',l} - \delta M_{\eta_{m_1}}} < +\infty.$$

It follows from Lemma 9.8 that we have $h_{m_1,m',l,\epsilon,j}\sqrt{e^{-u(-v_{\epsilon}(\Psi_{m_1}))-h_{m',l}-\delta M_{\eta_{m_1}}}$ is weakly convergent to $h_{m',l,\epsilon,j}\sqrt{e^{-u(-v_{\epsilon}(\Psi))-h_{m',l}-\delta \max{\{\psi+T,2\log{|F|}\}}}$. Hence we have

$$\int_{M_{j}} |h_{m',l,\epsilon,j}|^{2} e^{-u(-v_{\epsilon}(\Psi))-h_{m',l}-\delta \max\{\psi+T,2\log|F|\}} \\
\leq \liminf_{m_{1}\to+\infty} \int_{M_{j}} |h_{m_{1},m',l,\epsilon,j}|^{2} e^{-u(-v_{\epsilon}(\Psi_{m_{1}}))-h_{m',l}-\delta M_{\eta_{m_{1}}}} \\
\leq \liminf_{m_{1}\to+\infty} \int_{M_{j}} v_{\epsilon}''(\Psi_{m_{1}})|fF^{1+\delta}|^{2} e^{-u(-v_{\epsilon}(\Psi_{m_{1}}))-h_{m',l}-\delta M_{\eta_{m_{1}}}} \\
\leq \int_{M_{j}} v_{\epsilon}''(\Psi)|fF^{1+\delta}|^{2} e^{-u(-v_{\epsilon}(\Psi))-h_{m',l}-\delta \max\{\psi+T,2\log|F|\}} < +\infty.$$
(9.18)

Replace m by m_1 in (9.14) and let $m_1 \to +\infty$, we have

$$\bar{\partial} u_{m',l,\epsilon,j} + \sqrt{2\pi b \tilde{M} \tilde{\delta}_{m'}} h_{m',l,\epsilon,j} = \bar{\partial} \left((1 - v'_{\epsilon}(\Psi)) f F^{1+\delta} \right).$$
(9.19)

Step 5: Letting $m' \to +\infty$.

When $\Psi < -t_0 - \epsilon < -t_0$, we have $\psi - 2\log|F| < -T$ and then max $\{\psi + T, 2\log|F|\} = 2\log|F|$. Hence

$$\int_{M_j} v_{\epsilon}''(\Psi) |fF^{1+\delta}|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_{m',l} - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)} + \delta \max\left\{\psi + T, 2\log|F|\right\}} = \int_{M_j} v_{$$

Note that

$$v_{\epsilon}''(\Psi)|fF|^{2}e^{-u(-v_{\epsilon}(\Psi))-h_{m',l}} \leq \frac{2}{B} \left(\sup_{M_{j}} e^{-u(-v_{\epsilon}(\Psi))+l}|F|^{2} \right) \mathbb{I}_{\{\Psi < -t_{0}\}}|f|^{2}.$$

It follows from $\int_{\{\Psi < -t_0\} \cap \overline{M_i}} |f| < +\infty$ and dominated convergence theorem that

$$\lim_{m'\to+\infty} \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi))-h_{m',j}}$$
$$= \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi))-h_l} < +\infty.$$

Note that $h_{m',l} \leq h_{1,l}$ for any m' and $h_{1,l}$ is quasi-plurisubharmonic function on $\overline{M_i}$. Then

$$\inf_{m'} \inf_{M_j} e^{v_{\epsilon}(\Psi) - h_{m',l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi)) \ge C_j \inf_{m'} \inf_{M_j} e^{-h_{1,l}} > 0,$$

where $C_j := \inf_{M_j} e^{v_{\epsilon}(\Psi) - \delta \max\{\psi + T, 2 \log |F|\}} c(-v_{\epsilon}(\Psi))$ is a positive number.

It follows from inequality (9.17) that

$$\sup_{m'} \int_{M_j} |u_{m',l,\epsilon,j}|^2 < +\infty.$$

Since the closed unit ball of the Hilbert space is weakly compact, we can extract a subsequence $u_{m'',l,\epsilon,j}$ weakly convergent to $u_{l,\epsilon,j}$ in $L^2(M_j, K_M)$ as $m'' \to +\infty$.

Note that

$$\sup_{m''} \sup_{M_j} e^{v_{\epsilon}(\Psi) - h_{m'',l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi)) \le \left(\sup_{M_j} \frac{e^{v_{\epsilon}(\Psi)} c(-v_{\epsilon}(\Psi))}{|F|^{2\delta}} \right) e^l.$$

It follows from Lemma 9.8 that we have $u_{m'',l,\epsilon,j}\sqrt{e^{v_{\epsilon}(\Psi)-h_{m'',l}-\delta\max\{\psi+T,2\log|F|\}}c(-v_{\epsilon}(\Psi))}$ weakly convergent to $u_{l,\epsilon,j}\sqrt{e^{v_{\epsilon}(\Psi)-h_{l}-\delta\max\{\psi+T,2\log|F|\}}c(-v_{\epsilon}(\Psi))}$.

It follows from inequality (9.17) that

$$\int_{M_{j}} |u_{l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi) - h_{l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi))$$

$$\leq \liminf_{m_{1} \to +\infty} \int_{M_{j}} |u_{m'',l,\epsilon,j}|^{2} e^{v_{\epsilon}(\Psi) - h_{m'',l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi))$$

$$\leq \liminf_{m_{1} \to +\infty} \int_{M_{j}} v_{\epsilon}''(\Psi) |fF|^{2} e^{-u(-v_{\epsilon}(\Psi)) - h_{m'',l}}$$

$$= \int_{M_{j}} v_{\epsilon}''(\Psi) |fF|^{2} e^{-u(-v_{\epsilon}(\Psi)) - h_{l}} < +\infty.$$
(9.20)

As $\inf_{m''} \inf_{M_j} e^{-u(-v_{\epsilon}(\Psi))-h_{m'',l}-\delta \max\{\psi+T,2\log|F|\}} \geq \tilde{C}_j \inf_{M_j} e^{-h_{1,l}} > 0$, where $\tilde{C}_j = \inf_{M_j} e^{-u(-v_{\epsilon}(\Psi))-\delta \max\{\psi+T,2\log|F|\}}$ is a positive number. Then it follows from inequality (9.18) that we know

$$\sup_{m^{\prime\prime}} \int_{M_j} |h_{m^{\prime\prime},l,\epsilon,j}|^2 < +\infty.$$

Since the closed unit ball of the Hilbert space is weakly compact, we can extract a subsequence of $\{h_{m'',l,\epsilon,j}\}$ (also denote by $h_{m'',l,\epsilon,j}$) weakly convergent to

 $h_{l,\epsilon,j}$ in $L^2(M_j, \wedge^{n,1}T^*M)$ as $m'' \to +\infty$. Then it follows from Lemma 9.8 and $\lim_{m'' \to +\infty} \delta_{m''} = 0 \text{ that } \sqrt{2\pi b \tilde{M} \tilde{\delta}_{m''} h_{m'',l,\epsilon,j}} \text{ is weakly convergent to } 0.$ Replace m' by m'' in (9.19) and let m'' goes to $+\infty$, we have

$$\bar{\partial}u_{l,\epsilon,j} = \bar{\partial}\left((1 - v'_{\epsilon}(\Psi))fF^{1+\delta}\right).$$
(9.21)

Denote $F_{l,\epsilon,j} := -u_{l,\epsilon,j} + (1 - v'_{\epsilon}(\Psi))fF^{1+\delta}$. It follows from (9.21) and inequality (9.20) that we know $F_{l,\epsilon,j}$ is a holomorphic (n, 0) form on M_j and

$$\int_{M_{j}} |F_{l,\epsilon,j} - (1 - v_{\epsilon}'(\Psi)) f F^{1+\delta}|^{2} e^{v_{\epsilon}(\Psi) - h_{l} - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi))$$

$$\leq \int_{M_{j}} v_{\epsilon}''(\Psi) |fF|^{2} e^{-u(-v_{\epsilon}(\Psi)) - h_{l}} < +\infty.$$
(9.22)

Step 6: Letting $\epsilon \to 0$. Note that

$$v_{\epsilon}''(\Psi)|fF|^{2}e^{-u(-v_{\epsilon})(\Psi)-h_{l}} \leq \frac{2}{B}\sup_{\epsilon}\sup_{M_{j}}\left(e^{-u(-v_{\epsilon}(\Psi))+l}|F|^{2}\right)\mathbb{I}_{\{\Psi<-t_{0}\}}|f|^{2}.$$

It follows from $\int_{\overline{M_i} \cap \{\Psi < -t_0\}} |f|^2 < +\infty$ and dominated convergence theorem that

$$\lim_{\epsilon \to 0} \int_{M_j} v_{\epsilon}''(\Psi) |fF|^2 e^{-u(-v_{\epsilon}(\Psi)) - h_l}
= \int_{M_j} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |fF|^2 e^{-u(-v(\Psi)) - h_l}
\leq \left(\sup_{M_j} e^{-u(-v(\Psi))} \right) \int_{M_j} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |fF|^2 e^{-h_l}$$
(9.23)

When $\Psi < -t_0$, we know that $\max\{\psi + T, 2\log|F|\} = 2\log|F|$. Note that $h + \delta \max\{\psi + T, 2\log|F|\} = \varphi + \Psi = \varphi_{\alpha} + (1+\delta)\max\{\psi + T, 2\log|F|\} + \Psi.$ Hence

$$\int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B<\Psi<-t_{0}\}} |fF|^{2} e^{-h_{l}} \\
= \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B<\Psi<-t_{0}\}} |fF^{1+\delta}|^{2} e^{-h_{l}-\delta 2\log|F|} \\
\leq \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B<\Psi<-t_{0}\}} |fF^{1+\delta}|^{2} e^{-h-\delta \max\{\psi+T,2\log|F|\}} \\
= \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B<\Psi<-t_{0}\}} |f|^{2} e^{-\varphi_{\alpha}-\Psi} < +\infty.$$
(9.24)

Note that

$$\inf_{\epsilon} \inf_{M_j} e^{v_{\epsilon}(\Psi) - h_l - \delta \max\{\psi + T, 2\log|F|\}} c(-v_{\epsilon}(\Psi)) > 0.$$

We have

$$\sup_{\epsilon} \int_{M_j} |F_{l,\epsilon,j} - (1 - v'_{\epsilon}(\Psi))fF^{1+\delta}|^2 < +\infty.$$

We also note that

$$\sup_{\epsilon} \int_{M_j} |(1 - v'_{\epsilon}(\Psi)) f F^{1+\delta}|^2 \le (\sup_{M_j} |F|^{2(1+\delta)}) \int_{\overline{M_j} \cap \{\Psi < -t_0\}} |f|^2 < +\infty.$$

Then we know that

$$\sup_{\epsilon} \int_{M_j} |F_{l,\epsilon,j}|^2 < +\infty,$$

and there exists a subsequence of $\{F_{l,\epsilon,j}\}$ (also denoted by $\{F_{l,\epsilon,j}\}$) compactly convergent to a holomorphic (n, 0) form $F_{l,j}$ on M_j . It follows from Fatou's Lemma and inequalities (9.22), (9.23), (9.24) that

$$\begin{split} &\int_{M_{j}} |F_{l,j} - (1 - b(\Psi))fF^{1+\delta}|^{2}e^{v(\Psi) - h_{l} - \delta \max\{\psi + T, 2\log|F|\}}c(-v(\Psi)) \\ &\leq \liminf_{\epsilon \to 0} \int_{M_{j}} |F_{l,\epsilon,j} - (1 - v_{\epsilon}'(\Psi))fF^{1+\delta}|^{2}e^{v_{\epsilon}(\Psi) - h_{l} - \delta \max\{\psi + T, 2\log|F|\}}c(-v_{\epsilon}(\Psi)) \\ &\leq \liminf_{\epsilon \to 0} \int_{M_{j}} v_{\epsilon}''(\Psi)|fF|^{2}e^{-u(-v_{\epsilon}(\Psi)) - h_{l}} \\ &\leq \left(\sup_{M_{j}} e^{-u(-v(\Psi))}\right) \int_{M_{j}} \frac{1}{B}\mathbb{I}_{\{-t_{0} - B < \Psi < -t_{0}\}}|fF|^{2}e^{-h_{l}} \\ &\leq \left(\sup_{M_{j}} e^{-u(-v(\Psi))}\right) \int_{M_{j}} \frac{1}{B}\mathbb{I}_{\{-t_{0} - B < \Psi < -t_{0}\}}|f|^{2}e^{-\varphi_{\alpha} - \Psi} < +\infty. \end{split}$$

$$(9.25)$$

Step 7: Letting $l \to +\infty$.

It follows from $h_l \leq h_1$ for any $l \in \mathbb{Z}^+$ and h_1 is a plurisubharmonic function on M that

$$\inf \inf_{M_j} e^{v(\Psi) - h_l - \delta \max\{\psi + T, 2\log|F|\}} c(-v(\Psi)) \ge \hat{C}_j \inf_{M_j} e^{-h_1} > 0,$$

where $\hat{C}_j = \inf_{M_j} e^{v(\Psi) - \delta \max\{\psi + T, 2 \log |F|\}} c(-v(\Psi))$ is a positive number. By inequality (9.25), we have

$$\sup_{l} \int_{M_j} |F_{l,j} - (1 - b(\Psi))fF^{1+\delta}|^2 < +\infty.$$

Note that

$$\sup_{l} \int_{M_{j}} |(1 - b(\Psi))fF^{1+\delta}|^{2} \le \left(\sup_{M_{j}} |F|^{2(1+\delta)} \right) \int_{\overline{M_{j}} \cap \{\Psi < -t_{0}\}} |f|^{2} < +\infty.$$

Hence we know that

$$\sup_{l} \int_{M_j} |F_{l,j}|^2 < +\infty,$$

and there exists a subsequence of $\{F_{l,j}\}$ (also denoted by $\{F_{l,j}\}$) compactly convergent to a holomorphic (n, 0) form F_j on M_j . It follows from Fatou's Lemma and inequality (9.25) that

$$\int_{M_{j}} |F_{j} - (1 - b(\Psi))fF^{1+\delta}|^{2}e^{v(\Psi) - \varphi - \Psi}c(-v(\Psi))$$

$$= \int_{M_{j}} |F_{j} - (1 - b(\Psi))fF^{1+\delta}|^{2}e^{v(\Psi) - h - \delta\max\{\psi + T, 2\log|F|\}}c(-v(\Psi))$$

$$\leq \liminf_{l \to +\infty} \int_{M_{j}} |F_{l,j} - (1 - b(\Psi))fF^{1+\delta}|^{2}e^{v(\Psi) - h_{l} - \delta\max\{\psi + T, 2\log|F|\}}c(-v(\Psi))$$

$$\leq \left(\sup_{M_{j}} e^{-u(-v(\Psi))}\right) \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0} - B < \Psi < -t_{0}\}} |f|^{2}e^{-\varphi_{\alpha} - \Psi} < +\infty.$$
(9.26)

Step 8: Letting $j \to +\infty$. It is easy to see that

$$\left(\sup_{M_{j}} e^{-u(-v(\Psi))}\right) \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B < \Psi < -t_{0}\}} |f|^{2} e^{-\varphi_{\alpha}-\Psi}$$

$$\leq \left(\sup_{M} e^{-u(-v(\Psi))}\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_{0}-B < \Psi < -t_{0}\}} |f|^{2} e^{-\varphi_{\alpha}-\Psi} < +\infty.$$
(9.27)

For fixed j, as $e^{v(\Psi)-\varphi-\Psi}c(-v(\Psi))$ has a positive lower bound on any $\overline{M_j}$, we have for $j_1 > j$,

$$\sup_{j_1} \int_{M_j} |F_{j_1} - (1 - b(\Psi))fF^{1+\delta}|^2 < +\infty.$$

Combining with

$$\int_{M_j} |(1 - b(\Psi))fF^{1+\delta}|^2 \le \left(\sup_{M_j} |F^{1+\delta}|^2\right) \int_{M_j \cap \{\Psi < -t_0\}} |f|^2 < +\infty,$$

we know that for $j_1 > j$, $\int_{M_j} |F_{j_1}|^2$ is uniformly bounded with respect to j_1 .

By diagonal method, there exists a subsequence of $\{F_i\}$ (also denoted by $\{F_i\}$) compactly convergent to a holomorphic (n, 0) form \tilde{F} on M. Then it follows from Fatou's Lemma, inequality (9.26) and inequality (9.27) that

$$\int_{M} |\tilde{F} - (1 - b(\Psi))fF^{1+\delta}|^{2} e^{v(\Psi) - \varphi - \Psi} c(-v(\Psi))$$

$$\leq \liminf_{j \to +\infty} \int_{M_{j}} |F_{j} - (1 - b(\Psi))fF^{1+\delta}|^{2} e^{v(\Psi) - \varphi - \Psi} c(-v(\Psi))$$

$$\leq \liminf_{j \to +\infty} \left(\sup_{M_{j}} e^{-u(-v(\Psi))} \right) \int_{M_{j}} \frac{1}{B} \mathbb{I}_{\{-t_{0} - B < \Psi < -t_{0}\}} |f|^{2} e^{-\varphi_{\alpha} - \Psi}$$

$$\leq \left(\sup_{M} e^{-u(-v(\Psi))} \right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_{0} - B < \Psi < -t_{0}\}} |f|^{2} e^{-\varphi_{\alpha} - \Psi} < +\infty.$$
(9.28)

Step 9: ODE System.

Step 9: ODL system. Now we want to find η and ϕ such that $(\eta + g^{-1}) = e^{-v_{\epsilon}(\Psi_m)}e^{-\phi}\frac{1}{c(-v_{\epsilon}(\Psi_m))}$. As $\eta = s(-v_{\epsilon}(\Psi_m))$ and $\phi = u(-v_{\epsilon}(\Psi_m))$, we have $(\eta + g^{-1})e^{v_{\epsilon}(\Psi_m)}e^{\Phi} = \left((s + \frac{s'^2}{u''s - s''})e^{-t}e^u\right)\circ$

 $(-v_{\epsilon}(\Psi_m)).$

Summarizing the above discussion about s and u, we are naturally led to a system of ODEs:

$$1)(s + \frac{s'^2}{u''s - s''})e^{u - t} = \frac{1}{c(t)},$$

$$2)s' - su' = 1,$$
(9.29)

when $t \in (T, +\infty)$.

We solve the ODE system (9.29) and get $u(t) = -\log(\frac{1}{\delta}c(T)e^{-T} + \int_T^t c(t_1)e^{-t_1}dt_1)$ and $s(t) = \frac{\int_T^t (\frac{1}{\delta}c(T)e^{-T} + \int_T^{t_2} c(t_1)e^{-t_1}dt_1)dt_2 + \frac{1}{\delta^2}c(T)e^{-T}}{\frac{1}{\delta}c(T)e^{-T} + \int_T^t c(t_1)e^{-t_1}dt_1}$. It follows that $s \in C^{\infty}([T, +\infty))$ satisfies $s \ge \frac{1}{\delta}$ and $u \in C^{\infty}([T, +\infty))$ satisfies

u''s - s'' > 0.

As $u(t) = -\log(\frac{1}{\delta}c(T)e^{-T} + \int_T^t c(t_1)e^{-t_1}dt_1)$ is decreasing with respect to t, then it follows from $-T \ge v(t) \ge \max\{t, -t_0 - B_0\} \ge -t_0 - B_0$, for any $t \le 0$ that

$$\sup_{M} e^{-u(-v(\psi))} \le \sup_{t \in [T, t_0 + B]} e^{-u(t)} = \frac{1}{\delta} c(T) e^{-T} + \int_{T}^{t_0 + B} c(t_1) e^{-t_1} dt_1.$$
(9.30)

Combining with inequality (9.28), we have

$$\begin{split} &\int_{M} |\tilde{F} - (1 - b(\Psi))fF^{1+\delta}|^2 e^{v(\Psi) - \varphi - \Psi} c(-v(\Psi)) \\ &\leq \left(\frac{1}{\delta} c(T) e^{-T} + \int_{T}^{t_0 + B} c(t_1) e^{-t_1} dt_1\right) \int_{M} \frac{1}{B} \mathbb{I}_{\{-t_0 - B < \Psi < -t_0\}} |f|^2 e^{-\varphi_{\alpha} - \Psi} < +\infty. \end{split}$$

Lemma 2.1 is proved.

Acknowledgements. The first author and the second author were supported by National Key R&D Program of China 2021YFA1003103. The first author was supported by NSFC-11825101, NSFC-11522101 and NSFC-11431013.

References

- Q.A. Guan, Z.T. Mi and Z. Yuan, [1] S.J. Bao, Concavity property of minimal L^2 weights, integrals with Lebesgue measurable gain VII-negligible https://www.researchgate.net/publication/358215153
- [2] S.J. Bao, Q.A. Guan and Z. Yuan, Boundary points, minimal L^2 integrals and concavity property, arXiv:2203.01648.
- Sébastien Boucksom, Singularities of plurisubharmonic functions and multiplier ideals, electronically accessible at http://sebastien.boucksom.perso.math.cnrs.fr/notes/L2.pdf.
- [4] J.Y. Cao, Ohsawa-Takegoshi extension theorem for compact Kähler manifolds and applications, Complex and symplectic geometry, 19-38, Springer INdAM Ser., 21, Springer, Cham, 2017.
- [5] J.Y. Cao, J-P. Demailly and S. Matsumura, A general extension theorem for cohomology classes on non reduced analytic subspaces, Sci. China Math. 60 (2017), no. 6, 949-962, DOI 10.1007/s11425-017-9066-0.
- [6] B.Y. Chen, A degenerate Donnelly-Fefferman theorem and its applications. In: Byun J., Cho H., Kim S., Lee KH., Park JD. (eds) Geometric Complex Analysis. Springer Proceedings in Mathematics & Statistics, vol 246. Springer, Singapore. https://doi.org/10/1007/978-981-13-1672-2_6

- [7] T. Darvas, E. Di Nezza and H.C. Lu, Monotonicity of nonpluripolar products and complex Monge-Ampére equations with prescribed singularity, Anal. PDE 11 (2018), no. 8, 2049-2087.
- [8] T. Darvas, E. Di Nezza and H.C. Lu, The metric geometry of singularity types, J. Reine Angew. Math. 771 (2021), 137-170.
- [9] J.-P Demailly, Estimations L² pour l'opérateur ∂ d'un fibré vectoriel holomorphe semi-positif au-dessus d'une variété kählérienne complète.(French) L² estimates for the ∂-operator of a semipositive holomorphic vector bundle over a complete Kähler manifold, Ann. Sci. École Norm. Sup. (4) 15 (1982), no. 3, 457-511.
- [10] J.-P Demailly, Regularization of closed positive currents of type (1,1) by the flow of a Chern connection, Actes du Colloque en l'honneur de P.Dolbeault (Juin 1992),édité par H.Skoda et J.-M Trépreau, Aspect of Mathematics, Vol.E26, Vieweg,1994,105-126.
- [11] J.-P Demailly, On the Ohsawa-Takegoshi-Manivel L² extension theorem, Complex analysis and geometry (Paris, 1997), Progr. Math., 188, Birkhäuser, Basel, 2000, p47-82.
- [12] J.-P Demailly, Complex analytic and differential geometry, electronically accessible at https://www-fourier.ujf-grenoble.fr/ demailly/manuscripts/agbook.pdf.
- [13] J.-P Demailly, Multiplier ideal sheaves and analytic methods in algebraic geometry, School on Vanishing Theorems and Effective Result in Algebraic Geometry (Trieste, 2000),1-148,ICTP IECT.Notes, 6, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2001.
- [14] J.-P Demailly, Analytic Methods in Algebraic Geometry, Higher Education Press, Beijing, 2010.
- [15] J.-P Demailly, L. Ein and R. Lazarsfeld, A subadditivity property of multiplier ideals, Michigan Math. J. 48 (2000) 137-156.
- [16] J.-P Demailly and J. Kollár, Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds, Ann. Sci. Éc. Norm. Supér. (4) 34 (4) (2001) 525-556.
- [17] J.-P Demailly and T. Peternell, A Kawamata-Viehweg vanishing theorem on compact Kähler manifolds, J. Differential Geom. 63 (2) (2003) 231-277.
- [18] J.E. Fornæss, Several complex variables. arXiv:1507.00562
- [19] J.E. Fornæss and J.J. Wu, A global approximation result by Bert Alan Taylor and the strong openness conjecture in Cⁿ, J. Geom. Anal. 28 (2018), no. 1, 1-12.
- [20] J.E. Fornæss and J.J. Wu, Weighted approximation in C, Math. Z. 294 (2020), no. 3-4, 1051-1064.
- [21] H. Grauert and R. Remmert, Coherent analytic sheaves, Grundlehren der mathematischen Wissenchaften, 265, Springer-Verlag, Berlin, 1984.
- [22] Q.A. Guan, Genral concavity of minimal L^2 integrals related to multiplier sheaves, arXiv:1811.03261.v4.
- [23] Q.A. Guan, A sharp effectiveness result of Demailly's strong Openness conjecture, Adv.Math. 348 (2019) :51-80.
- [24] Q.A. Guan, A proof of Saitoh's conjecture for conjugate Hardy H2 kernels. J. Math. Soc. Japan 71 (2019), no. 4, 1173–1179.
- [25] Q.A. Guan and Z.T. Mi, Concavity of minimal L^2 integrals related to multiplier ideal sheaves, arXiv:2106.05089v2.
- [26] Q.A. Guan and Z.T. Mi, Concavity of minimal L^2 integrals related to multiplier ideal sheaves on weakly pseudoconvex Kähler manifolds, submitted.
- [27] Q.A. Guan, Z.T. Mi and Z. Yuan, Concavity property of minimal L² integrals with Lebesgue measurable gain II, https://www.researchgate.net/publication/354464147.
- [28] Q.A. Guan and Z. Yuan, Concavity property of minimal L² integrals with Lebesgue measurable gain, https://www.researchgate.net/publication/353794984.
- [29] Q.A. Guan and Z. Yuan, An optimal support function related to the strong openness property, arXiv:2105.07755v2.
- [30] Q.A. Guan and Z. Yuan, Effectiveness of strong openness property in L^p , arXiv:2106.03552v3.
- [31] Q.A. Guan and Z. Yuan, Twisted version of strong openness property in L^p , arXiv:2109.00353.
- [32] Q.A. Guan and X.Y. Zhou, A solution of an L² extension problem with an optimal estimate and applications, Ann. of Math. (2) 181 (2015), no. 3, 1139–1208.
- [33] Q.A. Guan and X.Y Zhou, A proof of Demailly's strong openness conjecture, Ann. of Math.(2) 182 (2015), no. 2, 605-616.

- [34] Q.A. Guan and X.Y Zhou, Effectiveness of Demailly's strong openness conjecture and related problems, Invent. Math. 202 (2015), no. 2, 635-676.
- [35] Q.A. Guan and X.Y. Zhou, Strong openness of multiplier ideal sheaves and optimal L² extension, Sci. China Math. (2017), 60:967-976.
- [36] Q.A. Guan and X.Y. Zhou, Restriction formula and subadditivity property related to multiplier ideal sheaves, J. Reine Angew. Math. 769, 1-33 (2020).
- [37] H. Guenancia, Toric plurisubharmonic functions and analytic adjoint ideal sheaves, Math. Z. 271 (3-4) (2012) 1011-1035.
- [38] L. Hörmander, An introduction to complex analysis in several variables, 3rd edn (North-Holland, Amsterdam, 1990).
- [39] M. Jonsson and M. Mustață, Valuations and asymptotic invariants for sequences of ideals, Annales de L'Institut Fourier A. 2012, vol. 62, no.6, pp. 2145-2209.
- [40] M. Jonsson and M. Mustaţă, An algebraic approach to the openness conjecture of Demailly and Kollár, J. Inst. Math. Jussieu (2013), 1–26.
- [41] D. Kim, Skoda division of line bundle sections and pseudo-division, Internat. J. Math. 27 (2016), no. 5, 1650042, 12 pp.
- [42] D. Kim and H. Seo, Jumping numbers of analytic multiplier ideals (with an appendix by Sebastien Boucksom), Ann. Polon. Math., 124 (2020), 257-280.
- [43] R. Lazarsfeld, Positivity in Algebraic Geometry. I. Classical Setting: Line Bundles and Linear Series. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 48. Springer-Verlag, Berlin, 2004;
 R. Lazarsfeld, Positivity in Algebraic Geometry. II. Positivity for vector bundles, and multiplier ideals. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 49. Springer-Verlag, Berlin, 2004.
- [44] J. D. McNeal, D. Varolin, L^2 estimate for the $\bar\partial$ operator. Bull.Math. Sci. 5 (2015), no.2, 179-249.
- [45] A. Nadel, Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature, Ann. of Math. (2) 132 (3) (1990) 549-596.
- [46] Y.T. Siu, The Fujita conjecture and the extension theorem of Ohsawa-Takegoshi, Geometric Complex Analysis, World Scientific, Hayama, 1996, pp.223-277.
- [47] Y.T. Siu, Invariance of plurigenera and torsion-freeness of direct image sheaves of pluricanonical bundles, in Finite or Infinite Dimensional Complex Analysis and Applications (Kluwer, Boston, MA, 2004), Adv. Complex Anal. Appl. 2, pp. 45-83.
- [48] Y.T. Siu, Multiplier ideal sheaves in complex and algebraic geometry, Sci. China Ser. A 48 (suppl.) (2005) 1-31.
- [49] Y.T. Siu, Dynamic multiplier ideal sheaves and the construction of rational curves in Fano manifolds, Complex Analysis and Digital Geometry, in: Acta Univ. Upsaliensis Skr. Uppsala Univ. C Organ. Hist., vol.86, Uppsala Universitet, Uppsala, 2009, pp.323-360.
- [50] G. Tian, On Kähler-Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, Invent. Math. 89 (2) (1987) 225-246.
- [51] X.Y Zhou and L.F.Zhu, An optimal L² extension theorem on weakly pseudoconvex Kähler manifolds, J. Differential Geom.110(2018), no.1, 135-186.
- [52] X.Y Zhou and L.F.Zhu, Optimal L² extension of sections from subvarieties in weakly pseudoconvex manifolds. Pacific J. Math. 309 (2020), no. 2, 475-510.
- [53] X.Y. Zhou and L.F. Zhu, Siu's lemma, optimal L2 extension and applications to twisted pluricanonical sheaves, Math. Ann. 377 (2020), no. 1-2, 675-722.

Qi'AN GUAN: School of Mathematical Sciences, Peking University, Beijing 100871, China.

Email address: guanqian@math.pku.edu.cn

ZHITONG MI: INSTITUTE OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING, CHINA

Email address: zhitongmi@amss.ac.cn

Zheng Yuan: School of Mathematical Sciences, Peking University, Beijing 100871, China.

 $Email \ address: \verb"zyuan@pku.edu.cn" \\$