Concentration Properties of Random Codes

Lan V. Truong, Giuseppe Cocco, Josep Font-Segura and Albert Guillén i Fàbregas

Abstract

This paper studies concentration properties of random codes. Specifically, we show that, for discrete memoryless channels, the error exponent of a randomly generated code with pairwise-independent codewords converges in probability to its expectation—the typical error exponent. For high rates, the result is a consequence of the fact that the random-coding error exponent and the sphere-packing error exponent coincide. For low rates, instead, the convergence is based on the fact that the union bound accurately characterizes the probability of error. The paper also zooms into the behavior at asymptotically low rates, and shows that the error exponent converges in distribution to a Gaussian-like distribution. Finally, we present several results on the convergence of the error probability and error exponent for generic ensembles and channels.

I. INTRODUCTION

The discrete memoryless channel (DMC) has been devoted a lot of interest in information theory ever since in [1] Shannon showed that for DMC there exist codes whose probability of error vanishes with the codewords length for rates below the channel capacity. Since then, one of the most active areas of research in Information Theory has been the study of properties of the probability of error. For rates below capacity, Fano [2] characterized the exponential decay of the error probability defining the error exponent as the negative normalized logarithm of the ensemble-average error probability, i.e., the random coding exponent (RCE). In [3], Gallager derived the RCE in a simpler way and introduced the idea of expurgation in order to obtain an improved exponent the at low rates. A lower bound on the error probability in the DMC, called sphere-packing bound, was first introduced in [4] and it was shown to coincide with the RCE for rates higher than a certain critical rate. Nakiboğlu in [5] recently derived sphere-packing bounds for some stationary memoryless channels using Augustin's method [6].

In [7], Barg and Forney studied the random-coding ensemble over the binary symmetric channel (BSC) with maximum likelihood decoding and showed that the error exponent of most random codes in the i.i.d. ensemble is close to the so-called typical random coding (TRC) exponent, strictly larger than the RCE at low rates. Upper and

L. V. Truong is with the Department of Engineering, University of Cambridge, Cambridge CB2 1PZ, U.K. (e-mail: lt407@cam.ac.uk). G. Cocco and J. Font-Segura are with Department of Information and Communication Technologies, Universitat Pompeu Fabra, Barcelona 08018, Spain (e-mail: giuseppe.cocco@upf.edu, josep.font@upf.edu). A. Guillén i Fàbregas is with the Department of Engineering, University of Cambridge, Cambridge CB2 1PZ, U.K. and the Department of Information and Communication Technologies, Universitat Pompeu Fabra, Barcelona 08018, Spain (e-mail: guillen@ieee.org).

This work has been funded in part by the European Research Council under ERC grant agreement 725411, by the Secretary of Universities and Research (Catalan Government) under a Beatriu de Pinós postdoctoral fellowship, and by the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement 801370.

This work has been presented in part at the 2021 IEEE Information Theory Workshop, Kanazawa, Japan.

lower bounds on the TRC for constant-composition codes and general DMCs were provided in [8]. For the same type of codes and channels, Merhav [9] determined the exact TRC error exponent and a wide class of stochastic decoders called generalized likelihood decoder (GLD), of which maximum-likelihood is a special case. Merhav derived the TRC exponent for spherical codes over coloured Gaussian channels [10] and for random convolutional code ensembles [11]. The error exponent of a random pairwise-independent constant-composition code with GLD was shown to converge in probability to the TRC in [12]. The convergence is non-symmetric: the lower tail decays exponentially while the upper tail decays doubly-exponentially. The latter was first established for a limited range of rates in [13]. The TRC was shown to be universally achievable with the likelihood mutual information decoder in [14]. For pairwise-independent ensembles and arbitrary channels, Cocco *et al.* showed in [15] that the probability that the exponent of a given code in the ensemble is smaller than a lower bound on the TRC exponent is vanishingly small.

The main motivation of our work is the fact that the aforementioned results highlight the importance of the statistical properties of the error probability and the error exponent across the random-coding ensemble. After describing the main performance metrics of random codes for reliable communication in the next section, namely the error probability and the error exponent, we use the notion of convergence in probability and convergence in distribution to obtain a number of concentration results of such performance metric, seen as sequences of random variables, as the blocklength tends to infinity. Since neither the error probability or the error exponent are sums of i.i.d. terms, our results are based on probability results beyond the central limit theorem, such as the Stein's method and a novel, modified Wasserstein metric. We anticipate here some of our main results in Sec. III, valid for the DMC and the i.i.d. and constant-composition ensembles.

- We show in Theorem 1 that the error exponent converges in probability the TRC exponent.
- In Theorems 2–4 we provide bounds on the rate of such convergence.
- For codes with a constant number of codewords, we obtain in Theorem 5 that the error exponent converges in distribution to a quasi-Gaussian, while for codes with a sub-exponential number of codewords it does converge to a Gaussian, as shown in Theorem 6.

For general channels under some additional conditions, we obtain in Sec. IV the following results.

- For any channel and capacity-achieving ensemble, Theorem 7 states that the error probability converges to the ensemble average.
- Theorems 8–11 discuss several convergence results relating random properties of the error probability, the random-coding error exponent and the TRC exponent.
- Sufficient conditions for the union bound on the error probability and any general function of the error probability to converge to a Gaussian are respectively described in Theorems 12–13.

The proofs of our main results, including our modified Wasserstein metric, are reported in Sec. V.

II. PRELIMINARIES

We consider the problem of transmitting M_n equiprobable messages over a DMC with transition probability Wand finite input and output alphabets \mathcal{X} and \mathcal{Y} , respectively. We employ a codebook $c_n = \{x_1, x_2, \cdots, x_{M_n}\}$ with $x_m \in \mathcal{X}^n$, for $m = 1, ..., M_n$. The conditional distribution of a channel output $y \in \mathcal{Y}^n$ given a transmitted codeword x is given by $W^n(y|x) = \prod_{i=1}^n W(y_i|x_i)$. We consider maximum-likelihood decoding, that is, the decoder produces an estimate of the transmitted codeword as $\hat{x} = \arg \max_{x \in c_n} W^n(y|x)$. The error probability of such code is

$$P_{\rm e}(\boldsymbol{c}_n) = \frac{1}{M_n} \sum_{m=1}^{M_n} \mathbb{P}\bigg[\bigcup_{\bar{m} \neq m} \{\boldsymbol{x}_m \to \boldsymbol{x}_{\bar{m}}\}\bigg],\tag{1}$$

where $\{x_m \to x_{\bar{m}}\} = \{y \in \mathcal{Y} : W^n(y|x_{\bar{m}}) \ge W^n(y|x_m)\}$ is the maximum-likelihood pairwise error event, i.e., the event of deciding in favor of codeword $x_{\bar{m}}$ when codeword x_m was transmitted. The error exponent of code c_n is defined as

$$E_n(c_n) = -\frac{1}{n} \log P_{\mathbf{e}}(c_n).$$
⁽²⁾

Let $R = \lim_{n\to\infty} \frac{1}{n} \log M_n$ be the rate of the code in bits per channel use. An error exponent E(R) is said to be achievable when there exists a sequence of codes $\{c_n\}_{n=1}^{\infty}$ such that $\liminf_{n\to\infty} E_n(c_n) \ge E(R)$. The channel capacity C is the supremum of the code rates R such that E(R) > 0.

We next consider the random generation of the codebook. Similarly to random variables, C_n denotes a random code, and c_n denotes a specific code in the ensemble. In particular, we consider the pairwise-independent random-coding ensemble, i.e., the set of random codes C_n whose codewords X_1, X_2, \dots, X_{M_n} are pairwise-independently generated. We consider the i.i.d. ensemble, in which each codeword is generated according to the distribution

$$Q^{\text{iid}}(\boldsymbol{x}) = \prod_{i=1}^{n} Q(x_i), \tag{3}$$

Q being the single-letter distribution and the constant-composition ensemble, in which each codeword is generated according to the distribution

$$Q^{\rm cc}(\boldsymbol{x}) = \frac{1}{|\mathcal{T}_n(Q_n)|} \mathbb{1}\{\boldsymbol{x} \in \mathcal{T}_n(Q_n)\}$$
(4)

where $\mathcal{T}_n(Q_n)$ is the type class of composition $Q_n \in \mathcal{P}_n(\mathcal{X})$, i.e., all *n*-length sequences whose empirical distribution is Q_n such that $\max_x |Q_n(x) - Q(x)| \leq \frac{1}{n}$ for a given distribution Q. For a given distribution or composition Q, we define the random-coding error exponent $E_{\rm rce}(R,Q)$ as

$$E_{\rm rce}(R,Q) = \lim_{n \to \infty} -\frac{1}{n} \log \mathbb{E}[P_{\rm e}(\mathcal{C}_n,Q)],\tag{5}$$

where $P_{e}(C_n, Q)$ denotes the error probability of the random code ensemble C_n parametrized by the distribution or composition Q and where the expectation is taken over the code ensemble. Eq. (5) suggests that $E_{rce}(R, Q)$ is the asymptotic exponent of the ensemble-average probability of error. For the i.i.d. ensemble, it is known that [3, Th. 1]

$$E_{\rm rce}^{\rm iid}(R,Q) = \max_{0 \le \rho \le 1} \left\{ E_0^{\rm iid}(\rho,Q) - \rho R \right\}$$
(6)

with

$$E_0^{\text{iid}}(\rho, Q) = -\log \sum_y \left(\sum_x Q(x) W(y|x)^{\frac{1}{1+\rho}}\right)^{1+\rho};$$
(7)

reamble we have that (6) remains valid but $\overline{U}^{ijd}(x, 0)$ is replaced by

while for the constant-composition ensemble, we have that (6) remains valid, but $E_0^{\text{iid}}(\rho, Q)$ is replaced by [16, Eq. (53)]

$$E_0^{\rm cc}(\rho,Q) = \sup_{a(x)} -\log \sum_y \left(\sum_x Q(x)W(y|x)^{\frac{1}{1+\rho}} e^{a(x)-\phi_a}\right)^{1+\rho}$$
(8)

where a(x) is an auxiliary function and $\phi_a = \sum_x Q(x)a(x)$. It is known that for any given Q, $E_{\text{rce}}^{\text{iid}}(R,Q) \leq E_{\text{rce}}^{\text{cc}}(R,Q)$ (see e.g. [17]).

While $E_{\rm rce}(R,Q)$ in (5) is the limiting exponential rate of decay of the ensemble average probability of error, the typical random-coding exponent $E_{\rm trc}(R,Q)$ is instead defined as the limiting expected error exponent over the ensemble, that is,

$$E_{\rm trc}(R,Q) = \lim_{n \to \infty} -\frac{1}{n} \mathbb{E} \big[\log P_{\rm e}(\mathcal{C}_n,Q) \big].$$
(9)

We observe that Jensen's inequality implies that the random-coding error exponent in (5) and the typical randomcoding error exponent in (9) satisfy $E_{\rm rce}(R,Q) \leq E_{\rm trc}(R,Q)$. The proofs of our results exploit the idea that there are two rate regimes; one where $E_{\rm rce}(R,Q) < E_{\rm trc}(R,Q)$ and the other where $E_{\rm rce}(R,Q) = E_{\rm trc}(R,Q)$. For DMCs with constant-composition codes [9] and for i.i.d. codes over the BSC [7], the typical error exponent can be further expressed in terms of the expurgated error exponent $E_{\rm ex}(R,Q)$ as

$$E_{\rm trc}(R,Q) = \max\{E_{\rm ex}(2R,Q) + R, E_{\rm rce}(R,Q)\}.$$
(10)

In the next sections, we derive concentration results of the error probability (1) and the error exponent (2) of sequences of random codes C_n in the asymptotic regime as $n \to \infty$. As sequence of random variables, we assume throughout the paper that $P_e(C_n)$ and $E_n(C_n)$ do not diverge as $n \to \infty$, leaving particular cases such as channels with positive zero-error capacity beyond the scope of this paper. We will use the notion of convergence in probability and convergence in distribution. A sequence of random variables $\{A_n\}_{n=1}^{\infty}$ converges to A in probability, denoted as $A_n \xrightarrow{(p)} A$ if for all $\delta > 0$ [18, Sec. 2.2],

$$\lim_{n \to \infty} \mathbb{P}[|A_n - A| > \delta] = 0.$$
⁽¹¹⁾

If $A_n = \frac{1}{n} \sum_{i=1}^{n} X_i$, where X_i , i = 1, ..., n are i.i.d. random variables, then $A = \mathbb{E}[X_1]$ and (11) reduces to the weak law of large numbers [18, Th. 2.2.3]. The weak law of large numbers is at the core of the asymptotic equipartition property, a widely used tool in information theory to establish the achievable rates using random coding [19]; it is well known that the asymptotic equipartition property is not sufficient to show the achievability of error exponents. Alternatively to (11), we say that a sequence of random variables $\{A_n\}_{n=1}^{\infty}$ converges to A in distribution, denoted as $A_n \xrightarrow{(d)} A$ if [18, Sec. 3.2]

$$\lim_{n \to \infty} \sup_{x \in \mathbb{R}} \left| \mathbb{P}[A_n \le x] - \mathbb{P}[A \le x] \right| = 0$$
(12)

for all continuous points x of $\mathbb{P}[A \leq x]$.

We first state in Sec. III our main results for the relevant case of i.i.d. and constant-composition ensembles over generic DMCs. Additional results are shown in Sec. IV for general channels with few additional assumptions on the ensemble or conditions on the statistical behavior of the error probability as $n \to \infty$. The proofs of our theorems are included in Sec. V, while most lemmas thereby used are proved in the Appendix. In this section, we introduce our main concentration results for DMCs. Our first result states the convergence in probability of the error exponent $E_n(\mathcal{C}_n)$ to the TRC exponent $E_{trc}(R)$. Since the exponent of the probability of error is not a sum of i.i.d. terms, the weak law of large numbers cannot be applied. This result holds for i.i.d. and constant-composition ensembles over DMCs with input distribution Q.

Theorem 1: For a general DMC channel, i.i.d. and constant-composition ensembles and rates $0 \le R < C$, it holds that

$$E_n(\mathcal{C}_n) \xrightarrow{(\mathbf{p})} E_{\mathrm{trc}}(R,Q).$$
 (13)

Proof: Sec. V-A.

Theorem 1 not only proves the achievability of the TRC exponent, but also shows that the probability of finding a code in the ensemble with higher or lower exponent than the TRC exponent tends to zero. The above concentration property gives more information about the error exponent behaviour of the ensemble than the traditional derivation of the random coding error exponent, which computes the exponent of the expected error probability. This way, the TRC emerges as the most likely error exponent for pairwise-independent random-coding ensembles as the block length n tends to infinity —if one wishes to improve the error exponent, one must improve the ensemble. The TRC exponent is lower than or equal to the expurgated exponent and can in some case be strictly smaller. This implies that the codes in the pairwise independent ensemble that achieve the expurgated exponent are not typical codes and are unlikely to be found by random generation.

The proof of Theorem 1 requires different techniques for the rate regimes $E_{\text{trc}}(R,Q) = E_{\text{rce}}(R,Q)$ and $E_{\text{trc}}(R,Q) > E_{\text{rce}}(R,Q)$. For the first regime, corresponding to high rates such that $R_{\text{crit}}(R,Q) \leq R \leq C$, where $R_{\text{crit}}(R,Q)$ is the critical rate, we exploit the fact that $E_{\text{trc}}(R,Q) = E_{\text{rce}}(R,Q)$ and the Levy's continuity theorem [20, Sec. XIII.1] to obtain that the moment-generating function of the random variable $E_n(\mathcal{C}_n)$ converges to the moment-generating function of a deterministic variable with value $E_{\text{trc}}(R,Q)$. In other words, $E_n(\mathcal{C}_n)$ converges in distribution to a constant $E_{\text{trc}}(R,Q)$, implying the convergence in probability in (13) for both the i.i.d. and the constant-composition ensembles.

For the second regime where $E_{\rm trc}(R,Q) > E_{\rm rce}(R,Q)$, corresponding to low rates, we shift the convergence analysis to the union upper bound on the error probability $P_{\rm e}^{\rm ub}(\mathcal{C}_n)$ in (20), a bound that is tight enough at low rates. Then, we use De Caen's inequality [21], a bound proved to be tight for the random-coding ensemble average error probability in [22], to argue that such convergence also happens for the exact error probability $P_{\rm e}(\mathcal{C}_n)$. In particular, we set $\delta = 3\varepsilon$ in (11) and write

$$\mathbb{P}[|E_n(\mathcal{C}_n) - E_{\text{trc}}(R)| > 3\varepsilon] \le \alpha_n + \beta_n + \gamma_n, \tag{14}$$

where α_n accounts for the convergence of the exact error exponent to that of the union bound

$$\alpha_n = \mathbb{P}\Big[\big| E_n(\mathcal{C}_n) - E_n^{\mathrm{ub}}(\mathcal{C}_n) \big| > \varepsilon \Big],$$
(15)

5

$$\beta_n = \mathbb{P}\left[\left|E_n^{\mathrm{ub}}(\mathcal{C}_n) - \left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)\right]\right)\right| > \varepsilon\right],\tag{16}$$

and the last term deals with the convergence of such ensemble average to the TRC, namely

$$\gamma_n = \mathbb{P}\left[\left| \left(-\frac{1}{n} \mathbb{E} \left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) \right] \right) - E_{\mathrm{trc}}(R, Q) \right| > \varepsilon \right].$$
(17)

For the i.i.d. ensemble, we exploit the symbol-wise independence to show that the three terms α_n , β_n and γ_n vanish as $n \to \infty$ in (14), hence obtaining our result in (13). For constant-composition codes, we are also able to obtain a vanishing α_n , β_n and γ_n exploiting the independence of the joint type between two pairs of codewords and consider the expression of the limiting ensemble average of $E_n^{ub}(\mathcal{C}_n)$ for constant-composition codes reported in [9].

Theorem 1 shows the converge of sequences of random variables to the statistical mean. A refined analysis to that of Theorem 1 consists of studying, separately, the probability tails involved in the definition of convergence in probability in (11). The work in [12], addressed this issue for the constant-composition ensemble over DMCs. Specifically, [12] showed an interesting asymmetry: the probability $\mathbb{P}[E_n(\mathcal{C}_n) < E_{\text{trc}}(R,Q)]$ decays exponentially, while $\mathbb{P}[E_n(\mathcal{C}_n) > E_{\text{trc}}(R,Q)]$ decays double-exponentially. This implies that, beyond the concentration property, it is significantly more difficult to find a code in the ensemble with exponent higher than $E_{\text{trc}}(R,Q)$.

We next derive some results on the convergence rate of the error exponent $E_n(\mathcal{C}_n)$ to the typical random-coding exponent $E_{\text{trc}}(R,Q)$.

Theorem 2: For the i.i.d. or constant-composition ensembles with rate $0 \le R \le C$ and any $\varepsilon > 0$, it holds that

$$\mathbb{P}\big[E_n(\mathcal{C}_n) < E_{\text{trc}}(R,Q) - \varepsilon\big] \stackrel{.}{\leq} 2^{-n\varepsilon},\tag{18}$$

that is an exponential decay in the coding blocklength. In addition, for any $0 < R \le C$ and $\varepsilon > 0$, it holds that

$$\mathbb{P}\big[E_n(\mathcal{C}_n) > E_{\mathrm{trc}}(R,Q) + \varepsilon\big] = O\bigg(\frac{1}{\sqrt{n}}\bigg).$$
(19)

Proof: Sec. V-B.

Theorem 2, that strengthens Theorem 1, implicitly assumes that $E_{\text{trc}}^{\text{ub}}(R,Q) = E_{\text{trc}}(R,Q)$ for all rates below capacity to obtain (18) and uses the Berry-Esseen theorem [23] to obtain (19). For the union bound to the error probability,

$$P_{\rm e}^{\rm ub}(\boldsymbol{c}_n) = \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}[\boldsymbol{x}_i \to \boldsymbol{x}_j],$$
(20)

we are able to refine the upper tail in (19) as follows.

Theorem 3: For all rates satisfying $E_{\text{trc}}(R,Q) > E_{\text{rce}}(R,Q)$, there exists some $\epsilon > 0$ such that the following holds:

$$\mathbb{P}\left[E_n^{\rm ub}(\mathcal{C}_n) \ge E_{\rm ex}(R) + \epsilon\right] \le 2^{-2^{n\epsilon}},\tag{21}$$

where $E_{\text{ex}}(R)$ is the expurgated error exponent [24].

6

For strictly zero rate, that is R = 0, the expurgated is tight. Therefore, we have the following result.

Theorem 4: For the i.i.d. or constant-composition ensembles with rate R = 0 and any $\varepsilon > 0$, we have that

$$\mathbb{P}\left[E_n(\mathcal{C}_n) \ge E_{\text{trc}}(0, Q) + \epsilon\right] \le 2^{-2^{-1}}.$$
(22)

Proof: Sec. V-D.

So far, we have introduced results related to the convergence in probability of the error exponent for pairwiseindependent random codes. In the remaining of the section, we discuss the concentration in distribution of the error exponent $E_n(C_n)$ to its ensemble-average $\mathbb{E}[E_n(C_n)]$ as $n \to \infty$ at the low rate regime. While Theorem 5, valid for an exactly constant number of messages, states that the random-coding error exponent converges to a Gaussian-like distribution, we let the number of messages M_n in Theorem 6 to grow sub-exponentially with n, yet at a minimum rate $M_n \gg \sqrt{n}$, and show that it converges to a Gaussian. As in Theorem 6 and Theorem 13, most of the following results use the Stein's method [25].

Theorem 5: Let $M_n = M$ be a constant number of messages, fixed for every n, and let $U_{ij} \sim \mathcal{N}(0,1)$, for $i = 1, \ldots, M$ and $j = 1, \ldots, M$ such that $i \neq j$, be a set of independent standard normal random variables. Then, the error exponent for both i.i.d. and constant-composition random-coding ensembles satisfies

$$\frac{E_n(\mathcal{C}_n) - \mathbb{E}[E_n(\mathcal{C}_n)]}{\sqrt{\operatorname{Var}(E_n(\mathcal{C}_n))}} \xrightarrow{(d)} \frac{\min_{i \neq j} U_{ij} - \mathbb{E}[\min_{i \neq j} U_{ij}]}{\sqrt{\operatorname{Var}(\min_{i \neq j} U_{ij})}}.$$
(23)

Proof: Sec. V-E.

The proof of Theorem 5 is based on the fact that $E_n(\mathcal{C}_n)$ is a minimization of a constant number of M(M-1) terms where each term is a sum of independent random variables in the i.i.d. ensemble, and a sum of dependent random variables with an additional vanishing term in the constant-composition ensemble. Hence, the central limit theorm [20, Ch. VIII] and the Levy's continuity theorem [23] can be applied. In fact, it is easy to see that (23) holds when the exponent of the pairwise error probability of two different codewords X_i and X_j , that is $-\frac{1}{n} \log \mathbb{P}[X_i \rightarrow X_j]$ forms a set of independent random variables. However, even for the i.i.d. ensemble, such variables are only pairwise-independent. To give an example, for the binary symmetric channel with only three codewords X_1 , X_2 and X_3 , given the Hamming distance between two pairs, the Hamming distance between the third pair is not independent on the previous ones. Theorem 5 argues that, despite such dependence, this becomes negligible as $n \to \infty$ when the number of codewords is constant.

We illustrate in Fig. 1 the histogram of the error exponent $E_n(\mathcal{C}_n)$ used over a binary symmetric channel (BSC) with bit-flipping probability p = 0.11, equiprobable bits and M = 4 codewords for a blocklength of n = 10,000. The histograms are obtained for the i.i.d. and constant-composition ensembles using the Monte Carlo method after 10^7 trials. For the sake of comparison, we also depict the asymptotic distribution of the random variable $\min_{i \neq j} U_{ij}$ in the right-hand side of (23) (solid), and a normal approximation with the same mean and variance (dashed). We observe that the histogram matches the Gaussian-like distribution predicted by Theorem 5, with a slightly asymmetric tail tilting.

8




Fig. 1. Distribution of the error exponent of the (a) i.i.d. and (b) constant-composition codes over the BSC with M = 4, n = 10,000, symmetric input distribution and composition, and p = 0.11. Histograms of $E_n(C_n)$ with 10^7 trials, dashed black lines are normal distributions, and solid blue lines are the distributions of $\min_{i \neq j} U_{ij}$.

We observe that the cumulative distribution function of the random variable in the right-hand side of (23) can be easily obtained using that, for any sequence of random variables of length L > 1, for example X_1, \ldots, X_L , we have that

$$\mathbb{P}[\min\{X_1, X_2, \cdots, X_L\} \ge t] = \mathbb{P}[X_1 \ge t] \cdots \mathbb{P}[X_L \ge t].$$
(24)

If such sequence is i.i.d. standard-normally distributed, then

$$\mathbb{P}[\min\{X_1, X_2, \cdots, X_L\} \ge t] = Q(t)^L, \tag{25}$$

where $Q(t) = \frac{1}{\sqrt{2\pi}} \int_t^\infty e^{-x^2/2} dx$ is the Gaussian tail function. It is obvious that the right-hand side of (25) does *not* correspond to a Gaussian cumulative distribution function.

Theorem 6: Let M_n be a subexponential number of messages, namely $\lim_{n\to\infty} \frac{1}{n} \log M_n = 0$, satisfying the condition

$$\sum_{n=1}^{\infty} \frac{1}{M_n(M_n-1)} < \infty.$$

$$\tag{26}$$

Then, the error exponent of codes in the i.i.d. and constant-composition random-coding ensembles satisfies

$$\frac{E_n(\mathcal{C}_n) - \mathbb{E}[E_n(\mathcal{C}_n)]}{\sqrt{\operatorname{Var}(E_n(\mathcal{C}_n))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0, 1).$$
(27)

Proof: Sec. V-F.

The proof of Theorem 6 is based on the fact that, for both i.i.d. and constant-composition, the error exponent $E_n(\mathcal{C}_n)$ is the minimum of an infinite number of terms, where each term converges to a Gaussian distribution.

For a constant number of messages $M_n = M$, the condition in Theorem 6 is not satisfied, and therefore the error exponent does not concentrate according to (27) but to (23) instead. The fact that M_n does not grow with n implies that the dependence between the codewords vanishes as $n \to \infty$, and therefore the independence of U_{ij} is preserved. On the contrary, for a (sub-exponentially) growing number of messages M_n , the dependence among the codewords, and therefore the correlation among U_{ij} , increases such that the random variables U_{ij} can be represented by a common Gaussian random variable U. One example of sub-exponential growth of the number of messages satisfying (26) is the polynomial function of n given, for some $\delta > 0$, by $M_n = \Omega(n^{\frac{1+\delta}{2}})$.

IV. GENERAL CHANNELS

In this section, we introduce a number of results related to the concentration of the error probability and error exponent for relatively general channels and ensembles. The first result is a direct consequence of elementary probability results such as Chebyshev's inequality or Jensen's inequality.

Theorem 7: For a general channel and general ensemble such that $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \to 0$ for $0 \leq R < C$ and $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \to 1$ for R > C, we have

$$P_{\mathbf{e}}(\mathcal{C}_n) - \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)] \xrightarrow{(\mathbf{p})} 0.$$
(28)

Proof: Sec. V-G.

The above result holds for general channels and general ensembles as long as the strong converse property is satisfied. For channels and ensembles only satisfying a weak converse, namely that $\lim_{n\to\infty} \mathbb{E}[P_e(\mathcal{C}_n)] > \epsilon$, then (28) is valid if $\mathbb{E}[P_e(\mathcal{C}_n)^2]/\mathbb{E}[P_e(\mathcal{C}_n)]^2 \to 1$ as $n \to \infty$. In fact, such condition also guarantees the convergence in probability of the error exponent $E_n(\mathcal{C}_n)$ to the TRC, obtained by a direct application of Markov's inequality for general channels and pairwise-independent ensembles. Since the next results are very general and only assume pairwise-independent codewords, we drop the single-letter input distribution Q in the notation of $E_{\text{trc}}(R)$ and $E_{\text{rce}}(R)$ for the rest of the section.

Theorem 8: For a general channel and a pairwise-independent ensemble, under the condition that

$$\frac{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)^2]}{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)]^2} \to 1,$$
(29)

we have

$$E_n(\mathcal{C}_n) \xrightarrow{(\mathbf{p})} E_{\mathrm{trc}}(R).$$
 (30)

Proof: Sec. V-H.

As a remark, the condition (29) might hold for all the rate less than or equal to the critical rate, but not too small. This idea is made more precise in the following result, based on [15, Th. 1] and the Paley-Zygmund inequality [26], valid for general channels and pairwise-independent ensembles.

Theorem 9: For a general channel and pariwise-independent ensemble with rate such that $E_{\text{trc}}(R) > E_{\text{rce}}(R)$, we have

$$\frac{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2}{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2]} \to 0.$$
(31)

Proof: Sec. V-I.

In words, at low rates where the typical random-coding error exponent is strictly larger than the random-coding error exponent, the second-order moment of the error probability vanishes slower than the squared first-order moment. This implies that $\operatorname{Var}(P_{e}(\mathcal{C}_{n}))$ vanishes slower than the squared ensemble average $\mathbb{E}[P_{e}(\mathcal{C}_{n})]^{2}$, suggesting that the error probability cannot converge to a Gaussian distribution this rate regime. Such intuition is formalized in the next result, based on Theorem 9 and Slutsky's theorem [23].

Theorem 10: For any code ensemble and channel such that $E_{\rm trc}(R) > E_{\rm rce}(R)$, it holds that

$$\frac{P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0, 1).$$
(32)

Proof: Sec. V-J.

Theorem 11: For a general channel and pairwise-independent ensemble such that the normalized error probability converges in distribution to the standard normal distribution, that is

$$\frac{P_{\rm e}(\mathcal{C}_n) - \mathbb{E}[P_{\rm e}(\mathcal{C}_n)]}{\sqrt{\operatorname{Var}(P_{\rm e}(\mathcal{C}_n))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0, 1), \tag{33}$$

we have that

$$E_n(\mathcal{C}_n) \xrightarrow{(\mathbf{p})} E_{\mathrm{trc}}(R).$$
 (34)

Proof: Sec. V-K.

We remark that condition (33) is sufficient, but not necessary. To show (34), we bound the tail probabilities of the error probability $P_{e}(C_n)$ around $2^{-nE_{trc}(R)}$ and relate such bounds with the standard normal distribution.

In the remaining of the section, we state two auxiliary results related to the convergence in distribution of the union bound to the error probability of a code c_n in (20), and the convergence in distribution of an arbitrary function of the error probability.

By applying [25, Th. 3.6], a result for the sum of random variables with local dependence, to the union bound 20, we obtain the following result.

Theorem 12: Let Y_{12} and γ^2 be two parameters respectively given by $Y_{12} = \mathbb{P}[\{X_1 \to X_2\}] - \mathbb{E}[\mathbb{P}[\{X_1 \to X_2\}]]$ and $\gamma^2 = \operatorname{Var}(\mathbb{P}[\{X_1 \to X_2\}])$. For general channels and i.i.d. ensembles such that

$$\frac{M_n}{\gamma^3} \mathbb{E}[|Y_{12}|^3] \to 0 \tag{35}$$

$$\frac{M_n}{\gamma^4} \mathbb{E}[|Y_{12}|^4] \to 0 \tag{36}$$

as $n \to \infty$, we have that

$$\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n) - \mathbb{E}[P_{\rm e}^{\rm ub}(\mathcal{C}_n)]}{\sqrt{\operatorname{Var}(P_{\rm e}^{\rm ub}(\mathcal{C}_n))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0,1).$$
(37)

Proof: Sec. V-L.

Despite the result is about an upper bound on the error probability, Theorem 12 gives sufficient conditions for convergence in probability of the probability of error, while Theorem 10 gives a sufficient condition that prevents this to happen. This implies that for all codes and channels such that the two conditions (35), (36) hold, the condition $E_{\rm trc}(R,Q) > E_{\rm rce}(R,Q)$ cannot be satisfied.

To obtain a similar result for the exact error probability $P_e(C_n)$, that is a sum of all-dependent random variables, we note that a simple application of Hölder's inequality suggests that [25, Th. 3.6] is too loose. While other Stein method-based approaches such as exchangeable pairs could be applied, it is actually very challenging to find a partner for the error probability. In addition, the error exponent $E_n(C_n)$ is *not* even a sum of random variables.

In the last result, we develop a general condition for the convergence in distribution of a random variable sequence to the standard normal random variable based on Stein's method. Our method is based on a modification of the Wasserstein metric as in the proof of Theorem 6, and requires channels and decoding rules to satisfy certain conditions. It is open to find which specific channels and (random) codebook ensembles such that these conditions hold.

Theorem 13: Let $g_n : [0,1] \to \mathbb{R}$ be an arbitrary sequence of functions. For general channels and random codebook ensembles, under the condition that

$$\mathbb{E}\left[\left|\frac{g_n(P_{\mathbf{e}}(\mathcal{C}_n)) - \mathbb{E}[g_n(P_{\mathbf{e}}(\mathcal{C}_n))]}{\sqrt{\operatorname{Var}(g_n(P_{\mathbf{e}}(\mathcal{C}_n)))}}\right|\right] \to 0,\tag{38}$$

and

 $\mathbb{E}\left[\left|\left(\frac{g_n(P_{\mathbf{e}}(\mathcal{C}_n)) - \mathbb{E}[g_n(P_{\mathbf{e}}(\mathcal{C}_n))]}{\sqrt{\operatorname{Var}(g_n(P_{\mathbf{e}}(\mathcal{C}_n)))}}\right)^2 - 1\right|\right] \to 0,\tag{39}$

the following holds:

$$\frac{g_n(P_{\mathbf{e}}(\mathcal{C}_n)) - \mathbb{E}[g_n(P_{\mathbf{e}}(\mathcal{C}_n))]}{\sqrt{\operatorname{Var}(g_n(P_{\mathbf{e}}(\mathcal{C}_n)))}} \xrightarrow{(d)} \mathcal{N}(0, 1).$$
(40)

Proof: Sec. V-M.

Summarizing, in Sec. III and Sec. IV we discussed a number of convergence results of the error probability $P_e(C_n)$ and the error exponent $E_n(C_n)$, important performance metrics of random codes used in the study of reliable communication. In the next section, we present the proofs for such results.

V. PROOFS OF THEOREMS

We begin by introducing some definitions used in the Stein's method [25]. We also introduce a novel, modified Wasserstein metric that is used throughout the section.

Definition 1: Define

$$\mathcal{V} = \left\{ h : \mathbb{R} \to \mathbb{R}_{+} : h(u) = \begin{cases} c, & x \le a \\ 0, & x \ge a+c \\ \text{linear between } a \text{ and } a+c, & \text{otherwise} \end{cases} \right\}.$$
(41)

Definition 2 (Probability metrics): For two random variables *X* and *Y*, the probability metrics have the following form:

$$d_{\mathcal{H}}(X,Y) = \sup_{h \in \mathcal{H}} \left| \mathbb{E}[h(X)] - \mathbb{E}[h(Y)] \right|,\tag{42}$$

$$\bar{d}_{\mathcal{H}}(X,Y) = \sup_{h \in \mathcal{H}} \min\left\{ \left| \mathbb{E}[h(X)] - \mathbb{E}[h(Y)] \right|, \left| \mathbb{E}[h(-X)] - \mathbb{E}[h(Y)] \right| \right\},\tag{43}$$

where \mathcal{H} is some family of "test" functions on \mathbb{R} .

We now details examples of metrics of this form along with some useful properties and relations:

- By taking H = {1{· ≤ u} u ∈ ℝ} in (43) and the probability metric d_H(X, Y), we obtain the Kolmogorov metric, which denote by d_K. By definition, the convergence in the Kolmogorov metric means the convergence in distribution.
- By taking H = {h : ℝ → ℝ : |h(u) h(v)| ≤ |u v|} and the probability metric d_H(X, Y), we obtain the Wasserstein metric, which we denote d_W.
- By taking H = {h ∈ V : c ≤ 4√2π} and the probability metric d_H(X, Y), we obtain a slightly modified Wasserstein metric d̃_{W,mod}.
- By taking H = {h ∈ V : c ≤ 4√2π} and the probability metric d
 H(X, Y), we obtain a modified Wasserstein metric¹, which we denote d{W,mod}.

The following auxiliary lemma is also very important in deriving the convergence in distributions in most of lemmas and theorems in this paper, whose proof can be found in the Appendix A.

Lemma 1: Assume that $U_n \xrightarrow{\text{(d)}} U$ as $n \to \infty$ for some random variable U, and $\mathbb{E}[U_n] = 0$ and $\operatorname{Var}(U_n) = 1$. Then, it holds that

$$\mathbb{E}[U] = 0, \qquad \operatorname{Var}(U) = 1. \tag{44}$$

Proof: Appendix A.

A. Proof of Theorem 1

We start by stating De Caen's inequality.

Lemma 2 (De Caen [27]): Let $\{A_i\}_{i \in \mathcal{I}}$ be finite family of events in a probability space (Ω, \mathbb{P}) . Then²

$$\mathbb{P}\left(\bigcup_{i\in\mathcal{I}}A_i\right)\geq\sum_{i\in\mathcal{I}}\frac{\mathbb{P}^2(A_i)}{\sum_{j\in\mathcal{I}}\mathbb{P}[A_i\cap A_j]}$$
(45)

Lemma 3: Under the condition that $E_{trc}(R,Q) = E_{rce}(R,Q)$ and for $\lambda > 0$, the following holds for i.i.d. and constant-composition ensembles.

$$\lim_{n \to \infty} \mathbb{E} \left[P_{\mathbf{e}}(\mathcal{C}_n, Q)^{\frac{\lambda}{n}} \right] = 2^{-\lambda E_{\mathrm{trc}}(R, Q)}.$$
(46)

Proof: Appendix B.

1) Proof of Theorem 1 for $E_{trc}(R,Q) = E_{rce}(R,Q)$: Let

$$\varphi(\lambda) = 2^{-\lambda E_{\rm trc}(R,Q)} \tag{47}$$

¹This definition of Wasserstein metric is a variant of the definition in [25], where we constraint the set \mathcal{H} to achieve a tighter bound. ²We make the convention $\frac{0}{0} = 0$, so that events of probability zero are not counted in (45). for all $\lambda > 0$, which is the Laplace transform of the constant random variable $-E_{\text{trc}}(R,Q)$. Let $\varphi_n(\lambda)$ be the Laplace transform of the distribution of $\frac{1}{n} \log P_e(\mathcal{C}_n, Q)$. Then, we have

$$\varphi_n(\lambda) = \mathbb{E}\left[2^{\lambda \frac{1}{n} \log P_{\mathrm{e}}(\mathcal{C}_n, Q)}\right]$$
(48)

$$= \mathbb{E}\Big[P_{\mathrm{e}}(\mathcal{C}_n, Q)^{\frac{\lambda}{n}}\Big].$$
(49)

Then, by Lemma 3, the following holds:

$$\lim_{n \to \infty} \varphi_n(\lambda) = 2^{-\lambda E_{\rm trc}(R,Q)}.$$
(50)

Applying the Levy's continuity theorem [20, Sec. XIII.1], we obtain from (50) that

$$-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n, Q) \xrightarrow{\rm (d)} E_{\rm trc}(R, Q).$$
(51)

However, we know that the convergence in distribution to a constant implies convergence in probability, i.e.

$$-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n, Q) \xrightarrow{({\rm p})} E_{\rm trc}(R, Q).$$
(52)

2) Proof of Theorem 1 for $E_{\text{trc}}(R,Q) > E_{\text{rce}}(R,Q)$: This section is devoted to the proof (13) for the range of rates for which $E_{\text{trc}}(R,Q) > E_{\text{rce}}(R,Q)$. We first need some definitions and lemmas. For this range of rates, the proof uses the union bound to the error probability (1) and shows that it gives a good estimate of the probability of error. The union bound is given by,

$$P_{\rm e}(c_n) \le P_{\rm e}^{\rm ub}(c_n),\tag{53}$$

where $P_{e}^{ub}(c_n)$ is defined in (20), and we define its finite-length error exponent as

$$E_n^{\rm ub}(c_n) = -\frac{1}{n} \log P_{\rm e}^{\rm ub}(c_n).$$
(54)

We denote by $E_{\text{trc}}(R,Q)$ and $E_{\text{rce}}(R,Q)$ respectively the typical error and the random coding error exponents for the fixed underlying distribution Q, and we define

$$d_{\rm B}(x,x') = -\log\left(\sum_{y}\sqrt{W(y|x)W(y|x')}\right)$$
(55)

to be the Bhattacharyya distance between symbols $x, x' \in \mathcal{X}$.

We assume that the DMC is such that

$$0 < D_{\rm b} = \max_{x,x'} d_{\rm B}(x,x') < \infty,$$
(56)

that is, we leave the cases where W(y|x)W(y|x') = 0 for for some x and x' and all y beyond the scope of the paper. This case would correspond to a positive zero-error capacity, where some symbols cannot be confused at the decoder.

First, we introduce some auxiliary results about the exponential decay f the pairwise error probability between two codewords, using the method of types. We let $\mathcal{P}_n(\mathcal{X} \times \mathcal{X})$ be the set of all joint types on $\mathcal{X} \times \mathcal{X}$, and $\mathcal{P}(\mathcal{X} \times \mathcal{X})$ be the set of all possible probability distributions on $\mathcal{X} \times \mathcal{X}$.

Lemma 4: For $R < R_{crit}$, the pairwise codeword error probability between two codewords x_i, x_j given their joint type $P_{XX'}$ satisfies

$$\mathbb{P}[\boldsymbol{x}_{i} \to \boldsymbol{x}_{j} | P_{XX'}] = g_{n}(P_{XX'}) \doteq 2^{-n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')}$$
(57)

for some function $g_n : \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) \to [0, 1]$.

Proof: Appendix C

Lemma 5: Given an i.i.d. random codebook ensemble. For each $P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})$, let $\mathcal{N}(P_{XX'})$ be the number of codeword pairs in a specific code such that their joint type is $P_{XX'}$. Let $Q_X = Q'_X = Q$. Define

$$\mathcal{V}_n = \left\{ \mathcal{N}(P_{XX'}) = 0, \quad \forall P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q_X') > 2R \right\}$$
(58)

which is the event that the (random) number of pairs $(i, j) \in [M_n] \times [M_n]$ such that $i \neq j$ and $(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})$ is equal to zero for each *n*-joint type $P_{XX'}$ with $D(P_{XX'} || Q_X Q'_X) > 2R$. Then, we have

$$\mathbb{P}\big[\mathcal{V}_n^c\big] \le 2^{-n\alpha(R)} \tag{59}$$

for some $\alpha(R) > 0$ for all $R \ge 0$.

Proof: Appendix D

Lemma 6: Assume that R > 0. Take an arbitrary $\nu \ge 0$ such that $\nu \le 2R$. Let $Q_X = Q_{X'} = Q$ and

$$\mathcal{P} = \left\{ P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) \le 2R - \nu \right\}$$
(60)

and

$$D_n = \frac{1}{M} \sum_{P_{XX'} \in \mathcal{P}} \mathcal{N}(P_{XX'}) g_n(P_{XX'}), \tag{61}$$

$$\bar{E}_{\rm trc}(\nu, R, Q) = \begin{cases} R + \sum_{x,x'} d_{\rm B}(x, x') P_{XX'}^*(x, x'), & \text{if} \quad D(P_{XX'}^* \| Q_X Q_X') = 2R - \nu, \\ E_{\rm rce}(R, Q), & \text{otherwise} \end{cases}$$
(62)

where $P_{XX'}^*$ is an optimizer of $\min_{P_{XX'} \in \mathcal{P}} D(P_{XX'} || Q_X Q_{X'}) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') - R$. Then, the following holds:

$$\mathbb{E}[D_n] \doteq 2^{-n\bar{E}_{\rm trc}(\nu,R)} \tag{63}$$

and

$$\frac{\operatorname{Var}(D_n)}{\left(\mathbb{E}[D_n]\right)^2} \stackrel{.}{\leq} 2^{-n\zeta(\nu,R)}$$
(64)

for some constant $\zeta(\nu, R)$ such that $\zeta(\nu, R) > 0$ if $0 < \nu \leq 2R$.

Proof: Appendix E

Lemma 7: Let

$$E_{\rm trc}^{\rm ub}(R,Q) = \lim_{n \to \infty} -\frac{1}{n} \mathbb{E}[\log P_{\rm e}^{\rm ub}(\mathcal{C}_n)].$$
(65)

Then, for $0 < R < R_{crit}(Q)$, the following holds:

$$E_{\rm trc}^{\rm ub}(R,Q) = \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q_X') \le 2R} D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} d_{\rm B}(x,x') P_{XX'}(x,x') - R \tag{66}$$

$$=\bar{E}_{\rm trc}(0,R,Q),\tag{67}$$

where \bar{E}_{trc} is defined in (62), Lemma 6.

Proof: Appendix F

Lemma 8: For the range of rates $0 \le R < R_{crit}(Q)$, any $\varepsilon > 0$ and for some $\kappa > 0$, it holds that

$$\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) > \frac{1}{2}2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)-\varepsilon)}\right] + \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+\varepsilon)}\right] \le \frac{1}{n^{1+\kappa}}.$$
(68)

Proof: Appendix G

Lemma 9: For all rate R such that $0 < R < R_{crit}(Q)$ and for some $\delta(R) > 0$, it holds that

$$0 \leq \frac{\mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})]}{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]} - 1 \leq 2^{-n\left(\delta(R) + E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) - E_{\mathrm{rce}}(R,Q)\right)}.$$
(69)

Proof: Appendix H

We are now equipped to prove Theorem 1 by observing that for any $\varepsilon > 0$, the convergence in probability of $E_n(\mathcal{C}_n)$ to $E_{\text{trc}}(R,Q)$ can be written and upper bounded as

$$\mathbb{P}\left[|E_{n}(\mathcal{C}_{n}) - E_{\mathrm{trc}}(R)| > 3\varepsilon\right] \leq \underbrace{\mathbb{P}\left[\left|E_{n}(\mathcal{C}_{n}) - E_{n}^{\mathrm{ub}}(\mathcal{C}_{n})\right| > \varepsilon\right]}_{\alpha_{n}} + \underbrace{\mathbb{P}\left[\left|E_{n}^{\mathrm{ub}}(\mathcal{C}_{n}) - \left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]\right)\right| > \varepsilon\right]}_{\beta_{n}} + \underbrace{\mathbb{P}\left[\left|\left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]\right) - E_{\mathrm{trc}}(R,Q)\right| > \varepsilon\right]}_{\gamma_{\mathrm{rc}}}.$$
(70)

We next show that the terms α_n , β_n and γ_n in (70) tend to zero as $n \to \infty$, implying the concentration result in (13).

3) First term of (70): The term α_n quantifies the deviation of the error exponent of the error probability (2) with that of the union bound (20). By the symmetry of the pairwise-independent i.i.d. random-coding ensemble, for any pair of codewords X_i and X_j with $i \neq j$ we have that

$$\mathbb{E}\big[\mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j]\big] = \mathbb{E}\big[\mathbb{P}[\boldsymbol{X}_1 \to \boldsymbol{X}_2]\big].$$
(71)

Similarly, for any triplet of codewords X_i , X_j and X_k with $j, k \neq i$ and $j \neq k$, it holds that

$$\mathbb{E}\Big[\mathbb{P}\big[\{\boldsymbol{X}_i \to \boldsymbol{X}_j\} \cap \{\boldsymbol{X}_i \to \boldsymbol{X}_k\}\big]\Big] = \mathbb{E}\Big[\mathbb{P}\big[\{\boldsymbol{X}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{X}_1 \to \boldsymbol{X}_3\}\big]\Big]$$
(72)

where in both (71) and (72), the expectations are calculated with respect to the i.i.d. ensemble codeword distribution $Q^n(\mathbf{x}) = \prod_{k=1}^n Q(x_k)$, where Q(x) is the single-letter input distribution. We next provide separate convergence of α_n for R = 0 and for $0 < R < R_{\text{crit}}(Q)$. For the case of R = 0, we first observe that the union bound (20) can be bounded from above as

$$P_{\rm e}^{\rm ub}(\mathcal{C}_n) = \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}[\boldsymbol{x}_i \to \boldsymbol{x}_j]$$
(73)

$$\leq (M_n - 1) \max_{i \neq j} \mathbb{P}[\boldsymbol{x}_i \to \boldsymbol{x}_j],$$
(74)

while the probability of error (1) can be lower bounded by

$$P_{\rm e}(\mathcal{C}_n) = \frac{1}{M_n} \sum_{i=1}^{M_n} \mathbb{P}\left[\bigcup_{j \neq i} \{\boldsymbol{x}_i \to \boldsymbol{x}_j\}\right]$$
(75)

$$\geq \frac{1}{M_n} \max_{i \neq j} \mathbb{P}[\boldsymbol{x}_i \to \boldsymbol{x}_j].$$
(76)

From (74) and (76), we have that the first term in the r.h.s. of (70) satisfies

$$\alpha_n = \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) > 2^{n\varepsilon} P_{\mathrm{e}}(\mathcal{C}_n)\right]$$
(77)

$$\leq \mathbb{P}\left[(M_n - 1) \max_{i \neq j} \mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j] > 2^{n\varepsilon} \frac{1}{M_n} \max_{i \neq j} \mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j] \right]$$
(78)

$$= \mathbb{P}\left[(M_n - 1) > 2^{n\varepsilon} \frac{1}{M_n} \right].$$
(79)

Since M_n is any sub-exponential sequence in n, the probability in (79) vanishes as $n \to \infty$ for $\varepsilon > 0$.

We now consider the case of $0 < R < R_{crit}(Q)$. We define the sequence a_n as

 \leq

$$a_n = 2^{-n(E_{\rm trc}^{\rm ub}(R,Q) + \frac{\varepsilon}{2})}.$$
(80)

Then, we have

$$\mathbb{P}\left[\left|E_{n}(\mathcal{C}_{n})-E_{n}^{\mathrm{ub}}(\mathcal{C}_{n})\right|>\varepsilon\right]=\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})>2^{\varepsilon n}P_{\mathrm{e}}(\mathcal{C}_{n})\right]$$
(81)

$$= \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - a_{n} - 2^{\varepsilon n} \left(P_{\mathrm{e}}(\mathcal{C}_{n}) - a_{n}\right) > (2^{\varepsilon n} - 1)a_{n}\right]$$
(82)

$$\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - a_{n} > \frac{1}{2}(2^{\varepsilon n} - 1)a_{n}\right] + \mathbb{P}\left[-2^{\varepsilon n}\left(P_{\mathrm{e}}(\mathcal{C}_{n}) - a_{n}\right) > \frac{1}{2}(2^{\varepsilon n} - 1)a_{n}\right],$$
(83)

where (83) follows from the fact that $\mathbb{P}[A+B>2C] = \mathbb{P}[\{A>C\} \cup \{B>C\}] \le \mathbb{P}[A>C] + \mathbb{P}[B>C].$

Now, observe that

$$\mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) - a_{n} > \frac{1}{2}(2^{\varepsilon n} - 1)a_{n}\right] = \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) > \frac{1}{2}(2^{\varepsilon n} + 1)a_{n}\right]$$
(84)

$$= \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) > \frac{1}{2}(2^{\varepsilon n} + 1)2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) + \varepsilon/2)}\right]$$
(85)

$$\leq \mathbb{P}\bigg[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) > \frac{1}{2} 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) - \varepsilon/2)}\bigg].$$
(86)

On the other hand, we also have

$$\mathbb{P}\left[-2^{\varepsilon n}\left(P_{\mathrm{e}}(\mathcal{C}_{n})-a_{n}\right) > \frac{1}{2}(2^{\varepsilon n}-1)a_{n}\right] \\
= \mathbb{P}\left[2^{\varepsilon n}\left(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})-P_{\mathrm{e}}(\mathcal{C}_{n})\right) - 2^{\varepsilon n}\left(P_{\mathrm{e}}(\mathcal{C}_{n})-a_{n}\right) > \frac{1}{2}(2^{\varepsilon n}-1)a_{n}\right] \\
\leq \mathbb{P}\left[2^{\varepsilon n}\left(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})-P_{\mathrm{e}}(\mathcal{C}_{n})\right) > \frac{1}{4}(2^{\varepsilon n}-1)a_{n}\right] \tag{87}$$

$$+ \mathbb{P}\bigg[-2^{\varepsilon n}\bigg(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - a_{n}\bigg) > \frac{1}{4}(2^{\varepsilon n} - 1)a_{n}\bigg].$$

$$(88)$$

Now, we know that

$$\mathbb{P}\left[-2^{\varepsilon n}\left(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})-a_{n}\right)>\frac{1}{4}(2^{\varepsilon n}-1)a_{n}\right]=\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})<\left(1-\frac{1}{4}\left(\frac{2^{\varepsilon n}-1}{2^{\varepsilon n}}\right)\right)a_{n}\right]$$

$$=\left[-\mathrm{ub}\left(a_{n}\right)-\left(1-\frac{1}{4}\left(\frac{2^{\varepsilon n}-1}{2^{\varepsilon n}}\right)\right)-\mathrm{e}\left(E^{\mathrm{ub}}(\mathcal{D}_{n})+e^{2}(n)\right)\right]$$
(89)

$$= \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < \left(1 - \frac{1}{4}\left(\frac{2^{\varepsilon n} - 1}{2^{\varepsilon n}}\right)\right)2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) + \varepsilon/2)}\right] \quad (90)$$

$$\leq \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) + \varepsilon/2)}\right].$$
(91)

In addition, we also have

$$\mathbb{P}\left[2^{\varepsilon n}(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - P_{\mathrm{e}}(\mathcal{C}_{n})) > \frac{1}{4}(2^{\varepsilon n} - 1)a_{n}\right] \stackrel{\cdot}{\leq} a_{n}^{-1}\mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - P_{\mathrm{e}}(\mathcal{C}_{n})]$$
(92)

$$=2^{(E_{\rm trc}^{\rm ub}(R,Q)+\varepsilon/2)n}\mathbb{E}[P_{\rm e}^{\rm ub}(\mathcal{C}_n)-P_{\rm e}(\mathcal{C}_n)],\tag{93}$$

where (92) follows from $P_{\rm e}^{\rm ub}(\mathcal{C}_n) \geq P_{\rm e}(\mathcal{C}_n)$ and Markov's inequality, and (93) follows from (80).

Now, for R>0 and $E_{\rm trc}^{\rm ub}(R,Q)>E_{\rm rce}(R,Q),$ from Lemma 9, we have

$$\mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n}) - P_{e}(\mathcal{C}_{n})] = \mathbb{E}[P_{e}(\mathcal{C}_{n})] \left(\frac{\mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n})]}{\mathbb{E}[P_{e}(\mathcal{C}_{n})]} - 1\right)$$
(94)

$$\leq 2^{-nE_{\rm rce}(R,Q)} \left(2^{-n\left(\delta(R) + E_{\rm trc}^{\rm ub}(R,Q) - E_{\rm rce}(R,Q)\right)} \right).$$
(95)

From (93) and (95), we obtain

$$\mathbb{P}\left[2^{\varepsilon n}(P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) - P_{\mathrm{e}}(\mathcal{C}_{n})) > \frac{1}{4}(2^{\varepsilon n} - 1)a_{n}\right] \stackrel{\cdot}{\leq} 2^{\left(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) + \varepsilon/2\right)n} 2^{-nE_{\mathrm{rce}}(R,Q)} \left[2^{-n\left(\delta(R) + E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) - E_{\mathrm{rce}}(R,Q)\right)}\right]$$

$$\tag{96}$$

$$\leq 2^{-n\left(\delta(R) - \varepsilon/2\right)}.$$
(97)

Hence, from (88), (91), and (97), we have

$$\mathbb{P}\bigg[-2^{\varepsilon n}\big(P_{\mathbf{e}}(\mathcal{C}_{n})-a_{n}\big) > \frac{1}{2}(2^{\varepsilon n}-1)a_{n}\bigg] \le \mathbb{P}\bigg[P_{\mathbf{e}}^{\mathbf{ub}}(\mathcal{C}_{n}) < 2^{-N(E_{\mathrm{tre}}^{\mathbf{ub}}(R,Q)+\varepsilon/2)}\bigg] + 2^{-n\big(\delta(R)-\varepsilon/2\big)}.$$
(98)

From (83), (86), and (98), we have

$$\mathbb{P}\left[\left|-\frac{1}{n}\log P_{e}(\mathcal{C}_{n})-\frac{-\log P_{e}^{ub}(\mathcal{C}_{n})}{n}\right| > \varepsilon\right] \\
\leq \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n})>\frac{1}{2}2^{-n(E_{trc}^{ub}(R,Q)-\varepsilon/2)}\right] + \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n})<2^{-n(E_{trc}^{ub}(R,Q)+\varepsilon/2)}\right] + 2^{-n\left(\delta(R)-\varepsilon/2\right)} \tag{99}$$

$$\leq \frac{1}{n^{1+\beta}} + 2^{-n\left(\delta(R) - \varepsilon/2\right)} \tag{100}$$

$$\rightarrow 0,$$
 (101)

for any $0 < \varepsilon < 2\delta(R)$, where (100) follows from Lemma 8 with β being a positive constant. Since $\mathbb{P}[|E_n(\mathcal{C}_n) - E_n^{ub}(\mathcal{C}_n)| > \varepsilon]$ is a non-increasing function in ε , (101) must hold for all $\varepsilon > 0$.

Furthermore, since $\mathbb{P}[|E_n(\mathcal{C}_n) - E_n^{ub}(\mathcal{C}_n)| > \varepsilon]$ is a non-increasing function in ε , for any $\varepsilon > 0$, there exists an $\varepsilon_0 \in (0, 2\delta(R))$ such that

$$\mathbb{P}\left[\left|E_{n}(\mathcal{C}_{n})-E_{n}^{\mathrm{ub}}(\mathcal{C}_{n})\right|>\varepsilon\right]\leq\frac{1}{n^{1+\beta}}+2^{-n\left(\delta(R)-\varepsilon_{0}/2\right)}$$
(102)

for some $\varepsilon_0 \in (0, 2\delta(R))$. It follows from (102) that

$$\sum_{n=1}^{\infty} \mathbb{P}\left[\left|E_n(\mathcal{C}_n) - E_n^{\mathrm{ub}}(\mathcal{C}_n)\right| > \varepsilon\right] < \infty.$$
(103)

Hence, by Borel-Cantelli's lemma [23], we have

$$E_n(\mathcal{C}_n) - E_n^{\mathrm{ub}}(\mathcal{C}_n) \xrightarrow{(\mathrm{a.s.})} 0,$$
 (104)

where $\xrightarrow{(a.s.)}$ denotes almost sure convergence as $n \to \infty$, that is, a sequence of random variables $\{A_n\}_{n=1}^{\infty}$ converge almost surely to A if

$$\mathbb{P}\left[\lim_{n \to \infty} A_n = A\right] = 1.$$
(105)

On the other hand, observe that

$$\left|E_{n}(\mathcal{C}_{n}) - E_{n}^{\mathrm{ub}}(\mathcal{C}_{n})\right| \leq -\frac{2\log P_{\mathrm{e}}(\mathcal{C}_{n})}{n}$$
(106)

$$\leq 2E_{\rm sp}(R),\tag{107}$$

where (107) follows from the fact that the error exponent of any sufficiently long code is upper bounded by the sphere-packing bound.

Hence, from (104) and (107), by the bounded convergence theorem, it holds that

$$\lim_{n \to \infty} \mathbb{E} \left[E_n(\mathcal{C}_n) - E_n^{\mathrm{ub}}(\mathcal{C}_n) \right] = 0.$$
(108)

This means that

$$E_{\rm trc}(R,Q) = \lim_{n \to \infty} E_n(\mathcal{C}_n) \tag{109}$$

$$=\lim_{n\to\infty} E_n^{\rm ub}(\mathcal{C}_n) \tag{110}$$

$$=E_{\rm trc}^{\rm ub}(R,Q).$$
(111)

4) Second term of (70): Using Chebyshev's inequality, we have

$$\mathbb{P}\left[\left|E_{n}^{\mathrm{ub}}(\mathcal{C}_{n}) - \left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]\right)\right| > \varepsilon\right] \\
\leq \frac{1}{\varepsilon^{2}}\operatorname{Var}\left(-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n}\right)$$
(112)

$$= \frac{1}{n^2 \varepsilon^2} \operatorname{Var} \left(-\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) \right)$$
(113)

$$=\frac{1}{n^2\varepsilon^2}\mathbb{E}\left[\left(-\log(M_n-1)-\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{M_n-1}\right)\right)^2\right]-\frac{1}{\varepsilon^2}\left(\frac{\mathbb{E}\left[-\log P_{\rm e}^{\rm ub}(\mathcal{C}_n)\right]}{n}\right)^2.$$
(114)

Now, define

$$\xi(p, n, R) = 2^{-n(E_{\rm trc}^{\rm ub}(R, Q) + R)}.$$
(115)

From (114), we obtain

$$\mathbb{P}\left[\left|E_{n}^{\mathrm{ub}}(\mathcal{C}_{n})-\left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]\right)\right| > \varepsilon\right] \\
\leq \frac{1}{n^{2}\varepsilon^{2}}\mathbb{E}\left[\left(-\log(M_{n}-1)-\log\xi(p,n,R)-\log\left(\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right)^{2}\right] \\
-\frac{1}{\varepsilon^{2}}\left(\frac{\mathbb{E}\left[-\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]}{n}\right)^{2}.$$
(116)

By Lemma 7, we know that

$$\lim_{n \to \infty} \frac{\mathbb{E}\left[-\log P_{\rm e}^{\rm ub}(\mathcal{C}_n)\right]}{n} = E_{\rm trc}^{\rm ub}(R,Q),\tag{117}$$

hence, it holds that

$$\limsup_{n \to \infty} \mathbb{P}\left[\left|E_{n}^{\mathrm{ub}}(\mathcal{C}_{n}) - \left(-\frac{1}{n}\mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right]\right)\right| > \varepsilon\right] \\
= \limsup_{n \to \infty} \frac{1}{\varepsilon^{2}}\mathbb{E}\left[\left(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) - \frac{1}{n}\log\left(\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right)^{2}\right] - \frac{\left(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)\right)^{2}}{\varepsilon^{2}} \quad (118) \\
\leq \frac{1}{\varepsilon^{2}}\left(\left(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)\right)^{2} - 2E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)\liminf_{n \to \infty} \mathbb{E}\left[\frac{1}{n}\log\left(\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right] \\
+ \limsup_{n \to \infty} \mathbb{E}\left[\left(\frac{1}{n}\log\left(\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right)^{2}\right]\right) - \frac{\left(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)\right)^{2}}{\varepsilon^{2}}, \quad (119)$$

where (119) follows from the sub-additivity of lim sup. Now, we need to estimate

$$\liminf_{n \to \infty} \mathbb{E}\left[\frac{1}{n} \log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n - 1)\xi(p, n, R)}\right)\right]$$

and

$$\limsup_{n \to \infty} \mathbb{E}\left[\left(\frac{1}{n} \log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n - 1)\xi(p, n, R)}\right)\right)^2\right].$$

First, we show that

$$\frac{1}{n}\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M-1)\xi(p,n,R)}\right) \xrightarrow{(\text{a.s.})} 0 \tag{120}$$

$$\mathbb{P}\left[\left|\frac{1}{n}\log\left(\frac{P_{e}^{ub}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right| > \nu\right] \\
= \mathbb{P}\left[\left|\log\left(\frac{P_{e}^{ub}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right| > n\nu\right]$$
(121)

$$= \mathbb{P}\left[\left|-\log\xi(p,n,R) + \log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{M_n - 1}\right)\right| > n\nu\right]$$
(122)

$$= \mathbb{P}\left[\left|n(E_{\rm trc}^{\rm ub}(R,Q) + R) + \log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{M_n - 1}\right)\right| > n\nu\right]$$
(123)

$$= \mathbb{P}\left[\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{M_{n}-1} > 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+R-\nu)}\right] + \mathbb{P}\left[\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{M_{n}-1} < 2^{-N(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+R+\nu)}\right]$$
(124)

$$\leq \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) > \frac{1}{2}2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)-\nu)}\right] + \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+\nu)}\right]$$
(125)

$$\leq \frac{1}{n^{1+\beta}},\tag{126}$$

for some constants $\beta > 0$, where (126) follows from Lemma 8.

From (126), we obtain

$$\sum_{n=1}^{\infty} \mathbb{P}\left[\left|\frac{1}{n}\log\left(\frac{P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)}\right)\right| > \nu\right] < \sum_{n=1}^{\infty} \frac{1}{n^{1+\beta}} < \infty$$
(127)

by using D'Alembert criterion.

This means that (120) holds, or

$$\frac{1}{n} \log \left(\frac{P_{e}^{ub}(\mathcal{C}_{n})}{(M_{n}-1)\xi(p,n,R)} \right) \xrightarrow{(a.s.)} 0$$
(128)

by Borel-Cantelli lemma [23].

Now, since $0 \leq \mathbb{P}(X_i \to X_j) \leq 1$ for all $i, j \in [M] : i \neq j$, it holds that

$$\frac{1}{n}\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n-1)\xi(p,n,R)}\right) = \frac{1}{n}\log\left(\frac{1}{M_n(M_n-1)\xi(p,n,R)}\sum_{i\neq j}\mathbb{P}(\boldsymbol{X}_i\to\boldsymbol{X}_j)\right)$$
(129)

$$\leq \frac{1}{n} \log\left(\frac{1}{\xi(p,n,R)}\right) \tag{130}$$

$$\leq E_{\rm trc}^{\rm ub}(R,Q) + R,\tag{131}$$

where (131) follows from (115). On the other hand, from the sphere-packing bound ³, it holds almost surely that

$$\frac{1}{n}\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n-1)\xi(p,n,R)}\right) \ge \frac{1}{n}\log\left(\frac{P_{\rm e}(\mathcal{C}_n)}{(M_n-1)\xi(p,n,R)}\right)$$
(132)

$$\stackrel{\cdot}{\geq} \frac{1}{n} \log \left(\frac{2^{-nE_{\rm sp}(R)}}{(M_n - 1)\xi(p, n, R)} \right)$$

$$(133)$$

$$= -E_{\rm sp}(R) - R + \left(E_{\rm trc}^{\rm ub}(R,Q) + R\right)$$
(134)

$$=E_{\rm trc}^{\rm ub}(R,Q) - E_{\rm sp}(R), \tag{135}$$

³In case that the sphere packing bound diverges, we can use $E_{ex}(R=0)$ as an upper bound, which is finite at R=0 unless the zero error capacity $C_0 > 0$.

where (133) follows from the sphere-packing bound [28], and (134) follows from (115) and $M_n \doteq 2^{nR}$.

From (131) and (135), $\frac{1}{n} \log \left(\frac{P_e^{\text{ub}}(\mathcal{C}_n)}{(M_n - 1)\xi(p, n, R)} \right)$ is bounded (both below and above).

Hence, by the bounded convergence theorem and the continuous mapping theorem [23], it holds that

$$\mathbb{E}\left[\frac{1}{n}\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n-1)\xi(p,n,R)}\right)\right] \to 0,\tag{136}$$

$$\mathbb{E}\left[\left(\frac{1}{n}\log\left(\frac{P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{(M_n-1)\xi(p,n,R)}\right)\right)^2\right] \to 0.$$
(137)

From (119), (136), and (137), we finally have

$$\lim_{n \to \infty} \mathbb{P}\left[\left| E_n^{\mathrm{ub}}(\mathcal{C}_n) - \left(-\frac{1}{n} \mathbb{E}\left[\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) \right] \right) \right| > \varepsilon \right] = 0$$
(138)

for any arbitrary $\varepsilon > 0$. This leads to

$$\lim_{n \to \infty} \mathbb{P}\left[\left| E_n^{\rm ub}(\mathcal{C}_n) - \left(-\frac{1}{n} \mathbb{E} \left[\log P_{\rm e}^{\rm ub}(\mathcal{C}_n) \right] \right) \right| > \varepsilon \right] = 0$$
(139)

by the fact that the probability measure is bounded from below by zero.

5) Third term of (70): By Lemma 7, it is known that

$$\mathbb{E}\left[\frac{-\log P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{n}\right] \to E_{\rm trc}^{\rm ub}(R,Q).$$
(140)

On the other hand, from (111) in Step 1, we know that

$$E_{\rm trc}^{\rm ub}(R,Q) = E_{\rm trc}(R,Q). \tag{141}$$

It follows from (140) and (141) that

$$\mathbb{P}\left[\left|\mathbb{E}\left[\frac{-\log P_{\rm e}^{\rm ub}(\mathcal{C}_n)}{n}\right] - E_{\rm trc}(R,Q)\right| > \varepsilon\right] \to 0.$$
(142)

In conclusion, as anticipated, the three terms of (70) tend to zero as $n \to \infty$, showing (13) for rates below the critical rate. Together with Subsection V-A1, we proved Theorem 1, which states the convergence in probability of the error exponent of the codes in the ensemble to the typical random-coding error exponent.

6) Extension to Constant Composition Codes: For the constant-composition code, for all the rate $R_{crit}(R,Q) \leq R \leq C$, the proof of Theorem 1 holds by using the Levy's continuity theorem since it is not hard to see that $E_{rce}(R,Q) = E_{trc}(R,Q)$ for this case. At all the rate $0 \leq R \leq E_{trc}(R,Q)$, Lemma 4 - Lemma 6 still hold since $\mathbf{1}\{(X_i, X_j) \in \mathcal{T}(Q_{XX'})\}$ and $\mathbf{1}\{(X_k, X_l) \in \mathcal{T}(\tilde{Q}_{XX'})\}$ are still pairwise-independent for the constant-composition code for all $\{i, j, k, l \in [M] : i \neq j, k \neq l\}$. In Lemma 7, the typical error exponent of the union bound should be replaced by E_{trc}^{ub} for the constant-composition code in [9]. To show that Theorem 1 still holds for the constant-composition code, we need to prove that the mapping from the error probability and the union bound in Lemma 8 and Lemma 9 still work. It is not hard to see that the proof of Lemma 8 still holds for the constant-composition code since its correctness depends on Lemma 4, Lemma 6 and the fact that \tilde{V}_{ij} 's are pairwise-independent where \tilde{V}_{ij} is defined (748). Lemma 9 still holds for the constant-composition code, i.e.,

Lemma 10: For any constant-composition code with type Q and for all the rate R such that $0 < R < R_{crit}$, it holds that

$$0 \le \frac{\mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)]}{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)]} - 1 \le 2^{-N\left(\delta(R) + E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) - E_{\mathrm{rce}}(R,Q)\right)}$$
(143)

for some $\delta(R) > 0$.

Proof: To prove Lemma 10, we use the same proof as Lemma 9 in Appendix H. It is easy to check that (796) still holds for the constant-composition code. In addition, the pairwise error probability only depends on the joint-type of the two codewords as in the i.i.d .case.

B. Proof of Theorem 2

First, we prove (18). Under the condition that $E_{\rm rce}(R,Q) = E_{\rm trc}(R,Q)$, observe that

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) < E_{\mathrm{trc}}(R,Q) - \varepsilon\right] = \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) > 2^{-n(E_{\mathrm{trc}}(R,Q) - \varepsilon)}\right]$$
(144)

$$\leq 2^{n(E_{\rm trc}(R,Q)-\varepsilon)} \mathbb{E}[P_{\rm e}(\mathcal{C}_n)] \tag{145}$$

$$\doteq 2^{n(E_{\rm trc}(R,Q)-\varepsilon)}2^{-nE_{\rm rce}(R,Q)}$$
(146)

$$=2^{-n\varepsilon},\tag{147}$$

where (145) follows from the Markov inequality, (146) follows from $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \doteq 2^{-nE_{rce}(R,Q)}$, (147) follows from $E_{rce}(R,Q) = E_{trc}(R,Q)$.

Now, for any s > 0, observe that

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) < E_{\mathrm{trc}}(R,Q) - \varepsilon\right] = \mathbb{P}\left[\frac{s}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) > -s(E_{\mathrm{trc}}(R) - \varepsilon)\right]$$
(148)

$$= \mathbb{P}\left[2^{\frac{s}{n}\log P_{e}(\mathcal{C}_{n})} > 2^{-s(E_{\mathrm{trc}}(R,Q)-\varepsilon)}\right]$$
(149)

$$\leq 2^{s(E_{\rm trc}(R,Q)-\varepsilon)} \mathbb{E}\left[2^{\frac{s}{n}\log P_{\rm e}(\mathcal{C}_n)}\right]$$
(150)

$$=2^{s(E_{\rm trc}(R,Q)-\varepsilon)}\mathbb{E}[(P_{\rm e}(\mathcal{C}_n))^{s/n}]$$
(151)

$$\leq 2^{s(E_{\rm trc}(R,Q)-\varepsilon)} \mathbb{E}[(P_{\rm e}^{\rm ub}(\mathcal{C}_n))^{s/n}].$$
(152)

On the other hand, for any $0 \le s \le n$ and $\lambda > 0$, we have

$$\mathbb{E}\left[\left(P_{e}^{ub}(\mathcal{C}_{n})\right)^{s/n}\right] = \mathbb{E}\left[\left(\frac{1}{M}\sum_{i\neq j}\mathbb{P}(\boldsymbol{X}_{i}\to\boldsymbol{X}_{j})\right)^{s/n}\right]$$
(153)

$$\leq \mathbb{E}\left[\sum_{i\neq j} \left(\frac{1}{M} \sum_{i\neq j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j)\right)^{s/n}\right]$$
(154)

$$= \frac{1}{M^{s/n}} \sum_{i \neq j} \mathbb{E}\left[\left(\mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \right)^{s/n} \right]$$
(155)

$$= \frac{M(M-1)}{M^{s/n}} \mathbb{E}\left[\left(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right)^{s/n}\right]$$
(156)

$$\leq M^{2-\frac{s}{n}} \mathbb{E}\left[\left(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right)^{s/n}\right]$$
(157)

$$= M^{2-\frac{s}{n}} \mathbb{E}\left[\left(\left[\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right]^{1+\lambda}\right)^{\frac{s}{n(1+\lambda)}}\right],\tag{158}$$

where (154) follows from

$$(x_1 + x_2 + \dots + x_n)^{\alpha} \le x_1^{\alpha} + x_2^{\alpha} + \dots + x_n^{\alpha}$$
(159)

for any $x_1, x_2, \dots, x_n \ge 0$ and $\alpha \in [0, 1]$, and (158) follows from $0 \le \mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2) \le 1$ and $\lambda > 0$, so $\left(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right)^{s/n} \le \left(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right)^{s/n} \le \left(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\right)^{s/n}$.

On the other hand, by Lemma 4, the pairwise codeword error probability $\mathbb{P}(X_1 \to X_2)$ given their joint type $Q_{XX'}$ satisfies

$$\mathbb{P}(X_1 \to X_2) \doteq 2^{-n \sum_{k=1}^n \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{1k}, X_{2k}) = (x,x')\}},$$
(160)

where

$$d_{\rm B}(x,x') = -\log\bigg(\sum_{y}\sqrt{W(y|x)W(y|x')}\bigg).$$
(161)

Hence, for any $0 \le s \le n$, we have

$$\mathbb{E}\Big[\big(\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)\big)^{\frac{s}{n}}\Big] = \mathbb{E}\Big[2^{\frac{s}{n}\log\mathbb{P}(\boldsymbol{X}_1 \to \boldsymbol{X}_2)}\Big]$$
(162)

$$\doteq \mathbb{E}\left[2^{-\frac{s}{n}\sum_{k=1}^{n}\sum_{x,x'}^{n}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right].$$
(163)

Now, since $\{\sum_{x,x'} d_B(x,x') \mathbf{1}\{(X_{1k}, X_{2k}) = (x,x')\}\}_{k=1}^n$ are i.i.d., hence by the SLLN, we have

$$\frac{1}{n} \sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{1k}, X_{2k}) = (x,x')\} \xrightarrow{(\mathrm{a.s.})} \sum_{x,x'} Q(x)Q(x')d_{\mathrm{B}}(x,x').$$
(164)

On the other hand, we have

$$0 \le \frac{1}{n} \sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{1k}, X_{2k}) = (x,x')\} \le \max_{x,x'} d_{\mathrm{B}}(x,x') < \infty.$$
(165)

Hence, by the bounded convergence theorem [23], we have

$$\mathbb{E}\left[2^{-\frac{s}{n}\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right] \to 2^{-s\sum_{x,x'}Q(x)Q(x')d_{\mathrm{B}}(x,x')}.$$
(166)

Similarly, for any fixed constant $\lambda \ge 0$, we have

$$\mathbb{E}\left[2^{-\frac{s}{n(1+\lambda)}\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right] \to 2^{-\frac{s}{1+\lambda}\sum_{x,x'}Q(x)Q(x')d_{\mathrm{B}}(x,x')}.$$
(167)

Hence, from (166) and (167), for any fixed constant $\lambda \ge 0$, it holds that

$$\mathbb{E}\left[2^{-\frac{s}{n}\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right]$$
$$=(1+o(1))\left(\mathbb{E}\left[2^{-\frac{s}{n(1+\lambda)}\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right]\right)^{1+\lambda}$$
(168)

$$= (1+o(1)) \left(\mathbb{E} \left[2^{-\frac{s}{n(1+\lambda)} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{11},X_{21})=(x,x')\}} \right] \right)^{n(1+\lambda)}.$$
(169)

From (158) and (169), we obtain

$$\mathbb{E}\left[\left(P_{e}^{ub}(\mathcal{C}_{n})\right)^{s/n}\right] \leq (1+o(1))M^{2-\frac{s}{n}}\left(\mathbb{E}\left[2^{-\frac{s}{n(1+\lambda)}\sum_{x,x'}d_{B}(x,x')\mathbf{1}\{(X_{11},X_{21})=(x,x')\}}\right]\right)^{n(1+\lambda)}.$$
(170)

Now, observe that

$$\mathbb{E}\left[2^{-\frac{s}{n(1+\lambda)}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{11},X_{21})=(x,x')\}}\right]$$

$$=\sum_{x,x'}\mathbb{P}((X_{11},X_{21})=(x,x'))\mathbb{E}\left[2^{-\frac{s}{n(1+\lambda)}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{11},X_{21})=(x,x')\}}\middle|(X_{11},X_{21})=(x,x')\right] \quad (171)$$

$$=\sum_{x,x'}Q(x)Q(x')2^{-\frac{s}{n(1+\lambda)}d_{\mathrm{B}}(x,x')}. \quad (172)$$

From (170) and (172), we obtain

$$\mathbb{E}\left[\left(P_{\rm e}^{\rm ub}(\mathcal{C}_n)\right)^{\frac{s}{n}}\right] \le (1+o(1))M^{2-\frac{s}{n}}\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{s}{n(1+\lambda)}}d_{\rm B}(x,x')\right)^{n(1+\lambda)}.$$
(173)

From (152) and (173), for any s such that $0 \le s \le n$ and any fixed constant $\lambda > 0$, we have

$$\mathbb{P}\left[-\frac{1}{n}\log P_{e}(\mathcal{C}_{n}) < E_{trc}(R,Q) - \varepsilon\right]$$

$$= \mathbb{P}\left[\frac{s}{n}\log P_{e}(\mathcal{C}_{n}) > -s(E_{trc}(R,Q) - \varepsilon)\right]$$
(174)
$$= \left[-\frac{s}{n}\log P_{e}(\mathcal{C}_{n}) - s(E_{trc}(R,Q) - \varepsilon)\right]$$

$$= \mathbb{P}\left[2^{\frac{s}{n}\log P_{e}(\mathcal{C}_{n})} > 2^{-s(E_{\mathrm{trc}}(R,Q)-\varepsilon)}\right]$$
(175)

$$\leq 2^{s(E_{\rm trc}(R,Q)-\varepsilon)} \mathbb{E}\left[2^{\frac{s}{n}\log P_{\rm e}(\mathcal{C}_n)}\right]$$
(176)

$$=2^{s(E_{\rm trc}(R,Q)-\varepsilon)}\mathbb{E}[(P_{\rm e}(\mathcal{C}_n))^{\frac{s}{n}}]$$
(177)

$$\leq (1+o(1))2^{s(E_{\rm trc}(R,Q)-\varepsilon)}M^{2-\frac{s}{n}}\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{s}{n(1+\lambda)}d_{\rm B}(x,x')}\right)^{n(1+\lambda)}.$$
(178)

From (178), by choosing s = n and using $M = 2^{nR}$, we have

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) < E_{\mathrm{trc}}(R,Q) - \varepsilon\right] \leq (1+o(1))2^{n\left(E_{\mathrm{trc}}(R,Q) + \left(2-\frac{s}{n}\right)R-\varepsilon\right)} \left(\sum_{x,x'} Q(x)Q(x')2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda}}\right)^{n(1+\lambda)}$$

$$(179)$$

$$= (1+o(1))2^{n(E_{\text{trc}}(R,Q)+R-\varepsilon)}2^{n(1+\lambda)\log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{a_{\mathbf{B}}(x,x')}{1+\lambda}}\right)}$$
(180)

$$= (1+o(1))2^{n\left[E_{\text{trc}}(R,Q)+R+(1+\lambda)\log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{\text{B}}(x,x')}{1+\lambda}}\right)\right]}2^{-n\varepsilon}.$$
(181)

Now, for $E_{\rm trc}(R,Q) \neq E_{\rm rce}(R,Q)$, from (111) and Lemma 7, observe that

$$E_{\rm trc}(R,Q) = \min_{P_{XX'}: D(P_{XX'} || Q_X Q_{X'}) \le 2R} D(P_{XX'} || Q_X Q_{X'}) + \sum_{x,x'} d_{\rm B}(x,x') P_{XX'}(x,x') - R.$$
(182)

Given the distribution Q and $Q_X = Q_{X'} = Q$, the optimization problem in (182) is convex in $\{P_{XX'}(x, x')\}_{x,x'}$ since the KL divergence is convex. By using standard KKT conditions, it is easy to see that (182) has as optimal solution:

$$P_{XX'} \in \{P_{XX'}^0, P_{XX'}^*\},\tag{183}$$

where

$$P_{XX'}^0(x,x') = \frac{Q(x)Q(x')2^{-d_{\rm B}(x,x')}}{\sum_{x,x'}Q(x)Q(x')2^{-d_{\rm B}(x,x')}},$$
(184)

and

$$P_{XX'}^*(x,x') = \frac{Q(x)Q(x')2^{-\frac{d_{\mathbf{B}}(x,x')}{1+\lambda_*}}}{\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{\mathbf{B}}(x,x')}{1+\lambda_*}}}.$$
(185)

Here, λ^* is the unique positive solution of the following equation:

$$2R = D(P_{XX'}^*(x, x') \| Q_X Q_{X'}).$$
(186)

Now, if $P_{XX'} = P_{XX'}^0$, then we have

$$E_{\rm trc}(R,Q) + R + \log\left(\sum_{x,x'} Q(x)Q(x')2^{-d_{\rm B}(x,x')}\right)$$

$$= D(P_{XX'}^{0} || Q_{X}Q_{X'}) + \sum_{x,x'} d_{\rm B}(x,x')P_{XX'}^{0}(x,x') + \log\left(\sum_{x,x'} Q(x)Q(x')2^{-d_{\rm B}(x,x')}\right)$$
(187)
$$= \sum_{x,x'} \frac{Q(x)Q(x')2^{-d_{\rm B}(x,x')}}{\sum_{x,x'} Q(x)Q(x')2^{-d_{\rm B}(x,x')}} \log \frac{2^{-d_{\rm B}(x,x')}}{\sum_{x,x'} Q(x)Q(x')2^{-d_{\rm B}(x,x')}} + \sum_{x,x'} d_{\rm B}(x,x')P_{XX'}^{0}(x,x') + \log\left(\sum_{x,x'} Q(x)Q(x')2^{-d_{\rm B}(x,x')}\right)$$
(188)

$$= -\sum_{x,x'} \frac{Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}}{\sum_{x,x'}Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}} d_{\mathrm{B}}(x,x') - \log\left(\sum_{x,x'}Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}\right) + \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}^{0}(x,x') + \log\left(\sum_{x,x'}Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}\right)$$
(189)
$$\sum_{x,x'} D_{\mathrm{B}}^{0}(x,x') + \log\left(\sum_{x,x'}Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}\right)$$
(189)

$$= -\sum_{x,x'} P_{XX'}^{0}(x,x') d_{\mathrm{B}}(x,x') - \log\left(\sum_{x,x'} Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}\right) + \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}^{0}(x,x') + \log\left(\sum_{x,x'} Q(x)Q(x')2^{-d_{\mathrm{B}}(x,x')}\right)$$
(190)

$$=0.$$
 (191)

Hence, by choosing $\lambda = 0$, from (181) and (191), we obtain

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) < E_{\rm trc}(R,Q) - \varepsilon\right] \stackrel{\scriptstyle{\scriptstyle{\leq}}}{\leq} 2^{-n\varepsilon}.$$
(192)

Similarly, for the case $P_{XX'} = P^*_{XX'}$, we have

$$E_{\rm trc}(R,Q) + R + (1+\lambda^*) \log\left(\sum_{x,x'} Q(x)Q(x')2^{-\frac{d_{\rm B}(x,x')}{1+\lambda^*}}\right)$$

= $D(P_{XX'}^* \|Q_X Q_{X'}) + \sum_{x,x'} d_{\rm B}(x,x')P_{XX'}^*(x,x') + (1+\lambda^*) \log\left(\sum_{x,x'} Q(x)Q(x')2^{-\frac{d_{\rm B}(x,x')}{1+\lambda^*}}\right)$ (193)
 $\leq (1+\lambda^*)D(P_{XX'}^* \|Q_X Q_{X'}) + \sum_{x,x'} d_{\rm B}(x,x')P_{XX'}^*(x,x') + (1+\lambda^*) \log\left(\sum_{x,x'} Q(x)Q(x')2^{-\frac{d_{\rm B}(x,x')}{1+\lambda^*}}\right)$

$$\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\begin{array}{c}
\end{array} \\
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\begin{array}{c}
\end{array} \\
\end{array} \\
\end{array} \\
\end{array}$$
(194) (194)

$$= (1+\lambda^{*}) \sum_{x,x'} \frac{Q(x)Q(x')2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda^{*}}}}{\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda^{*}}}} \log \frac{2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda^{*}}}}{\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda^{*}}}} + \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}^{*}(x,x') + (1+\lambda^{*})\log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{\mathrm{B}}(x,x')}{1+\lambda^{*}}}\right)$$
(195)

$$= -(1+\lambda^{*})\sum_{x,x'} \frac{Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}}{\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}} \frac{d_{B}(x,x')}{1+\lambda^{*}} - (1+\lambda^{*})\log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}\right) + \sum_{x,x'} d_{B}(x,x')P_{XX'}^{*}(x,x') + \log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}\right) = -\sum_{x,x'}P_{XX'}^{*}(x,x')d_{B}(x,x') - (1+\lambda^{*})\log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}\right) + \sum_{x,x'}d_{B}(x,x')P_{XX'}^{*}(x,x') + \log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}\right) + \sum_{x,x'}d_{B}(x,x')P_{XX'}^{*}(x,x') + \log\left(\sum_{x,x'}Q(x)Q(x')2^{-\frac{d_{B}(x,x')}{1+\lambda^{*}}}\right)$$

$$(197)$$

$$= 0.$$

$$(198)$$

From (181),(191), and (198), where we set $\lambda = 0$ for the first case and $\lambda = \lambda^*$ for the second one, we have

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) < E_{\rm trc}(R,Q) - \varepsilon\right] \le (1+o(1))2^{-n\varepsilon}$$
(199)

$$\doteq 2^{-n\varepsilon}.$$
 (200)

Finally, from (147) and (200), we obtain (18). This concludes our proof of (18) for the i.i.d. random codebook ensemble.

For the constant-composition codebook ensemble, to prove (18), we first prove the following lemma, which is somewhat similar to Lemma 16.

Lemma 11: Let X_1, X_2, \cdots, X_n be Bernoulli random variables on \mathbb{R} . In addition, there exists a set $\mathcal{V} \subset \mathbb{R}^n$ with cardinality $|\mathcal{V}|$ such that for all $x_1, x_2, \cdots, x_n \in \mathcal{V}$,

$$\mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \le (1 + o(1)) \prod_{k=1}^n P(x_k)$$
(201)

for some distribution P on $\{0, 1\}$, and

$$\mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \le 2^{-n\zeta}, \quad \forall (x_1, x_2, \cdots, x_n) \in \mathcal{V}^c$$
(202)

for some $\zeta > 0$. In addition,

$$2^{-n\zeta}|\mathcal{V}^c| \to 0. \tag{203}$$

Let $S_n = X_1 + X_2 + \cdots + X_n$. Then, for any t > 0 it holds that

$$\mathbb{E}_{\prod_{i=1}^{n} P(x_i)} \left[2^{-tS_n/n} \right] + 2^{-n\zeta} |\mathcal{V}^c|.$$
(204)

Proof: Appendix I.

It is known that $\{\mathbf{1}\{(\mathbf{X}_1, \mathbf{X}_2) \in \mathcal{T}_{Q_{XX'}}\}\}_{Q_{XX'}}$ satisfies all the conditions of Lemma 11. Hence, for any $0 \le s \le n$, from (163), we have

$$\mathbb{E}\left[\left(\mathbb{P}(\boldsymbol{X}_{1} \to \boldsymbol{X}_{2})\right)^{\overline{n}}\right] = \mathbb{E}\left[2^{\frac{s}{n}\log\mathbb{P}(\boldsymbol{X}_{1} \to \boldsymbol{X}_{2})}\right]$$
(205)

$$\doteq \mathbb{E}\left[2^{-\frac{s}{n}\sum_{k=1}^{n}\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{1k},X_{2k})=(x,x')\}}\right]$$
(206)

$$\doteq \mathbb{E}_{\mathbf{1}\{(\mathbf{X}_1, \mathbf{X}_2) \in \mathcal{T}_{Q_{XX'}}\}} \text{ is independent} \left[2^{-s \sum_{Q_{XX'}} \sum_{x,x'} d_{\mathrm{B}}(x,x') Q_{XX'}(x,x') \mathbf{1}\{(\mathbf{X}_1, \mathbf{X}_2) \in \mathcal{T}_{Q_{XX'}}\}} \right] + 2^{-n\zeta} |\mathcal{V}^c|.$$

$$(207)$$

The rest follows the same as the proof of (18) for the i.i.d. random codebook ensemble.

Now, we prove (19). For any i.i.d. and constant-composition random codebooks, it is easy to see that

$$P_{\mathbf{e}}(\mathcal{C}_n) \ge \max_{i \ne j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j).$$
(208)

Recall the definition of V_n in (404). It follows from (208) that

$$\mathbb{P}\left[-\frac{1}{n}\log P_{e}(\mathcal{C}_{n}) > E_{trc}(R,Q) + \varepsilon\right]$$

$$\leq \mathbb{P}\left[-\frac{1}{n}\log\max_{i\neq j}\mathbb{P}(\boldsymbol{X}_{i} \to \boldsymbol{X}_{j}) > E_{trc}(R,Q) + \varepsilon\right]$$
(209)

$$= \mathbb{P}\bigg[V_n > n\big(E_{\rm trc}(R,Q) + \varepsilon\big)\bigg]$$
(210)

$$= \mathbb{P}\left[\frac{V_n - \mathbb{E}[V_n]}{\sqrt{\operatorname{Var}(V_n)}} > \frac{n\left(E_{\operatorname{trc}}(R,Q) + \varepsilon - \frac{\mathbb{E}[V_n]}{n}\right)}{\sqrt{\operatorname{Var}(V_n)}}\right]$$
(211)

$$=Q\left(\frac{n\left(E_{\rm trc}(R,Q)+\varepsilon-\frac{\mathbb{E}[V_n]}{n}\right)}{\sqrt{{\rm Var}(V_n)}}\right)+O\left(\frac{1}{\sqrt{n}}\right)$$
(212)

$$= Q\left(\frac{\sqrt{n}\left(E_{\rm trc}(R,Q) + \varepsilon - E_{\rm trc}(Q,0)\right)}{\sqrt{\sum_{x,x'} d_{\rm B}^2(x,x')Q(x)Q(x') - \left(\sum_{x,x'} d_{\rm B}(x,x')Q(x)Q(x')\right)^2}}\right) + O\left(\frac{1}{\sqrt{n}}\right)$$
(213)

as $n \to \infty$ since $E_{\text{trc}}(R, Q) \ge E_{\text{trc}}(Q, 0)$, where (212) follows from Theorem 6, and (213) follows from (532) and the Berry-Esseen theorem [23].

C. Proof of Theorem 3

We start with some accessory results, then prove the main part of Theorem 3.

28

In the lemma below we use the simplified notation: $\mathcal{I}{i, j} = \mathcal{I}{(\mathbf{x}_i, \mathbf{x}_j) \in \mathcal{T}(P_{XX'})}, \mathcal{I}{.}$ being the indicator function, i.e., the two considered codewords have joint type $P_{XX'}$.

Lemma 12:

$$2^{-n2D(P_{XX'}||Q_XQ'_X)} \le \mathbb{E}[\mathcal{I}\{i,j\}\mathcal{I}\{i,k\}] \le 2^{-n[D(P_{XX'}||Q_XQ'_X)+\eta]}$$
(214)

where $0 \leq \eta \leq D(P_{XX'} || Q_X Q'_X)$.

Proof: Appendix J.

Lemma 13: For any $\epsilon > 0$ and for any joint type $P_{XX'}$ such that $D(P_{XX'}||Q_XQ'_X) \leq R - \epsilon$, $\forall \epsilon > 0$, the following holds:

$$\mathbb{P}\left[\mathcal{N}(P_{XX'}) \le 2^{-n\epsilon} \mathbb{E}[\mathcal{N}(P_{XX'})]\right] \le 2^{-2^{n\epsilon}}$$
(215)

Proof: Appendix K.

Using Lemma 12 and Lemma 13 we prove the following theorem, which states that the probability of finding a code for which the exponent of $P_{e}^{ub}(\mathcal{C}_{n})$ is larger than the expurgated exponent $E_{ex}(R)$ is double exponentially decaying in n.

Now we can prove the main part of theorem 3. We have that

$$P_{\rm e}^{\rm ub}(\mathcal{C}_n) \doteq \max_{P_{XX'}} \mathcal{N}(P_{XX'}) e^{-n[\sum_{x,x'} d_{\rm B}(x,x')P_{XX'}(x,x')+R]}$$
(216)

Let us refer to the maximizing joint type of (216) as $P^*_{XX'}$. We define the following complementary events:

$$A = \{P_{XX'}^* \in \mathcal{P}\}\tag{217}$$

$$\overline{A} = \{P_{XX'}^* \in \overline{\mathcal{P}}\}\tag{218}$$

where

$$\mathcal{P} = \{P_{XX'} | D(P_{XX'} \| Q_X Q'_X) \le 2R\}$$

while

$$\overline{\mathcal{P}} = \{P_{XX'} | D(P_{XX'} || Q_X Q'_X) > 2R\},\$$

 $Q_X Q'_X$ being the theoretical joint type. Consider a positive real number $E_2 > E_{\text{trc}}(R,Q)$. We have:

$$\mathbb{P}\left[-\frac{1}{n}\log P_{e}^{ub}(\mathcal{C}_{n}) \ge E_{2}\right] = \mathbb{P}\left[-\frac{1}{n}\log P_{e}^{ub}(\mathcal{C}_{n}) \ge E_{2}, A\right] + \mathbb{P}\left[-\frac{1}{n}\log P_{e}^{ub}(\mathcal{C}_{n}) \ge E_{2}, \overline{A}\right]$$
(219)

Now we proceed to bound from above both terms at the right hand side of (219).

1) First Term:

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) \ge E_{2}, A\right] = \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) \le 2^{-nE_{2}}, A\right]$$

$$[220]$$

$$= \mathbb{P}\left[\frac{1}{M_n} \sum_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{-n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \le 2^{-nE_2}, A\right]$$
(221)

$$= \mathbb{P}\left[\sum_{P_{XX'}} \mathcal{N}(P_{XX'}) e^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')} \le 2^{-n(E_2-R)}, A\right]$$
(222)

$$= \mathbb{P}\left[\max_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')} \le 2^{-n(E_2-R)}, A\right]$$
(223)

$$= \mathbb{P}\left[\max_{P_{XX'} \in \mathcal{P}} \mathcal{N}(P_{XX'}) 2^{-n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \le 2^{-n(E_2 - R)}, A\right]$$
(224)

$$\leq \mathbb{P}\left[\max_{P_{XX'} \in \mathcal{P}} \mathcal{N}(P_{XX'}) 2^{-n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \leq 2^{-n(E_2 - R)}\right]$$
(225)

$$= \mathbb{P}\left[\bigcap_{P_{XX'} \in \mathcal{P}} \left[\mathcal{N}(P_{XX'})e^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')} \le 2^{-n(E_2-R)}\right]\right\}$$
(226)

$$= \mathbb{P}\left[\bigcap_{P_{XX'} \in \mathcal{P}} \left[\mathcal{N}(P_{XX'}) \le 2^{-n(E_2 - R - \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x'))}\right]\right\}$$
(227)

where (224) comes from the definition of A while (225) comes from removing the event A. Let us now define \mathcal{P}' :

$$\mathcal{P}' = \{ P_{XX'} | D(P_{XX'} || Q_X Q'_X) \le R \},$$
(228)

and note that $\mathcal{P}' \subset \mathcal{P}$. Let us consider the term $2^{-n(E_2-R-\sum_{x,x'}d_B(x,x')P_{XX'}(x,x'))}$. We now look for a $P_{XX'} \in \mathcal{P}'$ such that this is smaller than the mean of the enumerator function, i.e, a $P_{XX'} \in \mathcal{P}'$ such that the following holds:

$$2^{-n(E_2 - R - \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x'))} \le 2^{n[2R - D(P_{XX'}||Q_XQ'_X) - \epsilon]}$$
(229)

$$E_2 - R - \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') \ge -[2R - D(P_{XX'}||Q_XQ'_X) - \epsilon]$$
(230)

$$E_2 - R - \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') \ge -2R + D(P_{XX'}||Q_XQ_X') + \epsilon$$
(231)

$$E_2 \ge -R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') + \epsilon.$$
(232)

Let us indicate the $P_{XX'}$ that minimizes (232) with $P'_{XX'}$. Minimizing the term at the right hand side of (232) we can set the value of E_2 to:

$$E_2 = \min_{P_{XX'} \in \mathcal{P}'} -R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x')) + \epsilon.$$
(233)

$$\min_{P_{XX'} \in \mathcal{P}'} -R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') + \epsilon$$

$$= \min_{P_{XX'} \in \mathcal{P}'} +R - 2R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') + \epsilon$$
(234)
(235)

$$> \min_{P_{XX'} \in \mathcal{P}} + R - 2R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') + \epsilon$$
(236)

$$= \min_{P_{XX'} \in \mathcal{Z}_{GGV}} + R - 2R + D(P_{XX'} || Q_X Q'_X) + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') + \epsilon$$
(237)

$$= \min_{P_{XX'} \in \mathcal{Z}_{GGV}} + R + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') + \epsilon$$
(238)

$$=E_{\rm trc}(R,Q)+\epsilon \tag{239}$$

where (236) follows from the fact that $\mathcal{P}' \subset \mathcal{P}$, (237) follows from the concavity of the objective function (minimum is on the border) while (239) follows from the definition of \mathcal{Z}_{GGV} . With this definition of E_2 we ensure that for at least one joint type the conditions for applying Lemma 13 (i.e., (229)) hold. Using the definition in (227) together with the statement of Lemma 13 we have:

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) \geq E_{2}, A\right] \stackrel{.}{\leq} \mathbb{P}\left[\bigcap_{P_{XX'} \in \mathcal{P}} \left[\mathcal{N}(P_{XX'}) \leq 2^{-n(E_{2}-R+\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x'))}\right]\right]\right]$$
(240)

$$\leq 2^{-2^{n[R-D(P'_{XX'})|Q_XQ'_X)]}}$$
(241)

$$\leq 2^{-2^{n\epsilon'}} \tag{242}$$

with $\epsilon' > 0$.

2) Second Term:

$$\mathbb{P}\left[-\frac{1}{n}\log P_{e}^{ub}(\mathcal{C}_{n}) \geq E_{2}, \overline{A}\right] = \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) \leq 2^{-nE_{2}}, \overline{A}\right]$$

$$[243)$$

$$= \mathbb{P}\left[\frac{1}{M_n} \sum_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \le 2^{-nE_2}, \overline{A}\right]$$
(244)

$$= \mathbb{P}\left[\sum_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \le 2^{-n(E_2 - R)}, A\right]$$
(245)

$$\doteq \mathbb{P}\left[\max_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{n \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')} \le 2^{-n(E_2-R)}, \overline{A}\right]$$
(246)

Consider (246). The event \overline{A} implies that the joint type maximizing the expression at the left hand side lays outside \mathcal{P} . This implies that any $P_{XX'}$ which lies inside \mathcal{P} leads to a value which is no greater than the maximum. Since

31

this is an implication of the events within brackets, its probability is larger than or equal to the one of (246). Thus we have:

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) \geq E_{2}, \overline{A}\right] \doteq \mathbb{P}\left[\max_{P_{XX'}} \mathcal{N}(P_{XX'}) 2^{n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')} \leq 2^{-n(E_{2}-R)}, \overline{A}\right]$$
(247)

$$\leq \mathbb{P}\left[\max_{P_{XX'}\in\mathcal{P}}\mathcal{N}(P_{XX'})2^{n\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}(x,x')} \leq 2^{-n(E_2-R)}\right]$$
(248)

$$\leq 2^{-2^{n\epsilon'}} \tag{249}$$

where (249) is because (248) has the same form as (225) and thus the same inequalities as for the first term hold. Finally, we note that from (233) we can further state the following:

$$E_2 = \min_{P_{XX'} \in \mathcal{P}'} -R + D(P_{XX'} || Q_X Q_X') + \sum_{x,x'} d_B(x, x') P_{XX'}(x, x') + \epsilon$$
(250)

$$= \min_{P_{XX'} \in \mathcal{P}'} - \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') + \epsilon$$
(251)

$$=E_{\rm ex}(R)+\epsilon\tag{252}$$

where (251) follows from the concavity of the objective function, which implies that the minimum is on the border of the region \mathcal{P}' , and from the definition of \mathcal{P}' while (252) is found by calculating the derivative of [24, Eq. (5.7.11)] with respect to the optimization variable ρ and, after some change of variable, equating to zero.

D. Proof of Theorem 4

Now let us consider the following inequality

$$P_{\rm e}^{\rm ub}(\mathcal{C}_n) \le M_n P_{\rm e}(\mathcal{C}_n) \tag{253}$$

which follows from upper-bounding the probability $\mathbb{P}[x_i \to x_j]$ in (20) by $\mathbb{P}\left[\bigcup_{j \neq i} \{x_i \to x_j\}\right]$ in (1). From Theorem 3 and using (253) we have

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\rm e}^{\rm ub}(\mathcal{C}_n) \ge E_{\rm ex}(R) + R + \epsilon\right] \le 2^{-2^{n\epsilon}}$$
(254)

and finally

$$\mathbb{P}\left[-\frac{1}{n}\log P_{\rm e}^{\rm ub}(\mathcal{C}_n) \ge E_{\rm ex}(0) + \epsilon\right] \le 2^{-2^{n\epsilon}}.$$
(255)

E. Proof of Theorem 5

1) i.i.d. ensemble: Observe that

$$\max_{i\neq j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \le P_{\mathbf{e}}(\mathcal{C}_n) \le \sum_{i=1}^{M_n} \sum_{j\neq i} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \le M_n(M_n - 1) \max_{i\neq j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j).$$
(256)

On the other hand, by Lemma 4, the pairwise codeword error probability $\mathbb{P}(X_i \to X_j)$ given their joint type $P_{XX'}$ satisfies

$$\mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \doteq 2^{-n \sum_{x,x'} d_{\mathrm{B}}(x,x') \hat{P}_{\boldsymbol{X}_i \boldsymbol{X}_j}(x,x')},$$
(257)

where $\hat{P}_{\boldsymbol{X}_{i}\boldsymbol{X}_{j}}$ is the *n*-joint type of $(\boldsymbol{X}_{1}, \boldsymbol{X}_{2})$, and

$$d_{\rm B}(x,x') = -\log\bigg(\sum_{y}\sqrt{W(y|x)W(y|x')}\bigg).$$
(258)

Observe that

$$\hat{P}_{\boldsymbol{X}_{i}\boldsymbol{X}_{j}}(x,x') = \frac{1}{n} \sum_{k=1}^{n} \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}.$$
(259)

It follows from (257) and (259) that

$$\mathbb{P}(\boldsymbol{X}_{i} \to \boldsymbol{X}_{j}) \doteq 2^{-\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x')} \mathbf{1}_{\{(X_{ik}, X_{jk}) = (x,x')\}}$$
(260)

for all $i, j \in [M_n], i \neq j$. Since M_n sub-exponential in n, from (256) and (260), we obtain

$$-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) \xrightarrow{(\rm a.s.)} \frac{V}{n},\tag{261}$$

where

$$V = \min_{i \neq j} Z_{ij} \tag{262}$$

with

$$Z_{ij} = -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\},$$
(263)

for all $i, j \in [M_n]$ and $i \neq j$.

Now, observe that

$$\mathbb{E}[Z_{ij}] = -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbb{P}\{(X_{ik}, X_{jk}) = (x,x')\}$$
(264)

$$= -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbb{P}(X_{ik} = x) \mathbb{P}(X_{jk} = x')$$
(265)

$$= -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x')Q(x)Q(x').$$
(266)

In addition, we have

$$\operatorname{Var}(Z_{ij}) = \sum_{k=1}^{n} \operatorname{Var}\left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right)$$
(267)

$$= n \left(\mathbb{E} \left[\left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1} \{ (X_{i1}, X_{j1}) = (x,x') \} \right)^2 \right] - \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') \right)^2 \right)$$
(268)

$$= n \left(\mathbb{E} \left[\sum_{x,x'} d_{\mathrm{B}}^2(x,x') \mathbf{1} \{ (X_{1i}, X_{2i}) = (x,x') \} \right] - \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') \right)^2 \right)$$
(269)

$$= n \left(\sum_{x,x'} d_{\mathrm{B}}^2(x,x') \mathbb{P}\{(X_{1i}, X_{2i}) = (x,x')\} - \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') \right)^2 \right)$$
(270)

$$= n \left(\sum_{x,x'} d_{\mathrm{B}}^2(x,x') Q(x) Q(x') - \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') \right)^2 \right).$$
(271)

for all $i \neq j$.

March 16, 2022

Now, define

$$T_{ij} := \frac{Z_{ij} - \mathbb{E}[Z_{ij}]}{\sqrt{\operatorname{Var}(Z_{ij})}}$$
(272)

$$=\frac{Z_{ij} - \mathbb{E}[Z_{12}]}{\sqrt{\operatorname{Var}(Z_{12})}},$$
(273)

where (273) follows from the fact that Z_{ij} 's are identically distributed.

Then, by CLT, it holds that

$$T_{ij} \xrightarrow{(\mathrm{d})} \mathcal{N}(0,1), \quad \forall i \neq j.$$
 (274)

On the other hand, for any fixed tuple $(\{\alpha_{ij}\}: i, j \in [M], i \neq j)$, we have

$$\sum_{i \neq j} \alpha_{ij} T_{ij} = \frac{\sum_{i \neq j} \alpha_{ij} \sum_{k=1}^{N} \sum_{x,x'} d_{\mathrm{B}}(x,x') \left(Q(x)Q(x') - \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\} \right)}{\sqrt{\mathrm{Var}(Z_{12})}}$$
(275)

$$=\sum_{k=1}^{N} \frac{\sum_{i\neq j} \alpha_{ij} \sum_{x,x'} d_{\mathrm{B}}(x,x') \left(Q(x)Q(x') - \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right)}{\sqrt{\mathrm{Var}(Z_{12})}}.$$
(276)

Now, by the i.i.d. random codebook generation, it holds that $\{V_k\}_{k=1}^n$ are i.i.d. random variables, where

$$V_{k} = \frac{\sum_{i \neq j} \alpha_{ij} \sum_{x,x'} d_{\mathrm{B}}(x,x') \left(Q(x)Q(x') - \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\} \right)}{\sqrt{\mathrm{Var}(Z_{12})}}.$$
(277)

In addition, since $(X_{i1}, X_{j1})_{i \neq j}$'s are pairwise independent, we have

$$\operatorname{Var}(V_{1}) = \frac{\sum_{i \neq j} \alpha_{ij}^{2} \sum_{x,x'} d_{\mathrm{B}}^{2}(x,x') \operatorname{Var}\left(\mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\} - Q(x)Q(x')\right))}{\operatorname{Var}(Z_{12})}$$
(278)

$$=\frac{\sum_{i\neq j}\alpha_{ij}^{2}\left[\sum_{x,x'}d_{\mathrm{B}}^{2}(x,x')Q(x)Q(x')-\left(\sum_{x,x'}d_{\mathrm{B}}(x,x')Q(x)Q(x')\right)^{2}\right]}{n\left[\sum_{x,x'}d_{\mathrm{B}}^{2}(x,x')Q(x)Q(x')-\left(\sum_{x,x'}d_{\mathrm{B}}(x,x')Q(x)Q(x')\right)^{2}\right]}$$
(279)

$$=\frac{\sum_{i\neq j}\alpha_{ij}^2}{n}.$$
(280)

Hence, it holds from (276) and (280) that

$$\sum_{i \neq j} \alpha_{ij} T_{ij} = \sqrt{\sum_{i \neq j} \alpha_{ij}^2} \left(\frac{\sum_{k=1}^n V_k}{\sqrt{n \operatorname{Var}(V_1)}} \right)$$
$$\xrightarrow{\text{(d)}} \mathcal{N}\left(0, \sum_{i \neq j} \alpha_{ij}^2\right), \tag{281}$$

where (281) follows from the CLT. Hence, the distribution of the vector $\{T_{ij} : i, j \in [M], i \neq j\}$ goes to the distribution of a jointly Gaussian random vector by the Levy's continuity theorem [23].

Now, it is known that the distribution of any Gaussian random vector (both p.d.f and c.d.f.) is defined by its mean and covariance matrix. Since the covariance matrix of the vector $\{T_{ij} : i, j \in [M], i \neq j\}$ is the identity matrix by the pairwise independence of T_{ij} , which originates from the pairwise independence of $\mathbb{P}(\mathbf{X}_i \to \mathbf{X}_j)'s$, hence, the limit distribution is the standard normal Gaussian vector with dimension M(M-1). This distribution is equal to the joint distribution of M(M-1) independent standard normal variables $\{U_{ij}\}_{i\neq j}$. Hence, by the continuous mapping theorem [23], it follows that

$$\min_{i \neq j} T_{ij} \xrightarrow{(d)} \min_{i \neq j} U_{ij}.$$
(282)

Now, let

$$T = \frac{V - \mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}},\tag{283}$$

then, we have

$$T = \min_{i \neq j} \tilde{T}_{ij},\tag{284}$$

where

$$\tilde{T}_{ij} = \frac{Z_{ij} - \mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}},$$
(285)

Hence, it is easy to see that

$$T = \min_{i \neq j} \frac{\sqrt{\operatorname{Var}(Z_{ij})}}{\sqrt{\operatorname{Var}(V)}} \tilde{T}_{ij} + \frac{\mathbb{E}[Z_{ij}] - \mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}}$$
(286)

$$= \min_{i \neq j} \frac{\sqrt{\operatorname{Var}(Z_{12})}}{\sqrt{\operatorname{Var}(V)}} \tilde{T}_{ij} + \frac{\mathbb{E}[Z_{22}] - \mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}}$$
(287)

$$=\frac{\sqrt{\operatorname{Var}(Z_{12})}}{\sqrt{\operatorname{Var}(V)}}\min_{i\neq j}\tilde{T}_{ij}+\frac{\mathbb{E}[Z_{22}]-\mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}},$$
(288)

where (287) follows from Z_{ij} are identically distributed by the i.i.d. random codebook generation.

Now, assume that

$$\lim_{n \to \infty} \sqrt{\frac{\operatorname{Var}(Z_{12})}{\operatorname{Var}(V)}} = \zeta,$$
(289)

$$\lim_{n \to \infty} \frac{\mathbb{E}[Z_{12}] - \mathbb{E}[V]}{\sqrt{\operatorname{Var}(V)}} = \beta,$$
(290)

for some $\zeta, \beta \in \mathbb{R}^4$. Then, by applying Slutsky's theorem, from (288) and (282), we have

$$T \xrightarrow{(\mathrm{d})} \zeta \min_{i \neq j} U_{ij} + \beta \tag{291}$$

where $U_{ij} \sim \mathcal{N}(0, 1)$ and U_{ij} 's are independent.

Since $\mathbb{E}[T] = 0$ and $\operatorname{Var}(T) = 1$, it follows from Lemma 1 that

$$T \xrightarrow{\text{(d)}} \frac{\min_{i \neq j} U_{ij} - \mathbb{E}[\min_{i \neq j} U_{ij}]}{\sqrt{\operatorname{Var}(\min_{i \neq j} U_{ij})}}.$$
(292)

Finally, from (261), (283) and (292), by appying Slutsky's theorem, we have

$$\frac{\frac{-\log P_{e}(\mathcal{C}_{n})}{n} - \mathbb{E}[\frac{-\log P_{e}(\mathcal{C}_{n})}{n}]}{\sqrt{\operatorname{Var}\left(\frac{-\log P_{e}(\mathcal{C}_{n})}{n}\right)}} \xrightarrow{(d)} \frac{\min_{i \neq j} U_{ij} - \mathbb{E}[\min_{i \neq j} U_{ij}]}{\sqrt{\operatorname{Var}(\min_{i \neq j} U_{ij})}}.$$
(293)

⁴The existence of these limits can be proved easily.

2) Constant-composition ensemble: In this part, we use Stein's method to derive some criteria that provide sufficient conditions for the convergence in distribution to the normal random variable of the error probabilities and error exponents for general random coding ensemble over general channels, including the zero rate where $M_n \to \infty$ as we mentioned. This includes other random codebooks than i.i.d. random codebook ensembles.

We start by showing that Theorem 5 also holds for the constant-composition codes. In order to do this, we need some extra lemmas.

Lemma 14: Let X be a random variable on some finite set \mathcal{X} . Assume that for a certain $\beta > 0$

$$\mathbb{P}\big[\{X \in \mathcal{A}_1\} \cap \{X \in \mathcal{A}_2\}\big] \le \beta \mathbb{P}[X \in \mathcal{A}_1] \mathbb{P}[X \in \mathcal{A}_2]$$
(294)

holds for any $A_1, A_2 \subset \mathcal{X}$. Then, for any sequence of sets $\{A_k\}_{k=1}^{n-1}$ such that $A_k \subset \mathcal{X}$ and there exists $A_i \cap A_j = \emptyset$ for some $i \neq j$ and $i, j \in [n]$, it holds that

$$\mathbb{P}\left[\bigcap_{k=1}^{n} \left\{ X \in \mathcal{A}_{k} \right\}\right] \leq \beta \prod_{k=1}^{n} \mathbb{P}\left[X \in \mathcal{A}_{k} \right].$$
(295)

Proof: Appendix L

Next, we have the following lemma.

Lemma 15: Let $\mathcal{T}_n(\mathcal{X} \times \mathcal{X})$ be the set of all *n*-joint-types in $\mathcal{X} \times \mathcal{X}$ and let $Q_{XX'}, \tilde{Q}_{XX'} \in \mathcal{T}_N(\mathcal{X} \times \mathcal{X})$ such that $Q_{XX'} \neq \tilde{Q}_{XX'}$. Let $\{\mathbf{X}_1, \mathbf{X}_2, \cdots, \mathbf{X}_M\}$ be codewords of a constant-composition code with type Q. For a fixed pair (i, j) with $i \neq j$ and $i, j \in [n]$, define $Z_{Q_{XX'}} = \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(Q_{XX'})\}$ and $Z_{\tilde{Q}_{XX'}} = \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(\tilde{Q}_{XX'})\}$. Then, it holds that

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = a\} \cap \{Z_{\tilde{Q}_{XX'}} = b\}\big] \le \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}\big[Z_{Q_{XX'}} = a\big] \mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = b\big] \bigg|,$$
(296)

for any $(a, b) \in \{0, 1\} \times \{0, 1\}$. Here,

$$I_{\min}(Q) = \min_{Q_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : I_{Q_{XX'}}(X;X') > 0} I_{Q_{XX'}}(X;X') > 0.$$
(297)

Proof: Appendix M

Corollary 1: Let $\mathcal{T}_n(\mathcal{X} \times \mathcal{X})$ be the set of all *n*-joint types in $\mathcal{X} \times \mathcal{X}$. For a constant-composition code and $i \neq j$ and $i, j \in [M]$, let

$$Z_{\tilde{Q}_{XX'}} = \mathbf{1}\left\{ (\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right\}$$
(298)

for all $Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X})$. Then, for any vector $\{z_{Q_{XX'}}\}_{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X})}$ such that there are at least two joint types $Q_{XX'}$ and $\tilde{Q}_{XX'}$ such that $\tilde{Q}_{XX'} \neq Q_{XX'}$ and $z_{Q_{XX'}} = z_{\tilde{Q}_{XX'}} = 1$, it holds that

$$\mathbb{P}\left[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\right] \le \frac{1}{1 - 2^{-nI_{\min}}(Q)} \prod_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \mathbb{P}\left[Z_{Q_{XX'}} = z_{Q_{XX'}}\right]$$
(299)

Proof: Since the vector $\{z_{Q_{XX'}}\}_{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X})}$ has at least two *n*-joint types $Q_{XX'}$ and $\tilde{Q}_{XX'}$ such that $\tilde{Q}_{XX'} \neq Q_{XX'}$ and $z_{Q_{XX'}} = z_{\tilde{Q}_{XX'}} = 1$, it holds that

$$\mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{\tilde{Q}_{XX'}} = z_{Q_{XX'}}\}\bigg] \le \mathbb{P}\bigg[\{Z_{\tilde{Q}_{XX'}} = z_{Q_{XX'}}\} \cap \{Z_{\tilde{Q}_{XX'}} = z_{\tilde{Q}_{XX'}}\}\bigg]$$
(300)

$$= \mathbb{P}\left[\{ Z_{\tilde{Q}_{XX'}} = 1 \} \cap \{ Z_{\tilde{Q}_{XX'}} = 1 \} \right]$$
(301)

$$= \mathbb{P}\left[\left\{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \right\} \cap \left\{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(\tilde{Q}_{XX'}) \right\} \right]$$
(302)

$$= 0.$$
 (303)

This concludes our proof of this corollary.

Now, we show the following fact which is based on Stein's method.

=

Lemma 16: Let f be a bounded function with bounded first and second derivative. Let X_1, X_2, \dots, X_n be zero-mean random variables on \mathbb{R} such that $\mathbb{E}|X_k^4| < \infty$ for all $k \in [n]$. In addition, assume there exists a function $f : \mathbb{Z}_+ \to \mathbb{R}^+$ such that $f(n) \to \infty$ as $n \to \infty$ and a set $\mathcal{V} \subset \mathbb{R}^n$ with cardinality $|\mathcal{V}|$ such that

$$\mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \le (1 + o(1)) \prod_{k=1}^n \mathbb{P}[X_k = x_k]$$
(304)

for all $x_1, x_2, \cdots, x_n \in \mathcal{V}$ and

$$\mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \le 2^{-n\zeta},$$
(305)

$$\sum_{k=1} |x_k|^4 \le g(n), \quad \forall (x_1, x_2, \cdots, x_n) \in \mathcal{V}^c$$
(306)

for some $\zeta > 0$. Assume also that:

$$2^{-n\zeta} |\mathcal{V}^c| g(n) \to 0 \quad \text{as} \quad n \to \infty, \tag{307}$$

$$ng(n) \to \infty$$
 as $n \to \infty$. (308)

Let $S_n = X_1 + X_2 + \dots + X_n$ and

$$\tilde{T} = \frac{S_n}{\sqrt{\operatorname{Var}(S_n)}}.$$
(309)

Then, under the condition that

$$\frac{1}{n^{3/2}} \sum_{i=1}^{n} \mathbb{E}[|X_i^3|] \to 0, \tag{310}$$

$$\frac{1}{n^2} \sum_{i=1}^n \mathbb{E}[X_i^4] \to 0, \tag{311}$$

we have

$$\tilde{T} \xrightarrow{(\mathrm{d})} \mathcal{N}(0,1).$$
 (312)

Proof: Appendix N.

Now, we return to proof Theorem 5. As in the i.i.d. case, we have

$$\frac{-\log P_{\rm e}(\mathcal{C}_n)}{n} \xrightarrow{\text{(a.s.)}} \frac{V}{n},\tag{313}$$

where

$$V = \min_{i \neq j} Z_{ij} \tag{314}$$

with

$$Z_{ij} = -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\},$$
(315)

for all $i, j \in [M]$ and $i \neq j$.

For the constant-composition code, we have

$$Z_{ij} = -\sum_{k=1}^{n} \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}$$
(316)

$$= -\sum_{x,x'} d_{\mathrm{B}}(x,x') \sum_{k=1}^{n} \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\} \sum_{Q_{XX'}} \mathbf{1}\{(X_i, X_j) \in \mathcal{T}(Q_{XX'})\}$$
(317)

$$= -n \sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') d_{\mathcal{B}}(x,x') \mathbf{1}\{(X_i, X_j) \in \mathcal{T}(Q_{XX'})\}.$$
(318)

Let

$$U_{Q_{XX'}} = \frac{\bar{Z}_{ij} - \mathbb{E}[\bar{Z}_{ij}]}{\sqrt{\operatorname{Var}(\bar{Z}_{ij})}}$$
(319)

where

$$\bar{Z}_{ij} = \frac{Z_{ij}}{n} = -\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_i, X_j) \in \mathcal{T}(Q_{XX'})\}.$$
(320)

Then, we have

$$\frac{Z_{ij} - \mathbb{E}[Z_{ij}]}{\sqrt{\operatorname{Var}(Z_ij)}} = \sum_{Q_{XX'}} U_{Q_{XX'}},\tag{321}$$

and

$$U_{Q_{XX'}}(Z_{Q_{XX'}}) = \frac{-\sum_{x,x'} Q_{XX'}(x,x') d_{\mathcal{B}}(x,x') Z_{Q_{XX'}} + \sum_{x,x'} Q_{XX'}(x,x') d_{\mathcal{B}}(x,x') \mathbb{P}(Z_{Q_{XX'}} = 1)}{\sqrt{\operatorname{Var}(\bar{Z}_{ij})}}$$
(322)

where

$$Z_{Q_{XX'}} = \mathbf{1}\{(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\}.$$
(323)

Let

$$\mathcal{V} = \mathcal{V}_1 \cup \mathcal{V}_2 \cup \mathcal{V}_3,\tag{324}$$

where

$$\mathcal{V}_{1} = \left\{ \{ z_{Q_{XX'}} \}_{Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X})} : \text{there are at least two different } n\text{-joint types } Q_{XX'} \text{ and } \tilde{Q}_{XX'} \\ \text{such that } z_{Q_{XX'}} = z_{\tilde{Q}_{XX'}} = 1 \right\},$$

$$(325)$$

and

$$\mathcal{V}_{2} = \left\{ \{ z_{Q_{XX'}} \}_{Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X})} : \text{there is exactly one } n\text{-joint types } Q_{XX'}^{*} \right\}$$
such that $z_{Q_{XX'}}^{*} = 1$ and $Q_{XX'}^{*} = Q_{X}^{*}Q_{X'}^{*} \right\},$
(326)

and

$$\mathcal{V}_{3} = \left\{ \{ z_{Q_{XX'}} \}_{Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X})} : \text{such that } z_{Q_{XX'}} = 1 \text{ for all } Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X}) \right\}.$$
(327)

Now, for any $\{z_{Q_{XX'}}\}_{Q_{XX'}\in\mathcal{V}},$ there are three subcases:

• $\{z_{Q_{XX'}}\}_{Q_{XX'}} \in \mathcal{V}_3$. Then, we have

$$\mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\bigg] = \mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{(\mathbf{X}_i, \mathbf{X}_j)\in\mathcal{T}(Q_{XX'})\}\bigg]$$
(328)

$$= 0$$
 (329)

$$\leq \prod_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \mathbb{P}[Z_{Q_{XX'}} = z_{Q_{XX'}}].$$
(330)

• $\{z_{Q_{XX'}}\}_{Q_{XX'}} \in \mathcal{V}_2$. Then, we have

$$\mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q^*_{XX'})\big] \doteq 1.$$
(331)

$$\mathbb{P}\left[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\right]$$
$$= \mathbb{P}\left[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(Q_{XX'})\}\right]$$
(332)

$$= \mathbb{P}\bigg[\{ Z_{Q_{XX'}^*} = 1 \} \cap \bigcap_{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^*} \{ Z_{Q_{XX'}} = 0 \} \bigg]$$
(333)

$$= \mathbb{P}\left[\{(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}^{*})\} \cap \bigcap_{Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^{*}} \{(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \notin \mathcal{T}(Q_{XX'})\}\right]$$
(334)

$$= \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q^*_{XX'}) \right].$$
(335)

On the other hand, it is known that

$$\prod_{\substack{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^* \\ = \prod_{\substack{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^*}} \mathbb{P}[(\mathbf{X}_i, \mathbf{X}_j) \notin \mathcal{T}(Q_{XX'})]$$
(336)

$$\doteq \prod_{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^*} \left(1 - 2^{-nI_{Q_{XX'}}(X;X')} \right)$$
(337)

$$\stackrel{\cdot}{\geq} \prod_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X}):Q_{XX'}\neq Q^*_{XX'}} \left(1-2^{-nI_{\min}(Q)}\right)$$
(338)

$$= \left(1 - 2^{-nI_{\min}(Q)}\right)^{|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})| - 1}$$
(339)

$$\geq 1 - \left(\left| \mathcal{T}_n(\mathcal{X} \times \mathcal{X}) \right| - 1 \right) 2^{-nI_{\min}(Q)}$$
(340)

$$= 1 + o(1),$$
 (341)

where (340) follows from $(1-x)^n \ge 1-nx$ for all $x \in (0,1)$, and (341) follows from $|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})| \le (n+1)^{|\mathcal{X}|}$. From (335) and (340), we have

$$\mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\bigg] \le (1+o(1))\prod_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \mathbb{P}\big[Z_{Q_{XX'}} = z_{Q_{XX'}}\big].$$
(342)

• $\{z_{Q_{XX'}}\}_{Q_{XX'}} \in \mathcal{V}_1$. Then, from Lemma 14 and Lemma 15, we obtain

$$\mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\bigg] \le (1+o(1))\prod_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \mathbb{P}\big[Z_{Q_{XX'}} = z_{Q_{XX'}}\big].$$
(343)

From case 1, case 2, and case 3, on \mathcal{V} , it holds

$$\mathbb{P}\bigg[\bigcap_{Q_{XX'}\in\mathcal{T}_N(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\bigg] \le (1+o(1))\prod_{Q_{XX'}\in\mathcal{T}_N(\mathcal{X}\times\mathcal{X})} \mathbb{P}\big[Z_{Q_{XX'}} = z_{Q_{XX'}}\big].$$
(344)

Now, for $\{z_{Q_{XX'}}\}_{Q_{XX'}} \in \mathcal{V}^c$, it holds that $z_{Q_{XX'}} = 1$ for exact one N-joint type $Q_{XX'}^* \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X})$ and $I_{Q_{XX'}^*}(X;X') \ge I_{\min}(Q) + o_N(1)$. Note that there are at most $|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})| \le (n+1)^{|\mathcal{X}|}$ such sequences. Hence, on this subset, we have

$$\mathbb{P}\left[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{Z_{Q_{XX'}} = z_{Q_{XX'}}\}\right]$$
$$= \mathbb{P}\left[\bigcap_{Q_{XX'}\in\mathcal{T}_n(\mathcal{X}\times\mathcal{X})} \{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(Q_{XX'})\}\right]$$
(345)

$$= \mathbb{P}\bigg[\{ Z_{Q_{XX'}^*} = 1 \} \cap \bigcap_{Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X}) : Q_{XX'} \neq Q_{XX'}^*} \{ Z_{Q_{XX'}} = 0 \} \bigg]$$
(346)

$$= \mathbb{P}\left[\{(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}^{*})\} \cap \bigcap_{Q_{XX'} \in \mathcal{T}_{n}(\mathcal{X} \times \mathcal{X}): Q_{XX'} \neq Q_{XX'}^{*}} \{(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \notin \mathcal{T}(Q_{XX'})\}\right]$$
(347)

$$= \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q^*_{XX'}) \right]$$
(348)

$$\leq 2^{-nI_{\min}(Q)}.\tag{349}$$

In addition, on \mathcal{V}^c , we have

$$\sum_{Q_{XX'}} \frac{U_{Q_{XX'}}^4}{\operatorname{Var}(\bar{Z}_{ij})} \le \left(\operatorname{Var}(\bar{Z}_{ij})\right)^{-1} \left(\sum_{Q_{XX'}} U_{Q_{XX'}}^4\right).$$
(350)

Now, observe that

$$\operatorname{Var}(\bar{Z}_{ij}) = \mathbb{E}\left[\left(\sum_{Q_{XX'}}\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')\mathbf{1}\{(\boldsymbol{X}_{i},\boldsymbol{X}_{j})\in\mathcal{T}(Q_{XX'})\}\right)^{2}\right] - \left(\mathbb{E}\left[\sum_{Q_{XX'}}\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')\mathbf{1}\{(\boldsymbol{X}_{i},\boldsymbol{X}_{j})\in\mathcal{T}(Q_{XX'})\}\right]\right)^{2}$$
(351)

$$= \mathbb{E} \left[\sum_{\tilde{Q}_{XX'}} \sum_{Q_{XX'}} \sum_{x,x'} \sum_{y,y'} \tilde{Q}_{XX'}(y,y') Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') d_{\mathrm{B}}(y,y') \right. \\ \left. \times \mathbf{1} \{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \} \mathbf{1} \{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(\tilde{Q}_{XX'}) \} \right] \\ \left. - \left(\mathbb{E} \left[\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') \mathbf{1} \{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \} \right] \right)^{2}$$
(352)

$$= \mathbb{E}\bigg[\sum_{Q_{XX'}} \sum_{x,x'} \sum_{y,y'} Q_{XX'}(y,y') Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') d_{\mathrm{B}}(y,y') \mathbf{1}\{(\boldsymbol{X}_{i},\boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'})\}\bigg] - \bigg(\mathbb{E}\bigg[\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') \mathbf{1}\{(\boldsymbol{X}_{i},\boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'})\}\bigg]\bigg)^{2}$$
(353)

$$= \sum_{Q_{XX'}} \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right] \sum_{x,x'} \sum_{y,y'} Q_{XX'}(y, y') Q_{XX'}(x, x') d_{\mathrm{B}}(x, x') d_{\mathrm{B}}(y, y') - \left(\mathbb{E}\left[\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x, x') d_{\mathrm{B}}(x, x') \mathbf{1}\{(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\} \right] \right)^2$$
(354)

$$= \sum_{Q_{XX'}} \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right] \left(\sum_{x,x'} Q_{XX'}(x, x') d_{\mathrm{B}}(x, x') \right)^2 - \left(\sum_{Q_{XX'}} \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right] \sum_{x,x'} Q_{XX'}(x, x') d_{\mathrm{B}}(x, x') \right)^2$$
(355)

$$= \sum_{Q_{XX'}} \mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \right] \left(\sum_{x,x'} Q_{XX'}(x, x') d_{B}(x, x') \right)^{2}$$
$$- \sum_{Q_{XX'}} \left(\mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \right] \right)^{2} \left(\sum_{x,x'} Q_{XX'}(x, x') d_{B}(x, x') \right)^{2}$$
$$- 2 \sum_{Q_{XX'}} \sum_{\tilde{Q}_{XX'}} \sum_{x,x'} \sum_{y,y'} Q_{XX'}(x, x') \tilde{Q}_{XX'}(y, y') d_{B}(x, x') d_{B}(y, y')$$
$$\times \mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \right] \mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(\tilde{Q}_{XX'}) \right].$$
(356)

Since $Q_{XX'} \neq \tilde{Q}_{XX'}$, we have

$$\mathbb{P}[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'})] \mathbb{P}[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(\tilde{Q}_{XX'})] \doteq 2^{-nI_{Q_{XX'}}(X;X')} 2^{-nI_{\tilde{Q}_{XX'}}(X;X')}$$
(357)

$$\leq 2^{-nI_{\min}(Q)}.\tag{358}$$

Hence, we have

$$\sum_{Q_{XX'}} \sum_{\tilde{Q}_{XX'}} \sum_{x,x'} \sum_{x,x'} \sum_{y,y'} Q_{XX'}(x,x') \tilde{Q}_{XX'}(y,y') d_{\mathcal{B}}(x,x') d_{\mathcal{B}}(y,y')$$

$$\times \mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(Q_{XX'}) \right] \mathbb{P}\left[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(\tilde{Q}_{XX'}) \right]$$
$$\leq 2^{-nI_{\min}(Q)} \sum_{Q_{XX'}} \sum_{\tilde{Q}_{XX'}} \sum_{x,x'} \sum_{y,y'} Q_{XX'}(x, x') \tilde{Q}_{XX'}(y, y') d_{\mathrm{B}}(x, x') d_{\mathrm{B}}(y, y')$$
(359)

$$\leq 2^{-nI_{\min}(Q)} \bigg(\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') d_{\mathrm{B}}(x,x') \bigg)^2$$
(360)

$$\leq 2^{-nI_{\min}(Q)} d_{\max}^2 \left(\sum_{Q_{XX'}} \sum_{x,x'} Q_{XX'}(x,x') \right)^2$$
(361)

$$\leq 2^{-nI_{\min}(Q)} d_{\max}^2 |\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|^2 \tag{362}$$

$$\leq 2^{-nI_{\min}(Q)} d_{\max}^2 (n+1)^{2|\mathcal{X}|},\tag{363}$$

where

$$d_{\max} = \max_{x,x'} \left[-\log\left(\sum_{y} \sqrt{W(y|x)W(y|x')}\right) \right].$$
(364)

From (356) and (363), we obtain

$$\operatorname{Var}(\bar{Z}_{ij}) \geq \sum_{Q_{XX'}} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})] \left(\sum_{x, x'} Q_{XX'}(x, x') d_{\mathsf{B}}(x, x')\right)^2 - 2^{-nI_{\min}(Q)} d_{\max}^2 (n+1)^{2|\mathcal{X}|}$$

$$= \sum_{Q_{XX'}} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})] \left(1 - \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})]\right) \left(\sum_{x, x'} Q_{XX'}(x, x') d_{\mathsf{B}}(x, x')\right)^2 - 2^{-nI_{\min}(Q)} d_{\max}^2 (n+1)^{2|\mathcal{X}|}$$

$$= \sum_{Q_{XX'}} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})] \left(1 - \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})]\right) \left(\sum_{x, x'} Q_{XX'}(x, x') d_{\mathsf{B}}(x, x')\right)^2 - 2^{-nI_{\min}(Q)} d_{\max}^2 (n+1)^{2|\mathcal{X}|}$$

$$(365)$$

$$= \sum_{Q_{XX'}} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})] \left(1 - \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})]\right) \left(\sum_{x, x'} Q_{XX'}(x, x') d_{\mathsf{B}}(x, x')\right)^2$$

$$(366)$$

$$\geq \sum_{Q_{XX'}:Q_{XX'}\neq Q_XQ_{X'}} \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right] \left(1 - \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'}) \right] \right) \left(\sum_{x,x'} Q_{XX'}(x, x') d_{\mathrm{B}}(x, x') \right)^2 - 2^{-nI_{\min}(Q)} d_{\max}^2(n+1)^{2|\mathcal{X}|}$$
(367)

$$\geq \sum_{\substack{Q_{XX'}:Q_{XX'}\neq Q_XQ_{X'}\\-2^{-nI_{\min}(Q)}d_{\max}^2(n+1)^{2|\mathcal{X}|}}} \mathbb{P}\left[(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(Q_{XX'})\right] \left(1 - 2^{-nI_{\min}(Q)}\right) \left(\sum_{x,x'} Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')\right)^2$$
(368)

$$\geq \left(1 - 2^{-nI_{\min}(Q)}\right) d_{\min}^2 \sum_{Q_{XX'}:Q_{XX'} \neq Q_X Q_{X'}} \mathbb{P}\left[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\right] - 2^{-nI_{\min}(Q)} d_{\max}^2 (n+1)^{2|\mathcal{X}|}$$
(369)

where

$$d_{\min} = \min_{x,x'} \left[-\log\left(\sum_{y} \sqrt{W(y|x)W(y|x')}\right) \right] > 0.$$
(370)

Now, observe that

$$\sum_{Q_{XX'}:Q_{XX'}\neq Q_X Q_{X'}} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})]$$
$$= 1 - \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_X Q_{X'})]$$
(371)

$$= 1 - \sum_{\{i,i=1\}} 2^{-nH(Q_X)} 2^{-nH(Q'_X)}$$
(372)

$$(\mathbf{x},\mathbf{x}')\in\mathcal{T}(Q_XQ_{X'})$$

$$= 1 - \left| \mathcal{T}(Q_X Q_{X'}) \right| 2^{-nH(Q_X)} 2^{-nH(Q'_X)}.$$
(373)

Now, by [29], it holds that

$$2^{nH(Q_XQ_{X'})}(n+1)^{-|\mathcal{X}|} \le \left|\mathcal{T}(Q_XQ_{X'})\right| \le 2^{nH(Q_XQ_{X'})}.$$
(374)

We can assume that $\left|\mathcal{T}(Q_XQ_{X'})\right| \neq 2^{nH(Q_XQ_{X'})},$ then

$$|\mathcal{T}(Q_X Q_{X'})| \le (\text{poly}(n))^{-1} 2^{nH(Q_X Q_{X'})}$$
(375)

where $poly(n) \neq 1$ where poly(n) is some polynomial in n. Then, from (373) and $Q_X = Q_{X'} = Q$, we obtain

$$\sum_{Q_{XX'}:Q_{XX'}\neq Q_XQ_{X'}} \mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\big] \ge 1 - (\operatorname{poly}(n))^{-1} \ge 1 - \alpha$$
(376)

for some $\alpha \in (0,1)$ and n sufficiently large.

Hence, from (369), (373), and (376), we have

$$\operatorname{Var}(\bar{Z}_{ij}) \ge \left(1 - 2^{-nI_{\min}(Q)}\right) d_{\min}^2(1 - \alpha) - 2^{-nI_{\min}(Q)} d_{\max}^2(n+1)^{2|\mathcal{X}|}$$
(377)

$$= \Omega(1). \tag{378}$$

From (350) and (378),

$$\sum_{Q_{XX'}} \frac{V_{Q_{XX'}}^4(Z_{Q_{XX'}})}{\operatorname{Var}(\bar{Z}_{ij})} \le \left(\operatorname{Var}(\bar{Z}_{ij})\right)^{-1} \left(\sum_{Q_{XX'}} V_{Q_{XX'}}^4(Z_{Q_{XX'}})\right)$$
(379)

$$= O(1) \sum_{Q_{XX'}} V_{Q_{XX'}}^4(Z_{Q_{XX'}})$$
(380)

where

$$V_{Q_{XX'}}(Z_{Q_{XX'}}) = -\sum_{x,x'} Q_{XX'}(x,x') d_{\mathcal{B}}(x,x') Z_{Q_{XX'}} + \sum_{x,x'} Q_{XX'}(x,x') d_{\mathcal{B}}(x,x') \mathbb{P}(Z_{Q_{XX'}} = 1).$$
(381)

Note that

$$\left|V_{Q_{XX'}}(Z_{Q_{XX'}})\right|^{4} \leq 2\left(\left|\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')Z_{Q_{XX'}}\right|^{4} + \left|\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')\mathbb{P}(Z_{Q_{XX'}}=1)\right|^{4}\right)$$
(382)

$$\leq 4d_{\max}^4$$
 (383)

for all $Q_{XX'} \in \mathcal{T}_n(\mathcal{X} \times \mathcal{X})$, and

$$|V_{Q_{XX'}}(Z_{Q_{XX'}})|^{3} \leq 2\left(\left|\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')Z_{Q_{XX'}}\right|^{3} + \left|\sum_{x,x'}Q_{XX'}(x,x')d_{\mathrm{B}}(x,x')\mathbb{P}(Z_{Q_{XX'}}=1)\right|^{4}\right)$$
(384)

$$\leq 4d_{\max}^3 \tag{385}$$

Hence, from (380) and (383), we have

$$\sum_{Q_{XX'}} \frac{V_{Q_{XX'}}^4(Z_{Q_{XX'}})}{\operatorname{Var}(\bar{Z}_{ij})} \le 4|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|d_{\max}^4$$
(386)

$$\leq 4(n+1)^{|\mathcal{X}|} d_{\max}^4. \tag{387}$$

From (349) and (387), we have

$$2^{-nI_{\min}(Q)} |\mathcal{V}^c| \sum_{Q_{XX'}} \frac{V_{Q_{XX'}}^4(Z_{Q_{XX'}})}{\operatorname{Var}(\bar{Z}_{ij})} \le 2^{-nI_{\min}(Q)} (n+1)^{|\mathcal{X}|} 4(n+1)^{|\mathcal{X}|} d_{\max}^4 \to 0$$
(388)

as $n \to \infty$.

Now, we have

$$\frac{1}{|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|^2} \sum_{Q_{XX'}} V_{Q_{XX'}}^4(Z_{Q_{XX'}}) \le \frac{1}{|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|} 4d_{\max}^4$$
(389)

$$\doteq \frac{1}{2^{nH(Q_{XX'})}} 4d_{\max}^4$$
 (390)

 $\rightarrow 0$ (391)

as $n \to \infty$, where (389) follows from (386).

Similarly, we have

$$\frac{1}{|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|^{3/2}} \sum_{Q_{XX'}} |V_{Q_{XX'}}(Z_{Q_{XX'}})|^3 \le \frac{1}{|\mathcal{T}_n(\mathcal{X} \times \mathcal{X})|^{3/2}} \sum_{Q_{XX'}} 4d_{\max}^3$$
(392)

$$\rightarrow 0.$$
 (393)

From the above facts and Lemma 16, we conclude that

$$T_{ij} = \frac{Z_{ij} - \mathbb{E}[Z_{ij}]}{\sqrt{\operatorname{Var}(Z_{ij})}} \xrightarrow{\text{(d)}} \mathcal{N}(0, 1).$$
(394)

Similarly, we can prove that if M is a constant, we have

$$\sum_{i \neq j} \alpha_{ij} T_{ij} \xrightarrow{\text{(d)}} \mathcal{N}(0, 1).$$
(395)

for any sequence $\{\alpha_{ij}\}_{i,j\in[M],i\neq j}$. Then, by using the same arguments as the proof of Lemma 5 for the rest, we obtain our result in (23).

F. Proof of Theorem 6

Our proof of this theorem is based on a modification of the Wasserstein metric, inspired by the classical Kolmogorov and Wasserstein metrics, that measures the distance between the distribution of the error exponent and that of the standard Gaussian. Such modification is needed to deal with an infinite number of terms as $n \to \infty$, a case where the classical Wasserstein metric upper bound fails to work [25, Prop. 2.4].

Recall the definitions of probability metrics in Definition 2. First, we prove the following fundamental lemma. Lemma 17: If $Z \sim \mathcal{N}(0, 1)$, then for any random variable T, it holds that

$$\left|\mathbb{P}(T \le x) - \mathbb{P}(Z \le x)\right| \le 2(8\pi)^{-1/4} \sqrt{d_{W,\text{mod}}(T,Z)} + \left|\mathbb{P}(T \le x) - \mathbb{P}(T \ge -x)\right|$$
(396)

for all $x \in \mathbb{R}$. In addition, if the distribution of T is tight⁵, for any $x \to 0$, which is a continuous point of the limit distribution of T, as $N \to \infty$, we have

$$\limsup_{N \to \infty} \left| \mathbb{P}(T \le x) - \mathbb{P}(Z \le x) \right| \le 2(8\pi)^{-1/4} \limsup_{N \to \infty} \sqrt{d_{W, \text{mod}}(T, Z)}.$$
(397)

Proof: Appendix O.

By using the definition of $d_{W,\text{mod}}$ and setting T = X and Y = Z where $Z \sim \mathcal{N}(0, 1)$, we obtain the following result, which is tighter than (or at least equal to) the upper bound of $d_K(T, Z)$ in [25, Prop. 2.4]. However, we note that the probability metric here is the modified Wasserstein metric. See the same arguments to achieve a similar result in [25, Prop. 2.4].

Lemma 18: For $h \in \mathcal{H}$, let f_h solve

$$f'_{h}(w) - wf_{h}(w) = h(w) - \mathbb{E}[h(Z)].$$
 (398)

If T is a random variable and Z has the standard normal distribution, then

$$d_{W,\text{mod}}(T,Z) \le \sup_{h \in \mathcal{H}} \min\left\{ \left| \mathbb{E} \left[f'_h(T) - Tf_h(T) \right] \right|, \left| \mathbb{E} \left[f'_h(-T) + Tf_h(-T) \right] \right| \right\}.$$
(399)

Proof: Left as exercise.

Now, we prove the following lemma.

Lemma 19: Assume that $T = \min\{T_1, T_2, \dots, T_L\}$ for some $L \in \mathbb{Z}^+$ and T_1, T_2, \dots, T_L are identically distributed random variables. Then, it holds that

$$d_{W,\text{mod}}(T,Z) \le \max\left\{ \sup_{h \in \mathcal{H}} \left| \mathbb{E}[f'_{h}(T_{1}) - T_{1}f_{h}(T_{1})] \right|, \sup_{h \in \mathcal{H}} \left| \mathbb{E}[f'_{h}(-T_{1}) + T_{1}f_{h}(-T_{1})] \right| \right\} + \sup_{h \in \mathcal{H}} \min\left\{ \mathbb{E}[h(T) - h(T_{1})], \mathbb{E}[h(-T_{1}) - h(-T)] \right\}.$$
(400)

Proof: Appendix P.

⁵A distribution on $(\mathbb{R}, \mathcal{B}(\mathbb{R}))$ is tight if for any fixed $\varepsilon > 0$, there exists $u, v \in \mathbb{R}$ such that $\mathbb{P}(u < T \le v) > 1 - \varepsilon$ [23].

Lemma 20: [25, Th. 3.2] Let X_1, X_2, \dots, X_n be independent mean zero random variables such that $\mathbb{E}[|X_i|^4] < \infty$ and $\mathbb{E}[X_i^2] = 1$. If $T = \sum_{i=1}^n X_i / \sqrt{n}$ and Z has the standard normal distribution, then

$$\max\left\{\sup_{h\in\mathcal{H}}\left|\mathbb{E}\left[f_{h}'(T) - Tf_{h}(T)\right]\right|, \sup_{h\in\mathcal{H}}\left|\mathbb{E}\left[f_{h}'(-T) + Tf_{h}(-T)\right]\right|\right\} \le \frac{1}{n^{3/2}}\sum_{i=1}^{n}\mathbb{E}[|X_{i}|^{3}] + \frac{\sqrt{2}}{n\sqrt{\pi}}\sqrt{\sum_{i=1}^{n}\mathbb{E}[X_{i}^{4}]}.$$
(401)

We can observe the fact (401) since T and -T are both the sums of independent random variables. Now, we are ready to prove Theorem 6. Observe that

$$\max_{i \neq j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \le P_{\mathbf{e}}(\mathcal{C}_n) \le \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) \le M_n(M_n - 1) \max_{i \neq j} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j).$$
(402)

Hence, for M_n sub-exponential in n, it holds that

$$-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) \xrightarrow{(\rm a.s.)} \frac{V_n}{n},\tag{403}$$

where

$$V_n = \min_{i \neq j} Z_{ij}(n) \tag{404}$$

with

$$Z_{ij}(n) = -\sum_{k=1}^{n} \sum_{x,x'} d_{\rm B}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\},\tag{405}$$

for all $i, j \in [M_n]$ and $i \neq j$ (See the proof for this fact from the proof of Theorem 5).

Define

$$T_{ij}(n) = \frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(n))}},$$
(406)

we have

$$\min_{i \neq j} T_{ij}(n) = \min_{i \neq j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(n))}}.$$
(407)

Now, for any $\varepsilon > 0$, let the event

$$\mathcal{E}_{n} = \left\{ \frac{1}{M_{n}(M_{n}-1)} \left| \sum_{i \neq j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(n))}} \right| \ge \varepsilon \right\}$$
(408)

for all $n \in \mathbb{Z}^+$. Then, we have

$$\sum_{n=1}^{\infty} \mathbb{P}(\mathcal{E}_n) \le \sum_{n=1}^{\infty} \mathbb{P}\left[\frac{1}{(M_n - 1)M_n} \left| \sum_{i \ne j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(n))}} \right| \ge \varepsilon \right]$$
(409)

$$\leq \sum_{n=1}^{\infty} \frac{1}{\varepsilon^2 M_n^2 (M_n - 1)^2} \operatorname{Var}\left(\sum_{i \neq j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(n))}}\right)$$
(410)

$$=\sum_{n=1}^{\infty} \frac{1}{\varepsilon^2 M_n^2 (M_n - 1)^2} \sum_{i \neq j} \operatorname{Var}\left(\frac{Z_{ij}(n) - \mathbb{E}[Z_{ij}(n)]}{\sqrt{\operatorname{Var}(Z_{ij}(N))}}\right)$$
(411)

$$=\sum_{n=1}^{\infty}\frac{1}{\varepsilon^2 M_n(M_n-1)}$$
(412)

$$<\infty,$$
 (413)

where (410) follows from Chebyshev's inequality, (411) follows from the pairwise independence of Z_{ij} 's, and (413) follows from the condition (26).

Hence, by Borel-Cantelli lemma, from (413), we have

$$\mathbb{P}\bigg[\bigcup_{n=1}^{\infty}\bigcap_{k=n}^{\infty}\mathcal{E}_{k}^{c}\bigg] = 1.$$
(414)

However, we have

$$\mathbb{P}\left[\bigcup_{n=1}^{\infty}\bigcap_{k=n}^{\infty}\mathcal{E}_{k}^{c}\right] = \mathbb{P}\left[\bigcup_{n=1}^{\infty}\bigcap_{k=n}^{\infty}\left\{\frac{1}{M_{n}(M_{n}-1)}\left|\sum_{i\neq j}\frac{Z_{ij}(k) - \mathbb{E}[Z_{ij}(k)]}{\sqrt{\operatorname{Var}(Z_{ij}(k))}}\right| < \varepsilon\right\}\right].$$
(415)

It follows from (414) and (415) that

$$\mathbb{P}\left[\left.\bigcup_{n=1}^{\infty}\bigcap_{k=n}^{\infty}\left\{\frac{1}{M_n(M_n-1)}\right|\sum_{i\neq j}\frac{Z_{ij}(k)-\mathbb{E}[Z_{ij}(k)]}{\sqrt{\operatorname{Var}(Z_{ij}(k))}}\right|<\varepsilon\right\}\right]=1,\tag{416}$$

or

$$\frac{1}{M_n(M_n-1)} \sum_{i \neq j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{12}(n)]}{\sqrt{\operatorname{Var}(Z_{12}(n))}} \xrightarrow{\text{(a.s.)}} 0, \tag{417}$$

as $n \to \infty$. Hence, there exists a subset \mathcal{A} such that $\mathbb{P}(\mathcal{A}) = 1$ and

$$\frac{1}{M_n(M_n-1)} \sum_{i \neq j} \frac{Z_{ij}(n) - \mathbb{E}[Z_{12}(n)]}{\sqrt{\operatorname{Var}(Z_{12}(n))}} \to 0$$
(418)

on \mathcal{A} .

Now, from Theorem 5, we have $T_{ij}(n), T_{i'j'}(n), T_{12}(n)$ are independent as $n \to \infty$ if $(i, j) \neq (i', j') \neq (1, 2)$. Then, for any $B_1, B_2 \in \mathcal{B}(R)$ (Borel sets in \mathbb{R}), as $n \to \infty$, we have

$$\mathbb{P}\left[\left\{T_{ij}(n) - T_{12}(n) \in B_{1}\right\} \cap \left\{T_{ij}(n) - T_{12}(n) \in B_{2}\right\}\right] \\
= \int_{\mathbb{R}} \mathbb{P}\left[\left\{T_{ij}(n) - T_{12}(n) \in B_{1}\right\} \cap \left\{T_{i'j'}(n) - T_{12}(n) \in B_{2}\right\} \middle| T_{12}(n) = \alpha\right] f_{T_{12}(n)}(\alpha) d\alpha \tag{419}$$

$$= \int_{\mathbb{R}} \mathbb{P}\left[\left\{T_{ij}(n) \in \alpha + B_1\right\} \cap \left\{T_{i'j'}(n) \in \alpha + B_2\right\} \middle| T_{12}(n) = \alpha\right] f_{T_{12}(n)}(\alpha) d\alpha$$
(420)

$$= \int_{\mathbb{R}} \mathbb{P}\left[\left\{T_{ij}(n) \in \alpha + B_1\right\} \cap \left\{T_{i'j'}(n) \in \alpha + B_2\right\}\right] f_{T_{12}(n)}(\alpha) d\alpha$$
(421)

$$= \int_{\mathbb{R}} \mathbb{P}[T_{ij}(n) \in \alpha + B_1] \mathbb{P}[T_{i'j'}(n) \in \alpha + B_2] f_{T_{12}}(\alpha) d\alpha$$
(422)

$$= \mathbb{P}[T_{ij}(n) - T_{12}(n) \in B_1] \mathbb{P}[T_{i'j'}(n) - T_{12}(n) \in B_2] + o(1),$$
(423)

i.e., $T_{ij}(n) - T_{12}(n)$ and $T_{i'j'}(n) - T_{12}(n)$ are asymptotically independent. This means that $\{T_{ij}(n) - T_{12}(n)\}$ are asymptotically pairwise independent. Hence, by using the same arguments to achieve (417), we have

$$\frac{1}{M_n(M_n-1)} \sum_{i \neq j} T_{ij}(n) - T_{12}(n) \xrightarrow{\text{(a.s.)}} 0, \tag{424}$$

as $n \to \infty$ (point-wise convergence). Then, there exist a subset \mathcal{B} such that $\mathbb{P}(\mathcal{B}) = 1$, and

$$\frac{1}{M_n(M_n-1)} \sum_{i \neq j} T_{ij}(n) - T_{12}(n) \to 0$$
(425)

on \mathcal{B} as $n \to \infty$.

Hence, we have $P_{e}(\mathcal{C}_{n}) = 1$, where $\mathcal{C} = \mathcal{A} \cap \mathcal{B}$. It follows that, for any $\varepsilon > 0$, on the set \mathcal{C} , as *n* sufficiently large (which depends on each realization of $\{T_{ij}(n)\}_{n=1}^{\infty}$'s), we have

$$-\varepsilon < \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) < \varepsilon,$$
(426)

$$-\varepsilon < \frac{1}{M_n(M_n-1)} \sum_{i \neq j} T_{ij}(n) - T_{12}(n) < \varepsilon.$$

$$(427)$$

The first step consists of showing that $\min \left\{ \mathbb{E}[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n))], \mathbb{E}[h(-T_{12}(n)) - h(-\min_{i \neq j} T_{ij}(n))] \right\} \rightarrow 0$ as $n \rightarrow \infty$. We carry out with two sub-steps, step 1a and step 1b.

1) Step 1a: To begin with, we prove that $\mathbb{E}[h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))] \to 0$ as $n \to \infty$ for all $h \in \{\mathcal{H} : a \ge 0\}$. We divide into different cases based on the value of a as following:

• Case 1: $\liminf_{n\to\infty} a > 0$.

Now, take an arbitrary small $\varepsilon > 0$ such that $\varepsilon < a$ as $n \to \infty$. From (426), we have $\min_{i \neq j} T_{ij}(n) < \varepsilon < a$. It follows that

$$h(\min_{i \neq j} T_{ij}(n)) = h\left(\frac{1}{M(M-1)} \sum_{i \neq j} T_{ij}(n)\right) = c$$
(428)

by the definition of \mathcal{H} .

Then, we have

$$h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) = \left[h(\min_{i \neq j} T_{ij}(n)) - h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) \right] + \left[h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) - h(T_{12}(n)) \right]$$
(429)

$$=h\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - h(T_{12}(n))$$
(430)

$$\leq \left| h \left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \right) - h(T_{12}(n)) \right|.$$
(431)

• Case 2: $a \ge 0$ and $\lim_{n \to \infty} a = 0$.

Then, if $\min_{i\neq j} T_{ij}(n) \leq a$ as $n \to \infty$, we have

$$h(\min_{i\neq j} T_{ij}(n)) - h\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j} T_{ij}(n)\right) = h(a) - h\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j} T_{ij}(n)\right)$$
(432)

$$\leq \left| a - \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \right| \tag{433}$$

$$\leq \max\left\{a, \frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right\}$$
(434)

$$\leq 2\varepsilon,$$
 (435)

where (433) follows from 1-Lipschitz property of h for all $h \in \mathcal{V}$.

On the other hand, if $a < \min_{i \neq j} T_{ij}(n) \le \frac{1}{M_n(M_n-1)} \sum_{i \neq j} T_{ij}(n) \le \varepsilon$ and $\liminf_{n \to \infty} c > 0$, we have

$$h(\min_{i \neq j} T_{ij}(n)) - h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) = \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) - \min_{i \neq j} T_{ij}(n)$$
(436)

$$\leq \varepsilon - a$$
 (437)

$$\leq \varepsilon.$$
 (438)

In addition, if $a < \min_{i \neq j} T_{ij}(n) \le \frac{1}{M_n(M_n-1)} \sum_{i \neq j} T_{ij}(n) \le \varepsilon$ and $\lim_{n \to \infty} c = 0$, we have

$$h(\min_{i \neq j} T_{ij}(n)) - h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) \le c$$
(439)

 $\leq \varepsilon.$ (440)

From (435), (438), and (440), it holds that

$$h(\min_{i \neq j} T_{ij}(n)) - h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) \le 2\varepsilon$$
(441)

for this case.

From (431) and (441), we have

$$h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \le 2\varepsilon + \left| h\left(\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n)\right) - h(T_{12}(n)) \right|$$
(442)

$$\leq 2\varepsilon + \left| \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) - T_{12}(n) \right|$$
 (443)

on \mathcal{C} .

From (443), on C, we have

$$\mathbb{E}\left[h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))\right] = \mathbb{E}\left[\left(h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))\right)\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\left(h(\min_{i\neq j} T_{ij}) - h(T_{12})\right)\mathbf{1}\{\mathcal{C}^c\}\right]$$
(444)

$$\leq \mathbb{E}\bigg[\big(h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))\big)\mathbf{1}\{\mathcal{C}\}\big] + c\mathbb{P}(\mathcal{C}^c)$$
(445)

$$= \mathbb{E}\bigg[\big(h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n))\big)\mathbf{1}\{\mathcal{C}\}\big].$$
(446)

Now, since $|(h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n)))\mathbf{1}\{\mathcal{C}\}| \leq c$, hence by the bounded convergence theorem [23], we have

$$\lim_{n \to \infty} \sup_{n \to \infty} \mathbb{E} \left[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \right]$$

$$\leq \mathbb{E} \left[\lim_{n \to \infty} h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \right]$$
(447)

$$= \mathbb{E}\left[\left(2\varepsilon + \left|\frac{1}{M_n(M_n - 1)}\sum_{i \neq j} T_{ij}(n) - T_{12}(n)\right|\right) \mathbf{1}\{\mathcal{C}\}\right] + c\mathbb{P}(\mathcal{C}^c)$$
(448)

$$\leq 2\varepsilon + \varepsilon \tag{449}$$

$$= 3\varepsilon \tag{450}$$

for any $\varepsilon > 0$, where (449) follows from (427).

From (450), by taking $\varepsilon \to 0$, we obtain

$$\limsup_{n \to \infty} \mathbb{E} \left[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \right] \le 0.$$
(451)

Since $h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n)) \ge 0$, by the fact that h is non-increasing for all $h \in \mathcal{V}$, from (451), we obtain

$$\lim_{n \to \infty} \mathbb{E} \Big[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \Big] = 0.$$
(452)

2) Step 1b: Next, we prove that $\mathbb{E}[h(-\min_{i\neq j} T_{ij}(n)) - h(-T_{12}(n))] \to 0$ as $n \to \infty$ for all $h \in \{\mathcal{H} : a < 0\}$. For all $h \in \mathcal{H}$, let $\tilde{h}(x) = h(-x)$ for all $x \in \mathbb{R}$. Then, we have

$$h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) = \tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} \{T_{ij}\}(n)).$$
(453)

Take an arbitrary $\varepsilon > 0$, under the condition (26), from (426) and (427), as $n \to \infty$, we obtain that

$$-\varepsilon < \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) < \varepsilon,, \qquad (454)$$

$$-\varepsilon < \frac{1}{M(M-1)} \sum_{i \neq j} T_{ij}(n) - T_{12}(n) < \varepsilon$$
(455)

on C where $P_{e}(C_n) = 1$.

Now, we show that $\mathbb{E}[\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))] \to 0$ as $n \to \infty$.

Similar to Step 1a, we divide into different cases based on a + c as following:

• Case 1: $\limsup_{n \to \infty} (a+c) < 0$.

Then, by taking $\varepsilon > 0$ small enough such that $\varepsilon < -\lim \sup_{n \to \infty} (a + c)$, from (454), as $n \to \infty$, we have

$$\min_{i \neq j} T_{ij}(n) \le \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \le \varepsilon < -(a + c).$$
(456)

Hence, it holds that

$$\tilde{h}\left(\max_{i\neq j} \tilde{T}_{ij}(n)\right) = \tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j} T_{ij}(n)\right) = 0.$$
(457)

It follows that on \mathcal{C} , as $n \to \infty$, we have

$$\tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(\min_{i\neq j}T_{ij}(n)) = 0.$$
(458)

• Case 2: $\lim_{n\to\infty} a + c = 0$.

Then, if $\min_{i\neq j} T_{ij}(n) \leq -(a+c)$, as $n \to \infty$, we have

$$\tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(\min_{i\neq j}\{T_{ij}(n)\}) = \tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(-(a+c))$$
(459)

$$\leq \left| \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) + (a + c) \right|$$
(460)

$$\leq \left| \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \right| + |a + c|$$
(461)

$$< 2\varepsilon.$$
 (462)

In addition, if $\min_{i \neq j} T_{ij}(n) \ge -(a+c)$ and $\limsup_{n \to \infty} a < 0$, we have $\lim_{n \to \infty} c = 0$. Hence,

$$\tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(\min_{i\neq j}T_{ij}(n)) \le \left|\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n) - \min_{i\neq j}T_{ij}(n)\right|$$
(463)

$$\leq \varepsilon + (a+c) \tag{464}$$

$$< 2\varepsilon.$$
 (465)

Finally, if if $\min_{i\neq j} T_{ij}(n) \ge -(a+c)$ and $\lim_{n\to\infty} a = 0$, we have

$$\tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(\min_{i\neq j}T_{ij}(n)) = \tilde{h}(-a) - \tilde{h}(\min_{i\neq j}T_{ij}(n))$$
(466)

$$\leq \left| -a - \min_{i \neq j} T_{ij}(n) \right| \tag{467}$$

$$\leq -a - (a + c) \tag{468}$$

$$< 2\varepsilon.$$
 (469)

From (462), (465), and (469), as $n \to \infty$, we have

_

$$\tilde{h}\left(\frac{1}{M_{n}(M_{n}-1)}\sum_{i\neq j}T_{ij}(n)\right) - \tilde{h}(\min_{i\neq j}\{T_{ij}(n)\}) < 2\varepsilon + \left|\tilde{h}(T_{12}(n)) - \tilde{h}\left(\frac{1}{M_{n}(M_{n}-1)}\sum_{i\neq j}T_{ij}(n)\right)\right|$$
(470)

on $\mathcal{C}.$

It follows from (458) and (470) that

$$\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n)) = \left[\tilde{h}(T_{12}(n)) - \tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i \neq j} T_{ij}(n)\right)\right] + \left[\tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i \neq j} T_{ij}(n)\right) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\right]$$
(471)

$$<\left|\tilde{h}(T_{12}(n)) - \tilde{h}\left(\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\right)\right| + 2\varepsilon$$
(472)

$$\leq 2\varepsilon + \left| T_{12}(n) - \frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \right|.$$
(473)

From (473), on C, we have

$$\mathbb{E}\left[\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\right] = \mathbb{E}\left[\left(\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\right)\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\left(\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\left(\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right) - \mathbb{E}\left[\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\left(\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right) - \mathbb{E}\left[\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(T_{12}(n) - \tilde{h}(\max_{i \neq j} T_{ij}(n)))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(T_{12}(n) - \tilde{h}(\max_{i \neq j} T_{ij}(n))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(\max_{i \neq j} T_{ij}(n) - \tilde{h}(\max_{i \neq j} T_{ij}(n))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(\max_{i \neq j} T_{ij}(n) - \tilde{h}(\max_{i \neq j} T_{ij}(n))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(\max_{i \neq j} T_{ij}(n) - \tilde{h}(\max_{i \neq j} T_{ij}(n))\mathbf{1}\{\mathcal{C}\}\right] + \mathbb{E}\left[\tilde{h}(\max_{i \neq j} T_{ij}(n) - \tilde{h}(\max_{i$$

$$\leq \mathbb{E}\left[\left(\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\right) \mathbf{1}\{\mathcal{C}\}\right] + c\mathbb{P}(\mathcal{C}^c)$$
(475)

$$= \mathbb{E}\left[\left(\tilde{h}(T_{12}(n) - \tilde{h}(\min_{i \neq j} T_{ij}(n))) \right) \mathbf{1}\{\mathcal{C}\} \right].$$
(476)

Now, since $|(\tilde{h}(\min_{i\neq j} T_{ij}(n)) - \tilde{h}(T_{12}(n)))\mathbf{1}\{\mathcal{C}\}| \leq c$, hence by the bounded convergence theorem [23], we have

$$\lim_{N \to \infty} \sup_{n \to \infty} \mathbb{E} \left[\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n)) - \right]$$

$$\leq \mathbb{E} \left[\lim_{n \to \infty} \tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n)) \right]$$
(477)

$$\leq \mathbb{E}\left[\lim_{n \to \infty} h(T_{12}(n)) - h(\min_{i \neq j} T_{ij}(n))\right]$$
(477)

$$= \mathbb{E}\left[\left(2\varepsilon + \left|\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n) - T_{12}(n)\right|\right)\mathbf{1}\{\mathcal{C}\}\right] + c\mathbb{P}(\mathcal{C}^c)$$
(478)

$$\leq 2\varepsilon + \varepsilon \tag{479}$$

$$= 3\varepsilon$$
 (480)

for any $\varepsilon > 0$, where (479) follows from (455).

From (480), by taking $\varepsilon \to 0$, we obtain

$$\limsup_{N \to \infty} \mathbb{E}\big[\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\big] \le 0.$$
(481)

Since $h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \ge 0$, from (481), we obtain

$$\lim_{n \to \infty} \mathbb{E}\left[\tilde{h}(T_{12}(n)) - \tilde{h}(\min_{i \neq j} T_{ij}(n))\right] = 0,$$
(482)

or

$$\lim_{n \to \infty} \mathbb{E} \left[h(-T_{12}(n)) - h(-\min_{i \neq j} T_{ij}(n)) \right] = 0.$$
(483)

From (452) and (483), we finally have

$$\lim_{n \to \infty} \min \left\{ \mathbb{E} \left[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \right], \mathbb{E} \left[h(-T_{12}(n)) - h(-\min_{i \neq j} T_{ij}(n)) \right] \right\} = 0$$
(484)

for all $h \in \mathcal{H}$.

3) Step 2: In this step, we show that $\lim_{n\to\infty} d_{W,\text{mod}}(\min_{i\neq j} T_{ij}, Z) = 0$. Indeed, from Lemma 20, we have

$$\sup_{h \in \mathcal{H}} \left| \mathbb{E} \left[f_h'(T_{ij}(n)) - T_{ij}(n) f_h(T_{ij}(n)) \right] \right| \le \frac{1}{n^{3/2}} \sum_{k=1}^n \mathbb{E} \left[|X_k|^3 \right] + \frac{\sqrt{2}}{n\sqrt{\pi}} \sqrt{\sum_{k=1}^n \mathbb{E} [X_k^4]}$$
(485)

$$= \frac{1}{\sqrt{n}} \mathbb{E}[|X_1|^3] + \frac{\sqrt{2}}{\sqrt{\pi n}} \sqrt{\mathbb{E}[X_1^4]}$$
(486)

where

$$X_{k} := \frac{-\sum_{x,x'} d_{\mathrm{B}}(x,x') \left(\mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\} - \mathbb{P}[(X_{ik}, X_{jk}) = (x,x')] \right)}{\sqrt{\operatorname{Var}\left(-\sum_{x,x'} d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right)}}, \quad \forall k \in [N].$$
(487)

Now, observe that

$$\operatorname{Var}(Z_{ij}(n)) = \operatorname{Var}\left(-\sum_{k=1}^{n}\sum_{x,x'}d_{\mathrm{B}}(x,x')\mathbf{1}\{(X_{ik},X_{jk}) = (x,x')\}\right)$$
(488)

$$=\sum_{k=1}^{n} \operatorname{Var}\left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik},X_{jk})=(x,x')\}\right)$$
(489)

$$= n \operatorname{Var}\left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right), \quad \forall k \in [n],$$
(490)

where (489) and (490) follow from the fact that (X_{ik}, X_{jk}) are i.i.d. given i, j.

Hence, we have

$$\operatorname{Var}\left(-\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right) = \mathbb{E}\left[\left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right)^{2}\right] - \left(\mathbb{E}\left[\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1}\{(X_{ik}, X_{jk}) = (x,x')\}\right]\right)^{2}$$
(491)

$$=\sum_{x,x'} d_{\mathrm{B}}^{2}(x,x') \mathbb{P}\big[(X_{ik},X_{jk})=(x,x')\big] - \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbb{P}\big[(X_{ik},X_{jk})=(x,x')\big]\right)^{2}$$
(492)

$$=\sum_{x,x'} d_{\rm B}^2(x,x')Q(x)Q(x') - \left(\sum_{x,x'} d_{\rm B}(x,x')Q(x)Q(x')\right)^2 = L_2.$$
(493)

In addition, we have

$$\mathbb{E}\left[\left|\sum_{x,x'} d_{\mathrm{B}}(x,x') \left(\mathbf{1}\{(X_{ik},X_{jk})=(x,x')\} - \mathbb{P}\left[(X_{ik},X_{jk})=(x,x')\right]\right)\right|^{3}\right]$$
(494)

$$\leq 4 \left(\mathbb{E} \left[\left| \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbf{1} \{ (X_{ik}, X_{jk}) = (x,x') \} \right|^3 \right] + \left| \sum_{x,x'} d_{\mathrm{B}}(x,x') \mathbb{P} \left[(X_{ik}, X_{jk}) = (x,x') \right] \right|^3 \right)$$
(495)

$$=4\left[\sum_{x,x'}d_{\rm B}^3(x,x')Q(x)Q(x') + \left(\sum_{x,x'}d_{\rm B}(x,x')Q(x)Q(x')\right)^3\right] = L_3,\tag{496}$$

where (495) follows from $(a+b)^3 \le 4(|a|^3+|b^3|).$

Similarly, we have

$$\mathbb{E}\left[\left|\sum_{x,x'} d_{\mathrm{B}}(x,x') \left(\mathbf{1}\{(X_{ik},X_{jk})=(x,x')\} - \mathbb{P}\left[(X_{ik},X_{jk})=(x,x')\right]\right)\right|^{4}\right]$$
(497)

$$\leq 8 \left[\sum_{x,x'} d_{\mathrm{B}}^4(x,x') Q(x) Q(x') + \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') \right)^4 \right] = L_4, \tag{498}$$

where we use $(a + b)^4 \le 8(a^4 + b^4)$ in (498).

Hence, from (486), (493), (496), and (498), we obtain

$$\sup_{h \in \mathcal{H}} \left| \mathbb{E} \left[f'_h(T_{ij}(n)) - T_{ij}(n) f_h(T_{ij}(n)) \right] \right| \le \frac{1}{\sqrt{n}} \left(\frac{L_3}{L_2^{3/2}} \right) + \sqrt{\frac{2}{\pi n}} \frac{L_4}{L_2^2}, \qquad \forall i \neq j.$$
(499)

Similarly, we also have

$$\sup_{h \in \mathcal{H}} \left| \mathbb{E} \left[f'_h(-T_{ij}(n)) + T_{ij}(n) f_h(-T_{ij}(n)) \right] \right| \le \frac{1}{\sqrt{n}} \left(\frac{L_3}{L_2^{3/2}} \right) + \sqrt{\frac{2}{\pi n}} \frac{L_4}{L_2^2}, \qquad \forall i \neq j.$$
(500)

52

Since $T_{ij}(n)$'s (for $i \neq j$) are identically distributed by the random codebook generation, it follows from Lemma 19 and (499) that for any $x \in \mathbb{R}$,

$$d_{W,\text{mod}}(\min_{i\neq j} T_{ij}, Z) \leq \max \left\{ \sup_{h\in\mathcal{H}} \left| \mathbb{E} \left[f'_{h}(T_{12}(n)) - T_{12}(n)f_{h}(T_{12}(n)) \right] \right|, \sup_{h\in\mathcal{H}} \left| \mathbb{E} \left[f'_{h}(-T_{12}(n)) + T_{12}(n)f_{h}(-T_{12}(n)) \right] \right| \right\} + \sup_{h\in\mathcal{H}} \min \left\{ \mathbb{E} [h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))], \mathbb{E} [h(-T_{12}(n)) - h(-\min_{i\neq j} T_{ij}(n))] \right\}$$
(501)
$$\leq \frac{1}{\sqrt{n}} \left(\frac{L_{3}}{L_{2}^{3/2}} \right) + \sqrt{\frac{2}{\pi n}} \frac{L_{4}}{L_{2}^{2}} + \sup_{h\in\mathcal{H}} \min \left\{ \mathbb{E} [h(\min_{i\neq j} T_{ij}(n)) - h(T_{12}(n))], \mathbb{E} [h(-T_{12}(n)) - h(-\min_{i\neq j} T_{ij}(n))] \right\} \to 0,$$
(502)

where (502) follows from (484).

4) Step 3: In the third step, we whow that $\lim_{n\to\infty} |\mathbb{P}(\min_{i\neq j} T_{ij}(n) \le x) - \mathbb{P}(\min_{i\neq j} T_{ij}(n) \ge -x)| = 0$ for all $x \in \mathbb{R}$ and x is a continuous point of the limiting distribution of $\min_{i\neq j} T_{ij}(n)$.

First, by the i.i.d. random codebook generation, observe that $T_{12}(n), T_{23}(n), \dots, T_{(M-1)M}(n)$ are i.i.d.. For any $x \in \mathbb{R}$, as $n, M_n \to \infty$, we have

$$\mathbb{P}\Big[\min_{i \neq j} T_{ij}(n) \ge x\Big] \le \mathbb{P}\Big[\min\{T_{12}(n), T_{34}(n), \cdots, T_{(M-1)M}\}(n) \ge x\Big]$$
(503)

$$= \mathbb{P}[T_{12}(n) \ge x] \mathbb{P}[T_{34}(n) \ge x] \cdots \mathbb{P}[T_{(M-1)M}(n) \ge x]$$
(504)

$$= \left(\mathbb{P}[T_{12}(n) \ge x]\right)^{\lfloor M/2 \rfloor}$$
(505)

$$\leq \left(Q(x) + o(1)\right)^{\lfloor M/2 \rfloor} \tag{506}$$

$$\leq \left(\frac{Q(\limsup_{N \to \infty} x)}{2}\right)^{\lfloor M/2 \rfloor}$$
(507)

$$\rightarrow 0,$$
 (508)

where (506) follows from the fact that $T_{12}(n)$ is the sum of n i.i.d. terms, so we can apply the CLT, and (507) follows from the fact that as $n \to \infty$, $Q(x) \ge Q(\limsup_{n \to \infty} x) > Q(\limsup_{n \to \infty} x)/2$ as n sufficiently large, and (508) follows from $M_n \to \infty$ under the condition (26).

Now, we divide into different cases depending on x:

• Case 1: $x \in \mathbb{R}$ and $\liminf_{n \to \infty} x < 0$.

Then, there exists an $\varepsilon > 0$ such that $x < -\varepsilon$ as n sufficiently large. Hence, under the condition (26), we have

$$\mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge -x) \le \mathbb{P}\left[\frac{1}{M_n(M_n - 1)} \sum_{i \neq j} T_{ij}(n) \ge -x\right]$$
(509)

$$\leq \mathbb{P}\left[\frac{1}{M_n(M_n-1)}\sum_{i\neq j}T_{ij}(n)\geq \varepsilon\right]$$
(510)

$$\rightarrow 0,$$
 (511)

where (511) follows from (417). It follows from (508) and (511) that

$$\lim_{n \to \infty} \mathbb{P}\left[\min_{i \neq j} T_{ij}(n) \ge x\right] - \mathbb{P}\left[\min_{i \neq j} T_{ij}(n) \ge -x\right] = 0,$$
(512)

which leads to

$$\lim_{n \to \infty} \left| \mathbb{P} \left[\min_{i \neq j} T_{ij}(n) \ge x \right] - \mathbb{P} \left[\min_{i \neq j} T_{ij}(n) \ge -x \right] \right| = 0.$$
(513)

• Case $\lim_{n\to\infty} x = 0$.

From (452), we have

$$\lim_{n \to \infty} \mathbb{E} \left[h(\min_{i \neq j} T_{ij}(n)) - h(T_{12}(n)) \right] = 0,$$
(514)

for all $h \in \mathcal{H}$: $\liminf_{n \to \infty} a \ge 0$.

Now, for any fixed constants (u, v) such that 0 < u < v, define $h_u(t) = \mathbf{1}\{t \le x\}$ and $h_v(t) = \mathbf{1}\{t \le x\}$ for all $t \in R$. Then, we have

$$\mathbb{P}\left[\min_{i\neq j} T_{ij}(n) \notin (u, v]\right) \\
= \mathbb{P}\left[\min_{i\neq j} T_{ij}(n) > v\right] + \mathbb{P}\left[\min_{i\neq j} T_{ij}(n) \le u\right] \\
\le \left|\mathbb{P}\left[\min_{i\neq j} T_{ij}(n) > v\right] - \mathbb{P}(T_{12}(n) > v)\right] + \left[\mathbb{P}\left[\min_{i\neq j} T_{ij}(n) \le u\right] - \mathbb{P}(T_{12}(n) \le u)\right]$$
(515)

$$\mathbb{E}\left[\mathbb{P}\left[\min_{i\neq j} T_{ij}(n) > v\right] - \mathbb{P}(T_{12}(n) > v)\right] + \left[\mathbb{P}\left[\min_{i\neq j} T_{ij}(n) \le u\right] - \mathbb{P}(T_{12}(n) \le u)\right]$$

$$+ \mathbb{P}(T_{12}(n) > v) + \mathbb{P}(T_{12}(n) \le u)$$
(516)

$$= \left| \mathbb{P}\left[\min_{i \neq j} T_{ij}(n) \le v\right] - \mathbb{P}(T_{12}(n) \le v) \right] + \left[\mathbb{P}\left[\min_{i \neq j} T_{ij}(n) \le u\right] - \mathbb{P}(T_{12}(n) \le u) \right]$$
$$+ \mathbb{P}(T_{12}(n) > v) + \mathbb{P}(T_{12}(n) \le u)$$
(517)

$$= \left| \mathbb{E} \Big[h_v(\min_{i \neq j} T_{ij}(n)) - h_v(T_{12}(n)) \Big] + \mathbb{E} \Big[h_u(\min_{i \neq j} T_{ij}(n)) - h_u(T_{12}(n)) \Big] \right| \\ + \mathbb{P}(T_{12}(n) > v) + \mathbb{P}(T_{12}(n) \le u)$$
(518)

$$= o(1) + \mathbb{P}(T_{12}(n) > v) + \mathbb{P}(T_{12}(n) \le u)$$
(519)

$$= o(1) + Q(v) + Q(u), \tag{520}$$

where (519) follows from (514), and (520) follows from $T_{12}(n) \xrightarrow{(d)} \mathcal{N}(0,1)$ by the CLT (it is a sum of i.i.d. random variables with finite variance). Hence, for any $\varepsilon > 0$, by letting u and v sufficiently large constants such that $Q(u) + Q(v) < \varepsilon/2$. Then, for N sufficiently large, we have

$$\mathbb{P}\big[\min_{i\neq j} T_{ij}(n) \in (u,v]\big) > 1 - \varepsilon.$$
(521)

This means that the probability distribution of $\min_{i \neq j} T_{ij}(n)$ is tight. Then, [23], the distribution of $\min_{i \neq j} T_{ij}(n)$ is convergent to some distribution of a random variable Y. Hence, by Lemma 17, we have

$$\limsup_{n \to \infty} \left| \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le x) - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge -x) \right| \le 2(8\pi)^{-1/4} \limsup_{n \to \infty} \sqrt{d_{W, \text{mod}}(\min_{i \neq j} T_{ij}(n), Z)} = 0,$$
(522)

where (522) follows from Step 3.

• For any $x \ge 0$, then $-x \le 0$. Hence, from two cases above, we have

$$\lim_{n \to \infty} \left| \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le -x) - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge x) \right| = 0.$$
(523)

However, for any continuous point x of the limiting distribution of $\min_{i\neq j} T_{ij}(n)$, we have

$$\lim_{n \to \infty} \left| \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le -x) - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge x) \right|$$
(524)

$$= \lim_{n \to \infty} \left| (1 - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge -x)) - (1 - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le x)) \right|$$
(525)

$$= \lim_{n \to \infty} \left| \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le x) - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge -x) \right|.$$
(526)

From (523) and (526), we obtain

$$\lim_{n \to \infty} \left| \mathbb{P}(\min_{i \neq j} T_{ij}(n) \le x) - \mathbb{P}(\min_{i \neq j} T_{ij}(n) \ge -x) \right| = 0.$$
(527)

5) Step 4: The last step proves that $T_n = \frac{V_n - \mathbb{E}[V_n]}{\sqrt{\operatorname{Var}(V_n)}} \xrightarrow{(d)} \mathcal{N}(0, 1)$. From Lemma (502), Lemma 17, and Step 3, we have

$$\mathbb{P}\big[\min_{i \neq j} T_{ij}(n) \le x\big] - \mathbb{P}\big[Z \le x\big] \to 0$$
(528)

as $n \to \infty$ for any continuous point $x \in \mathbb{R}$ of the limiting distribution of $\min_{i \neq j} T_{ij}(n)$ (point-wise convergence), or

$$\min_{i \neq j} T_{ij}(n) \xrightarrow{(d)} Z = \mathcal{N}(0, 1).$$
(529)

Now, observe that

$$T_n = \min_{i \neq j} \tilde{T}_{ij}(n), \tag{530}$$

where

$$\tilde{T}_{ij}(n) = \frac{Z_{ij}(n) - \mathbb{E}[V_n]}{\sqrt{\operatorname{Var}(V_n)}}.$$
(531)

Hence, by using Lemma 1 and (by using the same arguments to achieve (292) from (529) as in the proof of Theorem 5), we obtain

$$T_n \xrightarrow{(d)} \mathcal{N}(0,1).$$
 (532)

Finally, from (403) and (532), by applying Slutsky's theorem, we have

$$\frac{\frac{-\log P_{e}(\mathcal{C}_{n})}{n} - \mathbb{E}[\frac{-\log P_{e}(\mathcal{C}_{n})}{n}]}{\sqrt{\operatorname{Var}\left(\frac{-\log P_{e}(\mathcal{C}_{n})}{n}\right)}} \xrightarrow{(d)} \mathcal{N}(0,1).$$
(533)

G. Proof of Theorem 7

Consider first the case $0 \le R < C$. Since the random variable $P_e(\mathcal{C}_n)$ takes values in [0, 1], we have that

$$\operatorname{Var}[P_{\mathrm{e}}(\mathcal{C}_{n})] = \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})^{2}] - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]^{2}$$
(534)

$$\leq \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2] \tag{535}$$

$$\leq \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)] \to 0 \tag{536}$$

56

where (536) follows from the assumption that $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \to 0$ for $0 \leq R < C$. Applying Chebyshev's inequality we have that

$$\mathbb{P}\Big[|P_{e}(\mathcal{C}_{n}) - \mathbb{E}[P_{e}(\mathcal{C}_{n})]| \ge \delta\Big] \le \frac{\operatorname{Var}[P_{e}(\mathcal{C}_{n})]}{\delta^{2}}$$
(537)

$$\leq \frac{\mathbb{E}[P_{\rm e}(\mathcal{C}_n)]}{\delta^2} \to 0 \tag{538}$$

where (538) follows from (536) and is valid for any given $\delta > 0$.

Now let us consider the case R > C. The following hold:

$$\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)] \to 1 \tag{539}$$

$$\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2 \to 1 \tag{540}$$

$$\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2] \ge \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2 \tag{541}$$

$$\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2] \to 1 \tag{542}$$

$$\operatorname{Var}[P_{\mathbf{e}}(\mathcal{C}_n)] \to 0 \tag{543}$$

where (539) follows from the theorem assumption, (540) follows from (539), (541) follows from Jensen's inequality, (542) follows from (540) and (541) and the fact that $\mathbb{E}[P_{e}(\mathcal{C}_{n})^{2}] \leq 1$, while (543) follows from (542) and (540) and the additivity of limits.

Finally, using Chebyshev's inequality again we find that

$$\mathbb{P}\Big[|P_{\mathbf{e}}(\mathcal{C}_n) - \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]| \ge \delta\Big] \le \frac{\operatorname{Var}[P_{\mathbf{e}}(\mathcal{C}_n)]}{\delta^2} \to 0$$
(544)

for any $\delta > 0$.

H. Proof of Theorem 8

First, by the condition (29), we observe that

$$\frac{\operatorname{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}{\left(\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right)^{2}} = \frac{\mathbb{E}[P_{\mathrm{e}}^{2}(\mathcal{C}_{n})] - \left(\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right)^{2}}{\left(\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right)^{2}}$$
(545)

$$= \frac{\mathbb{E}[P_{e}^{2}(\mathcal{C}_{n})]}{\left(\mathbb{E}[P_{e}(\mathcal{C}_{n})]\right)^{2}} - 1$$
(546)

$$\rightarrow 0.$$
 (547)

On the other hand, we know that $\frac{\operatorname{Var}(P_{\mathrm{e}}(\mathcal{C}_n))}{\left(\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)]\right)^2} \to \infty$ if $E_{\mathrm{rce}}(R) < E_{\mathrm{trc}}(R)$. Hence, from (547), we must have

$$E_{\rm trc}(R) = E_{\rm rce}(R). \tag{548}$$

Now, for any $\varepsilon > 0$, we have

$$\mathbb{P}\left[\left|-\frac{\log P_{\mathrm{e}}(\mathcal{C}_{n})}{N}-E_{\mathrm{trc}}(R)\right|>\varepsilon\right]=\mathbb{P}\left[\left\{P_{\mathrm{e}}(\mathcal{C}_{n})<2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right\}\cup\left\{P_{\mathrm{e}}(\mathcal{C}_{n})>2^{-n(E_{\mathrm{trc}}(R)-\varepsilon)}\right\}\right]$$
(549)

=

$$\mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] + \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) > 2^{-n(E_{\mathrm{trc}}(R)-\varepsilon)}\right]$$
(550)

$$\leq \mathbb{P}\bigg[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\bigg] + 2^{n(E_{\mathrm{trc}}(R)-\varepsilon)}\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]$$
(551)

$$\leq \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] + 2^{n(E_{\mathrm{trc}}(R)-\varepsilon)}2^{-n(E_{\mathrm{rce}}(R)-\varepsilon/2)}$$
(552)

$$= \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] + 2^{-n\varepsilon/2}$$
(553)

for *n* sufficiently large, where (551) follows from Markov's inequality, and (552) follows from $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \doteq 2^{-nE_{rce}(R)}$, so $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \leq 2^{-n(E_{rce}(R)-\varepsilon/2)}$ for *n* sufficiently large.

Now, observe that

$$\mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] = \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})] < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)} - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right]$$
(554)

$$= \mathbb{P}\bigg[-\left(P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right) > \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})] - 2^{-n(E_{\mathrm{trc}}(R) + \varepsilon)}\bigg].$$
 (555)

Now, since $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \doteq 2^{-nE_{rce}(R)} = 2^{-nE_{trc}(R)}$ by (548), so $\mathbb{E}[P_{e}(\mathcal{C}_{n})] - 2^{-n(E_{trc}(R)+\varepsilon)} > 0$ for *n* sufficiently large. It follows from (555) that

$$\mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] \leq \mathbb{P}\left[\left|P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right| > \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})] - 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right]$$
(556)

$$\approx \frac{\operatorname{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}{\left(\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]\right)^{2}}$$
(557)

$$\rightarrow 0,$$
 (558)

where (557) follows Markov's inequality and the fact that $\mathbb{E}[P_{e}(\mathcal{C}_{n})] - 2^{-n(E_{tre}(R)+\varepsilon)} = \Theta(\mathbb{E}[P_{e}(\mathcal{C}_{n})])$, and (558) follows from (547).

From (553) and (558), we obtain

$$\mathbb{P}\left[\left|-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) - E_{\rm trc}(R)\right| > \varepsilon\right] \to 0,$$
(559)

or equivalently,

$$-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) \xrightarrow{({\rm p})} E_{\rm trc}(R).$$
(560)

I. Proof of Theorem 9

From [15, Th. 1] and from (111), for n sufficiently large we have:

$$\mathbb{P}\left[P_{\mathbf{e}}(\mathcal{C}_{n}) \geq \gamma_{n}^{\rho} \min_{\rho \in [1,\infty)} E[P_{e}(\mathcal{C}_{n})^{\frac{1}{\rho}}]^{\rho}\right] = \mathbb{P}\left[P_{\mathbf{e}}(\mathcal{C}_{n}) \geq 2^{-n(E_{\mathrm{trc}}(R) - \epsilon_{n})}\right]$$
(561)

$$\leq \frac{1}{\gamma_n} \tag{562}$$

where $\gamma_n \to \infty$, $\frac{\log \gamma_n}{n} \to 0$ and $\epsilon_n \to 0$. On the other hand, the Paley-Zygmund inequality [26] implies that, for large enough n:

$$\mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) \geq \delta_{n} \mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n})\right]\right] = \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) \geq 2^{-n(E_{\mathrm{rce}}(R) + \epsilon_{n}')}\right]$$
(563)

$$\geq (1 - \delta_n)^2 \frac{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2}{E[P_{\mathbf{e}}(\mathcal{C}_n)^2]}$$
(564)

where we choose a sequence δ_n that goes to zero subexponentially, i.e., $\epsilon'_n \to 0$ and $0 < \delta_n < 1 \ \forall n$. Let n_0 be such that $\Delta E > \epsilon_n$, $\forall n > n_0$. Note that such an n_0 must exist from the definition of limit for ϵ_n . Now consider the following chain of inequalities for a large enough $n, n > n_0$:

$$2^{-n(E_{\rm trc}(R)-\epsilon_n)} = 2^{-n(E_{\rm rce}(R)+\Delta E-\epsilon_n)}$$
(565)

$$\leq 2^{-n(E_{\rm rce}(R) + \Delta E - \epsilon_{n_0})} \tag{566}$$

$$<2^{-n(E_{\rm rce}(R)+\epsilon_n')}\tag{567}$$

where (565) is from the theorem statement, (566) is valid from a certain n onwards from the definition of limit for ϵ_n , while (567) is because $\Delta E - \epsilon_{n_0}$ is a positive constant and, for large enough n, $\epsilon'_n < \Delta E - \epsilon_{n_0}$. Now, using (567), (561) and (563) we have:

$$(1 - \delta_n) \frac{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2}{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2]} \le \mathbb{P}\left[P_{\mathbf{e}}(\mathcal{C}_n) \ge \delta_n \mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]\right]$$
(568)

$$= \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) \ge 2^{-n(E_{\mathrm{rce}}(R) + \epsilon'_{n})}\right]$$
(569)

$$\leq \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) \geq 2^{-n(E_{\mathrm{trc}}(R) - \epsilon_{n})}\right]$$
(570)

$$= \mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) \geq \gamma_{n}^{\rho} \min_{\rho \in [1,\infty)} \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})^{\frac{1}{\rho}}]^{\rho}\right]$$
(571)

$$\leq \frac{1}{\gamma_n} \tag{572}$$

where(570) follows from (567). Finally, notice that, by definition, the following inequalities hold:

$$(1 - \delta_n) \to 1$$

 $\frac{1}{\gamma_n} \to 0$

that imply:

$$\frac{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)]^2}{\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)^2]} \to 0$$

J. Proof of Theorem 10

Under the condition $E_{\rm trc}(R) > E_{\rm rce}(R)$, it holds by Theorem 9

$$\frac{\mathbb{E}[P_{e}(\mathcal{C}_{n})]}{\sqrt{\operatorname{Var}P_{e}(\mathcal{C}_{n})}} \to 0.$$
(573)

Now, assume that

$$\frac{P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0, 1).$$
(574)

Then, from (573) and (574) and Slutsky's theorem [23], it holds that

$$\frac{P_{\mathrm{e}}(\mathcal{C}_{n})}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0,1), \tag{575}$$

which is a contradiction since the LHS of (575) is a non-negative random variable.

K. Proof of Theorem 11

First, if $\liminf_{n\to\infty} \frac{\mathbb{E}[P_e^2(\mathcal{C}_n)]}{\left(\mathbb{E}[P_e(\mathcal{C}_n)]\right)^2} > 1$, then it holds that

$$\nu = \limsup_{n \to \infty} \frac{\mathbb{E}[P_{e}(\mathcal{C}_{n})]}{\sqrt{\operatorname{Var}(P_{e}(\mathcal{C}_{n}))}} < \infty.$$
(576)

Then, for n sufficiently large, we have

$$\frac{P_{\rm e}(\mathcal{C}_n) - \mathbb{E}[P_{\rm e}(\mathcal{C}_n)]}{\sqrt{\operatorname{Var}(P_{\rm e}(\mathcal{C}_n))}} \ge \frac{P_{\rm e}(\mathcal{C}_n)}{\sqrt{\operatorname{Var}(P_{\rm e}(\mathcal{C}_n))}} - \nu$$
(577)

$$\geq -\nu,$$
 (578)

hence

$$\frac{P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}} \xrightarrow{(\mathrm{d})} \mathcal{N}(0, 1).$$
(579)

Hence, the condition (33) implies that

$$\frac{\mathbb{E}[P_{e}^{2}(\mathcal{C}_{n})]}{\left(\mathbb{E}[P_{e}(\mathcal{C}_{n})]\right)^{2}} \to 1,$$
(580)

which leads to

$$\frac{\mathbb{E}[P_{e}(\mathcal{C}_{n})]}{\sqrt{\operatorname{Var}(P_{e}(\mathcal{C}_{n}))}} \to \infty.$$
(581)

Now, for any $\varepsilon > 0$, we have

$$\mathbb{P}\left[\left|-\frac{1}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) - E_{\mathrm{trc}}(R)\right| > \varepsilon\right]$$
$$= \mathbb{P}\left[\left\{P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right\} \cup \left\{P_{\mathrm{e}}(\mathcal{C}_{n}) > 2^{-n(E_{\mathrm{trc}}(R)-\varepsilon)}\right\}\right]$$
(582)

$$\leq \mathbb{P}\bigg[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\bigg] + \mathbb{P}\bigg[P_{\mathrm{e}}(\mathcal{C}_{n}) > 2^{-n(E_{\mathrm{trc}}(R)-\varepsilon)}\bigg].$$
(583)

Furthermore, under the condition $\frac{\mathbb{E}[P_{e}(C_{n})]}{\sqrt{\operatorname{Var}(P_{e}(C_{n}))}} \to \infty$, we must have

$$E_{\rm trc}(R) = E_{\rm rce}(R). \tag{584}$$

Observe that

$$\mathbb{P}\bigg[P_{\mathrm{e}}(\mathcal{C}_{n}) > 2^{-n(E_{\mathrm{trc}}(R)-\varepsilon)}\bigg] \le 2^{n(E_{\mathrm{trc}}(R)-\varepsilon)}\mathbb{E}\big[P_{\mathrm{e}}(\mathcal{C}_{n})\big]$$
(585)

$$\doteq 2^{n(E_{\rm trc}(R)-\varepsilon)} 2^{-nE_{\rm rce}(R)} \tag{586}$$

$$=2^{-n\varepsilon}.$$
(587)

On the other hand, from (33), for $Z \sim \mathcal{N}(0, 1)$, we have

$$\mathbb{P}\left[P_{\mathrm{e}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}\right] \leq \mathbb{P}\left[\frac{P_{\mathrm{e}}(\mathcal{C}_{n}) - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}} \leq \frac{2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)} - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}}\right]$$
(588)

$$= \mathbb{P}\left[Z \le \frac{2^{-n(E_{\text{trc}}(R)+\varepsilon)} - \mathbb{E}[P_{\text{e}}(\mathcal{C}_n)]}{\sqrt{\text{Var}(P_{\text{e}}(\mathcal{C}_n))}}\right] + o_n(1),$$
(589)

where (589) follows from the condition (33).

Now, since $2^{-n(E_{\text{trc}}(R)+\varepsilon)} - \mathbb{E}[P_{e}(\mathcal{C}_{n})] < 0$ for n sufficiently large since $\mathbb{E}[P_{e}(\mathcal{C}_{n})] \doteq 2^{-nE_{\text{rce}}(R)}$ and $E_{\text{trc}}(R) \ge E_{\text{rce}}(R)$, it holds that

$$\mathbb{P}\left[Z \le \frac{2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)} - \mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})]}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}}\right] = Q\left(\frac{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_{n})] - 2^{-n(E_{\mathrm{trc}}(R)+\varepsilon)}}{\sqrt{\mathrm{Var}(P_{\mathrm{e}}(\mathcal{C}_{n}))}}\right) \to 0$$
(590)

as $n \to \infty$ since

$$\frac{\mathbb{E}[P_{e}(\mathcal{C}_{n})] - 2^{-n(E_{trc}(R) + \varepsilon)}}{\sqrt{\operatorname{Var}(P_{e}(\mathcal{C}_{n}))}} \to \infty.$$
(591)

From (583), (587), and (591), we obtain

$$\mathbb{P}\left[\left|-\frac{1}{n}\log P_{\rm e}(\mathcal{C}_n) - E_{\rm trc}(R)\right| > \varepsilon\right] \to 0$$
(592)

or (34) holds.

L. Proof of Theorem 12

We first introduce a result which is developed in [25] for sum of random variables with local dependence.

Definition 3: We say that a collection of random variables (Y_1, Y_2, \dots, Y_n) has dependency neighbourhoods $\mathcal{N}_i \subset \{1, 2, \dots, n\}, i = 1, 2, \dots, n$, if $i \in \mathcal{N}_i$ and Y_i is independent of $\{Y_j\}_{j \notin \mathcal{N}_i}$.

Lemma 21: [25, Th. 3.6] Let Y_1, Y_2, \dots, Y_n be random variables such that $\mathbb{E}[Y_i^4] < \infty$, $\mathbb{E}[Y_i = 0]$, $\sigma^2 = Var(\sum_{i=1}^n Y_i)$, and define $T = \sum_{i=1}^n Y_i/\sigma$. Let the collection (Y_1, Y_2, \dots, Y_n) have dependency neighborhoods $\mathcal{N}_i, i = 1, 2, \dots, n$, and also define $D = \max_{1 \le i \le n} |\mathcal{N}_i|$. Then for Z a standard normal random variable,

$$d_W(T,Z) \le \frac{D^2}{\sigma^3} \sum_{i=1}^n \mathbb{E}|Y_i|^3 + \frac{\sqrt{28}D^{3/2}}{\sqrt{\pi}\sigma^2} \sqrt{\sum_{i=1}^n \mathbb{E}[Y_i^4]}.$$
(593)

Observe that

$$P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) - \mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)] = \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} Y_{ij}$$
(594)

$$= \frac{2}{M_n} \sum_{i=1}^{M_n} \sum_{i < j \le M_n} Y_{ij},$$
(595)

where

$$Y_{ij} = \mathbb{P}\big[\{\boldsymbol{X}_i \to \boldsymbol{X}_j\}\big] - \mathbb{E}\big[\mathbb{P}\big[\{\boldsymbol{X}_i \to \boldsymbol{X}_j\}\big]\big], \qquad \forall i, j \in [M_n] \times [M_n].$$
(596)

For i.i.d. random coding ensembles, $\{Y_{ij}\}_{1 \le i < j \le M_n}$ are pairwise independent and identically distributed by the symmetry of the random codebook ensemble. Hence, we have

$$\sigma^2 = \operatorname{Var}\left(\sum_{i=1}^{M_n} \sum_{j \neq i} Y_{ij}\right)$$
(597)

$$= 4 \operatorname{Var}\left(\sum_{1 \le i < j \le M_n} Y_{ij}\right)$$
(598)

$$=4\sum_{1\le i< j\le M_n} \operatorname{Var}(Y_{ij})$$
(599)

$$= 2M_n(M_n - 1)\gamma^2.$$
 (600)

In addition, it is easy to see that $D \leq 2(M_n - 1)$. Hence, by Lemma 21, we have

$$d_{W}\left(\frac{P_{e}^{ub}(\mathcal{C}_{n}) - \mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n})]}{\sqrt{\operatorname{Var}(P_{e}^{ub}(\mathcal{C}_{n})}}, Z\right) \leq \frac{4(M_{n}-1)^{2}}{(2M_{n}(M_{n}-1)\gamma^{2})^{3/2}} \left(\frac{M_{n}(M_{n}-1)}{2}\right) \mathbb{E}[|Y_{12}|^{3}] + \frac{\sqrt{28}(2(M_{n}-1))^{3/2}}{\sqrt{\pi}(2M_{n}(M_{n}-1)\gamma^{2})} \sqrt{\frac{M_{n}(M_{n}-1)}{2}} \mathbb{E}[|Y_{12}|^{4}]$$
(601)

$$\leq \frac{M_n}{\gamma^3} \mathbb{E}[|Y_{12}|^3] + \sqrt{\frac{28}{\pi}} \sqrt{M_n \mathbb{E}[|Y_{12}|^4]},\tag{602}$$

which tends to zero if both (35) and (36) happen simultaneously.

M. Proof of Theorem 13

We first state two auxiliary lemmas.

Lemma 22: If $Z \sim \mathcal{N}(0, 1)$, then for any random variable T, it holds that

$$d_K(T,Z) \le 2(8\pi)^{-1/4} \sqrt{\tilde{d}_{W,\text{mod}}(T,Z)}.$$
 (603)

Proof: The proof is similar to the first part of the proof of Lemma 17 in Appendix O, so we omit this proof.

Lemma 23: If T is a random variable such that $\mathbb{E}[T] = 0$ and $\operatorname{Var}(T) = 1$, and Z has the standard normal distribution, then

$$d_K(T,Z) < 14(8\pi)^{-1/4} \sqrt{\mathbb{E}[|T|] + \mathbb{E}[|T^2 - 1|]}.$$
(604)

Proof: Appendix Q.

Theorem 13 is a direct application of Lemma 23 by setting $T = g_n(P_e(C_n))$, gives a criterion for the convergence in distribution of the error exponent and any function of the error probability, in general.

APPENDIX

A. Proof of Lemma 1

Since $U_n \xrightarrow{(d)} U$, by Skorokhod's representation theorem [23], there exists a probability space (Ω, \mathcal{F}, P) and two random variables V_n and V such that $V_n \sim U_n$ and $V \sim U$ such that $V_n \xrightarrow{(a.s.)} V$ on (Ω, \mathcal{F}, P) . Now, for any fixed $\varepsilon \in (0, 1)$, observe that

$$\mathbb{E}_{P}[|V_{n}|^{1+\varepsilon}] \le \left(\mathbb{E}_{P}[|V_{n}|^{2}]\right)^{(1+\varepsilon)/2} \tag{605}$$

$$= 1 < \infty, \tag{606}$$

where (605) follows from the concavity of the function $f(x) = x^{(1+\varepsilon)/2}$ for any $\varepsilon \in (0, 1)$, and (606) follows from $\mathbb{E}_P[V_n^2] = \mathbb{E}[U_n^2] = 1$. From (606), it follows that V_n is uniformly integrable on (Ω, \mathcal{F}, P) [23]. Hence, we have

$$\mathbb{E}[U] = \mathbb{E}_P[V] \tag{607}$$

$$=\lim_{n\to\infty}\mathbb{E}_P[V_n]\tag{608}$$

$$=\lim_{n\to\infty}\mathbb{E}[U_n]\tag{609}$$

$$= 0.$$
 (610)

On the other hand, for any fixed n, we have

$$|V_n|^2 \mathbf{1}\{|V_n| > \alpha\} \le |V_n|^2, \tag{611}$$

which satisfies

$$\mathbb{E}_{P}[|V_{n}|^{2}] = \mathbb{E}[|U_{n}|^{2}] = 1.$$
(612)

Hence, by the dominated convergence theorem [30], we have

$$\lim_{\alpha \to \infty} \mathbb{E}_P \left[|V_n|^2 \mathbf{1}\{|V_n| > \alpha\} \right] = \mathbb{E}_P \left[\lim_{\alpha \to \infty} |V_n|^2 \mathbf{1}\{|V_n| > \alpha\} \right]$$
(613)

$$= 0$$
 (614)

uniformly in n.

On the other hand, for each fixed α , we have

$$\lim_{n \to \infty} \mathbb{E}_P\left[|V_n|^2 \mathbf{1}\{|V_n| > \alpha\}\right] = \lim_{n \to \infty} \mathbb{E}_P\left[V_n^2\right] - \mathbb{E}_P\left[|V_n|^2 \mathbf{1}\{|V_n| \le \alpha\}\right]$$
(615)

$$=1-\lim_{n\to\infty}\mathbb{E}_P\big[|V_n|^2\mathbf{1}\{|V_n|\le\alpha\}\big],\tag{616}$$

where (616) follows from (612). Note that

$$|V_n|^2 \mathbf{1}\{|V_n| \le \alpha\} \le \alpha^2,\tag{617}$$

and

$$|V_n|^2 \mathbf{1}\{|V_n| \le \alpha\} \xrightarrow{\text{(a.s.)}} V^2 \mathbf{1}\{|V| \le \alpha\}$$
(618)

by the continuous mapping theorem [23] and $V_n \xrightarrow{(a.s.)} V$. Hence, by the dominated convergence theorem [23], we also have

$$\lim_{n \to \infty} \mathbb{E}_P\left[|V_n|^2 \mathbf{1}\{|V_N| \le \alpha\}\right] = \mathbb{E}_P\left[|V|^2 \mathbf{1}\{|V| \le \alpha\}\right]$$
(619)

point-wise in α . Hence, from (616) and (619), we have

$$\lim_{n \to \infty} \mathbb{E}_P\left[|V_n|^2 \mathbf{1}\{|V_n| > \alpha\}\right] = 1 - \mathbb{E}_P\left[|V|^2 \mathbf{1}\{|V| \le \alpha\}\right]$$
(620)

point-wise in α .

From (614) and (620), by Moore-Osgood theorem [31] on the interchange of limits, it holds that

$$\lim_{\alpha \to \infty} \lim_{n \to \infty} \mathbb{E}_P \left[|V_n|^2 \mathbf{1}\{ |V_n| \le \alpha\} \right] = \lim_{n \to \infty} \lim_{\alpha \to \infty} \mathbb{E}_P \left[|V_n|^2 \mathbf{1}\{ |V_n| \le \alpha\} \right]$$
(621)

$$=0,$$
 (622)

which leads to $\{V_n\}_{n=1}^\infty$ be uniformly integrable. Hence, it holds that

$$\lim_{n \to \infty} \mathbb{E}_P[V_n^2] = \mathbb{E}_P[V^2] \tag{623}$$

$$=\mathbb{E}[U^2].\tag{624}$$

From (612) and (624), we obtain

$$\mathbb{E}[U^2] = 1. \tag{625}$$

B. Proof of Lemma 3

First, we prove that for any $\alpha > 1$ and $\lambda > 0$, the following holds:

$$\mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n},Q)^{\frac{\lambda}{\alpha n}}\right]^{\frac{\lambda n}{\lambda}} \leq \mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n},Q)^{\frac{\lambda}{n}}\right]^{\frac{n}{\lambda}}.$$
(626)

Indeed, let

$$r = \frac{\lambda}{\alpha n}, \qquad p = \frac{\lambda}{n}, \qquad q = \frac{\lambda}{(\alpha - 1)n}.$$
 (627)

Then, it holds that

$$\frac{1}{r} = \frac{1}{p} + \frac{1}{q}$$
(628)

and $p, q, r \in (0, \infty)$ if $\alpha > 1$. By applying the generalized Hölder's inequality [3], [30], we have

$$\left(\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_{n},Q)^{r}]\right)^{\frac{1}{r}} \leq \left(\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_{n},Q)^{p}]\right)^{\frac{1}{p}} \left(\mathbb{E}[1^{q}]\right)^{\frac{1}{q}}$$
(629)

$$= \left(\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n, Q)^p]\right)^{\frac{1}{p}},\tag{630}$$

implying that (626) holds.

Since (626) holds for any $\alpha > 1$, we have

$$\mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n},Q)^{\frac{\lambda}{n}}\right]^{\frac{n}{\lambda}} \geq \lim_{\alpha \to \infty} \mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n},Q)^{\frac{\lambda}{\alpha n}}\right]^{\frac{\alpha n}{\lambda}}$$
(631)

$$=2^{\mathbb{E}[\log P_{\mathrm{e}}(\mathcal{C}_{n},Q)]} \tag{632}$$

where (632) follows from the identity $\mathbb{E}[\log X] = \lim_{x \to \infty} \log \mathbb{E}[X^{\frac{1}{x}}]^x$ for any given X > 0. From the definition of $E_{\text{trc}}(R,Q)$ in (9) and the definition of limit, we have that for every $\epsilon > 0$ there exists an n_0 such that for $n > n_0$,

$$\left| -\frac{1}{n} \mathbb{E}[\log P_{\mathrm{e}}(\mathcal{C}_n, Q)] - E_{\mathrm{trc}}(R, Q) \right| < \epsilon.$$
(633)

Therefore, from (632) we have that

$$\mathbb{E}\left[P_{\mathrm{e}}(\mathcal{C}_{n},Q)^{\frac{\lambda}{n}}\right]^{\frac{n}{\lambda}} \ge 2^{-n(1-\varepsilon)E_{\mathrm{trc}}(R,Q)}.$$
(634)

Thus, from (634) and (626), by letting $\varepsilon \to 0$, it holds that

$$\liminf_{n \to \infty} \mathbb{E} \left[P_{\mathbf{e}}(\mathcal{C}_n, Q)^{\frac{\lambda}{n}} \right] \ge 2^{-\lambda E_{\mathrm{trc}}(R, Q)}.$$
(635)

Now, by the concavity of the function $x^{\frac{\lambda}{n}}$ on $(0,\infty)$, we have by Jensen's inequality that

$$\limsup_{n \to \infty} \mathbb{E} \left[P_{\mathbf{e}}(\mathcal{C}_n, Q)^{\frac{\lambda}{n}} \right] \le \limsup_{n \to \infty} \mathbb{E} \left[P_{\mathbf{e}}(\mathcal{C}_n, Q) \right]^{\frac{\lambda}{n}}$$
(636)

$$=2^{-\lambda E_{\rm rce}(R,Q)}.$$
(637)

where (637) follows from the fact that $\mathbb{E}[P_{e}(\mathcal{C}_{n}, Q)] \leq 2^{-nE_{rce}(R,Q)}$ [3], [28].

From (635) and (637), under the condition that $E_{\rm trc}(R,Q) = E_{\rm rce}(R,Q)$, it holds that

$$\lim_{n \to \infty} \mathbb{E} \left[P_{\mathbf{e}}(\mathcal{C}_n, Q)^{\frac{\lambda}{n}} \right] = 2^{-\lambda E_{\mathrm{rce}}(R, Q)}.$$
(638)

This concludes the proof of this lemma.

C. Proof of Lemma 4

The upper bound follows from Bhattacharyya bound. Now, by [32], it holds that

$$\mathbb{E}[P_{\mathbf{e}}(\mathcal{C}_n)] \doteq 2^{-nE_{\mathrm{rce}}(R,Q)} \tag{639}$$

for all $R < R_{crit}(Q)$. In addition, at this range of rate, the Bhattacharyya bound achieves the Gallager's random coding bound $E_{rce}(R,Q)$. Hence, from (639), we have

$$\mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)] \doteq 2^{-nE_{\mathrm{rce}}(R,Q)}$$
(640)

for all $R < R_{\text{crit}}(Q)$, where $P_{\text{e}}^{\text{ub}}(\mathcal{C}_n)$ is the union bound on $P_{\text{e}}(\mathcal{C}_n)$.

Now, for all rate $R < R_{\text{crit}}(Q)$, $E_{\text{rce}}(R,Q) = R_0(Q) - R$, where R_0 is the cut-off rate corresponding to the underlying distribution Q, i.e.,

$$R_0(Q) = -\log\left(\sum_y \left(\sum_x Q(x)\sqrt{W(y|x)}\right)^2\right).$$
(641)

Let $Q_X = Q'_X = Q$. By using standard KKT conditions for convex optimization, it is not hard to prove that

$$R_0(Q) = \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} P_{XX'}(x,x') d_{\mathrm{B}}(x,x').$$
(642)

Hence, from (642), we obtain

$$\mathbb{E}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})\right] \doteq 2^{nR} 2^{-n\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} \left(D(P_{XX'} \| Q_{X} Q_{X'}') + \sum_{x,x'} P_{XX'}(x,x') d_{\mathrm{B}}(x,x')\right)}$$
(643)

$$\doteq M_n \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} 2^{-nD(P_{XX'} \| Q_{XX'})} 2^{-n\sum_{x,x'} P_{XX'}(x,x') d_{\mathrm{B}}(x,x')}.$$
(644)

Now, let $\mathcal{N}(P_{XX'})$ be the number of codeword pairs which have the same join type $P_{XX'}$. Then, it holds that

$$\mathcal{N}(P_{XX'}) = \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbf{1}\{(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(P_{XX'})\},\tag{645}$$

which leads to

$$\mathbb{E}[\mathcal{N}(P_{XX'})] = \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(P_{XX'})]$$
(646)

$$= M_n (M_n - 1) 2^{-nD(P_{XX'} || Q_{XX'})}.$$
(647)

From (644) and (647), we have

$$(M_n - 1)\mathbb{E}\big[P_{\mathbf{e}}^{\mathrm{ub}}(\mathcal{C}_n)\big] \doteq \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} \mathbb{E}[\mathcal{N}(P_{XX'})] 2^{-n\sum_{x,x'} P_{XX'}(x,x')d_{\mathrm{B}}(x,x')}.$$
(648)

On the other hand, observe that

$$(M_n - 1)\mathbb{E}\big[P_{\mathbf{e}}^{\mathrm{ub}}(\mathcal{C}_n)\big] = \mathbb{E}\bigg[\sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}\big(\boldsymbol{X}_i \to \boldsymbol{X}_j\big)\bigg].$$
(649)

From (648) and (649), we obtain

$$\mathbb{E}\left[\sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j)\right] = \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} 2^{-nD(P_{XX'} \| Q_{XX'})} 2^{-n\sum_{x,x'} P_{XX'}(x,x')d_{\mathrm{B}}(x,x')}.$$
 (650)

Since (650) holds for all random i.i.d. codebook ensembles, hence for any fixed type $P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})$, by choosing a sub-random codebook ensemble which contains all the codewords with the same joint type $P_{XX'}$, we obtain

$$\mathbb{P}\left[\boldsymbol{X}_{i} \to \boldsymbol{X}_{j} \middle| (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(P_{XX'}) \right] \doteq 2^{-n \sum_{x,x'} P_{XX'}(x, x') d_{\mathrm{B}}(x, x')},$$
(651)

or (57) holds.

D. Proof of Lemma 5

Observe that

$$\mathbb{P}[\mathcal{V}_n^c] = \mathbb{P}\bigg[\sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} || Q_X Q'_X) > 2R} \mathcal{N}(P_{XX'}) \ge 1\bigg]$$
(652)

$$\leq \mathbb{E}\bigg[\sum_{P_{XX'}\in\mathcal{P}_n(\mathcal{X}\times\mathcal{X}):D(P_{XX'}\|Q_XQ'_X)>2R}\mathcal{N}(P_{XX'})\bigg]$$
(653)

$$= \mathbb{E}\bigg[\sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) > 2R} \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg]$$
(654)

$$= \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} || Q_X Q'_X) > 2R} \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(P_{XX'})]$$
(655)

$$\leq \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) > 2R} \sum_{i=1}^{M_n} \sum_{j \neq i} 2^{-nD(P_{XX'} || Q_X Q'_X)}$$
(656)

$$\leq \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) > 2R} \sum_{i=1}^{M_n} \sum_{j \neq i} 2^{-nD(P_{XX'} || Q_X Q'_X)}$$
(657)

$$= M_n(M_n - 1) \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} || Q_X Q'_X) > 2R} 2^{-nD(P_{XX'} || Q_X Q'_X)}$$
(658)

$$\leq 2^{2nR} \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) > 2R} 2^{-nD(P_{XX'} || Q_X Q'_X)}$$
(659)

$$\leq 2^{2nR} \sum_{\substack{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) > 2R}} 2^{-n(2R+\alpha(R))}$$
(660)

$$\dot{<} 2^{-n\alpha(R)} \tag{661}$$

for some $\alpha(R) > 0$, where (657) follows from [29], (660) follows from the fact that the number of possible joint types on $\mathcal{X} \times \mathcal{X}$ is sub-exponential in n.

E. Proof of Lemma 6

Define

$$\tilde{V}_{ij} = \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R - \nu} \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\} g_n(P_{XX'}).$$
(662)

Then, we have

$$D_n = \frac{1}{M_n} \sum_{\substack{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu}} \mathcal{N}(P_{XX'}) g_n(P_{XX'})$$
(663)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \sum_{\substack{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu}} \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\} g_n(P_{XX'})$$
(664)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \tilde{V}_{ij}.$$
 (665)

It is easy to see that $\{\tilde{V}_{ij}\}_{i,j=1}^{M_n}$ are pairwise independent. Hence, from (665), we have

$$\operatorname{Var}(D_{n}) = \frac{1}{M_{n}^{2}} \sum_{i=1}^{M_{n}} \sum_{j \neq i} \operatorname{Var}(\tilde{V}_{ij}).$$
(666)

Observe that

$$\begin{aligned} \operatorname{Var}(\tilde{V}_{ij}) &= \operatorname{Var}\left(\sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) \leq 2R - \nu} \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\} g_n(P_{XX'})\right) \end{aligned}$$
(667)

$$&= \mathbb{E}\bigg[\sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) \leq 2R - \nu} \left(\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\right) \\ &- \mathbb{E}\bigg[\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg]\bigg)^2 g_n^2(P_{XX'})\bigg] \end{aligned}$$
(668)

$$&= \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) \leq 2R - \nu} \left(\mathbb{E}\bigg[\bigg(\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg)^2\bigg] \\ &- \bigg(\mathbb{E}\bigg[\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg]\bigg)^2\bigg)g_n^2(P_{XX'}) \end{aligned}$$
(669)

$$&\leq \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) \leq 2R - \nu} \bigg(\mathbb{E}\bigg[\bigg(\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg)^2\bigg] \\ &- \bigg(\mathbb{E}\bigg[\mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\}\bigg]\bigg)^2\bigg)g_n^2(P_{XX'}) \end{aligned}$$
(670)

$$&= \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) \leq 2R - \nu} \mathbb{P}\bigg[\big(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\bigg]\bigg(1 - \mathbb{P}\bigg[(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\bigg]\bigg)g_n^2(P_{XX'}) \end{aligned}$$

$$\in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(\mathcal{P}_{XX'} || Q_X Q_X) \le 2R - \nu$$
(671)

$$\leq 2^{n \max_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \leq 2R - \nu} - \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x')} \\ \times \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \leq 2R - \nu} \mathbb{P}\bigg[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(P_{XX'}) \bigg] g_n(P_{XX'}),$$
(672)

where (668) follows from the pairwise independence of $\{(X_i, X_j)\}_{i \neq j}$, and (672) follows from (746).

Hence, from (666) and (672), we obtain

$$\operatorname{Var}(D_{n}) = \frac{1}{M_{n}^{2}} \sum_{i=1}^{M_{n}} \sum_{j \neq i} \operatorname{Var}(\tilde{V}_{ij})$$

$$\leq 2^{-2nR} \times 2^{n \max_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \parallel Q_{X}Q'_{X}) \leq 2R-\nu} - \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')}$$

$$\times \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \parallel Q_{X}Q'_{X}) \leq 2R-\nu} \sum_{i=1}^{M_{n}} \sum_{j \neq i} \mathbb{P}\Big[(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \in \mathcal{T}(P_{XX'}) \Big] g_{n}(P_{XX'}) \tag{673}$$

 $= 2^{-2nR} \times 2^{n \max_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R-\nu} - \sum_{x,x'} d_{\mathbf{B}}(x,x') P_{XX'}(x,x')}$

$$\times \sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu} \mathbb{E}[\mathcal{N}(P_{XX'})]g_n(P_{XX'})$$
(674)

$$= 2^{-2nR} \times 2^{n \max_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R-\nu} - \sum_{x,x'} d_{\mathbf{B}}(x,x') P_{XX'}(x,x')}$$

$$\times \left(\sum_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} || Q_X Q'_X) \le 2R - \nu} \mathbb{E}[\mathcal{N}(P_{XX'})]g_n(P_{XX'})\right)$$
(675)

$$= M_n 2^{-2nR} \times 2^{-n \min_{P_{XX'} \in \mathcal{P}(X \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') \right)} \mathbb{E}[D_n]$$
(676)

$$= 2^{-nR} \times 2^{-n\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) = 2R-\nu} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')\right)} \mathbb{E}[D_n],$$
(677)

where (677) follows from the fact that the optimizer of the linear programming is in the boundary of the convex constraint set [33].

On the other hand, from (665), we have

$$\mathbb{E}[D_n] = \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \mathbb{E}[\tilde{V}_{ij}]$$

$$M_n$$
(678)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) \le 2R - \nu} \mathbb{P}\Big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(P_{XX'}) \Big] g_n(P_{XX'})$$
(679)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q_X') \le 2R - \nu} 2^{-nD(P_{XX'} \| Q_X Q_X')} g_n(P_{XX'})$$
(680)

$$= (M_n - 1) \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q_X') \le 2R - \nu} 2^{-nD(P_{XX'} \| Q_X Q_X')} g_n(P_{XX'})$$
(681)

$$\doteq (M-1) \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q_X') \le 2R - \nu} 2^{-nD(P_{XX'} \| Q_X Q_X')} 2^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')}$$
(682)

$$\doteq (M_n - 1)2^{-n \min_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu} \left(D(P_{XX'} \parallel Q_X Q'_X) + \sum_{x, x'} d_B(x, x') P_{XX'}(x, x') \right)$$
(683)

$$\doteq (M_n - 1) 2^{-n \min_{P_{XX'} \in \mathcal{P}(X \times X) : D(P_{XX'} \parallel Q_X Q'_X) \le 2R - \nu} \left(D(P_{XX'} \parallel Q_X Q'_X) + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') \right)},$$
(684)

where (680) follows from [29], (682) follows from (746), and (684) follows from the fact that $\mathcal{P}_n(\mathcal{X} \times \mathcal{X})$ is dense in $\mathcal{P}(\mathcal{X} \times \mathcal{X})$.

Similarly, observe that

$$\mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n})] = \frac{1}{M_{n}} \sum_{i=1}^{M_{n}} \sum_{j \neq i} \mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_{i} \to \boldsymbol{X}_{j}]\right]$$
(685)

$$= (M_n - 1) \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})} \mathbb{E}[\mathcal{N}(P_{XX'})]g_n(P_{XX'})$$
(686)

$$= (M_n - 1) \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})} 2^{-nD(P_{XX'} \parallel Q_X Q'_X)} 2^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')}$$
(687)

$$\doteq 2^{-N \min_{P_{XX'} \in \mathcal{P}_n(X \times \mathcal{X})} \left(D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') - R \right) }$$
(688)

$$= 2^{-n \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} \left(D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') - R \right)},$$
(689)

where (689) follows from the fact $\mathcal{P}_n(\mathcal{X} \times \mathcal{X})$ is dense in $\mathcal{P}(\mathcal{X} \times \mathcal{X})$.

It follows from (689) that

$$E_{\rm rce}(R,Q) + R = \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} D(P_{XX'} \| Q_X Q'_X) + \sum_{x,x'} d_{\rm B}(x,x') P_{XX'}(x,x')$$
(690)

Now, we have

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \parallel Q_X Q'_X) \ge 2R - \nu} D(P_{XX'} \parallel Q_X Q'_X) + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') \\
\geq \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X})} \min_{P(R)} ||Q_X Q'_X| \ge 2R - \nu + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') \tag{691}$$

$$= \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) = 2R - \nu} 2R - \nu + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x')$$
(692)

where (692) follows from the fact that the optimizer of the linear programming is on the boundary of the convex constraint set [33].

Now, following the standard approach to solve the linear programming in (692) we have the following Lagrangian:

$$\mathcal{L}(P_{XX'},\lambda) = 2R - \nu + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') + \lambda \big(D(P_{XX'} \| Q_X Q_X') - 2R + \nu \big).$$
(693)

By setting,

$$0 = \frac{\partial \mathcal{L}(P_{XX'}, \lambda)}{\partial P_{XX'}}(x, x') = d_{\mathrm{B}}(x, x') + \lambda \bigg(\log \frac{P_{XX'}(x, x')}{Q_X Q'_X} + \log e\bigg),\tag{694}$$

we have

$$P_{XX'}(x,x') = \frac{1}{e} Q_X(x) Q_{X'}(x') 2^{-\frac{d_{\mathbf{B}}(x,x')}{\lambda}}.$$
(695)

Since

$$\sum_{x,x'} P_{XX'}(x,x') = 1,$$
(696)

we must choose λ such that

$$\frac{1}{e} \sum_{x,x'} Q_X(x) Q_{X'}(x') 2^{-\frac{d_{\mathrm{B}}(x,x')}{\lambda}} = 1.$$
(697)

Since $d_{\rm B}(x,x') = -\log \sum_{y \in \mathcal{Y}} \sqrt{W(y|x)W(y|x')} \ge 0$, from (697) we must have $\lambda < 0$, otherwise the LHS of (697) is less than or equal 1/e < 1.

With the choice $P_{XX'}$ in (695), we have

$$2R - \nu + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{X,X'}(x,x')$$

= $2R - \nu + \frac{1}{e} \sum_{x,x'} d_{\mathrm{B}}(x,x') Q(x) Q(x') 2^{-\frac{d_{\mathrm{B}}(x,x')}{\lambda}}$ (698)

$$=2R-\nu-\frac{1}{e}\sum_{x,x'}Q(x)Q(x')\log\left(\sum_{y\in\mathcal{Y}}\sqrt{W(y|x)W(y|x')}\right)\left(\sum_{y\in\mathcal{Y}}\sqrt{W(y|x)W(y|x')}\right)^{\frac{1}{\lambda}}.$$
(699)

Now, from (697), we have

$$1 = \frac{1}{e} \sum_{x,x'} Q_X(x) Q_{X'}(x') 2^{-\frac{d_{\mathrm{B}}(x,x')}{\lambda}}$$
(700)

$$= \frac{1}{e} \sum_{x,x'} Q_X(x) Q_{X'}(x') \left(\sum_{y \in \mathcal{Y}} \sqrt{W(y|x)W(y|x')} \right)^{\frac{1}{\lambda}}$$
(701)

$$\leq \frac{1}{e} \left(\sum_{x,x'} Q(x)Q(x') \sum_{y \in \mathcal{Y}} \sqrt{W(y|x)W(y|x')} \right)^{\frac{1}{\lambda}},\tag{702}$$

where (702) follows from the concavity of the function $x^{1/\lambda}$ for $\lambda < 0$ and $x \in [0, 1]$.

On the other hand, since $\lambda < 0$, the function $-(\log x)x^{1/\lambda}$ is convex in $x \in [0, 1]$. Hence, by Jensen's inequality, from (699), we have

$$2R - \nu + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{X,X'}(x,x')$$

$$> 2R - \nu - \frac{1}{e} \log\left(\sum_{x,x'} Q(x)Q(x') \sum_{y \in \mathcal{Y}} \sqrt{W(y|x)W(y|x')}\right) \left(\sum_{x,x'} \sum_{y \in \mathcal{Y}} Q(x)Q(x') \sqrt{W(y|x)W(y|x')}\right)^{\frac{1}{\lambda}}$$

$$(703)$$

$$=2R-\nu+\left[-\log\left(\sum_{y\in\mathcal{Y}}\left(\sum_{x\in\mathcal{X}}Q(x)\sqrt{W(y|x)}\right)^2\right)\right]\left[\frac{1}{e}\left(\sum_{x,x'}\sum_{y\in\mathcal{Y}}Q(x)Q(x')\sqrt{W(y|x)W(y|x')}\right)^{\frac{1}{\lambda}}\right]$$
(704)

$$= 2R - \nu + R_0(Q) \left[\frac{1}{e} \left(\sum_{x,x'} \sum_{y \in \mathcal{Y}} Q(x)Q(x')\sqrt{W(y|x)W(y|x')} \right)^{\frac{1}{\lambda}} \right]$$
(705)

$$\geq 2R - \nu + R_0(Q) \tag{706}$$

$$\geq R + E_{\rm rce}(R,Q) \tag{707}$$

for $\nu \leq 2R$, where in (703), the equality does not hold by the condition (56), (705) follows from [28] where $R_0(Q)$ is the cut-off rate of the DMC under the underlying distribution Q, (706) follows from (702), and (707) follows from [28].

From (692) and (707), we have

$$\min_{\substack{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \ge 2R - \nu}} D(P_{XX'} \| Q_X Q'_X) + \sum_{x,x'} d_B(x, x') P_{XX'}(x, x')$$

$$\geq \min_{\substack{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) = 2R - \nu}} 2R - \nu + \sum_{x,x'} d_B(x, x') P_{XX'}(x, x')$$

$$> E_{\text{rce}}(R, Q) + R$$
(708)

if $0 \le \nu \le 2R$.

Therefore, from (690) and (708), we must have

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q_X') < 2R - \nu} D(P_{XX'} \parallel Q_X Q_X') + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') = E_{\mathrm{rce}}(R,Q) + R.$$
(709)

It follows that

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R - \nu} D(P_{XX'} \| Q_X Q'_X) + \sum_{x,x'} d_{\mathcal{B}}(x, x') P_{XX'}(x, x')$$

$$= \begin{cases} 2R - \nu + \sum_{x,x'} d_{\mathcal{B}}(x, x') P^*_{XX'}(x, x'), & \text{if } D(P^*_{XX'} \| Q_X Q'_X) = 2R - \nu, \\ R + E_{\text{rce}}(R, Q), & \text{otherwise,} \end{cases}$$

$$(710)$$

where (710) follows from (709). Here, $P_{XX'}^*$ is an optimizer of $\min_{P_{XX'} \in \mathcal{P}} D(P_{XX'} || Q_X Q_{X'}) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') - R$.

From (684) and (710), we obtain (63).

Furthermore, from (677) and (63), we obtain

$$\frac{\operatorname{Var}(D_{n})}{\left(\mathbb{E}[D_{n}]\right)^{2}} \stackrel{.}{\leq} \begin{cases} 2^{-n\nu}, & D(P_{XX'}^{*}|Q_{X}Q_{X}') = 2R - \nu\\ 2^{-n\left(\min_{P_{XX'} \in \mathcal{P}(X \times X): D(P_{XX'}||Q_{X}Q_{X}'}) = 2R - \nu + \sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x') - E_{\mathrm{rce}}(R,Q)\right)}, & \text{otherwise} \end{cases}$$
(711)

 $\doteq 2^{-\zeta(\nu,R)n} \tag{712}$

for

$$\zeta(\nu, R) = \begin{cases} \nu, & \text{if} \quad D(P_{XX'}^* | Q_X Q_X') = 2R - \nu \\ \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} | | Q_X Q_X') = 2R R + \sum_{x,x'} d_{\mathcal{B}}(x, x') P_{XX'}(x, x') - E_{\text{rce}}(R, Q), & \text{otherwise} \end{cases} > 0,$$
(713)

which is followed from (708) and $\nu > 0$.

This concludes our proof of Lemma 6.

F. Proof of Lemma 7

Recall the definition of $\mathcal{N}(P_{XX'})$ in Lemma 5. For a pair of codewords with $(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})$, it holds that

$$\mathbb{P}(\boldsymbol{X}_i \to \boldsymbol{X}_j) = g_n(P_{XX'}). \tag{714}$$

$$P_{e}^{ub}(\mathcal{C}_{n}) = \frac{1}{M_{n}} \sum_{i=1}^{M_{n}} \sum_{j \neq i} \mathbb{P}(\boldsymbol{X}_{i} \to \boldsymbol{X}_{j})$$
(715)

$$=\frac{1}{M_n}\sum_{P_{XX'}}\mathcal{N}(P_{XX'})g_n(P_{XX'})$$
(716)

where (716) follows from (714).

$$B_n = \frac{1}{M_n} \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})} \mathcal{N}(P_{XX'}) g_n(P_{XX'}), \tag{717}$$

$$\tilde{D}_n = \frac{1}{M_n} \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_X Q'_X) \le 2R} \mathcal{N}(P_{XX'}) g_n(P_{XX'}),$$
(718)

$$D_{n} = \frac{1}{M_{n}} \sum_{P_{XX'} \in \mathcal{P}_{n}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} || Q_{X}Q'_{X}) \le 2R - \nu} \mathcal{N}(P_{XX'}) g_{n}(P_{XX'}).$$
(719)

Since $\mathbb{E}[\tilde{D}_n] - \mathbb{E}[D_n] \to 0$ as $\nu \to 0$ and that $\mathbb{E}[\tilde{D}_n]$ and $\mathbb{E}[D_n]$ are exponentially decaying in n, for ν small enough, it holds that

$$\mathbb{E}[D_n] \le \mathbb{E}[\tilde{D}_n] \le \mathbb{E}[D_n] 2^{\varepsilon n/2}.$$
(720)

Recall the typical set defined in Lemma 5. For any given $\varepsilon > 0$, observe that

$$\mathbb{P}\left[\left|-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n}+\frac{1}{n}\log\left(\frac{1}{M_{n}}\sum_{P_{XX'}\in\mathcal{P}_{n}(\mathcal{X}\times\mathcal{X}):D(P_{XX'}\|Q_{X}Q_{X}')\leq 2R}\mathbb{E}\left[\mathcal{N}(P_{XX'})\right]g_{n}(P_{XX'})\right)\right|>\varepsilon\right]$$
(721)

$$= \mathbb{P}\left[\left|\log\frac{B_n}{\mathbb{E}[\tilde{D}_n]}\right| > \varepsilon n\right]$$
(722)

$$\leq \mathbb{P}\left[\left|\log\frac{B_n}{\mathbb{E}[\tilde{D}_n]}\right| > \varepsilon n \left|\mathcal{V}_n\right] \mathbb{P}[\mathcal{V}_n] + \mathbb{P}[\mathcal{V}_n^c]$$
(723)

$$= \mathbb{P}\left[\left|\log\frac{\tilde{D}_n}{\mathbb{E}[\tilde{D}_n]}\right| > \varepsilon n \left|\mathcal{V}_n\right] \mathbb{P}[\mathcal{V}_n] + \mathbb{P}[\mathcal{V}_n^c]$$
(724)

$$\leq \mathbb{P}\left[\left|\log\frac{\tilde{D}_{n}}{\mathbb{E}[\tilde{D}_{n}]}\right| > \varepsilon n\right] + \mathbb{P}[\mathcal{V}_{n}^{c}]$$
(725)

$$= \mathbb{P}\big[\tilde{D}_n > \mathbb{E}[\tilde{D}_n]2^{\varepsilon n}\big] + \mathbb{P}\big[\tilde{D}_n < \mathbb{E}[\tilde{D}_n]2^{-\varepsilon n}\big] + \mathbb{P}[\mathcal{V}_n^c]$$
(726)

$$\leq \frac{\mathbb{E}[D_n]}{2^{\varepsilon n} \mathbb{E}[\tilde{D}_n]} + \mathbb{P}\left[D_n < \mathbb{E}[D_n]2^{-\varepsilon/2n}\right] + \mathbb{P}[\mathcal{V}_N^c]$$
(727)

$$=2^{-\varepsilon n} + \mathbb{P}\big[D_n < \mathbb{E}[\tilde{D}_n]2^{-\varepsilon n}\big] + 2^{-n\alpha(R)}$$
(728)

$$\leq 2^{-\varepsilon n} + \mathbb{P}\left[D_n < \mathbb{E}[D_n]2^{-(\varepsilon/2)n}\right] + 2^{-n\alpha(R)}$$
(729)

where (723) follows from $\mathbb{P}(A) = \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^c)\mathbb{P}(B^c) \leq \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(B^c)$, (724) follows from the fact that given \mathcal{V}_n , it holds that $B_n = D_n$, (725) follows from $\mathbb{P}(A|B)\mathbb{P}(B) \leq \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^c)\mathbb{P}(B^c) = \mathbb{P}(A)$, (727) follows from Markov's inequality, (728) follows from $D_n \leq \tilde{D}_n$ and from Lemma 5, (729) follows from $\mathbb{E}[\tilde{D}_n] \leq \mathbb{E}[D_n]2^{\varepsilon n/2}$ for ν sufficiently small by (720).

Now, we have

$$\mathbb{P}\big[D_n < \mathbb{E}[D_n]2^{-(\varepsilon/2)n}\big] = \mathbb{P}\big[D_n - \mathbb{E}[D_n] < \mathbb{E}[D_n]\big(2^{-(\varepsilon/2)n} - 1\big)\big]$$
(730)

$$\leq \mathbb{P}\big[\big|D_n - \mathbb{E}[D_n]\big| > \mathbb{E}[D_n]\big(1 - 2^{-(\varepsilon/2)n}\big)\big] \tag{731}$$

$$\stackrel{\cdot}{\leq} \frac{\operatorname{Var}(D_n)}{\left(\mathbb{E}[D_n]\right)^2}$$
(732)

$$\leq 2^{-n\beta(\nu,R)}, \tag{733}$$

where (733) follows from Lemma 6.

From (729) and (733), for any $\varepsilon > 0$ and R > 0, we have

$$\mathbb{P}\left[\left|-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n}+\frac{1}{n}\log\left(\frac{1}{M_{n}}\sum_{P_{XX'}:D(P_{XX'}|\|Q_{X}Q_{X}')\leq 2R}\mathbb{E}\left[\mathcal{N}(P_{XX'})\right]g_{n}(P_{XX'})\right)\right|>\varepsilon\right]$$
(734)

$$\leq 2^{-\varepsilon n} + 2^{-n\beta(\nu,R)} + 2^{-n\alpha(R)}.$$
(735)

It follows from (735) that

$$\sum_{n=1}^{\infty} \mathbb{P}\left[\left|-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n}+\frac{1}{n}\log\left(\frac{1}{M_{n}}\sum_{P_{XX'}:D(P_{XX'}|\|Q_{X}Q_{X}')\leq 2R}\mathbb{E}\left[\mathcal{N}(P_{XX'})\right]g_{n}(P_{XX'})\right)<\infty.$$
(736)

Hence, by Borel-Cantelli's lemma [23], we have

$$-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n} + \frac{1}{n} \log \left(\frac{1}{M_{n}} \sum_{P_{XX'}: D(P_{XX'} \parallel Q_{X}Q'_{X}) \leq 2R} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_{N}(P_{XX'}) \right) \xrightarrow{(\mathrm{a.s.})} 0 \tag{737}$$

On the other hand, we have

$$-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}))}{n} + \frac{1}{n} \log \left(\frac{1}{M_{n}} \sum_{P_{XX'}: D(P_{XX'} \parallel Q_{X}Q'_{X}) \leq 2R} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_{n}(P_{XX'}) \right) \right|$$

$$\leq -\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n})}{n} + \left| -\frac{1}{n} \log \left(\frac{1}{M_{n}} \sum_{P_{XX'}: D(P_{XX'} \parallel Q_{X}Q'_{X}) \leq 2R} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_{n}(P_{XX'}) \right) \right|$$
(738)

$$\leq -\frac{1}{n}\log P_{\mathrm{e}}(\mathcal{C}_{n}) + \left| -\frac{1}{n}\log\left(\frac{1}{M_{n}}\sum_{P_{XX'}:D(P_{XX'}||Q_{X}Q_{X}'|) \leq 2R} \mathbb{E}\left[\mathcal{N}(P_{XX'})\right]g_{n}(P_{XX'})\right) \right|$$
(739)

$$\leq E_{\rm sp}(R) + \left| -\frac{1}{n} \log \left(\frac{1}{M_n} \sum_{P_{XX'}: D(P_{XX'} \parallel Q_X Q'_X) \leq 2R} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_n(P_{XX'}) \right) \right|, \tag{740}$$

$$\leq E_{\rm sp}(R) + \left| -\frac{1}{n} \log\left(\frac{\mathbb{E}[D_n]}{M_n}\right) \right|,\tag{741}$$

$$\leq E_{\rm sp}(R) + R + \min_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q'_X) = 2R} \sum_{x,x'} d_{\rm B}(x,x') P_{XX'}(x,x')$$
(742)

$$\leq E_{\rm sp}(R) + R + D_{\rm b} < \infty,\tag{743}$$

where (740) follows from [28], (741), and (742) follows from Lemma 6, where (743) follows with the fact that $d_{\rm B}(x, x') \leq D_{\rm b} < \infty$ for all x, x' by the condition (56).

From (737), (743), and the bounded convergence theorem [23], we have

$$\lim_{n \to \infty} \mathbb{E} \left[-\frac{\log P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)}{n} + \frac{1}{n} \log \left(\frac{1}{M_n} \sum_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_n(P_{XX'}) \right) \right] = 0.$$
(744)

Now, by Lemma 6, we have

$$\lim_{n \to \infty} -\frac{1}{n} \log \left(\frac{1}{M_n} \sum_{\substack{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R}} \mathbb{E} \left[\mathcal{N}(P_{XX'}) \right] g_n(P_{XX'}) \right)$$
$$= \min_{\substack{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R}} \left(D(P_{XX'} \| Q_X Q'_X) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x') - R \right).$$
(745)

Hence, we obtain (66) from (744) and (745). Note that (67) can be achieved from (66) by using (710) with $\nu = 0$.

G. Proof of Lemma 8

Let $g_n(P_{XX'})$ be the pairwise error probability given the joint type $P_{XX'}$ for $P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})$. By Lemma 4, the pairwise error probability can be expressed as

$$g_n(P_{XX'}) \doteq 2^{-n\sum_{x,x'} d_{\mathcal{B}}(x,x')P_{XX'}(x,x')},\tag{746}$$

where

$$d_{\rm B}(x,x') = -\log\bigg(\sum_{y\in\mathcal{Y}}\sqrt{W(y|x)W(y|x')}\bigg).$$
(747)

Now, let

$$\tilde{V}_{ij} = \sum_{P_{XX'}: D(P_{XX'} || Q_X Q'_X) \le 2R} \mathbf{1}\{(X_i, X_j) \in \mathcal{T}(P_{XX'})\} g_n(P_{XX'}).$$
(748)

Now, let

$$\tilde{P}_{e}^{ub}(\mathcal{C}_{n}) = \frac{1}{M_{n}} \sum_{\substack{P_{XX'}: D(P_{XX'} || Q_{X}Q_{X}') \le 2R \\ M}} \mathcal{N}(P_{XX'}) g_{N}(P_{XX'})$$
(749)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \sum_{P_{XX'}: D(P_{XX'} || Q_X Q'_X) \le 2R} \mathbf{1}\{(\mathbf{X}_i, \mathbf{X}_j) \in \mathcal{T}(P_{XX'})\} g_n(P_{XX'})$$
(750)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \tilde{V}_{ij},$$
(751)

$$A_n = \mathbb{E}[\tilde{P}_{e}^{ub}(\mathcal{C}_n)].$$
(752)

From Lemma 6 and Lemma 7, we have

$$A_n = 2^{-nE_{\rm trc}^{\rm ub}(R,Q)}.$$
(753)

Now, recall the definition of the typical set \mathcal{V}_n in Lemma 5. Observe that

$$\mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) > \frac{1}{2}2^{-n(E_{trc}^{ub}(R,Q)-\varepsilon)}\right] + \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) < 2^{-n(E_{trc}^{ub}(R,Q)+\varepsilon)}\right] \\
= \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) - A_{n} > \frac{1}{2}2^{-n(E_{trc}^{ub}(R,Q)-\varepsilon)} - A_{n}\right] \\
+ \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) - A_{n} < 2^{-n(E_{trc}^{ub}(R,Q)+\varepsilon)} - A_{n}\right]$$
(754)

$$\leq \mathbb{P}\left[\left|P_{e}^{ub}(\mathcal{C}_{n}) - A_{n}\right| > \frac{1}{2}2^{-n(E_{trc}^{ub}(R,Q) - \varepsilon)} - A_{n}\right] + \mathbb{P}\left[\left|P_{e}^{ub}(\mathcal{C}_{n}) - A_{n}\right| > A_{n} - 2^{-n(E_{trc}^{ub}(R,Q) + \varepsilon)}\right]$$

$$(755)$$

$$\leq \mathbb{P}\bigg[\left| P_{e}^{ub}(\mathcal{C}_{n}) - A_{n} \right| > \frac{1}{2} 2^{-N(E_{trc}^{ub}(R,Q)-\varepsilon)} - A_{n} |\mathcal{V}_{n}] \mathbb{P}[\mathcal{V}_{n}] + \mathbb{P}(\mathcal{V}_{n}^{c}) \\ + \mathbb{P}\bigg[\left| P_{e}^{ub}(\mathcal{C}_{n}) - A_{n} \right| > A_{n} - 2^{-n(E_{trc}^{ub}(R,Q)+\varepsilon)} |\mathcal{V}_{n}] \mathbb{P}[\mathcal{V}_{n}] + \mathbb{P}(\mathcal{V}_{n}^{c}) \\ \leq \mathbb{P}\bigg[\left| \tilde{P}_{e}^{ub}(\mathcal{C}_{n}) - A_{n} \right| > \frac{1}{2} 2^{-n(E_{trc}^{ub}(R,Q)-\varepsilon)} - A_{n} |\mathcal{V}_{n}] \mathbb{P}[\mathcal{V}_{n}] + \mathbb{P}(\mathcal{V}_{n}^{c})$$
(756)

$$+ \mathbb{P}\left[\left|\tilde{P}_{e}^{ub}(\mathcal{C}_{n}) - A_{n}\right| > A_{n} - 2^{-n(E_{trc}^{ub}(R,Q) + \varepsilon)} |\mathcal{V}_{n}\right] \mathbb{P}[\mathcal{V}_{n}] + \mathbb{P}(\mathcal{V}_{n}^{c})$$

$$\tag{757}$$

$$\leq \mathbb{P}\left[\left|\tilde{P}_{e}^{ub}(\mathcal{C}_{n}) - A_{n}\right| > \frac{1}{2}2^{-n(E_{trc}^{ub}(R,Q)-\varepsilon)} - A_{n}\right] + \mathbb{P}(\mathcal{V}_{n}^{c}) \\ + \mathbb{P}\left[\left|\tilde{P}_{e}^{ub}(\mathcal{C}_{n}) - A_{n}\right| > A_{n} - 2^{-n(E_{trc}^{ub}(R,Q)+\varepsilon)}\right] + \mathbb{P}(\mathcal{V}_{n}^{c})$$

$$(758)$$

$$\leq \frac{\operatorname{Var}(P_{e}^{\operatorname{ub}}(\mathcal{C}_{n}))}{2^{-2nE_{\operatorname{tre}}^{\operatorname{ub}}(R,Q)}} + 2^{-\alpha(R)n}$$

$$(759)$$

$$=2^{2nE_{\rm trc}^{\rm ub}(R,Q)}\frac{1}{M_n^2}\sum_{i=1}^{M_n}\sum_{j\neq i}\operatorname{Var}(\tilde{V}_{ij})+2^{-\alpha(R)n}$$
(760)

$$=2^{2nE_{\rm trc}^{\rm ub}(R,Q)}2^{-2nR}2^{n\max_{P_{XX'}:D(P_{XX'}||Q_XQ'_X)\leq 2R}-\sum_{x,x'}d_{\rm B}(x,x')P_{XX'}(x,x')}2^{nR}2^{-nE_{\rm trc}^{\rm ub}(R,Q)}+2^{-\alpha(R)N}$$
(761)

$$=2^{-n(R-E_{\rm trc}^{\rm ub}(R,Q)+\min_{P_{XX'}:D(P_{XX'}||Q_XQ'_X)\leq 2R}\sum_{x,x'}d_{\rm B}(x,x')P_{XX'}(x,x')}+2^{-\alpha(R)n},$$
(762)

where (756) follows from $\mathbb{P}(A) = \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(A|B^c)\mathbb{P}(B^c) \leq \mathbb{P}(A|B)\mathbb{P}(B) + \mathbb{P}(B^c)$, (759) follows from Chebyshev's inequality, (684), and Lemma 5, (760) follows from the pairwise independence of \tilde{V}_{ij} , and (761) follows from (677) and $M_n \doteq 2^{nR}$.

Now, for the case $E_{\rm trc}^{\rm ub}(R,Q)=E_{\rm rce}(R,Q),$ we must have

$$R + \min_{P_{XX'}: D(P_{XX'} \parallel Q_X Q_X') \le 2R} d_{\mathcal{B}}(x, x') P_{XX'}(x, x') = \min_{P_{XX'}: D(P_{XX'} \parallel Q_X Q_X') = 2R} R + \sum_{x, x'} d_{\mathcal{B}}(x, x') P_{XX'}(x, x')$$
(763)

$$\geq \min_{P_{XX'}: D(P_{XX'} \parallel Q_X Q'_X) \geq 2R} R + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x')$$
(764)

$$> E_{\rm rce}(R,Q)$$
 (765)

$$=E_{\rm trc}^{\rm ub}(R,Q),\tag{766}$$

where (765) follows from (708) by setting $\nu = 0$.

Hence, for this case, from (762), we have

$$\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) > \frac{1}{2}2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)-\varepsilon)}\right] + \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+\varepsilon)}\right] \le 2^{-n\Delta}$$
(767)

for some $\Delta > 0$.

Now, we consider the case $E_{\text{trc}}^{\text{ub}}(R,Q) > E_{\text{rce}}(R,Q)$. First of all we need the following lemma, which extends [7, Th. 2.1] to the DMC case.

Lemma 24: In a DMC, for all rates for which $E_{\text{trc}}^{\text{ub}}(R,Q) = E_{\text{rce}}(R,Q) + \Delta E$, $\Delta E > 0$ the probability that a code of length n and rate R from the RCE has a codeword pair with empirical joint type $P_{XX'}$ such that $D(P_{XX'}||Q_XQ'_X) > 2R$ goes to 0 exponentially fast as $n \to \infty$.

Furthermore, if $P_{XX'}$ is such that $D(P_{XX'}||Q_XQ'_X) \leq 2R$, the probability that the number of codeword pairs with joint type $P_{XX'}$ satisfies $\mathcal{N}(P_{XX'}) \doteq 2^{-n(2R-D(P_{XX'})|Q_XQ'_X))}$ goes to one exponentially fast as $n \to \infty$.

Proof: Let $\mathcal{N}(P_{XX'})$ be the number of codeword pairs with joint type $P_{XX'}$. Then, we have

$$\mathbb{P}\big[\mathcal{N}(P_{XX'}) \ge 1\big] \le \mathbb{E}\big[\mathcal{N}(P_{XX'})\big] \tag{768}$$

$$=\frac{M(M-1)}{2}\Pr\left[P_{XX'}\right] \tag{769}$$

$$\doteq e^{n2R} e^{-nD(P_{XX'} \| Q_X Q_X')}$$
(770)

$$=2^{-n\left(D(P_{XX'}\|Q_XQ'_X)-2R\right)}$$
(771)

$$\doteq 2^{-n\left(D(P_{XX'} \| Q_X Q'_X) - 2R\right)} \to 0,$$
(772)

where (768) follows from the Markov's inequality, $\mathbb{P}\{P_{XX'}\}$ in (769) indicates the probability to find a codeword pair with joint type $P_{XX'}$ and (770) follows from [19, Th. 11.1.4]

Similarly, it is not hard to see that

$$\operatorname{Var}(\mathcal{N}(P_{XX'})) \doteq 2^{-n\left(D(P_{XX'} \| Q_X Q'_X) - 2R\right)}.$$
(773)

The second statement can be proven by observing that by Chebyshev's inequality and (773), for any positive number Δ :

$$\mathbb{P}\left[|\mathcal{N}(P_{XX'}) - E\left[\mathcal{N}_n(P_{XX'})\right]| > 2^{-n(D(P_{XX'}||Q_XQ'_X) - 2R + \Delta)}\right] \le 2^{-n(2R - D(P_{XX'}||Q_XQ'_X) - 2\Delta)}.$$
(774)

Furthermore, since $\mathbb{E}[\mathcal{N}(P_{XX'})] \doteq 2^{-n(2R-D(P_{XX'}||Q_XQ'_X))}$, from (774) the probability that the number of codeword pairs with joint type $P_{XX'}$ satisfies $\mathcal{N}(P_{XX'}) \doteq 2^{-n(2R-D(P_{XX'}||Q_XQ'_X))}$ goes to one exponentially fast as $n \to \infty$. As a closing remark, note that the region defined by the inequality $D(P_{XX'}||Q_XQ'_X) \le 2R$ is the equivalent for general DMC to the Gilbert-Varshamov region in [7].

Now, for the case $E^{\rm ub}_{\rm trc}(R,Q)>E_{\rm rce}(R,Q),$ observe that

$$\mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) \geq \frac{1}{2}e^{-n[E_{trc}^{ub}(R,Q)-\epsilon]}\right] + \mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) \leq e^{-n[E_{trc}^{ub}(R,Q)+\epsilon]}\right]$$
(775)

$$\leq \frac{1}{n^{1+\kappa}} + \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n) \leq e^{-n[E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+\epsilon]}\right]$$
(776)

where (776) follows from [15, Eq. (22)] with $\gamma_n = n^{1+\kappa'}$ for some $\kappa' > 0$. Next, we bound the second term in (776) for large values of n.

Let $\Delta > 0$ small enough and define

$$P_{XX'}^* = \operatorname*{arg\,min}_{P_{XX'}:D(P_{XX'} \parallel Q_X Q_X') \le 2R} R + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x'), \tag{777}$$

$$\mathcal{A} = \left[|\mathcal{N}(P_{XX'}^*) - E\left[\mathcal{N}(P_{XX'}^*)\right]| > 2^{-n(D(P_{XX'}^*)|Q_XQ_X') - 2R + \Delta)} \right].$$
(778)

Then, on \mathcal{A}^c , by Lemma 24, we have

$$2\mathcal{N}(P_{XX'}^*)g_N(P_{XX'}^*) \doteq 2^{-n(D(P_{XX'}^*)|Q_XQ_X')-2R)}g_N(P_{XX'}^*)$$
(779)

$$\doteq 2^{-n\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}^*(x,x')}, \tag{780}$$

where (779) follows from Lemma 4, and (780) follows from Lemma 7 and Lemma 6 which proves that

 $D(P_{XX'}^* \| Q_X Q_{X'}) = 2R$

for the case $E_{\rm trc}(R,Q) > E_{\rm rce}(R,Q)$.

Then, we have

$$\mathbb{P}\left[P_{e}^{ub}(\mathcal{C}_{n}) \leq 2^{-n(E_{trc}^{ub}(R,Q)+\epsilon)}\right]$$
$$= \mathbb{P}\left[\frac{2}{M}\sum_{P_{XX'}}\mathcal{N}(P_{XX'})g_{N}(P_{XX'}) \leq 2^{-n(E_{trc}^{ub}(R,Q)+\epsilon)}\right]$$
(781)

$$\leq \mathbb{P}\left[\frac{2}{M}\mathcal{N}(P_{XX'}^*)g_N(P_{XX'}^*) \leq 2^{-n(E_{\mathrm{tre}}^{\mathrm{ub}}(R,Q)+\epsilon)}\right]$$
(782)

$$\leq \mathbb{P}\left[\frac{2}{M}\mathcal{N}(P_{XX'}^{*})g_{N}(P_{XX'}^{*}) \leq 2^{-n(R+\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}^{*}(x,x')+\epsilon)}\right]$$
(783)

$$\leq \mathbb{P}\left[2\mathcal{N}(P_{XX'}^{*})g_{N}(P_{XX'}^{*}) \leq 2^{-n(\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}^{*}(x,x')+\epsilon)}\right]$$
(784)

$$= \mathbb{P}\left[2\mathcal{N}(P_{XX'}^*)g_N(P_{XX'}^*) \le 2^{-n(\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}^*(x,x')+\epsilon)} \middle| \mathcal{A}\right] \mathbb{P}(\mathcal{A}) \\ + \mathbb{P}\left[2\mathcal{N}(P_{XX'}^*)g_N(P_{XX'}^*) \le 2^{-n(\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}^*(x,x')+\epsilon)} \middle| \mathcal{A}^c\right] \mathbb{P}(\mathcal{A}^c)$$

$$(785)$$

$$\leq \mathbb{P}(\mathcal{A}) + \mathbb{P}\left[2\mathcal{N}(P_{XX'}^*)g_N(P_{XX'}^*) \leq 2^{-n(\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}^*(x,x')+\epsilon)} \middle| \mathcal{A}^c\right]$$
(786)

$$\leq 2^{-n\Delta'} + 0 \tag{787}$$

$$=2^{-n\Delta'},\tag{788}$$

for *n* sufficiently large, where (783) follows from Lemma 7 and $E_{\rm trc}(R,Q) > E_{\rm rce}(R,Q)$, (787) follows from Lemma 24 and (780), respectively.

Hence, for this case, we have

$$\mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) > \frac{1}{2}2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)-\varepsilon)}\right] + \mathbb{P}\left[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_{n}) < 2^{-n(E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)+\varepsilon)}\right] \le \frac{1}{n^{1+\kappa'}} + 2^{-n\Delta'}$$
(789)

 $\text{for some }\kappa'>0 \text{ and }\Delta'>0.$

Finally, from (767) and (789), our proof is concluded.

H. Proof of Lemma 9

Observe that

$$\mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n})] \ge \mathbb{E}[P_{e}(\mathcal{C}_{n})]$$
(790)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \mathbb{E}\left[\mathbb{P}\left(\bigcup_{j \neq i} \{\boldsymbol{X}_i \to \boldsymbol{X}_j\}\right)\right]$$
(791)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \mathbb{E}\left[\mathbb{E}\left[\mathbf{1}\left\{\bigcup_{j \neq i} \{\mathbf{X}_i \to \mathbf{X}_j\}\right\}\right]\right]$$
(792)

$$\geq \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \frac{\left(\mathbb{E}\left[\mathbb{E}[\mathbf{1}\{\mathbf{X}_i \to \mathbf{X}_j\}]\right]\right)^2}{\mathbb{E}\left[\mathbb{E}[\mathbf{1}\{\mathbf{X}_i \to \mathbf{X}_j\}]\right] + \sum_{k \neq i,j} \mathbb{E}\left[\mathbb{E}[\mathbf{1}\{\{\mathbf{X}_i \to \mathbf{X}_j\} \cap \{\mathbf{X}_i \to \mathbf{X}_k\}\}]\right]}$$
(793)

$$= \frac{1}{M_n} \sum_{i=1}^{M_n} \sum_{j \neq i} \frac{\left(\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j]\right]\right)^2}{\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j]\right] + \sum_{k \neq i,j} \mathbb{E}\left[\mathbb{P}\left[\{\boldsymbol{X}_i \to \boldsymbol{X}_j\} \cap \{\boldsymbol{X}_i \to \boldsymbol{X}_k\}\right]\right]}$$
(794)

$$=\frac{(M_n-1)\left(\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_1\to\boldsymbol{X}_2]\right]\right)^2}{\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_1\to\boldsymbol{X}_2]\right]+(M_n-2)\mathbb{E}\left[\mathbb{P}[\{\boldsymbol{X}_1\to\boldsymbol{X}_2\}\cap\{\boldsymbol{X}_1\to\boldsymbol{X}_3\}]\right]}$$
(795)

$$= \frac{P_{e}^{ub}(\mathcal{C}_{n})\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_{1} \to \boldsymbol{X}_{2}]\right]}{\mathbb{E}\left[\mathbb{P}[\boldsymbol{X}_{1} \to \boldsymbol{X}_{2}]\right] + (M_{n} - 2)\mathbb{E}\left[\mathbb{P}[\{\boldsymbol{X}_{1} \to \boldsymbol{X}_{2}\} \cap \{\boldsymbol{X}_{1} \to \boldsymbol{X}_{3}\}]\right]},$$
(796)

where (793) follows from Caen's inequality in Lemma 2 by, for each fixed *i*, setting $\mathcal{I}_i = \{j \in [M] \setminus \{i\} : j \neq i\}, A_j^{(i)} = \{\mathbf{X}_i \to \mathbf{X}_j\}$ with the probability measure defined as $\mathbb{P}(A_j^{(i)}) = \mathbb{E}[\mathbb{E}[\mathbf{1}\{\mathbf{X}_i \to \mathbf{X}_j\}]] = \mathbb{E}[\mathbb{P}[\mathbf{X}_i \to \mathbf{X}_j]]$, where the inner expectation is over the BSC channel randomness and the outer one is over the random codebook ensemble. This is the probability of event $\{\mathbf{X}_i \to \mathbf{X}_j\}$ on the a product probability space generated from channel statistics and random codebook generations. By the symmetry of the codebook generation, it is easy to see that $\mathbb{P}(A_j^{(i)}) = \mathbb{E}[\mathbb{P}[\mathbf{X}_i \to \mathbf{X}_j]] = \mathbb{E}[\mathbb{P}[\mathbf{X}_1 \to \mathbf{X}_2]] = \mathbb{P}(A_2^{(1)})$ for all $j \neq i$.

From (796), it holds that

$$1 \le \frac{\mathbb{E}[P_{\mathrm{e}}^{\mathrm{ub}}(\mathcal{C}_n)]}{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)]}$$
(797)

$$\leq 1 + (M_n - 2) \frac{\mathbb{E}[\mathbb{P}[\{\boldsymbol{X}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{X}_1 \to \boldsymbol{X}_3\}]]}{\mathbb{E}[\mathbb{P}[\boldsymbol{X}_1 \to \boldsymbol{X}_2]]}.$$
(798)

Recall the definition of $d_{\mathrm{B}}(x, x')$ in (746). Assume that $x_1 \in \mathcal{T}(P_X)$ for some $P_X \in \mathcal{P}_n(\mathcal{X})$, which is a fixed vector. Then, given $(x_1, x_2) \in \mathcal{T}(P_{XX'})$ and $(x_1, x_3) \in \mathcal{T}(P_{XX''})$ where $P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})$ and $P_{XX''} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X})$, it holds that

$$\mathbb{P}\{\{\boldsymbol{x}_{1} \to \boldsymbol{x}_{2}\} \cap \{\boldsymbol{x}_{1} \to \boldsymbol{x}_{3}\} | (\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \in \mathcal{T}(P_{XX'}), (\boldsymbol{x}_{1}, \boldsymbol{x}_{3}) \in \mathcal{T}(P_{XX''})\} \\ \leq \min\left\{\mathbb{P}\{\boldsymbol{x}_{1} \to \boldsymbol{x}_{2} | (\boldsymbol{x}_{1}, \boldsymbol{x}_{2}) \in \mathcal{T}(P_{XX'})\}, \mathbb{P}\{\boldsymbol{x}_{1} \to \boldsymbol{x}_{3} | (\boldsymbol{x}_{1}, \boldsymbol{x}_{3}) \in \mathcal{T}(P_{XX''})\}\right\}$$
(799)

$$\leq \min\left\{2^{-n\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}(x,x')}, 2^{-n\sum_{x,x''}d_{\mathrm{B}}(x,x'')P_{XX''}(x,x'')}\right\}$$
(800)

$$=2^{-n\max\left\{\sum_{x,x'}d_{\mathrm{B}}(x,x')P_{XX'}(x,x'),\sum_{x,x''}d_{\mathrm{B}}(x,x')P_{XX''}(x,x')\right\}},$$
(801)

which does not depend on x_1, x_2, x_3 , where (800) follows from Lemma 4.

=

In addition, we have

$$\mathbb{P}\left[(\boldsymbol{x}_1, \boldsymbol{X}_2) \in \mathcal{T}(P_{XX'})\right] = \sum_{\boldsymbol{x}_2} \mathbb{P}(\boldsymbol{x}_2) \mathbf{1}\{(\boldsymbol{x}_1, \boldsymbol{x}_2) \in \mathcal{T}(P_{XX'})\}$$
(802)

$$= \sum_{\boldsymbol{x}_2} 2^{-n \left(H(P'_X) + D(P_{X'} || Q) \right)} \mathbf{1} \{ (\boldsymbol{x}_1, \boldsymbol{x}_2) \in \mathcal{T}(P_{XX'}) \}$$
(803)

$$=2^{-n\left(H(P'_{X})+D(P_{X'}\|Q)\right)}\sum_{\boldsymbol{x}_{2}}\mathbf{1}\{(\boldsymbol{x}_{1},\boldsymbol{x}_{2})\in\mathcal{T}(P_{XX'})\}$$
(804)

$$=2^{-n\left(H(P_X')+D(P_{X'}\|Q)\right)}\frac{|\mathcal{T}(P_{XX'}|)}{|\mathcal{T}(P_X)|}$$
(805)

$$\doteq 2^{-n \left(H(P_X') + D(P_{X'} \| Q) \right)} \frac{2^{NH(P_{XX'})|}}{2^{NH(P_X)|}}$$
(806)

$$=2^{-n\left(I_P(X;X')+D(P'_X||Q)\right)},$$
(807)

where (803) and (806) follow from [29].

Similarly, we also have

$$\mathbb{P}\left[\left(\boldsymbol{x}_{1}, \boldsymbol{X}_{3}\right) \in \mathcal{T}(P_{XX^{"}})\right] = 2^{-n\left(I_{P}(X; X^{"}) + D(P_{X}^{"} \| Q)\right)}.$$
(808)

Hence, we have

$$\mathbb{P}\left[\left\{ (\boldsymbol{x}_{1}, \boldsymbol{X}_{2}) \in \mathcal{T}(P_{XX'}) \right\} \cap \left\{ (\boldsymbol{x}_{1}, \boldsymbol{X}_{3}) \in \mathcal{T}(P_{XX''}) \right\} \middle| \boldsymbol{X}_{1} = \boldsymbol{x}_{1} \right]$$
$$= \mathbb{P}\left[\left\{ (\boldsymbol{x}_{1}, \boldsymbol{X}_{2}) \in \mathcal{T}(P_{XX'}) \right\} \cap \left\{ (\boldsymbol{x}_{1}, \boldsymbol{X}_{3}) \in \mathcal{T}(P_{XX''}) \right\} \right]$$
(809)

$$= \mathbb{P}\left[(\boldsymbol{x}_1, \boldsymbol{X}_2) \in \mathcal{T}(P_{XX'}) \right] \mathbb{P}\left[(\boldsymbol{x}_1, \boldsymbol{X}_3) \in \mathcal{T}(P_{XX''}) \right]$$
(810)

$$=2^{-n\left(I_P(X;X')+D(P'_X||Q)\right)}2^{-n\left(I_P(X;X'')+D(P'_X||Q)\right)}$$
(811)

$$=2^{-n(I_P(X;X')+I_P(X;X'')+D(P'_X||Q)+D(P_{X''}||Q))},$$
(812)

where (811) follows from (807) and (808).

It follows from (801) and (812) that

$$\mathbb{E}_{\boldsymbol{X}}[\mathbb{P}[\{\boldsymbol{x}_{1} \to \boldsymbol{X}_{2}\} \cap \{\boldsymbol{x}_{1} \to \boldsymbol{X}_{3}\}]] \\
= \sum_{P_{X'|X}} \sum_{P_{X'|X}} \mathbb{E}\left[\mathbb{P}\{\{\boldsymbol{x}_{1} \to \boldsymbol{X}_{2}\} \cap \{\boldsymbol{x}_{1} \to \boldsymbol{X}_{3}\} \middle| (\boldsymbol{x}_{1}, \boldsymbol{X}_{2}) \in \mathcal{T}_{P_{XX'}}, (\boldsymbol{x}_{1}, \boldsymbol{X}_{3}) \in \mathcal{T}_{P_{XX''}}\} \\
\times \mathbb{P}\left[\{(\boldsymbol{x}_{1}, \boldsymbol{X}_{2}) \in \mathcal{T}(P_{XX'})\} \cap \{(\boldsymbol{x}_{1}, \boldsymbol{X}_{3}) \in \mathcal{T}(P_{XX''})\} \middle| \boldsymbol{X}_{1} = \boldsymbol{x}_{1}\right]\right]$$

$$\leq \sum_{P_{X'|X}} \sum_{P_{X''|X}} 2^{-n \max\{\sum_{x,x'} d_{B}(x,x')P_{XX'}(x,x'), \sum_{x,x''} d_{B}(x,x'')P_{XX''}(x,x'')\}}$$
(813)

$$\times 2^{-n(I_P(X;X')+I_P(X;X'')+D(P'_X||Q)+D(P''_X||Q))}$$
(814)

$$\leq \sum_{P_{X'|X}} \sum_{P_{X''|X}} 2^{-\frac{n}{2} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') + \sum_{x,x''} d_{\mathrm{B}}(x,x') P_{XX''}(x,x'') \right)} \times 2^{-n(I_{P}(X;X') + I_{P}(X;X'') + D(P'_{X} ||Q) + D(P_{X''} ||Q))}$$
(815)

$$\times 2^{-n(I_P(X;X')+I_P(X;X'')+D(P'_X ||Q)+D(P_{X''} ||Q))}$$
(815)

$$= \left(\sum_{P_{X'|X}} 2^{-\frac{n}{2} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')\right)} 2^{-n\left(I_{P}(X;X')+D(P'_{X}||Q)\right)}\right) \times \left(\sum_{P_{X''}|X} 2^{-\frac{n}{2} \left(\sum_{x,x''} d_{\mathrm{B}}(x,x'')P_{XX''}(x,x'')\right)} 2^{-n\left(I_{P}(X;X'')+D(P_{X''}||Q)\right)}\right)$$
(816)

$$= \left(\sum_{P_{X'|X}} 2^{-\frac{n}{2} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x')\right)} 2^{-n \left(I_{P}(X;X') + D(P'_{X} \| Q)\right)}\right)^{2},$$
(817)

where (815) follows from $\max\{a, b\} \ge \frac{a+b}{2}$.

It follows from (817) that

$$\mathbb{E}[\mathbb{P}[\{\boldsymbol{X}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{X}_1 \to \boldsymbol{X}_3\}]]$$

= $\sum_{\boldsymbol{x}_1} \mathbb{P}(\boldsymbol{x}_1) \mathbb{E}[\mathbb{P}[\{\boldsymbol{x}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{x}_1 \to \boldsymbol{X}_3\}] | \boldsymbol{X}_1 = \boldsymbol{x}_1]$ (818)

$$=\sum_{\boldsymbol{x}_1} \mathbb{P}(\boldsymbol{x}_1) \mathbb{E}[\mathbb{P}[\{\boldsymbol{x}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{x}_1 \to \boldsymbol{X}_3\}]]$$
(819)

$$=\sum_{P_X}\sum_{\boldsymbol{x}_1\in\mathcal{T}(P_X)}\mathbb{P}(\boldsymbol{x}_1)\mathbb{E}[\mathbb{P}[\{\boldsymbol{x}_1\to\boldsymbol{X}_2\}\cap\{\boldsymbol{x}_1\to\boldsymbol{X}_3\}]]$$
(820)

$$= \sum_{P_X} \sum_{\boldsymbol{x}_1 \in \mathcal{T}(P_X)} 2^{-n(D(P_X || Q) + H(P_X))} \mathbb{E}[\mathbb{P}[\{\boldsymbol{x}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{x}_1 \to \boldsymbol{X}_3\}]]$$
(821)

$$\leq \sum_{P_{X}} \sum_{\boldsymbol{x}_{1} \in \mathcal{T}(P_{X})} 2^{-n(D(P_{X} \| Q) + H(P_{X}))} \left(\sum_{P_{X'|X}} 2^{-\frac{n}{2} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') \right)} 2^{-n \left(I_{P}(X;X') + D(P'_{X} \| Q) \right)} \right)^{2}$$

$$(822)$$

$$\leq \sum_{P_X} 2^{-nD(P_X \parallel Q)} \bigg(\sum_{P_{X'\mid X}} 2^{-\frac{n}{2} \left(\sum_{x,x'} d_{\mathrm{B}}(x,x')P_{XX'}(x,x') \right)} 2^{-n \left(I_P(X;X') + D(P'_X \parallel Q) \right)} \bigg)^2, \tag{823}$$

where (819) follows from the independence of codewords in the random codebook ensemble, (821) follows from [29].

Now, from (772) in the proof of Lemma 24, for all joint type $P_{XX'}$ such that $D(P_{XX'} || Q_X Q_{X'}) > 2R$, it holds that

$$\sum_{n=1}^{\infty} \mathbb{P}\Big[\mathcal{N}(P_{XX'}) \ge 1\Big] \le \sum_{n=1}^{\infty} 2^{-n\left(D(P_{XX'} \| Q_X Q'_X) - 2R\right)} < \infty.$$
(824)

From (824) and Borel-Cantelli's lemma [23], it holds almost surely that $\mathcal{N}(P_{XX'}) = 0$ for all joint type $P_{XX'}$ such that $D(Q_{XX'} || Q_X Q_{X'}) > 2R$.

Hence, from (817) and the above fact with noting the number of types or conditional types are sub-exponential in N, we have

$$\mathbb{E}[\mathbb{P}[\{X_1 \to X_2\} \cap \{X_1 \to X_3\}]] \\ \doteq 2^{-n\left(\min_{P_{XX'} \in \mathcal{P}_n(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \parallel Q_X Q'_X) \le 2R} D(P_X \parallel Q) + 2\left(I_P(X; X') + D(P'_X \parallel Q)\right) + \sum_{x,x'} d_B(x,x') P_{XX'}(x,x')\right)}$$

$$(825)$$

$$= 2^{-n(\min_{P_{XX'} \in \mathcal{P}(X \times X): D(P_{XX'} \| Q_X Q_X') \le 2R D(|X| \| Q) + 2(P(X, X) + D(|X| \| Q)) + \sum_{x,x'} \alpha_{B}(w, w) + X_{XX'}(w, w))},$$
(826)

where (825) follows from the sub-exponential number of possible *n*-types in $\mathcal{X} \times \mathcal{X}$ [29], and (826) follows from the fact that $\mathcal{P}_n(\mathcal{X} \times \mathcal{X})$ is dense in $\mathcal{P}(\mathcal{X} \times \mathcal{X})$.

Now, note that $Q_X = Q'_X = Q$, so we have

$$I_P(X;X') = D(P_{XX'} || P_X P_X')$$
(827)

$$= D(P_{XX'} \| Q_X Q'_X) - D(P_X \| Q) - D(P'_X \| Q).$$
(828)

It follows that

$$D(P_X \| Q) + 2 (I_P(X; X') + D(P'_X \| Q))$$

= $D(P_X \| Q) + 2 (D(P_{XX'} \| Q_X Q'_X) - D(P_X \| Q))$ (829)

$$= 2D(P_{XX'} || Q_X Q'_X) - D(P_X || Q)$$
(830)

$$\geq D(P_{XX'} \| Q_X Q_X'),\tag{831}$$

where (831) follows from the data processing for KL divergence (or log-sum inequality [34]).

Hence, we have

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R} D(P_X \| Q) + 2 \left(I_P(X; X') + D(P'_X \| Q) \right) + \sum_{x,x'} d_B(x, x') P_{XX'}(x, x')$$

$$\geq \min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q'_X) \le 2R} D(P_{XX'} \| Q_X Q'_X) + \sum_{x,x'} d_B(x, x') P_{XX'}(x, x')$$

$$= E_{\text{trc}}(R, Q) + R,$$
(833)

where (832) follows from (831), and (833) follows from Lemma 7. Note that (832) becomes equality if and only if $P_{XX'}(x, x') = Q(x)Q(x')$ for all $x, x' \in \mathcal{X} \times \mathcal{X}$. However, at $P_{XX'} = Q_X Q'_X$, we have

$$\min_{P_{XX'} \in \mathcal{P}(X \times X) : D(P_{XX'} \| Q_X Q'_X) \le 2R} D(P_X \| Q) + 2 \left(I_P(X; X') + D(P'_X \| Q) \right) + \sum_{x, x'} d_B(x, x') P_{XX'}(x, x')$$

$$\sum_{X, x' \in \mathcal{P}(X \times X) : D(P_X | Q_X Q'_X) \le 2R} D(P_X \| Q) + 2 \left(I_P(X; X') + D(P'_X \| Q) \right) + \sum_{x, x'} d_B(x, x') P_{XX'}(x, x')$$
(224)

$$= \sum_{x,x'} d_{\rm B}(x,x')Q(x)Q(x')$$
(834)

$$= -\sum_{x,x'} \log\left(\sum_{y\in\mathcal{Y}} \sqrt{W(y|x)W(y|x')}\right) Q(x)Q(x')$$
(835)

$$> -\log\left(\sum_{x,x'}\sum_{y\in\mathcal{Y}}\sqrt{W(y|x)W(y|x')}Q(x)Q(x')\right)$$
(836)

$$= -\log\left(\sum_{y\in\mathcal{Y}}\sum_{x,x'}\sqrt{W(y|x)W(y|x')}Q(x)Q(x')\right)$$
(837)

$$= -\log\left(\sum_{y\in\mathcal{Y}}\left(\sum_{x}\sqrt{W(y|x)}Q(x)\right)^{2}\right)$$
(838)

$$= -\log\left(\sum_{y\in\mathcal{Y}}\left(\sum_{x}\sqrt{W(y|x)}Q(x)\right)^{2}\right)$$
(839)

$$=R_0(Q) \tag{840}$$

$$=E_{\rm rce}(R,Q)+R,\tag{841}$$

where (836) follows from the convexity of the function $-\log x$ with noting that the equality does not happen by the condition (836), and (840) follows from [28, Eq. (8.45)] with $R_0(Q)$ is the cut-off rate of the DMC at the distribution Q.

Therefore, from (833) and (841), we have

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}): D(P_{XX'} \| Q_X Q_X') \le 2R} D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} d_{\mathcal{B}}(x,x') P_{XX'}(x,x') > E_{\text{trc}}(R,Q) + R$$
(842)

for the case $E_{\rm trc}^{\rm ub}(R,Q) = E_{\rm rce}(R,Q).$

Now, for the case $E_{\text{trc}}^{\text{ub}}(R,Q) > E_{\text{rce}}(R,Q)$, (833) happens at the optimizer $P_{XX'}^*$ satisfying $D(P_{XX'}^* || Q_X Q_X') = 2R$, which leads to $P_{XX'}^* \neq Q_X Q_X'$ if R > 0, so the equality can not happen in (831).

In summary, at R > 0 and a fixed underlying distribution Q, it holds that

$$\min_{P_{XX'} \in \mathcal{P}(\mathcal{X} \times \mathcal{X}) : D(P_{XX'} \| Q_X Q_X') \le 2R} D(P_{XX'} \| Q_X Q_X') + \sum_{x,x'} d_{\mathrm{B}}(x,x') P_{XX'}(x,x') > E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q) + R.$$
(843)

Hence, it holds from (826) and (56) that

$$\mathbb{E}\left[\mathbb{P}\left[\{\boldsymbol{X}_1 \to \boldsymbol{X}_2\} \cap \{\boldsymbol{X}_1 \to \boldsymbol{X}_3\}\right] \le 2 \times 2^{-\left(R + E_{\mathrm{trc}}^{\mathrm{ub}}(R,Q)\right)n} 2^{-\delta(R)n}$$
(844)

for some constant $\delta(R) > 0$.

Now, on the other hand, we know that

$$\mathbb{E}[\mathbb{P}[\boldsymbol{X}_1 \to \boldsymbol{X}_2]] = \frac{1}{M_n(M_n - 1)} \sum_{i=0}^{M_n - 1} \sum_{j \neq i} \mathbb{E}[\mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j]]$$
(845)

$$= \frac{1}{M_n - 1} \mathbb{E}\left[\frac{1}{M_n} \sum_{i=0}^{M_n - 1} \sum_{j \neq i} \mathbb{P}[\boldsymbol{X}_i \to \boldsymbol{X}_j]\right]$$
(846)

$$=\frac{\mathbb{E}[P_{\rm e}^{\rm ub}(\mathcal{C}_n)]}{M_n - 1} \tag{847}$$

$$\geq \frac{\mathbb{E}[P_{\mathrm{e}}(\mathcal{C}_n)]}{M_n - 1} \tag{848}$$

$$\doteq 2^{-n(E_{\rm rce}(R,Q)-R)}.$$
(849)

From (798), (844), and (849), we obtain

$$0 \le \frac{\mathbb{E}[P_{e}^{ub}(\mathcal{C}_{n})]}{\mathbb{E}[P_{e}(\mathcal{C}_{n})]} - 1$$
(850)

$$\doteq 2^{-N(\delta(R) + E_{\rm trc}^{\rm ub}(R,Q) - E_{\rm rce}(R,Q))}.$$
(851)

I. Proof of Lemma 11

Observe that

$$\mathbb{E}\left[2^{-tS_n/n}\right] = \sum_{x_1, x_2, \cdots, x_n} 2^{-\frac{t(x_1 + x_2 + \cdots + x_n)}{n}} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n]$$

$$= \sum_{x_1, x_2, \cdots, x_n} 2^{-\frac{t(x_1 + x_2 + \cdots + x_n)}{n}} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \mathbf{1}\{(x_1, x_2, \cdots, x_n) \in \mathcal{V}\}$$

$$+ \sum_{x_1, x_2, \cdots, x_n} 2^{-\frac{t(x_1 + x_2 + \cdots + x_n)}{n}} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \mathbf{1}\{(x_1, x_2, \cdots, x_n) \in \mathcal{V}^c\}$$

$$(853)$$

$$\leq \sum_{x_1, x_2, \cdots, x_n} 2^{-\frac{t(x_1 + x_2 + \cdots + x_n)}{n}} \prod_{k=1}^n P(x_k) \mathbf{1}\{(x_1, x_2, \cdots, x_n) \in \mathcal{V}\}$$

$$+\sum_{x_1,x_2,\cdots,x_n} 2^{-\frac{t(x_1+x_2+\cdots+x_n)}{n}} \mathbf{1}\{(x_1,x_2,\cdots,x_n) \in \mathcal{V}^c\}$$
(854)

$$\leq \sum_{x_1, x_2, \cdots, x_n} 2^{-\frac{t(x_1 + x_2 + \cdots + x_n)}{n}} \prod_{k=1}^n P(x_k) + \sum_{x_1, x_2, \cdots, x_n} 2^{-n\zeta} \mathbf{1}\{(x_1, x_2, \cdots, x_n) \in \mathcal{V}^c\}$$
(855)

$$= \mathbb{E}_{\prod_{i=1}^{n} P(x_i)} \left[2^{-tS_n/n} \right] + 2^{-n\zeta} |\mathcal{V}^c|.$$
(856)

J. Proof of Lemma 12

We have that

$$\mathbb{E}[\mathcal{I}\{i,j\}\mathcal{I}\{i,k\}] = \mathbb{P}\{(X_i, X_j) \in \mathcal{T}(P_{XX'}), (X_i, X_k) \in \mathcal{T}(P_{XX'})\}$$
(857)

$$= \sum_{\boldsymbol{x}_i} \mathbb{P}\{X_i = \boldsymbol{x}_i\} \mathbb{P}\{(\boldsymbol{x}_i, X_j) \in \mathcal{T}(P_{XX'}), (\boldsymbol{x}_i, X_k) \in \mathcal{T}(P_{XX'})\}$$
(858)

$$= \sum_{\boldsymbol{x}_i} \mathbb{P}\{X_i = \boldsymbol{x}_i\} \mathbb{P}\{(\boldsymbol{x}_i, X_j) \in \mathcal{T}(P_{XX'})\} \mathbb{P}\{(\boldsymbol{x}_i, X_k) \in \mathcal{T}(P_{XX'})\}$$
(859)

$$= \sum_{\boldsymbol{x}_i} \mathbb{P}\{X_i = \boldsymbol{x}_i\} \mathbb{P}\{(\boldsymbol{x}_i, X_j) \in \mathcal{T}(P_{XX'})\}^2$$
(860)

$$=\sum_{P_X} \mathcal{N}(P_X) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_X)\} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\}^2$$
(861)

$$\doteq \max_{P_X} \mathcal{N}(P_X) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_X)\} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\}^2$$
(862)

where in (858) we conditioned to codeword X_i being equal to a given realization x_i , (859) is because X_j and X_k are independent, (860) is because they are also identically distributed, in (861) we grouped codewords X_i according to their type P_X and used the fact that $\mathbb{P}\{(x_i, X_j) \in \mathcal{T}(P_{XX'})\}$ takes the same value when x_i has the same type. Expression (862) is hard to calculate because of the term $\mathbb{P}\{(x, X_j) \in \mathcal{T}(P_{XX'}) | x \in \mathcal{T}(P_X)\}$. Therefore we find a lower bound and an upper bound on Eqn. (861). The lower bound is:

$$\sum_{P_X} \mathcal{N}(P_X) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_X)\} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\}^2$$
$$\geq \left(\sum_{P_X} \mathcal{N}(P_X) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_X)\} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\}\right)^2$$
(863)

$$= \mathbb{P}\{(X_i, X_j) \in \mathcal{T}(P_{XX'})\}^2$$
(864)

$$\doteq 2^{-n2D(P_{XX'}||Q_XQ'_X)} \tag{865}$$

while the upper bound is:

$$\sum_{P_{X}} \mathcal{N}(P_{X}) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_{X})\} \mathbb{P}\{(\boldsymbol{x}, X_{j}) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_{X})\}^{2}$$

$$\leq \sum_{P_{X}} \mathcal{N}(P_{X}) \mathbb{P}\{\boldsymbol{x} \in \mathcal{T}(P_{X})\} \mathbb{P}\{(\boldsymbol{x}, X_{j}) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_{X})\}$$

$$\times \cdot \max_{P_{X}} \mathbb{P}\{(\boldsymbol{x}, X_{j}) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_{X})\}$$
(866)

$$= \mathbb{P}\{(X_i, X_j) \in \mathcal{T}(P_{XX'})\} \max_{P_X} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\}$$
(867)

$$= 2^{-n[D(P_{XX'}||Q_XQ'_X)+\eta]}$$
(868)

where

$$\eta = -\frac{1}{n} \log \max_{P_X} \mathbb{P}\{(\boldsymbol{x}, X_j) \in \mathcal{T}(P_{XX'}) | \boldsymbol{x} \in \mathcal{T}(P_X)\} \le D(P_{XX'} || Q_X Q_X')$$

and the inequality follows from (865).

K. Proof of Lemma 13

The proof is based on [35, Th. 10]. A similar proof of an equivalent result is presented for the case of constant composition codes in [12]. However, there are several differences with our case. First of all our Lemma 12 gives a bound rather than a dot equality, which has implications on the minimum exponent starting from which a double exponential decay is found. Other differences with [12] are indicated in the following.

Let us define the quantities:

$$a = 2^{-n\epsilon} \tag{869}$$

$$\Delta = \mathbb{E}[\mathcal{N}(P_{XX'})] \doteq 2^{n(2R - D(P_{XX'}||Q_XQ'_X))}.$$
(870)

Let us consider a graph in which vertices are indicated with pairs (e.g., (k, l) is a vertex). Two vertices (k, l) and (i, j) are connected if exactly one index in both pairs coincide. Let us indicate with $(k, l) \sim (i, j)$ the case in which vertices (k, l) and (i, j) are connected. Let us also indicate with $[M]^2_*$ the set $\{i, j \in \{1, 2, ..., M_n\} : i \neq j\}$. Let us define and bound the following quantity:

$$\Theta = \frac{1}{2} \sum_{(i,j)\in[M]^2_*} \sum_{(k,l)\in[M]^2_*,(k,l)\sim(i,j)} \mathbb{E}[\mathcal{I}\{i,j\}\mathcal{I}\{i,k\}]$$
(871)

$$\doteq \frac{1}{2} 2^{2nR} (2^{nr} + 2^{nr} - 2) \mathbb{E}[\mathcal{I}\{1, 2\} \mathcal{I}\{1, 3\}]$$
(872)

$$\leq \frac{1}{2} 2^{2nR} (2^{nr} + 2^{nr} - 2) 2^{-nD(P_{XX'}||Q_XQ'_X)}$$
(873)

$$= \frac{1}{2} 2^{n(3R - D(P_{XX'}||Q_XQ'_X))}$$
(874)

where (872) and (873) follow from Lemma 12 as well as from the fact that codewords are i.i.d. and noticing that there are about M_n^2 codeword pairs (i, j) and, for each of them, there are exactly $2^{nR} - 1$ connected vertices. Note that in [12] a dot equality rather than an inequality is present.

Now let us define and, where needed, bound the following three quantities:

$$\Omega = \max_{(i,j)\in[M]^2_*} \sum_{(k,l)\in[M]^2_*, (k,l)\sim(i,j)} \mathbb{E}[\mathcal{I}\{i,k\}]$$
(875)

$$\doteq (2^{nr} + 2^{nr} - 2)2^{-nD(P_{XX'}||Q_XQ'_X)}$$
(876)

$$= 2^{n(R-D(P_{XX'}||Q_XQ'_X))}$$
(877)

$$\frac{\Delta}{6\Omega} \doteq \frac{2^{n[2R-D(P_{XX'}||Q_XQ'_X)]}}{2^{n[R-D(P_{XX'}||Q_XQ'_X)]}} = 2^{nR}$$
(878)

$$\frac{\Delta^2}{8\Theta + 2\Delta} \doteq \frac{2^{n[4R - 2D(P_{XX'}||Q_XQ'_X)]}}{e^{n[3R - D(P_{XX'}||Q_XQ'_X)]} + 2^{n[2R - D(P_{XX'}||Q_XQ'_X)]}}$$
(879)

$$=\frac{2^{n[2R-D(P_{XX'}||Q_XQ'_X)]}}{2^{nR}+1}$$
(880)

$$\doteq 2^{n[R-D(P_{XX'}||Q_XQ'_X)]}$$
(881)

(882)

Using the definitions above and the result in [35, Th. 10] we obtain:

$$\mathbb{P}\left[\mathcal{N}(P_{XX'}) \le 2^{-n\epsilon} \mathbb{E}[\mathcal{N}(P_{XX'})]\right] \doteq \mathbb{P}\left[\mathcal{N}(P_{XX'}) \le 2^{n[2R-D(P_{XX'}||Q_XQ'_X)-\epsilon]}\right]$$
(883)

$$\leq \exp\left\{-\min\left(2^{n[R-D(P_{XX'}||Q_XQ'_X)]}, 2^{nR}\right)\right\}$$
(884)

$$= \exp\left\{-2^{n[R-D(P_{XX'}||Q_XQ'_X)]}\right\}$$
(885)

which concludes the proof of the lemma.

L. Proof of Lemma 14

We prove by induction. The condition (294) shows that (295) holds for n = 2, $\forall A_1, A_2 \subset \mathcal{X}$. Now, assume that (295) holds for some $n \ge 2$. We need to show that (295) holds for n + 1. Indeed, let $A_1, A_2, \dots, A_n, A_{n+1} \in \mathcal{X}$ such that there exists $i, j \in [n]$ with $i \ne j$ such that $A_i \cap A_j = \emptyset$.

Now, by reordering $\{A_k\}_{k=1}^n$, we can assume without loss of generality that $i, j \in [n]$. This leads to $\bigcap_{k=1}^n A_k = \emptyset$. Observe that

$$\mathbb{P}\left[\bigcap_{k=1}^{n+1} \left\{ X \in \mathcal{A}_k \right\}\right] = \mathbb{P}\left[\bigcap_{k=1}^{n} \left\{ X \in \mathcal{A}_k \right\}\right] \mathbb{P}\left[X \in \mathcal{A}_{n+1} \middle| X \in \bigcap_{k=1}^{n} \mathcal{A}_k\right]$$
(886)

$$\leq \mathbb{P}\left[\bigcap_{k=1}^{n} \left\{X \in \mathcal{A}_{k}\right\}\right] \mathbb{P}\left[X \in \mathcal{A}_{n+1}\right]$$
(887)

$$\leq \beta \prod_{k=1}^{n} \mathbb{P}[X \in \mathcal{A}_{k}] \mathbb{P}[X \in \mathcal{A}_{n+1}]$$
(888)

$$=\beta\prod_{k=1}^{n+1}\mathbb{P}[X\in\mathcal{A}_k].$$
(889)

This concludes our proof by induction.

M. Proof of Lemma 15

We consider four cases:

• Case 1: a = 1, b = 1. Then, we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = a\} \cap \{Z_{\tilde{Q}_{XX'}} = b\}\big] = 0.$$
(890)

On the other hand, we have

$$\mathbb{P}\big[Z_{Q_{XX'}} = 1\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 1\big] = \mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\big]\mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(\tilde{Q}_{XX'})\big]$$
(891)

$$\doteq 2^{-nI_{Q_{XX'}}(X;X')} 2^{-nI_{\tilde{Q}_{XX'}}(X;X')}, \tag{892}$$

where (892) follows from [12]. Hence, we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = 1\} \cap \{Z_{\tilde{Q}_{XX'}} = 1\}\big] \le \mathbb{P}\big[Z_{Q_{XX'}} = 1\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 1\big]$$
(893)

$$\leq \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}[Z_{Q_{XX'}} = 1] \mathbb{P}[Z_{\tilde{Q}_{XX'}} = 1],$$
(894)

hence, (296) holds for this case.

• Case 2: a = 1, b = 0. Then, we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = 1\} \cap \{Z_{\tilde{Q}_{XX'}} = 0\}\big] = \mathbb{P}[\{(X_i, X_j) \in \mathcal{T}(Q_{XX'})\} \cap \{\{(X_i, X_j) \notin \mathcal{T}(\tilde{Q}_{XX'})\}\}]$$
(895)

$$= \mathbb{P}[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})]$$
(896)

$$\doteq 2^{-nI_{Q_{XX'}}(X;X')}.$$
(897)

On the other hand, we have

$$\mathbb{P}\big[Z_{Q_{XX'}} = 1\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 0\big] = \mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \in \mathcal{T}(Q_{XX'})\big]\mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \notin \mathcal{T}(\tilde{Q}_{XX'})\big]$$
(898)

$$\doteq 2^{-nI_Q} XX'}(X;X') \left(1 - 2^{-nI_{\bar{Q}}} XX'}(X;X')\right).$$
(899)

Now, there are two sub-cases. If $I_{\tilde{Q}_{XX'}}(X;X')=0,$ from (897) and (899) we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = 1\} \cap \{Z_{\tilde{Q}_{XX'}} = 0\}\big] = \mathbb{P}\big[Z_{Q_{XX'}} = 1\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 0\big]$$
(900)

$$\leq \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}[Z_{Q_{XX'}} = 1] \mathbb{P}[Z_{\tilde{Q}_{XX'}} = 0].$$
(901)

On the other hand, if $I_{\tilde{Q}_{XX^{\prime}}}(X;X^{\prime})>0,$ we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}}=1\} \cap \{Z_{\tilde{Q}_{XX'}}=0\}\big] = \frac{1}{1-2^{-nI_{\tilde{Q}_{XX'}}(X;X')}} \mathbb{P}\big[Z_{Q_{XX'}}=1\big] \mathbb{P}\big[Z_{\tilde{Q}_{XX'}}=0\big]$$
(902)

$$\leq \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}[Z_{Q_{XX'}} = 1] \mathbb{P}[Z_{\tilde{Q}_{XX'}} = 0],$$
(903)

where (903) follows from $I_{\tilde{Q}_{XX'}}(X;X') \ge I_{\min}(Q)$.

• Case 3: a = 0, b = 1. Similarly as case 2, it holds that

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = 1\} \cap \{Z_{\tilde{Q}_{XX'}} = 0\}\big] = \mathbb{P}\big[Z_{Q_{XX'}} = 1\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 0\big]$$
(904)

$$\leq \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}[Z_{Q_{XX'}} = 1] \mathbb{P}[Z_{\tilde{Q}_{XX'}} = 0].$$
(905)

• Case 4: a = 0, b = 0. Then, we have

$$\mathbb{P}\left[\{Z_{Q_{XX'}} = 0\} \cap \{Z_{\tilde{Q}_{XX'}} = 0\}\right]$$
$$= \mathbb{P}\left[\{(\boldsymbol{X}_i, \boldsymbol{X}_j) \notin \mathcal{T}(Q_{XX'})\} \cap \{\{(\boldsymbol{X}_i, \boldsymbol{X}_j) \notin \mathcal{T}(\tilde{Q}_{XX'})\}\}\right]$$
(906)

$$= \mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \notin \mathcal{T}(Q_{XX'}) \big] + \mathbb{P}\big[(\boldsymbol{X}_i, \boldsymbol{X}_j) \notin \mathcal{T}(\tilde{Q}_{XX'}) \big]$$

$$-\mathbb{P}\left[\left\{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \notin \mathcal{T}(Q_{XX'}) \right\} \cup \left\{ \left\{ (\boldsymbol{X}_{i}, \boldsymbol{X}_{j}) \notin \mathcal{T}(Q_{XX'}) \right\} \right\} \right]$$
(907)
= $\left(1 - 2^{-nI_{Q_{XX'}}(X;X')}\right) + \left(1 - 2^{-nI_{\bar{Q}_{XX'}}(X;X')}\right)$

$$-\left(1 - \mathbb{P}\left[\left\{\left(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}\right) \in \mathcal{T}(Q_{XX'})\right\} \cap \left\{\left(\boldsymbol{X}_{i}, \boldsymbol{X}_{j}\right) \in \mathcal{T}(\tilde{Q}_{XX'})\right\}\right]\right)$$
(908)

$$= \left(1 - 2^{-nI_{Q_{XX'}}(X;X')}\right) + \left(1 - 2^{-nI_{\tilde{Q}_{XX'}}(X;X')}\right) - 1$$
(909)

$$=1-2^{-nI_{Q_{XX'}}(X;X')}-2^{-nI_{\tilde{Q}_{XX'}}(X;X')}.$$
(910)

On the other hand, we also have

$$\mathbb{P}[Z_{Q_{XX'}} = 0] \mathbb{P}[Z_{\bar{Q}_{XX'}} = 0]$$

= $(1 - 2^{-nI_{Q_{XX'}}(X;X')})(1 - 2^{-nI_{\bar{Q}_{XX'}}(X;X')})$ (911)

$$= 1 - 2^{-nI_{Q_{XX'}}(X;X')} - 2^{-nI_{\bar{Q}_{XX'}}(X;X')} + 2^{-nI_{Q_{XX'}}(X;X')} 2^{-nI_{\bar{Q}_{XX'}}(X;X')}.$$
(912)

From (910) and (912), we have

$$\mathbb{P}\big[\{Z_{Q_{XX'}} = 0\} \cap \{Z_{\tilde{Q}_{XX'}} = 0\}\big] \le \mathbb{P}\big[Z_{Q_{XX'}} = 0\big]\mathbb{P}\big[Z_{\tilde{Q}_{XX'}} = 0\big]$$
(913)

$$\leq \frac{1}{1 - 2^{-nI_{\min}}(Q)} \mathbb{P}[Z_{Q_{XX'}} = 1] \mathbb{P}[Z_{\tilde{Q}_{XX'}} = 0].$$
(914)

From the four cases above, we finally obtain (296)

$$\frac{1}{n^{3/2}} \sum_{i=1}^{n} \mathbb{E}[|X_i^3|] \to 0, \tag{915}$$

$$\frac{1}{n^2} \sum_{i=1}^{n} \mathbb{E}[X_i^4] \to 0, \tag{916}$$

N. Proof of Lemma 16

The proof is based on a modification of a proof based on Stein's method in [25, Lemma 3.2]. Without loss of generality, (or by scaling), we can assume that $\sum_{i=1}^{n} \mathbb{E}[X_i^2] = n$.

Let

$$T = \frac{S_n}{\sqrt{n}},\tag{917}$$

and

$$T_i = \frac{1}{\sqrt{n}} \sum_{j \neq i} X_j, \quad \forall i \in [n].$$
(918)

Observe that

$$\mathbb{E}[Tf(T)] = \frac{1}{\sqrt{n}} \mathbb{E}\left[\sum_{i=1}^{n} X_i(f(T) - f(T_i)) - (T - T_i)f'(T)\right] \\ + \frac{1}{\sqrt{n}} \mathbb{E}\left[\sum_{i=1}^{n} X_i(T - T_i)f'(T)\right] + \frac{1}{\sqrt{n}} \mathbb{E}\left[\sum_{i=1}^{n} X_if(T_i)\right].$$
(919)

Now, we have

$$\mathbb{E}[Tf(T) - f'(T)] = \frac{1}{\sqrt{n}} \mathbb{E}\bigg[\sum_{i=1}^{n} X_i \big(f(T) - f(T_i) - (T - T_i)f'(T)\big)\bigg] \\ + \frac{1}{\sqrt{n}} \mathbb{E}\bigg[\sum_{i=1}^{n} X_i (T - T_i)f'(T)\bigg] + \frac{1}{\sqrt{n}} \mathbb{E}\bigg[\sum_{i=1}^{n} X_i f(T_i)\bigg] - \mathbb{E}[f'(T)]$$
(920)
$$\le \bigg|\frac{1}{\sqrt{n}} \mathbb{E}\bigg[\sum_{i=1}^{n} X_i \big(f(T) - f(T_i) - (T - T_i)f'(T)\big)\bigg]\bigg|$$

$$+\left|\frac{1}{\sqrt{n}}\mathbb{E}\left[\sum_{i=1}^{n}X_{i}f(T_{i})\right]\right|+\left|\mathbb{E}\left[f'(T)\left(1-\frac{1}{\sqrt{n}}\sum_{i=1}^{n}X_{i}(T-T_{i})\right)\right]\right|$$
(921)

$$\leq \frac{\|f''\|}{2\sqrt{n}} \sum_{i=1}^{n} \mathbb{E}|X_i(T-T_i)^2| + \frac{1}{\sqrt{n}} \left| \mathbb{E} \left[\sum_{i=1}^{n} X_i f(T_i) \right] \right| + \frac{\|f'\|}{n} \mathbb{E} \left| \sum_{i=1}^{n} (1-X_i^2) \right|$$
(922)

$$\leq \frac{\|f''\|}{2\sqrt{n}} \sum_{i=1}^{n} \mathbb{E}|X_i^3| + \frac{1}{\sqrt{n}} \left| \mathbb{E}\left[\sum_{i=1}^{n} X_i f(T_i)\right] \right| + \frac{\|f'\|}{n} \mathbb{E}\left|\sum_{i=1}^{n} (1 - X_i^2)\right|.$$
(923)

Now, observe that

$$\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\mathbb{E}[X_if(T_i)] = \frac{1}{\sqrt{n}}\sum_{x_1,x_2,\cdots,x_n}\mathbb{P}[X_1 = x_1, X_2 = x_2,\cdots,X_n = x_n]\sum_{i=1}^{n}x_if(t_i)$$
(924)

$$= \frac{1}{\sqrt{n}} \sum_{x_1, x_2, \cdots, x_n \in \mathcal{V}} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \sum_{i=1}^n x_i f(t_i)$$
(925)

$$+\frac{1}{\sqrt{n}}\sum_{x_1,x_2,\cdots,x_n\in\mathcal{V}^c}\mathbb{P}[X_1=x_1,X_2=x_2,\cdots,X_n=x_n]\sum_{i=1}^n x_if(t_i)$$
(926)

$$\leq \left(\frac{1}{1-2^{-f(n)}}\right) \frac{1}{\sqrt{n}} \sum_{x_1, x_2, \cdots, x_n \in \mathcal{V}} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \sum_{i=1}^n x_i f(t_i) \quad (927)$$

$$+\frac{1}{\sqrt{n}}\sum_{x_1,x_2,\cdots,x_n\in\mathcal{V}^c}\mathbb{P}[X_1=x_1,X_2=x_2,\cdots,X_n=x_n]\sum_{i=1}^n x_if(t_i)$$
(928)

$$= \frac{1}{\sqrt{n}} \sum_{x_1, x_2, \cdots, x_n \in \mathcal{V}^c} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \sum_{i=1}^n x_i f(t_i).$$
(929)

From (929), we obtain

$$\left|\frac{1}{\sqrt{n}}\sum_{i=1}^{n}\mathbb{E}[X_{i}f(T_{i})]\right| \leq \frac{1}{\sqrt{n}}\sum_{x_{1},x_{2},\cdots,x_{n}\in\mathcal{V}^{c}}\mathbb{P}[X_{1}=x_{1},X_{2}=x_{2},\cdots,X_{n}=x_{n}]\left|\sum_{i=1}^{n}x_{i}f(t_{i})\right|$$
(930)

$$\leq \frac{\|f\|_{\infty}}{\sqrt{n}} \sum_{x_1, x_2, \cdots, x_N \in \mathcal{V}^c} \mathbb{P}[X_1 = x_1, X_2 = x_2, \cdots, X_n = x_n] \left| \sum_{k=1}^n x_k \right|$$
(931)

$$=\frac{\|f\|_{\infty}}{\sqrt{n}}|\mathcal{V}^{c}|\sup_{(x_{1},x_{2},\cdots,x_{n})\in\mathcal{V}^{c}}\left|\sum_{i=1}^{n}x_{i}\right|$$
(932)

$$\leq \frac{\|f\|_{\infty}}{\sqrt{n}} |\mathcal{V}^{c}| \sup_{(x_{1}, x_{2}, \cdots, x_{n}) \in \mathcal{V}^{c}} \sum_{k=1}^{n} x_{k}^{4}$$
(933)

$$\leq \frac{\|f\|_{\infty}}{\sqrt{n}} |\mathcal{V}^c|g(n) \to 0 \tag{934}$$

as $n \to \infty$.

Similarly, we have

$$\frac{1}{n} \mathbb{E} \left[\left| \sum_{i=1}^{n} (1 - X_{i}^{2}) \right| \right] \leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n} \right) \sum_{x_{1}, x_{2}, \cdots, x_{n}} \prod_{i=1}^{n} \mathbb{P}(X_{i} = x_{i}) \sum_{i=1}^{n} (1 - x_{i}^{2}) \right| \\
+ \frac{1}{n} |\mathcal{V}^{c}| \sup_{(x_{1}, x_{2}, \cdots, x_{n}) \in \mathcal{V}^{c}} \left| \sum_{i=1}^{n} (1 - x_{i}^{2}) \right| \\
= \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n} \right) \sum_{x_{1}, x_{2}, \cdots, x_{n}} \prod_{i=1}^{n} \mathbb{P}(X_{i} = x_{i}) \sum_{i=1}^{n} (1 - x_{i}^{2}) \right| \\
+ \frac{1}{n} |\mathcal{V}^{c}| \sup_{(x_{1}, x_{2}, \cdots, x_{n}) \in \mathcal{V}^{c}} \max \left\{ \sum_{i=1}^{n} x_{i}^{2}, 1 \right\} \tag{936}$$

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}(X_i = x_i) \sum_{i=1}^n (1 - x_i^2) \Big| \\ + \frac{1}{n} |\mathcal{V}^c| \sup_{(x_1, x_2, \cdots, x_n) \in \mathcal{V}^c} \max\left\{ \sqrt{n \sum_{i=1}^n x_i^4}, 1 \right\}$$
(937)

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}(X_i = x_i) \sum_{i=1}^n (1 - x_i^2) \Big| \\ + \frac{1}{n} |\mathcal{V}^c| \max\left\{\sqrt{ng(n)}, 1\right\}$$
(938)

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}(X_i = x_i) \sum_{i=1}^n (1 - x_i^2) \Big| \\ + |\mathcal{V}^c|g(n) \left(\frac{1}{ng(n)} \max\left\{\sqrt{ng(n)}, 1\right\}\right)$$
(939)

$$= \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n}\right) \sum_{\substack{x_1, x_2, \cdots, x_n \ i=1}} \prod_{i=1}^n \mathbb{P}(X_i = x_i) \sum_{i=1}^n (1 - x_i^2) \bigg| + o(1)$$
(940)

$$\leq \left(\frac{1}{1-2^{-f(n)}}\right) \frac{\|f''\|_{\infty}}{2n^{3/2}} \sum_{k=1}^{n} \mathbb{E}[|X_k|^3] + o(1), \tag{941}$$

where (941) follows from [25, Proof of Lemma 3.4].

Furthermore, we also have

$$\frac{1}{n^{3/2}} \sum_{i=1}^{n} \mathbb{E}\left[|X_{i}|^{3}\right] \leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_{1}, x_{2}, \cdots, x_{n}} \prod_{i=1}^{n} \mathbb{P}[X_{i} = x_{i}] \left(\sum_{i=1}^{n} |x_{i}|^{3}\right) \\
+ \frac{1}{n^{3/2}} |\mathcal{V}^{c}| \sup_{(x_{1}, x_{2}, \cdots, x_{n}) \in \mathcal{V}^{c}} \left(\sum_{i=1}^{n} |x_{i}|^{3}\right) \\
= \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_{1}, x_{2}, \cdots, x_{n}} \prod_{i=1}^{n} \mathbb{P}[X_{i} = x_{i}] \left(\sum_{i=1}^{n} |x_{i}|^{3}\right) \tag{942}$$

March 16, 2022

$$+\frac{1}{n^{1/2}}|\mathcal{V}^{c}|\sup_{(x_{1},x_{2},\cdots,x_{n})\in\mathcal{V}^{c}}\left(\frac{1}{n}\sum_{i=1}^{n}|x_{i}|^{3}\right)$$

$$(943)$$

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1} \mathbb{P}[X_i = x_i] \left(\sum_{i=1}^{n} |x_i|^3\right) \\ + \frac{1}{n^{1/2}} |\mathcal{V}^c| \sup_{(x_1, x_2, \cdots, x_n) \in \mathcal{V}^c} \left(\frac{1}{n} \sum_{i=1}^n |x_i|^4\right)^{3/4}$$
(944)

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}[X_i = x_i] \left(\sum_{i=1}^n |x_i|^3\right) \\ + \frac{1}{n^{1/2}} |\mathcal{V}^c| \left(\frac{g(n)}{n}\right)^{3/4}$$
(945)

$$\leq \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}[X_i = x_i] \left(\sum_{i=1}^n |x_i|^3\right) \\ + \frac{1}{n^{1/2}} |\mathcal{V}^c| g(n) \frac{1}{(g(n)n^5)^{1/4}}$$
(946)

$$= \frac{1}{1 - 2^{-f(n)}} \left(\frac{1}{n^{3/2}}\right) \sum_{x_1, x_2, \cdots, x_n} \prod_{i=1}^n \mathbb{P}[X_i = x_i] \left(\sum_{i=1}^n |x_i|^3\right) + o(1)$$
(947)

$$\leq \left(\frac{1}{1-2^{-f(n)}}\right) \frac{\|f''\|_{\infty}}{n} \sqrt{\sum_{k=1}^{n} \mathbb{E}[X_k^4]},\tag{948}$$

where (944) follows from the concavity of the function $x^{3/4}$ on $(0,\infty)$.

From (923), (934), (940), and (948), we obtain

$$\left|\mathbb{E}[f'(T) - Tf(T)]\right| \le \left(\frac{1}{1 - 2^{-f(n)}}\right) \frac{\|f''\|_{\infty}}{2n^{3/2}} \sum_{k=1}^{n} \mathbb{E}[|X_k|^3] + \left(\frac{1}{1 - 2^{-f(n)}}\right) \frac{\|f''\|_{\infty}}{n} \sqrt{\sum_{k=1}^{n} \mathbb{E}[X_k^4] + o(1)} \to 0$$
(949)

as $n \to \infty$ under the conditions (310) and (311).

Then, by [25, Th. 3.1], we conclude that

$$T \xrightarrow{(d)} \mathcal{N}(0,1),$$
 (950)

under the conditions (310) and (311). Now, since \tilde{T} is a scaling of T, hence by Slutsky's theorem, it holds that

$$\tilde{T} \xrightarrow{(\mathrm{d})} \left(\lim_{n \to \infty} \sqrt{\frac{\operatorname{Var}(S_n)}{\sum_{i=1}^{n} \mathbb{E}[X_i^2]}} \right) \mathcal{N}(0, 1).$$
(951)

Now, since $\mathbb{E}[\tilde{T}] = 0$ and $\operatorname{Var}(\tilde{T}) = 1$, by applying Lemma 1 with $\operatorname{Var}(Z) = 1$, we must have

$$\lim_{n \to \infty} \sqrt{\frac{\operatorname{Var}(S_n)}{\sum_{i=1}^n \mathbb{E}[X_i^2]}} = 1,$$
(952)

which leads to

$$\tilde{T} \xrightarrow{(d)} \overset{(d)}{\to} \mathcal{N}(0,1).$$
 (953)

O. Proof of Lemma 17

This proof is based on the proof of [25, Prop. 1.2]. Consider the function $h_x(w) = \mathbf{1}\{w \le x\}$, and the 'smooth' $h_{x,\varepsilon}(w)$ defined to be one for $w \le x$, zero for $w > x + \varepsilon$, and linear between them. Then, it is clear that $h_{x,\varepsilon} \in \mathcal{V}$ with a = x and $c = \varepsilon$.

First, observe that $\varepsilon h_{x,\varepsilon}(w)$ is 1-Lipschitz and

$$\left\|\varepsilon h_{x,\varepsilon}\right\|_{\infty} \le \varepsilon. \tag{954}$$

Hence, it holds that

$$4\sqrt{2\pi}h_{x,4\sqrt{2\pi}} \in \mathcal{H} = \{h \in \mathcal{V} : c \le 4\sqrt{2\pi}\},\tag{955}$$

so \mathcal{H} in the definition of Wasserstein metric (cf. Definition 2) is a non-empty set, and $d_W(T, Z)$ is well-defined.

Furthermore, by definition of $d_{W, \text{mod}}(T, Z)$, it holds that

$$d_{W,\text{mod}}(T,Z) \le \sup_{h \in \mathcal{H}} \mathbb{E}[|h(Z)|] + \mathbb{E}[|h(T)|]$$
(956)

$$\leq 2\|h\|_{\infty} \tag{957}$$

$$= 2c \tag{958}$$

$$\leq 8\sqrt{2\pi}.\tag{959}$$

Now, by setting $\varepsilon = (2\pi)^{1/4} \sqrt{2d_{W, \text{mod}}(T, Z)}$, it holds that

$$\left\|\varepsilon h_{x,\varepsilon}\right\|_{\infty} \le (2\pi)^{1/4} \sqrt{2d_{W,\mathrm{mod}}(T,Z)}$$
(960)

$$\leq 4\sqrt{2\pi},\tag{961}$$

where (960) follows from (954), and (961) follows from (959). This means that $\varepsilon h_{x,\varepsilon} \in \mathcal{H}$ since $\varepsilon h_{x,\varepsilon} \in \mathcal{V}$ as mentioned above.

Then, we have

$$\mathbb{E}[h_x(T)] - \mathbb{E}[h_x(Z)] = \mathbb{E}[h_x(T)] - \mathbb{E}[h_{x,\varepsilon}(Z)] + \mathbb{E}[h_{x,\varepsilon}(Z)] - \mathbb{E}[h_x(Z)]$$
(962)

$$\leq \mathbb{E}[h_{x,\varepsilon}(T)] - \mathbb{E}[h_{x,\varepsilon}(Z)] + \mathbb{E}[h_{x,\varepsilon}(Z)] - \mathbb{E}[h_x(Z)]$$
(963)

$$= \frac{1}{\varepsilon} \left(\mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)] \right) + \left| \mathbb{E}[h_{x,\varepsilon}(Z)] - \mathbb{E}[h_x(Z)] \right|$$
(964)

$$\leq \frac{1}{\varepsilon} \left| \mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)] \right| + \left| \mathbb{E}[h_{x,\varepsilon}(Z)] - \mathbb{E}[h_x(Z)] \right|.$$
(965)

Similarly, by choosing $h_{x,\varepsilon}(\omega)$ to be 1 when $\omega \le x - \varepsilon$, 0 when $\omega \ge x$, and linear between them, which is also a function in \mathcal{V} , we can show that

$$\mathbb{E}[h_x(Z)] - \mathbb{E}[h_x(T)] \le \frac{1}{\varepsilon} \left| \mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)] \right| + \left| \mathbb{E}[h_{x,\varepsilon}(Z)] - \mathbb{E}[h_x(Z)] \right|$$
(966)

$$\leq \frac{1}{\varepsilon} \left| \mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)] \right| \\ + \int^{x+\varepsilon} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^2}{2}\right) \left(h_{x,\varepsilon}(z) - h_x(z)\right) dz$$
(967)

$$\int_{x} \sqrt{2\pi} \left(2 \right)^{\varepsilon} \left(\frac{\varepsilon}{2} \right)^{\varepsilon} = \frac{1}{\varepsilon} \left| \mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)] \right| + \frac{\varepsilon}{2\sqrt{2\pi}}.$$
(968)

From (965) and (966), we obtain

$$\left|\mathbb{P}(T \le x) - \mathbb{P}(Z \le x)\right| \le \frac{1}{\varepsilon} \left|\mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)]\right| + \frac{\varepsilon}{2\sqrt{2\pi}}.$$
(969)

Similarly, we also have

$$\left|\mathbb{P}(-T \le x) - \mathbb{P}(Z \le x)\right| \le \frac{1}{\varepsilon} \left|\mathbb{E}[\varepsilon h_{x,\varepsilon}(-T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)]\right| + \frac{\varepsilon}{2\sqrt{2\pi}}.$$
(970)

It follows from (969) and (970) that

$$\sup_{x \in \mathbb{R}} \min\{\left|\mathbb{P}(T \le x) - \mathbb{P}(Z \le x)\right|, \left|\mathbb{P}(-T \le x) - \mathbb{P}(Z \le x)\right|\}$$

$$\leq \sup_{h \in \mathcal{H}} \min\left\{\frac{1}{\varepsilon}\left|\mathbb{E}[\varepsilon h_{x,\varepsilon}(T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)]\right|, \frac{1}{\varepsilon}\left|\mathbb{E}[\varepsilon h_{x,\varepsilon}(-T)] - \mathbb{E}[\varepsilon h_{x,\varepsilon}(Z)]\right|\right\} + \frac{\varepsilon}{2\sqrt{2\pi}}$$
(971)

$$=\frac{1}{\varepsilon}d_{W,\mathrm{mod}}(T,Z) + \frac{\varepsilon}{2\sqrt{2\pi}}$$
(972)

$$= (8\pi)^{-1/4} \sqrt{d_{W,\text{mod}}(T,Z)},\tag{973}$$

where (972) follows from $\varepsilon h_{x,\varepsilon} \in \mathcal{H}$, and (973) follows from our setting $\varepsilon = (2\pi)^{1/4} \sqrt{2d_{W,\text{mod}}(T,Z)}$ above.

Now, for any $x \in \mathbb{R}$, we have

$$\sup_{x \in \mathbb{R}} \min \left\{ \left| \mathbb{P}(T \le x) - \mathbb{P}(Z \le x) \right|, \left| \mathbb{P}(T \le -x) - \mathbb{P}(Z \le x) \right| \right\}$$
$$\geq \min \left\{ \left| \mathbb{P}(T \le x) - \mathbb{P}(Z \le x) \right|, \left| \mathbb{P}(T \le -x) - \mathbb{P}(Z \le x) \right| \right\}$$
(974)

$$\geq \min\left\{ \left| \mathbb{P}(T \leq x) - \mathbb{P}(Z \leq x) \right|, \left| \mathbb{P}(T \leq x) - \mathbb{P}(Z \leq x) \right| - \left| \mathbb{P}(T \leq x) - \mathbb{P}(T \geq -x) \right| \right\}$$
(975)

$$\geq \min\left\{\left|\mathbb{P}(T \leq x) - \mathbb{P}(Z \leq x)\right|, \left|\mathbb{P}(T \leq x) - \mathbb{P}(Z \leq x)\right|\right\} - \left|\mathbb{P}(T \leq x) - \mathbb{P}(T \geq -x)\right|,\tag{976}$$

where (975) follows from the triangle inequality.

From (973) and (976), we obtain (396).

Now, if the distribution of T is tight, then there exists a distribution \tilde{Y} such that $T \xrightarrow{(d)} \tilde{Y}$ [23]. Then, if x is a continuous point of $\mathbb{P}(\tilde{Y} \leq x)$ such that $x \to 0$ as $n \to \infty$, we have

$$\lim_{n \to \infty} \min \left\{ \left| \mathbb{P}(T \le x) - \mathbb{P}(Z \le x) \right|, \left| \mathbb{P}(T \le -x) - \mathbb{P}(Z \le x) \right| \right\}$$
$$= \lim_{n \to \infty} \min \left\{ \left| \mathbb{P}(\tilde{Y} \le x) - \mathbb{P}(Z \le x) \right|, \left| \mathbb{P}(\tilde{Y} \le -x) - \mathbb{P}(Z \le x) \right| \right\}$$
(977)

$$= \min\left\{ \left| \mathbb{P}(\tilde{Y} \le 0) - \mathbb{P}(Z \le 0) \right|, \left| \mathbb{P}(\tilde{Y} \le 0) - \mathbb{P}(Z \le 0) \right| \right\}$$
(978)

$$= \lim_{n \to \infty} \left| \mathbb{P}(T \le 0) - \mathbb{P}(Z \le 0) \right|, \tag{979}$$

where (977) follows from $\lim_{N\to\infty} \min\{A_n, B_n\} = \min\{\lim_{n\to\infty} A_n, \lim_{n\to\infty} B_n\}$ if both the limits $\lim_{n\to\infty} A_n$ and $\lim_{N\to\infty} B_n$ exist.

Hence, we obtain (397) from (979) and (973).

P. Proof of Lemma 19

By Lemma 18, we have

$$d_{W,\text{mod}}(T,Z) \le \sup_{h \in \mathcal{H}} \left\{ \left| \mathbb{E} \left[f'_h(T) - Tf_h(T) \right] \right|, \left| \mathbb{E} \left[f'_h(-T) + Tf_h(-T) \right] \right| \right\}.$$
(980)

Now, observe that

$$\mathbb{E}[f'_h(T) - Tf_h(T)] = \mathbb{E}[h(T)] - \mathbb{E}[h(Z)]$$
(981)

$$= \mathbb{E}[h(T_1)] - \mathbb{E}[h(Z)] + \mathbb{E}[h(T) - h(T_1)]$$
(982)

$$= \mathbb{E}[f'_h(T_1) - T_1 f_h(T_1)] + \mathbb{E}[h(T) - h(T_1)],$$
(983)

where (981) and (983) follow from (398).

It follows that

$$\left| \mathbb{E} [f'_h(T) - Tf_h(T)] \right| = \left| \mathbb{E} [f'_h(T_1) - T_1 f_h(T_1)] + \mathbb{E} [h(T) - h(T_1)] \right|$$
(984)

$$\leq \left| \mathbb{E}[f_{h}'(T_{1}) - T_{1}f_{h}(T_{1})] \right| + \left| \mathbb{E}[h(T) - h(T_{1})] \right|$$
(985)

$$= \left| \mathbb{E}[f_h'(T_1) - T_1 f_h(T_1)] \right| + \mathbb{E}[h(T) - h(T_1)]$$
(986)

where (986) follows from $T \leq T_1$ and h is non-increasing.

Similarly, we have

$$\left| \mathbb{E} \big[f_h'(-T) + T f_h(-T) \big] \right| = \left| \mathbb{E} [f_h'(-T_1) + T_1 f_h(-T_1)] + \mathbb{E} [h(-T) - h(-T_1)] \right|$$
(987)

$$\leq \left| \mathbb{E}[f_h'(-T_1) + T_1 f_h(-T_1)] \right| + \mathbb{E}[h(-T_1) - h(T)],$$
(988)

where (988) follows from $T \leq T_1$ and h is non-increasing.

From (986) and (988), for all $h \in \mathcal{H}$, we have

$$\min\left\{\left|\mathbb{E}[f_{h}'(T) - Tf_{h}(T)]\right|, \left|\mathbb{E}[f_{h}'(-T) + Tf_{h}(-T)]\right|\right\} \le \max\left\{\left|\mathbb{E}[f_{h}'(T_{1}) - T_{1}f_{h}(T_{1})]\right|, \left|\mathbb{E}[f_{h}'(T_{1}) - T_{1}f_{h}(T_{1})]\right|\right\} + \min\left\{\mathbb{E}[h(T) - h(T_{1})], \mathbb{E}[h(-T_{1}) - h(-T)]\right\}$$
(989)

Finally, we obtain (400) from (989).

Q. Proof of Lemma 23

The proof of this lemma is based on the proof of the [25, Th. 3.1]. Given $h \in \mathcal{H}$, we choose f_h be a solution of the following ODE equation:

$$f'_{h}(w) - wf_{h}(w) = h(w) - \Phi(h)$$
(990)

where $\Phi(h) = \mathbb{E}[h(Z)]$ with $Z \sim \mathcal{N}(0, 1)$, then we have

$$f_h(w) = e^{\frac{w^2}{2}} \int_w^\infty e^{-\frac{t^2}{2}} (\Phi(h) - h(t)) dt$$
(991)

$$= -e^{\frac{w^2}{2}} \int_{-\infty}^{w} e^{-\frac{t^2}{2}} (\Phi(h) - h(t)) dt.$$
(992)

Now, it is easy to prove the following facts from (992) (see [25]):

$$\|f_h\|_{\infty} \le 2\|h'\|_{\infty} = 2,\tag{993}$$

$$\|f_{h}^{'}\|_{\infty} \leq \sqrt{\frac{2}{\pi}} \|h'\|_{\infty} = \sqrt{\frac{2}{\pi}},\tag{994}$$

$$\|f_h^{"}\|_{\infty} \le 2\|h'\|_{\infty} = 2.$$
(995)

Now, assume that $\mathbb{E}[T] = 0$ and $\mathbb{E}[T^2 = 1]$.

Furthermore, for any $h \in \mathcal{H}$, from (990), it holds that

$$|f_{h}'(T) - Tf_{h}(T)| = |h(T) - \Phi(h)|$$
(996)

$$= |h(T) - \mathbb{E}[h(Z)]| \tag{997}$$

$$\leq 2\|h\|_{\infty} \tag{998}$$

$$\leq 8\sqrt{2\pi}.\tag{999}$$

Furthermore, from (990), we also have

$$d_W(T,Z) = \sup_{h \in \mathcal{H}} \left| \mathbb{E}[h(T)] - \mathbb{E}[h(Z)] \right|$$
(1000)

$$\leq \sup_{f_h:h\in\mathcal{H}} \left| \mathbb{E}[Tf_h(T) - f'_h(T)] \right|.$$
(1001)

Now, for all $f_h : h \in \mathcal{H}$, observe that

$$Tf_{h}(T) - f'_{h}(T) = T(f_{h}(T) - f_{h}(0) - Tf'_{h}(0)) + Tf_{h}(0) + (T^{2} - 1)f'_{h}(0) + (f'_{h}(0) - f'_{h}(T)).$$
(1002)

It follows from (1002) that

$$\mathbb{E}\left[Tf_{h}(T) - f_{h}^{'}(T)\right] = \mathbb{E}\left[T\left(f_{h}(T) - f_{h}(0) - Tf_{h}^{'}(0)\right)\right] + f_{h}(0)\mathbb{E}[T] + f_{h}^{'}(0)\mathbb{E}\left[T^{2} - 1\right] + \mathbb{E}[f_{h}^{'}(0) - f_{h}^{'}(T)]$$
(1003)

$$= \mathbb{E}\left[T\left(f_h(T) - f_h(0) - Tf'_h(0)\right)\right] + \mathbb{E}[f'_h(0) - f'_h(T)],$$
(1004)

where (1004) follows from the fact that $\mathbb{E}[T]=0$ and $\mathbb{E}[T^2]=1.$

Hence, from (1001) and (1004), we have

$$\tilde{d}_{W,\text{mod}}(T,Z) \le \sup_{f_h:h\in\mathcal{H}} \left| \mathbb{E}[Tf_h(T) - f'_h(T)] \right|$$
(1005)

$$\leq \sup_{f_h:h\in\mathcal{H}} \mathbb{E}\big[\big|T\big(f_h(T) - f_h(0) - Tf_h'(0)\big)\big|\big] + \mathbb{E}\big[\big|f_h'(0) - f_h'(T)\big|\big].$$
(1006)

Now, observe that

$$\left|T\left(f_{h}(T) - f_{h}(0) - Tf_{h}^{'}(0)\right)\right| = \left|Tf_{h}(T) - f_{h}^{'}(T) + f_{h}^{'}(T) - Tf_{h}(0) - T^{2}f_{h}^{'}(0)\right|$$
(1007)

$$= \left| Tf_h(T) - f'_h(T) + f'_h(T) - Tf_h(0) - f'_h(0) + (1 - T^2)f'_h(0) \right|$$
(1008)

$$\leq \left| Tf_h(T) - f'_h(T) \right| + \left| f'_h(T) \right| + \left| Tf_h(0) \right| + \left| f'_h(0) \right| + \left| f'_h(0) (T^2 - 1) \right| \right)$$
(1009)

$$\leq 8\sqrt{2\pi} + 2\sqrt{\frac{2}{\pi}} + 2|T| + \sqrt{\frac{2}{\pi}}|T^2 - 1|$$
(1010)

$$= \left(8 + \frac{2}{\pi}\right)\sqrt{2\pi} + 2|T| + \sqrt{\frac{2}{\pi}}|T^2 - 1|,$$
(1011)

where (1010) follows from (993), (994) and (999).

Hence, we have

$$T\left(f_{h}(T) - f_{h}(0) - Tf_{h}'(0)\right)|] = \min\left\{\left(8 + \frac{2}{\pi}\right)\sqrt{2\pi} + 2|T|, |T\left(f_{h}(T) - f_{h}(0) - Tf_{h}'(0)\right)|\right\}$$
(1012)
$$\left(\left(-2\right)\sqrt{2\pi} + 2|T|, |T\left(f_{h}(T) - f_{h}(0) - Tf_{h}'(0)\right)|\right\}$$

$$\leq \min\left\{\left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}, \left|T\left(f_h(T) - f_h(0) - Tf'_h(0)\right)\right|\right\} + 2|T| + \sqrt{\frac{2}{\pi}}|T^2 - 1|,$$
(1013)

where (1013) follows from $\min\{A + B, C\} \le \min\{A, C\} + B$ for all $A, B, C \ge 0$. It follows from (1013) that

$$\mathbb{E}\left[\left|T\left(f_{h}(T) - f_{h}(0) - Tf_{h}^{'}(0)\right)\right|\right] \leq \mathbb{E}\left[\min\left\{\left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}, \left|T\left(f_{h}(T) - f_{h}(0) - Tf_{h}^{'}(0)\right)\right|\right\}\right] + 2\mathbb{E}\left[|T|\right].$$
(1014)

Now, by Taylor's expansion, for some $\eta \in (0, -|T|) \cup (0, |T|)$, we have

$$f_h(T) - f_h(0) - Tf'_h(0) = \frac{1}{2}f'_h(\eta)T^2,$$
(1015)

so

$$\left|T\left(f_{h}(T) - f_{h}(0) - Tf_{h}'(0)\right)\right| = \frac{1}{2} \left|T^{3}f_{h}''(\eta)\right|$$
(1016)

$$\leq \frac{1}{2} \|f_h^{"}\|_{\infty} |T^3| \tag{1017}$$

$$\leq |T^3|.\tag{1018}$$

Hence, from (1014) and (1018), we obtain

$$\mathbb{E}\left[\left|T\left(f_{h}(T) - f_{h}(0) - Tf_{h}'(0)\right)\right|\right] \le \mathbb{E}\left[\min\left\{\left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}, |T|^{3}\right\}\right] + 2\mathbb{E}\left[|T|\right] + \sqrt{\frac{2}{\pi}}|T^{2} - 1|.$$
(1019)

Similarly, by Taylor's expansion, for some $\theta \in (0,-|T|) \cup (0,|T|),$ we have

$$f'_{h}(T) - f'_{h}(0) = f^{"}_{h}(\theta)T,$$
(1020)

so

$$\mathbb{E}\left[\left|f_{h}^{'}(T) - f_{h}^{'}(0)\right|\right] = \mathbb{E}\left[\left|f_{h}^{"}(\theta)T\right|\right]$$
(1021)

$$\leq \mathbb{E}\big[|f_h^{"}(\theta)||T|\big] \tag{1022}$$

$$\leq \|f_h^{"}\|_{\infty} \mathbb{E}\big[|T|\big] \tag{1023}$$

$$\leq 2\mathbb{E}[|T|]. \tag{1024}$$

Finally, from (1013), (1019), and (1024), we have

$$\tilde{d}_{W,\text{mod}}(T,Z) \le \sup_{f_h:h\in\mathcal{H}} \mathbb{E}\left[\left| T \left(f_h(T) - f_h(0) - T f'_h(0) \right) \right| \right] + \mathbb{E}\left[\left| f'_h(0) - f'_h(T) \right| \right]$$
(1025)

$$\leq \mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, \left|T\right|^{3}\right\}\right] + 2\mathbb{E}\left[\left|T\right|\right] + \sqrt{\frac{2}{\pi}}|T^{2}-1|.$$
(1026)

Now, observe that

$$\mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|^3\right\}\right]$$

$$=\mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|^3\right\} \left||T| \le 1\right]\mathbb{P}[|T| \le 1]$$

$$+\mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|^3\right\} \left||T| > 1\right]\mathbb{P}[|T| > 1]$$

$$\leq \mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|^3\right\} \left||T| \le 1\right]\mathbb{P}[|T| \le 1]$$

$$(1027)$$

$$= \mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|\right\}||T| \le 1\right]\mathbb{E}\left[|T| \ge 1\right]$$

$$+ \mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi}, |T|\right\}||T| > 1\right]\mathbb{E}\left[|T| > 1\right]$$

$$(1028)$$

$$\leq \mathbb{E}\left[\left|T\right| \left| \left|T\right| \leq 1\right] \mathbb{P}\left[\left|T\right| \leq 1\right] + \mathbb{E}\left[\left|T\right| \leq 1\right] + \mathbb{E}\left[\left|T\right| \leq 1\right] + \mathbb{E}\left[\left|T\right| \geq 1\right] + \mathbb{E}\left[\left|T\right| = 1$$

$$+\mathbb{E}\left[\min\left\{\left(8+\frac{2}{\pi}\right)\sqrt{2\pi},\left|T\right|^{3}\right\}\left||T|>1\right]\mathbb{P}[|T|>1]\right]$$
(1029)

$$\leq \mathbb{E}\left[|T|\right] + \mathbb{E}\left[\min\left\{\left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}, |T|^3\right\} ||T| > 1\right] \mathbb{P}[|T| > 1]$$

$$(1030)$$

$$\leq \mathbb{E}\left[|T|\right] + \left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}\mathbb{P}[|T| > 1]$$
(1031)

$$\leq \mathbb{E}\left[|T|\right] + \left(8 + \frac{2}{\pi}\right)\sqrt{2\pi}\mathbb{E}[|T|] \tag{1032}$$

$$\leq \left(10 + \frac{1}{\pi}\right)\sqrt{2\pi}\mathbb{E}[|T|],\tag{1033}$$

where (1028) follows from $|T|^3 \leq |T|$ for all $|T| \leq 1$, (1030) follows from $\mathbb{E}[X] = \mathbb{E}[X|A]\mathbb{P}(A) + \mathbb{E}[X|A^c]\mathbb{P}(A^c) \geq \mathbb{E}[X|A]\mathbb{P}(A)$ for all non-negative random variable X, and (1032) follows from Markov's inequality.

From (1026) it follows that

$$\tilde{d}_{W,\text{mod}}(T,Z) \le \left(\left(10 + \frac{1}{\pi} \right) \sqrt{2\pi} + \left(4 + \sqrt{\frac{2}{\pi}} \right) \right) \mathbb{E}[|T|] + \sqrt{\frac{2}{\pi}} \mathbb{E}[|T^2 - 1|]$$
(1034)

$$<40\mathbb{E}[|T|] + \sqrt{\frac{2}{\pi}\mathbb{E}[|T^2 - 1|]}.$$
 (1035)

By combining Lemma 17 and (1035), we have

$$d_K(T,Z) < 2(8\pi)^{-1/4} \sqrt{40\mathbb{E}[|T|]} + \sqrt{\frac{2}{\pi}} \mathbb{E}[|T^2 - 1|]$$
(1036)

$$\leq 14(8\pi)^{-1/4}\sqrt{\mathbb{E}[|T|] + \mathbb{E}[|T^2 - 1|]}.$$
(1037)

This concludes our last proof.

REFERENCES

- [1] C. E. Shannon, "A mathematical theory of communication," Bell System Technical Journal, vol. 27, pp. 379-423, 1948.
- [2] R. M. Fano, Transmission of Information. New York: Wiley, 1961.
- [3] R. G. Gallager, "Simple derivation of the coding theorem and some applications," IEEE Trans. Inf. Theory, vol. 11, pp. 3–18, Jan 2008.
- [4] C. E. Shannon, R. G. Gallager, and E. R. Berlekamp, "Lower bounds to error probability for coding in discrete memoryless channels I-II," *Information and Control*, vol. 10, pp. 65–103, 522–552, 1967.
- [5] B. Nakiboğlu, "The sphere packing bound for memoryless channels," Problems of Information Transmission, vol. 56, pp. 201–244, 2020.
- [6] B. Nakiboğlu, "The Augustin capacity and center," Problems of Information Transmission, vol. 55, no. 4, pp. 299-342, 2019.
- [7] A. Barg and G. D. Forney, "Random codes: minimum distances and error exponents," *IEEE Trans. Inf. Theory*, vol. 48, no. 9, pp. 2568–2573, 2002.
- [8] A. Nazari, A. Anastasopoulos, and S. S. Pradhan, "Error exponent for multiple-access channels: Lower bounds," *IEEE Trans. Inf. Theory*, vol. 60, no. 9, pp. 5095–5115, 2014.
- [9] N. Merhav, "Error exponents of typical random codes," IEEE Trans. Inf. Theory, vol. 64, no. 9, pp. 6223-6235, 2018.
- [10] N. Merhav, "Error exponents of typical random codes for the colored Gaussian channel," *IEEE Trans. Inf. Theory*, vol. 65, no. 12, pp. 8164–8179, 2019.
- [11] N. Merhav, "Error exponents of typical random trellis codes," IEEE Trans. Inf. Theory, vol. 66, no. 4, pp. 2067–2077, 2019.
- [12] R. Tamir, N. Merhav, N. Weinberger, and A. Guillén i Fàbregas, "Large deviations behavior of the logarithmic error probability of random codes," *IEEE Trans. Inf. Theory*, vol. 66, no. 11, pp. 6635–6659, 2020.
- [13] R. Ahlswede and G. Dueck, "Good codes can be produced by a few permutations," *IEEE Trans. Inf. Theory*, vol. 28, no. 3, pp. 430–443, 1982.
- [14] R. T. (Averbuch) and N. Merhav, "Universal decoding for the typical random code and for the expurgated code," *IEEE Trans. Inf. Theory*, 2022.
- [15] G. Cocco, A. Guillén i Fàbregas, and J. Font-Segura, "A dual-domain achievability of the typical error exponent," in *IEEE Int. Symp. Inf. Theory*, (Melbourne, Australia), 2021.
- [16] J. Scarlett, A. Martinez, and A. Guillén i Fàbregas, "Mismatched decoding: Finite-Length bounds, error exponents and approximations," *IEEE Trans. Inf. Th.*, 2014.
- [17] R. G. Gallager, "Fixed composition arguments and lower bounds to the error probability, lecture notes MIT 6.441," 1994.
- [18] R. Durrett, Probability: Theory and Examples. Cambridge Univ. Press, 4th ed., 2010.
- [19] T. Cover and J. Thomas, Elements of Information Theory. John Wiley and Sons, 2006.
- [20] W. Feller, An Introduction to Probability Theory and Its Applications. John Wiley and Sons, 2nd ed., 1971.
- [21] A. Cohen and N. Merhav, "Lower bounds on the error probability of block codes based on improvements on de caen's inequality," *IEEE Trans. Inf. Theory*, vol. 50, no. 2, pp. 290–310, 2004.
- [22] J. Scarlett, A. Martinez, and A. Guillén i Fàbregas, "Ensemble-tight error exponents for mismatched decoders," in *Proc. of Allerton Conference*, 2012.

- [23] P. Billingsley, Probability and Measure. Wiley-Interscience, 3rd ed., 1995.
- [24] R. G. Gallager, Information Theory and Reliable Communication. USA: John Wiley & Sons, Inc., 1968.
- [25] N. Ross, "Fundamentals of Stein's method," Probability Surveys, vol. 8, no. none, pp. 210 293, 2011.
- [26] R. Paley and A. Zygmund, "On some series of functions, (3)," *Mathematical Proceedings of the Cambridge Philosophical Society*, vol. 28, no. 2, pp. 190–205, 1932.
- [27] D. de Caen, "A lower bound on the probability of a union," Discrete Mathematics, vol. 169, no. 1, pp. 217-220, 1997.
- [28] S. M. Moser, Advanced Topics in Information Theory. Lecture Notes. 2019.
- [29] I. Csiszár, "The method of types," IEEE Trans. Inf. Th., vol. 44, no. 6, pp. 2505-23, 1998.
- [30] H. Royden and P. Fitzpatrick, Real Analysis. Pearson, 4th ed., 2010.
- [31] J. Stewart, Multivariate Calculus. Cengate Learning, 6th ed., 2008.
- [32] R. Gallager, "The random coding bound is tight for the average code (corresp.)," *IEEE Trans. Inf. Theory*, vol. 19, no. 2, pp. 244–246, 1973.
- [33] S. Boyd and L. Vandenberghe, Convex Optimization. Cambridge University Press, 2004.
- [34] T. Cover, "Comments on broadcast channels," IEEE Trans. Inf. Th., vol. 44, no. 6, pp. 2524-30, 1998.
- [35] S. Janson, "New versions of Suen's correlation inequality," Random Struct. Algorithms, vol. 13, pp. 467–483, Oct. 1998.