

# ALGEBRAIC $K$ -THEORY AND ALGEBRAIC COBORDISM OF ALMOST MATHEMATICS

YUKI KATO

ABSTRACT. Faltings [12]; Gabber and Ramero [13] introduced almost mathematics. In another way, almost mathematics can be characterized bilocalization abelian category of modules mentioned in Quillen’s unpublished note [31]. Applying the concept of Quillen’s bilocalization to Gabber and Ramero’s work, this paper establishes the almost version of algebraic  $K$ -theory and cobordism and proves that the  $K$ -theory of almost perfect complexes is equivalent to the Thomason–Trobath  $K$ -theory whose weak equivalences are almost quasi-isomorphisms. Furthermore, on the category of zero-section stable integral perfectoid algebras with finite syntomic topology, we obtain that almost algebraic cobordism holds tilting equivalence.

## 1. INTRODUCTION

Faltings [12] first introduced almost mathematics, proving almost purity. More concisely, Gabber and Ramero established [13] almost ring theory in their textbook: theories of almost modules, almost algebras, and almost homotopical algebra. While almost mathematics has various applications to arithmetic geometry, for example, Scholze’s work perfectoid geometry [32], Quillen [31] mentioned linear algebra over non-unital rings which is the same as almost mathematics. Quillen’s work is more conceptual in the sense of using categorical language: in his work, almost mathematics is characterized as bilocalization of an abelian category of modules.

In almost mathematics, the theory of derived categories is a bit complicated. For instance, exact sequences in the almost-world are not exact in the corresponding module category in general. By considering the derived category of a bilocalization of the category of modules, this paper provides algebraic  $K$ -theory and cobordism of almost mathematics. More precisely, we define the derived category (or stable  $\infty$ -category) of almost perfect complexes, enabling us to obtain the  $K$ -theory of almost modules, and compare the Thomason–Trobath  $K$ -theory of almost perfect complexes and perfect complex whose weak equivalences are almost quasi-isomorphisms.

Furthermore, defining almost finite syntomic algebras, we can get the defining algebraic cobordism in almost mathematics similar to Elmanto, Hoyois, Khan, Sosnilo, and Yakerson’s work [10] via framed correspondences.

---

*Date:* April 26, 2022.

*Key words and phrases.*  $\mathbb{A}^1$ -homotopy theory, almost mathematics, perfectoid algebra.

This paper's another result is the tilting equivalence between the almost algebraic cobordism of zero-section stable integral perfectoid algebras by a similar argument to the author's previous work [19] about the algebraic cobordism of non-unital integral perfectoid algebras (Theorem 4.20). This result lets us expect some analogy between non-unital and almost algebra. Here the condition zero-section stability is defined as follows: Let  $i : \{0\} \rightarrow \Delta^\infty = \operatorname{colim} \Delta^n$  denote the canonical injection and  $\pi : \Delta^\infty \rightarrow \Delta^0$  the projection. One has a Quillen adjunction

$$\pi_* \circ i_* : (\operatorname{Set}_\Delta)_{/\Delta^0} \rightleftarrows (\operatorname{Set}_\Delta)_{/\Delta^0} : \pi^* \circ i^*$$

of simplicial model categories. A simplicial set  $X$  is *zero-section stable* if the morphism  $\pi \circ i \circ \eta : X \rightarrow \pi_*(i_*(X))$  induced by the unit map  $\eta : X \rightarrow (i^* \circ \pi^*)((\pi_* \circ i_*)(X))$  is a weak equivalence. Let  $\mathbb{A}_S^\bullet$  denote the cosimplicial motivic space defined by  $\mathbb{A}_S^\bullet([n]) = \mathbb{A}_S^n$  for each  $n \geq 0$ . Then the singular functor  $\operatorname{Hom}_S(\mathbb{A}^\bullet, -) : \mathbf{MS} \rightarrow \mathbf{MS}$  admits a left adjoint. We say that  $X$  is locally zero-section stable if the simplicial sheaf  $\operatorname{Hom}_S(\mathbb{A}_S^\bullet, X)$  is stalk-wise zero-section stable. Let  $\operatorname{MSp}^0$  denote the full subcategory of zero-section motivic spectra. Then the inclusion functor:  $\operatorname{MSp}^0 \rightarrow \operatorname{MSp}$  admits a left adjoint  $Z_0 : \operatorname{MSp} \rightarrow \operatorname{MSp}^0$  which is defined to be the homotopy colimit

$$Z_0(X) = \varinjlim_n \overbrace{(\pi_* \circ i_*) \circ (\pi_* \circ i_*) \circ \cdots \circ (\pi_* \circ i_*)}^n(X).$$

We call  $Z_0(X)$  the *zero-section stabilization* of  $X$ .

This paper is organized as follows: In Section 2, we provide a brief review of Quillen's work [31] bilocalization by the Serre subcategory of almost zero modules. Section 3 defines almost perfect complexes by using derived functors of categorical equivalences between full subcategories spanned by local objects and spanned by colocal objects. Via the equivalence, we prove the comparison theorem of almost  $K$ -theories (Theorem 3.15). In Section 4, on the viewpoint of non-unital algebra, we study the left adjoint of the localization functor from the category of algebras to one of almost algebras. We define almost finite syntomic algebras, enabling us to establish the almost version of algebraic cobordism by following [10]. In the final part of this paper, localizing the stable  $\infty$ -category of motivic spectra by zero-section stable finite syntomic morphisms, we prove the tilting equivalence of the almost algebraic cobordism for zero-section stable integral perfectoid algebras.

## 2. BILOCALIZATION OF ABELIAN CATEGORY OF MODULES

In this paper, we fix an unital commutative ring  $V$ . In this section; we review Quillen's unpublished note [31] in the almost mathematics language, giving the proof of the propositions.

Let  $\mathfrak{m}$  be an idempotent ideal of  $V$ . A  $V$ -module  $M$  is said to be *almost zero* if  $\mathfrak{m}M = 0$ . A  $V$ -homomorphism  $f : M \rightarrow N$  of  $V$ -modules is called *an almost isomorphism* if both the kernel and the cokernel of  $f$  are almost zero.

**Lemma 2.1** ([31]). *Let  $V$  be an unital commutative ring and  $\mathfrak{m}$  an idempotent ideal. Then  $\mathfrak{m} \otimes_V M = 0$  if and only if  $M$  is almost zero.*

**proof.** The only if direction is clear. Assume  $\mathfrak{m}M = 0$ . For  $a \otimes x \in \mathfrak{m} \otimes_V M$  ( $a \in \mathfrak{m}$ ,  $x \in M$ ), there exists  $(a_i b_i) \subset \mathfrak{m}^2$  such that  $a = \sum_i a_i b_i$ . Hence one has  $a \otimes x = (\sum_i a_i b_i) \otimes x = \sum_i a_i b_i \otimes x = \sum_i a_i \otimes b_i x = 0$ .  $\square$

**Proposition 2.2** ([31]). *Let  $V$  be an unital ring,  $\mathfrak{m}$  an idempotent ideal of  $V$ , and  $M$  a  $V$ -module. Then the canonical morphisms  $\mu : \mathfrak{m} \otimes_V M \rightarrow M$  and  $\mu' : M \rightarrow \text{Hom}_V(\mathfrak{m}, M)$  are both almost isomorphisms.*

**proof.** Since the abelian category  $\text{Mod}_V$  is both enough projective and injective, the kernels and the cokernels;  $\text{Tor}_i^V(V/\mathfrak{m}, M)$  and  $\text{Ext}_V^i(V/\mathfrak{m}, M)$  for  $i = 0, 1$ , are killed by  $\mathfrak{m}$ .  $\square$

**Definition 2.3.** In the situation of Proposition 2.2, we say that  $M$  is *firm* if  $\mu$  is an isomorphism and  $M$  is *closed* if  $\mu'$  is an isomorphism.

In particular, an injective  $V$ -module  $Q$  is closed if and only if it is  $\mathfrak{m}$ -torsion free (i.e.  $\text{Hom}_V(V/\mathfrak{m}, Q) = 0$ ).

**Corollary 2.4** ([31]). *Let  $\mathfrak{m}$  be an idempotent ideal of an unital ring  $V$ . Then the map  $\mu_{\mathfrak{m}} : \mathfrak{m} \otimes_V \mathfrak{m} \rightarrow \mathfrak{m}$  is an almost isomorphism and  $\mu_{\mathfrak{m}} \otimes \mathfrak{m} : \mathfrak{m} \otimes_V \mathfrak{m} \otimes_V \mathfrak{m} \rightarrow \mathfrak{m} \otimes_V \mathfrak{m}$  already an isomorphism.*

**proof.** This follows from Lemma 2.1 and Proposition 2.2.  $\square$

**Corollary 2.5** ([31] Proposition 4.1 and Proposition 5.3). *Let  $V$  be an unital ring and  $\mathfrak{m}$  an idempotent ideal of  $V$ . Write  $\tilde{\mathfrak{m}} = \mathfrak{m} \otimes_V \mathfrak{m}$ . For any  $V$ -module  $M$ ,  $\tilde{\mathfrak{m}} \otimes_V M$  is firm and  $\text{Hom}(\tilde{\mathfrak{m}}, M)$  closed.*

**proof.** By Corollary 2.4, the counit  $\mu_{\tilde{\mathfrak{m}} \otimes_V M} : \mathfrak{m} \otimes_V \tilde{\mathfrak{m}} \otimes_V M \rightarrow \tilde{\mathfrak{m}} \otimes_V M$  is an isomorphism, inducing canonical isomorphisms  $\text{Hom}(\tilde{\mathfrak{m}}, M) \simeq \text{Hom}(\mathfrak{m} \otimes_V \tilde{\mathfrak{m}}, M) \simeq \text{Hom}(\mathfrak{m}, \text{Hom}(\tilde{\mathfrak{m}}, M))$ .  $\square$

Let  $M$  be a closed module and an injective hull  $M \rightarrow Q$ . Then, by the right exactness of  $\text{Hom}_V(\mathfrak{m}, -)$ , one has an injection  $M \rightarrow \text{Hom}_V(\tilde{\mathfrak{m}}, Q)$ . Then  $\text{Ext}_V^1(N, \text{Hom}_V(\tilde{\mathfrak{m}}, Q)) = \text{Ext}_V^1(N, \mathbb{R}\text{Hom}_V(\tilde{\mathfrak{m}}, Q)) \simeq \text{Ext}_V^1(N \otimes_V^{\mathbb{L}} \tilde{\mathfrak{m}}, Q) = 0$  implies that  $\text{Hom}_V(\tilde{\mathfrak{m}}, Q)$  is injective. Hence, a  $V$ -module is closed if and only if it is a kernel of a homomorphism of closed injective modules.

Let  $\mathcal{S}$  denote the Serre subcategory of  $\text{Mod}_V$  spanned by almost zero modules. We say that a  $V$ -module  $M$  is  $\mathcal{S}$ -local (resp.  $\mathcal{S}$ -colocal) if for any almost isomorphism  $f : N_1 \rightarrow N_2$ , the induced map

$$f_* : \text{Hom}_V(M, N_1) \rightarrow \text{Hom}_V(M, N_2) \quad (\text{resp. } f^* : \text{Hom}_V(N_2, M) \rightarrow \text{Hom}_V(N_1, M))$$

is an isomorphism. The following well-known proposition is crucial for almost mathematics. We explain the proof:

**Proposition 2.6** ([31] Proposition 4.1 and Proposition 5.3). *Let  $V$  be an unital ring,  $\mathfrak{m}$  an idempotent ideal of  $V$  and  $M$  a  $V$ -module. Let  $\mathcal{S}$  denote the Serre subcategory of almost zero modules of  $\text{Mod}_V$ . Then the following conditions are equivalent:*

- (1) *The  $V$ -module  $M$  is  $\mathcal{S}$ -local.*
- (2) *The  $V$ -module  $M$  is firm.*
- (3) *The  $V$ -module  $M$  satisfies that  $\text{Ext}_V^0(M, N) = \text{Ext}_V^1(M, N) = 0$  for any almost zero module  $N$ .*

*Similarly, the following conditions are equivalent:*

- (1)' *The  $V$ -module  $M$  is  $\mathcal{S}$ -colocal.*
- (2)' *The  $V$ -module  $M$  is closed.*
- (3)' *The  $V$ -module  $M$  satisfies that  $\text{Ext}_V^0(N, M) = \text{Ext}_V^1(N, M) = 0$  for any almost zero module  $N$ .*

**proof.** Assume that  $M$  is  $\mathcal{S}$ -local. By Proposition 2.2, the induced map

$$\mu_* : \text{Hom}(M, \mathfrak{m} \otimes_V M) \rightarrow \text{Hom}(M, M)$$

is bijective. Hence  $\mu : \mathfrak{m} \otimes_V M \rightarrow M$  is surjective, in particular  $M = \mathfrak{m}M$ . Since  $M$  is a direct factor of  $\mathfrak{m} \otimes_V M$ , the composition  $M \rightarrow \mathfrak{m} \otimes_V M \rightarrow \tilde{\mathfrak{m}} \otimes_V M$  is split. A direct factor  $M$  of the firm module  $\tilde{\mathfrak{m}} \otimes_V M$  is also firm.

Assume that  $M$  is firm. Let  $N$  be an almost zero module. We prove that  $\text{Ext}_V^i(M, N) = 0$  for  $i = 0, 1$ . The condition  $\mathfrak{m}N = 0$  implies that  $N$  is also  $V/\mathfrak{m}$ -module. The adjunction between the derived categories:

$$(V/\mathfrak{m}) \otimes_V^{\mathbb{L}} - : D(V) \rightleftarrows D(V/\mathfrak{m}) : \mathbb{R}\text{Hom}_V(V/\mathfrak{m}, -)$$

induces a canonical equivalence  $\mathbb{R}\text{Hom}_V((V/\mathfrak{m}) \otimes_V^{\mathbb{L}} M, N) \simeq \mathbb{R}\text{Hom}_V(M, N)$  and the spectral sequence

$$E_2^{pq} = \text{Ext}_{V/\mathfrak{m}}^p(\text{Tor}_q^V(V/\mathfrak{m}, M), N) \implies \text{Ext}_V^{p+q}(M, N)$$

where  $E_2^{pq} = 0$  for  $q = 0, 1$ . Hence one has  $\text{Ext}_V^0(M, N) = \text{Ext}_V^1(M, N) = 0$ .

Assume that condition (3). Let  $f : N_1 \rightarrow N_2$  be a morphism of  $V$ -modules with almost zero modules  $N' = \text{Ker}f$  and  $N'' = \text{Coker}f$ . Write  $N = \text{Im}f$  and consider an exact sequences:

$$0 \rightarrow N' \rightarrow N_1 \rightarrow N \rightarrow 0 \quad \text{and} \quad 0 \rightarrow N \rightarrow N_2 \rightarrow N'' \rightarrow 0.$$

We obtain that the induced map  $f_* : \text{Hom}_V(M, N_1) \rightarrow \text{Hom}_V(M, N) \rightarrow \text{Hom}_V(M, N_2)$  is composition of isomorphisms by the assumption (3).

Next, assume that  $M$  is  $\mathcal{S}$ -colocal. Then the almost isomorphism  $\mathfrak{m} \rightarrow V$  induces an isomorphism  $M \rightarrow \text{Hom}_V(\mathfrak{m}, M)$ . Therefore  $M$  is closed.

If  $M$  is closed, there exist an injective resolution  $0 \rightarrow M \rightarrow Q^0 \rightarrow Q^1 \rightarrow Q^2 \rightarrow \dots$ , where  $Q^i$  is closed for  $i = 0, 1$ . By the canonical isomorphism  $\text{Hom}_V(N, \text{Hom}_V(\mathfrak{m}, M)) \simeq \text{Hom}_V(N \otimes_V$

$\mathfrak{m}, M$ ), for any almost zero module  $N$ ,  $\mathrm{Hom}_V(N, Q^i) \simeq \mathrm{Hom}_V(N \otimes_V \mathfrak{m}, Q^i) = 0$  ( $i = 0, 1$ ). Therefore we obtain  $\mathrm{Ext}_V^i(N, M) = 0$  for  $i = 0, 1$ .

By the similar argument of the above (3) to (1), the implication (3)' to (1)' holds.

□

**Theorem 2.7** ([31] Theorem 4.5 and Theorem 5.6). *Let  $V$  be an unital ring and  $\mathfrak{m}$  an idempotent ideal of  $V$ . Let  $\mathcal{S}$  denote the Serre subcategory of almost zero modules of  $\mathrm{Mod}_V$ . Then the localization functor  $(-)^a : \mathrm{Mod}_V \rightarrow \mathrm{alMod}_V$  is also a colocalization. Furthermore, the left adjoint is isomorphic to the functor  $\tilde{\mathfrak{m}} \otimes_V - : \mathrm{alMod}_V \rightarrow \mathrm{Mod}_V$  whose essential image is the full subcategory of film modules, and the right adjoint is isomorphic to  $\mathrm{Hom}_V(\tilde{\mathfrak{m}}, -) : \mathrm{alMod}_V \rightarrow \mathrm{Mod}_V$  whose essential image is the full subcategory of closed modules.*

□

Let  $V$  be an unital ring and  $\mathfrak{m}$  an idempotent ideal of  $V$ . Let  $A$  be a  $V$ -algebra and  $(-)^a : \mathrm{Mod}_A \rightarrow \mathrm{alMod}_A$  denote the bilocalization of  $\mathrm{Mod}_A$  by the Serre subcategory of almost zero  $A$ -modules.

**Corollary 2.8.** *Let  $V$  be an unital ring and  $\mathfrak{m}$  an idempotent ideal of  $V$ . Let  $\mathrm{Mod}_A^{\mathrm{fi}}$  denote the full subcategory of  $\mathrm{Mod}_A$  spanned by firm  $A$ -modules and  $\mathrm{Mod}_A^{\mathrm{cl}}$  by closed  $A$ -modules. Then those adjunctions*

$$\begin{aligned} \tilde{\mathfrak{m}} \otimes_V (-) : \mathrm{Mod}_A^{\mathrm{fi}} &\rightleftarrows \mathrm{alMod}_A : (-)^a \\ (-)^a : \mathrm{Mod}_A^{\mathrm{cl}} &\rightleftarrows \mathrm{alMod}_A : \mathrm{Hom}_V(\tilde{\mathfrak{m}}, -) \end{aligned}$$

are categorical equivalences. Assume that  $\tilde{\mathfrak{m}}$  is a flat  $V$ -module. The functor  $\tilde{\mathfrak{m}} \otimes_V (-)$  induces a categorical equivalence of the derived categories:

$$\begin{aligned} \tilde{\mathfrak{m}} \otimes_V (-) : D(\mathrm{alMod}_A) &\rightleftarrows D(\mathrm{Mod}_A^{\mathrm{fi}}) : (-)^a \\ \tilde{\mathfrak{m}} \otimes_V (-) : D(\mathrm{Mod}_A^{\mathrm{cl}}) &\rightleftarrows D(\mathrm{Mod}_A^{\mathrm{fi}}) : \mathbb{R}\mathrm{Hom}_V(\tilde{\mathfrak{m}}, -). \end{aligned}$$

□

Write  $(-)_! = \tilde{\mathfrak{m}} \otimes_V \mathrm{Hom}_V(\tilde{\mathfrak{m}}, (-)) : \mathrm{alMod}_A \rightarrow \mathrm{Mod}_A^{\mathrm{fi}}$ . Since  $\tilde{\mathfrak{m}}$  is flat, the functor  $(-)_!$  is left exact. Furthermore, if  $f : M \rightarrow N$  is almost surjective, one has  $(\mathrm{Coker} f)_! = 0$ , being  $(-)_!$  is right exact. Hence  $(-)_!$  is an exact functor.

### 3. ALMOST ALGEBRAIC $K$ -THEORY

**3.1. Almost perfect complexes.** In this paper, we always assume that  $\tilde{\mathfrak{m}}$  is a flat  $V$ -module. The full subcategory  $\mathrm{Mod}_A^{\mathrm{cl}}$  is enough injective. Then the derived category of  $\mathrm{alMod}_A$  is defined to be the localization of the category of complexes of closed  $A$ -modules by homotopy equivalences. That is  $D(\mathrm{alMod}_A) \simeq \mathrm{Ho}(\mathrm{Mod}_A^{\mathrm{cl-inj}})$ .

**Definition 3.1.** An  $A$ -module  $M$  is *almost finitely generated* (resp. *almost finitely presented*) if for any filtered inductive system  $(N_\alpha)$  of firm  $A$ -module, the canonical map

$$\varinjlim \mathrm{Hom}_{\mathrm{Mod}_A^{\mathrm{fi}}}(\tilde{\mathfrak{m}} \otimes_V M, N_\alpha) \rightarrow \mathrm{Hom}_{\mathrm{Mod}_A^{\mathrm{fi}}}(\tilde{\mathfrak{m}} \otimes_V M, \varinjlim N_\alpha)$$

is almost injective (resp. an almost isomorphism).

**Definition 3.2.** An  $A$ -module  $P$  is *almost projective* if the functor  $\mathrm{Hom}_{\mathrm{alMod}_A}(P, -) : \mathrm{alMod}_A \rightarrow \mathrm{alMod}_A$  is exact, equivalently,  $\mathrm{Hom}_{\mathrm{Mod}_A^{\mathrm{fi}}}(\tilde{\mathfrak{m}} \otimes_V P, -) : \mathrm{Mod}_A^{\mathrm{fi}} \rightarrow \mathrm{Mod}_A$  is exact.

Any projective  $A$ -module is also almost projective. Let  $\mathrm{alPMod}_A$  denote the full subcategory of  $\mathrm{Mod}_A$  spanned by those objects  $\tilde{\mathfrak{m}} \otimes_V P$  where  $P$  is finitely generated projective morphisms.

**Proposition 3.3.** *The firm module  $\tilde{\mathfrak{m}} \otimes_V A$  is a compact object of  $D(\mathrm{Mod}_A^{\mathrm{fi}})$ .*  $\square$

**Definition 3.4.** Let  $\mathrm{APerf}(A)$  denote the full triangulated subcategory of  $D(\mathrm{Mod}_A^{\mathrm{fi}})$  generated by  $\tilde{\mathfrak{m}} \otimes_V A$ . An *almost perfect complex*  $E$  is an object of  $D(\mathrm{alMod}_A)$  satisfying  $\tilde{\mathfrak{m}} \otimes_V E \in \mathrm{APerf}(A)$ . That is. The category  $\mathrm{Perf}^{\mathrm{al}}(A)$  of almost perfect  $A$ -complexes is defined to be the pullback  $D(\mathrm{alMod}_A) \times_{D(\mathrm{alMod}_A)} \mathrm{APerf}(A)$  of triangulated categories.

**Definition 3.5.** Let  $A$  be an almost  $V$ -algebra and  $K^{\mathrm{al}}(A)$  denote the  $K$ -theory spectrum of the triangulated category of  $D(\mathrm{APerf}(R))$ . We call  $K^{\mathrm{al}}(A)$  the *almost  $K$ -theory spectrum* of  $A$ .

We will characterize almost  $K$ -theory.

**Proposition 3.6.** *The exact functor  $\tilde{\mathfrak{m}} \otimes_V (-) : \mathrm{Mod}_A \rightarrow \mathrm{Mod}_A^{\mathrm{fi}}$  induces an essential surjective functor from the triangulated category of perfect  $A$ -complexes to the one of almost perfect  $A$ -complexes.*  $\square$

**Lemma 3.7.** *Let  $P$  be a finitely generated projective  $A_*$ -module. Then  $P$  is closed.*

**proof.** Clearly  $A_*^n$  is closed for any integer  $n \geq 0$ . By the assumption,  $P$  is a direct factor of some  $A_*^n$ , implying that  $P$  is closed.  $\square$

By Proposition 2.6,  $\tilde{\mathfrak{m}} \otimes_V (-)$  and  $\mathrm{Hom}_V(\tilde{\mathfrak{m}}, -)$  invert almost isomorphisms, implying that one has  $\tilde{\mathfrak{m}} \otimes_V (\mathrm{Hom}_V(\tilde{\mathfrak{m}}, A)) \simeq \tilde{\mathfrak{m}} \otimes_V A$  and  $\mathrm{Hom}_V(\tilde{\mathfrak{m}}, \tilde{\mathfrak{m}} \otimes_V A) \simeq \mathrm{Hom}_V(\tilde{\mathfrak{m}}, A)$ .

*Remark 3.8.* While the triangulated category  $D(\mathrm{Perf}(A_*))$  is contained in  $D(\mathrm{Mod}_A^{\mathrm{cl}})$ , which is distinct to  $D(\mathrm{Mod}_{A_*})$ . An  $A_*$ -module  $\tilde{\mathfrak{m}} \otimes_V A_*$  is not closed in general. Indeed, if  $A_*$  is an unital ring,  $\mathrm{Hom}_V(\tilde{\mathfrak{m}}, \tilde{\mathfrak{m}} \otimes_V A_*) \simeq A_* \neq \tilde{\mathfrak{m}} \otimes_V A_* \not\simeq 1$ .

**3.2. Localization by almost quasi-isomorphisms.** We assume that  $\mathfrak{m}$  is flat. Then the product map  $\mu : \tilde{\mathfrak{m}} \rightarrow \mathfrak{m}$  is an isomorphism. Let  $f : E \rightarrow F$  be a morphism of  $A$ -complexes. Then  $f$  is said to be an *almost quasi-isomorphism* if the induced morphism  $\tilde{\mathfrak{m}} \otimes_V f : \tilde{\mathfrak{m}} \otimes_V E \rightarrow \tilde{\mathfrak{m}} \otimes_V F$  is a quasi-isomorphism.

**Proposition 3.9.** *Let  $f : E \rightarrow F$  be a morphism of  $A$ -complexes. Then the following conditions are equivalent:*

- (1) *The morphism  $f$  is an almost quasi-isomorphism.*
- (2) *The mapping cone:  $\text{cone} f$  is almost acyclic. That is, all of  $\text{cone} f$ 's homology are killed by  $\mathfrak{m}$ .*
- (3) *The firm complex  $\tilde{\mathfrak{m}} \otimes_V \text{cone} f$  is acyclic.*

**proof.** The proof is straightforward. □

Similarly, a morphism  $f : E \rightarrow F$  is a *mod- $\mathfrak{m}$ -quasi-isomorphism* if the induced morphism  $\bar{f} : E/\mathfrak{m}E \rightarrow F/\mathfrak{m}F$  is a quasi-isomorphism.

**Proposition 3.10.** *Let  $f : E \rightarrow F$  be a morphism of  $A$ -complexes. Then the following conditions are equivalent:*

- (1) *The morphism  $f$  is a mod- $\mathfrak{m}$ -quasi-isomorphism.*
- (2) *The mapping cone:  $\text{cone} f$  is mod- $\mathfrak{m}$ -acyclic. That is,  $\text{cone} f/\mathfrak{m}\text{cone} f$  is acyclic.* □

Let  $E$  be a complex of  $A$ -modules and consider the following exact sequence:

$$0 \rightarrow \mathfrak{m} \otimes_V E \rightarrow E \rightarrow E/\mathfrak{m}E \rightarrow 0.$$

Then the followings hold:

**Proposition 3.11.** *Let  $E$  be a complex of  $A$ -modules.*

- (1) *The complex  $E$  is almost acyclic if and only if the projection:  $E \rightarrow E/\mathfrak{m}E$  is a quasi-isomorphism.*
- (2) *The complex  $E$  is mod- $\mathfrak{m}$ -acyclic if and only if the injection:  $\mathfrak{m} \otimes_V E \rightarrow E$  is a quasi-isomorphism.* □

Let  $A$  be a  $V$ -algebra and  $\text{Perf}(A)$  denote the triangulated category of perfect complex of  $A$ -modules. Choosing classes of weak equivalences on  $\text{Perf}(A)$  defines the following  $K$ -theory spectra:

**Definition 3.12.** We define  $K$ -theory spectra as the following:

- The almost Thomason–Trobaugh  $K$ -theory  $K^{\text{aTT}}(A)$  is the Waldhausen  $K$ -theory of  $\text{Perf}(A)$  whose weak-equivalences are almost quasi-isomorphisms.
- The Thomason–Trobaugh  $K$ -theory  $K^{\text{TT}}(A^{\mathfrak{m}})$  is the Waldhausen  $K$ -theory of the category  $\text{Perf}^{\mathfrak{m}}(A)$  spanned by almost acyclic perfect  $A$ -complexes.
- The mod- $\mathfrak{m}$  Thomason–Trobaugh  $K$ -theory  $K^{\text{TT}}(A, \mathfrak{m})$  is the Waldhausen  $K$ -theory of  $\text{Perf}(A)$  whose weak-equivalences are mod- $\mathfrak{m}$  quasi-isomorphisms.

- The Thomason–Trobaugh  $K$ -theory  $K^{\text{TT}}(\mathfrak{m}A)$  is the Waldhausen  $K$ -theory of the category  $\text{Perf}(A)$  spanned by mod- $\mathfrak{m}$  acyclic perfect  $A$ -complexes.

To compare almost  $K$ -theories mentioned, we recall the following the approximation theorem:

**Theorem 3.13** ([38], Theorem 1.6.7 (Approximation theorem)). *Let  $\mathcal{C}$  and  $\mathcal{D}$  be small saturated Waldhausen categories. Let  $F : \mathcal{C} \rightarrow \mathcal{D}$  be an exact functor. If the following conditions hold:*

- (1) *The category  $\mathcal{C}$  is closed under cylinder objects.*
- (2) *A morphism in  $f$  is a weak equivalence in  $\mathcal{C}$  if and only if  $F(f)$  is a weak equivalence in  $\mathcal{D}$ .*
- (3) *The functor  $F$  is a coCartesian fibration. That is, for any object  $c$  of  $\mathcal{C}$  and any map  $x : F(c) \rightarrow d$  in  $\mathcal{D}$ , there exists a map  $a : c \rightarrow c'$  and a weak equivalence  $x' : F(c') \rightarrow d$  such that  $x = x' \circ F(a)$ .*

*Then the induced morphism  $K(F) : K(\mathcal{C}) \rightarrow K(\mathcal{D})$  is a weak equivalence.* □

**Lemma 3.14.** *Those triangulated categories in Definition 3.12 admit cylinder objects.*

**proof.** Since  $\mathfrak{m} \otimes_V -$  and  $- \otimes_V \mathfrak{m}$  are exact and commute with shift and the mapping cone functor, the all of  $\text{Perf}(A)$ ,  $\text{APerf}(A)$ ,  $\text{Perf}^{\text{al}}(A)$ , and  $\text{Perf}(A)$  admit the usual mapping cylinder complexes  $\text{cyl}(f : E \rightarrow F) = \text{cone}(g : \text{cone}f[-1] \rightarrow E)$  satisfying the cylinder axioms. □

The following theorem holds without the flatness of  $\mathfrak{m}$ :

**Theorem 3.15.** *Assume that  $\tilde{\mathfrak{m}}$  is a flat  $V$ -module. Then the functors  $\text{Perf}(A_*) \rightarrow \text{Perf}^{\text{al}}(A) \rightarrow \text{APerf}(A)$  induces weak equivalences*

$$K(\tilde{\mathfrak{m}} \otimes_V (-)) : K^{\text{aTT}}(A_*) \rightarrow K(\text{Perf}^{\text{al}}(A)) \rightarrow K^{\text{al}}(A)$$

*of  $K$ -theory spectra.*

**proof.** Clearly,  $\tilde{\mathfrak{m}} \otimes_V (-)$  is exact and sends all almost quasi-isomorphisms to quasi-isomorphism of almost perfect complexes. By definition of almost quasi-isomorphisms, the condition (2) in Lemma 3.13 is satisfied. By the definition of  $\text{Perf}^{\text{al}}(A)$  and  $\text{APerf}(A)$ , the condition (3) in Lemma 3.13 holds, implying that the canonical morphism  $K(\text{Perf}^{\text{al}}(A)) \rightarrow K^{\text{al}}(A)$  is an equivalence. Let  $E$  be a perfect  $A$ -complex and  $F$  be a firm almost perfect complex. Given an morphism  $f : \tilde{\mathfrak{m}} \otimes_V E \rightarrow F \simeq \tilde{\mathfrak{m}} \otimes_V F$ , we obtain a morphism  $\text{Hom}(\tilde{\mathfrak{m}}, f) : \text{Hom}_V(\tilde{\mathfrak{m}}, E) \rightarrow \text{Hom}_V(\tilde{\mathfrak{m}}, F)$ . Then  $\text{Hom}_V(\tilde{\mathfrak{m}}, E)$  and  $\text{Hom}_V(\tilde{\mathfrak{m}}, F)$  are perfect complexes of  $A_*$ -modules. Therefore,  $K^{\text{aTT}}(A_*) \rightarrow K^{\text{al}}(A)$  is also an equivalence. □

By the Fibration theorem of Waldhausen  $K$ -theory [38, Proposition 1.6.4], we obtain the following:

**Theorem 3.16.** *Let  $A$  be an almost  $V$ -algebra. Assume that  $\mathfrak{m}$  is flat. Then the sequence*

$$K^{\mathrm{TT}}(A_*^{\mathfrak{m}}) \rightarrow K^{\mathrm{TT}}(A_*) \rightarrow K^{\mathrm{al}}(A)$$

and

$$K^{\mathrm{TT}}(\mathfrak{m}A_*) \rightarrow K^{\mathrm{TT}}(A_*) \rightarrow K^{\mathrm{TT}}(A_*, \mathfrak{m})$$

are homotopy fiber sequences of  $K$ -theory spectra.  $\square$

#### 4. FINITE SYNTOMIC ALGEBRAIC COBORDISM OF ALMOST ALGEBRAS

Fix a base ring  $V$  which is unital and commutative and an almost  $V$ -algebra  $A$ . Let  $\mathfrak{m}$  be an idempotent ideal of  $V$ . Assume that  $\mathfrak{m}$  is a flat  $V$ -module. Then the canonical map  $\mu_{\mathfrak{m}} : \tilde{\mathfrak{m}} \rightarrow \mathfrak{m}$  is bijective.

**4.1. Definition of finite syntomic motivic model structure of the category of simplicial presheaves.** Let  $\mathcal{X}$  be a Grothendieck site with an interval object  $I$ . We assume that  $\mathcal{X}$  has enough points: That is, a morphism  $f : X \rightarrow Y$  in  $\mathcal{X}$  is an isomorphism if  $f_x : X_x \rightarrow Y_x$  is an isomorphism of sets for any point  $x : * \rightarrow \mathcal{X}$  where the functor  $(-)_x : \mathcal{X} \rightarrow \mathbf{Sets}$  denotes the right adjoint of the induced functor  $x_* : \mathbf{Sets} \rightarrow \mathcal{X}$ . A simplicial object  $U_{\bullet} : \Delta^{\mathrm{op}} \rightarrow \mathcal{X}$  with an augmentation  $\pi : U_{\bullet} \rightarrow X \in \mathcal{X}$  is a *hypercov*er of  $X$  if the following conditions are hold:

- For any  $n \geq 0$ ,  $U_{\bullet}([n])$  is a coproduct of compact objects represented by small objects of  $\mathcal{X}$ .
- The augmentation  $\pi : U_{\bullet} \rightarrow X$  is a stalk-wise trivial Kan fibration: That is,  $\pi_x : U_{x,\bullet} \rightarrow *$  is a trivial Kan fibration for any point  $x : * \rightarrow X$ .

The category  $\mathbf{Set}_{\Delta}$  of simplicial set has a proper combinatorial simplicial model structure, called *Kan–Quillen model structure*. Then the injective model structure of the category  $\mathbf{Set}_{\Delta}^{\mathcal{X}^{\mathrm{op}}}$  of simplicial presheaves on  $\mathcal{X}$  is also proper combinatorial. Let  $\mathbf{MS}_{\mathcal{X}}^{\Delta}$  denote the Bousfield localization of  $\mathbf{Set}_{\Delta}^{\mathcal{X}^{\mathrm{op}}}$  defined as follows: A simplicial presheaf  $F$  is *motivic local* if  $F$  satisfies the following conditions:

- The presheaf  $F$  is stalk-wise fibrant.
- For any hypercover  $\pi : U_{\bullet} \rightarrow X$  of  $X \in \mathcal{X}$ , the induced map  $F(f) : F(X) \rightarrow F(|U_{\bullet}|)$  is a weak equivalence. Here the functor  $| - | : \mathbf{Fun}(\Delta^{\mathrm{op}}, \mathcal{X}) \rightarrow \mathcal{X}$  denotes the geometric realization of simplicial objects.
- The canonical map  $\mathbf{1} : I \rightarrow *$  induces a weak equivalence  $F(U) \rightarrow F(U \times I)$  for any  $U \in \mathcal{X}$ .

A map  $f : F \rightarrow G$  of simplicial presheaves on  $\mathcal{X}$  is a *motivic equivalence* if the induced map

$$f^* : \mathrm{Hom}(G, Z) \rightarrow \mathrm{Hom}(F, Z)$$

is a weak homotopy equivalence of simplicial sets for each motivic local presheaf  $Z$ . By [1, p.56, Corollary 4.55], the injective model structure of  $\text{Set}_\Delta^{\text{cop}}$  is proper combinatorial and symmetric monoidal. Therefore the Bousfield localization  $\mathbf{MS}_X^\Delta$  of  $\mathbf{sSet}^{\text{cop}}$  is also proper.

We apply the finite syntomic site on the category  $\mathbf{Sch}_V^{\text{fp}}$  of finitely presented schemes over  $\text{Spec}V$ , and the interval object  $\mathbb{A}_V^1$ . We let  $\mathbf{MS}_{\text{FSyn}}$  denote the  $\infty$ -category determined by the simplicial model category  $\mathbf{MS}_{\mathbf{Sch}_V^{\text{fp}}}^\Delta$  and we call objects of  $\mathbf{MS}_{\text{FSyn}}$  (*finite syntomic*) *motivic spaces*. Moreover,  $\mathbf{MSp}_{\text{FSyn}}$  denotes the stable  $\infty$ -category of (finite syntomic) motivic spectra defined by

$$\mathbf{MSp}_{\text{FSyn}} = \varprojlim(\cdots \xrightarrow{\Omega_{\mathbb{P}_+^1}} (\mathbf{MS}_{\text{FSyn}})_* \xrightarrow{\Omega_{\mathbb{P}_+^1}} (\mathbf{MS}_{\text{FSyn}})_*),$$

where  $(\mathbf{MS}_{\text{FSyn}})_*$  the  $\infty$ -category of pointed motivic spaces and  $\Omega_{\mathbb{P}_+^1}(-) = \text{Map}_{(\mathbf{MS}_{\text{FSyn}})_*}(\mathbb{P}_+^1, -)$  is the pointed  $\mathbb{P}^1$ -loop functor. The stable  $\infty$ -category  $\mathbf{MSp}_{\text{FSyn}}$  is the full subcategory of the model category  $\text{Spt}_{\mathbb{P}_+^1}(\mathbf{MS}_{\text{FSyn}}^\Delta)$  spanned by  $\mathbb{P}_+^1$ -stable fibrant objects. (See [17, Section 10.2]).

**Theorem 4.1** ([19] Corollary 3.9). *The localization  $L_{\text{FSyn}} : \mathbf{MSp} \rightarrow \mathbf{MSp}$  by the family of zero-section stable finite syntomic surjective morphisms induces a categorical equivalence of stable  $\infty$ -categories  $L_{\text{FSyn}}(\mathbf{MSp}) \rightarrow \mathbf{MSp}^{\text{FSyn}}$ .  $\square$*

**4.2. The left adjoint of the localization  $(-)^a : \text{CAlg}(\text{Mod}_A) \rightarrow \text{CAlg}(\text{alMod}_A)$  from the non-unital algebra viewpoint.** We will review almost mathematics from the theory of non-unital algebra. We assume that  $1 \notin \mathfrak{m}$ . Let  $A$  be an almost  $V$ -algebra and  $B$  an  $A$ -algebra. Then  $B_{\mathfrak{I}} := \tilde{\mathfrak{m}} \otimes_V \text{Hom}_A(\tilde{\mathfrak{m}}, B)$  is a (non-unital)  $B$ -algebra. Consider the unitalization functor

$$V \oplus (-) : \text{CAlg}^{\text{nu}}(\text{Mod}_A) \rightarrow \text{CAlg}(\text{Mod}_A)$$

Then the direct sum  $V \oplus B_{\mathfrak{I}}$  has a canonical unital ring structure defined by

$$(v, b) \cdot (v', b') = (vv', vb' + v'b + bb')$$

for  $v, v' \in V$  and  $b, b' \in B$ . Note the functor  $V \oplus - : \text{CAlg}^{\text{nu}}(\text{Mod}_A) \rightarrow \text{CAlg}(\text{Mod}_A)$  induces a categorical equivalence between  $\text{CAlg}^{\text{nu}}(\text{Mod}_A)$  and the category of augmented commutative rings  $\text{CAlg}(\text{Mod}_A)_{/V}$ . The right adjoint is the augmented ideal functor  $\text{Ker}(- \rightarrow V) : \text{CAlg}(\text{Mod}_A)_{/V} \rightarrow \text{CAlg}^{\text{nu}}(\text{Mod}_A)$ , becoming the quasi-inverse of  $V \oplus -$ .

Let  $B \leftarrow A \rightarrow C$  be a diagram of commutative non-unital rings. Then the coCartesian product  $B \oplus_A C$  has a canonical non-unital commutative algebra structure. The following proposition justifies that  $B \oplus_A C$  is the colimit of the diagram  $B \leftarrow A \rightarrow C$  of non-unital algebras:

**Proposition 4.2.** *Let  $V$  be an unital commutative ring and  $B \leftarrow A \rightarrow C$  a diagram of commutative non-unital  $V$ -algebras. Then the unit of the adjunction*

$$V \oplus (-) : \text{CAlg}^{\text{nu}}(\text{Mod}_V) \rightleftarrows \text{CAlg}(\text{Mod}_V)_{/V} : \text{Ker}(- \rightarrow V)$$

induces an isomorphism:

$$V \oplus (B \oplus_A C) \rightarrow (V \oplus B) \otimes_{V \oplus A} (V \oplus C)$$

of unital  $V$ -algebras.

**proof.** The proof is straightforward: By the following commutative diagram:

$$\begin{array}{ccccccc} 0 & \longrightarrow & B & \longrightarrow & (V \oplus B) \otimes_V (V \oplus C) & \longrightarrow & V \oplus C \longrightarrow 0 \\ & & \parallel & & \parallel & & \parallel \\ 0 & \longrightarrow & K & \longrightarrow & (V \oplus B) \otimes_V (V \oplus C) & \longrightarrow & V \longrightarrow 0 \end{array}$$

where both horizontal sequences are exact, one can easily check that the argumentation ideal  $K$  of  $(V \oplus B) \otimes_V (V \oplus C)$  is isomorphic to  $B \oplus C$ . Therefore the canonical map  $V \oplus (B \oplus C) \rightarrow (V \oplus B) \otimes_V (V \oplus C)$  is an isomorphism, inducing a coequalizer sequence:

$$V \oplus A \rightrightarrows V \oplus (B \oplus C) \longrightarrow (V \oplus B) \otimes_{V \oplus A} (V \oplus C) \longrightarrow 0.$$

Hence there exists a canonical isomorphism:  $V \oplus (B \oplus_A C) \rightarrow (V \oplus B) \otimes_{V \oplus A} (V \oplus C)$   $\square$

Let  $B_{!!}$  denote the coCartesian product  $V \oplus_{\tilde{m}} B_!$ . Then one has an exact sequence

$$\tilde{m} \rightarrow V \oplus B_! \rightarrow B_{!!} \rightarrow 0,$$

where the left map is almost injective. After tensoring with  $\tilde{m}$ , indeed, one has a trivial split exact sequence:

$$0 \rightarrow \tilde{m} \rightarrow \tilde{m} \oplus (\tilde{m} \otimes_V B_!) \rightarrow \tilde{m} \otimes_V B_{!!} \rightarrow 0,$$

implying that  $\tilde{m} \otimes_V B_! \rightarrow \tilde{m} \otimes_V B_{!!}$  is an isomorphism. Therefore the functor  $(-)_!! : \text{CAlg}(\text{Mod}_A) \rightarrow \text{CAlg}(\text{alMod}_{A!!})$  is left adjoint to the localization  $(-)^a : \text{CAlg}(\text{Mod}_{A!!}) \rightarrow \text{CAlg}(\text{alMod}_A)$ .

If  $\tilde{m} \rightarrow \tilde{m} \oplus (\tilde{m} \otimes_V B_!)$  is exactly injective,  $B$  is called an *exact* almost  $V$ -algebra [13, Definition 2.2.27]. In the case  $\tilde{m} \simeq \mathfrak{m}$ , in particular, that  $\mathfrak{m}$  is flat,  $B$  is always exact [13, Remark 2.2.28].

**4.3. Some of almost homotopical algebra.** We will recall and prove some result of almost homotopical algebra.

**Definition 4.3.** Let  $A$  be an almost  $V$ -algebra. The *Jacobson radical* is defined to be  $\text{Jac}(A) = \text{Jac}(A_*)^a \subset A$ .

Clearly  $I \rightarrow \text{Jac}(A)$  is almost injective if and only if  $\tilde{m} \otimes_V I \subset \text{Jac}(A_*)$ .

**Definition 4.4.** Let  $A$  be an almost  $V$ -algebra and  $I$  an ideal of  $A$ . We say that  $I$  is *tight* if there exists a finitely generated ideal  $\mathfrak{m}_0 \subset \mathfrak{m}$  and an integer  $n \geq 0$  such that  $I^n \subset \mathfrak{m}_0 A$ .

**Proposition 4.5** ([13] Corollary 5.1.17.). *Let  $A$  be an almost  $V$ -algebra,  $I$  a tight ideal of  $A$  contained in the Jacobson radical of  $A$ . Then  $I_*$  is contained in the Jacobson radical of  $A_*$ .*  $\square$

**Lemma 4.6** ([13] Lemma 5.1.7.). *Let  $A$  be an almost  $V$ -algebra,  $I$  a tight ideal of  $A$  which is contained in the Jacobson radical of  $A$ . If  $M$  is an almost finitely generated  $A$ -module satisfying  $IM = M$ , then exactly  $M = 0$ .*  $\square$

**Corollary 4.7** ([13] Corollary 5.1.8.). *Let  $A$  be an almost  $V$ -algebra,  $I$  a tight ideal of  $A$  which is contained in the Jacobson radical of  $A$ . Let  $f : M \rightarrow N$  be a homomorphism of almost finitely generated projective  $A$ -modules. If  $f \otimes_A A/I : M/IM \rightarrow N/IN$  is an isomorphism. Then  $f$  is also an isomorphism.*  $\square$

**Theorem 4.8** ([13] Theorem 5.3.24.). *Let  $A$  be an almost  $V$ -algebra,  $I$  a tight ideal of  $A$  which is contained in the Jacobson radical of  $A$ . Write  $A_n = A/I^n$  for each  $n \geq 1$ , and  $A_\infty = \varprojlim A_n$ . Then*

$$\varprojlim : 2 - \varprojlim \text{alPMod}_{A_n} \rightarrow \text{alPMod}_{A_\infty}$$

*is a categorical equivalence.*  $\square$

**Theorem 4.9.** *Let  $A$  be an almost  $V$ -algebra and  $I$  a tight ideal which is contained in the Jacobson radical of  $A$ . For  $n \geq 1$ , set  $A_n = A/I^n$ , and  $A_\infty = \varprojlim A_n$ . Then the canonical adjunction  $(- \otimes_A^\mathbb{L} A_n) : D(A_\infty) \rightleftarrows 2 - \varprojlim D(A_n) : \mathbb{R} \varprojlim$  induces a categorical equivalence*

$$\mathbb{R} \varprojlim : 2 - \varprojlim \text{APerf}(A_n) \rightarrow \text{APerf}(A_\infty)$$

*between stable  $\infty$ -categories.*

**proof.** Let  $E$  be an almost perfect  $A_\infty$ -complex. Since  $E$  is a dualizable object of the derived category  $D(A_\infty^a)$ ,  $E \otimes_{A_\infty}^\mathbb{L} (-)$  preserves all small limits. Therefore the canonical morphism  $E_* \rightarrow \mathbb{R} \varprojlim (E \otimes_{A_\infty}^\mathbb{L} A_n)_*$  is a quasi-isomorphism.

Conversely, let  $(E_n)$  be an inverse system of complex of almost finitely generated projective  $A_n$ -modules. Since  $(A_*)_\infty \rightarrow A_*/(I_*)^n$  is surjective for each  $n \geq 1$ ,  $\varprojlim E_n \rightarrow E_n$  is also surjective for each  $n \geq 1$ . Therefore, clearly, the canonical morphism  $(\varprojlim E_n) \otimes_{A_\infty} A_n \rightarrow E_n$  is a quasi-isomorphism for each  $n \geq 1$ . Hence the morphism  $(\varprojlim E_n) \otimes_{A_\infty} A_n \rightarrow (E_n)$  is an inverse system of quasi-isomorphisms.  $\square$

#### 4.4. Almost finite syntomic morphisms.

**Definition 4.10.** A morphism  $f : A \rightarrow B$  of almost  $V$ -algebras is *almost syntomic* if the following conditions are satisfied:

- (1) An  $A$ -algebra  $B$  is a almost finitely projective  $A$ -module.
- (2) The almost relative cotangent complex  $L_{B/A} := B_{!!} \otimes_{(V \oplus B)_{!!}} L_{(V \oplus B)_{!!}/(V \oplus A)_{!!}}$  is almost perfect and tor-amplitude in  $[-1, 0]$  in the category  $D(\text{Mod}_{B_{!!}})$ .

Fix a regular cardinal  $\kappa$ . We write  $\mathbf{MS} = \mathbf{MS}_{\text{FSyn}}$  and let  $\mathbf{MS}^\kappa$  denote full subcategory spanned by all  $\kappa$ -small objects of  $\mathbf{MS}$ . Referring the results [10, Theorem 3.4.1 and Lemma 3.5.1] and [9, Lemma 5.1.3], we define the almost version of algebraic cobordism:

**Definition 4.11.** Let  $A$  be an almost  $V$ -algebra. Let  $\text{alFSyn}(A)$  denote the largest groupoid of the category  $\text{Alg}_A^{\text{alFSyn}}$ , and set  $\text{alMGL} = \Sigma_+^\infty \text{alFSyn}^\kappa(-)$ , where  $\text{alFSyn}^\kappa$  is the right Kan extension of the restriction  $\text{alFSyn}|_{(\mathbf{MS}^\kappa)^{\text{op}}}$  along the inclusion  $(\mathbf{MS}^\kappa)^{\text{op}} \rightarrow \mathbf{MS}^{\text{op}}$ . We say that the motivic spectrum  $\text{alMGL}$  is the *almost algebraic cobordism*.

Let  $V$  be a commutative ring and  $\omega \in V$  a non-zero divisor. For  $n \geq 0$ , set  $A_n = A/\omega^{n+1}A$ . Write  $\overline{A}_n = \omega^n A/\omega^{n+1}A$ . Since  $\overline{A}_n$  is square-zero non-unital ring, the  $A$ -linear inclusion:  $\overline{A}_n \rightarrow A_n$  is a ring homomorphism as non-unital rings, inducing a ring homomorphism:

$$(4.1) \quad \varphi_n : (\overline{A}_n)_{!!} \rightarrow (A_n)_{!!}$$

Hence  $(A_n)_{!!}$  has an  $(\overline{A}_n)_{!!}$ -algebra structure.

By the canonical isomorphism  $\overline{A}_1 \rightarrow \overline{A}_n$ , the following holds.

**Lemma 4.12.** *The linear map  $(\omega^{n-1} \cdot \text{Id})_{!!} : (A_n)_{!!} \rightarrow (A_n)_{!!}$  induces an isomorphism:  $(\overline{A}_1)_{!!} \rightarrow (\overline{A}_n)_{!!}$  of  $A_{!!}$ -algebras.  $\square$*

**4.5. The almost algebraic cobordism of perfectoid algebras.** We recall the definition of perfectoid algebra.

**Definition 4.13.** Let  $K$  be a complete non-Archimedean non-discrete valuation field of rank 1, and  $K^\circ$  denote the subring of powerbounded elements. We say that  $K$  is a *perfectoid field* if the Frobenius  $\Phi : K^\circ/p \rightarrow K^\circ/p$  is surjective, where  $p$  is a positive prime integer which is equal to the characteristic of the residue field of  $K^\circ$ .

In this section, we fix a perfectoid field  $K$  whose valuation ring  $K^\circ$  is mixed characteristic  $(0, p)$ . We put  $V = K^\circ$  and  $\mathfrak{m} = K^{\circ\circ}$ , where  $K^{\circ\circ} = \{x \in K \mid |x| < 1\}$  is the maximal ideal of  $K^\circ$ . Then  $\mathfrak{m}$  is the set of topologically nilpotent elements, being an idempotent ideal. We fix a pseudouniformizer  $\omega \in V$  with  $|p| \leq |\omega| < 1$ . In this case, the ideal  $\mathfrak{m} = \varinjlim_{n \geq 1} \omega^{1/p^n} V$ , which is a filtered colimit of free  $V$ -modules, is flat by Lazard's theorem.

**Definition 4.14.** An integral perfectoid  $V$ -algebra is an  $\omega$ -adic complete flat  $V$ -algebra  $A$  on which Frobenius induces an isomorphism  $\Phi : A/\omega^{\frac{1}{p}}A \rightarrow A/\omega A$ .

For any  $V$ -algebra  $B$ , let  $B^{\flat}$  denote the tilting algebra  $\varprojlim_{x \rightarrow x^p} B/\omega B$  of  $B$ . The tilting ideal  $\mathfrak{m}^{\flat} \subset V^{\flat}$  is a flat  $V^{\flat}$ -module as  $\mathfrak{m}$  is. Therefore, we can apply almost mathematics to perfectoid algebras.

**Proposition 4.15.** *Let  $K$  be a perfectoid field with the valuation ring  $V$  whose residue field is of characteristic  $p > 0$ . Let  $\omega$  be a pseudouniformizer and  $m = \varinjlim_{m \geq 1} \omega^{p^{-m}} V$ . Let  $A$  be an integral perfectoid  $V$ -algebra and write  $A_n = A/\omega^n A$ . Then  $(A_n)_{!!}$  is an ind-finite syntomic  $(\overline{A_1})_{!!}$ -algebra for  $n \geq 1$ .*

**proof.** We may assume  $A_* = A$  and  $A_{!} = m \otimes_V A$ . The injection  $(\overline{A_n})_{!} \rightarrow (A_n)_{!}$  is induced by the inductive system of (non-unital) ring homomorphisms:

$$\omega^{n+\frac{1}{p^m}} A / \omega^{n+1+\frac{1}{p^m}} A \rightarrow \omega^{\frac{1}{p^m}} A / \omega^{n+1+\frac{1}{p^m}} A,$$

which induces a finite syntomic homomorphism

$$\varphi_{n,m} : V \oplus (\omega^{\frac{1}{p^m}})^{np^m+1} A / \omega^{n+1+\frac{1}{p^m}} A \rightarrow V \oplus \omega^{\frac{1}{p^m}} A / \omega^{n+1+\frac{1}{p^m}} A$$

for each  $m \geq 1$ . Since the functor  $(-)_{!!}$  preserves all small colimit, by definition of almost relative cotangent complex,  $(A_n)_{!!}$  is ind-finite syntomic over  $(\overline{A_1})_{!!}$  by Lemma 4.12 for each  $n \geq 1$ .  $\square$

For an almost  $V$ -algebra  $A$ , let  $\text{Alg}_A^{\text{alFSyn}}$  denote the category of almost finite syntomic  $A$ -algebras. The results in Section 4.3 provides us the following:

**Proposition 4.16.** *Let  $A$  be an almost  $V$ -algebra and  $I$  a tight ideal which is contained in the Jacobson radical of  $A$ . Let  $(A_n)_{n \geq 1}$  be an inverse system of commutative rings and write  $A = \varprojlim A_n$ . The functor  $(-\otimes_A A_n)_{n \geq 1} : \text{Alg}_A^{\text{alFSyn}} \rightarrow 2\text{-}\varprojlim_n \text{Alg}_{A_n}^{\text{alFSyn}}$  is a categorical equivalence.*

**proof.** Let  $(B_n)$  be an inverse system of almost finite syntomic  $A_n$ -algebras. Then, by the assumption,  $B = \varprojlim A_n$  is almost finite, projective  $A$ -algebra. We show that the relative cotangent complex  $L_{B/A}$  is perfect and tor-amplitude in  $[-1, 0]$ . Since  $B \rightarrow B_n$  is unramified, the sequence  $A \rightarrow B \rightarrow B_n$  induces a quasi-isomorphism  $\tilde{m} \otimes_V L_{B/A} \otimes_B^{\mathbb{L}} B_n \rightarrow \tilde{m} \otimes_V L_{B_n/A}$ . Similarly, the sequence  $A \rightarrow A_n \rightarrow B_n$  induces a quasi-isomorphism  $\tilde{m} \otimes_V L_{B_n/A} \rightarrow \tilde{m} \otimes_V L_{B_n/A_n}$ . Therefore  $\tilde{m} \otimes_V L_{B/A} \otimes_B^{\mathbb{L}} B_n \rightarrow \tilde{m} \otimes_V L_{B_n/A_n}$  is a quasi-isomorphism. Since each  $B_n$  is an almost finitely generated projective  $A$ -module,  $\tilde{m} \otimes_V B \otimes_A^{\mathbb{L}} A_n \rightarrow \tilde{m} \otimes_V B_n$  is an isomorphism for each  $n$ . Then  $\tilde{m} \otimes_V L_{B/A} \otimes_A^{\mathbb{L}} A_n \rightarrow \tilde{m} \otimes_V L_{B/A} \otimes_B^{\mathbb{L}} (B \otimes_A A_n) \rightarrow \tilde{m} \otimes_V L_{B/A} \otimes_B^{\mathbb{L}} B_n$  is a composition of quasi-isomorphisms for each  $n$ . By Theorem 4.9,  $\tilde{m} \otimes_V L_{B/A} \rightarrow \varprojlim \tilde{m} \otimes_V L_{B_n/A_n}$  is a quasi-isomorphism and they are almost perfect complexes of  $B$ -modules. By the Milnor exact sequence

$$0 \rightarrow \varprojlim^1 H_{\bullet+1}(\tilde{m} \otimes_V L_{B_n/A_n}) \rightarrow H_{\bullet}(\varprojlim \tilde{m} \otimes_V L_{B/A}) \rightarrow \varprojlim H_{\bullet}(\tilde{m} \otimes_V L_{B_n/A_n}) \rightarrow 0,$$

the projective limit  $\tilde{m} \otimes_V L_{B_n/A_n}$  is tor-amplitude in  $[-1, 0]$  in  $D(\text{Mod}_{B_n})$ .  $\square$

**Corollary 4.17.** *Let  $A$  be an almost  $V$ -algebra and  $I$  a tight ideal which is contained in the Jacobson radical of  $A$ . Let  $(A_n)_{n \geq 1}$  be an inverse system of commutative rings and write  $A = \varprojlim A_n$ . The induced map  $\text{alFSyn}^k(\text{Spec} A) \rightarrow \text{alFSyn}^k(\varinjlim \text{Spec} A_n)$  is a weak equivalence of spaces.*

**proof.** We may assume that the functor  $\text{Spec}A$  is small. By Proposition 4.16, one has a chain of weak equivalences:

$$\begin{aligned} \text{alFSyn}(\text{Spec}A) &\simeq \text{Map}_{\text{Cat}}(\Delta^0, \text{alFSyn}_A) \simeq \text{Map}_{\text{Cat}}(\Delta^0, 2 - \varprojlim \text{alFSyn}_{A_n}) \\ &\simeq \varprojlim \text{Map}_{\text{Cat}}(\Delta^0, \text{alFSyn}_{A_n}) \simeq \varprojlim \text{alFSyn}(\text{Spec}A_n), \end{aligned}$$

where  $\text{Cat}$  denotes the large category of small categories that mapping spaces are the largest groupoids of the functor categories. Therefore, one has an equivalences:  $\text{alFSyn}^\kappa(\text{Spec}A) \simeq \varprojlim \text{alFSyn}^\kappa(\text{Spec}A_n) \simeq \text{alFSyn}^\kappa(\varinjlim \text{Spec}A_n)$ .  $\square$

*Remark 4.18.* Proposition 4.16 and its corollary holds under the only the assumption that  $\varprojlim : 2 - \varprojlim \text{alPMod}_{A_n} \rightarrow \text{alPMod}_A$  is a categorical equivalence.

**Lemma 4.19.** *On the condition of Proposition 4.15, let  $A_n^+$  denote the cokernel of*

$$\mathfrak{m}/\omega^n \mathfrak{m} \rightarrow V/\omega^n \oplus (\mathfrak{m} \otimes_V A/\omega^n A)$$

*for each  $n$ . Then there exists a canonical isomorphism  $(A_n)_{!!} \rightarrow A_n^+$  for each  $n \geq 1$ .*

**proof.** Consider the following commutative diagram:

$$\begin{array}{ccccc} \mathfrak{m} & \longrightarrow & \mathfrak{m}/\omega^n \mathfrak{m} & \longrightarrow & \mathfrak{m} \otimes_V A_n \\ \downarrow & & \downarrow & & \downarrow \\ V & \longrightarrow & V/\omega^n & \longrightarrow & A_n^+, \end{array}$$

where both squares are coCartesian. Therefore the morphism  $(A_n)_{!!} \rightarrow A_n^+$  is canonically an isomorphism.  $\square$

Set  $\text{alFSyn}^b(A) = \text{alFSyn}(A^b)$  for any perfectoid algebra  $A$ , and write  $\text{alMGL}^b = \Sigma_+^\infty \text{alFSyn}^b$ . Finally, we prove the tilting equivalence of almost algebraic cobordism:

**Theorem 4.20.** *Let  $V$  be a mixed characteristic perfectoid valuation ring with unit and  $V^b$  denote the tilting. Then these spectra  $\text{alMGL}$  and  $\text{alMGL}^b$  are equivalent on zero-section stable affine integral perfectoid schemes over  $V$ .*

**proof.** By Lemma 4.19, there exists a zig-zag of isomorphisms

$$(A_1)_{!!} \rightarrow (A_1^+)_{!!} \leftarrow (A_1^b)_{!!}^+ \leftarrow (A_1^b)_{!!}$$

of commutative rings, where the middle map is a canonical isomorphism. By Proposition 4.15, for each  $n \geq 1$ ,  $\text{alMGL}(Z_0(A_n))$  (resp.  $\text{alMGL}(Z_0(A_n^b))$ ) is weakly equivalent to the same  $\text{alMGL}(A_1)$  (resp.  $\text{alMGL}(A_1^b)$ ), implying that  $\mathbb{R} \varprojlim \text{alMGL}(Z_0(A_n))$  and  $\mathbb{R} \varprojlim \text{alMGL}(Z_0(A_n^b))$  are weakly equivalent, where  $Z_0$  denotes the zero-section stabilization. Note that the zero-section stabilization preserves small colimits. The zero-section stability of  $\varinjlim \text{Spec}A_n$  implies

that one has a chain of equivalences  $\mathbb{L} \varinjlim \mathrm{Spec} A_n \simeq \mathbf{Z}_0(\mathbb{L} \varinjlim \mathrm{Spec} A_n) \simeq \mathbb{L} \varinjlim \mathbf{Z}_0(\mathrm{Spec} A_n)$ . Similarly, one has  $\mathbb{L} \varinjlim \mathrm{Spec} A_n^b \simeq \mathbb{L} \varinjlim \mathbf{Z}_0(\mathrm{Spec} A_n^b)$ . Consider the diagram of fiber sequences:

$$\begin{array}{ccccc} \mathrm{alMGL}(\mathbb{L} \varinjlim \mathrm{Spec} A_n) & \longrightarrow & \mathrm{alMGL}(\oplus_n \mathbf{Z}_0(\mathrm{Spec} A_n)) & \xrightarrow[\oplus t_n^*]{\mathrm{Id}} & \mathrm{alMGL}(\oplus_n \mathbf{Z}_0(\mathrm{Spec} A_n)) \\ \downarrow & & \downarrow & & \downarrow \\ \mathbb{R} \varprojlim \mathrm{alMGL}(\mathbf{Z}_0(A_n)) & \longrightarrow & \prod_n \mathrm{alMGL}(\mathbf{Z}_0(A_n)) & \xrightarrow[\prod t_n]{\mathrm{Id}} & \prod_n \mathrm{alMGL}(\mathbf{Z}_0(A_n)), \end{array}$$

where the middle and right vertical arrows are weak equivalences. Since each  $\mathrm{Spec} A_n$  is cofibrant and closed immersion  $t_n^* : \mathrm{Spec} A_n \rightarrow \mathrm{Spec} A_{n+1}$  a cofibration with respect to the motivic model structure, the inductive system  $(\mathrm{Spec} A_n)_{n \geq 1}$  is injectively cofibrant. Therefore  $\varinjlim \mathrm{Spec} A_n \rightarrow \mathbb{L} \varinjlim \mathrm{Spec} A_n$  is a motivic equivalence. Furthermore, by Proposition 4.16, one has weak equivalences of spectra  $\mathrm{alFSyn}^k(\mathrm{Spec} A) \simeq \mathrm{alFSyn}^k(\varinjlim \mathrm{Spec} A_n)$  and  $\mathrm{alFSyn}^k(\mathrm{Spec} A^b) \simeq \mathrm{alFSyn}^k(\varinjlim \mathrm{Spec} A_n^b)$ . Hence  $\mathrm{alMGL}$  and  $\mathrm{alMGL}^b$  are weakly equivalent on the category of integral perfectoid  $V$ -algebras.  $\square$

#### REFERENCES

- [1] C Barwick, *On left and right model categories and left and right Bousfield localizations*, Homology, Homotopy Appl. **12** (2010), no. 2, 245–320.
- [2] J. E. Bergner, *A model category structure on the category of simplicial categories*, Trans. Amer. Math. Soc. **359** (2007), no. 5, 2043–2058, DOI 10.1090/S0002-9947-06-03987-0. MR2276611 (2007i:18014)
- [3] D. Dugger, S. Hollander, and D. C. Isaksen, *Hypercovers and simplicial presheaves*, Math. Proc. Cambridge Philos. Soc. **136** (2004), no. 1, 9–51, DOI 10.1017/S0305004103007175. MR2034012
- [4] D. Dugger and D. C. Isaksen, *Topological hypercovers and  $\mathbb{A}^1$ -realizations*, Math. Z **246** (2004), 667–689.
- [5] W. G. Dwyer and D. M. Kan, *Calculating simplicial localizations*, J. Pure Appl. Algebra **18** (1980), no. 1, 17–35, DOI 10.1016/0022-4049(80)90113-9. MR578563
- [6] ———, *Simplicial localizations of categories*, J. Pure Appl. Algebra **17** (1980), no. 3, 267–284, DOI 10.1016/0022-4049(80)90049-3. MR579087
- [7] ———, *Function complexes in homotopical algebra*, Topology **19** (1980), no. 4, 427–440, DOI 10.1016/0040-9383(80)90025-7. MR584566
- [8] Elden Elmanto, Marc Hoyois, Adeel A. Khan, Vladimir Sosnilo, and Maria Yakerson, *Framed transfers and motivic fundamental classes*, J. Topol. **13** (2020), no. 2, 460–500, DOI 10.1112/topo.12134. MR4092773
- [9] ———, *Motivic infinite loop spaces*, Camb. J. Math. **9** (2021), no. 2, 431–549.
- [10] ———, *Modules over algebraic cobordism*, Forum of Mathematics, Pi **8** (2020), 1–44, DOI 10.1017/fmp.2020.13.
- [11] D. Gepner and V. Snaith, *On the motivic spectra representing algebraic cobordism and algebraic K-theory*, Documenta Mathematica **14** (2009), 359–396.
- [12] Faltings Gerd, *p-adic Hodge theory*, Jarnal of american mathematical society **1** (1988), no. 1, 255–299, DOI 10.2307/1990970.
- [13] Ofer Gabber and Lorenzo Ramero, *Almost ring theory*, Lecture Notes in Mathematics, vol. 1800, Springer-Verlag, Berlin, 2003. MR2004652

- [14] P.G. Goerss and J.F. Jardine, *Simplicial homotopy theory*, Progress in Mathematics, vol. 174, Birkhäuser Verlag, Basel, 1999. MR1711612 (2001d:55012)
- [15] P. Hu, *S-modules in the category of schemes*, Mem. Amer. Math. Soc. **161** (2003), no. 767.
- [16] J. F. Jardine, *Motivic symmetric spectra*, Documenta Mathematica **5** (2000), 445–553 (electronic).
- [17] ———, *Local homotopy theory*, Springer Monographs in Mathematics, Springer, New York, 2015. MR3309296
- [18] A. Joyal, *Quasi-categories and Kan complexes*, J. Pure Appl. Algebra **175** (2002), no. 1-3, 207–222. Special volume celebrating the 70th birthday of Professor Max Kelly.
- [19] Y. Kato, *Finite syntomic topology and algebraic cobordism of non-unital algebras*, Preprint, available at arXiv:2203.05331 (2022).
- [20] J. Lurie, *Higher topos theory*, Annals of Mathematics studies, vol. 170, Princeton University Press, 2009.
- [21] ———, *Higher algebra*, Preprint, available at [www.math.ias.edu/lurie](http://www.math.ias.edu/lurie) (2017).
- [22] M. Levine and F. Morel, *Algebraic cobordism*, Springer Monographs in Mathematics, Springer, Berlin, 2007. MR2286826
- [23] M. Levine and R. Pandharipande, *Algebraic cobordism revisited*, Invent. Math. **176** (2009), no. 1, 63–130, DOI 10.1007/s00222-008-0160-8. MR2485880
- [24] Saunders Mac Lane and Ieke Moerdijk, *Sheaves in geometry and logic*, Universitext, Springer-Verlag, New York, 1994. A first introduction to topos theory; Corrected reprint of the 1992 edition. MR1300636
- [25] F. Morel and V. Voevodsky,  *$\mathbf{A}^1$ -homotopy theory of schemes*, Institut des Hautes Études Scientifiques. Publications Mathématiques **90** (1999), 45–143 (2001).
- [26] J. Nuiten, *Localizing  $\infty$ -categories with hypercovers*, Preprint, arXiv:1612.03800 (2016).
- [27] M. Spitzweck and P. A. Østvær, *Motivic twisted K-theory*, Preprint, arXiv:1008.4915 (2010).
- [28] I. Panin, K. Pimenov, and O. Röndigs, *On the relation of Voevodsky’s algebraic cobordism to Quillen’s K-theory*, Invent. Math. **175** (2009), no. 2, 435–451. MR2470112
- [29] D. Quillen, *Homotopical algebra*, Lecture Notes in Mathematics, No. 43, Springer-Verlag, Berlin-New York, 1967.
- [30] ———, *Higher algebraic K-theory. I*, Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972), Springer, Berlin, 1973, pp. 85–147. Lecture Notes in Math., Vol. 341.
- [31] ———, *Module theory over non-unital rings*, available at: <https://ncatlab.org/nlab/files/QuillenModulesOverRngs.pdf> (1996).
- [32] P. Scholze, *Perfectoid spaces*, Publ. math., I.H.E.S. **116** (2012), 245–313, DOI 10.1112/S0010437X06002624.
- [33] Stefan Schwede, *Symmetric spectra*, 2012.
- [34] The Stacks project authors, *The Stacks project*, available at: <https://stacks.math.columbia.edu> (2022).
- [35] R. W. Thomason and T. Trobaugh, *Higher algebraic K-theory of schemes and of derived categories*, The Grothendieck Festschrift, Vol. III, Progr. Math., vol. 88, Birkhäuser Boston, Boston, MA, 1990, pp. 247–435. MR1106918 (92f:19001)
- [36] V. Voevodsky, *Homology of schemes*, Selecta Math. (N.S.) **2** (1996), no. 1, 111–153. MR1403354
- [37] ———,  *$\mathbf{A}^1$ -homotopy theory*, Documenta Mathematica **Extra Vol. I** (1998), 579–604 (electronic).
- [38] F. Waldhausen, *Algebraic K-theory of spaces*, Algebraic and geometric topology (New Brunswick, N.J., 1983), Lecture Notes in Math., vol. 1126, Springer, Berlin, 1985, pp. 318–419.
- [39] C. Weibel, *Homotopy algebraic K-theory*, Contemp. Math., vol. 83, 1989, pp. 461–488.

NATIONAL INSTITUTE OF TECHNOLOGY, UBE COLLEGE, 2-14-1, TOKIWADAI, UBE, YAMAGUCHI, JAPAN 755-8555.  
*Email address:* [ykato@ube-k.ac.jp](mailto:ykato@ube-k.ac.jp)