# A unified approach to embeddings of a line in 3-space

Drew Lewis\*

March 17, 2022

#### Abstract

While the general question of whether every closed embedding of an affine line in affine 3-space can be rectified remains open, there have been several partial results proved by several different means. We provide a new approach, namely constructing (strongly) residual coordinates, that allows us to give new proofs of all known partial results, and in particular generalize the results of Bhatwadekar-Roy and Kuroda on embeddings of the form  $(t^n, t^m, t^l + t)$ .

### 1 Introduction

Let k be an algebraically closed field of characteristic zero throughout. The embedding problem, one of the central problems in affine algebraic geometry, asks whether every closed embedding  $\mathbb{A}^m \hookrightarrow \mathbb{A}^n$  is equivalent to the standard embedding (such embeddings are called *rectifiable*). In general, this is a very difficult problem that has led to several fruitful avenues of research. If  $n \ge 2m + 2$ , then every embedding is rectifiable due to a general result of Srinivas [15]. Two remaining cases have attracted the most attention: the case where n = m + 1, and the case where m = 1. In the former case, it is often referred to as the Abhyankar-Sathaye embedding conjecture, and can be reformulated as asking whether every hyperplane in  $\mathbb{A}^n$  is a coordinate; as this is not the focus of this paper, we simply suggest the papers [13, 12, 18, 7] to the interested reader, and note that it remains open for  $n \ge 3$ .

In this paper, we are interested in the m = 1 case, i.e. embeddings  $\mathbb{A} \hookrightarrow \mathbb{A}^n$ . Note that the result of Srinivas [15] provides an affirmative answer in the case  $n \ge 4$ . In the case n = 2, an affirmative answer was provided by Abyhankar and Moh [2], and independently, Suzuki [16]. Thus, in the m = 1 case, the only remaining open case is embeddings  $\mathbb{A} \hookrightarrow \mathbb{A}^3$ , so we direct our attention there.

Abhyankar [1] conjectured (among other things) that the family of embeddings  $(t^n, t^{n+1}, t^{n+2} + t)$  are not rectifiable. Craighero [5, 6] showed for n = 3

<sup>\*</sup>Department of Mathematics and Statistics, University of South Alabama. Email address: drewlewis@southalabama.edu

and n = 4, these embeddings are, in fact, rectifiable; but it remains an open question for  $n \ge 5$ . Later, Bhatwadekar and Roy [3] considered the more general family of embeddings  $(t^n, t^m, t^l + t)$ , and showed that many of these are rectifiable. More recently, Kuroda [9] showed that another subset of this family are rectifiable. These results are stated precisely in Section 3 below.

All of these papers use different approaches. In [5], Craighero explicitly writes down the rectifying automorphism, while in [6] he explicitly computes a polynomial, and as this polynomial is linear in a variable, appeals to a well-known result of Sathaye [13] to see it can be extended to a rectifying automorphism. Bhatwadekar and Roy [3] make use of the Dedekind conductor, while Kuroda [9] constructs rectifying automorphisms from exponentials of locally nilpotent derivations similar to the Nagata map.

In this paper, we take a single approach that we apply to recover all of these results and extend some of them. To broadly explain our approach, first note that an embedding  $\mathbb{A} \to \mathbb{A}^3$  corresponds to a surjection of polynomial rings  $k[x, y, z] \to k[t]$ . It is well-known that an embedding is rectifiable if and only if there is a coordinate of k[x, y, z] that maps to t. Our approach is to construct (strongly) residual coordinates that map to t, and then use a criterion of [11] to see that they are coordinates (see Corollary 4 below).

In the next section, we review some standard definitions and results from the field, and then describe our general approach. Then in Section 3, we precisely state and give new proofs of the results of [5, 6, 3, 9]; in particular, we also prove generalizations of the results of Bhatwadekar and Roy [3] and Kuroda [9] in Theorems 8 and 12, respectively.

### 2 Preliminaries

Let us begin by quickly recalling some fairly standard definitions; more detailed explanations can be found in [17] or [19].

- We use  $GA_n(k)$  to denote the general automorphism group  $Aut_k \mathbb{A}^n$ . This group is anti-isomorphic to the automorphism group of the polynomial ring  $k^{[n]}$ .
- $EA_n(k)$  denotes the subgroup of  $GA_n(k)$  generated by elementary automorphisms, i.e. those fixing n-1 variables.
- A polynomial  $f \in k^{[n]}$  is called a *coordinate* (or *variable* by some authors) if there exist  $f_2, \ldots, f_n$  such that  $(f, f_2, \ldots, f_n) \in GA_n(k)$ .
- An embedding  $\phi : \mathbb{A} \hookrightarrow \mathbb{A}^3$  is called *rectifiable* if there exists  $\theta \in GA_3(k)$  such that  $\theta \phi = (t, 0, 0)$ .
- For any morphism  $\phi : \mathbb{A}^m \to \mathbb{A}^n$ , we use  $\phi^*$  to denote the corresponding map  $\phi^* : k^{[n]} \to k^{[m]}$ .

The following is a well-known algebraic characterization of rectifiable embeddings.

**Lemma 1.** Let  $\phi : \mathbb{A} \hookrightarrow \mathbb{A}^n$  be an embedding. If  $f \in k^{[n]}$  is a coordinate with  $\phi^*(f) = t$ , then  $\phi$  is rectifiable.

Proof. Let  $\theta = (f, f_2, \dots, f_n) \in \operatorname{GA}_n(k)$ . Since  $\phi^*(f) = t$ , we have  $\theta \phi = (t, g_2(t), \dots, g_n(t))$  for some polynomials  $g_2, \dots, g_n \in k[t]$ . Let  $\psi = (x_1, x_2 - g_2(x_1), \dots, x_n - g_n(x_1)) \in \operatorname{GA}_n(k)$ . Then  $\psi \theta \phi = (t, 0, \dots, 0)$ , so  $\phi$  is rectifiable.

This lemma underlies our approach in this paper: given an embedding, we aim to construct a coordinate that the embedding maps to t. To construct coordinates, we rely up on the idea of (strongly) residual coordinates.

### 2.1 Strongly residual coordinates and associated embeddings

**Definition 1.** A polynomial  $f \in k[x]^{[n]}$  is called a *residual coordinate* if it becomes a coordinate modulo x - c for each  $c \in k$ . A polynomial is called a *strongly residual coordinate* if it becomes a coordinate modulo x, and also a coordinate over  $k[x, x^{-1}]$ .

Note that strongly residual coordinates are residual coordinates. A special case of the Dolgachev-Weisfeiler conjecture asserts that all residual coordinates are coordinates. This remains open for  $n \ge 3$ , with the Vénéreau polynomial  $y+x(xz-y(yu+z^2))$  being a famous example of a (strongly) residual coordinate whose status as a coordinate remains unresolved<sup>1</sup>.

To illustrate our methods, let us consider the well known construction of the Nagata automorphism. Let  $\alpha = (x, y, z - \frac{y^2}{x}) \in GA_2(k[x, x^{-1}])$ , and set  $\beta = (x, y + x^2z, z) \in GA_2(k[x])$ . It is straightforward to compute that

$$\alpha^{-1}\beta\alpha = (x, y + x(xz - y^2), z + 2y(xz - y^2) + x(xz - y^2)^2).$$

This resulting element of  $GA_2(k[x])$  is known as the Nagata automorphism; after many years, it was famously shown [14] to not be generated by elementary and affine elements of  $GA_3(k)$ . Since it is a conjugation of an elementary automorphism (over  $k[x, x^{-1}]$ ), it can also be obtained as an exponential of a locally nilpotent derivation, namely  $(xz - y^2) \left(x \frac{\partial}{\partial y} + 2y \frac{\partial}{\partial z}\right)$ ; Kuroda exploits this fact in [9]. However, if we instead consider  $\alpha_0 = (x, y, z - \frac{y^2}{x^2})$  and  $\beta_0 = (x, y + x^3 z, z)$ , then  $\alpha_0^{-1}\beta_0\alpha_0 \notin GA_2(k[x])$ ; however, letting  $\gamma = (x, y, z + \frac{2y^3}{x})$ , then a straightforward computation verifies that  $\gamma \alpha_0^{-1} \beta_0 \alpha_0 \in GA_2(k[x])$ . Note that since we are no longer simply conjugating, it is not an exponential of a locally nilpotent derivation.

We now generalize this approach; the following is a special case of Theorem 13 of [11], but we also present a short direct proof here.

 $<sup>^{1}</sup>$ In fact, the Vénéreau polynomial is also a hyperplane, meaning it is a potential counterexample to the Abhyankar-Sathaye conjecture as well. We refer the reader to [8, 10, 4] for further reading on the Vénéreau polynomial.

**Theorem 2.** Let  $\alpha, \beta \in EA_2(k[x, x^{-1}])$  be of the form  $\alpha = (x, y, z + x^{-l}P(y))$  for some  $P \in k[x][y]$  and  $\beta = (x, y + x^a(x^l z)^b, z)$  for some  $a, b \in \mathbb{N}$ . Then there exists  $R \in k[x][y]$  such that setting  $\gamma = (x, y, z + x^{-l}R(y))$ , then  $\gamma\beta\alpha \in GA_2(k[x])$ . Moreover, if  $P(y) \in (y^c)$  for some  $c \in \mathbb{N}$ , then  $R(y) \in (y^c)$ .

*Proof.* We begin by proving the following:

**Claim 3.** Let  $F \in k[x]^{[2]}$  and  $G \in k[x]^{[1]}$ , and suppose

$$\phi = (x, y + x^a (x^l z + P(y))^b, z + x^a z F(x^l z, y) + x^{-l} G(y)) \in GA_2(k[x, x^{-1}]).$$

Then setting  $\theta = (x, y, z - x^{-l}G(y)) \in EA_2(k[x, x^{-1}])$ , there exist  $\tilde{F} \in k[x]^{[2]}$ and  $\tilde{G} \in k[x]^{[1]}$  such that

$$\theta\phi = (x, y + x^a (x^l z + P(y))^b, z + x^a z \tilde{F}(x^l z, y) + x^{-l+a} \tilde{G}(y)) \in GA_2(k[x, x^{-1}]).$$

Moreover,  $\tilde{G}(y) \in (y^c)k[x, y]$  as well.

*Proof.* This follows from direct computation. Note that  $\theta$  fixes x and y, so we need only compute

$$(\theta\phi)^*(z) = z + x^a z F(x^l z, y) + x^{-l} \left( G(y) - G(y + x^a (x^l z + P(y))^b) \right).$$

The key observation is to apply Taylor's formula to obtain

$$G(y) - G(y + x^{a}(x^{l}z + P(y))^{b}) = -\sum_{i=1}^{n} \frac{1}{i!} G^{(i)}(y) (x^{a}(x^{l}z + P(y))^{b})^{i}.$$

Applying a binomial expansion to  $(x^l z + P(y))^{bi}$ , we see that there exists  $\tilde{F}_0 \in k[x]^{[2]}$  such that

$$G(y) - G(y + x^{a}(x^{l}z + P(y))^{b}) = -\sum_{i=1}^{l} \frac{1}{i!} G^{(i)}(y) x^{ai}(P(y))^{bi} + x^{l+a} z \tilde{F}_{0}(x^{l}z, y).$$

Then we observe that setting

$$\tilde{G}(y) = -\sum_{i=1}^{l} \frac{1}{i!} G^{(i)}(y) x^{a(i-1)} (P(y))^{bi} \text{ and } \tilde{F}(x^l z, y) = F(x^l z, y) + \tilde{F}_0(x^l z, y),$$

we obtain the desired result.

Now, beginning with  $\beta \alpha$  and repeatedly applying the claim, we may inductively produce a sequence  $\theta_0, \ldots, \theta_r$  with  $\theta_i = (x, y, z - x^{-l+ai}G_i(y))$  (with each  $G_i(y) \in (y^c)$ ) such that

$$\theta_r \cdots \theta_0 \beta \alpha = (x, y + x^a (x^l z + P(y))^b, z + x^a z \tilde{F}_r(x^l z, y) + x^{-l+a(r+1)} \tilde{G}_r(y))$$

for some  $\tilde{F}_r(x^l z, y)$ ,  $\tilde{G}_r(x^l z, y) \in k[x][x^l z, y]$ . Let r be minimal so that  $a(r+1) \geq l$ , and set  $\gamma = \theta_r \cdots \theta_0$ . Then letting  $R(y) = -\sum_{i=0}^r x^{ai} G_i(y) \in (y^c)$ , we see  $\gamma = (x, y, z + x^{-l}R(y))$  with  $\gamma\beta\alpha \in \text{GA}_2(k[x, x^{-1}])$  and  $\gamma\beta\alpha \in (k[x, y, z])^3$ . It is well known (e.g. Proposition 1.1.7 in [17]) that this implies that  $\gamma\beta\alpha \in \text{GA}_2(k[x])$  as required.

**Corollary 4.** Let  $\phi : \mathbb{A} \hookrightarrow \mathbb{A}^3$  be an embedding. Suppose  $\alpha, \beta \in EA_2(k[x, x^{-1}])$  are of the form  $\alpha = (x, y, z + x^{-l}P(y))$  and  $\beta = (x, y + xQ(x^lz), z)$  for some  $P, Q \in k[x]^{[1]}$ , and that  $\phi^* \alpha^* \beta^*(y) = t$ . Then  $\phi$  is rectifiable.

*Proof.* Apply the previous theorem to produce  $\gamma = (x, y, z + x^{-l}R(y))$  such that  $\gamma\beta\alpha \in GA_2(k[x])$ . Let  $f = (\gamma\beta\alpha)^*(y)$ , so f is a coordinate. Since  $\gamma^*(y) = y$ , we have  $t = \phi^*\alpha^*\beta^*(y) = \phi^*\alpha^*\beta^*\gamma^*(y) = \phi^*(\gamma\beta\alpha)^*(y) = \phi^*(f)$ . Lemma 1 then completes the proof.

Remark 1. An elementary example of how this corollary can be used can be found in Theorem 6.

### 2.2 A useful combinatorial lemma

Here we prove a combinatorial lemma that we use in the proof of Lemma 10 below; the reader may prefer to skip the proof for now, and proceed to Section 3. This is used only in the proof of Lemma 10 below. For an integer m, let us

write  $\delta_m = \begin{cases} 1 & m \text{ is even} \\ 0 & m \text{ is odd} \end{cases}$ .

**Lemma 5.** Let  $m \in \mathbb{N}$ . Then there exist  $\alpha_0, \ldots, \alpha_{\lfloor \frac{m}{2} \rfloor}, \beta \in k$  such that, in k[s],

$$\sum_{i=0}^{\left\lfloor \frac{m}{2} \right\rfloor} \alpha_i s^i (1+s)^{m-2i} = 1 + \beta \delta_m s^{\frac{m}{2}} + s^m.$$

*Proof.* First, suppose m = 2k + 1 is odd. We induct on k, with the k = 0 case being trivial. For k > 0, we first expand  $(1 + s)^{2k+1}$  noting the symmetry of coefficients:

$$(1+s)^{2k+1} = 1+s^{2k+1} + \sum_{j=1}^{k} \binom{2k+1}{j} (s^j + s^{2k+1-j}) = 1+s^{2k+1} + \sum_{j=1}^{k} \binom{2k+1}{j} s^j (1+s^{2k+1-2j})$$

For  $1 \leq j \leq k$ , by the induction hypothesis we choose  $\alpha_{j,0}, \ldots, \alpha_{k-j}$  such that

$$1 + s^{2k+1-2j} = \sum_{i=0}^{k-j} \alpha_{j,i} s^i (1+s)^{2k+1-2j-2i}.$$

Then

$$(1+s)^{2k+1} = 1 + s^{2k+1} + \sum_{j=1}^{k} \binom{2k+1}{j} s^j \sum_{i=0}^{k-j} \alpha_{j,i} s^i (1+s)^{2k+1-2j-2i}$$

Letting  $\alpha_{0,i} = -\binom{2k+1}{i}$  for  $0 \le i \le k$ , we then have

$$1 + s^{2k+1} = -\sum_{j=0}^{k} \sum_{i=0}^{k-j} {\binom{2k+1}{j}} \alpha_{j,i} s^{i+j} (1+s)^{2k+1-2(i+j)}$$
$$= \sum_{r=0}^{k} s^{r} (1+s)^{2k+1-2r} \left( -\sum_{j=0}^{r} {\binom{2k+1}{j}} \alpha_{j,r-j} \right)$$

Now, suppose m = 2k is even. We again induct on k, with k = 0 being trivial. For k > 0 we proceed similarly by expanding  $(1 + s)^{2k+1}$ , noting however that there is an odd number of terms now.

$$(1+s)^{2k} = 1 + \binom{2k}{k} s^k + s^{2k} + \sum_{j=1}^{k-1} \binom{2k}{j} (s^j + s^{2k-j}) = 1 + \binom{2k}{k} s^k + s^{2k} + \sum_{j=1}^{k-1} \binom{2k}{j} s^j (1+s^{2k-2j}) = 1 + \binom{2k}{k} s^k + s^{2k} + \sum_{j=1}^{k-1} \binom{2k}{j} s^j (1+s^{2k-2j}) = 1 + \binom{2k}{k} s^k + \frac{2k}{j} s^{2k-2j} + \frac{2k}{j} s^{2k-$$

For  $1 \leq j \leq k-1$ , by the induction hypothesis we choose  $\alpha_{j,0}, \ldots, \alpha_{k-j}, \beta_j$  such that

$$1 + s^{2k-2j} = \beta_j s^{k-j} + \sum_{i=0}^{k-j} \alpha_{j,i} s^i (1+s)^{2k-2j-2i}.$$

Then

$$(1+s)^{2k} = 1 + \binom{2k}{k}s^k + s^{2k} + \sum_{j=1}^{k-1}\binom{2k}{j}s^j \left(\beta_j s^{k-j} + \sum_{i=0}^{k-j}\alpha_{j,i}s^i(1+s)^{2k-2j-2i}\right)$$
$$= 1 + s^k \left(\binom{2k}{k} + \sum_{j=1}^{k-1}\binom{2k}{j}\beta_j\right) + s^{2k} + \sum_{j=1}^{k-1}\sum_{i=0}^{k-j}\binom{2k}{j}\alpha_{j,i}s^{i+j}(1+s)^{2k-2(i+j)}$$

Now, set  $\alpha_{0,i} = -\binom{2k}{i}$  for  $0 \le i \le k$ , and set  $\beta = \binom{2k}{k} + \sum_{j=1}^{k-1} \binom{2k}{j} \beta_j$ . Then

$$1 + \beta s^{k} + s^{2k} = -\sum_{j=0}^{k} \sum_{i=0}^{k-j} \binom{2k}{j} \alpha_{j,i} s^{i+j} (1+s)^{2k-2(i+j)}$$
$$= \sum_{r=0}^{k} s^{r} (1+s)^{2k-2r} \left( -\sum_{j=0}^{r} \binom{2k}{j} \alpha_{j,r-j} \right).$$

## 3 A unified approach to known results on embedding

In this section we summarize all results known to us on embeddings  $\mathbb{A} \hookrightarrow \mathbb{A}^3$ , and show how they can all be proved via Corollary 4. For convenience, we will

adopt the notation  $X = \phi^*(x)$ ,  $Y = \phi^*(y)$ , and  $Z = \phi^*(z)$ . So for example, when considering the Abhyankar embeddings  $(t^n, t^{n+1}, t^{n+2} + t)$ , we will write  $X = t^n$ ,  $Y = t^{n+1}$ , and  $Z = t^{n+2} + t$ . As an elementary example, consider the n = 2 Abhyankar embedding  $(t^2, t^3, t^4 + t)$ . It is straightforward to see that  $Z - X^2 = t$ , and the polynomial  $z - x^2$  is a coordinate (of the elementary automorphism  $(x, y, z - x^2)$ ), so by Lemma 1 the embedding  $(t^2, t^3, t^4 + t)$  is rectifiable.

### 3.1 Craighero's results

The n = 3 and n = 4 cases of Abhyankar's conjecture were resolved by Craighero [5, 6]; we are able to give short proofs here.

**Theorem 6** (Craighero). The embedding  $\phi = (t^3, t^4, t^5 + t)$  is rectifiable.

*Proof.* We begin by computing

$$Y - \frac{Z^2}{X^2} + 2 = t^4 - \frac{t^{10} + 2t^6 + t^2}{t^6} + 2 = -t^{-4}.$$

Let  $\alpha = \left(x, y - \frac{z^2}{x^2} + 2, z\right)$  and  $\beta = (x, y, z + x(x^2y))$ . Then we compute

$$\phi^* \alpha^* \beta^* (z) = \phi^* \alpha^* (z + x(x^2 y))$$
  
=  $\phi^* \left( z + x^3 \left( y - \frac{z^2}{x^2} + 2 \right) \right)$   
=  $Z + X^3 \left( Y - \frac{Z^2}{X^2} + 2 \right)$   
=  $(t^5 + t) + t^9 (-t^{-4})$   
=  $t.$ 

Thus, by Corollary 4, the embedding is rectifiable.

**Theorem 7** (Craighero). The embedding  $\phi = (t^4, t^5, t^6 + t)$  is rectifiable.

*Proof.* We first observe that  $Z^2 - X^3 = 2t^7 + t^2$ . Next, letting  $a, b \in k$  be arbitrary for the moment, we compute

$$Y + a\frac{Z^3}{X^2} + bXZ - 3a = t^5 + \frac{a\left(t^{18} + 3t^{13} + 3t^8 + t^3\right)}{t^8} + b(t^{10} + t^5) - 3a$$
$$= (a+b)t^{10} + (1+3a+b)t^5 + at^{-5}.$$

Setting  $a = -\frac{1}{2}$  and  $b = \frac{1}{2}$  causes these first two coefficients to vanish, so we see

$$Y - \frac{1}{2}\frac{Z^3}{X^2} + \frac{1}{2}XZ + \frac{3}{2} = -\frac{1}{2}t^{-5}.$$

Thus,

$$\left(Y - \frac{1}{2}\frac{Z^3}{X^2} + \frac{1}{2}XZ + \frac{3}{2}\right) + \frac{1}{4}\frac{Z^2 - X^3}{X^3} = -\frac{1}{2}t^{-5} + \frac{1}{4}\frac{2t^7 + t^2}{t^{12}} = \frac{1}{4}t^{-10}.$$

Now, let

$$\alpha = \left(x, y - \frac{1}{2}\frac{z^3}{x^2} + \frac{1}{2}xz + \frac{3}{2} + \frac{1}{4}\frac{z^2 - x^3}{x^3}, z\right)$$
  
$$\beta = (x, y, z + x(-4x^3y))$$

Then we see

$$\begin{split} \phi^* \alpha^* \beta^*(z) &= Z - 4X^4 \left( Y - \frac{1}{2} \frac{Z^3}{X^2} + \frac{1}{2} XZ + \frac{3}{2} + \frac{1}{4} \frac{Z^2 - X^3}{X^3} \right) \\ &= t^6 + t - 4(t^{16}) \left( -\frac{1}{4} t^{-10} \right) \\ &= t. \end{split}$$

Thus  $\phi$  is rectifiable by Corollary 4.

### 3.2 Generalizing Bhatwadekar-Roy's results

The next results were due to Bhatwadekar and Roy [3], who studied embeddings of the form  $(t^n, t^{an+1}, t + t^l)$  for positive integers a, n and l > n. This more general class includes the Abhyankar embeddings mentioned above. Interstingly, in positive characteristic they were able to show that all such embeddings are rectifiable; in the characteristic zero case we are interested in, they obtained two positive results: when  $l \equiv -1 \pmod{n}$ , and when n = 4. We first generalize the former in Theorem 8, and then provide a new proof of the latter in Theorem 11.

**Theorem 8.** Let  $m, n \in \mathbb{N}$  be coprime positive integers. Write  $m \equiv c \pmod{n}$ for some 0 < c < n, and set d = n - c. Write  $\lambda_1 c = \mu_1 n - 1$  and  $\lambda_2 d = \mu_2 n - 1$ for  $\lambda_1, \lambda_2 \in \mathbb{N}$  and minimal  $\mu_1, \mu_2 \in \mathbb{N}$ . If  $b > \min\{\mu_1, \mu_2\}$ , then the embedding  $\phi = (t^n, t^m, t + t^{bn-1})$  is rectifiable.

Remark 2. When c = 1, we obtain one of the results of Bhatwadekar and Roy in [3].

*Example* 1. Let n = 5 and m = 7, so c = 2, in which case  $\mu_1 = 1$ . Then we see the embedding  $(t^5, t^7, t + t^9)$  is rectifiable.

More generally, we have

**Corollary 9.** Let  $n \in \mathbb{N}$  be odd. Then for any b > 1, the embedding  $(t^n, t^{n+2}, t+t^{bn-1})$  is rectifiable.

*Proof.* Note that c = 2, so since n is odd, n - 1 is even and thus  $\mu_1 = 1$ .  $\Box$ 

In order to prove Theorem 8, we first prove the following lemma.

**Lemma 10.** Suppose  $\phi = (t^n, t^m, t^{bn-1})$  is an embedding. Then for any  $r \in \mathbb{N}$  there exits  $p \in k[x, z]$  such that  $\phi^*(p) = t^r + t^{(bn-1)r}$ .

*Proof.* We begin by applying Lemma 5. Recall that  $\delta_r = \begin{cases} 1 & r \text{ is even} \\ 0 & r \text{ is odd} \end{cases}$ ; then by Lemma 5, there exist  $\alpha_0, \ldots, \alpha_{\lfloor \frac{r}{2} \rfloor}, \beta \in k$  such that, for any  $s \in k[t]$ ,

$$\sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \alpha_i s^i (1+s)^{r-2i} = 1 + \beta \delta_r s^{\frac{r}{2}} + s^r.$$
(1)

Now we set  $p = \sum_{i=0}^{\lfloor \frac{r}{2} \rfloor} \alpha_i x^{bi} z^{r-2i} - \beta \delta_r x^{b\frac{r}{2}} \in k[x, z]$ , and we compute

$$\phi^{*}(p) = \sum_{i=0}^{\left\lfloor \frac{r}{2} \right\rfloor} \alpha_{i}(t^{n})^{bi}(t+t^{bn-1})^{r-2i} - \beta \delta_{r}(t^{n})^{b\frac{r}{2}}$$
$$= \sum_{i=0}^{\left\lfloor \frac{r}{2} \right\rfloor} \alpha_{i}t^{nbi+r-2i}(1+t^{bn-2})^{r-2i} - \beta \delta_{r}t^{bn\frac{r}{2}}$$
$$= t^{r} \left( \sum_{i=0}^{\left\lfloor \frac{r}{2} \right\rfloor} \alpha_{i}\left(t^{bn-2}\right)^{i}(1+t^{bn-2})^{r-2i} - \beta \delta_{r}\left(t^{bn-2}\right)^{\frac{r}{2}} \right)$$

Substituting  $s = t^{bn-2}$  into (1) above, we obtain

$$\phi^*(p) = t^r \left(1 + \left(t^{bn-2}\right)^r\right) = t^r + t^{(bn-1)r}.$$

Proof of Theorem 8. Write m = an + c for some  $a \in \mathbb{N}$ . We begin by supposing  $a \geq bd - 1$ , so a + 1 = bd + h for some  $h \in \mathbb{N}$ . In this case, we can write m = (a + 1)n - d = (bn - 1)d + hn. By Lemma 10, there exists  $p \in k[x, z]$  such that  $\phi^*(p) = t^d + t^{(bn-1)d}$ . Then, setting  $\theta = (x, -y + x^h p(x, z), z) \in GA_3(k)$ , we see that  $\theta\phi = (t^n, t^{hn+d}, t + t^{bn-1})$ , and this is rectifiable if and only if  $\phi$  is rectifiable. Note that since  $b \geq 2$ , hn + d < (bn - 1)d + hn = m. Thus, by repeating this process we may continue until we have a < bd - 1.

So now it suffices to assume a < bd - 1. We proceed in two cases; first, we suppose  $b > \mu_2$ . By Lemma 10, there exists  $p \in k[x, z]$  such that  $\phi^*(p) = t^d + t^{(bn-1)d}$ . We then set  $\alpha = \left(x, y - \frac{p(x,z)}{x^{bd-a-1}}, z\right)$  and  $\beta = \left(x, y, z + x^{b-\mu_2} \left(x^{bd-a-1}y\right)^{\lambda_2}\right)$ ,

and compute

$$\begin{split} \phi^* \alpha^* \beta^*(z) &= Z + X^{b-\mu_2} (X^{bd-a-1}Y - p(X,Z))^{\lambda_2} \\ &= t + t^{bn-1} + (t^n)^{b-\mu_2} \left( (t^n)^{bd-a-1} t^{an+c} - \left( t^d + t^{(bn-1)d} \right) \right)^{\lambda_2} \\ &= t + t^{bn-1} + t^{n(b-\mu_2)} \left( t^{bdn-n+c} - t^d - t^{(bn-1)d} \right)^{\lambda_2} \\ &= t + t^{bn-1} + t^{nb-\lambda_2d-1} (-t^{\lambda_2d}) \\ &= t. \end{split}$$

Then by Corollary 4,  $\phi$  is rectifiable.

Now, we must consider the case  $b > \mu_1$ . Then by Lemma 10, find  $q \in k[x, z]$  such that  $\phi^*(q) = t^c + t^{(bn-1)c}$ , and as above,  $p \in k[x, z]$  such that  $\phi^*(p) = t^d + t^{(bn-1)d}$ .

Now set 
$$\alpha = \left(x, y - \frac{p(x,z)}{x^{bd-a-1}} + \frac{q}{x^{bc+bd-a-2}}, z\right)$$
 and  $\beta = \left(x, y, z - x^{b-\mu_1} \left(x^{bc+bd-a-2}y\right)^{\lambda_1}\right)$ , and compute

$$\begin{split} \phi^* \alpha^* \beta^*(z) &= Z - X^{b-\mu_1} (X^{bc+bd-a-2}Y - X^{bc-1}p(X,Z) + q(X,Z))^{\lambda_1} \\ &= t + t^{bn-1} - (t^n)^{b-\mu_1} \left( (t^n)^{bc+bd-a-2}t^{an+c} - (t^n)^{bc-1} \left( t^d + t^{(bn-1)d} \right) + \left( t^c + t^{(bn-1)c} \right) \right)^{\lambda_1} \\ &= t + t^{bn-1} - t^{n(b-\mu_1)} \left( t^{bcn+bdn-2n+c} - t^{bcn-n+d} - t^{bnc-n+(bn-1)d} + t^c + t^{(bn-1)c} \right)^{\lambda_1} \\ &= t + t^{bn-1} - t^{nb-\lambda_1c-1} (t^{\lambda_1c}) \\ &= t. \end{split}$$

Then again by Corollary 4,  $\phi$  is rectifiable.

We next give a new proof of the other result of [3].

**Theorem 11** (Bhatwadekar-Roy).  $\phi = (t^4, t^{4a+1}, t^m + t)$  is rectifiable for any  $a, m \in \mathbb{N}$ .

*Proof.* We divide the proof into four cases, based on the residue of m modulo 4. Three cases are relatively straightforward, while the case  $m \equiv 2 \pmod{4}$  generalizes our proof of Theorem 7 above.

**Case 1:**  $m \equiv 1 \pmod{4}$ . Write m = 4k + 1 for some  $k \in \mathbb{N}$ . Without loss of generality, we may assume a < k; for if  $a \ge k$ , we may apply the map  $(x, -y + x^{a-k}z, z)$  to produce the embedding  $(t^4, t^{4(a-k)+1}, t^m + t)$ , and repeat until a < k. But in this case, observe that  $Z - X^{k-a}Y = t$ , and  $z - x^{k-a}y$  is a coordinate.

**Case 2:**  $m \equiv 2 \pmod{4}$ . Write m = 4k + 2 for some  $k \in \mathbb{N}$ . This is the hardest case, but proceeds in the same way as our proof of Theorem 7. First, we observe that

$$Z^2 - X^{2k+1} = 2t^{4k+3} + t^2.$$

$$Y + c\frac{Z^3}{X^{2k+1-a}} + dX^a Z - 3cX^{a-k} = (c+d)t^{4a+4k+2} + (1+3c+d)t^{4a+1} + c\frac{1}{t^{8k-4a+1}}$$

Setting  $c = -\frac{1}{2}$  and  $d = \frac{1}{2}$  causes these first two coefficients to vanish, so we see

$$Y - \frac{1}{2}\frac{Z^3}{X^{2k+1-a}} + \frac{1}{2}X^a Z + \frac{3}{2}X^{a-k} = -\frac{1}{2}\frac{1}{t^{8k-4a+1}}$$

Thus

$$Y - \frac{1}{2}\frac{Z^3}{X^{2k+1-a}} + \frac{1}{2}X^a Z + \frac{3}{2}X^{a-k} + \frac{1}{4}\frac{Z^2 - X^{2k+1}}{X^{3k-a+1}} = \frac{1}{4}\frac{1}{t^{12k-4a+2}}$$

Now, let

$$\alpha = \left(x, \left(y - \frac{1}{2}\frac{z^3}{x^{2k+1-a}} + \frac{1}{2}x^a z + \frac{3}{2}x^{a-k}\right) + \frac{1}{4}\frac{z^2 - x^3}{x^{3k-a+1}}, z\right)$$

Note that if a > 3k + 1, then  $\alpha \in EA_3(k)$ . And in this case, we have 12k-4a+2 = 4(3k-a+1)-2 < 0; so then we have  $\alpha \phi = (t^4, \frac{1}{4}t^{4a-(12k+2)}, t^m+t)$ , with 4a - (12k+2) < 4a + 1. Now, if  $k \ge a - 3k - 1$ , we can set  $\beta = (x, y, z - 4x^{4k-a+1}y) \in EA_3(k)$ , and compute  $\phi^* \alpha^* \beta^*(z) = t$ , and as  $\beta \alpha \in GA_3(k)$ ,  $\phi$  is rectifiable by Lemma 1. If instead k < a - 3k - 1, we set  $\beta = (x, -4y + x^{a-4k-1}z, z) \in GA_3(k)$ , and compute that  $\beta \alpha \phi = (t^4, t^{4(a-4k-1)+1}, t^m + t)$ .

This process can be repeated, so we may now proceed assuming without loss of generality that  $a \leq 3k + 1$ . We construct  $\alpha$  as above (but now,  $\alpha \in \text{EA}_2(k[x, x^{-1}]))$ , and further define

$$\beta = \left(x, y, z + x^k \left(-4x^{3k-a+1}y\right)\right)$$

Then we see

$$\begin{split} \phi^* \alpha^* \beta^*(z) &= Z - 4X^{4k-a+1} \left( Y - \frac{1}{2} \frac{Z^3}{X^{2k+1-a}} + \frac{1}{2} X^a Z + \frac{3}{2} X^{a-k} + \frac{1}{4} \frac{Z^2 - X^{2k+1}}{X^{3k-a+1}} \right) \\ &= t^{4k+2} + t - 4(t^{16k-4a+4}) \left( \frac{1}{4} t^{-12k+4a-2} \right) \\ &= t. \end{split}$$

Thus  $\phi$  is rectifiable by Corollary 4.

**Case 3:**  $m \equiv 3 \pmod{4}$ . Write m = 4k + 3 for some  $k \in \mathbb{N}$ . Without loss of generality, we may assume  $a \leq 3k + 2$ ; for if a > 3k + 2, we set  $\alpha = (x, -y + x^{a-2-3k}z^3 - 3x^{a-2k-1}z, z) \in GA_3(k)$  and compute

$$\begin{split} \phi^* \alpha^*(y) &= -Y + X^{a-2-3k} Z^3 - 3X^{a-2k-1} Z \\ &= -t^{4a+1} + t^{4(a-2-3k)} (t^{4k+3} + t)^3 - 3t^{4a-8k-4} (t^{4k+3} + t) \\ &= -t^{4a+1} + t^{4a-8-12k} \left( t^{12k+9} + 3t^{8k+7} + 3t^{4k+5} + t^3 \right) - 3t^{4a-4k-1} - 3t^{4a-8k-3} \\ &= t^{4a-12k-5} \end{split}$$

This process can be repeated until  $a \leq 3k + 2$ .

Now, assuming  $a \leq 3k + 2$ , we set  $\alpha = \left(x, y - \frac{z^3}{x^{3k+2-a}} - 3x^{a-2k-1}z, z\right)$  and  $\beta = (x, y, z + x^k(-x^{3k+2-a}y))$ , and compute  $\phi^* \alpha^* \beta^*(z) = Z - X^K \left(Z^3 - X^{3k-a+2}Y - 3X^{k+1}Z\right)$   $= t + t^{4k+3} - t^{4k} \left((t^{4k+3} + t)^3 - t^{4(3k-a+2)}(t^{4a+1}) - 3t^{4k+4}(t^{4k+3} + t)\right)$   $= t + t^{4k+3} - t^{4k} \left(t^{12k+9} + 3t^{8k+7} + 3t^{4k+5} + t^3 - t^{12k+9} - 3t^{8k+7} - 3t^{4k+5}\right)$   $= t + t^{4k+3} - t^{4k} \left(t^3\right)$ = t.

Thus  $\phi$  is rectifiable by Corollary 4.

**Case 4:**  $m \equiv 0 \pmod{4}$ . In this case, m = 4k for some  $k \in \mathbb{N}$ , so  $Z - X^k = t$ , and  $z - x^k$  is a coordinate.

### 3.3 Generalizing Kuroda's result

This section is devoted to proving the following theorem.

**Theorem 12.** Let  $n, a, c, l, s \in \mathbb{N}$  such that cl < a. Then the embedding  $(t^n, t^{an+c}, t+t^{(an+c)s-ln})$  is rectifiable.

Remark 3. Kuroda [9] proved the special case of c = 1 and a = 2l + m for some nonnegative integer m (in which case the assumption cl < a is satisfied automatically).

Proof of Theorem 12. Let  $\alpha = \left(x, y, z - \frac{y^s}{x^l}\right)$  and  $\beta = \left(x, y - x^{a-cl}(x^l z)^c, z\right)$ We compute

$$\begin{split} \phi^* \alpha^* \beta^* (y) &= Y - X^{a-cl} \left( X^l Z - Y^s \right)^c \\ &= t^{an+c} - t^{n(a-cl)} \left( t^{ln} (t + t^{(an+c)s-ln}) - t^{(an+c)s} \right)^c \\ &= t^{an+c} - t^{n(a-cl)} \left( t^{ln+1} \right)^c \\ &= t^{an+c} - t^{n(a-cl)+c(ln+1)} \\ &= 0 \end{split}$$

It is also easy to see that  $\phi^* \alpha^* \beta^*(z) = t$ , so we have  $\beta \alpha \phi = (t^n, 0, t)$ . Now, from Theorem 2, we can produce  $\gamma = (x, y, z + \frac{R(y)}{x^l})$  such that  $\gamma \beta \alpha \in GA_2(k[x])$ , and  $R(y) \in (y^s)k[x, y]$ , so that  $(\gamma \beta \alpha \phi)^*(z) = (\beta \alpha \phi)^*(z + \frac{R(y)}{x^l}) = t$ . Then letting  $f = (\gamma \beta \alpha)^*(z)$ , we have f is a coordinate with  $\phi^*(f) = t$ , so  $\phi$  is rectifiable by Lemma 1.

Letting c = 1, a = 3, l = 2, we obtain the following:

**Corollary 13.** For any  $n \in \mathbb{N}$ , the embedding  $(t^n, t^{3n+1}, t^{4n+2} + t)$  is rectifiable.

Letting c = 2, a = 5, l = 2, we obtain the following:

**Corollary 14.** For any  $n \in \mathbb{N}$ , the embedding  $(t^n, t^{5n+2}, t^{8n+4} + t)$  is rectifiable.

### References

- S. S. Abhyankar, On the semigroup of a meromorphic curve (Part 1), Intl. Symp. on Algebraic Geometry, Kyoto (1977), pp. 249-414.
- [2] S. Abhyankar and T.T. Moh, Embeddings of the line in the plane, J. Reine Angew. Math. 276 (1975), 148–166.
- [3] S. Bhatwadekar and A. Roy, Some results on embedding of a line in 3-space, J. Algebra 142 (1991), 101–109.
- [4] J. Blanc and P.-M. Poloni, Bivariables and Vénéreau polynomials, arXiv:2004.10739.
- [5] P. C. Craighero, About Abhyankar's conjectures on space lines, Rendiconti del Seminario Matematico della Università di Padova 74 (1985), 115–122.
- [6] P. C. Craighero, A remark on Abhyankar's space lines, Rendiconti del Seminario Matematico della Università di Padova, 80 (1988), 87–93.
- [7] S. Kaliman, Polynomials with general C<sup>2</sup>-fibers are variables, Pac. J. Math., 203 (2002), 161−190.
- [8] S. Kaliman, S. Vénéreau, M. Zaidenberg, Simple birational extensions of the polynomial algebra C<sup>3</sup>, Trans. Amer. Math. Soc., **356** (2) (2004), 509– 555.
- S. Kuroda, The Nagata type polynomial automorphisms and rectifiable space lines, Comm. Alg., 42 (2014), 4451–4455.
- [10] D. Lewis, Vénéreau-type polynomials as potential counterexamples, J. Pure and Applied Algebra, 217 (5) (2013), 946–957.
- [11] D. Lewis, Strongly residual coordinates over A[x], Automorphisms in Birational and Affine Geometry (2014), 407–430.
- P. Russell and A. Sathaye, On finding and cancelling variables in k[X, Y, Z], J. Algebra, 57 (1979), 151–166.
- [13] A. Sathaye, On linear planes, Proc. Amer. Math. Soc. 56 (1976), 1–7.
- [14] I. Shestakov and U. Umirbaev, The tame and the wild automorphisms of polynomial rings in three variables, J. Amer. Math. Soc. 17 (2004), 197– 227.
- [15] V. Srinivas, On the embedding dimension of an affine variety, Math. Ann., 289 (1991), 125–132.
- [16] M. Suzuki, Propiétés topologiques des polynomes de deux variables complexes, et automorphismes algéarique de l'espace C<sup>2</sup>, J. Math. Soc. Japan, 26 (1974), 241−257.

- [17] A. van den Essen, Polynomial automorphisms and the Jacobian conjecture (2000). Progress in Mathematics, 190. Birkhäuser Verlag, Basel.
- [18] D. Wright, Cancellation of variables of the form  $bT^n a$ , J. Algebra, **52** (1978), 94–100.
- [19] D. Wright, *Polynomial automorphism groups*, Polynomial automorphisms and related topics (2007), Publishing House for Science and Technology, Hanoi, 1–19.