INFINITELY MANY FAMILIES OF SASAKI-EINSTEIN METRICS ON SPHERES

YUCHEN LIU, TARO SANO, AND LUCA TASIN

ABSTRACT. We show that there exist infinitely many families of Sasaki-Einstein metrics on every odd-dimensional standard sphere of dimension at least 5. We also show that the same result is true for all odd-dimensional exotic spheres that bound parallelizable manifolds.

1. Introduction

An odd-dimensional compact Riemannian manifold (M,g) is called Sasakian if its (punctured) metric cone $(M\times\mathbb{R}^{>0},\bar{g})$ is Kähler, where $\bar{g}=r^2g+dr^2$ is the conical metric on $M\times\mathbb{R}^{>0}$ with r the coordinate of $\mathbb{R}^{>0}$. Denote by $X:=M\times\mathbb{R}^{\geq 0}/M\times\{0\}$ the cone over M. Moreover, a Sasakian manifold (M,g) is called Sasaki-Einstein if g has constant Ricci curvature, or equivalently, if \bar{g} is a Ricci-flat Kähler cone metric on X. The Reeb vector field ξ on a Sasakian manifold (M,g) is given by $\xi:=J(r\partial_r)$ where J is the integrable complex structure on X.

It is an important problem in differential geometry to find Sasaki-Einstein metrics on odd dimensional manifolds. As a starting case, the case of spheres and exotic spheres attracted much attention. In [BGK05], Boyer, Galicki, and Kollár constructed a large (finite) number of inequivalent families of Sasaki-Einstein metrics on all homotopy spheres in dimension 4n + 1, or in dimension 7. Using computer programming, [BGKT05] generalizes this result to dimensions 11 and 15 for homotopy spheres that bound parallelizable manifolds. In [CS19], Collins and Székelyhidi found infinitely many families of inequivalent Sasaki-Einstein metrics on S^5 . In [PW21], Park and Won completely determined which simply connected rational homology 5-spheres admit Sasaki-Einstein metrics.

In this paper, we show the existence of infinitely many families of Sasaki-Einstein metrics on a large class of homotopy spheres in all (odd) dimensions. Our main result is the following.

Theorem 1.1. Let $n \geq 3$ be an integer. Then any (2n-1)-dimensional homotopy sphere which bounds parallelizable manifolds admits infinitely many inequivalent families of Sasaki-Einstein metrics.

As a result, Theorem 1.1 implies the following two conjectures of Collins-Székelyhidi and Boyer-Galicki-Kollár.

Corollary 1.2 ([CS19, Conjecture 9.2]). There are infinitely many families of Sasaki-Einstein metrics on every odd-dimensional standard sphere of dimension at least 5.

Corollary 1.3 ([BGK05, Conjecture 4]). All odd-dimensional homotopy spheres that bound parallelizable manifolds admit Sasaki-Einstein metrics.

The strategy to prove Theorem 1.1 is based on a method developed in [BG01, BGK05, BGKT05, CS19], that we now outline (see the book [BG08] for a detailed exposition on Sasakian geometry).

Consider the link $L(\mathbf{a}) := Y(\mathbf{a}) \cap S^{2n+1}(1)$ of a Brieskorn-Pham singularity

$$Y(\mathbf{a}) := (z_0^{a_0} + z_1^{a_1} + \dots + z_n^{a_n} = 0) \subset \mathbb{C}^{n+1}$$

where $n \geq 3$ and $\mathbf{a} = (a_0, \dots, a_n) \in \mathbb{Z}_{>1}^{n+1}$. Then $L(\mathbf{a})$ is a smooth (2n-1)-manifold which bounds a parallelizable manifold. Under certain numerical conditions on \mathbf{a} , the link $L(\mathbf{a})$ is a homotopy (2n-1)-sphere and so it is an element of the group bP_{2n} of equivalence classes of oriented homotopy (2n-1)-spheres that bound parallelizable manifolds. See Section 2.

Let $d := \text{lcm}(a_0, \dots, a_n)$ and $d_i := \frac{d}{a_i}$ for $i = 0, \dots, n$. Let $\xi := \sum_{i=0}^n d_i z_i \partial_{z_i}$ be a Reeb vector field on $Y(\mathbf{a})$ of weight (d_0, \dots, d_n) which generates the corresponding \mathbb{C}^* -action on $Y(\mathbf{a})$. The following result shows that the Lichnerowicz obstruction [GMSY07, (3.23)] is not only necessary but also sufficient for the existence of Sasaki-Einstein metrics on $L(\mathbf{a})$.

Theorem 1.4. Notation as above. Assume that $a_0 \le a_1 \le \cdots \le a_n$. Then $(Y(\mathbf{a}), \xi)$ is a K-polystable (resp. K-semistable) Fano cone singularity if and only if

(1)
$$1 < \sum_{i=0}^{n} \frac{1}{a_i} < (resp. \le) 1 + \frac{n}{a_n}.$$

Moreover, the link $L(\mathbf{a})$ admits a Sasaki-Einstein metric if the above strict inequalities hold.

We note that [GK07, Theorem 1] studied the case where a_0, \ldots, a_n are pairwise coprime and [ST24, Proposition 6.1] studied the case where the corresponding weighted hypersurface is well-formed.

The diffeomorphism type of $L(\mathbf{a})$ is determined by the Arf invariant if n is odd and by the signature $\tau(\mathbf{a})$ of the Milnor fiber if n is even. In Section 3, we show that for appropriate choices of \mathbf{a} we get each possible value in bP_{2n} infinitely many times, proving in this way Theorem 1.1. The hard case is when n is even: the problem of computing $\tau(\mathbf{a})$ is reduced to showing that a certain function counting integral points of a polytope is a quasi-polynomial. We then use this to generalize a classical example of Brieskorn.

In Section 4 we look at the moduli space of Sasaki-Einstein metrics proving the following.

Corollary 1.5. For $m \geq 3$, let $\Sigma^{4m-1} \in bP_{4m}$ be a homotopy sphere. Then the dimensions of the families of Sasaki-Einstein metrics on Σ^{4m-1} and S^{4m-3} are both unbounded.

This gives a negative answer to a question in [BG06, Section 6].

In Subsection 4.1, we show that the contact structures induced by our construction on each homotopy sphere Σ^{4m-1} $(m \geq 3)$ belong to infinitely many families.

Acknowledgements. We would like to thank Charles Boyer, János Kollár, Yoshihiko Matsumoto, Yuji Odaka, Andrea Petracci, Chenyang Xu, Zhouli Xu, and Ruixiang Zhang for helpful discussions and comments. We would like to thank the referees for helpful comments.

The first named author is partially supported by NSF Grant DMS-2148266 (formerly DMS-2001317). The second named author is partially supported by JSPS KAKENHI Grant Numbers JP17H06127, JP19K14509. The third named author is a member of the GNSAGA group of INdAM and is supported by the project PRIN 2020KKWT53.

2. Brieskorn-Pham singularities

2.1. Topological background on Brieskorn manifolds. We fix some notation. Let $n \geq 3$ be an integer and let $\mathbf{a} = (a_0, \dots, a_n) \in \mathbb{Z}_{>0}^{n+1}$ such that each $a_i > 1$. Let

$$Y(\mathbf{a}) := (z_0^{a_0} + z_1^{a_1} + \dots + z_n^{a_n} = 0) \subset \mathbb{C}^{n+1}$$

and its link $L(\mathbf{a}) := Y(\mathbf{a}) \cap S^{2n+1}(1)$ which is called a *Brieskorn manifold*. We know that $L(\mathbf{a})$ is a smooth (2n-1)-manifold [Mil68, Corollary 2.9]. Let $M(\mathbf{a})$ be the Milnor fiber of the polynomial $f(z) = z_0^{a_0} + z_1^{a_1} + \cdots + z_n^{a_n}$, that is, the fiber of the locally trivial fibration $\varphi \colon S^{2n+1}(1) \setminus L(\mathbf{a}) \to S^1(1)$ defined by $\varphi(z) = \frac{f(z)}{|f(z)|}$. Since the Milnor fiber is pararellizable ([Mil68, Theorem 5.1], [Dim92, (1.23) Proposition]), we know that $L(\mathbf{a})$ always bounds a parallelizable manifold.

Definition 2.1. To each **a** as above, one associates a graph $G(\mathbf{a})$ with n+1 vertices labeled by a_0, \ldots, a_n . Two vertices a_i and a_j are connected in $G(\mathbf{a})$ if and only if $\gcd(a_i, a_j) > 1$. Let $G(\mathbf{a})_{\text{ev}}$ be the connected component of $G(\mathbf{a})$ that contains all even integers. Note that $G(\mathbf{a})_{\text{ev}}$ may contain odd integers as well.

The following theorem gives a characterization of when $L(\mathbf{a})$ is a homotopy (2n-1)-sphere, that is, a smooth (2n-1)-manifold homotopy equivalent to the (2n-1)-sphere. (By the topological generalized Poincaré conjecture, we can replace "homotopy equivalent" in the above with "homeomorphic".)

Theorem 2.2. [Bri66, Satz 1 (ii)] Let $n \geq 3$, \mathbf{a} , $L(\mathbf{a})$ and $G(\mathbf{a})$ be as above. The link $L(\mathbf{a})$ is homeomorphic to the (2n-1)-sphere if and only if either of the following holds.

- (1) $G(\mathbf{a})$ contains at least two isolated points, or
- (2) $G(\mathbf{a})$ contains a unique odd isolated point, and $G(\mathbf{a})_{\text{ev}}$ has an odd number of vertices with $\gcd(a_i, a_j) = 2$ for any distinct $a_i, a_j \in G(\mathbf{a})_{\text{ev}}$.

Remark 2.3. When $L(\mathbf{a})$ is homeomorphic to the (2n-1)-sphere, we call $L(\mathbf{a})$ a *Brieskorn sphere*. When $n \geq 3$, every diffeomorphism type of a homotopy (2n-1)-sphere which bounds parallelizable manifolds can be realized by a Brieskorn sphere $L(\mathbf{a})$ for some \mathbf{a} ([Bri66], [BG08, Theorem 9.4.8]).

2.2. **Diffeomorphism types of Brieskorn spheres.** We continue to use the notations in 2.1.

Let Θ_{2n-1} be the Kervaire-Milnor groups consisting of equivalence classes of oriented homotopy (2n-1)-spheres that are equivalent under oriented homotopism (see e.g. [KM63]). We are interested in the subgroup $bP_{2n} < \Theta_{2n-1}$ consisting of equivalence classes of oriented homotopy (2n-1)-spheres that bound parallelizable manifolds.

If n=2m $(m \geq 2)$ is even, we have the following theorem on the diffeomorphism type of $L(\mathbf{a})$.

Theorem 2.4. Let n = 2m $(m \ge 2)$ and $\mathbf{a} \in \mathbb{Z}_{>0}^{n+1}$ satisfying the condition in Theorem 2.2 so that $L(\mathbf{a}) \in bP_{4m}$.

(i) ([KM63, p.526], [Lev85, Corollary 3.20]) The group bP_{4m} is cyclic of order

$$|bP_{4m}| = a_m 2^{2m-2} (2^{2m-1} - 1) \cdot \text{numerator} \left(\frac{4B_m}{m}\right),$$

where $a_m = \begin{cases} 1 & (m \in 2\mathbb{Z}) \\ 2 & (m \notin 2\mathbb{Z}) \end{cases}$ and B_m is the m-th Bernoulli number.

(ii) ([KM63, Theorem 7.5], [Bri66, Satz 3]) The class of $L(\mathbf{a})$ in bP_{4m} is $\frac{\tau(\mathbf{a})}{8}$ with a suitable choice of the generator of bP_{4m} , where $\tau(\mathbf{a})$ is is the signature of the intersection pairing on H^n of the Milnor fiber $M(\mathbf{a})$. Moreover, $\tau(\mathbf{a})$ can be described as follows.

$$\tau(\mathbf{a}) = \# \left\{ \mathbf{x} = (x_0, \dots, x_{2m}) \in \mathbb{Z}^{2m+1} \mid 0 < x_i < a_i \text{ and } 0 < \sum_{i=0}^{2m} \frac{x_i}{a_i} < 1 \mod 2 \right\}$$
$$- \# \left\{ \mathbf{x} = (x_0, \dots, x_{2m}) \in \mathbb{Z}^{2m+1} \mid 0 < x_i < a_i \text{ and } 1 < \sum_{i=0}^{2m} \frac{x_i}{a_i} < 2 \mod 2 \right\}.$$

If n = 2m+1 ($m \ge 1$) is odd and **a** satisfies the condition in Theorem 2.2, then $L(\mathbf{a})$ is a homotopy (4m+1)-sphere and belongs to bP_{4m+2} . The group bP_{4m+2} has order 1 or 2, and it is trivial for $4m+1 \in \{1,5,13,29,61\}$, and order 2 for $4m+1 \notin \{1,5,13,29,61,125\}$ thanks to the almost completely solved Kervaire invariant problem [Bro69, HHR16].

The Kervaire sphere is a homotopy (4m+1)-sphere whose Arf invariant equals 1. If $bP_{4m+2} \cong \mathbb{Z}_2$ then the Kervaire sphere is the only non-trivial

element. Brieskorn also proved the following (see e.g. [BG08, Theorem 9.4.3]).

Theorem 2.5. Let n = 2m+1 $(m \ge 1)$ and \mathbf{a} as above so that $L(\mathbf{a}) \in bP_{4m+2}$. Then $L(\mathbf{a})$ is the Kervaire sphere if and only if all of the following holds:

- the condition (2) of Theorem 2.2 holds.
- the one isolated point, say a_0 , satisfies $a_0 \equiv \pm 3 \mod 8$.
- $G(\mathbf{a}) = G(\mathbf{a})_{\text{ev}} \cup \{a_0\}.$

We remark that our method can only produce Sasaki-Einstein metrics on homotopy spheres in bP_{2n} . The quotient group Θ_{2n-1}/bP_{2n} is the hard part of the group of homotopy spheres, which almost completely reduces to the computation of stable homotopy groups of spheres. For a recent account on this topic, see [IWX23, Section 1.4].

2.3. **K-stability.** We refer the definition of Fano cone singularities and K-polystability/semistability to [LWX21, Section 2.2].

Throughout the paper, we always work with quasi-regular Fano cone singularities (Y, ξ) , i.e. Y is an affine variety with klt singularities with the vertex $y \in Y$, and ξ is a rational Reeb vector generating a good \mathbb{C}^* -action on Y [LWX21, Definition 2.12].

In this case, we have the quotient morphism $Y \setminus \{y\} \to X$ by the \mathbb{C}^* -action and the branch divisor Δ_X so that (X, Δ_X) is a log Fano pair [LWX21, Definition 2.13].

We shall use the following.

Theorem 2.6. Let (Y,ξ) be a Fano cone singularity and (X,Δ_X) be its \mathbb{C}^* -quotient as above. Then the following are equivalent.

- (i) (Y, ξ) is K-polystable (resp. K-semistable).
- (ii) (X, Δ_X) is K-polystable (resp. K-semistable).

(See e.g. [LWX21, Definition 2.23, 2.28] for the definitions of K-polystability/K-semistability of Fano cone singularities and log Fano pairs.)

Assume that $Y \setminus \{y\}$ is smooth. Then the K-polystability of (Y, ξ) is equivalent to the following.

(iii) Y admits a Ricci-flat Kähler cone metric and the link L_Y of Y admits a Sasaki-Einstein metric with Reeb vector field ξ .

Proof. (i) \Leftrightarrow (ii): This follows from the correspondence of special test configurations of (Y, ξ) and (X, Δ_X) as in [Li21, 2.5] and the equality of generalized Futaki invariants [LWX21, Lemma 2.30] (cf. [Li21, Theorem 2.9]).

Now assume that $Y \setminus \{y\}$ is smooth.

Then the last equivalence follows from [CS19, Theorem 1.1] and [BG08, Corollary 11.1.8], for example (See also [AJL21, Li21]). \Box

We prove Theorem 1.4 by modifying the argument in [ST24, Proposition 6.1].

Proof of Theorem 1.4. Let $(Y(\mathbf{a}), \xi)$ be the Fano cone singularity induced by the Reeb vector field ξ on $Y(\mathbf{a})$. Let (X, Δ_X) be the log Fano pair induced by the \mathbb{C}^* -action on $Y(\mathbf{a})$ as above. Then $(Y(\mathbf{a}), \xi)$ is K-polystable (K-semistable) if and only if (X, Δ_X) is K-polystable (K-semistable) by Theorem 2.6. Hence we shall show that (X, Δ_X) is K-polystable (K-semistable) if and only if the inequality (1) holds.

We see that

$$\Delta_X = \sum_{j=0}^{n} \left(1 - \frac{1}{g_j} \right) H_j$$

for $g_j := \gcd(d_0, \dots, \hat{d}_j, \dots, d_n)$ and $H_j := (z_j = 0) \cap X$. Note that X is isomorphic as a variety to the quasi-smooth weighted hypersurface

$$(y_0^{a_0'} + \dots + y_n^{a_n'} = 0) \subset \mathbb{P}\left(\frac{d_0 g_0}{g}, \dots, \frac{d_n g_n}{g}\right) =: \mathbb{P}',$$

where $a_i' := a_i/g_i$ and $g := g_0 \cdots g_n$.

Let $\Pi: \mathbb{P}' \to \mathbb{P}^n$ be the Galois cover determined by

$$[y_0:\cdots:y_n]\mapsto [y_0^{a_0'}:\cdots:y_n^{a_n'}].$$

Let $L := (w_0 + \dots + w_n = 0) \subset \mathbb{P}^n$ and $\pi := \Pi|_X \colon X \to H$ be the induced Galois cover. Then we have the ramification formula

$$K_X = \pi^* \left(K_L + \sum_{i=0}^n \left(1 - \frac{1}{a_i'} \right) L_i \right)$$

for $L_i := L \cap (w_i = 0)$ for i = 0, ..., n. By these and $a'_j H_j = \pi^*(L_j)$, we see that

(2)
$$K_X + \Delta_X = \pi^* \left(K_L + \sum_{i=0}^n \left(1 - \frac{1}{a_i} \right) L_i \right).$$

By this, we see that the log Fano condition is equivalent to $1 < \sum_{i=0}^{n} \frac{1}{a_i}$. We also see that the K-polystability (resp. K-semistability) of (X, Δ_X) is equivalent to that of $(L, \sum_{i=0}^{n} (1 - \frac{1}{a_i})L_i)$) by the above equality (2) and [LZ22, Theorem 1.2], [Zhu21, Corollary 4.13]. Hence the statement is reduced to the following.

Claim. $\left(L, \sum_{i=0}^{n} (1 - \frac{1}{a_i})L_i\right)$ is K-polystable (resp. K-semistable) if and only if

$$1 + \frac{n}{a_n} - \sum_{i=0}^n \frac{1}{a_i} > \text{ (resp. } \ge) 0.$$

Proof of Claim. By [Fuj21, Corollary 1.6], the K-polystability (resp. K-semistability) is equivalent to

(3)
$$k \sum_{i=0}^{n} \left(1 - \frac{1}{a_i} \right) - n \sum_{j=1}^{k} \left(1 - \frac{1}{a_{i_j}} \right) > \text{ (resp. } \ge) 0$$

for all $1 \le k \le n-1$ and $0 \le i_1 < \cdots < i_k \le n$. We obtain the claim since we have

L.H.S. of (3) =
$$k - k \sum_{i=0}^{n} \frac{1}{a_i} + n \sum_{j=1}^{k} \frac{1}{a_{i_j}} = k \left(1 - \sum_{i=0}^{n} \frac{1}{a_i} + \frac{n}{k} \sum_{j=1}^{k} \frac{1}{a_{i_j}} \right)$$

$$\geq k \left(1 - \sum_{i=0}^{n} \frac{1}{a_i} + \frac{n}{a_n} \right).$$

Finally, the existence of Sasaki-Einstein metrics on $L(\mathbf{a})$ follows from Theorem 2.6.

Remark 2.7. Note that, for $Y(\mathbf{a})$ as in Theorem 1.4, we have

$$I_{\mathbf{a}} := I_{Y(\mathbf{a})} := \sum_{i=0}^{n} \frac{d}{a_i} - d = d \left(\sum_{i=0}^{n} \frac{1}{a_i} - 1 \right).$$

Hence the condition in Theorem 1.4 is equivalent to $0 < I_{Y_a} < nd_n$ as in [ST24, Proposition 6.1].

We shall use the following propositions to distinguish the Sasaki-Einstein metrics on the links of weighted homogeneous hypersurface singularities.

Proposition 2.8. ([BGK05, Corollary 22]) For i = 1, 2, let

$$Y_i := (F_i = 0) \subset \mathbb{C}^m$$

be an affine hypersurface defined by a weighted homogeneous polynomial F_i with weights $\mathbf{w}_i = (w_{i1}, \dots, w_{im}) \in \mathbb{Z}_{>0}^m$. Let

$$X_i := (Y_i \setminus \{0\})/\mathbb{C}^*$$

be the quotient by the \mathbb{C}^* -action on Y_i of weights \mathbf{w}_i . Assume that X_i has a Kähler-Einstein orbifold metric and no orbifold holomorphic contact structure. Let L_i be the corresponding link with the Einstein metric g_i .

Then (L_1, g_1) is isometric to (L_2, g_2) if and only if the following conditions holds.

- (i) $\mathbf{w}_1 = \mathbf{w}_2$ holds up to permutations.
- (ii) There exists an automorphism $\tau \in \operatorname{Aut}(\mathbb{C}^m, \mathbf{w}_1)$ as in [BGK05, 15] such that $\tau(Y_1) = Y_2$ or its complex conjugate \overline{Y}_2 .

Proposition 2.9. Let $0 \in Y = (F = 0) \subset \mathbb{C}^{n+1}$ be an isolated hypersurface singularity defined by a weighted homogeneous polynomial F of degree d for the \mathbb{C}^* -action of weights $\mathbf{w} = (w_0, \dots, w_n) \in \mathbb{Z}_{>0}^{n+1}$. Let $X := (Y \setminus \{0\}) / \mathbb{C}^*$ be the orbifold defined by the \mathbb{C}^* -action.

Then X admits no orbifold holomorphic contact structure if one of the following holds:

- (i) n is odd;
- (ii) n = 2m is even and $m \nmid I_Y := \sum_{i=0}^n w_i d$.

Proof. This follows from [BGK05, §16].

2.4. Normalized volumes.

Definition 2.10 ([Li18]). Let $y \in Y$ be a complex n-dimensional klt singularity. Let v be a real valuation of $\mathbb{C}(Y)$ centered at y, i.e. $v : \mathbb{C}(Y)^{\times} \to \mathbb{R}$ is a valuation satisfying $v(\mathbb{C}^{\times}) = 0$ and $v(\mathfrak{m}_y) > 0$. We define the normalized volume of v as

$$\widehat{\text{vol}}(v) := A_Y(v)^n \cdot \text{vol}(v),$$

where $A_Y(v)$ is the log discrepancy of v according to [JM12, BdFFU15], and vol(v) is the volume of v according to [ELS03]. We define the local volume of the singularity $y \in Y$ as

$$\widehat{\text{vol}}(y, Y) := \min_{v} \widehat{\text{vol}}(v),$$

where v runs over all real valuations of $\mathbb{C}(Y)$ centered at y. The existence of a vol-minimizing valuation was proved by Blum [Blu18].

Li's motivation of introducing the normalized volume was to study the local K-stability theory for klt singularities. We refer the readers to the surveys [LLX20, Zhu23] for more backgrounds and discussions. Here we will focus on the case of K-polystable Fano cones.

Theorem 2.11. Let (Y,ξ) be a K-polystable Fano cone singularity with $y \in Y$ the cone vertex. Then $\widehat{\text{vol}}(y,Y) = \widehat{\text{vol}}(\text{wt}_{\xi})$, where wt_{ξ} is the valuation induced by the Reeb vector field ξ (see e.g. [LWX21, (2.6)]).

If moreover $Y \setminus \{y\}$ is smooth, then

$$\widehat{\operatorname{vol}}(y,Y) = n^n \cdot \frac{\operatorname{Vol}(L_Y, g_{L_Y})}{\operatorname{Vol}(S^{2n-1}, g_{\operatorname{st}})}.$$

Here g_{L_Y} is the Sasaki-Einstein metric on the link L_Y of Y whose existence follows from Theorem 2.6, g_{st} is the standard metric on the round sphere S^{2n-1} , and Vol denotes the Riemannian volume.

Proof. The first statement follows from [LX18, Theorem 1.3]. The second statement follows from the well-known equality $\widehat{\text{vol}}(\text{wt}_{\xi}) = n^n \cdot \frac{\text{Vol}(L_Y, g_{L_Y})}{\text{Vol}(S^{2n-1}, g_{\text{st}})}$, see e.g. [GMSY07, MSY08, DS17, HS17, LX18].

Example 2.12. Let $(Y(\mathbf{a}), \xi)$ be a K-polystable Brieskorn-Pham singularity with $a_0 \leq \cdots \leq a_n$. Then by [GMSY07, (3.18)] and Theorem 2.11 we have

$$\widehat{\operatorname{vol}}(0, Y(\mathbf{a})) = \frac{d(d_0 + \dots + d_n - d)^n}{d_0 \dots d_n} = \left(\sum_{i=0}^n \frac{1}{a_i} - 1\right)^n \cdot \prod_{i=0}^n a_i.$$

By Theorem 1.4, we have $0 < \sum_{i=0}^{n} \frac{1}{a_i} - 1 \le \frac{n}{a_n}$ which implies $a_0 \le n$ and

(4)
$$\widehat{\text{vol}}(0, Y(\mathbf{a})) \le n^n \prod_{i=0}^{n-1} \frac{a_i}{a_n} \le n^n \frac{a_0}{a_n} \le \frac{n^{n+1}}{a_n}.$$

Remark 2.13. Let \mathcal{M} be the set of Riemannian metrics on a given compact smooth manifold M of dimension n. For a Riemannian manifold (M, g), we have the total scalar curvature functional on \mathcal{M}

$$S(g) := \int_{M} s_g \operatorname{vol}_g,$$

where s_g is the scalar curvature and vol_g is the Riemannian volume form of g [Bes87, 4.16 Definition]. By [Bes87, 4.21 Theorem], we know that g is Einstein if it is a critical point of the functional $\frac{S(g)}{\operatorname{Vol}(g)^{\frac{n-2}{n}}}$, where $\operatorname{Vol}(g) := \int_M \operatorname{vol}_g$ is the Riemannian volume. Note that s_g is (the Einstein)

 $\operatorname{Vol}(g) := \int_M \operatorname{vol}_g$ is the Riemannian volume. Note that s_g is (the Einstein) constant if g is Einstein since $\operatorname{Ric}_g = s_g \cdot g$ in this case. By these, we see that $s_g \cdot \operatorname{Vol}(g)^{\frac{2}{n}}$ is constant on a continuous family of Einstein metrics. (See also [Bes87, 12.52 Corollary].)

Hence, in order to find an infinitely many families of Einstein metrics on a given manifold, it is enough to find a family of Einstein metrics with infinitely many values of $s_g \cdot \operatorname{Vol}(g)^{\frac{2}{n}}$. Note that, if (M,g) is Sasaki-Einstein, then $s_g = \dim M - 1$ (See e.g. [BG08, Lemma 11.1.5]). By these, Theorem 2.11 and Example 2.12, it is enough to exhibit K-polystable Fano cone singularities $(Y(\mathbf{a}), \xi)$ with infinitely many values of normalized volumes $\widehat{\operatorname{vol}}(0, Y(\mathbf{a}))$ to show Theorem 1.1 and Corollary 1.2. By the inequality (4), it suffices to find \mathbf{a} with arbitrarily large a_n .

3. Infinite sequence of K-polystable Fano cones

3.1. **Dimension** 4m+1. In this case, n=2m+1 with $m\geq 1$. By the prime number theorem, for every sufficiently large prime number p_n , there are at least n-2 distinct prime numbers p_2, \dots, p_{n-1} in the interval $(\frac{(n-2)}{2(n-1)}p_n, \frac{1}{2}p_n)$. Let

$$a_0 = a_1 = 2$$
, $a_i = 2p_i$ for $2 \le i \le n - 1$, and $a_n = p_n$.

We can check that

$$1 < \sum_{i=0}^{n} \frac{1}{a_i} = 1 + \sum_{i=2}^{n-1} \frac{1}{2p_i} + \frac{1}{p_n} < 1 + (n-2) \cdot \frac{(n-1)}{(n-2)p_n} + \frac{1}{p_n} = 1 + \frac{n}{a_n}.$$

Thus by Theorem 1.4 the Fano cone $(Y(\mathbf{a}), \xi(\mathbf{a}))$ is K-polystable, i.e. $L(\mathbf{a})$ admits a Sasaki-Einstein metric. Moreover, the condition of Theorem 2.2(2) holds for the link $L(\mathbf{a})$. Thus $L(\mathbf{a})$ is a standard sphere if $p_n \equiv \pm 1 \mod 8$, and is a Kervaire sphere if $p_n \equiv \pm 3 \mod 8$. By Dirichlet's theorem, there are infinitely many prime numbers p_n satisfying each of the congruency conditions. Thus, by Remark 2.13, we find infinite families of Sasaki-Einstein metrics on both the standard sphere and the Kervaire sphere of dimension 4m+1. Note that since n is odd, $(Y(\mathbf{a}) \setminus \{0\}) / \mathbb{C}^*$ admits no holomorphic orbifold contact structure and so these metrics are pairwise non-isometric by Proposition 2.8.

- 3.2. Dimension 4m-1. In this case, n=2m with $m \geq 2$.
- 3.2.1. Standard spheres. Let $p, q \geq 2$ be two positive integers. Let

$$\mathbf{a} = (a_0, \dots, a_{n-2}, a_{n-1}, a_n) = (n-1, \dots, n-1, p, q).$$

By Theorem 2.2, $L(\mathbf{a})$ is a homotopy (4m-1)-sphere if n-1, p, and q are pairwise coprime. Since $\sum_{i=0}^{n} \frac{1}{a_i} = 1 + \frac{1}{p} + \frac{1}{q}$, by Theorem 1.4 we know that $L(\mathbf{a})$ admits a Sasaki-Einstein metric if (n-1)p > q > n-1 and (n-1)q > p > n-1. Note that, by $d := \text{lcm}(a_0, \ldots, a_n) = (n-1)pq$ and Remark 2.7, we have

$$I_{\mathbf{a}} = (n-1)pq\left(\frac{1}{p} + \frac{1}{q}\right) = (n-1)(p+q).$$

For simplicity, denote by s := n - 1 = 2m - 1. First of all, we show that $\tau(\mathbf{a})$ is an integer combination of γ_i for $1 \le i \le s$, where

$$\gamma_j := \# \left\{ (x, y) \in \mathbb{Z}^2 \mid 0 < x < p, \ 0 < y < q, \ \text{ and } 0 < \frac{x}{p} + \frac{y}{q} < \frac{j}{s} \right\}.$$

Note that $\frac{x}{p} + \frac{y}{q}$ will never be of the form $\frac{j}{s}$ since s, p, q are coprime. Then we may group $\mathbf{x} \in \mathbb{Z}^{n+1}$ by the sum $\sigma = \sum_{i=0}^{n-2} x_i$, and for each value of σ the corresponding terms in $\tau(\mathbf{a})$ is an integer combination of $\gamma_j - \gamma_{j-1}$ for $1 \leq j \leq 2s$. Furthermore, the involution $(x, y) \mapsto (p - x, q - y)$ yields $\gamma_j - \gamma_{j-1} = \gamma_{2s-j+1} - \gamma_{2s-j}$, so we only need $1 \leq j \leq s$.

Next, we deduce a formula of γ_j for special values of p and q. Let k be a positive integer, and let p = sk + 1, $q = sp - 1 = s^2k + s - 1$. Clearly s, p, and q are pairwise coprime satisfying $s , so <math>L(\mathbf{a})$ is a Sasaki-Einstein homotopy sphere. Then $\frac{x}{p} + \frac{y}{q} < \frac{j}{s}$ implies that

$$x < \frac{jp}{s} = \frac{j(sk+1)}{s} = jk + \frac{j}{s}.$$

Hence $1 \le x \le jk$. For each such x, we have

$$y < q\left(\frac{j}{s} - \frac{x}{p}\right) = (sp - 1)\frac{pj - sx}{sp} = pj - sx - \frac{pj - sx}{sp}.$$

Hence $1 \le y \le pj - sx - 1$ as $\frac{pj - sx}{sp} \in (0, 1]$. Therefore,

$$\gamma_j = \sum_{x=1}^{jk} (pj - sx - 1) = \frac{jk}{2} (sjk + 2j - s - 2).$$

Thus γ_j is divisible by $\frac{k}{2}$ assuming k is even. As a result, $\frac{\tau(\mathbf{a})}{8}$ is divisible by $\frac{k}{16}$ assuming $16 \mid k$. Thus for every k that is a multiple of $16|bP_{4m}|$, we have that $\frac{\tau(\mathbf{a})}{8}$ is divisible by $|bP_{4m}|$, i.e. $L(\mathbf{a})$ is diffeomorphic to the

standard sphere. Thus we get an infinite sequence of p,q where $L(\mathbf{a})$ is a Sasaki-Einstein standard sphere. Since

$$\frac{s+1}{2} = \frac{n}{2} \nmid I_{Y_{\mathbf{a}}} = (n-1)(p+q) = s(sk+k+1) = s(k(s+1)+1),$$

 $(Y(\mathbf{a}) \setminus \{0\}) / \mathbb{C}^*$ admits no holomorphic contact structure by Proposition 2.9. This implies that the metrics we found are pairwise non-isometric by Proposition 2.8 and gives infinitely many inequivalent families by Remark 2.13.

3.2.2. Exotic spheres. For $k \in \mathbb{Z}_{>0}$, let $\mathbf{a}_2 := (2, \dots, 2, 3, 6k \pm 1) \in \mathbb{Z}^{n+1}$. Brieskorn ([Bri66], [BG08, Example 9.4.7]) found that $L(\mathbf{a}_2)$ satisfies

$$\frac{\tau(\mathbf{a}_2)}{8} = (-1)^m k.$$

So if we run over all $k \in \mathbb{Z}_{>0}$ we get all (4m-1)-spheres in bP_{4m} . Let us fix $k \in \{1, ..., |bP_{4m}|\}$. In the following, we study the sequence

$$a_0 = a_1 = 2$$
, $a_2 = \cdots = a_{n-2} = p$, $a_{n-1} = p + 1$, $a_n = p + l$,

where $p, l \geq 2$ are integers such that p is even, and p, p+1, and p+l are pairwise coprime. In other words, p is even, gcd(p, l) = 1 and gcd(p+1, l-1) = 1. Such p, p + 1, p + l can be realized by taking, for any $q \in \mathbb{Z}_{>0}$,

$$l := 6k - 3$$
(or $6k - 1$), $p := ql(l - 1) + 2$.

Let $\mathbf{a}_{p,l} := (a_0, \dots, a_n) = (2, 2, p, \dots, p, p+1, p+l) \in \mathbb{Z}^{n+1}$. For $p \gg 0$ (w.r.t. n and l), we have

$$1 < \sum_{i=0}^{n} \frac{1}{a_i} = 1 + \frac{n-3}{p} + \frac{1}{p+1} + \frac{1}{p+l} < 1 + \frac{n}{p+l} = 1 + \frac{n}{a_n},$$

thus see that $L(\mathbf{a}_{n,l})$ admits a Sasaki-Einstein metric by Theorem 1.4. Let us recall the definition of a quasi-polynomial.

Definition 3.1. A function $f: \mathbb{Z}_{>0} \to \mathbb{Z}$ is a quasi-polynomial if there exist $s \in \mathbb{Z}_{>0}$ and polynomials p_0, \ldots, p_{s-1} such that $f(n) = p_i(n)$ when $i \equiv n$

The main result of this section is the following.

Proposition 3.2. Under the above assumptions, $\tau(\mathbf{a}_{p,l})$ is a quasi-polynomial in p of period l(l-1).

Corollary 3.3. $\tau(\mathbf{a}_{p,l}) \equiv (-1)^m k \mod |bP_{4m}|$ for infinitely many p. In particular, any homotopy sphere in bP_{4m} admits infinitely many inequivalent families of Sasaki-Einstein metrics.

Proof of Corollary 3.3. Take $k \in \{1, \ldots, |bP_{4m}|\}$ and l = 6k-3 (or l = 6k-1). Then for any $p \equiv 2 \mod ml(l-1)|bP_{4m}|$, that is, p = ql(l-1) + 2, where $q = q'm|bP_{4m}|$ for some $q' \in \mathbb{Z}_{>0}$, we get that $\tau(\mathbf{a}_{p,l}) \equiv (-1)^m k \mod |bP_{4m}|$ by Proposition 3.2.

We shall confirm that the metrics on $L(\mathbf{a}_{p,l})$ are pairwise non-isometric. Note that

$$d = \text{lcm}(a_0, \dots, a_n) = p(p+1)(p+l)$$

and

$$I_{\mathbf{a}} = d\left(\sum_{i=0}^{n} \frac{1}{a_i} - 1\right) = (n-3)(p+1)(p+l) + p(p+l) + p(p+1)$$
$$= n(p+1)(p+l) - 3(p+1)(p+l) + p(2p+l+1)$$
$$\equiv -p^2 - 2pl - 2p - 3l \mod m.$$

If l=6k-3, then $I_{\bf a}\equiv -42k+13 \mod m$, while if l=6k-1, then $I_{\bf a}\equiv -42k-1$. Hence, for a fixed m and k, we can always choose l=6k-3 or l=6k-1 so that $m\nmid I_{\bf a}$. This implies that $X=(Y({\bf a})\setminus\{0\})/\mathbb{C}^*$ admits no holomorphic contact structure by Proposition 2.9 and so the metrics we found are pairwise non-isometric by Proposition 2.8. Remark 2.13 implies the existence of infinitely many families.

Proof of Theorem 1.1. Just combine Subsection 3.1 and Corollary 3.3. \square

The rest of the section is devoted to the proof of Proposition 3.2. We will repeatedly use the following lemma.

Lemma 3.4. Let f(x,y) be a quasi-polynomial in x and y of period l (that is, for $0 \le k, k' < l$, there exists a polynomial $f_{kk'}(x,y)$ such that $f(x,y) = f_{kk'}(x,y)$ when $x \equiv k$, $y \equiv k' \mod l$).

Then $\sum_{x=0}^{s} f(x,y)$ is a quasi-polynomial in s and y of period l.

Proof. In the proof, we may assume that $y \equiv k' \mod l$ for a fixed k' so that, if $x \equiv k \mod l$, then

$$f(x,y) = \sum_{i=0}^{m} \sum_{j=0}^{m} a_{ij}^{k} x^{i} y^{j}$$

for some m and constants a_{ij}^k for $i, j = 0, \ldots, m$ and $k = 0, \ldots, l-1$. Write

$$\sum_{x=0}^{s} f(x,y) = \sum_{k=0}^{l-1} \sum_{\substack{0 \le x \equiv k \\ \text{mod } l}}^{s} f(x,y) = \sum_{k=0}^{l-1} \sum_{\substack{0 \le x \equiv k \\ \text{mod } l}}^{s} \left(\sum_{i=0}^{m} \sum_{j=0}^{m} a_{ij}^{k} x^{i} y^{j} \right)$$

$$= \sum_{j=0}^{m} \sum_{i=0}^{m} \sum_{k=0}^{l-1} a_{ij}^{k} y^{j} \sum_{\substack{0 \le x \equiv k \\ \text{mod } l}}^{s} x^{i}.$$

It is hence enough to prove that

$$\sum_{\substack{0 \le x \equiv k \\ \text{mod } I}}^{s} x^{r}$$

is a quasi-polynomial in s of period l for any non-negative integer n. We have

$$\sum_{\substack{0 \leq x \equiv k \\ \text{mod } l}}^s x^n = \sum_{d=0}^{\left \lfloor \frac{s-k}{l} \right \rfloor} (dl+k)^n = \sum_{d=0}^{\left \lfloor \frac{s-k}{l} \right \rfloor} \sum_{i=0}^n \binom{n}{i} (dl)^i k^{n-i} = \sum_{i=0}^n \binom{n}{i} l^i k^{n-i} \sum_{d=0}^{\left \lfloor \frac{s-k}{l} \right \rfloor} d^i.$$

For any $i=0,\ldots,n$ the sum $\sum_{d=0}^{\left\lfloor\frac{s-k}{l}\right\rfloor}d^i$ is a polynomial in $\left\lfloor\frac{s-k}{l}\right\rfloor$ of degree i+1 by Faulhaber's formula (See e.g. [Bea96, (2.2)]) and $\left\lfloor\frac{s-k}{l}\right\rfloor$ is a quasi-polynomial in s of period l. Hence we obtain the required quasipolynomialness.

For the proof of Proposition 3.2 we first need two preliminary lemmas on counting integral points in polytopes.

Lemma 3.5. Let p and l be positive integers and let j be a non-negative integer. Set $R := \left| \frac{lj}{p} \right|$ and

$$\gamma_j := \# \left\{ (x, y) \in \mathbb{Z}_{\geq 0}^2 \mid \frac{x}{p+1} + \frac{y}{p+l} \leq \frac{j}{p} \right\}.$$

Then

(5)
$$\gamma_{j} = \frac{1}{2} \left(j - \left\lfloor \frac{R}{l} \right\rfloor \right) \left(j + \left\lfloor \frac{R}{l} \right\rfloor + 1 \right) + (j+1)(R+1) + \sum_{r=0}^{R} \min \left\{ \left\lfloor \frac{R}{l} \right\rfloor, \left\lfloor \frac{1}{l-1} (j-(p+1)r + R) \right\rfloor \right\}.$$

Proof. As $\frac{x}{p+1} + \frac{y}{p+l} \le \frac{j}{p}$, we have $x \le \frac{j(p+1)}{p} = j + \frac{j}{p}$, i.e. $x \le j + \left\lfloor \frac{j}{p} \right\rfloor$. For each x, we have

$$y \le \left(\frac{j}{p} - \frac{x}{p+1}\right)(p+l) = j - x + \frac{lj}{p} - \frac{(l-1)x}{p+1}.$$

In other words, $y \leq j - x + \left\lfloor \frac{lj}{p} - \frac{(l-1)x}{p+1} \right\rfloor$. Therefore,

$$\gamma_{j} = \sum_{x=0}^{j+\left\lfloor \frac{j}{p} \right\rfloor} (j-x) + 1 + \left\lfloor \frac{lj}{p} - \frac{(l-1)x}{p+1} \right\rfloor$$
$$= \frac{1}{2} \left(j - \left\lfloor \frac{j}{p} \right\rfloor \right) \left(j + \left\lfloor \frac{j}{p} \right\rfloor + 1 \right) + \sum_{x=0}^{j+\left\lfloor \frac{j}{p} \right\rfloor} 1 + \left\lfloor \frac{lj}{p} - \frac{(l-1)x}{p+1} \right\rfloor.$$

For any non-negative integer r we have

$$\left\lfloor \frac{lj}{p} - \frac{(l-1)x}{p+1} \right\rfloor \ge r \Longleftrightarrow \frac{lj}{p} - \frac{(l-1)x}{p+1} \ge r \Longleftrightarrow x \le \frac{p+1}{l-1} \left(\frac{lj}{p} - r \right)$$

$$\iff x \le j + \left\lfloor \frac{1}{l-1} \left(j - (p+1)r + \frac{lj}{p} \right) \right\rfloor = j + \left\lfloor \frac{1}{l-1} (j - (p+1)r + R) \right\rfloor.$$

Clearly $j + \left\lfloor \frac{1}{l-1}(j-(p+1)r+R) \right\rfloor \ge 0$ for every $0 \le r \le R$. Also note that $\left\lfloor \frac{j}{p} \right\rfloor = \left\lfloor \frac{R}{l} \right\rfloor$. We have

$$\sum_{x=0}^{j+\left\lfloor\frac{j}{p}\right\rfloor} 1 + \left\lfloor\frac{lj}{p} - \frac{(l-1)x}{p+1}\right\rfloor = \sum_{r=0}^{R} \# \left\{0 \le x \le j + \left\lfloor\frac{R}{l}\right\rfloor + \left\lfloor\frac{lj}{p} - \frac{(l-1)x}{p+1}\right\rfloor \ge r\right\}$$

$$= \sum_{r=0}^{R} \# \left\{0 \le x \le j + \left\lfloor\frac{R}{l}\right\rfloor + x \le j + \left\lfloor\frac{1}{l-1}(j-(p+1)r+R)\right\rfloor\right\}$$

$$= \sum_{r=0}^{R} j + 1 + \min\left\{\left\lfloor\frac{R}{l}\right\rfloor, \left\lfloor\frac{1}{l-1}(j-(p+1)r+R)\right\rfloor\right\}$$

$$= (j+1)(R+1) + \sum_{r=0}^{R} \min\left\{\left\lfloor\frac{R}{l}\right\rfloor, \left\lfloor\frac{1}{l-1}(j-(p+1)r+R)\right\rfloor\right\}.$$

Putting all together, we get the desired formula.

Lemma 3.6. Let p be positive integer and let l, η be a non-negative integer. Then

$$\delta_{\eta} := \# \left\{ \mathbf{x} \in \mathbb{Z}^n \mid x_i \ge 0 \text{ and } \sum_{i=1}^{n-1} \frac{x_i}{p} + \frac{x_n}{p+l} \le \eta \right\}$$

is a quasi-polynomial in p of period l (meaning that δ_{η} is a polynomial in p if l=0,1).

Proof. If l=0, then this follows from Ehrhart's theorem (see e.g. [BR07, Theorem 3.23]), so we can assume $l \geq 1$. Denote by $\sigma := \sum_{i=1}^{n-1} x_i$. Then for any given integer $j \in [0, \eta p]$, the range of $y := x_n$ such that $\sigma = \eta p - j$ is

$$y \le (p+l)\frac{j}{p} = j + \frac{lj}{p}$$
, i.e. $y \le j + R$,

where $R = \left\lfloor \frac{lj}{p} \right\rfloor$. For each σ , the number of solutions of (x_1, \dots, x_{n-1}) is $F(\sigma) := \binom{\sigma + \stackrel{\mathsf{L}}{n-2}}{n-2}$. Thus

$$\delta_{\eta} = \sum_{j=0}^{\eta p} F(\eta p - j) \cdot (j + R + 1)$$

$$= \sum_{j=0}^{\eta p} F(\eta p - j) \cdot (j + 1) + \sum_{k=0}^{\eta l - 1} R \cdot \sum_{j=e_{p}}^{e_{k}^{+}} F(\eta p - j) + F(0) \cdot \eta l,$$

where e_R^{\pm} are defined as

$$e_R^- := \left\lceil \frac{pR}{l} \right\rceil, \quad e_R^+ := \left\lfloor \frac{p(R+1) - 1}{l} \right\rfloor.$$

By Lemma 3.4, the first term in δ_{η} is a polynomial in p and in the second term, $\sum_{j=e_R^-}^{e_R^+} F(\eta p - j)$ is a polynomial in $\eta p - e_R^-$ and $\eta p - e_R^+$. The latter are quasi-polynomials in p of period l. Thus δ_{η} is a quasi-polynomial in p of period l.

For $j, r, R \in \mathbb{Z}_{>0}$, let

$$\gamma_R'(j) := \frac{1}{2} \left(j - \left\lfloor \frac{R}{l} \right\rfloor \right) \left(j + \left\lfloor \frac{R}{l} \right\rfloor + 1 \right) + (j+1)(R+1),$$

$$\gamma_{r,R}''(j) := \min \left\{ \left\lfloor \frac{R}{l} \right\rfloor, \left\lfloor \frac{1}{l-1} (j - (p+1)r + R) \right\rfloor \right\}$$

so that $\gamma_j = \gamma_R'(j) + \sum_{r=0}^R \gamma_{r,R}''(j)$ for $R = \left| \frac{lj}{p} \right|$ as in Lemma 3.5.

Proof of Proposition 3.2. The proof has several steps. Denote by $\sigma := \sum_{i=2}^{n-2} x_i$, $x := x_{n-1}$, and $y := x_n$. We start with the following reduction.

Claim. It is enough to show that, for each integer $1 \le \eta \le n-1$, the number

$$\beta_{\eta} := \# \left\{ \mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid x_i \ge 0 \text{ and } \sum_{i=2}^n \frac{x_i}{a_i} \le \eta \right\}$$

is a quasi-polynomial in p with period l(l-1).

Proof of Claim. Since $a_0 = a_1 = 2$, we have

$$\tau(\mathbf{a}_{p,l}) = \# \left\{ \mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid 0 < x_i < a_i \text{ and } 1 < \sum_{i=2}^n \frac{x_i}{a_i} < 2 \mod 2 \right\}$$
$$- \# \left\{ \mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid 0 < x_i < a_i \text{ and } 0 < \sum_{i=2}^n \frac{x_i}{a_i} < 1 \mod 2 \right\}.$$

For each $1 \le \eta \le n-1$, let

$$\alpha_{\eta} := \# \left\{ \mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid 0 < x_i < a_i \text{ and } 0 < \sum_{i=2}^n \frac{x_i}{a_i} \le \eta \right\}.$$

Clearly,

$$\alpha_{\eta} - \alpha_{\eta - 1} = \#\{\mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid 0 < x_i < a_i \text{ and } \eta - 1 < \sum_{i=2}^n \frac{x_i}{a_i} < \eta\},$$

as

$$\sum_{i=2}^{n} \frac{x_i}{a_i} = \frac{\sigma}{p} + \frac{x}{p+1} + \frac{y}{p+l}$$

is not an integer by coprimeness for p+1, p+l together with 0 < x < p+1 or 0 < y < p+l. Thus $\tau(\mathbf{a}_{p,l}) = \sum_{\eta=1}^{n-1} (-1)^{\eta} (\alpha_{\eta} - \alpha_{\eta-1})$. This reduces the problem to show that α_{η} is a quasi-polynomial in p of period l(l-1) for each $1 \le \eta \le n-1$.

We define

$$\beta'_{\eta} := \# \left\{ \mathbf{x} = (x_2, \dots, x_n) \in \mathbb{Z}^{n-1} \mid x_i > 0 \text{ and } \sum_{i=2}^n \frac{x_i}{a_i} \le \eta \right\}.$$

If $U:=\{\mathbf{x}\in\mathbb{Z}^{n-1}\mid x_i>0 \text{ and } \sum_{i=2}^n\frac{x_i}{a_i}\leq\eta\}$, and $U_i:=\{\mathbf{x}\in U\mid x_i\geq a_i\}$ for $i=2,\ldots,n$, then $\alpha_\eta=\#(U\setminus\bigcup_{i=2}^nU_i)$. By inclusion-exclusion, we know that α_η is an integer combination of $\#U=\beta_\eta'$ and $\#\cap_{i\in I}U_i$ for subsets $I\subset\{2,3,\cdots,n\}$. By replacing x_i with x_i-a_i , we see that $\#\cap_{i\in I}U_i=\beta_{\eta-\#I}$. Thus α_η is an integer combination of β_\bullet and β_\bullet' , so it suffices to show quasipolynomialness of β 's. Furthermore, using inclusion-exclusion again we can express $\beta_\eta-\beta_\eta'$ as a integer combination of lower-dimensional counting numbers of the form β_η (or δ_η as in Lemma 3.6). Hence, using induction on the dimension we reduce to only showing that β_η are quasi-polynomials in p and the claim is proven.

We now focus on β_{η} . If we fix $\sigma \geq 0$, then the number of solutions $\sum_{i=2}^{n-2} x_i = \sigma$ with each $x_i \in \mathbb{Z}_{\geq 0}$ equals $\binom{\sigma+n-4}{n-4}$. For each $0 \leq j \leq \eta p$, let γ_j be as in Lemma 3.5. Denote by $F(x) := \binom{x+n-4}{n-4}$.

Thus we have

$$\beta_{\eta} = \sum_{j=0}^{\eta p} F(\eta p - j) \gamma_{j}.$$

For each $0 \le R \le \eta l - 1$, the range of j is $e_R^- \le j \le e_R^+$ with

(6)
$$e_R^- := \left\lceil \frac{pR}{l} \right\rceil, \quad e_R^+ := \left\lfloor \frac{p(R+1) - 1}{l} \right\rfloor.$$

Clearly e_R^{\pm} are linear quasi-polynomials in p of period l and $e_{R+1}^- = e_R^+ + 1$. Hence we have

$$\beta_{\eta} = \sum_{j=0}^{\eta p} F(\eta p - j) \gamma_{j} = F(0) \gamma_{\eta p} + \sum_{R=0}^{\eta l - 1} \sum_{j=e_{R}^{-}}^{e_{R}^{+}} F(\eta p - j) \gamma_{j}.$$

By Pick's theorem (See e.g. [BR07, Theorem 2.18]), $\gamma_{\eta p}$ is a polynomial in p. Thus the problem reduces to showing quasi-polynomialness of $\sum_{R=0}^{\eta l-1} \sum_{j=e^-}^{e^+_R} F(\eta p-j) \gamma_j$ which is equal to

$$\sum_{R=0}^{\eta l-1} \sum_{j=e_R^-}^{e_R^+} F(\eta p - j) \gamma_R'(j) + \sum_{R=0}^{\eta l-1} \sum_{j=e_R^-}^{e_R^+} \sum_{r=0}^R F(\eta p - j) \gamma_{r,R}''(j).$$

The former sum is a quasi-polynomial in p as $F(\eta p-j)\gamma_R'(j)$ is a polynomial in j for a fixed R, thus, by Lemma 3.4, $\sum_{j=e_R^-}^{e_R^+} F(\eta p-j)\gamma_R'(j)$ is a polynomial in e_R^{\pm} which are quasi-polynomials in p. We only need to focus on the latter sum with γ''_j . Suppose $r \in [0, \eta l - 1]$ is fixed, and R ranges in $[r, \eta l - 1]$. Then j ranges in $[e_r^-, \eta p - 1]$. Note that

$$\sum_{R=0}^{\eta l-1} \sum_{j=e_R^-}^{e_R^+} \sum_{r=0}^R F(\eta p - j) \cdot \gamma_{r,R}''(j) = \sum_{r=0}^{\eta l-1} \sum_{R=r}^{\eta l-1} \sum_{j=e_R^-}^{e_R^+} F(\eta p - j) \cdot \gamma_{r,R}''(j).$$

Thus it reduces to showing quasi-polynomialness (in p with period l(l-1)) of the following function

(7)
$$\sum_{j=e_r^-}^{\eta p-1} F(\eta p - j) \cdot \gamma_{r,R}''(j) = \sum_{R=r}^{\eta l-1} \sum_{j=e_R^-}^{e_R^+} F(\eta p - j) \cdot \gamma_{r,R}''(j).$$

 $\gamma_{r,R}''(j)$ can be described by the following.

Lemma 3.7. (1) If
$$j \leq pr$$
, then $\lfloor \frac{R}{l} \rfloor \geq \lfloor \frac{1}{l-1}(j-(p+1)r+R) \rfloor$.
(2) If $j \geq pr$, then $\lfloor \frac{R}{l} \rfloor \leq \lfloor \frac{1}{l-1}(j-(p+1)r+R) \rfloor$.

Proof. We look at the difference

$$l(l-1)\left(\frac{R}{l} - \frac{1}{l-1}(j-(p+1)r + R)\right) = l(p+1)r - lj - R.$$

If
$$j \leq pr$$
, then $R = \left\lfloor \frac{lj}{p} \right\rfloor \leq lr$, so $l(p+1)r - lj - R \geq l(p+1)r - lpr - lr = 0$.
If $j \geq pr$, then $R \geq lr$, so similarly $l(p+1)r - lj - R \leq 0$.

Clearly $e_r^- = \lceil \frac{pr}{l} \rceil \le pr$. From Lemma 3.7 we know that (7) can be expressed as

If
$$r \leq \eta - 1$$
:
$$\sum_{j=e_r^-}^{pr-1} F(\eta p - j) \cdot \left\lfloor \frac{R}{l} \right\rfloor + \sum_{j=pr}^{\eta p-1} F(\eta p - j) \cdot \left\lfloor \frac{1}{l-1} (j - (p+1)r + R) \right\rfloor.$$
If $r \geq \eta$:
$$\sum_{j=e_r^-}^{\eta p-1} F(\eta p - j) \cdot \left\lfloor \frac{R}{l} \right\rfloor.$$

Note that $pr-1=e^+_{lr-1}$, $pr=e^-_{lr}$, $\eta p-1=e^+_{\eta l-1}$, and that, if $r\leq \eta-1$ (resp. $r\geq \eta$), then (7) becomes

$$\sum_{R=r}^{lr-1} Q_1(p,R) + \sum_{R=lr}^{\eta l-1} Q_2(p,R) \quad \left(\text{resp. } \sum_{R=r}^{\eta l-1} Q_1(p,R)\right),$$

where

$$Q_1(p,R) := \sum_{j=e_R^-}^{e_R^+} F(\eta p - j) \cdot \left\lfloor \frac{R}{l} \right\rfloor$$
 and

$$Q_2(p,R) := \sum_{j=e_R^-}^{e_R^+} F(\eta p - j) \cdot \left[\frac{1}{l-1} (j - (p+1)r + R) \right].$$

Hence it suffices to show quasi-polynomialness of $Q_1(p,R)$ and $Q_2(p,R)$ (in p), where $r,R \in [0,\eta l-1]$ are both fixed in Q_1 , and $r \in [0,\eta-1]$ and $R \in [lr,\eta l-1]$ are both fixed in Q_2 . Hence we write $Q_1(p) := Q_1(p,R)$ and $Q_2(p) := Q_2(p,R)$.

By Lemma 3.4, $Q_1(p)$ is clearly a polynomial in terms of p and e_R^+ , e_R^- , both being a quasi-polynomial in p of period l. Thus $Q_1(p)$ is a quasi-polynomial in p of period l.

Since $F(\eta p - j) \cdot \left\lfloor \frac{1}{l-1}(j - (p+1)r + R) \right\rfloor$ is a quasi-polynomial in j and p of period l-1, by using Lemma 3.4 again, we see that $Q_2(p)$ is a quasi-polynomial in e_R^{\pm} and p, thus $Q_2(p)$ is a quasi-polynomial in p of period l(l-1).

The proof is finished.
$$\Box$$

We end this section with a natural conjecture.

Conjecture 3.8. Let n be a positive integer. For each $1 \leq i \leq n$, let b_i, c_i be non-negative integers satisfying $(b_i, c_i) \neq (0, 0)$. Define a function $P: \mathbb{Z}_{\geq 0} \to \mathbb{Z}$ as

$$P(t) := \# \left\{ \mathbf{x} \in \mathbb{Z}^n \mid x_i \ge 0 \text{ and } \sum_{i=1}^n \frac{x_i}{b_i t + c_i} \le 1 \right\}.$$

In other words, P(t) is the number of lattice points in the closed simplex as the convex hull of $\{0\} \cup \{(b_i t + c_i) \mathbf{e}_i\}_{1 \leq i \leq n}$ where $(\mathbf{e}_1, \dots, \mathbf{e}_n)$ is the standard basis of \mathbb{Z}^n . Then P(t) is a quasi-polynomial for $t \in \mathbb{Z}_{>0}$.

As we saw earlier, the proof of Proposition 3.2 reduces to showing a special case of Conjecture 3.8.

If all (b_i, c_i) are proportional, then Conjecture 3.8 follows from a classical theorem of Ehrhart (see e.g. [BR07, Theorem 3.23]). For the general case, Conjecture 3.8 is proved under the extra assumption of $t \gg 1$ in [CLS10].

4. Dimension of the moduli spaces

By modifying examples in 3.2.2, we obtain a family of Sasaki-Einstein metrics of arbitrarily large dimension on homotopy spheres.

Proposition 4.1. Let $n \geq 5$ be an integer. Let $p \gg 0$ and $l \in \mathbb{Z}_{\geq 2}$ be as in 3.2.2 so that p is even and p, p + 1, p + l are pairwise coprime. Let

$$Y := (z_0^2 + z_1^2 + F_p + z_{n-1}^{p+1} + z_n^{p+l} = 0) \subset \mathbb{C}^{n+1},$$

where $F_p \in \mathbb{C}[z_2, \dots, z_{n-2}]$ is a general homogeneous polynomial of degree p, that is, Y is a deformation of $Y(\mathbf{a}_{p,l})$ for

$$\mathbf{a}_{p,l} = (a_0, \dots, a_n) = (2, 2, p, \dots, p, p+1, p+l).$$

Let ξ be the Reeb field of weights (d_0,\ldots,d_n) , where $d_i=d/a_i$ for

$$d := p(p+1)(p+l)$$

and $j = 0, \ldots, n$.

Then (Y, ξ) is K-polystable.

Remark 4.2. Note that Y gives a semi-universal deformations of $Y(\mathbf{a}_{p,l})$ with the \mathbb{C}^* -action.

Proof. Let (X, Δ_X) be the log pair corresponding to the orbifold Y/\mathbb{C}^* for the \mathbb{C}^* -action induced by ξ . We check that

$$X \simeq (z_0^2 + z_1^2 + F_p + y_{n-1} + y_n = 0) \subset \mathbb{P}\left(\frac{p}{2}, \frac{p}{2}, 1, \dots, 1, p, p\right)$$

and $\Delta_X = (1 - \frac{1}{p+1})H_{n-1} + (1 - \frac{1}{p+1})H_n$ for $H_{n-1} := (y_{n-1} = 0)$ and $H_n := (y_n = 0)$. As in Theorem 1.4, it is enough to show the K-polystability of (X, Δ_X) .

$$X_p := (y_0 + y_1 + F_p(y_2, \dots, y_{n-2}) + y_{n-1} + y_n = 0) \subset \mathbb{P}(p, p, 1, \dots, 1, p, p),$$

where y_0, \dots, y_n are the coordinates of $\mathbb{P}(p, p, 1, \dots, 1, p, p)$. Let $\phi_p \colon X \to X_p$ be the Galois cover determined by

$$[z_0:z_1:z_2:\cdots:z_{n-2}:y_{n-1}:y_n]\mapsto [z_0^2:z_1^2:z_2:\cdots:z_{n-2}:y_{n-1}:y_n].$$

Let

$$\Delta_p := \frac{1}{2} \left(H_0' + H_1' \right) + \left(1 - \frac{1}{p+1} \right) H_{n-1}' + \left(1 - \frac{1}{p+l} \right) H_n',$$

where $H'_i := (y_i = 0) \subset X_p$. Then we have

$$K_X + \Delta_X = \phi_p^* (K_{X_p} + \Delta_p),$$

thus the K-polystability of (X, Δ_X) is reduced to that of (X_p, Δ_p) . In fact, we shall show:

Claim. (i) The automorphism group of (X_p, Δ_p) is finite.

(ii) Moreover, the pair (X_p, Δ_p) is uniformly K-stable.

Proof of Claim. For both statements, we may assume that $F_p = y_2^p + \cdots + y_{n-2}^p$ by the upper-semicontinuity and the openness of the uniform K-stability [BL22, BLX22].

(i) This is a slight modification of the proof of [BGK05, Proposition 37]. We need to show that there are no continuous families of isomorphisms of the form

$$\tau_t(y_i) = y_i + \sum_{j \ge 1} t^j g_{ij}(y_0, \dots, y_n)$$

satisfying $\tau_t(y_0y_1) = y_0y_1$, $\tau_t(y_j) = y_j$ for j = n - 1, n and

$$\tau_t(y_0 + y_1) + \sum_{i=2}^{n-2} \tau_t(y_i)^p + \tau_t(y_{n-1} + y_n) = y_0 + y_1 + \sum_{i=2}^{n-2} y_i^p + y_{n-1} + y_n$$

since τ_t induces elements of $\operatorname{Aut}(X_p, \Delta_p)$. The last equality implies

$$\tau_t(y_0 + y_1) = y_0 + y_1,$$

thus we check $\tau_t(y_i) = y_i$ for i = 0, 1 and

$$\sum_{i=2}^{n-2} \tau_t(y_i)^p = \sum_{i=2}^{n-2} y_i^p.$$

By the same argument as in the proof of [BGK05, Proposition 37] and $p \ge 3$, we see that $\tau_t(y_i) = y_i$ for i = 2, ..., n-2. This is a contradiction.

(ii) We use the same symbols L, L_i, \ldots as in Theorem 1.4. Let

$$L := (w_0 + w_1 + \dots + w_n = 0) \subset \mathbb{P}^n$$

with the Galois cover

$$\pi_p \colon X_p \to L; \ [y_0 : \cdots : y_n] \mapsto [y_0 : y_1 : y_2^p : \cdots : y_{n-2}^p : y_{n-1} : y_n]$$

and let

$$\Delta_L := \sum_{i=0}^n \left(1 - \frac{1}{a_i} \right) L_i.$$

Note that (L, Δ_L) is K-stable and $K_{X_p} + \Delta_p = \pi_p^*(K_L + \Delta_L)$ as in the proof of Theorem 1.4. By this and [LZ22, Theorem 1.2], [Zhu21, Corollary 4.13], we see that (X_p, Δ_p) is K-polystable. This implies that (X_p, Δ_p) is K-stable by (i) and hence uniformly K-stable by [LXZ22].

Proof of Corollary 1.5. Let (d_0, \ldots, d_n) and d be as in Proposition 4.1 and let $\mathbb{P} := \mathbb{P}(d_0, \dots, d_n)$. By Proposition 4.1 and [BGK05, (15.2)], we see that the connected component containing the family of weighted hypersurfaces in \mathbb{P} induced by Y as in Proposition 4.1 has dimension at least

$$h^{0}(\mathbb{P}, \mathcal{O}(d)) - \sum_{i=0}^{n} h^{0}(\mathbb{P}, \mathcal{O}(d_{i}))$$

$$= \left(3 + 2\left(\frac{\frac{p}{2} + n - 4}{n - 4}\right) + \left(\frac{p + n - 4}{n - 4}\right) + 2\right)$$

$$- \left(2\left(2 + \left(\frac{\frac{p}{2} + n - 4}{n - 4}\right)\right) + (n - 3)^{2} + 2\right)$$

$$= \left(\frac{p + n - 4}{n - 4}\right) - (n - 3)^{2} - 1.$$

This is unbounded for $p \gg 0$ and n > 4. This and Proposition 2.8 imply that the dimension of a family of Sasaki-Einstein metrics on the link $L(\mathbf{a}_{p,l})$ induced by Y is unbounded.

When $n=2m\geq 6$ is even, for a given $\Sigma\in bP_{4m}$, there are infinitely many p and l such that $L(\mathbf{a}_{p,l})$ is diffeomorphic to Σ by Corollary 3.3. When $n=2m-1\geq 5$ is odd, the link $L(\mathbf{a}_{p,l})$ is a standard sphere S^{4m-3} by the criterion in 2.1. By these, we finish the proof of Corollary 1.5.

Remark 4.3. The construction in the proof of Corollary 1.5 does not provide families of unbounded dimensions on S^5 , homotopy 7-spheres, or the Kervaire spheres.

4.1. Mean Euler characteristic. Consider the links $L(\mathbf{a}_{p,l})$ with the contact structure induced by the construction of Subsection 3.2.2. We are going to compute the mean Euler characteristic χ_m of $L(\mathbf{a}_{p,l})$. As explained in [BMvK16, Section 4.5] and [KvK16, Section 5.8], this is an invariant of the contact structure. For the notation we refer to [BMvK16, KvK16].

Link	Period	χ^{S^1}	Frequency
$L(2,2,p\ldots,p,p+1,p+l)$	p(p+1)(p+l)	n	1
L(2,2,p+1,p+l)	2(p+1)(p+l)	3	p/2 - 1
L(p+1, p+l)	(p+1)(p+l)	1	p/2
$L(2,2,p\ldots,p,p+l)$	p(p+l)	n-1	p
$L(2,2,p\ldots,p,p+1)$	p(p + 1)	n-1	p+l-1
L(2,2,p+l)	2(p+l)	2	p(p+1)/2 - 3p/2
L(2,2,p+1)	2(p+1)	2	p(p+l)/2 - 3p/2 - l + 1
$L(2,2,p,\ldots,p)$	p	$\chi_p^{S^1}$	(p+1)(p+l) - 2p - l
L(2,2)	2	2	ϕ_2

where

$$\phi_2 = \frac{p(p+1)(p+l)}{2} + 2p - p^2 - \frac{pl}{2}$$

and $\chi_p^{S^1}$ is a polynomial in p with leading term p^{n-4} (to compute it one can use [KvK16, Theorem 3.10]).

Note that all orbits except L(2,2) and $L(2,2,p,\ldots,p)$ are rational homology spheres by Brieskorn graph theorem [BG05, Theorem 39], so the computation of χ^{S^1} follows by [FSvK12, Lemma 4.5].

The Maslov index of the principal orbit is ([BMvK16, Formula 4.9])

$$\mu_P = 2p(p+1)(p+l)\left(1 + \frac{n-3}{p} + \frac{1}{p+1} + \frac{1}{p+l} - 1\right)$$
$$= 2\left((n-3)(p+1)(p+l) + p(2p+l+1)\right).$$

Hence

$$\chi_m = -\frac{2\phi_2 + \chi_p^{S^1}((p+1)(p+l) - 2p - l)}{\mu_P} - \frac{2(p(p+l)/2 - 3p/2 - l + 1) + 2(p(p+1)/2 - 3p/2)}{\mu_P} - \frac{(n-1)(p+l-1) + (n-1)p + p/2 + 3(p/2 - 1) + n}{\mu_P}.$$

Since $\chi_p^{S^1}$ has leading term p^{n-4} , χ_m assumes infinitely many different values for $p \gg 0$. This implies that the corresponding contact structures are not isomorphic.

References

- [AJL21] Vestislav Apostolov, Simon Jubert, and Abdellah Lahdili. Weighted K-stability and coercivity with applications to extremal Kähler and Sasaki metrics. *Geom. Topol.*, to appear, 2021. arXiv:2104.09709.
- [BdFFU15] S. Boucksom, T. de Fernex, C. Favre, and S. Urbinati. Valuation spaces and multiplier ideals on singular varieties. In *Recent advances in algebraic* geometry, volume 417 of *London Math. Soc. Lecture Note Ser.*, pages 29–51. Cambridge Univ. Press, Cambridge, 2015.
- [Bea96] A. F. Beardon. Sums of powers of integers. Amer. Math. Monthly, 103(3):201– 213, 1996.
- [Bes87] Arthur L. Besse. Einstein manifolds, volume 10 of Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]. Springer-Verlag, Berlin, 1987.
- [BG01] Charles P. Boyer and Krzysztof Galicki. New Einstein metrics in dimension five. J. Differential Geom., 57(3):443–463, 2001.
- [BG05] Charles P. Boyer and Krzysztof Galicki. Sasakian geometry, hypersurface singularities, and Einstein metrics. *Rend. Circ. Mat. Palermo (2) Suppl.*, (75):57–87, 2005.
- [BG06] Charles P. Boyer and Krzysztof Galicki. Sasakian geometry and Einstein metrics on spheres. In *Perspectives in Riemannian geometry*, volume 40 of

- CRM Proc. Lecture Notes, pages 47–61. Amer. Math. Soc., Providence, RI, 2006.
- [BG08] Charles P. Boyer and Krzysztof Galicki. Sasakian geometry. Oxford Mathematical Monographs, Oxford University Press, Oxford, 2008.
- [BGK05] Charles P. Boyer, Krzysztof Galicki, and János Kollár. Einstein metrics on spheres. Ann. of Math. (2), 162(1):557–580, 2005.
- [BGKT05] Charles P. Boyer, Krzysztof Galicki, János Kollár, and Evan Thomas. Einstein metrics on exotic spheres in dimensions 7, 11, and 15. Experiment. Math., 14(1):59-64, 2005.
- [BL22] Harold Blum and Yuchen Liu. Openness of uniform K-stability in families of \mathbb{Q} -Fano varieties. Ann. Sci. Éc. Norm. Supér. (4), 55(1):1–41, 2022.
- [Blu18] Harold Blum. Existence of valuations with smallest normalized volume. Compos. Math., 154(4):820–849, 2018.
- [BLX22] Harold Blum, Yuchen Liu, and Chenyang Xu. Openness of K-semistability for Fano varieties. Duke Math. J., 171(13):2753-2797, 2022.
- [BMvK16] Charles P. Boyer, Leonardo Macarini, and Otto van Koert. Brieskorn manifolds, positive Sasakian geometry, and contact topology. Forum Math., 28(5):943–965, 2016.
- [BR07] Matthias Beck and Sinai Robins. Computing the continuous discretely. Undergraduate Texts in Mathematics. Springer, New York, 2007. Integerpoint enumeration in polyhedra.
- [Bri66] Egbert Brieskorn. Beispiele zur Differentialtopologie von Singularitäten. Invent. Math., 2:1-14, 1966.
- [Bro69] William Browder. The Kervaire invariant of framed manifolds and its generalization. Ann. of Math. (2), 90:157–186, 1969.
- [CLS10] Sheng Chen, Nan Li, and Steven V. Sam. Generalized Ehrhart polynomials. In 22nd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2010), volume AN of Discrete Math. Theor. Comput. Sci. Proc., pages 239–246. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2010.
- [CS19] Tristan C. Collins and Gábor Székelyhidi. Sasaki-Einstein metrics and Kstability. Geom. Topol., 23(3):1339-1413, 2019.
- [Dim92] Alexandru Dimca. Singularities and topology of hypersurfaces. Universitext. Springer-Verlag, New York, 1992.
- [DS17]Simon Donaldson and Song Sun. Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, II. J. Differential Geom., 107(2):327–371, 2017.
- [ELS03] Lawrence Ein, Robert Lazarsfeld, and Karen E. Smith. Uniform approximation of Abhyankar valuation ideals in smooth function fields. Amer. J. Math., 125(2):409-440, 2003.
- [FSvK12] Urs Frauenfelder, Felix Schlenk, and Otto van Koert. Displaceability and the mean Euler characteristic. Kyoto J. Math., 52(4):797-815, 2012.
- [Fuj21] Kento Fujita. K-stability of log Fano hyperplane arrangements. J. Algebraic Geom., 30(4):603-630, 2021.
- [GK07] Alessandro Ghigi and János Kollár. Kähler-Einstein metrics on orbifolds and Einstein metrics on spheres. Comment. Math. Helv., 82(4):877–902, 2007.
- [GMSY07] Jerome P. Gauntlett, Dario Martelli, James Sparks, and Shing-Tung Yau. Obstructions to the existence of Sasaki-Einstein metrics. Comm. Math. Phys., 273(3):803-827, 2007.
- [HHR16] M. A. Hill, M. J. Hopkins, and D. C. Ravenel. On the nonexistence of elements of Kervaire invariant one. Ann. of Math. (2), 184(1):1–262, 2016.
- Hans-Joachim Hein and Song Sun. Calabi-Yau manifolds with isolated conical [HS17] singularities. Publ. Math. Inst. Hautes Études Sci., 126:73–130, 2017.

- [IWX23] Daniel C. Isaksen, Guozhen Wang, and Zhouli Xu. Stable homotopy groups of spheres: from dimension 0 to 90. Publ. Math. Inst. Hautes Études Sci., 137:107-243, 2023.
- [JM12] Mattias Jonsson and Mircea Mustață. Valuations and asymptotic invariants for sequences of ideals. *Ann. Inst. Fourier (Grenoble)*, 62(6):2145–2209, 2012.
- [KM63] Michel A. Kervaire and John W. Milnor. Groups of homotopy spheres. I. Ann. of Math. (2), 77:504–537, 1963.
- [KvK16] Myeonggi Kwon and Otto van Koert. Brieskorn manifolds in contact topology. Bull. Lond. Math. Soc., 48(2):173–241, 2016.
- [Lev85] J. P. Levine. Lectures on groups of homotopy spheres. In Andrew Ranicki, Norman Levitt, and Frank Quinn, editors, Algebraic and Geometric Topology, pages 62–95, Berlin, Heidelberg, 1985. Springer Berlin Heidelberg.
- [Li18] Chi Li. Minimizing normalized volumes of valuations. *Math. Z.*, 289(1-2):491–513, 2018.
- [Li21] Chi Li. Notes on weighted Kähler-Ricci solitons and application to Ricci-flat Kähler cone metrics. 2021. arXiv:2107.02088.
- [LLX20] Chi Li, Yuchen Liu, and Chenyang Xu. A guided tour to normalized volume. In Geometric analysis—in honor of Gang Tian's 60th birthday, volume 333 of Progr. Math., pages 167–219. Birkhäuser/Springer, Cham, 2020.
- [LWX21] Chi Li, Xiaowei Wang, and Chenyang Xu. Algebraicity of the metric tangent cones and equivariant K-stability. J. Amer. Math. Soc., 34(4):1175–1214, 2021.
- [LX18] Chi Li and Chenyang Xu. Stability of valuations: higher rational rank. *Peking Math. J.*, 1(1):1–79, 2018.
- [LXZ22] Yuchen Liu, Chenyang Xu, and Ziquan Zhuang. Finite generation for valuations computing stability thresholds and applications to K-stability. *Ann. of Math.* (2), 196(2):507–566, 2022.
- [LZ22] Yuchen Liu and Ziwen Zhu. Equivariant K-stability under finite group action.

 Internat. J. Math., 33(1):Paper No. 2250007, 21, 2022.
- [Mil68] John Milnor. Singular points of complex hypersurfaces, volume No. 61. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1968.
- [MSY08] Dario Martelli, James Sparks, and Shing-Tung Yau. Sasaki-Einstein manifolds and volume minimisation. Comm. Math. Phys., 280(3):611–673, 2008.
- [PW21] Jihun Park and Joonyeong Won. Simply connected Sasaki-Einstein rational homology 5-spheres. *Duke Math. J.*, 170(6):1085–1112, 2021.
- [ST24] Taro Sano and Luca Tasin. On K-stability of Fano weighted hypersurfaces. Algebr. Geom., 11(2):296–317, 2024.
- [Zhu21] Ziquan Zhuang. Optimal destabilizing centers and equivariant K-stability. Invent. Math., 226(1):195–223, 2021.
- [Zhu23] Ziquan Zhuang. Stability of klt singularities. 2023. arXiv:2307.10525.

Department of Mathematics, Northwestern University, Evanston, IL 60208, USA

Email address: yuchenl@northwestern.edu

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, KOBE UNIVERSITY, 1-1, ROKKODAI, NADA-KU, KOBE 657-8501, JAPAN

Email address: tarosano@math.kobe-u.ac.jp

Dipartimento di Matematica F. Enriques, Università degli Studi di Milano, Via Cesare Saldini 50, 20133 Milano, Italy

Email address: luca.tasin@unimi.it