Enumerative aspect of symplectic log Calabi-Yau divisors and almost toric fibrations

Tian-Jun Li, Jie Min, Shengzhen Ning

March 17, 2022

Abstract

In this paper we are interested in the isotopy classes of symplectic log Calabi-Yau divisors in a fixed symplectic rational surface. We give several equivalent definitions and prove the stability, finiteness and rigidity results. Motivated by the problem of counting toric actions, we obtain a general counting formula of symplectic log Calabi-Yau divisors in a restrictive region of c_1 -nef cone. A detailed count in the case of 2- and 3-point blow-ups of complex projective space for all symplectic forms is also given. In our framework the complexity of the combinatorics of analyzing Delzant polygons is reduced to the arrangement of homology classes. Then we study its relation with almost toric fibrations. We raise the problem of realizing all symplectic log Calabi-Yau divisors by some almost toric fibrations and verify it together with another conjecture of Symington in a special region.

Contents

1	Introduction				
	1.1	Space of symplectic log Calabi-Yau divisors	2		
	1.2	Stability and rigidity	4		
	1.3	Finiteness and counting	5		
	1.4	Relation with almost toric fibrations	6		
2	Isot	opy class space of symplectic log Calabi-Yau divisors	7		
	2.1	Symplectomorphism and symplectic Torelli theorem	7		
	2.2	Operations and minimal models	11		
	2.3	$\mathcal{HLCY}(X,\omega)$, realizability and stability	13		
	2.4	Finiteness	18		
	2.5	Tautness and toric symplectic log Calabi-Yau divisors	21		
3	c_1 -n	ef cone and counting symplectic log Calabi-Yau divisors	24		
	3.1	c_1 -nef subcone and counting	24		
	3.2	Count of minimal models	25		
	3.3	Counting symplectic log Calabi-Yau divisors in M_2 and M_3	26		
	3.4	Toric regions	31		
	3.5	General counting formulas for restrictive reduced symplectic classes	34		

4	Symplectic log Calabi-Yau divisors and almost toric fibrations		
	4.1	Toric manifolds and Delzant polytope	39
	4.2	Moment polytope and toric symplectic log Calabi-Yau divisors	40
	4.3	Comparison with the counting results of Karshon-Kessler-Pinsonnault	43
	4.4	Almost toric fibrations	44
	4.5	Realizing log Calabi-Yau divisors by ATF	45
	4.6	Connectedness of toric fibrations	50
	4.7	Realization and connectedness in the restrictive region	53
\mathbf{A}	App	pendix	58
	A.1	More discussions on tautness	58
	A.2	Details of counting toric divisors in M_3	61
	A.3	Details of computing toric regions for M_5	64

1 Introduction

Let X be a smooth rational surface and let $D \subset X$ be an effective reduced anti-canonical divisor. Such pairs (X, D) are called anti-canonical pairs or log Calabi-Yau surfaces and has been extensively studied since Looijenga ([Loo81]). The mirror symmetry aspects of anticanonical pairs were first studied by Gross, Hacking and Keel in [GHK15a] and [GHK15b]. Then it was studied by many people in for example [Pas14], [Kea18] and [HK20]. The moduli space of anti-canonical pairs have been studied in [Loo81] and [GHK15b], where Torelli type results were proved. See [Fri16] for an excellent survey. As an symplectic analogue of anticanonical pairs, the notion of symplectic log Calabi-Yau pairs was introduced and studied in [LM16], [LM21] and [LMM20], with applications to contact structures and symplectic fillings.

Instead of considering the pair of a symplectic manifold and a divisor, in this paper we are interested in the space of isotopy classes of symplectic log Calabi-Yau divisors in a fixed symplectic rational surface. We will give several different but equivalent definitions, prove stability, rigidity and finiteness properties. Making use of those results we count the number of elements in it and investigate its relation to toric actions and almost toric fibrations. Note that in this paper although the word "space" is frequently used, we will only study its properties as a set. In other word, no additionally topological or algebraic structures are added. Hopefully this will not cause confusion.

1.1 Space of symplectic log Calabi-Yau divisors

A symplectic divisor refers to a connected configuration of finitely many closed embedded symplectic surfaces $D = \bigcup C_i$ in a symplectic 4-manifold such that all intersections are positively transversal and there are no triple intersections. For a more detailed discussion on symplectic divisors, see [LM19] or [LM21].

A symplectic log Calabi-Yau pair (X, D, ω) is a closed symplectic 4-manifold (X, ω) together with a nonempty symplectic divisor $D = \bigcup C_i$ representing the Poincare dual of $c_1(X, \omega)$. It's an easy observation ([LM18]) that D is either a torus or a cycle of spheres. In the former case, (X, D, ω) is called an elliptic log Calabi-Yau pair. In the later case, it's called a symplectic Looijenga pair and can only happen when (X, ω) is rational. As a consequence, we have $b^+(Q_D) = 0$ or 1, where Q_D is the intersection matrix of D. The symplectic divisor D is then called a **symplectic log Calabi-Yau divisor** in (X, ω) .

Now fixing a symplectic rational surface (X, ω) , we denote by $p\mathcal{LCY}(X, \omega)$ the set of symplectic log Calabi-Yau divisors in (X, ω) , where the prefix 'p' stands for 'pre'. The symplectomorphism group $Symp(X, \omega)$ naturally acts on $p\mathcal{LCY}(X, \omega)$. The following space we mainly study in this paper can be viewed as the set of path components of $p\mathcal{LCY}(X, \omega)$ modulo the action of $Symp(X, \omega)$.

Definition 1.1. The isotopy class space of symplectic log Calabi-Yau divisors is

 $\mathcal{LCY}(X,\omega) = (p\mathcal{LCY}(X,\omega)/Symp(X,\omega))/\sim,$

where $[D] \sim [D']$ with $[D], [D'] \in p\mathcal{LCY}(X, \omega)/Symp(X, \omega)$ if there exist representatives $D, D' \in p\mathcal{LCY}(X, \omega)$ and an symplectic isotopy D_t in (X, ω) with $D_0 = D$ and $D_1 = D'$.

Denote by $p\mathcal{LCY}^{\perp}(X,\omega)$ the subset of ω -orthogonal divisors in $p\mathcal{LCY}(X,\omega)$, that is, the divisors whose irreducible components all intersect ω -orthogonally.

Theorem 1.2. The following sets are different but equivalent ways to describe the isotopy class space of symplectic log Calabi-Yau divisors in (X, ω) .

$$\begin{split} \mathcal{LCY}(X,\omega) =& p\mathcal{LCY}(X,\omega) / strict \ symplectic \ deformation \ equivalence \\ \cong & p\mathcal{LCY}(X,\omega) / strict \ homological \ equivalence \\ \cong & p\mathcal{LCY}(X,\omega) / strict \ H^2 \text{-} automorphism \\ \cong & p\mathcal{LCY}^{\perp}(X,\omega) / symplectomorphism \end{split}$$

Here the notions of strict homological equivalence and strict symplectic deformation equivalence were defined in [LM16] and will be recalled in Definition 2.1. By a strict H^2 automorphism we mean an integral isometry of the lattice $H^2(X;\mathbb{Z})$ preserving the Poincare dual of the homology classes of components of D and the symplectic class $[\omega]$ (see Proposition 2.11). And by symplectomorphism we mean the restriction of the action of $Symp(X,\omega)$ on the subset $p\mathcal{LCY}^{\perp}(X,\omega) \subset p\mathcal{LCY}(X,\omega)$.

Theorem 1.2 is a meta-theorem, made up of three equivalences. The first equivalence is the main result of [LM16] (see also Theorem 2.2). The second equivalence is proved in Proposition 2.11, which can be viewed as an improvement of the first equivalence. The last equivalence is established in Proposition 2.10, where we prove a version of symplectic isotopy extension theorem for symplectic divisors, adapting an argument of Siebert and Tian ([ST05]). This theorem can be seen as a symplectic version of the Torelli theorem in [Fri16].

For two symplectic log Calabi-Yau pairs (X, ω_1, D_1) and (X, ω_2, D_2) with the same underlying manifold but different symplectic forms, there is also a notion of (non-strict) symplectic deformation equivalence introduced in [LM16], see also Definition 2.1. In this paper we also introduce the following space of (non-strict) deformation classes associated to a manifold X.

Definition 1.3. For a closed smooth 4-manifold X, the deformation class space of symplectic log Calabi-Yau divisors is

$$\widetilde{\mathcal{LCY}}(X) = \Big(\bigsqcup_{\substack{\omega \text{ symplectic forms on } X}} p\mathcal{LCY}(X,\omega)\Big)/\sim,$$

where \sim is the symplectic deformation equivalence.

Note that if $D_1, D_2 \in p\mathcal{LCY}(X, \omega)$ are strict deformation equivalent, then the pairs (X, ω, D_1) and (X, ω, D_2) are deformation equivalent. Therefore there is a natural map (not always injective) $\mathcal{LCY}(X, \omega) \to \widetilde{\mathcal{LCY}}(X)$ for each symplectic form ω .

Recall that a toric action on a symplectic 4-manifold (X, ω) is a group homomorphism $\rho: T \to Ham(X, \omega)$, where $T \cong (S^1)^2$ is a 2-torus. We define the space of toric actions

$$\mathcal{T}(X,\omega) = \{\rho: T^2 \to Ham(X,\omega)\} / \sim^t,$$

where \sim^t is the equivalence of toric actions on (X, ω) (Definition 4.1). For any toric action on a closed symplectic 4-manifold, we obtain a natural symplectic log Calabi-Yau divisor called the boundary divisor, as preimage of the boundary of moment polygon. To further inverstigate this relation between symplectic log Calabi-Yau divisors and toric actions on symplectic rational surfaces, we introduce the notion of toric symplectic log Calabi-Yau divisors. For this purpose, given $D \in p\mathcal{LCY}(X, \omega)$ we denote the length of D by k(D) and define $q(D) = 12 - k(D) - D^2$ (this is called the "charge" in [Fri16]).

Definition 1.4. $D \in p\mathcal{LCY}(X, \omega)$ is called a toric symplectic log Calabi-Yau divisor if q(D) = 0. Denote by

$$t\mathcal{LCY}(X,\omega) = \{ D \in p\mathcal{LCY}(X,\omega) \mid q(D) = 0 \} / \sim \subset \mathcal{LCY}(X,\omega)$$

the isotopy class space of toric symplectic log Calabi-Yau divisors, where \sim is the strict symplectic deformation equivalence. Moreover, let $\widetilde{tLCY}(X) \subset \widetilde{LCY}(X)$ be the subset consisting only deformation classes of toric symplectic log Calabi-Yau divisors.

The boundary divisors of a toric action is easily seen to be a toric symplectic log Calabi-Yau divisor. And in fact, the converse is also true up to strict symplectic deformation equivalence. In Section 4.2, we prove the following equivalence.

Theorem 1.5. Let (X, ω) be a symplectic rational surface. The natural map

$$\mathcal{T}(X,\omega) \to t\mathcal{LCY}(X,\omega)$$

taking a toric action to its boundary divisor is a bijection.

1.2 Stability and rigidity

To give a better description of the isotopy class space $\mathcal{LCY}(X, \omega)$, we also introduce a more combinatorial space $\mathcal{HLCY}(X, \omega)$ consisting of **homological log Calabi-Yau divisors** (see Definition 2.16). There is a natural map

$$\mathcal{LCY}(X,\omega) \to \mathcal{HLCY}(X,\omega)$$

by sending a geometric divisor to its underlying homological configuration. In Proposition 2.24 we prove this map is bijective. So we can actually identify $\mathcal{LCY}(X,\omega)$ and $\mathcal{HLCY}(X,\omega)$. It turns out working over $\mathcal{HLCY}(X,\omega)$ will simplify our later discussions on finiteness, counting and almost toric fibrations.

As an immediate consequence, we observe the following stability phenomenon. For a symplectic 4-manifold (X, ω) , denote by \mathcal{S}_{ω} the set of elements of $H_2(X; \mathbb{Z})$ that can be represented by ω -symplectic spheres. It turns out the isotopy class space of symplectic log Calabi-Yau divisors is determined by ω through \mathcal{S}_{ω} .

Theorem 1.6. Let X be a rational surface. Suppose ω and ω' are two symplectic forms with $c_1(X,\omega) \cdot [\omega] > 0$, $c_1(X,\omega') \cdot [\omega'] > 0$ and $\mathcal{S}_{\omega} = \mathcal{S}_{\omega'}$. Then

$$\mathcal{LCY}(X,\omega) = \mathcal{LCY}(X,\omega').$$

For each symplectic log Calabi-Yau divisor $D \subset (X, \omega)$ of length k, we can associate two invariants of D, its self-intersection sequence $s(D) = ([C_i]^2)_{i=1}^k \in \mathbb{Z}^k$ and its symplectic area sequence $a(D) = (\omega([C_i]))_{i=1}^k \in (\mathbb{R}^+)^k$. In the holomorphic category, a labeled toric anticanonical pair (defined in [Fri16]) is called **taut** if its isomorphism class is determined by its self-intersection sequence. Similarly we define a labeled symplectic log Calabi-Yau divisor to be **def-taut** (resp. **iso-taut**) if its symplectic deformation class (resp. isotopy class) is determined by s(D) (resp. (s(D), a(D))). Such notion can be seen as a type of rigidity for symplectic log Calabi-Yau divisors.

We show in Lemma 2.37 that toric symplectic log Calabi-Yau divisors are both def-taut and iso-taut, which will be used to prove Proposition 4.4. However general symplectic log Calabi-Yau divisors might lose the tautness. In the Appendix A.1 we give a criterion of deftautness for several families of $b^+ = 1$ divisors discussed in [LMM20]. These results are not used in the current paper but are related to the problems of classifying symplectic fillings of torus bundles. We hope there will be applications in the future.

1.3 Finiteness and counting

Karshon, Kessler and Pinsonnault proved the finiteness of toric actions in [KKP07] and gave an upper bound for the number of toric actions in [KKP15] and [KKP]. Inspired by their results and the identification of $\mathcal{T}(X,\omega)$ with $t\mathcal{LCY}(X,\omega)$ discussed above, we study the enumerative aspect of symplectic log Calabi-Yau divisors. The strategy of [KKP15] and [KKP] is to understand the combinatorics of Delzant polygons. In our framework, the combinatorics is reduced to the arrangement of homology classes by investigating the space $\mathcal{HLCY}(X,\omega)$.

First we observe that the isotopy class space of symplectic log Calabi-Yau divisors

$$|\mathcal{LCY}(X,\omega)| < \infty$$

is finite for any (X, ω) in Corollary 2.34. So it makes sense to count the number $|\mathcal{LCY}(X, \omega)|$.

Since we are only interested in symplectic log Calabi-Yau divisors up to strict homological equivalence, it suffices to count in the normalized reduced symplectic cone, which is the fundamental domain inside the symplectic cone under the action of orientation-preserving diffeomorphisms. A reduced symplectic class $[\omega]$ on $M_l = \mathbb{CP}^2 \# l \mathbb{CP}^2$ is determined by a reduced vector $(\delta_1, \ldots, \delta_l)$ and write $\mathcal{LCY}(l; \delta_1, \ldots, \delta_l) := \mathcal{LCY}(M_l, \omega)$. Note that the existence of symplectic log Calabi-Yau divisors implies $c_1 \cdot \omega > 0$. So we may focus on a subregion satisfying $\delta_1 + \cdots + \delta_l < 3$, which is called the c_1 -nef cone. (see Section 3.1).

By enumerating all homological log Calabi-Yau divisors, we get a detailed count for $M_2 = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$ with every reduced symplectic form in Proposition 3.5 and Corollary 3.6. For general symplectic log Calabi-Yau divisors, we have

$$|\mathcal{LCY}(2;\delta_1,\delta_2)| = 7(\lceil \frac{\delta_1}{1-\delta_1} \rceil + \lceil \frac{\delta_1-\delta_2}{1-\delta_1} \rceil) + 12,$$

except $|\mathcal{LCY}(2; \delta_1, \delta_2)| = 13$ when $\delta_1 = \delta_2 < \frac{1}{2}$. For toric symplectic log Calabi-Yau divisors, we have

$$|t\mathcal{LCY}(2;\delta_1,\delta_2)| = \lceil \frac{\delta_1}{1-\delta_1} \rceil + \lceil \frac{\delta_1-\delta_2}{1-\delta_1} \rceil.$$

For the case of $M_3 = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$, we also give a detailed counting result for toric symplectic log Calabi-Yau divisors in Proposition 3.7.

However in general it's quite difficult to give a detailed count for M_l when l is getting large, even for toric ones. Therefore we explore the region in the c_1 -nef cone which admits at least one toric action. This is already a very intricate and interesting question. In section 3.4 we give the complete answer of the toric regions for $l \leq 5$ and some geometric descriptions of the toric regions for general l.

In section 3.5, we introduce the notion of **restrictive** region, which is a smaller region than the toric region. It turns out that in this region, the counting problem is simplified dramatically and indeed we are able to give a general counting formula (Proposition 3.16), though quite involved, in this special region. As a by-product we recover the upper bound in [KKP15].

1.4 Relation with almost toric fibrations

An almost toric fibration of a symplectic 4-manifold (X, ω) is a Lagrangian fibration $\pi : X \to B$ with only nodal and elliptic singularities, which can be viewed as a generalization of toric actions. Almost toric fibrations play an important role in the study of exotic Lagrangians ([Via14], [Via16]) and symplectic embeddings ([CV20], [CGHMP21]). Similar to the case of toric fibrations, the preimage $\pi^{-1}(\partial B)$ of the base is a symplectic divisor representing the Poincare dual of $c_1(X, \omega)$ (Proposition 8.2 of [Sym03]), i.e. a symplectic log Calabi-Yau divisor. Moreover, the operations (nodal trade, almost toric blow up, toric blow up) on almost toric fibrations have their analogues (smoothing, non-toric blow up, toric blow up) on symplectic log Calabi-Yau divisors. Therefore we naturally want to understand their relations. Let $\mathcal{ATF}(X, \omega)$ denote all the almost toric fibrations on (X, ω) . The first issue is the following realization problem:

Conjecture 1.7 (Realization). The map $\Phi : \mathcal{ATF}(X, \omega) \to \mathcal{LCY}(X, \omega)$ is surjective.

Unlike the space of toric actions $\mathcal{T}(X,\omega)$ which contains the equivalent classes, the space $\mathcal{ATF}(X,\omega)$ here denotes all the almost toric fibrations. Thus the next step is to find a suitable equivalence relation \sim^{atf} such that a bijection between $\mathcal{ATF}(X,\omega)/\sim^{atf}$ and $\mathcal{LCY}(X,\omega)$ can be established.

Note that an affirmative answer to Conjecture 1.7 will imply every symplectic rational manifold with $\omega \cdot c_1 > 0$ admits almost toric fibrations, which is also a question we don't know the answer to. Based on the detailed counting results for small rational manifolds in the previous sections, we are able to prove Conjecture 1.7 holds for $M = S^2 \times S^2$ and $\mathbb{CP}^2 \# l \overline{\mathbb{CP}^2}$ with $l \leq 3$ with arbitrary symplectic forms (Proposition 4.17 and 4.19).

In [Sym03], Symington observed that different toric fibrations on $(S^2 \times S^2, \omega)$ can be changed to each other via nodal trades, nodal slides and mutations, which gives a path of almost toric fibrations connecting toric fibrations. This observation led to the following conjecture. **Conjecture 1.8** (Connectedness). Any two toric fibrations of (M, ω) can be connected by a path of almost toric fibrations of (M, ω) .

Based on Symington's observation and the detailed counting results for toric divisors in small rational manifolds, we can prove it for $M = S^2 \times S^2$ and $\mathbb{CP}^2 \# l\mathbb{CP}^2$ with $l \leq 3$ (Proposition 4.17).

It's hard to generalize the approaches of proving these two conjectures for small rational manifolds to the cases for arbitrary (X, ω) due to the difficulty of enumerating $\mathcal{LCY}(X, \omega)$ and $t\mathcal{LCY}(X, \omega)$. However, if $[\omega]$ is restrictive in the sense of section 3.5, we are fortune enough to verify these two conjectures. In other words, besides the cases for small rational manifolds $(M_l \text{ for } l \leq 3)$, there exists an open region inside the c_1 -nef cone supporting these two conjectures. The proofs are shown in Section 4.7.

Acknowledgments: We are grateful to Cheuk Yu Mak, Liya Ouyang and Shuo Zhang for helpful conversations. We also thank Yael Karshon for sharing with us a draft of their unpublished paper. All authors are supported by NSF grant DMS 1611680.

2 Isotopy class space of symplectic log Calabi-Yau divisors

2.1 Symplectomorphism and symplectic Torelli theorem

The goal of this section is to prove Theorem 1.2. It will be a combination of Theorem 2.2, Proposition 2.10 and Proposition 2.11. We begin with some notions and results in [LM16].

Definition 2.1. Let X be a closed oriented 4-manifold. Symplectic divisors (D^0, ω^0) and (D^1, ω^1) in X with the same number of components, are symplectic deformation equivalent if they are connected by a family of symplectic divisors (D^t, ω^t) , up to an orientation-preserving diffeomorphism. They are called homologically equivalent if there is an orientation-preserving diffeomorphism $\Phi: X \to X$ such that $\Phi_*[C_i^0] = [C_i^1]$ for all $j = 1, \ldots, k$.

Fix a closed symplectic 4-manifold (X, ω) . Symplectic divisors D^0 and D^1 in (X, ω) are strictly symplectic deformation equivalent if they are connected by a family of symplectic divisor D^t , up to a symplectomorphism of (X, ω) . They are strictly homologically equivalent if there is a homological equivalence fixing $[\omega]$.

The main result of [LM16] states that homological equivalence and symplectic deformation equivalence introduced above are equivalent for symplectic log Calabi-Yau divisors.

Theorem 2.2 ([LM16],[LM18]). Two symplectic log Calabi-Yau divisors (D^0, ω^0) and (D^1, ω^1) in X are symplectic deformation equivalent if and only if they are homologically equivalent. In particular, each symplectic deformation class contains a Kahler pair. Moreover, two symplectic log Calabi-Yau divisors D^0 and D^1 in (X, ω) are strictly symplectic deformation equivalent if and only if they are strictly homologically equivalent.

By the above theorem, we have the following equivalent definition for the isotopy class space of symplectic log Calabi-Yau divisors

 $\mathcal{LCY}(X,\omega) = p\mathcal{LCY}(X,\omega) / \text{strict symplectic deformation equivalence} \\ = p\mathcal{LCY}(X,\omega) / \text{strict homological equivalence}$

Remark 2.3. Definition 2.1 and Theorem 2.2 were originally stated for symplectic log Calabi-Yau pairs (X, ω, D) in [LM16]. The original statements restrict to the current ones by fixing X or (X, ω) .

For symplectic orthogonal divisors, the equivalences introduced in the previous subsection can be shown to be equivalent to symplectomorphism. This is a singular analogue of the following result of Siebert-Tian, saying that in the case of a single symplectic surface, a symplectic isotopy of the surface extends to a Hamiltonian isotopy of the ambient symplectic 4-manifold.

Proposition 2.4 (Proposition 0.3 of [ST05]). A smooth isotopy $\Sigma_t \subset (X^4, \omega)$ of a symplectic surface can be generated by a Hamiltonian isotopy, i.e. $\Sigma_t = \phi_{H_t}^t(\Sigma_0)$. Given an open subset $U \subset X^4$ in which $\Sigma_t \cap U = \Sigma_0 \cap U$ is fixed, we may moreover assume that $H_t|_V = 0$ holds in any relatively compact subset V in U.

It is observed in [DRGI16] that the argument of Siebert-Tian works for symplectic divisors when the symplectic isotopy is constant near the intersection points. Such an isotopy is an instance of a strong *D*-symplectic isotopy of marked symplectic divisors introduced in [LM16].

Definition 2.5. A marked symplectic divisor consists of a 5-tuple

$$\Theta = (X, \omega, D, \{p_j\}, \{I_j\})$$

such that

- $D \subset (X, \omega)$ is an ω -orthogonal divisor,
- p_i , called centers of marking, are points on D (intersection points of D allowed)
- $I_j : (B(\delta_j), \omega_{std}) \to (X, \omega)$, called coordinates of marking, are symplectic embeddings sending the origin to p_j and with $I_j^{-1}(D) = \{z = 0\} \cap B(\delta_j)$ (resp. $I_j^{-1}(D) = (\{z = 0\} \cup \{w = 0\}) \cap B(\delta_j))$ if p_j is a smooth (resp. an intersection) point of D, where (z, w) is the complex coordinate for $B(\delta_j)$. Moreover, we require that the image of I_j are disjoint.

Every ω -orthogonal symplectic divisor can be regarded as a marked divisor where $\{p_j\}$ consists of all intersection points and I_j exists since D is ω -orthogonal. In the rest of this subsection, we will always take an ω -orthogonal symplectic divisor as a marked divisor in such way.

Definition 2.6. Let $\Theta = (X, \omega, D, \{p_j\}, \{I_j\})$ be a marked divisor. A D-symplectic isotopy of Θ is a 4-tuple $(X, \omega_t, D, \{p_j\})$ such that ω_t is a smooth family of cohomologous symlectic forms on X with $\omega_0 = \omega$ and D being ω_t -symplectic for all t.

A strong D-symplectic isotopy is a 5-tuple $(X, \omega_t, D, \{p_j\}, \{I_{j,t}\})$ such that

- the 4-tuple (X, ω_t, D, p_j) is a D-symplectic isotopy of Θ ,
- D is ω_t -orthogonal,
- $I_{j,t}: (B(\epsilon_j), \omega_{std}) \to (X, \omega_t)$ are symplectic embeddings sending the origin to p_j , $I_{j,0} = I_j$ and $I_{j,t}^{-1}(D) = \{z = 0\} \cap B(\epsilon_j)$ (resp. $I_{j,t}^{-1}(D) = (\{z = 0\} \cup \{w = 0\}) \cap B(\epsilon_j))$ if p_j is a smooth point (resp. an intersection point) of D, for some $\epsilon_j < \delta_j$, where (z, w) is the complex coordinate for $B(\delta_j)$.

Lemma 2.7 ([LM16]). If $(X, \omega, D, \{p_j\}, \{I_j\})$ and $(X', \omega', D', \{p'_j\}, \{I'_j\})$ are D-symplectic isotopy, then they are strong D-symplectic isotopy.

In order to fix a neighborhood of the intersections points during the isotopy, we also need the following result which extends a path of symplectic ball embeddings to a Hamiltonian isotopy.

Lemma 2.8 (Theorem 3.3.1 of [MS17]). Let (X, ω) be a closed symplectic manifold. Let $0 \leq \lambda < r$ and let $\psi_0, \psi_1 : B(r) \to X$ be symplectic embeddings that are joined by a smooth family of symplectic embeddings $\psi_t : B(\lambda) \to X$. Then there exists a Hamiltonian isotopy $\{\phi_t\}$ of X and a constant $\lambda < \rho < r$ such that $\phi_0 = id$, $\phi_1 \circ \psi_0|_{B(\rho)} = \psi_1|_{B(\rho)}$ and $\phi_t \circ \psi_0|_{B(\lambda)} = \psi_t$.

Combining the above two lemmas, we can modify the isotopy to the one considered in [DRGI16] and apply the argument of [ST05].

Lemma 2.9. Two strong D-symplectic isotopic ω -orthogonal symplectic divisors are symplectomorphic. In particular, they are Hamiltonian diffeomorphic if $H^1(X; \mathbb{R}) = 0$.

Proof. Let $(X, \omega_t, D, \{p_j\}, \{I_{j,t}\})$ be a strong *D*-symplectic isotopy. By Moser stability, there is an isotopy $\psi_t : X \to X$ such that $\psi_0 = id$ and $\psi_t^* \omega_t = \omega_0$. Let $D_t = \psi_t^{-1}(D)$ be the image of *D* in (X, ω_0) . Denote by $\tilde{I}_{j,t} = \psi_t^{-1} \circ I_{j,t} : B(\epsilon_j) \to (X, \omega_0)$ the family of symplectic ball embeddings such that $(\tilde{I}_{j,t})^{-1}(D_t) = (\{z = 0\} \cup \{w = 0\}) \cap B(\epsilon_j)$. Note that ψ_t is actually Hamiltonian if $H^1(X; \mathbb{R}) = 0$. So it suffices to show (X, ω, D_0) and (X, ω, D_1) are Hamiltonian diffeomorphic.

By Lemma 2.8, there is a Hamiltonian isotopy ϕ_t of (X, ω_0) such that $\phi_t \circ I_{j,0} = I_{j,t}$, after possibly shrinking to a smaller ϵ_j . Replacing D_t by $\phi_t^{-1}(D_t)$, we may assume that $D_t \cap \tilde{I}_{j,0}(B(\epsilon_j))$ is fixed during the isotopy. At a smooth point p on D_{t_0} , choose a complex Darboux coordinate (z = x + iy, w = u + iv). For t close to t_0 , there are functions f_t, g_t such that D_t is the graph $w = f_t(z) + ig_t(z)$. Then following the argument of Siebert-Tian, we define

$$H_t = -(\partial_t g_t) \cdot (u - f_t) + (\partial_t f_t) \cdot (v - g_t),$$

which generates the isotopy. At each intersection point $p_j \in D_{t_0}$, take the Darboux ball $\tilde{I}_{j,0}: B(\epsilon_j) \to (X, \omega_0)$ and define $H_t = 0$ in this chart, which generates the constant isotopy.

Since each H_t vanishes along D_{t_0} , locally defined H_t can be patched via a partition of unity of D_{t_0} to a function defined near D_{t_0} and then extend to M arbitrarily. Finally extend the construction to all $t \in [0, 1]$ via a partition of unity in t.

Proposition 2.10. Two ω -orthogonal symplectic divisors D and D' in (X, ω) are strictly symplectic deformation equivalent if and only if they are symplectomorphic. Moreover, two general symplectic divisors are strictly symplectic deformation equivalent if and only if they are symplectomorphic after a small symplectic isotopy locally supported near the intersection points.

Proof. Up to symplectomorphism, there is a symplectic isotopy D_t with $D_0 = D$ and $D_1 = D'$. Regard D and D' as marked symplectic divisors with markings centered at the intersection points. Extend the isotopy D_t smoothly to an ambient isotopy $\psi_t : X \to X$ so that $\psi_t^{-1}(D_t) = D$ and let $\omega_t = \psi_t^* \omega$. Then $(X, \omega_t, D, \{p_i\})$ is a D-symplectic isotopy. By Lemma 2.7, (X, ω, D) and (X, ω_1, D) are strong *D*-symplectic isotopic and thus symplectomorphic by Lemma 2.9. So *D* is symplectomorphic to *D'* by composing with ψ_1 .

Two general symplectic divisors can be perturbed to be symplectic orthogonal by a symplectic isotopy locally supported near the intersection points (Lemma 2.3 of [Gom95]). After the local perturbation, they are still strictly symplectic deformation equivalent and thus symplectomorphic.

In the holomorphic cateogory, we have the a version of Torelli theorem which basically says two log Calabi-Yau pairs are isomorphic if there is an integral isometry of the second cohomology, which maps the homology of one divisor to the other (Theorem 8.5 of [Fri16]). Here an integral isometry means an automorphism preserving the intersection form. In this subsection, we give a symplectic analogue of this Torelli theorem.

On a closed 4-manifold X, each diffeomorphism induces an automorphism of the lattice of the second integral cohomology, which gives a natural map $\text{Diff}(X) \to \text{Aut}(H^2(X;\mathbb{Z}))$. We denote the image of this map by D(X). This map is not always surjective, i.e. not every integral isometry of the second cohomology is generated by a diffeomorphism. For rational surfaces with Euler characteristic $\chi \leq 12$, every integral isometry is induced from a diffeomorphism ([Wal64]). But this is not true when $\chi > 12$ by [FM88].

Theorem 2.2 can be regarded as a symplectic analogue of the Torelli theorem, but only in the weak sense. Firstly, it only concerns symplectic deformation equivalence as compared to isomorphism in the holomorphic case. Secondly, as discussed above, not every integral isometry of the second cohomology is generated by a diffeomorphism. However, we observed that Theorem 2.2 can be improved to the following much stronger analogue of the Torelli theorem using Proposition 2.10 and a characterization of D(X) in [LL02].

Proposition 2.11 (Symplectic Torelli Theorem). Let D and D' be two symplectic log Calabi-Yau divisors in (X, ω) with component C_i and C'_i respectively, and an integral isometry

$$\gamma: H^2(X; \mathbb{Z}) \to H^2(X; \mathbb{Z})$$

such that $\gamma(PD[C'_i]) = PD[C_i]$, and $\gamma([\omega]) = [\omega]$. Then D and D' are strictly symplectic deformation equivalent. They are symplectomorphic if they are both ω -orthogonal.

Proof. Notice that $[\omega] \cdot c_1(X, \omega) > 0$ by the existence of symplectic log Calabi-Yau divisors. Since $\gamma([\omega]) = [\omega]$ and $\gamma(c_1(X, \omega)) = c_1(X, \omega)$, we have $\gamma \in D(X)$ by Theorem 2.8 (1) of [LL02], i.e. there is a diffeomorphism $\phi : X \to X$ such that $\phi^* = \gamma$. Since ϕ^* preserves the symplectic class $[\omega]$, it must be orientation-preserving and therefore a strict homological equivalence. Then the two divisors are strictly symplectic deformation equivalent by Theorem 2.2. If they are both ω -orthogonal, then they are actually symplectomorphic by Proposition 2.10.

An automorphism of $H^2(X; \mathbb{Z})$ as in Proposition 2.11 is called a **strict** H^2 -**automorphism**. Denote by $p\mathcal{LCY}^{\perp}(X,\omega) \subset p\mathcal{LCY}(X,\omega)$ the subspace of ω -orthogonal symplectic divisors in (X,ω) . Then the two main results of this subsection, Proposition 2.10 and 2.11, give two more equivalent definitions of the isotopy class space of symplectic log Calabi-Yau divisors

$$\begin{aligned} \mathcal{LCY}(X,\omega) &= p\mathcal{LCY}^{\perp}(X,\omega) / \text{Symp}(X,\omega) \\ &= p\mathcal{LCY}(X,\omega) / \text{strict } H^2 \text{-automorphism.} \end{aligned}$$

2.2 Operations and minimal models

In this subsection, we review various facts about the operations and minimal models of symplectic log Calabi-Yau pairs studied in [LM16], which will become useful later.

For a symplectic divisor $D \subset (X, \omega)$, a **toric blow-up** (resp. **non-toric blow-up**) of D is the total (resp. proper) transform of a symplectic blow-up centered at an intersection point (resp. a smooth point) of D. A **toric blow-down** refers to blowing down an exceptional sphere contained in D, intersecting exactly once with exactly two other irreducible components of D. A **non-toric blow-down** refers to blowing down an exceptional sphere not contained in D, intersecting exactly one irreducible component of D. These operations preserve the symplectic log Calabi-Yau condition and have analogues in the holomorphic category.

Remark 2.12. Theorem 2.2 implies the following.

- Let D_1, D_2 be two strictly symplectic deformation equivalent symplectic log Calabi-Yau divisors in (X, ω) such that a nodal point $p_1 \in D_1$ corresponds to $p_2 \in D_2$ and a component $C_1 \subset D_1$ corresponds to $C_2 \subset D_2$. For i = 1, 2, let \tilde{D}_i be the non-toric blow-up of D_i , of the same size, at some point on C_i . Then \tilde{D}_1 and \tilde{D}_2 are strictly symplectic deformation equivalent. The same holds if \tilde{D}_i is toric blow-up of D_i , of the same size, at p_i .
- Similarly, let e₁ be a toric/non-toric exceptional class of D₁ and let D
 ₁ be the blowdown of D₁. Then there is a corresponding toric/non-toric exceptional class e₂ of D₂ under the strict homology equivalence, such that the corresponding blow-down D
 ₂ of D₂ is strictly symplectic deformation equivalent to D
 ₁.

Definition 2.13. A symplectic log Calabi-Yau pair (X, D, ω) is called minimal if (X, ω) is minimal, or (X, D, ω) is a symplectic Looijenga pair with $X = \mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$.

Through a maximal sequence of non-toric blow-downs and then a maximal sequence of toric blow-downs, we get a minimal pair from any symplectic log Calabi-Yau pair (X, D, ω) and is called a minimal model of (X, D, ω) .

We recall here the homology types of minimal symplectic log Calabi-Yau pairs (modulo cyclic symmetry and homological equivalence). The following result stated in [LM16] (but with no detailed account) is obtained by considering the intersection numbers of the components and adjunction formula. We will give a detailed explanation of this in the proof of Lemma 2.19 in the next section.

Theorem 2.14 ([LM16]). Any minimal symplectic log Calabi-Yau pair (X, D, ω) is homological equivalent to one of the following.

• Case (A): X is a symplectic ruled surface with base genus 1. D is a torus.

- Case (B): $X = \mathbb{CP}^2, c_1 = 3h$.
- (B1) D is a torus,
- (B2) D consists of a h-sphere and a 2h-sphere, or
- (B3) D consists of three h-spheres.

• Case (C): $X = S^2 \times S^2$, $c_1 = 2f_1 + 2f_2$, where f_1 and f_2 are the homology classes of the two factors.

(C1) D is a torus. (C2) k(D) = 2 and $[C_1] = bf_1 + f_2, [C_2] = (2 - b)f_1 + f_2.$ (C3) k(D) = 3 and $[C_1] = bf_1 + f_2, [C_2] = f_1, [C_3] = (1 - b)f_1 + f_2.$ (C4) k(D) = 4 and $[C_1] = bf_1 + f_2, [C_2] = f_1, [C_3] = -bf_1 + f_2, [C_4] = f_1.$ The graphs in (C1), (C2), (C3) and (C4) are given respectively by



• Case (D): $X = \mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$, $c_1 = f + 2s$, where f and s are the fiber class and section class with $f \cdot f = 0$, $f \cdot s = 1$ and $s \cdot s = 1$.

(D1) D cannot be a torus because it would not be minimal.

(D2) k(D) = 2, and either $([C_1], [C_2]) = (af + s, (1 - a)f + s)$ or $([C_1], [C_2]) = (2s, f)$.

(D3) k(D) = 3 and $[C_1] = af + s, [C_2] = f, [C_3] = -af + s.$

 $(D4) \ k(D) = 4 \ and \ [C_1] = af + s, [C_2] = f, [C_3] = -(a+1)f + s, [C_4] = f.$

The graphs in (D2), (D3) and (D4) are given respectively by



We now introduce the **smoothing** operation, which corresponds to the nodal trade operation of the almost toric fibrations (see Section 4.4). Geometrically, given $D = C_1 \cup \cdots C_k \cup C_{k+1} \subset (X, \omega)$ in the space of symplectic log Calabi-Yau divisors $p\mathcal{LCY}(X, \omega)$, let p be the intersection point of C_k and C_{k+1} and $U \subset X$ an arbitrarily small neighborhood of p. Then one can always perform the local surgery to get another $D' = C_1 \cup \cdots C_{k-1} \cup C'_k \subset (X, \omega)$ in the space $p\mathcal{LCY}(X, \omega)$ such that outside U, C'_k coincides with $C_k \cup C_{k+1}$ by smoothing the singular point p. Homologically, this operation transfers the homological sequence $([C_1], \cdots, [C_k], [C_{k+1}])$ representing D into $([C_1], \cdots, [C_{k-1}], [C'_k] = [C_k] + [C_{k+1}])$ representing D'. Remember that $\mathcal{LCY}(X, \omega)$ is equal to $p\mathcal{LCY}(X, \omega)$ modulo strict symplectic deformation equivalence and $\widetilde{\mathcal{LCY}}(X)$ is equal to the union of $p\mathcal{LCY}(X, \omega)$ with ω being all the symplectic forms modulo symplectic deformation equivalence. By passing to the quotient space, we can define two elements in the space $\mathcal{LCY}(X, \omega)$ or $\widetilde{\mathcal{LCY}}(X)$ can be related by the smoothing operation if and only if there exists representatives in their equivalent classes which can be related by the smoothing operation. Thus we can view smoothing as the operation on the space $\mathcal{LCY}(X, \omega)$ or $\widetilde{\mathcal{LCY}}(X)$.

For example, consider the symplectic log Calabi-Yau pair $(S^2 \times S^2, C_1 \cup C_2 \cup C_3 \cup C_4, \omega)$ in case (C4) of Theorem 2.14. We can get case (C3)(or (C2)/(C1)) if we perform the smoothing operation at $C_2 \cap C_3$ (or $C_2 \cap C_3, C_3 \cap C_4/C_1 \cap C_2, C_2 \cap C_3, C_3 \cap C_4, C_4 \cap C_1$).

We have the following observation:

Lemma 2.15. Assuming X is rational, then any element in $\widetilde{\mathcal{LCY}}(X)$ can be obtained by performing smoothing operations on some element in $\widetilde{\mathcal{LCY}}(X)$.

Proof. It is easy to see the conclusion holds for the minimal cases. From Theorem 2.14 we see that cases (B3), (C4), (D4) are toric, and all the other cases in (B), (C), (D) can be obtained by smoothing the toric ones. Therefore, by the Lemma 3.4 of [LM16], we only need to show if $(\omega, D) \in \widetilde{\mathcal{LCV}}(X)$ comes from smoothing an element in $\widetilde{\mathcal{LCV}}(X)$, then its toric or non-toric blow-up will come from smoothing some element in $\widetilde{\mathcal{LCV}}(X \# \mathbb{C}P^2)$.

We may assume $D = C_1 \cup \cdots \cup C_{k-1} \cup C_k$ is obtained from smoothing the toric divisor $(A_{11} \cup \cdots \cup A_{1a_1}) \cup \cdots \cup (A_{k1} \cup \cdots \cup A_{ka_k})$ at the intersection points $A_{il} \cap A_{i(l+1)}$ such that $[C_i] = \sum_{p=1}^{a_i} [A_{ip}]$ for all $1 \leq i \leq k$. For the toric or non-toric blow up of D, without loss of generality, suppose their homology sequences are $([C_1] - E, E, [C_2] - E, \cdots, [C_k])$ and $([C_1] - E, [C_2], \cdots, [C_k])$ respectively, where E is the exceptional class representing the generator of $\mathbb{C}P^2$. Then we can perform small toric blow-up on the toric divisor $(A_{11} \cup \cdots \cup A_{1a_1}) \cup \cdots \cup (A_{k1} \cup \cdots \cup A_{ka_k})$ at $A_{1a_1} \cap A_{21}$ to get another toric divisor $(B_{11} \cup \cdots \cup B_{1a_1}) \cup e \cup (B_{21} \cup \cdots \cup B_{2a_2}) \cup \cdots$ on $X \# \mathbb{C}P^2$ whose homological sequence is

$$([A_{11}], \cdots, [A_{1a_1}] - E, E, [A_{21}] - E, \cdots, [A_{k1}], \cdots, [A_{ka_k}])$$

So performing the smoothing operation at the intersection points $B_{il} \cap B_{i(l+1)}$ will lead to the toric blow-up of D and if we continue to do one more smoothing operation at $e \cap B_{21}$ we will get the non-toric blow-up of D.

2.3 $\mathcal{HLCY}(X,\omega)$, realizability and stability

In this section we introduce a more combinatorical description of the isotopy class space of symplectic log Calabi-Yau divisors, which is given by the underlying homological configurations. In the rest of this paper, whenever we talk about symplectic log Calabi-Yau divisors, these two descriptions will be used interchangeably.

Fix a closed oriented 4-manifold X. Let $T = (V, E, \{A_{\alpha}\}_{\alpha \in V})$ be a decorated graph with an underlying finite connected undirected graph (V, E). Then there is the map

$$\phi: E \to \{\{x, y\} | x, y \in V, x \neq y\}$$

Each vertex α is labeled by a homology class $A_{\alpha} \in H_2(X; \mathbb{Z})$, satisfying $A_{\alpha} \cdot A_{\beta} = \#\phi^{-1}(\{\alpha, \beta\})$ for $\alpha \neq \beta$. Such *T* is called a **homological configuration** in *X* and we write $[T] = \sum_{\alpha \in V} A_{\alpha}$. Two homological configurations $(V, E, \{A_{\alpha}\})$ and $(V', E', \{A'_{\beta}\})$ are called equivalent if there is a graph isomorphism $f : (V, E) \to (V', E')$ such that $A_{\alpha} = A_{f(\alpha)}$ for all $\alpha \in V$. We will always be talking about homological configurations up to equivalences. Note that for a fixed symplectic structure ω , we can associate to each vertex v_{α} an ω -symplectic genus

$$g_{\alpha} = 1 + \frac{1}{2}(A_{\alpha}^2 - c_1(X, \omega) \cdot A_{\alpha}),$$

which we also take as part of T. When the decorated graph T is a cycle, namely $V = \{1, \dots, l\}, E = \{e_1, \dots, e_l\}$ with $l \ge 2$, $\phi(e_l) = \{1, l\}, \phi(e_i) = \{i, i+1\}$ for $1 \le i \le l-1$ or $V = \{1\}, E = \emptyset$, we call it a **cyclic homological configuration**. We usually denote a cyclic homological configuration by a sequence of homology classes $T = (A_1, \dots, A_l)$ and (anti)cyclic permutations of such sequences produce equivalent cyclic homological configurations.

Definition 2.16. For any symplectic rational manifold (X, ω) , we define the space of **pre**homological log Calabi-Yau divisor $p\mathcal{HLCY}(X, \omega)$ to be the set of equivalent classes of cyclic homological configurations $T = (V, E, \{A_1, \dots, A_l\})$ satisfying that:

- Each $A_i \in H_2(X; \mathbb{Z})$ has an ω -symplectic surface representative;
- $[T] = PD(c_1(X, \omega))$

There is a natural action of the the symplectomorphism group $Symp(X,\omega)$ on $p\mathcal{HLCY}(X,\omega)$, factoring through D(X). And the space of **homological log Calabi-Yau divisor** $\mathcal{HLCY}(X,\omega)$ is defined to be $p\mathcal{HLCY}(X,\omega)/Symp(X,\omega)$.

Remark 2.17. If ω and ω' are two symplectic forms on X with the same class $[\omega] = [\omega']$, the set of homology classes which can be represented by ω -symplectic surfaces and ω' -symplectic surfaces are the same ([ALLP19]). Therefore we can further define $\mathcal{PHLCY}(X, u)$ to be any $\mathcal{PHLCY}(X, \omega)$ with $u = [\omega]$. Moreover, from [LW12] we know that the homological action $Symp(X, \omega)/Symp_h(X, \omega) = D_{K_{\omega}, [\omega]}$ by the symplectomorphism group also only depends on the symplectic class $[\omega]$. As a result, we could also define $\mathcal{HLCY}(X, u) = \mathcal{HLCY}(X, \omega)$ for any $u = [\omega]$.

Our goal in this section is to identify the geometric object, the isotopy class space $\mathcal{LCY}(X, \omega)$ with the homological object, the space $\mathcal{HLCY}(X, \omega)$. There is a natural map

$$F: \mathcal{LCY}(X,\omega) \to \mathcal{HLCY}(X,\omega),$$

which sends the geometric divisor $D = \bigcup_{i=1}^{k} C_i$ to its homological configuration $T = ([C_1], \dots, [C_k])$, and we say T has a **realization** by D. We would often write (D) instead of T when it is the underlying homological configuration of the divisor D. Obviously the homological configurations obtained by the image of F satisfy all the conditions in Definition 2.16. We will prove F is a bijection in the rest of this section.

Lemma 2.18. F is well-defined and injective.

Proof. If two geometric divisors are strictly deformation equivalent, then up to a symplectomorphism, they will be isotopic and thus have the same homological configurations. Now since we have taken the quotient of $\text{Symp}(X, \omega)$ in the definition of $\mathcal{HLCY}(X, \omega)$, there is no ambiguity of the image of F. We see that F is well-defined.

If we have two geometric divisors $D = \bigcup_{i=1}^{k} C_i$ and $D' = \bigcup_{i=1}^{k} C'_i$ in X whose image under F are the same, then by definition, we have a symplectomorphism $h \in \text{Symp}(X, \omega)$ such that $h_*([C_i]) = [C'_i]$ for all $1 \leq i \leq k$. This implies (X, ω, D) and (X, ω, D') are strictly homologically equivalent and thus will be identified in the space $\mathcal{LCY}(X, \omega)$ according to our definition. This shows that F is injective.

The proof of surjectivity requires more work. Firstly we prove it for minimal models, that is, when $X = \mathbb{CP}^2$, $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$ or $S^2 \times S^2$. We use the standard basis $\{H\}$, $\{H, E\}$, $\{F, B\}$ of their second homology groups. Note that it's a classical result that any symplectic forms can be diffeomorphic to the ones satisfying $\omega(H) > 0$, $\omega(H) > \omega(E) > 0$, $\omega(F) \ge \omega(B) > 0$ and the first Chern classes are 3H, 3H - E, 2F + 2B. Therefore we may make these assumptions in the following lemma. **Lemma 2.19.** For $X = \mathbb{CP}^2$ with symplectic form ω such that $\omega(H) > 0$ and $c_1(X, \omega) = 3H$,

$$p\mathcal{HLCY}(X,\omega) = \mathcal{HLCY}(X,\omega) = \{(3H), (2H,H), (H,H,H)\}$$

For $X = \mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$ with symplectic form ω such that $\omega(H) > \omega(E) > 0$ and $c_1(X, \omega) = 3H - E$, elements of $p \mathcal{HLCY}(X, \omega)$ are of the following forms:

- 1. (3H E)
- 2. $(2H, H E), ((a + 1)H aE, (-a + 2)H + (a 1)E)_{a \in \mathbb{Z}_+}$
- 3. $(aH + (-a + 1)E, H E, (-a + 2)H + (a 1)E)_{a \in \mathbb{Z}_+}$
- 4. $(aH + (-a + 1)E, H E, (-a + 1)H + aE, H E)_{a \in \mathbb{Z}_+}$

For $X = S^2 \times S^2$ with symplectic form ω such that $\omega(F) \ge \omega(B) > 0$ and $c_1(X, \omega) = 2B + 2F$, elements of $p\mathcal{HLCY}(X, \omega)$ are of the following forms:

- 1. (2F + 2B)
- 2. $(2F+B,B), ((F+bB,F+(2-b)B)_{b\in\mathbb{Z}_+})$
- 3. $(F + bB, B, F + (1 b)B)_{b \in \mathbb{Z}_+}$
- 4. $(F + (b-1)B, B, F + (1-b)B, B)_{b \in \mathbb{Z}_+}$

As a result, F is surjective when X is one of these three manifolds.

Proof. The case when l = 1 is always trivial, next we will assume $l \ge 2$. Note that when $l \ge 2$, by condition, $A_i \cdot (c_1 - A_i) = A_i \cdot (A_1 + \cdots + A_{i-1} + A_{i+1} + \cdots + A_l) = 2$. So by adjunction we see that all the classes are actually represented by symplectic surfaces of genus $g_i = 0$.

For \mathbb{CP}^2 , the result is obvious since we know the classes of symplectic surfaces must be kH with positive k.

For $X = \mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$, suppose A = xH + yE with $A^2 = t$ has a symplectic sphere representative, then by adjunction formula we have that

$$\begin{cases} x^2 - y^2 = t \\ t = 3x + y - 2 \end{cases}$$

It follows that $\begin{cases} x = \frac{t+1}{2} \\ y = \frac{-t+1}{2} \end{cases}$ or $\begin{cases} x = \frac{t}{4} + 1 \\ y = \frac{t}{4} - 1 \end{cases}$. In the former case we get A = aH + (1-a)E

with odd self intersection 2a - 1 = t, and in the later cases we get A = (k + 1)H + (k - 1)E with even self intersection 4k = t. Note that the intersection between two odd classes $a_1H + (1 - a_1)H$ and $a_2H + (1 - a_2)E$ is $a_1 + a_2 - 1$, the intersection between two even classes $(k_1 + 1)H + (k_1 - 1)E$ and $(k_2 + 1)H + (k_2 - 1)E$ is $2(k_1 + k_2)$, and the intersection between an odd class aH + (1 - a)E and an even class (k + 1)H + (k - 1)E is 2ak - k + 1. Using the fact that the homological configuration is cyclic, we have some simple observations.

• When l > 3, A_i , A_{i+1} and A_{i+2} can not all be odd classes.

- When l > 2, if A_i is an even class, A_{i-1}, A_{i+1} must be odd classes and $A_i = H E$.
- If $A_i = H E$, all the other A_j with $j \neq i + 1, i 1$ must also be H E.

Now we could easily exclude the cases when the length l > 4 by considering the arrangement of odd classes and even classes according to the first two bullets above. Then it's possible to list all the homological configurations for $l \leq 4$ which will coincide with our statement.

The analysis for $S^2 \times S^2$ is similar to $\mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$. By adjunction formula, if xF + yB has a self intersection *t*-spherical symplectic representative, then we have $\begin{cases} 2xy = t \\ t = 2x + 2y - 2 \end{cases}$.

Since we made the assumption $\omega(F) \ge \omega(B) \ge 0$ it follows that all the possible classes are either F + bB for arbitrary b or bF + B for non negative b. Note that the intersection between $F + b_1B$ and $F + b_2B$ or $b_1F + B$ and $b_2F + B$ is $b_1 + b_2$, and the intersection between $F + b_1B$ and $b_2F + B$ is $1 + b_1b_2$. We then see that:

- When l > 3, if $A_i = bF + B$, then $b \le 1$ and all the other A_j with $j \ne i + 1, i 1$ must be B if b = 0 or F B if b = 1
- If $A_i = F + b_1 B$, $A_{i+1} = F + b_2 B$, $A_{i+2} = F + b_3 B$, then l > 3 and $b_1 = b_3 = 0$, $b_2 = 1$.
- If $A_i = B$ or F, then all the other A_j with $j \neq i + 1, i 1$ must be B or F.

Again, the above observations are enough for us to exclude l > 4 and list all the possible cases for $l \leq 4$.

Note that the homological configurations (H, H, H), (aH + (-a + 1)E, H - E, (-a + 1)H + aE, H - E), (F + (b - 1)B, B, F + (1 - b)B, B) can be realized as the boundary divisors of the toric actions (moment map's preimage of the boundary of Delzant polygon). And all the other homological configurations come from the smoothings of the toric ones. Therefore the claim of F is surjective has been verified for these three cases.

We need some results before proving the surjectivity for all rational manifolds. A symplectic form ω on X is called T-positive if $\omega(A_{\alpha}) > 0$ for all α . The following result is crucial for our proof of realizability.

Lemma 2.20 ([DLW18]). Let (X, ω) be a symplectic manifold with $b^+ = 1$ and T a homological configuration realized by a symplectic divisor D. Then D is ω -stable, that is, for any T-positive symplectic form ω' deformation equivalent to ω , there is a symplectic divisor D' realizing T with respect to ω' .

Lemma 2.21. Suppose (X, ω) is the symplectic blow up of (X', ω') with the canonical identification $H_2(X, \mathbb{Z}) = H_2(X', \mathbb{Z}) \oplus \mathbb{Z}E$ and the homological configuration $(A_1, \dots, A_l) \in \mathcal{HLCY}(X', \omega')$ has a realization. Then if $(A_1, \dots, A_i - E, \dots, A_l)$ or $(A_1, \dots, A_i - E, E, A_{i+1} - E, \dots, A_l)$ is in $\mathcal{HLCY}(X, \omega)$, they also have a realization.

Proof. For the realization (X', ω', D') of (A_1, \dots, A_l) we could always perform small blow up to obtain (X'', ω'', D'') realizing $(A_1, \dots, A_i - E', \dots, A_l)$ or $(A_1, \dots, A_i - E', E', A_{i+1} - E', \dots, A_l)$. Here we identify $H_2(X'', \mathbb{Z}) = H_2(X', \mathbb{Z}) \oplus \mathbb{Z}E'$. Choose a diffeomorphism $f : X \to X''$ such that $f_*(E) = E'$ and $f_*|_{H_2(X',\mathbb{Z})}$ is identity. We see that $(A_1, \dots, A_i - E, \dots, A_l)$ or $(A_1, \dots, A_i - E, E, A_{i+1} - E, \dots, A_l)$ has a realization under the symplectic form $(f^{-1})^*\omega''$. It is well known (for example, Theorem 3.5 and Theorem 3.18 of [Li08]) that the diffeomorphism group acts transitively on the space of symplectic forms in a fixed class and any two symplectic forms on a rational manifold are deformation equivalent up to a diffeomorphism fixing the symplectic class. We can pick $g \in \text{Diff}^+(X)$ such that $g^*(f^{-1})^*\omega''$ is deformation equivalent to ω . Since $g_* \in D_{K_{(f^{-1})}*\omega''}, [(f^{-1})*\omega''] \cong \text{Symp}(X, (f^{-1})^*\omega'')/Symp_h(X, (f^{-1})^*\omega''),$ we could pick $h \in \text{Symp}(X, (f^{-1})^*\omega'')$ such that h_*g_* is identity. It then follows that

$$(A_1, \cdots, A_i - E, \cdots, A_l) = h_* g_*(A_1, \cdots, A_i - E, \cdots, A_l) \text{ or}$$
$$(A_1, \cdots, A_i - E, E, A_{i+1} - E, \cdots, A_l) = h_* g_*(A_1, \cdots, A_i - E, E, A_{i+1} - E, \cdots, A_l)$$

has a realization under the form $g^*((f^{-1})^*\omega'')$. Of course the assumption that the blown-up configuration is in $\mathcal{HLCY}(X,\omega)$ guarantees the ω -positivity. By lemma 2.20 we see that it then also has a realization under the form ω .

Remark 2.22. Note that one can run the above argument for a single class to get the following useful statement that will be used later: with the same assumption in Lemma 2.21, if A can be represented by ω' -symplectic surface and A - E has positive ω -area, then it can be represented by ω -symplectic surface.

As we want to prove by induction, the following useful result will enable us to perform the blow down process which reduces the arbitrary case to the case of minimal models.

Lemma 2.23 ([Pin08]). Let (X, ω) be a symplectic manifold with $X \neq \mathbb{CP}^2 \# \overline{\mathbb{CP}^2}$, then for any ω -tame almost complex structure J, all symplectic exceptional classes of minimal area have J-holomorphic representatives.

Proposition 2.24. F is surjective for all (X, ω) . Therefore $\mathcal{LCY}(X, \omega) = \mathcal{HLCY}(X, \omega)$.

Proof. Suppose $X = \mathbb{CP}^2 \# n \overline{\mathbb{CP}^2}$ with $n \ge 2$ and let E be the minimal area symplectic excetional class. For $(A_1, \dots, A_l) \in \mathcal{HLCY}(X, \omega)$, the case when l = 1 is trivial and we firstly assume $l \ge 3$. By adjunction formula, $E \cdot (A_1 + \dots + A_l) = 1$. Since each A_i has a symplectic representative and thus a *J*-holomorphic representative for some tame *J*, by Lemma 2.23, we have $E \cdot A_i \ge 0$ or $A_i = cE$ for some positive integer *c*. Note that the symplectic genus of A_i being non-negative forces c = 1. Thus either $E \cdot A_i = 1$ for some *i* and $E \cdot A_j = 0$ for all $j \ne i$, or $E = A_i < 0$ for some *i*.

In the former case, we could pick a tame J such that the symplectic surface representing A_i is J-holomorphic. By lemma 2.23, E also has a J-holomorphic representative. By positivity of intersection, they intersect transversally at one point. Thus if we blow down the exceptional sphere C_i representing E, we get another symplectic manifold (X', ω') with $A_i - E$ having ω' representative. For other A_j , one can also find exceptional sphere C_j representing E such that they don't intersect. Since all symplectic exceptional spheres are symplectic isotopic, and there exists an ambient symplectic isotopy inducing that, there must be a representative of A_j not intersecting C_i . Thus A_j also have ω' representative in X'. Now we see that $(A_1, \dots, A_i - E, \dots, A_l) \in \mathcal{HLCY}(X', \omega')$. By induction and lemma 2.21, (A_1, \dots, A_l) has a realization.

For the later case, we have $A_{i-1} \cdot E = A_{i+1} \cdot E = 1$ and $A_j \cdot E = 0$ for $j \neq i-1, i, i+1$. Now by a similar blow down argument as above, $(A_1, \dots, A_{i-1} + E, A_{i+1} + E, \dots, A_l) \in \mathcal{HLCY}(X', \omega')$ for some blow down manifold (X', ω') . Again, the result follows from induction and lemma 2.21.

Now assume l = 2, the homological configuration becomes (K - E, E). Since $n \ge 2$, we could always find another exceptional class E' such that $E' \cdot E = 0$ and thus $E' \cdot (K - E) = 1$. By Theorem 1.2.7 of [MO15], there exists some tame almost complex structure J such that both E' and K - E has J-holomorphic representative. By positivity of intersection, these two curves meet transversally at one point. If we blow down the E' curve to get the manifold (X', ω') , it follows that $(K - E - E', E) \in p\mathcal{HLCY}(X', \omega')$. Now the result also follows from lemma 2.21.

From Proposition 2.24 we also see that $\mathcal{LCY}(X,\omega)$ should be controlled by the set of symplectic sphere classes in (X, ω) , which leads to the stability result (Theorem 1.6).

Proof of Theorem 1.6. By Proposition 2.24, it suffices to show $\mathcal{HLCY}(X,\omega) = \mathcal{HLCY}(X,\omega')$. Note that by Proposition 4.1 of [LL01] if $\mathcal{S}_{\omega}^{-1} = \mathcal{S}_{\omega'}^{-1}$ we must have $K_{\omega} = K_{\omega'} = K$. For any homological configuration $T = (A_1, \ldots, A_k) \in p\mathcal{HLCY}(X, \omega)$, we would have $\sum A_i =$ $c_1(X,\omega) = c_1(X,\omega')$ and each $A_i \in \mathcal{S}_{\omega} = \mathcal{S}_{\omega'}$. So $T \in p\mathcal{HLCY}(X,\omega')$ and $p\mathcal{HLCY}(X,\omega) = c_1(X,\omega')$ $p\mathcal{HLCY}(X,\omega') \subset H_2(X;\mathbb{Z})$. By Proposition 4.1 of [LW12], $D_{K,\omega}$ is generated by reflections along classes in $\mathcal{L}_{K,\omega}$. Note that

$$\mathcal{L}_{K,\omega} \bigsqcup \mathcal{S}_{\omega}^{-2} \bigsqcup (-\mathcal{S}_{\omega}^{-2}) = \mathcal{L}_{K,\omega'} \bigsqcup \mathcal{S}_{\omega'}^{-2} \bigsqcup (-\mathcal{S}_{\omega'}^{-2}),$$
$$\mathcal{L}_{K,\omega'} \text{ and thus } \mathcal{HLCY}(X,\omega) = \mathcal{HLCY}(X,\omega').$$

we must have $\mathcal{L}_{K,\omega} = \mathcal{L}_{K,\omega'}$ and thus $\mathcal{HLCY}(X,\omega) = \mathcal{HLCY}(X,\omega')$.

2.4**Finiteness**

In the holomorphic category, we have the following finiteness result.

Theorem 2.25 ([Fri16]). There are only finitely many deformation types of holomorphic log Calabi-Yau pairs (Y, D) with the same self-intersection sequence.

Since there is always a Kahler pair in a symplectic deformation class of symplectic log Calabi-Yau pairs ([LM16]), we have the following analogous finiteness result in the symplectic category.

Theorem 2.26 ([LMM20]). There are only finitely many symplectic deformation types of symplectic log Calabi-Yau pairs with the same self-intersection sequence.

Now we show that such finiteness also holds for strict symplectic deformation classes. We need to make a digression first.

Having established the equivalence between $\mathcal{LCY}(X,\omega)$ and $\mathcal{HLCY}(X,\omega)$, we also want to describe it in a more convenient way. The finiteness of $\mathcal{LCY}(X,\omega)$ will be shown by that description in this section. In order to do that we need to introduce the concepts of reduced cones and blowup forms for rational manifolds. Denote by M_l the *l*-fold blow-up $\mathbb{CP}^2 \# l\overline{\mathbb{CP}}^2$. Fix a standard basis $\{H, E_1, \ldots, E_l\}$ of $H_2(M_l; \mathbb{Z})$. A class $\lambda H - \sum \delta_i E_i$ is called **reduced** if

$$\delta_1 \ge \delta_2 \ge \cdots \ge \delta_k > 0 \quad and \quad \lambda \ge \delta_1 + \delta_2 + \delta_3.$$

We also say a second cohomology class is reduced if its Poincare dual is reduced.

Definition 2.27. A symplectic form ω is reduced if $[\omega]$ is reduced, in which case $[\omega]$ is called a reduced symplectic class and $(\lambda, \delta_1, \ldots, \delta_k)$ is called the reduced vector of $[\omega]$. The space $\tilde{P}_l = \tilde{P}(M_l)$ of reduced symplectic classes is called the reduced symplectic cone. The normalized reduced symplectic cone $P_l = P(M_l)$ is the space of reduced symplectic classes on M_l with $\lambda = 1$.

For $M'_1 = S^2 \times S^2$, let $B = [S^2 \times pt]$, $F = [pt \times S^2]$. The reduced symplectic classes are those of the forms satisfying $\omega(F) = \lambda, \omega(B) = \mu$ with $\lambda \ge \mu > 0$ and it is normalized if $\lambda = 1$. The crucial fact we need about reduced symplectic cones is that, for symplectic rational surfaces, the reduced symplectic cone is the fundamental domain of the action of orientation-preserving diffeomorphism on the K_0 -symplectic cone ([Tau95], [LL95]). This means whenever we have a symplectic rational manifold (X, ω) in hand, then up to scaling, we may always assume it is $(M_l, \omega_{\delta_1, \dots, \delta_l})$ with reduced or (M'_1, ω'_{μ}) .

A blowup form on M_l is a symplectic form for which there exist pairwise disjoint embedded symplectic spheres in the class H, E_1, \dots, E_l . The result in [KK17] shows that every normalized reduced class encoded by $(\delta_1, \dots, \delta_l)$ with $1 > \delta_1^2 + \dots + \delta_l^2$ contains a blowup form. Therefore, given $(M_l, \omega_{\delta_1, \dots, \delta_l})$ with some reduced blowup form, we could blow down those disjoint exceptional spheres to obtain a sequence of symplectic manifolds $(M_{l-1}, \omega_{\delta_1, \dots, \delta_{l-1}}), \dots, (M_1, \omega_{\delta_1})$, all of which are equipped with reduced blowup forms. Note that in $(M_2, \omega_{\delta_1, \delta_2}), H - E_1 - E_2$ also has exceptional spherical representative, whose blow down manifold is $(M'_1, \omega'_{1-\delta_1})$ after scaling. Note that in this case the inclusion $H_2(M'_1, \mathbb{Z}) \hookrightarrow$ $H_2(M_2, \mathbb{Z})$ sends F to $H - E_2$ and B to $H - E_1$ so that $H_2(M_2, \mathbb{Z}) = H_2(M'_1, \mathbb{Z}) \oplus \mathbb{Z} \langle H - E_1 - E_2 \rangle$.

Remark 2.28. Since any symplectic form is diffeomorphic to some blowup form and the spaces of divisors (both \mathcal{LCY} and \mathcal{HLCY}) for diffeomorphic symplectic forms can be identified naturally, our strategy for proving the finiteness in this section and counting results later is by only considering the spaces under reduced blowup forms.

Now we need the following results:

Lemma 2.29. ([KK17]) If $l \ge 3$, E_l has the minimal area among all symplectic exceptional classes.

Lemma 2.30. ([Zha17]) For any tame J on M_2 , if E_2 has no J-holomorphic representative, then E_1 and $H - E_1 - E_2$ must have J-holomorphic representatives.

Using the above lemmas, by the same argument in Proposition 2.24 it can be shown that:

Lemma 2.31. Assume $\omega_{\delta_1,\dots,\delta_l}$ is a reduced blowup form on M_l . Suppose $(A_1,\dots,A_k) \in p\mathcal{HLCY}(M_l,\omega_{\delta_1,\dots,\delta_l})$, when $k \ge 3, l \ge 2$, one of the followings occurs:

- There is exactly one $1 \leq i \leq k$ such that $A_i \notin H_2(M_{l-1}; \mathbb{Z})$, in which case $A_i + E_l \in H_2(M_{l-1}; \mathbb{Z})$ and $(A_1, \dots, A_{i-1}, A_i + E_l, A_{i+1}, \dots, A_k) \in p\mathcal{HLCY}(M_{l-1}, \omega_{\delta_1, \dots, \delta_{l-1}});$
- There is exactly one $1 \leq i \leq k$ such that $A_i = E_l$, in which case $A_j + E_l \in H_2(M_{l-1}; \mathbb{Z})$ if $|i-j| \equiv 1 \pmod{k}, A_j \in H_2(M_{l-1}; \mathbb{Z})$ for all the other $j \neq i$ and $(A_1, \cdots, A_{i-2}, A_{i-1} + E_l, A_i + E_l, A_{i+1}, \cdots, A_k) \in p\mathcal{HLCY}(M_{l-1}, \omega_{\delta_1, \cdots, \delta_{l-1}}).$

When k = 2, if none of the above cases happens, $(A_1, A_2) = (3H - E_1 - \cdots - E_{l-1} - 2E_l, E_l)$.

Proof. By lemma 2.29, unless l = 2 and E_2 is not of minimal area, E_l can always be represented by *J*-holomorphic sphere for any tame almost complex structure *J* due to lemma 2.23, in which case the result will follow from the argument in Proposition 2.24. So we only need to consider the case when l = 2 and E_2 is not of minimal area.

If E_2 is in one of the component of (A_1, \dots, A_k) , it will fall into the second bullet without any pseudoholomorphic argument. So we further assume there is no A_i equal to E_2 . We want to show any component $A_i = aH - bE_1 - cE_2$ has non negative intersection with E_2 so that we can apply Theorem 1.2.7 of [MO15] to find tame J_i such that E_2, A_i both have J_i -holomorphic representatives. If $A_i^2 < 0$, according to the classification of negative symplectic sphere classes in M_2 in Proposition 3.4 of [LL20] (See also Proposition 3.4),it's immediate that $E_2 \cdot A_i \ge 0$. So next we assume $A_i^2 \ge 0$. Firstly we pick an arbitrary tame J such that A_i can be represented by J-holomorphic sphere. By lemma 2.30, if E_2 has no J-holomorphic representative, then $E_1, H - E_1 - E_2$ must have. By positivity of intersection and the assumption that $A_i^2 \ge 0$, it follows that $E_1 \cdot A_i \ge 0$ and $(H - E_1 - E_2) \cdot A_i \ge 0$ for arbitrary i. Since $E_1 \cdot (A_1 + \dots + A_k) = (H - E_1 - E_2) \cdot (A_1 + \dots + A_k) = 1$ by adjunction, we can conclude that $1 \ge E_1 \cdot A_i \ge 0$ and $1 \ge (H - E_1 - E_2) \cdot A_i \ge 0$. Now we have $1 \ge b \ge 0$, $1 \ge a - b - c \ge 0$ and $3a - b - c \ge 2$ by adjunction. Now it's easy to see these three inequalities imply $c \ge 0$.

Therefore, even if E_2 is not of minimal area, we are still able to find J_i such that E_2, A_i both have J_i -holomorphic representatives. And we can use the argument in Proposition 2.24 to get the conclusion.

With the help of this lemma and the list in Lemma 2.19, we can write down all the homology classes ingredients for homological log Calabi-Yau divisors under the reduced blowup form on M_l with pairwise disjoint symplectic spherical classes H, E_1, \dots, E_l :

The homology classes appearing in the homological log Calabi-Yau divisors must belong to

$$\mathcal{H}_{l} = \begin{cases} kH - (k-1)E_{1} - \sum_{i=2}^{l} \epsilon_{i}E_{i}, k \in \mathbb{Z}, \epsilon_{i} \in \{0,1\} \\ H - E_{1} - \sum_{i=2}^{l} \epsilon_{i}E_{i}, \epsilon_{i} \in \{0,1\} \\ E_{p} - \sum_{i=p+1}^{l} \epsilon_{i}E_{i}, 2 \leq p \leq l, \epsilon_{i} \in \{0,1\} \\ 2H - \sum_{i=2}^{l} \epsilon_{i}E_{i}, \epsilon_{i} \in \{0,1\} \\ 3H - E_{1} - \dots - E_{l} \\ 3H - E_{1} - \dots - E_{p-1} - 2E_{p} - \sum_{i=p+1}^{l} \epsilon_{i}E_{i}, 2 \leq p \leq l, \epsilon_{i} \in \{0,1\} \end{cases}$$

The homology classes appearing in the toric homological log Calabi-Yau divisors must belong to

$$t\mathcal{H}_{l} = \begin{cases} kH - (k-1)E_{1} - \sum_{i=2}^{l} \epsilon_{i}E_{i}, k \in \mathbb{Z}, \epsilon_{i} \in \{0,1\} \\ H - E_{1} - \sum_{i=2}^{l} \epsilon_{i}E_{i}, \epsilon_{i} \in \{0,1\} \\ E_{p} - \sum_{i=p+1}^{l} \epsilon_{i}E_{i}, 2 \leq p \leq l, \epsilon_{i} \in \{0,1\} \end{cases}$$

Moreover, by Remark 2.22 and induction, we see that $A \in \mathcal{H}_l$ can be represented by a symplectic surface if and only if $\omega_{\delta_1,\dots,\delta_l}(A) > 0$. The immediate corollaries are the followings:

Corollary 2.32. For each symplectic rational surface (X, ω) with $c_1(X, \omega) \cdot [\omega] > 0$, there is a unique elliptic symplectic log Calabi-Yau divisor in $\mathcal{LCY}(X, \omega)$.

Corollary 2.33. Under the reduced blowup form ω on M_l with pairwise disjoint symplectic spherical classes H, E_1, \dots, E_l , the cyclic homological configuration T given by (A_1, \dots, A_k) is in $\mathcal{HLCY}(M_l, \omega)$ if and only if:

- $A_i \in \mathcal{H}_l$, for all $1 \leq i \leq k$;
- $[T] = PD(c_1(M_l, \omega));$
- ω is *T*-positive.

Corollary 2.34. $\mathcal{LCY}(X, \omega)$ is finite.

Proof. Since the length of the divisors in $\mathcal{LCY}(X, \omega)$ has an upper bound, it's enough to show for a fixed ω , only fintely many classes $A \in \mathcal{H}_l$ are in the homological configurations. This is implied by the obvious inequality $\omega \cdot c_1(X, \omega) \ge \omega(A) > 0$, thus a bound of k.

2.5 Tautness and toric symplectic log Calabi-Yau divisors

In the holomorphic category, a labeled anticanonical pair is called taut if its isomorphism class is determined by the self-intersection sequence. The tautness of holomorphic anticanonical pairs is used to prove Torelli Theorem in [Fri16]. Now we introduce two kinds of symplectic analogy of tautness as follows. A label on a symplectic log Calabi-Yau divisor $D \in p\mathcal{LCY}(X, \omega)$ is a bijection from $\{1, \dots, k\}$ to the irreducible components of D such that each two consecutive numbers are mapped to two components which intersect. For simplicity, we will just use (C_1, \dots, C_k) to denote a labeled symplectic log Calabi-Yau divisor. Two labeled divisors $(D = (C_1, \dots, C_k), \omega)$ and $(D' = (C'_1, \dots, C'_k), \omega')$ with the same number of components are symplectic deformation equivalent if they are connected by a family of symplectic divisors $(D^t = (C_1^t, \dots, C_k^t), \omega^t)$, up to an orientation-preserving diffeomorphism. Clearly, two unlabeled divisors are symplectic deformation equivalent in the sense of Definition 2.1 if and only if there exist labels on them such that they are symplectic deformation equivalent as labeled divisors. Similarly, we can also define the strict deformation equivalence for labeled divisors.

Definition 2.35. A labeled symplectic log Calabi-Yau divisor $(D = (C_1, \dots, C_k), \omega)$ on X is called **def-taut** if all the labeled symplectic log Calabi-Yau divisors $(D' = (C'_1, \dots, C'_k), \omega')$ on X with $[C_i]^2 = [C'_i]^2$ for all $1 \le i \le k$ are symplectic deformation equivalent to D. It's called **iso-taut** if all the labeled symplectic log Calabi-Yau divisors $(D' = (C'_1, \dots, C'_k), \omega')$ on X with $[C_i]^2 = [C'_i]^2, \omega(C_i) = \omega'(C'_i)$ for all $1 \le i \le k$ are strictly symplectic deformation equivalent to D.

Next we will focus on the iso-tautness which will be used in the proof of Proposition 4.4. We put the discussion of def-tautness into the appendix since there is no direct application in the remaining of this paper.

Recall that a symplectic Looijenga pair (X, ω, D) is called a **toric symplectic log Calabi-**Yau pair if q(D) = 0, where $q(D) = 12 - k(D) - D^2 = 12 - 3k - \sum_{i=1}^{k} s_i$. When we fix the ambient symplectic manifold (X, ω) , we also call D a **toric symplectic log Calabi-Yau** divisor. From the definition we see that being toric only depends on the self-intersection sequence s. In the following we discuss some basic properties of toric symplectic log Calabi-Yau divisors.

The following lemma shows that this definition of being toric coincides with that in [Fri16].

Lemma 2.36. A symplectic log Calabi-Yau divisor $D \subset (X, \omega)$ is toric if and only if it is an iterated toric blow-up of toric symplectic log Calabi-Yau divisors in \mathbb{CP}^2 , $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ and $S^2 \times S^2$.

Proof. The 'if' part is obvious since q(D) is preserved under toric blow-ups and q(D) = 0 for minimal models B3, C4 and D4. Now suppose D is toric. By minimal reduction ([LM16]), we can do a maximal sequence of non-toric blow-downs and then a maximal sequence of toric blow-downs to get a minimal model (X', ω', D') . Note that non-toric blow-down decreases q(D), we have $q(D') \leq q(D) = 0$. By checking the minimal models in Theorem 2.14, we see that all of them have $q(D') \geq 0$. So we must have q(D') = 0 and D is a toric blow-up of D'. The only possible D' are B3, C4 and D4, which are toric ones among minimal models.

The above lemma implies that a Kahler log Calabi-Yau pair (X, ω, D) is a toric symplectic log Calabi-Yau pair if and only if (X, D) is a toric anticanonical pair in the holomorphic category.

A toric anticanonical divisor is taut while a general anticanonical divisor might not be taut in the holomorphic category ([Fri16]). Such rigidity of toric divisors persists to hold in the symplectic category.

Lemma 2.37. Any labeled toric symplectic log Calabi-Yau divisor is both def-taut and isotaut.

Proof. Let $D, D' \subset (X, \omega)$ be two toric symplectic log Calabi-Yau divisors with the same selfintersection sequence s(D) = s(D'). Then there are two Kahler log Calabi-Yau divisors $(\overline{\omega}, \overline{D})$ and $(\overline{\omega}', \overline{D}')$ in X symplectic deformation equivalent to (ω, D) and (ω, D') respectively by Theorem 2.2. The toric Kahler log Calabi-Yau divisors are taut (in the holomorphic sense) by Lemma 2.15 of [Fri16] and there is an isomorphism $\phi : (X, \overline{D}) \to (X, \overline{D}')$. The isomorphism ϕ is in particular an orientation-preserving diffeomorphism and thus a homological equivalence. So (X, ω, D) is also def-taut as a symplectic Calabi-Yau pair.

Suppose (X, ω, D) and (X', ω', D') satisfy (s, a) = (s', a'), then by above discussion there is a diffeomorphism $\Phi : X \to X'$ such that $\Phi_*[C_i] = [C'_i]$. Then we have $\Phi^*[\omega'] \cdot [C_i] = \Phi^*[\omega'] \cdot (\Phi_*)^{-1}[C'_i] = [\omega'] \cdot [C'_i] = \sum a'_i = \sum a_i = [\omega] \cdot [C_i]$, i.e $(\Phi^*[\omega'] - [\omega]) \cdot [C_i]$ for all *i*. Then $\Phi^*[\omega'] = [\omega]$ and Φ is a strictly homological equivalence, which follows from the fact that $\{[C_i]\}$ contains a basis of $H_2(X;\mathbb{Z})$.

We include a proof of this fact for completeness. For each of the minimal models B3, C4 and D4, we can see that the homology classes of the components generate the second homology. We proceed by induction and assume that $\{[C_i]\}$ generates $H_2(X;\mathbb{Z})$ for every toric symplectic log Calabi-Yau pair (X, ω, D) . Denote by (X', ω', D') a toric blow-up of (X, ω, D) . Then $H_2(X';\mathbb{Z}) = H_2(X;\mathbb{Z}) \oplus \mathbb{Z}E$ where E is the exceptional class, such that

$$\{[C'_1], \dots, [C'_{k+1}]\} = \{[C_1], \dots, [C_{k-1}] - E, E, [C_k] - E\},\$$

which is linearly equivalent to $\{[C_1], \ldots, [C_{k-1}], [C_k], E\}$, and thus generates $H_2(X'; \mathbb{Z})$.

Lemma 2.38. Let D be a toric symplectic log Calabi-Yau divisor, we have $b^+(D) = 1$.

Proof. If Q_D is negative definite, then $q(D) \ge 3$ ([Fri16], Corollary 1.3) and cannot be toric. If Q_D is negative semi-definite, then D is a cycle of -2 spheres. By Lemma 2.36, D is obtained

from (1, 1, 1) or (n, 0, -n, 0) by toric blow-ups. In particular, there must be at least one -1 sphere in D. So a symplectic circular spherical divisor D cannot be negative semi-definite. So we must have $b^+(D) = 1$.

Remark 2.39. A symplectic circular spherical divisor $D \subset (X, \omega)$ is an embedded normal crossing symplectic divisor which is topologically a cycle of spheres (see [LMM20] for definition). It follows from Theorem 1.3 of [LMM20] that such D must be log Calabi-Yau if $b^+(D) = 1$. So a symplectic circular spherical divisor D is a toric symplectic log Calabi-Yau divisor if and only if $b^+(D) = 1$ and q(D) = 0.

Another way to characterize toric symplectic log Calabi-Yau divisors is through their boundaries. To see this, we need to recall some notions on divisor neighborhoods in [LM19]. Let $(D = \bigcup_{i=1}^{k} C_i, \omega)$ be a symplectic divisor. A closed regular neighborhood P(D) of D is called a **concave/convex plumbing** if it is a strong symplectic cap/filling of its boundary. A concave plumbing is also called a **divisor cap** of its boundary. Let Q_D be the intersection matrix of D and $a = (C_i \cdot [\omega]) \in (\mathbb{R}_+)^k$ be the area sequence of D. A symplectic divisor D is said to satisfy positive (resp. negative) **GS criterion** if there exists $z \in (\mathbb{R}_+)^k$ (resp. $(\mathbb{R}_{\leq 0})^k$) such that $Q_D z = a$.

Theorem 2.40 ([LM19],[GS09]). Let (D, ω) be an ω -orthogonal symplectic divisor. Then D has a concave (resp. convex) plumbing if (D, ω) satisfies the positive (resp. negative) GS criterion. Moreover, the induced contact boundary is unique up to contactomorphism.

We will not recall the detailed construction here, but refer the readers to [GS09], [LM19] and [LM21].

Lemma 2.41. A symplectic log Calabi-Yau divisor $D \subset (X, \omega)$ has boundary diffeomorphic to T^3 if and only if it is toric.

Proof. By [GL16], we have $Coker(Q_D) = Coker(A(D) - I)$, where A(D) is the monodromy of boundary torus bundle of D. If Q_D is negative definite, then $A(D) \neq I$ and Y_D cannot be T^3 . If Q_D is negative semi-definite, then all components of D has self-intersection -2. One can compute the monodromy to be $\binom{k+1}{-k}$, which is conjugate to $\binom{1}{-k}$, where k is the length of D. So the boundary cannot be T^3 .

If $b^+(Q_D) = 1$, then D has self-intersection sequence s(D) to be one listed in Theorem 1.3 of [LMM20], up to toric blow-ups and blow-downs. Such equivalence is called a toric equivalence and preserves q(D). We can compute their q(D) by hand as follows.

- For s(D) = (1, p) or (-1, -p) with p = 1, 2, 3, we have $q(D) = 6 \pm (1 + p) \ge 2$.
- For s(D) = (1, 1, p) with $p \le 1$, we have $q(D) = 3 (2 + p) \ge 0$ and q(D) = 0 if and only if p = 1.
- For s(D) = (0, p) with $p \leq 4$, we have $q(D) = 6 p \ge 2$.
- For s(D) = (1, p) with $p \leq -1$, we have $q(D) = 6 1 p \geq 6$.
- The sequence $s(D) = (1, 1 p_1, p_2, \dots, p_{l-1}, 1 p_l)$ with $p_i \ge 2, l \ge 2$ is called blownup if it is non-toric blow-up of a divisor D', where D' is toric equivalent to D'' with s(D'') = (1, 1, 1). Since non-toric blow-up increases q(D), so we have q(D) > q(D') = 0.

So q(D) = 0 if and only if D is toric equivalent to (1, 1, 1). And by computing the monodromy of Y_D listed above, we see that Y_D is T^3 if and only if D is toric equivalent to (1, 1, 1).

3 c_1 -nef cone and counting symplectic log Calabi-Yau divisors

Now we investigate the enumerative aspect of symplectic log Calabi-Yau divisors. From now on, we make the convention that the homology class $H - \sum_{i \in I} E_i$ can be denoted by H_I . For example, $H - E_1 - E_2 - E_3$ can be written as H_{123} .

3.1 c_1 -nef subcone and counting

To ensure the existence of log Calabi-Yau divisors, the symplectic forms must satisfy $c_1(M_l, \omega) \cdot [\omega] > 0$. In the case of symplectic rational surfaces, $c_1(M_l, \omega)$ is unique up to orientationpreserving diffeomrophism ([LL01]). Since the reduced symplectic cone \tilde{P}_l is the fundamental domain inside the symplectic cone under the action of orientation-preserving diffeomorphisms, we have a unique first Chern class for symplectic classes in \tilde{P}_l . We denote this class by $c_1(M_l)$ and define the c_1 -nef subcone to be

$$\tilde{N}_l = \tilde{N}(M_l) = \{A \in H^2(M_l; \mathbb{R}) | PD(A) \text{ is reduced, } A \cdot c_1(M_l) > 0\},\$$

and the normalized c_1 -nef subcone N_l to contain those normalized reduced classes in \tilde{N}_l . We can see below that \tilde{N}_l is indeed a subcone of the reduced symplectic cone \tilde{P}_l and thus N_l a subcone of P_l .

Lemma 3.1. For $A \in \tilde{N}_l$, we have $A \cdot A > 0$ and A is a symplectic class.

Proof. When $l \leq 9$, it is previously known that any reduced class has positive square. We give a brief argument here. Let

$$A = H - \delta_1 E_1 - \dots - \delta_9 E_9$$

be a normalized reduced class in M_l , where $\delta_i = 0$ for i > l. Then we have $A \cdot (H - E_i - E_j - E_k) \ge 0$ for any distinct i, j, k since $1 \ge \delta_i + \delta_j + \delta_k$. Let

$$B = \delta_1(H - E_1 - E_2 - E_3) + \delta_4(H - E_4 - E_5 - E_6) + \delta_7(H - E_7 - E_8 - E_9),$$

and we have $A \cdot B \ge 0$. Consider

$$A - B = (1 - \delta_1 - \delta_4 - \delta_7)H + (\delta_1 - \delta_2)E_2 + (\delta_1 - \delta_3)E_3 + (\delta_4 - \delta_5)E_5 + (\delta_4 - \delta_6)E_6 + (\delta_7 - \delta_8)E_8 + (\delta_7 - \delta_8)E_9.$$

Then we have $A \cdot (A - B) \ge 0$ and thus $A \cdot A \ge 0$. Since $A \cdot c_1 > 0$, we must have $A \cdot A > 0$ by light cone lemma ([McD97]).

When $l \ge 10$, we use Lagrangian multiplier to find minimum of $A \cdot A$. Let $f(\delta_1, \ldots, \delta_l) = A \cdot A = 1 - \sum \delta_i^2$ and $g(\delta_1, \ldots, \delta_l) = A \cdot c_1(X) = 3 - \sum \delta_i$. By Lagrangian multiplier, we know f attains minimum when $\delta_1 = \cdots = \delta_l = \delta$. Since $A \cdot c_1(X) > 0$, we have $\delta < \frac{3}{l}$. Then $A \cdot A \ge \min(f) = 1 - l\delta^2 > 1 - \frac{9}{l} > 0$ as $l \ge 10$.

Since A is reduced, we have \hat{A} is a symplectic class by Proposition 4.9 of [LL01].

Since $c_1(M_l)$ is fixed by orientation-preserving diffeomorphisms, the c_1 -nef cone N_l is also preserved. This provides a natural and simplified domain for the existence of log Calabi-Yau divisors. One advantage of this c_1 -nef cone over the larger reduced symplectic cone is that c_1 -nef cone is a convex linear cone with finitely many faces while the reduced symplectic cone can have boundaries cut out by quadratic equations ([LLW19]).

We are interested in counting the number of symplectic log Calabi-Yau divisors in a symplectic rational surface (M_k, ω) with $[\omega] \in N_k$, the normalized c_1 -nef subcone. Recall Corollary 2.32 shows that there is always a unique elliptic log Calabi-Yau divisor in the c_1 -nef cone. So we will only consider cycles of spheres in the following subsections and automatically plus 1 to each counting result.

Proposition 2.24 implies that counting symplectic log Calabi-Yau divisors is equivalent to counting log Calabi-Yau homological configurations up to symplectomorphisms. But for a symplectic class in the interior of the reduced symplectic cone, the only orientation-preserving diffeomorphism preserving the symplectic class is the identity. So we have the following convenient lemma.

Lemma 3.2. For a symplectic class $[\omega]$ in the interior of the reduced symplectic cone, we have that $D_{K_0,[\omega]} = \{id\}$. In particular, two symplectic log Calabi-Yau divisors $D = \bigcup C_i$ and $D' = \bigcup C'_i$ are strictly symplectic deformation equivalent if and only if $[C_i] = [C'_i]$ for all i, up to a cyclic or anti-cyclic relabelling of the components. In other words, $p\mathcal{HLCY}(X,\omega) = \mathcal{HLCY}(X,\omega)$.

The linear boundaries of the reduced cone are given by equations of the form $\delta_i = \delta_j$ or $\delta_i + \delta_j + \delta_k = 1$. Then the class $E_i - E_j$ or $H - E_i - E_j - E_k$ is represented by a Lagrangian sphere ([LW12]) and the Dehn twist along this Lagrangian sphere is a symplectomorphism generating a strict homological equivalence between some symplectic log Calabi-Yau divisors, which causes repetitions of log Calabi-Yau homological configurations.

So in order to count symplectic log Calabi-Yau divisors, it suffices to count Calabi-Yau homological configurations for each symplectic structure and subtract repetitions along the boundary of the normalized reduced symplectic cone. This is the strategy we will employ in the rest of this section.

3.2 Count of minimal models

We start by counting symplectic log Calabi-Yau divisors in \mathbb{CP}^2 , $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ and $S^2 \times S^2$. They are all minimal and their count can be read off from Theorem 2.14 or Lemma 2.19 as follows.

On \mathbb{CP}^2 , the only normalized symplectic class is $[\omega_{FS}]$ such that $\omega_{FS}(H) = 1$. On $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, we consider the normalized symplectic form ω_{δ} with $\omega_{\delta}(H) = 1, \omega_{\delta}(E) = \delta < 1$. Similarly on $S^2 \times S^2$, we consider ω_{μ} with $\omega_{\mu}(B) = 1, \omega_{\mu}(F) = \mu \ge 1$. Then we have the following counts.

Lemma 3.3.

$$\begin{aligned} |\mathcal{LCY}(\mathbb{CP}^2, \omega_{FS})| &= 3, \\ |\mathcal{LCY}(S^2 \times S^2, \omega_{\mu})| &= \begin{cases} 1 + (1 + [\mu + 1]) + [\mu] + [\mu] = 3[\mu] + 3, when \ \mu > 1\\ 1 + [\mu + 1] + [\mu] + [\mu] = 3[\mu] + 2, when \ \mu = 1 \end{cases} \\ |\mathcal{LCY}(\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2, \omega_{\delta})| &= 1 + (1 + [\frac{1}{1 - \delta}]) + [\frac{1}{1 - \delta}] + [\frac{\delta}{1 - \delta}] = 3[\frac{\delta}{1 - \delta}] + 4. \end{aligned}$$

Proof. Their $p\mathcal{HLCY}$ spaces has been listed in Lemma 2.19. Note that those homological configurations can be realized if and only if all the homology classes have positive symplectic areas.

For \mathbb{CP}^2 , there is nothing to check since the symplectomorphism must act trivially on the homology group.

For $S^2 \times S^2$, if $\mu > 1$, then there is no Lagrangian (-2)-sphere so that any symplectomorphism must act trivially on the homology group ([LW12]). This implies the space $p\mathcal{HLCY}$ is the same as the space \mathcal{HLCY} . So the count for the type 1, 2, 3, 4 in Lemma 2.19 should be $1, 1 + [\mu + 1], [\mu], [\mu]$ respectively. And the total count is $3[\mu] + 3$. When $\mu = 1$, the homological action by symplectomorphism group is generated by switching F and B. Thus (2F + B, B), (F + 2B, F) will be identified after passing from $p\mathcal{HLCY}$ to its quotient \mathcal{HLCY} . In this case, the count should be $3[\mu] + 2$.

For $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, the symplectomorphism must act trivially on the homology group. Thus we only need to count the space $p\mathcal{HLCY}$. The count for the type 1, 2, 3, 4 in Lemma 2.19 should be $1, 1 + \lfloor \frac{1}{1-\delta} \rfloor, \lfloor \frac{1}{1-\delta} \rfloor, \lfloor \frac{\delta}{1-\delta} \rfloor$ respectively. So the total count is given by $3\lfloor \frac{\delta}{1-\delta} \rfloor + 4$.

3.3 Counting symplectic log Calabi-Yau divisors in M_2 and M_3

The remaining rational symplectic surfaces are $M_l = \mathbb{CP}^2 \# l \overline{\mathbb{CP}}^2$, $l \ge 2$ with normalized reduced symplectic form ω given by the reduced basis $(\delta_1, \ldots, \delta_l)$, where $[\omega]$ lies in the normalized c_1 -nef cone N_l . In the subsection, we give a detailed count of symplectic log Calabi-Yau divisors in M_2 and toric symplectic log Calabi-Yau divisors in M_3 for all symplectic forms in the c_1 -nef cone. Our counting in M_2 and also M_3 depends crucially on the following homological classification of negative symplectic spheres of log Calabi-Yau divisors by Corollary 2.33. It also follows from the Proposition 3.4 in [LL20].

Proposition 3.4. All possible negative symplectic spherical classes in M_2 and M_3 can be enumerated as follows.

For M_2 , we have that for integer $k \ge 1$,

- (-1) classes are $E_1, E_2, H E_1 E_2,$
- (-2k-1) class is $-kH + (k+1)E_1$,
- (-2k) class is $-(k-1)H + kE_1 E_2$.

For M_3 , we have that for integer $k \ge 1$,

- (-1) classes are $E_1, E_2, E_3, H E_1 E_2, H E_1 E_3, H E_2 E_3;$
- (-2) classes are $E_1 E_2, E_1 E_3, E_2 E_3, H E_1 E_2 E_3;$
- (-2k-1) classes are $-kH + (k+1)E_1, -(k-1)H + kE_1 E_2 E_3;$
- (-2k-2) classes are $-kH + (k+1)E_1 E_2, -kH + (k+1)E_1 E_3.$

Moreover, these classes can be realized by symplectic spheres if and only if their pairings with the symplectic class are positive. The normalized c_1 -nef cone N_2 coincides with the reduced symplectic cone P_2 and is the convex hull in \mathbb{R}^2 of the points $O = (0,0), M = (\frac{1}{2}, \frac{1}{2})$ and A = (1,0) with line segments OA and MA removed. The cone N_2 is cut by hyperplanes corresponding to negative sphere classes Proposition 3.4 into infinitely many regions, which we describe as follows.

classes Proposition 3.4 into infinitely many regions, which we describe as follows. Consider points $P_i = (\frac{i}{i+1}, \frac{1}{i+1})$ on edge MA and $Q_i = (\frac{i}{i+1}, 0)$ on edge OA, where $i \ge 1$ is an integer. And we also let $Q_0 = O$. The region $\triangle P_i P_{i+1} Q_i$ contains the interior of this triangle and the interior of the edge $Q_i P_{i+1}$. The region $\triangle Q_{i-1} Q_i P_i$ contains the interior of this triangle and the interior of the edge $Q_i P_i$. Then we have

$$N_k = \left(\bigsqcup_{i=1}^{\infty} \triangle P_i P_{i+1} Q_i\right) \sqcup \left(\bigsqcup_{i=1}^{\infty} \triangle Q_{i-1} Q_i P_i\right) \sqcup OM.$$

Proposition 3.5. The count of log Calabi-Yau divisors $\mathcal{LCY}(2; 1, \delta_1, \delta_2)$ in $(M_2; 1, \delta_1, \delta_2)$ is given by

- (1) 14i + 19 in region $\triangle P_i P_{i+1} Q_i, i \ge 1$,
- (2) 14i + 26 in region $\triangle Q_i Q_{i+1} P_{i+1}, i \ge 0$,
- (3) 13 in the interior of the line segment OM.

In other words, in the interior of reduced symplectic cone, we have

$$|\mathcal{LCY}(2;1,\delta_1,\delta_2)| = 7(\lceil \frac{\delta_1}{1-\delta_1} \rceil + \lceil \frac{\delta_1-\delta_2}{1-\delta_1} \rceil) + 12,$$

On the boundary of reduced symplectic cone, that is, when $0 < \delta_1 = \delta_2 < \frac{1}{2}$, we have $|\mathcal{LCY}(2; 1, \delta_1, \delta_2)| = 13$. The results are shown in Figure 1.

Proof. For $n \ge 1$, we can enumerate all possible self-intersection sequences for log Calabi-Yau divisors in M_2 as follows:

- (1) (n+3,-n),
- (2) (n+2,-1,-n), (n+1,0,-n),
- (3) (0, n, -1, -n), (0, n 1, 0, -n), (n + 1, -1, -1, -n),
- $(4) \ (0, n-1, -1, -1, -n),$
- (5) (3,0), (2,1), (1,0,0), (1,-1,1),
- (6) (7).

For each normalized reduced vector $(1, \delta_1, \delta_2)$, we denote by $F_n(2; 1, \delta_1, \delta_2), n \ge 1$, the total count of log Calabi-Yau divisors with self-intersection sequence to be one of (1) - (4) and denote by $F_0(2; 1, \delta_1, \delta_2)$ the count corresponding to (5). We will also denote by #s the count of log Calabi-Yau divisors with self-intersection sequence s. By Proposition 3.4, we consider cases n = 2k and n = 2k + 1 with $k \ge 1$ for each s listed above.

For n = 2k, we have that all possible homology types are

- (1) $((k+2)H (k+1)E_1, -(k-1)H + kE_1 E_2)$
- (2) $((k+2)H (k+1)E_1 E_2, E_2, -(k-1)H + kE_1 E_2),$ $((k+1)H - kE_1, H - E_1, -(k-1)H + kE_1 - E_2)$
- (3) $(H E_1, (k + 1)H kE_1 E_2, E_2, -(k 1)H + kE_1 E_2),$ $(H - E_1, kH - (k - 1)E_1, H - E_1, -(k - 1)H + kE_1 - E_2),$ $((k + 1)H - kE_1, H - E_1 - E_2, E_2, -(k - 1)H + kE_1 - E_2)$

(4)
$$(H - E_1, kH - (k - 1)E_1, H - E_1 - E_2, E_2, -(k - 1)H + kE_1 - E_2)$$

These homology types can be realized by log Calabi-Yau divisors if and only if the symplectic area of each component is positive. So we have

$$F_{2k}(2; 1, \delta_1, \delta_2) = \begin{cases} 7 & \text{if } -(k-1) + k\delta_1 - \delta_2 > 0\\ 0 & \text{otherwise} \end{cases}$$

For n = 2k + 1, all possible homology types are

(1)
$$((k+3)H - (k+2)E_1 - E_2, -kH + (k+1)E_1)$$

- (2) $((k+2)H (k+1)E_1, H E_1 E_2, -kH + (k+1)E_1),$ $((k+2)H - (k+1)E_1 - E_2, H - E_1, -kH + (k+1)E_1)$
- (3) $(H E_1, (k + 1)H kE_1, H E_1 E_2, -kH + (k + 1)E_1),$ $(H - E_1, (k + 1)H - kE_1 - E_2, H - E_1, -kH + (k + 1)E_1),$ $((k + 2)H - (k + 1)E_1 - E_2, E_2, H - E_1 - E_2, -kH + (k + 1)E_1)$

(4)
$$(H - E_1, (k+1)H - kE_1 - E_2, E_2, H - E_1 - E_2, -kH + (k+1)E_1)$$

So we have

$$F_{2k+1}(2; 1, \delta_1, \delta_2) = \begin{cases} 7 & \text{if } -k + (k+1)\delta_1 > 0\\ 0 & \text{otherwise} \end{cases}$$

For n = 1, we have the following homology types:

- (1) $(4, -1) = (2H, H E_1 E_2), (3H 2E_p E_q, E_p),$ (2) $(3, -1, -1) = (2H - E_i, E_i, H - E_1 - E_2),$ $(2, 0, -1) = (2H - E_1 - E_2, H - E_i, E_i),$
- (3) $(0, 1, -1, -1) = (H E_i, H, H E_1 E_2, E_i),$ $(0, 0, 0, -1) = (H - E_p, H - E_q, H - E_p, E_p),$ $(2, -1, -1, -1) = (2H - E_1 - E_2, E_1, H - E_1 - E_2, E_2),$
- (4) $(0, 0, -1, -1, -1) = (H E_2, H E_1, E_1, H E_1 E_2, E_2),$

where $i \in \{1, 2\}$ and $\{p, q\} = \{1, 2\}$. So

$$F_1(2; 1, \delta_1, \delta_2) = \begin{cases} 13, & \text{if } \delta_1 > \delta_2; \\ 8, & \text{if } \delta_1 = \delta_2. \end{cases}$$

For those s in (5), we could also list their homology types as follows:

- $(3,0) = (2H E_p, H E_q)$
- $(2,1) = (2H E_1 E_2, H)$
- $(1,0,0) = (H, H E_1, H E_2)$
- $(1, -1, 1) = (H, H E_1 E_2, H)$

where $\{p,q\} = \{1,2\}$. So

$$F_0(2; 1, \delta_1, \delta_2) = \begin{cases} 5, & \text{if } \delta_1 > \delta_2; \\ 4, & \text{if } \delta_1 = \delta_2. \end{cases}$$

Also by Corollary 2.32, there is always a unique divisor in (7). Then we have

$$|\mathcal{LCY}(2;1,\delta_1,\delta_2)| = 1 + F_0(2;1,\delta_1,\delta_2) + F_1(2;1,\delta_1,\delta_2) + \sum_{n=2}^{\infty} F_n(2;1,\delta_1,\delta_2)$$

The result follows by recognizing

Also we get the formula because when $\delta_1 > \delta_2$ we have

$$\left\lceil \frac{\delta_1}{1-\delta_1} \right\rceil + \left\lceil \frac{\delta_1 - \delta_2}{1-\delta_1} \right\rceil = \begin{cases} 2i+1 & \text{if } -i + (i+1)\delta_1 > 0, -i + (i+1)\delta_1 - \delta_2 \leqslant 0, \\ 2i+2 & \text{if } -i + (i+1)\delta_1 - \delta_2 > 0, -(i+1) + (i+2)\delta_1 \leqslant 0. \end{cases}$$





Figure 1: Counting of \mathcal{LCY} for M_2 .

Figure 2: Counting of $t\mathcal{LCY}$ for M_2 .

By only counting the toric log Calabi-Yau divisors in the proof of the above proposition, we get the following count of toric divisors.

Corollary 3.6. The count of toric log Calabi-Yau divisors on $(M_2; 1, \delta_1, \delta_2)$ is given by

- (1) 2i + 1 in the region $\triangle Q_i P_i P_{i+1}$;
- (2) 2i + 2 in the region $\triangle Q_i Q_{i+1} P_{i+1}$;
- (3) 1 in the interior of edge $OM = Q_0 P_1$.

In other words, we have

$$|t\mathcal{LCY}(M_2; 1, \delta_1, \delta_2)| = \left\lceil \frac{\delta_1}{1 - \delta_1} \right\rceil + \left\lceil \frac{\delta_1 - \delta_2}{1 - \delta_1} \right\rceil$$

The results are shown in Figure 2.

Proof. The toric symplectic log Calabi-Yau divisors has self-intersection sequence (0, n - 1, -1, -n). For n = 2k, the only homology type is $(H - E_1, kH - (k - 1)E_1, H - E_1 - E_2, E_2, -(k - 1)H + kE_1 - E_2)$. So the number of toric symplectic log Calabi-Yau divisors is

$$tF_{2k}(2; 1, \delta_1, \delta_2) = 1$$
 if $-(k-1) + k\delta_1 - \delta_2 > 0, \delta_1 + \delta_2 < 1.$

For n = 2k + 1, the only homology type is $(H - E_1, (k + 1)H - kE_1 - E_2, E_2, H - E_1 - E_2, -kH + (k + 1)E_1)$. Then

$$tF_{2k+1}(2; 1, \delta_1, \delta_2) = 1$$
 if $-k + (k+1)\delta_1 > 0, \delta_1 + \delta_2 < 1.$

For n = 1, the only homology type is $(H - E_2, H - E_1, E_1, H - E_1 - E_2, E_2)$. Then

$$tF_1(2; 1, \delta_1, \delta_2) = 1$$
 if $\delta_1 > \delta_2, \delta_1 + \delta_2 < 1$

Above counts are 0 everywhere else. We get the results by summing over the counts. \Box

Next we give a detailed count of toric symplectic log Calabi-Yau divisors in M_3 for all symplectic forms in the normalized c_1 -nef cone N_3 .

The normalized c_1 -nef cone N_3 is the same as the reduced symplectic cone P_3 and is the convex hull in \mathbb{R}^3 of the points $O = (0, 0, 0), M = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}), A = (1, 0, 0)$ and $B = (\frac{1}{2}, \frac{1}{2}, 0)$ with the triangle OAB removed. By Proposition 3.4, there is a pattern of four walls in N_3 given by $-(k-1)H + kE_1, -(k-1)H + kE_1 - E_2 - E_3, -kH + (k+1)E_1 - E_2, -kH + (k+1)E_1 - E_3$. They never intersect with each other except on the edges. The cone N_3 is again cut by these walls into infinitely many regions, which we describe as follows.

walls into infinitely many regions, which we describe as follows. Consider points $P_i = (\frac{i+1}{i+3}, \frac{1}{i+3}, \frac{1}{i+3})$ on edge MA, $Q_i = (\frac{i}{i+1}, 0, 0)$ on edge OA and $R_i = (\frac{i+1}{i+2}, \frac{1}{i+2}, 0)$ on edge AB, where $i \ge 0$ is an integer. The regions $Q_{i-1}R_{i-1}P_{2i-1}P_{2i-2}, \quad Q_{i-1}Q_iR_{i-1}P_{2i-1}, \quad Q_iR_{i-1}P_{2i-1}P_{2i}, \quad Q_iR_{i-1}R_iP_{2i}$

contain the interior subcones and the interior of their faces

$$Q_{i-1}R_{i-1}P_{2i-2}, \quad Q_iR_{i-1}P_{2i-1}, \quad Q_iR_{i-1}P_{2i}, \quad Q_iR_iP_{2i}$$

respectively; the regions

$$Q_{i-1}P_{2i-2}P_{2i-1}, \quad Q_{i-1}Q_iP_{2i-1}, \quad Q_iP_{2i-1}P_{2i}, R_{i-1}P_{2i-2}P_{2i-1}, \quad R_{i-1}P_{2i-1}P_{2i}, \quad R_{i-1}R_iP_{2i-2}P_{2i-1}, \quad R_{i-1}P_{2i-2}P_{2i-1}, \quad R_{i-1}P_{2i-2}P_{2i-2}P_{2i-1}, \quad R_{i-1}P_{i-1}P_{i-2}P_{i-2}P_{i-2}P_{i-2}P_{i-2}P_{i-2}P_{i-2}P_{i-2}P_{i-2}$$



Figure 3: Counting of $t\mathcal{LCY}$ for M_3 in the interior and boundary of the c_1 -nef cone.

contain the interior triangles and the interior of their edges

 $Q_{i-1}P_{2i-1}, \quad Q_iP_{2i-1}, \quad Q_iP_{2i}, \quad R_{i-1}P_{2i-1}, \quad R_{i-1}P_{2i}, \quad R_iP_{2i}$

respectively; the region $P_{i-1}P_i$ contains the interior of this edge and the point P_i .

These regions form the entire normalized reduced symplectic cone $P_3 = N_3$, so we have listed all the possible symplectic classes on M_3 . We claim the following counting results, which follows from the same method for M_2 cases. The proof is quite long, thus we put it in the appendix A.2.

Proposition 3.7. Let $i \ge 1$ be an integer. The number of toric symplectic log CY divisors on the rational surface M_3 is

(1) 10i - 2 in the region $Q_{i-1}R_{i-1}P_{2i-1}P_{2i-2}$ (2) 10i in the region $Q_{i-1}Q_iR_{i-1}P_{2i-1}$

(3) 10i + 3 in the region $Q_i R_{i-1} P_{2i-1} P_{2i}$ (4) 10i + 5 in the region $Q_i R_{i-1} R_i P_{2i}$

- (5) 4i 1 in the region $Q_{i-1}P_{2i-2}P_{2i-1}$ (6) 4i in the region $Q_{i-1}Q_iP_{2i-1}$
- (7) 4i + 1 in the region $Q_i P_{2i-1} P_{2i}$ (8) 4i in the region $R_{i-1} P_{2i-2} P_{2i-1}$
- (9) 4i + 2 in the region $R_{i-1}P_{2i-1}P_{2i}$ (10) 4i + 3 in the region $R_{i-1}R_iP_{2i}$
- (11) i + 1 in the region $P_{i-1}P_i$ (12) 3 in the region $OBM = Q_0R_0P_0$

(13) 2 in the region $BM = R_0P_0$ (14) 1 in the region $OM = Q_0P_0$ and the monotone point $M = P_0 = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3})$. The results are shown in Figure 3.

3.4 Toric regions

Instead of giving a counting formula of toric symplectic log Calabi-Yau divisors, we could also investigate the symplectic classes for which they could exist. This is called the **toric region** T_l of the manifold M_l . By Theorem 1.5 that will be proved in the next section, this will also be the part of the normalized c_1 -nef cone where there exists a toric action. From the counting results for M_3, M_2 and minimal models, we already know the following:

Corollary 3.8. When $l \leq 3$, the toric region T_l will be the whole normalized reduced symplectic cone P_l .

Now we discuss the cases for rational surfaces M_4, M_5, M_6 . We start with the following simple observation:

Fact 3.9. For $4 \leq l \leq 6$, the toric regions T_l contains the interior of normalized reduced symplectic cones P_l .

Proof. Consider the following homological configurations for M_4, M_5, M_6 respectively:

$$(H_{134}, E_4, E_1 - E_4, H_{12}, E_2, H_{23}, E_3)$$
$$(H_{134}, E_4, E_1 - E_4, H_{12}, E_2 - E_5, E_5, H_{235}, E_3)$$
$$(H_{134}, E_4, E_1 - E_4, H_{125}, E_5, E_2 - E_5, H_{236}, E_6, E_3 - E_6)$$

Note that they can be realized by Corollary 2.33 since it is easy to see that each homology class in the above sequences has positive pairing with any $[\omega]$ in the interior of P_l . Also the realization of them are toric log Calabi-Yau since $q(D) = 12 - k(D) - D^2 = 12 - (l+3) - (9-l) = 0$.

Recall that the normalized reduced symplectic cone P_4 for M_4 is the convex hull of $O = (0, 0, 0, 0), M = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}), A = (1, 0, 0, 0), B = (\frac{1}{2}, \frac{1}{2}, 0, 0)$ and $C = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0)$ not containing the face OABC.

Proposition 3.10. The toric region T_4 for M_4 is equal to P_4 deleting two edges MO, MA.

Proof. If the homological sequence for M_4 in Fact 3.9 can not be realized by some $[\omega] \in P_l$, then either $\omega(H_{134}) = 0$ or $\omega(E_1 - E_4) = 0$. They lead to the restrictions $\delta_1 + \delta_2 + \delta_3 = 1$, $\delta_2 = \delta_3 = \delta_4$ or $\delta_1 = \delta_2 = \delta_3 = \delta_4$, which exactly correspond to the edge MA or MO. Thus we only need to show the nonexistence of toric symplectic log Calabi-Yau divisors on MA and MO.

Firstly by considering the toric blow up operation on the self-intersection sequences of M_3 , we can classify all the possible self-intersection sequences for M_4 which are:

$$\{A_n = (-2, -1, -2, -n, -1, -1, n-2)\}_{n=1}^{\infty} \{B_n = (-1, -2, -1, -n-3, -1, -1, n)\}_{n=1}^{\infty} \{C_n = (-1, -2, -2, -1, -n-3, 0, n)\}_{n=1}^{\infty} \{D_n = (-1, -3, -1, -2, n, 0, -n-2)\}_{n=-\infty}^{\infty} \{E_n = (-3, -1, -2, -2, n, 0, -n-1)\}_{n=-\infty}^{\infty}$$

Next by Proposition 3.4 of [LL20], for $k \ge 1$, S_{ω}^{-2k-2} and S_{ω}^{-2k-3} are subsets of the following sets respectively, with the only restrictions on positive pairing with ω :

$$\begin{split} \mathcal{S}_{\omega}^{-2k-2} &\subset \{-(k-1)H + kE_1 - E_2 - E_3 - E_4, -kH + (k+1)E_1 - E_2, \\ &- kH + (k+1)E_1 - E_3, -kH + (k+1)E_1 - E_4\} \\ \mathcal{S}_{\omega}^{-2k-3} &\subset \{-kH + (k+1)E_1 - E_2 - E_3, -kH + (k+1)E_1 - E_2 - E_4, \\ &- kH + (k+1)E_1 - E_3 - E_4, -(k+1)H + (k+2)E_1\} \end{split}$$

For $\mathcal{S}_{\omega}^{-1}$, $\mathcal{S}_{\omega}^{-2}$ and $\mathcal{S}_{\omega}^{-3}$, their possible choices are as follows, again with the postive pairing condition as the only restriction:

$$\begin{split} \mathcal{S}_{\omega}^{-1} &\subset \{E_k, H - E_i - E_j | 1 \leqslant k \leqslant 4, 1 \leqslant i < j \leqslant 4\} \\ \mathcal{S}_{\omega}^{-2} &\subset \{E_p - E_m, H - E_i - E_j - E_k | 1 \leqslant p < m \leqslant 4, 1 \leqslant i < j < k \leqslant 4\} \\ \mathcal{S}_{\omega}^{-3} &\subset \{-H + 2E_1, H - E_1 - E_2 - E_3 - E_4, E_i - E_j - E_k | 1 \leqslant i < j < k \leqslant 4\} \end{split}$$

Now we can explain the nonexistence of toric symplectic log Calabi-Yau divisor when the symplectic class is on the edge OM or MA, which is the consequence of the lack of some (-2)- and (-3)-spheres.

Along the edge MO, we have

$$S_{\omega}^{-2} = \{H - E_i - E_j - E_k | 1 \le i < j < k \le 4\}$$

$$S_{\omega}^{-3} = \{H - E_1 - E_2 - E_3 - E_4\}.$$

From this we can see that any two elements in S_{ω}^{-2} or S_{ω}^{-3} have negative intersection number, which excludes the existence of toric symplectic log Calabi-Yau divisors of type A_n, C_n, D_n, E_n since the any two (-2)- or (-3)-components in the divisor must have intersection number 1 if they are adjacent in the sequence or 0 otherwise. Moreover, B_n type is also impossible since $S_{\omega}^{\leq -4} = \emptyset$ for $[\omega] \in MO$.

Along the edge MA, we have

$$S_{\omega}^{-2} = \{E_1 - E_2, E_1 - E_3, E_1 - E_4, H - E_2 - E_3 - E_4\}$$

$$S_{\omega}^{-3} = \{-H + 2E_1, E_1 - E_2 - E_3, E_1 - E_2 - E_4, E_1 - E_3 - E_4\}.$$

Still any two elements in S_{ω}^{-2} or S_{ω}^{-3} have negative intersection number, which excludes divisors of type A_n, C_n, D_n, E_n . And B_n type is also excluded because although $S_{\omega}^{\leq -4}$ may not be empty, the intersection numbers between the elements in S_{ω}^{-2} and $S_{\omega}^{\leq -4}$ are negative. However the (-2)-component and (-n-3)-component in B_n must have intersection number 0 since they are nonadjacent.

A similar but more complicated analysis can be done for M_5 . The normalized reduced symplectic cone P_5 for M_5 is the convex hull of

$$O = (0, 0, 0, 0, 0), M = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}), A = (1, 0, 0, 0, 0),$$
$$B = (\frac{1}{2}, \frac{1}{2}, 0, 0, 0), C = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0, 0), D = (\frac{1}{3}, \frac{1}{3}, \frac{1}{3}, \frac{1}{3}, 0),$$

not containing the face *OABCD*. Let's introduce a special point $X = (\frac{1}{2}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4}, \frac{1}{4})$ on the edge *MA*. We have the following result, the proof of which is quite similar to the case for M_4 and contained in the appendix A.3.

Proposition 3.11. The toric region T_5 for M_5 is equal to P_5 deleting the closed faces MOD, MAD and MOX which is a quarter of the face MOA.

Remark 3.12. After we establish the equivalence between toric divisors and toric actions later, we can also check the above results from the perspective of performing equivariant blow up, by analysing the possibilities of chopping the corners of Delzant polygons.

Next we give some qualitative characterizations of the toric regions for general M_l .

Lemma 3.13. The toric region T_l is star-shaped in the directions $\delta_2, \ldots, \delta_l$, i.e. for any $(1; \delta_1, \delta_2, \ldots, \delta_l) \in T_l$, $(1; \delta_1, \lambda_2 \delta_2, \ldots, \lambda_l \delta_l)$ is also in T_l for all $0 < \lambda_l \leq \cdots \leq \lambda_2 \leq 1$.

Proof. Let $a := [\omega_{(1;\delta_1,\delta_2,...,\delta_k)}]$ and $a_{\lambda} := [\omega_{(1;\delta_1,\lambda_2\delta_2,...,\lambda_l\delta_l)}]$. We could prove this lemma directly by using Corollary 2.33 but we can actually show a stronger result: $S_a \subset S_{a_{\lambda}}$. By Theorem 1.6, This implies $t\mathcal{HLCY}(M_l, a) \subset t\mathcal{HLCY}(M_l, a_{\lambda})$.

For $A = aH - \sum b_i E_i \in S_a$, we see that $A \in S_{a_\lambda}$ if and only if $a_\lambda \cdot A > 0$ by Lemma 2.20.

• If a > 0, then $b_i \ge 0$ for all *i* by Lemma 3.3 of [Che20], which implies

$$a_{\lambda} \cdot A = a - b_1 \delta_1 - \sum \lambda_i b_i \delta_i > a - b_1 \delta_1 - \sum b_i \delta_i > 0.$$

• If a < 0, then A must be of the form $-|a|H + (|a|+1)E_1 - E_{j_2} - \cdots - E_{j_s}$ by Lemma 3.4 of [Che20]. Then we have

$$a_{\lambda} \cdot A = -|a| + (|a|+1)\delta_1 - \lambda(\delta_{j_2} + \dots + \delta_{j_s}) \ge -|a| + (|a|+1)\delta_1 - (\delta_{j_2} + \dots + \delta_{j_s}) > 0.$$

• If a = 0, then by adjunction formula we have $\sum b_i(b_i - 1) = 2$. So there is one k such that $b_k = -1$ or 2 and $b_j = 0$ or 1 for $j \neq k$. Since $a \cdot A = \sum b_i \delta_i > 0$, we must have $b_k = -1$ and j < k if $b_j = 1$. Then

$$a_{\lambda} \cdot A = \lambda_k \delta_k - \sum_{j > k} \lambda_j b_j \delta_j \ge \lambda_k (\delta_k - \sum_{j > k} b_j \delta_j) = \lambda_k (a \cdot A) > 0.$$

Proposition 3.14. Toric region T_l is connected for each l.

Proof. For $l \leq 2$, $T_l = P_l$ is the entire normalized reduced cone and is thus connected. So we only consider the case $l \geq 3$. The region

$$T_l^{\varepsilon} := \{ (1; \delta_1, \dots, \delta_l) \in P_l \, | \, \delta_{i-1} > \delta_i, \delta_i < \varepsilon, i \ge 2 \}$$

is contained in the toric region T_l for sufficiently small ε since

$$(H - E_1, E_1 - E_2, E_2 - E_3, \dots, E_{k-1} - E_k, E_k, H - E_1 - E_2 - \dots - E_k, H)$$

is a toric log Calabi-Yau divisor in this region. Note that every $(1; \delta_1, \delta_2, \ldots, \delta_l)$ in T_l is connected to $(1; \delta_1, \lambda_2 \delta_2, \ldots, \lambda_l \delta_l) \in T_l^{\varepsilon}$ for sufficiently small $0 < \lambda_l < \cdots < \lambda_2 \leq 1$. So we conclude that T_l is connected.

3.5 General counting formulas for restrictive reduced symplectic classes

When the blow up times are getting higher, both the chamber structure on the normalized c_1 -nef cone and the homological action of symplectomorphism group become quite involved. It's hopeless to give a comprehensive description of the counting according to the division of the symplectic cone as Proposition 3.7 for all higher blow ups. Nevertheless, when we only focus on a restrictive part of reduced symplectic classes, it's possible to write down a general counting formula.

In this section, by a **restrictive** reduced symplectic class, we mean a reduced symplectic class encoded by $\vec{\delta} = (\delta_1, \dots, \delta_l)$ which staconetisfies the extra assumptions:

$$1 > \delta_1 + \dots + \delta_l$$

$$\delta_1 > \delta_2$$
$$\delta_k > \delta_{k+1} + \dots + \delta_l$$

for all $2 \le k \le l-1$. There are two advantages of counting in this region. Firstly we don't need to consider the homological action of the symplectomorphism group by Lemma 3.2 since this region is contained in the interior of the reduced symplectic cone. Secondly, the classes $H - \sum_{i=1}^{l} \epsilon_i E_i$ and $E_i - \sum_{j>i} \epsilon_j E_j$ can be used freely to create the homological configurations.

Now we introduce the strategy for this general counting formula, which is different from the counting of M_2 , M_3 in the previous sections. There, we firstly determine all the possible self-intersection sequence, according to which we then list all the possible homological configurations by Corollary 2.33 and consider the quotient by homological action of the symplectomorphism group. We also make use of Propostion 3.4 which makes our counting more efficient. However, for the counting issue of general M_l , we are going to apply Lemma 2.31 directly. The homological configurations in M_l come from the homological toric and non-toric blow ups of either homological configurations in M_1 or $(3H - E_1 - \cdots - E_{j-1} - 2E_j, E_j)$ in M_j . So we will actually count the number of the possible blow up patterns modulo some symmtries resulting the same homological configurations in M_l .

Note that by the restrictive condition, the only possible classes having negative symplectic area in \mathcal{H}_l (Corollary 2.33) are

$$kH - (k-1)E_1 - \sum_{i=2}^{l} \epsilon_i E_i, k \in \mathbb{Z}, \epsilon_i \in \{0, 1\}$$

with non positive k. We need some notations to record the number of such k that can appear in the homological configurations:

Let

$$\mathcal{G}_{l} = \left\{ g : \{2, \cdots l\} \to \{0, 1\} \right\}$$
$$\psi(\vec{\delta}, g) = \left\lceil \frac{\delta_{1} - \sum_{i=2}^{l} g(i)\delta_{i}}{1 - \delta_{1}} \right\rceil$$

We also need the following set of functions to record the length of the divisor at each step:

$$\mathcal{F}_{l}^{a} = \left\{ f : \{1, 2, \cdots l\} \to \mathbb{Z}_{+} \mid f(1) = a, f(i) - f(i-1) = 0 \text{ or } 1, \text{ for all } 2 \le i \le l \right\}$$

In particular, for the purpose of merely counting toric divisors, we denote by F_l^a the function in \mathcal{F}_l^a such that $F_l^a(i) = a + i - 1$ for all *i*.

For any $g \in \mathcal{G}_l, f \in \mathcal{F}_l^a$ and reduced vector $\vec{\delta} = (\delta_1, \cdots, \delta_l)$, define

$$\phi(f,g) = \prod_{i=2}^{l} ((1-g(i))f(i-1) - (-1)^{g(i)}(f(i) - f(i-1) + 1))$$

This number will represent the number of blow up patterns such that the length at each step is given by f and there exists the class $kH - (k-1)E_1 - \sum_{i=2}^l g(i)E_i$ in the ultimate configuration. (See its alternative expression in the proof of Lemma 3.15).

For the convenience of the statement, in the following we always use \vec{t} to denote an ordered finite set (t_1, \dots, t_k) with all t_i being positive integers. Let

$$|\vec{t}| = t_1 + \dots + t_k$$

 $a^{\vec{t}} = a^{t_1}(a+1)^{t_2} \cdots (a+k-1)^{t_k}$

where a is an integer. It follows that we have the relation:

Lemma 3.15.

$$\sum_{f \in \mathcal{F}_l^a, g \in \mathcal{G}_l} \phi(f, g) = 2 \sum_{|\vec{t}| = l-1} a^{\vec{t}}$$

Proof. Note that

$$((1-g(i))f(i-1)-(-1)^{g(i)}(f(i)-f(i-1)+1)) = \begin{cases} 2 & \text{if } g(i) = 1, f(i) - f(i-1) = 1\\ 1 & \text{if } g(i) = 1, f(i) - f(i-1) = 0\\ f(i-1) - 2 & \text{if } g(i) = 0, f(i) - f(i-1) = 1\\ f(i-1) - 1 & \text{if } g(i) = 0, f(i) - f(i-1) = 0 \end{cases}$$

If we fix some $f \in \mathcal{F}_l^a$ and sum over $g \in \mathcal{G}_l$, we will get

$$\sum_{g \in \mathcal{G}_l} \phi(f,g) = f(1)f(2) \cdots f(l-1)$$

Therefore

$$\sum_{f \in \mathcal{F}_l^a, g \in \mathcal{G}_l} \phi(f, g) = \sum_{f \in \mathcal{F}_l^a} \sum_{g \in \mathcal{G}_l} \phi(f, g) = 2 \sum_{|\vec{t}| = l-1} a^{\vec{t}}$$

To circumvent too many sigma notations in the general counting formulas, for any integer a and $g \in \mathcal{G}_l$, we also introduce $a^{\vec{g}}$ to denote:

$$\sum_{f\in \mathcal{F}_l^a}\phi(f,g)$$

Proposition 3.16. With the extra restrictive assumption on the reduced symplectic class encoded by $\vec{\delta} = (\delta_1, \dots, \delta_l)$, we have

$$\begin{aligned} |\mathcal{LCY}(M_l, \omega_{\delta_1, \cdots, \delta_l})| &= 1 + 2^l + \sum_{|\vec{t}| = l-1} (2^{\vec{t}} + 3^{\vec{t}}) + \sum_{1 \le |\vec{t}| \le j-1} 2^{\vec{t}} \\ &+ \sum_{g \in \mathcal{G}_l} \left[1 + \frac{1}{2} (2^{\vec{g}} + 4^{\vec{g}}) + 3^{\vec{g}} \right] \psi(\vec{\delta}, g) \end{aligned}$$

$$|t\mathcal{LCY}(M_l,\omega_{\delta_1,\cdots,\delta_l})| = \sum_{g\in\mathcal{G}_l} \frac{1}{2}\phi(F_l^4,g)\psi(\vec{\delta},g)$$

Moreover, when $\vec{\delta}$ doesn't satisfy the extra restrictive assumption, the above formulas will give a strictly upper bound of the counting.

Proof. Note that the assumption guarantees $\vec{\delta}$ is in the interior of the normalized reduced cone. Thus there is no difference among \mathcal{LCY} , \mathcal{HLCY} and $p\mathcal{HLCY}$. Given any $(A_1, \dots, A_k) \in$ $\mathcal{HLCY}(M_l, \omega_{\delta_1, \dots, \delta_l})$ with $l \ge 2$, by Lemma 2.31 we observe that it must be either the iterated toric or non-toric homological blow up of some element in $p\mathcal{HCLY}(M_1) = \bigcup_{\delta < 1} p\mathcal{HCLY}(M_1, \omega_{\delta})$ or $(3H - E_1 - \cdots - E_{j-1} - 2E_j, E_j)$ for some $2 \le j \le l$. We call them the **germ** of (A_1, \dots, A_k) . Conversely, blowing up different germs must give different divisors M_l . Only different toric or non-toric blow up patterns on the same germ may lead to the same divisor in M_l due to the cyclic symmetry of the circular divisors.

Therefore the counting is divided into the following parts according to the germ:

• The germ is some divisor $((k + 1)H - kE_1, H_1, -kH + (k + 1)E_1, H_1)$ with $k \ge 0$ in $p\mathcal{HCLY}(M_1)$. The assumption ensures that after l-1 times homological blow up, only $-kH + (k+1)E_1 - \sum_{i=2}^l g(i)E_i$ may not in $\mathcal{S}_{\omega_{\delta}}$ for some $g \in \mathcal{G}_l$. Now we use the function f to denote the length at each stage. So $\phi(f,g)$ actually denotes the number of blow up patterns which follows the length function f and gives $-kH + (k+1)E_1 - \sum_{i=2}^l g(i)E_i$ in the final homological configuration. And the sum over all possible f, g and integers k, which is given by $\sum_{f \in \mathcal{F}_l^4, g \in \mathcal{G}_l} \phi(f,g)\psi(\vec{\delta},g)$, leads to the count of blow up patterns starting from length 4 divisor in M_1 .

Observe that whenever the blow up pattern is not performing l-1 non-toric blow up on $(k+1)H-kE_1$ and $-kH+(k+1)E_1$, there exist exactly two patterns resulting in the same divisor in M_l due to the cyclic symmetry of $((k+1)H-kE_1, H_1, -kH+(k+1)E_1, H_1)$. The ultimate count should then be

$$\sum_{g \in \mathcal{G}_l} \sum_{f \in \mathcal{F}_l^4} (\frac{1}{2}(\phi(f,g) - 1) + 1)\psi(\vec{\delta},g)$$

When counting only toric divisors, eliminating the symmetry is simply by dividing by 2. So we could obtain the toric counting formula

$$|t\mathcal{LCY}(M_l,\omega_{\delta_1,\cdots,\delta_l})| = \sum_{g\in\mathcal{G}_l} \frac{1}{2}\phi(F_l^4,g)\psi(\vec{\delta},g)$$

• The germ is some divisor $((k+2)H - (k+1)E_1, -kH + (k+1)E_1, H_1)$ with $k \ge -1$ in $p\mathcal{HCLY}(M_1)$. Note that if $k \ge 0$, there is no cyclic symmetry so that the count of patterns exactly gives the count of divisors, which is

$$\sum_{g \in \mathcal{G}_l} \sum_{f \in \mathcal{F}_l^3} \phi(f,g) \psi(\vec{\delta},g)$$

When k = -1, the germ is (H_1, H, H) . Observe that exactly two patterns generate the same divisor unless the pattern is performing l-1 non-toric blow ups on H_1 or performing toric blow up once between H and H to get $H - E_j, E_j, H - E_j$ and non-toric blow ups on H_1 and E_j . The number of those patterns should be $1 + \sum_{j=2}^{l} 2^{l-j} = 2^{l-1}$. The count of divisors thus should be

$$\frac{1}{2} \bigg[-2^{l-1} + \sum_{f \in \mathcal{F}_l^3, g \in \mathcal{G}_l} \phi(f,g) \bigg] + 2^{l-1}$$

• The germ is some divisor $((k+3)H - (k+2)E_1, -kH + (k+1)E_1)$ with $k \ge -1$ or $(2H, H - E_1)$ in $p\mathcal{HCLY}(M_1)$. Observe that exactly two patterns generate the same divisor unless the pattern is performing l-1 non-toric blow ups. The count for $((k+3)H - (k+2)E_1, -kH + (k+1)E_1)$ with $k \ge 0$ is

$$\sum_{g\in \mathcal{G}_l} \sum_{f\in \mathcal{F}_l^2} (\frac{1}{2}(\phi(f,g)-1)+1)\psi(\vec{\delta},g)$$

The count for (2H, H - E) and (2H - E, H) should both be

$$\sum_{g \in \mathcal{G}_l} \sum_{f \in \mathcal{F}_l^2} \left(\frac{1}{2} (\phi(f,g) - 1) + 1 \right)$$

• The germ is $(3H - E_1 - \dots - E_{j-1} - 2E_j, E_j)$ for some $2 \le j \le l$. Similar to the cases of (2H, H - E) and (2H - E, H) above, the count should be

$$\sum_{j=2}^{l} \left[\sum_{f \in \mathcal{F}_{l-j+1}^2, g \in \mathcal{G}_{l-j+1}} \frac{1}{2} \phi(f,g) + 2^{l-j-1} \right]$$

Adding the above four parts and one elliptic divisor together and applying Lemma 3.15, we get our desired counting formula. Moreover, we have enumerated all the possible homological types in the above analysis. If $[\omega]$ is not in the restrictive region, the number of divisors will be strictly less than the number given by the formula due to the lack of some classes $H - \sum_{i=1}^{l} \epsilon_i E_i$ or $E_i - \sum_{j>i} \epsilon_j E_j$.

Remark 3.17. When l = 2, 3, this extra assumption only implies $\vec{\delta}$ is in the interior of the normalized reduced cone. We can compare the counting results and the above formulas:

For the toric case, when l = 2, \mathcal{G}_2 only contains two elements and $\frac{1}{2}\phi(F_2^4, g)$ is always equal to 1. This corresponds to Corollary 3.6: $\left[\frac{\delta_1}{1-\delta_1}\right] + \left[\frac{\delta_1-\delta_2}{1-\delta_1}\right]$. When l = 3,

$$\frac{1}{2}\phi(F_3^4,g) = \begin{cases} 2 & g(2) = 0, g(3) = 1\\ 2 & g(2) = 1, g(3) = 1\\ 3 & g(2) = 0, g(3) = 0\\ 3 & g(2) = 1, g(3) = 0 \end{cases}$$

This gives the counting $3(\lceil \frac{\delta_1}{1-\delta_1} \rceil + \lceil \frac{\delta_1 - \delta_2}{1-\delta_1} \rceil) + 2(\lceil \frac{\delta_1 - \delta_3}{1-\delta_1} \rceil + \lceil \frac{\delta_1 - \delta_2 - \delta_3}{1-\delta_1} \rceil)$ in the interior of the normalized reduced cone, which coincides with Proposition 3.7 cases (1)-(4). For the general LCYs, when l = 2, we see that

$$\sum_{\substack{|\vec{t}|=1}} 2^{\vec{t}} = 2$$
$$\sum_{\substack{|\vec{t}|=1}} 3^{\vec{t}} = 3$$

$$1 + \frac{1}{2}(2^{\vec{g}} + 4^{\vec{g}}) + 3^{\vec{g}} = 1 + 3 + 3 = 7 \quad for \ both \ g \in \mathcal{G}_2$$

This recovers $7(\lceil \frac{\delta_1}{1-\delta_1} \rceil + \lceil \frac{\delta_1 - \delta_2}{1-\delta_1} \rceil) + 12$ in Proposition 3.5.

4 Symplectic log Calabi-Yau divisors and almost toric fibrations

In this section we will relate the almost toric fibrations with the symplectic log Calabi-Yau divisors studied in the previous sections. Let's start with the toric actions, which of course could be viewed as a special type of almost toric fibrations.

4.1 Toric manifolds and Delzant polytope

An action of the 2-torus $\mathbf{T} \cong (S^1)^2$ on a symplectic 4-manifold (X, ω) is a homomorphism

$$\rho: \mathbf{T} \to \operatorname{Symp}(X, \omega)$$

from the torus to the symplectomorphism group of (X, ω) , such that the map $\rho^{\sharp} : \mathbf{T} \times X \to X$, defined by $\rho^{\sharp}(t, x) = \rho(t)(x)$, is smooth. An effective **T**-action with generating vector fields ξ_1, ξ_2 is Hamiltonian if there exists a moment map $\mu : X \to \mathbb{R}^2$ such that each component satisfies $d\mu_j = -\iota_{\xi_j}\omega$, for j = 1, 2. Such a **T**-action is called a toric action and could be seen as a smooth injective map $\rho : \mathbf{T} \to Ham(X, \omega)$. The 4-tuple $(X, \omega, \mathbf{T}, \mu)$ is called a symplectic toric manifold.

Definition 4.1. Two toric actions ρ_1, ρ_2 on a symplectic manifold (X, ω) are equivalent if there exists a symplectomorphism $\phi : X \to X$ and an automorphism $h : \mathbf{T} \to \mathbf{T}$ such that the diagram

$$\begin{array}{ccc} \mathbf{T} \times X & \stackrel{\rho_1^{\sharp}}{\longrightarrow} X \\ (h,\phi) & & \downarrow \phi \\ \mathbf{T} \times X & \stackrel{\rho_2^{\sharp}}{\longrightarrow} X \end{array}$$

is commutative.

Symplectic toric manifolds are interesting objects to study because much of their geometry and topology are determined by the combinatorial information of their moment image.

Definition 4.2. A Deltant polytope Δ in \mathbb{R}^2 is a polytope satisfying:

- (1) simplicity, i.e. there are 2 edges meeting at each vertex;
- (2) rationality, i.e. the edges meeting at each vertex p are of the form $p + tu_i, t \ge 0, u_i \in \mathbb{Z}^2$;
- (3) smoothness, i.e. for each vertex p, the corresponding u_1, u_2 can be choosen to be a \mathbb{Z} -basis of \mathbb{Z}^2 .

Recall that the moment image of a symplectic toric manifold is a Delzant polytope, which we call the moment polytope. Delzant's theorem ([Del88]) classifies equivalence classes of symplectic toric manifolds in terms of combinatorial data given by Delzant polytopes. In particular we have:

Proposition 4.3 ([KKP07]). Two toric action on (X, ω) are equivalent if and only if their moment map images are $AGL(2,\mathbb{Z})$ -congruent.

Let Δ be the moment polytope of a symplectic toric manifold $(X, \omega, \mathbb{T}, \mu)$ and e be an edge in the boundary $\partial \Delta$. Then the preimage $\mu^{-1}(e)$ is a symplectic sphere C_e . So the preimage $D = \mu^{-1}(\partial \Delta)$ is a cycle of symplectic spheres intersecting ω -orthogonally, which we call it the boundary divisor of Δ . Actually the Poincare dual of D is $c_1(X, \omega)$ ([Sym03], Proposition 8.2), so it is a symplectic log Calabi-Yau divisor.

We can read off the symplectic area and self-intersection of each component of D from the moment polytope as follows. Let $\bar{e} = re_0$ be a vector based at origin representing e, with r > 0 and e_0 a primitive vector. Then the symplectic area of C_e is $\omega([C_e]) = r$, which is also called the affine length of e. Suppose e_1, e_2 are the two edges intersecting e and denote by n_1, n_2, n the inward unit normal vector along e_1, e_2, e respectively. Then the self-intersection of C_e is equal to the unique integer s such that $n_1 + n_2 - sn = 0$.

4.2 Moment polytope and toric symplectic log Calabi-Yau divisors

The main result of the subsection is the following construction of a moment polytope from a toric symplectic log Calabi-Yau pair, which leads to the proof of Theorem 1.5 at the end of this subsection.

Proposition 4.4. Given an ω -orthogonal toric symplectic Calabi-Yau divisor $D \subset (X, \omega)$, there is a toric action on (X, ω) such that D is the boundary divisor. In particular, the map

$$\mathcal{T}(X,\omega) \to t\mathcal{LCY}(X,\omega)$$

taking a toric action to its boundary divisor is a surjection.

Define a set of primitive vectors in \mathbb{R}^2 as follows. Let $d_1 = \begin{pmatrix} 0 \\ -1 \end{pmatrix}$, $d_2 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and define $d_i = -s_{i-1}d_{i-1} - d_{i-2}$ for $i = 3, \ldots, k$. Such set of vectors is called a **generating set** associated to s = s(D). Recall for real numbers b_1, \ldots, b_k , we define the continued fraction as

$$[b_1, \dots, b_k] = b_1 - \frac{1}{b_2 - \frac{1}{b_3 - \frac{1}{\frac{\vdots}{b_{k-1} - \frac{1}{b_k}}}}}$$

If we write $d_i = \begin{pmatrix} x_i \\ y_i \end{pmatrix}$ with $gcd(x_i, y_i) = 1$, then it's easy to check $-\frac{x_i}{y_i} = [s_2, \dots, s_{i-1}]$ for $i = 3, \dots, k$. So the vector d_i is determined up to sign by the continued fraction $[s_2, \dots, s_{i-1}]$.

Remark 4.5. Special care is needed when either x_i or y_i is 0. We take the computation of continued fraction to be entirely formal so that the appearance of 0 doesn't affect the outcome. If $x_i = 0$, then y_i must be ± 1 and vice versa. This is because $x \pm \frac{1}{y} = \frac{0}{a}$ implies |x| = |y| = |a| = 1 and $x \pm \frac{1}{y} = \frac{a}{0}$ implies that y = 0 and |a| = 1. So each continued fraction corresponds uniquely to a primitive vector.

All indices below are taken to be modulo k.

Lemma 4.6. When D toric, we have $d_{i-1} + s_i d_i + d_{i+1} = 0$ for all i = 1, ..., k.

Proof. It suffices to prove the case for i = 1, k. Note that

$$d_{k-1} + s_k d_k + d_1 = 0 \quad \Leftrightarrow \quad [s_2, \dots, s_k] = 0$$
$$d_k + s_1 d_1 + d_2 = 0 \quad \Leftrightarrow \quad [s_1, \dots, s_{k-1}] = 0$$

We claim that when D is toric, $[s_{i+1}, s_{i+2}, \ldots, s_k, s_1, \ldots, s_{i-1}] = 0$ for all *i*. It is easily checked to be true when s = (1, 1, 1) or s = (0, n, 0, -n). So it suffices to prove that the continued fraction is invariance under toric blow-up, i.e.

$$[b_1,\ldots,b_i-1,-1,b_{i+1}-1,\ldots,b_k] = [b_1,\ldots,b_k].$$

Since continued fractions splits like $[b_1, \ldots, b_k] = [[b_1, \ldots, b_j], [b_{j+1}, \ldots, b_k]]$, it suffices to prove [x, y] = [x - 1, -1, y - 1], which is an easy computation.

Lemma 4.7. Let D be toric symplectic log Calabi-Yau divisor, a its area vector and $\{d_i\}$ its generating set. Then we have

- (1) $\sum a_i d_i = 0$,
- (2) $\det(d_i, d_{i+1}) := \det \begin{pmatrix} x_i & x_{i+1} \\ y_i & y_{i+1} \end{pmatrix} = 1.$
- (3) The angle between two consecutive vectors d_i, d_{i+1} is less than π . In particular, d_{i+1} is determined by either one of d_i, d_{i+2} and the continued fraction $[s_2, \ldots, s_i]$.

Proof. Since $b^+(D) = 1$, there exists $z \in \mathbb{R}^k$ such that $Q_D z = a$ ([LMM20], Proposition 5.13), i.e. $a_i = z_{i-1} + s_i z_i + z_{i+1}$ for all *i*. Then we have

$$\sum a_i d_i = \sum (z_{i-1}d_i + z_{i-1}d_i + z_i s_i d_i)$$

= $\sum z_i (d_{i-1} + s_i d_i + d_{i+1}) = 0.$

Note when i = 1, we have det $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = 1$. Then (2) follows from

$$\det(d_i, d_{i+1}) = \det(d_i, -s_i d_i - d_{i-1}) = \det(d_i, -d_{i-1}) = \det(d_{i-1}, d_i)$$

For (3) suppose θ is the angle between d_i, d_{i+1} , then we have $\sin \theta = \frac{\det(d_i, d_{i+1})}{||d_i|| \cdot ||d_{i+1}||} > 0$ and thus $\theta < \pi$. Recall that d_i is determined up to sign by $[s_2, \ldots, s_i]$. The sign ambiguity disappears when d_i is determined because of the angle restriction.

Lemma 4.8. Let \tilde{D} with self-intersection sequence $\tilde{s} = (s_1, \ldots, s_i - 1, -1, s_{i+1} - 1, \ldots, s_k)$ be the toric blow-up of D with self-intersection sequence $s = (s_1, \ldots, s_k)$. Then the corresponding generating set $\{\tilde{d}_i\}$ has the property that $\tilde{d}_j = d_j$ for $j \leq i$, $\tilde{d}_{i+1} = d_i + d_{i+1}$ and $\tilde{d}_j = d_{j-1}$ for $j \geq i + 1$. In particular, if D is toric, then its associated generating set of vectors winds around the origin exactly once.

Proof. The fact that $\tilde{d}_j = d_{j-1}$ for $j \ge i+3$ follows easily from the blow-up invariance of continued fraction proved in Lemma 4.6 and $\tilde{d}_j = d_j$ for $j \le i$ is trivial. Since $[s_2, \ldots, s_{i-1}, s_i - 1, -1] = [s_2, \ldots, s_i]$ and $\tilde{d}_{i+3} = d_{i+2}$, we have $\tilde{d}_{i+2} = d_{i+1}$ and $\tilde{d}_{i+1} = \tilde{d}_i + \tilde{d}_{i+2} = d_i + d_{i+1}$. For the second statement, let s be the self-intersection sequence of D. As seen above, toric blow-up inserts a new vector to the generating set in between two vectors and doesn't add to the winding number. So it suffices to see when s = (1, 1, 1) or (0, n, 0, -n), the associated vectors winds around the origin exactly once. But this is trivial because $k \le 4$ and angle between consecutive vectors is less than π .

Proof of Proposition 4.4. One can build a moment polytope for the induced toric action from divisor D as follows. Let $Q_1 = (0,0)$ and $Q_{i+1} = Q_i + a_i d_i$ for $i = 1, \ldots, k - 1$. Lemma 4.7 (1)(3) and Lemma 4.8 imply that the polygonal chain with vertices Q_1, \ldots, Q_k encloses a simple convex polygon P. This polygon P is actually a Delzant polygon by Lemma 4.7 (2). Then there is a symplectic toric manifold (X', ω') with moment polytope P and boundary divisor D'. From the construction we have (s(D'), a(D')) = (s(D), a(D)). So (X, ω, D) is strictly homological equivalent to (X', ω', D') by Lemma 2.37. Since D is ω -orthogonal and D' is ω' -orthogonal, by Proposition 2.10, there is a symplectomorphism between (X, ω, D) and (X', ω', D') . Composing this symplectomorphism with the moment map of (X', ω') , we get a toric action on (X, ω) such that D is the boundary divisor by construction.

Recall that

$$\mathcal{T}(X,\omega) = \{\rho: T^2 \to Ham(X,\omega)\} / \sim^3$$

is the set of equivalence classes of toric actions on (X, ω) and

$$t\mathcal{LCY}(X,\omega) = \{D \in p\mathcal{LCY}(X,\omega) | D \text{ is a toric divisor} \} / \sim^{s}$$

is the set of strictly symplectic deformation classes of toric symplectic log Calabi-Yau divisors in (X, ω) . Now we are ready to establish the correspondence between $\mathcal{T}(X, \omega)$ and $t\mathcal{LCY}(X, \omega)$.

Proof of Theorem 1.5. Consider the map $f : \mathcal{T}(X,\omega) \to t\mathcal{LCY}(X,\omega)$, where for each toric action $\rho: T^2 \to \text{Symp}(X,\omega)$ with moment map $\mu: X \to \mathbb{R}^2$ define $f(\rho)$ to be the boundary divisor D of its moment polygon $\mu(X)$. D is a cycle of symplectic spheres of length at least 3 and the homology class [D] is Poincare dual to the first Chern class $c_1(X,\omega)$. So (X,ω,D) is a symplectic Looijenga pair. Take a small collar neighborhood R of the boundary in $\mu(X)$, which lifts to a neighborhood P(D) of the divisor D. We might assume the origin (0,0) is in the interior of $\mu(X)$ by an affine translation. The outward radial vector field on \mathbb{R}^2 lifts to a Liouville vector field near the boundary $\partial P(D)$, which points into P(D). So $(P(D),\omega)$ is a concave neighborhood of D, which means D satisfies the positive GS criterion ([LM19]) and must have $b^+(Q_D) = 1$ ([LMM20]). The preimage of $\mu(X) - Int(R)$ is diffeomorphic to

 $D^2 \times T^2$. The boundary $\partial P(D)$ is T^3 and thus q(D) = 0 by Lemma 2.41. So D is a toric symplectic log Calabi-Yau divisor.

Let ρ, ρ' be two toric actions on (X, ω) with boundary divisors $D = f(\rho)$ and $D' = f(\rho')$. choose a cyclic labeling of both boundary divisors so that $D = f(\rho) = \bigcup C_i$ and $D' = f(\rho') = \bigcup C'_i$. Denote by (s, a) and (s', a') the self-intersection and area vectors of D and D' with respect some labeling. If ρ, ρ' are equivalent then their moment map images are $AGL(2,\mathbb{Z})$ -congruent ([KKP07]), which implies that (s, a) and (s', a') differ by cyclic and anti-cyclic permutations [KKP15]. So D and D' are strictly symplectic deformation equivalent by Lemma 2.37 and f is well-defined.

Suppose $D, D' \in t\mathcal{LCY}(X, \omega)$ are strictly homological equivalent. Note (s, a) and (s', a') depend only on the homology classes of components of D and D'. So they are the same up to cyclicly or anti-cyclicly relabeling the components. So the constructed moment polytopes in Proposition 4.4 are the same and the corresponding symplectic toric manifolds are equivariantly symplectomorphic. So f is injective.

Again by Proposition 4.4, f is also surjective and this finishes the proof.

With Theorem 1.5, properties of symplectic log Calabi-Yau divisors translate to properties of toric actions. Note that the proof of Theorem 1.6 also implies the stability of toric symplectic log Calabi-Yau divisors. Combined with Theorem 1.5 we get the following stability result for toric actions. Recall that S_{ω} denotes the set of ω -symplectic sphere classes.

Corollary 4.9. Let X be a rational surface. Suppose ω and ω' are two symplectic forms with $S_{\omega} = S_{\omega'}$. Then

$$\mathcal{T}(X,\omega) = \mathcal{T}(X,\omega').$$

4.3 Comparison with the counting results of Karshon-Kessler-Pinsonnault

In [KKP07], [KKP15] and [KKP], Karshon, Kessler and Pinsonnault first considered the question of counting inequivalent toric actions on symplectic four-manifolds. Their strategy relies on analyzing the combinatorics of Delzant polygons. Now since we have established the equivalence between toric divisors and toric actions, we could compare our counting results of divisors with their results.

Firstly, combining Theorem 1.5 with Corollary 2.34, we recover the following finiteness result of toric actions by Karshon, Kessler and Pinsonnault.

Corollary 4.10 ([KKP07]). A fixed symplectic rational surface only admits finitely many inequivalent toric actions.

By the general toric counting formula we could confirm the upper bound of toric actions

given by Karshon-Kessler-Pinsonnault. This is because according to Proposition 3.16:

$$\begin{split} |t\mathcal{LCY}(M_l,\omega_{\delta_1,\cdots,\delta_l})| &= \sum_{g\in\mathcal{G}_l} \frac{1}{2}\phi(F_l^4,g)\psi(\vec{\delta},g) \\ &= \sum_{g\in\mathcal{G}_l,g(2)=0} \frac{1}{2}\phi(F_l^4,g)\psi(\vec{\delta},g) + \sum_{g\in\mathcal{G}_l,g(2)=1} \frac{1}{2}\phi(F_l^4,g)\psi(\vec{\delta},g) \\ &\leqslant \sum_{g\in\mathcal{G}_l,g(2)=0} \frac{1}{2}\phi(F_l^4,g)\left[\frac{\delta_1}{1-\delta_1}\right] + \sum_{g\in\mathcal{G}_l,g(2)=1} \frac{1}{2}\phi(F_l^4,g)\left[\frac{\delta_1-\delta_2}{1-\delta_1}\right] \\ &= \frac{1}{2}(F_l^4(1)-2)F_l^4(2)\cdots F_l^4(l-1)\left[\frac{\delta_1}{1-\delta_1}\right] + \frac{1}{2}2F_l^4(2)\cdots F_l^4(l-1)\left[\frac{\delta_1-\delta_2}{1-\delta_1}\right] \\ &= \frac{(l+2)!}{4!}\left[\frac{\delta_1}{1-\delta_1}\right] + \frac{(l+2)!}{4!}\left[\frac{\delta_1-\delta_2}{1-\delta_1}\right] \end{split}$$

Corollary 4.11 ([KKP15]). The number of toric actions on (M_l, ω) is at most

$$(\lceil \frac{\delta_1}{1-\delta_1}\rceil + \lceil \frac{\delta_1-\delta_2}{1-\delta_1}\rceil) \cdot \frac{(l+2)!}{4!},$$

where $[\omega] = H - \delta_1 E_1 - \cdots - \delta_l E_l$ is a normalized reduced symplectic class.

Remark 4.12. By Proposition 3.7, for $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$ with $\delta_1 = \delta_2 > \delta_3$, the count is 3. Explicitly, up to equivalence, these are

$$(H - E_2 - E_3, E_3, E_2 - E_3, H - E_1 - E_2, E_1, H - E_1)$$

(E₃, H - E₂ - E₃, E₂, H - E₁ - E₂, E₁, H - E₁ - E₃)
(H - E₂, E₂ - E₃, E₃, H - E₁ - E₂ - E₃, E₁, H - E₁)

However, the count in [KKP] in this case is 5 by Corollary 8.7 (2) and Example 8.9 (3). The error comes from not taking into account the symmetry of switching E_1 and E_2 . Note there is a symplectomorphism switching E_1 and E_2 since $\delta_1 = \delta_2$. In other words, 5 is the result of counting the toric ones in pLCY but 3 is the result of counting the toric ones in LCY.

4.4 Almost toric fibrations

In this subsection, we introduce the basics of almost toric fibrations and their associated almost toric base diagrams. We refer to [Sym03], [LS10] and [Eva21] for details.

Definition 4.13 ([LS10], Definition 2.2). An almost toric fibration of a symplectic 4-manifold (X, ω) is a Lagrangian fibration $\pi : X \to B$ with only nodal and elliptic singularities. A toric fibration is a Lagrangian fibration induced by an effective Hamiltonian torus action. We denote by $\mathcal{ATF}(X, \omega)$ the set of almost toric fibrations on (X, ω) and write $\mathcal{ATF}(X) := \bigcup \mathcal{ATF}(X, \omega)$, where the union ranges over all symplectic forms ω of X.

The set B_0 of regular values of π carries an integral affine structure and there is a monodromy circling around a nodal singularity. With a suitable choice of cuts, we get an integral affine immersion $B - \bigcup$ cuts $\rightarrow \mathbb{R}^2$. The image P of this immersion, together with nodal rays and nodes, is called an **almost toric base diagram** representing the almost toric fibration, as an analogue to the moment polygon in the case of toric fibrations. The nodal rays are dotted rays representing the image of the cuts while the nodes are marks on the rays and represent focus-focus singularities.

There are three important surgery operations on the base diagram that fix the symplectomorphism type of the manifold. The first is a **nodal trade**, which replaces a neighborhood of a corank 2 elliptic singularity with a local model of a focus-focus singularity. The effect on the base diagram is that we insert a nodal ray from the toric vertex, with a node representing the focus-focus singularity. The second is a **nodal slide**, which moves the node along the ray on the base diagram (see Figure 4). Nodal trades and slides change the almost toric fibration but not the underlying manifold. The third operation is called a **mutation** with respect to a nodal ray. It changes the base diagram in the following way. The base diagram is sliced into two parts by the nodal ray. One part is unchanged while the other part is acted on by an affine transformation in $AGL(2,\mathbb{Z})$. Mutations change only the base diagram but not the almost toric fibration (see Figure 5).



Figure 4: Nodal trade and nodal slide.

Figure 5: Mutation.

Similar to the case of toric fibrations, the preimage $\pi^{-1}(\partial B)$ of the base is a symplectic divisor representing the Poincare dual of $c_1(X, \omega)$ (Proposition 8.2 of [Sym03]), i.e. a symplectic log Calabi-Yau divisor. We call it the boundary divisor of π .

As in toric fibrations, one can do toric blow-ups at elliptic corank 2 singularities of an almost toric fibration, which amounts to chopping corners centered at toric vertices on the base diagram (see Figure 6). There is another fibration compatible way to blow up, called an **almost toric blow-up**. Consider an edge on the base diagram, which we may assume is in the (1,0)-direction by an $SL(2;\mathbb{Z})$ transformation. An almost toric blow-up removes a right triangle with edge length ε and then adds a node at the top of the triangle with two dashed lines representing the cut (see Figure 7). Since each edge represents a symplectic sphere, this has the effect of a non-toric blow-up at the corresponding symplectic sphere with size ε .

The smoothing operation introduced in Section 2.2 can also be realized as a nodal trade on the base diagram. Near the toric vertex, the boundary divisor comprises two transversely intersecting symplectic discs. After a nodal trade, the boundary divisor becomes a symplectic annulus which is a smoothing of the pair of discs.

4.5 Realizing log Calabi-Yau divisors by ATF

For each almost toric fibration on a symplectic rational surface (X, ω) , its boundary divisor is a sympletic log Calabi-Yau divisor D, which has a homology type (D). So we have a map

$$\Phi: \mathcal{ATF}(X,\omega) \to \mathcal{LCY}(X,\omega),$$





Figure 6: Toirc blow up.

Figure 7: Almost toric blow up.

from the set of almost toric fibrations on (X, ω) to the set of symplectic log Calabi-Yau divisors.

In Engel's thesis ([Eng18], Section 5.1 Part 1), he showed that every holomorphic log Calabi-Yau divisor is realized as the boundary divisor of some almost toric fibration. Since every deformation class of log Calabi-Yau divisor is realized by some holomorphic divisor, we have the following result.

Proposition 4.14 ([Eng18]). The map $\tilde{\Phi} : \mathcal{ATF}(X) \to \widetilde{\mathcal{LCY}}(X)$ is surjective.

Remark 4.15. The above Proposition can also be derived from Lemma 2.15. This is simply because every element in $t \mathcal{LCY}(X)$ has a toric realization by Theorem 1.5 and the smoothing operations in the category of \mathcal{LCY} corresponds to the nodal trade operations in the category of \mathcal{ATF} .

Motivated by Engel's result, we raise the following conjecture on the realization of symplectic log Calabi-Yau divisors by almost toric fibrations.

Conjecture 4.16. The map $\Phi : \mathcal{ATF}(X, \omega) \to \mathcal{LCY}(X, \omega)$ is surjective.

Based on the observation that all log Calabi-Yau divisors in small rational manifolds are smoothing of toric log Calabi-Yau divisors and the fact that smoothing of boundary divisors can be realized by nodal trade in almost toric fibrations, we prove the following cases of Conjecture 4.16.

Proposition 4.17. The map $\mathcal{ATF}(X,\omega) \to \mathcal{LCY}(X,\omega)$ is surjective for $X = \mathbb{CP}^2, S^2 \times S^2$, $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ and $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$.

Proof. It suffices to show every homological configuration $(D) \in \mathcal{HLCY}(X, \omega)$ can be obtained from smoothing a toric homological configuration $(\overline{D}) \in t\mathcal{HLCY}(X, \omega)$. Then the corresponding almost toric fibration is obtained from the toric fibration of \overline{D} by nodal trades. Again it suffices to show this for normalized reduced symplectic classes. Also note that for $\mathbb{CP}^2, S^2 \times S^2$ and $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, all homological log Calabi-Yau divisors are listed in Theorem 2.14.

(1) Consider $X = \mathbb{CP}^2$ with symplectic class $[\omega_{FS}] = H$. Then

$$\mathcal{HLCY}(X,\omega_{FS}) = \{(3H), (2H,H), (H,H,H)\}$$

and each of them is a smoothing of (H, H, H).

(2) Consider $X = S^2 \times S^2$ with a standard basis $\{B, F\}$ and a normalized reduced symplectic class $[\omega_{\mu}] = B + \mu F$ with $\mu \ge 1$. Then $t\mathcal{HLCY}(X, \omega_{\mu}) = \{(B+kF, F, B-kF, F) \mid \mu > k \ge 0\}$ and the non-toric homological log Calabi-Yau divisors are

$$\{(B+(k+1)F,F,B-kF),(B+(k+2)F,B-kF) \mid \mu > k \ge 0\} \cup \{(B+F,B+F),(2B+2F)\}.$$

So it is clear they can be obtained from toric ones via smoothing.

(3) The case for $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ is similar to the case of $S^2 \times S^2$.

(4) Consider $X = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$ with a standard basis $\{H, E_1, E_2\}$ and a normalized reduced symplectic class $[\omega]$. Suppose $(D) \in \mathcal{HLCY}(X, \omega)$ a homological configuration which contains a (-n)-component for some $n \ge 2$. By Proposition 3.5, there is exactly one toric homological configuration (\overline{D}) with a (-n)-component. Suppose n = 2k for $k \ge 1$. Then the homological configuration (\overline{D}) is given by

$$(H - E_1, kH - (k - 1)E_1, H - E_1 - E_2, E_2, -(k - 1)H + kE_1 - E_2).$$

Note that smoothing of (\overline{D}) gives all the possible homological log Calabi-Yau divisors with a (-n)-component, as listed in the proof of Proposition 3.5:

- $((k+1)H kE_1, H E_1 E_2, E_2, -(k-1)H + kE_1 E_2),$ $(H - E_1, kH - (k-1)E_1, H - E_1, -(k-1)H + kE_1 - E_2),$ $(H - E_1, (k+1)H - kE_1 - E_2, E_2, -(k-1)H + kE_1 - E_2)$
- $((k+2)H (k+1)E_1 E_2, E_2, -(k-1)H + kE_1 E_2),$ $((k+1)H - kE_1, H - E_1, -(k-1)H + kE_1 - E_2)$
- $((k+2)H (k+1)E_1, -(k-1)H + kE_1 E_2).$

In particular, (D) is a smoothing of (\overline{D}) . The case of n = 2k - 1 for $k \ge 1$ is similar. They are all smoothings of the toric configuration

$$(H - E_1, (k+1)H - kE_1 - E - 2, E_2, H - E_1 - E_2, -kH + (k+1)E_1).$$

The remaining homological log Calabi-Yau divisors are

- $(2H E_2, H E_1),$
- $(3H E_1 2E_2, E_2),$
- $(H, H E_1 E_2, H),$

which are $[\omega]$ -positive for any normalized reduced symplectic class $[\omega]$. They are smoothings of the toric homological configuration

$$(H - E_2, H - E_1, E_1, H - E_1 - E_2, E_2),$$

which is also $[\omega]$ -positive for all normalized reduced symplectic class.

Remark 4.18. Not all symplectic log Calabi-Yau divisors can be obtained from smoothing toric ones. There could be some symplectic classes outside the toric cone so that they don't admit toric divisors. For example, $M_4 = \mathbb{CP}^2 \# 4\overline{\mathbb{CP}}^2$ with reduced symplectic class $\delta_1 = \delta_2 =$ $\delta_3 = \delta_4 < \frac{1}{4}$. There is no toric divisor in this symplectic class, but there are LCYs such as $(H - E_1 - \cdots - E_4, H, H)$. Moreover, even if $[\omega]$ is in the toric cone, there could exsit some LCYs which don't come from the smoothing. For instance, $M_3 = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$ with reduced symplectic class $(\delta_1, \delta_2, \delta_3) = (\frac{6}{15}, \frac{5}{15}, \frac{4}{15})$ which is in the region (8) of Proposition 3.7 for i = 1. So the count of toric LCY is 4:

$$(E_1 - E_2, E_2, H_{12}, H_3, E_3, H_{13})$$

$$(E_1 - E_3, E_3, H_{13}, H_2, E_2, H_{12})$$
$$(E_2 - E_3, E_3, H_{23}, H_1, E_1, H_{12})$$
$$(H_{12}, E_1, H_{13}, E_3, H_{23}, E_2)$$

However $(H_{12}, E_2 - E_3, E_1 - E_2, H_1, H)$ is a LCY under this symplectic class, which can not be obtained by smoothing the above toric LCYs.

As remarked above, the smoothing operation, which corresponds to nodal trade in ATF, is not enough to generate all the realization of LCY by ATF, even for M_3 . Nevertheless, when we take almost toric blow up into consideration, we are able to construct ATF realization for divisors in M_3 .

Proposition 4.19. The map $\mathcal{ATF}(X, \omega) \to \mathcal{LCY}(X, \omega)$ is surjective for $X = M_3$.

Proof. Suppose $D \in \mathcal{LCY}(M_3, \omega)$, $[\omega]$ is normalized and reduced with $\omega(E_i) = \delta_i$. Then D is either the toric blow up or non-toric blow up of some $D' \in \mathcal{LCY}(M_2, \omega')$. By the previous proposition, D' has ATF realization as the nodal trade of the toric one

$$(-kH + (k+1)E_1, H_1, (k+1)H - kE_1 - E_2, E_2, H_{12})$$

for some $k \in \mathbb{Z}$. Thus we can assume D' is realized by the almost toric base diagram which is a Delzant polygon with branches at some vertices which could be sufficiently short by nodal slides. See Figure 8.



Figure 8: Possible ATF realizations for M_2 .

In the following, we will assume for $k \ge 0$. The case when k < 0 is totally similar and will be omitted. Let's firstly consider the case when D is the non-toric blow up on the component C' of D'. Note that exactly one of the following two cases will happen:

• If C' doesn't come from the smoothing of two (or more) toric components, or comes from the smoothing involving the component H_1 or $(k+1)H - kE_1 - E_2$ in the toric divisors, we don't need to do mutations. By the reducedness conditions that $\delta_3 \leq \delta_2, \delta_3 \leq 1 - \delta_1 - \delta_2$, there always exists enough space for performing almost toric blow up, which realizes the non-toric blow up operation for LCYs. See Figure 9.



Figure 9: The case when the blow up size is smaller than the length of one edge.

If C' comes from the smoothing involving the component H₁₂ in the toric divisors, when 1 = δ₁ + δ₂ + δ₃ we can use two types of mutations to create enough space for almost toric blow up. To be more precise, when C' comes from the smoothing involving both H₁₂ and E₂, we can apply the mutation trick shown in the left graph of Figure 10; when C' comes from the smoothing involving both H₁₂ and -kH + (k + 1)E₁, we can apply the mutation trick shown in the right graph of Figure 10.



Figure 10: Two kinds of mutations creating enough space for almost toric blow up.

Next we assume D is the toric blow up of D', and we want to perform corner chopping of size δ_3 at some vertex P in the almost toric base diagram.

Since ω is reduced, we have the following easy observation: the size δ_3 can not be larger than the sum of affine lengths of two adjacent edges in the Delzant polygon. This would imply it suffices to consider the following two cases:

- If two edges having P as endpoint both have lengths larger than δ_3 , we could just perform nodal slides such that the branches don't intersect the corner of size δ_3 we want to chop.
- If one of two edges having P as endpoint is the component H_{12} with $1 = \delta_1 + \delta_2 + \delta_3$ or E_2 with $\delta_2 = \delta_3$, then we can apply the reflection along H_{123} or $E_2 - E_3$ (note that the conditions on $\vec{\delta}$ guarantee the reflection can be realized by symplectomorphisms) to the divisor D to obtain an equivalent one. In these two cases, D contains no component of H_{12} or E_2 which implies there should be no E_3 component in the equivalent divisor given by the reflection. It follows that we can also view D as the non-toric blow up of some divisor in M_2 . Therefore the above discussion for non-toric blow up cases gives the realization.
- If one of two edges having P as endpoint is the component $-kH + (k+1)E_1$ with $-k + (k+1)\delta_1 \leq \delta_3$, then the other edge having P as endpoint is either H_{12} or H_1 .



Figure 11: A special situa-

tion that mutation is not al-Figure 12: Two kinds of mutations creating enough space for toric lowed.

In the former case, there might a very special situation when $k = 0, \delta_1 = \delta_2 = \delta_3$. It's shown in Figure 11 that the mutation is forbidden since the eigenray will intersect with another vertex. Nevertheless, we could apply the reflection along $E_1 - E_3$ to reduce it to the non-toric blow up cases as what we did in the last paragraph. Other than this special situation, we could always do the mutations to create enough space for corner chopping. The same is true for the latter case. These are shown in Figure 12.

4.6 Connectedness of toric fibrations

In [Sym03], Symington observed that different toric fibrations on $(S^2 \times S^2, \omega)$ can be changed to each other via nodal trades, nodal slides and mutations, which gives a path of almost toric fibrations connecting toric fibrations. This observation led to the following conjecture.

Conjecture 4.20 ([Sym03]). Any two toric fibrations of (M, ω) can be connected by a path of almost toric fibrations of (M, ω) .

Based on the counting results for toric divisors, we prove Symington's conjecture for \mathbb{CP}^2 , $S^2 \times S^2$, $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, $\mathbb{CP}^2 \# 2 \overline{\mathbb{CP}}^2$ and $\mathbb{CP}^2 \# 3 \overline{\mathbb{CP}}^2$. We also prove a weaker version of this conjecture for general rational surfaces.

From a toric fibration π_0 , we could perform the following series of operations to get a path π_t of almost toric fibrations such that π_1 is another toric fibration. We perform first a nodal trade at a vertex to get a node, slide the node along the eigenray further away from the original vertex, then perform a mutation along this eigenray, which creates a new vertex where the eigenray intersects some edge of the base diagram. If the new vertex is smooth (i.e. satisfies the smoothness condition in Definition 4.2), then one can slide the node towards this vertex and use a nodal trade to get back to a toric fibration. The process is illustrated in Figure 13, which we call a **toric mutation** at the chosen vertex. In effect, it cuts the moment polygon into two parts along the eigenray at this vertex, applies an $AGL(2,\mathbb{Z})$ transformation to one part and glues it back to form a new moment polygon. See Section 2 of [CV20] for a more detailed account.

Proposition 4.21. Symington's conjecture is true for \mathbb{CP}^2 , $S^2 \times S^2$, $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$, $\mathbb{CP}^2 \# 2 \overline{\mathbb{CP}}^2$ and $\mathbb{CP}^2 \# 3 \overline{\mathbb{CP}}^2$.



Figure 13: Toric mutation (= nodal trade + mutation + nodal trade) gives a path of almost toric fibrations.

Proof. It suffices to consider normalized reduced symplectic forms on these rational surfaces.

(1) For $X = \mathbb{CP}^2$, there is only one toric fibration.

(2) For $X = S^2 \times S^2$ with standard basis $\{B, F\}$ and a normalized reduced symplectic class $[\omega] = B + \mu F$, the possible Delzant polygons are $\beta(2n)$ as in Figure 14, where $n \in \mathbb{Z}_{\geq 0}$. By toric mutation along the eigenray at the lower right vertex, we could always connect $\beta(2n)$ and $\beta(2n+2)$.



Figure 14: Toric mutations for Delzant polygons of $S^2 \times S^2$.

(3) For $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ with standard basis $\{H, E\}$ and a normalized reduced symplectic class $[\omega] = H - \mu E$, the possible Delzant polygons are $\alpha(2n+1)$ as in Figure 15, where $n \in \mathbb{Z}_{\geq 0}$. Again, by toric mutation we can connect $\alpha(2n+1)$ and $\alpha(2n+3)$.



Figure 15: Toric mutations for Delzant polygons of $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$.

(4) For $X = \mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$ with a normalized reduced symplectic class $[\omega] = H - \delta_1 E_1 - \delta_2 E_2$. By Corollary 3.6, the toric fibrations are given by the family $\gamma(n)$ for $n \in \mathbb{Z}_{\geq 0}$ shown in Figure 16. When n = 2k, $\gamma(2k)$ denotes $((k+1)H - kE_1 - E_2, H_1, -kH + (k+1)E_1, H_{12}, E_2)$; when n = 2k + 1, $\gamma(2k + 1)$ denotes $((k + 1)H - kE_1, H_1, -kH + (k + 1)E_1 - E_2, E_2, H_{12})$. We see that by toric mutation along the eigenray at the lower right vertex, we can connect $\gamma(2k)$ and $\gamma(2k + 2)$ or $\gamma(2k + 1)$ and $\gamma(2k + 3)$. Moreover, by toric mutation at the lower left vertex, we can connect $\gamma(0)$ and $\gamma(1)$.



Figure 16: Toric mutations for Delzant polygons of $\mathbb{CP}^2 \# 2\overline{\mathbb{CP}}^2$.

(5) For $X = \mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$, we claim if the toric fibrations come from chopping different corners of the same pentagon $\gamma(n)$, then by toric mutation, they can be connected. If the corners are adjacent, this is shown in Figure 25 in the next section; if not, this is also true simply because in a pentagon whenenver we can perform corner chopping at two non-adjacent corners, we can also perform corner chopping at another corner adjacent to both of them.

Therefore, it suffices to consider the family $\theta(n)$ for $n \in \mathbb{Z}_{\geq 0}$, where $\theta(2k)$ denotes $((k + 1)H - kE_1 - E_2 - E_3, E_3, H_{13}, -kH + (k + 1)E_1, H_{12}, E_2)$ and $\theta(2k + 1)$ denotes $((k + 1)H - kE_1 - E_3, E_3, H_{13}, -kH + (k + 1)E_1 - E_2, E_2, H_{12})$. Thus the Delzant polygons of $\theta(n)$ is obtained by chopping the lower right corner of the Delzant polygons of $\gamma(n)$ in Figure 16. The reason why $\theta(0)$ and $\theta(1)$ can be connected is the same as (4). Moreover, note that if there exists toric fibration $\theta(2k + 2)$, the toric fibration $\theta'(2k)$ given by $((k + 1)H - kE_1 - E_2, H_{13}, E_3, -kH + (k + 1)E_1 - E_3, H_{12}, E_2)$ should also exist by the assumption that ω is reduced. In the Figure 17 we show that $\theta'(2k)$ can be connected to $\theta(2k + 2)$ and $\theta'(2k)$ can be connected to $\theta(2k + 3)$ and $\theta(2k + 1)$ can be connected. Consequently all the possible toric fibrations $\theta(n)$ can all be connected.



Figure 17: Toric mutations for Delzant polygons of $\mathbb{CP}^2 \# 3\overline{\mathbb{CP}}^2$.

4.7 Realization and connectedness in the restrictive region

Recall that the restrictive reduced symplectic classes defined in Section 3.5 form a region in the normalized reduced symplectic cone, which is cut out by the inequalities:

$$1 > \delta_1 + \dots + \delta_l$$
$$\delta_1 > \delta_2$$
$$\delta_k > \delta_{k+1} + \dots + \delta_l$$

for all $2 \le k \le l-1$. One can think of this region as an explicit description of blow up with sufficiently small sizes which has many good properties. In this section we will prove Conjecture 4.16 and 4.20 hold in this region.

Let's firstly discuss Conjecture 4.16. From the counting result Proposition 3.16 we see that in the restrictive region, all the divisors come from the toric and non-toric blow ups of some germs. Our strategy is to choose some appropriate ATF realizations of those germs and then realize each toric or non-toric blow up by toric or almost toric blow up on the almost toric base diagram. In the proof of Proposition 4.19 we need to apply toric mutations to overcome the issue that there might be the edge whose length is less than the blow up size (this happens only when $\delta_2 = \delta_3$ or $1 = \delta_1 + \delta_2 + \delta_3$). However, when the restrictive condition is added, we will see there is no need to do toric mutations if we select almost toric base diagrams for germs wisely. So we next only need to check whether there is enough space on the diagram to perform toric or almost toric blow up.

It's worth noting that for an almost toric blow up, there are different presentations of excising an affine triangle of on the edge which differ by $AGL(2,\mathbb{Z})$ transformations (See Figure 18). The only requirement is the directions of any two edges form a \mathbb{Z} -basis. We say that an almost toric triangle of size δ can be embedded along an edge if it is possible to excise a size δ affine triangle in one of such forms.



Figure 18: Different embeddings of almost toric triangles.

Lemma 4.22. Suppose there are distinct positive numbers $\delta_1 > \cdots > \delta_k, \delta_1 > \delta_{1,1} > \cdots > \delta_{1,\alpha_1}, \cdots, \delta_k > \delta_{k,1} > \cdots > \delta_{k,\alpha_k}$ such that if we order them in the decreasing manner, they satisfy the restrictive condition. Let Δ be the region obtained by performing corner choppings of sizes $\delta_1, \cdots, \delta_k$ on the moment map image Δ_0 of the unit open ball in \mathbb{C}^2 . Then there is a way to embed almost toric triangles of sizes $\delta_{i,1}, \cdots, \delta_{i,\alpha_i}$ along the edge coming from corner chopping of size δ_i for all $1 \leq i \leq k$ such that they are pairwisely disjoint.

Proof. Assume Δ_0 is the region given by $\{(x, y) | x \ge 0, y \ge 0, x + y < 1\}$. Let $\vec{u}_0 = (1, 0)$ be a vector. After chopping the corner of size δ_1 , we will get a new region $\Delta_1 = \{(x, y) | x \ge 0, y \ge 0, \delta_1 \le x + y < 1\}$, two vertices $A_1 = (0, \delta_1), B_1 = (\delta_1, 0)$ and a unit affine length vector

 $\vec{u}_1 = \delta_1^{-1} \overline{A_1 B_1}$. For the remaining corner chopping, it makes sense to say whether it is on the left or right of the first corner (A_1 side or B_1 side). Let $L_1, R_1 \subset \{2, \dots, k\}$ denote the left and right ones respectively. Now by the restrictive condition, it's always possible to choose two points P_1, Q_1 on the segment A_1B_1 such that the affine lengths of $\overline{A_1P_1}, \overline{P_1Q_1}, \overline{Q_1B_1}$ are greater than $\sum_{i \in L_1} (\delta_i + \delta_{i,1} + \dots + \delta_{i,\alpha_i}), \delta_{1,1} + \dots + \delta_{1,\alpha_1}, \sum_{i \in R_1} (\delta_i + \delta_{i,1} + \dots + \delta_{i,\alpha_i})$ respectively. We then define the parallelogram region Θ_1 attached to the segment A_1B_1 by $\{z \in \mathbb{R}^2 \mid \overline{Q_1z} = a\vec{u}_0 + b\overline{Q_1P_1}, a \in [0, \delta_{1,1}], b \in [0, 1]\}$. Note that by our choice of P_1, Q_1, Θ_1 will be inside the region Δ_1 and we can embed the pairwisely disjoint almost toric triangles of sizes $\delta_{1,1}, \dots, \delta_{1,\alpha_1}$ into Θ_1 . See Figure 19.

After performing the second corner chopping of size δ_2 , we will get the region Δ_2 . Similarly as above, we can define two new vertices A_2, B_2 (A_2 is on the left of B_2), unit affine length vector $\vec{u}_2 = \delta_2^{-1} \overline{A_2 B_2}$, left and right remaining corner chopping indices $L_2, R_2 \subset \{3, \dots, k\}$, P_2, Q_2 on the segment $A_2 B_2$ satisfying the affine lengths of $\overline{A_2 P_2}, \overline{P_2 Q_2}, \overline{Q_2 B_2}$ are greater than $\sum_{i \in L_2} (\delta_i + \delta_{i,1} + \dots + \delta_{i,\alpha_i}), \delta_{2,1} + \dots + \delta_{2,\alpha_2}, \sum_{i \in R_2} (\delta_i + \delta_{i,1} + \dots + \delta_{i,\alpha_i})$ respectively. The selection of the parallelogram region Θ_2 attached to the segment $A_2 B_2$ needs more words. When $2 \in L_1$, that is, the second corner chopping is at A_1 , we define Θ_2 to be $\{z \in \mathbb{R}^2 \mid \overline{Q_2 z} = a \vec{u}_1 + b \overline{Q_2 P_2}, a \in [0, \delta_{2,1}], b \in [0, 1]\}$; when $2 \in R_1, \Theta_2$ will be $\{z \in \mathbb{R}^2 \mid \overline{Q_2 z} = a \vec{u}_0 + b \overline{Q_2 P_2}, a \in [0, \delta_{2,1}], b \in [0, 1]\}$. See Figure 20.



Figure 19: Embedding the first parallelogram Figure 20: Two cases of embedding the second parallelogram region Θ_1 .

We could repeat this procedure for all the regions $\Delta_3, \dots, \Delta_k = \Delta$ obtained by the remaining corner choppings. The only thing that needs to be mentioned is the choice of Θ_j . Assume from Δ_{j-1} to Δ_j , the new edge $A_j B_j$ come from the chopping the corner of Δ_{j-1} whose right edge has the unit affine length direction $\vec{u}_{\phi(j)}$ ($\phi(j)$ can be an arbitrary positive integer less than j). We will let $\Theta_j = \{z \in \mathbb{R}^2 \mid \overrightarrow{Q_j z} = a \vec{u}_{\phi(j)} + b \overrightarrow{Q_j P_j}, a, b \in [0, 1]\}$. Note that this choice is consistent with the requirement of Δ_2 in the above paragraph.

Therefore we get these parallelogram regions $\Theta_1, \dots, \Theta_k$. It's easy to see from the restrictive condition and our choice that Θ_j will not intersect the corner we will chop away from Δ_j to get Δ_{j+1} . So these parallelogram regions actually all live in the final region $\Delta_k = \Delta$. To finish the proof of this lemma, it suffices to show these Θ_j 's are pairwisely disjoint by our construction. This can be verified by the observation that, for any i < j, Θ_i and Θ_j will be contained in two different half planes divided by the line ℓ of the opposite sides of P_jQ_j in the parallelogram Θ_j . Note that we can consider the line of the segment ℓ' and the intersection point S between ℓ and ℓ' . By our choice of P_i, Q_i , if Θ_i is on the left (resp. right) of Θ_j , then S must be on the left of P_i (resp. right of Q_i). See Figure 21.



Figure 21: Θ_j is on the left or right of Θ_i . In both cases the line ℓ will always separate two parallelogram regions.

Now let's choose the almost toric base diagrams for the germs in the proof of Proposition 3.16. We can put $((k+1)H - kE_1, H_1, -kH + (k+1)E_1, H_1), ((k+2)H - (k+1)E_1, -kH + (k+1)E_1, H_1), ((k+3)H - (k+2)E_1, -kH + (k+1)E_1)$ with $k \ge 0$ together since their almost toric base diagrams only differ by nodal trades, see Figure 22. The remaining germs are $(H, H, H_1), (2H - E_1, H), (2H, H_1)$ and $(3H - E_1 - \cdots - 2E_j, E_j)$ whose almost toric base diagrams are shown in Figure 23. Note that we have given the ATF realizations we will use for all the germs.

Proposition 4.23. For any rational surface (M_l, ω) , if $[\omega]$ is in the restrictive region, then the map $\Phi : \mathcal{ATF}(M_l, \omega) \to \mathcal{LCY}(M_l, \omega)$ is surjective.

Proof. For the first diagram in Figure 22, let $X_1, X_2, X_3, X_4 \subset \{2, \dots, l\}$ be the set of indices of non-toric blow ups over the four edges and $Y_1, Y_2, Y_3, Y_4 \subset \{2, \dots, l\}$ be the set of indices of toric blow ups at four corners. Again, similar to the proof of Lemma 4.22, by the restrictive condition we can embed pairwisely disjoint parallelogram regions Φ_p with affine width $\sum_{i \in X_p} \delta_i$, affine height $\max_{i \in X_p} \delta_i$ and triangles Ψ_p of sizes $\sum_{i \in Y_p} \delta_i$ for all $1 \leq p \leq 4$. Then we apply Lemma 4.22 to the regions inside Ψ_p to realize all the remaining non-toric blow ups over the edges with indices in Y_p by almost toric blow ups. It follows that we are able to create the almost toric base diagram for the ultimate divisor, see Figure 24. The other two cases in Figure 22 and the cases in 23 can be settled in the same way. Observe that in each of those cases, the restrictive condition guarantees each edge has enough length to perform blow ups so that we don't need to do toric mutations. Also, by nodal slides, we can make the nodal points close enough the vertices so that the nodal rays don't intersect with those parallelogram or triangle regions.





Figure 23: ATF realizations of the remaining germs.

Figure 22: ATF realizations of some germs.



Figure 24: Arrangement of Ψ_p , Φ_p regions. When we zoom into the ψ_p region we will get the diagram that appears in Lemma 4.22.

Now let's prove Conjecture 4.20 in the restrictive region.

Lemma 4.24. Suppose $\vec{\delta}' = (\delta_1, \dots, \delta_k)$ and $\vec{\delta} = (\delta_1, \dots, \delta_k, \delta_{k+1})$ are in the restrictive regions. Let $\pi' : (X, \omega_{\vec{\delta}'}) \to \Delta'$ be a toric fibration. Then if $\pi_i : (X \# \overline{\mathbb{CP}}^2, \omega_{\vec{\delta}}) \to \Delta_i$ (i = 1, 2) are two toric fibrations obtained from performing corner chopping at different corners of Δ' , then they can be connected by a path of almost toric fibrations.

Proof. By the assumption of $\vec{\delta} = (\delta_1, \dots, \delta_k, \delta_{k+1})$ being restrictive, it's immediate to see from the list in $t\mathcal{H}_k$ before Corollary 2.33 that the only possible homology class of the boundary divisor's component of Δ' with symplectic area $\leq \delta_{k+1}$ must be

$$-aH + (a-1)E_1 - \sum_{i=2}^k \epsilon_i E_i, a \in \mathbb{Z}_+, \epsilon_i \in \{0,1\}$$

This implies there is at most one edge of Δ' whose length is not larger than δ_{k+1} . Consequently it suffices to show that corner chopping at two adjacent corners can be connected. And this can be achieved by toric mutation as in Figure 25.



Figure 25: Adjacent corner choppings differ by a mutation.

Figure 26: The eigenray ℓ divides the plane into two parts and must touch another edge.

Lemma 4.25. Under the assumption of Lemma 4.24 with additional hypothesis $k \ge 3$, if $\pi': (X, \omega_{\vec{x}}) \to \Delta'$ can be connected to another toric fibration by mutation along an eigenray ℓ , then it's always possible to chop some corner with size δ_{k+1} not intersecting ℓ .

Proof. Suppose ℓ divides the plane into two open half-planes S_1, S_2 , which contains k_1, k_2 vertices of the Delzant polygon Δ' respectively. Then it's immediate to see that $k_1 + k_2 = k + 2$ since ℓ does not pass through another vertex when the mutation is allowed. This is shown in Figure 26. Remember that the restrictive condition guarantees there is at most one edge of Δ' whose length is not larger than δ_{k+1} . Assuming ℓ intersect the edge L, a triangle of size δ_{k+1} can always embedded at the corners of those $k_1 + k_2$ vertices which are not the endpoints of L or the shortest edges. Note that there will be at least $k + 2 - 2 - 2 = k - 2 \ge 1$ such corners where we can perform corner chopping of size δ_{k+1} without intersecting ℓ .

Proposition 4.26. For rational surface $(M_l, \omega_{(\delta_1, \dots, \delta_l)})$, if $(\delta_1, \dots, \delta_l)$ is in the restrictive region, then any two toric fibrations can be connected by a path of almost toric fibrations. As a result, on (X, ω) with arbitrary ω , any two toric fibrations are symplectic deformation equivalent to some toric fibrations on (X, ω') (in the sense of their boundary divisors) which can be connected through a path of almost toric fibrations.

Proof. By Proposition 4.21, we only need to look at the cases when $l \ge 4$. By induction we will assume any two toric fibrations on $(M_{l-1}, \omega_{(\delta_1, \dots, \delta_{l-1})})$ can be connected by toric mutations and prove the case for l. For any two toric fibrations π, π' on $(M_l, \omega_{(\delta_1, \dots, \delta_l)})$ which come from performing the corner chopping of two toric fibrations $\overline{\pi}, \overline{\pi}'$ on $(M_{l-1}, \omega_{(\delta_1, \dots, \delta_{l-1})})$ respectively, there will be a sequence of toric fibrations $\pi_1 = \overline{\pi}, \pi_2, \cdots, \pi_{n-1}, \pi_n = \overline{\pi'}$ on $(M_{l-1}, \omega_{(\delta_1, \cdots, \delta_{l-1})})$

such that π_i and π_{i+1} can be connected by performing one toric mutation for all $1 \leq i \leq n-1$. By Lemma 4.25, there exist toric fibrations $\tilde{\pi}_i, \tilde{\tilde{\pi}}_i$ on $(M_l, \omega_{(\delta_1, \dots, \delta_l)})$ which are toric blow up of π_i, π_{i+1} and can be connected. As a result, by Lemma 4.24 we see that $\tilde{\pi}_1, \tilde{\tilde{\pi}}_{n-1}$ can be connected and π, π' can be connected to $\tilde{\pi}_1, \tilde{\tilde{\pi}}_{n-1}$ respectively. Therefore we finally obtain that π can be connected to π' .

Now given two toric fibrations on (X, ω) with toric divisors D_1, D_2 , by Lemma 3.13, there are always symplectic deformations (ω_i^t, D_i^t) such that $\omega_i^0 = \omega$ and $\omega_i^1 = \omega'$ (i = 1, 2) with ω' satisfying the restrictive condition. Then the the toric fibrations on X corresponding to $(\omega', D_1^1), (\omega', D_2^1)$ can be connected by a path of almost toric fibrations.

A Appendix

A.1 More discussions on tautness

In this subsection, we study the def-tautness. Hopefully the result can be applied to study the problem of classifying symplectic fillings of torus bundles in the future.

Labeled holomorphic anticanonical pairs are in general not taut ([Fri16]). The following example of Golla and Lisca shows that the labeled symplectic log Calabi-Yau divisors are also in general not def-taut.

Example A.1 (Proposition 3.6 of [GL16]). The cycle of spheres with self-intersections

$$(1, -2, -3, -3, -2, -3, -2)$$

admits two different symplectic embeddings (with possibly different symplectic structures) in $X = \mathbb{CP}^2 \# 9 \overline{\mathbb{CP}}^2$ as symplectic log Calabi-Yau divisors (ω_1, D_1) and (ω_2, D_2) . Golla and Lisca showed that their complements have different intersection forms and thus not homotopy equivalent. So the two divisors are not symplectic deformation equivalent (in both labeled and unlabeled senses).

The example given by Golla and Lisca is included in the classification of log Calabi-Yau divisors with $b^+ = 1$ up to toric equivalence in Theorem 1.4 (4) of [LMM20]. We can actually give a criterion of def-tautness for this type of divisors. Recall that in Theorem 1.4 (4) of [LMM20], a blown-up $(1, 1 - p_1, -p_2, \dots, -p_{l-1}, 1 - p_l)$ means this self-intersection sequence is obtained by toric blow up of (1, 1, 1) and then followed by non-toric blow up. We could give an equivalent description of this type of sequences following [Lis08]. Let

$$\mathcal{C} = \{ (n_1, \cdots, n_k, n_{k+1}) : \sum_{i=1}^{k+1} n_i = 3(k-1), [n_1, \cdots, n_k] = 0, (n_1, \cdots, n_k) \text{ is admissible} \}$$
$$\mathcal{C}' = \{ (n_1, \cdots, n_k, n_{k+1}, a_1, \cdots, a_{k+1}) : (n_1, n_2, \cdots, n_k, n_{k+1}) \in \mathcal{C}, a_i \in \mathbb{Z}_{\ge 0}, n_i + a_i \ge 2 \}$$

where $[n_1, \dots, n_k]$ denotes the continued fraction and 'admissible' means each denominator appearing in the continued fraction is positive. By Lemma 2.2 in [Lis08], there is a surjective map

$$\Phi: \mathcal{C}' \to \{\text{blown-up } (1, 1-p_1, -p_2, \cdots, -p_k, 1-p_{k+1}), p_i \ge 2\}$$

sending $(n_1, \dots, n_k, n_{k+1}, a_1, \dots, a_{k+1})$ to $(1, 1 - (n_1 + a_1), -(n_2 + a_2), \dots, -(n_k + a_k), 1 - (n_{k+1} + a_{k+1}))$. For instance, (1, -2, -3, -3, -2, -3, -2) in the above example is equal to $\Phi(2, 1, 3, 2, 1, 3, 1, 2, 0, 0, 2, 0)$ or $\Phi(3, 1, 3, 1, 3, 1, 0, 2, 0, 1, 0, 2)$.

Proposition A.2. A labeled symplectic log Calabi-Yau divisor $(D = (C_0, C_1, \dots, C_l), \omega)$ with blown-up self-intersection sequence $(1, 1 - p_1, -p_2, \dots, -p_{l-1}, 1 - p_l)$ such that all $p_i \ge 2$ is def-taut if and only if the preimage of the self-intersection sequence under Φ has only one element.

Proof. Firstly we show that the homology class of any component of self intersection 1 in a symplectic log Calabi-Yau divisor with length ≥ 2 can be mapped to the standard line class H by some $\gamma \in D(X)$, where D(X) is the image of $Diff^+(X) \to Aut(H^2(X;\mathbb{Z}))$. Note that with standard basis H, E_1, \dots, E_l , the reflections along $E_i - E_j, H - E_i - E_j - E_k$ are in D(X). By Corollary 2.33, we may assume the self intersection 1-component is one of the following

$$kH - (k-1)E_1 - \sum_{i=2}^{l} \epsilon_i E_i, k \in \mathbb{Z}, \epsilon_i \in \{0, 1\}$$
$$2H - \sum_{i=2}^{l} \epsilon_i E_i, \epsilon_i \in \{0, 1\}$$
$$3H - E_1 - \dots - E_{p-1} - 2E_p - \sum_{i=p+1}^{l} \epsilon_i E_i, 2 \le p \le l, \epsilon_i \in \{0, 1\}$$

For the second and the third cases, we can firstly perform the reflections along some $E_1 - E_i$ for the *i* such that $\epsilon_i = 1$ or $E_1 - E_p$, so that these homology classes can be transformed into the first type. For the first type, we could perform the reflections along $H - E_1 - E_i - E_j$ for *i*, *j* such that $\epsilon_i = \epsilon_j = 1$ to decrease *k* by 1. By repeating this procedure we will get k = 1and the homology class must be *H*.

Since the symplectic deformation equivalent class will not changed by a diffeomorphism, by the above claim we may just assume $[C_0] = H$. By Proposition 4.4 in [Lis08], it follows that $[C_1]$ has the form of $H - \sum_{i=2}^{l} \epsilon_i E_i, \epsilon_i \in \{0, 1\}, [C_i]$ with 1 < i < l has the form of $E_{\alpha_i} - \sum_{\beta > \alpha_i} \epsilon_\beta E_\beta, \epsilon_\beta \in \{0, 1\}$. Define $a_i = \#\{E_j \mid E_j \cdot [C_i] = 1, E_j \cdot [C_k] = 0$ for other $k\}$. It follows that $\Phi(p_1 - a_1, \dots, p_l - a_l, a_1, \dots, a_l) = (1, 1 - p_1, -p_2, \dots, -p_{l-1}, 1 - p_l)$. Therefore each homological configuration will correspond to some element in the preimage of Φ according to the number of non-toric blow up times at each position. Conversely, if there are two different elements in the preimage, then the homological configurations corresponding to them can not be only differed by a homological action of diffeomorphism. Since otherwise, the diffeomorphism maps H to H, which implies its homological action must be some permutations of E_i 's. It's easy to see the value a_i will not change after the permutations of E_i 's. As a result, the two divisors corresponding to those two homological configurations are not symplectic deformation equivalent by Theorem 2.2 but have the same self-intersection sequence, which means the labeled divisor is not def-taut.

For the completeness, we also determine the def-tautness for all the other self-intersection sequences with $b^+ = 1$ in Theorem 1.4 of [LMM20]. Those are (1, p), (-1, -p) with p = 1, 2, 3, (1, 1, p) with $p \leq 1, (0, p)$ with $p \leq 4$ and (1, p) with $p \leq 1$.

Proposition A.3. All the labeled symplectic log Calabi-Yau divisors with the above selfintersection sequences are def-taut. *Proof.* One could use the same argument as the proof of Proposition A.2 to show that if $A \in \mathcal{H}_l$ and $A \neq 3H - E_1 - \cdots - E_l$, then when $A^2 = 0$, it could be transformed to $H - E_1$ (or F when the manifold is $S^2 \times S^2$) by reflections and when $A^2 = -1$ it could be transformed to E_1 by reflections unless $A = H - E_1 - E_2$ with l = 2. Note that those reflections along $E_i - E_j, H - E_i - E_j - E_k$ preserve $c_1(M_l) = 3H - E_1 - \cdots - E_l$. So when the length of the sequence is 2, if we could transform one component by reflections, the other one will be transformed as well. This shows that labeled divisors of self-intersection sequences (1, p), (0, p)are def-taut, and labeled divisors of self-intersection sequences (-1, -p) are def-taut unless the manifold is M_2 , in which case p = -4. Therefore we have checked all the length 2 sequences above are def-taut. For the remaining case (1, 1, p), note that if we transform one of the self-intersection 1-component into H, the other self-intersection 1-component must also be H. This is simply because $A \cdot H = 1, A^2 = 1$ has the only solution A = H. This shows the homological configurations of the labeled divisors of self-intersection sequences (1, 1, p)can always be transformed into $(H, H, H - E_1 - \cdots - E_l)$ which implies labeled divisors of self-intersection sequences (1, 1, p) are def-taut.

Proposition A.4. The labeled symplectic log Calabi-Yau divisors with self-intersection sequence (s_1, \dots, s_k) are def-taut if and only if the labeled symplectic log Calabi-Yau divisors with the toric blow up self-intersection sequences (which are $(-1, s_1 - 1, \dots, s_k - 1), (s_1 - 1, -1, s_2 - 1, \dots, s_k), \dots, (s_1, \dots, s_{k-1} - 1, -1, s_k - 1), (s_1 - 1, \dots, s_k - 1, -1)$) are def-taut.

Proof. We only need to consider $(-1, s_1 - 1, \dots, s_k - 1)$, the others are same. Assume the divisors of self-intersection sequences $(s_1, \dots, s_k), (-1, s_1 - 1, \dots, s_k - 1)$ are in M_{l-1}, M_l with reduced blowup forms. We adopt the canonical identification $H_2(M_l; \mathbb{Z}) = H_2(M_{l-1}; \mathbb{Z}) \oplus \mathbb{Z}E_l$ and always identify symplectic deformation equivalence and homological equivalence by Theorem 2.2. We will use the following convenient result, which is actually implied in Section 4.1.2 of [LW12] : let K_{l-1}, K_l be the standard canonical classes, if γ, γ' are integral isometries of $H_2(M_{l-1}; \mathbb{Z}), H_2(M_l; \mathbb{Z})$ respectively and $\gamma' = \gamma \oplus id_{E_l}$, then $\gamma \in D_{K_{l-1}}(M_{l-1})$ if and only if $\gamma' \in D_{K_l}(M_l)$.

On the one hand, if the labeled divisors of self-intersection sequence $(-1, s_1 - 1, \dots, s_k - 1)$ are def-taut and $D = (C_1, \dots, C_k), D' = (C'_1, \dots, C'_k)$ are two labeled divisors of self-intersection sequence (s_1, \dots, s_k) , then there must exist divisors on M_l with homological configurations $(E_l, [C_1] - E_l, \dots, [C_k] - E_l), (E_l, [C'_1] - E_l, \dots, [C'_k] - E_l)$ by Corollary 2.33. And there is some homological action $\gamma \in D_{K_l}(M_l)$ such that $\gamma(E_l, [C_1] - E_l, \dots, [C_k] - E_l) = (E_l, [C'_1] - E_l, \dots, [C'_k] - E_l)$. $\gamma(E_l) = E_l$ implies $\gamma([C_1], \dots, [C_k]) = ([C'_1], \dots, [C'_k])$ and $\gamma|_{H_2(M_{l-1};\mathbb{Z})}$ is an isomorphism of the lattice $H_2(M_{l-1};\mathbb{Z})$ and thus belongs to $D_{K_{l-1}}(M_{l-1})$ by the claim. This shows D, D' are symplectic deformation equivalent.

On the other hand, if the labeled divisors of self-intersection sequence (s_1, \dots, s_k) are def-taut and $D = (C_0, C_1, \dots, C_k), D' = (C'_0, C'_1, \dots, C'_k)$ are two labeled divisors of self-intersection sequence $(-1, s_1 - 1, \dots, s_k - 1)$. As the proof of the previous Proposition, unless l = 2 we may assume $[C_0] = [C'_0] = E_l$. It then follows that $([C_1] + E_l, \dots, [C_k] + E_l), ([C'_1] + E_l, \dots, [C'_k] + E_l)$ are homological configurations with classes in $H_2(M_{l-1}; \mathbb{Z})$. So there is some $\gamma \in D_{K_{l-1}}(M_{l-1})$ such that $\gamma([C_1] + E_l, \dots, [C_k] + E_l) = ([C'_1] + E_l, \dots, [C'_k] + E_l)$. Thus by the claim $\gamma' = \gamma \oplus id_{E_l} \in D_{K_l}(M_l)$ will satisfy $\gamma'([C_0], [C_1], \dots, [C_k]) = ([C'_0], [C'_1], \dots, [C'_k])$. This shows that D, D' are symplectic deformation equivalent.

Combining the above Propositions with Theorem 1.4 of [LMM20], we already determined

all the def-taut labeled divisors with $b^+ = 1$.

Remark A.5. There are some differences between the symplectic def-tautness and holomorphic tautness. From the definition it's easy to see the holomorphic tautness implies symplectic def-tautness, but the converse is not true. In [Fri16] the sequence (4,0) is considered to be not taut, but in our notion it is def-taut. This is because (4,0) is the (only) self-intersection sequence having embeddings into two different manifolds $S^2 \times S^2$ or $\mathbb{CP}^2 \# \overline{\mathbb{CP}}^2$ and the definition of tautness in the holomorphic category does not fix the ambient manifolds. Other than this, the self-intersection sequence (2,2) is not holomorphic taut since it has realizations in $S^2 \times S^2$ with two non-isomorphic complex structures (namely \mathbb{F}_0 and \mathbb{F}_2 , see [Fri16]). However it's not hard to check the def-tautness for divisors with self-intersection sequence (2,2).

Remark A.6. For the purpose of completely determining the diffeomorphism types of symplectic fillings of torus bundles, one should define the def-tautness for unlabeled divisors. However it's difficult to obtain the toric blow up invariance for unlabeled def-tautness analogous to Proposition A.4 due to the cyclic symmetry.

A.2 Details of counting toric divisors in M_3

Proof of Proposition 3.7: The proof is very similar to that of Proposition 3.5. We start by listing the possible self-intersection sequences to be

- (1) (-n, 0, n-2, -1, -2, -1),
- (2) (-n, 0, n-1, -2, -1, -2),
- (3) (-n, -1, -1, n-2, -1, -1),

$$(4) \ (-1, -1, -1, -1, -1, -1),$$

where $n \ge 2$. Also, to make the presentation easier, we write $H_i = H - E_i$, $H_{ij} = H - E_i - E_j$ and $H_{ijk} = H - E_i - E_j - E_k$, for distinct i, j, k.

We denote by $tF_n(3; 1, \delta_1, \delta_2, \delta_3)$ the number of toric log Calabi-Yau divisors in (1)-(3) and by $tF_1(3; 1, \delta_1, \delta_2, \delta_3)$ the number of toric log Calabi-Yau divisors in (4). We could make use of Proposition 3.4 to enumerate the homology classes with self-intersections listed above.

Let's first consider (1) – (3). If n = 2k + 1 with $k \ge 1$, then the possible homology types are

1.
$$(-(k-1)H + kE_1 - E_2 - E_3, H_1, kH - (k-1)E_1, H_{12}, E_2 - E_3, E_3),$$

 $(-kH + (k+1)E_1, H_1, (k+1)H - kE_1 - E_2 - E_3, E_3, E_2 - E_3, H_{12}),$

2.
$$(-kH + (k+1)E_1, H_1, (k+1)H - kE_1 - E_2, E_2 - E_3, E_3, H_{123}),$$

3.
$$(-(k-1)H + kE_1 - E_2 - E_3, E_2, H_{12}, kH - (k-1)E_1, H_{13}, E_3), (-kH + (k+1)E_1, H_{12}, E_2, (k+1)H - kE_1 - E_2 - E_3, E_3, H_{13}).$$

Then we have

$$tF_{2k+1}(3;1,\delta_1,\delta_2,\delta_3) = \begin{cases} 5 & \text{if } -k + (k+1)\delta_1 > 0, \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 2 & \text{if } -k + (k+1)\delta_1 > 0, \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } -k + (k+1)\delta_1 > 0, \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 1 & \text{if } -k + (k+1)\delta_1 > 0, \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } \delta_1 + \delta_2 + \delta_3 - 1 < -k + (k+1)\delta_1 \leqslant 0, \delta_2 > \delta_3\\ 1 & \text{if } \delta_1 + \delta_2 + \delta_3 - 1 < -k + (k+1)\delta_1 \leqslant 0, \delta_2 = \delta_3\\ 0 & \text{if } -k + (k+1)\delta_1 \leqslant \delta_1 + \delta_2 + \delta_3 - 1 \end{cases}$$

If n = 2k + 2 with $k \ge 1$, then the possible homology types are

1.
$$(-kH + (k+1)E_1 - E_i, H_1, (k+1)H - kE_1 - E_j, E_j, H_{123}, E_i)$$
, with $\{i, j\} = \{2, 3\}$
2. $(-kH + (k+1)E_1 - E_2, H_1, (k+1)H - kE_1, H_{123}, E_3, E_2 - E_3)$
3. $(-kH + (k+1)E_1 - E_i, E_i, H_{1i}, (k+1)H - kE_1 - E_j, E_j, H_{1j})$, with $\{i, j\} = \{2, 3\}$

Then we have

$$tF_{2k+2}(3;1,\delta_1,\delta_2,\delta_3) = \begin{cases} 5 & \text{if } -k + (k+1)\delta_1 - \delta_2 > 0, \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 2 & \text{if } -k + (k+1)\delta_1 - \delta_2 > 0, \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } -k + (k+1)\delta_1 - \delta_2 > 0, \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 1 & \text{if } -k + (k+1)\delta_1 - \delta_2 > 0, \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } \delta_3 - \delta_2 < -k + (k+1)\delta_1 - \delta_2 \leqslant 0, \delta_1 + \delta_2 + \delta_3 < 1\\ 1 & \text{if } \delta_3 - \delta_2 < -k + (k+1)\delta_1 - \delta_2 \leqslant 0, \delta_1 + \delta_2 + \delta_3 = 1\\ 0 & \text{if } -k + (k+1)\delta_1 - \delta_2 \leqslant \delta_3 - \delta_2 \end{cases}$$

If n = 2, the possible homology types are

- 1. $(E_i E_j, H_i, H_k, E_k, H_{123}, E_j), \{i, j, k\} = \{1, 2, 3\}, i < j$ 2. $(E_1 - E_2, H_1, H, H_{123}, E_3, E_2 - E_3)$
- 3. $(E_i E_j, E_j, H_{ij}, H_k, E_k, H_{ik}), \{i, j, k\} = \{1, 2, 3\}, i < j$

Then we have

$$tF_2(3;1,\delta_1,\delta_2,\delta_3) = \begin{cases} 7 & \text{if } \delta_1 > \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 3 & \text{if } \delta_1 > \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } \delta_1 > \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 1 & \text{if } \delta_1 > \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 2 & \text{if } \delta_1 = \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 < 1\\ 1 & \text{if } \delta_1 = \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 = 1\\ 0 & \text{if } \delta_1 = \delta_2 = \delta_3 \end{cases}$$

If n = 1, there is only one possible homology type $(H_{12}, E_1, H_{13}, E_3, H_{23}, E_2)$ and is always realized. So we have

$$tF_1(3; 1, \delta_1, \delta_2, \delta_3) = 1$$

Now the results follow from

$$|t\mathcal{LCY}(3;1,\delta_1,\delta_2,\delta_3)| = \sum_{n=1}^{\infty} tF_n(3;1,\delta_1,\delta_2,\delta_3),$$

and analyzing the bounds on δ_1 , δ_2 , δ_3 in each region. For example, in region $Q_{i-1}R_{i-1}P_{2i-1}P_{2i-2}$, one just have to check the followings:

1. $-k + (k+1)\delta_1 > 0$ if and only if k < i2. $\delta_1 + \delta_2 + \delta_3 - 1 < -k + (k+1)\delta_1 \leq 0$ will never happen for any k3. $-k + (k+1)\delta_1 - \delta_2 > 0$ if and only if k < i4. $\delta_3 - \delta_2 < -k + (k+1)\delta_1 - \delta_2 \leq 0$ will never happen for any k5. $\delta_1 > \delta_2 > \delta_3, \delta_1 + \delta_2 + \delta_3 < 1$

In the region $Q_{i-1}Q_iP_{2i-1}$, one can check that:

1. $-k + (k+1)\delta_1 > 0$ if and only if k < i2. $\delta_1 + \delta_2 + \delta_3 - 1 < -k + (k+1)\delta_1 \leq 0$ if and only if k = i3. $-k + (k+1)\delta_1 - \delta_2 > 0$ if and only if k < i4. $\delta_3 - \delta_2 < -k + (k+1)\delta_1 - \delta_2 \leq 0$ will never happen for any k5. $\delta_1 > \delta_2 = \delta_3, \delta_1 + \delta_2 + \delta_3 < 1$

Therefore, the count in the region $Q_{i-1}R_{i-1}P_{2i-1}P_{2i-2}$ is given by

$$|t\mathcal{LCY}(3;1,\delta_1,\delta_2,\delta_3)| = \sum_{n=1}^{\infty} tF_n(3;1,\delta_1,\delta_2,\delta_3) = 1 + 7 + \sum_{k=1}^{i-1} 5 + \sum_{k=1}^{i-1} 5 = 10i - 2$$

And the count in the region $Q_{i-1}Q_iP_{2i-1}$ is given by

$$|t\mathcal{LCY}(3;1,\delta_1,\delta_2,\delta_3)| = \sum_{n=1}^{\infty} tF_n(3;1,\delta_1,\delta_2,\delta_3) = 1 + 2 + \sum_{k=1}^{i-1} 2 + 1 + \sum_{k=1}^{i-1} 2 = 4i$$

These two examples just correspond to the case (1) and (6) in our statement.

A.3 Details of computing toric regions for M_5

Proof of Proposition 3.11: Firstly note that if the homological sequence for M_5 in Fact 3.9 can not be realized as a toric symplectic log Calabi-Yau divisor, then one of $\omega(H_{134}), \omega(E_1 - E_4), \omega(E_2 - E_5), \omega(H_{235})$ must be zero. Thus we only need to focus on the regions MAD, MOD, MOA where the toric divisors might not exist.

Secondly, we consider the following toric homological sequence:

$$(E_2, E_1 - E_2 - E_3, E_3, H_{135}, E_5, H_{45}, E_4, H_{124})$$

In the interior of region XOA, we see that ω has positive pairing with all the homology classes in the above sequence. Again by Proposition 2.24, the homological sequence can be realized as a toric symplectic Calabi-Yau divisor. So the interior of region XOA is also contained in the toric cone.

Next we explain the nonexistence in regions MOD, MAD and MOX. In the M_5 case we have the following more complicated classification for self-intersection sequences:

$$\begin{split} A_n^1 &= (-1, -2, -2, -1, n, -1, -1, -n - 4), A_n^2 &= (-1, -3, -1, -2, n, -1, -1, -n - 3) \\ A_n^3 &= (-3, -1, -2, -2, n, -1, -1, -n - 2), B_n^1 &= (-1, -2, -1, n, -1, -2, -1, -n - 4) \\ B_n^2 &= (-2, -1, -2, n, -1, -2, -1, -n - 3), B_n^3 &= (-2, -1, -2, n, -2, -1, -2, -n - 2) \\ C_n^1 &= (-4, -1, -2, -2, -2, n, 0, -n - 1), C_n^2 &= (-3, -2, -1, -3, -2, n, 0, -n - 1) \\ C_n^3 &= (-3, -1, -3, -1, -3, n, 0, -n - 1), D_n^1 &= (-2, -1, -4, -1, -2, n, 0, -n - 2) \\ D_n^2 &= (-1, -4, -1, -2, -2, n, 0, -n - 2), D_n^3 &= (-3, -1, -2, -3, -1, n, 0, -n - 2) \\ E_n^1 &= (-2, -1, -3, -2, -1, n, 0, -n - 3), E_n^2 &= (-1, -3, -1, -3, -1, n, 0, -n - 3) \\ F_n^1 &= (-1, -2, -2, -2, -1, n, 0, -n - 4) \end{split}$$

Similarly as in the case for M_4 , we can apply the following information about the negative symplectic sphere classes by Proposition 3.4 of [LL20].

In region MOD:

$$\mathcal{S}_{\omega}^{-2} \subset \{E_p - E_5, H_{ijk} | 1 \le p \le 4, 1 \le i < j < k \le 5\}$$

$$\mathcal{S}_{\omega}^{-3} \subset \{H_{ijkl} | 1 \le i < j < k < l \le 5\}, \mathcal{S}_{\omega}^{-4} \subset \{H_{12345}\}, \mathcal{S}_{\omega}^{\le -5} = \emptyset$$

In region MAD:

$$\begin{split} \mathcal{S}_{\omega}^{-2} &\subset \{E_1 - E_p, E_q - E_5, H_{234}, H_{235}, H_{345}, H_{125}, H_{135}, H_{145}, H_{245} | 2 \leqslant p \leqslant 5, 1 \leqslant q \leqslant 4\} \\ \mathcal{S}_{\omega}^{-3} &\subset \{-H + 2E_1, H_{2345}, E_1 - E_p - E_q | 2 \leqslant p < q \leqslant 5\} \\ \mathcal{S}_{\omega}^{-4} &\subset \{E_1 - E_i - E_j - E_k, -H + 2E_1 - E_p | 2 \leqslant i < j < k \leqslant 5, 2 \leqslant p \leqslant 5\} \\ \text{For any } k \geq 1: \\ \mathcal{S}_{\omega}^{-2k-3} &\subset \{-(k-1)H + kE_1 - E_2 - E_3 - E_4 - E_5, -kH + (k+1)E_1 - E_2 - E_i, -(k+1)H + (k+2)E_1, -kH + (k+1)E_1 - E_p - E_q | 3 \leqslant i, p, q \leqslant 5, p \neq q\} \\ \mathcal{S}_{\omega}^{-2k-4} &\subset \{-(k+1)H + (k+2)E_1 - E_2, -kH + (k+1)E_1 - E_2 - E_p - E_q, -kH + (k+1)E_1 - E_3 - E_4 - E_5, -(k+1)H + (k+2)E_1 - E_i | 3 \leqslant i, p, q \leqslant 5, p \neq q\} \end{split}$$

In region MOX:

$$\begin{split} \mathcal{S}_{\omega}^{-2} &\subset \{E_1 - E_p, H_{ijk} | 2 \leqslant p \leqslant 5, 1 \leqslant i < j < k \leqslant 5\} \\ \mathcal{S}_{\omega}^{-3} &\subset \{-H + 2E_1, H_{ijkl}, E_1 - E_p - E_q | 1 \leqslant i < j < k < l \leqslant 5, 2 \leqslant p < q \leqslant 5\} \\ \mathcal{S}_{\omega}^{-4} &\subset \{H_{12345}, E_1 - E_i - E_j - E_k, -H + 2E_1 - E_p | 2 \leqslant i < j < k \leqslant 5, 2 \leqslant p \leqslant 5\} \\ \mathcal{S}_{\omega}^{\leqslant -5} &= \emptyset \end{split}$$

Finally by checking all the possible self-intersection sequences, one can carefully exclude all the possibilities by obeying the rules that any two adjacent homology classes have intersection number 1 and any two nonadjacent ones have intersection number 0. \Box

References

- [ALLP19] Silvia Anjos, Jun Li, Tian-Jun Li, and Martin Pinsonnault, *Stability of the symplectomorphism group of rational surfaces*, arXiv preprint arXiv:1911.00961 (2019).
- [CGHMP21] Dan Cristofaro-Gardiner, Tara S. Holm, Alessia Mandini, and Ana Rita Pires, On infinite staircases in toric symplectic four-manifolds, Preprint arXiv:2004.13062 (2021).
- [Che20] Weimin Chen, Finite group actions on symplectic Calabi-Yau 4-manifolds with $b_1 > 0$, J. Gökova Geom. Topol. GGT 14 (2020), 1–54. MR 4206325
- [CV20] Roger Casals and Renato Vianna, *Sharp ellipsoid embeddings and toric mutations*, Preprint arXiv:2004.13232 (2020).
- [Del88] Thomas Delzant, Hamiltoniens périodiques et images convexes de l'application moment, Bull. Soc. Math. France 116 (1988), no. 3, 315–339. MR 984900
- [DLW18] Josef G. Dorfmeister, Tian-Jun Li, and Weiwei Wu, Stability and existence of surfaces in symplectic 4-manifolds with $b^+ = 1$, J. Reine Angew. Math. **742** (2018), 115–155. MR 3849624
- [DRGI16] Georgios Dimitroglou Rizell, Elizabeth Goodman, and Alexander Ivrii, Lagrangian isotopy of tori in $S^2 \times S^2$ and $\mathbb{C}P^2$, Geom. Funct. Anal. **26** (2016), no. 5, 1297–1358. MR 3568033
- [Eng18] Philip Engel, Looijenga's conjecture via integral-affine geometry, J. Differential Geom. 109 (2018), no. 3, 467–495. MR 3825608
- [Eva21] Jonathan David Evans, *Lectures on lagrangian torus fibrations*, Preprint arXiv:2110.08643 (2021).
- [FM88] Robert Friedman and John W. Morgan, On the diffeomorphism types of certain algebraic surfaces. I, J. Differential Geom. 27 (1988), no. 2, 297–369. MR 925124
- [Fri16] Robert Friedman, On the geometry of anticanonical pairs, Preprint arXiv:1502.02560 (2016).

- [GHK15a] Mark Gross, Paul Hacking, and Sean Keel, Mirror symmetry for log Calabi-Yau surfaces I, Publ. Math. Inst. Hautes Études Sci. 122 (2015), 65–168. MR 3415066
- [GHK15b] _____, Moduli of surfaces with an anti-canonical cycle, Compos. Math. 151 (2015), no. 2, 265–291. MR 3314827
- [GL16] Marco Golla and Paolo Lisca, On Stein fillings of contact torus bundles, Bull.
 Lond. Math. Soc. 48 (2016), no. 1, 19–37. MR 3455745
- [Gom95] Robert E. Gompf, A new construction of symplectic manifolds, Ann. of Math.
 (2) 142 (1995), no. 3, 527–595. MR 1356781
- [GS09] David T. Gay and András I. Stipsicz, Symplectic surgeries and normal surface singularities, Algebr. Geom. Topol. 9 (2009), no. 4, 2203–2223. MR 2558309
- [HK20] Paul Hacking and Ailsa Keating, *Homological mirror symmetry for log calabi*yau surfaces, Preprint arXiv:2005.05010 (2020).
- [Kea18] Ailsa Keating, Homological mirror symmetry for hypersurface cusp singularities, Selecta Math. (N.S.) **24** (2018), no. 2, 1411–1452. MR 3782425
- [KK17] Yael Karshon and Liat Kessler, *Distinguishing symplectic blowups of the complex projective plane*, Journal of Symplectic Geometry **15** (2017), no. 4, 1089–1128.
- [KKP] Yael Karshon, Liat Kessler, and Martin Pinsonnault, Symplectic blowups of the complex projective plane and counting toric actions.
- [KKP07] _____, A compact symplectic four-manifold admits only finitely many inequivalent toric actions, J. Symplectic Geom. 5 (2007), no. 2, 139–166. MR 2377250
- [KKP15] _____, Counting toric actions on symplectic four-manifolds, C. R. Math. Acad. Sci. Soc. R. Can. **37** (2015), no. 1, 33–40. MR 3364303
- [Li08] Tian-Jun Li, *The space of symplectic structures on closed 4-manifolds*, arXiv: 0805.2931 (2008).
- [Lis08] Paolo Lisca, On symplectic fillings of lens spaces, Trans. Amer. Math. Soc. 360 (2008), no. 2, 765–799. MR 2346471
- [LL95] T. J. Li and A. Liu, Symplectic structure on ruled surfaces and a generalized adjunction formula, Math. Res. Lett. 2 (1995), no. 4, 453–471. MR 1355707
- [LL01] Tian-Jun Li and Ai-Ko Liu, Uniqueness of symplectic canonical class, surface cone and symplectic cone of 4-manifolds with $B^+ = 1$, J. Differential Geom. 58 (2001), no. 2, 331–370. MR 1913946
- [LL02] Bang-He Li and Tian-Jun Li, Symplectic genus, minimal genus and diffeomorphisms, Asian J. Math. 6 (2002), no. 1, 123–144. MR 1902650

- [LL20] Jun Li and Tian-Jun Li, Symplectic (-2)-spheres and the symplectomorphism group of small rational 4-manifolds, Pacific J. Math. 304 (2020), no. 2, 561–606. MR 4062781
- [LLW19] Jun Li, Tian-Jun Li, and Weiwei Wu, Symplectic (-2)-spheres and the symplectomorphism group of small rational 4-manifolds, ii, arXiv preprint arXiv:1911.11073 (2019).
- [LM16] Tian-Jun Li and Cheuk Yu Mak, Symplectic log Calabi-Yau surface deformation class, Adv. Theor. Math. Phys. 20 (2016), no. 2, 351–379. MR 3541847
- [LM18] _____, Geometry of symplectic log Calabi-Yau pairs, ICCM Not. 6 (2018), no. 2, 42–50. MR 3961489
- [LM19] _____, Symplectic divisorial capping in dimension 4, J. Symplectic Geom. 17 (2019), no. 6, 1835–1852. MR 4057728
- [LM21] Tian-Jun Li and Jie Min, Local geometry of symplectic divisors with applications to contact torus bundles, Preprint arXiv:2101.05981 (2021).
- [LMM20] Tian-Jun Li, Cheuk Yu Mak, and Jie Min, *Circular spherical divisors and their contact topology*, arXiv:2002.10504 (2020).
- [Loo81] Eduard Looijenga, Rational surfaces with an anticanonical cycle, Ann. of Math.
 (2) 114 (1981), no. 2, 267–322. MR 632841
- [LS10] Naichung Conan Leung and Margaret Symington, Almost toric symplectic fourmanifolds, J. Symplectic Geom. 8 (2010), no. 2, 143–187. MR 2670163
- [LW12] Tian-Jun Li and Weiwei Wu, Lagrangian spheres, symplectic surfaces and the symplectic mapping class group, Geom. Topol. 16 (2012), no. 2, 1121–1169. MR 2946805
- [McD97] Dusa McDuff, Lectures on Gromov invariants for symplectic 4-manifolds, Gauge theory and symplectic geometry (Montreal, PQ, 1995), NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., vol. 488, Kluwer Acad. Publ., Dordrecht, 1997, With notes by Wladyslav Lorek, pp. 175–210. MR 1461573
- [MO15] Dusa McDuff and Emmanuel Opshtein, Nongeneric J-holomorphic curves and singular inflation, Algebr. Geom. Topol. **15** (2015), no. 1, 231–286. MR 3325737
- [MS17] Dusa McDuff and Dietmar Salamon, *Introduction to symplectic topology*, third ed., Oxford Graduate Texts in Mathematics, Oxford University Press, Oxford, 2017. MR 3674984
- [Pas14] James Pascaleff, Floer cohomology in the mirror of the projective plane and a binodal cubic curve, Duke Math. J. 163 (2014), no. 13, 2427–2516. MR 3265556
- [Pin08] Martin Pinsonnault, Maximal compact tori in the Hamiltonian group of 4dimensional symplectic manifolds, J. Mod. Dyn. 2 (2008), no. 3, 431–455. MR 2417479

 [Sym03] Margaret Symington, Four dimensions from two in symplectic topology, Topo ogy and geometry of manifolds (Athens, GA, 2001), Proc. Sympos. Pure Math vol. 71, Amer. Math. Soc., Providence, RI, 2003, pp. 153–208. MR 2024634 [Tau95] Clifford Henry Taubes, The Seiberg-Witten and Gromov invariants, Math. ReLett. 2 (1995), no. 2, 221–238. MR 1324704 [Via14] Renato Vianna, On exotic Lagrangian tori in CP², Geom. Topol. 18 (2014 no. 4, 2419–2476. MR 3268780 [Via16] Renato Ferreira de Velloso Vianna, Infinitely many exotic monotone Lagrangia tori in CP², J. Topol. 9 (2016), no. 2, 535–551. MR 3509972 [Wal64] C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. 39 (1964 131–140. MR 163323 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[ST05]	Bernd Siebert and Gang Tian, On the holomorphicity of genus two Lefschetz fibrations, Ann. of Math. (2) 161 (2005), no. 2, 959–1020. MR 2153404
 [Tau95] Clifford Henry Taubes, The Seiberg-Witten and Gromov invariants, Math. Red Lett. 2 (1995), no. 2, 221–238. MR 1324704 [Via14] Renato Vianna, On exotic Lagrangian tori in CP², Geom. Topol. 18 (2014 no. 4, 2419–2476. MR 3268780 [Via16] Renato Ferreira de Velloso Vianna, Infinitely many exotic monotone Lagrangia tori in CP², J. Topol. 9 (2016), no. 2, 535–551. MR 3509972 [Wal64] C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. 39 (1964 131–140. MR 163323 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math. Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[Sym03]	Margaret Symington, <i>Four dimensions from two in symplectic topology</i> , Topology and geometry of manifolds (Athens, GA, 2001), Proc. Sympos. Pure Math., vol. 71, Amer. Math. Soc., Providence, RI, 2003, pp. 153–208. MR 2024634
 [Via14] Renato Vianna, On exotic Lagrangian tori in CP², Geom. Topol. 18 (2014 no. 4, 2419–2476. MR 3268780 [Via16] Renato Ferreira de Velloso Vianna, Infinitely many exotic monotone Lagrangia tori in CP², J. Topol. 9 (2016), no. 2, 535–551. MR 3509972 [Wal64] C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. 39 (1964 131–140. MR 163323 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[Tau95]	Clifford Henry Taubes, <i>The Seiberg-Witten and Gromov invariants</i> , Math. Res. Lett. 2 (1995), no. 2, 221–238. MR 1324704
 [Via16] Renato Ferreira de Velloso Vianna, Infinitely many exotic monotone Lagrangia tori in CP², J. Topol. 9 (2016), no. 2, 535–551. MR 3509972 [Wal64] C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. 39 (1964 131–140. MR 163323 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[Via14]	Renato Vianna, On exotic Lagrangian tori in $\mathbb{CP}^2,$ Geom. Topol. 18 (2014), no. 4, 2419–2476. MR 3268780
 [Wal64] C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. 39 (1964 131–140. MR 163323 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math. Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[Via16]	Renato Ferreira de Velloso Vianna, Infinitely many exotic monotone Lagrangian tori in \mathbb{CP}^2 , J. Topol. 9 (2016), no. 2, 535–551. MR 3509972
 [Zha17] Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851 	[Wal64]	C. T. C. Wall, Diffeomorphisms of 4-manifolds, J. London Math. Soc. ${\bf 39}$ (1964), 131–140. MR 163323
	[Zha17]	Weiyi Zhang, The curve cone of almost complex 4-manifolds, Proc. Lond. Math. Soc. (3) 115 (2017), no. 6, 1227–1275. MR 3741851

School of Mathematics, University of Minnesota, Minneapolis, MN, US E-mail address: tjli@math.umn.edu

Max Planck Institute for Mathematics, Bonn, Germany E-mail address: minxx1270umn.edu

School of Mathematics, University of Minnesota, Minneapolis, MN, US E-mail address: ning00400umn.edu