MODULI OF UNSTABLE BUNDLES OF HN-LENGTH TWO WITH FIXED ALGEBRA OF ENDOMORPHISMS

L. BRAMBILA-PAZ AND ROCÍO RÍOS SIERRA

ABSTRACT. Let X be a smooth irreducible complex projective curve of genus $g \geq 2$ and $U_{\mu_1}(n, d)$ the moduli scheme of indecomposable vector bundles over X with fixed Harder-Narasimhan type $\sigma = (\mu_1, \mu_2)$. In this paper, we give necessary and sufficient conditions for a vector bundle $E \in U_{\mu_1}(n, d)$ to have $\mathbb{C}[x_1, \ldots, x_k]/(x_1, \ldots, x_k)^2$ as its algebra of endomorphisms. Fixing the dimension of the algebra of endomorphisms we obtain a stratification of $U_{\mu_1}(n, d)$ such that each stratum $U_{\mu_1}(n, d, k)$ is an algebraic variety, moreover, a coarse moduli space. A particular case of interest is when the unstable bundles are simple. In that case the moduli space is fine. Topological properties of $U_{\mu_1}(n, d, k)$ will depend on the generality of the curve X. Such results differ from the corresponding results for the moduli space of stable bundles, where non-emptiness, dimension etc. are independent of the curve.

1. INTRODUCTION

Let X be a complex smooth projective variety. It is well known that the moduli space of semistable bundles with fixed invariants exists, and that sometimes is fine, i.e. there exists a universal family. Some moduli spaces of unstable bundles have been constructed by adding some extra information: the moduli spaces of unstable vector bundles of rank 2 and 3 were constructed in [14] and in [39], respectively, when dim X = 1 and the algebra of endomorphisms is fixed; in [41] when X is the projective plane and in [3] when $X = \mathbb{P}^3(\mathbb{C})$, in both cases they consider the degree of instability. J. M. Drezet studied in [19] the case of very unstable bundles when dim X > 2. In [27] the authors construct the moduli spaces of pure sheaves with fixed Harder-Narasimhan type which have some additional data called a *m*-rigidification. The moduli spaces of unstable sheaves via non-reductive GIT has been studied by V. Hoskins, G. Berczi, J. Jackson and F. Kirwan in [5].

Unless otherwise stated we assume now that X is a smooth irreducible complex projective curve of genus $g \ge 2$. The aim of this paper is to study unstable bundles over X using their algebra of endomorphisms. The advantage of using algebra of endomorphisms lies in the fact that it allow us to construct coarse moduli spaces, and even fine moduli spaces.

Date: April 26, 2022.

²⁰¹⁰ Mathematics Subject Classification. 14H60, 14J60.

Both authors acknowledges the support of CONACYT, in particular of CONACYT grant 251938.

In order to state our results let us recall that any vector bundle over X has a unique filtration (called Harder-Narasimhan filtration)

$$0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$$

such that for $0 \le i \le m - 1$, E_{i+1}/E_i is semistable and

(1.1)
$$\mu(E_1) > \mu(E_2/E_1) > \dots > \mu(E_m/E_{m-1}).$$

The sequence of slopes $\sigma = (\mu(E_1), \mu(E_2/E_1), \cdots, \mu(E_m/E_{m-1}))$ is called the *Harder-Narasimhan type (HN-type for short)* of *E*. The moduli space of vector bundles of HN-type $\sigma = (\mu(E_1))$ is the moduli space of semistable bundles. It was constructed by Mumford [36] in 1960's using Geometric Invariant Theory and by M.S. Narasimhan and S. Seshadri [38] using representation theory. For a treatment of a more general case we refer the reader to [40], [20], [32] and [33]. The moduli space of some non-simple semistable vector bundles with a fixed algebra of endomorphisms was constructed in [10], [11] and [12].

Denote by $U_{\mu_1}(n, d)$ the set of indecomposable vector bundles of rank n and degree d of coprime-type $\sigma = (\mu_1, \mu_2)$ (see Definition 2.1). If $\mu_1 - \mu_2 > 2g - 1$, $U_{\mu_1}(n, d) = \emptyset$ (see Proposition 2.6). For $0 < \mu_1 - \mu_2 \le 2g - 1$, $U_{\mu_1}(n, d)$ has a projective scheme structure that makes it an moduli scheme (see Theorem 4.6).¹

Our purpose is to use one more numerical invariant to describe a stratification of $U_{\mu_1}(n,d)$ such that each stratum is an algebraic variety, and a coarse moduli space. Such invariant is the dimension, as a \mathbb{C} -vector space, of the algebra of global endomorphisms. First we determine the structure of the algebra of endomorphism of vector bundles in $U_{\mu_1}(n,d)$. If $\mu_1 = \frac{d_1}{n_1}$, write $d_2 = d - d_1$ and $n_2 = d - d_1$. We prove that (see Corollary 3.2)

if
$$E \in U_{\mu_1}(n,d)$$
 then $End(E) = \mathbb{C}[x_1, \dots, x_k]/(x_1, \dots, x_k)^2$

where $0 \le k \le \frac{d_1 n_2 - d_2 n_1}{2} + n_1 n_2$.

Set $A_k := \mathbb{C}[x_1, \ldots, x_k]/(x_1, \ldots, x_k)^2$ and $A_0 := \mathbb{C}$. For any integer $0 \le k \le \frac{d_1n_2-d_2n_1}{2} + n_1n_2$ we will denote by $U_{\mu_1}(n, d, k)$ the set

$$U_{\mu_1}(n, d, k) := \{ E \in U_{\mu_1}(n, d) : End(E) \cong A_k \}.$$

Fixing dim End(-) as invariant we prove that a flattening stratification of an Ext^1 sheaf over a convenient variety Y (see Theorems 4.1) gives the existence of a schematic stratification of the variety Y with a universal property. We use such sub schemes and the ideas of twisted Brill-Noether theory (see [24]) to give an algebraic structure to $U_{\mu_1}(n, d, k)$ and prove the following theorems. By $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ we mean the twisted Brill-Noether locus of product of two stable bundles with at least k section and by $\mathcal{Y}_k \subset B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ those with exactly k sections (see (4.2) and Section 4 for the definition of the twisted Brill-Noether locus $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ and of the number h^1).

¹At the time of writing this article, some results on vector bundles of type $\sigma = (\mu_1, \mu_2)$ were obtained independently, and by different methods, in [25] and [28]

Theorem 1.1. (Theorems 4.6 and Corollary 4.7) $\mathcal{U}_{\mu_1}(n, d, k)$ is a coarse moduli space and $\mathcal{U}_{\mu_1}(n, d, 0)$ is a fine moduli space. Moreover, if \mathcal{Y}_k is irreducible and smooth of dimension ρ , then $U_{\mu_1}(n, d, k)$ is irreducible and smooth of dimension $\rho + h^1 - 1$. In particular, if $U_{\mu_1}(n, d, k)$ is non-empty, $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ is non-empty.

Corollary 1.2. (Corollary 4.8) If \mathcal{Y}_k is irreducible and smooth then $H^i(\mathcal{U}_{\mu_1}(n, d, k), \mathbb{C}) \cong H^i(\mathcal{Y}_k, \mathbb{C})$ for $i \ge 0$.

Non-emptiness and topological properties of $\mathcal{U}_{\mu_1}(n, d, k)$ are given in the following theorems. Of particular interest are the vector bundles of HN-type $\sigma = (\frac{d-a}{n-1}, a)$, where a is an integer. That is, indecomposable unstable bundles that are extensions of a line bundle by a semistable bundle. In this case the results are a reformulation of the known results of the Brill-Noether theory in terms of the moduli of unstable bundles. The expected dimension $\beta(g, n_1, d_1, n_2, d_2, k)$ of $U_{\mu_1}(n, d, k)$ is given in Section 4. Recall that the *Brill-Noether loci* are defined as

$$B(n, d, k) := \{ G \in M(n, d) : h^0(G) \ge k \},\$$

where M(n, d) is the moduli space of stable vector bundles of degree d and rank n over X.

Theorem 1.3. (Theorem 5.1) Assume 0 < d - an < 2(n - 1) and $(n - 1, d - an, k) \neq (n - 1, n - 1, n - 1)$. Then for $\mu_1 = \frac{d-a}{n-1}$, $U_{\mu_1}(n, d, k)$ is non-empty if and only if $k \leq n - 1 + \frac{d-n(a+1)+1}{g}$. Moreover, if $U_{\mu_1}(n, d, k)$ is non-empty then it is irreducible and smooth of the expected dimension.

There are special results for general and Petri curves;

Theorem 1.4. (Theorem 5.4) Let (g, n - 1, d - na, k) be integers that satisfies the conditions given in Theorem 2.8,(5),(6) and (7). For general curve, $U_{\mu_1}(n, d, k)$ is non-empty and has an irreducible component of the expected dimension. Moreover, if X is a Petri curve of genus $g \ge 3, n \ge 5$ and $g \ge 2n - 4$ then $U_{\mu_1}(n, d, k)$ is non-empty.

As in the Brill-Noether theory for vector bundles, it is possible that for special curves the above conclusion not holds, and where the Brill-Noether locus, and hence the moduli space, is not even reduced. Thus, the above results differ from the corresponding results for the moduli space of stable bundles, where non-emptiness, dimension etc. are independent of the curve.

To our best knowledge the following theorems give also new results in the Brill-Noether and twisted Bril-Noether theory (see [7]).

Theorem 1.5. (Theorem 5.6) Assume that $B(n_1, d_1, n_1 + a)$ is non-empty with a > 0. If $2n_1 < d_1 < a(g+1)$ and $d_2 > 2gm$ then for any $0 \le k \le (d_2 + m(1-g))(n_1 + a) - (d_2n_1 + d_1m + mn_1(1-g))$, $\mathcal{Y}_k \subset B^k(U_1, U_2^*)$ is non-empty. Moreover, if $\mu_1 = \frac{d_1}{n_1}$ then $U_{\mu_1}(n, d, k)$ is non-empty, where $n = n_1 + n_2$ and $d = d_1 + d_2$.

For Perti curves we prove

Theorem 1.6. (Theorem 5.7) Let X be a Petri curve of genus $g \ge 3$ and $(\mathcal{O}(D), V)$ a general generated linear system of degree $d_2 \ge g + 1$ and dim $V = n_2 + 1$ with $n_2 \le 4$ or if $n_2 \ge 5$ then $g \ge 2n_2 - 4$. Assume that $B(n_1, d_1, t)$ is non-empty and $\frac{d_2}{n_2} < \frac{d_1}{n_1}$. For any $0 \le k \le n_2 t - n_1 d_2$, $\mathcal{Y}_k \subset B^k(U_1, U_2^*)$ is non-empty and if $n = n_2 + n_1$, $d = d_2 + d_1$ and $\mu_1 = \frac{d_1}{n_1}$, $U_{\mu_1}(n, d, k)$, is non-empty.

In the remainder of the last section we give necessary and sufficient conditions for $U_{\mu_1}(n, d, k)$ be smooth on $E \in U_{\mu_1}(n, d, k)$. Let $0 \subset E_1 \subset E$ be the HN-filtration of E in $U_{\mu_1}(n, d, k)$. Write $E/E_1 = F_1$. We use the following diagram

$$\begin{array}{ccc} H^{1}(End(E_{1})) \oplus H^{1}(End(F_{1})) & \xrightarrow{d\Phi} & H^{1}(End(E_{1} \otimes F_{1}^{*})) \\ & \eta_{E} \searrow & \beta \downarrow \\ & & H^{0}(E_{1} \otimes F_{1}^{*})^{*} \otimes H^{1}(E_{1} \otimes F_{1}^{*}) \end{array}$$

to define η_E as $\eta_E := \beta \circ d\Phi$ and prove

Theorem 1.7. (Theorem 5.9) $U_{\mu_1}(n, d, k)$ is smooth at E and of the expected dimension if and only if η_E is surjective.

In Section 2 we review some of the standard facts on unstable bundles. Section 3 will be concerned with the algebra of endomorphisms of unstable bundles. In Sections 4 and 5 our main results are stated and proved.

Acknowledgments: The first author gratefully acknowledges the many helpful suggestions of Peter E. Newstead during the preparation of the paper. Her thanks are also to Alfonso Zamora Saiz for drawing the author's attention to some unclear points. The first author is a member of the international research group VBAC (Vector Bundles on Algebraic Curves).

Notation The rank and degree of a vector bundle E are denoted by rk(E) and d(E) respectively and the slope as the rational number $\mu(E) := \frac{d(E)}{rk(E)}$. We will write the projection in the i-factor as p_i and by $\langle E \xrightarrow{f} F \rangle$ the linear space generated by the function $f: E \to F$. The cohomology groups $H^i(X, E)$ as $H^i(E)$ and its dimension as $h^i(E)$. Given an exact sequences

$$\rho: 0 \to G \to E \to F \to 0,$$

- by ρ^* we mean the dual sequence $\rho^*: 0 \to F^* \to E^* \to G^* \to 0$,
- by $(M \otimes \rho)$ the sequence ρ tensor by the vector bundle M i.e.

 $(M \otimes \rho) : 0 \to M \otimes G \to M \otimes E \to M \otimes F \to 0.$

• By $H^*(\rho)$ we mean the cohomology sequence of ρ

$$0 \to H^0(X,G) \to H^0(X,E) \to H^0(X,F) \xrightarrow{\delta} H^1(X,G) \to H^1(X,E) \to \dots$$

• To shorten notation, sometimes we write $E \in H^1(X, F^* \otimes G)$ instead of $(\rho : 0 \to G \to E \to F \to 0) \in H^1(X, F^* \otimes G)$

MODULI OF UNSTABLES

2. Unstable bundles of HN-lenght 2

From now on, X will be a smooth irreducible complex projective curve of genus $g \ge 2$. Recall that a vector bundle E over X is semistable if for all proper subbundle $F \subset E$ the slopes satisfy the following inequality

$$\mu(F) \le \mu(E).$$

The vector bundle E is stable if the inequality is strict and unstable if it is not semistable.

In [23] it was proved that any vector bundle over X has a unique filtration, called *Harder-Narasimhan filtration*,

$$(2.1) 0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$$

such that for $1 \leq i \leq m$,

• E_i/E_{i-1} is semistable and

(2.2)
$$\mu(E_1) > \mu(E_2/E_1) > \dots > \mu(E_m/E_{m-1}).$$

To shorten notation, we write HN-filtration instead of Harder-Narasimhan filtration and for abbreviation, we write F_i instead of the quotient E_{i+1}/E_i , when no confusion can arise. Note that $F_0 = E_1$. The HN-max and HN-min of E are defined as

 $\mu_{max}(E) := \mu(E_1)$ and $\mu_{min}(E) := \mu(F_{m-1}).$

The vector bundle E_1 is called the maximal destabilizing subbundle.

The following definitions will be used throughout all the paper.

Definition 2.1. Let $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ be the Harder-Narasimhan filtration of E.

- The HN-type is the sequence of slopes $\sigma = (\mu(E_1), \dots, \mu(E_m/E_{m-1})).$
- The number m is called the *HN*-length of the HN-filtration.
- The HN-filtration is called of *simple type* if each E_i and F_i are simple for $i = 1, \ldots, m-1$.
- The HN-filtration is called of *coprime type* if the numbers in each slope $\mu(F_i)$ and $\mu(E_i)$ are coprime for $i = 1, \ldots, m 1$.
- The HN-filtration is called *HN-indecomposable* if each E_i and F_i is indecomposable $i = 1, \ldots, m 1$.
- E is called HN-general (respectively HN-special) if all the quotient F_i are general (respectively special) in the Brill-Noether theory.
- The extension $\rho_i : 0 \to E_i \to E_{i+1} \to F_i \to 0$ is called the HN(i)-extension and the sequence of extensions $\rho = (\rho_1, \dots, \rho_{m-1})$ is called the *HN-sequences of E*.

Remark 2.2. Let $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ be the HN-filtration of E. Note that:

(1) the condition to being of simple/coprime type or HN-indecomposable is for $i = 1, \ldots, m-1$. Therefore, $E_m = E$ is not necessarily simple or indecomposable.

- (2) Simple type \implies HN-indecomposable type. However, HN-indecomposable \Rightarrow simple type.
- (3) If $i \neq 0$, E_i is unstable and $0 = E_0 \subset E_1 \subset \cdots \subset E_i$ is the HN-filtration of E_i . Moreover, if the filtration of E is of coprime, simple, HN-indecomposable, or general type, the same holds for the HN-filtration of any E_i . This will allow us to make induction on the HN-length.
- (4) For each E_i we have an exact sequence

(2.3)
$$\rho_i: 0 \to E_{i-1} \to E_i \to F_{i-1} \to 0.$$

Thus, E is constructed by a successive sequence of extensions.

In this section we will restrict our attention to the case of vector bundles of HN-type $\sigma = (\mu_1, \mu_2)$. Note that the value of μ_1 fix the value of μ_2 and viceversa. Indeed, if $\mu_1 = \frac{d_1}{n_1}$ then $\mu_2 = \frac{d(E)-d_1}{rk(E)-n_1}$. Our aim is to describe some properties and parameterize vector bundles of HN-type $\sigma = (\mu_1, \mu_2)$.

Let

(2.4)
$$(\rho_1: 0 \to E_1 \xrightarrow{i} E_2 \xrightarrow{p} F_1 \to 0) \in Ext^1(F_1, E_1) \cong H^1(X, F_1^* \otimes E_1)$$

be an extension of two semistable bundles with $\mu_1 := \mu(E_1) > \mu(F_1)$. It follows that E_2 is unstable and $0 \subset E_1 \subset E_2$ is its HN-filtration, with $E_2/E_1 = F_1$.

Remark 2.3. Note that given two extensions $(\rho_1 : 0 \to E_1 \xrightarrow{i} E_2 \xrightarrow{p} F_1 \to 0)$ and $(\rho'_1 : 0 \to E_1 \xrightarrow{i} E'_2 \xrightarrow{p} F_1 \to 0)$ in $Ext^1(F_1, E_1), E_2 \cong E'_2$ if and only if $\rho_1 \sim \rho'_1$ in $Ext^1(F_1, E_1)/(Aut(E_1) \times Aut(F_1))$. Moreover, if E_1 and F_1 are simple, $E_2 \cong E'_2$ iff $\rho = \lambda \rho'$ with $\lambda \in \mathbb{C}^*$.

To prove our first results we use the following lemmas.

Lemma 2.4. Let $\rho_1 : 0 \to E_1 \xrightarrow{i} E_2 \xrightarrow{p} F_1 \to 0$ be an extension of two semistable bundles with $\mu_1 := \mu(E_1) > \mu(F_1)$. If $\mu(E_1) - \mu(F_1) > 2g - 2$, $\rho_1 = 0$. Moreover, if ρ_1 is non-trivial then

- (1) $H^0(X, E_1^* \otimes E_1) = H^0(X, E_1^* \otimes E_2)$. Moreover, if E_1 is simple, $h^0(E_1^* \otimes E_2) = 1$ and $H^0(E_1^* \otimes E_2) = \langle E_1 \xrightarrow{i} E_2 \rangle$.
- (2) $H^0(F_1^* \otimes F_1) = H^0(E_2^* \otimes F_1)$. Moreover, if F_1 is simple, $h^0(E_2^* \otimes F_1) = 1$ and $H^0(E_2^* \otimes F_1) = \langle E_2 \xrightarrow{p} F_1 \rangle$.
- (3) If F_1 is simple then $H^0(F_1^* \otimes E_1) = H^0(F_1^* \otimes E_2)$.

Proof. The inequality $\mu(E_1) - \mu(F_1) > 2g - 2$ implies that $\mu(F_1^* \otimes E_1) > 2g - 2$. Hence, since $F_1^* \otimes E_1$ is semistable, $H^1(X, F_1^* \otimes E_1) = 0$.

Since $H^0(X, E_1^* \otimes F_1) = 0$, (1) and (2) follow from the cohomology sequences

(2.5)
$$H^*(E_1^* \otimes (\rho_1)) : 0 \to H^0(X, E_1^* \otimes E_1) \to H^0(X, E_1^* \otimes E_2) \to H^0(X, E_1^* \otimes F_1) \xrightarrow{\delta_0} \cdots$$

and

$$(2.6) \quad H^*((\rho_1)^* \otimes F_1) : 0 \to H^0(X, F_1^* \otimes F_1) \to H^0(X, E_2^* \otimes F_1) \to H^0(X, E_1^* \otimes F_1) \xrightarrow{\delta_1} \cdots$$

(3).- From the cohomology sequence $H^*(F_1^* \otimes (\rho_1))$

$$(2.7) \quad 0 \to H^0(X, F_1^* \otimes E_1) \to H^0(X, F_1^* \otimes E_2) \to H^0(X, F_1^* \otimes F_1) \xrightarrow{\phi_2} H^1(X, F_1^* \otimes E_1) \cdots$$

 $H^0(X, F_1^* \otimes E_1) = H^0(X, F_1^* \otimes E_2)$, since F_1 is simple and $\rho_1 \neq 0$.

Lemma 2.5. Assume E_1 and F_1 are semistable with E_1 simple and F_1 indecomposable. Then E_2 is indecomposable if and only if $\rho_1 \neq 0$.

Proof. One direction is trivial. Assume $\rho_1 \neq 0$. Suppose, contrary to our claim, that $E_2 = G_1 \oplus G_2$ and the extension

$$0 = \rho : 0 \to G_1 \xrightarrow{j} E_2 \xrightarrow{q} G_2 \to 0$$

splits. Denote by $\gamma: G_2 \to E_2$ the splitting morphism, i.e. $q \circ \gamma = id_{G_2}$. We can certainly assume that $\mu(G_2) \leq \mu(E_2)$, for if not, we replace G_2 by G_1 .

The inclusion $i: E_1 \to E_2$ in the extension ρ_1 induces the following diagram

We have different cases.

Case (1)-. $H \neq 0$ and $M \neq 0$.

In this case there is a contradiction to Lemma 2.4,(1) since the morphism $\gamma \circ \nu \circ \sigma$: $E_1 \rightarrow E_2$ is not a scalar multiple of the inclusion $i: E_1 \rightarrow E_2$.

Case (2)-. $H = E_1$.

The diagram

induces the following diagram

The semistability of F_1 implies that $\mu(G_1) \leq \mu(F_1) < \mu(E_2)$, hence that $\mu(E_2) < \mu(G_2)$, which is a contradiction to our assumption.

Case (3)-. H = 0.

In this case we have the following diagram

If $0 \neq \eta := p \circ \gamma : G_2 \to F_1$,

$$\ell \circ \eta = \ell \circ p \circ \gamma = q \circ \gamma = id_{G_2}$$

and hence F_1 is decomposable which is a contradiction, since F_1 is indecomposable.

If $\eta = 0$ then $G_2 \subset E_2 \subset G_1$, which is also a contradiction since $E_2 = G_1 \oplus G_2$.

Therefore, E_2 is indecomposable, as claimed.

Let $T = Ext^1(F_1, E_1) = H^1(X, F_1^* \otimes E_1)$. It is well known (see [37]) that $\mathbb{P}(T)$ parameterize non-trivial extensions $\rho_1 : 0 \to E_1 \xrightarrow{i} E \xrightarrow{p} F_1 \to 0$ and that if \mathbb{H} is the hyperplane bundle over $\mathbb{P}(T)$ the extension

$$0 \to p_1^*(E_1) \otimes p_2^*(\mathbb{H}) \to \mathcal{E} \to p_1^*(F_1) \to 0$$

that corresponds to the identity under the isomorphism

$$End(T) \cong H^1(X \times \mathbb{P}(T), p_1^*(Hom(F_1, E_1)) \otimes p_2^*(\mathbb{H}))$$

has universal properties. Thus, Lemma 2.5 leads us to the following result.

Proposition 2.6. If E_1 and F_1 are simple semistable vector bundles with $\mu(E_1) > \mu(F_1)$ then the pair $(\mathbb{P}(T), \mathcal{E})$ is a fine moduli space for indecomposable unstable bundles Ewith E_1 as the maximal destabilizing subbundle of E and $F_1 = E/E_1$. Moreover, if $\mu(E_1) - \mu(F_1) > 2g - 2$, $\mathbb{P}(T) = \emptyset$.

From now on we tacitly assume that $0 < \mu(E_1) - \mu(F_1) \leq 2g - 2$. By Riemann-Roch

(2.8)
$$\dim \mathbb{P}(T) = h^0(F_1^* \otimes E_1) - \tilde{d} + \tilde{r}(g-1) - 1,$$

where $\tilde{d} := deg(E_1)rk(F_1) - deg(F_1)rk(E_1)$ and $\tilde{r} := rk(E_1)rk(F_1)$. The dimension of $\mathbb{P}(T)$ depends on $h^0(F_1^* \otimes E_1)$. Actually, dim $\mathbb{P}(T) > 0$ if and only if $h^0(F_1^* \otimes E_1) > \tilde{d} + \tilde{r}(1-g) + 1$. Since $F_1^* \otimes E_1$ is semistable and $0 < \mu(E_1) - \mu(F_1) \le 2g - 2$, the study of the dimension $h^0(F_1^* \otimes E_1)$ belongs to the Brill-Noether Theory for vector bundles.

We will touch only a few aspects of the theory. Recall that a vector bundle G is special if $h^0(G) \cdot h^1(G) \neq 0$, otherwise is called general. The *Brill-Noether loci* are defined as

$$B(n, d, k) := \{ G \in M(n, d) : h^0(G) \ge k \},\$$

where M(n, d) is the moduli space of stable vector bundles of degree d and rank n over X. Moreover, the Brill-Noether loci induce the following filtration:

$$M(n,d) \supseteq B(n,d,1) \supseteq \cdots \supseteq B(n,d,k) \supseteq B(n,d,k+1) \supseteq \cdots$$

The number

$$\rho(g, n, d, k) := n^2(g-1) + 1 - k(k - d + n(g-1))$$

is called the Brill-Noether number and is the expected dimension of B(n, d, k). The Brill-Noether loci are also defined on the moduli space of S-equivalent semistable bundles (see [13]).

Remark 2.7. Denote by χ the number $\chi := d + n(1-g)$. Note that if $k \leq \chi$, B(n, d, k) = M(n, d). Thus, to have proper Brill-Noether loci we will assume that $k > \chi$.

For the convenience of the reader we summarize the main results of these concepts and the relevant material for us without proofs, thus making our exposition self-contained. See [22] for a survey of the main results on the Brill-Noether Theory. Recall that X is a *Petri curve* if the multiplication map $\mu : H^0(X, L) \otimes H^0(X, K_X \otimes L^*) \to H^0(X, K_X)$ is injective for every line bundle L on X (see [2]).

Theorem 2.8. Let G be a semistable vector bundle of rank n and degree d on X.

(1) If G is general,

$$h^{0}(G) = \begin{cases} 0 & \text{if } 0 < \mu(G) < g - 1, \\ d + r(1 - g) & \text{if } g - 1 \le \mu(G) < 2(g - 1). \end{cases}$$

- (2) If B(n, d, k) is proper, the general point E of every proper component of B(n, d, k)has $h^0(E) = k$.
- (3) (Clifford's Theorem for vector bundles [13, Theorem 2.1]) If G is special,

$$h^0(G) \le \frac{d}{2} + n.$$

(4) ([13] and [35]) For any curve X, if $0 < \mu(G) < 2$,

 $B(n, d, k) \neq \emptyset$ if and only if n < d + (n - k)g and $(n, d, k) \neq (n, n, n)$.

Moreover, if B(n, d, k) is not empty, then it is irreducible, of dimension $\rho(g, n, d, k)$ and Sing(B(n, d, k)) = B(n, d, k + 1).

(5) ([15]) For general curve B(n, d, k) is not empty if d = nd' + d'' and $n \le d'' + (n-k)g$ and the following conditions are satisfied

$$0 < d'' < 2n, \ d' \ge \frac{(s-1)(s+g)}{s}, \ with \ 1 < s < g, \ (d'',k) \ne (n,n).$$

(6) ([22]) If X is generic and $d = nd_1 + d_2$, $k = nk_1 + k_2$, $d_2 < n$, $k_2 < n$ for some positive integers d_i and k_i then $B(n, d, k) \neq \emptyset$ and has one component of the expected dimension if one of the following conditions is satisfied:

$$\begin{array}{ll} g - (k_1 + 1)(g - d_1 + k_1 - 1) \geq 1, & 0 = d_2 \geq k_2 \\ g - k_1(g - d_1 + k_1 + 1) > 1, & d_2 = k_2 = 0 \\ g - (k_1 + 1)(g - d_1 + k_1) \geq 1, & d_2 < k_2. \end{array}$$

(7) ([17]) Let X be a Petri curve of genus $g \ge 3, n \ge 5$ and $g \ge 2n - 4$. Then B(n, d, n + 1) is non-empty.

Remark 2.9. The non-emptiness of the locus B(n, d, n+1) has been proved with different relation among n, d and g (see [17], [16] or [9] and the bibliography there). For simplicity we just quote one of the results in Theorem 2.8,(7).

According to the above theorem, in some cases the non emptiness and dimension of the moduli space $\mathbb{P}(T)$ differ from general or special curves. Theorem 2.8 will be relevant in Section 5 when we consider families of extensions of semistable bundles.

The remainder of this section will be devoted to the general properties of unstable bundles.

It is well know that $H^0(E) = 0$ for semistable vector bundles E of negative degree. However, unstable bundles of negative degree can have sections. Indeed, let F be a bundle of positive degree with $H^1(F) \neq 0$. Any extension $\rho: 0 \to \mathcal{O} \to G \to F^* \to 0 \in H^1(F)$ defines an unstable bundle G with at least one section. In the following proposition we proof for unstable bundles some of the well known properties of semistable bundles. They follow directly from the properties of the HN-filtration. Maybe some are already known but we prefer to include them in one proposition.

Proposition 2.10. Let E be an indecomposable unstable vector bundle of rank n and degree d and $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ its Harder-Narasimhan filtration.

- (1) If $\mu_{min}(E) > 2g 1$ then $H^1(X, E) = 0$ and E is (globally) generated. Moreover, indecomposable unstable bundles with fix HN-filtration are bounded.
- (2) If E is HN-special, $h^0(E) \leq \frac{d}{2} + n$.
- (3) If E is HN-general and $\mu_{min}(E) > g + 1$, $H^1(X, E) = 0$ and $h^0(E) = d(E) + rk(E)(1-g)$. Moreover, if $0 < \mu_{max}(E) < g$, $h^0(E) = 0$.

Proof. Let $\rho_i: 0 \to E_{i-1} \to E_i \to F_{i-1} \to 0$ be the HN(i)-extension of E and

(2.9)
$$0 \to H^0(X, E_{i-1}) \to H^0(X, E_i) \to H^0(X, F_{i-1}) \xrightarrow{\delta}$$

(2.10)
$$\stackrel{\delta}{\to} H^1(X, E_{i-1}) \to H^1(X, E_i) \to H^1(X, F_{i-1}) \to 0$$

its cohomology sequence. Recall that we have the inequalities

$$\mu_{max}(E) = \mu(E_1) > \mu(F_1) > \dots > \mu(F_{m-1}) = \mu_{min}(E) > 2g - 1.$$

The proofs are by induction on the HN-length of E. We give the proof for the case m = 2 using the HN(2)-extension.

(1).- Since $\mu(E_1) > \mu(F_1) = \mu_{min}(E_2) > 2g - 1$, (2.10) shows that $H^1(E_2) = 0$, by the semistability of E_1 and F_1 . As E_1 and F_1 are generated we have that E_2 is generated. That are bounded follows from [32] (see also [27]).

(2).- We conclude from (2.9) that $h^{0}(E_{2}) \leq h^{0}(E_{1}) + h^{0}(F_{1})$, hence that

$$h^{0}(E_{1}) + h^{0}(F_{1}) \le \frac{d(E_{1})}{2} + rk(E_{1}) + \frac{d(F_{1})}{2} + rk(F_{1})$$

(see Theorem 2.8(2)) and finally that

$$h^0(E_2) \le \frac{d(E_2)}{2} + rk(E_2)$$

(3).- Follows also from (2.9) and (2.10), Riemann-Roch and the assumption that HN-filtration is HN-general.

From Remark 2.2 we can apply induction. Analysis similar to that in the proof of m = 2 (now using the HN(i)-extension), shows that (1), (2) and (3) are satisfied for any m > 2. The details are left to the reader.

It is known (see [26, Lemma 1.3.3]) that if E and E' are two unstable vector bundles with $\mu_{min}(E) > \mu_{max}(E')$ then Hom(E, E') = 0. Our aim is to study Hom(E, E) = End(E) for unstable bundles.

3. Algebra of endomorphisms

In this section, we determine the structure of the algebra of endomorphisms End(-) of a HN-indecomposable vector bundle of rank n, degree d and of HN-length 2. Using the HN-filtration, we give also an upper bound for dim End(-) when the HN-length > 2.

Recall from [1], that if E is indecomposable vector bundle then

$$(3.1) End(E) \cong (Id_E) \oplus Nil(E),$$

where Nil(E) is the subset $Nil(E) \subset End(E)$ consisting of all nilpotent global endomorphisms of E. In [12], the following upper bound

(3.2)
$$\dim End(E) \le 1 + \frac{n(n-1)}{2}$$

was given for indecomposable semistable bundles of rank n. The bound for indecomposable bundles of HN-length 2 is established by our next proposition.

Proposition 3.1. Let E_2 be a indecomposable unstable bundle and $0 = E_0 \subset E_1 \subset E_2$ be its HN-filtration. Then, if E_1 and $F_1 = E_2/E_1$ are indecomposable then

(3.3)
$$\dim End(E_2) \le 1 + \frac{n(n-1)}{2} + h^0(F_1^* \otimes E_1),$$

where $F_1 = E_2/E_1$.

Proof. Let $\rho_1 : 0 \to E_1 \xrightarrow{\iota} E_2 \xrightarrow{p} F_1 \to 0$ be the HN(2)-sequence of E_2 . Assume $n_1 := rk(E_1)$ and $n_2 := rk(F_1)$. We will use the following sequences

$$(3.4) \quad 0 \to H^0(X, E_1 \otimes E_2^*) \to H^0(X, E_2 \otimes E_2^*) \to H^0(X, F_1 \otimes E_2^*) \xrightarrow{\delta_0} H^1(X, E_1 \otimes E_2^*) \cdots$$

$$(3.5) \quad 0 \to H^0(X, F_1^* \otimes F_1) \to H^0(X, E_2^* \otimes F_1) \to H^0(X, E_1^* \otimes F_1) \xrightarrow{\delta_1} H^1(X, F_1^* \otimes F_1) \cdots$$

$$(3.6) \quad 0 \to H^0(X, F_1^* \otimes E_1) \to H^0(X, E_2^* \otimes E_1) \to H^0(X, E_1^* \otimes E_1) \xrightarrow{\delta_2} H^1(X, F_1^* \otimes E_1)$$

that are part of the cohomology sequences of $H^*((\rho_1) \otimes E_2^*)$, $H^*((\rho_1)^* \otimes F_1)$ and of $H^*((\rho_1)^* \otimes E_1)$, respectively.

From (3.4)

(3.7)
$$\dim End(E_2) \le h^0(E_1 \otimes E_2^*) + h^0(F_1 \otimes E_2^*).$$

The semistability of the F_1 and E_1 and the inequalities (2.2) imply that $H^0(X, E_1^* \otimes F_1) = 0$, hence from (3.5)

(3.8)
$$H^{0}(X, F_{1} \otimes E_{2}^{*}) = H^{0}(X, F_{1}^{*} \otimes F_{1}) = End(F_{1}).$$

From (3.6)

$$h^{0}(E_{1} \otimes E_{2}^{*}) \leq h^{0}(F_{1}^{*} \otimes E_{1}) + h^{0}(E_{1}^{*} \otimes E_{1})$$

since the extension $\delta_2(Id)$ is the element classifying the extension $(\rho_1)^*$ and ρ_1^* is non-trivial. Therefore, since E_1 and F_1 are indecomposables

$$\dim End(E_2) \leq h^0(E_1 \otimes E_2^*) + h^0(F_1 \otimes E_2^*) \\ = h^0(E_1 \otimes E_2^*) + \dim End(F_1) \\ \leq h^0(F_1^* \otimes E_1) + \dim End(E_1) + \dim End(F_1) \\ \leq h^0(F_1^* \otimes E_1) + \frac{n_1(n_1-1)}{2} + 1 + \frac{n_2(n_2-1)}{2} + 1 \quad (\text{see } (3.2)) \\ \leq h^0(F_1^* \otimes E_1) + 1 + \frac{(n_1^2+n_2^2)}{2} - \frac{n_1+n_2}{2} + 1 \\ \leq h^0(F_1^* \otimes E_1) + 1 + \frac{n(n-1)}{2} + 1 - n_1n_2 \\ \leq h^0(F_1^* \otimes E_1) + 1 + \frac{n(n-1)}{2} \end{aligned}$$

as claimed.

Corollary 3.2. Let E_2 be a indecomposable bundle of rank n of HN-length 2. If E_1 and F_1 are simple then dim $End(E_2) = 1 + h^0(F_1^* \otimes E_1)$ and

$$End(E_2) \cong \mathbb{C}[x_1,\ldots,x_k]/(x_1,\ldots,x_k)^2,$$

where $k = h^0(F_1^* \otimes E_1)$. Moreover, E_2 is simple if and only if $h^0(F_1^* \otimes E_1) = 0$.

г			ч	
L.			1	
	_	_		

Proof. The vector bundle E_2 is indecomposable of simple type, hence Lemma 2.4, (2) shows that $h^0(F_1 \otimes E_2^*) = 1$ and from (3.4) we conclude that $\delta_0(\lambda p) = 0$. Thus,

dim
$$End(E_2) = h^0(E_1 \otimes E_2^*) + h^0(F_1 \otimes F_1^*) = h^0(E_1 \otimes E_2^*) + 1.$$

It follows from (3.6) that $\delta_2(Id_{E_1}) \neq 0$, hence that $h^0(F_1^* \otimes E_1) = h^0(E_1 \otimes E_2^*)$, and finally that

dim
$$End(E_2) = 1 + h^0(F_1^* \otimes E_1).$$

From (3.1) we deduce that dim $Nil(E) = h^0(F_1^* \otimes E_1)$. The map

$$\varphi: H^0(F_1^* \otimes E_1) \to Nil(E)$$

defined as $\sigma \mapsto \iota \circ \sigma \circ p$ is a well defined injective homomorphism. Hence, $H^0(F_1^* \otimes E_1) \cong Nil(E)$. Moreover, $(\varphi(\sigma))^2 = 0$ and

$$(\varphi(\sigma_1)) \circ (\varphi(\sigma_2)) = (\varphi(\sigma_2)) \circ (\varphi(\sigma_1)) = 0.$$

We thus get $End(E) \cong \mathbb{C}[x_1, \ldots, x_k]/(x_1, \ldots, x_k)^2$ where $k = h^0(F_1^* \otimes E_1)$. Moreover, E_2 is simple if and only if $h^0(F_1^* \otimes E_1) = 0$ which is our claim.

Remark 3.3. Note that if $F_1^* \otimes E_1$ is special $h^0(F_1^* \otimes E_1) \cdot h^1(F_1^* \otimes E_1) \neq 0$ and if $0 \neq \rho : 0 \rightarrow E_1 \rightarrow E_2 \rightarrow F_1 \rightarrow 0 \in H^1(X, F_1^* \otimes E_1), E_2$ is indecomposable and dim $End(E_2) > 1$. Moreover, if $F_1^* \otimes E_1$ is no special. $h^0(F_1^* \otimes E_1) \cdot h^1(F_1^* \otimes E_1) = 0$, consequently, either E_2 is decomposable or is simple.

The next propositions is a fairly straightforward generalization for unstable bundles of HN-length m > 2.

Proposition 3.4. Let E be an indecomposable unstable vector bundle of rank n. If the HN-filtration $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ is HN-indecomposable then

(3.9)
$$\dim End(E) \le 1 + \frac{n(n-1)}{2} + \sum^{m-1} h^0(F_i^* \otimes E_i)$$

For the proof we will use the following lemma.

Lemma 3.5. For all $0 < i < j \le m$

- (1) $H^0(X, F_i^* \otimes F_i) = 0$ and
- (2) for all $0 < i \le j \le m$, $H^0(X, E_i^* \otimes F_j) = 0$.

Proof. (1) follows from the semistability of the quotients F_i 's and the inequalities (2.2).

(2) The semistability of E_1 and F_1 and the inequalities (2.2) imply that

$$H^{0}(X, F_{1}^{*} \otimes F_{j}) = H^{0}(X, E_{1}^{*} \otimes F_{j}) = 0,$$

Hence, from the cohomology sequence

 $H^*((\rho_1)^* \otimes F_j) : 0 \to H^0(X, F_1^* \otimes F_j) \to H^0(X, E_2^* \otimes F_j) \to H^0(X, E_1^* \otimes F_j) \to \cdots$

we conclude that $H^0(X, E_2^* \otimes F_j) = 0$. In the same manner we can see that $H^0(X, E_i^* \otimes F_j) = 0$ for all $i \leq j \leq m$. The detailed verification being left to the reader.

Proof of Proposition 3.4 The proof is by induction on the HN-length. The first step of induction is Proposition 3.1. From the cohomology sequence $H^*(\rho_{m-1} \otimes E_m^*)$

(3.10)
$$\dim End(E_m) \le h^0(E_{m-1} \otimes E_m^*) + h^0(F_{m-1} \otimes E_m^*).$$

We can now proceed analogously to the proof of Proposition 3.1. From the cohomology sequence $H^*((\rho_{m-1})^* \otimes F_{m-1})$ and Lemma 3.5,(2) it follows that

$$H^0(F_{m-1} \otimes E_m^*) = H^0(F_{m-1}^* \otimes F_{m-1}).$$

From the cohomology sequence $H^*((\rho_{m-1})^* \otimes E_{m-1})$ we have the inequality

$$h^{0}(E_{m-1} \otimes E_{m}^{*}) \leq h^{0}(End(E_{m-1})) + h^{0}(F_{m-1}^{*} \otimes E_{m-1}).$$

Recall that E_m is HN-indecomposable.

Now (3.10) becomes

$$\dim End(E_m) \leq h^0(End(E_{m-1})) + h^0(F_{m-1}^* \otimes E_{m-1}) + h^0(F_{m-1}^* \otimes F_{m-1})$$

$$\leq \frac{rk(E_{m-1})(rk(E_{m-1})-1)}{\sum^{m-2} h^0(F_i^* \otimes E_i)} + \frac{rk(F_{m-1})(rk(F_{m-1})-1)}{2} + 2 + \frac{rk(E_m)(rk(E_m)-1)}{2} + \frac{rk(E_m)(rk(E_m)-1)}{$$

which is the desired conclusion.

The following result may be proved in much the same way as Corollary 3.2.

Corollary 3.6. Let E be an indecomposable unstable vector bundle of rank n of simple type. If $0 = E_0 \subset E_1 \subset \cdots \subset E_m = E$ is its HN-filtration then

(3.11)
$$\dim End(E) = 1 + h^0(F_{m-1}^* \otimes E_{m-1})$$

Moreover, $h^0(F_{m-1}^* \otimes E_{m-1}) \le \sum_{0}^{m-2} h^0(F_{m-1}^* \otimes F_i).$

Proof. We give only the main ideas of the proof. We can proceed analogously to the proof of Corollary 3.2. The equality dim $End(E) = 1 + h^0(F_{m-1}^* \otimes E_{m-1})$ follows from Lemma 3.5 and the cohomology of the exact sequences

$$0 \to E_{m-1} \otimes E^* \to E \otimes E^* \to F_{m-1} \otimes E^* \to 0,$$

$$0 \to F_{m-1}^* \otimes F_{m-1} \to E^* \otimes F_{m-1} \to E_{m-1}^* \otimes F_{m-1} \to 0,$$

$$0 \to F_{m-1}^* \otimes E_{m-1} \to E^* \otimes E_{m-1} \to E_{m-1}^* \otimes E_{m-1} \to 0,$$

since E is of simple type, and hence HN-indecomposable.

The basic idea of the proof of the bound for $h^0(F_{m-1}^* \otimes E_{m-1})$ is to take the cohomology of the sequences

$$0 \to E_{i-1} \otimes F_{m-1}^* \to E_i \otimes F_{m-1}^* \to F_{i-1} \otimes F_{m-1}^* \to 0,$$

and proceed by induction. The details are left to the reader.

MODULI OF UNSTABLES

4. Moduli spaces

In this section Proposition 2.6 is given in a more general setting. Our aim is to consider families of semistable bundles. Recall that if the vector bundles E_1 and F_1 have automorphisms the \mathbb{C}^* -action does not identify all the isomorphic classes of the vector bundles in $H^1(X, E_1 \otimes F_1^*)$. The simple vector bundles over X with fixed rank and degree possess a coarse moduli space (see [30, Corollary 6.5]), but there is no universal family. It is possibly that it is non-separated and by the work of M. Artin is an algebraic space. Therefore, to construct a moduli scheme of indecomposable vector bundles of HN-length 2 we will consider those of coprime type $\sigma = (\mu_1, \mu_2)$. Note that if μ_1 is given by $\mu_1 = \frac{d_1}{n_1}$ then $\mu_2 = \frac{d-d_1}{n-n_1} := \frac{d_2}{n_2}$. From now on we tacitly assume that $\mu_1 = \frac{d_1}{n_1} > \frac{d_2}{n_2} = \mu_2$ and denote by $U_{\mu_1}(n, d)$ the set of such bundles. Let

$$U_{\mu_1}(n,d,k) := \{ E \in U_{\mu_1}(n,d) : \dim End(E) = 1+k \}.$$

The moduli space $M(n_i, d_i)$ of semistable vector bundles of rank n_i and degree d_i for 1 = 1, 2 with $gcd(n_i, d_i) = 1$ carries a universal bundle $\mathcal{U}_i \to X \times M(n_i, d_i)$. In general, \mathcal{U}_i is determined up to tensoring by a line bundle lifted from $M(n_i, d_i)$. In this paper it will be fixed, unless otherwise stated. Denote by p_{ij} the projection of $X \times M(n_1, d_1) \times M(n_2, d_2)$ in the ij-factors. We will denote by \mathcal{R}_{μ_1} the 1st- direct image sheaf

$$\mathcal{R}_{\mu_1} := \mathcal{R}^1_{p_{23}}(p_{12}^*\mathcal{U}_1 \otimes p_{13}^*\mathcal{U}_2^*)$$

over $M(n_1, d_1) \times M(n_2, d_2)$.

With the notation $M_i := M(n_i, d_i)$, the following diagram summarise the notation.

The following theorem follows from [21], [31], Proposition 2.5 and Lemma 3.5.

Theorem 4.1. If $\mu_1 - \frac{d_2}{n_2} \ge 2g - 1$, $U_{\mu_1}(n, d) = \emptyset$ and if $\mu_1 - \frac{d_2}{n_2} < 2g - 1$, $U_{\mu_1}(n, d)$ has a projective scheme structure that makes it an moduli scheme. Moreover, there is a natural isomorphism between $U_{\mu_1}(n, d)$ and $Proj(\mathcal{R}_{\mu_1})$.

Proof. Let $E \in U_{\mu_1}(n, d)$. Note that being of coprime type implies that E is an extension of two stable bundles, that is

$$\rho_1: 0 \to E_1 \to E \to F_1 \to 0.$$

Since $Ext^{2}(p_{13}^{*}\mathcal{U}_{2}, p_{12}^{*}\mathcal{U}_{1}) = 0$, it follows that $\mathcal{R}_{p_{23}}^{1}(p_{12}^{*}\mathcal{U}_{1} \otimes p_{13}^{*}\mathcal{U}_{2}^{*})|_{(E_{1},F_{1})} \to H^{1}(X, E_{1} \otimes F_{1}^{*})$ is an isomorphism. From [21], [31], the coherent sheaf $\mathcal{R}_{\mu_{1}} := \mathcal{R}_{p_{23}}^{1}(p_{12}^{*}\mathcal{U}_{1} \otimes p_{13}^{*}\mathcal{U}_{2}^{*})$

parameterizes the classes of extensions of two stable bundles $(E_1, F_1) \in M_1 \times M_2$. From Lemma 3.3,(1) E is indecomposable if and only if $\rho_2 \neq 0$. Therefore, the theorem follows from Remark 2.3.

Remark 4.2. Note that the coherent sheaf \mathcal{R}_{μ_1} could be defined even if $gcd(n_0, d_0) \neq 1$. Indeed, if $gcd(n_i, d_i) \neq 1$, there exists an étale covering \widetilde{M}_i of M_i such that a universal bundle $\widetilde{\mathcal{U}}_i$ exists on $X \times \widetilde{M}_i$ (see [37, Proposition 2.4]). Thus, the coherent sheaf $\widetilde{\mathcal{R}}_{\mu_1} := \mathcal{R}_{p_{23}}^1(p_{12}^*\widetilde{\mathcal{U}}_1 \otimes p_{13}^*\widetilde{\mathcal{U}}_2^*)$ will parameterize extensions of two semistable bundles $(E_1, F_1) \in \widetilde{M}_1 \times \widetilde{M}_2$. However, in this case we can not apply Lemma 3.3,(1) (see Remark 2.3) since semistable bundles can be non-simple.

In general, the coherent sheaf $\mathcal{R}_{\mu_1} = \mathcal{R}_{p_{23}}^1(p_{12}^*\mathcal{U}_1 \otimes p_{13}^*\mathcal{U}_2^*)$ is not locally free. We will give a flattening stratification of $M_1 \times M_2$ for \mathcal{R}_{μ_1} . To give the schematic semi-continuity stratification of $M_1 \times M_2$ we will use the algebra of endomorphisms and the twisted Brill-Noether theory. For a recent account of the theory we refer the reader to [24]. We follow [2] in the construction of the determinantal varieties that we need.

Let D be an effective divisor on X of degree $d_0 >> 0$ such that for any $(E_1, F_1) \in M_1 \times M_2$, $H^1(X, E_1 \otimes F_1^* \otimes \mathcal{O}(D)) = 0$. Let

$$0 \to H^0(X, E_1 \otimes F_1^*) \to H^0(X, E_1 \otimes F_1^* \otimes \mathcal{O}(D)) \xrightarrow{\phi} H^0(X, (E_1 \otimes F_1^*)|_D) \to H^1(X, E_1 \otimes F_1^*) \to 0$$

be the cohomology sequence of

$$0 \to E_1 \otimes F_1^* \to E_1 \otimes F_1^* \otimes \mathcal{O}(D) \to (E_1 \otimes F_1^*)_{|_D} \to 0.$$

Thus, $h^0(E_1 \otimes F_1^*) \ge k$ if and only if $rk(\phi) < h^0(E_1 \otimes F_1^* \otimes \mathcal{O}(D)) - k$.

We want to vary (E_1, F_1) in $M_1 \times M_2$. Let $\Gamma := D \times M_1 \times M_2$ be the product divisor in $X \times M_1 \times M_2$. We have the following sequence

$$0 \to p_{12}^* \mathcal{U}_1 \otimes p_{13}^* \mathcal{U}_2^* \to p_{12}^* \mathcal{U}_1 \otimes p_{13}^* \mathcal{U}_2^* \otimes \mathcal{O}(\Gamma) \to \mathcal{N} \to 0,$$

over $X \times M_1 \times M_2$, where \mathcal{N} is the quotient $p_{12}^* \mathcal{U}_1 \otimes p_{13}^* \mathcal{U}_2^* \otimes \mathcal{O}(\Gamma)/p_{12}^* \mathcal{U}_1 \otimes p_{13}^* \mathcal{U}_2^*$.

Since $H^1(E_1 \otimes F_1^* \otimes \mathcal{O}(D)) = 0$, the direct image induces the complex $\phi : \mathcal{K}_0 \to \mathcal{K}_1$ of locally free sheaves over $M_1 \times M_2$ where $\mathcal{K}^0 := \mathcal{R}^0_{p_{23}}(p_{12}^*\mathcal{U}_1 \otimes p_{13}^*\mathcal{U}_2^* \otimes \mathcal{O}(\Gamma))$ is of rank $d(E_1 \otimes F_1^* \otimes \mathcal{O}(D)) + n_1 n_2 (1-g)$ and $\mathcal{K}^1 := \mathcal{R}^0_{p_{23}}(\mathcal{N})$. We follow the notation of [24] and denote by $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ the kth-determinantal variety of the complex $\phi : \mathcal{K}_0 \to \mathcal{K}_1$. That is,

(4.2)
$$Supp(B^{k}(\mathcal{U}_{1},\mathcal{U}_{2}^{*})) := \{(E_{1},F_{1}) \in M_{1} \times M_{2} : h^{0}(E_{1} \otimes F_{1}^{*}) \geq k\}.$$

Remark 4.3. When no confusion can arise and to simplify notation, from now on we use the following notation:

• $n=n_1+n_2$,

•
$$d = d_1 + d_2$$

• $n_0 = n_1 n_2$,

- d₀ = n₂d₁ − n₁d₂,
 h⁰ = k and
- $h^1 = k d_0 + n_0(g 1).$

Hence, as kth-determinantal variety, the expected dimension of $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ is the number

$$\rho(g, n_1, d_1, n_2, d_2, k) := \dim(M_1 \times M_2) - h^0 \cdot h^1 \\
= (n_1^2 + n_2^2)(g - 1) + 2 - k(k - d_0 + n_0(g - 1)).$$

and $B^{k+1}(\mathcal{U}_1, \mathcal{U}_2^*) \subseteq Sing B^k(\mathcal{U}_1, \mathcal{U}_2^*)$. Moreover, the filtration

$$M_1 \times M_2 \supset B^1(\mathcal{U}_1, \mathcal{U}_2^*) \supset \cdots \supset B^k(\mathcal{U}_1, \mathcal{U}_2^*) \supset B^{k+1}(\mathcal{U}_1, \mathcal{U}_2^*) \supset \cdots \cdots$$

define a stratification by closed subsets. We will denote by \mathcal{Y}_k the stratum

$$\mathcal{Y}_k := B^k(\mathcal{U}_1, \mathcal{U}_2^*) - B^{k+1}(\mathcal{U}_1, \mathcal{U}_2^*).$$

That is,

$$\mathcal{Y}_k = \{ (E_1, F_1) \in M_1 \times M_2 : h^0(E_1 \otimes F_1^*) = k \}.$$

For any $(E_1, F_1) \in \mathcal{Y}_k$,

$$\dim H^1(X, E_1 \otimes F_1^*) = k - d(E_1 \otimes F_1^*) + n_1 n_2(g-1) = k - d_0 + n_0(g-1).$$

Therefore, the restriction of \mathcal{R}_{μ_1} to \mathcal{Y}_k , denoted by $\mathcal{R}_{\mu_1}(\mathcal{Y}_k) \to \mathcal{Y}_k$, is locally free of rank $k - d_0 + n_0(g - 1)$. Let

 $\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_k)) \to \mathcal{Y}_k$

be the projective bundle associated to $\mathcal{R}_{\mu_1}(\mathcal{Y}_k) \to \mathcal{Y}_k$.

The expected dimension of $\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_k))$ is

$$\beta(g, n_1, d_1, n_2, d_2, k) := \dim(M_1 \times M_2) - h^0 h^1 + h^1 - 1 = \dim(M_1 \times M_2) - h^1 (h^0 - 1) - 1 = (n_1^2 + n_2^2)(g - 1) + 1 - (k - 1)(k - d_0 + n_0(g - 1)).$$

Remark 4.4. \mathcal{Y}_0 is the locus where $h^0(E_1 \otimes F_1^*) = 0$, and hence, the indecomposable bundles $E \in H^1(X, E_1 \otimes F_1^*)$ are simple (see Corollary 3.2). Under the assumption that $M_1 \times M_2 \neq B^1(\mathcal{U}_1, \mathcal{U}_2^*)$,

$$\mathcal{Y}_0 = M_1 \times M_2 - B^1(\mathcal{U}_1, \mathcal{U}_2^*)$$

is an open set and, in consequence,

$$\dim \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0)) = \dim M_1 + \dim M_2 + h^1 - 1.$$

Remark 4.5. (1) If $p_2 : B^k(\mathcal{U}_1, \mathcal{U}_2^*) \subset M_1 \times M_2 \to M_2$ is the projection then for $F_1 \in p_2(B^k(\mathcal{U}_1, \mathcal{U}_2^*))$ the inverse image $p_2^{-1}(F_1)$ is the locus

$$B^{k}(\mathcal{U}_{1}, F_{1}^{*}) := \{ E_{1} \in M_{1} : h^{0}(E_{1} \otimes F_{1}^{*}) \ge k \}$$

(2) In particular, if $M_2 = Pic^0(X)$ and $F_1 = \mathcal{O}_X$ then $B^k(\mathcal{U}_1, \mathcal{O}_X)$ is the Brill-Noether locus $B(n_1, d_1, k)$. In this case we denote \mathcal{Y}_k as Y_k . That is,

$$Y_k = B(n_1, d_1, k) - B(n_1, d_1, k+1).$$

- (3) In general, the product of two stable bundles is semistable. However, from [6, Lemma 3.5] if on of the bundles is general then $E \otimes F^*$ is stable, or if $(n_0, d_0) = 1$. Thus, in this case if $B^k(\mathcal{U}_1, \mathcal{U}_2^*) \neq \emptyset$ then $B(n_0, d_0, k) \neq \emptyset$.
- (4) If $M_1 \times M_2 \neq B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ and $B^k(\mathcal{U}_1, \mathcal{U}_2^*) \neq \emptyset$ then $\mathcal{Y}_k \neq \emptyset$.

Using the notation of Remark 4.3 we can now formulate one of our main results. For any $0 \le k \le \frac{d_0}{2} + n_0$, let $U_{\mu_1}(n, d, k)$ be the set

$$U_{\mu_1}(n, d, k) := \{ E \in U_{\mu_1}(n, d) : \dim End(E) = 1 + k \}.$$

Theorem 4.6. $U_{\mu_1}(n, d, k) = \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_k))$ is coarse moduli space and $U_{\mu_1}(n, d, 0)$ is a fine moduli space. Moreover, if \mathcal{Y}_k is irreducible and smooth of dimension ρ , then $U_{\mu_1}(n, d, k)$ is irreducible and smooth of dimension $\rho + h^1 - 1$.

Proof. From what has already been proved and Corollary 3.2 it follows that

$$U_{\mu_1}(n,d,k) = \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_k))$$

If $E \in U_{\mu_1}(n, d, 0)$, E is simple and hence from Corollary 3.2 $h^0(E_1 \otimes F_1^*) = 0$. Thus, $(\mathcal{R}^0_{p_{23}}(p_{12}^*\mathcal{U}_1 \otimes p_{13}^*\mathcal{U}_2^*))|_{\mathcal{Y}_0} = 0$. Hence, from [31, Corollary 4.5], $\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0))$ parameterize a universal extension

$$0 \to q^*(p_{12}^*(\mathcal{U}_1)) \otimes p_2^* \mathbb{H} \to \mathcal{E} \to q^* p_{13}^*(\mathcal{U}_2) \to 0,$$

where $q : X \times \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0)) \to X \times \mathcal{Y}_0$ is the induced map, $p_2 : X \times \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0)) \to \mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0))$ is the projection and \mathbb{H} the hyperplane bundle over $\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0))$. The universal properties of the family \mathcal{E} imply that the pair $(\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_0)), \mathcal{E})$ is the fine moduli space for $U_{\mu_1}(n, d, 0)$.

If \mathcal{Y}_k is irreducible and smooth of dimension ρ , a straightforward computation shows $U_{\mu_1}(n, d, k)$ is irreducible and smooth of dimension $\rho + h^1 - 1$, since $\mathbb{P}(\mathcal{R}_{\mu_1}(\mathcal{Y}_k))$ is a projective bundle with fibre $\mathbb{P}(H^1(X, E_1 \otimes F_1^*))$ at $(E_1, F_1) \in \mathcal{Y}_k$.

Corollary 4.7. If $U_{\mu_1}(n, d, k)$ is non-empty, $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ is non-empty.

Corollary 4.8. If \mathcal{Y}_k is irreducible and smooth then $H^i(\mathcal{U}_{\mu_1}(n, d, k), \mathbb{C}) \cong H^i(\mathcal{Y}_k, \mathbb{C})$ for $i \geq 0$.

Remark 4.9. The moduli space of simple bundles of type $\sigma = (\mu(E_1), \ldots, \mu(E_m/E_{m-1}))$, for m > 2, are considered in [8].

Remark 4.10. Theorem 4.6 expresses the equivalence of the existence and topology of the $U_{\mu_1}(n, d, k)$ and that of twisted Brill-Noether loci. For some values of (g, n, d, k), non emptiness, dimension, and irreducibility of $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$, and of \mathcal{Y}_k , are known for general curve (see [24]). Thus, as in the Brill-Noether and twisted Brill-Noether theory for vector bundles, it is possible that for special curves the moduli space $U_{\mu_1}(n, d, k)$ is even reduced. Thus, the above results differ from the corresponding results for the moduli space of stable bundles, where non-emptiness, dimension etc. are independent of the curve.

MODULI OF UNSTABLES

5. Non-emptiness of $U_{\mu_1}(n, d, k)$

In this section we will prove non emptiness of $U_{\mu_1}(n, d, k)$ for some values of (g, n, d, k). First we consider the case of bundles of type $\sigma = (\mu_1, \mu_2)$ where $\mu_1 = \frac{d-a}{n-1}$ and $\mu_2 = a$ is an integer. That is, the quotient of the maximal destabilizing subbundle is a line bundle of degree a. In this case \mathcal{Y}_k is a subset of the Brill-Noether locus B(n-1, d-an, k), and will be denoted as Y_k .

The next theorems are applications the known results on B(n-1, d-an, k) and Theorem 4.6.

Theorem 5.1. Let n, d, a and k be positive integers. Assume 0 < d - an < 2(n - 1) and $(n - 1, d - an, k) \neq (n - 1, n - 1, n - 1)$. Then for $\mu_1 = \frac{d - a}{n - 1}$, $U_{\mu_1}(n, d, k)$ is non-empty if and only if $k \leq n - 1 + \frac{d - n(a+1) + 1}{g}$. Moreover, if $U_{\mu_1}(n, d, k)$ is non-empty then it is irreducible and smooth of the expected dimension.

Proof. The theorem follows from Theorem 4.6, since from [13, Theorem A] and [35, A-1 Théorème], $Y_k \subset B(n-1, d-an, k)$ has the required properties.

Remark 5.2. Note that under the assumptions of Theorem 5.1, the positivity of the expected dimension $\rho(n-1, d-an, 1, a, k)$ does not imply non emptiness of $U_{\mu_1}(n, d, k)$.

For g = 2 we have a complete description.

Corollary 5.3. Assume g = 2. If 0 < d - an < 2(n - 1) and $0 \le k \le \frac{d - an}{2} + \frac{n - 1}{2}$ then $U_{\mu_1}(n, d, k)$ is irreducible and smooth of the expected dimension. If $d - an \ge 2(n - 1)$, $U_{\mu_1}(n, d, k) = \emptyset$.

We now rephrase Theorem 2.8,(5),(6) and (7) as follows.

Theorem 5.4. Let (g, n - 1, d - na, k) be integers that satisfies the conditions given in Theorem 2.8,(5),(6) and (7). For general curve, $U_{\mu_1}(n, d, k)$ is non-empty and has an irreducible component of the expected dimension and if X is a Petri curve of genus $g \ge 3, n \ge 5$ and $g \ge 2n - 4$ then $U_{\mu_1}(n, d, n)$ is non-empty.

In the above case, to the best of our knowledge, it is know that there exists an irreducible component of the expected dimension, however, in general, it is not known that B(n, d, k) is irreducible.

Let us now consider the general case. We will use the notation given in Remark 4.3. Recall that $B^k(\mathcal{U}_1, \mathcal{U}_2^*) \subset M_1 \times M_2$. Denote by M_{12} the moduli space of stable bundles $M(n_0, d_0)$. From what has already been proved, we conclude that if $B(n_0, d_0, k) = \emptyset$ then $U_{\mu_1}(n, d, k) = \emptyset$. However, if $B(n_0, d_0, k) \neq \emptyset$ does not imply that $U_{\mu_1}(n, d, n) \neq \emptyset$. Let $\Phi: M_1 \times M_2 \to M_{12}$ be the morphism defined as $(E, F) \mapsto E \otimes F^*$. This gives

$$\Phi(B^{\kappa}(\mathcal{U}_1,\mathcal{U}_2^*)) \subset B(n_0,d_0,k).$$

We want to describe $\Phi(B^k(\mathcal{U}_1, \mathcal{U}_2^*))$ in some cases.

Remark 5.5. Let G be a vector bundle of rank m and degree d_2 generated by linear subspace $V \subset H^0(X, E)$ of dimension $k_2 = m + n_2$. Let $D^*_{E,V}$ be the kernel of the evaluation map (or the syzygy bundle). Let us introduce the temporary notation F_1^* for $D^*_{E,V}$. That is, F_1^* has rank $rk(F_1) = n_2$, degree $d(F_1^*) = -d_2$ and fits into the following exact sequence

$$(5.1) 0 \to F_1^* \to V \otimes \mathcal{O} \to G \to 0.$$

Tensor (5.1) with a vector bundle E_1 of rank n_1 and degree d_1 with $h^0(E_1) \ge k_1$. The injectivity of the multiplication map $\mu_{V,E_1} : V \otimes H^0(E_1) \to H^0(G \otimes E_1)$ is measure by $h^0(F_1^* \otimes E_1)$. Indeed, if $H^0(F_1^* \otimes E_1) = 0$, μ_{V,E_1} is injective. From the cohomology sequence

$$0 \to H^0(F_1^* \otimes E_1) \to V \otimes H^0(E_1) \to H^0(G \otimes E_1) \to$$

we obtain the inequality

(5.2)
$$\dim V \cdot h^0(F) - h^0(G \otimes E_1) \le h^0(F_1^* \otimes E_1)$$

Thus, if

$$(5.3) k \le k_1 k_2 - h^0 (G \otimes E_1)$$

then $k \leq h^0(F_1^* \otimes E_1)$.

With the above notation, $n_0 = rk(F_1^* \otimes E_1) = n_1n_2$ and $d_0 = d(F_1^* \otimes E_1) = d_1n_2 - d_2n_1$. Let $\tilde{n} := rk(G \otimes E_1) = mn_1$ and $\tilde{d} := d(G \otimes E_1) = d_2n_1 + d_1m$. Our interest is to apply Remark 5.5 to situations in which E_1 is general in $M(n_1, d_1) = M_1$, $F_1 \in M(n_2, d_2) = M_2$ and $k \ge 0$, in particular when μ_{V,E_1} is not injective. In that case, $(E_1, F_1^*) \in B^k(U_1, U_2^*)$ and $E_1 \otimes F_1^* \in B(n_0, d_0, k)$. The following theorems give existence of some Brill-Noether and twisted Brill-Noether loci, and therefore the non emptiness of $U_{\mu_1}(n, d, k)$. The proofs are different of those of [24] and some values, to our best knowledge, are not been include in [24] nor in Theorem 2.8 (see [7]).

Theorem 5.6. Assume that $B(n_1, d_1, n_1+a)$ is non-empty with a > 0. If $2n_1 < d_1 < a(g+1)$ and $d_2 > 2gm$ then for any $0 \le k \le (d_2+m(1-g))(n_1+a)-(d_2n_1+d_1m+mn_1(1-g))$, $\mathcal{Y}_k \subset B^k(U_1, U_2^*)$ is non-empty. Moreover, if $\mu_1 = \frac{d_1}{n_1}$ then $U_{\mu_1}(n, d, k)$ is non-empty, where $n = n_1 + n_2$ and $d = d_1 + d_2$.

Proof. Let $G \in M(m, d_2)$ and $E_1 \in B(n_1, d_1, n_1 + a)$. We want to prove that $0 \le k_1k_2 - h^0(G \otimes E_1)$, where $k_1 = n_1 + a$ and $k_2 = d_2 + m(1 - g)$.

If $d_2 > 2gm$, G is generated and from [18, Theorem 1.2] we have the exact sequence

(5.4)
$$0 \to F_1^* \to H^0(G) \otimes \mathcal{O} \to G \to 0$$

with $F_1 \in M(n_2, d_2)$ where $n_2 := h^0(G) - m = d_2 - mg$. From Remark 5.5 and (5.3)

(5.5)
$$h^{0}(G) \cdot (n_{1} + a) \leq h^{0}(G) \cdot h^{0}(E_{1}) - h^{0}(G \otimes E_{1}) \leq h^{0}(E_{1} \otimes F_{1}^{*}).$$

If $d_1 < a(g+1)$ and $d_2 > 2gm$ then $\frac{d_1}{a} + g - 1 < 2g < \frac{d_2}{m}$. Hence, (5.6) $d_1m + am(g-1) < ad_2$. Now add in both sides of 5.6 $d_2n_1 + mn_1(g-1)$ to obtain

$$d_2n_1 + d_1m + (n_1 + a)m(g - 1) < d_2(n_1 + a) + n_1m(g - 1).$$

Hence,

(5.7)
$$0 < (n_1 + a)(d_2 + m(1 - g)) - (d_2n_1 + d_1m + mn_1(1 - g)) = k_1k_2 - h^0(G \otimes E_1).$$

This gives $0 \le k \le h^0(E_1 \otimes F_1^*)$, for any $0 \le k \le (d_2 + m(1-g))(n_1 + a) - (d_2n_1 + d_1m + mn_1(1-g))$. It follows that $B^k(U_1, U_2^*)$ is non-empty and, in consequence, $U_{\mu_1}(n, d, k)$ is non-empty.

Theorem 5.7. Let X be a Petri curve of genus $g \ge 3$ and $(\mathcal{O}(D), V)$ a general generated linear system of degree $d_2 \ge g + 1$ and dim $V = n_2 + 1$ with $n_2 \le 4$ or if $n_2 \ge 5$ then $g \ge 2n_2 - 4$. Assume that $B(n_1, d_1, t)$ is non-empty and $\frac{d_2}{n_2} < \frac{d_1}{n_1}$. For any $0 \le k \le n_2 t - n_1 d_2$, $\mathcal{Y}_k \subset B^k(U_1, U_2^*)$ is non-empty and if $n = n_2 + n_1$, $d = d_2 + d_1$ and $\mu_1 = \frac{d_1}{n_1}$, $U_{\mu_1}(n, d, k)$, is non-empty.

Proof. Under the above conditions there exist an exact sequence

(5.8)
$$0 \to F_1^* \to V \otimes \mathcal{O} \to \mathcal{O}(D) \to 0$$

such that $F_1 \in M(n_2, d_2)$ (see [9], [16] and [17]).

Let $E_1 \in B(n_1, d_1, n_1 + a)$ such that $h^0(E_1) = t \ge n_1 + a$. From Remark 5.5 and (5.3)

(5.9)
$$(n_2+1) \cdot t - h^0(\mathcal{O}(D) \otimes E_1) \le h^0(F_1^* \otimes E_1).$$

From the assumption $\frac{d_2}{n_2} < \frac{d_1}{n_1}$ and the exact sequence

$$(5.10) 0 \to E_1 \to \mathcal{O}(D) \otimes E_1 \to (E_1)_D \to 0$$

we deduce that

(5.11)
$$tn_2 - d_2n_1 < (n_2 + 1) \cdot t - h^0(\mathcal{O}(D) \otimes E_1) \le h^0(F_1^* \otimes E_1).$$

Therefore, for any $0 \le k \le tn_2 - d_2n_1$, $\mathcal{Y}_k \subset B^k(U_1, U_2^*)$ is non-empty and if $n = n_2 + n_1$, $d = d_2 + d_1$ and $\mu_1 = \frac{d_1}{n_1}$, $U_{\mu_1}(n, d, k)$, is non-empty, as claimed.

Remark 5.8. Under the hypothesis of Theorems 5.6 and 5.7 we can choose the d_2, n_2, n_1 and d_1 such that $(n_0, d_0) = 1$, and hence $B(n_0, d_0, k)$ is non-empty.

We conclude now with a description of a smooth point of $U_{\mu_1}(n, d, k)$. Recall that $U_{\mu_1}(n, d, k)$ is a projective bundle over \mathcal{Y}_k . Let $0 \subset E_1 \subset E$ be the HN-filtration of $E \in U_{\mu_1}(n, d, k)$. Write $E/E_1 = F_1$. Let us describe the tangent bundle of $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ at a point (E_1, F_1^*) . We abbreviate $B^k(\mathcal{U}_1, \mathcal{U}_2^*)$ to \mathcal{B}_{12} . Since $\mathcal{B}_{12} \subset M_1 \times M_2$, for any $z := (E_1, F_1) \in \mathcal{B}_{12}$ we have the sequence

(5.12)
$$0 \to T_z \mathcal{B}_{12} \to T_{E_1} M_1 \oplus T_{F_1} M_2 \to N \to 0$$

where N is the normal bundle. Moreover, from the morphism $\Phi: M_1 \times M_2 \to M_{12}$ we get

(5.13)
$$T_{E_1}M_1 \oplus T_{F_1}M_2 \xrightarrow{d\Phi} T_z M_{1,2}$$

That is,

$$H^1(X, End(E_1)) \oplus H^1(X, End(F_1)) \xrightarrow{d\Phi} H^1(X, End(E_1 \otimes F_1^*))$$

1.4

Moreover, from the dual of the multiplication map

$$H^0(X, E_1 \otimes F_1^*) \otimes H^0(X, (E_1 \otimes F_1^*)^* \otimes K) \to H^0(X, End(E_1 \otimes F_1^*) \otimes K)$$

we have the morphism

$$H^1(X, End(E_1 \otimes F_1^*)) \xrightarrow{\beta} H^0(X, E_1 \otimes F_1^*)^* \otimes H^1(X, E_1 \otimes F_1^*).$$

Thus,

$$\begin{array}{ccc} H^1(X, End(E_1)) \oplus H^1(X, End(F_1)) & \stackrel{d\Phi}{\to} & H^1(X, End(E_1 \otimes F_1^*)) \\ & \eta \searrow & & \beta \downarrow \\ & & & H^0(X, E_1 \otimes F_1^*)^* \otimes H^1(X, E_1 \otimes F_1^*) \end{array}$$

where $\eta = \beta \circ d\Phi$ and the image of η is precisely the normal bundle. Hence from [24, Proposition 3.5] and Theorem 4.6 we conclude

Theorem 5.9. $U_{\mu_1}(n, d, k)$ is smooth at E and of expected dimension

 $\beta(g, n_1, d_1, n_2, d_2, k)$ if and only if η is surjective.

References

- Atiyah, M. F.: Complex analityc connections in fibres bundles, Trans. Amer. Math. Soc. 85 (1957), 181–207.
- [2] Arbarello, E., Cornalba, M., Griffiths, P.A., Harris, J.: Geometry of algebraic curves. In: Vol. I, Volume 267 Springer, New York (1985).
- [3] Bănică, C.: Sur les fibrés instables de rang 2 sur $\mathbb{P}^3(\mathbb{C})$, Arch. Math. (Basel) 43 :3 (1984), 250–257.
- [4] Bănică, C., Putinar, M., Schumacher, G.: Variation der globalen Ext in Deformationen kompakter komplexer Rfiume. Math. Ann. 250 (1980), 135–155.
- [5] Bérczi G., Hoskins V., Jackson J., Kirwan F.: Moduli of unstable sheaves. In preparation.
- [6] Biswas, I., Brambila-Paz, L., Newstead, P.E.: Stability and deformations of generalised Picard sheaves, arXiv:1812.09732
- [7] Brambila-Paz, L, Newsteasd, P. E.: New examples of twisted Brill-Noether loci, in preparation.
- [8] Brambila-Paz, L.: On the moduli space of simple unstable bundles, In preparation.
- [9] Brambila-Paz, L.: Non-emptiness of moduli spaces of coherent systems, Internat. J. Math. 18, 777– 799.
- [10] Brambila-Paz, L.: Nonsimple semistable vector bundles over a curve. Math. Z. 212 (1993), no. 3, 301–311.
- [11] Brambila-Paz, L.: Moduli of endomorphisms of semistable vector bundles over a compact Riemann surface. Glasgow Math. J. 32 (1990), no. 1, 1–12.
- [12] Brambila-Paz, L.: Algebras of endomorphisms of semistable vector bundles of rank 3 over a Riemann surface. J. Algebra 123 (1989), no. 2, 414–425.
- Brambila-Paz, L., Grzegorczyk, I., Newstead, P.E.: Geography of Brill-Noether loci for small slopes. J. Algebraic Geom. 6 (1997), no.4, 645–669.
- [14] Brambila-Paz L., Mata-Gutiérrez O.: Un teorema de Torelli para haces vectoriales inestables. Aportaciones Matemáticas-Memorias de la Sociedad Matemática Mexicana 46 (2013), 03–16.
- [15] Brambila-Paz, L., Mercat, V., Newstead, P.E., Ongay, F.: Nonemptiness of Brill-Noether loci. Internat. J. Mat. 11 (2000), no.6, 737–760.
- [16] Bhosle, U.N., Brambila-Paz, L., Newstead, P.E.: On coherent systems of type (n, d, n + 1) on Petri curves. Manuscr. Math. 126(4) (2008), 409–441.
- [17] Bhosle, U.N., Brambila-Paz, L., Newstead, P.E.: On linear series and a conjecture of D. C. Butler International Journal of Mathematics Vol.26 (2015), no.02.

- [18] Butler, D. C.: Normal generation of vector bundles over a curve. J. Diff. Geom. 39 (1994), 1-34.
- [19] Drézet, Jean-Marc: Déformations des extensions larges de faisceaux, Pacific J. Math. 220 (2015), 201–297.
- [20] Gieseker, D.: On the Moduli of Vector Bundles on an Algebraic Surface, Annals of Mathematics Second Series, Vol. 106, No. 1 (1977), 45–60.
- [21] Grothendieck, A., Dieudonn, I.A.: "Éliments de géométrie algébrique I," Grundlehren 166, Springer, New York, 1971.
- [22] Grzegorczyk, I., Teixidor i Bigas, M.: Brill-Noether theory for stables vector bundles. Cambridge University Press, (2009), 29–50.
- [23] Harder, G., Narasimhan, M.S.: On the Cohomology Groups of Moduli Spaces of Vector Bundles on Curves. Ann. of Math. 212 (1975), 215–248.
- [24] Hitching, G.H., Hoff, M., Newstead, P.E.: Nonemptiness and smoothness of twisted Brill–Noether loci. Annali di Matematica Pura ed Applicata (2021) 200:685–709.
- [25] Hoskins, V., Jackson J.: Quotients by parabolic groups and moduli spaces of unstables objects. arXiv:2111.07429v1 [Math.AG] (2021).
- [26] Huybrechts, D. and Lehn M.: The Geometry of Moduli Spaces of Sheaves, Second Edition, Cambridge University Press (2010).
- [27] Hoskins, V., Kirwan, F.: Quotients of unstable subvarieties and moduli spaces of sheaves of fixed Harder-Narasimhan type. Proc. Lond. Math. Soc. (3) 105 (2012), no. 4, 852–890.
- [28] Jackson, Joshua: Moduli spaces of unstables objects: Sheaves of Harder-Narasimhan lenght 2. arXiv:2111.07428v1 [Math.AG] (2021).
- [29] Kempf, G.R.: Instability in invariant theory, Ann. of Math. 108 (1978), 299–316.
- [30] S. Kosarew and C. Okonek, Global moduli spaces and simple holomorphic bundles, Publ. RIMS, Kyoto Univ. 25 (1989), 1–19.
- [31] Lange, H.: Universal Families of Extensions Journal of algebra 83 (1983), 101–112.
- [32] Maruyama, M.: Moduli of stable sheaves I, J. Math. Kyoto Univ. 17 (1977), 91–126.
- [33] Maruyama, M.: Moduli of stable sheaves II, J. Math. Kyoto Univ. 18 (1978), 577-614.
- [34] Mercat, V.: Le problème de Brill-Noether et le théorème de Teixidor. Manuscr. Math., 98(1) (1999), 75-85.
- [35] Mercat, V.: Le problème de Brill-Noether pour des fibrès stables de petite pente. J. Reine Angew. Math. 506 (1999), 1–41.
- [36] Mumford, D.: Projective invariants of projective structures and applications. Proc. Internat. Congr. Mathematicians (Stockholm, 1962) pp. 526–530 Inst. Mittag-Leffler, Djursholm.
- [37] Narasimhan, M.S., Ramanan, S.: Deformations of the moduli space of vector bundles over an algebraic curve. Ann. Math. (2) (1975), 101:391–417.
- [38] Narasimhan, M.S., Seshadri, C.S.: Stable and unitary vector bundles on a compact Riemann surface. Ann. of Math. 82 (1965), 540–567.
- [39] Ríos-Sierra R.: Moduli of unstable bundles of rank 3. In Preparation
- [40] Simpson, C.T.: Moduli of representations of the fundamental group of a smooth projective variety, Publ. Math. I.H.E.S. 79 (1994), 47–129.
- [41] Strømme, S.A.: Deforming vector bundles on the projective plane. Math. Ann. 263 (1983), no. 3, 385–397.
- [42] Teixidor i Bigas, M.: Brill–Noether theory for stable vector bundles. Duke Math. J., 62(2) (1991), 385–400.

CIMAT, APDO. POSTAL 402,C.P. 36240, GUANAJUATO, MEXICO Email address: lebp@cimat.mx

CIMAT, Apdo. Postal 402, C.P. 36240, Guanajuato, Mexico

Email address: rocio.rios@cimat.mx