

# A NOTE ABOUT RATIONAL SURFACES AS UNIONS OF AFFINE PLANES

JORGE CARAVANTES<sup>1</sup>, J. RAFAEL SENDRA<sup>1</sup>, DAVID SEVILLA<sup>2</sup> AND CARLOS VILLARINO<sup>1</sup>

ABSTRACT. We prove that any smooth rational projective surface over the field of complex numbers has an open covering consisting of 3 subsets isomorphic to affine planes.

Since all smooth rational curves are isomorphic to  $\mathbb{P}^1$ , they can be seen as the union of two affine lines. In dimension two, as a consequence of the structure Theorem 1.2 below, all rational surfaces admit a covering of open subsets isomorphic to the affine plane. However, up to the authors' knowledge, no general results are known on the minimal number of open subsets of such a covering, while some advances are known by computer algebraists in terms of surjectivity of parametrizations [BR95, SSV17, CSSV18, CSSV21]. In this short note we prove that all projective smooth rational surfaces behave like the projective plane in this aspect.

## 1. MAIN RESULT

**Theorem 1.1.** *Let  $X$  be a projective smooth rational surface over the complex field. Then, there are three open subsets  $U_0, U_1, U_2 \subset X$  such that:*

- (1)  $U_0 \cup U_1 \cup U_2 = X$ .
- (2) For all  $i = 0, 1, 2$ ,  $U_i$  is isomorphic to the affine plane.

To prove Theorem 1.1, we will use the following well-known result:

**Theorem 1.2.** *(see e.g. [Bea83, Theorem V.10]) Every non-singular rational surface can be obtained by repeatedly blowing up either  $\mathbb{P}^2$  or the projective bundle  $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-n))$  (the Hirzebruch surface  $\Sigma_n$ ), for  $n \neq 1$ .*

By Theorem 1.2, there exists a chain of morphisms  $\pi = \pi_1 \circ \dots \circ \pi_r : X \rightarrow M$  such that  $M$  is either  $\mathbb{P}^2$  or a Hirzebruch surface and  $\pi_i : X_i \rightarrow X_{i-1}$  is the blowup of a smooth surface at a single point. Let  $E$  be the exceptional divisor of  $\pi$  and  $E_i$  the exceptional divisor of  $\pi_i$ . Then,  $\pi(E) \subset M$  is a finite set of closed points and  $\pi_i(E_i)$  is one closed point. Moreover,  $E_i \simeq \mathbb{P}^1$  and  $E$  is a finite union of smooth rational curves (in fact,  $E_1$  and the proper transforms of all the  $E_2, \dots, E_r$ ). We begin by proving Theorem 1.1 for  $X = M$  with care for the centers of the blowups:

**Lemma 1.3.** *In the above conditions, there exist three open subsets  $U_0^0, U_1^0, U_2^0$  such that:*

- (1)  $U_0^0 \cup U_1^0 \cup U_2^0 = M$ .
- (2) For all  $i = 0, 1, 2$ ,  $U_i^0$  is isomorphic to the affine plane.
- (3)  $\pi(E) \subset U_0^0 \cap U_1^0 \cap U_2^0$ .

---

2010 *Mathematics Subject Classification.* Primary 14E05, 14M20; Secondary 14A10.

The authors are partially supported by the grant PID2020-113192GB-I00/AEI/ 10.13039/501100011033 (Mathematical Visualization: Foundations, Algorithms and Applications) from the Spanish State Research Agency (Ministerio de Ciencia e Innovación). J. Caravantes and C. Villarino belong to the Research Group ASYNACS (Ref. CT-CE2019/683). D. Sevilla is a member of the research group GADAC and is partially supported by Junta de Extremadura and Fondo Europeo de Desarrollo Regional (GR21055).

*Proof.* The case  $M = \mathbb{P}^2$  is well-known. Since  $\pi(E)$  is finite and we work over an infinite field, one can choose three different projective lines  $L_1, L_2$  and  $L_3$  in  $\mathbb{P}^2$  such that  $\pi(E) \cap (L_1 \cup L_2 \cup L_3) = \emptyset = L_1 \cap L_2 \cap L_3$ .

If  $M$  is a Hirzebruch surface, then it is the projective bundle of a rank two vector bundle  $\mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-m)$  over  $\mathbb{P}^1$ . This means that there is a surjective morphism  $p : M \rightarrow \mathbb{P}^1$  such that, for any point  $P \in \mathbb{P}^1$ ,  $p^{-1}(\mathbb{P}^1 - \{P\}) \simeq (\mathbb{P}^1 - \{P\}) \times \mathbb{P}^1$ . Then, since we work over an infinite field, one can choose a closed point  $P_0 \in \mathbb{P}^1 - p(\pi(E))$  with its isomorphism  $q_0 : p^{-1}(\mathbb{P}^1 - \{P_0\}) \rightarrow \mathbb{A}^1 \times \mathbb{P}^1$ . Then, we choose a line  $L_0 = \mathbb{A}^1 \times \{Q_0\}$ , such that  $q_0(\pi(E)) \cap L_0$  is empty. With this choice,  $U_0^0 = q_0^{-1}(\mathbb{A}^1 \times \mathbb{P}^1 - L_0)$  is isomorphic to  $\mathbb{A}^2$  and contains  $\pi(E)$ .

Then,  $M - U_0^0$  is the union of two rational curves  $C_1 := p^{-1}(P_0)$  and  $C_2 := \overline{q_0^{-1}(L_0)}$ . Choosing  $P_1 \in \mathbb{P}^1 - (p(\pi(E)) \cup \{P_0\})$  (again, the complement of a finite set), together with the isomorphism  $q_1 : p^{-1}(\mathbb{P}^1 - \{P_1\}) \rightarrow \mathbb{A}^1 \times \mathbb{P}^1$ , we have that  $p^{-1}(\mathbb{P}^1 - \{P_1\})$  contains  $C_1$  and  $C_2$  with the exception of the point  $R_1 := C_2 \cap p^{-1}(P_1)$  (the intersection of a section  $\mathbb{A}^1 \rightarrow \mathbb{A}^1 \times \mathbb{P}^1$  with a fiber). We now choose a line  $L_1 = \mathbb{A}^1 \times \{Q_1\}$  such that:

- $Q_1 \in \mathbb{P}^1$  is not in the second projection of  $q_1(\pi(E)) \in \mathbb{A}^1 \times \mathbb{P}^1$ ; and
- $L_1 \neq q_1(C_2)$  (i.e. we are asking a constant section not to coincide with a given one, which is an open condition for  $Q_1$ ), so the intersection of the two curves is finite.

Then,  $U_1^0 = q_1^{-1}(\mathbb{A}^1 \times \mathbb{P}^1 - L_1)$  is isomorphic to  $\mathbb{A}^2$  and contains  $\pi(E)$ .

Now,  $M - (U_0^0 \cup U_1^0) = (C_1 \cup C_2) - U_1^0$  is the finite set  $A := \{R_1\} \cup q_1^{-1}(L_1 \cap q_1(C_2))$ . Finally, we have again the complement of a finite set to choose  $P_2 \in \mathbb{P}^1 - (p(\pi(E)) \cup p(A) \cup \{P_0, P_1\})$  with the isomorphism  $q_2 : p^{-1}(\mathbb{P}^1 - \{P_2\}) \rightarrow \mathbb{A}^1 \times \mathbb{P}^1$ , so we have  $A \subset p^{-1}(\mathbb{P}^1 - \{P_2\})$ . We now choose  $L_2 = \mathbb{A}^1 \times \{Q_2\}$  such that  $Q_2 \in \mathbb{P}^1$  is not in the second projection of the finite set  $q_2(A \cup \pi(E)) \subset \mathbb{A}^1 \times \mathbb{P}^1$ , and we define  $U_2^0 = q_2^{-1}(\mathbb{A}^1 \times \mathbb{P}^1 - L_2) \simeq \mathbb{A}^2$ . Then  $\pi(E) \subset U_2^0$  and, since  $A \subset U_2^0$ , we have that  $U_0^0 \cup U_1^0 \cup U_2^0 = M$ .  $\square$

*Remark 1.4.* Let  $\text{Bl}_P(\mathbb{A}^2)$  be the blowup of the affine plane at a point  $P$ . Then, there exists one morphism  $\pi_l : U_l \simeq \mathbb{A}^2 \mapsto \mathbb{A}^2$  for each affine line  $l \subset \mathbb{A}^2$  passing through  $P$ , such that:

- (1) for  $l_1 \neq l_2$ ,  $U_{l_1} \cup U_{l_2} = \text{Bl}_P(\mathbb{A}^2)$ .
- (2) if  $E_{\mathbb{A}^2}$  is the exceptional divisor of  $\text{Bl}_P(\mathbb{A}^2)$ , for any line  $l$  passing through  $p$ ,  $E_{\mathbb{A}^2} - U_l$  consists in one point, given by the isomorphism between  $E_{\mathbb{A}^2}$  and the  $\mathbb{P}^1$  of all lines through  $P$ .
- (3) the restriction  $\pi_l|_{U_l - E_{\mathbb{A}^2}}$  is an isomorphism between  $U_l - E_{\mathbb{A}^2}$  and  $\mathbb{A}^2 - l$ .

**Lemma 1.5.** *Let  $X$  be a smooth rational surface such that there exist three open subsets  $U_0, U_1, U_2 \subset X$  with*

- (1)  $U_0 \cup U_1 \cup U_2 = X$ .
- (2) For all  $i = 0, 1, 2$ ,  $U_i$  is isomorphic to the affine plane.

*Consider a finite set  $A_1 \subset U_0 \cap U_1 \cap U_2$ . Let  $P \in (U_0 \cap U_1 \cap U_2) - A_1$  be a point and consider  $\pi : Y \rightarrow X$  to be the blowup of  $X$  at  $P$ . Consider also a finite set  $A_2$  in the exceptional divisor  $E = \pi^{-1}(P) \subset Y$ . Then, there are three open subsets  $U'_0, U'_1, U'_2 \subset Y$  such that*

- (1)  $U'_0 \cup U'_1 \cup U'_2 = Y$ .
- (2) For all  $i = 0, 1, 2$ ,  $U'_i$  is isomorphic to the affine plane.
- (3) Both  $A_2$  and the proper transform of  $A_1$  are contained in  $U'_0 \cap U'_1 \cap U'_2$

*Remark 1.6.* In the conditions of Lemma 1.5, note that for any  $i, j = 0, 1, 2$ ,  $i \neq j$ ,  $X - (U_i \cup U_j)$  is a Zariski closed subset of a projective surface which is contained in  $U_k \simeq \mathbb{A}^2$ , with  $i \neq k \neq j$ . Since it is a projective scheme in an affine space, it must be finite.

*Proof.* Taking into account Remark 1.4, consider a line  $l_0 \subset U_0 \simeq \mathbb{A}^2$  through  $P$  such that

- The intersection of  $l_0$  with the finite set  $X - (U_1 \cup U_2)$  (see Remark 1.6) is empty.
- $A_1 \cap l_0 = \emptyset$ .
- The intersection point of the proper transform of  $l_0$  with the exceptional divisor is not in  $A_2$ .

Then, we define  $U'_0$  to be the open subset  $U_{l_0}$  of the blowup of  $U_0$  at  $P$ .  $U_0$  is isomorphic to the affine plane, as said in Remark 1.4, and  $Y - U'_0$  consists in the proper transform of  $l_0 \cup (X - U_0)$ . Therefore, it is one-dimensional.

Now, we choose a line  $l_1 \subset U_1 \simeq \mathbb{A}^2$  such that the following open conditions are satisfied:

- (1) The intersection of  $l_1$  with the finite set  $X - (U_0 \cup U_2)$  (see Remark 1.6) is empty.
- (2) the intersection multiplicity of  $l_1$  and  $l_0$  at  $P$  is 1 (note that  $l_1$  is smooth at  $P$ , so we are asking that  $l_1$  is not the tangent line at  $P$  to  $l_0$ , when we see them in  $U_1$ ).
- (3)  $l_1$  does not contain any point in  $l_0 \cap (X - U_2)$  (note that  $l_0$  is irreducible and  $P \in l_0 \cap U_2$ , so such intersection is finite).
- (4)  $A_1 \cap l_1 = \emptyset$ .
- (5) The intersection point of the proper transform of  $l_1$  with the exceptional divisor is not in  $A_2$ .

Since we work over an infinite field, these conditions define a nonempty Zariski open subset to choose  $l_1$  from. Now, we define  $U'_1$  to be  $U_{l_1} \simeq \mathbb{A}^2$ . The whole exceptional divisor is in  $U'_0 \cup U'_1$ . Then,  $Y - (U'_0 \cup U'_1)$  is the proper transform of the finite sets  $B_1 = [l_0 \cup (X - U_0)] \cap l_1$  and  $B_2 = X - (U_0 \cup U_1)$ .

Note that  $P \notin B_1 \cup B_2 \subset U_2$ , so we choose a last line  $l_2 \subset U_2 \simeq \mathbb{A}^2$  such that

- the intersection of  $l_2$  with the finite set  $X - (U_0 \cup U_1)$  (see Remark 1.6) is empty,
- $(A_1 \cup B_1 \cup B_2) \cap l_2 = \emptyset$ , and
- the intersection point of the proper transform of  $l_2$  with the exceptional divisor is not in  $A_2$ .

Defining  $U'_2 = U_{l_2}$ , one concludes the proof. □

*Remark 1.7.* It is likely that a generalisation of Lemma 1.5 to higher dimension is possible. However, it is not yet known if all rational varieties of dimension greater than 2 are covered by open subsets isomorphic to open subsets of  $\mathbb{A}^n$  (see [Gro89] for the original question). These varieties are known as plain [BHSVU08] or uniformly rational [BB14] and it is possible that the main result can be extended to higher dimension for this type of varieties.

*Proof. (of Theorem 1.1)* By Lemma 1.3, we have that  $M = X_0 = U_0^0 \cup U_1^0 \cup U_2^0$  with  $U_i^0 \simeq \mathbb{A}^2$  and  $\pi(E) \subset U_0^0 \cap U_1^0 \cap U_2^0$ . Now we apply Lemma 1.5 to  $\pi_i : X_i \rightarrow X_{i-1}$ , choosing

$$A_1 = [\pi_i \circ \pi_{i+1}(E_{i+1}) \cup \dots \cup \pi_i \circ \dots \circ \pi_r(E_r)] - \{P_i\}$$

(i.e. the points to be the center of future blowups outside  $\{P_i\}$ ) and

$$A_2 = [\pi_{i+1}(E_{i+1}) \cup \dots \cup \pi_{i+1} \circ \dots \circ \pi_r(E_r)] \cap E_i$$

(i.e. the points to be center of future blowups in  $E_i$ ). Note that any curve contracted by  $\pi_{i+1} \circ \dots \circ \pi_i$  is contracted to a point in  $\pi_i^{-1}(A_1) \cup A_2$ . We then get  $U_0^i, U_1^i, U_2^i$  from  $U_0^{i-1}, U_1^{i-1}, U_2^{i-1}$  all isomorphic to  $\mathbb{A}^2$  and covering  $X_i$ , with all centers of future blowups in the intersection of the three open subsets. Then  $U_0^r, U_1^r$  and  $U_2^r$  are the three open subsets in the statement. □

## REFERENCES

- [BB14] Fedor Bogomolov and Christian Böhning. On uniformly rational varieties. In *Topology, geometry, integrable systems, and mathematical physics*, volume 234 of *Amer. Math. Soc. Transl. Ser. 2*, pages 33–48. Amer. Math. Soc., Providence, RI, 2014.
- [Bea83] Arnaud Beauville. *Complex algebraic surfaces*, volume 68 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 1983. Translated from the French by R. Barlow, N. I. Shepherd-Barron and M. Reid.
- [BHSVU08] Gábor Bodnár, Herwig Hauser, Josef Schicho, and Orlando Villamayor U. Plain varieties. *Bulletin of the London Mathematical Society*, 40(6):965–971, 11 2008.
- [BR95] Chandrajit L. Bajaj and Andrew V. Royappa. Finite representations of real parametric curves and surfaces. *Internat. J. Comput. Geom. Appl.*, 5(3):313–326, 1995.
- [CSSV18] Jorge Caravantes, J. Rafael Sendra, David Sevilla, and Carlos Villarino. On the existence of birational surjective parametrizations of affine surfaces. *J. Algebra*, 501:206–214, 2018.
- [CSSV21] Jorge Caravantes, J. Rafael Sendra, David Sevilla, and Carlos Villarino. Covering rational surfaces with rational parametrization images. *Mathematics*, 9(4), 2021.
- [Gro89] M. Gromov. Oka’s principle for holomorphic sections of elliptic bundles. *Journal of the American Mathematical Society*, 2(4):851–897, 1989.
- [SSV17] J. Rafael Sendra, David Sevilla, and Carlos Villarino. Covering rational ruled surfaces. *Math. Comput.*, 86(308):2861–2875, 2017.

<sup>1</sup> UNIVERSIDAD DE ALCALÁ, DPTO. DE FÍSICA Y MATEMÁTICAS, 28871 ALCALÁ DE HENARES (MADRID), SPAIN,

<sup>2</sup> CENTRO U. DE MÉRIDA, UNIVERSIDAD DE EXTREMADURA, AV. SANTA TERESA DE JORNET 38, 06800 MÉRIDA (BADAJOZ), SPAIN