

ON BRUHAT-TITS THEORY OVER A HIGHER DIMENSIONAL BASE

VIKRAMAN BALAJI AND YASHONIDHI PANDEY

To our parents and Apurv Pandey

ABSTRACT. Let k be a perfect field. Assume that the characteristic of k satisfies certain tameness assumptions (1.1). Let $\mathcal{O}_n := k[z_1, \dots, z_n]$ and set $K_n := \text{Fract } \mathcal{O}_n$. Let G be an almost-simple, simply-connected affine Chevalley group scheme with a maximal torus T and a Borel subgroup B . Given a n -tuple $\mathbf{f} = (f_1, \dots, f_n)$ of concave functions on the root system of G as in Bruhat-Tits [BT1], [BT2], we define n -bounded subgroups $P_{\mathbf{f}} \subset G(K_n)$ as a direct generalization of Bruhat-Tits groups for the case $n = 1$. We show that these groups are *schematic*, i.e. they are valued points of smooth *quasi-affine* (resp. *affine*) group schemes with connected fibres and *adapted to the divisor with normal crossing* $z_1 \cdots z_n = 0$ in the sense that the restriction to the generic point of the divisor $z_i = 0$ is given by f_i (resp. sums of concave functions given by points of the apartment). This provides a higher-dimensional analogue of the Bruhat-Tits group schemes with natural specialization properties. In §9, under suitable assumptions on k §2.2, we extend all these results for a $n + 1$ -tuple $\mathbf{f} = (f_0, \dots, f_n)$ of concave functions on the root system of G replacing \mathcal{O}_n by $\mathcal{O}[x_1, \dots, x_n]$ where \mathcal{O} is a complete discrete valuation ring with a perfect residue field k of characteristic p . In particular, if x_0 is the uniformizer of \mathcal{O} , then the group scheme is adapted to the divisor $x_0 \cdots x_n = 0$. In the last part of the paper, we give applications in char zero to constructing certain natural group schemes on wonderful embeddings of groups and also certain families of 2-parahoric group schemes on minimal resolutions of surface singularities that arose in [Bal].

CONTENTS

1. Introduction	2
Part 1. THE BOUNDED GROUPS	9
2. Towards the n -parahoric Lie algebra bundle	9
3. Examples and Remarks	14
Part 2. SCHEMATIZATION	24
4. The n -parahoric Lie-algebra bundle \mathcal{R} on an affine space	24
5. Schematization of n -parahoric groups	27
6. Schematization of n -bounded groups P_{Ω}	35
7. Schematization of n -bounded groups $P_{\mathbf{f}}$	44
8. Schematization of n -Moy-Prasad groups	46
9. Schematization in the mixed characteristic case	50
Part 3. SOME APPLICATIONS IN $\text{char}(k) = 0$ EQUICARACTERISTIC CASE	53
10. BT-group schemes on wonderful embeddings in $\text{char}(k) = 0$	53

2000 *Mathematics Subject Classification.* 14L15, 14M27, 14D20.

Key words and phrases. Bruhat-Tits group scheme, parahoric groups, concave functions, wonderful compactifications, degenerations.

11. 2BT group schemes and degenerations of torsors	58
12. Appendix on parabolic and equivariant bundles	62
References	64

1. INTRODUCTION

Let G be an almost-simple, simply-connected affine Chevalley group scheme. Let \mathcal{O} be a complete discrete valuation ring with a perfect residue field k . Let $\mathcal{O}_n := k[[z_1, \dots, z_n]]$ and let $K_n := \text{Fract } \mathcal{O}_n$, or more generally, we consider $\mathcal{O}[[x_1, \dots, x_n]]$, when the residue characteristic p satisfies some tameness assumptions (1.1). The object of this paper is to define and study an optimal class of subgroups $P \subset G(K_n)$ which we term as *n-bounded subgroups*. These are natural generalisations of the classical bounded groups and parahoric groups due to Bruhat and Tits who dealt with the case when $n = 1$ ([BT2], [BT1]).

We show that these n -bounded groups are *schematic* in the sense of Bruhat and Tits, i.e., there exist smooth, finite type group schemes over $\text{Spec } \mathcal{O}_n$ with connected fibres, such that the group of sections gives back the n -bounded groups. We will call these group schemes *nBT-group schemes* (1.4). The *nBT-group schemes* we construct are always *quasi-affine*, with a large, and perhaps most significant class, being *affine* (1.3). This provides natural analogues of the Bruhat-Tits group schemes over \mathcal{O}_n with good specialization properties.

In [BT2, 3.9.4], one finds general remarks on how one could possibly generalize [BT2, Theorem 3.8.1] to a base of dimension 2. Somewhat later in the mid-nineties, there was an approach to Bruhat-Tits buildings over higher dimensional local fields made by A. Parshin [Par] with possible motivation from questions in arithmetic. Our approach has its origins in geometry stemming from the study of degenerations of the stack Bun_G of G -bundles over smooth projective curves [Bal]. Unlike Parshin, our aim is not towards generalization of the notion of a Bruhat-Tits building. Rather, we follow the general spirit of the Bruhat-Tits approach [BT2], namely, to define subgroups of $G(K_n)$ characterisable by group schemes over $\text{Spec } \mathcal{O}_n$.

Recall that a *parabolic vector bundle* in the sense of C.S. Seshadri is a vector bundle with additional data defined on pairs (X, D) , where X is *smooth quasi-projective scheme* and D a *simple normal crossing divisor* (see §12). Hitherto, these have been *objects* of study primarily occurring as points in certain moduli spaces. In the present paper, the notion of a parabolic bundle helps us get a technical tool to address the issue of extending affine group schemes across subsets of codimension bigger than 1. These connections came up first in the paper [BS], where under the assumptions of characteristic 0 and for group schemes which are generically split (1.1), it was shown [BS, Theorem 5.2.7], that *any parahoric Bruhat-Tits group scheme over a complete discrete valuation ring \mathcal{O} can be realized "invariant direct images", i.e., obtained by taking Galois invariants of Weil restriction of scalars of a reductive group scheme, on a ramified cover of \mathcal{O} (see §12.2). It is easily checked (see §9) that this can be made to work under some mild tameness assumptions on residue field characteristics for arbitrary complete DVRs (see §1.1). This result and its underlying philosophy is the cornerstone of some of the basic results of this paper.*

Our first step is to define \mathfrak{n} -parahoric subgroups of $G(K_n)$. Let T be a maximal torus contained in a Borel B of G over k . Let $X(T) = \text{Hom}(T, \mathbb{G}_m)$ be the group of characters of T and $Y(T) = \text{Hom}(\mathbb{G}_m, T)$ be the group of all one-parameter subgroups of T . Let \mathfrak{g} denote the Lie-algebra. We denote $\Phi^+, \Phi^- \subset \Phi$ the set of positive and negative roots with respect to B . Let $S = \{\alpha_1, \dots, \alpha_\ell\}$ denote the set of simple roots of G , where ℓ is the rank of G . Let α^\vee denote the coroot corresponding to $\alpha \in S$. We also choose an *épinglage* or a *pinning* of G , i.e. in addition to S , we also choose for every $r \in S$, a k -isomorphism u_r of the additive group \mathbb{G}_a to the unipotent group U_r of G associated to the root r . Let $\{u_r\}_{r \in \Phi}$ be their canonical extensions satisfying the Chevalley relations.

Let $\mathbb{S} := S \cup \{\alpha_0\}$ denote the set of affine simple roots. Let \mathcal{A}_T denote the affine apartment corresponding to T . It can be identified with the affine space $\mathbb{E} := Y(T) \otimes_{\mathbb{Z}} \mathbb{R}$ together with its origin 0 .

For a point $\theta \in \mathcal{A}_T$ and a root $r \in \Phi$, let $r(\theta) := \langle r, \theta \rangle$. Set

$$m_r(\theta) := -\lfloor r(\theta) \rfloor. \quad (1.0.1)$$

Let \mathcal{O} be a complete discrete valuation ring with residue field k , with uniformizer z and field of fractions K . Classically ([BT1]), in terms of generators a parahoric subgroup P_θ of $G(K)$ is defined as:

$$P_\theta := \langle T(\mathcal{O}), U_r(z^{m_r(\theta)} \mathcal{O}), r \in \Phi \rangle. \quad (1.0.2)$$

More generally, for a *enclosed* bounded subset Ω in the affine apartment \mathcal{A}_T ([BT1, 2.4.6] or [Cou, §3]), set

$$m_r(\Omega) := -\lfloor \inf_{\theta \in \Omega} r(\theta) \rfloor = \lceil \sup_{\theta \in \Omega} -r(\theta) \rceil \quad (1.0.3)$$

Note that there is a notion of a *closure* of a bounded subset and the function $m_r(\Omega)$ does not change by taking closures. So unless otherwise mentioned, in this article we always work with enclosed bounded subsets. A bounded subgroup P_Ω of $G(K)$ associated to a bounded subset Ω of \mathcal{A}_T is defined as follows:

$$P_\Omega := \langle T(\mathcal{O}), U_r(z^{m_r(\Omega)} \mathcal{O}), r \in \Phi \rangle. \quad (1.0.4)$$

Following [BT1], [BT2], we define a larger class of bounded groups as follows:

DEFINITION 1.1. *A subgroup $Q \subset G(K)$ will be called schematic if there exists a smooth \mathcal{O} -group scheme \mathfrak{G} , with connected fibres and of finite type with $\mathfrak{G}_K \simeq G \times K$ such that $Q = \mathfrak{G}(\mathcal{O})$.*

*Throughout this paper we work under the assumption that all the \mathfrak{n} -Bruhat-Tits group schemes we study are connected and the generic fibre G_K is a product $G \times_{\text{Spec } K} \text{Spec } K$; we call this the “generically split case”. As we have mentioned above, parahoric subgroups are all *schematic*. However, Bruhat and Tits study a larger class of schematic subgroups of $G(K)$ which are defined by *concave functions* (in fact they work with the class of quasi-concave functions).*

Let $\tilde{\Phi} := \Phi \cup \{0\}$. Recall [BT1, 6.4.3, page 133], that a function $f : \tilde{\Phi} \rightarrow \mathbb{R}$ is said to be *concave* if whenever $r_i \in \tilde{\Phi}$ are such that $\sum_i r_i \in \tilde{\Phi}$, then

$$f\left(\sum_i r_i\right) \leq \sum_i f(r_i). \quad (1.0.5)$$

In this paper we generalise the theory for all concave functions over higher dimensional bases. We hasten to add that Bruhat-Tits theory works for general connected reductive groups, while we make rather simplifying assumptions on G in the spirit of

[BT2, §3.2 page 52] that it is almost simple, simply connected and *generically split*. Even under these assumptions, the problem seems sufficiently complex.

A natural class of 2-parahoric group schemes (1.4) came up naturally in the work of the senior author while constructing flat degenerations of the stack of G -torsors on smooth projective curves, when we allow the curve to degenerate to stable curves (cf. [Bal]). A special 2-parahoric group scheme comes up in the paper [PZ] (see 9.4 below and also [Lou]). We note that BT-group schemes on discrete valuations rings arising from general concave functions have appeared in the papers of J-K.Yu [Yu1], [Yu], and Reeder-Yu [RY]. Further, a very special class has found applications in the papers of Moy-Prasad (see [MP, 2.6]) and Schneider-Stuhler [SS].

Even in the most geometric contexts i.e. without the complications of rationality questions, the larger class of concave functions are needed to express [Bal] or the present paper (see (11.1) and (11.2)). Indeed, the description of the *closed fibres* of the n -parahoric group schemes (1.4) over \mathcal{O}_n , at points of depth bigger than 1, needs perforce the Bruhat-Tits theory of bounded groups associated to concave functions (1.5). The interesting feature is that, the concave functions which appear in this situation *do not in general arise from bounded subsets of the apartment*, but are from the most general class considered by F. Bruhat and J. Tits (see examples (3.10), (3.12), (3.11), (3.2.35) and Remark 11.2).

In view of [BT2, §3.2 second paragraph page 52 and §3.2.15], we believe that Bruhat and Tits may have already envisioned the theory over higher dimensional bases (cf. [BT2, §3.9.4, page 76]) as well. We revisit the latter in §5.9.

1.1. Assumptions on the residue field k . In this introduction, we assume that k is *algebraically closed*. We however note that, by assuming only perfectness of k together with a few assumptions on the existence of roots of unity (see §2.2) the results of the present paper continue to hold.

We will assume throughout that the characteristic p of k is coprime to the order of the center of G and the coefficients of the highest root.

- This assumption suffices when the concave function is of type I, with no further restrictions on the type of the group G .
- For groups of type A_n, G_2, F_4, E_6 , this suffices for concave functions of type II as well.
- Let h_G be the Coxeter number of G . Let $m(G)$ be the dimension of the minimal faithful representation of G . When the concave function is strictly of type III, we further need p to be greater than h_G as well as be coprime to $m(G)$. The second condition more precisely means the following: p is coprime to 7 for G_2 , coprime to 13 for F_4 , coprime to 3 for E_6 , coprime to 14 for E_7 , coprime to 31 for E_8 and finally coprime to $2n + 1$ for other classical groups of rank n .
- For the schematization of n -Moy-Prasad groups we refer to §8.1.

1.2. Higher dimensions and statement of main results. We begin with a few key definitions in terms of which the present paper is organised.

Types of Concave functions.

DEFINITION 1.2. (1) For $j = 1, \dots, n$, let points θ_j in the apartment \mathcal{A}_T be given. Let $\mathbf{f}_\theta = \{f_{\theta_j}\}_{j=1}^n$ be the set of concave functions associated to them by the function $r \mapsto m_r(\theta)$ (1.0.1). These will be called n -concave functions of type I.

- (2) For $j = 1, \dots, n$, let bounded subsets $\Omega_j \subset \mathcal{A}_r$ be given. Let $\mathbf{f}_\Omega = \{f_{\Omega_j}\}_{j=1}^n$ be the set of concave functions associated to them by $r \mapsto m_r(\Omega)$ (1.0.3). These will be called \mathbf{n} -concave functions of type II.
- (3) Let $\mathbf{f} := (\dots, f_j, \dots) : \Phi \rightarrow \mathbb{R}^n$ be a \mathbf{n} -concave function on Φ defined by concave functions $\{f_j\}$ (1.0.5). These will be called \mathbf{n} -concave functions of type III.

As mentioned earlier, there are examples of concave functions of type III which are not of type II (this is stated in [BT1, 6.4.4, page 133], however see 3.11 below for explicit examples) and similarly of type II not of type I.

Bounded groups in higher dimensions. We begin by stating our results in the equicharacteristic case. Recall $\mathcal{O}_n := k\llbracket z_1, \dots, z_n \rrbracket$ and let $K_n := \text{Fract } \mathcal{O}_n$. For each set of n rational points $\boldsymbol{\theta} = (\theta_1, \dots, \theta_n) \in \mathcal{A}^n$, in terms of generators we define a \mathbf{n} -parahoric subgroup of $G(K_n)$ as

$$P_{\boldsymbol{\theta}} := \langle T(\mathcal{O}_n), U_r \left(\prod_{1 \leq i \leq n} z_i^{m_r(\theta_i)} \mathcal{O}_n \right), r \in \Phi \rangle. \quad (1.2.1)$$

More generally, for $\Omega = (\Omega_1, \dots, \Omega_n)$ where $\Omega_i \subset \mathbb{E}$ are *enclosed* bounded subsets ([BT1, 2.4.6]), we can analogously define \mathbf{n} -bounded subgroups of $G(K_n)$:

$$P_{\Omega} := \langle T(\mathcal{O}_n), U_r \left(\prod_{1 \leq i \leq n} z_i^{m_r(\Omega_i)} \mathcal{O}_n \right), r \in \Phi \rangle. \quad (1.2.2)$$

Even more generally, let $f_i : \Phi \rightarrow \mathbb{R}$ be concave functions (1.0.5), and let $\mathbf{f} := (\dots, f_i, \dots) : \Phi \rightarrow \mathbb{R}^n$ be the \mathbf{n} -concave function defined by them. To the \mathbf{n} -concave function \mathbf{f} we can associate a \mathbf{n} -bounded subgroup of $G(K_n)$ as follows:

$$P_{\mathbf{f}} := \langle T(\mathcal{O}_n), U_r \left(\prod_{1 \leq i \leq n} z_i^{f_i(r)} \mathcal{O}_n \right), r \in \Phi \rangle. \quad (1.2.3)$$

Similarly, one may define the \mathbf{n} -bounded Lie subalgebra $\text{Lie}(P_{\mathbf{f}})$ of $\mathfrak{g}(K_n)$.

As in (1.1), we can call a subgroup $Q \subset G(K_n)$ a *schematic* subgroup if there exists a connected, smooth \mathcal{O}_n -group scheme \mathfrak{G} of finite type with $\mathfrak{G}_{K_n} \simeq G \times_{\text{Spec } \mathbb{Z}} K_n$ such that $Q = \mathfrak{G}(\mathcal{O}_n)$.

We remark that an important aspect of Bruhat-Tits group schemes over discrete valuation rings is the structure of a *big cell* which extends the big cell of the generic fibre (cf. [BT2, page 51, 3.1.3] and (1.3) (5) below). This aspect also generalizes to the \mathbf{n} -Bruhat-Tits group schemes.

We now state our main theorems in the equicharacteristic case. This corresponds to the layout of the paper.

THEOREM 1.3. *Let p satisfy the hypothesis of §(1.1) (or more generally §2.2). For a \mathbf{n} -concave function \mathbf{f} of each type in (1.2), there exists a smooth group scheme $\mathfrak{G}_{\mathbf{f}}$ of finite type on \mathbb{A}_k^n with connected fibres, whose restriction to $\text{Spec } \mathcal{O}_n$ comes with an isomorphism h_{K_n} of the generic fibre \mathfrak{G}_{K_n} with G_{K_n} and such that the following hold:*

- (1) *The group scheme $\mathfrak{G}_{\mathbf{f}}$ is affine when \mathbf{f} is of type I and in general it is quasi-affine in types II and III.*
- (2) *The \mathbf{n} -bounded subgroup $P_{\mathbf{f}}$ is schematic. More precisely, we have an identification $\mathfrak{G}_{\mathbf{f}}(\mathcal{O}_n) = P_{\mathbf{f}}$.*
- (3) *We have a natural isomorphism $\text{Lie}(\mathfrak{G}_{\mathbf{f}})(\mathcal{O}_n)$ with $\text{Lie}(P_{\mathbf{f}})$.*
- (4) *Let us denote the canonical images under h_{K_n} of the maximal torus T_{K_n} and the root groups $\{U_{r, K_n}\}_{r \in \Phi}$ in the generic fibre \mathfrak{G}_{K_n} by the same notations.*

- There exists closed subgroup schemes $\{\mathfrak{U}_{r,\mathbf{f}} \subset \mathfrak{G}_{\mathbf{f}}\}_{r \in \Phi}$ (resp. $\mathcal{T}_{\mathbf{f}} \subset \mathfrak{G}_{\mathbf{f}}$) on $\text{Spec } \mathcal{O}_n$, with generic fibre isomorphic to the root groups $\{U_{r,K_n}\}_{r \in \Phi}$ (resp. T_{K_n}).
- (5) There exists a big cell \mathfrak{B}_{Φ} of $\mathfrak{G}_{\mathbf{f}}$, such that for any choice of total ordering on Φ^+ (resp. Φ^-), the morphism induced by multiplication

$$\left(\prod_{r \in \Phi^+} \mathfrak{U}_{r,\mathbf{f}} \right) \times \mathcal{T}_{\mathbf{f}} \times \left(\prod_{r \in \Phi^-} \mathfrak{U}_{r,\mathbf{f}} \right) \rightarrow \mathfrak{G}_{\mathbf{f}} \quad (1.2.4)$$

is an open immersion with image \mathfrak{B}_{Φ} . Moreover, \mathfrak{B}_{Φ} restricts to the big cell \mathfrak{B}_{K_n} of the generic fibre \mathfrak{G}_{K_n} over $\text{Spec}(K_n)$.

- (6) More generally, on a base which is a smooth quasi-projective variety with a divisor with simple normal crossings, given a data of type I concave functions at the height 1 primes associated to the components of the divisor, in (5.4), it is shown that there exists a smooth affine group scheme with connected fibres on X together with a big cell structure which interpolates this datum on the divisors.
- (7) Let \mathbf{f} be a special n -concave function of type III such that each term f_i is a sum of concave functions of type I. Then the associated group scheme $\mathfrak{G}_{\mathbf{f}}$ is in fact affine (5.9).

The pair $(\mathfrak{G}_{\mathbf{f}}, h_{K_n})$ is uniquely determined upto a unique isomorphism, by the properties (4) and (5).

DEFINITION 1.4. The group scheme $\mathfrak{G}_{\mathbf{f}}$ is called the n BT-group scheme associated to a n -concave function \mathbf{f} . An n -parahoric group scheme is an n BT-group scheme associated to an n -concave function of type I.

THEOREM 1.5. The group scheme $\mathfrak{G}_{\mathbf{f}}$ associated to the n -concave function \mathbf{f} has the following structure at points of depth bigger than 1.

- (a) Let $I \subset \{1, \dots, n\}$ be a non-empty subset. For a general point $\xi \in \cap_{i \in I} H_i$, where the H_i are the coordinate hyperplanes in $\text{Spec } \mathcal{O}_n$, the fibre $\mathfrak{G}_{\xi, \xi}$ is isomorphic to the closed fibre of the Bruhat-Tits group scheme $\mathfrak{G}_{\mathbf{f}_I}$ on $\text{Spec } \mathcal{O}$ associated to the 1-concave function $\mathbf{f}_I : \Phi \rightarrow \mathbb{R}$, given by $r \mapsto \sum_{i \in I} f_i$.
- (b) More precisely, let $\mathbb{A}_I \subset \mathbb{A}^n$ be the “sub-diagonal” obtained by setting the coordinates $x_i = t$ for $i \in I$. Let $\text{Spec } A_I$ be the local ring at the generic point of the divisor $t = 0$ in \mathbb{A}_I . Then the restriction of $\mathfrak{G}_{\mathbf{f}}$ to $\text{Spec } A_I$ is isomorphic to the Bruhat-Tits group scheme $\mathfrak{G}_{\mathbf{f}_I}$ associated to the concave function \mathbf{f}_I .

Let \mathcal{O} be a complete dvr in mixed characteristics. Extending the methods for group schemes over \mathbb{A}_k^n to the case $\mathbb{A}_{\mathcal{O}}^n$, in §9 we prove the following:

THEOREM 1.6. For $n+1$ -concave function $\mathbf{f} = (f_0, \dots, f_n)$, where f_0 is prescribed at the uniformizer of \mathcal{O} , the main theorems THEOREM 1.3 and THEOREM 1.5 hold over $\mathbb{A}_{\mathcal{O}} = \mathbb{A}_{\mathcal{O}}^n$ under the tameness assumption (1.1) (in fact under assumptions in §2.2).

1.3. An outline of the strategy of proof. We now outline briefly the strategy of the proof in this paper. The general strategy is an *induction on the order of complexity* of concave functions on root systems. The simplest one is a function f_{θ} (1.0.1) which is a concave function of type I; the next in the order is the concave function f_{Ω} (1.0.3) associated to a bounded subset Ω in the apartment, i.e. of type II. The third in the order of complexity is the general concave function f of type III which need not arise as an f_{Ω} .

In other words, we build the family of nBT-group schemes $\{\mathcal{G}_r\}$ on $\text{Spec } \mathcal{O}_n$ associated to n-concave function $\mathbf{f}: \Phi \rightarrow \mathbb{R}^n$ in three tiers. This is carried out by a bootstrapping process. The first step §5 is to build the n-parahoric group scheme (1.4) associated to a collection of n points in the affine apartment i.e. a n-concave function $\mathbf{f} = \{f_{\theta_j}\}$ of type I. This case falls into a general method of proof where the ideas arising from the notion of parabolic vector bundles plays a key role. It has its origins in the paper [BS] and the end result is somewhat surprising in that, unlike what Bruhat-Tits expect, namely only a *quasi-affine* group scheme (see [BT2, 3.9.4] and §5.9 below), one in fact gets an *affine* n-parahoric group scheme on $\text{Spec } \mathcal{O}_n$ along with its cell-structure. The pair of the n-parahoric group scheme and its cell-structure become the base for the subsequent constructions. We then naturally move on to the case when instead of points in the apartment, we are given a collection of n bounded subsets in the affine apartment and so work with a n -tuple of *concave* functions arising from these bounded subsets. The construction (in §6) of the corresponding nBT-group scheme on $\text{Spec } \mathcal{O}_n$ builds on the construction of n-parahoric group scheme. We assume in §6 that G is one of A_n, G_2, F_4 , or E_6 where results from [BT3] and [GY1], [GY2] play the important role of reducing schematic constructions over DVRs for enclosed bounded subsets to their extremal vertices. The proof here is somewhat involved, especially in extending the cell structure and the schematic group law across subsets of codimension two. The end result is that, much as expected, we get a *quasi-affine* nBT-group scheme associated to the n-concave functions of type II. Finally, in §7, we turn our attention to the case of G with no constraints, and also a general n-concave function. This case is then reduced to the previous case, i.e. of n-concave functions of type II. This reduction follows from the *not so obvious* observation (3.21), that an *optimal concave function* on groups of type A_n always arises as one coming from a bounded subset of the apartment. This remark together with the power of the main results of [BT2] helps us to realise the nBT-group scheme associated to an n -tuple of concave functions; these group schemes are *a fortiori* quasi-affine as expected. A surprise and a possible opening, is offered by a special class of concave functions of type III, which occur as sums of concave functions of type I. Here, our methods yield the *affineness* of the group scheme (5.9). This leads us to a natural question namely: “Characterise n -tuple of optimal concave functions for which the associated nBT-group schemes are *affine*”.

Recall that in the equicharacteristic case when the residue field characteristic satisfies mild tameness assumptions, by [BS, Theorem 5.2.7] all parahoric subgroup schemes are recovered by the invariant direct image functor (i.e., obtained by taking Galois invariants of Weil restriction of scalars). Following [BS] closely, in §8 we show that Moy-Prasad group schemes over DVRs which generalize parahorics can be obtained by invariant direct image together with the simplest types of dilatations. Over higher dimensions, this generalizes suitably using dilatations results of [MRR]. We use results of Bruhat-Tits [BT2] to show in §9 that the generalization of [BS, Theorem 5.2.7] to mixed characteristics reduces to a verification on the big cell which was done in §8 following [BS] closely.

1.4. Other related results and proof strategies. In [L], Bruhat-Tits group schemes are constructed by using extension of *birational group laws* and the theorem of Artin-Weil [SGAD, Expose XVIII]. The possibility of such an approach is briefly mentioned in [BT2, §3.1.7]. In the very recent paper [Lou, Theorem 3.2.5], smooth and separated

group schemes over arbitrary Noetherian bases has been constructed. This last result assumes that the *schematic root datum* extends to the base ring A . However, in general proving the key properties such as *quasi-affineness* of the group schemes does not seem immediate by these approaches. In the present paper, for $k[z_1, \dots, z_n]$ or $\mathcal{O}[[z_1, \dots, z_n]]$ where \mathcal{O} is a discrete valuation ring, in the course of our proofs for each of the three types of concave functions, we actually show that the schematic root datum extends.

In the context of the present paper on higher dimensional bases, the first construction goes back to [BT2, 3.9.4] where for $A = k[x, y]$ or $A = A_1[[z]]$, for A_1 a Dedekind ring, a 2BT-group scheme is constructed by generalising [BT2, 3.8.1].

In the special setting of a 2-concave function $\mathbf{f} = (0, f)$ over $A = A_1[[z]]$, for A_1 a Dedekind ring, we refer to [PZ] and 9.4 below. The very recent paper [Lou] also gives a completely different proof of the key *affineness result* of these group schemes over $A = A_1[[z]]$; further, it goes on to construct and study affine Grassmannians for these group schemes.

1.5. Other directions. In a slightly different direction, recall that Moy-Prasad groups (see [MP, 2.6]) can be realized as schematic subgroups arising from BT-group schemes arising from a small variant of the concave functions in [BT1]. As a natural extension of our approach, in §8 we define n -Moy-Prasad groups and the corresponding schematization. Their schematisation over discrete valuation rings was derived as a consequence of a general approach by J.K.Yu in [Yu]. To adapt these groups to our theory we need to realize the Moy-Prasad groups as *invariant direct images* (5.2) of certain group schemes from ramified covers [BS] (see §8). This involves generalising the notion of an *unit group* to one corresponding to the variant of concave function mentioned above. As a consequence, we show that n -Moy-Prasad group schemes are affine just like the group scheme $\mathcal{G}_{\mathbf{f}}$ when \mathbf{f} is of type I.

As further applications of this approach, we construct higher BT *group schemes* on certain basic spaces. More precisely, we construct and describe

- an ℓ BT-group scheme on the De Concini-Procesi wonderful compactification [DP] \mathbf{X} of the adjoint group G_{ad} , where $\ell = \text{rank}(G)$,
- an $(\ell + 1)$ BT-group scheme on the loop “wonderful embedding” \mathbf{X}^{aff} of the adjoint affine Kac-Moody group $G_{\text{ad}}^{\text{aff}}$, constructed by Solis [So]. The novelty is that the base here is an *ind-scheme*.
- A family of 2BT-group schemes on the minimal resolution of singularities of normal surface singularities in the context of [Bal].

Acknowledgments. This paper had its origins in math arXiv [2101.09212] which however had an inaccuracy in the description of the group scheme at points of codimension higher than one. This was pointed to us by J. Heinloth. We thank him for this. The contents of the former manuscript is now subsumed in §10 of this paper. We also thank M. Brion and Patrick Polo for their numerous questions and comments on the paper. Finally, we thank the anonymous referee for his comments and questions which have led to a great improvement of the manuscript.

FREQUENTLY OCCURRING NOTATIONS AND CONVENTIONS

Throughout this paper we follow the classical notations and conventions as in the foundational papers [BT2] and [BT1] of Bruhat-Tits, and, that of the

treatise “Neron Models” [BLR] by S. Bosch, W. Lutkebohmert and M. Raynaud.

Φ	Root system relative to T
\mathcal{A}	the affine apartment of T
$\mathbf{f} = (f_1, \dots, f_n)$	n -concave-function on $\Phi \cup \{0\}$ (1.2)
$\mathbf{\Omega} = (\Omega_1, \dots, \Omega_n)$	n -tuple of bounded subsets of \mathcal{A}
$\mathbf{\theta} = (\theta_1, \dots, \theta_n)$	n -tuple of points in the affine apartment
\mathbf{A}	The affine space \mathbb{A}_k^n upto §8 or \mathbb{A}_σ^n as in §9.2.
\mathbf{A}_0	The complement of the coordinate hyperplanes in \mathbf{A}
(X, D)	Smooth scheme with a s.n.c divisor
X', U	Open subschemes of X whose codimension is ≥ 2 . Also called big open subsets.
$P_{\mathbf{f}}$ (1.2.3)	n -bounded groups associated to a n -concave function \mathbf{f} (1.2)
$P_{\mathbf{\Omega}}$ (1.2.2)	n -bounded groups associated to a n -bounded subset $\mathbf{\Omega}$
$P_{\mathbf{f}}^{\text{diag}}$ (3.5) and (3.1.2)	the specialisation to the diagonal
$\mathfrak{G}_{\mathbf{f}}$ (resp. $\mathfrak{G}_{\mathbf{\Omega}}, \mathfrak{G}_{\mathbf{\theta}}$)	n -Bruhat-Tits group scheme associated to an n -concave function \mathbf{f} (resp. n -bounded subset $\mathbf{\Omega}$, n -points $\mathbf{\theta}$)
Γ	Galois group of a Kawamata cover
p_*^Γ (5.2)	the <i>invariant direct image</i> i.e., Weil restrictions and Γ -invariants
\mathcal{R}	a Lie algebra bundle

Part 1. THE BOUNDED GROUPS

2. TOWARDS THE n -PARAHORIC LIE ALGEBRA BUNDLE

The first two subsections are in vigour throughout this article. The remaining ones recall a “loop” approach to parahoric groups for application in §4 and §10.

2.1. Lie-data of G/k . In this paragraph, we assume the group-theoretical notions of G already introduced in the introduction and give only those that we need further in this article. Let \mathfrak{a}_0 be the unique Weyl alcove of G whose closure contains 0 and which is contained in the dominant Weyl chamber corresponding to B . Under the natural pairing between $Y(T) \otimes_{\mathbb{Z}} \mathbb{Q}$ and $X(T) \otimes_{\mathbb{Z}} \mathbb{Q}$, the integral basis elements dual to S are called the fundamental co-weights $\{\omega_{\alpha}^{\vee} | \alpha \in S\}$. Let c_{α} be the coefficient of α in the highest root. The vertices of the Weyl alcove \mathfrak{a}_0 are exactly 0 and

$$\theta_{\alpha} := \frac{\omega_{\alpha}^{\vee}}{c_{\alpha}}, \alpha \in S. \quad (2.1.1)$$

We will call the lattice generated by the θ_{α} as the Tits lattice.

Let $\mathrm{SL}(m(G))$ be the minimal faithful representation of G . We fix a maximal split torus T_{SL} of $\mathrm{SL}(m(G))$. Let $\mathcal{A}_{T_{\mathrm{SL}}}$ denote its apartment. We fix an alcove \mathfrak{a} in $\mathcal{A}_{T_{\mathrm{SL}}}$.

DEFINITION 2.1. *For any $\theta \in Y(T) \otimes \mathbb{Q}$ let d_{θ} be the least positive integer such that $d_{\theta} \cdot \theta$ lies in $Y(T)$. For $\theta \in Y(T_{\mathrm{SL}}) \otimes \mathbb{Q}$, let d_{θ} be defined similarly.*

Thus, if e_{α} is the order of ω_{α}^{\vee} in the quotient of the co-weight lattice by $Y(T)$, it follows that for θ_{α} , the number $d_{\alpha} := d_{\theta_{\alpha}}$ is:

$$d_{\alpha} = e_{\alpha} \cdot c_{\alpha}. \quad (2.1.2)$$

An affine simple root $\alpha \in S$ may be viewed as an affine functional on \mathcal{A}_T . Any non-empty subset $\mathbb{I} \subset S$ defines the facet $\Sigma_{\mathbb{I}} \subset \overline{\mathfrak{a}_0}$ where exactly the α 's not lying in the subset \mathbb{I} vanish. So the set S corresponds to the interior of the alcove and the vertex θ_{α} of the alcove corresponds to $\alpha \in S$. Conversely any facet $\Sigma \subset \overline{\mathfrak{a}_0}$ defines non-empty subset $\mathbb{I}_{\Sigma} \subset S$. For $\emptyset \neq \mathbb{I} \subset S$, the barycenter of $\Sigma_{\mathbb{I}}$ is given by

$$\theta_{\mathbb{I}} := \frac{1}{|\mathbb{I}|} \sum_{\alpha \in \mathbb{I}} \theta_{\alpha}. \quad (2.1.3)$$

2.2. Assumptions on the residue field k and points in the apartments compatible with k . The following assumptions will be in vigour throughout the article. We assume k is perfect to be able to apply [BT2]. Let $p := \mathrm{char}(k)$.

Characteristic Assumptions: These are needed for the existence of a Kawamata cover (see §12.1). We will assume that p is coprime to d_{α} (2.1.2). Notice that since G is simply-connected this assumption is implied by §1.1. Indeed, since $Y(T)$ equals the coroot lattice, Z_G identifies with the quotient of the coweight lattice by $Y(T)$. Further, whenever we work with a finite set of rational points $\{\theta_1, \dots, \theta_n\}$ of \mathcal{A}_T , or $\mathcal{A}_{T_{\mathrm{SL}}}$ we assume that for $1 \leq i \leq n$, p coprime to the integers d_{θ_i} (2.1). Let h_G denote the Coxeter number of G . Let $m(G)$ denote the dimension of the minimal faithful representation of G . When the concave function is strictly of type III, we further need $p \geq h_G$ as well as p be coprime to $m(G)$.

Assumptions on the existence of primitive roots of unity: These are needed to make §12.2 work. The residue field k is such that for every facet Σ of \mathfrak{a}_0 (see §2.1) there exists a rational point $\theta \in \Sigma$ such that d_{θ} (2.1) is coprime to $\mathrm{char}(k)$ and k contains the primitive d_{θ} th roots of unity.

- For a general G , this suffices when the concave function is of type I.
- For A_n, G_2, F_4, E_6 , this suffices for type II as well.
- when the concave function is strictly of type III, in addition to the condition for \mathfrak{a}_0 , we need to assume the analogous condition for \mathfrak{a} (see §2.1).

Whenever we work with a finite set of rational points $\{\theta_1, \dots, \theta_n\}$ of \mathcal{A}_T , or $\mathcal{A}_{T_{\text{SL}}}$, we assume that for $1 \leq i \leq n$, k contains the primitive d_{θ_i} th roots of unity.

We observe any facet $\Sigma \subset \mathcal{A}_T$ of dimension contains a rational one parameter subgroup θ of T which is a convex combination of the vertices of Σ in a way that for $d \in \mathbb{N}$ coprime to p , the multiple $d\theta$ becomes a one parameter subgroup of T . We see this as follows. Since the alcove \mathbf{a}_0 is a fundamental domain for the action of the affine Weyl group we may suppose that Σ is a facet $\Sigma_{\mathbb{I}}$ of \mathbf{a}_0 . Say $\mathbb{I} = \{\theta_0, \dots, \theta_j\}$ are the vertices of $\Sigma_{\mathbb{I}}$ and $\theta_0 = \theta_{\alpha}$. Then since d_{α} (2.1.2) are always strictly greater than one, it follows we may take

$$\theta = \frac{\theta_0 + \sum_{i=1}^j (d_{\alpha}^i - d_{\alpha}^{i-1})\theta_i}{d_{\alpha}^j}. \quad (2.2.1)$$

We remark that if L denotes the LCM of the coefficients of the highest root, ℓ is the rank of T , then for type I, it suffices that k has $(|Z_G|L)^{\ell}$ primitive roots of unity. For type III, k must further have primitive $m(G)^{m(G)}$ roots of unity.

2.3. Loop groups and their parahoric subgroups. *We have included this section where we introduce the language of loop groups to describe parahoric groups. We however add that this terminology is restricted to §4 and §10 of this paper.*

The loop group LG is the group functor which associates to a k -algebra R the group $G(R((z)))$; this is representable by an ind-scheme over k . Similarly the loop Lie algebra functor $L\mathfrak{g}$ is given by $L\mathfrak{g}(R) = \mathfrak{g}(R((z)))$. We can similarly define the positive loops (also called *jet groups* in the literature) $L^+(G)$ to be the subfunctor of LG defined by $L^+(G)(R) := G(R[[z]])$. The positive loop construction extends more generally to any group scheme $\mathcal{G} \rightarrow \text{Spec}(A)$ and any vector bundle $\mathfrak{F} \rightarrow \text{Spec}(A)$ whose sheaf of sections carries a Lie-bracket.

Let $L^{\times}G = \mathbb{G}_m \times LG$ where the *rotational torus* \mathbb{G}_m acts on LG by acting on the uniformizer via the *loop rotation action* as follows: $u \in \mathbb{G}_m(R)$ acts on $\gamma(z) \in LG(R) = G(R((z)))$ by $u\gamma(z)u^{-1} = \gamma(uz)$. A maximal torus of $L^{\times}G$ is $T^{\times} = \mathbb{G}_m \times T$. A $\eta \in \text{Hom}(\mathbb{G}_m, T^{\times}) \otimes_{\mathbb{Z}} \mathbb{Q}$ over $R(s)$ can be viewed as a rational 1-PS, i.e., a 1-PS $\mathbb{G}_m \rightarrow T^{\times}$ over $R(w)$ where $w^n = s$ for some $n \geq 1$. In this case for $\gamma(z) \in L^{\times}G(R)$ we will view $\eta(s)\gamma(z)\eta(s)^{-1}$ as an element in $L^{\times}G(R(w))$. Hence, by the condition

$$\lim_{s \rightarrow 0} \eta(s)\gamma(z)\eta(s)^{-1} \text{ exists in } L^{\times}G(R) \quad (2.3.1)$$

on $\gamma(z) \in L^{\times}G(R)$, we mean that there exists a $n \geq 1$ such that for $w^n = s$, we have

$$\eta(s)\gamma(z)\eta(s)^{-1} \in L^{\times}G(R[[w]]). \quad (2.3.2)$$

This is the underlying principle in [BS] in a more global setting. Let $\eta = (a, \theta)$ for a a strictly positive rational number and θ a rational 1-PS of T . Then, we have

$$\eta(s)\gamma(z)\eta(s)^{-1} = \theta(s)\gamma(s^a z)\theta(s)^{-1}. \quad (2.3.3)$$

We note further that for any $0 < d \in \mathbb{N}$, setting $\eta = (\frac{a}{d}, \frac{\theta}{d})$ we have

$$\eta(s)\gamma(z)\eta(s)^{-1} = \theta(s^{\frac{1}{d}})\gamma(s^{\frac{a}{d}}z)\theta(s^{\frac{1}{d}})^{-1}. \quad (2.3.4)$$

So for $\eta = \frac{1}{d}(1, \theta)$ observe that the statement " $\lim_{s \rightarrow 0} \eta(s)\gamma(z)\eta(s)^{-1}$ exists" is a condition which is equivalent to

$$\theta(s)\gamma(s)\theta(s)^{-1} \in L^{\times}G(R[[w]]). \quad (2.3.5)$$

In other words, the condition of existence of limits is independent of d , and we may further set $z = s$ in $\gamma(z)$. More generally, if $a > 0$ the condition for (a, θ) and $(1, \frac{\theta}{a})$ are equivalent. We may write this condition on $\gamma(z) \in L^\times G(R)$ or $LG(R)$ as

$$" \lim_{s \rightarrow 0} \theta(s) \gamma(s) \theta(s)^{-1} \text{ exists in } L^\times G(R) \text{ or } LG(R) \text{ if } \gamma(z) \in LG(R) " \quad (2.3.6)$$

For $r \in \Phi$, let $u_r : \mathbb{G}_a \rightarrow G$ denote the root subgroup. If for some $b \in \mathbb{Z}$ and $t(z) \in R[[z]]$ we take $\gamma(z) := u_r(z^b t(z)) \in Lu_r(R)$, and $\eta := (1, \theta)$, then

$$\eta(s) u_r(z^b t(z)) \eta(s)^{-1} = \theta(s) u_r((sz)^b t(sz)) \theta(s)^{-1} = u_r(s^{r(\theta)} (sz)^b t(sz)). \quad (2.3.7)$$

So for $\eta = (1, \theta)$ the condition that the limit exists is equivalent to

$$r(\theta) + b \geq 0 \iff [r(\theta) + b] = [r(\theta)] + b \geq 0 \iff b \geq -[r(\theta)]. \quad (2.3.8)$$

We note the independence of the above implications on the number d occurring in the equation $\eta := \frac{1}{d}(1, \theta)$.

Let $\pi_1 : T^\times \rightarrow \mathbb{G}_m$ be the first projection. For any rational 1-PS $\eta : \mathbb{G}_m \rightarrow T^\times$ we say η is positive if $\pi_1 \circ \eta > 0$, negative if $\pi_1 \circ \eta < 0$ and non-zero if it is either positive or negative. In this paper, we will only need to work with η which are positive.

Any non-zero $\eta = (a, \theta)$ defines the following positive-loop functors from the category of k -algebras to the category of groups and Lie-algebras:

$$P_\eta^\times(R) := \{\gamma \in L^\times G(R) \mid \lim_{s \rightarrow 0} \eta(s) \gamma(z) \eta(s)^{-1} \text{ exists in } L^\times G(R)\}, \quad (2.3.9)$$

$$\mathfrak{P}_\eta^\times(R) := \{h \in L^\times \mathfrak{g}(R) \mid \lim_{s \rightarrow 0} Ad(\eta(s))(h(z)) \text{ exists in } L^\times \mathfrak{g}(R)\}, \quad (2.3.10)$$

$$P_\eta(R) := \{\gamma \in LG(R) \mid \lim_{s \rightarrow 0} \eta(s) \gamma(z) \eta(s)^{-1} \text{ exists in } LG(R)\}, \quad (2.3.11)$$

$$\mathfrak{P}_\eta(R) := \{h \in L\mathfrak{g}(R) \mid \lim_{s \rightarrow 0} Ad(\eta(s))(h(z)) \text{ exists in } L\mathfrak{g}(R)\}, \quad (2.3.12)$$

$$\text{Thus,, } P_\eta := P_\eta^\times \cap (1 \times LG) \text{ and } \mathfrak{P}_\eta := \mathfrak{P}_\eta^\times \cap (0 \oplus L\mathfrak{g}). \quad (2.3.13)$$

A *parahoric* subgroup of $L^\times G$ (resp. LG) is a subgroup that is conjugate to P_η^\times (resp. P_η) for some η . In this paper, we will mostly be using only the case when $\eta = \frac{1}{d}(1, \theta)$. In this case, letting $L\mathfrak{g}(R) = \mathfrak{g}(R((s)))$ as in (2.3.6), we may reformulate (2.3.13) as

$$\mathfrak{P}_\eta(R) = \{h \in L\mathfrak{g}(R) \mid \lim_{s \rightarrow 0} Ad(\theta(s))(h(s)) \text{ exists in } L\mathfrak{g}(R)\}. \quad (2.3.14)$$

Let \mathfrak{t} denote the Cartan sub-algebra associated to T and let $\mathfrak{g}_r \subset \mathfrak{g}$ be the root subspace associated to U_r . Then in terms of generators the Lie-algebra functor associated to the group functor P_θ (1.0.2) is given by

$$\text{Lie}(P_\theta)(R) = \langle \mathfrak{t}(R[[z]]), \mathfrak{g}_r(z^{m_r(\theta)} R[[z]]), r \in \Phi \rangle. \quad (2.3.15)$$

By the conditions (2.3.9) and (2.3.13), using (2.3.8), we may express P_η in terms of generators as in (1.0.2) and its Lie-algebra functor $\text{Lie}(P_\eta)$ as in (2.3.15). Thus, for any η of the type $(1, \theta)$, we get the equality of Lie-algebra functors:

$$\text{Lie}(P_\eta) = \mathfrak{P}_\eta. \quad (2.3.16)$$

This can be seen by (2.3.10) and (2.3.13) and then replacing the conjugation in (2.3.3) by $Ad(\eta(s))$.

For a rational 1-PS θ of T , with $\eta = (1, \theta)$ we will have the notation:

$$\mathfrak{P}_\theta := \mathfrak{P}_\eta. \quad (2.3.17)$$

2.4. Pro-groups and pro-Lie algebras. Recall that to each facet $\Sigma_{\mathbb{I}} \subset \overline{\mathfrak{a}_0}$, Bruhat-Tits theory associates a parahoric group scheme $\mathfrak{G}_{\mathbb{I}}$ on $\text{Spec } \mathcal{O}$, which is smooth and affine with connected fibres and whose generic fibre is $G \times_{\text{Spec } k} \text{Spec } K$. Let $L^+(\mathfrak{G}_{\mathbb{I}})$ be the group functor:

$$L^+(\mathfrak{G}_{\mathbb{I}})(R) := \mathfrak{G}_{\mathbb{I}}(R[[z]]). \quad (2.4.1)$$

This is represented by a pro-algebraic group over k which are subgroups of the loop group LG . For $\eta = (1, \theta)$ where θ is any rational 1-PS lying in $\Sigma_{\mathbb{I}}$, we therefore have the following identifications:

$$L^+(\mathfrak{G}_{\mathbb{I}})(k) = P_{\eta} \quad \text{and} \quad \text{Lie}(L^+(\mathfrak{G}_{\mathbb{I}})(k)) = \mathfrak{P}_{\eta}. \quad (2.4.2)$$

2.5. n-parahoric loop groups in higher dimensions. In this subsection we give a loop-theoretic reformulation of the definitions in §2.3 to the higher dimensional base. As always, let $\mathcal{O}_n := k[[z_1, \dots, z_n]]$ and let R denote an arbitrary k -algebra. We denote the field of Laurent polynomials by $K_n = k((z_1, \dots, z_n)) = k[[z_1, \dots, z_n]][\dots, z_i^{-1}, \dots]$. The n -loop group $L_n G$ on a k -algebra R is given by $L_n G(R) := G(R((z_1, \dots, z_n)))$. Similarly the n -loop Lie algebra $L_n \mathfrak{g}$ is given by $L_n \mathfrak{g}(R) := \mathfrak{g}(R((z_1, \dots, z_n)))$.

Let $L_n^{\times} G := \mathbb{G}_m^n \times L_n G$ where the *rotational torus* \mathbb{G}_m^n acts on $L_n G$ by acting on the uniformizer via the *loop rotation action*. This goes as follows: an element $\mathbf{u} := (\dots, u_i, \dots) \in \mathbb{G}_m^n(R)$ acts on $\gamma(z_1, \dots, z_n) \in L_n G(R) = G(R((z_1, \dots, z_n)))$ by

$$\mathbf{u}\gamma(z)\mathbf{u}^{-1} := \gamma(\dots, u_i z_i, \dots). \quad (2.5.1)$$

A maximal torus of $L_n^{\times} G$ is $T^{\times} = \mathbb{G}_m^n \times T$. A $\eta \in \text{Hom}(\mathbb{G}_m, T^{\times}) \otimes_{\mathbb{Z}} \mathbb{Q}$ over $R(s)$ can be viewed as a rational 1-PS. For example if $\eta_i = ((0, \dots, a_i, \dots, 0), \theta_i)$ for $(0, \dots, a_i, \dots, 0) \in \mathbb{Q}^n$ and θ_i a rational 1-PS of T , then we have as before

$$\eta_i(s)\gamma(z_1, \dots, z_n)\eta_i(s)^{-1} = \theta_i(s)\gamma(z_1, \dots, s^{a_i} z_i, \dots, z_n)\theta_i(s)^{-1}. \quad (2.5.2)$$

For η_1, \dots, η_n , as before we can make sense of the condition on $\gamma(z) \in L_n^{\times} G(R)$

$$\text{“}\lim_{s_i \rightarrow 0} \prod \eta_i(s_i)\gamma(z_1, \dots, z_n) \prod \eta_i(s_i)^{-1} \text{ exists in } L_n^{\times} G(R)\text{”}.$$

Similarly, if $\theta = (\theta_1, \theta_i, \dots, \theta_n) \in (Y(T) \otimes \mathbb{Q})^n$ denotes an n -tuple of rational 1-PS's of T , and if $m_r(\theta) = -\lfloor r(\theta) \rfloor$ (1.0.1), in terms of generators we have n -parahoric group $P_{\theta}(R)$ associated to θ as in (1.2.1).

Defining $\eta_i := ((0, \dots, 1, \dots, 0), \theta_i)$ as before, we can describe the n -parahoric groups as follows:

$$P_{\eta_1, \dots, \eta_n}(R) := \{\gamma(z) \in L_n^{\times} G(R) \mid \lim_{s_i \rightarrow 0} \prod \eta_i(s_i) \cdot \gamma(z_1, \dots, z_n) \cdot \prod \eta_i(s_i)^{-1} \text{ exists}\} \quad (2.5.3)$$

$$\text{and in fact } P_{\theta}(R) = P_{\eta_1, \dots, \eta_n}(R) \cap (1 \times L_n G)(R). \quad (2.5.4)$$

Thus facets correspond to n -tuple of usual facets. Further, let $\lambda \in \mathcal{A}_T$. Then

$$\lambda(z_i)P_{\theta}\lambda(z_i)^{-1} = P_{(\theta_1, \dots, \theta_i + \lambda, \dots, \theta_n)}. \quad (2.5.5)$$

So conjugacy classes of parahorics correspond to n -tuples of non-empty subsets of \mathbb{S} .

We can similarly describe the n -parahoric Lie-algebras. In particular,

$$\mathfrak{P}_{\eta_i}^{\times}(R) = \{h \in L_n \mathfrak{g}(R) \mid \lim_{s_i \rightarrow 0} \text{Ad}(\prod \eta_i(s_i))(h(z)) \text{ exists in } L_n^{\times} \mathfrak{g}(R)\}, \quad (2.5.6)$$

$$\text{Thus, } \mathfrak{P}_{\theta}(R) := \mathfrak{P}_{\eta}^{\times}(R) \cap (0 \oplus L_n \mathfrak{g})(R). \quad (2.5.7)$$

2.6. Standard parahoric subgroups. The *standard parahoric subgroups* of $G(K)$ are parahoric subgroups of the distinguished hyperspecial parahoric subgroup $G(\mathcal{O})$. These are realized as inverse images under the evaluation map $ev : G(\mathcal{O}) \rightarrow G(k)$ of standard parabolic subgroups of G . In particular, the standard *Iwahori subgroup* \mathcal{I} is the group $\mathcal{I} = ev^{-1}(B)$.

3. EXAMPLES AND REMARKS

The first example is illustrative of the fact that the operations of imposing regularity conditions and setting variables to be equal do not commute.

Example 3.1. Let $G = \mathrm{SL}_2$, $\mathcal{O} = \mathcal{O}_2$ and $K := K_2$. Let $\Theta = (\theta_1, \theta_2) \in \mathbb{Q}^2$. Then

$$\begin{pmatrix} s_1^{\theta_1} s_2^{\theta_2} & 0 \\ 0 & s_1^{-\theta_1} s_2^{-\theta_2} \end{pmatrix} \begin{pmatrix} X_{11}(s_1 z_1, s_2 z_2) & X_{12}(s_1 z_1, s_2 z_2) \\ X_{21}(s_1 z_1, s_2 z_2) & X_{22}(s_1 z_1, s_2 z_2) \end{pmatrix} \begin{pmatrix} s_2^{-\theta_2} s_1^{-\theta_1} & 0 \\ 0 & s_2^{\theta_2} s_1^{\theta_1} \end{pmatrix} \quad (3.0.1)$$

$$= \begin{pmatrix} X_{11}(s_1 z_1, s_2 z_2) & X_{12}(s_1 z_1, s_2 z_2) s_1^{2\theta_1} s_2^{2\theta_2} \\ X_{21}(s_1 z_1, s_2 z_2) s_1^{-2\theta_1} s_2^{-2\theta_2} & X_{22}(s_1 z_1, s_2 z_2) \end{pmatrix} \subset \mathrm{SL}_2(K(s_1, s_2)). \quad (3.0.2)$$

If we put the regularity condition " $\lim_{s_i \rightarrow 0}$ exists", we get

$$P_{\Theta} = \begin{pmatrix} \mathcal{O} & \mathcal{O} z_1^{-\lfloor 2\theta_1 \rfloor} z_2^{-\lfloor 2\theta_2 \rfloor} \\ \mathcal{O} z_1^{-\lfloor -2\theta_1 \rfloor} z_2^{-\lfloor -2\theta_2 \rfloor} & \mathcal{O} \end{pmatrix}. \quad (3.0.3)$$

Now setting $t := z_1 = z_2$, $B := k[[t]]$ and $\Theta \in (0, \frac{1}{2})^2$ we get

$$P_{\Theta}^{diag} = \begin{pmatrix} B & B \\ t^2 B & B \end{pmatrix}.$$

On the other hand, setting $t := z_1 = z_2$ and $s := s_1 = s_2$ in (3.0.2) we get

$$\begin{pmatrix} X_{11}(st) & X_{12}(st) s^{2\theta_1 + 2\theta_2} \\ X_{21}(st) s^{-2\theta_1 - 2\theta_2} & X_{22}(st) \end{pmatrix}. \quad (3.0.4)$$

Now putting regularity condition " $\lim_{s \rightarrow 0}$ exists", we get $\begin{pmatrix} B & B t^{-\lfloor 2\theta_1 + 2\theta_2 \rfloor} \\ B t^{-\lfloor -2\theta_1 - 2\theta_2 \rfloor} & B \end{pmatrix}$.

Notice for $\Theta \in (0, \frac{1}{2})^2$ that when $0 < \theta_1 + \theta_2 < \frac{1}{2}$, we get $\begin{pmatrix} B & B \\ tB & B \end{pmatrix}$ while when $\frac{1}{2} \leq \theta_1 + \theta_2 < 1$, we get $\begin{pmatrix} B & t^{-1}B \\ t^2B & B \end{pmatrix}$. In either case, for $\Theta \in (0, \frac{1}{2})^2$ we see that the operations of imposing regularity conditions and setting variables to be equal do not commute.

Example 3.2. This example shows that the multiplication law in higher dimensional parahorics is more involved than in the case of discrete valuation rings.

Let $G := \mathrm{SL}_3$, $\mathcal{O} := \mathcal{O}_2$ and $K := K_2$. In this example we study $\Theta := (\frac{\omega_{\alpha_1}^{\vee}}{3}, \frac{\omega_{\alpha_2}^{\vee}}{3})$. The co-weights are given by the points $(\frac{2}{3}, \frac{-1}{3}, \frac{-1}{3}), (\frac{1}{3}, \frac{1}{3}, \frac{-2}{3}) \in \mathcal{A}_T \subset \mathbb{R}^3$ i.e $\omega_{\alpha_1}^{\vee}(t) = \mathrm{diag}(t^{\frac{2}{3}}, t^{\frac{-1}{3}}, t^{\frac{-1}{3}})$ and $\omega_{\alpha_2}^{\vee}(t) = \mathrm{diag}(t^{\frac{1}{3}}, t^{\frac{1}{3}}, t^{\frac{-2}{3}})$. We have

$$\Theta(s_1, s_2) = \mathrm{diag}(s_1^{\frac{2}{9}} s_2^{\frac{1}{9}}, s_1^{\frac{-1}{9}} s_2^{\frac{1}{9}}, s_1^{\frac{-1}{9}} s_2^{\frac{2}{9}}). \quad (3.0.5)$$

Conjugating the 3×3 -matrix $(E_{ij}(s_1 z_1, s_2 z_2))$ by $\Theta(s_1, s_2)$ we get

$$\begin{pmatrix} E_{11}(s_1 z_1, s_2 z_2) & E_{12}(s_1 z_1, s_2 z_2) s_1^{\frac{1}{3}} & E_{13}(s_1 z_1, s_2 z_2) s_1^{\frac{1}{3}} s_2^{\frac{1}{3}} \\ E_{21}(s_1 z_1, s_2 z_2) s_1^{-\frac{1}{3}} & E_{22}(s_1 z_1, s_2 z_2) & E_{23}(s_1 z_1, s_2 z_2) s_2^{\frac{1}{3}} \\ E_{31}(s_1 z_1, s_2 z_2) s_1^{-\frac{1}{3}} s_2^{-\frac{1}{3}} & E_{32}(s_1 z_1, s_2 z_2) s_2^{-\frac{1}{3}} & E_{33}(s_1 z_1, s_2 z_2) \end{pmatrix}. \quad (3.0.6)$$

Imposing regularity conditions forces E_{12} , E_{13} and E_{23} to be regular and $E_{21}(z_1, z_2)$ (resp. $E_{23}(z_1, z_2)$) to have a zero of order at least one with respect to z_1 (resp. z_2) and $E_{31}(z_1, z_2)$ to have a zero of order at least one with respect to both z_1 and z_2 . So we get

$$P_{\Theta} = \left\{ \begin{pmatrix} F_{11}(z_1, z_2) & F_{12}(z_1, z_2) & F_{13}(z_1, z_2) \\ z_1 F_{21}(z_1, z_2) & F_{22}(z_1, z_2) & F_{23}(z_1, z_2) \\ z_1 z_2 F_{31}(z_1, z_2) & z_2 F_{32}(z_1, z_2) & F_{33}(z_1, z_2) \end{pmatrix} \mid F_{ij}(z_1, z_2) \in \mathcal{O}_2 \right\}. \quad (3.0.7)$$

We notice that

$$z_1 z_2 F_{31} = z_1 z_2 (g_{31} h_{11} + g_{32} h_{21} + g_{33} h_{31}) \quad \text{but} \quad (3.0.8)$$

$$z_1 F_{21} = z_1 (g_{21} h_{11} + g_{22} h_{21} + g_{23} h_{31} z_2). \quad (3.0.9)$$

In other words, the multiplication law for the entry (3,1) is like that of $SL_3(\mathcal{O}_2)$ but that of (2,1) has an extra z_2 . Similarly the law for the entries (2,2) and (1,1) are unlike the one-dimensional case.

3.1. Some remarks on specializations of n-parahoric groups and examples. The purpose of this subsection is to show that specialization of n-parahoric groups leads to a strictly larger cadre than that of parahoric groups.

Example 3.3. This example shows the simplest case of relating higher dimensional parahorics to lower ones. We place ourselves in the context of the previous example. If we set $s_2, z_2 := 1$ or $s_1, z_1 := 1$, we get the two standard maximal parabolics. If we set $s_1, z_1, s_2, z_2 := t$ all variables as t , then conjugation is by $\text{diag}(t^{\frac{1}{3}}, 1, t^{-\frac{1}{3}})$. This gives the standard Iwahori (2.6).

PROPOSITION 3.4. *Let $\theta \in \mathcal{A}_T$ and let $m_r(\theta) := -\lfloor r(\theta) \rfloor$ (1.0.1). Then the function $m_{\theta} : \tilde{\Phi} \rightarrow \mathbb{R}$ defined by $m_{\theta}(r) := m_r(\theta)$ is concave.*

Proof. Let $r_i \in \tilde{\Phi}$ be such that $\sum_i r_i \in \tilde{\Phi}$ as well. We need to check that

$$m_{\theta}(\sum_i r_i) = m_{\sum_i r_i(\theta)} \leq \sum_i m_{r_i}(\theta). \quad (3.1.1)$$

This follows by setting $a_i := r_i(\theta)$ and observing $\lfloor a_1 + \dots + a_n \rfloor \geq \lfloor a_1 \rfloor + \lfloor a_2 \rfloor + \dots + \lfloor a_n \rfloor$.
Q.E.D

DEFINITION 3.5. *For $\mathbf{f} := (f_1, \dots, f_n)$ an n-concave function, by the specialization of $P_{\mathbf{f}} \subset G(k((z_1, \dots, z_n)))$ to the diagonal*

$$P_{\mathbf{f}}^{\text{diag}} \subset G(k((t))) \quad (3.1.2)$$

we mean the bounded subgroup obtained by setting $z_1 = z_2 = \dots = z_n := t$.

In particular we have the specializations P_{Ω}^{diag} (of P_{Ω}), and P_{Θ}^{diag} (of P_{Θ}), associated to $\Omega = (\Omega_1, \dots, \Omega_n)$ and $\Theta := (\theta_1, \dots, \theta_n)$ respectively.

Since sums of concave functions are concave, in terms of the concave function $\Phi \rightarrow \mathbb{Z}$ given by $r \mapsto \sum_i m_r(\theta_i)$, we get

$$P_{\Theta}^{\text{diag}} = \langle T(k[[t]]), U_r(t^{\sum_i m_r(\theta_i)} k[[t]]), r \in \Phi \rangle. \quad (3.1.3)$$

We set

$$\mathfrak{P}_{\Theta}^{\text{diag}} := \text{Lie}(\mathbb{P}_{\Theta}^{\text{diag}}). \quad (3.1.4)$$

Remark 3.6. We now make some remarks comparing the *bounded subgroup* $\mathbb{P}_{\Theta}^{\text{diag}}$ with the *parahoric subgroup* $\mathbb{P}_{\Sigma_i \theta_i}$, both subgroups of $G(K)$. Recall that $m_r(\theta) = -\lfloor r(\theta) \rfloor$ (1.0.1). So let us compare $m_r(\Sigma_i \theta_i)$ with $\sum_i m_r(\theta_i)$. For simplicity we assume that all θ_i lie in the dominant Weyl chamber so that for all positive roots r we have $r(\theta_i) > 0$.

Consider the case when for all $r \in R^+$ we have $m_r(\Sigma_i \theta_i) = \sum_i m_r(\theta_i)$. This occurs for instance when all θ_i lie in the alcove \mathfrak{a}_0 and their sum lies in $\mathfrak{a}_0 \setminus \text{far wall}$ or when the θ_i lie in the lattice defined by the θ_{α_i} (2.1.1). For alcove vertices θ_{α_i} an example of such a $\Theta = (\theta_1, \dots, \theta_\ell)$ is given by assigning:

$$\theta_i := \frac{\theta_{\alpha_i}}{(\ell + 1)}. \quad (3.1.5)$$

In this case, for any $r \in \Phi^+$ although $r(\theta_i) > 0$, but nonetheless $m_r(\Sigma_i \theta_i) = 0$ and $m_r(\theta_i) = 0$ and further, $m_{-r}(\theta_i) = 1$ and $m_{-r}(\Sigma_i \theta_i) = 1$. Then the order of poles that the group entries of $\mathbb{P}_{\Theta}^{\text{diag}}$ and $\mathbb{P}_{\Sigma_i \theta_i}$ can allow at r are the same. However, the order of zeros of elements of $\mathbb{P}_{\Theta}^{\text{diag}}$ at $-r$ is the number S_r of distinct simple roots $\alpha \in S$ occurring with strictly positive coefficient in the expression of r . Thus,

$$\mathbb{P}_{\Theta}^{\text{diag}} = \langle T(k[[t]]), U_r(R[[t]]), U_{-r}(t^{S_r} k[[t]]), r \in \Phi^+ \rangle. \quad (3.1.6)$$

Further, the bounded subgroup $\mathbb{P}_{\Sigma_i \theta_i}$ is not a $G(K)$ -conjugate of a subgroup of $\mathbb{P}_{\Theta}^{\text{diag}}$ in this case unless $|S| = 1$.

Now consider the case wherein, for $r \in R^+$, we have $m_r(\Sigma_i \theta_i) < \sum_i m_r(\theta_i)$. This occurs for instance when $G = \text{SL}_2$, $n = 2$ and $r(\theta_1) = r(\theta_2) = a_1 = a_2 = \frac{1}{2}$. In this case, $m_r(\theta_i) = 0$ but $m_r(\Sigma_i \theta_i) = -1$. Then the order of pole which the elements of $\mathbb{P}_{\Sigma_i \theta_i}$ can allow is strictly greater than those of the elements of $\mathbb{P}_{\Theta}^{\text{diag}}$ at r . On the other hand, $m_{-r}(\theta_i) = 1$ and $m_{-r}(\Sigma_i \theta_i) = 1$ as well. So the order of zeros of $\mathbb{P}_{\Sigma_i \theta_i}$ at $-r$ is at least one but that of $\mathbb{P}_{\Theta}^{\text{diag}}$ at $-r$ is at least two. So reasoning on the order of the poles and zeros, it follows that $\mathbb{P}_{\Sigma_i \theta_i}$ is not a $G(K)$ -conjugate of a subgroup of $\mathbb{P}_{\Theta}^{\text{diag}}$.

3.2. Constructing concave functions of type III from type I. The purpose of this subsection is to illustrate the phenomenon that one may begin with a group scheme datum of type I at the generic points of the divisors, but the type of the BT-group scheme at points of depth bigger than 1 could strictly be even of type III. The computations should be seen in the setting of Theorem 1.5.

To illustrate the issue, we work with (3.1.5), which are the simplest non-trivial cases; these also highlight the role of the root system. We make explicit constructions for A_2 , B_2 and G_2 cases with Θ given by (3.1.5). The contrast between the B_2 and G_2 cases highlights the critical role that the root system of the group plays in the analysis.

Recall that for a simple root α we denote by θ_{α} (2.1.1), the vertices of the alcove.

Example 3.7 (A_2). The set of positive roots is $\Phi^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2\}$. We work with $\theta_1 := \theta_{\alpha_1}/3 = \omega_1^\vee/3$ and $\theta_2 := \theta_{\alpha_2}/3 = \omega_2^\vee/3$. Recall $m_r(\theta) := -\lfloor r(\theta) \rfloor$.

$$r(\theta_1) = \begin{cases} 1/3, 0, 1/3 & \text{if } r \in \Phi^+ \\ -1/3, 0, -1/3 & \text{if } r \in \Phi^- \end{cases}. \quad (3.2.1)$$

$$m_r(\theta_1) = \begin{cases} 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 0, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.2)$$

Similarly for θ_2 . We get:

$$r(\theta_1 + \theta_2) = \begin{cases} 1/3, 1/3, 2/3 & \text{if } r \in \Phi^+ \\ -1/3, -1/3, -2/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.3)$$

$$m_r(\theta_1 + \theta_2) = \begin{cases} 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.4)$$

The function $m: r \mapsto \sum m_r(\theta_j)$:

$$m(r) = \begin{cases} 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.5)$$

which computes $P_{\mathfrak{g}}^{diag}$. Let $\Omega \subset \mathcal{A}_T$ be the bounded subset $\Omega = \{b, b'\}$, where $b = \theta_1 + \theta_2$ and $b' = 2\theta_1 + 2\theta_2$. Recall that $m_r(\Omega) = -[\text{Inf}_{\theta \in \Omega} r(\theta)]$. For the subset Ω define the concave function $f_{\Omega}(r) := m_r(\Omega)$ (1.0.3).

$$r(b) = \begin{cases} 1/3, 1/3, 2/3 & \text{if } r \in \Phi^+ \\ -1/3, -1/3, -2/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.6)$$

$$r(b') = \begin{cases} 2/3, 2/3, 4/3 & \text{if } r \in \Phi^+ \\ -2/3, -2/3, -4/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.7)$$

$$m_r(\Omega) = \begin{cases} 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.8)$$

Hence, $m(r) = f_{\Omega}(r)$ for every $r \in \Phi$, i.e., of type II. In particular, the group scheme \mathfrak{G}_m associated to the concave function on $\text{Spec}(A)$ is realized by a bounded subset of the affine apartment.

Before we discuss the next set of examples, we prove a lemma regarding the concave function f_{Ω} defined above. Observe that if

$$f_{\Omega}(r) = a \iff a - 1 < -\inf_{\theta \in \Omega} r(\theta) \leq a \iff 1 - a > \inf_{\theta \in \Omega} r(\theta) \geq -a \quad (3.2.9)$$

The last statement can be interpreted as saying

- (1) $\forall \theta \in \Omega, r(\theta) \geq -a$
- (2) $\exists \theta' \in \Omega$ such that $1 - a > r(\theta')$

LEMMA 3.8. Let Ω be a non-empty bounded region in the affine apartment. Suppose that there is a root r such that $f_{\Omega}(-r) = -f_{\Omega}(r)$. Set $n := f_{\Omega}(r)$. Let $H_r(n)$ be the hyperplane in the affine apartment defined by $H_r(n) := \{\theta \mid r(\theta) = -n\}$. Then,

$$\Omega \subset H_r(n). \quad (3.2.10)$$

Proof. It is immediate from property (1) above. Q.E.D

Example 3.9 (B_2 first example). The set of positive roots is $\Phi^+ = \{\beta, \alpha, \beta + \alpha, \beta + 2\alpha\}$. We work with $\theta_1 := \theta_{\beta}/3 = \omega_1^{\vee}/3$ and $\theta_2 := \theta_{\alpha}/3 = \omega_2^{\vee}/6$. We have

$$r(\theta_1) = \begin{cases} 1/3, 0, 1/3, 1/3 & \text{if } r \in \Phi^+ \\ -1/3, 0, -1/3, -1/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.11)$$

$$m_r(\theta_1) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 0, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.12)$$

$$r(\theta_2) = \begin{cases} 0, 1/6, 1/6, 1/3 & \text{if } r \in \Phi^+ \\ 0, -1/6, -1/6, -1/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.13)$$

$$m_r(\theta_2) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 0, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.14)$$

We get:

$$r(\theta_1 + \theta_2) = \begin{cases} 1/3, 1/3, 2/3, 2/3 & \text{if } r \in \Phi^+ \\ -1/3, -1/3, -2/3, -2/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.15)$$

$$m_r(\theta_1 + \theta_2) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.16)$$

The function $m: r \mapsto \sum m_r(\theta_j)$:

$$m(r) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 2, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.17)$$

which computes P_{θ}^{diag} . Let $\Omega \subset \mathcal{A}_T$ be the bounded subset $\Omega = \{b, b'\}$, where $b = \theta_1 + \theta_2$ and $b' = (2-e)\theta_1 + (2+e')\theta_2$, where $0 < e, e' < 1$. Recall that $m_r(\Omega) = -\lfloor \text{Inf}_{\theta \in \Omega} r(\theta) \rfloor$. For the subset Ω define the concave function $f_{\Omega}(r) := m_r(\Omega)$.

$$r(b) = \begin{cases} 1/3, 1/6, 1/2, 2/3 & \text{if } r \in \Phi^+ \\ -1/3, -1/3, -1/2, -2/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.18)$$

$$r(b') = \begin{cases} 2/3 - e/3, 1/3 + e'/6, 1 + (e' - 2e)/6, 4/3 + (e' - e)/3 & \text{if } r \in \Phi^+ \\ -2/3 + e/3, -1/3 - e'/6, -1 - (2e - e')/6, -4/3 - (e - e')/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.19)$$

$$m_r(\Omega) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 2, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.20)$$

Hence, $m(r) = f_{\Omega}(r)$ for every $r \in \Phi$, i.e., again of type II.

Example 3.10 (B_2 second example). The second example here has special significance for §11. This time we work with $\theta_1 := \theta_{\alpha}/2 = \omega_{\alpha}^{\vee}/4$ and $\theta_2 := \theta_{\alpha} = \omega_{\alpha}^{\vee}/2$. Then we get

$$m_r(\theta_1) = \begin{cases} 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 0, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.21)$$

and

$$m_r(\theta_2) = \begin{cases} 0, 0, 0, -1 & \text{if } r \in \Phi^+ \\ 0, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.22)$$

The function $m: r \mapsto \sum m_r(\theta_j)$ is given by:

$$m(r) = \begin{cases} 0, 0, 0, -1 & \text{if } r \in \Phi^+ \\ 0, 2, 2, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.23)$$

Claim: The concave function m (3.2.23) is not of type II, i.e., there does not exist any bounded region $\Omega \subset \mathcal{A}$ such that $m = f_{\Omega}$. If not, then there is an Ω such that $f_{\Omega}(\beta) = -f_{\Omega}(-\beta) = 0$ and hence $\Omega \subset H(0)$. Again, since $f_{\Omega}(\alpha) = 0, f_{\Omega}(-\alpha) = 2$ (3.2.23), it follows that $\forall \theta \in \Omega$, by (1) before (3.8), we get the condition:

$$0 \leq \alpha(\theta) \leq 2, \forall \theta \in \Omega. \quad (3.2.24)$$

Similarly, $f_{\Omega}(\alpha + \beta) = 0, f_{\Omega}(-(\alpha + \beta)) = 2$ (3.2.23), forces:

$$0 \leq (\alpha + \beta)(\theta) \leq 2, \forall \theta \in \Omega \quad (3.2.25)$$

and $f_{\Omega}(2\alpha + \beta) = -1, f_{\Omega}(-(2\alpha + \beta)) = 2$ forces:

$$(2\alpha + \beta)(\theta) \leq 2, \forall \theta \in \Omega. \quad (3.2.26)$$

On the other hand, since $f_\Omega(-\alpha) = 2$, (2) says that $\exists \theta' \in \Omega$ such that

$$1 < \alpha(\theta'). \quad (3.2.27)$$

Notice that since $\beta(\theta') = 0$, the two inequalities (3.2.26) and (3.2.27) are not compatible.

Example 3.11 (G_2 first example). The set of positive roots is $\Phi^+ = \{\alpha, \beta, \alpha + \beta, 2\alpha + \beta, 3\alpha + \beta, 3\alpha + 2\beta\}$. We work with $\theta_1 := \theta_\alpha/3 = \omega_1^\vee/9$ and $\theta_2 := \theta_\beta/3 = \omega_2^\vee/6$. We have

$$r(\theta_1) = \begin{cases} 1/9, 0, 1/9, 2/9, 1/3, 1/3 & \text{if } r \in \Phi^+ \\ -1/9, 0, -1/9, -2/9, -1/3, -1/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.28)$$

$$r(\theta_2) = \begin{cases} 0, 1/6, 1/6, 1/6, 1/6, 1/3 & \text{if } r \in \Phi^+ \\ 0, -1/6, -1/6, -1/6, -1/6, -1/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.29)$$

$$m_r(\theta_1) = \begin{cases} 0, 0, 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 0, 1, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.30)$$

$$m_r(\theta_2) = \begin{cases} 0, 0, 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 0, 1, 1, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.31)$$

We get:

$$r(\theta_1 + \theta_2) = \begin{cases} 1/9, 1/6, 5/18, 7/18, 1/2, 2/3 & \text{if } r \in \Phi^+ \\ -1/9, -1/6, -5/18, -7/18, -1/2, -2/3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.32)$$

$$m_r(\theta_1 + \theta_2) = \begin{cases} 0, 0, 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 1, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.33)$$

The function $m: r \mapsto \sum m_r(\theta_j)$:

$$m(r) = \begin{cases} 0, 0, 0, 0, 0, 0 & \text{if } r \in \Phi^+ \\ 1, 1, 2, 2, 2, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.34)$$

which computes $P_{\mathfrak{o}}^{diag}$.

Claim: In contrast, the concave function m (3.2.34) is not of type II, i.e., there does not exist any bounded region $\Omega \subset \mathcal{A}$ such that $m = f_\Omega$. This is explicit here but see also [BT1, 6.4.4, page 133].

To prove this claim, let us suppose to the contrary that there exists such a bounded subset $\Omega \subset \mathcal{A}_T$. Let $\theta \in \Omega$, be given by

$$\theta := (x\omega_1^\vee, y\omega_2^\vee).$$

Then we have

$$r(\theta) = \begin{cases} x, y, x+y, 2x+y, 3x+y, 3x+2y & \text{if } r \in \Phi^+ \\ -x, -y, -x-y, -2x-y, -3x-y, -3x-2y & \text{if } r \in \Phi^- \end{cases} \quad (3.2.35)$$

For $r \in \Phi^+$, since $-\lfloor \inf_{\theta \in \Omega}(r, \theta) \rfloor = 0$ so $\inf_{\theta \in \Omega}(r, \theta) \in [0, 1)$. Thus for $r = \alpha$ and β , for all points θ in Ω we have

$$x, y \in [0, 1).$$

Now let us assume that $\inf_{\theta \in \Omega}(r, \theta)$ is realised at (x_1, y_1) for $r = -(\alpha + \beta)$ and at (x_2, y_2) for $r = -(3\alpha + 2\beta)$. In particular, $x_1, x_2, y_1, y_2 \in [0, 1)$.

We have $-\lfloor -(x_1 + y_1) \rfloor = 2$. So $-2 \leq -x_1 - y_1 < -1$. Thus

$$1 < x_1 + y_1 \leq 2 \quad (3.2.36)$$

$$x_1 > 0. \quad (3.2.37)$$

By the condition for $-3\alpha - 2\beta$, we have $-3x_2 - 2y_2 \leq -3x_1 - 2y_1$. Thus

$$3x_1 + 2y_1 \leq 3x_2 + 2y_2. \quad (3.2.38)$$

By $-[-3x_2 - 2y_2] = 2$, we have $-1 > -3x_2 - 2y_2 \leq -2$. Thus

$$1 < 3x_2 + 2y_2 \leq 2. \quad (3.2.39)$$

We get the following contradiction

$$2 \stackrel{(3.2.36)}{<} 2x_1 + 2y_1 \stackrel{(3.2.37)}{<} 3x_1 + 2y_1 \stackrel{(3.2.38)}{\leq} 3x_2 + 2y_2 \stackrel{(3.2.39)}{\leq} 2.$$

This completes the proof of the claim.

Example 3.12 (G_2 second example). The second example here has special significance for §11. We work now with $\theta_1 := \theta_\alpha = \omega_\alpha^\vee/3$ and $\theta_2 := 2\theta_\alpha = 2\omega_\alpha^\vee/3$. Then we get

$$m_r(\theta_1) = \begin{cases} 0, 0, 0, 0, -1, -1 & \text{if } r \in \Phi^+ \\ 1, 0, 1, 1, 1, 1 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.40)$$

$$m_r(\theta_2) = \begin{cases} 0, 0, 0, -1, -2, -2 & \text{if } r \in \Phi^+ \\ 1, 0, 1, 2, 2, 2 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.41)$$

The function $m: r \mapsto \sum m_r(\theta_j)$ is given by:

$$m(r) = \begin{cases} 0, 0, 0, -1, -3, -3 & \text{if } r \in \Phi^+ \\ 2, 0, 2, 3, 3, 3 & \text{if } r \in \Phi^- \end{cases} \quad (3.2.42)$$

Claim: The concave function m (3.2.42) is not of type II. If not, then there is an Ω such that $f_\Omega(\beta) = -f_\Omega(-\beta) = 0$ and hence by (3.8) we have $\Omega \subset H_\beta(0)$. Thus

$$\beta(\theta) = 0 \forall \theta \in \Omega. \quad (3.2.43)$$

Similarly, $f_\Omega(3\alpha + 2\beta) = 3, f_\Omega(-(3\alpha + 2\beta)) = -3$ (3.2.42), forces:

$$(3\alpha + 2\beta)(\theta) = 3, \forall \theta \in \Omega \quad (3.2.44)$$

and (3.2.43) and (3.2.44) gives us

$$\alpha(\theta) = 1 \forall \theta \in \Omega. \quad (3.2.45)$$

Now since $f_\Omega(-\alpha) = 2$, by (2) before (3.8), we know that there exists a $\theta' \in \Omega$ such that $\alpha(\theta') > 1$, which is incompatible with (3.2.45).

Example 3.13. (The case of constant concave function). Let $a \geq 0$ and $f(r) = a$ for all $r \in \Phi$. In this case, when G is of type A_n, B_n or C_n , there exists regions Ω such that $f_\Omega = f$. For G_2 when $a \geq 3$, there does not exist a region.

3.3. Various cones of concave functions. We consider the question: "how to construct general concave functions from points in the apartment?" It turns out that this can always be done in the case A_n (see (3.21) and the remarks before it). Let Φ be an irreducible root system. Let

$$\Phi^\Sigma := \{(\alpha, \beta) \in \Phi^2 \mid \alpha + \beta \in \Phi\}. \quad (3.3.1)$$

For $(\alpha, \beta) \in \Phi^\Sigma$, we get functionals $\psi_{(\alpha, \beta)}$ and ψ_α defined on $\{f: \Phi \rightarrow \mathbb{R}\} \rightarrow \mathbb{R}$ by $f \mapsto f(\alpha) + f(\beta) - f(\alpha + \beta)$ and $f \mapsto f(\alpha) + f(-\alpha)$ respectively.

Let \mathcal{C} denote the space of concave functions $\{f: \Phi \rightarrow \mathbb{R}\}$. It can be expressed as

$$\mathcal{C} = \cap \{\psi_{(\alpha, \beta)} \geq 0\} \cap \{\psi_\alpha \geq 0\}. \quad (3.3.2)$$

Being an intersection of half-spaces, it is a polyhedral cone and further since the intersection of the open half-spaces is non-empty, we see that \mathcal{C} is of dimension $|\Phi|$.

Let \mathcal{C}_Ω denote the cone of concave functions defined by bounded regions $\Omega \subset \mathcal{A}_T$.

PROPOSITION 3.14. *Let Φ be an irreducible root system. The dimension of \mathcal{C}_Ω equals $|\Phi|$.*

Proof. Recall a bounded subset $\Omega \subset \mathcal{A}_T$, defines the concave function

$$f_\Omega : \Phi \rightarrow \mathbb{R} \quad r \mapsto -[\inf_{\theta \in \Omega} r(\theta)]. \quad (3.3.3)$$

For $r \in \Phi$, let H_r be the hyperplane in \mathcal{A}_T where r vanishes. Consider the unit sphere S in \mathcal{A}_T . We fix an isomorphism of T with a product of \mathbb{G}_m . This gives a basis of $Y(T)$ and $X(T)$. Now we may identify \mathcal{A}_T with its dual space so that for a root $r \in \Phi$ and $\theta \in \mathcal{A}_T$, the evaluation $r(\theta)$ equals the inner product of θ with the image of r . Thus, after this identification, every root $r \in \Phi$ defines a canonical unit vector normal to H_r meeting S in p_r . Since Φ is irreducible, $H_r = H_s \implies r = \pm s$. Further, $-p_r = p_{-r}$. So $r \mapsto p_r$ defines a bijection. To prove the proposition, we now consider the bounded subset

$$\Omega := \{p_r | r \in \Phi\}. \quad (3.3.4)$$

So $|\Omega| = |\Phi|$. For any $r \in \Phi$, $\inf_{\theta \in \Omega} r(\theta)$ is attained at p_{-r} . We now let points vary in radial directions determined by $\{p_r\}$. In small open neighbourhoods of p_r , if $\Omega' := \{p'_r\}$ denotes a neighbouring set of points, we again see that $\inf_{\theta \in \Omega'} r(\theta)$ is attained at p'_{-r} . This completes the proof.

Q.E.D

3.4. Recovering the bounded group P_Ω over a dvr in terms of n-parahoric groups.

Let $\Omega \subset \mathcal{A}$ be a bounded subset of the apartment. Without loss of generality, we may suppose that it is compact.

We make a small modification in the notation just for this lemma. Let $A_1 = L[t]$ be a polynomial ring over a field L not necessarily algebraically closed. Let $K_1 = \text{Fract}(A_1)$. Similarly, let $A_N = L[z_1, \dots, z_N]$ and $K_N = \text{Fract}(A_N)$. Let

$$P_\Omega := \langle T(L[t]), U_r(t^{m_r(\Omega)} L[t]) \quad r \in \Phi \rangle, \quad (3.4.1)$$

and for $\Theta = (\theta_1, \dots, \theta_N)$,

$$P_\Theta := \langle T(A_N), U_r(\prod_{1 \leq j \leq N} z_j^{-\lfloor r, \theta_j \rfloor} A_N) \quad r \in \Phi \rangle. \quad (3.4.2)$$

The next lemma shows the natural relationship between n-bounded groups associated to bounded subsets Ω in the affine apartment and N-parahoric groups. We give it here for its general interest.

LEMMA 3.15. *Let $N = |\Phi|$ and let the inclusion $G(K_1) \subset G(K_N)$ be obtained by mapping the uniformizer t to $\prod_{r \in \Phi} z_r$. Given a bounded domain Ω , there exist an N -tuple $\Theta = (\dots, \theta_r, \dots)_{r \in \Phi}$ of points such that taking intersection in $G(K_N)$ we have the equality*

$$P_\Omega = G(K_1) \cap P_\Theta. \quad (3.4.3)$$

Proof. For every root $r \in \Phi$, we may choose a $\theta_r \in \Omega$ such that we have

$$(r, \theta_r) \leq (r, \theta) \quad \forall \theta \in \Omega. \quad (3.4.4)$$

Set $L[\Phi] := L[z_r, r \in \Phi]$ and consider the N-parahoric given by

$$\Theta = (\dots, \theta_j, \dots)_{j \in \Phi}.$$

Thus, by definition we have

$$P_{\mathbf{0}} = \langle T(L[\Phi]), U_r \left(\prod_{j \in \Phi} z_j^{-\lfloor (r, \theta_j) \rfloor} L[\Phi] \right) \mid r \in \Phi \rangle. \quad (3.4.5)$$

For $r \in \Phi$, consider the group $U_r \left(\prod_{j \in \Phi} z_j^{-\lfloor (r, \theta_j) \rfloor} L[\Phi] \right)$. By (3.4.4), we see that

$$(r, \theta_r) \leq (r, \theta_j) \quad \forall j \in \Phi. \quad (3.4.6)$$

Thus $-\lfloor (r, \theta_r) \rfloor \geq -\lfloor (r, \theta_j) \rfloor$ for all $j \in \Phi$. In other words we have

$$-\lfloor (r, \theta_r) \rfloor = -\lfloor \inf_{\theta \in \Omega} (r, \theta) \rfloor = m_r(\Omega). \quad (3.4.7)$$

Now $K_1 = L(t)$ and $A_1 = L[t]$, so letting $t = \prod_{r \in \Phi} z_r$, for every root $r \in \Phi$ we get the desired order of zeros and poles. Further $T(L[\Phi]) \cap T(K_1) = T(A_1)$. Since the group law is inherited from $G(K_1)$, the intersection is the parahoric subgroup P_{Ω} .

Q.E.D

For $1 \leq i \leq n$, let $f_i : \Phi \rightarrow \mathbb{R}$ be concave functions. Let $\mathbf{F} := (\dots, f_i, \dots) : \Phi \rightarrow \mathbb{R}^n$. In (1.2.3), we have defined the n -concave subgroup $P_{\mathbf{F}}$ of $G(K_n)$ in terms of generators.

LEMMA 3.16. *Let $G(K_1) \subset G(K_n)$ by mapping the uniformizer t to $\prod_{1 \leq i \leq n} z_i$. Taking intersection in $G(K_n)$ we have the equality*

$$P_{\sup_i f_i} = G(K_1) \cap P_{\mathbf{F}}, \quad (3.4.8)$$

Proof. The proof is almost identical to that of Lemma 3.15.

Q.E.D

Let $\mathcal{C}_{\Omega}^{ext}$ denote the cone defined by (1): taking sums of concave functions of the form $\sum_{i=1}^m f_{\Omega_i}$ where we let $m \geq 1$ and bounded regions $\Omega_i \subset \mathcal{A}_T$ vary, (2): changing the uniformizers by the substitution $t = z^n$ or $t^n = z$ and (3): by taking supremums.

We have the obvious inclusions

$$\mathcal{C}_{\Omega} \subset \mathcal{C}_{\Omega}^{ext} \subset \mathcal{C} \quad (3.4.9)$$

Example 3.17. *The first inclusion is strict for the concave function (3.2.34) by 3.11.*

For irreducible reduced root systems, the notions of quasi-concave and concave functions coincide. Let f be any concave function defined more generally on Φ . Let E be the set of points in \mathcal{A}_T fixed by G_f . For a root $\alpha \in \Phi$, let

$$S(\alpha) := \{(\alpha_i, \lambda_i)_{\{1 \leq i \leq r\}} \in \cup_{r \geq 1} (\Phi \times \mathbb{Q}_+^*)^r \mid \sum_{1 \leq i \leq r} \lambda_i \alpha_i = \alpha\} \quad (3.4.10)$$

denote all possible ways of writing α in the rational positive cone of some subset of roots in Φ . Let f' be the function on Φ defined by:

$$f'(\alpha) := \inf_{S(\alpha)} \left[\sum_{1 \leq i \leq r} \lambda_i f(\alpha_i) \right]. \quad (3.4.11)$$

By [Cou, Prop 3.3], we have $f' = f_E$ for some subset E of the apartment.

Example 3.18. For the case $G = \mathrm{SL}_2$, the three cones coincide because any concave function is of the form f_{Ω} (see (3.2.1)). Indeed, by [Cou, Prop 3.3] for $G = \mathrm{SL}_2$, we have $S(\alpha) = \{(\alpha, 1)\}$ and $S(-\alpha) = \{(-\alpha, 1)\}$. Thus for $G = \mathrm{SL}_2$, the subset E determines f in return.

PROPOSITION 3.19. *We now return to the case $G = G_2$. For the concave function m (3.2.34), m' equals m except for $-\alpha_1 - \alpha_2$ where it is one. The region is the dilation of the alcove \mathbf{a}_0 by a factor of two i.e. enclosed by origin, $2\theta_{\alpha_1}$ and $2\theta_{\alpha_2}$.*

Proof. We give an indirect proof. Since \mathfrak{m} does not come from a region, we have the inequality $\mathfrak{m}' < \mathfrak{m}$. Now we can check that the concave function coming from the dilation of \mathfrak{a}_0 by a factor of two is as stated above. Since there cannot be any other concave function between them, this must be \mathfrak{m}' . Q.E.D

Remark 3.20. It would be interesting to understand the inclusion $\mathcal{C}_\Omega^{ext} \subset \mathcal{C}$ better. Given bounded regions $\{\Omega_1, \dots, \Omega_n\}$, consider the concave function f taking r to $\sum m_r(\Omega_i)$. It would be interesting to understand the region corresponding to f' as a function of $\{\Omega_i\}$.

Recall (1.2.3), if $f: \Phi \rightarrow \mathbb{R}$ is a concave function then we can associate the bounded subgroup $P_f := \langle T(\mathcal{O}), U_r(z^{f(r)}\mathcal{O}), r \in \Phi \rangle$ of $G(K)$. Note that $\mathfrak{G}_f(\mathcal{O}) = P_f$.

Following ([BT2, 4.5.2] and [Cou, page 540]) we say that a concave function f is *optimal* if for any concave function $f': \Phi \rightarrow \mathbb{R}$, such that $f' > f$, we have a strict inclusion $P_{f'} \subsetneq P_f$. However, in the context of the present paper, it is easy to see that if f is a concave function then so is $\lceil f \rceil$ which is in fact optimal. Moreover, the bounded groups $\mathfrak{G}_f(\mathcal{O})$ coincides with $\mathfrak{G}_{\lceil f \rceil}(\mathcal{O})$. So we may as well work with optimal concave functions.

In [BT3, Remarques 3.9.3], it is remarked that on root systems of type A_n , any optimal concave function is of the form f_Ω for a *enclosed* bounded subset Ω of the apartment(see [BT1, 2.4.6] for the definition of "enclosed" subsets). For the sake of completeness we give below a proof of this statement.

PROPOSITION 3.21. *Let G be of type A_n . Any optimal concave function f is of type II, i.e., of the form f_Ω for a enclosed bounded subset Ω of the apartment. Thus we have $\mathcal{C}_\Omega = \mathcal{C}$ (3.20).*

Proof. For the convenience of the reader we recast the notions of [Cou], which were developed for connected reductive groups over local fields, into the setting of this paper. Recall that the residue field of our local rings are algebraically closed (see). For a root $\alpha \in \Phi$, the $c(\alpha)$ and $v(\alpha)$ may be assumed to be 0 and 1 respectively and further an optimal concave function takes only integral values. By [Cou, Prop 2.3], concavity is equivalent to pseudo-concavity since our group is split over K . A function is of the form f_Ω if and only if it is strongly concave according to [Cou, Prop 3.3].

For a root $\alpha \in \Phi$, let $S'(\alpha) \subset S(\alpha)$ (3.4.10) be the restricted set such that

- the α_i are linearly independent;
- for every $i \neq j$, we have $(\alpha_i, \alpha_j) \geq 0$; we can assume that $\alpha_i + \alpha_j$ is not a root;
- for every i , $\lambda_i < 1$.

In [Cou, Lemma 5.1] Courtes gives an equivalent reformulation of strong concavity in terms of a technical numerical inequality which should be checked for each element in $S'(\alpha)$ where α varies in Φ . We claim that when $G = A_n$, then this numerical condition is vacuously true because $S'(\alpha) = \emptyset$. We see this as follows.

Let $\alpha = \epsilon_l - \epsilon_m$ be a root of A_n . Since $\alpha = \sum_{i=1}^r \lambda_i \alpha_i = \epsilon_l - \epsilon_m$ and $0 < \lambda_i < 1$ we see that $r \geq 2$ and there exist at least two indices $1 \leq i_1 < i_2 \leq r$ such that $\alpha_{i_1} = \epsilon_l - \epsilon_{j_1}$ and $\alpha_{i_2} = \epsilon_l - \epsilon_{j_2}$. We suppose firstly that j_1 is different from m . In this case, there must exist another index $1 \leq i_3 \leq r$ such that $\alpha_{i_3} = \epsilon_{j_1} - \epsilon_{j_2}$. But then $\alpha_{i_1} + \alpha_{i_3} = \epsilon_l - \epsilon_{j_2}$ is either a root of A_n or $\alpha_{i_1} = -\alpha_{i_3}$. Both these possibilities violate the conditions for $\{(\lambda_{i_1}, \alpha_{i_1}), (\lambda_{i_2}, \alpha_{i_2})\}$ to be a subset of an element in $S'(\alpha)$. Now suppose that j_1 equals m . Then α_{i_1} equals α . So we have $\sum_{i=1}^r \lambda_i \alpha_i - \lambda_{i_1} \alpha_{i_1} = (1 - \lambda_{i_1}) \alpha = (1 - \lambda_{i_1})(\epsilon_l - \epsilon_m) \neq 0$. Repeating the above argument, we find that α_{i_2} equals α as well. But then α_{i_1} and α_{i_2}

are the same. This violates the conditions for $\{(\lambda_{i_1}, \alpha_{i_1}), (\lambda_{i_2}, \alpha_{i_2})\}$ to be a subset of an element in $S'(\alpha)$. Q.E.D

Remark 3.22. Let $f : \tilde{\Phi} \rightarrow \mathbb{Z}$ be a concave function. We have $f(r) = f(0+r) \leq f(0) + f(r)$. Thus, $0 \leq f(0)$. Let us fix a value $f(0)$ of $0 \in \tilde{\Phi}$. For each simple root $\alpha \in S$, by the condition $f(0) = f(\alpha + -\alpha) \leq f(\alpha) + f(-\alpha)$, we have

$$(f(\alpha), f(-\alpha)) \in \{(x, y) \in \mathbb{Z}^2 \mid x + y \geq f(0)\}.$$

On the other hand, the condition $f(r_1) = f(r_1 + r_2 - r_2) \leq f(r_1 + r_2) + f(-r_2)$ implies

$$f(r_1) - f(r_2) \leq f(r_1 + r_2) \leq f(r_1) + f(r_2).$$

Thus, for an integral concave function f we have infinitely many choices for values on $S^\pm \cup \{0\}$ but only finitely many choices for non-simple roots.

Part 2. SCHEMATIZATION

4. THE n -PARAHORIC LIE-ALGEBRA BUNDLE \mathcal{R} ON AN AFFINE SPACE

The aim of this section is to present an *ab initio* construction of the n -parahoric Lie algebra bundle on the affine n -space together with a full description of the fibres at points of higher depth. This played the initially important role for us to guess the definition of the n -parahoric group. The reader who is interested only in the group scheme constructions may skip this section.

For $1 \leq i \leq n$, let $H_i \subset \mathbf{A} := \mathbb{A}_k^n$ denote the coordinate hyperplanes. Let $\mathbf{A}_0 \subset \mathbf{A}$ be the open subscheme which is the complement of the union of hyperplanes. Let $\Theta := (\theta_1, \dots, \theta_n) \in \mathcal{A}^n$ be a point of the n -apartment satisfying the conditions of §2.2. The aim of this section is to construct a n -parahoric Lie algebra bundle \mathcal{R} on $\mathbf{A} := \mathbb{A}_k^n$ such that its restriction to the generic points of the hyperplane H_i are the parahoric Lie algebra bundles associated to the weight θ_i and to describe it on the whole of \mathbf{A} .

4.1. Elementary remarks on reflexive sheaves. We begin with three elementary lemmas which should be well known.

LEMMA 4.1. *Let E be a locally free sheaf on an irreducible smooth scheme X . Let $\xi \in X$ be the generic point and let $W \subset E_\xi$ be an \mathcal{O}_ξ -submodule. Then there exists a unique coherent subsheaf $F \subset E$ such that $F_\xi = W$ and $Q := \text{Coker}(F \hookrightarrow E)$ is torsion-free. Moreover F is a reflexive sheaf.*

Proof. Define a sheaf \tilde{F} by the following condition: its section on an affine open $U \subset X$ are given by $\tilde{F}(U) := E(U) \cap W$. Then it is easily seen that \tilde{F} defines a coherent subsheaf $F \subset E$ and it is the maximal coherent subsheaf of E whose fiber over ξ is W . To check Q is torsion-free, let T be the torsion submodule of Q . Let $K := \text{Ker}(E \rightarrow Q/T)$. Then since $\xi \notin \text{Supp}(T)$, $K_\xi = W$ and hence, by the maximality of F we have $K = F$. Since E is locally free, we have $F^{\vee\vee}/F \hookrightarrow E^{\vee\vee}/F = E/F$. But since $F^{\vee\vee}/F$ is only torsion, and $E/F = Q$ is torsion-free, it follows that F is automatically a reflexive sheaf. Q.E.D

LEMMA 4.2. *Let X be as above and $i : U \hookrightarrow X$ an open subset such that $X \setminus U$ has codimension ≥ 2 in X . Let F_U be a reflexive sheaf on U . Then $i_*(F_U)$ is a reflexive sheaf on X which extends F_U .*

Proof. By [EGA, Corollaire 9.4.8], there exists a coherent \mathcal{O}_X -module F_1 such that $F_1|_U \simeq F_U$. Set $F := F_1^{**}$ to be the double dual. Then F is reflexive and also since F_U is reflexive, $F|_U \simeq F_U$. Hence, we have $i_*(F_U) = F$ [H, Proposition 1.6]. Q.E.D

LEMMA 4.3. *Let X be integral and factorial. Then any reflexive sheaf of rank 1 is locally free.*

Proof. ([H, Proposition 1.9]) Let \mathcal{F} be a reflexive sheaf of rank 1. Then since X is normal, there is a *open* $U \subset X$ such that $\text{codim}_X(U) \geq 2$, and \mathcal{F}_U is locally free. Since X is locally factorial, we have $\text{Pic}(X) \rightarrow \text{Pic}(U)$ is bijective. Hence, there is a locally free sheaf \mathcal{L} of rank 1 on X such that $\mathcal{L}_U = \mathcal{F}_U$. It is clear that $\mathcal{L} \simeq \mathcal{F}$ on X . Q.E.D

4.2. The construction of \mathcal{R} .

THEOREM 4.4. *Associated to Θ , there exists a canonical Lie-algebra bundle \mathcal{R} on \mathbf{A} which extends the trivial bundle with fiber \mathfrak{g} on $\mathbf{A}_0 \subset \mathbf{A}$; furthermore, in formal neighbourhood $V := \text{Spec}(\hat{\mathcal{O}}_n)$ of the origin in the affine n -space \mathbf{A} , we have the identification of functors from the category of k -algebras to k -Lie-algebras:*

$$L_n^+(\mathcal{R}|_V) = L_n^+(\mathfrak{A}_\Theta^+). \quad (4.2.1)$$

Proof. Let $b_{i\alpha} \in \mathbb{Q}$ be defined by the relations

$$\theta_i := \sum_{\alpha \in S} b_{i\alpha} \omega_\alpha^\vee \quad \text{for } i \leq n. \quad (4.2.2)$$

Let $d \in \mathbb{N}$ be such that $db_{i\alpha} \in \mathbb{Z}$ for all $1 \leq i \leq n$ and $\alpha \in S$.

Consider an inclusion of k -algebras $B_0 \subset B$, where

$$B := k[y_i^{\pm 1}]_{1 \leq i \leq n}, \quad B_0 := k[x_i^{\pm 1}]_{1 \leq i \leq n} \quad (4.2.3)$$

and $\{y_i^d = x_i\}_{1 \leq i \leq n}$. Let $\mathbf{Y}^0 := \text{Spec}(B)$, and $\mathbf{A}_0 = \text{Spec}(B_0)$, and let $p: \mathbf{Y}^0 \rightarrow \mathbf{A}_0$ be the natural morphism. We define the ‘‘roots’’ map:

$$\tau: \mathbf{Y}^0 \rightarrow \mathbf{A}_0, \text{ as} \quad (4.2.4)$$

$$\tau^\#((x_i)) := (y_i). \quad (4.2.5)$$

Note that as a map between tori, τ is an isomorphism. Let $T_{ad} = \text{Spec} k[z_\alpha^{\pm 1}]$. We define the action map:

$$\alpha: \mathbf{Y}^0 \rightarrow T_{ad}, \text{ as} \quad (4.2.6)$$

$$\alpha^\#((z_\alpha)) := \left(\prod_{1 \leq i \leq n} y_i^{db_{i\alpha}} \right). \quad (4.2.7)$$

We consider the map

$$\text{Ad} \circ \alpha: \mathbf{Y}^0 \times \mathfrak{g} \rightarrow \mathbf{Y}^0 \times \mathfrak{g} \quad (4.2.8)$$

$$(\mathbf{t}, x) \mapsto (\mathbf{t}, \text{Ad}(\alpha(\mathbf{t}))(x)). \quad (4.2.9)$$

We define the embedding of modules

$$j: \mathfrak{g}(B_0) \hookrightarrow \mathfrak{g}(B) \xrightarrow{\Gamma(\text{Ad} \circ \alpha)} \mathfrak{g}(B) \quad (4.2.10)$$

where the second map is the one induced by (4.2.8) on sections. The T -weight space decomposition of \mathfrak{g} induces a T -weight space decomposition on $\mathfrak{g}(B_0)$ by viewing the latter as sections of the trivial bundle on \mathbf{A}_0 with fibers \mathfrak{g} .

Let $B^+ = k[y_i]_{1 \leq i \leq n}$ and $B_0^+ := k[x_i]_{1 \leq i \leq n}$. Taking intersection as Lie submodules of $\mathfrak{g}(B)$ we define the following B_0^+ -submodule of $\mathfrak{g}(B_0)$:

$$\mathcal{R}(B_0^+) := j^{-1}(j(\mathfrak{g}(B_0)) \cap \mathfrak{g}(B^+)). \quad (4.2.11)$$

In words, it is the set of elements in $\mathfrak{g}(B_0)$ which lie in $\mathfrak{g}(B^+)$ under j . Taking E as the bundle defined by the free B_0^+ -module $\mathfrak{g}(B^+)$ and the submodule $W := j(\mathfrak{g}(B_0))$ in (4.1),

we observe that \mathcal{R} is a reflexive sheaf on the affine embedding $\mathbf{A}_0 \hookrightarrow \mathbf{A} := \text{Spec}(B_0^+) (\simeq \mathbb{A}^n)$. Further, this is a sheaf of Lie algebras with its Lie bracket induced from $\mathfrak{g}(B)$.

We now check that \mathcal{R} is locally-free on \mathbf{A} . Observe that the intersection (4.2.11) also respects T_{ad} -weight space decomposition on the sections. More precisely, for any root $r \in \Phi$, since by definition we have $\mathcal{R}(B_0^+)_r = j^{-1}(j(\mathfrak{g}_r(B_0)) \cap \mathfrak{g}_r(B^+))$, and so we get the following equalities:

$$j^{-1}(j(\mathfrak{g}(B_0)) \cap \mathfrak{g}_r(B^+)) = j^{-1}(j(\mathfrak{g}_r(B_0)) \cap \mathfrak{g}(B^+)) = \mathcal{R}(B_0^+)_r. \quad (4.2.12)$$

Since \mathcal{R} is reflexive, there is an open subset $U \subset \mathbf{A}$ with complement of codimension at least two such that the restriction $\mathcal{R}' := \mathcal{R}|_U$ is locally free. Clearly U contains \mathbf{A}_0 and the generic points ζ_α of the divisors H_α . This gives a decomposition on \mathcal{R}' obtained by restriction from \mathcal{R} . The locally free sheaf \mathcal{R}' is a direct sum of the trivial bundle $\text{Lie}(T_{\text{ad}}) \times U$ (coming from the 0-weight space) and the invertible sheaves coming from the root decomposition. Now since invertible sheaves extend as invertible sheaves across codimension ≥ 2 (4.3), the reflexivity of \mathcal{R} implies that this direct sum decomposition of \mathcal{R}' extends to \mathbf{A} . Whence \mathcal{R} is locally free.

We introduce some convenient notation to check the main assertion of the theorem. Recall $\lambda_i : \mathbb{G}_m \rightarrow \mathbf{A}_0$ is the i -th axis of \mathbf{A}_0 . For example, λ_i may be expressed in n -many coordinates as follows:

$$\lambda_i(\mathbf{x}) = (1, \dots, \mathbf{x}, \dots, 1) \quad (4.2.13)$$

with \mathbf{x} at the coordinate corresponding to $1 \leq i \leq n$. The map $\tau : \mathbf{Y}^0 \rightarrow \mathbf{A}_0$ is abstractly an isomorphism of tori and we can therefore consider the 1-PS $\lambda_i : \mathbb{G}_m \rightarrow \mathbf{Y}^0$ defined by $\lambda_i := \tau^{-1} \circ \lambda_i$. We observe the following:

$$p \circ \lambda_i = \lambda_i, \quad (4.2.14)$$

$$\mathfrak{a} \circ (\lambda_1, \dots, \lambda_n) = (\theta_1, \dots, \theta_n) = \Theta. \quad (4.2.15)$$

Recall that $V = \text{Spec } \mathcal{O}_n$, denotes a formal neighbourhood of the origin. Let us check the isomorphism $L_n^+(\mathcal{R}|_V) \simeq L_n^+(\mathfrak{Y}_\Theta)$ by first evaluating at k -valued points. Since we have $d \cdot \Theta$ is a n -tuple of 1-PS's of \mathbf{A}_0 , we see that $\Theta(\dots, \mathbf{x}_i, \dots) = \Theta(\dots, \mathbf{y}_i^d, \dots) = (d\Theta)(\dots, \mathbf{y}_i, \dots)$ has become integral in $\{y_i\}$.

By (4.2.11), a section δ of \mathcal{R} over V , is first of all given by a local section δ_K over the generic point $\text{Spec } K_n$ of $\text{Spec } \mathcal{O}_n$. More precisely, this is firstly an element in $j(\mathfrak{g}(K))$. Whence this element may be written as

$$\delta_K = \text{Ad}\left(\mathfrak{a} \prod_{1 \leq i \leq n} \lambda_i(\mathbf{x}_i)\right)(x_K) = \text{Ad}\left(\prod_{1 \leq i \leq n} \theta_i(\mathbf{x}_i)\right)(x_K).$$

Therefore, the membership of δ in $\mathfrak{g}(B^+)$ (4.2.11) gets interpreted as follows:

$$\text{Ad}\left(\prod_{1 \leq i \leq n} \theta_i(\mathbf{x}_i)\right)(x_K) \in L\mathfrak{g}(B^+). \quad (4.2.16)$$

But this is exactly the Lie-algebra version of the conditions "lim $_{s \rightarrow 0}$ exists" condition for $\eta = (1, \theta)$ in the observations (2.3.5) and (2.5), for each individual \mathbf{x}_i . In our situation we have assumed $R = k$ and so by (4.2.15) the above is equivalent to the membership

$$\delta \in \mathfrak{Y}_\Theta. \quad (4.2.17)$$

This proves the assertion for k -valued points. The above proof goes through for all k -algebras because the underlying module structure is already defined on k , and so is Lie-bracket. This completes the proof of the theorem.

Q.E.D

COROLLARY 4.5. *Let notations be as in (4.4). Then $\lambda := \sum_{i \leq n} \lambda_i$ is a 1-PS of \mathbf{A}_0 defining the diagonal curve D whose limit point is the origin. Let \mathcal{D} be a formal neighbourhood of the origin in D . We get the following identification (see (3.1.4) for notations):*

$$L^+(\mathcal{R}|_{\mathcal{D}}) = L^+(\mathfrak{P}_{\mathfrak{o}}^{diag}). \quad (4.2.18)$$

of the specialized objects.

Let $\lambda = \sum k_i \lambda_i$ where $k_i \geq 0$ but not all are zero. Let \mathcal{D} be the formal neighbourhood of origin associated to the curve defined by λ . Let $f : \Phi \rightarrow \mathbb{Z}$ be the concave function defined by the assignment $r \mapsto \sum k_i m_r(\theta_i)$. Then for any k -algebra R we have

$$L^+(\mathcal{R}|_{\mathcal{D}})(R) = \langle \mathfrak{t}(R[[t]]), \mathfrak{u}_r(t^{f(r)} R[[t]]), r \in \Phi \rangle. \quad (4.2.19)$$

Proof. Note that restriction to \mathcal{D} followed by taking global sections of \mathcal{R} sets all variables z_i to t in the sections of \mathcal{R} on U . Now the first assertion follows by the definition (3.1.4). The restriction of \mathcal{R} to \mathcal{D} in the second case followed by taking global sections corresponds to setting $z_i := t^{k_i}$ in the sections of \mathcal{R} on U . Q.E.D

Remark 4.6. In the setting of Theorem 4.4, we may view $p : \text{Spec}(B^+) \rightarrow \text{Spec}(B_0^+)$ as a ramified covering space of affine toric varieties induced by an inclusion of lattices. The Galois group for this covering is the dual of the quotient of lattices. The computation in 4.4 can be seen as a higher dimensional analogue of [BS].

5. SCHEMATIZATION OF \mathfrak{n} -PARAHORIC GROUPS

In this section and the next couple of sections, we will freely use the theorems from SGA3 [SGAD], the paper of Bruhat-Tits ([BT2]) and M. Raynaud's book [Ray]. We begin by recalling a few essential facts about Weil restrictions of scalars and related things.

5.1. Weil restrictions and Lie algebras. Let X be an arbitrary k -scheme. For an affine (or possibly ind-affine) group scheme $\mathcal{H} \rightarrow X$, we denote $\text{Lie}(\mathcal{H})$ the sheaf of Lie-algebras on X whose sections on $U \rightarrow X$ are given by

$$\text{Lie}(\mathcal{H})(U) = \ker(\mathcal{H}(U \times k[\epsilon]) \rightarrow \mathcal{H}(U)). \quad (5.1.1)$$

LEMMA 5.1. *Let $p : \tilde{X} \rightarrow X$ be a finite flat map of noetherian schemes. Let $\text{Res}_{\tilde{X}/X}$ denote the "Weil restriction of scalars" functor. Let $\mathcal{H} \rightarrow \tilde{X}$ be an affine group scheme. Then we have a natural isomorphism*

$$\text{Lie}(\text{Res}_{\tilde{X}/X} \mathcal{H}) \simeq \text{Res}_{\tilde{X}/X} \text{Lie}(\mathcal{H}). \quad (5.1.2)$$

When p is also Galois with Galois group Γ , under tameness assumptions (see 1.1), we have a natural isomorphism

$$\text{Lie}((\text{Res}_{\tilde{X}/X} \mathcal{H})^\Gamma) \simeq (\text{Res}_{\tilde{X}/X} \text{Lie}(\mathcal{H}))^\Gamma. \quad (5.1.3)$$

Proof. See [CGP, Page 533] and [Ed, 3.1, page 293] respectively. Q.E.D

Notation 5.2. Following [BLR], we will denote the Weil restriction of scalars as p_* . The functor which composes with taking group invariants will be denoted by p_*^Γ . This is often called the "invariant direct image" functor following [BS] The conclusion of (5.1) can be expressed as:

$$\text{Lie}(p_*^\Gamma(\mathcal{H})) \simeq p_*^\Gamma(\text{Lie}(\mathcal{H})) \quad (5.1.4)$$

5.2. Some constructions from EGA. We recall briefly some facts about vector group schemes from [SGAD]. For a coherent S -module \mathcal{F} , we recall [SGAD, Exp I, 4.6, 4.6.3, 4.6.5] (see also [BT2, 1.4.1]), the functor from the category of S -schemes to abelian groups given by:

$$W : S' \mapsto \Gamma(S', \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{O}_{S'}). \quad (5.2.1)$$

When \mathcal{F} is locally free, this is represented by a smooth group scheme $W(\mathcal{F})$ defined by the affine scheme $\text{Sym}(\mathcal{F}^\vee)$. Its Lie algebra is canonically isomorphic to \mathcal{F} . In fact, there is a canonical isomorphism

$$W(\mathcal{F}) \simeq W(\text{Lie}(W(\mathcal{F}))). \quad (5.2.2)$$

Following [SGAD], we will call $W(\mathcal{F})$ a *vector group scheme*, noting that

$$\mathcal{F} \mapsto W(\mathcal{F}) \quad (5.2.3)$$

gives a functor from the category of locally free \mathcal{O}_S -modules of finite rank to the category of vector S -groups. Observe that over a field a vector group is simply \mathbb{G}_a^n for some n . We now state a few important properties satisfied by the functor W :

- (a) If \mathcal{F}' is locally free S -submodule of \mathcal{F} , then $W(\mathcal{F}')$ is a closed subgroup scheme of $W(\mathcal{F})$ if and only if \mathcal{F}' is a direct summand of \mathcal{F} (see [DG, page 148] and [BT2, 1.4.1]).
- (b) In particular, if $S = \text{Spec } \Lambda$ with fractional field F , a projective \mathcal{O}_S -module \mathcal{L} of rank 1 corresponds to an S -group scheme $W(\mathcal{L})$ whose generic fibre is isomorphic to the additive group $\mathbb{G}_{a,F}$.
- (c) It is easy to check that Weil restrictions and taking invariants commutes with the functor W .

5.3. Restriction of \mathcal{R} to formal neighbourhoods of generic points of coordinate hyperplanes H_i . Let C_i be the coordinate curve defined by the vanishing of λ_j §(4.2) for $j \neq i$. Let U_i be the formal neighbourhood of the origin in C_i . For $1 \leq i \leq n$, let θ_i be the point in the apartment corresponding to the hyperplane H_i .

For each θ_i , let $d_i \in \mathbb{N}$ be such that $d\theta_i \in Y(T)$. We first suppose that θ_i lies in the fundamental domain of $Y(T)$ in \mathcal{A}_T . Recall (see §12.2) that by [BS, Proposition 5.1.2] there exists a ramified cover $q : U'_i \rightarrow U_i$ of ramification index d_i and Galois group Γ , together with a Γ -equivariant G -torsor E_i such that the adjoint group scheme $\mathcal{H}_i = E_i(G)$ has simply-connected fibers isomorphic to G and we have the *canonical* identification of U -group schemes:

$$q_*^\Gamma(\mathcal{H}_i) \simeq \mathfrak{G}_{\theta_i}. \quad (5.3.1)$$

For the general case, move the points θ_i into the fundamental domain for the action of $Y(T)$ on \mathcal{A}_T by elements $h_i \in Y(T)$. These $h_i \in Y(T)$ are uniquely determined. By an abuse of notation, we may view them as $h_i \in T(K)$ as well. We now consider the modified embedding of G in the generic fibre of $\mathfrak{G}_{h_i, \theta_i}$ given by a conjugation by h_i . Then the *schematic closure* via this embedding in $\mathfrak{G}_{h_i, \theta_i}$ gives the group scheme \mathfrak{G}_{θ_i} . Now (5.3.1) generalises without any restriction on θ_i . Henceforth in this paper, we fix the identification of U -group schemes thus obtained:

$$\phi(h_i) : q_*^\Gamma(\mathcal{H}_i) \simeq \mathfrak{G}_{\theta_i}. \quad (5.3.2)$$

We therefore get the following useful corollary to (4.4).

COROLLARY 5.3. *As sheaves of Lie algebras, we have an isomorphism*

$$\mathrm{Lie}(\phi(h_i)) : q_*^\Gamma(E_i(\mathfrak{g})) \simeq \mathcal{R}|_{U_i}. \quad (5.3.3)$$

Proof. This is an immediate consequence of (4.4) and (5.1). Q.E.D

5.4. Group schemes on smooth varieties with normal crossing divisors. We now prove a result on constructing group schemes on varieties with normal crossing divisors under some general hypotheses. This plays a role in this paper as a useful substitute for the Artin-Weil theorem on existence of group scheme structures stemming from birational group laws. Let $\Theta := (\theta_1, \dots, \theta_n)$ be a n -tuple of rational points in the affine apartment \mathcal{A}_T satisfying conditions of §2.2. Let k be a *perfect field* with characteristic assumptions as in §2.2.

THEOREM 5.4. *Let X be a smooth quasi-projective scheme over k , and let $D = \sum_{i=1}^n D_i$ be a reduced normal crossing divisor. Let $\{\zeta_i\}_{i=1}^n$ be the set of generic points of the components $\{D_i\}_{i=1}^n$ and let $A_i = \mathcal{O}_{X, \zeta_i}$, $X_i := \mathrm{Spec}(\hat{A}_i)$ and let $X_o := X - D$. Let \mathfrak{G}_{θ_i} be the parahoric group scheme on X_i for each i associated to the point θ_i of the affine apartment. Let \mathcal{R} denote a Lie algebra bundle on X which is such that $\mathcal{R}|_{X_o} \simeq \mathfrak{g} \times X_o$ and $\mathcal{R}|_{X_i} \simeq \mathrm{Lie}(\mathfrak{G}_{\theta_i})$ for each i . Then there is a smooth, affine group scheme \mathfrak{G} on X with connected fibres, unique upto isomorphism, such that*

- (1) $\mathfrak{G}|_{X_o} \simeq G \times X_o$,
- (2) $\mathfrak{G}|_{X_i} \simeq \mathfrak{G}_{\theta_i}$, for each i ,
- (3) $\mathrm{Lie}(\mathfrak{G}) \simeq \mathcal{R}$.

Proof. Let $L_i := \mathrm{Fract}(A_i)$ (resp. $\hat{L}_i := \mathrm{Fract}(\hat{A}_i)$) for each i . By (5.3.2), for each i there exist a Γ_i -equivariant G -torsor E_i on a ramified cover $q_i : \tilde{X}_i \rightarrow X_i$ such that the adjoint group scheme $\mathcal{H}_i = E_i(G)$ has simply-connected fibers isomorphic to G and we have the identification of A_i -group schemes:

$$\phi(h_i) : q_{i,*}^\Gamma(\mathcal{H}_i) \simeq \mathfrak{G}_{\theta_i}. \quad (5.4.1)$$

Furthermore, by (5.3), we also get the identifications:

$$\mathrm{Lie}(\phi(h_i)) : q_{i,*}^\Gamma(E_i(\mathfrak{g})) \simeq \mathrm{Lie}(\mathfrak{G}_{\theta_i}). \quad (5.4.2)$$

Let $\mathfrak{G}_o := G \times X_o$. Let $X' \subset X$ be an open subset containing X_o whose complement in X has codimension ≥ 2 . We restrict the Lie algebra bundle $\mathcal{R} = \mathcal{R}_X$ further to X' . By assumption, over the open subset $X_o \subset X'$, we have $\mathcal{R} \simeq X_o \times \mathfrak{g}$. Observe that $\mathrm{Aut}(\mathfrak{g}) = \mathrm{Aut}(G)$ since G is simply connected. Further, since the local parahoric group schemes \mathfrak{G}_{θ_i} are assumed to be *generically split*, it follows that the transition functions of \mathcal{R} give for each i an isomorphism

$$\tau_i : \mathfrak{G}_o \times_{X_o} \hat{L}_i \simeq \mathfrak{G}_{\theta_i} \times_{X_i} \hat{L}_i$$

We can now apply [BLR, 6.2, Proposition D.4(b)], to each $\tau_i : \mathfrak{G}_o \times_{X_o} L_i \times_{L_i} \hat{L}_i \simeq \mathfrak{G}_{\theta_i} \times_{X_i} \hat{L}_i$ to obtain by “descent” an affine, finite type group scheme \mathfrak{G}'_{θ_i} over $\mathrm{Spec}(A_i)$ for each i , such that $\mathfrak{G}'_{\theta_i} \times_{A_i} \hat{A}_i \simeq \mathfrak{G}_{\theta_i}$ and $\mathfrak{G}'_{\theta_i} \times_{A_i} L_i \simeq \mathfrak{G}_o \times_{X_o} L_i$. *By an abuse of notation, we will denote the group scheme \mathfrak{G}'_{θ_i} simply by \mathfrak{G}_{θ_i} .*

Since the group schemes \mathfrak{G}_{θ_i} on $\mathrm{Spec}(A_i)$ are of finite type, they can be extended (by clearing denominators) to open sub-schemes X_{f_i} of X . By a further shrinking of this neighbourhood X_{f_i} of ζ_i one can glue it to \mathfrak{G}_o along the intersection. Since the

group schemes are smooth and affine, we obtain the connected, smooth, affine group scheme

$$\mathfrak{G}' \rightarrow X'. \quad (5.4.3)$$

Transporting structures by (5.3) (see also §12), the Lie algebra bundle \mathcal{R} gets canonical parabolic structures at the generic points ζ_i of the components $\{D_i\}_i$. By (12.1), for a (generic) parabolic vector bundle with prescribed rational weights such as \mathcal{R} , we firstly get a global Kawamata cover (12.1) $p: Z \rightarrow X$ ramified over D with ramification prescribed by the weights $\{d_i\}$ (2.1.2), with Galois group Γ which “realizes the local ramified covers q_i at the points ζ_i ” i.e. the isotropy subgroup of Γ at ζ_i is Γ_i .

We *claim* that we in fact get an Γ -equivariant vector bundle V on Z such that

$$p_*^\Gamma(V) \simeq \mathcal{R}. \quad (5.4.4)$$

Let $Z' := p^{-1}(X')$. As the first step, following the reasoning in §12, it follows by the dimension 1 strategy that there exists a Γ -equivariant vector bundle V' on Z' such that $p_*^\Gamma(V') \simeq \mathcal{R}|_{X'}$.

Furthermore, in formal neighbourhoods over the generic points ζ_i of the components D_i , we may identify V' with $E_i(\mathfrak{g})$ and Γ with Γ_i . Also over $Z_o := p^{-1}(X_o)$ we can identify V' with $\mathfrak{g} \times Z_o$. The transition functions of $\mathcal{R}|_{X'}$ can be lifted to give transition functions for V' and we get a canonical Γ -equivariant Lie algebra bundle structure on V' over Z' (with fibre type \mathfrak{g}) such that $p_*^\Gamma(V') \simeq \mathcal{R}|_{X'}$ as Lie algebra bundles.

Since the weights θ_i are all chosen to lie in the same affine apartment \mathcal{A}_T , the local representations for the stabilizers $\Gamma_i \subset \Gamma$ in G take values in the maximal torus T . This determines a uniform Cartan decomposition on the fibres of the Lie algebra bundle V' on Z' (compare the proof of Theorem 4.4, where one has similar arguments to check that \mathcal{R} is locally-free on \mathbf{A}). This gives a decomposition of the underlying vector bundle V' as a direct sum of line bundles.

Now since invertible sheaves extend as invertible sheaves across codimension ≥ 2 (4.3), and since $\text{codim}(Z \setminus Z') \geq 2$, the reflexivity of the double dual V of V' implies that this direct sum decomposition of V' extends term by term to Z . Whence the double dual V is locally free. The equivariant structure on V' canonically gives an equivariant structure on V and it is easy to check that (5.4.4) holds. This completes the proof of the claim.

The structure of Lie algebra bundle on V' is such that the Killing form is non-degenerate everywhere on Z' . By a Hartogs-type argument (see for example [EGA IV, Theorem 5.10.5] or [H1, Theorem 3.8, page 44]), the Lie bracket $[\cdot, \cdot]$ on V' extends to a Lie bracket on the locally free sheaf V with the Killing form being non-degenerate on the whole of Z . *In other words, V is now a locally free sheaf of Lie algebras on the whole of Z with semisimple fibres; these fibres are isomorphic to the Lie algebra \mathfrak{g} by the rigidity of semisimple Lie algebras.* We denote this bundle of Lie algebras by $V_{\mathfrak{g}}$.

By construction V' has the structure of an equivariant Lie algebra bundle. The $[\cdot, \cdot]$ on $V_{\mathfrak{g}}$ is a section of $V_{\mathfrak{g}} \otimes V_{\mathfrak{g}}^* \otimes V_{\mathfrak{g}}^*$. For each $g \in \Gamma$, the sections $g.[\cdot, \cdot]$ and $[\cdot, \cdot]$ agree on Z' . Hence they agree everywhere on Z , i.e. the underlying equivariant structure on $V_{\mathfrak{g}}$ gives $V_{\mathfrak{g}}$ the structure of an *equivariant Lie algebra bundle*. The transition functions of this equivariant Lie algebra bundle lies in $\text{Aut}(\mathfrak{g}) = \text{Aut}(G)$. Hence we obtain an equivariant $\text{Aut}(G)$ -torsor on Z . Note that \mathcal{E} is only étale locally trivial. Observe also that the associated Lie algebra bundle $\mathcal{E}(\mathfrak{g}) := \mathcal{E} \times^{\text{Aut}(G)} \mathfrak{g}$ is isomorphic to $V_{\mathfrak{g}}$ as Lie algebra bundles.

Let $\mathcal{H} := \mathcal{E} \times^{\text{Aut}(G)} G$ denote the associated fibre space for the canonical action of $\text{Aut}(G)$ on G . Since the group $\text{Aut}(G)$ acts on G as group automorphisms, it preserves the group structure and hence \mathcal{H} is a group scheme. Further, since \mathcal{E} is étale locally trivial, by étale descent, it follows that \mathcal{H} is an *affine* smooth group scheme on Z . Since \mathcal{E} is an equivariant $\text{Aut}(G)$ -torsor, the group scheme \mathcal{H} is furthermore an equivariant group scheme on Z for the action of the Galois group Γ and so we define

$$\mathfrak{G}_X := p_*^\Gamma(\mathcal{H}). \quad (5.4.5)$$

By the properties of Weil restriction of scalars and taking invariants, it follows that \mathfrak{G}_X is a *smooth, affine group scheme*. Also $\mathfrak{G}_X|_{X'} = \mathfrak{G}'$.

Note that we have the identification $\text{Lie}(\mathcal{H}) \simeq V_{\mathfrak{g}}$ as Lie algebra bundles on Z . Since by (5.1), the “invariant direct image” functor commutes with taking Lie algebras, we moreover get isomorphisms of locally-free sheaves of Lie algebras

$$\text{Lie}(\mathfrak{G}_X) \simeq \text{Lie}(p_*^\Gamma(\mathcal{H})) \simeq p_*^\Gamma(\text{Lie}(\mathcal{H})) \simeq p_*^\Gamma(V_{\mathfrak{g}}) \simeq \mathcal{R}. \quad (5.4.6)$$

This proves the third claim in the theorem.

Q.E.D

5.5. The “big cell” structure on \mathfrak{G} . Recall that the choice of the set of positive roots Φ^+ determines a *big cell* of G over k .

By (5.4.3), the group scheme $\mathfrak{G}_{X'}$ is obtained by gluing the parahoric group schemes \mathfrak{G}_{θ_i} with $G \times X_o$ using the identity map id_G . Therefore, the parahoric group schemes \mathfrak{G}_{θ_i} come with a canonical *big cell* which glues with the standard big cell of the split group scheme $G \times X_o$. This gives a big cell $\mathfrak{B}_{X'} = \mathfrak{U}_{X'}^+ \times \mathcal{T}_{X'} \times \mathfrak{U}_{X'}^- \subset \mathfrak{G}_{X'}$. This also gives the decomposition:

$$\text{Lie}(\mathfrak{G}_{X'}) = \text{Lie}(\mathfrak{U}_{X'}^+) \oplus \text{Lie}(\mathcal{T}_{X'}) \oplus \text{Lie}(\mathfrak{U}_{X'}^-). \quad (5.5.1)$$

In the next proposition we show the existence of an open subscheme \mathfrak{B}_Φ of \mathfrak{G} obtained in (5.4). This open subscheme is analogous to the *big cell in a split reductive group scheme* [SGAD, 4.1, Expose XXII]. Following [BT2], we will call this open subscheme a “*big cell*” of \mathfrak{G} . The *big cell* in particular provides a neighbourhood of the identity section of the group scheme \mathfrak{G} .

PROPOSITION 5.5. *We work in the setting of (5.4) and make the choice of transition function for the gluing to be the $\text{id}_G \in \text{Aut}(G)$. Let Φ^+ be a system of positive roots in Φ . Let $\mathfrak{B}_{\mathcal{H}}$ denote the big cell of \mathcal{H} . Let*

$$\mathfrak{B}_\Phi := p_*^\Gamma(\mathfrak{B}_{\mathcal{H}})$$

denote the invariant direct image of $\mathfrak{B}_{\mathcal{H}}$.

Then \mathfrak{B}_Φ restricts to the standard big cell of G over the open subset X_o . In fact, there exist unipotent subgroup schemes \mathfrak{U}^\pm and a toral subgroup scheme \mathcal{T} of \mathfrak{G} such that the morphism

$$\mathfrak{U}^- \times \mathcal{T} \times \mathfrak{U}^+ \rightarrow \mathfrak{G} \quad (5.5.2)$$

induced by multiplication to \mathfrak{G} is an open immersion with image \mathfrak{B}_Φ .

Proof. Notations are as in the proof of (5.4). Since the Galois action of Γ is via the fixed maximal torus T , we observe that the Cartan decomposition of the Lie algebra bundle is preserved under the functor that takes Γ -invariants. Notice that the functor W of §5.2 commutes with Weil restrictions and taking invariants. Hence the root group schemes and the toral subgroup scheme of the reductive group scheme \mathcal{H} can be

pushed down by p_*^Γ to give the corresponding root group schemes and toral group scheme below. The group structure on \mathcal{H} induces one on the product of the root groups for the positive and negative roots (with a fixed and prescribed order) to give closed unipotent subgroup schemes on \mathcal{H} . These again carry the action of the Galois group Γ .

Hence these unipotent subgroup schemes gets pushed down by p_*^Γ to give closed subgroup schemes \mathcal{U}^\pm of \mathfrak{G} . Further, Weil restrictions and taking invariants preserves the product structure and this shows that the *big cell* structure on the reductive group scheme \mathcal{H} goes down by Weil restriction and invariants, to give the required *big cell* structure on \mathfrak{G} . Q. E. D

5.6. Bruhat-Tits group scheme associated to concave functions. Let $\tilde{\Phi} := \Phi \cup \{0\}$. Recall that a function $f : \tilde{\Phi} \rightarrow \mathbb{R}$ is a concave function such that $f(0) = 0$, (1.0.5). We recall a few salient features of the main results from [BT2, 4.6.2, page 124] and [Yu, Theorem 0.1] in our special setting, where G_K is a generically split, simple, simply connected group over K , i.e. $G_K \simeq G \times \text{Spec}(K)$.

Bruhat-Tits [BT2, 4.6.2, page 124]: Let \mathcal{P}_f be the subgroup of $G(K)$ generated by $U_{r,f(r)}$ and the subgroup $T(A)$.

There exists a canonical affine smooth group scheme \mathfrak{G}_f on $\text{Spec } \mathcal{O}$ with generic fibre G_K such that:

- (1) $\mathfrak{G}_f(A) = \mathcal{P}_f$
- (2) the multiplication morphism:

$$\left(\prod_{r \in \Phi^-} \mathcal{U}_{r,f(r)} \right) \times \mathcal{T}_{f,\mathcal{O}} \times \left(\prod_{r \in \Phi^+} \mathcal{U}_{r,f(r)} \right) \rightarrow \mathfrak{G}_f \quad (5.6.1)$$

is an open immersion. The group scheme $\mathcal{T}_{f,\mathcal{O}}$ is the schematic closure of T_K in \mathfrak{G}_f and since T_K is a split torus, $\mathcal{T}_{f,\mathcal{O}}$ is a connected multiplicative group scheme [BLR, Example 5, page 291]. By computing the tangent space at the identity section which lies in the image of the immersion, i.e., in the *big cell* \mathfrak{B}_f of \mathfrak{G}_f , we see that:

$$\text{Lie}(\mathfrak{G}_f) = \bigoplus_{r \in \Phi} \text{Lie}(\mathcal{U}_{r,f(r)}) \oplus \text{Lie}(\mathcal{T}_{f,\mathcal{O}}) \quad (5.6.2)$$

5.7. On the Lie algebra bundle \mathcal{R} in (5.4). We now prove a small but significant variant of (5.4). In the hypothesis in (5.4), we have assumed the existence of a n -parahoric Lie algebra bundle \mathcal{R} on X . We will give sufficient general conditions for the existence of a Lie algebra bundle alone.

PROPOSITION 5.6. *Let the notations be as in (5.4). Let $\mathbf{f} = \{f_j\}_{j=1}^n$ be a collection of n -concave functions. Let \hat{A}_i be the completion of A_i at the maximal ideal. Suppose that we are given an assignment of BT-group schemes \mathfrak{G}_{f_j} on $X_j := \text{Spec } \hat{A}_j$ for each j . Then there exists a Lie algebra bundle \mathcal{R} on (X, D) such that $\mathcal{R}|_{X_o} \simeq \mathfrak{g} \times X_o$, the Cartan decomposition on \mathfrak{g} extends to \mathcal{R} and $\mathcal{R}|_{X_j} \simeq \text{Lie}(\mathfrak{G}_{f_j})$ for each j .*

Proof. The notations are as in (5.4) and its proof. But unlike (5.4), we do not begin with a Lie algebra bundle. So we are free to choose the initial gluing data according to our requirements. Since the BT-group schemes for us are always, connected, smooth and generically split, we can choose (as in (5.5)) the id_G as the maps for gluing the product group scheme $\mathfrak{G}_o = G \times X_o$ with \mathfrak{G}_{f_i} .

Exactly as in the proof of (5.4), by a descent argument, we get smooth, affine group schemes \mathfrak{G}_{f_i} now on $\text{Spec } A_i$ for each i and hence we get a smooth, affine group scheme \mathfrak{G}' on X' . We now define:

$$\mathcal{R}' := \text{Lie}(\mathfrak{G}') \quad (5.7.1)$$

which is a Lie algebra bundle on $X' \subset X$.

Since the gluing functions are id_G , exactly as in the proof of (5.5), we also have an extended *big cell* structure on the group scheme \mathfrak{G}' over the whole of X' . As in (5.5.1), we see that the Cartan decomposition extends to give one for \mathcal{R}' . In particular this gives a direct sum decomposition of \mathcal{R}' in terms of line bundles.

Now define \mathcal{R} to be the reflexive closure of \mathcal{R}' (4.2). Since we are on an irreducible smooth scheme X , by (4.3), reflexive closures of invertible sheaves remain invertible sheaves. Thus \mathcal{R} is also a locally free sheaf on X . The Lie bracket on \mathcal{R}' now extends by a Hartogs type argument and we get a Lie algebra bundle structure on \mathcal{R} with the desired properties.

Q. E. D

5.8. Construction of the n-parahoric group scheme. The notations in this section are as in the previous one, namely $\mathbf{A} \simeq \mathbb{A}^n$ and $\mathbf{A}_0 \subset \mathbf{A}$ is the complement of the coordinate hyperplanes. Recall that $\mathcal{O}_n = k[[z_1, \dots, z_n]]$ and $K_n = \text{Fract}(\mathcal{O}_n)$.

Let $\{H_i\}$ be the hyperplanes of \mathbf{A} . Let ζ_i 's denote the generic points of the divisors H_i 's. Let

$$A_i = \mathcal{O}_{\mathbf{A}, \zeta_i} \quad (5.8.1)$$

be the dvr's obtained by localizing at the height 1-primes given by the ζ_i 's and let $X_i := \text{Spec}(A_i)$. Base changing by the local morphism $X_i \rightarrow \mathbf{A}$, we have a Lie algebra bundle $\mathcal{R}|_{X_i}$ for each i . Moreover, the Lie algebra bundle \mathcal{R} on \mathbf{A} (4.4) gives canonical gluing data to glue $\mathcal{R}|_{X_i}$ with the trivial bundle $\mathfrak{g} \times \mathbf{A}_0$.

The following first case is the key to proving the main result (1.3).

THEOREM 5.7. *The main theorem (1.3) holds when the n-tuple of concave functions is of type I (1.2).*

Proof. We begin by making constructions on \mathbf{A} . The existence of the Lie algebra bundle \mathcal{R} on \mathbf{A} follows by applying (5.6) to the n-concave function of type I defined by the point $\Theta = (\theta_1, \dots, \theta_n)$. Since we are in the type I case, we could have obtained the Lie algebra bundle directly from (4.4) as well. Again since we are in type I, the existence of the smooth *affine* group scheme \mathfrak{G}_Θ with connected fibres, is a consequence of (5.4); its cell structure and properties (4) and (5) of (1.3) follow from (5.5).

We now restrict the group scheme \mathfrak{G}_Θ to $\text{Spec } \mathcal{O}_n \subset \mathbf{A}$ and continue to denote the restricted group scheme by \mathfrak{G}_Θ . Since the base is smooth and the group scheme is also smooth, by a theorem of Weil [BLR, Theorem 1, p 109], it follows that the global sections are determined on any open subset U whose complement is of codimension ≥ 2 . Let us call such an open subset a *big open subset*. Sections over U are obtained by intersecting the sections of the parahoric groups defined at the height 1 primes in $G(K_n)$. It follows immediately that the sections over the big open subset U coincides with the n-parahoric group P_Θ proving (2) of (1.3).

We now discuss *the uniqueness of the group scheme*; we note that although we do this here in the setting of type I, the arguments are identical and will imply the same for the most general type III case, once the construction along with the properties (4) and (5) of (1.3) is carried out in §7.

If \mathfrak{H} is a connected, affine smooth group scheme over \mathcal{O}_n which is generically split and such that it has closed toral and root subgroup schemes as in (4) and a *big cell* as in (5) of (1.3), then by, [BT2, 1.2.13, 1.2.14] we get the desired isomorphism of $\mathfrak{G}_\mathfrak{o}$ and \mathfrak{H} , showing the uniqueness.

A subtler feature regarding the uniqueness can be extracted from these arguments. Once the smooth connected group scheme $\mathfrak{G}_\mathfrak{o}$ is abstractly constructed, the big cell structure over the big open subset U (which contains all points of height 1) can be obtained at each of the height 1 primes as in [BT2]. So the cell structure coincides with the *big cell* of the parahoric group schemes \mathfrak{G}_{θ_j} for each j and hence by the uniqueness of the group schemes over discrete valuation rings [BT2, 3.8.3], we get the identification of the group schemes $\mathfrak{G}_\mathfrak{o}$ and \mathfrak{H} over U . Now, we can appeal to the deeper results in [Ray, Cor IX, 1.5, page 130] to extend the isomorphism of group schemes from U to $\text{Spec } \mathcal{O}_n$.

Q.E.D

Remark 5.8. In continuation of (5.6), the description of the diagonal restriction of $\mathfrak{G}_\mathfrak{o}$ can be obtained directly from the facts that it is affine by (5.7) and the group $\mathfrak{G}_\mathfrak{o}(\mathcal{O}_n)$ of \mathcal{O}_n -valued points is the n -parahoric group $P_\mathfrak{o}$. Let notations be as in (1.5). Let $\Delta := \mathbb{A}_I$ for $I = \{1, \dots, n\}$ be the diagonal curve in the affine space and \mathcal{O} be the ring of functions on Δ at the origin. Hence the restriction $\mathfrak{G}_\mathfrak{o}|_\Delta$ is such that the space of its sections $\mathfrak{G}_\mathfrak{o}|_\Delta(\mathcal{O})$ is identified with $\mathfrak{G}_\mathfrak{o}(\mathcal{O}_n)$ in which we set all variables as equal. Thus we get the subgroup $P_\mathfrak{o}^{\text{diag}} \subset G(k((t)))$ (3.1.2). This forces by the étouffé property of BT-group schemes that $\mathfrak{G}_\mathfrak{o}|_\Delta$ is the BT-group scheme given by the concave function $m_\mathfrak{o}$ where $m_\mathfrak{o}(r) := \sum_i m_r(\theta_i)$. We remark that the above discussion holds good if we work with sub-diagonal curves \mathbb{A}_I as (1.5)(b), which is defined by any sub-collection of the hyperplanes indexed by a non-empty $I \subset \{1, \dots, n\}$.

COROLLARY 5.9. *Consider a n -tuple \mathbf{f} of concave functions each of which is a sum of those of type I. Then the group scheme $\mathfrak{G}_\mathbf{f}$ is affine.*

Proof. Let $\mathbf{f} = (f_1, \dots, f_n)$ where $f_i = \sum_j f_{\theta_{ij}}$. Set $\Theta = (\theta_{ij})$. Then $\mathfrak{G}_\mathfrak{o}$ is affine by (5.7). Note that this group scheme constructed out of N -parahoric group schemes is over an affine space of dimension N , for $N \gg 0$ and obtained from the data of the $\{\theta_{ij}\}$'s.

Observe that the group scheme $\mathfrak{G}_\mathbf{f}$ can be seen as a restriction to a sub-diagonal in the larger affine space of $\mathfrak{G}_\mathfrak{o}$. Now the assertion of affineness follows by (5.8). Q.E.D

5.9. Revisiting two «contre-exemples» in Bruhat-Tits [BT2]. In this subsection we take a closer look at a couple of counter-examples in Bruhat-Tits [BT2, 3.2.15]. The examples highlight how the procedure developed in §3 of [BT2] of taking schematic closures in certain linear groups could give schemes which are neither *flat* nor *group schemes*. This phenomenon happens when the base scheme is \mathbb{A}^2 . By discarding certain bad components in the fibre over $o \in \mathbb{A}^2$ they extract a 2BT-group scheme whose *affineness* is left in doubt.

The base ring $\mathcal{O} = k[\xi, \eta]$, the group is $G = \text{SL}_2$, the concave function f is defined by $f(a) = \xi \cdot \mathcal{O}$ and $f(-a) = \eta \cdot \mathcal{O}$, the representation $\rho = \text{id} \times \text{Ad}$ on the module $K^2 \times \mathfrak{g}$. The \mathcal{O} -module M is a free module generated by the basis of K^2 and the elements

$\eta.e_{-a}, h, \xi.e_a$. We view them as $\begin{pmatrix} 0 & 0 \\ -\eta & 0 \end{pmatrix}, \begin{pmatrix} 1/2 & 0 \\ 0 & -1/2 \end{pmatrix}$ and $\begin{pmatrix} 0 & \xi \\ 0 & 0 \end{pmatrix}$. Thus we have:

$$\rho \begin{pmatrix} x & y \\ z & t \end{pmatrix} = \begin{pmatrix} x & y \\ z & t \end{pmatrix}, \begin{pmatrix} t^2 & -\eta.zt & \xi\eta^{-1}z^2 \\ -2\eta.yt & xt+yz & -2\xi.xz \\ \eta.\xi^{-1}y^2 & -\xi.xy & x^2 \end{pmatrix} \quad (5.9.1)$$

The schematic closure of $\rho(G)$ in $\mathfrak{GL}(M)$ is \mathfrak{G} . The fibre over $o \in \mathbb{A}^2$ is:

$$\mathfrak{G}_o = \mathfrak{G}_o^1 \cup \mathfrak{G}_o^2 \quad (5.9.2)$$

where \mathfrak{G}_o^1 has dimension 4 and \mathfrak{G}_o^2 dimension 3. Hence \mathfrak{G} is not flat over \mathcal{O} , while $\mathfrak{H} := \mathfrak{G} \setminus \mathfrak{G}_o^1$ is a *flat group scheme* albeit in the words of [BT2, page 60], «*mais probablement pas affine*», i.e. *but probably not affine*.

The fibre of \mathfrak{H} at o is obtained from [BT2, 3.2.14(2)], which is obtained by a diagonal restriction.

We now reinterpret this example as the simplest case of Theorem (5.7), for $G = \mathrm{SL}_2$. The surprising picture which emerges from our approach is that we even get the *affineness* of their group scheme \mathfrak{H} as a consequence.

In our setting of (3.2) we have $\Phi^+ = \{a\}$ and in terms of the alcove vertices (2.1.1) we have $\theta_1 := \theta_a/2 = \omega^\vee/2$ and $\theta_2 := -\theta_a/2 = -\omega^\vee/2$. With $m_r(\theta) := -\lfloor r(\theta) \rfloor$, we get,

$$a(\theta_1) = \begin{cases} 1/2 & \text{if } a \in \Phi^+ \\ -1/2 & \text{if } a \in \Phi^- \end{cases} \quad (5.9.3)$$

$$m_a(\theta_1) = \begin{cases} 0 & \text{if } a \in \Phi^+ \\ 1 & \text{if } a \in \Phi^- \end{cases} \quad (5.9.4)$$

Similarly for θ_2 . We get:

$$a(\theta_2) = \begin{cases} -1/2 & \text{if } a \in \Phi^+ \\ 1/2 & \text{if } a \in \Phi^- \end{cases} \quad (5.9.5)$$

$$m_a(\theta_2) = \begin{cases} 1 & \text{if } a \in \Phi^+ \\ 0 & \text{if } a \in \Phi^- \end{cases} \quad (5.9.6)$$

By (5.7), if $\theta = (\theta_1, \theta_2)$, the group scheme \mathfrak{G}_θ is precisely \mathfrak{H} and also a *smooth, affine* group scheme over \mathcal{O} . Furthermore, the diagonal restriction is the affine group scheme $\mathfrak{G}_{\mathfrak{m}_\theta}$ given by the concave function: $\mathfrak{m}_\theta : r \mapsto \sum m_r(\theta_j)$:

$$\mathfrak{m}_\theta(a) = 1 \quad \text{for all } a \in \Phi, \quad (5.9.7)$$

which computes \mathbb{P}_θ^{diag} and recovers the concave function considered in [BT2, 3.2.14].

6. SCHEMATIZATION OF \mathfrak{n} -BOUNDED GROUPS \mathbb{P}_Ω

The notations in this section are as in the previous one, namely $\mathbf{A} \simeq \mathbb{A}^n$ and $\mathbf{A}_0 \subset \mathbf{A}$ is the complement of the coordinate hyperplanes. In this entire section we work with G being one of A_n, G_2, F_4 , or E_6 .

We begin by remarking that in the foundational paper, [BT1] Bruhat and Tits study bounded groups associated to concave functions on the $\Phi \cup \{0\}$. In the sequel [BT2], the main result is to prove that these groups are *schematic*.

A class of concave functions, which play a basic role even in the general study, are the ones which come associated to bounded subsets Ω of the affine apartment. In this section, we extend the above theory of \mathfrak{n} -parahoric groups (with their schematic

properties) to the bounded groups associated to products of bounded regions $\Omega := (\Omega_1, \dots, \Omega_n)$ in the apartment.

In the following key remark we recall a few basic facts from [BT3].

Remark 6.1. Let $G = SL(n)$ for some n . Let Ω be a non-empty *enclosed* bounded subset of the apartment \mathcal{A} . It is the convex hull of a finite subset $\{\theta_1, \dots, \theta_s\}$ of points in \mathcal{A} . In terms of the points θ_j 's, the BT-group scheme \mathfrak{G}_Ω on $\text{Spec } \mathcal{O}$ is precisely the schematic closure of the image of G in $\prod_{j=1}^s \mathfrak{G}_{\theta_j}$ under the diagonal map. This is a consequence of the discussion in [BT3, 3.9, 1,2,3], together with [BT3, §5, 5.3], where it is shown that the group scheme \mathfrak{G}_Ω over $\text{Spec } \mathcal{O}$ can be realised as the *closed image* in $\prod_{j=1}^s \mathfrak{G}_{\theta_j}$ of the diagonal morphism of G in the generic fibre $G^\mathfrak{s}$ of $\prod_{j=1}^s \mathfrak{G}_{\theta_j}$.

We further remark that the stated result for $SL(n)$ also works by the results [GY1, Theorem 10.1] for G_2 and [GY2, Theorem 9.1] for F_4 and E_6 .

Remark 6.2. An important aspect of the remark made above which will be used in what follows is to view the group scheme $\prod_{j=1}^s \mathfrak{G}_{\theta_j}$ as an *invariant direct image* from a suitable ramified cover (see §12.2). Now the group scheme \mathfrak{G}_{θ_j} arises from an invariant direct image if θ_j belongs to the alcove \mathfrak{a}_0 (see §2.1) by [BS]. This generalizes to the case θ_j belongs to the fundamental domain of $Y(T)$ by (8.1) and §9.1. For the general case, we may have to move the points $\{\theta_1, \dots, \theta_s\}$ of points in apartment \mathcal{A} into the fundamental domain for the action of $Y(T)$. Let $h_j \in T(K)$ be elements such that the point $\{h_1.\theta_1, \dots, h_s.\theta_s\}$ now lies in the s -fold product of the fundamental domain of $Y(T)$. We now consider the modified diagonal embedding of G in the generic fibre of $\prod_{j=1}^s \mathfrak{G}_{h_j.\theta_j}$ which is given by a simultaneous conjugation by the h_j 's, i.e., $g \mapsto (h_1.g.h_1^{-1}, \dots, h_s.g.h_s^{-1})$. Then the *schematic closure* via this embedding in $\prod_{j=1}^s \mathfrak{G}_{h_j.\theta_j}$ gives the group scheme \mathfrak{G}_Ω upto an isomorphism. Moreover, the new ambient group scheme $\prod_{j=1}^s \mathfrak{G}_{h_j.\theta_j}$ comes as an invariant direct image from a single cover.

6.1. The schematic hull $\mathbb{S}_{\Omega, \Lambda}$ (6.1.1) of n -bounded set Ω and its big-cell structure. For simplicity of notation, let Λ stand for the polynomial ring $k[z_1, \dots, z_n]$, and F stand for $\text{Fract}(\Lambda)$, throughout this proof.

Let $\Omega = (\Omega_1, \dots, \Omega_n)$, where we may assume that Ω_i are *enclosed* bounded subsets of the affine apartment \mathcal{A} . Let Ω_i be the convex hull of points $\{\theta_{i1}, \dots, \theta_{is}\}$ of \mathcal{A}_T . Note that *a priori* the number s of points also varies with i . By (6.1) and (6.2), we may suppose that

- (1) these points lie in a single fundamental domain of $Y(T)$ in \mathcal{A}_T and
- (2) that over a dvr $\text{Spec } \mathcal{O}$, we realise each \mathfrak{G}_{Ω_i} as the schematic closure of G in

$$\prod_{j=1}^s \mathfrak{G}_{\theta_{ij}}.$$

By choosing s to be the maximum of terms for varying $i = 1, \dots, n$, and by repeating the θ_{ij} 's when the number is lesser than s , we may assume that all the group schemes \mathfrak{G}_{Ω_i} are schematic closure of a product of s parahoric group schemes on $\text{Spec } \mathcal{O}$.

Fix a $j \in \{1, \dots, s\}$. For each hyperplane H_i ($i = 1, \dots, n$), we attach the point θ_{ij} . For this configuration of points in the apartment, we have by Theorem 5.7, a n -parahoric group scheme $\mathfrak{G}_{\theta_{1j}, \dots, \theta_{nj}}$ on \mathbf{A} . We set

$$\mathbb{S}_{\Omega, \Lambda} := \prod_{1 \leq j \leq s} \mathfrak{G}_{\theta_{1j}, \dots, \theta_{nj}} \tag{6.1.1}$$

to be the s -fold product of n -parahoric group schemes on \mathbf{A} . Thus $\mathbb{S}_{\Omega, \Lambda}$ is a *smooth, affine group scheme* on $\text{Spec } \Lambda = \mathbf{A}$ with generic fibre being the split group scheme $G^s \times \text{Spec } F$. Since each n -parahoric group scheme $\mathfrak{G}_{\theta_{1j}, \dots, \theta_{nj}}$ is realized as a Weil restriction of scalars from a finite flat cover of \mathbf{A} (5.7), so is the group scheme $\mathbb{S}_{\Omega, \Lambda}$.

DEFINITION 6.3. *We call the product group scheme $\mathbb{S}_{\Omega, \Lambda}$ (6.1.1) the schematic hull of the n -bounded set Ω and will denote it simply as \mathbb{S}_Λ .*

This group scheme \mathbb{S}_Λ will play a central role in all that follows.

Recall each of the n -parahoric group schemes $\mathfrak{G}_{\theta_{1j}, \dots, \theta_{nj}}$ occurring in (6.1.1) is equipped with a *big cell* by (5.7) (see §5.5). Taking their product, we define the *big cell* of the group scheme \mathbb{S}_Λ . Let us denote it as the image of the product morphism

$$\mathfrak{B}^- \times \mathcal{T} \times \mathfrak{B}^+ \rightarrow \mathbb{S}_\Lambda. \quad (6.1.2)$$

Let $\mathfrak{U}_{r,j}$ denote the root group of $\mathfrak{G}_{\theta_{1j}, \dots, \theta_{nj}}$ corresponding to $r \in \Phi$. For $r \in \Phi$, let us set

$$\mathfrak{B}_r := \prod_{1 \leq j \leq s} \mathfrak{U}_{r,j}. \quad (6.1.3)$$

Then each \mathfrak{B}_r under the induced group law from \mathbb{S}_Λ is abelian.

6.2. The diagonal embedding. Let U be an open subset of $\text{Spec } \Lambda$ which contains all primes of height ≤ 1 . At the completion of the local rings at the height 1 prime ideal associated to the generic point of each H_i indexed by $i = 1, \dots, n$, we assign the Bruhat-Tits group scheme \mathfrak{G}_{Ω_i} . By definition, these group schemes are also generically split.

We let $\mathfrak{G}_{\Omega, F}$ denote the *image of the diagonal embedding* of $G \times \text{Spec } F$ in $G^s \times \text{Spec } F$. As in (5.6) and in the proof of (5.4), by gluing (using the map id_G) with the \mathfrak{G}_{Ω_j} , the group scheme $\mathfrak{G}_{\Omega, F}$ extends to each of the generic points of the coordinate hyperplanes. Let $\mathfrak{G}_{\Omega, U}$ denote this smooth, affine group scheme on U . We begin with the closed embedding $\mathfrak{G}_{\Omega, F} \subset \mathbb{S}_F$, where $\mathbb{S}_F = \mathbb{S}_\Lambda|_{\text{Spec } F}$. Let

$$\mathfrak{Y}_\Lambda := \overline{\mathfrak{G}_{\Omega, F}} \quad (6.2.1)$$

denote the *schematic closure* of $\mathfrak{G}_{\Omega, F}$ in \mathbb{S}_Λ .

LEMMA 6.4. *The group scheme $\mathfrak{G}_{\Omega, U}$ is in fact the schematic closure \mathfrak{Y}_U of $\mathfrak{G}_{\Omega, F}$ in \mathbb{S}_U .*

Proof. Since U contains all primes of height ≤ 1 , we see that the schematic closure \mathfrak{Y}_U of $\mathfrak{G}_{\Omega, F}$ in \mathbb{S}_U is in fact a *flat* group scheme on U and since \mathbb{S}_U is affine, it is in fact *affine* as well.

Since $\mathfrak{G}_{\Omega, F}$ is smooth and hence reduced, \mathfrak{Y}_Λ is in fact the closure of $\mathfrak{G}_{\Omega, F}$ in \mathbb{S}_Λ with its reduced scheme structure. Since the base is of dimension ≥ 2 , this schematic closure however need not be a group scheme (see [BT2, 3.2.15]).

To prove the contention of the lemma, we need to verify that the restriction of \mathfrak{Y}_U to the completion of the local rings of each height 1 prime associated to H_i is in fact isomorphic to the BT-group scheme \mathfrak{G}_{Ω_i} . This follows from Remarks 6.1 and 6.2.

Q. E. D

6.3. n -bounded-groups and schematization. The schematization of the n -bounded analogue is now carried out in two steps. As in the construction of the n -parahoric, we construct the Lie algebra bundle \mathcal{R}_Ω first (see (4.4)). This will be eventually turn out to be $\text{Lie}(\mathfrak{G}_\Omega)$.

PROPOSITION 6.5. *Let notations be as in (4.4) and (5.6). The trivial bundle with fiber \mathfrak{g} on \mathbf{A}_0 extends to \mathbf{A} preserving the Cartan decomposition as a canonical Lie-algebra bundle \mathcal{R}_Ω . Further, for each ζ_i , the restriction $\mathcal{R}_\Omega|_{X_i}$ is isomorphic to \mathcal{R}_{Ω_i} .*

Proof. This is special case of (5.6). Q.E.D

We will denote the Cartan decomposition of \mathcal{R} as follows:

$$\mathcal{R}_\Omega = \text{Lie}(T) \otimes \mathcal{O}_\Lambda \oplus_{r \in \Phi} \mathcal{R}_r. \quad (6.3.1)$$

The group scheme construction on the higher dimensional base has two obstructions. Firstly, schematic closures need not be flat when the base is not a Dedekind domain (see 5.9), and secondly, there is no immediate analogue of (5.4). This is because the bounded subsets Ω_i need not be singletons, and so the local BT-group schemes at the height 1 primes do not necessarily come from ramified covers.

Towards this end, we begin with a useful lemma.

LEMMA 6.6. (cf. [BT2, page 17-18]) *Let K be an infinite field and $A \subset K$ be a subring such that K is the quotient field of A . Let $f : X \rightarrow Y$ be a morphism of schemes over A and let $i : Z \hookrightarrow Y$ be a closed subscheme. Suppose that X is A -flat and that $f_K : X_K \rightarrow Y_K$ factorises through Z_K . Then f factorises through Z .*

Proof. By going to an affine cover of Y , we may assume that X and Y are A -affine with coordinate rings $\mathcal{O}[X]$ and $\mathcal{O}[Y]$. Since X is A -flat, the map $j_X : \mathcal{O}[X] \rightarrow K \otimes_{\mathcal{O}} \mathcal{O}[X]$ is injective. Let $a \in I(Z)$ be an element in the ideal defining Z in Y . Then $j_X \circ f^*(a) = f_K^* \circ j_Y(a) = 0$ since f_K has a factorisation. By the injectivity of j_X , we get $f^*(a) = 0$. Q.E.D

6.4. CONTINUATION OF PROOF OF THEOREM 1.3.

THEOREM 6.7. *The main theorem (1.3) holds when the n -tuple of concave functions is of type II (1.2) and when G is of type A_n, G_2, F_4 and E_6 (see (6.1)).*

Proof. We will break up the rather long and technical proof into two parts. Part (I) constructs the candidate for *big cell* in the type II case. Part (II) then constructs the group scheme by using this cell structure. In this sense, the key issue is the extension of the *schematic root datum*.

Part (I) of the proof: We begin with the closed subgroup scheme $\mathfrak{G}_{\Omega, F}$ of \mathbb{S}_F (obtained by the diagonal embedding of G in G^s).

The closed subgroup scheme $\mathfrak{G}_{\Omega, F} \hookrightarrow \mathbb{S}_F$ over $\text{Spec } F$ is such that each factor of the product $\mathfrak{U}_{\Omega, F}^- \times T_{\Omega, F} \times \mathfrak{U}_{\Omega, F}^+$, embeds factor by factor in the corresponding term in the cell of \mathbb{S}_F . The full cell $\mathfrak{C}_{\Omega, F} := \text{Im}(\mathfrak{U}_{\Omega, F}^- \times T_{\Omega, F} \times \mathfrak{U}_{\Omega, F}^+ \rightarrow \mathfrak{G}_{\Omega, F})$ therefore sits as a flat closed subscheme of the open subscheme

$$\text{Im}\left(\mathfrak{B}_F^- \times \mathcal{T}_F \times \mathfrak{B}_F^+ \rightarrow \mathbb{S}_F\right). \quad (6.4.1)$$

As before, let U be an open subset of $\text{Spec } \Lambda$ which contains all primes of height ≤ 1 . By (6.5) we have a locally free Lie algebra bundle \mathcal{R}_Ω on $\text{Spec } \Lambda = \mathbf{A}$. This gives an extension of $\text{Lie}(\mathfrak{G}_{\Omega, F})$ as a sub Lie-algebra bundle \mathcal{R}_Ω of $\text{Lie}(\mathbb{S}_\Lambda)$. Under the Cartan decomposition, we get the following refinement of sub-bundles

$$\mathcal{R}_{r, \Omega} \hookrightarrow \text{Lie}(\mathfrak{B}_r). \quad (6.4.2)$$

In the proof, as a preliminary step we will construct a *big cell* $\mathcal{C}_{\Omega, \Lambda}$ as a subscheme of the group scheme \mathbb{S}_Λ . This is achieved in (6.4.9) below. This scheme $\mathcal{C}_{\Omega, \Lambda}$ will be such that when restricted to the open subset U of $\text{Spec } \Lambda$, we have an open immersion

$$\mathcal{C}_U \hookrightarrow \mathfrak{G}_{\Omega, U}. \quad (6.4.3)$$

We recall that by (6.2.1) and the proof of (6.4), the scheme \mathfrak{Y}_Λ is in fact the *schematic closure* of $\mathfrak{G}_{\Omega, U}$ in \mathbb{S}_Λ . The main aim of this first part of the proof is not only to show the existence of a *big cell* $\mathcal{C}_{\Omega, \Lambda}$ in (6.4.9), but to show in fact that (6.8) may be applied to it because it satisfies the following key property:

Claim 1. *There is an open immersion from the scheme $\mathcal{C}_{\Omega, \Lambda}$ into \mathfrak{Y}_Λ .*

We begin with the *big cell* $\mathcal{C}_{\Omega, F} := \text{Im}(\mathfrak{U}_{\Omega, F}^+ \times T_{\Omega, F} \times \mathfrak{U}_{\Omega, F}^- \rightarrow \mathfrak{G}_{\Omega, F})$. This gives the decomposition:

$$\text{Lie}(\mathfrak{G}_{\Omega, F}) = \text{Lie}(\mathfrak{U}_{\Omega, F}^+) \oplus \text{Lie}(T_{\Omega, F}) \oplus \text{Lie}(\mathfrak{U}_{\Omega, F}^-).$$

The goal now is to extend $\mathcal{C}_{\Omega, F}$ to a *big cell* $\mathcal{C}_{\Omega, \Lambda}$, as an open subscheme of \mathfrak{Y}_Λ . We prove this in several steps.

STEP 1: TORAL EXTENSION TO Λ AND CLOSED EMBEDDING IN \mathfrak{Y}_Λ : Since we work with the *generically split* case, the maximal torus $T_{\Omega, F} \subset \mathfrak{G}_{\Omega, F}$ is *split*. We now work with U which contains all primes of height ≤ 1 . The torus $T_{\Omega, F}$ canonically extends to a smooth closed multiplicative subgroup scheme $\mathcal{T}_{\Omega, U}$ of $\mathfrak{G}_{\Omega, U}$ over U . This can be explicitly constructed as in [BT2, Section 4.4.18 and 4.6.2] (or by using the identity component of the Néron-lft model from [BLR, Proposition 6, page 292], as in [L, Proposition 3.2, page 32]).

By [Ray, Proposition IX. 2.4], it follows that the inclusion $\mathcal{T}_{\Omega, U} \hookrightarrow \mathfrak{G}_{\Omega, U} \hookrightarrow \mathbb{S}_U$ extends uniquely to a smooth group scheme $\mathcal{T}_{\Omega, \Lambda}$ together with a closed immersion $\mathcal{T}_{\Omega, \Lambda} \hookrightarrow \mathbb{S}_\Lambda$ as a sub-torus of the group scheme \mathbb{S}_Λ . Since $\mathcal{T}_{\Omega, \Lambda}$ is *flat and hence torsion-free*, it follows that $\mathcal{T}_{\Omega, \Lambda}$ is the schematic closure of $\mathcal{T}_{\Omega, U}$ in \mathbb{S} . Again, since \mathfrak{Y}_Λ is the schematic closure of $\mathfrak{G}_{\Omega, U}$ in \mathbb{S} , we conclude that the embedding $\mathcal{T}_{\Omega, \Lambda} \hookrightarrow \mathbb{S}_\Lambda$ factors via a closed embedding $\mathcal{T}_{\Omega, \Lambda} \hookrightarrow \mathfrak{Y}_\Lambda$.

STEP 2: UNIPOTENT AND BIG CELL EXTENSION TO U : Next, we need to extend the unipotent group schemes $\mathfrak{U}_{\Omega, F}^\pm$, firstly as closed subschemes of $\mathfrak{G}_{\Omega, U}$. This is easy since these are precisely the schematic closures of the root groups $\mathfrak{U}_{r, F}^\pm$, for $r \in \Phi$, and hence of $\mathfrak{U}_{\Omega, F}^\pm$ in $\mathfrak{G}_{\Omega, U}$; we rely obviously on Bruhat-Tits theory over discrete valuation rings. Call these extended group schemes $\mathfrak{U}_{\Omega, U}^\pm$ and observe that since we work over height 1 primes, these are *flat* over U .

Thus we have the *big cell*

$$\mathcal{C}_U := \text{Im}(\mathfrak{U}_{\Omega, U}^- \times \mathcal{T}_{\Omega, U} \times \mathfrak{U}_{\Omega, U}^+ \hookrightarrow \mathfrak{G}_{\Omega, U}) \quad (6.4.4)$$

in $\mathfrak{G}_{\Omega, U}$ which has the property that at each of the height 1 primes, it restricts to the big cell in the usual Bruhat-Tits group schemes $\mathfrak{G}_{\Omega, j}$.

STEP 3: UNIPOTENT EXTENSION TO $\text{Spec } \Lambda$ AS CLOSED SUBSCHEMES OF \mathfrak{Y}_Λ : We next ensure that the group schemes $\mathfrak{U}_{\Omega, U}^\pm$ extend as subschemes of the schematic closure \mathfrak{Y}_Λ to ensure that a *big cell* $\mathcal{C}_{\Omega, \Lambda}$ can be openly embedded in \mathfrak{Y}_Λ .

We do this in three sub-steps.

STEP 3.1: We first extend the unipotent group schemes abstractly from U to the whole of $\text{Spec } \Lambda$. For each $r \in \Phi$, consider the *fractional invertible ideal* of Λ i.e., a

finitely generated projective sub Λ -module of F given by:

$$f(r) := z_1^{m_r(\Omega_1)} \cdots z_n^{m_r(\Omega_n)} \Lambda \subset F. \quad (6.4.5)$$

By [BT2, 3.2.6] this ideal defines smooth Λ -group schemes $\mathcal{U}_{r,\Omega,\Lambda}$ extending the root groups $U_{r,F}$ as well as the root groups $\mathcal{U}_{r,\Omega,U}$ over U .

STEP 3.2: We now prove that the group schemes $\mathcal{U}_{r,\Omega,\Lambda}$ are closed subgroup schemes of the abelian subgroup schemes \mathfrak{B}_r (6.1.3) of \mathbb{S}_Λ and therefore closed subschemes of \mathfrak{Y}_Λ . We do so by showing in fact that they are *schematic closures* of the root groups $\mathcal{U}_{r,\Omega,U}$ in \mathfrak{B}_r . Our method is to reduce verifications by §5.2 to an inclusion of Lie-algebra bundles on Λ . This check further reduces to U .

STEP 3.2.1: By §5.2, the following group schemes arise from the W -construction of their Lie-algebra bundles:

$$\mathcal{U}_{r,\Omega,U} \simeq W(\text{Lie}(\mathcal{U}_{r,\Omega,U})) \quad \mathfrak{B}_r \simeq W(\text{Lie}(\mathfrak{B}_r)). \quad (6.4.6)$$

Note that the root groups $\mathcal{U}_{r,\Omega,U}$ are closed subgroup schemes (over U) of the restrictions $\mathfrak{B}_{r,U}$ to U . Moreover, \mathfrak{B}_r are themselves closed abelian subgroup schemes of \mathbb{S}_Λ (see (6.1.3)). Thus, by [SGAD, Expose I, Proposition 4.6.6] (recalled in §5.2), we conclude that the corresponding rank 1 Lie algebra bundle $\text{Lie}(\mathcal{U}_{r,\Omega,U})$ is a direct summand of the Lie algebra bundle $\text{Lie}(\mathfrak{B}_r|_U)$.

STEP 3.2.2: We claim that $\text{Lie}(\mathcal{U}_{r,\Omega,U})$ extends to $\text{Spec } \Lambda$ as the line bundle \mathcal{R}_r of (6.3.1). Indeed, since \mathcal{R}_r restricts to $\text{Lie}(\mathcal{U}_{r,\Omega,U})$ on U , it follows that by taking their reflexive closures inside \mathcal{R}_Ω (see (6.5)), each $\text{Lie}(\mathcal{U}_{r,\Omega,U})$ extends to $\text{Spec } \Lambda$ as the line bundle \mathcal{R}_r . This extension \mathcal{R}_r is a summand of $\text{Lie}(\mathfrak{B}_r)$ because their restriction to U has this property.

Moreover, these are extensions as Lie algebra subbundles of \mathcal{R}_Ω because by the Hartogs lemma the Lie bracket extends across codimension bigger than two.

STEP 3.2.3: By STEP 3.1, the root group schemes $\mathcal{U}_{r,\Omega,U}$ have already the extensions as unipotent group schemes given by $\mathcal{U}_{r,\Omega,\Lambda}$. Again, since the Lie algebra bundles \mathcal{R}_r and $\text{Lie}(\mathcal{U}_{r,\Omega,\Lambda})$ agree on U , we have an isomorphism of Lie algebra bundles:

$$\mathcal{R}_r \simeq \text{Lie}(\mathcal{U}_{r,\Omega,\Lambda}). \quad (6.4.7)$$

This therefore identifies $\text{Lie}(\mathcal{U}_{r,\Omega,\Lambda})$ as a Lie subalgebra bundle of \mathcal{R}_Ω as well as a *direct summand* of $\text{Lie}(\mathfrak{B}_r)$.

STEP 3.2.4: We observe that $\mathcal{U}_{r,\Omega,\Lambda}$ arises from the functor W construction of §5.2. More precisely, we have scheme-theoretic isomorphisms of vector group schemes

$$\mathcal{U}_{r,\Omega,\Lambda} = W(\text{Lie}(\mathcal{U}_{r,\Omega,\Lambda})). \quad (6.4.8)$$

Since $\text{Lie}(\mathcal{U}_{r,\Omega,\Lambda})$ are direct summands of $\text{Lie}(\mathfrak{B}_r)$, by §(5.2) it follows that $\mathcal{U}_{r,\Omega,\Lambda}$ are closed subgroup schemes of \mathfrak{B}_r and therefore of \mathbb{S}_Λ . It follows in particular that the inclusions $\mathcal{U}_{r,\Omega,\Lambda} \subset \mathfrak{Y}_\Lambda$ are *closed embeddings* as claimed.

STEP 3.3: We have arrived at the setting of [BT2, 2.2.2]. For $r \in \Phi^\pm$, we denote the scheme-theoretic product of group schemes $\mathcal{U}_{r,\Omega,\Lambda}$ as the Λ -scheme $\mathcal{U}_{\Omega,\Lambda}^\pm$. Since each $\mathcal{U}_{r,\Omega,\Lambda}$ is the schematic closure of $\mathcal{U}_{r,\Omega,U}$ in \mathbb{S}_Λ , it follows that $\mathcal{U}_{\Omega,\Lambda}^\pm$ are the *schematic closures* of $\mathcal{U}_{\Omega,U}^\pm$ in \mathbb{S}_Λ . The $\mathcal{U}_{\Omega,U}^\pm$ are smooth unipotent subgroup schemes of \mathbb{S}_Λ over U and since $\mathcal{U}_{\Omega,\Lambda}^\pm$ are flat, by [BT2, 1.2.7] it follows that $\mathcal{U}_{\Omega,\Lambda}^\pm$ are closed subgroup schemes of \mathbb{S}_Λ . Hence we get the smooth unipotent group scheme structure on $\mathcal{U}_{\Omega,\Lambda}^\pm$ as well as realise them as closed subschemes of \mathfrak{Y}_Λ .

This completes the proof of STEP 3.3, STEP 3.2, and STEP 3.1 and as a consequence of STEP 3.

STEP 4: CLOSED EMBEDDING OF BIG CELL IN \mathfrak{Y}_Λ : We finally need to ensure that the open immersion $j_U : \mathfrak{U}_{\Omega,U}^- \times \mathcal{T}_{\Omega,U} \times \mathfrak{U}_{\Omega,U}^+ \hookrightarrow \mathfrak{G}_{\Omega,U}$ extends as an open immersion $j_\Lambda : \mathfrak{U}_{\Omega,\Lambda}^- \times \mathcal{T}_{\Omega,\Lambda} \times \mathfrak{U}_{\Omega,\Lambda}^+ \hookrightarrow \mathfrak{Y}_\Lambda$.

More precisely, we show that the schematic closure of the image of the multiplication map of $\mathfrak{U}_{\Omega,\Lambda}^- \times \mathcal{T}_{\Omega,\Lambda} \times \mathfrak{U}_{\Omega,\Lambda}^+$ in \mathbb{S}_Λ is actually \mathfrak{Y}_Λ and then derive the open immersion property as a consequence.

By using the discussion preceding (6.4.1), and taking closures term by term, we see that $\mathfrak{U}_{\Omega,\Lambda}^- \times \mathcal{T}_{\Omega,\Lambda} \times \mathfrak{U}_{\Omega,\Lambda}^+$ is a flat closed subscheme of $\mathfrak{B}^- \times \mathcal{T} \times \mathfrak{B}^+$ which is itself an open subscheme of \mathbb{S}_Λ . In other words, the product defines an isomorphism of schemes of $\mathfrak{U}_{\Omega,\Lambda}^- \times \mathcal{T}_{\Omega,\Lambda} \times \mathfrak{U}_{\Omega,\Lambda}^+$ onto a flat closed subscheme

$$\mathfrak{C}_{\Omega,\Lambda} \subset \text{Im}(\mathfrak{B}^- \times \mathcal{T} \times \mathfrak{B}^+) \quad (6.4.9)$$

and moreover, $\text{Im}(\mathfrak{B}^- \times \mathcal{T} \times \mathfrak{B}^+)$ is an open subscheme of \mathbb{S}_Λ .

Let $\bar{\mathfrak{C}}$ be the schematic closure of $\mathfrak{C}_{\Omega,\Lambda}$ in \mathbb{S}_Λ . We *claim* that $\bar{\mathfrak{C}} = \mathfrak{Y}_\Lambda$. By what we have checked above, it is clear that $\bar{\mathfrak{C}}$ is contained in \mathfrak{Y}_Λ .

Conversely, the big cell $\mathfrak{C}_{\Omega,F}$ is an open dense subscheme of $\mathfrak{G}_{\Omega,F}$, hence $\bar{\mathfrak{C}} \supset \mathfrak{G}_{\Omega,F}$, implying that $\bar{\mathfrak{C}}$ contains \mathfrak{Y}_Λ . Hence, by applying [BT2, 1.2.6], we conclude that $\mathfrak{C}_{\Omega,\Lambda}$ is an *open dense subscheme* of its schematic closure which has been identified with \mathfrak{Y}_Λ .

This completes the proof of Claim 1 and hence of Part I of the argument.

Part II of the proof. We now state and give the proof of Bruhat and Tits [BT2, Proposition 2.2.10], minimally tweaked and tailored for our purposes. The result is referred to in a couple of places in their work (see [BT2, 3.9.4]), where they talk of possibility of an extension of their theory to 2BT-group schemes; but for us it is indispensable. In [BT2, Proposition 2.2.10], the role of the group scheme $\mathfrak{GL}(M)$ is played by the group scheme $\mathbb{S}_{\Omega,\Lambda}$ (6.3) in our situation. Hence we almost “reproduce” their rather long and technical proof in its entirety, with suitable changes.

PROPOSITION 6.8. ([BT2, Proposition 2.2.10]) *Let \mathfrak{H} be the image of $\mathfrak{C}_{\Omega,\Lambda} \times \mathfrak{C}_{\Omega,\Lambda}$ in \mathbb{S}_Λ by the product morphism π of \mathbb{S}_Λ . Then \mathfrak{H} is open in \mathfrak{Y}_Λ and is a subgroup scheme of \mathbb{S}_Λ , with fibre $\mathfrak{G}_{\Omega,U}$ over U , flat, of finite type, containing $\mathfrak{U}_{\Omega,\Lambda}^-$, $\mathcal{T}_{\Omega,\Lambda}$ and $\mathfrak{U}_{\Omega,U}^+$ as closed subgroup schemes and containing $\mathfrak{C}_{\Omega,\Lambda}$ as an open subscheme (a big cell). Furthermore, the group scheme \mathfrak{H} is a quasi-affine smooth group scheme over Λ .*

Proof. For simplicity of notation, we will do away with all *multiple greek letters* in subscripts.

As a preliminary step so as to follow the rest of the argument in [BT2], we need to check that $\mathfrak{U}_\Lambda^- \cdot \mathcal{T}_\Lambda$ and $\mathcal{T}_\Lambda \cdot \mathfrak{U}_\Lambda^+$ are group schemes.

For this we consider the morphism:

$$\mathfrak{U}_\Lambda^- \times \mathcal{T}_\Lambda \rightarrow \mathbb{S}_\Lambda \quad (6.4.10)$$

given by $(u, t) \mapsto t.u.t^{-1}$. By [BT2, 4.4.19], this gives an action of T_U on \mathfrak{U}_U^- and hence over U , the image lands in \mathfrak{U}_U^- . Since \mathfrak{U}_Λ^- is a closed subscheme of \mathbb{S}_Λ , and since U contains all points of height ≤ 1 , it follows that the morphism given by conjugation has its image in \mathfrak{U}_Λ^- . In other words, $\mathfrak{U}_\Lambda^- \cdot \mathcal{T}_\Lambda = \mathcal{T}_\Lambda \cdot \mathfrak{U}_\Lambda^-$ in \mathbb{S}_Λ , i.e., it is

a subgroup scheme [DG, 3.10, page 166]. Similarly, $\mathcal{T}_\Lambda \mathcal{U}_\Lambda^+$ is also a group scheme.

As $\mathcal{C} \times \mathcal{C}$ is flat and $\pi(\mathcal{C}_U \times \mathcal{C}_U) \subset \mathfrak{G}_U$, by (6.6) (where \mathfrak{Y} plays the role of Z), we have the inclusion $\mathfrak{H} \subset \mathfrak{Y}$.

By what we have seen in Part I above, \mathcal{C} is an open subset of \mathfrak{Y} . So the inverse image Γ of \mathcal{C} by $\pi : \mathcal{C} \times \mathcal{C} \rightarrow \mathfrak{Y}$ is therefore an open subset of $\mathcal{C} \times \mathcal{C}$. Since $\mathcal{U}_\Lambda^- \mathcal{T}_\Lambda$ and $\mathcal{T}_\Lambda \mathcal{U}_\Lambda^+$ are groups, we get the inclusions:

$$\Gamma \supset ((\mathcal{U}_\Lambda^- \times \mathcal{T}_\Lambda \times \{1\}) \times (\mathcal{U}_\Lambda^- \times \mathcal{T}_\Lambda \times \{1\})) \cup ((\{1\} \times \mathcal{T}_\Lambda \times \mathcal{U}_\Lambda^+) \times (\{1\} \times \mathcal{T}_\Lambda \times \mathcal{U}_\Lambda^+)). \quad (6.4.11)$$

Let $p \in \text{Spec } \Lambda$ be a closed point and let k be the residue field of the local ring Λ_p . The fibre Γ_p of Γ is an open subset of $\mathcal{C}_p \times \mathcal{C}_p$, dense in $\mathcal{C}_p \times \mathcal{C}_p$: in fact, if $u, u' \in \mathcal{U}_p^-$ and $z, z' \in \mathcal{T}_p$, the set V of $(v, v') \in \mathcal{U}_p^+ \times \mathcal{U}_p^+$ such that $((u, z, v), (u', z', v')) \in \Gamma_p$ is an open subset of $\mathcal{U}_p^+ \times \mathcal{U}_p^+$, and is non-empty since $(1, 1) \in V$ by (6.4.11), therefore dense in $\mathcal{U}_p^+ \times \mathcal{U}_p^+$ since \mathcal{U}_p^+ is connected.

Let H_p be the closure of \mathcal{C}_p in \mathbb{S}_p . As \mathcal{C}_p is open in \mathfrak{Y}_p , *a fortiori*, \mathcal{C}_p is open in H_p . Moreover, Γ_p is dense in $H_p \times H_p$, and $\pi(H_p \times H_p) \subset H_p$ since by definition $\pi(\Gamma_p) \subset \mathfrak{Y}_p$. In other words, H_p is a subgroup scheme of \mathbb{S}_p . As \mathcal{C}_p is open dense in H_p , we have $H_p = \mathcal{C}_p \cdot \mathcal{C}_p$, hence

$$H_p = \mathfrak{H} \cap \mathbb{S}_p. \quad (6.4.12)$$

We also have $H_p = \mathcal{C}_p^{-1} \cdot \mathcal{C}_p = \mathcal{U}_p^+ \mathcal{T}_p \mathcal{U}_p^- \mathcal{U}_p^- \mathcal{T}_p \mathcal{U}_p^+ = \mathcal{U}_p^+ \mathcal{C}_p$, and hence

$$H_p = \bigcup_{a \in \mathcal{U}_p^+(k)} a \cdot \mathcal{C}_p. \quad (6.4.13)$$

We now show that \mathfrak{H} is open in \mathfrak{Y} and is flat. Firstly, \mathfrak{H} is a *constructible* subset of \mathfrak{Y} . To show that it is open, (since Λ is noetherian), it suffices to show that if $x \in \mathfrak{H}$ and $y \in \mathfrak{Y}$ are such that $x \in \{y\}$, then $y \in \mathfrak{H}$ ([DG, Cor 3.4, p.76]).

Let p be the projection of x in $\text{Spec } \Lambda$. Since the smooth group scheme \mathbb{S}_Λ is obtained by a Weil restriction of scalars (see the paragraph just before (6.3)), it behaves well under base change. Let $j : \text{Spec } \Lambda_p \rightarrow \text{Spec } \Lambda$ be the induced morphism. As the local ring Λ_p is flat over Λ , the schematic closures of \mathfrak{G}_U and $\mathcal{C} \times_\Lambda \Lambda_p$ in $j^*(\mathbb{S}_\Lambda)$ coincide with $j^*(\mathfrak{Y})$ ([BLR, Proposition 2, page 56], [DG, page 56]). Hence we may reduce to the case when Λ is a local ring with maximal ideal \mathfrak{p} .

Let B_x be the local ring at x in \mathbb{S}_Λ and let \tilde{B} be the strict henselization of B_x . Let $x' \in \text{Spec } \tilde{B}$ be the closed point and let $\alpha : \text{Spec } \tilde{B} \rightarrow \text{Spec } B_x$ be the canonical morphism: we have $\alpha(x') = x$. For us the residue field at x is algebraically closed and hence by (6.4.13), there exists an $a \in \mathcal{U}_p^+(k)$ such that $x \in a \cdot \mathcal{C}_p$; since \mathcal{U}_Λ^+ is smooth, by the Hensel lemma, there exists a section $\hat{a} \in \mathcal{U}_\Lambda^+(\tilde{B})$ of \mathcal{U}_Λ^+ above $\text{Spec } \tilde{B}$ such that $\hat{a}(x') = a$.

We now do a base change by $\Lambda \rightarrow \tilde{B}$. Let $\tilde{\mathfrak{X}}$ (resp. X) be a Λ -scheme (resp. a U -scheme, where U has been base changed to the local ring at p), set $\tilde{\mathfrak{X}} := \tilde{\mathfrak{X}} \times_{\text{Spec } \Lambda} \text{Spec } \tilde{B}$ (resp. $\tilde{X} := X \times_U \tilde{U}$, with $\tilde{U} = U \times_{\text{Spec } \Lambda} \text{Spec } \tilde{B}$). If $f : \tilde{\mathfrak{X}} \rightarrow \tilde{\mathfrak{Z}}$ is a morphism of Λ -schemes, set $\tilde{f} = f \times \text{id} : \tilde{\mathfrak{X}} \rightarrow \tilde{\mathfrak{Z}}$. Finally, let $\beta : \tilde{\mathbb{S}} \rightarrow \mathbb{S}_\Lambda$ be the first projection. As \tilde{B} is flat over B_x and B_x is flat over Λ (as \mathbb{S}_Λ is flat), \tilde{B} is flat over Λ and we see as above that $\tilde{\mathfrak{Y}}$ is the schematic closure of both $\tilde{\mathcal{C}}$ and $\tilde{\mathfrak{G}}_U$ in $\tilde{\mathbb{S}}$. On the other hand, the left-translation by the section \hat{a} is an automorphism of the scheme $\tilde{\mathbb{S}}$ and the restriction \hat{a}_U of \hat{a} to \tilde{U}

belongs to $\mathcal{U}_p^+(\tilde{U}) \subset \tilde{\mathcal{G}}_U(\tilde{U})$; thus we have $\hat{a}.\tilde{\mathcal{G}}_U = \hat{a}_U.\tilde{\mathcal{G}}_U = \tilde{\mathcal{G}}_U$ and $\hat{a}.\tilde{\mathcal{Y}} = \tilde{\mathcal{Y}}$. As $\tilde{\mathcal{C}}$ is an open in $\tilde{\mathcal{Y}}$, it follows that $\hat{a}.\tilde{\mathcal{C}}$ is *open in* $\tilde{\mathcal{Y}}$.

As \tilde{B} is faithfully flat over B_x , the morphism α is surjective. We have $x \in \overline{\{y\}}$, thus $y \in \text{Spec } B_x$ and there exists $y' \in \text{Spec } \tilde{B}$ such that $\alpha(y') = y$; moreover, $x' \in \overline{\{y'\}}$ since the closed point of $\text{Spec } \tilde{B}$ belongs to the closure of any other point of $\text{Spec } \Lambda$. Set $x'' := (\alpha \times \text{id})(x') = (x, x')$ and $y'' := (\alpha \times \text{id})(y') = (y, y')$; then $x'' \in \hat{a}.\tilde{\mathcal{C}}$, $y'' \in \tilde{\mathcal{Y}}$ and $x'' \in \overline{\{y''\}}$. As $\hat{a}.\tilde{\mathcal{C}}$ is open in $\tilde{\mathcal{Y}}$, it follows that

$$y'' \in \hat{a}.\tilde{\mathcal{C}} = \tilde{\pi} \circ (\hat{a} \times \text{id})(\text{Spec } \tilde{B} \times_{\tilde{B}} \tilde{\mathcal{C}}) \subset \tilde{\pi}(\tilde{\mathcal{C}} \times_{\tilde{B}} \tilde{\mathcal{C}}). \quad (6.4.14)$$

We then deduce that

$$y = \beta(y'') \in \beta \circ \tilde{\pi}(\tilde{\mathcal{C}} \times_{\tilde{B}} \tilde{\mathcal{C}}) = \pi \circ (\beta \times \beta)(\tilde{\mathcal{C}} \times_{\tilde{B}} \tilde{\mathcal{C}}) \subset \pi(\mathcal{C} \times \mathcal{C}) = \mathfrak{H}, \quad (6.4.15)$$

and hence that \mathfrak{H} is *open in* \mathcal{Y} .

Moreover, $\hat{a}.\tilde{\mathcal{C}}$ is flat over $\text{Spec } \tilde{B}$. As it is a neighbourhood of x in $\mathfrak{H}_{\tilde{B}}$, it follows that the local ring at x in $\mathfrak{H}_{\tilde{B}}$ is flat over $\text{Spec } \tilde{B}$ and by faithfully flat descent, the local ring of x in \mathfrak{H} is flat over Λ . Hence \mathfrak{H} is *flat*.

It follows that, \mathfrak{H} , being an open subscheme of \mathcal{Y} , is a subscheme of \mathbb{S}_Λ (which is no reason for it to be affine!). It is a subgroup scheme of \mathbb{S}_Λ . In fact, the restriction to \mathfrak{H} of the product π of \mathbb{S}_Λ sends $\mathfrak{H}_U \times \mathfrak{H}_U$ into \mathcal{G}_U , and hence, since \mathfrak{H} is flat, it factorises by a morphism $\mathfrak{H} \times \mathfrak{H} \rightarrow \mathcal{Y}$ (6.6). But set-theoretically we have $\pi(\mathfrak{H} \times \mathfrak{H}) \subset \mathfrak{H}$ since the fibres of \mathfrak{H} are subgroups (6.4.13). Since \mathfrak{H} is an open in \mathcal{Y} , it implies that the morphism $\pi : \mathfrak{H} \times \mathfrak{H} \rightarrow \mathcal{Y}$ factorises through \mathfrak{H} . A similar argument for the *inverse* morphism shows that \mathfrak{H} is a subgroup scheme.

Now \mathfrak{H} , is an open subscheme of \mathcal{Y} which is closed in the schematic hull \mathbb{S}_Λ . Further, by the (6.1.1), \mathbb{S}_Λ is affine and hence \mathfrak{H} is *quasi-affine*. That it has the remaining stated properties is easy to check. Q.E.D

Remark 6.9. The quasi-affineness can also be deduced as follows. Since the fibre of \mathfrak{H} over U is \mathcal{G}_U which is affine, and since \mathfrak{H} , being the image of $\mathcal{C} \times \mathcal{C}$, has connected fibres, by [Ray, Corollary VII, 2.2, p. 112], we conclude that \mathfrak{H} is *quasi-affine*.

Remark 6.10. When the characteristic of the residue field is zero, the group scheme \mathfrak{H} is *flat* and hence smooth. Over residue fields of positive characteristics, the smoothness of \mathfrak{H} is ensured by the smoothness of the torus $\mathcal{T}_{\Omega, \Lambda}$ since the unipotent groups $\mathcal{U}_{\Omega, \Lambda}^\pm$ are smooth. By the argument in the paragraph following (5.5.1), the torus on the big open subset extends as a torus $\mathcal{T}_{\Omega, \Lambda}$ and is therefore smooth even in the general case.

To complete the proof of THEOREM 6.7, we now set

$$\mathcal{G}_\Omega := \mathfrak{H}. \quad (6.4.16)$$

This group scheme clearly satisfies the properties in the statement of the theorem (1.3). Indeed, in general it is quasi-affine, its sections can be computed by restriction to U , (4) and (5) follow from Part (I) of the proof and the proof of the characterisation and uniqueness of the group scheme follows exactly as in (5.7). Q.E.D

6.5. PROOF OF THEOREM 1.5.

Proof. Since \mathcal{G}_Ω is quasi-affine by (6.7), as in (5.8) we can identify the restriction of \mathcal{G}_Ω to a sub-diagonal ((1.5) (b)) with the group scheme $\mathcal{G}_{f_{(i_1, \dots, i_m)}}$ where $f_{i_\ell} := f_{\Omega_{i_\ell}}$. The

description of the closed fibre follows from [BT2, 4.6.5, 4.6.12] applied to the concave function $f_{(i_1, \dots, i_m)} = \sum_{\ell=1}^m f_{i_\ell}$. Q.E.D

7. SCHEMATIZATION OF \mathfrak{n} -BOUNDED GROUPS P_f

The notations in this section are as in the previous one, namely $\mathbf{A} \simeq \mathbb{A}^n$ and $\mathbf{A}_0 \subset \mathbf{A}$ is the complement of the coordinate hyperplanes. In this section, we return to the general case of an almost-simple group G . We complete the picture by proving THEOREM 1.3 for n -concave functions of type III. More precisely, we construct the group scheme over \mathbf{A} , given the data of n -concave functions $\mathbf{f} = \{f_j\}$ on the root system Φ of G , along with an assignment of the BT-group schemes $\{\mathfrak{G}_{f_j}\}_{j=1}^n$ at the generic points of each of the coordinate hyperplanes $H_j \subset \mathbf{A}$.

Although the strategy of the proof is broadly a reduction to the proof of $\mathrm{SL}(n)_\Omega$ case, in the present case of (G, \mathbf{f}) there are two crucial differences. First, unlike the case of the concave function f_Ω , we do not have any way of assigning points in the affine apartment (see §(3.11) for an example) to get a product \mathfrak{n} -parahoric group scheme \mathbb{S}_Λ (see §(6.1)) for G which gave us an ambient group scheme to carry out the strategy. And second, as it will become clearer by the discussion below, the characteristic of the residue field needs to be weakened to $p \geq h_G$.

As a first step, we recall some of the main results of [BT2]. These will play the key role in this section. The setting of [BT2, §3], K is an infinite field containing a subring A having K as its field of fractions. The aim of that paragraph (see [BT2, §3, 3.1.3]) is to show that if G has a *donnée radicielles schématique* $\mathcal{D} = (\mathcal{L}, (\mathcal{U}_a)_{a \in \Phi})$ (see [BT2, 3.1.1]) or a *schematic root datum*, then there exists an A -group scheme \mathfrak{G} extending $G \times \mathrm{Spec} K$ and with big-cell given by \mathcal{D} . This is done by showing that for any faithful representation $G \hookrightarrow \mathrm{GL}(V)$, there is an A -lattice $M \subset V_K$ such that the BT-group scheme \mathfrak{G} is the schematic closure of G_K in $\mathfrak{GL}(M)$. In [BT2, §4, 4.5.4], when K is equipped with a valuation, it is shown that given any concave function f on Φ , one can construct a *schematic root datum* for G . We apply these results to our context as follows. Since G is almost-simple, we may assume that we have a factorisation $G \hookrightarrow \mathrm{SL}(V)$. Hence, \mathfrak{G}_f is the schematic closure of G_K in the schematic closure of $\mathrm{SL}(V)_K$ in $\mathfrak{GL}(M)$. By choosing a \mathcal{O} -basis for M , it is easy to see that we get a concave function h on the root system Φ_V of $\mathrm{SL}(V)$ such that the schematic closure of $\mathrm{SL}(V)_K$ in $\mathfrak{GL}(M)$ is precisely the BT-group scheme $\mathrm{SL}(V)_h$.

By (3.21) and the remarks preceding it, we may assume that h is an *optimal concave function* on Φ_V and hence it is of the form h_Ω for a bounded subset Ω in the affine apartment of $\mathrm{SL}(V)$.

Thus, we are reduced to the following situation: the BT-group scheme \mathfrak{G}_f is the schematic closure of G_K in $\mathrm{SL}(V)_\Omega$. This can be carried out for each of the $\{f_j\}_{j=1}^n$ and we get bounded subsets $\mathbf{\Omega} = (\Omega_1, \dots, \Omega_n)$ in the affine apartment of $\mathrm{SL}(V)$.

By THEOREM 6.7 taking $\mathbf{\Omega} = (\Omega_1, \dots, \Omega_n)$ and the group $\mathrm{SL}(V)$, we obtain a \mathfrak{n} BT-group scheme $\mathfrak{GL}_{\mathbf{\Omega}}$ on \mathbf{A} . Although only quasi-affine, this group scheme will play the role of the schematic hull \mathbb{S}_Λ (see (6.1.1)) of §6. Further, on U whose complement in \mathbf{A} has codimension ≥ 2 , we have an inclusion of group schemes:

$$\mathfrak{G}_{f,U} \subset \mathfrak{GL}_{\mathbf{\Omega},U}. \tag{7.0.1}$$

The proof strategy for proving THEOREM 6.7 now almost applies. Let us explain the difference. In that proof, it was shown in STEP 3.2 that the root groups $\mathfrak{U}_{r,\Omega,\Lambda}$ are the

schematic closures of $\mathfrak{U}_{r,\Omega,U}$ in \mathbb{S}_Λ by the intermediary of *vector subgroup* \mathfrak{B}_r of \mathfrak{S}_Λ . This vector group exists because of the specific *diagonal embedding* into the ambient group scheme \mathbb{S}_Λ . In the present setting however our chosen representation, namely $G \subset \mathrm{SL}(V)$, is arbitrary and cannot be expected to have the desirable property that we have for every root $r \in \Phi$, a vector subgroup scheme of $\mathrm{SL}(n)_\Omega$ whose restriction to U contains $\mathfrak{U}_{r,\mathbf{f},U}$. Consequently, to replace the arguments of STEP 3.2, we recover the group law of unipotent group schemes by the Baker-Campbell-Hausdorff formula. We do this as follows. If $\{c_\alpha\}$ denote the coefficients of the higher root, recall the Coxeter number of G h_G is defined as $1 + \sum c_\alpha$. Let \mathfrak{N} be a nilpotent Lie algebra bundle on \mathbf{A} . By [Se, Section 2.1] (see also [BDP, 2.2] and other references there), *when the characteristic $p \geq h_G$* , the Baker-Campbell-Hausdorff group law equips the scheme $W(\mathfrak{N})$ with the structure of a *smooth, affine, unipotent group scheme with connected fibres* together with an isomorphism $\exp : \mathfrak{N} \rightarrow \mathrm{Lie}(W(\mathfrak{N}))$ of nilpotent Lie algebra bundles. Conversely, if \mathfrak{U} is an affine, smooth, unipotent group scheme with connected fibres, we have a unique isomorphism of unipotent group schemes $\exp : W(\mathrm{Lie}(\mathfrak{U})) \rightarrow \mathfrak{U}$. Note that the Baker-Campbell-Hausdorff formula realizes the possible non-abelian unipotent group scheme structures on \mathfrak{U} . In the absence of vector group schemes \mathfrak{B}_r , this is the key substitute. Let us now go over arguments that replace STEP 3.2.

As in STEP 3.2 we need to prove that the extended root group schemes $\mathfrak{U}_{r,\mathbf{f},\mathbf{A}}$ are closed subgroup schemes of $\mathfrak{S}\mathfrak{L}_\Omega$ on \mathbf{A} . We begin by noting that over the open subset $U \subset \mathbf{A}$, the root group schemes $\mathfrak{U}_{r,\mathbf{f},U}$ are closed subgroup schemes of the restriction to U of one of the big cells, say \mathfrak{B}^+ of $\mathfrak{S}\mathfrak{L}_\Omega$. By THEOREM 6.7, \mathfrak{B}^+ of $\mathfrak{S}\mathfrak{L}_\Omega$ is affine. We set $\mathfrak{N}^\pm := \mathrm{Lie}(\mathfrak{B}^\pm)$. This is a nilpotent Lie algebra bundle on \mathbf{A} . When $p \geq h_G$, we have the natural isomorphism:

$$\exp : W(\mathfrak{N}^\pm) \simeq \mathfrak{B}^\pm \quad (7.0.2)$$

of affine, smooth, connected unipotent group schemes. Since $\mathfrak{U}_{r,\mathbf{f},U} \subset \mathfrak{B}_U^+$ is a closed subgroup scheme, we may review this inclusion in terms of the “W”-functor as follows:

$$W(\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U})) \subset W(\mathrm{Lie}(\mathfrak{B}_U^+)). \quad (7.0.3)$$

By this closed embedding of affine U -schemes, we get the Lie algebra subbundle $\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U}) \subset \mathrm{Lie}(\mathfrak{B}_U^+)$ structure. This is in fact a *direct summand* of locally free sheaves on U . Now taking the *reflexive closure* $\overline{\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U})}$ of the rank one Lie algebra bundle $\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U})$, we see that $\overline{\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U})}$ is a locally free *direct summand* of \mathfrak{N}^+ over \mathbf{A} .

As in STEP 3.2, the reflexive closure $\overline{\mathrm{Lie}(\mathfrak{U}_{r,\mathbf{f},U})}$ gets identified with the line Lie algebra bundle \mathfrak{R}_r . It is therefore a *direct summand* of \mathfrak{N}^+ over \mathbf{A} . This implies that as affine schemes, we have a closed embedding:

$$W(\mathfrak{R}_r) \subset W(\mathfrak{N}^+) \simeq \mathfrak{B}^+ \quad (7.0.4)$$

Setting $\mathfrak{U}_{r,\mathbf{f},\mathbf{A}} := W(\mathfrak{R}_r)$, it follows that the scheme underlying $\mathfrak{U}_{r,\mathbf{f},\mathbf{A}}$ is a closed subscheme of \mathfrak{B}^+ . By the Baker-Campbell-Hausdorff formula, this inclusion is in fact one of closed subgroup schemes over \mathbf{A} and hence of $\mathfrak{S}\mathfrak{L}_\Omega$ on \mathbf{A} . This completes the proof of the key difference we mentioned above. In particular, at this stage the *big cell* \mathfrak{B}_Φ can also be easily seen to exist with the standard properties exactly as in the proof of THEOREM 6.7.

With these changes in place, the second part of the proof of THEOREM 6.7 may be applied. We get the required nBT-group scheme $\mathfrak{G}_{\mathbf{f},\mathbf{A}}$ associated to \mathbf{f} on \mathbf{A} . It is open inside a closed subscheme, say \mathcal{X} , of $\mathfrak{S}\mathfrak{L}_\Omega$. Recall $\mathfrak{S}\mathfrak{L}_\Omega$ is open inside a closed

subscheme say C of \mathbb{S}_Λ . Let $\overline{\mathcal{Z}}$ denote the closure of \mathcal{Z} in C . The intersection of $\overline{\mathcal{Z}}$ with \mathfrak{GL}_n in C is \mathcal{Z} . It follows that \mathcal{Z} is open in $\overline{\mathcal{Z}}$. Therefore, $\mathfrak{G}_{f,\Lambda}$ is open inside $\overline{\mathcal{Z}}$. Now $\overline{\mathcal{Z}}$ is affine because \mathbb{S}_Λ is affine. We conclude $\mathfrak{G}_{f,\Lambda}$ is quasi-affine (this can also be checked using [Ray, Corollary VII, 2.2, p. 112]). In conclusion, when $\text{char}(k) \geq h_G$ and satisfies the hypothesis of §(2.2) we have constructed $\mathfrak{G}_{f,\Lambda}$ is a *smooth, quasi-affine* group scheme with all the properties stated in THEOREM 1.3 for n -concave functions of type III.

8. SCHEMATIZATION OF n -Moy-Prasad GROUPS

In this section, we discuss Moy-Prasad groups over complete discrete valuation rings in the equal characteristic case. The aim is to recover these anew in the spirit of [BS]. This then gives the generalization to the higher dimensional bases.

8.1. Tameness Assumptions. In this section the given data $(\Theta, \mathbf{e}') := ((\theta_1, \mathbf{e}'_1), \dots, (\theta_n, \mathbf{e}'_n))$ will consist of a n -tuple of rational one parameter subgroups θ_i and positive rational numbers e'_i . We suppose that there exists a positive number d coprime to the characteristic of the residue field such that $d\theta_i$ is a one parameter subgroups and de'_i are positive numbers.

8.2. The unit group. Let $\mathcal{O} := k[[z]]$ and K be its quotient field. Let $B := k[[\omega]]$ and L be its quotient field, where $\omega^d = z$. We view $N := \text{Spec}(B)$ as a Galois cover over $D := \text{Spec}(\mathcal{O})$ with $\text{Gal}(N/D) = \mu_d$. Let $p: N \rightarrow D$ be the quotient morphism. Let $\rho: \mu_d \rightarrow G$ be a representation. Consider the twisted μ_d -action on $N \times G$ defined by

$$\gamma(u, g) = (\gamma u, \rho(\gamma)g) \quad \text{for } u \in N, \gamma \in \mu_d. \quad (8.2.1)$$

This defines a (μ_d, G) bundle on N . We will denote it by E .

Define the *unit group* of E as an automorphisms of the (μ_d, G) -torsor E :

$$\mathbb{U}_E := \text{Aut}_{(\mu_d, G)}(E). \quad (8.2.2)$$

Let $e \geq 0$ be a non-negative integer. Consider the thickened fibre $j_e: E|_{\text{Spec}(B/\omega^e B)} \hookrightarrow E$. Define $\mathbb{U}_E^{(e)}$ to be:

$$\mathbb{U}_E^{(e)} := \ker\left(\text{Aut}_{(\mu_d, G)}(E) \rightarrow \text{Aut}_{(\mu_d, G)}(j_e^*(E))\right) \quad (8.2.3)$$

i.e. the subgroup of (μ_d, G) automorphisms of E over N inducing identity on the restriction of E to $\text{Spec}(B/\omega^e B)$. For $e=0$, this is just the unit group \mathbb{U}_E of E .

Consider the adjoint group scheme of E

$$\mathfrak{G}_E \simeq G \times N \quad (8.2.4)$$

which is the constant group scheme together with a Γ -action.

It is well known ([Yu, 2.8, page 7]) that for each $e \geq 0$, by a *dilatation* of \mathfrak{G}_E , there exists a smooth group scheme $\mathfrak{G}_E^{(e)}$ such that

$$\mathfrak{G}_E^{(e)}(B) = \Gamma(\omega^e, \mathfrak{G}_E) \quad (8.2.5)$$

where $\Gamma(\omega^e, \mathfrak{G}_E)$ is the e^{th} - congruence subgroup defined by

$$\Gamma(\omega^e, \mathfrak{G}_E) = \ker(\mathfrak{G}_E(B) \rightarrow \mathfrak{G}_E(B/\omega^e B)).$$

8.3. The representation $\rho : \Gamma \rightarrow T$ and rational one parameter subgroups θ of T . Following [BS, Lemma 2.2.8], make the identification

$$\text{Hom}(\mu_d, T) = \frac{Y(T)}{dY(T)} \simeq \frac{1}{d} Y(T) \text{mod}(Y(T)). \quad (8.3.1)$$

We will interpret this identification explicitly to suit our specific needs. Given a one-parameter subgroup

$$\Delta : \mathbb{G}_m \longrightarrow T, \quad (8.3.2)$$

by projecting it onto $\frac{Y(T)}{dY(T)}$ and using (8.3.1), this 1 PS Δ gives a rational one-parameter subgroup $\theta_\Delta \in \frac{1}{d} Y(T) \text{mod}(Y(T))$. By (8.3.1) again, this gives a homomorphism

$$\rho_\Delta : \mu_d \rightarrow T. \quad (8.3.3)$$

In the reverse construction, given a homomorphism $\rho \in \text{Hom}(\mu_d, T)$, we choose a rational one-parameter subgroup θ_ρ determined by the class of ρ (we will need to make a choice here). Then θ_ρ defines a one-parameter subgroup

$$\Delta = d\theta_\rho : \mathbb{G}_m \rightarrow T. \quad (8.3.4)$$

Remark 8.1. The point θ_ρ determined above by the choice of Δ firstly gives a point of the Weyl alcove \mathbf{a}_0 (see §2.1). This point gives a precise *weight* as obtained by the process of invariant push-forward in [BS]. More precisely, the point to be noted is that, this construction $\rho \mapsto \theta_\rho$ gives a point of the *fundamental domain* for the action of $Y(T)$ and not just the affine Weyl group. Further, the proofs in §8.5 and that of Theorem 9.1 show that [BS, Theorem 2.3.1] and [BS, Proposition 5.1.2] hold more generally for θ_ρ in the fundamental domain of $Y(T)$ and not just the affine Weyl group.

8.4. Brief recall of Moy-Prasad groups. Moy-Prasad groups have been defined in ([MP, 2.6]). Using a Chevalley basis of G , here below we give an equivalent reformulation which is suitable for our setup. Let $\mathcal{A} = \text{Hom}(\mathbb{G}_m, T) \otimes \mathbb{R}$ denote the apartment corresponding to T . For a point $\theta \in \mathcal{A}$ recall that the parahoric group $P_\theta \subset \text{G}(K)$ is defined as

$$P_\theta = \langle T(\mathcal{O}), U_r(z^{-\lfloor r(\theta) \rfloor} \mathcal{O}), r \in \Phi \rangle \quad (8.4.1)$$

and for a non-negative integer $e' \geq 0$, the Moy-Prasad group is defined as

$$G_{\theta, e'} := \langle T(1 + z^{[e']} \mathcal{O}), U_r(z^{-\lfloor r(\theta) - e' \rfloor} \mathcal{O}), r \in \Phi \rangle \quad (8.4.2)$$

While considering parahoric or Moy-Prasad groups, it suffices to assume that $\theta \in \mathcal{A}_T$ is a rational point and $e' \in \mathbb{Q}_{\geq 0}$. This follows because for any $(r, n) \in \Phi \times \mathbb{Z}$, the functional $\mathcal{A}_T \times \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}$ defined by $(x, e') \mapsto r(x) + n - e'$ is defined over rationals.

8.5. Invariant-direct image of dilatations of reductive group schemes. The following theorem follows the proofs in [BS, Theorem 2.3.1] and [BS, Proposition 5.1.2] closely.

THEOREM 8.2. *Let $\mathfrak{G}_{\theta, e'}$ be the Bruhat-Tits group scheme satisfying $\mathfrak{G}_{\theta, e'}(\mathcal{O}) = G_{\theta, e'}$ (see 8.4.2). Then (see 8.2.5) we have a natural isomorphism of group schemes:*

$$p_*^{\mu_d}(\mathfrak{G}_E^{(e)}) \simeq \mathfrak{G}_{\theta_\rho, e'}. \quad (8.5.1)$$

Proof. Note that the \mathcal{O} -valued points are enough to determine the group scheme since $\mathfrak{G}_{\theta, e'}$ is étouffé in the sense of [BT2, page 37].

We begin with a few observations. Now an element $\phi_0 \in U_E$ corresponds to a μ_d -equivariant automorphism of the (μ_d, G) -bundle, i.e. $\phi_0 : N \times G \rightarrow N \times G$ together with

the equivariance. In other words, $\phi_0 \in U_E^{(e)}$ corresponds to a regular morphism $\phi : N \rightarrow G$ (i.e. $\phi \in G(B)$), given by

$$\phi_0(u, g) = (u, \phi(u)g), \quad (8.5.2)$$

which satisfies the following conditions

1) μ_d -equivariance i.e.

$$\phi(\gamma u) = \rho(\gamma)\phi(u)\rho(\gamma)^{-1} \quad u \in N, \gamma \in \mu_d \quad (8.5.3)$$

2) When $\underline{e} \geq \underline{1}$, the map ϕ restricted to $\text{Spec}(B/\omega^e B)$ is identity, i.e.

$$\phi \in \ker(G(B) \rightarrow G(B/\omega^e B)). \quad (8.5.4)$$

Thus, as in [BS, 2.3.1.5], we can identify U_E with ϕ satisfying (8.5.3) and the subgroup $U_E^{(e)} \subset U_E$ be the subgroup of U_E given by the above two conditions.

From here onwards, we follow closely the proof of [BS, Theorem 2.3.1].

For each $\phi \in U_E$ define

$$\psi(\omega) = \Delta^{-1} \cdot \phi \cdot \Delta. \quad (8.5.5)$$

Then $\psi(\gamma \cdot u) = \psi(u), \forall \gamma \in \mu_d, u \in N$. It therefore descends to a rational function $\tilde{\psi} : \text{Spec}(K) \rightarrow G$ such that

$$\tilde{\psi}(z) = \psi(\omega). \quad (8.5.6)$$

Now for $\underline{e} \geq \underline{1}$, by (8.5.4), we have $\phi(o) = 1$ which lies in G° , where $G^\circ \subset G$ is the big cell in G . Since N is an formal neighbourhood of the origin o , it follows that $\phi(N) \subset G^\circ$.

For $\underline{e} = \underline{0}$, in the following we will assume that we work with a ϕ such that $\phi(N) \subset G^\circ$. The set of such ϕ may be called the big cell of U_E and we will denote it by the symbol \mathfrak{U} .

The condition $\phi(N) \subset G^\circ$ allows us to describe $\phi : N \rightarrow G^\circ$ uniquely as a tuple of the following morphisms

$$\{ \{ \phi_r : N \rightarrow U_r \}_{r \in \Phi}, \phi_t : N \rightarrow T \},$$

where $U_r \subset G$ is the root subgroup corresponding to $r \in \Phi$. Let us view ϕ_r and ϕ_t as elements of $G(B) \subset G(L)$.

We may write

$$\phi_r(\omega) = \psi_r(\omega)\omega^{r(\Delta)} = \tilde{\psi}_r(z)\omega^{r(\Delta)}. \quad (8.5.7)$$

Since $\phi_r(\omega)$ has a zero of order at least e in ω ,

$$\frac{\phi_r(\omega)}{\omega^e} = \tilde{\psi}_r(z)\omega^{r(\Delta)-e},$$

and the right side defines a regular function in ω i.e. it extends to a regular function on N . So $\tilde{\psi}_r(z)$ can have a pole in ω of order at most $r(\Delta) - e$. Thus in z it can have a pole of order at most

$$\lceil \frac{r(\Delta) - e}{d} \rceil.$$

i.e. $= \lceil r(\frac{\Delta}{d}) - \frac{e}{d} \rceil \stackrel{(8.3.4)}{=} \lceil r(\theta_\rho) - e' \rceil$, where $e' = \frac{e}{d}$. This forces the containment

$$\tilde{\psi}_r(z) \in U_r(z^{-\lceil r(\theta_\rho) - e' \rceil} \mathcal{O}), \quad (8.5.8)$$

and hence $\tilde{\psi}_r$ belongs to $G_{\theta_\rho, e'}$. On the other hand, since $\phi_t(u) \in T$ for $u \in N$, it follows that $\phi_t(\gamma \cdot u) = \rho(\gamma)\phi_t(u)\rho(\gamma)^{-1} = \phi_t(u)$. Hence $\phi_t : N \rightarrow T$ is μ_d -invariant. Also by

(8.5.5) it follows that $\phi_t = \psi_t$. So it descends to a regular function on D and this function is $\tilde{\psi}_t : D \rightarrow T$. Now $\phi_t(u) - 1$ has a zero of order at least e in ω . So by definition of e' we see that $\tilde{\psi}_t - 1$ has a zero in z of order at least

$$\lceil \frac{e}{d} \rceil = \lceil e' \rceil.$$

Hence $\tilde{\psi}_t$ also belongs to $G_{\theta_\rho, e'}$.

Since ψ and ϕ are related by a conjugation in $G(L)$ by $\Delta(\omega)$, ψ like ϕ can also be described by ψ_r and ψ_t uniquely. So the same holds for $\tilde{\psi}$. Hence since $\tilde{\psi}_r$ and $\tilde{\psi}_t$ belong to $G_{\theta_\rho, e'}$, therefore so does $\tilde{\psi} \in G_{\theta_\rho, e'}$.

For $e \geq 1$ ¹, we get the following isomorphism of groups in the spirit of [BS, Theorem 2.3.1]

$$U_E^{(e)} \simeq G_{\theta_\rho, e'}. \quad (8.5.10)$$

Combining these with (8.5.3) and (8.5.4), we deduce that for $e \geq 1$:

$$p_*^{\mu_d}(\mathfrak{G}_E^{(e)})(A) = \mathfrak{G}_E^{(e)}(B)^{\mu_d} = U_E^{(e)} \simeq G_{\theta_\rho, e'} \simeq \mathfrak{G}_{\theta_\rho, e'}(A).$$

Q.E.D

Remark 8.3. Note that the existence and defining properties for BT-group schemes which realize Moy-Prasad groups were constructed in [Yu]. They arise from an enlargement of the definition of concave functions and constructing BT group schemes associated to them. We observe that the above realization of these group schemes as *invariant direct images* can be interpreted as giving a different construction of these group schemes, namely as invariant direct images of congruence subgroup schemes from ramified covers. We have done this in the simple case field of characteristic zero, and this can be extended to "good" characteristics for the group G .

8.6. The n -Moy-Prasad-group scheme. The proof of (5.7) does not quite apply as it is. Unlike the scenario of (5.7), on the Kawamata cover we no longer have a constant group scheme with fibre G but have the *congruence subgroup schemes*.

THEOREM 8.4. *We have an n -Moy-Prasad-group scheme $\mathfrak{G}_{(\Theta, \mathbf{e}')}^{(\Theta, \mathbf{e}'')}$ which is smooth and affine on $\text{Spec}(\mathcal{O})$ which is given by the data of $(\Theta, \mathbf{e}') := ((\theta_1, \mathbf{e}'_1), \dots, (\theta_n, \mathbf{e}'_n))$. It satisfies the properties as in (1.3).*

Proof. For the given data $(\Theta, \mathbf{e}') := ((\theta_1, \mathbf{e}'_1), \dots, (\theta_n, \mathbf{e}'_n))$, under assumptions of §8.1, setting $\Theta = (\theta_1, \dots, \theta_n)$, we have a Kawamata cover $q : Z \rightarrow \mathbf{A}$ which recovers the n -parahoric group scheme \mathfrak{G}_Θ on \mathbf{A} as the invariant push-forward $q_*^\Gamma(\mathcal{H})$, where \mathcal{H} is a group scheme on Z with fibres G . Recall q is ramified with varying ramification indices d_i . In fact, by our tameness hypothesis, the ramification indices d_i are such that $e_i := d_i e'_i$ are positive integers which are coprime to the residue characteristic. Without loss of generality, we may suppose that the e_i 's are *increasing with i* .

Set $e_0 = 0$. Now for j going from 1 to n , we use the main result in [MRR] to construct an iterated dilatation of \mathcal{H} along each component $q^*(\cup_{i \geq j} D_i)_{\text{red}}$ of the reduced divisor in Z thickened by $e_j - e_{j-1}$. We get a new smooth, affine group scheme $\tilde{\mathcal{H}}$ on Z . By the

¹For $e = 0$, we get an isomorphism

$$j : \mathfrak{U} \rightarrow \mathfrak{B} \quad (8.5.9)$$

of the big cell \mathfrak{U} of U_E with that of G_{θ_ρ} which we will denote by \mathfrak{B} .

universal property of dilatation it follows that $q_*^\Gamma(\tilde{\mathcal{H}})$ is the required n-Moy-Prasad-group scheme on \mathbf{A} . It is smooth and affine since $\tilde{\mathcal{H}}$ is so and because the invariant direct image functor preserves these properties. Q.E.D

9. SCHEMATIZATION IN THE MIXED CHARACTERISTIC CASE

Let $(\mathcal{O}, \mathfrak{m})$ be a complete discrete valuation ring with residue field k of characteristic p and uniformizer z . We want to extend all the results of the previous sections to the mixed characteristics case. This will be possible under assumptions in §2.2.

9.1. Extension of [BS, Theorem 2.3.1] and [BS, Proposition 5.1.2]. Let $\theta \in \mathcal{A}_T$ be a rational point satisfying the assumptions in §2.2. As in §2.2, we assume that d is coprime to $\text{char}(k)$ such that $d\theta \in Y(T)$. Consider the Eisenstein polynomial $f(x) = x^d - z \in A[x]$. Let $B := \mathcal{O}[x]/f(x)$. Thus z is totally ramified in B and the residue field k_B has d -distinct roots of unity. Let ω be a uniformizer in B such that $\omega^d = z$. Let ζ be a primitive d -th root of unity in B .

Let K be its quotient field of \mathcal{O} and L the quotient field of B . We view $N := \text{Spec}(B)$ as a Galois cover over $D := \text{Spec}(\mathcal{O})$ with $\text{Gal}(N/D) = \mu_d$. Let $p : N \rightarrow D$ be the quotient morphism. Let E be the trivial principal G bundle $N \times G$ together with the twisted μ_d action as in (8.2.1). This makes E into a (μ_d, G) bundle on N . Let \mathfrak{G}_E be its adjoint group scheme (see 8.2.4). Let $\mathfrak{G}_{\theta_\rho}$ be the Bruhat-Tits group scheme over \mathcal{O} associated to it as in Theorem (8.2). Analogous to (1.0.2), it will be convenient to denote

$$P_\theta = \langle T(\mathcal{O}), U_r(z^{m_r(\theta)} \mathcal{O}), r \in \Phi \rangle \quad (9.1.1)$$

both in the equal and mixed characteristic cases. This abuse of notation in the mixed characteristic case should not cause any confusion because of the structure theorem for complete discrete valuation rings.

Following the discussion in §8 under the tameness assumptions, let us prove the required generalization of [BS, Theorem 2.3.1] which is (8.2) for $e' = 0$.

THEOREM 9.1. *We have an isomorphism of group schemes:*

$$p_*^{\mu_d}(\mathfrak{G}_E) \simeq \mathfrak{G}_{\theta_\rho}. \quad (9.1.2)$$

Proof. Let us state at the outset that by [BT2, §1, page 13] the references we will cite are valid for an infinite field K containing a subring A such that K is its quotient field.

The \mathcal{O} group scheme $\mathfrak{G}_{\theta_\rho}$ is connected and smooth because it is a parahoric group scheme. Weil restriction of scalars followed by taking invariants preserves connectedness. Further, Weil restrictions of smooth group schemes are smooth. Moreover, under the tameness assumptions, it is checked easily that the fixed point subscheme is also smooth. Thus, the \mathcal{O} group scheme $p_*^{\mu_d}(\mathfrak{G}_E)$ is connected and smooth. As in 8.5.3, the restriction of $p_*^{\mu_d}(\mathfrak{G}_E)$ to $\text{Spec } K$ consists of $\phi : \text{Spec } L \rightarrow G$ satisfying the μ_d -equivariance i.e.

$$\phi(\gamma u) = \rho(\gamma)\phi(u)\rho(\gamma)^{-1} \quad u \in \text{Spec } L, \gamma \in \mu_d \quad (9.1.3)$$

Conjugating ϕ by Δ (see 8.3.4), as it also happened in 8.5.5, the conjugate $\Delta^{-1}\phi\Delta$ descends to a rational function $\tilde{\psi} : \text{Spec } K \rightarrow G$ as in 8.5.6. This defines on $\text{Spec } K$ a morphism

$$j_K : p_*^{\mu_d}(\mathfrak{G}_E)|_K \rightarrow \mathfrak{G}_{\theta_\rho}|_K. \quad (9.1.4)$$

By evaluating on sections, it may be verified that j_K is an isomorphism.

Recall that in the proof of (8.2), we defined the big cell \mathfrak{U} of the unit group U_E (see 8.2.2) to be those $\phi : \text{Spec } B \rightarrow G$ whose image lies in the big cell G° of G . This defines the big cell of $p_*^{\mu_d}(\mathfrak{G}_E)$. We will denote it as \mathfrak{U} as well. Taking \mathfrak{U} to be the open neighbourhood of identity of $p_*^{\mu_d}(\mathfrak{G}_E)$, we exhibited in 8.5.9 the isomorphism $j : \mathfrak{U} \rightarrow \mathfrak{B}$ onto the big cell \mathfrak{B} of $\mathfrak{G}_{\theta\rho}$.

Now by [BT2, §1.2.13], j_K extends uniquely to a \mathcal{O} -morphism

$$j : p_*^{\mu_d}(\mathfrak{G}_E) \rightarrow \mathfrak{G}_{\theta\rho}$$

of group schemes. Since j restrict to an isomorphism from \mathfrak{U} to \mathfrak{B} between big cells, by [BT2, §1.2.14] it follows that j is an isomorphism of $p_*^{\mu_d}(\mathfrak{G}_E)$ onto an open subgroup scheme of $\mathfrak{G}_{\theta\rho}$. Thus j is an isomorphism because $\mathfrak{G}_{\theta\rho}$ is connected. Q.E.D

The preceding theorem has two consequences. Firstly, [BS, Proposition 5.1.2] recalled in (5.3.2) generalizes following [BS, §4.5]. Secondly, (8.2) extends without any extra conditions imposed by e' . This is so because in the proof of (8.2) we only need to have distinct roots of unity corresponding to the point θ in the apartment.

Remark 9.2. We observe that for the results of sections §5, §6, §7, the assumption of algebraic closedness of the residue field k is not essential. The existence of a Kawamata cover (see §12.1) is needed only for affine spaces and this can be achieved by hand under assumptions of §2.2. We however always assume that k is *perfect* so as to be able to apply [BT2]. One needs to have extensions k_b as above with the desired roots of unity for the above extensions of [BS, Theorem 2.3.1] and [BS, Proposition 5.1.2]. This is possible under assumptions of §2.2.

9.2. The extension of THEOREM 5.7. In this paragraph, we will denote the uniformizer z of \mathcal{O} by x_0 . Let $\mathbf{A}_\theta := \mathbb{A}_\theta^n = \text{Spec } \mathcal{O}[x_1, \dots, x_n]$. For $0 \leq i \leq n$, let H_i be the coordinate hyperplanes defined by the vanishing of x_i and let ζ_i be their generic points.

- (1) If we let $(X, D) = (\mathbf{A}_\theta, H)$, then the analogue of (5.6) holds without any change for an arbitrary $(n+1)$ -concave-function \mathbf{f} . *This statement holds for any characteristic of the residue field.* In the following we denote the Lie algebra bundle by \mathcal{R} .
- (2) Observe that since \mathcal{O} is assumed to be complete, the formal power series ring $\mathcal{O}[[x_1, \dots, x_n]]$ is the completion of $\mathcal{O}[x_1, \dots, x_n]$ at $\mathfrak{m}[x_1, \dots, x_n]$. We begin with a natural extension of the notion of a $(n+1)$ -bounded group associated to a $(n+1)$ -concave function \mathbf{f} on the root system Φ of G by defining

$$P_{\mathbf{f}} := \langle T(\mathcal{O}[[x_1, \dots, x_n]]), U_r \left(\prod_{0 \leq i \leq n} x_i^{f_i(r)} \mathcal{O}[[x_1, \dots, x_n]] \right), r \in \Phi \rangle. \quad (9.2.1)$$

- (3) Suppose we are given an ordered $(n+1)$ -tuple $(\Sigma_0, \dots, \Sigma_n)$ of facets of the apartment \mathcal{A}_T . Then by §2.2, we may choose a point $\Theta := (\theta_0, \dots, \theta_n)$ in $(\Sigma_0, \dots, \Sigma_n)$ such that if d_i is minimum integer such that $d_i \theta_i$ lies in $Y(T)$, then each d_i is coprime to $\text{char}(k)$ and k contains primitive d_i -th roots of unity. More generally, suppose we are given a point $\Theta := (\theta_0, \dots, \theta_n) \in \mathcal{A}_T^n$ satisfying assumptions in §2.2. Let $\widehat{\mathcal{O}}$ be a complete discrete valuation ring containing \mathcal{O} where x_0 ramifies to order a multiple of d_0 and it contains the primitive d_i -th roots of unity for $0 \leq i \leq n$.
- (4) Let $X' \subset \mathbf{A}_\theta$ denote the open subset of height 1 prime ideals as in the paragraph after (5.4.2). Taking \mathbf{f} to be the concave function determined by θ in (1), let \mathcal{R}' denote the restriction of \mathcal{R} to X' .
- (5) We are now in the setting of (5.4) with the only difference that \mathbf{A}_θ replaces \mathbb{A}_k^{n+1} .

- (6) Let $\mathbf{B}_\theta := \text{Spec } \tilde{\mathcal{O}}[\mathbf{y}_1, \dots, \mathbf{y}_n]$ where $y_i^{d_i} = x_i$. Let $q : \mathbf{B}_\theta \rightarrow \mathbf{A}_\theta$ be the natural projection map with Galois group Γ . This plays the role of an explicit Kawamata covering. We set $\mathbf{A}_0 := \mathbf{A}_\theta \setminus H$, $\mathbf{B}_0 := q^{-1}(\mathbf{A}_0)$, $Z' := q^{-1}(X')$ and $\tilde{X}_i := q^{-1}(X_i)$. Let $q_i : \tilde{X}_i \rightarrow X_i$ be the restriction of q . Let Γ_i denote the Galois group of q_i . We are now in the setting of §5.3.
- (7) By applying §9.1 to the uniformizer x_0 and the generalization of [BS, Proposition 5.1.2] for the rest of the height 1 primes given by the variables x_j , we get a smooth, affine group scheme \mathcal{H}' on Z' such that the Weil restrictions gives $q_*^\Gamma(\mathcal{H}') = \mathcal{G}'$ on X' .
- (8) Let $V' = \text{Lie}(\mathcal{H}')$ be the Lie algebra bundle on Z' as in the discussion after (5.4.4). Then, V' is a Lie algebra bundle with fibre type \mathfrak{g} . Furthermore, the Cartan decomposition on \mathcal{H}' lifts to give an equivariant Cartan decomposition of the Lie algebra bundle V' on Z' . Hence its reflexive closure V also gets an equivariant structure and is also locally free.
- (9) As in the proof of (5.4), now that we have the Lie algebra bundle V with fibre \mathfrak{g} , we get a group scheme \mathcal{H} on \mathbf{B}_θ and by taking invariant direct images, the group scheme $\mathcal{G} := q_*^\Gamma(\mathcal{H})$ extending \mathcal{G}' to the whole of \mathbf{A}_θ . This proves the extension of Theorem 5.7 in the setting of mixed characteristics with the tameness assumptions.
- (10) We also note that as in (5.5), the big cell structure also descends to give one for \mathcal{G} on the whole of \mathbf{A}_θ .
- (11) Finally, we note that the arguments in (5.7) for the uniqueness of the group scheme carry over to the mixed characteristic situation as well.

9.3. The main theorem for concave functions. By (9.2 (1)), we see that there is a Lie algebra bundle on (\mathbf{A}_θ, H) associated to a $(n+1)$ -concave function on Φ . The results in §6 now work under the tameness assumptions (1.1) or more generally under the assumptions on $\text{char}(k)$ of §2.2. We get therefore the extension of THEOREM 1.3 to (\mathbf{A}_θ, H) (see (6.10)). Note that this is for the groups $G = \text{SL}(n), G_2, F_4, E_6$.

For the rest of the groups by the arguments in §7, we need a faithful representation $G \hookrightarrow \text{GL}(V)$ to start with and so if we choose a faithful representation of minimal dimension $m(G)$, then for these groups, we need to choose p to be coprime to $m(G)$ and bigger than the Coxeter number h_G of G (see §7). The numbers $m(G)$ are known and can be obtained from the dimension formulae of the fundamental representations. They are as follows: $m(E_6) = 27$, $m(E_7) = 56$ and for B_n, C_n and D_n it is $2n+1$ (resp. $2n, 2n$). In summary we have completed the proof of THEOREM 1.6.

9.4. The example of a 2BT-group scheme in the work of Pappas and Zhu [PZ]. As we mentioned in the introduction, an example of a 2BT-group scheme comes up in [PZ, Theorem 4.1].

There are broadly two parts to [PZ, Theorem 4.1], the first being the existence of the 2BT-group scheme on \mathbb{A}_θ^1 and second proving that the resulting group scheme is *affine*. Although the theorem [PZ, Theorem 4.1] states the case of a parahoric group scheme along one axis, the proof considers the more general case of the 2-concave function $\mathbf{f} = (0, f)$, where f is a concave function on the root system Φ . The proof of the affineness follows ideas from [Yu]. We are unable to comprehend it when f is given by a non-hyperspecial point of the apartment and G is split.

We first briefly describe the setting in [PZ, Section 4] so as to place their result in our context. We will then indicate how this result can be derived as a special case of our

results. We note however that the setting in [PZ] is for the general case of a quasi-split group G , while we consider only the split case.

In the notation of 9.2 above, the setting in [PZ] is for the case $n = 1$, i.e., when $x_1 = u$. The axes we consider correspond to setting x_0 and x_1 as zero (here by $x_0 = 0$ we mean going modulo the uniformizer), but, Pappas-Zhu consider the axes given by setting u as the uniformizer ϖ and the one given by setting $\varpi = 0$. The result [PZ, Theorem 4.1] is stated for the case when the assignment along the x_0 -axis is the split group (see [PZ, 4.2.2(a)]) while the assignment in the diagonal $\varpi = u$ -direction is given by a parahoric group associated to a point θ of the affine apartment. Thus, in our setting this case corresponds to the 2-concave function which is $\mathbf{f} = (0, f)$, where f is a concave function f_θ on the root system Φ along the u -direction and the 0-concave function gives the split group scheme along the ϖ -direction. Observe that this gives the concave function $f = 0 + f$ in the $u = \varpi$ -direction.

The key point is that only *a single* concave function f is involved. By [BT2, §4, 4.5.4], one obtains a schematic root datum coming from f . We now work in the setting of [BT2, 3.9.4]. Once a faithful representation $G \hookrightarrow \mathrm{GL}(V)$ is fixed, then the schematic root datum arising out of the choice of f entails the choice of a lattice $M \subset V_\kappa$. The arguments in [BT2, 3.9.4] furnishes the group scheme $\mathcal{G}_{\theta_0, f}$ (see [PZ, paragraph 2, page 175]). Thus, the 2BT-group scheme over \mathbb{A}_θ^1 which comes in [PZ] is obtained as a direct consequence of [BT2, 3.9.4]. However, the method of construction of [BT2, 3.9.4 and 2.2.10] gives *a priori* only a *quasi-affine group scheme*. So the burden of [PZ, Section 4] lies in proving the *affineness* of the group scheme $\mathcal{G}_{\theta_0, f}$. We emphasize that all this is for a single concave function.

We now compare this with the results of the present paper. Firstly, [PZ, Theorem 4.1], as stated is for the 2-concave function $(0, f_\theta)$ and this is a special case of THEOREM 5.7 as described in 9.2. Thus we derive directly both the *existence and affineness* of the required group scheme and in the more general context of an $(n + 1)$ -concave function over \mathbb{A}_θ^n which comes by prescribing $n + 1$ points in the affine apartment. We note however that the methods in the present paper force certain mild tameness assumptions.

Part 3. SOME APPLICATIONS IN $\mathrm{CHAR}(k) = 0$ EQUICARACTERISTIC CASE

10. BT-GROUP SCHEMES ON WONDERFUL EMBEDDINGS IN $\mathrm{CHAR}(k) = 0$

10.1. The wonderful compactification \mathbf{X} . Let $\mathbf{X} := \overline{G_{\mathrm{ad}}}$ be the wonderful compactification of G_{ad} . Let $\{D_\alpha | \alpha \in S\}$ denote the irreducible smooth *divisors* of \mathbf{X} . Let $D := \cup_{\alpha \in S} D_\alpha$. Then $\mathbf{X} \setminus G_{\mathrm{ad}} = D$. The pair (\mathbf{X}, D) is the primary example of a $(G_{\mathrm{ad}} \times G_{\mathrm{ad}})$ -homogenous pair [Br2, Brion]. Let P_I be the standard parabolic subgroup defined by subsets $I \subset S$, the notation being such that the Levi subgroup $L_{I, \mathrm{ad}}$ containing T_{ad} has root system with basis $S \setminus I$. Recall that the $(G_{\mathrm{ad}} \times G_{\mathrm{ad}})$ -orbits in \mathbf{X} are indexed by subsets $I \subset S$ and have the following description:

$$Z_I = (G_{\mathrm{ad}} \times G_{\mathrm{ad}}) \times_{P_I \times P_I^-} L_{I, \mathrm{ad}}.$$

Then by [Br1, Proposition A1], each Z_I contains a unique base point z_I such that $(B \times B^-) \cdot z_I$ is dense in Z_I and there is a 1-PS λ of T satisfying $P_I = P(\lambda)$ and $\lim_{t \rightarrow 0} \lambda(t) = z_I$. The closures of these λ define curves $C_I \subset \mathbf{X}$ which meet the strata Z_I transversally at z_I . In particular, if $I = \{\alpha\}$ a singleton, then the divisor D_α is the orbit closure \bar{Z}_I and the 1-PS can be taken to be the fundamental co-weight ω_α^\vee . The closure of the 1-PS

$\omega_\alpha^\vee : \mathbb{G}_m \rightarrow G_{\text{ad}}$ defines the curve $C_\alpha \subset \mathbf{X}$ transversal to the divisor D_α at the point z_α . For a non-empty $I \subset S$, the λ defining C_I can be taken to be $\sum_{\alpha \in I} \omega_\alpha^\vee$.

10.2. The ℓ BT-group scheme $\mathfrak{G}_\mathbf{X}^\circ$ on \mathbf{X} . Before stating the main result of this section, we make a few remarks which might help the reader. Let $\mathbf{Y}_0 \subset \mathbf{Y}$ the affine toric varieties associated to the Weyl chamber and the fan of Weyl chambers. The basic underlying principle in these constructions is that the combinatorial data encoded in the triple consisting of the Weyl chamber, the fan of Weyl chambers, and the Tits building, is geometrically replicated by the inclusion $\mathbf{Y}_0 \subset \mathbf{Y} \subset \mathbf{X}$. The "wonderful" Bruhat-Tits group scheme which arises on \mathbf{X} has its local Weyl-chamber model on the affine toric variety \mathbf{Y}_0 . Indeed, in this case, the Kawamata cover is even explicit (4.6). In particular, the theorem below, can be executed for \mathbf{Y}_0 but this will give the group scheme associated to the data coming from the Weyl chamber alone.

We now state a consequence of our main theorem in the context of the wonderful compactification of G_{ad} which is a primary example of a smooth variety with normal crossing divisors.

THEOREM 10.1. *Let $\Theta = (\theta_\alpha \mid \alpha \in S)$ where $\theta_\alpha \in \mathcal{A}$ are arbitrary points in the apartment of T satisfying conditions of §2.2. There exists an affine "wonderful" Bruhat-Tits group scheme $\mathfrak{G}_\mathbf{X}^\circ$ on \mathbf{X} satisfying the following classifying properties:*

- (1) *At the height 1 primes associated to the canonical boundary divisors associated to the maximal parabolic subgroups $P_\alpha \subset G, \alpha \in S$, the group scheme $\mathfrak{G}_\mathbf{X}^\circ$ is isomorphic to the parahoric group scheme $\mathfrak{G}_{\theta_\alpha}$.*
- (2) *For $\emptyset \neq I \subset S$ the restriction of $\mathfrak{G}_\mathbf{X}^\circ$ to the formal neighbourhood U_I of z_I in the coordinate curves C_I §(10.1) is isomorphic to the Bruhat-Tits group scheme associated to the concave function $m_{\theta_I} : \Phi \rightarrow \mathbb{Z}$ given by $r \mapsto \sum_{\alpha \in I} m_r(\theta_\alpha)$.*
- (3) *Let $\Phi_\theta := \{r \in \Phi \mid r(\theta) \in \mathbb{Z}\}$. The closed fibre of $\mathfrak{G}_{m_{\theta_I}}$ modulo its unipotent radical has root system generated by the intersection of Φ_{θ_α} for $\alpha \in I$.*

Proof. By Proposition 5.6 we have a Lie-algebra bundle \mathcal{R} on \mathbf{X} prescribed by the first property. Thus by Theorem 5.4, we get a group scheme $\mathfrak{G}_\mathbf{X}^\circ$ on \mathbf{X} which by construction satisfies the first property. The context of the second property is that of formal neighbourhoods which is also that of §5.8. As remarked before the theorem, the group scheme constructed on \mathbf{X} has a local Weyl-chamber model on the affine toric variety \mathbf{Y}_0 . Thus we may restrict to \mathbf{Y}_0 . Whence the second property follows from Theorem 5.7. Recall [BT2, 4.6.12, page 129], that the reductive quotient of the closed fibre of \mathfrak{G}_f has root system given by $\{r \in \Phi \mid [f](-r) = -[f](r)\}$. For the case of the concave function $m_r(\theta)$ this set is Φ_θ and for m_{θ_I} it will be the intersection of Φ_{θ_α} for $\alpha \in I$. Q.E.D

COROLLARY 10.2. *Let θ_α be the non-zero vertices of the Weyl alcove (2.1.1). Let L be the LCM of the coefficients of the highest root and ℓ the semisimple rank of G . Then setting Θ as $(\theta_\alpha \mid \alpha \in S), (\theta_\alpha / (\ell + 1) \mid \alpha \in S)$ and $(L\theta_\alpha \mid \alpha \in S)$ gives a group scheme whose restriction to every coordinate curve C_I has closed fibre with root system generated by the highest root α_0 and $S \setminus I$, by $S \setminus I$ or that of G respectively.*

Proof. The structure of the set $\{r \in \Phi \mid [f](-r) = -[f](r)\}$ and the formula (2.1.1), shows that the root system of the reductive quotient of the closed fibre equals the set of roots whose simple root coefficients, coming from the subset I of S , equals an integral multiple of the corresponding term of the highest root. In the first case, this is the set of

roots generated by the highest root α_0 and $S \setminus I$. In the second case, it is the set of roots generated by $S \setminus I$. In the third case, all roots in Φ satisfy this property. Q.E.D

10.3. The wonderful embedding \mathbf{X}^{aff} . We continue to use the notations as in previous sections. Let G^{aff} denote the Kac-Moody group associated to the affine Dynkin diagram of G . Recall that G^{aff} is given by a central extension of $L^\times G$ by \mathbb{G}_m . Analogous to the wonderful compactification of G_{ad} , P. Solis in [So] has constructed a wonderful embedding \mathbf{X}^{aff} for $G_{\text{ad}}^{\text{aff}} := G^{\text{aff}}/Z(G^{\text{aff}}) = \mathbb{G}_m \times LG/Z(G)$. It is an ind-scheme containing $G_{\text{ad}}^{\text{aff}}$ as a dense open ind-scheme and carrying an equivariant action of $L^\times G \times L^\times G$.

Let $T_{\text{ad}} := T/Z(G)$, $T_{\text{ad}}^\times := \mathbb{G}_m \times T_{\text{ad}} \subset G_{\text{ad}}^{\text{aff}}$, where \mathbb{G}_m is the rotational torus. In \mathbf{X}^{aff} , the closure $\mathbf{Y}^{\text{aff}} := \overline{T_{\text{ad}}^\times}$ gives a torus-embedding. It is covered by the affine Weyl group W^{aff} -translates of the affine torus embedding $\mathbf{Y}_0^{\text{aff}} := \overline{T_{\text{ad},0}^\times} \simeq \mathbb{A}^{\ell+1}$ given by the negative Weyl alcove.

10.4. On the torus-embedding $\mathbf{Y}_0^{\text{aff}}$. Recall that $Z := \mathbf{Y}_0^{\text{aff}} \setminus T_{\text{ad}}^\times$ is a union $\cup_{\alpha \in \mathbb{S}} H_\alpha$ of $\ell+1$ many standard coordinate hyperplanes meeting at normal crossings. For $\alpha \in \mathbb{S}$, let the ζ_α 's denote the generic points of the divisors H_α 's. Let

$$A_\alpha = \mathcal{O}_{\mathbf{Y}_0^{\text{aff}}, \zeta_\alpha} \tag{10.4.1}$$

be the dvr's obtained by localizing at the height 1-primes given by the ζ_α 's. Let K_α be the quotient field of A_α . Let $Y_\alpha := \text{Spec}(A_\alpha)$. Note that we can identify the open subset $\text{Spec}(K_\alpha)$ with $T_{\text{ad}}^\times \cap Y_\alpha$. Let

$$Y' := T_{\text{ad}}^\times \cup_\alpha Y_\alpha. \tag{10.4.2}$$

The complement $\mathbf{Y}_0^{\text{aff}} \setminus Y'$ can again be realized as a colimit of open subsets of $\mathbf{Y}_0^{\text{aff}}$ whose codimension is at least 2 in $\mathbf{Y}_0^{\text{aff}}$.

10.5. Construction of a finite-dimensional Lie algebra bundle J on $\mathbf{Y}_0^{\text{aff}}$ together with parabolic structures. This construction is exactly analogous to the construction of \mathcal{R} on \mathbf{Y} in §4. For $\alpha \in \mathbb{S}$, let $\theta_\alpha \in \mathcal{A}$ be arbitrary points in the apartment satisfying conditions of §2.2. We let T_{ad}^\times , $\mathbf{Y}_0^{\text{aff}}$, $(1, \omega_\alpha^\vee)$ and \mathbb{S} play the role of \mathbf{A}_0 , \mathbf{A} , λ_i and $\{1, \dots, n\}$. More precisely, we set

$$\eta_\alpha := ((0, \dots, 1, \dots, 0), \theta_\alpha), \quad \alpha \in \mathbb{S} \tag{10.5.1}$$

with the unique 1 in the α -th coordinate. Consider the loop rotation action given by (2.5.1) and (2.5.2). We may prove the following theorem exactly like (4.4).

THEOREM 10.3. *Let $\Theta := (\theta_\alpha \mid \alpha \in \mathbb{S})$ where θ_α are arbitrary points in the apartment \mathcal{A} of T satisfying conditions of §2.2. Let U be a formal neighbourhood of origin in $\mathbf{Y}_0^{\text{aff}}$. There exists a canonical Lie-algebra bundle J on $\mathbf{Y}_0^{\text{aff}}$ which extends the trivial bundle with fiber \mathfrak{g} on $T_{\text{ad}}^\times \subset \mathbf{Y}_0^{\text{aff}}$ and with notations as in (2.5.7) we have the identification of functors from the category of k -algebras to k -Lie-algebras:*

$$L^+(\mathfrak{P}_\Theta) = L^+(\mathcal{R}|_U). \tag{10.5.2}$$

We may prove the following corollary like (4.5).

COROLLARY 10.4. Let $\lambda = \sum_{\alpha \in \mathbb{S}} k_\alpha (1, \theta_\alpha)$ be a non-zero dominant 1-PS of T_{ad}^\times where not all k_α are zero. Let \mathcal{D} be the formal neighbourhood of origin in $\mathbf{Y}_0^{\text{aff}}$ associated to the curve defined by λ . Let $f : \Phi \rightarrow \mathbb{Z}$ be the concave function defined by the assignment $r \mapsto \sum_{\alpha \in \mathbb{S}} k_\alpha m_r(\theta_\alpha)$. Then for any k -algebra R we have

$$L^+(\mathcal{R}|_{\mathcal{D}})(R) = \langle \mathfrak{t}(R[[t]]), \mathfrak{u}_r(t^{f(r)} R[[t]]), r \in \Phi \rangle. \quad (10.5.3)$$

COROLLARY 10.5. The Lie algebra bundle $J_{\mathbf{Y}^{\text{aff}}}$ gets canonical parabolic structures (§12) at the generic points ξ_α of the W^{aff} -translates of the divisors $H_\alpha \subset \mathbf{Y}_0^{\text{aff}}$, $\alpha \in \mathbb{S}$.

Proof. We prescribe ramification indices d_α on the divisor H_α such that $d_\alpha \theta_\alpha$ belongs to $Y(T)$. (Thus in the case d_α are the alcove vertices (2.1.1), then d_α are given by (2.1.2)). Then the identification (5.3.3) of the Lie algebra structures of $J_{\mathbf{Y}^{\text{aff}}}$ and the parahoric Lie algebra structures on the localizations of the generic points of H_α allows us to endow parabolic structures at the generic points of the divisors. Q. E. D

10.6. The parahoric group scheme on the torus embedding \mathbf{Y}^{aff} .

THEOREM 10.6. Let $\Theta = (\theta_\alpha \mid \alpha \in \mathbb{S})$ where $\theta_\alpha \in \mathcal{A}$ are arbitrary points in the apartment of T satisfying conditions of §2.2. There exists an affine and smooth “wonderful” Bruhat-Tits group scheme $\mathfrak{G}_{\mathbf{Y}^{\text{aff}}}^\Theta$ on \mathbf{Y} together with a canonical isomorphism $\text{Lie}(\mathfrak{G}_{\mathbf{Y}^{\text{aff}}}^\Theta) \simeq J$. It further satisfies the following classifying property:

For any point $h \in \mathbf{Y}^{\text{aff}} \setminus T_{ad}^\times$, let $\mathfrak{l} \subset \mathbb{S}$ be a subset such that $h \in \cap_{\alpha \in \mathfrak{l}} H_\alpha$. Let $C_{\mathfrak{l}} \subset \mathbf{Y}^{\text{aff}}$ be a smooth curve with generic point in T_{ad}^\times and closed point h . Let $U_{\mathfrak{l}} \subset C_{\mathfrak{l}}$ be a formal neighbourhood of h . Then, the restriction $\mathfrak{G}_{\mathbf{Y}^{\text{aff}}}^\Theta|_{U_{\mathfrak{l}}}$ is isomorphic to the Bruhat-Tits group scheme associated to the concave function $m_{\Theta, \mathfrak{l}} : \Phi \rightarrow \mathbb{Z}$ given by $r \mapsto \sum_{\alpha \in \mathfrak{l}} m_r(\theta_\alpha)$.

Proof. Recall that \mathbf{Y}^{aff} is covered by affine spaces $Y_w \simeq \mathbb{A}^{\ell+1}$ parametrized by the affine Weyl group W^{aff} . Each Y_w is the translate of $\mathbf{Y}_0^{\text{aff}}$. The translates of the divisors H_α meet each Y_w in the standard hyperplanes on $\mathbb{A}^{\ell+1}$ and Thus, we can prescribe the same ramification data at the hyperplanes on each of the Y_w 's. On the other hand, although we have simple normal crossing singularities, we do not have an analogue of the Kawamata covering lemma for schemes such as \mathbf{Y}^{aff} . The lemma is known only in the setting of quasi-projective schemes. So to construct the group scheme, we employ a different approach using the naturality of the constructions for gluing.

We observe firstly that, the formalism of Kawamata coverings applies in the setting of the affine spaces $Y_w \subset \mathbf{Y}^{\text{aff}}$. Let $p_w : Z_w \rightarrow Y_w$ be the associated Kawamata cover (12.1) with Galois group Γ_w . In Corollary 10.5 we observed that the Lie-algebra bundle J has a canonical parabolic structure. Letting Y_w play the role of \mathbf{Y}_0 and using all arguments in the proof of Theorem 10.1 we obtain $\mathcal{H}_w \rightarrow Z_w$ which is Γ_w -group scheme with fibers isomorphic to G whose invariant direct image is a group scheme \mathfrak{G}_w such that $\text{Lie}(\mathfrak{G}_w) = J|_{Y_w}$. The induced parabolic structure on $J|_{Y_w}$ is the restriction of the one on J . Indeed, by (10.5) these parabolic structures are essentially given at the local rings at the generic points ξ_α of the divisors H_α and hence these parabolic structures on J agree on the intersections $Y_{uv} := Y_u \cap Y_v$.

Let $Z_{uv} := p_u^{-1}(Y_{uv})$. Let \tilde{Z}_{uv} be the normalization of a component of $Z_{uv} \times_{Y_{uv}} Z_{vu}$. Then \tilde{Z}_{uv} serves as Kawamata cover (12.1) of Y_{uv} (see [V, Corollary 2.6, page 56]). We

consider the morphisms $\tilde{Z}_{uv} \rightarrow Z_u$ (resp. $\tilde{Z}_{uv} \rightarrow Z_v$) and let $\mathcal{H}_{u,\tilde{Z}}$ (resp. $\mathcal{H}_{v,\tilde{Z}}$) denote the pull-backs of \mathcal{H}_u (resp. \mathcal{H}_v) to \tilde{Z}_{uv} .

Let Γ denote the Galois group for $\tilde{Z}_{uv} \rightarrow Y_{uv}$. Then by Lemma 5.1, the invariant direct images of the equivariant Lie algebra bundles $\text{Lie}(\mathcal{H}_{u,\tilde{Z}})$ and $\text{Lie}(\mathcal{H}_{v,\tilde{Z}})$ coincide with the Lie algebra structure on J restricted to the Y_{uv} and also as isomorphic parabolic bundles. Therefore we have a natural isomorphism of equivariant Lie-algebra bundles

$$\text{Lie}(\mathcal{H}_{u,\tilde{Z}}) \simeq \text{Lie}(\mathcal{H}_{v,\tilde{Z}}). \quad (10.6.1)$$

As in the proof of Theorem 10.1, this gives a canonical identification of the equivariant group schemes $\mathcal{H}_{u,\tilde{Z}}$ and $\mathcal{H}_{v,\tilde{Z}}$ on \tilde{Z}_{uv} . Since the invariant direct image of both the group schemes $\mathcal{H}_{u,\tilde{Z}}$ and $\mathcal{H}_{v,\tilde{Z}}$ are the restrictions $\mathfrak{G}_{u,Y_{uv}}$ and $\mathfrak{G}_{v,Y_{uv}}$, it follows that on $Y_{uv} = Y_u \cap Y_v$ we get a canonical identification of group schemes:

$$\mathfrak{G}_{u,Y_{uv}} \simeq \mathfrak{G}_{v,Y_{uv}}. \quad (10.6.2)$$

These identifications are canonically induced from the gluing data of the Lie algebra bundle J for the cover Y_w 's. Therefore the cocycle conditions are clearly satisfied and the identifications (10.6.2) glue to give the group scheme $\mathfrak{G}_{\mathbf{Y}^{\text{aff}}}$ on \mathbf{Y}^{aff} . The verification of the classifying property follows exactly as in the proof of Theorem 10.1.

Q. E. D

10.7. The $(\ell + 1)$ BT group scheme on \mathbf{X}^{aff} . Let \mathbf{X}^{aff} be as in §10.3. The situation in this subsection is somewhat distinct from the previously discussed cases. Unlike the scheme \mathbf{X} , the space \mathbf{X}^{aff} is an *ind-scheme* and so we give the details of the construction of the group scheme.

We begin with a generality. Let \mathbb{X} be an ind-scheme. By an open-subscheme $i: \mathbb{U} \hookrightarrow \mathbb{X}$ we mean an ind-scheme such that for any $f: \text{Spec}(A) \rightarrow \mathbb{X}$, the natural morphism $\mathbb{U} \times_{\mathbb{X}} \text{Spec}(A) \rightarrow \text{Spec}(A)$ is an open immersion. For a sheaf \mathbb{F} on \mathbb{U} by $i_*(\mathbb{F})$ we mean the sheaf associated to the pre-sheaf on the “big site” of \mathbb{X} , whose sections on $f: \text{Spec}(A) \rightarrow \mathbb{X}$ are given by $\mathbb{F}(\mathbb{U} \times_{\mathbb{X}} \text{Spec}(A))$.

The ind-scheme \mathbf{X}^{aff} has a certain open subset \mathbf{X}_0 whose precise definition is somewhat technical [So, Page 705]. Let us mention the properties relevant for us.

Recall that $\mathbf{X}^{\text{aff}} = (G_{\text{ad}}^{\text{aff}} \times G_{\text{ad}}^{\text{aff}}) \mathbf{X}_0$ and in fact $\mathbf{X}^{\text{aff}} = (G_{\text{ad}}^{\text{aff}} \times G_{\text{ad}}^{\text{aff}}) \mathbf{Y}^{\text{aff}}$. Further, the torus-embedding \mathbf{Y}^{aff} is covered by $\mathbf{Y}_w^{\text{aff}} \simeq \mathbb{A}^{\ell+1}$ which are W^{aff} -translates of $\mathbf{Y}_0^{\text{aff}}$, where $\mathbf{Y}_0^{\text{aff}} = \mathbf{Y}^{\text{aff}} \cap \mathbf{X}_0$. We remark that, analogous to the case of $\mathbf{Y}_0 \subset \mathbf{Y} \subset \mathbf{X}$, just as the toric variety \mathbf{Y}_0 was associated to the negative Weyl chamber, the toric variety $\mathbf{Y}_0^{\text{aff}}$ is associated to the negative Weyl alcove.

So let us denote by 0 the neutral element of W^{aff} also. Let $U^{\pm} \subset B^{\pm}$ be the unipotent subgroups. Let $\mathcal{U}^{\pm} := ev^{-1}(U^{\pm})$ where $ev: G(\mathcal{O}) \rightarrow G(k)$ is the evaluation map. Further by [So, Prop 5.3]

$$\mathbf{X}_0 = \mathcal{U} \times \mathbf{Y}_0^{\text{aff}} \times \mathcal{U}^-. \quad (10.7.1)$$

Let $\mathbf{X}_w := \mathcal{U} \times \mathbf{Y}_w^{\text{aff}} \times \mathcal{U}^-$. These cover $\mathcal{U} \times \mathbf{Y}^{\text{aff}} \times \mathcal{U}^-$. For $g \in G_{\text{ad}}^{\text{aff}} \times G_{\text{ad}}^{\text{aff}}$, let

$$\mathbf{X}_g := g\mathbf{X}_0 \quad \mathbf{X}_{g,w} := g\mathbf{X}_w \quad \mathbf{Y}_{g,w}^{\text{aff}} := g\mathbf{Y}_w^{\text{aff}}. \quad (10.7.2)$$

Note that we have the projection $\mathbf{X}_{g,w} \rightarrow \mathbf{Y}_{g,w}^{\text{aff}}$ which is a $\mathcal{U} \times \mathcal{U}^-$ -bundle.

By [So, Theorem 5.1] the ind-scheme \mathbf{X}^{aff} has divisors D_α for $\alpha \in \mathbb{S}$ such that the complement of their union is $\mathbf{X}^{\text{aff}} \setminus G_{\text{ad}}^{\text{aff}}$. The next proposition shows the existence of a finite dimensional Lie algebra bundle on \mathbf{X}^{aff} which is analogous to the bundle \mathcal{R} over \mathbf{X} .

PROPOSITION 10.7. *There is a finite dimensional Lie algebra bundle \mathbf{R} on \mathbf{X}^{aff} which extends the trivial Lie algebra bundle $G_{\text{ad}}^{\text{aff}} \times \mathfrak{g}$ on the open dense subset $G_{\text{ad}}^{\text{aff}} \subset \mathbf{X}^{\text{aff}}$ and whose restriction to \mathbf{Y}^{aff} is J .*

Proof. Since the projection $\mathbf{X}_{g,w} \rightarrow \mathbf{Y}_{g,w}^{\text{aff}}$ is a $\mathcal{U} \times \mathcal{U}^-$ -bundle, the transition functions of J and its restrictions to tubular neighbourhoods of its divisors may be used to construct a locally free sheaf J' on an open subset \mathbf{X}' containing the union of $G_{\text{ad}}^{\text{aff}}$ and the height one prime ideals of \mathbf{X}^{aff} using the transition functions of J . Let \mathbf{R} denote its push-forward to \mathbf{X}^{aff} . To check that the push-forward is locally free, without loss of generality we may consider its restriction to \mathbf{X}_0 . But on \mathbf{X}_0 , the pushforward of J' restricts to the pull-back of a Lie-algebra bundle J on \mathbf{Y}^{aff} constructed in (10.3), which completes the argument. The rest of the properties follows immediately. Q.E.D

THEOREM 10.8. *Let $\Theta = (\theta_\alpha \mid \alpha \in \mathbb{S})$ where $\theta_\alpha \in \mathcal{A}$ are arbitrary points in the apartment of T satisfying conditions of §2.2. There exists an affine and smooth “wonderful” Bruhat-Tits group scheme $\mathfrak{G}_{\mathbf{X}^{\text{aff}}}^\Theta$ on \mathbf{X}^{aff} together with a canonical isomorphism $\text{Lie}(\mathfrak{G}_{\mathbf{X}^{\text{aff}}}^\Theta) \simeq \mathbf{R}$. It further satisfies the following classifying property:*

For any point $h \in \mathbf{X}^{\text{aff}} \setminus G_{\text{ad}}^{\text{aff}}$, let $\mathbb{1} \subset \mathbb{S}$ be defined by the condition $h \in \cap_{\alpha \in \mathbb{1}} D_\alpha$. Let $C_1 \subset \mathbf{X}$ be a smooth curve with generic point in $G_{\text{ad}}^{\text{aff}}$ with closed point h . Let $U_h \subset C_1$ be a formal neighbourhood of h . Then, the restriction $\mathfrak{G}_{\mathbf{X}^{\text{aff}}}^\Theta|_{U_h}$ is isomorphic to the Bruhat-Tits group scheme associated to the concave function $m_{\Theta_1} : \Phi \rightarrow \mathbb{Z}$ given by $r \mapsto \sum_{\alpha \in \mathbb{1}} m_r(\theta_\alpha)$.

Proof. We begin by observing that $\mathbf{X}_{g,w}$ and $\mathbf{R} \rightarrow \mathbf{X}^{\text{aff}}$ play the role of $\mathbf{Y}_{g,w}^{\text{aff}}$ and $J \rightarrow \mathbf{Y}^{\text{aff}}$ in the proof of Theorem 10.6. Therefore the group scheme $\mathfrak{G}_{g,w}$ glue together and we obtain the global group scheme $\mathfrak{G}_{\mathbf{X}^{\text{aff}}}^\Theta$. The verification of the classifying property follows exactly as in the proof of Theorem 10.1. Q.E.D

11. 2BT GROUP SCHEMES AND DEGENERATIONS OF TORSORS

In this section we assume $\text{char}(k) = 0$. Our aim is to revisit certain smooth, affine group schemes on the minimal resolution of normal surface singularities. We work with the complete local rings and in this discussion stick to the A_n -type singularity alone and satisfy ourselves with a few remarks on the other types in a brief remark. The senior author used these constructions in [Bal] where 2BT-group schemes occur first. The purpose was to construct degenerations of the moduli spaces of principal G -bundles on smooth projective curves, when the curve is made to degenerate to an irreducible nodal curve.

The picture and notations we use is as in [Bal]. The novel feature is that the constructions were made using the geometric McKay correspondence of González-Sprinberg and Verdier ([GV]), the role of which is intriguing.

Let $N_d = \text{Spec} \frac{k[[t]][x,y]}{(x,y-t^d)}$. We recall that N_d is a *normal* surface with an isolated singularity of type A_d . By the generality of A_d -type singularities, one can realize N_d as a quotient $\sigma : D \rightarrow N_d$ of $D := \text{Spec} \frac{k[[t]][u,v]}{(u,v-t)}$ by the cyclic group $\mu_d = \langle \gamma \rangle$, where $x = u^d$, $y = v^d$, ζ is a primitive d^{th} -root of unity and μ_d acts on D as follows:

$$\gamma.(u, v) = (\zeta.u, \zeta^{d-1}.v). \quad (11.0.1)$$

We consider the following basic diagram for all $d > 0$ (see [GV]):

$$\begin{array}{ccc} D^{(d)} & \xrightarrow{f} & \mathbb{N}^{(d)} \\ q \downarrow & & \downarrow p_d \\ 0 \in D & \xrightarrow{\sigma} & N_d \ni c \end{array} \quad (11.0.2)$$

where $p_d : \mathbb{N}^{(d)} \rightarrow N_d$ is the *minimal resolution of singularities* of N_d obtained by successively blowing up the singularity, with the exceptional divisor $E^{(d)} = p_d^{-1}(c)$ having $d-1$ rational components, and

$$D^{(d)} := (D \times_{N_d} \mathbb{N}^{(d)})_{\text{red}}. \quad (11.0.3)$$

The *closed fibre* $F^{(d)}$ of the canonical morphism $\mathbb{N}^{(d)} \rightarrow \text{Spec}(k[[t]])$ looks like:

$$F^{(d)} = E^{(d)} \cup E(1) \cup E(2), \quad (11.0.4)$$

where $E(1)$ and $E(2)$ are the inverse images of the branches corresponding to $x=0$ and $y=0$ in N_d .

Thus, $F^{(d)} \subset \mathbb{N}^{(d)}$ is a normal crossing divisor with $d+1$ components. By [GV, Proposition 2.4] the morphism

$$f : D^{(d)} \rightarrow \mathbb{N}^{(d)} \quad (11.0.5)$$

is *finite and flat, the minimal platicateur in the sense of Grothendieck* [GV, Cor.7, page 448]. Since $\mathbb{N}^{(d)}$ is smooth, this implies that f is ramified at the generic point of each of the $d-1$ rational components of the exceptional divisor $E^{(d)} = p_d^{-1}(c) \subset F^{(d)}$.

Let

$$\mathcal{T}_D \simeq D \times^{\rho} G \quad (11.0.6)$$

be the trivial (μ_d, G) -torsor on D (11.0.2) with a μ_d -structure given by a homomorphism $\rho : \mu_d \rightarrow G$. This gives a homomorphism $\rho : \mu_d \rightarrow T$ into the maximal torus T of G . We fix once for all an isomorphism $T \simeq \mathbb{G}_m^{\ell}$. Thus ρ determines an ordered pair of integers modulo d called the *type* $\tau = (a_1, a_2, \dots, a_{\ell})$. More precisely, we have a μ_d -action on $D \times G$, given by

$$\gamma.(u, v, g) = (\zeta.u, \zeta^{-1}.v, \rho(\gamma).g). \quad (11.0.7)$$

Let z_1 (resp. z_2) be the two points in $D^{(d)}$ above the origin $0 \in D$ where the normalization of the curve $u.v=0$ meets the fibre $q^{-1}(0)$. We observe that since the action of μ_d is *balanced* at these two marked points i.e., the action of a generator ζ on the tangent spaces to each branch are inverses to each other. For the corresponding dual action in the neighbourhood D'_0 (the component with local coordinate v), the action is by ζ^{-1} . If we begin with a representation $\rho : \mu_d \rightarrow G$ of local type τ at a point in a branch, then

the corresponding local type for the dual action at the point in the second branch is denoted by $\bar{\tau}$.

Consider the adjoint group scheme $\mathcal{T}_D(G) := \mathcal{T}_D \times^{G, \text{Ad}} G$ on D , where G acts on itself by inner conjugation. We define the equivariant group scheme:

$$E(G, \tau) := q^*(\mathcal{T}_D(G)) \quad (11.0.8)$$

on $D^{(d)}$ of local type τ in the sense that it comes with a μ_d -action via a representation $\rho : \mu_d \rightarrow G$. Since the morphism $f : D^{(d)} \rightarrow \mathbb{N}^{(d)}$ is also *finite and flat* we can take the Weil restriction of scalars:

$$f_*(E(G, \tau)_{D^{(d)}}) := \text{Res}_{D^{(d)}/\mathbb{N}^{(d)}}(E(G, \tau)_{D^{(d)}}), \quad (11.0.9)$$

and since $E(G, \tau)_{D^{(d)}} \rightarrow D^{(d)}$ is a smooth (affine) group scheme, the basic properties of Weil restriction of scalars ([Ed, Lemma 2.2]) show that $f_*(E(G, \tau))$ is a smooth group scheme on $\mathbb{N}^{(d)}$ together with a μ_d -action. By taking invariants under the action of μ_d and noting that we are over characteristic zero, by [Ed, Prop 3.4], we obtain the smooth (affine) group scheme obtained by taking *invariant direct images*:

$$\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G = (f_*^{\mu_d})(E(G, \tau)_{D^{(d)}}), \quad (11.0.10)$$

on $\mathbb{N}^{(d)}$ (see [BS, Definition 4.1.3]). The 2BT-group scheme of type τ with generic fibre G of singularity type A_d associated to θ_τ is defined to be the affine group scheme $\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G$ (11.0.10) on the regular surface $\mathbb{N}^{(d)}$. This process defines a distinguished collection of 2BT-group scheme $\left\{ \mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G \right\}_\tau$ indexed by the type τ .

11.1. The McKay correspondence revisited. We recall the geometric interpretation of the McKay correspondence given by González-Sprinberg and Verdier ([GV]). Let $\text{Irr}^o(\mu_d) \subset \text{Irr}(\mu_d)$ be the nontrivial irreducible representations of μ_d and let $\text{Irr}(E^{(d)})$ denote the set of irreducible rational components of the exceptional divisor of the minimal resolution $p_d : \mathbb{N}^{(d)} \rightarrow N_d$ (11.0.2). Let ψ be a non-trivial character of $\mu_d = \langle \gamma \rangle$. Then ψ corresponds to $\zeta \mapsto \zeta^s$, where ζ is the primitive d^{th} -root of 1 chosen above, and $1 \leq s \leq d-1$. Let L_ψ be the equivariant line bundle on D where μ_d acts on $D \times k$ as $\gamma \cdot (u, v, a) = (\zeta \cdot u, \zeta^{d-1} \cdot v, \zeta^s \cdot a)$, $a \in k$. A μ_d -invariant section \mathcal{h} of this line bundle is given by the relation $\{\gamma \cdot \mathcal{h}\}(u, v) = \mathcal{h}(\gamma \cdot (u, v)) = \zeta^s \mathcal{h}(u, v)$, and hence the μ_d -invariant sections are generated by u^s and v^{d-s} . Let $\mathcal{L}_\psi := f_*^{\mu_d}(q^*(L_\psi))$ be the induced line bundle on $\mathbb{N}^{(d)}$. This is a line bundle since f is finite and flat.

McKay correspondence following González-Sprinberg and Verdier. There is a bijection $\text{Irr}^o(\mu_d) \rightarrow \text{Irr}(E^{(d)})$, $\psi \mapsto E_\psi$, such that for any $E_j \in \text{Irr}(E^{(d)})$, we have

$$c_1(\mathcal{L}_\psi) \cdot E_j = \begin{cases} 0 & \text{if } E_j \neq E_\psi \\ 1 & \text{if } E_j = E_\psi. \end{cases} \quad (11.1.1)$$

The above statement implies that the first Chern class $c_1(\mathcal{L}_\psi)$ can be represented by an *effective* divisor $\delta_\psi \subset \mathbb{N}^{(d)}$ which meets $F^{(d)}$ transversally at a unique point which lies in E_ψ .

We interpret this on the side of the surface $D^{(d)}$. Consider the reduced fibre $\tilde{E} := q^{-1}(0)_{\text{red}}$. The group μ_d -fixes the divisor $q^{-1}(0)$ and hence its reduced subscheme \tilde{E} .

There is a smooth curve $\delta'_\psi \in D^{(d)}$ which meets the divisor $q^{-1}(0) \subset D^{(d)}$ at a unique component \tilde{E}_ψ of \tilde{E} . Locally at this point of \tilde{E}_ψ the transversal curve δ'_ψ is given by the invariant section \mathfrak{h} of $q^*(\mathcal{L}_\psi)$. The action of μ_d on \mathfrak{h} shows that the group μ_d acts on the local uniformizer z_ψ of the component \tilde{E}_ψ by $\gamma \cdot z_\psi \mapsto \zeta^s z_\psi$.

Let us explain how to a representation $\rho: \mu_d \rightarrow T$ of type τ , we associate a rational point θ_τ of \mathcal{A}_T in the fundamental domain of $Y(T)$ as in [BS], and not just in the alcove \mathbf{a}_0 . Let $\rho: \mu_d \rightarrow T$ map the generator γ of μ_d to $(\zeta_d^{a_1}, \dots, \zeta_d^{a_\ell})$ where ζ_d is a primitive d th root of unity and the a_i are integers uniquely determined modulo d . We say ρ is of type $\tau = (a_1, a_2, \dots, a_\ell)$. This determines a rational point in $\frac{1}{d}Y(T) \bmod Y(T)$ and therefore a rational point θ_τ in the fundamental domain of $Y(T)$ in \mathcal{A}_T . Let $F^{(d)} = E^{(d)} \cup E(1) \cup E(2)$, be as in (11.0.4) and let the two nodal end points be $\mathbf{z}_i, i = 1, 2$ and the nodes where the rational components meet at $\mathbf{y}_s, s = 1, \dots, d-2$ for $d \geq 3$.

Consider the canonical morphism $\mathbb{N}^{(d)} \rightarrow \text{Spec}(k[[t]])$. Let $\mathbb{N}_t^{(d)} \subset \mathbb{N}^{(d)}$ be the open subset defined by the non-vanishing of t . For $s = 1, \dots, d-2$, let $C_s \subset \mathbb{N}^{(d)}$ be a smooth curve with generic point in $\mathbb{N}_t^{(d)}$ and containing \mathbf{y}_s as a closed point. Let U_s be the formal neighbourhood of \mathbf{y}_s in C_s . Similarly, let $C_{\mathbf{z}_i}$ for $i = 1, 2$ denote smooth curves with generic point in $\mathbb{N}_t^{(d)}$ and containing \mathbf{z}_i as a closed point. Let $U_{\mathbf{z}_i}$ be the formal neighbourhood of \mathbf{z}_i in $C_{\mathbf{z}_i}$.

THEOREM 11.1. *The group scheme $\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G$ is affine. It has the following description at the generic points of the rational curves, at $U_{\mathbf{z}_i}$ and at U_s :*

- (1) *Let ψ denote the character which takes $\zeta \rightarrow \zeta^s$. We read the tuple $(s.a_1, s.a_2, \dots, s.a_\ell)$ modulo d . Let $\tau_s := (s.a_1, s.a_2, \dots, s.a_\ell)$. At the generic point of the rational component E_ψ , the local type of the restriction of $\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G$ is given by the point θ_{τ_s} in the fundamental domain of $Y(T)$.*
- (2) *The restriction of $\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G$ to $U_{\mathbf{z}_i}$ for $i = 1, 2$ is isomorphic to the parahoric group scheme $\mathfrak{G}_{\theta_\tau}$ and $\mathfrak{G}_{\theta_{\bar{\tau}}}$ corresponding to τ and $\bar{\tau}$ respectively.*
- (3) *The restriction of $\mathcal{H}_{\tau, \mathbb{N}^{(d)}}^G$ to U_s is isomorphic to the Bruhat-Tits group scheme \mathfrak{G}_{f_s} where $f_s: \Phi \rightarrow \mathbb{R}$ is the concave function given by*

$$f_s(r) := m_r(\theta_{\tau_s}) + m_r(\theta_{\tau_{s+1}}). \quad (11.1.2)$$

Proof. This is an immediate consequence of (5.7) and (5.4). Q.E.D

Remark 11.2. Let us assume that the residue field k contains primitive 8-th (resp. 3rd) root of unity. Let $\{\alpha, \beta\}$ denote the short and long roots of B_2 (resp. G_2). Let us fix an isomorphism of T with \mathbb{G}_m^2 . Since the coroot lattice equals $Y(T)$, let us take the coroot homomorphisms $\alpha^\vee, \beta^\vee: \mathbb{G}_m \rightarrow T$ as the first and second coordinate maps. Let us take ρ of type τ_1 such that θ_{τ_1} equals $\frac{\theta_\alpha}{2}$ for G of type B_2 (resp. θ_α for G of type G_2). Then τ_2 gives θ_α (resp. $2\theta_\alpha$). By example 3.10 (resp. 3.12), we see that for $s = 1$, the f_s is of type III. This shows that the limiting objects of [Bal] could become torsors under BT group schemes of the most general type.

12. APPENDIX ON PARABOLIC AND EQUIVARIANT BUNDLES

In this section we recall and summarise some results on parabolic bundles and equivariant bundles on Kawamata covers. These play a central role in the constructions of the Bruhat-Tits group schemes made above. Consider a pair (X, D) , where X is a smooth quasi-projective variety and $D = \sum_{j=0}^{\ell} D_j$ is a *reduced normal crossing divisor* with non-singular components D_j intersecting each other transversely. The basic examples we have in mind in the note are discrete valuation rings with its closed point, the wonderful compactification \mathbf{A} with its boundary divisors or affine toric varieties.

Let E be a locally free sheaf on X . Let $n_j, j = 0, \dots, \ell$ be positive integers attached to the components D_j . Let ξ be a generic point of D .

Let $E_\xi := E \otimes_{\mathcal{O}_X} \mathcal{O}_{X, \xi}$ and $\bar{E}_\xi := E_\xi / \mathfrak{m}_\xi E_\xi$, \mathfrak{m}_ξ the maximal ideal of $\mathcal{O}_{X, \xi}$.

DEFINITION 12.1. *A (generic) parabolic structure on E consists of the following data:*

- A flag $\bar{E}_\xi = F^1 \bar{E}_\xi \supset \dots \supset F^{r_j} \bar{E}_\xi$, at the generic point ξ of each of the components D_j of D ,
- Weights d_s / n_j , with $0 \leq d_s < n_j$ attached to $F^s \bar{E}_\xi$ such that $d_1 < \dots < d_{r_j}$

By saturating the flag datum on each of the divisors we get for each component D_j a filtration:

$$E_{D_j} = F_j^1 \supset \dots \supset F_j^{r_j} \quad (12.0.1)$$

of sub-sheaves on D_j . Define the coherent subsheaf \mathcal{F}_j^s , where $0 \leq j \leq \ell$ and $1 \leq s \leq r_j$, of E by:

$$0 \rightarrow \mathcal{F}_j^s \rightarrow E \rightarrow E_{D_j} / F_j^s \rightarrow 0 \quad (12.0.2)$$

the last map is by restriction to the divisors.

We wish to emphasise that this definition, where the quasi-parabolic structure is given only at the generic points of the divisors, is somewhat different from what is done in [Bi].

In [MS], working over curves, two essential things are shown:

- (1) given a parabolic bundle E as above, there is a ramified covering of $p : (Y, \tilde{D}) \rightarrow (X, D)$ with suitable ramification data and Galois group Γ together with a Γ -equivariant vector bundle V on Y such that the invariant direct image sheaf $p_*^\Gamma(V)$ equals E and furthermore V also recovers the parabolic structure, namely the filtrations \mathcal{F}_j^s and weights, on E .
- (2) Conversely, if we begin with an equivariant bundle V on Y then the invariant direct image $p_*^\Gamma(V)$ gives a vector bundle E on X with parabolic structures at the generic points of the height 1 prime ideals.

In this appendix we work with the higher-dimensional and generic variant we have defined above and establish analogous results under suitable conditions. This is the key fact that is used in the present paper.

For the sake of completeness, we give a self-contained *ad hoc* argument for this construction which is more in the spirit of the present note. The data given in (12.1) is as in [MS] which deals with points on curves. *Note that in our setting we have rational weights.* Under these conditions, we have a natural functor

$$p_*^\Gamma : \{\Gamma\text{-equivariant bundles on } Y\} \rightarrow \{\text{parabolic bundles on } X\}. \quad (12.0.3)$$

Under suitable conditions relevant to this paper this functor is in fact a surjection. See [Bi] for other conditions where we can obtain a surjection.

As the notation suggests, this is achieved by taking invariant direct images. Since $p : Y \rightarrow X$ is finite and flat, if V is locally free on Y then so is $p_*^\Gamma(V)$. An equivariant bundle V on Y is defined in an formal neighbourhood of a generic point ζ of a component by a representation of the isotropy group Γ_ζ and locally (in the formal sense) the action of Γ_ζ is the product action. This is called the local type in [Ses] or [MS]. By appealing to the 1-dimensional case, we obtain a canonical (generic) parabolic structure on E , i.e., parabolic structures at the generic points ξ of the components D_j in X .

Conversely, given a (generic) parabolic bundle E on (X, D) , to get the Γ -equivariant bundle V on Y , such that $p_*^\Gamma(V) = E$, we heuristically proceed as follows. If we did know the existence of such a V then we can consider the inclusion $p^*(E) \subset V$. Taking its dual (and since $V/p^*(E)$ is torsion, taking duals is an inclusion), we get:

$$V^* \hookrightarrow (p^*(E))^* = p^*(E^*). \quad (12.0.4)$$

By the 1-dimensional case (obtained by restricting to the height 1-primes at the generic points) we see that the quotient $T_\zeta := p^*(E^*)_\zeta / V_\zeta^*$ is a torsion $\mathcal{O}_{Y, \zeta}$ -module and T_ζ is completely determined by the parabolic structure on E .

The (generic) parabolic structure on E therefore determines canonical quotients:

$$p^*(E^*)_\zeta \rightarrow T_\zeta \quad (12.0.5)$$

for each generic point ζ of components in Y above the components D_j in X .

The discussion above suggest how one would construct such a V ; we begin with these quotients (12.0.5). Then we observe that there is a maximal coherent subsheaf $V' \subset p^*(E^*)$ such that

$$p^*(E^*)_\zeta / V_\zeta' = T_\zeta. \quad (12.0.6)$$

Away from $p^{-1}(D)$ in Y , the inclusion $V' \hookrightarrow p^*(E^*)$ is an isomorphism. Dualizing again, we get an inclusion $p^*(E) \hookrightarrow (V')^*$ which is an isomorphism away from $p^{-1}(D)$. Set $W = (V')^*$. Since W is the dual of a coherent sheaf, it is *reflexive*.

In situations as in (5.4), we find examples of vector bundles with Lie algebra structures which satisfy the surjectivity of (12.0.3).

12.1. Kawamata Coverings. Suppose we are given positive integers n_0, \dots, n_ℓ . Let X be a smooth quasi-projective variety over a perfect field k whose characteristic $\text{char}(k)$ coprime to n_0, \dots, n_ℓ . Let D be a simple or reduced normal crossing divisor with decomposition $D = \sum_{i=0}^\ell D_i$ into its smooth components intersecting transversally.

By a ‘‘Kawamata Covering’’ of X we mean the existence of of a connected smooth quasi-projective variety Z over k and a Galois covering morphism

$$\kappa : Z \rightarrow X \quad (12.1.1)$$

such that the reduced divisor $\kappa^* D := (\kappa^* D)_{\text{red}}$ is a normal crossing divisor on Z and furthermore, $\kappa^* D_i = n_i \cdot (\kappa^* D_i)_{\text{red}}$. Let Γ denote the Galois group for κ .

The ‘‘Covering Lemma’’ of Y. Kawamata (see [V, Lemma 2.5, page 56]) says that there is one such under the assumption of k being algebraically closed.

We however note that if the base is reasonably simple, such as an affine space \mathbb{A}_k^n , then such a covering will exist with just the assumption of k being perfect.

The isotropy group of any point $z \in Z$, for the action of Γ on Z , will be denoted by Γ_z . It is easy to see that the stabilizer at generic points of the irreducible components of $(\kappa^* D_i)_{\text{red}}$ are cyclic of order n_i .

Remark 12.2. (In positive characteristics) The Kawamata covering lemma is seen to hold under *tameness assumptions*, i.e., if the characteristic p is coprime to the n_i 's, then it holds [V, Lemma 2.5, page 56].

12.2. The group scheme situation. Let (X, D) be as above. Let ξ be the generic point of a component D_j of D and let $A := \mathcal{O}_{X, \xi}$ and K be the quotient field of A . We always assume that these group scheme are generically split. Let us begin by stating some results from [BS] in the simplest situation. Then we will state the more general case.

Let $\mathfrak{G}_{\mathfrak{a}_0}$ be a Bruhat-Tits group scheme on $\text{Spec}(A)$ associated to a vertex $\theta_{\mathfrak{a}}$ of the Weyl alcove \mathfrak{a}_0 (see §2.1). Let $B = \mathcal{O}_{Y, \zeta}$ where ζ is the generic point of a component of Y above D_j and let L be the quotient field of B . We assume that the local ramification data for the Kawamata covering has numbers d_{α} (2.1.2). Let Γ_{ζ} be the stabilizer of Γ at $\zeta \in Y$. The results of [BS, Proposition 5.1.2 and Remark 2.3.3] show that there exists an equivariant group scheme \mathcal{H}_B on $\text{Spec}(B)$ with fiber isomorphic to the simply-connected group G and such that $\text{Res}_{B/A}(\mathcal{H}_B)^{\Gamma_{\zeta}} \simeq \mathfrak{G}_{\mathfrak{a}_0}$.

Although in [BS] the above is stated for points in the alcove \mathfrak{a}_0 but the proofs reveal that actually they hold more generally. Under assumptions on the residue field (see §1.1 and §2.2), the proofs of these results generalise in a straightforward way for an arbitrary complete DVR (see §9.1). We may further assume (see Remark (8.1)) that θ is a rational point of \mathcal{A}_T in the fundamental domain of $Y(T)$ and not just a point in the alcove \mathfrak{a}_0 (see §2.1).

Suppose that we have a group scheme $\mathfrak{G}_{X'}$ on an open $X' \subset X$ which includes all the height 1 primes coming from the divisors D_j 's with the following properties:

- Away from the divisor $D \subset X$, \mathfrak{G} is the constant group scheme with fiber G .
- The restrictions $\mathfrak{G}|_{\text{Spec}(A)}$ at the generic points ξ are isomorphic to the Bruhat-Tits group scheme \mathfrak{G}_{θ} for varying ξ and θ varying in the fundamental domain of $Y(T)$ in \mathcal{A}_T .

In other words, $\mathfrak{G}_{X'}$ is obtained by a gluing of the constant group schemes with \mathfrak{G}_{θ} 's along $\text{Spec}(K)$ by an automorphism of constant group scheme G_K .

Now consider the inverse image of the constant group scheme $p^*(G_{X-D}) \simeq G \times p^*(X-D)$. Then using the gluing on X' , we can glue the constant group scheme $p^*(G_{X-D})$ with the local group schemes \mathcal{H}_B for each generic point ζ to obtain a group scheme $\mathcal{H}_{Y'}$ on $Y' = p^{-1}(X')$ such that:

$$\text{Res}_{Y'/X'}(\mathcal{H}_{Y'})^{\Gamma} \simeq \mathfrak{G}_{X'}. \quad (12.2.1)$$

REFERENCES

- [BS] V.Balaji and C.S. Seshadri, Moduli of parahoric \mathfrak{G} -torsors on a compact Riemann surface, *J. Algebraic Geometry*, 24, (2015), 1-49.
- [Bal] V.Balaji, Torsors on semistable curves and degenerations, *Proceedings of the Indian Academy of Sciences, Mathematical Sciences*, Volume 132, Article number: 27 (2022).
- [BDP] V. Balaji, P. Deligne and A.J. Parameswaran, On complete reducibility in characteristic p , *Épjournal de Géométrie Algébrique*, Volume 1 (September 1, 2017).
- [Bi] I. Biswas, Chern classes for parabolic bundles. *J. Math. Kyoto Univ.* 37 (1997), no. 4, 597–613. doi:10.1215/kjm/1250518206.
- [BLR] S. Bosch, W.Lutkebohmert and M.Raynaud, *Neron Models*, Ergebnisse 21, Springer Verlag, (1990).

- [Br1] M. Brion, The behaviour at infinity of the Bruhat decomposition, *Comm. Math. Helv.* 73 (1998)137-174.
- [Br2] M. Brion, Log homogeneous varieties, Actas del XVI Coloquio Latinoamericano de Álgebra, 1–39, Biblioteca de la Revista Matemática Iberoamericana, Madrid, 2007.
- [BT1] F.Bruhat and J.Tits : Groupes réductifs sur un corps local, I, *Publications Mathématiques de l’IHÉS*, **41** (1972) pp 5-251.
- [BT2] F.Bruhat and J.Tits, Groupes réductifs sur un corps local II: Schémas en groupes. Existence d’une donnée radicielle valuée, *Publications Mathématiques de l’IHÉS* 60 (1984) 5-184.
- [BT3] F. Bruhat and J. Tits, Schémas en groupes et immeubles des groupes classiques sur un corps local. *Bulletin Soc. Math. France* 112 (1984), 259–301.
- [DP] C. De Concini and C. Procesi, Complete symmetric varieties, in Invariant theory (Montecatini, 1982), pp. 1–44, Lecture Notes in Math. 996, Springer, 1983.
- [DS] C. De Concini and T. Springer, Compactification of Symmetric Varieties, *Transform. Groups* 4 (1999), no. 2-3, pp. 273-300.
- [CGP] B. Conrad, O.Gabber and Gopal Prasad, Pseudo-reductive groups, *Cambridge University Press*, New Mathematical Monographs, No 17.(2010), 2nd Edition.
- [Cou] Francois Courtes, Strongly concave functions on root systems, *Mathematica Slovaca* 61, (2011), No. 4, 539-560.
- [SGAD] M. Demazure and A. Grothendieck, editors, Schemas en groupes. I, II, III, Seminaire de Geometrie Algebrique du Bois Marie 1972/64, Lecture Notes in Math., vols. 151 153, Springer Verlag, Berlin and New York, 1970.
- [DG] M. Demazure and P. Gabriel, *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*. Avec un appendice Corps de classes local par Michiel Hazewinkel. Masson & Cie, Paris; North Holland Publishing Co., Amsterdam, 1970. MR 0302656
- [Ed] B. Edixhoven, Néron models and tame ramification, *Compositio Mathematica*, **81**, (1992), 291-306.
- [GY1] W. T. Gan and J.-K. Yu, Schémas en groupes et immeubles des groupes exceptionnels sur un corps local. Première partie: le groupe G_2 , *Bull. Soc. math. France*, 131 (3), 2003, p. 307-358.
- [GY2] W. T. Gan and J.-K. Yu, Schémas en groupes et immeubles des groupes exceptionnels sur un corps local. Deuxième partie: les groupes F_4 et E_6 , *Bull. Soc. math. France*, 133 (2), 2005, p. 159-197.
- [GV] G. González-Sprinberg and J.-L.Verdier, Construction géométrique de la correspondance de McKay, *Annales scientifiques de l’École Normale Supérieure* 16.3 (1983), 409-449.
- [EGA] A. Grothendieck, Éléments de Géométrie Algébrique, *Publications Mathématiques de l’IHÉS* 4, (1960), 5-228.
- [EGA IV] A. Grothendieck, Éléments de Géométrie Algébrique, *Publications Mathématiques de l’IHÉS* 24, (1965), 5-231.
- [H] R. Hartshorne, Stable Reflexive Sheaves, *Mathematische Annalen* 254 (1980), 121-176.
- [H1] R. Hartshorne, Local Cohomology. Springer, Lecture Notes 41, 1967.
- [KKMS] G. Kempf, F. Knudsen, D. Mumford, and B. Saint-Donat, Toroidal Embeddings I, Springer Lecture Notes 339, 1973.
- [L] E.Landvogt, A Compactification of the Bruhat-Tits Building. Lecture Notes in Mathematics, vol. 1619. Springer, Berlin (1996).
- [Lou] Lourenço J. Grassmanniennes affines tordues sur les entiers. *Forum of Mathematics, Sigma*. 2023;11:e12. doi:10.1017/fms.2023.4
- [MRR] Arnaud Mayeux, Timo Richarz, Matthieu Romagny. Néron blowups and low-degree cohomological applications. 2020. fihal-02490010v2f
- [MS] V. Mehta and C.S. Seshadri, Moduli of vector bundles on curves with parabolic structures, *Math. Ann*, **248**, (1980), 205-239.
- [MP] Allen Moy and Gopal Prasad, Unrefined minimal K -types for p -adic groups, *Invent Math*, 116, (12.1994).
- [PZ] G. Pappas, X. Zhu, Local models of Shimura varieties and a conjecture of Kottwitz, *Invent. Math.* 194 (2013), 147-254.
- [Par] A. N. Parshin, Higher Bruhat-Tits buildings and vector bundles on an algebraic surface, Algebra and Number Theory (Proc. Conf. at Inst. Exp. Math. , Univ. Essen, 1992), de Gruyter, Berlin, 1994, 165-192.
- [Ray] M. Raynaud, Faisceaux amples sur les schemas en groupes et les espaces homogenes, Lecture Notes in Math., vol. 119, Springer Verlag, Berlin and New York, 1970.
- [RY] M.Reeder and J.-K.Yu Epipelagic Representations and Invariant theory. *J. Amer. Math. Soc.*, 27(2), (2014) 437–477.

- [Se] J-P. Serre, *Exemples de plongements des groupes $PSL_2(\mathbb{F}_p)$ dans des groupes de Lie simples*, *Invent. Math.*, **124** (1996), 525-562. MR 1369427
- [Ses] C.S. Seshadri, *Moduli of π -vector bundles over an algebraic curve*, *Questions On algebraic Varieties*, C.I.M.E, Varenna, (1969), 139-261.
- [So] P. Solis, *A wonderful embedding of the loop group*. *Advances in Mathematics* 313, , (2017), 689-717.
- [SS] Schneider, Peter; Stuhler, Ulrich. *Representation theory and sheaves on the Bruhat-Tits building*. *Publications Mathématiques de l'IHÉS*, Volume 85 (1997), pp. 97-191.
- [V] E. Vieweg, *Projective Moduli for Polarized Manifolds*, , *Ergebnisse*, Springer(1995).
- [Yu] J-K Yu, *Smooth models associated to concave functions in Bruhat-Tits theory*, *Autour des schémas en groupes*. Vol. III, 2015, pp. 227-258.
- [Yu1] J-K Yu, *Construction of tame supercuspidal representations*, *J. Amer. Math. Soc.* 14 (2001), no. 3, 579–622 (electronic).

CHENNAI MATHEMATICAL INSTITUTE, PLOT NUMBER H1, SIPCOT IT PARK, SIRUSERI, CHENNAI, 603103
INDIA

Email address: `balaji@cmi.ac.in`

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, MOHALI KNOWLEDGE CITY, SECTOR 81, SAS
NAGAR, MANAULI PO 140306, INDIA

Email address: `ypandey@iisermohali.ac.in`, `yashonidhipandey@yahoo.co.uk`,
`pandeyyashonidhi@gmail.com`