RDP DEL PEZZO SURFACES WITH GLOBAL VECTOR FIELDS IN ODD CHARACTERISTIC

GEBHARD MARTIN AND CLAUDIA STADLMAYR

ABSTRACT. We classify RDP del Pezzo surfaces with global vector fields over arbitrary algebraically closed fields of characteristic $p \neq 2$. In characteristic 0, every RDP del Pezzo surface X is *equivariant*, that is, $\operatorname{Aut}_X = \operatorname{Aut}_{\widetilde{X}}$, where \widetilde{X} is the minimal resolution of X, hence the classification of RDP del Pezzo surfaces with global vector fields is equivalent to the classification of weak del Pezzo surfaces with global vector fields.

In this article, we show that if $p \neq 2, 3, 5, 7$, then it is still true that every RDP del Pezzo surface is equivariant. We classify the non-equivariant RDP del Pezzo surfaces in characteristic p = 3, 5, 7, giving explicit equations for every such RDP del Pezzo surface in all possible degrees. As an application, we construct regular non-smooth RDP del Pezzo surfaces over imperfect fields of characteristic 7, thereby showing that the known bound $p \leq 7$ for the characteristics, where such a surface can exist, is sharp.

CONTENTS

1.	Introduction	1	
2.	An application: Regular inseparable twists of RDP del Pezzo surfaces		
3.	Preliminaries on (RDP) del Pezzo surfaces	4	
4.	Group scheme actions on anti-canonical models	6	
5.	On equivariant resolutions	7	
6.	Automorphism schemes of equivariant RDP del Pezzo surfaces	10	
7.	Finding (-1) -curves in the equivariant locus	11	
8.	Automorphism schemes of non-equivariant RDP del Pezzo surfaces	13	
	8.1. In characteristic 7	13	
	8.2. In characteristic 5	14	
	8.3. In characteristic 3	17	
Ap	Appendix: Classification tables		
Ref	ferences	33	

1. INTRODUCTION

We are working over an algebraically closed field k of characteristic $p \ge 0$. Let X be a del Pezzo surface with at worst rational double points and let $\pi : \tilde{X} \to X$ be its minimal resolution, so that, by definition, \tilde{X} is a weak del Pezzo surface. Since $-K_X$ is ample, Aut_X is an affine group scheme of finite type, hence its group of automorphisms $\operatorname{Aut}(X) := \operatorname{Aut}_X(k)$ is infinite if and only if the automorphism scheme Aut_X is positive-dimensional. Moreover, by Blanchard's Lemma [Bri17, Theorem 7.2.1], and since X is the anticanonical model of \tilde{X} , there is a closed immersion of group schemes $\pi_* : \operatorname{Aut}_{\tilde{X}} \hookrightarrow \operatorname{Aut}_X$. We call X equivariant, if π_* is an isomorphism. Summarizing, for all characteristics, there is the following chain of implications:

(1.1)
$$|\operatorname{Aut}(X)| = |\operatorname{Aut}(\widetilde{X})| = \infty \implies H^0(\widetilde{X}, T_{\widetilde{X}}) \neq 0 \implies H^0(X, T_X) \neq 0$$

Over the complex numbers, every RDP del Pezzo surface X is equivariant, so in particular we have $H^0(X, T_X) = H^0(\widetilde{X}, T_{\widetilde{X}})$, and by Cartier's theorem Aut^0_X is smooth, hence it is positive-dimensional

Date: March 18, 2022.

²⁰²⁰ Mathematics Subject Classification. 14E07, 14J26, 14J50, 14L15, 14J17, 14B05, 14G17.

if and only if $H^0(X,T_X) \neq 0$. In other words, in characteristic 0, all implications in (1.1) are in fact equivalences.

In our recent article [MS20], we obtained the classification of weak del Pezzo surfaces with global vector fields over algebraically closed fields of arbitrary characteristic (over the complex numbers, an independent proof was given by Cheltsov and Prokhorov in [CP21]). By (1.1), this includes the classification of all RDP del Pezzo surfaces with infinite automorphism group, but we note that if p = 2, 3, there are RDP del Pezzo surfaces with finite automorphism group whose minimal resolution has global vector fields, so the first implication in (1.1) is not an equivalence precisely if p = 2, 3. In other words, we have

(1.2)
$$|\operatorname{Aut}(X)| = |\operatorname{Aut}(\widetilde{X})| = \infty \xrightarrow{p \neq 2.3} H^0(\widetilde{X}, T_{\widetilde{X}}) \neq 0 \implies H^0(X, T_X) \neq 0$$

and a classification of \widetilde{X} with $H^0(\widetilde{X}, T_{\widetilde{X}}) \neq 0$ in all characteristics.

Now, the missing piece is a classification of RDP del Pezzo surfaces X with $H^0(X, T_X) \neq 0$. As a first step, we extend Hirokado's [Hir19] results on the liftability of vector fields to resolutions of rational double points to group scheme actions on RDP del Pezzo surfaces as follows:

Theorem 1.1 (= Theorem 6.1). Let X be an RDP del Pezzo surface and let $\pi : \widetilde{X} \to X$ be its minimal resolution. Assume that one of the following conditions holds:

- (1) $p \notin \{2, 3, 5, 7\},\$
- (2) p = 7 and X does not contain an RDP of type A_6 .
- (3) p = 5 and X does not contain an RDP of type A_4 or E_8^0 .
- (4) p = 3 and X does not contain an RDP of type $A_2, A_5, A_8, E_6^0, E_6^1, E_7^0, E_8^0$ or E_8^1 . (5) p = 2 and X does not contain an RDP of type $A_1, A_3, A_5, A_7, D_n^r, E_6^0, E_7^0, E_7^1, E_7^2, E_8^3, E_8^1, E_8^2$ or E_8^3 .

Then, $\operatorname{Aut}_X = \operatorname{Aut}_{\widetilde{X}}$, and thus, in particular, $H^0(\widetilde{X}, T_{\widetilde{X}}) = H^0(X, T_X)$. Therefore, $H^0(X, T_X) \neq 0$ if and only if X is the anti-canonical model of one of the surfaces in the classification tables of [MS20].

Thus, in order to classify RDP del Pezzo surfaces X with global vector fields, we may restrict our attention to RDP del Pezzo surfaces containing a configuration Γ of RDPs excluded in Theorem 6.1. In Theorem 7.1, we give a criterion for X to be the blow-up of an RDP del Pezzo surface of higher degree with the same configuration Γ . In the language of the Minimal Model Program, this means that we give a sufficient criterion for the existence of a $K_{\tilde{X}}$ -negative extremal ray on X which lies in the orthogonal complement of the exceptional locus over Γ . Using Blanchard's Lemma, this allows us to set up an inductive argument for the classification of non-equivariant RDP del Pezzo surfaces X with RDP configuration Γ . This strategy will be carried out in Sections 8.1, 8.2, and 8.3, for characteristic p = 7, p = 5, and p = 3, respectively. The following theorem is obtained by combining Theorems 8.3, 8.6, and 8.8.

Theorem 1.2. Let X be an RDP del Pezzo surface and let $\pi : \widetilde{X} \to X$ be its minimal resolution. Assume that $H^0(X, T_X) \neq 0$. Then, the following hold:

- (1) If p = 7 and X contains an RDP of type A_6 , then X is one of the 2 surfaces in Table 4.
- (2) If p = 5 and X contains an RDP of type A_4 or E_8^0 , then X is one of the 9 surfaces in Table 5.
- (3) If p = 3 and X contains an RDP of type $A_2, A_5, A_8, E_6^0, E_6^1, E_7^0, E_8^0$ or E_8^1 , then X is a member of one of the 56 families of surfaces in Tables 6, 7, 8, and 9.

For each of these surfaces, we have $h^0(X, T_X) > h^0(\widetilde{X}, T_{\widetilde{X}})$ and in particular $\operatorname{Aut}_X \neq \operatorname{Aut}_{\widetilde{X}}$.

Remark 1.3. The reason why we do not treat the case p = 2 is due to the sheer amount of RDP del Pezzo surfaces with global vector fields in characteristic 2. Indeed, by Theorem 6.1, it is unclear whether Aut $_{\tilde{X}}$ = Aut $_X$ as soon as X has a single node and in fact, even for the quadratic cone $\{x_0^2 - x_1 x_2 = 0\} \subseteq \mathbb{P}^3$ in characteristic 2 it is not true that every vector field lifts to its minimal resolution; consider for example $x_3\partial_{x_0}$. However, in principle, our approach would also work if p=2.

Comparing Tables 4, 5, 6, 7, 8, and 9 with the classification in [MS20], we see that in characteristics p =3,5 and 7, there exists an RDP del Pezzo surface X with $H^0(X, T_X) \neq 0$ whose minimal resolution admits no non-trivial global vector fields. In other words, we have the following picture, where the implications from right to left hold *only* in the indicated characteristics:

(1.3)
$$|\operatorname{Aut}(X)| = |\operatorname{Aut}(\widetilde{X})| = \infty \xrightarrow{p \neq 2,3 \atop \longleftarrow} H^0(\widetilde{X}, T_{\widetilde{X}}) \neq 0 \xrightarrow{p \neq 2,3,5,7} H^0(X, T_X) \neq 0$$

Acknowledgements. Research of the first named author was supported by the DFG Research Grant MA 8510/1-1. The second named author gratefully acknowledges funding by the DFG Sachbeihilfe LI 1906/5-1 "Geometrie von rationalen Doppelpunkten" and support by the doctoral program TopMath and the TUM Graduate School.

2. AN APPLICATION: REGULAR INSEPARABLE TWISTS OF RDP DEL PEZZO SURFACES

A twisted form of a k-scheme X over a field extension $L \supseteq k$ is a scheme Y over L such that $Y_{\bar{L}} \cong X_{\bar{L}}$, where \bar{L} is an algebraic closure of L. If X is a proper scheme over k, then smoothness of Aut_X is intimately related with properties of twisted forms of X, as the following proposition shows. Even though this proposition should be well-known, we include the proof for the convenience of the reader.

Proposition 2.1. Let $k \subseteq L$ be a field extension. Let X be a proper scheme over k and let Y be a twisted form of X over L. Assume that Aut_X is smooth. Then, the following hold:

- (1) If L is separably closed, then $Y \cong X_L$.
- (2) If \mathcal{P} is a property of schemes that is stable under field extensions and local in the étale topology, then, if X satisfies \mathcal{P} , also Y satisfies \mathcal{P} .

Proof. Let us first prove Claim (1). As explained for example in [Mil16, p.134], an isomorphism $\bar{\varphi} : Y_{\bar{L}} \cong X_{\bar{L}}$ gives rise to a Čech cocycle on the fppf site of Spec *L*, hence to an element in $\check{H}^1_{\text{fppf}}(\text{Spec } L, \text{Aut}_{X_L})$. By [Mil16, Chapter III: Theorem 4.3.(b), Corollary 4.7], the smoothness of Aut_X implies that

$$H^1_{\text{fppf}}(\operatorname{Spec} L, \operatorname{Aut}_{X_L}) = H^1_{\text{\acute{e}t}}(\operatorname{Spec} L, \operatorname{Aut}_{X_L})$$

and since L is separably closed, the latter is trivial. Hence, Y and X_L are already isomorphic over L.

For Claim (2), let L^{sep} be the separable closure of L in \overline{L} . By (1), we have $X_{L^{sep}} \cong Y_{L^{sep}}$. Since X is proper, this isomorphism is defined over a finite subextension $L \subseteq L' \subseteq L^{sep}$, so that $X_{L'} \cong Y_{L'}$. The morphism Spec $L' \to$ Spec L is finite and étale. Hence, by our assumptions on \mathcal{P} , if X satisfies \mathcal{P} , then $X_{L'}$ satisfies \mathcal{P} , and thus also Y satisfies \mathcal{P} .

Choosing for \mathcal{P} the property that the singular locus is non-empty and specializing to the case where X is an RDP del Pezzo surface, we obtain the following:

Corollary 2.2. Let X be an RDP del Pezzo surface over k. Let $k \subseteq L$ be a field extension and let Y be a twisted form of X over L. If X has at least one singular point and Y is regular, then Aut_X is non-smooth.

At a first glance, these twists seem rather hard to get a grip on geometrically, but it turns out that they can be written down explicitly if one has explicit descriptions of X and Aut_X. For example, consider $\mu_p \subseteq \text{PGL}_{n,k}$ embedded diagonally with weights $(a_0, a_1, \ldots, a_{n-1}, 1)$. We can write μ_p as the kernel of the surjective homomorphism

$$f: \quad \mathbb{G}_m^n \quad \to \mathbb{G}_m^n$$
$$(u_0, \dots, u_n) \quad \mapsto (u_0 u_n^{-a_0}, \dots, u_{n-1} u_n^{-a_{n-1}}, u_n^p).$$

By [Mil16, Proposition 4.5] and Hilbert 90, for every field extension $k \subseteq L$, this yields a short exact sequence of abelian groups

$$0 \to \mathbb{G}_m^n(L) \xrightarrow{f(L)} \mathbb{G}_m^n(L) \xrightarrow{d} \check{H}^1_{\text{fppf}}(\text{Spec } L, \mu_p) \to 0.$$

Here, for $g \in \mathbb{G}_m^n(L)$, the element d(g) is defined by choosing $\bar{g} \in \mathbb{G}_m^n(\bar{L})$ such that $f(\bar{L})(\bar{g}) = g_{\bar{L}}$ and setting d(g) to be the image of the cocycle $(\bar{g} \otimes 1)^{-1}(1 \otimes \bar{g}) \in \mu_p(\bar{L} \otimes_L \bar{L})$. If $X \subseteq \mathbb{P}_k^n$ is a subvariety stabilized by μ_p , then $\bar{g}^{-1}(X_{\bar{L}}) \subseteq \mathbb{P}_{\bar{L}}^n$ is defined over L and the cocycle one associates to this twisted form is in the same class as d(g). In other words, we can realize every twist of X corresponding to an element of $\check{H}^1_{\text{fppf}}(\text{Spec }L,\mu_p)$ by choosing $\bar{g} \in \mathbb{G}_m^n(\bar{L})$ such that $f(\bar{L})(\bar{g})$ is defined over L and translating X along

Example 2.3. Consider the quartic curve $Q = \{x^3y + y^3z + z^3x = 0\} \subseteq \mathbb{P}^2_k$ in characteristic 7. It is stable under the μ_7 -action with weights (4,2,1). Let L = k(t) and consider $\bar{g} = (t^{4/7}, t^{2/7}, t^{1/7}) \in \mathbb{G}_m^2(\overline{L}) \subseteq$ $PGL_3(L)$. Then,

$$\bar{g}^{-1}(Q) = \{t^2x^3y + ty^3z + tz^3x = 0\} = \{tx^3y + y^3z + z^3x = 0\}.$$

Spreading out over $\mathbb{A}_k^1 - \{0\} = \operatorname{Spec} k[t, t^{-1}]$, we obtain a fibered surface S over $\mathbb{A}_k^1 - \{0\}$ which is easily checked to be smooth using the Jacobian criterion. Its generic fiber is a twisted form of Q over k(t) and this twisted form is regular, because it is the generic fiber of a flat morphism between smooth k-schemes. Taking the double cover of $(\mathbb{A}^1 - \{0\}) \times \mathbb{P}^2$ branched over S, we obtain a smooth fibered threefold T over $\mathbb{A}^1 - \{0\}$ whose generic fiber is the regular del Pezzo surface Y of degree 2 given by the equation

$$\{w^2 = tx^3y + y^3z + z^3x\} \subseteq \mathbb{P}_{k(t)}(1, 1, 1, 2).$$

As before, Y is regular, being the generic fiber of a flat morphism of smooth k-schemes. Observe, however, that Y is not smooth, because $Y_{\overline{k(t)}} \cong X$, where X is the del Pezzo surface of degree 2 with a singularity of type A_6 given in our Table 4.

Remark 2.4. Example 2.3 shows that the bound $p \leq 7$ given in [BT20, Proposition 5.2] for the characteristics in which non-smooth regular RDP del Pezzo surfaces can exist is sharp. Using the approach explained in the beginning of Example 2.3, it is not hard to construct similar examples if p = 2, 3, 5, but since this is not the topic of this article, we leave these constructions to the interested reader. Finally, we note that it is no mere coincidence that the Klein quartic in characteristic 7 appears in this context and refer the reader to [Stö04] for a closer study of this curve and its regular twists.

3. PRELIMINARIES ON (RDP) DEL PEZZO SURFACES

In this section, we recall the definition of RDP del Pezzo surfaces and weak del Pezzo surfaces, which occur as minimal resolutions of RDP del Pezzo surfaces, as well as their basic properties.

Definition 3.1. Let X and \widetilde{X} be projective surfaces.

- X is a *del Pezzo surface* if it is smooth and $-K_X$ is ample.
- X is a *weak del Pezzo surface* if it is smooth and -K_X is big and nef.
 X is an *RDP del Pezzo surface* if all its singularities are rational double points and -K_X is ample.

In all the above cases, the number $\deg(X) = K_X^2$ (resp. $\deg(\widetilde{X}) = K_{\widetilde{X}}^2$) is called the *degree of* X (resp. \widetilde{X}).

Recall that $1 \leq \deg(X) = \deg(\widetilde{X}) \leq 9$ and that every weak del Pezzo surface of degree d and different from $\mathbb{P}^1 \times \mathbb{P}^1$ and the second Hirzebruch surface \mathbb{F}_2 can be realized as a blow-up of \mathbb{P}^2 in 9 - d (possibly infinitely near) points in almost general position.

As mentioned in the beginning of this section, weak del Pezzo surfaces arise as the minimal resolutions of RDP del Pezzo surfaces and, conversely, every RDP del Pezzo surface X is the anti-canonical model of a weak del Pezzo surface X. The linear systems $|-nK_{\tilde{X}}|$ are well studied (see e.g. [BT20, Proposition 2.14, Theorem 2.15] for proofs in positive characteristic). We denote the morphism induced by a linear system |D| by $\varphi_{|D|}$ and recall that a curve singularity is called *simple* if its completion is isomorphic to one of the normal forms given in [GK90, Section 1].

Theorem 3.2. Let \widetilde{X} be a weak del Pezzo surface of degree d.

- (1) If $d \geq 3$, then $\varphi_{|-K_{\widetilde{X}}|}$ factors as $\widetilde{X} \xrightarrow{\pi} X \xrightarrow{\varphi_{|-K_X|}} \mathbb{P}^d$, where $\varphi_{|-K_X|}$ is a closed immersion that realizes X as a surface of degree d.
- (2) If d = 2, then $\varphi_{|-K_{\widetilde{X}}|}$ factors as $\widetilde{X} \xrightarrow{\pi} X \xrightarrow{\varphi_{|-K_X|}} \mathbb{P}^2$, where $\varphi_{|-K_X|}$ is finite flat of degree 2. If $p \neq 2$, then $\varphi_{|-K_X|}$ is branched over a quartic curve Q with simple singularities.

(3) If d = 1, then the $\varphi_{|-2K_{\widetilde{X}}|}$ factors as $\widetilde{X} \xrightarrow{\pi} X \xrightarrow{\varphi_{|-2K_X|}} \mathbb{P}(1,1,2) \subseteq \mathbb{P}^3$, where $\mathbb{P}(1,1,2)$ is the quadratic cone and $\varphi_{|-2K_X|}$ is finite flat of degree 2. If $p \neq 2$, then $\varphi_{|-2K_X|}$ is branched over a sextic curve S with simple singularities.

Next, we recall the notion of a *marking* of a weak del Pezzo surface $\widetilde{X} \notin \{\mathbb{P}^1 \times \mathbb{P}^1, \mathbb{F}_2\}$ and explain how to describe the negative curves on \widetilde{X} in terms of such a marking.

- A marking of \widetilde{X} is an isomorphism $\phi : I^{1,9-d} \xrightarrow{\sim} \operatorname{Pic}(\widetilde{X})$, where $I^{1,9-d}$ is the lattice of rank 10 d with quadratic form given by the diagonal matrix $(1, -1, \ldots, -1)$ with respect to a basis e_0, \ldots, e_{9-d} .
- A realization π : X̃ → P² of X̃ as an iterated blow-up of P² induces a marking φ with φ(e₀) = π^{*}O_{P²}(1) and φ(e_i) is the class of the preimage in X̃ of the *i*-th point blown up by π. A marking that arises in this way is called *geometric*.
- If ϕ is a geometric marking of \widetilde{X} , then $\phi^{-1}(K_{\widetilde{X}}) = (-3, 1, \dots, 1) =: k_{9-d}$.
- The lattice E_{9-d} is defined as $\langle k_{9-d} \rangle^{\perp} \subseteq \mathrm{I}^{1,9-d}$.
 - For d = 1, 2, 3, the lattices E_{9-d} are precisely the three exceptional irreducible root lattices.
 - For d = 4, 5, 6, there are identifications $E_5 = D_5, E_4 = A_4, E_3 = A_2 \oplus A_1$.
 - For d = 7, 8, the lattices E_{9-d} are no root lattices. Every maximal root lattice contained in E_2 is isomorphic to A_1 , and E_1 does not contain any (-2)-vectors.

Following [Dol12, Section 8.2], we let

$$\operatorname{Exc}_{9-d} := \{ v \in \mathbf{I}^{1,9-d} \mid v^2 = -1, v.k_{9-d} = -1 \} \subseteq \mathbf{I}^{1,9-d}$$

be the subset of *exceptional vectors*. Let \mathcal{R} be a set of linearly independent (-2)-vectors in E_{9-d} , and define the cone

$$C_{\mathcal{R}} := \{ v \in \mathbf{I}^{1,9-d} \otimes \mathbb{R} \mid v.w \ge 0 \text{ for all } w \in \mathcal{R} \}.$$

For a sublattice Λ of E_{9-d} , we denote the Weyl group of Λ by $W(\Lambda)$. That is, $W(\Lambda)$ is the subgroup of the orthogonal group $O(I^{1,9-d})$ generated by reflections along (-2)-vectors in Λ . With this notation, $W(\Lambda)$ preserves Exc_{9-d} and, for $\Lambda = \langle \mathcal{R} \rangle$, $C_{\mathcal{R}}$ is a fundamental domain for the action of $W(\Lambda)$ on $I^{1,9-d} \otimes \mathbb{R}$.

Lemma 3.3. With the above notation, we have the following description of certain sets of (-1)-curves on \tilde{X} :

(1) If \mathcal{R} is the pre-image of the set of all (-2)-curves on \widetilde{X} under a geometric marking ϕ , then ϕ induces a bijection

$$\{(-1)\text{-}curves \text{ on } X\} \iff C_{\mathcal{R}} \cap \operatorname{Exc}_{9-d} \cong \operatorname{Exc}_{9-d}/W(\Lambda).$$

(2) If $\mathcal{R}' \subseteq \mathcal{R}$ with $\Lambda' := \langle \mathcal{R}' \rangle$, then ϕ induces a bijection

$$\{(-1)\text{-curves on } \widetilde{X} \text{ disjoint from } \phi(\mathcal{R}')\} \iff C_{\mathcal{R}} \cap \operatorname{Exc}_{9-d} \cap (\Lambda')^{\perp} = C_{\mathcal{R}} \cap \operatorname{Exc}_{9-d}^{W(\Lambda')}.$$

(3) If, moreover, Λ' is a sum of connected components of Λ , then ϕ induces a bijection

$$\{(-1)\text{-}curves \text{ on } \widetilde{X} \text{ disjoint from } \phi(\mathcal{R}')\} \iff (\operatorname{Exc}_{9-d}^{W(\Lambda')})/W(\Lambda)$$

Proof. For (1) and (2), see [Dol12, Lemma 8.2.22 and Proposition 8.2.34]. To prove (3), we note that we have an orthogonal decomposition $\Lambda = \Lambda' \oplus \Lambda''$, where $\Lambda = \langle \mathcal{R} \rangle$ and $\Lambda'' = \langle \mathcal{R} \setminus \mathcal{R}' \rangle$. Therefore, the $W(\Lambda)$ -action preserves $\operatorname{Exc}_{9-d} \cap (\Lambda')^{\perp} = \operatorname{Exc}_{9-d}^{W(\Lambda')}$ and, by (1), we can write

$$C_{\mathcal{R}} \cap \operatorname{Exc}_{9-d} \cap (\Lambda')^{\perp} = (C_{\mathcal{R}} \cap \operatorname{Exc}_{9-d})^{W(\Lambda')} \cong (\operatorname{Exc}_{9-d}^{W(\Lambda')})/W(\Lambda).$$

GEBHARD MARTIN AND CLAUDIA STADLMAYR

4. GROUP SCHEME ACTIONS ON ANTI-CANONICAL MODELS

The purpose of this section is to recall some basic facts about group scheme actions on (blow-ups of) normal projective surfaces and to describe the automorphism scheme of an RDP del Pezzo surface in terms of the anti-(bi-)canonical morphisms recalled in Theorem 3.2.

Quite generally, the key tool to control the behaviour of group scheme actions under birational morphisms is Blanchard's Lemma [Bri17, Theorem 7.2.1].

Theorem 4.1. (Blanchard's Lemma) Let $\pi : \widetilde{X} \to X$ be a morphism of proper schemes with $\pi_* \mathcal{O}_{\widetilde{X}} = \mathcal{O}_X$. Then, π induces a homomorphism of group schemes $\pi_* : \operatorname{Aut}^0_{\widetilde{X}} \to \operatorname{Aut}^0_X$. If π is birational, then π_* is a closed immersion.

Given an action of a group scheme G on a scheme X and a closed subscheme $Z \subseteq X$, we let $\operatorname{Stab}_G(Z) \subseteq \operatorname{Aut}_X$ be the *stabilizer subgroup scheme* of Z. The following proposition describes the image of π_* if π is a blow-up of a closed point on a normal surface with at worst rational double points.

Proposition 4.2. Let X be a normal surface with at worst rational double points and let $\pi : \widetilde{X} \to X$ be the blow-up of X in a closed point P. Then, $\pi_*(\operatorname{Aut}^0_{\widetilde{Y}}) = (\operatorname{Stab}_{\operatorname{Aut}_X}(P))^0$.

Proof. By [Mar20, Proposition 2.7], it suffices to find an infinitesimal rigid subscheme $E \subseteq \tilde{X}$ whose schematic image is P. Thus, let E be the exceptional divisor of π , with scheme structure given by the inverse image ideal sheaf of P. In particular, E is a Cartier divisor on \tilde{X} .

If P is smooth, then E is a (-1)-curve, hence E is infinitesimally rigid. If P is not smooth, then P is a rational double point. In particular, π factors the minimal resolution $\pi': \widetilde{X}' \to X$ and $\pi'^* \omega_X \cong \omega_{\widetilde{X}'}$. Since \widetilde{X} has at worst rational double points as well and $\widetilde{X}' \to \widetilde{X}$ is its minimal resolution, we obtain $\pi^* \omega_X \cong \omega_{\widetilde{X}}$ using the projection formula. Now, ω_X is trivial in a neighborhood of P, so $\omega_{\widetilde{X}}$ is trivial in a neighborhood of E. Thus, the normal sheaf $\mathcal{N}_{E/\widetilde{X}}$ of E in \widetilde{X} coincides with ω_E by adjunction.

On the other hand, the rational double point $P \in X$ has multiplicity 2 and embedding dimension 3, hence E is isomorphic to a (possibly non-reduced) conic in \mathbb{P}^2 . As such, it satisfies $\omega_E \cong \mathcal{O}_{\mathbb{P}^2}(-1)|_E$. Hence,

$$h^{0}(E, \mathcal{N}_{E/\widetilde{X}}) = h^{0}(E, \mathcal{O}_{\mathbb{P}^{2}}(-1)|_{E}) = 0,$$

so, by [Ser06, Proposition 3.2.1.(ii)], E is infinitesimally rigid. This finishes the proof.

If X is an RDP del Pezzo surface, then we have the following description of Aut_X in terms of the anticanonical morphisms $\varphi_{|-nK_X|}$ of X.

Proposition 4.3. Let X be an RDP del Pezzo surface of degree d. Then, $\varphi_{|-nK_X|}$ is Aut_X -equivariant for all $n \ge 0$ and the following hold.

- (1) If $d \ge 3$, then $\operatorname{Aut}_X = \operatorname{Stab}_{\operatorname{PGL}_{d+1}}(X)$.
- (2) If d = 2 and $p \neq 2$, then there is an exact sequence of group schemes

$$1 \to \mathbb{Z}/2\mathbb{Z} \to \operatorname{Aut}_X \to \operatorname{Stab}_{\operatorname{PGL}_3}(Q) \to 1,$$

where Q is the branch quartic of the anti-canonical morphism $X \to \mathbb{P}^2$.

(3) If d = 1 and $p \neq 2$, then there is an exact sequence of group schemes

$$1 \to \mathbb{Z}/2\mathbb{Z} \to \operatorname{Aut}_X \to \operatorname{Stab}_{\operatorname{Aut}_{\mathbb{P}(1,1,2)}}(S) \to 1,$$

where S is the branch sextic of the anti-bi-canonical morphism $X \to \mathbb{P}(1, 1, 2) \subseteq \mathbb{P}^3$.

Proof. By [Bri18, Remark 2.15.(iv)], the line bundles $\omega_X^{\otimes(-n)}$ admit natural Aut_X-linearizations for all $n \ge 0$ and hence the natural action of Aut_X on the space of global sections of $\omega_X^{\otimes(-n)}$ induces a homomorphism $f_n : \operatorname{Aut}_X \to \operatorname{PGL}_{N+1}$ for $N = \dim(H^0(X, \omega_X^{\otimes(-n)})) - 1$ making the rational map $\varphi_{|-nK_X|} : X \dashrightarrow \mathbb{P}^N$ Aut_X-equivariant.

If $d \ge 3$, then the anti-canonical map is an embedding by Theorem 3.2, hence f_1 is a monomorphism. By [Mar20, Lemma 2.5], f_1 factors through $\operatorname{Stab}_{\operatorname{PGL}_{d+1}}(X)$. Conversely, restricting the G-action on \mathbb{P}^d to X yields a left-inverse g_1 : $\operatorname{Stab}_{\operatorname{PGL}_{d+1}}(X) \to \operatorname{Aut}_X$ to f_1 . Since X is not contained in a proper linear subspace of \mathbb{P}^d and the fixed locus of a subgroup scheme of PGL_{d+1} is a linear subspace, g_1 has to be a monomorphism. Hence f_1 is an isomorphism.

If d = 2 and $p \neq 2$, then the anti-canonical map is a finite flat cover of \mathbb{P}^2 branched over a quartic curve $Q \subseteq \mathbb{P}^2$ by Theorem 3.2. Let K be the kernel of f_1 . Restricting the action of K on X to the generic point of X yields a $k(\mathbb{P}^2)$ -linear action of $K_{k(\mathbb{P}^2)}$ on the degree 2 field extension k(X) of $k(\mathbb{P}^2)$. Since $p \neq 2$, the field extension $k(\mathbb{P}^2) \subseteq k(X)$ is Galois, which shows $K = \mathbb{Z}/2\mathbb{Z}$. Since K is normal in Aut_X, the action of Aut_X on X preserves the fixed locus X^{K} , hence, by [Mar20, Lemma 2.5], the induced action of Aut_X on \mathbb{P}^2 preserves the scheme-theoretic image of X^K under $\varphi_{|-K_X|}$, which is nothing but Q. Hence, f_1 factors through $\operatorname{Stab}_{\operatorname{PGL}_3}(Q)$. In order to show faithful flatness of $f'_1 : \operatorname{Aut}_X \to \operatorname{Stab}_{\operatorname{PGL}_3}(Q)$, we write X as $\{w^2 = Q(x, y, z)\} \subseteq \mathbb{P}(1, 1, 1, 2)$. For every k-algebra R and every automorphism σ of \mathbb{P}^2_R preserving $Q_R = \{Q(x, y, z) = 0\} \subseteq \mathbb{P}^2_R$, that is, mapping Q(x, y, z) to $\lambda Q(x, y, z)$ for some $\lambda \in \mathbb{R}^{\times}$, we can pass to the faithfully flat ring extension $R' := R[\sqrt{\lambda}]$ of R and lift σ to an automorphism of $X_{R'}$ by mapping w to $\pm \sqrt{\lambda}w$. Hence, f'_1 is faithfully flat and thus the sequence in (2) is exact.

If d = 1 and $p \neq 2$, we can apply essentially the same argument as in the previous paragraph to the anti-bi-canonical morphism $\varphi_{|-2K_X|}: X \to \mathbb{P}^3$: Indeed, the argument for $K = \mathbb{Z}/2\mathbb{Z}$ is exactly the same as in the previous paragraph. To prove faithful flatness of $f'_2 : \operatorname{Aut}_X \to \operatorname{Stab}_{\operatorname{Aut}_{\mathbb{P}}(1,1,2)}(S)$, we can write X as a hypersurface in $\mathbb{P}(1, 1, 2, 3)$ by identifying the quadratic cone with $\mathbb{P}(1, 1, 2)$ and then argue as above. \Box

5. ON EQUIVARIANT RESOLUTIONS

It is an immediate consequence of the uniqueness of the minimal resolution \bar{X} of a projective surface X that the action of the automorphism group $\operatorname{Aut}_X(k)$ on X lifts to \widetilde{X} . Over the complex numbers, this implies that the action of the automorphism scheme Aut_X lifts to X. In general, this is no longer true in positive characteristic. In this section, we will study this phenomenon.

Definition 5.1. Let $\pi : \tilde{X} \to X$ be a proper birational morphism of schemes.

- (1) The morphism π is called T_X -equivariant if the natural map $\pi_*T_{\widetilde{X}} \to T_X$ is an isomorphism.
- (2) Assume additionally that X is proper and $\pi_* \mathcal{O}_{\widetilde{X}} = \mathcal{O}_X$. Then, π is called Aut_X^0 -equivariant if the closed immersion $\pi_* : \operatorname{Aut}^0_{\widetilde{X}} \hookrightarrow \operatorname{Aut}^0_X$ induced by Blanchard's Lemma is an isomorphism.

Remark 5.2. Note that T_X -equivariance is local on X and implies $H^0(\widetilde{X}, T_{\widetilde{X}}) \cong H^0(X, T_X)$. If π is T_X -equivariant, then $\pi_* : \operatorname{Aut}^0_{\widetilde{Y}} \hookrightarrow \operatorname{Aut}^0_X$ is an isomorphism on tangent spaces.

The study of the T_X -equivariance of the minimal resolution of a rational double point has been initiated by Wahl [Wah75] and extended to all positive characteristics by Hirokado [Hir19]. There, T_X -equivariance is simply called "equivariance". For the convenience of the reader, we will recall the classification of RDPs whose minimal resolution is not T_X -equivariant (see [Hir19, Theorem 1.1]).

Proposition 5.3. Let $\pi : \widetilde{X} \to X$ be the minimal resolution of a rational double point (X, x). Then, π is not T_X equivariant if and only if (X, x) is of type

- (1) A_n with $p \mid (n+1)$,

- (2) E_8^0 if p = 5, (3) $E_6^0, E_6^1, E_7^0, E_8^0, E_8^1$ if p = 3, or (4) $D_n^r, E_6^0, E_7^0, E_7^1, E_7^2, E_7^3, E_8^0, E_8^1, E_8^2, E_8^3$ if p = 2.

In the next sections, we would like to apply the notions of Aut_X^0 -equivariance and T_X -equivariance to RDP del Pezzo surfaces and their partial resolutions.

Definition 5.4. Let X be a proper surface. A *partial resolution* of X is a proper birational morphism $\pi: X \to X$ such that the minimal resolution of X factors through π .

Proposition 5.5. Let X be a normal proper surface and let $\pi : \widetilde{X} \to X$ be a partial resolution of X. Assume that there exists an open subset $U \subseteq X$ such that $\pi^{-1}(U) \to U$ is an isomorphism and all singularities in $X \setminus U$ admit a T_X -equivariant minimal resolution. Then, π is T_X -equivariant.

Proof. Let U_{sing} be the set of singular points in U and let $V = X \setminus U_{sing}$. Let $\psi : V' \to V$ be the minimal resolution of V. In particular, $V' \supseteq (\psi)^{-1}(U \setminus U_{sing}) \to (U \setminus U_{sing}) \subseteq U$ is an isomorphism, which we can use to glue V' and U along the smooth locus $(U \setminus U_{sing})$ of U to a projective surface X' together with a proper birational morphism $\pi' : X' \to X$. Since π is a partial resolution of X, π' factors through π by construction.

Moreover, by the assumption on the singularities in $X \setminus U$ and Remark 5.2, π' is T_X -equivariant, that is, the composition

$$\pi'_*T_{X'} \to \pi_*T_{\widetilde{X}} \to T_X$$

is an isomorphism, hence the second map is surjective and thus an isomorphism, since T_X satisfies the (S_2) -condition. In particular, if additionally $\operatorname{Aut}_{\widetilde{X}}^0$ is smooth, this implies that π is Aut_X^0 -equivariant by Proposition 5.6.

In some situations, Aut_X^0 -equivariance can be deduced immediately from the simpler notion of T_X -equivariance, which is the content of the following proposition.

Proposition 5.6. Let $\pi : \widetilde{X} \to X$ be a birational morphism of proper k-schemes. If $\operatorname{Aut}_{\widetilde{X}}^0$ is smooth and π is T_X -equivariant, then π is Aut_X^0 -equivariant.

Proof. If π is T_X -equivariant, then

$$\dim \operatorname{Aut}_{\widetilde{X}}^{0} \leq \dim \operatorname{Aut}_{X}^{0} \leq \dim_{k} H^{0}(X, T_{X}) = \dim_{k} H^{0}(\widetilde{X}, T_{\widetilde{X}})$$

and since $\operatorname{Aut}_{\widetilde{X}}^0$ is smooth, all inequalities above are in fact equalities. Thus, Aut_X^0 is smooth and of the same dimension as $\operatorname{Aut}_{\widetilde{X}}^0$. Hence, we must have $\operatorname{Aut}_{\widetilde{X}}^0 = \operatorname{Aut}_X^0$, that is, π is Aut_X^0 -equivariant.

Remark 5.7. In particular, if, in the situation of Proposition 5.5, we assume in addition that $\operatorname{Aut}_{\widetilde{X}}^0$ is smooth, then π is Aut_X^0 -equivariant.

To the best of our knowledge, the question whether partial resolutions of a proper normal surface with rational double points are Aut_X^0 -equivariant has not been studied. In the following, we prove Aut_X^0 -equivariance for A_n -singularities with $n and bound the failure of <math>\operatorname{Aut}_X^0$ -equivariance for A_{p-1} -singularities. While this does not cover all rational double points and not even all A_n -singularities, it will come in handy for the calculation of the automorphism schemes of non-equivariant RDP del Pezzo surfaces in Section 8.

Proposition 5.8. Let X be a proper surface. Let $\pi : \tilde{X} \to X$ be a partial resolution of X and assume that the only singularities over which π is not an isomorphism are A_n -singularities with $n \leq p - 1$. Then,

$$\operatorname{length}\left(\operatorname{Aut}_{X}^{0}/\operatorname{Aut}_{\widetilde{X}}^{0}\right) \leq p^{m},$$

where m is the number of A_{p-1} -singularities on X over which π is not an isomorphism. In particular, if m = 0, then π is Aut_X -equivariant.

Proof. It suffices to prove the statement if $\pi : X \to X$ is not an isomorphism only over a single singularity P of type A_n . By Proposition 4.2, it suffices to show that $G := \operatorname{Aut}_X^0$ fixes P if n < p-1 and that the stabilizer of P has index 1 or p in G if n = p - 1. To see this, we equip the singular locus X_{sing} of X with a scheme structure using Fitting ideals and we let Y be the irreducible component of X_{sing} containing P. Since the scheme structure on X_{sing} is canonical and G is connected, G preserves Y, so we get a homomorphism $\varphi : G \to \operatorname{Aut}_Y$. To prove the proposition, it suffices to show that the stabilizer of P in Aut_Y has index 1 or p, with the latter only occurring for n = p - 1.

Since an A_n -singularity is given in a formal neighborhood by the equation $z^{n+1} + xy$, Y is isomorphic to

$$Y_n := \text{Spec} \left(k[z]/(z^{n+1}) \right) \quad \text{if} \quad n < p-1$$
$$Y_{p-1} := \text{Spec} \left(k[z]/(z^p) \right) \quad \text{if} \quad n = p-1.$$

Now, we calculate Aut_{Y_i} by computing its R-valued points for an arbitrary local k-algebra R. An element of $\operatorname{Aut}_{Y_i}(R)$ is an R-linear automorphism φ of $R[z]/(z^{i+1})$, hence it is determined by where it sends z. Let $a_0, \ldots, a_i \in R$ such that $\varphi(z) = \sum_{j=0}^i a_j z^j$. Let \mathfrak{m} be the maximal ideal of R, so that (\mathfrak{m}, z) is the maximal ideal of $R[z]/(z^{i+1})$. Since φ is an automorphism, it maps (\mathfrak{m}, z) to itself, hence $a_0 \in \mathfrak{m}$. If $a_1 \in \mathfrak{m}$, then the coefficient of z in every $\varphi(z^j)$ is in \mathfrak{m} , so z would not lie in the image of φ , which is absurd. Hence, $a_1 \in R \setminus \mathfrak{m} = R^{\times}$. Next, we know that $\varphi(z^{i+1}) = 0$. Since the degree 0 term of $\varphi(z^{i+1})$ is a_0^{i+1} , we have $a_0^{i+1} = 0$. The degree j term of $\varphi(z^{i+1})$ is of the form $\binom{i+1}{j}a_0^{i+1-j}a_1^j + a_0^{i+2-j}b_j$ for some $b_j \in R$. If i < p-1, then $p \nmid \binom{i+1}{j}$ for all j > 0, so we only get $a_0^p = 0$.

Conversely, given a sequence (a_0, \ldots, a_i) in R with $a_1 \in R^{\times}$ and $a_0 = 0$ if $i (resp. <math>a_0^p = 0$ if i = p), the morphism induced by $z \mapsto \sum_{j=0}^{i} a_j z^j$ is an automorphism, since it is well-defined and its inverse is given by $z \mapsto (\sum_{j=1}^{i} a_j z^{j-1})^{-1} z - a_0$.

Summarizing, we have natural identifications

$$\operatorname{Aut}_{Y_i}(R) = \{ (0, a_1, \dots, a_i) \mid a_j \in R, a_1 \in R^{\times} \} \text{ for } i
$$\operatorname{Aut}_{Y_{p-1}}(R) = \{ (a_0, a_1, \dots, a_{p-1}) \mid a_j \in R, a_1 \in R^{\times}, a_0^p = 0 \}.$$$$

In both cases, the corresponding automorphism of Y_i preserves $P \times \operatorname{Spec} R \subseteq Y_i \times \operatorname{Spec} R$ if and only if $a_0 = 0$, since the ideal of $P \times \operatorname{Spec} R$ is (z). Hence, the index of the stabilizer of P in Aut_{Y_i} is 1 if i < p-1, and p if i = p - 1. This finishes the proof.

Remark 5.9. The strategy of proof of Proposition 5.8 would, in principle, also apply to other rational double points. However, there are two obstacles to overcome:

- (1) The automorphism scheme of the singular locus is more complicated for more general RDPs, since the singular locus has a more complicated scheme structure in general. For example, if p = 5 and X admits an RDP of type E_8^0 , then, in a neighborhood of this singularity, the singular locus of X looks like Spec $k[[x, y]]/(x^2, y^5)$. This also makes the calculation of the stabilizer of the closed point more complicated.
- (2) If π : X̃ → X is the minimal resolution of an RDP surface, then Aut⁰_{X̃} is the intersection of all stabilizers of all singularities that occur in the blow-ups making up π. For example, if p = 3 and X admits a single RDP of type A₄, then the argument of Proposition 5.8 shows that Aut⁰_{X′} = Aut⁰_X, where X′ is the blow-up of the closed point of P, but X′ has a singularity of type A₂, so the approach of Proposition 5.8 only shows that length (Aut⁰_{X̃} / Aut⁰_{X̃}) ≤ 3 even though we would expect the two group schemes to be equal by Proposition 5.3.

One case where the argument of Proposition 5.8 goes through essentially unchanged is if p = 3 and the morphism $\pi : \widetilde{X} \to X$ is the minimal resolution of an RDP of type A_5 . In this case, π factors as a composition of three blow-ups $\widetilde{X} \to X'' \to X'$, where X' has an A_3 -singularity and X'' has an A_1 -singularity. Then, the argument of Proposition 5.8 shows that $\operatorname{Aut}^0_{\widetilde{X}} = \operatorname{Aut}^0_{X''} = \operatorname{Aut}^0_{X'}$ and length $(\operatorname{Aut}^0_X / \operatorname{Aut}^0_{\widetilde{X}}) \leq 3$, hence length $(\operatorname{Aut}^0_X / \operatorname{Aut}^0_{\widetilde{X}}) \leq 3$.

6. AUTOMORPHISM SCHEMES OF EQUIVARIANT RDP DEL PEZZO SURFACES

In [MS20], we classified all weak del Pezzo surfaces \widetilde{X} with global vector fields and calculated the identity component $\operatorname{Aut}^0_{\widetilde{X}}$ of their automorphism schemes. In particular, if X is a projective surface, whose minimal resolution $\pi: \widetilde{X} \to X$ is Aut_X^0 -equivariant and such that \widetilde{X} is a weak del Pezzo surface, then $\operatorname{Aut}_X^0 = \operatorname{Aut}_{\widetilde{X}}^0$ and thus, if Aut_X^0 is non-trivial, then \widetilde{X} appears in the classification tables of [MS20].

In the following, we will observe that all RDP del Pezzo surfaces in characteristic $p \ge 11$ fall into the above category and we will give a list of possible candidates for exceptions in small characteristics.

Theorem 6.1. Let X be an RDP del Pezzo surface over an algebraically closed field of characteristic p and let $\pi: X \to X$ be its minimal resolution. Assume that one of the following conditions holds:

- (1) $p \notin \{2, 3, 5, 7\}.$
- (2) p = 7 and X does not contain an RDP of type A_6 .
- (3) p = 5 and X does not contain an RDP of type A_4 or E_8^0 .
- (4) p = 3 and X does not contain an RDP of type $A_2, A_5, A_8, E_6^0, E_6^1, E_7^0, E_8^0$ or E_8^1 . (5) p = 2 and X does not contain an RDP of type $A_1, A_3, A_5, A_7, D_n^r, E_6^0, E_7^0, E_7^1, E_7^2, E_8^3, E_8^1, E_8^2$ or E_8^3 .

Then, $\operatorname{Aut}_X = \operatorname{Aut}_{\widetilde{X}}$, and thus, in particular, $H^0(\widetilde{X}, T_{\widetilde{X}}) = H^0(X, T_X)$. Therefore, $H^0(X, T_X) \neq 0$ if and only if X is the anti-canonical model of one of the surfaces in the classification tables of [MS20].

Proof. By Proposition 5.5 and Proposition 5.6, the theorem holds for those RDP del Pezzo surfaces that satisfy the following two conditions:

- (a) all singularities of X admit a T_X -equivariant minimal resolution, and
- (b) $\operatorname{Aut}^{0}_{\widetilde{V}}$ is smooth.

By Proposition 5.3, Condition (a) holds if we exclude the types of RDPs in the statement of the theorem.

Once we exclude those types of RDPs, then, by Tables 1 - 6 in [MS20], Condition (b) is also satisfied unless we are in one of the following three cases, where Γ is the RDP configuration on X and $d = \deg(X)$:

- (i) $p = 3, d = 2, \Gamma = A_6$ (ii) $p = 3, d = 2, \Gamma = D_6$
- (iii) $p = 2, d = 3, \Gamma = A_4$

In [MS20], these exceptions correspond to cases 2J, 2K, and 3N, respectively, and there is a unique weak del Pezzo surface of each of these types. In all cases, we have $H^0(\tilde{X}, T_{\tilde{X}}) = 1$, hence $H^0(X, T_X) = 1$ by Proposition 5.3, so the only remaining statement we have to show in these three cases is that π is Aut⁰_xequivariant. We will check this via explicit calculation:

(i) Assume p = 3. Consider the surface

$$X := \{w^2 = x^2 z^2 + x y^2 z + y^4 + x^3 y\} \subseteq \mathbb{P}(1, 1, 1, 2)$$

and let Q be the branch quartic of the induced double cover $X \to \mathbb{P}^2$. An elementary calculation shows that X admits an A_6 -singularity over [0:0:1] and no other singularity. By Proposition 4.3, we have $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{PGL}_3}(Q)^0$. The diagonal μ_3 -action with weights (0, 1, 2) on \mathbb{P}^2 preserves Q, hence $\mu_3 \subseteq \operatorname{Aut}_X^0$. Since π is T_X -equivariant by Proposition 5.3, \widetilde{X} must be the surface of type 2J of [MS20], hence $\operatorname{Aut}_{\widetilde{X}}^0 = \mu_3$. Since $H^0(X, T_X) = 1$, we have $\operatorname{Aut}_X^0[F] = \mu_3$, where $\operatorname{Aut}_X^0[F]$ denotes the kernel of Frobenius on Aut_X^0 . Hence, μ_3 is normal in $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{PGL}_3}(Q)^0$ and thus $\operatorname{Stab}_{\operatorname{PGL}_3}(Q)^0$ preserves the eigenspaces of the μ_3 -action, hence $\operatorname{Stab}_{\operatorname{PGL}_3}(Q)^0$ acts diagonally. With this restriction, it is easy to compute that $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{PGL}_3}(Q)^0 = \mu_3$. Therefore, π is $\operatorname{Aut}_{X}^{0}$ -equivariant, which is what we wanted to show.

(ii) Assume p = 3. Consider the surface

$$X := \{w^2 = x(x^3 + y^3 + xyz)\} \subseteq \mathbb{P}(1, 1, 1, 2)$$

and let Q be the branch quartic of the induced double cover $X \to \mathbb{P}^2$. Note that Q is the union of a nodal cubic and one of its nodal tangents, with the node located at [0:0:1], hence X has a D_6 -singularity at [0:0:1:0] and no other singularities. The diagonal μ_3 -action with weights (0,1,2) preserves Q, hence $H^0(X, T_X) \neq 0$, and thus \widetilde{X} is the surface of type 2K of [MS20]. The rest of the argument is the same as in the previous Case (i), and shows that π is Aut_X^0 -equivariant.

(iii) Assume p = 2. Consider the surface

$$X := \{x_0 x_1 x_3 + x_1^2 x_2 + x_0 x_2^2 + x_0^2 x_2\} \subseteq \mathbb{P}^3$$

which is a cubic surface with a single singularity, which is of type A_4 , at [0:0:0:1] (see e.g. [Roc96, Case B]). It admits a diagonal μ_2 -action with weights (0, 1, 0, 1), hence $H^0(X, T_X) \neq 0$ and therefore, as π is T_X -equivariant by Proposition 5.3, X is the anti-canonical model of the surface of type 3N in [MS20]. Straightforward calculation, again using that μ_2 is normal in $\text{Stab}_{PGL_4}(X)^0$, shows that $\text{Aut}_X^0 = \text{Stab}_{PGL_4}(X)^0 = \mu_2$, hence π is Aut_X^0 -equivariant.

7. Finding (-1)-curves in the equivariant locus

In view of Theorem 6.1, in order to classify RDP del Pezzo surfaces with global vector fields in odd characteristic, it remains to study RDP del Pezzo surfaces X which are not T_X -equivariant. Let Γ' be the configuration of rational double points on X which are not T_X -equivariant and let $\pi : \widetilde{X} \to X$ the minimal resolution of X.

In this section, we will describe a criterion for X to be the anti-canonical model of a blow-up of an RDP del Pezzo surface X' of higher degree containing Γ' such that $X' \to X$ is an isomorphism around Γ' . On the corresponding minimal resolutions, this will amount to finding (-1)-curves away from the configuration of (-2)-curves over Γ' . In other words, we are trying to find (-1)-curves on \widetilde{X} that map to the T_X -equivariant locus of π . In Section 8, this criterion will allow us to give a complete classification of non-equivariant RDP del Pezzo surfaces with global vector fields by setting up an inductive argument depending on the degree of the surface.

Theorem 7.1. Let X_d be an RDP del Pezzo surface of degree $d \le 8$ and let Γ' be a configuration of rational double points on X_d . Assume that its minimal resolution \tilde{X}_d is a blow-up of \mathbb{P}^2 and let $\Lambda' \subseteq \operatorname{Pic}(\tilde{X}_d)$ be the sublattice generated by the components of the exceptional locus over Γ' . Then the following are equivalent:

- (1) There exists a (-1)-curve on \widetilde{X}_d whose image in X_d does not pass through Γ' .
- (2) X_d is the anti-canonical model of a blow-up in a smooth point P of an RDP del Pezzo surface X_{d+1} of degree (d+1) containing Γ' such that $X_d \dashrightarrow X_{d+1}$ is an isomorphism around Γ' .
- (3) The map $\Lambda' \hookrightarrow \langle K_{\widetilde{X}_d} \rangle^{\perp} \cong E_{9-d}$ factors through an embedding $E_{8-d} \hookrightarrow E_{9-d}$.

Proof. First, we show $(1) \Rightarrow (2)$. Let \widetilde{C} be the (-1)-curve whose existence is asserted in (1). Contracting \widetilde{C} , we obtain a weak del Pezzo surface \widetilde{X}_{d+1} of degree (d+1) such that \widetilde{X}_d is the blow-up of \widetilde{X}_{d+1} in a smooth point \widetilde{P} . Let X_{d+1} be the anti-canonical model of \widetilde{X}_{d+1} and let P be the image of \widetilde{P} in X_{d+1} . By our choice of \widetilde{C} , all components of the exceptional locus over Γ' stay (-2)-curves in \widetilde{X}_{d+1} , hence X_{d+1} contains Γ' .

Since \widetilde{X}_d is a weak del Pezzo surface, \widetilde{P} cannot lie on a (-2)-curve (otherwise the strict transform of such a curve would have negative intersection with $-K_{\widetilde{X}_d}$, which is impossible as $-K_{\widetilde{X}_d}$ is nef), hence P is a smooth point on X_{d+1} . Thus, blowing up $P \in X_{d+1}$, we obtain a surface Y_d with the same singularities as X_{d+1} . In particular, Y_d has only rational double points as singularities and its minimal resolution is \widetilde{X}_d . Therefore, pullback of sections induces isomorphisms $H^0(Y_d, -nK_{Y_d}) \cong H^0(\widetilde{X}_d, -nK_{\widetilde{X}_d})$ for all $n \ge 0$, where the surjectivity follows from the fact that Y_d is normal. Thus, the anti-canonical model of Y_d coincides with X_d . The situation is summarized in the following Figure 1.

Note that $X_d \dashrightarrow Y_d \to X_{d+1}$ is an isomorphism in a neighborhood of Γ' , since \widetilde{C} is disjoint from the exceptional locus over Γ' .



Figure 1. Contracting a (-1)-curve disjoint from the singular locus

Next, we show (2) \Rightarrow (3). We have $\langle K_{\widetilde{X}_d} \rangle^{\perp} \cong E_{9-d}$ and $\langle K_{\widetilde{X}_{d+1}} \rangle^{\perp} \cong E_{8-d}$. Since X_{d+1} contains Γ' , the embedding $\Lambda' \hookrightarrow \operatorname{Pic}(\widetilde{X}_d)$ factors through the pullback map $\operatorname{Pic}(\widetilde{X}_{d+1}) \hookrightarrow \operatorname{Pic}(\widetilde{X}_d)$, which maps $\langle K_{\widetilde{X}_{d+1}} \rangle^{\perp}$ to $\langle K_{\widetilde{X}_d} \rangle^{\perp}$. Hence (3) follows.

Finally, to show that (3) \Rightarrow (1), we identify $\operatorname{Pic}(\widetilde{X}_d)$ and $I^{1,9-d}$ via a geometric marking. We have to show that there is a (-1)-curve \widetilde{C} on \widetilde{X}_d that does not meet the set \mathcal{R}' of exceptional curves over Γ' . Let Λ be the sublattice of $\operatorname{Pic}(\widetilde{X}_d)$ spanned by the classes of all (-2)-curves. Since Λ' is a sum of connected components of Λ , Lemma 3.3 shows that it suffices to prove

$$(\operatorname{Exc}_{9-d}^{W(\Lambda')})/W(\Lambda) \neq \emptyset.$$

Clearly, this is the case if and only if $\operatorname{Exc}_{9-d}^{W(\Lambda')} \neq \emptyset$. Since $\Lambda' \hookrightarrow E_{9-d}$ factors through an embedding $E_{8-d} \hookrightarrow E_{9-d}$, we have

$$\operatorname{Exc}_{9-d}^{W(E_{8-d})} \subseteq \operatorname{Exc}_{9-d}^{W(\Lambda')},$$

so it suffices to show that $\operatorname{Exc}_{9-d}^{W(E_{8-d})} \neq \emptyset$. Since the action of $W(E_{9-d})$ on $\operatorname{Pic}(\widetilde{X}_d)$ preserves Exc_{9-d} , the condition $\operatorname{Exc}_{9-d}^{W(E_{8-d})} \neq \emptyset$ depends on the embedding $E_{8-d} \hookrightarrow E_{9-d}$ only up to conjugation by elements of $W(E_{9-d})$ and up to automorphisms of E_{8-d} .

If $d \leq 5$, then E_{8-d} is a root lattice. By [Dyn52, Table 11] and [Mar03, Exercise 4.2.1, 4.6.2], the embedding $\iota : E_{8-d} \hookrightarrow E_{9-d}$ is unique up to the action of $O(E_{9-d})$. Since $O(E_{9-d})$ is generated by $\{\pm id\}$ and $W(E_{9-d})$ in every case (see e.g. [Mar03, Proposition 4.2.2, Theorem 4.3.3, 4.5.2, 4.5.3]), ι is unique up to the action of $W(E_{9-d})$ and up to automorphisms of E_{8-d} . Therefore, in order to show that $\operatorname{Exc}_{9-d}^{W(E_{8-d})} \neq \emptyset$, it suffices to show that there exists *some* \widetilde{X}_d containing a configuration of (-2)-curves of type E_{8-d} and such that a (-1)-curve disjoint from this configuration exists. This is known and can be seen for example in [MS20, Figures 23, 48, 57, 63, and 61].

If $d \ge 7$, then E_{8-d} does not contain any (-2)-vectors, hence $\Lambda' = 0$ and the implication $(3) \Rightarrow (1)$ holds, since \widetilde{X}_d is a blow-up of \mathbb{P}^2 by assumption.

Finally, if d = 6, then the maximal root lattice contained in $E_{8-d} = E_2$ is A_1 . Thus, we may assume $\Lambda' = A_1$, for otherwise we can argue as in the previous Case $d \ge 7$. Up to the action of $O(E_3) = \{\pm id\} \times W(E_3)$, there are two embeddings of A_1 into $E_3 = A_2 \oplus A_1$. It is easy to check that $\iota : A_1 \hookrightarrow E_3$ factors through E_2 if and only if ι factors through the A_2 -summand of E_3 and then ι is unique up to the action of $W(E_3)$ and up to automorphisms of A_1 . Hence, similar to what we did in the case $d \le 5$, it suffices to find some \tilde{X}_6 containing a (-2)-curve and a disjoint (-1)-curve. Again, this is known, see [MS20, Figure 24].

Corollary 7.2. Let X_d be an RDP del Pezzo surface of degree $d \le 8$, let Γ' be a configuration of rational double points on X_d , and let Λ' be the root lattice associated to Γ' . If Γ' occurs on an RDP del Pezzo surface of degree (d + 1) and satisfies one of the following conditions:

(1) $d \neq 4, 2, 1,$ (2) $d = 4 \text{ and } \Lambda' \neq A_3,$ (3) $d = 2 \text{ and } \Lambda' \notin \{A_5 + A_1, A_5, A_3 + 2A_1, A_3 + A_1, 4A_1, 3A_1\},$ (4) $d = 1 \text{ and } \Lambda' \notin \{A_7, 2A_3, A_5 + A_1, A_3 + 2A_1, 4A_1\}.$ Then X_d is the anti-canonical model of a blow-up in a smooth point P of an RDP del Pezzo surface X_{d+1} of degree (d+1) containing Γ' such that $X_d \dashrightarrow X_{d+1}$ is an isomorphism around Γ' .

Proof. The embeddings of root lattices into E_6, E_7 , and E_8 have been classified by Dynkin [Dyn52, Table 11] and for the embeddings of root lattices into $E_3 = A_2 \oplus A_1, E_4 = A_4$, and $E_5 = D_5$, we refer the reader to [Mar03, Exercise 4.2.1, 4.6.2]. It follows from these classifications that if an embedding of Λ' into E_{9-d} exists, then this embedding is unique (up to the action of $O(E_{9-d})$), except precisely in the cases excluded in (2), (3) and (4). Hence, if one embedding $\Lambda' \hookrightarrow E_{9-d}$ factors through an embedding $E_{8-d} \hookrightarrow E_{9-d}$, then *every* embedding factors through an embedding $E_{8-d} \hookrightarrow E_{9-d}$. If Γ' occurs on some RDP del Pezzo surface of degree (d + 1), then an embedding of Λ' with such a factorization exists and the claim follows from Theorem 7.1.

8. AUTOMORPHISM SCHEMES OF NON-EQUIVARIANT RDP DEL PEZZO SURFACES

Throughout this section, X denotes an RDP del Pezzo surface and $\pi : \tilde{X} \to X$ is its minimal resolution. In this section, we will prove Theorem 1.2, that is, we will classify all X with $H^0(X, T_X) \neq 0$ over a field of characteristic $p \in \{3, 5, 7\}$ and such that X contains one of the RDPs excluded in Theorem 6.1. We will treat the cases p = 7, p = 5, and p = 3, in Sections 8.1, 8.2, and 8.3, respectively. This will complete the classification of all RDP del Pezzo surfaces with global vector fields in odd characteristic.

The strategy of proof is as follows: First, for each degree $1 \le d \le 9$, we give the list of RDP configurations Γ that can occur on an RDP del Pezzo surface X of degree d and that contain at least one RDP whose minimal resolution is not T_X -equivariant. Then, starting with the highest possible degree and working our way down with Theorem 7.1, we classify those X containing Γ and satisfying $H^0(X, T_X) \ne 0$. In each step, we give explicit equations and calculate Aut^0_X using Proposition 4.2, Proposition 4.3, and Proposition 5.8.

Notation 8.1. If $d \ge 3$, we use the notation x_0, \ldots, x_d for the coordinates of \mathbb{P}^d . If d = 2, we use the notation x, y, z and w for the coordinates of $\mathbb{P}(1, 1, 1, 2)$, where w has weight 2. Finally, if d = 1, we use the notation s, t, x and y for the coordinates of $\mathbb{P}(1, 1, 2, 3)$, where x has weight 2 and y has weight 3. In Tables 4, 5, 6, 7, 8, and 9, we describe Aut_X^0 as follows:

- We only describe the *R*-valued points of Aut_X^0 , where *R* is an arbitrary local *k*-algebra. By [MS20, Lemma 3.5], this suffices to describe the scheme structure of Aut_X^0 completely. We do this by either describing a general *R*-valued point as a matrix or by describing the image of $[x_0 : \ldots : x_n]$ (resp. [x : y : z : w], resp. [s : t : x : y]) under a general *R*-valued automorphism of *X*.
- We often describe Aut_X^0 as the group scheme $\langle G_1, G_2 \rangle$ generated by subgroup schemes G_1 and G_2 of Aut_X^0 . By this we mean that we describe Aut_X^0 , using Proposition 4.3, as the smallest subgroup scheme of PGL_{d+1} (resp. $\operatorname{Aut}_{\mathbb{P}(1,1,2)}$ if d = 2, resp. $\operatorname{Aut}_{\mathbb{P}(1,1,2,3)}$ if d = 1) containing both G_1 and G_2 .
- We use the variables λ or λ_i for *R*-valued points of \mathbb{G}_m and μ_{p^n} (where $\lambda^{p^n} = 1$), and the variables ε or ε_i for *R*-valued points of \mathbb{G}_a and α_{p^n} (where $\varepsilon^{p^n} = 0$).

8.1. In characteristic 7. By Theorem 6.1, we have to list all RDP configurations containing A_6 that can occur on an RDP del Pezzo surface in characteristic 7.

Lemma 8.2. If p = 7, deg(X) = d, and X contains an A₆-singularity, then d and the configuration Γ of RDPs on X is one of the cases in Table 1.

d	Г	$\subseteq \langle k_{9-d} \rangle^{\perp}$
2	A_6	$\subseteq E_7$
1	$A_6, A_6 + A_1$	$\subseteq E_8$

Table 1. Non-equivariant RDP configurations in characteristic 7

Proof. Since A_6 has rank 6 and discriminant 7, it does not embed into E_{9-d} with $d \ge 3$. By [Dyn52, Table 11], the only root lattice containing A_6 and embedding into E_7 is A_6 itself and there are precisely two root lattices containing A_6 and embedding into E_8 , namely A_6 and $A_6 + A_1$.

Theorem 8.3. Assume that p = 7 and X contains an RDP of type A_6 . Then, $H^0(X, T_X) \neq 0$ if and only if X is given by an equation as in Table 4. Moreover, Aut_X^0 is as in Table 4, so that $\operatorname{Aut}_{\widetilde{X}}^0 \subsetneq \operatorname{Aut}_X^0$ and even $h^0(X, T_X) > h^0(\widetilde{X}, T_{\widetilde{X}})$.

Proof. By Table 1, we have $deg(X) = d \le 2$. Assume $H^0(X, T_X) \ne 0$.

If d = 2, then, by Theorem 3.2, the anti-canonical system of X realizes X as a double cover of \mathbb{P}^2 branched over a quartic curve Q with a simple singularity of type A_6 . Over the complex numbers, there is a unique such Q (see [BG81, Proposition 1.3.II]) and the argument carries over without change to characteristic 7. Now, an elementary calculation shows that the Klein quartic equation

$$x^{3}y + y^{3}z + z^{3}x = 0$$

defines such a Q with an A_6 -singularity at [1:2:-3]. Thus, X is given by the equation in Table 4. Clearly, the μ_7 -action described in Table 4 preserves X. Since $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial by [MS20], Proposition 5.8 implies that $\operatorname{Aut}^0_X = \mu_7$.

If d = 1, then, by Theorem 3.2, the anti-bi-canonical system of X realizes X as a double cover of the quadratic cone in \mathbb{P}^3 branched over a sextic curve S. By Table 1, the RDP configuration on X is either A_6 or $A_6 + A_1$. By Corollary 7.2, X is the anti-canonical model of a blow-up Y_1 in a smooth point P of the surface X_2 of Case d = 2. By Proposition 5.8, the morphism $Y_1 \to X$ is Aut_X^0 -equivariant, since it is an isomorphism around the A_6 -singularity. Hence, by Proposition 4.2, we have $\operatorname{Aut}_X^0 = \operatorname{Aut}_{Y_1}^0 = \operatorname{Stab}_{\operatorname{Aut}_{X_2}^0}(P)^0 = \operatorname{Stab}_{\mu_7}(P)^0$. So, since μ_7 is simple, Y_1 is the blow-up of X_2 in a smooth fixed point P of the μ_7 -action on X_2 , and $\operatorname{Aut}_X^0 = \mu_7$.

Next, we prove the uniqueness of X. We may assume that X_2 is given by the equation in Table 4. The fixed points of the μ_7 -action are the three points [1:0:0:0], [0:1:0:0] and [0:0:1:0]. The automorphism $x \mapsto y \mapsto z \mapsto x$ of X_2 permutes these fixed points. So, Y_1 is unique and hence so is X.

Therefore, X is the unique RDP del Pezzo surface of degree 1 with an A_6 -singularity and non-zero global vector fields. Now, the equation

$$y^2 = x^3 + ts^3x + t^5s$$

defines such an RDP del Pezzo surface of degree 1 in $\mathbb{P}(1, 1, 2, 3)$ with an A_6 -singularity at [1 : -3 : 1 : 0] and, additionally, an A_1 -singularity at [1 : 0 : 0 : 0]. Clearly, the μ_7 -action described in Table 4 preserves X. Hence, this is the surface we were looking for.

Remark 8.4. We remark that by [MS20], in both cases of Theorem 8.3 the minimal resolution X of X does not admit any non-trivial global vector fields. In particular, $\pi : \tilde{X} \to X$ is not T_X -equivariant.

8.2. In characteristic 5. By Theorem 6.1, we have to list all RDP configurations containing A_4 or E_8^0 that can occur on an RDP del Pezzo surface in characteristic 5.

Lemma 8.5. If p = 5, deg(X) = d, and X contains a singularity of type A_4 or E_8^0 , then d and the configuration Γ of RDPs on X is one of the cases in Table 2.

Proof. Since A_4 has rank 4 and discriminant 5, it does not embed into E_{9-d} with $d \ge 6$, and the only root lattice containing A_4 and embedding into E_{9-d} with $d \in \{4, 5\}$ is A_4 itself. The other cases can be found in [Dyn52, Table 11].

Theorem 8.6. Assume that p = 5 and X contains an RDP of type A_4 or E_8^0 . Then, $H^0(X, T_X) \neq 0$ if and only if X is given by an equation as in Table 5. Moreover, Aut_X^0 is as in Table 5, so that $\operatorname{Aut}_{\widetilde{X}}^0 \subsetneq \operatorname{Aut}_X^0$ and even $h^0(X, T_X) > h^0(\widetilde{X}, T_{\widetilde{X}})$.

d	Г	$\subseteq \langle k_{9-d} \rangle^{\perp}$
5	A_4	$\subseteq A_4$
4	A_4	$\subseteq D_5$
3	$A_4, A_4 + A_1$	$\subseteq E_6$
2	$A_4, A_4 + A_1, A_4 + A_2$	$\subseteq E_7$
1	$A_4, A_4 + A_1, A_4 + 2A_1, A_4 + A_2, A_4 + A_2 + A_1, A_4 + A_3, 2A_4, E_8^0$	$\subseteq E_8$

Table 2. Non-equivariant RDP configurations in characteristic 5

Proof. By Table 2, we have $d \ge 5$.

If d = 5, then X is a quintic surface in \mathbb{P}^5 . By Table 2, the RDP configuration on X is A_4 . By the same argument as in characteristic 0 (going through the possible configurations of four (possibly infinitely) near points in \mathbb{P}^2), there is a unique quintic surface in \mathbb{P}^5 containing an A_4 -singularity. It is given by the equations in Table 5 (see [Der14, Section 3.3., p.657]) with singular point at [0:0:0:0:0:1]. The α_5 -action given in Table 5 preserves X and does not preserve the singular point, hence $\alpha_5 \cap \operatorname{Aut}^0_{\widetilde{X}} = \{\operatorname{id}\}$. By Proposition 5.8, this implies that $\operatorname{Aut}^0_X = \langle \alpha_5, \operatorname{Aut}^0_{\widetilde{Y}} \rangle$.

If d = 4, then X is a quartic surface in \mathbb{P}^4 . By Table 2, the RDP configuration on X is A_4 . By Corollary 7.2, X is the anti-canonical model of a blow-up Y_4 in a smooth point P of the surface X_5 of Case d = 5.

Such an X is in fact unique: By [MS20], we have dim $\operatorname{Aut}_{X_5}^0 = \dim \operatorname{Aut}_{\widetilde{X}_5}^0 = 4$, and since the orbit of P is at most 2-dimensional, the stabilizer of P is positive-dimensional, so $H^0(\widetilde{X}, T_{\widetilde{X}}) \neq 0$. By [MS20], there is a unique quartic del Pezzo surface with an A_4 -singularity and whose minimal resolution has global vector fields, hence X is unique.

In Table 5, we give equations for such a surface (which we took from [Der14]), hence this is our X. The singular point is located at [0:0:0:1]. The α_5 -action given in Table 5 preserves X and does not preserve the singular point. Again, by Proposition 5.8, this implies that $\operatorname{Aut}_X^0 = \langle \alpha_5, \operatorname{Aut}_{\widetilde{Y}}^0 \rangle$.

If d = 3, then X is a cubic surface in \mathbb{P}^3 . By Table 2, the RDP configuration on X is A_4 or $A_4 + A_1$. By Corollary 7.2, X is the anti-canonical model of a blow-up Y_3 in a smooth point P of the surface X_4 of Case d = 4.

Next, we show that there are at most two non-isomorphic such X. By [MS20], we have dim $\operatorname{Aut}_{X_4}^0 = \operatorname{dim} \operatorname{Aut}_{\widetilde{X}_4}^0 = 2$. There is at most one 2-dimensional orbit on X_4 , hence there is at most one X whose minimal resolution does not admit global vector fields. On the other hand, by [MS20], there is precisely one X whose minimal resolution does have global vector fields.

In Table 5, we give two (non-isomorphic) equations for cubic surfaces with an A_4 -singularity, distinguished by their RDP configuration Γ , hence these are the two possible X:

- If Γ = A₄, the singular point is located at [-2: -1: 2: 1]. We describe a μ₅-action on X in Table
 In this case, Aut⁰_{X̃} is trivial by [MS20], hence Aut⁰_X = μ₅ by Proposition 5.8.
- (2) If $\Gamma = A_4 + A_1$, the A_4 -singularity is [0:0:0:1] while the A_1 -singularity is [1:0:0:0]. We describe an $\alpha_5 \rtimes \mathbb{G}_m$ -action on X in Table 5. In this case, $\operatorname{Aut}^0_{\widetilde{X}} = \mathbb{G}_m$ by [MS20], hence $\operatorname{Aut}^0_X = \alpha_5 \rtimes \mathbb{G}_m$ by Proposition 5.8.

If d = 2, then X is a double cover of \mathbb{P}^2 branched over a quartic curve Q. By Table 2, the possible RDP configurations on X are A_4 , $A_4 + A_1$, and $A_4 + A_2$. By Corollary 7.2, X is the anti-canonical model of a blow-up Y_2 in a smooth point P of an RDP del Pezzo surface X_3 of degree 3 with an A_4 -singularity. Since $Y_2 \to X_3$ and $Y_2 \to X$ are isomorphisms around the A_4 -singularities, Proposition 5.8 yields $\operatorname{Aut}^0_{Y_2} = \operatorname{Stab}_{\operatorname{Aut}^0_{X_3}}(P)^0$. Hence, for each of the surfaces X_3 in Case d = 3, we have to determine the points with non-trivial stabilizer.

• If the RDP configuration on X_3 is A_4 , then the points with non-trivial stabilizer under the action of $\operatorname{Aut}_{X_3}^0 = \mu_5$ are [1:0:0:0], [0:1:0:0], [0:0:1:0], and [0:0:0:1] and these fixed points

are permuted by the automorphism $x_0 \mapsto x_1 \mapsto x_2 \mapsto x_3 \mapsto x_0$. Hence, there is a unique choice for P up to isomorphism.

- If the RDP configuration on X_3 is $A_4 + A_1$, then there are four lines on X_3 : The lines $\ell_1 = \{x_0 = x_1 = 0\}, \ell_2 = \{x_0 = x_2 = 0\}, \ell_3 = \{x_1 = x_2 = 0\}$ pass through the A_4 -singularity at [0:0:0:1] and the line $\ell_4 = \{x_2 = x_3 = 0\}$ passes through the A_1 -singularity at [1:0:0:0], but not through A_4 . Moreover, ℓ_2 and ℓ_4 intersect in [0:1:0:0]. Straightforward calculation shows that the points with non-trivial stabilizer in $X_3 \setminus \{\ell_1, \ell_2, \ell_3\}$ are precisely those lying on the hyperplane $H = \{x_3 = 0\}$. The intersection $X_3 \cap H$ is the union of the conic $C = \{x_3 = x_0x_2 + x_1^2 = 0\}$ and the line ℓ_4 . Hence, either $P \in C \setminus (\ell_1 \cup \ell_2 \cup \ell_3) = C \setminus \{[1:0:0:0], [0:0:1:0]\}$ or $P \in \ell_4 \setminus (\ell_1 \cup \ell_2 \cup \ell_3) = \ell_4 \setminus \{[1:0:0:0], [0:1:0:0]\}$. The group scheme $\operatorname{Aut}_{X_3}^0$ acts transitively on both loci, hence there are only two choices for P up to isomorphism. In both cases, one checks that $\operatorname{Aut}_{Y_2}^0 = \operatorname{Stab}_{\operatorname{Aut}_{X_3}}(P)^0 = \mu_5$. One of the two choices of P can be reduced to a previous case as follows:
 - If $P \in C \setminus \{[1:0:0:0], [0:0:1:0]\}$, then the RDP configuration on X is $A_4 + A_1$. The strict transform C' of C on the blow-up X' of X in the A_1 -singularity is a (-1)-curve that passes through the (-2)-curve over A_1 and which is disjoint from A_4 . If we contract C', we obtain an RDP del Pezzo surface X'_3 of degree 3 which contains an A_4 -singularity as its only singularity and such that $H^0(X'_3, T_{X'_3}) \neq 0$ (by Proposition 5.5 and Blanchard's lemma). Hence X'_3 is isomorphic to the cubic surface with RDP configuration A_4 in Table 2, and thus X coincides with the surface we constructed in the previous bullet point.

Summarizing, there are at most two RDP del Pezzo surfaces of degree 2 which contain an A_4 -singularity and which admit a non-trivial global vector field. Moreover, $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{Aut}_{X_3}^0}(P) = \mu_5$ in both cases. In Table 5, we give two equations of such surfaces, distinguished by their RDP configuration Γ :

- (1) If $\Gamma = A_4 + A_1$, the A_4 -singularity is [-2:1:2:0], the A_1 -singularity is [0:1:0:0], and the corresponding $\operatorname{Aut}_X^0 = \mu_5$ -action is as in Table 5
- (2) If $\Gamma = A_4 + A_2$, the A_4 -singularity is [2:1:-1:0], the A_2 -singularity is [1:0:0:0], and the corresponding $\operatorname{Aut}_X^0 = \mu_5$ -action is as in Table 5.

If d = 1, then X is a double cover of the quadratic cone in \mathbb{P}^3 branched over a sextic curve S. We consider separately the three cases where X contains a single A_4 -singularity (and possibly equivariant RDPs of other types), two A_4 -singularities, and an E_8^0 -singularity, respectively:

- (a) X contains a single A_4 -singularity. By Corollary 7.2, X is the anti-canonical model of a blow-up Y_1 in a smooth point P of an RDP del Pezzo surface X_2 with an A_4 -singularity. By Proposition 5.8, we have $\operatorname{Aut}_X^0 = \operatorname{Aut}_{Y_1}^0 = \operatorname{Stab}_{\operatorname{Aut}_{X_2}}^0(P)^0$. Since $\operatorname{Aut}_{X_2}^0 = \mu_5$, we thus have to determine the fixed points of the μ_5 -action on X_2 .
 - If the RDP configuration on X_2 is $A_4 + A_2$, then the fixed points of the μ_5 -action are [1:0:0:0], [0:1:0:0], and [0:0:1:0], and we recall that [1:0:0:0] is the A_2 -singularity. Hence, there are two choices for P. In fact, one of them does not occur, as the following argument shows:
 - If P = [0:0:1:0], let Q be the branch quartic of $X_2 \to \mathbb{P}^2$. The line $\ell = \{y = 0\}$ is tangent to Q at [1:0:0] and [0:0:1], hence the preimage of ℓ in X_2 consists of two smooth rational curves C, C' meeting in P and the A_2 -singularity. In the minimal resolution \widetilde{Y}_1 of Y_1 , their strict transforms \widetilde{C} and \widetilde{C}' are (-2)-curves, which, together with the two exceptional curves over the A_2 -singularity, form an A_4 -configuration of (-2)-curves. Thus, X contains two A_4 -singularities, contradicting our assumption.
 - If the RDP configuration on X₂ is A₄ + A₁, then the fixed points of the µ₅-action are [1 : 0 : 0 : 1], [1 : 0 : 0 : -1], [0 : 1 : 0 : 0], and [0 : 0 : 1 : 0]. The first two points are interchanged by w → -w and the point [0 : 1 : 0 : 0] is the A₁-singularity on X₂, hence we have two choices for P up to isomorphism. Let Q be the branch quartic of X₂ → P² and recall that the

 A_4 -singularity is located at [-2:1:2:0]. We will now show that both choices for P lead to the same surface as the one in the previous bullet point:

- If P = [1 : 0 : 0 : 1], then the image [1 : 0 : 0] of P in P² lies on the two bitangents ℓ₁ = {y = 0} and ℓ₂ = {z = 0} of Q. Let C_i, C'_i be the two irreducible components of the preimage of ℓ_i for i = 1, 2 and assume that P lies on C₁ and C₂. On Y₁, the strict transforms C̃₁ and C̃₂ are (-2)-curves, while the strict transforms C̃₁' and C̃₂' remain (-1)-curves. Thus, the RDP configuration on X is A₄ + A₂ + A₁, where the A₂ is obtained from the A₁ of Y₁ by also contracting C̃₂, and the A₁ arises from the contraction of C̃₁. Therefore, if we contract the image of C′₁ in the surface Y′₁ obtained from Y₁ by contracting C₂, we obtain an RDP del Pezzo surface of degree 2 with global vector fields and containing an RDP configuration of type A₄ + A₂, hence this case is reduced to the previous bullet point.
- If P = [0:0:1:0], then the image [0:0:1] of P in P² lies on the bitangent l = {y = 0} and P is the non-transversal intersection point of the two irreducible components C and C' of the preimage of l in X₂. Thus, the strict transforms C̃ and C̃' of C and C' on Ỹ₁ are (-2)-curves, hence the RDP configuration on X is A₄ + A₂ + A₁, where the A₂ is obtained by contracting C̃ and C̃'. The preimage D of the line {x = 0} in X₂ is a cuspidal cubic with cusp at [0:1:0:0]. On X̃₂, the strict transform of D is a smooth rational curve of self-intersection 0 by adjunction. Hence, the strict transform D̃ of D on Ỹ₁ is a (-1)-curve which passes through the exceptional curve over the A₁-singularity. Contracting the image of D̃ on the surface Y₁' obtained from Y₁ by contracting C and C' and resolving the A₁-singularity, we obtain an RDP del Pezzo surface of degree 2 with global vector fields containing an RDP configuration of type A₄ + A₂, hence this case is also reduced to the previous bullet point.

Summarizing, there is at most one RDP del Pezzo surface of degree 1 with global vector fields and containing a single A_4 -singularity. In Table 5, we give an equation of such a surface, hence this is our X. The singularities of X are as follows: A_4 at [1:-2:2:0], A_2 at [0:1:0:0], and A_1 at [1:0:0:0]. The μ_5 -action we describe in Table 5 preserves the equation, hence $\operatorname{Aut}_X^0 = \mu_5$.

- (b) X contains two A₄-singularities. By [Lan94, Theorem 4.1.], there is a unique RDP del Pezzo surface of degree 1 with RDP configuration A₄ + A₄, namely the Weierstrass model of the rational elliptic surface with singular fibers of type I₅, I₅, II. Its equation is given in Table 5. The α₅ × μ₅-action we describe in Table 5 preserves the equation. By [MS20], Aut⁰_{X̃} is trivial, hence Aut⁰_X = α₅ × μ₅ follows from Proposition 5.8.
- (c) X contains an E₈⁰-singularity. By [Lan94], there are two RDP del Pezzo surfaces of degree 1 with an RDP of type E₈. The one whose equation we give in Table 5 has an E₈⁰-singularity, while the other one has an E₈¹-singularity (see [Sta21, Table 1]). The α₅ ⋊ G_m-action we describe in Table 5 preserves the equation. We leave it to the reader to check that Aut⁰_X = Stab_{Aut_{P(1,1,2)}(S)⁰ = α₅ ⋊ G_m.}

8.3. In characteristic 3. By Theorem 6.1, we have to list all RDP configurations containing A_2, A_5, A_8, E_6^0 , E_6^1, E_7^0, E_8^0 or E_8^1 that can occur on an RDP del Pezzo surface.

Lemma 8.7. If p = 3, $\deg(X) = d$, and X contains a singularity of type $A_2, A_5, A_8, E_6^0, E_6^1, E_7^0, E_8^0$ or E_8^1 , then d and the configuration Γ of RDPs on X is one of the cases in Table 3.

Proof. The maximal root lattice contained in E_2 is isomorphic to A_1 , hence none of the root lattices in the statement of the lemma embed into E_{9-d} with $d \ge 7$. The list for $4 \le d \le 6$ follows from [Mar03, Exercise 4.2.1, 4.6.2] and the one for $7 \le d \le 9$ follows from [Dyn52, Table 11] (note that the lattice $A_6 + A_2$ in

d	Γ	$\subseteq \langle k_{9-d} \rangle^{\perp}$
6	$A_2, A_2 + A_1$	$\subseteq A_2 + A_1$
5	$A_2, A_2 + A_1$	$\subseteq A_4$
4	$A_2, A_2 + A_1, A_2 + 2A_1$	
3	$A_2, A_2 + A_1, A_2 + 2A_1, 2A_2, 2A_2 + A_1, A_5, 3A_2, A_5 + A_1, E_6^0, E_6^1,$	
2	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\subseteq E_7$
1	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	$\subseteq E_8$

Table 3. Non-equivariant RDP configurations in characteristic 3

Dynkin's table of root lattices in E_8 should be $E_6 + A_2$), where we marked those RDP configurations whose associated root lattice embeds in two non-conjugate ways into E_{9-d} by a prime '.

Theorem 8.8. Assume that p = 3 and X contains an RDP of type $A_2, A_5, A_8, E_6^0, E_6^1, E_7^0, E_8^0$, or E_8^1 . Then, $H^0(X, T_X) \neq 0$ if and only if X is given by an equation as in Tables 6, 7, 8 and 9. Moreover, Aut_X^0 is as given in these tables, so that $\operatorname{Aut}_{\widetilde{X}}^0 \subsetneq \operatorname{Aut}_X^0$ and even $h^0(X, T_X) > h^0(\widetilde{X}, T_{\widetilde{X}})$.

Proof. By Table 3, we have $d \leq 6$.

If d = 6, then X is a sextic surface in \mathbb{P}^6 . The RDP configuration Γ on X is either A_2 or $A_2 + A_1$ by Table 3. In both cases, X is uniquely determined by its singularities, by the same argument as in characteristic 0, that is, by checking the possible configurations of infinitely near points in \mathbb{P}^2 :

- (a) If $\Gamma = A_2$ then, by [Der14, Section 3.2.], X is given by the equations in Table 6. The A_2 -singularity is [1:0:0:0:0:0:0] and we give an α_3 -action on X which does not preserve the singularity in Table 6. By Proposition 5.8, this implies $\operatorname{Aut}_X^0 = \langle \alpha_3, \operatorname{Aut}_{\widetilde{X}}^0 \rangle$.
- (b) If Γ = A₂ + A₁, then, by [KN09, Appendix, p.3], X is given by the equation in Table 6. The A₂-singularity is [0 : 1 : 0 : 0 : 0 : 0 : 0] and the A₁-singularity is [0 : 0 : 0 : 0 : 0 : 1]. We give an α₃-action on X which does not fix the A₂-singularity in Table 6. By Proposition 5.8, this implies Aut⁰_X = ⟨α₃, Aut⁰_X⟩.

If d = 5, then X is a quintic surface in \mathbb{P}^5 . By Table 3, the RDP configuration Γ on X is either A_2 or $A_2 + A_1$. As in the Case d = 6, X is uniquely determined by Γ , as can be seen by checking the possible configurations of four infinitely near points in \mathbb{P}^2 (see e.g. [Dol12, Section 8.5.1, p.430]).

- (a) If $\Gamma = A_2$, then, by [Der14, Section 3.3.], X is given by the equation in Table 6. The A_2 -singularity is [0:0:1:0:0:0] and we give an α_3 -action on X which does not preserve the singularity in Table 6. By Proposition 5.8, this implies $\operatorname{Aut}_X^0 = \langle \alpha_3, \operatorname{Aut}_{\widetilde{X}}^0 \rangle$.
- (b) If $\Gamma = A_2 + A_1$, then by [KN09, Appendix, p.5] is given by the equation in Table 6. The A_2 -singularity is [0:1:0:0:0:0:0] and the A_1 -singularity is [0:0:0:0:0:1]. We give an α_3 -action on X which does not fix the A_2 -singularity in Table 6. By Proposition 5.8, this implies $\operatorname{Aut}_X^0 = \langle \alpha_3, \operatorname{Aut}_{\widetilde{X}}^0 \rangle$.

If d = 4, then X is a quartic surface in \mathbb{P}^4 . By Table 3, the RDP configuration on X is A_2 , $A_2 + A_1$, or $A_2 + 2A_1$. By Corollary 7.2, X is the anti-canonical model of a blow-up Y_4 in a smooth point P of an RDP del Pezzo surface X_5 of degree 5 with an A_2 -singularity.

Next, we show that there are at most three non-isomorphic such X. By [MS20], there are at most two X whose minimal resolution has global vector fields. Assume that $\operatorname{Aut}^0_{\widetilde{X}} = {\operatorname{id}}$. Then, by Proposition

4.2 and Proposition 5.8, the stabilizer of P is 0-dimensional, hence dim $\operatorname{Aut}_{\tilde{X}_5}^0 = \operatorname{dim} \operatorname{Aut}_{X_5}^0 \leq 2$. Hence, by [MS20], X_5 is the RDP del Pezzo surface of degree 5 whose RDP configuration is A_2 . For this surface, dim $\operatorname{Aut}_{X_5}^0 = 2$, hence there is at most one 2-dimensional orbit on X_5 , so there is at most one X whose minimal resolution does not have global vector fields. In Table 6, we give equations for three such surfaces, distinguished by their RDP configuration Γ :

- (1) If $\Gamma = A_2$, the A_2 -singularity is at [1:1:1:1]. We give a μ_3 -action on X in Table 6, while the group scheme $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial by [MS20]. By Proposition 5.8, this implies $\operatorname{Aut}^0_X = \mu_3$.
- (2) If $\Gamma = A_2 + A_1$, the A_2 -singularity is [1:0:0:0:0] while the A_1 -singularity is [0:0:0:0:1]. We give a $\alpha_3 \rtimes \mathbb{G}_m$ -action on X in Table 6, while $\operatorname{Aut}^0_{\widetilde{X}} = \mathbb{G}_m$ by [MS20], hence $\operatorname{Aut}^0_X = \alpha_3 \rtimes \mathbb{G}_m$ by Proposition 5.8.
- (3) If $\Gamma = A_2 + 2A_1$, the A_2 -singularity is [0:0:0:0:1] and the two A_1 -singularities are [0:1:0:0:0] and [0:0:1:0:0]. We give a $\alpha_3 \rtimes \mathbb{G}_m^2$ -action on X in Table 6, while $\operatorname{Aut}_{\widetilde{X}}^0 = \mathbb{G}_m^2$ by [MS20], hence $\operatorname{Aut}_{X}^0 = \alpha_3 \rtimes \mathbb{G}_m^2$ by Proposition 5.8.

If d = 3, then X is a cubic surface in \mathbb{P}^3 . By Table 3, X contains either a single A_2 , two A_2 s, three A_2 s, one A_5 , one E_6^0 , or one E_6^1 . In the following, we consider these six cases separately.

- (a) X contains a single A₂-singularity. By Corollary 7.2, X is the anti-canonical model of a blow-up Y₃ in a smooth point P of an RDP del Pezzo surface X₄ with an A₂-singularity. More precisely, since X --→ X₄ is an isomorphism around the only non-equivariant RDP on X and all other singularities of X are A₁-singularities by Table 3, Proposition 5.8 and Proposition 4.2 imply that Aut⁰_X = Aut⁰_{Y₃} = Stab_{Aut⁰_{X4}}(P)⁰. In particular, P is a point with non-trivial stabilizer on one of the surfaces X₄ in Table 6. Before we go on, note that by [MS20], the group scheme Aut⁰_{X̃} is trivial, hence the stabilizer of P is 0-dimensional.
 - Assume the RDP configuration on X_4 is $A_2 + 2A_1$. In this case, $\operatorname{Aut}_{X_4}^0$ is 2-dimensional, so there is at most one 2-dimensional orbit for this action, hence there is at most one choice for P up to isomorphism.
 - Assume the RDP configuration on X_4 is $A_2 + A_1$. The surface X_4 contains the six lines $\ell_1 = \{x_0 = x_2 = x_3 = 0\}, \ell_2 = \{x_0 = x_2 = x_4 = 0\}, \ell_3 = \{x_0 = x_3 = x_1 + x_4 = 0\}, \ell_4 = \{x_1 = x_2 = x_3 = 0\}, \ell_5 = \{x_1 = x_2 = x_4 = 0\}, \ell_6 = \{x_1 = x_3 = x_4 = 0\}$. The lines ℓ_4, ℓ_5 , and ℓ_6 pass through the A_2 -singularity at [1:0:0:0:0:1] and the others do not. The lines ℓ_1 and ℓ_4 pass through the A_1 -singularity at [0:0:0:0:1] and the others do not. Using the description of the Aut $_{X_4}^0$ -action in Table 6, one easily checks that the points with non-trivial and 0-dimensional stabilizer on X_4 are precisely those on ℓ_2 and those on ℓ_3 , except for [0:1:0:0:0], [0:0:1:0:0], [0:0:0:1:0], [0:0:1:0], and <math>[0:1:0:0:-1]. Since the automorphism $x_2 \leftrightarrow x_3, x_4 \leftrightarrow -x_4 x_1$ interchanges the two lines ℓ_2 and ℓ_3 and $\mathbb{G}_m \subseteq \text{Aut}_{X_4}$ acts transitively on the locus of points with 0-dimensional stabilizer on each of ℓ_2 and ℓ_3 , there is a unique choice for P up to isomorphism. We will now reduce this case to the previous bullet point.
 - Without loss of generality, assume that $P \in \ell_2 \setminus \{[0:1:0:0:0], [0:0:0:1:0]\}$. Then, the strict transform of ℓ_2 on Y_3 is a (-2)-curve and the RDP configuration on X is $A_2 + 2A_1$. Since ℓ_3 is disjoint from ℓ_2 , it remains a (-1)-curve on X, hence we can contract it to obtain a realization of X as a blow-up of an RDP del Pezzo surface X'_4 with RDPs of type $A_2 + 2A_1$. Hence, this case is reduced to the previous bullet point.
 - Assume the RDP configuration on X₄ is A₂. Using the description of the Aut⁰_{X₄} = μ₃-action given in Table 6, we see that the points P with non-trivial stabilizer on X₄ are [1 : 0 : 0 : 0 : 0 : 0], [0 : 1 : 0 : 0], [0 : 0 : 1 : 0], [0 : 0 : 0 : 1 : 0], and [0 : 0 : 0 : 0 : 1]. The surface X₄ admits the two involutions x₀ ↔ x₁ and (x₀, x₁) ↔ (x₂, x₃). Hence, blowing up any of the first four points leads to the same surface. In fact, we already treated the resulting surface, as the following argument shows:

- Assume without loss of generality that P = [1:0:0:0:0]. There are two lines on X_4 passing through P, namely $\ell_1 = \{x_1 = x_2 = x_4 = 0\}$ and $\ell_2 = \{x_1 = x_3 = x_4 = 0\}$, and both of these lines do not pass through the A_2 -singularity at [1:1:1:1:1]. Their strict transforms on Y_3 are disjoint (-2)-curves. Thus, the RDP configuration on X is $A_2 + 2A_1$. Now, the conic $C = \{x_1 = x_2 + x_3 = x_0x_4 - x_3^2 = 0\}$ meets ℓ_1 and ℓ_2 transversally at P and does not pass through [1:1:1:1:1], hence we can contract the image of the strict transform of C in X and obtain an RDP del Pezzo surface X'_4 with RDP configuration $A_2 + 2A_1$ and with global vector fields, hence X'_4 is the surface in the first bullet point.

Summarizing, there are at most two RDP del Pezzo surfaces of degree 3 with global vector fields and containing a single A_2 -singularity. Moreover, $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{Aut}_X^0}(P)^0 = \mu_3$. In Table 7, we give equations for two such surfaces, distinguished by their RDP configuration Γ :

- (1) If $\Gamma = A_2$, the A_2 -singularity is at [1:1:-1:-1]. Clearly, the μ_3 -action we give preserves the equation.
- (2) If $\Gamma = A_2 + 2A_1$, the A_2 -singularity is at [1:-1:-1:1] and the two A_1 -singularities are at [0:1:0:0] and [0:0:1:0]. Again, the μ_3 -action we give preserves the equation.
- (b) X contains two A_2 -singularities. By Table 3, the RDP configuration Γ on X is $2A_2$ or $2A_2 + A_1$. Simplifying the normal form of Roczen given in [Roc96], we obtain the equations given in Table 7. In particular, there is a 1-dimensional family of X with $\Gamma = 2A_2$ and a unique X with $\Gamma = 2A_2 + A_1$.
 - (1) If Γ = 2A₂, the two A₂-singularities are at [0 : 0 : 1 : 0] and [0 : 0 : 0 : 1]. The two α₃-actions and the Aut⁰_{X̄} = G_m-action we give in Table 7 preserve the equation. Each of the α₃-actions fixes one of the A₂-singularities and does not preserve the other one, hence Proposition 5.8 shows Aut⁰_{X̄} = ⟨α₃, α₃, G_m⟩.
 - (2) If $\Gamma = 2A_2 + A_1$, the two A_2 -singularities are at [0:0:1:0] and [0:0:0:1] and the A_1 singularity is at [0:1:0:0]. The two α_3 -actions and the $\operatorname{Aut}^0_{\widetilde{X}} = \mathbb{G}_m$ -action we describe in Table 7 preserve the equation. By the same argument as in (1), we have $\operatorname{Aut}^0_X = \langle \alpha_3, \alpha_3, \mathbb{G}_m \rangle$.
- (c) X contains three A_2 -singularities. Again, we can simplify the normal form of Roczen given in [Roc96] and obtain the equation in Table 7, which admits A_2 -singularities at [0:1:0:0], [0:0:1:0], [0:0:1:0], [0:0:0:1:0], and <math>[0:0:0:1]. In Table 7, we give an action of $\alpha_3^3 \rtimes \mathbb{G}_m^2$ on X. By [MS20], we have $\operatorname{Aut}_{\widetilde{X}}^0 = \mathbb{G}_m^2$. Each of the three factors of the α_3^3 -action preserves two of the A_2 -singularities and moves the other one, hence Proposition 5.8 yields $\operatorname{Aut}_X^0 = \alpha_3^3 \rtimes \mathbb{G}_m^2$.
- (d) X contains an A_5 -singularity. As above, we simplify Roczen's normal form [Roc96] to the two equations in Table 7. Let Γ be the RDP configuration on X:
 - (1) If $\Gamma = A_5$, then the A_5 -singularity is at [0:0:0:1]. The α_3 -action we describe preserves the equation and does not fix the A_5 -singularity. By [MS20], we have $\operatorname{Aut}_{\widetilde{X}}^0 = \mathbb{G}_a \rtimes \mu_3$ and we describe this action in terms of the equation in Table 7. By Remark 5.9, we have $\operatorname{Aut}_X^0 = \langle \alpha_3, \mathbb{G}_a \rtimes \mu_3 \rangle$.
 - (2) If Γ = A₅ + A₁, then the A₅-singularity is at [0 : 0 : 0 : 1] and the A₁-singularity is at [0 : 0 : 1 : 0]. The α₃-action we describe preserves the equation. By [MS20], we have Aut⁰_{X̃} = G_a ⋊ G_m and we describe this action in terms of the equation in Table 7. By Remark 5.9, we have Aut⁰_{X̃} = ⟨α₃, G_a ⋊ G_m⟩.
- (e) X contains an E_6^0 -singularity. By [Roc96, Case C3], X is given by the equation in Table 7 and the singularity is at [0:0:0:1]. In Table 7, we give an action of a group scheme G of order 27 such that no subgroup scheme of G lifts to \tilde{X} , as well as the action of $\operatorname{Aut}_{\tilde{X}}^0 = \mathbb{G}_a^2 \rtimes \mathbb{G}_m$ in terms of the equation. We leave it to the reader to check that these two actions generate $\operatorname{Aut}_X^0 = \operatorname{Stab}_{PGL_4}(X)^0$.
- (f) X contains an E¹₆-singularity. By [Roc96], X is given by the equation in Table 7 and the singularity is at [0 : 0 : 0 : 1]. In Table 7, we give a μ₃-action that does not preserve the singular point, as well as the action of Aut⁰_{X̃} = G²_a in terms of the equation. We leave it to the reader to check that these two actions generate Aut⁰_X = Stab_{PGL4}(X)⁰.

If d = 2, then X is a double cover of \mathbb{P}^2 branched over a quartic curve Q. In this case, we will take a slightly different approach which will turn out to be more economical than using Corollary 7.2. Namely, we classify all possible Q with global vector fields. If Q admits a global vector field, then it also admits an additive or multiplicative global vector field. This vector field is induced by a vector field on \mathbb{P}^2 . Up to conjugation, there are two non-zero vector fields D on \mathbb{P}^2 with $D^3 = D$ and two non-zero vector fields with $D^3 = 0$. They correspond to the following four matrices in Jordan normal form in the Lie algebra of PGL₃:

$$\begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

Integrating the corresponding vector fields (see e.g. [Tzi17, Proposition 3.1]), we obtain the following four μ_3 - and α_3 -actions on \mathbb{P}^2 :

(a) $\mu_3 : [x:y:z] \mapsto [x:\lambda y:\lambda^2 z]$ (b) $\mu_3 : [x:y:z] \mapsto [x:y:\lambda z]$ (c) $\alpha_3 : [x:y:z] \mapsto [x + \varepsilon y:y:z]$ (d) $\alpha_3 : [x:y:z] \mapsto [x + \varepsilon y - \varepsilon^2 z: y + \varepsilon z:z]$

For each of these actions, we will now classify the quartics that are invariant under it:

(a) There are three types of quartics which are invariant under this μ_3 -action, namely

$$ax^{4} + bx^{2}yz + cxy^{3} + dxz^{3} + ey^{2}z^{2}$$

$$ax^{3}y + bx^{2}z^{2} + cxy^{2}z + dy^{4} + eyz^{3}$$

$$ax^{3}z + bx^{2}y^{2} + cxyz^{2} + dy^{3}z + ez^{4}$$

These families are identified by permuting x, y and z, so it suffices to study the first one. Now, we simplify this equation and identify the singularities. We sort the classification according to the number of coefficients that are 0. Since Q is reduced, at least 2 coefficients are non-zero. Except in the case where c = d = 0, we can scale three of the non-zero coefficients to 1.

 If three of the coefficients are 0, the other two can be scaled to 1. The fact that Q is reduced leaves us with the following three cases, after using the symmetry y ↔ z:

(8.1)
$$x^4 + y^2 z^2$$

$$(8.2) x^2yz + xy^3$$

 $(8.3) x^2yz + y^2z^2$

In Case (8.1), X has two A_3 -singularities, one at [0:1:0:0] and one at [0:0:1:0]. In particular, X contains only equivariant RDPs, so it does not appear in Table 8.

In Case (8.2), X has a D_6 -singularity at [0:0:1:0] and an A_1 -singularity at [1:0:0:0]. As before, X contains only equivariant RDPs, so it does not appear in Table 8.

In Case (8.3), X has an A_1 -singularity at [1:0:0:0] and two A_3 -singularities at [0:1:0:0] and [0:0:1:0]. Again, X contains only equivariant RDPs, so it does not appear in Table 8.

 If two of the coefficients are 0, we get the following cases, again using scaling and the symmetry *y* ↔ *z*:

(8.4)
$$x^4 + ax^2yz + y^2z^2$$

$$(8.5) x^4 + x^2yz + xy^3$$

$$\begin{array}{c} (8.6) \\ x^4 + y^2 z^2 + x y^3 \\ z^2 + z^2 z^2 \end{array}$$

$$(8.7) x^2 yz + y^2 z^2 + x y^2$$

$$(8.8) x^2 yz + xy^3 + xz^3$$

 $(8.9) xy^3 + xz^3 + y^2z^2$

In Case (8.4), we must have $a^2 \neq 1$, otherwise Q is a double conic. Then, Q is the union of two conics, tangent at the points [0:1:0] and [0:0:1]. The singularities of X over these two

points are two A_3 -singularities, hence X contains only equivariant RDPs and does not appear in Table 8.

In Case (8.5), X has a D_6 -singularity at [0:0:1:0], hence X contains only equivariant RDPs and does not appear in Table 8.

In Case (8.6), X has an A₃-singularity at [0:0:1:0] and an A₂-singularity at [1:-1:0:0]. By [MS20], Aut⁰_{\tilde{X}} is trivial, hence Proposition 5.8 implies Aut⁰_X = μ_3 .

In Case (8.7), X has an A_3 -singularity at [0:0:1:0], an A_2 -singularity at [1:1:1:0], and an A_1 -singularity at [1:0:0:0]. By [MS20], $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial, hence Proposition 5.8 implies $\operatorname{Aut}^0_X = \mu_3$.

In Case (8.8), X has an A_5 -singularity at [0:1:-1:0] and an A_1 -singularity at [1:0:0:0]. By [MS20], $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial, hence Proposition 5.8 implies $\operatorname{Aut}^0_X = \mu_3$.

In Case (8.9), X has an E_6^1 -singularity at [1 : 0 : 0 : 0]. It is elementary to check that $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\mathrm{PGL}_3}(Q) = \mu_3$ in this case. Alternatively, observe that this case does not occur in the classification of invariant quartics for the actions (b), (c), and (d) below, so $\operatorname{Aut}_X^0[F] = \mu_3$, where $\operatorname{Aut}_X^0[F]$ is the Frobenius kernel of Aut_X^0 . In particular, μ_3 is normal in Aut_X^0 , so the Aut_X^0 -action preserves the eigenspaces of the μ_3 -action, hence the induced action on \mathbb{P}^2 is diagonal. This simplifies the calculation of the stabilizer of Q considerably.

If only one coefficient is 0, then we get the following cases, again using scaling and the symmetry y ↔ z:

$$(8.10) x^4 + ax^2yz + xy^3 + xz^3$$

$$(8.11) x^4 + a^3 x^2 yz + xy^3 + y^2 z^2$$

$$(8.12) x^4 + xy^3 + xz^3 + ay^2z^2$$

$$(8.13) ax^2yz + xy^3 + xz^3 + y^2z^2$$

In Case (8.10), X has an A_5 -singularity at [0:1:-1:0]. By [MS20], $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial, so $\operatorname{Aut}^0_X = \mu_3$ by Remark 5.9.

Consider Case (8.11). If $a^2 = 1$, then X has an A_6 -singularity at [0:0:1:0], hence X contains only equivariant RDPs and does not occur in Table 8. If $a^2 \neq 1$, then X has an A_3 -singularity at [0:0:1:0] and an A_2 -singularity at $[a^2 - 1:a^4 + a^2 + 1:a^3:0]$. In this case, $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial by [MS20], so $\operatorname{Aut}^0_X = \mu_3$ by Proposition 5.8.

In Case (8.12), X has two A_2 -singularities, one at [1:0:-1:0] and one at [1:-1:0:0]. We leave it to the reader to check that $Aut_X^0 = \mu_3$ in this case.

In Case (8.13), X has an A_1 -singularity at [1 : 0 : 0 : 0], and additional singularities at $[u : au^2 : 1]$, where u is a solution of $a^3u^6 - au^3 + 1 = 0$. If a = 1, then u = -1 is unique and the resulting singularity of X is of type A_5 . If $a \neq 1$, then X has two singularities of type A_2 . In both cases, $\operatorname{Aut}_{\widetilde{X}}^0$ is trivial by [MS20], so $\operatorname{Aut}_X^0 = \mu_3$ if a = 1 by Remark 5.9. If $a \neq 1$, one can check $\operatorname{Aut}_X^0 = \mu_3$ directly.

• If no coefficient is 0, we get the following case

$$x^4 + xy^3 + xz^3 + ax^2yz + by^2z^2$$

Here, X has singularities at the points $[bu : au^2 : b]$, where u is a solution of $a^3u^6 + (b^3 - a^2b^2)u^3 + b^3 = 0$. If $(b^3 - a^2b^2)^2 = a^3b^3$, then x is unique and the resulting singularity of X is of type A_5 . If $(b^3 - a^2b^2)^2 \neq a^3b^3$, then X has two singularities of type A_2 . In both cases, $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial by [MS20], so $\operatorname{Aut}^0_X = \mu_3$ by Proposition 5.8 and Remark 5.9.

(b) There are three types of quartics which are invariant under this μ_3 -action, namely

$$f_2(x,y)z^2 \ f_3(x,y)z + f_0z^4 \ f_4(x,y) + f_1(x,y)z^3$$

(8.14)

where the f_i are homogeneous polynomials of degree i in x and y. All quartics in the first family contain a double line, hence they lead to non-normal X. For the same reason, we have $f_3, f_4 \neq 0$ in the latter two families. In the third family, we have $f_1 \neq 0$, for otherwise Q has a quadruple point and the corresponding singularity on X is not an RDP. The GL₂-action on x, y normalizes the μ_3 -action, hence acts on the space of invariant quartics. Similarly, substitutions of the form $z \mapsto z + \beta x + \gamma y$ with $\beta, \gamma \in k$ act on the space of invariant quartics. Using these substitutions, and keeping in mind that Q must be reduced, we obtain the following simplified normal forms:

- $(8.17) x^2yz + z^4$
- $(8.19) y^4 + x^2y^2 + xz^3$
- $(8.20) x^2y^2 + xz^3$
- $(8.21) x^3y + xz^3$

In Case (8.15), X has a D_4 -singularity at [0:0:1:0] and three A_1 -singularities, at [1:0:0:0], [0:1:0:0], and [1:-1:0:0]. In particular, X contains only equivariant RDPs, so it does not occur in Table 8.

In Case (8.16), X has an A_2 -singularity at [1:1:1:0] and three A_1 -singularities, at [1:0:0:0], [0:1:0:0], and [1:-1:0:0]. By [MS20], $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial, hence $\operatorname{Aut}^0_X = \mu_3$ by Proposition 5.8.

In Case (8.17), X has a D_5 -singularity at [0:1:0:0] and an A_1 -singularity at [1:0:0:0]. In particular, X contains only equivariant RDPs, so it does not occur in Table 8.

In Case (8.18), X has a singularity of type E_6^0 at [1:0:0:0]. In Table 8, we give an action of a group scheme G of length 27 which preserves X and such that no subgroup scheme of G lifts to \tilde{X} . Moreover, we give an action of $\operatorname{Aut}_{\tilde{X}}^0 = \mathbb{G}_m$ on X. As in the corresponding case in degree 3, we leave it to the reader to check that these two actions generate Aut_X^0 .

In Case (8.19), X has three A_2 -singularities, at [1:0:0:0], [1:-1:1:0], and [1:1:1:0]. In Table 8, we describe an action of $\alpha_3^2 \rtimes \mu_3$ on X. By [MS20], $\operatorname{Aut}^0_{\widetilde{X}}$ is trivial, hence $\operatorname{Aut}^0_X = \alpha_3^2 \rtimes \mu_3$ by Proposition 5.8.

In Case (8.20), X has an A_5 -singularity at [0:1:0:0] and an A_2 -singularity at [1:0:0:0]. In Table 8, we give an action of $\alpha_3^2 \rtimes \mathbb{G}_m$ on X. By [MS20], we have $\operatorname{Aut}_{\widetilde{X}}^0 = \mathbb{G}_m$, so Proposition 5.8 and Remark 5.9 show that $\operatorname{Aut}_X^0 = \alpha_3^2 \rtimes \mathbb{G}_m$.

In Case (8.21), X admits a singularity of type E_7^0 at [0:1:0:0]. In Table 8, we give an action of α_3 on X that does not fix the E_7^0 -singularity as well as the action of $\operatorname{Aut}_{\widetilde{X}}^0 = \mathbb{G}_a \rtimes \mathbb{G}_m$. We leave it to the reader to check that these two actions generate Aut_X^0 .

(c) We can write the equation of Q as

$$f_0x^4 + x^3f_1(y,z) + x^2f_2(y,z) + xf_3(y,z) + f_4(y,z),$$

where the f_i are homogeneous of degree *i* in *y* and *z*. Applying the α_3 -action, we obtain

$$f_0 x^4 + \varepsilon f_0 x^3 y + x^3 f_1(y, z) + (x^2 - \varepsilon xy + \varepsilon^2 y^2) f_2(y, z) + (x + \varepsilon y) f_3(y, z) + f_4(y, z).$$

By considering the non-zero term of highest degree in x, we see that this is a multiple of the original equation if and only if it is equal to it. Comparing the coefficients of ε^2 and ε , we see that this happens if and only if $f_0 = f_2 = f_3 = 0$. Hence, Q is of the form

$$x^{3}f_{1}(y,z) + f_{4}(y,z).$$

But then Q is invariant under the μ_3 -action $[x : y : z] \mapsto [\lambda x : y : z]$ and hence, after interchanging x and z, Q is as in Cases (8.18), (8.19), (8.20), and (8.21).

(d) We write the equation of Q as $\sum a_{ijk}x^iy^jz^k$. As in Case (c), one checks that Q is preserved by the α_3 -action if and only if its equation is preserved by the α_3 -action. Applying the α_3 -action and comparing the coefficients of ε and ε^2 , respectively, we see that Q is α_3 -invariant if and only if the following two conditions are satisfied:

$$\begin{aligned} a_{400}x^3y + a_{310}x^3z - a_{220}x^2yz + a_{211}x^2z^2 - a_{220}xy^3 - a_{211}xy^2z - (a_{202} + a_{121})xyz^2 \\ &+ a_{112}xz^3 + a_{130}y^4 + (a_{121} + a_{040})y^3z + a_{112}y^2z^2 + (a_{103} - a_{022})yz^3 + a_{013}z^4 = 0 \\ &- a_{400}x^3z + a_{220}x^2z^2 - a_{220}xy^2z + (a_{121} + a_{202})xz^3 + a_{220}y^4 + (a_{211} - a_{130})y^3z \\ &+ (a_{121} + a_{202})y^2z^2 + (a_{022} - a_{103})z^4 = 0 \end{aligned}$$

The form $(a_{121} + a_{202})g \approx (a_{022} - a_{103}) \approx$ In other words, Q is α_3 -invariant if and only if it is given by an equation of the form

$$a(xz + y^2)^2 + bz^2(xz + y^2) + cx^3z + dy^3z + ez^4.$$

The substitutions of the form $x \mapsto \beta^2 x + \beta \gamma y + \delta z, y \mapsto \beta y + \gamma z$ with $\beta \in k^{\times}$ and $\gamma, \delta \in k$ normalize the α_3 -action, hence they preserve the space of α_3 -invariant quartics. If $a \neq 0$, we can scale it to a = 1 and use δ to kill b. If a = 0, then $b \neq 0$, otherwise Q contains a triple line. Then, we can assume b = 1. Using the other substitutions, we arrive at one of the following five simplified normal forms:

$$(8.22) (xz+y^2)^2 + z^4$$

$$(8.23) (xz+y^2)^2+y^3.$$

$$(8.24) (xz+y^2)^2 + x^3z + a^6z^4$$

(8.25)
$$z^2(xz+y^2) + y^3z$$

In Case (8.22), X has an A_7 -singularity at [1:0:0:0], hence X contains only equivariant RDPs and does not occur in Table 8.

In Case (8.23), X has an A_4 -singularity at [1:0:0:0] and an A_2 -singularity at [0:0:1:0]. By [MS20], Aut⁰_{\tilde{Y}} is trivial, hence Aut⁰_X = α_3 by Proposition 5.8.

In Case (8.24), X is singular precisely at $[-a^2 : \pm a : 1]$. If a = 0, then this singularity is of type A_5 and, by [MS20] and Remark 5.9, we have $\operatorname{Aut}_X^0 = \alpha_3$. If $a \neq 0$, then both singularities are of type A_2 . Direct calculation shows that $\operatorname{Aut}_X^0 = \alpha_3$.

In Case (8.25), X has an RDP of type E_7^1 at [1:0:0:0]. In particular, X contains only equivariant RDPs and does not occur in Table 8.

Finally, in Case (8.26), has an RDP of type A_5 at [0:1:0:0]. By [MS20], $\operatorname{Aut}_{\widetilde{X}}^0$ is trivial, hence, by Remark 5.9, we have $\operatorname{Aut}_X^0 = \alpha_3$. Note that this surface is not isomorphic to the one in Case (8.24) (with a = 0), since here, Q contains a line, while in the other case, Q is irreducible.

If d = 1, then X is a double cover of the quadratic cone in \mathbb{P}^3 branched over a sextic curve S. In particular, X is given by an equation in Weierstrass form

$$y^{2} = x^{3} + a_{2}(s,t)x^{2} + a_{4}(s,t)x + a_{6}(s,t),$$

where a_i is a homogeneous polynomial of degree i in s and t, and S is given by the equation

(8.27)
$$x^3 + a_2(s,t)x^2 + a_4(s,t)x + a_6(s,t) = 0$$

in $\mathbb{P}(1,1,2)$. By Proposition 4.3, we have $\operatorname{Aut}_X^0 = \operatorname{Stab}_{\operatorname{Aut}_{\mathbb{P}(1,1,2)}}(S)^0$. Since $\mathbb{P}(1,1,2)$ is an Aut_X -equivariant RDP del Pezzo surface of degree 8, Theorem 6.1 and [MS20] show that $\operatorname{Aut}_{\mathbb{P}(1,1,2)} = (\mathbb{G}_a^3 \rtimes \operatorname{GL}_2)/(\mathbb{Z}/2\mathbb{Z})$, which acts via substitutions of the form

$$\begin{array}{rccc} x & \mapsto & x + f_2(s,t) \\ s & \mapsto & \alpha s + \beta t \\ t & \mapsto & \gamma s + \delta t \end{array}$$

where f_2 is a homogeneous polynomial of degree 2 in s and t, and $\alpha, \beta, \gamma, \delta$ are scalars such that $\alpha \delta - \beta \gamma$ is invertible.

Now, assume Aut_X^0 is non-trivial. Then, it contains $G \in \{\alpha_3, \mu_3\}$. If $G = \mu_3$, we claim that there are only three embeddings of G into $\operatorname{Aut}^0_{\mathbb{P}(1,1,2)} = \mathbb{G}^3_a \rtimes \operatorname{GL}_2$. First, by counting the possible Jordan normal forms, observe that there are only three conjugacy classes of embeddings of G into GL_2 . Then, applying [Ray66, Théorème 5.1.1 (ii) (b)], we see that every such embedding lifts uniquely, up to conjugation by \mathbb{G}_{a}^{3} . to an embedding of μ_3 into $\mathbb{G}_a^3 \rtimes \mathrm{GL}_2$. Hence, if $G = \mu_3$, we may assume that it acts in one of the following three ways on $\mathbb{P}(1, 1, 2)$:

- (a) $\mu_3 : [s:t:x] \mapsto [s:\lambda t:x],$
- (b) $\mu_3 : [s:t:x] \mapsto [\lambda s:\lambda t:x],$ (c) $\mu_3 : [s:t:x] \mapsto [\lambda s:\lambda^{-1}t:x].$

Next, assume that $G = \alpha_3$ and the image of G in GL_2 is trivial. Then, the embedding of G into $\mathbb{G}_a^3 \rtimes GL_2$ is given by a homomorphism $\alpha_3 \to \mathbb{G}^3_a$, which in turn corresponds to a choice of homogeneous polynomial f_2 of degree 2 in s and t. According to whether this polynomial has one double zero or two simple zeroes, we can conjugate the embedding of α_3 using the GL₂-action to get one of the following two α_3 -actions:

(d)
$$\alpha_3 : [s:t:x] \mapsto [s:t:x+\varepsilon s^2],$$

(e)
$$\alpha_3 : [s:t:x] \mapsto [s:t:x + \varepsilon st].$$

Finally, assume that $G = \alpha_3$ and the image of G in GL₂ is non-trivial. After conjugating by elements of GL₂, we can assume that the image of G in GL₂ acts as $(s,t) \mapsto (s + \varepsilon t, t)$. An action of α_3 on $\mathbb{P}(1,1,2)$ that lifts this embedding of α_3 into GL₂ acts on x via

$$x \mapsto x + p(\varepsilon)s^2 + q(\varepsilon)st + r(\varepsilon)t^2$$

where p, q, r are polynomials of degree 2 in ε satisfying the following conditions:

$$p(0) = q(0) = r(0) = 0$$
$$p(\varepsilon + \varepsilon') = p(\varepsilon) + p(\varepsilon')$$
$$q(\varepsilon + \varepsilon') = q(\varepsilon) + q(\varepsilon') - p(\varepsilon)\varepsilon'$$
$$r(\varepsilon + \varepsilon') = r(\varepsilon) + r(\varepsilon') + q(\varepsilon)\varepsilon' + p(\varepsilon)\varepsilon'^{2}$$

Solving this system of equations, we obtain p = 0, $q(\varepsilon) = \alpha \varepsilon$ and $r(\varepsilon) = \beta \varepsilon - \alpha \varepsilon^2$ for scalars $\alpha, \beta \in k$. Conjugating with the substitution $x \mapsto x - \alpha s^2 + \beta st$, we obtain that our α_3 -action is conjugate to the following:

(f) $\alpha_3 : [s:t:x] \mapsto [s+\varepsilon t:t:x].$

We will now classify the sextics as in Equation (8.27) which are reduced with only simple singularities and invariant under one of the above actions. In particular, note that if all the a_i are scalar multiples of the *i*-th power of the same linear polynomial in s and t, then S has a non-simple singularity, so it will not appear in our list. Calculating Aut_X^0 is straightforward here, using our description of $\operatorname{Aut}_{\mathbb{P}(1,1,2)}$ above, so it will be left to the reader without further mention. The results can be found in Table 9:

(a) The sextic S is invariant if and only if the t-degree of every monomial that occurs in the equation of S is divisible by 3. Note that the substitutions

$$\begin{array}{rccc} x & \mapsto & x + \alpha s^2 \\ s & \mapsto & \beta s \\ t & \mapsto & \gamma t + \delta s \end{array}$$

act on the space of sextics satisfying the condition on the t-degree, hence we can apply them to arrive at the following normal forms for S:

(8.28)
$$x^3 + s^4 x + t^6$$

(8.29) $x^3 + s^4 x + s^3 t^3$

(8.30)
$$x^3 + st^3x$$

$$(8.31) x^3 + st^3x + as^3t^3 + t^6$$

$$(8.32) x^3 + st^3x + s^3t^3$$

$$(8.33) x^3 + s^2 x^2 + s^3 t^3$$

$$(8.34) x3 + s2x2 + a3s3t3 + t6$$

$$(8.35) x3 + s2x2 + st3x + a3s3t3 + b3t6$$

In Case (8.28), X has an E_6^0 -singularity at [0:1:-1:0]. In Case (8.29), X has an E_8^1 -singularity at [0:1:0:0].

In Case (8.30), X has an E_7^0 -singularity at [1:0:0:0] and an A_1 -singularity at [0:1:0:0].

In Case (8.31), if $a \neq 0$, then X has an E_6^0 -singularity at [1:0:0:0]. If a = 0, then X has an E_7^0 -singularity at [1:0:0:0].

In Case (8.32), X has an E_6^0 -singularity at [1:0:0:0] and an A_1 -singularity at [0:1:0:0].

In Case (8.33), X has an E_6^1 -singularity at [0:1:0:0] and an A_2 -singularity at [1:0:0:0].

In Case (8.34), X is singular at [1:0:0:0], [1:-a:0:0], and [0:1:-1:0]. If $a \neq 0$, then all singular points are A_2 -singularities. If a = 0, then the first two combine to an A_5 -singularity, while the latter stays an A_2 -singularity.

Consider Case (8.35). Here, X is singular at [1:0:0:0] and $[u:1:u^{-1}:0]$, where u is a solution of $au^2 + (b-1)u + 1 = 0$. If b = 0, then X has an additional A_1 -singularity at [0:1:0:0]. If the first three singular points are distinct, which happens if and only if $a \neq 0$, $(b-1)^2$, then they are A_2 -singularities. Now, consider the case where a = 0 or $a = (b-1)^2$, but not both: Then [1:0:0:0] is an A_5 -singularity and $[u:1:u^{-1}:0]$ an A_2 -singularity if $0 = a \neq (b-1)^2$, and the other way round if $0 \neq a = (b-1)^2$. Note that the substitution $t \mapsto t + (b-1)s$, $x \mapsto x + (b-1)^3s^2$ maps the family with $a = (b-1)^2$ to the one with a = 0, which is why only the latter occurs in Table 9. Finally, if a = 0 and b = 1, then X has an A_8 -singularity at [1:0:0:0].

(b) The sextic S is invariant if and only if the degrees of the a_i are divisible by 3. This happens if and only if $a_2 = a_4 = 0$. Using a substitution from $Aut_{\mathbb{P}(1,1,2)}$, we obtain the following normal forms:

(8.36)
$$x^3 + s^5 t$$

(8.37)
$$x^3 + s^4 t^2$$

$$(8.38) x^3 + s^4 t^2 + s^2 t^4$$

In Case (8.36), X has an E_8^0 -singularity at [0:1:0:0]. In Case (8.37), X has an E_6^0 -singularity at [0:1:0:0] and an A_2 -singularity at [1:0:0:0]. In Case (8.38), X has four A_2 -singularities, at [1:0:0:0], [0:1:0:0], [1:-1:0:0], and [1:1:0:0].

(c) The sextic S is invariant if and only if for every monomial in the equation of S, the difference between the s- and t-degree is divisible by 3. We may assume that not both a_2 and a_4 are zero, otherwise we get Cases (8.36), (8.37), and (8.38). Note that the substitutions

$$\begin{array}{rccc} x & \mapsto & x + \alpha st \\ s & \mapsto & \beta s \\ t & \mapsto & \gamma t \end{array}$$

normalize our μ_3 -action, hence we can apply them to arrive at the following normal forms for S:

(8.39)
$$x^3 + s^2 t^2 x$$

$$(8.40) x^3 + s^2 t^2 x + t^6 + a^3 s^6$$

(8.41)
$$x^3 + stx^2 + s^6$$

$$(8.42) x^3 + stx^2 + a^3s^6 + s^3t^3$$

$$(8.43) x3 + stx2 + a3s6 + b3s3t3 + t6$$

In Case (8.39), X has two D_4 -singularities, one at [1:0:0:0] and one at [0:1:0:0], hence X contains only equivariant RDPs and does not occur in Table 9.

Consider Case (8.40). If a = 0, then X has a D_4 -singularity at [1:0:0:0] and an A_2 -singularity at [0:1:-1:0]. If $a \neq 0$, then X has two A_2 -singularities, one at [1:0:-a:0] and one at [0:1:-1:0].

In Case (8.41), X has an RDP of type D_7 at [0:1:0:0]. In particular, X contains only equivariant RDPs and does not occur in Table 9.

Consider Case (8.42). If a = 0, then X has RDPs of type D_4 at [1:0:0:0] and [0:1:0:0], hence X contains only equivariant RDPs and does not occur in Table 9. If $a \neq 0$, then X has an RDP of type D_4 at [0:1:0:0] and an A_2 -singularity at [1:-a:0:0].

Consider Case (8.43). If a = 0, we can interchange s and t to reduce to one of the previous two cases. Here, X has singularities at [1 : u : 0 : 0], where u is a solution of $u^2 + bu + a = 0$. If $b^2 = a$, then the unique singularity of X is an A_5 -singularity. If $b^2 \neq a$, then X has two A_2 -singularities.

- (d) If $a_2 \neq 0$ or $a_4 \neq 0$, then S cannot be α_3 -invariant. Hence, $a_2 = a_4 = 0$. But then S is given by one of the equations in (b), so we are done.
- (e) By the same argument as in Case (d), we can reduce this to Case (b).
- (f) The sextic S is α_3 -invariant if and only if each a_i is invariant under the α_3 -action. This happens if and only if the s-degree of each monomial in the equation of S is divisible by 3. Interchanging the roles of s and t, we can therefore reduce this Case to Case (a). In fact, this explains why each of the surfaces in Case (a) admits an α_3 -action of this form.

GEBHARD MARTIN AND CLAUDIA STADLMAYR

Appendix: C	LASSIFICATION	TABLES
-------------	---------------	--------

d	singularities	equation of X	Aut_X^0
2	A_6	$w^2 = x^3y + y^3z + z^3x$	$\mu_7: [\lambda x: \lambda^4 y: \lambda^2 z: w]$
1	$A_6 + A_1$	$y^2 = x^3 + ts^3x + t^5s$	$\mu_7: [\lambda s: \lambda^4 t: x: y]$

Table 4. Non-equivariant RDP del Pezzo surfaces with vector fields in characteristic 7

d	RDPs	equation(s) of X	Aut_X^0
5	A_4	$\begin{array}{rcrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$\alpha_5: \begin{pmatrix} \alpha_5, \operatorname{Aut}_{\widetilde{X}}^0 \rangle \text{ with } \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & -2\varepsilon^2 & 2\varepsilon^3 & \varepsilon & 2\varepsilon^4 \\ 0 & 0 & 1 & 2\varepsilon & 0 & -\varepsilon^2 \\ 0 & 0 & 0 & 1 & 0 & -\varepsilon \\ 0 & 0 & \varepsilon & \varepsilon^2 & 1 & -2\varepsilon^3 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}$
4	A_4	$\begin{array}{rcl} x_0 x_1 - x_2 x_3 & = & 0 \\ x_0 x_4 + x_1 x_2 + x_3^2 & = & 0 \end{array}$	$\langle lpha_5, \operatorname{Aut}^0_{\widetilde{X}} angle ext{ with } \ lpha_5 : egin{pmatrix} 1 & -arepsilon^3 & -2arepsilon & 2arepsilon^2 &$
	A_4	$x_0^2 x_1 + x_1^2 x_2 + x_2^2 x_3 + x_3^2 x_0 = 0$	$\mu_5: [x_0: \lambda x_1: \lambda^4 x_2: \lambda^3 x_3]$
3	$A_4 + A_1$	$x_0 x_1 x_3 + x_0 x_2^2 + x_1^2 x_2 = 0$	$\alpha_{5} \rtimes \mathbb{G}_{m} \text{ with}$ $\alpha_{5} : \begin{pmatrix} 1 \varepsilon \varepsilon^{2} -2\varepsilon^{3} \\ 0 1 2\varepsilon -\varepsilon^{2} \\ 0 0 1 -\varepsilon \\ 0 0 0 1 \end{pmatrix}$ $\mathbb{G}_{m} : [x_{0} : \lambda x_{1} : \lambda^{2} x_{2} : \lambda^{3} x_{3}]$
2	$A_4 + A_1$	$w^2 = x^4 + xy^2z + yz^3$	$\mu_5: [x:\lambda y:\lambda^3 z:w]$
	$A_4 + A_2$	$w^2 = xy^3 + yz^3 + x^2z^2$	$\mu_5:[\lambda^*x:\lambda y:\lambda^*z:w]$
1	$A_4 + A_2 + A_1$	$\frac{y^2 = x^3 + s^3 tx + s^2 t^4}{2}$	$\mu_5:[s:\lambda t:\lambda^3 x:\lambda^2 y]$
1	$2A_4$	$y^2 = x^3 + t^4x + s^3t$	$\alpha_5 \rtimes \mu_5 : [\lambda s + \varepsilon t : t : x : y]$
	E_8°	$y^2 = x^3 + s^3 t$	$\alpha_5 \rtimes \mathbb{G}_m : [\lambda s + \varepsilon t : \lambda^{-s} t : x : y]$

Table 5. Non-equivariant RDP del Pezzo surfaces with vector fields in characteristic 5

d	RDPs	equation(s) of X	Aut_X^0
6	A_2	$\begin{array}{rcrcrcrcrcrcrcrcrcrcrcrcrcrcrcrcrcrcrc$	$\alpha_{3}, \operatorname{Aut}_{\widetilde{X}}^{0} \rangle \text{ with } \\ \left(\begin{array}{cccccccccccccccccccccccccccccccccccc$
	$A_2 + A_1$	$\begin{array}{rcrcrcrcrc} x_0^2 - x_1 x_5 &=& 0\\ x_0 x_2 - x_1 x_4 &=& 0\\ x_0 x_3 - x_2 x_4 &=& 0\\ x_0 x_4 - x_2 x_5 &=& 0\\ x_0 x_5 - x_2 x_6 &=& 0\\ x_1 x_3 - x_2^2 &=& 0\\ x_3 x_5 - x_4^2 &=& 0\\ x_3 x_6 - x_4 x_5 &=& 0\\ x_4 x_6 - x_5^2 &=& 0 \end{array}$	$egin{array}{lll} &\langle lpha_3, \operatorname{Aut}^0_{\widetilde{X}} angle ext{with} \ & 1 & -arepsilon & 0 & 0 & 0 & 0 \ 0 & 1 & 0 & 0 & 0 & 0 & 0 \ 0 & 0 & 1 & 0 & 0 & 0 & 0 \ 0 & 0 & 0 & 1 & 0 & 0 & 0 \ 0 & 0 & -arepsilon & 0 & 1 & 0 & 0 & 0 \ arepsilon & arepsilon^2 & arepsilon^2 & arepsilon & arepsilon^2 & ar$
5	A_2	$\begin{array}{rcrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrrr$	$egin{array}{lll} &\langle lpha_3, { m Aut}_{\widetilde{X}}^0 angle { m with} \ &1 &arepsilon & 0 & -arepsilon^2 & -arepsilon & -arepsilon^2 \ 0 & 1 & -arepsilon^2 & arepsilon & 0 & 0 \ 0 & 0 & 1 & 0 & 0 & 0 \ 0 & 0 & arepsilon $
	$A_2 + A_1$	$\begin{array}{rcrcrcrcrcrc} x_0^2 - x_1 x_4 &=& 0\\ x_0 x_2 - x_1 x_3 &=& 0\\ x_0 x_3 - x_2 x_4 &=& 0\\ x_0 x_4 - x_2 x_5 &=& 0\\ x_3 x_5 - x_4^2 &=& 0 \end{array}$	$egin{aligned} & & \langle lpha_3, \operatorname{Aut}^0_{\widetilde{X}} angle ext{ with } \ & & \left(egin{aligned} 1 & -arepsilon & 0 & 0 & 0 \ 0 & 1 & 0 & 0 & 0 \ 0 & 0 & 1 & 0 & 0 & 0 \ 0 & 0 & -arepsilon & 1 & 0 & 0 \ arepsilon & $
	A_2	$\begin{array}{rcl} x_0x_1 + x_2x_4 + x_3x_4 &=& 0\\ x_0x_4 + x_1x_4 + x_2x_3 &=& 0 \end{array}$	$\mu_3: [x_0: x_1: \lambda x_2: \lambda x_3: \lambda^2 x_4]$
4	$A_2 + A_1$	$ \begin{array}{rcl} x_0 x_1 - x_2 x_3 &=& 0\\ x_1 x_2 + x_2 x_4 + x_3 x_4 &=& 0\\ \end{array} $	$\alpha_{3} \rtimes \mathbb{G}_{m} \text{ with} \\ \alpha_{3} : \begin{pmatrix} 1 & 0 & 0 & 0 & 0 \\ -\varepsilon^{2} & 1 & \varepsilon & -\varepsilon & 0 \\ -\varepsilon & 0 & 1 & 0 & 0 \\ \varepsilon & 0 & 0 & 1 & 0 \\ -\varepsilon^{2} & 0 & -\varepsilon & 0 & 1 \end{pmatrix} \\ \mathbb{G}_{m} : [\lambda^{2}x_{0} : x_{1} : \lambda x_{2} : \lambda x_{3} : x_{4}] $
	$A_2 + 2A_1$	$\begin{array}{rcl} x_0^2 - x_3 x_4 &=& 0 \\ x_0 x_3 - x_1 x_2 &=& 0 \end{array}$	$ \begin{array}{c} \alpha_{3} \rtimes \mathbb{G}_{m}^{2} \text{ with} \\ \alpha_{3} : \begin{pmatrix} 1 & 0 & 0 & 0 & -\varepsilon \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ \varepsilon & 0 & 0 & 1 & \varepsilon^{2} \\ 0 & 0 & 0 & 0 & 1 \end{array} $ $ \mathbb{G}_{m}^{2} : [x_{0} : \lambda_{1}x_{1} : \lambda_{2}x_{2} : \lambda_{1}\lambda_{2}x_{3} : (\lambda_{1}\lambda_{2})^{-1}x_{4}] $

Table 6. Non-equivariant RDP del Pezzo surfaces of degree at least 4 with vector fields in characteristic 3

d	RDPs	equation(s) of X	Aut_X^0
	A_2	$x_0^2 x_1 + x_0 x_1^2 + x_2^2 x_3 + x_2 x_3^2 = 0$	$\mu_3: [x_0:x_1:\lambda x_2:\lambda x_3]$
	$A_2 + 2A_1$	$x_0^2 x_1 + x_0^2 x_2 + x_0 x_3^2 + x_1 x_2 x_3 = 0$	$\mu_3: [x_0:\lambda x_1:\lambda x_2:\lambda^2 x_3]$
			$\langle lpha_3, lpha_3, \mathbb{G}_m angle$ with
	240	$x_0^3 + x_1 x_2 x_3 + x_0 x_1^2 + a x_0^2 x_1 = 0$	$\alpha_3: [x_0 + \varepsilon x_2: x_1: x_2: a\varepsilon x_0 - \varepsilon x_1 - a\varepsilon^2 x_2 + x_3]$
	2/12	with $a^2 \neq 1$	$\alpha_3: [x_0 + \varepsilon x_3: x_1: a\varepsilon x_0 - \varepsilon x_1 + x_2 - a\varepsilon^2 x_3: x_3]$
			$\mathbb{G}_m: [x_0: x_1: \lambda x_2: \lambda^{-1} x_3]$
			$\langle lpha_3, lpha_3, \mathbb{G}_m angle$ with
	$9A_{2} \perp A_{1}$	$x^3 + x_1 x_2 x_3 + x^2 x_4 = 0$	$\alpha_3: [x_0 + \varepsilon x_2: x_1: x_2: \varepsilon x_0 - \varepsilon^2 x_2 + x_3]$
	$2A_2 + A_1$	$x_0 + x_1 x_2 x_3 + x_0 x_1 = 0$	$\alpha_3: [x_0 + \varepsilon x_3: x_1: a\varepsilon x_0 + x_2 - \varepsilon^2 x_3: x_3]$
			$\mathbb{G}_m: [x_0: x_1: \lambda x_2: \lambda^{-1} x_3]$
			$\alpha_3^3 \rtimes \mathbb{G}_m^2$ with
	$3A_2$	$x_0^3 + x_1 x_2 x_3 = 0$	$\alpha_3^3: [x_0+\varepsilon_1x_1+\varepsilon_2x_2+\varepsilon_3x_3:x_1:x_2:x_3]$
			$\mathbb{G}_m^2: [x_0:\lambda_1x_1:\lambda_2x_2:(\lambda_1\lambda_2)^{-1}x_3]$
		$x_0^3 + x_0 x_2 x_3 + x_1^2 x_2 + x_2^3 = 0$	$\langle lpha_3, \mathbb{G}_a times \mu_3 angle$ with
3	4 -		$\alpha_3: [x_0+\varepsilon x_1-\varepsilon^2 x_3:x_1+\varepsilon x_3:x_2:x_3]$
ა	A15		$\mathbb{G}_a: [x_0:\varepsilon x_0 + x_1: x_2: -\varepsilon^2 x_0 + \varepsilon x_1 + x_3]$
			$\mu_3: [x_0:\lambda x_1:\lambda x_2:\lambda^2 x_3]$
		$x_0^3 + x_0 x_2 x_3 + x_1^2 x_2 = 0$	$\langle lpha_3, \mathbb{G}_a times \mathbb{G}_m angle$ with
	$A_r + A_1$		$lpha_3: [x_0+arepsilon x_1-arepsilon^2 x_3: x_1+arepsilon x_3: x_2: x_3]$
	21 <u>9</u> + 211		$\mathbb{G}_a: [x_0:\varepsilon x_0+x_1:x_2:-\varepsilon^2 x_0+\varepsilon x_1+x_3]$
			$\mathbb{G}_m: [x_0:\lambda x_1:\lambda x_2:\lambda^2 x_3]$
			$\langle G, \mathbb{G}_a^2 times \mathbb{G}_m \rangle$ with
			$\mathbb{G}_a: [x_0 + \varepsilon x_2: x_1: x_2: -\varepsilon^3 x_2 + x_3]$
			$\mathbb{G}_a: [x_0: x_1+arepsilon x_2: x_2:arepsilon^3 x_1-arepsilon^2 x_2+x_3]$
	E_{6}^{0}	$x_0^3 + x_1^2 x_2 + x_2^2 x_3 = 0$	$\mathbb{G}_m: [x_0:\lambda x_1:x_2:\lambda^{-2}x_2:\lambda^4x_3]$
			and G non-commutative, $ G = 27$, acting as
			$[x_0 + \varepsilon_1 x_1 + \varepsilon_2 x_3 : x_1 + \varepsilon_1^3 x_3 : -\varepsilon_1^3 x_1 + x_2 + \varepsilon_1^6 x_3 : x_3]$
			where $\varepsilon_1^9 = \varepsilon_2^3 = 0$
			$\langle \mu_3, \mathbb{G}^2_a angle$ with
	$E_6^1 x_0^3 +$	$x_0^3 + x_1^3 + x_0x_1x_2 + x_2^2x_3 = 0$	$\mu_3:[\lambda x_0:\lambda^2 x_1:x_2:x_3]$
		$_{0}$ $u_{0} + u_{1} + u_{0}u_{1}u_{2} + u_{2}u_{3} = 0$	$\mathbb{G}_a: [x_0 - \varepsilon x_2: x_1: x_2: \varepsilon x_1 + \varepsilon^3 x_2 + x_3]$
			$\mathbb{G}_a: [x_0: x_1 - \varepsilon x_2: x_2: \varepsilon x_0 + \varepsilon^3 x_2 + x_3]$

Table 7. Non-equivariant RDP del Pezzo surfaces of degree 3 with vector fields in characteristic 3

d	RDPs	equation(s) of X	Aut_X^0
	$A_2 + 3A_1$	$w^2 = z(xy(x+y) + z^3)$	$\mu_3: [x:y:\lambda z:\lambda^{-1}w]$
	$A_2 + A_3$	$w^2 = x^4 + a^3 x^2 yz + xy^3 + y^2 z^2$ with $a^2 \neq 1$	$\mu_3: [x: \lambda y: \lambda^{-1}z: w]$
	$A_2 + A_3 + A_1$	$w^2 = x^2 yz + xy^3 + y^2 z^2$	$\mu_3: [x: \lambda y: \lambda^{-1}z: w]$
	$A_2 + A_4$	$w^2 = (xz + y^2)^2 + y^3 z$	$\alpha_3: [x + \varepsilon y - \varepsilon^2 z : y + \varepsilon z : z : w]$
	$2A_2$	$w^{2} = x^{4} + xy^{3} + xz^{3} + ax^{2}yz + by^{2}z^{2}$ with $(b^{3} - a^{2}b^{2})^{2} \neq a^{3}b^{3}, b \neq 0$	$\mu_3: [x:\lambda y:\lambda^{-1}z:w]$
	$2A_2$	$w^2 = (xz + y^2)^2 + x^3z + a^6z^4$ with $a \neq 0$	$\alpha_3: [x + \varepsilon y - \varepsilon^2 z : y + \varepsilon z : z : w]$
	$2A_2 + A_1$	$w^{2} = ax^{2}yz + xy^{3} + xz^{3} + y^{2}z^{2}$ with $a \neq 0, 1$	$\mu_3: [x: \lambda y: \lambda^{-1}z: w]$
	$3A_2$	$w^2 = y^4 + x^2 y^2 + xz^3$	$\alpha_3^2 \rtimes \mu_3 : [x:y:\varepsilon_1 x + \varepsilon_2 y + \lambda z:w]$
	A_5	$w^{2} = x^{4} + xy^{3} + xz^{3} + ax^{2}yz + by^{2}z^{2}$ with $(b^{3} - a^{2}b^{2})^{2} = a^{3}b^{3}, b \neq 0$	$\mu_3: [x:\lambda y:\lambda^{-1}z:w]$
2	A_5	$w^{2} = x^{4} + ax^{2}yz + xy^{3} + xz^{3}$ with $a \neq 0$	$\mu_3: [x:\lambda y:\lambda^{-1}z:w]$
	A_5	$w^2 = (xz + y^2)^2 + x^3 z$	$\alpha_3:[x+\varepsilon y-\varepsilon^2 z:y+\varepsilon z:z:w]$
	A_5	$w^2 = z(z(xz + y^2) + x^3)$	$\alpha_3: [x + \varepsilon y - \varepsilon^2 z : y + \varepsilon z : z : w]$
	$A_5 + A_1$	$w^2 = x^2yz + xy^3 + xz^3$	$\mu_3: [x:\lambda y:\lambda^{-1}z:w]$
	$A_5 + A_1$	$w^2 = x^2 yz + xy^3 + xz^3 + y^2 z^2$	$\mu_3: [x:\lambda y:\lambda^{-1}z:w]$
	$A_5 + A_2$	$w^2 = x^2 y^2 + x z^3$	$\alpha_3^2 \rtimes \mathbb{G}_m : [x : \lambda^3 y : \varepsilon_1 x + \varepsilon_2 y + \lambda^2 z : \lambda^3 w]$
			$\langle G, \mathbb{G}_m \rangle$ with
		$\overline{}$	$\mathbb{G}_m : [x : \lambda^3 y : \lambda^4 z : \lambda^5 w]$
	$E_{ m 6}$	$w^2 = y^2 + xz^3$	and G non-commutative, $ G = 27$, acting as
			$[x:y-\varepsilon_1^*x:\varepsilon_2x+\varepsilon_1y+z:w]$ where $c^9=c^3=0$
	E_{a}^{1}	$w^2 - (u^3 + z^3)x + u^2 z^2$	$\frac{1}{ \mu_2 \cdot [r \cdot \lambda \mu \cdot \lambda^{-1} \tau \cdot \mu]}$
		$\omega = (g + z) x + g z$	$\frac{\mu_3 \cdot [x \cdot \lambda g \cdot \lambda - z \cdot w]}{\langle \alpha_2 \mathbb{G}_{\pi_2} \times \mathbb{G}_{\pi_2} \rangle \text{ with }}$
	<u>_</u>		$\alpha_3: [x:y:z+\varepsilon y:w]$
	E_{7}^{0}	$E_7^0 \qquad \qquad w^2 = x^3 y + x z^3$	$\mathbb{G}_{a}: [x: y + \varepsilon^{3}x: z - \varepsilon x: w]$
			$\mathbb{G}_m:[x:\lambda^6y:\lambda^2z:\lambda^3w]$

Table 8. Non-equivariant RDP del Pezzo surfaces of degree 2 with vector fields in characteristic 3

d	RDPs	equation(s) of X	Aut_X^0
	A + D	$y^2 = x^3 + stx^2 + a^3s^6 + s^3t^3$	$u_{\alpha} \cdot [\lambda_{\varepsilon} \cdot \lambda^{-1} t \cdot x \cdot y]$
	$A_2 + D_4$	with $a \neq 0$	$\mu_3 \cdot [\lambda s \cdot \lambda \iota \cdot \iota \cdot g]$
	$A_2 + D_4$	$y^2 = x^3 + s^2 t^2 x + t^6$	$\mu_3: [\lambda s: \lambda^{-1}t: x: y]$
	$2A_{0}$	$y^2 = x^3 + stx^2 + a^3s^6 + b^3s^3t^3 + t^6$	$u_{2} \cdot [\lambda_{S} \cdot \lambda^{-1} t \cdot r \cdot y]$
	2112	with $a \neq 0, b^2 \neq a$	$\mu_3 \cdot [\Lambda 5 \cdot \Lambda - b \cdot a \cdot g]$
	$2A_2$	$y^2 = x^3 + s^2 t^2 x + a^3 s^6 + t^6$	$u_2 \cdot [\lambda_s \cdot \lambda^{-1}t \cdot x \cdot y]$
		with $a \neq 0$	μ3. [//θ. // δ. ω. 9]
	$3A_2$	$y^2 = x^3 + s^2 x^2 + st^3 x + a^3 s^3 t^3 + b^3 t^6$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
		with $a \notin \{0, (b-1)^2\}, b \neq 0$	
	$3A_2$	$y^2 = x^3 + s^2 x^2 + a^3 s^3 t^3 + t^6$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
		with $a \neq 0$	
	$3A_2 + A_1$	$y^2 = x^3 + s^2 x^2 + st^3 x + a^3 s^3 t^3$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
		with $a \notin \{0, 1\}$	
	$4A_2$	$y^2 = x^3 + s^4 t^2 + s^2 t^4$	$\alpha_3^3 \rtimes \mu_3 : [\lambda s : \lambda t : x + \varepsilon_1 s^2 + \varepsilon_2 s t + \varepsilon_3 t^2 : y]$
	A_5	$y^2 = x^3 + stx^2 + b^5s^5 + b^5s^5t^5 + t^6$	$\mu_3: [\lambda s: \lambda^{-1}t:x:y]$
		with $b \neq 0$	
	$A_{5} + A_{2}$	$y^2 = x^3 + s^2 x^2 + st^3 x + b^3 t^3$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
		with $b \neq 0, 1$	
1	$A_5 + A_2$	$y^2 = x^3 + s^2 x^2 + t^3$	$\alpha_3 \rtimes \mu_3 : [s: \varepsilon s + \lambda \iota : x: y]$
	$A_5 + A_2 + A_1$	$y^{-} = x^{-} + s^{-}x^{-} + st^{-}x$	$\alpha_3 \times \mu_3 : [s : \varepsilon s + \lambda \iota : x : y]$
	E_{6}^{0}	$y = x^2 + st^2x + ds^2t^2 + t^2$ with $a \neq 0$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
	E^0	$will a \neq 0$ $u^2 - x^3 + e^4 x + t^6$	$u_{s} \cdot [s \cdot \lambda t \cdot r \cdot y]$
	$E_6^0 + A_1$	$\frac{g - x + 5 x + t}{y^2 - r^3 + st^3r + s^3t^3}$	$\frac{\mu_3 \cdot [s \cdot \lambda t \cdot x \cdot y]}{(\lambda_2 \times \mu_2 \cdot [\lambda_2^6 s \cdot \varepsilon s + \lambda t \cdot x + (1 - \lambda^3) s^2 \cdot y]}$
	26 + 21	$y = x + \delta t + \delta t$	$\frac{\langle G \mathbb{G}_{m} \rangle}{\langle G \mathbb{G}_{m} \rangle}$ with
		$u^2 - r^3 + s^4 t^2$	$\mathbb{C}_{\mathrm{rm}} : [\lambda s \cdot \lambda^{-2}t \cdot x \cdot y]$
	$E_{2}^{0} + A_{2}$		and G non-commutative $ G = 81$ acting as
	-0 + -2	y = x + s t	$\mathbb{G}_{\text{rm}} : [s \cdot t - \varepsilon_{1}^{3}s \cdot x + \varepsilon_{1}s^{2} + \varepsilon_{2}st + \varepsilon_{2}t^{2} \cdot u]$
			with $\varepsilon_1^3 = \varepsilon_2^9 = \varepsilon_3^3 = 0$
	$E_{c}^{1} + A_{2}$	$u^2 = x^3 + s^2 x^2 + s^3 t^3$	$\frac{\alpha_3 \rtimes \mu_3 : [s:\varepsilon s + \lambda t:x:u]}{\alpha_3 \rtimes \mu_3 : [s:\varepsilon s + \lambda t:x:u]}$
	E_{7}^{0}	$\frac{y^2 = x^3 + st^3x + t^6}{y^2 = x^3 + st^3x + t^6}$	$\alpha_3 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x : y]$
	$E_7^0 + A_1$	$\frac{y^2 = x^3 + st^3x}{y^2 = x^3 + st^3x}$	$\frac{\alpha_3 \rtimes \mathbb{G}_m : [\lambda^{-3}s : \varepsilon s + \lambda t : x : y]}{\alpha_3 \rtimes \mathbb{G}_m : [\lambda^{-3}s : \varepsilon s + \lambda t : x : y]}$
	A_8	$y^2 = x^3 + s^2 x^2 + st^3 x + t^6$	$\frac{\alpha_0 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x + \varepsilon^3 s^2 : y]}{\alpha_0 \rtimes \mu_3 : [s : \varepsilon s + \lambda t : x + \varepsilon^3 s^2 : y]}$
		<i>y</i>	$\langle \alpha_2^2, \mathbb{G}_q \rtimes \mathbb{G}_m \rangle$ with
		0 2 5	$\mathbb{G}_a:[s:t-a^3s:x+as^2:y]$
	E_{8}^{0}	$E_8^0 y^2 = x^3 + s^5 t$	$\mathbb{G}_m:[\lambda s:\lambda^{-5}t:x:y]$
			$\alpha_3^2: [s:t:x+arepsilon_1 st+arepsilon_2 t^2:y]$
			$\mathbb{G}_a \rtimes \mu_3$ with
	E_{8}^{1}	$y^2 = x^3 + s^4 x + s^3 t^3$	$\mathbb{G}_a:[s:t-(a^3+a)s:x+as^2:y]$
	~		$\mu_3:[s:\lambda t:x:y]$

Table 9. Non-equivariant RDP del Pezzo surfaces of degree 1 with vector fields in characteristic 3

REFERENCES

- [BG81] J. W. Bruce and P. J. Giblin. A stratification of the space of plane quartic curves. Proc. London Math. Soc. (3), 42(2):270– 298, 1981.
- [Bri17] M. Brion. Some structure theorems for algebraic groups. Proc. Symp. Pure Math., 94:53–125, 2017.
- [Bri18] M. Brion. Notes on automorphism groups of projective varieties. Lecture Notes. Summer school and Workshop on Varieties and Group Actions, Warsaw, September 2018.
- [BT20] F. Bernasconi and H. Tanaka. On del Pezzo fibrations in positive characteristic. Journal of the Institute of Mathematics of Jussieu, page 1–43, 2020.
- [CP21] I. Cheltsov and Y. Prokhorov. Del Pezzo surfaces with infinite automorphism groups. Algebr. Geom., 8(3):319–357, 2021.
- [Der14] U. Derenthal. Singular del Pezzo surfaces whose universal torsors are hypersurfaces. Proc. Lond. Math. Soc. (3), 108(3):638-681, 2014.
- [Dol12] I. V. Dolgachev. Classical algebraic geometry. Cambridge University Press, Cambridge, 2012. A modern view.
- [Dyn52] E. B. Dynkin. Semisimple subalgebras of semisimple Lie algebras. Mat. Sbornik N.S., 30(72):349-462, 1952.
- [GK90] G.-M. Greuel and H. Kröning. Simple singularities in positive characteristic. Math. Z., 203(2):339–354, 1990.
- [Hir19] M. Hirokado. Further Evaluation of Wahl Vanishing Theorems for Surface Singularities in Characteristic *p. Michigan Mathematical Journal*, 68(3):621 – 636, 2019.
- [KN09] T. Kikuchi and T. Nakano. On the projective embeddings of Gorenstein toric del Pezzo surfaces. *Illinois J. Math.*, 53(4):1051–1059, 2009.
- [Lan94] W. E. Lang. Extremal rational elliptic surfaces in characteristic *p*. II. Surfaces with three or fewer singular fibres. *Ark. Mat.*, 32(2):423–448, 1994.
- [Mar03] J. Martinet. Perfect lattices in Euclidean spaces, volume 327 of Grundlehren der Mathematischen Wissenschaften. Fundamental Principles of Mathematical Sciences. Springer-Verlag, Berlin, 2003.
- [Mar20] G. Martin. Infinitesimal automorphisms of algebraic varieties and vector fields on elliptic surfaces, arxiv:2004.07227. *ArXiv e-prints*, 2020.
- [Mil16] J.S. Milne. Etale Cohomology (PMS-33). Princeton University Press, 2016.
- [MS20] G. Martin and C. Stadlmayr. Weak del Pezzo surfaces with global vector fields, arxiv:2007.03665. *ArXiv e-prints*, july 2020.
- [Ray66] M. Raynaud. Groupes algébriques unipotents. Extensions entre groupes unipotents et groupes de type multiplicatif. In Schémas en Groupes (Sém. Géométrie Algébrique, Inst. Hautes Études Sci., 1963/64), Fasc. 5b, Exposé XVII, page 96. Inst. Hautes Études Sci., Paris, 1966.
- [Roc96] M. Roczen. Cubic surfaces with double points in positive characteristic. In Algebraic geometry and singularities (La Rábida, 1991), volume 134 of Progr. Math., pages 375–382. Birkhäuser, Basel, 1996.
- [Ser06] E. Sernesi. Deformations of algebraic schemes, volume 334 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2006.
- [Sta21] C. Stadlmayr. Which rational double points occur on del Pezzo surfaces? Épijournal de Géométrie Algébrique, Volume 5(17), november 2021.
- [Stö04] K. Stöhr. On Bertini's theorem in characteristic p for families of canonical curves in $\mathbb{P}^{(p-3)/2}$. Proceedings of the London Mathematical Society, 89(2):291–316, 2004.
- [Tzi17] N. Tziolas. Quotients of schemes by α_p or μ_p actions in characteristic p > 0. Manuscripta Math., 152(1-2):247–279, 2017.
- [Wah75] J. M. Wahl. Vanishing theorems for resolutions of surface singularities. Invent. Math., 31(1):17-41, 1975.

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY *Email address*: gmartin@math.uni-bonn.de

TU MÜNCHEN, ZENTRUM MATHEMATIK - M11, BOLTZMANNSTRASSE 3, 85748 GARCHING BEI MÜNCHEN, GERMANY *Email address*: claudia.stadlmayr@ma.tum.de