

Granular gas of inelastic and rough Maxwell particles

Gilberto M. Kremer^{1†} and Andrés Santos^{2*†}

¹Departamento de Física, Universidade Federal do Paraná,
Curitiba, Brazil.

²Departamento de Física and Instituto de Computación
Científica Avanzada (ICCAEx), Universidad de Extremadura,
Badajoz, E-06006, Spain.

*Corresponding author(s). E-mail(s): andres@unex.es;

Contributing authors: kremer@fisica.ufpr.br;

†These authors contributed equally to this work.

Abstract

The most widely used model for granular gases is perhaps the inelastic hard-sphere model (IHSM), where the grains are assumed to be perfectly smooth spheres colliding with a constant coefficient of normal restitution. A much more tractable model is the inelastic Maxwell model (IMM), in which the velocity-dependent collision rate is replaced by an effective mean-field constant. This simplification has been taken advantage of by many researchers to find a number of exact results within the IMM. On the other hand, both the IHSM and IMM neglect the impact of roughness—generally present in real grains—on the dynamic properties of a granular gas. This is remedied by the inelastic rough hard-sphere model (IRHSM), where, apart from the coefficient of normal restitution, a constant coefficient of tangential restitution is introduced. In parallel to the simplification carried out when going from the IHSM to the IMM, we propose in this paper an inelastic rough Maxwell model (IRMM) as a simplification of the IRHSM. The tractability of the proposed model is illustrated by the exact evaluation of the collisional moments of first and second degree, and the most relevant ones of third and fourth degree. The results are applied to the evaluation of the rotational-to-translational temperature ratio and the velocity cumulants in the homogeneous cooling state.

Keywords: Granular gas, Inelastic collisions, Rough particles, Maxwell model

2 Inelastic and rough Maxwell particles

1 Introduction

Granular gases are typically modeled as a system of agitated inelastic hard spheres [1, 2]. In the basic inelastic hard-sphere model (IHSM) of granular gases, the particles are assumed to be smooth (i.e., with no rotational degrees of freedom) and the collisions are characterized by a constant coefficient of *normal* restitution, $0 < \alpha \leq 1$. If the number density of the gas is low enough, the most relevant physical quantity is the one-particle velocity distribution function (VDF), which obeys the Boltzmann equation appropriately adapted to incorporate the inelastic nature of collisions. Nevertheless, the fact that the collision rate in the IHSM is proportional to the relative velocity of the two colliding particles prevents the associated collisional moments from being expressible in terms of a finite number of velocity moments, thus hampering the possibility of deriving analytical results.

The above difficulty is also present for molecular gases (where collisions are elastic, i.e., $\alpha = 1$). In that case, it can be overcome by assuming that the gas is made of Maxwell molecules, that is, particles interacting via a repulsive force inversely proportional to the fifth power of distance [3–8]. This makes the collision rate independent of the relative velocity, so that the collisional moments become bilinear combinations of velocity moments of the same or lower degree. If collisions are inelastic ($\alpha < 1$), one can still construct the so-called inelastic Maxwell model (IMM) by assuming an effective mean-field collision rate independent of the velocity [9–43].

The original IHSM and the simpler IMM capture many of the most relevant features of granular gases [2, 44]. On the other hand, both models left out the roughness of real grains, which gives rise to frictional collisions and energy transfer between the translational and rotational degrees of freedom. A convenient way of modeling this roughness effect is by augmenting the IHSM by means of a constant coefficient of *tangential* restitution $-1 \leq \beta \leq 1$ [45–86]. The resulting granular-gas model can be referred to as the inelastic rough hard-sphere model (IRHSM). Needless to say, the IRHSM is even much more difficult to tackle with than the IHSM of frictionless, smooth particles. Therefore, in order to incorporate the influence of roughness and the associated rotational degrees of freedom on the dynamical properties and yet have a tractable model, it seems natural to construct an inelastic rough Maxwell model (IRMM), which would play a role with respect to the IRHSM similar to that played by the well-known IMM with respect to the IHSM (see Fig. 1). To the best of our knowledge, such an IRMM has not been proposed or worked out before.

The aim of this paper is to construct a tractable IRMM and illustrate its main features by evaluating exactly all the collisional velocity moments of first and second degree, as well as the most relevant ones of third and fourth degree. As expected, the results reduce to the known ones in the smooth limit (IMM) [32, 40]. We also display the results for the conservative case of elastic and perfectly rough particles ($\alpha = \beta = 1$), what constitutes the Maxwell analog of the Pidduck gas [87]. Moreover, we have applied the exact knowledge of the




Fig. 1 Sketch of four granular-gas models: IHSM, IMM, IRHSM, and IRMM.

isotropic second- and fourth-degree collisional moments to the homogeneous cooling state (HCS), as described by the model.

The organization of the paper is as follows. In Sec. 2, after summarizing the basic properties of the Boltzmann equation for a granular gas of inelastic and rough particles, and recalling the IRHSM, our IRMM is written down. The details of the collision rules are given in Appendix A. The core of the paper is presented in Sec. 3, where the exact structure of the collisional moments is displayed in Table 1, the explicit expressions of the coefficients in terms of inelasticity and roughness being moved to Appendix B, while some particularizations and consistency tests are shown in Appendix C. Next, Sec. 4 is devoted to the application of the results to the HCS. Finally, the paper is closed in Sec. 5 with some concluding remarks.

2 Kinetic theory of inelastic and rough particles

2.1 Boltzmann equation

We consider a granular gas made of inelastic and rough particles of diameter σ , mass m , and moment of inertia I . Let us denote by \mathbf{v} and $\boldsymbol{\omega}$ the translational and angular velocities, respectively, and introduce the short-hand notation $\{\mathbf{v}, \boldsymbol{\omega}\} \rightarrow \boldsymbol{\xi}$ and $d\mathbf{v}d\boldsymbol{\omega} \rightarrow d\boldsymbol{\xi}$. A direct encounter between two particles 1 and 2 is characterized by the precollisional velocities $(\boldsymbol{\xi}_1; \boldsymbol{\xi}_2)$, the postcollisional velocities $(\boldsymbol{\xi}'_1; \boldsymbol{\xi}'_2)$, and the collision unit vector $\hat{\boldsymbol{\sigma}}$. For a restitution encounter the pre- and postcollisional velocities are denoted by $(\boldsymbol{\xi}^*_1; \boldsymbol{\xi}^*_2)$ and $(\boldsymbol{\xi}_1; \boldsymbol{\xi}_2)$, respectively, and the collision vector by $\hat{\boldsymbol{\sigma}}^* = -\hat{\boldsymbol{\sigma}}$.

The collision rules are described in Appendix A in terms of the coefficients of normal restitution (α) and tangential restitution (β) defined by Eq. (A5). In particular, $\hat{\boldsymbol{\sigma}}^* \cdot \mathbf{g} = -\alpha(\hat{\boldsymbol{\sigma}}^* \cdot \mathbf{g}^*) = -\hat{\boldsymbol{\sigma}} \cdot \mathbf{g}$, where $\mathbf{g} = \mathbf{v}_1 - \mathbf{v}_2$ denotes the center-of-mass relative velocity. The absolute value of the Jacobian of the transformation $(\boldsymbol{\xi}^*_1; \boldsymbol{\xi}^*_2) \rightarrow (\boldsymbol{\xi}_1; \boldsymbol{\xi}_2)$ becomes

$$\left| \frac{\partial(\boldsymbol{\xi}_1^*; \boldsymbol{\xi}_2^*)}{\partial(\boldsymbol{\xi}_1; \boldsymbol{\xi}_2)} \right| = \frac{1}{\alpha\beta^2}. \quad (1)$$

4 *Inelastic and rough Maxwell particles*

As a consequence,

$$d\xi_1^* d\xi_2^* = \frac{1}{\alpha\beta^2} d\xi_1 d\xi_2. \quad (2)$$

In the absence of external forces or torques, the Boltzmann equation for granular gases is given by

$$\frac{\partial f}{\partial t} + \mathbf{v} \cdot \nabla f = J[\xi|f, f], \quad (3)$$

where $f(\mathbf{r}, \xi, t)$ is the one-particle VDF and $J[\xi|f, f]$ is the bilinear collision operator. Quite generally, it can be written as

$$J[\xi_1|f, f] = \sigma^2 \int d\xi_2 \int d\hat{\sigma} \left[\mathcal{F}(\hat{\sigma}^* \cdot \mathbf{g}^*) \frac{f_1^* f_2^*}{\alpha\beta^2} - \mathcal{F}(\hat{\sigma} \cdot \mathbf{g}) f_1 f_2 \right], \quad (4)$$

where use has been made of Eq. (2) and, as usual the notation $f_1 = f(\xi_1)$, $f_2 = f(\xi_2)$, $f_1^* = f(\xi_1^*)$, $f_2^* = f(\xi_2^*)$ has been employed. Moreover, the function $\mathcal{F}(x)$ is proportional to the collision rate, its precise form defining the chosen kinetic model.

The so-called weak form of the Boltzmann equation is obtained by multiplying both sides of Eq. (3) by an arbitrary trial function $\Psi(\mathbf{r}, \xi, t)$ and integrating over ξ . This yields

$$\partial_t (n\langle\Psi\rangle) + \nabla \cdot (n\langle\mathbf{v}\Psi\rangle) - n\langle(\partial_t + \mathbf{v} \cdot \nabla)\Psi\rangle = n\mathcal{J}[\Psi], \quad (5)$$

where

$$n(\mathbf{r}, t) = \int d\xi f(\mathbf{r}, \xi, t) \quad (6)$$

is the local number density,

$$\langle\Psi\rangle = \frac{1}{n(\mathbf{r}, t)} \int d\xi \Psi(\mathbf{r}, \xi, t) f(\mathbf{r}, \xi, t) \quad (7)$$

is the local average value of Ψ , and

$$\begin{aligned} \mathcal{J}[\Psi] &\equiv \frac{1}{n} \int d\xi \Psi(\mathbf{r}, \xi, t) J[\xi|f, f] \\ &= \frac{\sigma^2}{2n} \int d\xi_1 \int d\xi_2 \int d\hat{\sigma} \mathcal{F}(\hat{\sigma} \cdot \mathbf{g}) f_1 f_2 (\Psi'_1 + \Psi'_2 - \Psi_1 - \Psi_2) \end{aligned} \quad (8)$$

is the collisional production term of Ψ . In the second step of Eq. (8), we have used the relationship (2) and the standard symmetry properties of the collision term.

In particular, the flow velocity (\mathbf{u}), the mean angular velocity ($\boldsymbol{\Omega}$) and the granular temperatures (T_t , T_r , and T) correspond to $\Psi = \mathbf{v}$, $\Psi = \boldsymbol{\omega}$, $\Psi = V^2$,

$\Psi = \omega^2$, and $\Psi = mV^2 + I\omega^2$, respectively, where $\mathbf{V} = \mathbf{v} - \mathbf{u}$ denotes the peculiar velocity. Thus,

$$\mathbf{u} = \langle \mathbf{v} \rangle, \quad \boldsymbol{\Omega} = \langle \boldsymbol{\omega} \rangle, \quad T_t = \frac{m}{3} \langle V^2 \rangle, \quad T_r = \frac{I}{3} \langle \omega^2 \rangle, \quad T = \frac{1}{2} (T_t + T_r). \quad (9)$$

The associated collisional production terms can be written as

$$\mathcal{J}[\mathbf{v}] = 0, \quad \mathcal{J}[\boldsymbol{\omega}] = -\zeta_\Omega \boldsymbol{\Omega}, \quad \mathcal{J}[V^2] = -\zeta_t \frac{3T_t}{m}, \quad \mathcal{J}[\omega^2] = -\zeta_r \frac{3T_r}{I}, \quad (10a)$$

$$\mathcal{J}[mV^2 + I\omega^2] = -6\zeta T, \quad \zeta = \frac{T_t}{2T} \zeta_t + \frac{T_r}{2T} \zeta_r = \frac{\zeta_t + \theta \zeta_r}{1 + \theta}. \quad (10b)$$

Here, ζ_Ω is a “de-spinning” rate coefficient, ζ_t and ζ_r are energy production rates, ζ is the cooling rate, and $\theta \equiv T_r/T_t$ is the rotational-to-translational temperature ratio.

2.2 The inelastic rough hard-sphere model (IRHSM)

If the gas is modeled by the IRHSM, one has $\mathcal{F}(x) = \Theta(x)x$, where $\Theta(x)$ is the Heaviside step function. In that case, the collision operator becomes

$$J_{\text{HS}}[\boldsymbol{\xi}_1 | f, f] = \sigma^2 \int d\boldsymbol{\xi}_2 \int_+ d\hat{\boldsymbol{\sigma}} (\hat{\boldsymbol{\sigma}} \cdot \mathbf{g}) \left(\frac{f_1^* f_2^*}{\alpha^2 \beta^2} - f_1 f_2 \right), \quad (11)$$

where the subscript (+) in the integration over $\hat{\boldsymbol{\sigma}}$ denotes the constraint $\hat{\boldsymbol{\sigma}} \cdot \mathbf{g} > 0$ and we have taken into account that $\Theta(\hat{\boldsymbol{\sigma}}^* \cdot \mathbf{g}^*) (\hat{\boldsymbol{\sigma}}^* \cdot \mathbf{g}^*) = \alpha^{-1} \Theta(\hat{\boldsymbol{\sigma}} \cdot \mathbf{g}) (\hat{\boldsymbol{\sigma}} \cdot \mathbf{g})$. Analogously, Eq. (8) becomes

$$\mathcal{J}_{\text{HS}}[\Psi] = \frac{\sigma^2}{2n} \int d\boldsymbol{\xi}_1 \int d\boldsymbol{\xi}_2 \int_+ d\hat{\boldsymbol{\sigma}} (\hat{\boldsymbol{\sigma}} \cdot \mathbf{g}) f_1 f_2 (\Psi'_1 + \Psi'_2 - \Psi_1 - \Psi_2). \quad (12)$$

If $\Psi(\boldsymbol{\xi})$ is a polynomial and thus $\langle \Psi \rangle$ is a velocity moment, the collisional moment $\mathcal{J}_{\text{HS}}[\Psi]$ involves the full VDF or, equivalently, all the higher-degree moments. As a consequence, the infinite hierarchy of moments given by Eq. (5) cannot be solved, even in spatially uniform states, unless an approximate closure is applied. For instance, if f is approximated by a two-temperature Maxwellian, one finds [2, 75]

$$\zeta_\Omega^{\text{HS}} \simeq \frac{5\nu_{\text{HS}}}{3} \frac{\tilde{\beta}}{\kappa}, \quad \nu_{\text{HS}} \equiv \frac{16}{5} n \sigma^2 \sqrt{\pi T_t/m}, \quad (13a)$$

$$\zeta_t^{\text{HS}} \simeq \frac{5\nu_{\text{HS}}}{3} \left[\tilde{\alpha}(1 - \tilde{\alpha}) + \tilde{\beta}(1 - \tilde{\beta}) - \frac{\tilde{\beta}^2}{\kappa} \theta(1 + X) \right], \quad X \equiv \frac{I\Omega^2}{3T_r}, \quad (13b)$$

$$\zeta_r^{\text{HS}} \simeq \frac{5\nu_{\text{HS}}}{3} \frac{\tilde{\beta}}{\kappa} \left[\left(1 - \frac{\tilde{\beta}}{\kappa} \right) (1 + X) - \frac{\tilde{\beta}}{\theta} \right], \quad (13c)$$

6 *Inelastic and rough Maxwell particles*

$$\zeta^{\text{HS}} \simeq \frac{5}{12} \frac{\nu_{\text{HS}}}{1 + \theta} \left[1 - \alpha^2 + \frac{1 - \beta^2}{1 + \kappa} \theta \left(\frac{\kappa}{\theta} + 1 + X \right) \right]. \quad (13d)$$

Here, κ is the reduced moment of inertia [see Eq. (A4)] and $\tilde{\alpha}$, $\tilde{\beta}$ are auxiliary quantities [see Eq. (A6)].

2.3 The inelastic rough Maxwell model (IRMM)

The above problem is also present in the conventional case of elastic collisions. However, if the collision rate is assumed to be a constant (Maxwell model), the collisional moments involve moments of a degree equal to or lower than the degree of the moment $\langle \Psi \rangle$ [4–8].

We now apply the same philosophy to a granular gas of inelastic and rough particles and propose the IRMM by choosing $\mathcal{F}(x) = \text{const}$ in Eq. (4), i.e.,

$$J_{\text{M}}[\xi_1 | f, f] = \frac{\nu_{\text{M}}}{4\pi n} \int d\xi_2 \int d\hat{\sigma} \left(\frac{f_1^* f_2^*}{\alpha \beta^2} - f_1 f_2 \right), \quad (14)$$

where ν_{M} is an effective collision frequency. Analogously, Eq. (8) becomes

$$\mathcal{J}_{\text{M}}[\Psi] = \frac{\nu_{\text{M}}}{8\pi n^2} \int d\xi_1 \int d\xi_2 \int d\hat{\sigma} f_1 f_2 (\Psi'_1 + \Psi'_2 - \Psi_1 - \Psi_2). \quad (15)$$

3 Collisional moments

Using the collision rules (A1) and (A7) in Eq. (15), it is possible to evaluate the collisional moments in terms of velocity moments. After some tedious work, we obtained all the collisional moments of first and second degree, plus the most relevant ones of third and fourth degree. Their structure is shown in Table 1. We have followed the criterion of using the Greek letters χ , φ , and ψ for the coefficients in $\mathcal{J}_{\text{M}}[\Psi]$ associated with scalar, vector, and tensor quantities $\Psi(\xi)$, respectively. Moreover, an overline is used if $\Psi(\xi)$ contains the inner product $\mathbf{V} \cdot \boldsymbol{\omega}$. The exact explicit expressions for those coefficients in terms $\tilde{\alpha}$, $\tilde{\beta}$, and κ are given in Appendix B.

The particularization of the coefficients to the IMM ($\alpha < 1$, $\beta = -1$) and to the Pidduck model ($\alpha = \beta = 1$) [87] is presented in Appendix C, where some consistency tests are performed..

The basic production rates are those defined by Eqs. (10). According to Table 1,

$$\zeta_{\Omega}^{\text{M}} = \nu_{\text{M}} \varphi_{01|01} = \frac{4\nu_{\text{M}}}{3} \frac{\tilde{\beta}}{\kappa}, \quad (16a)$$

$$\zeta_t^{\text{M}} = \nu_{\text{M}} \left[\chi_{20|20} + \frac{4\theta}{\kappa} \chi_{20|02} (1 + X) \right]$$

Table 1 Collisional moments according to the IRMM, Eqs. (14) and (15).

$\Psi(\xi)$	$-\nu_{\text{M}}^{-1} \mathcal{J}_{\text{M}}[\Psi]$
\mathbf{V}	0
ω	$\varphi_{01 01}\langle\omega\rangle$
V^2	$\chi_{20 20}\langle V^2 \rangle + \chi_{20 02}\sigma^2 (\langle \omega^2 \rangle + \langle \omega \rangle^2)$
$V_i V_j - \frac{V^2}{3}\delta_{ij}$	$\psi_{20 20}(\langle V_i V_j \rangle - \frac{\langle V^2 \rangle}{3}\delta_{ij}) + \psi_{20 02}\sigma^2 (\langle \omega_i \omega_j \rangle + \langle \omega_i \rangle \langle \omega_j \rangle - \frac{\langle \omega^2 \rangle + \langle \omega \rangle^2}{3}\delta_{ij})$
ω^2	$\chi_{02 02} (\langle \omega^2 \rangle + \langle \omega \rangle^2) + \frac{\chi_{02 20}}{\sigma^2} \langle V^2 \rangle$
$\omega_i \omega_j - \frac{\omega^2}{3}\delta_{ij}$	$\psi_{02 02} (\langle \omega_i \omega_j \rangle + \langle \omega_i \rangle \langle \omega_j \rangle - \frac{\langle \omega^2 \rangle + \langle \omega \rangle^2}{3}\delta_{ij}) + \frac{\psi_{02 20}}{\sigma^2} (\langle V_i V_j \rangle - \frac{\langle V^2 \rangle}{3}\delta_{ij})$
$V_i \omega_j$	$\psi_{11 11}\langle V_i \omega_j \rangle$
$V^2 V_i$	$\varphi_{30 30}\langle V^2 V_i \rangle + \varphi_{30 12}\sigma^2 [2\langle \omega^2 V_i \rangle - \langle (\mathbf{V} \cdot \omega) \omega_i \rangle + 4\langle \omega \rangle \cdot \langle \omega V_i \rangle - \langle \omega \rangle \cdot \langle \mathbf{V} \omega_i \rangle - \langle \mathbf{V} \cdot \omega \rangle \langle \omega_i \rangle]$
$\omega^2 V_i$	$\frac{\varphi_{12 30}}{\sigma^2} \langle V^2 V_i \rangle + \varphi_{12 12}^{(1)} \langle \omega^2 V_i \rangle + \varphi_{12 12}^{(2)} \langle (\mathbf{V} \cdot \omega) \omega_i \rangle + \varphi_{12 12}^{(3)} \langle \omega \rangle \cdot \langle \omega V_i \rangle + \varphi_{12 12}^{(4)} (\langle \omega \rangle \cdot \langle \mathbf{V} \omega_i \rangle - \langle \mathbf{V} \cdot \omega \rangle \langle \omega_i \rangle)$
$(\mathbf{V} \cdot \omega) \omega_i$	$\overline{\varphi}_{12 12}^{(1)} \langle (\mathbf{V} \cdot \omega) \omega_i \rangle + \overline{\varphi}_{12 12}^{(2)} \langle \omega^2 V_i \rangle + \overline{\varphi}_{12 12}^{(3)} \langle \omega \rangle \cdot \langle \mathbf{V} \omega_i \rangle + \overline{\varphi}_{12 12}^{(4)} \langle \omega \rangle \cdot \langle \omega V_i \rangle + \overline{\varphi}_{12 12}^{(5)} \langle \mathbf{V} \cdot \omega \rangle \langle \omega_i \rangle$
V^4	$\chi_{40 40}^{(1)} \langle V^4 \rangle + \chi_{40 40}^{(2)} \langle V^2 \rangle^2 + \chi_{40 40}^{(3)} \langle \mathbf{VV} \rangle : \langle \mathbf{VV} \rangle + \chi_{40 04}\sigma^4 (\langle \omega^4 \rangle + \langle \omega^2 \rangle^2 + 2\langle \omega \omega \rangle : \langle \omega \omega \rangle + 4\langle \omega^2 \omega \rangle \cdot \langle \omega \rangle) + \chi_{40 22}^{(1)} \sigma^2 [2\langle V^2 \omega^2 \rangle + 2\langle V^2 \rangle \langle \omega^2 \rangle - \langle (\mathbf{V} \cdot \omega)^2 \rangle - \langle \mathbf{VV} \rangle : \langle \omega \omega \rangle - 2\langle (\mathbf{V} \cdot \omega) \mathbf{V} \rangle \cdot \langle \omega \rangle + 4\langle V^2 \omega \rangle \cdot \langle \omega \rangle] + \chi_{40 22}^{(2)} \sigma^2 (\langle \mathbf{V} \cdot \omega \rangle^2 + \langle \mathbf{V} \omega \rangle : \langle \mathbf{V} \omega \rangle - 4\langle \mathbf{V} \omega \rangle : \langle \omega \mathbf{V} \rangle)$
ω^4	$\frac{\chi_{04 40}}{\sigma^4} (\langle V^4 \rangle + \langle V^2 \rangle^2 + 2\langle \mathbf{VV} \rangle : \langle \mathbf{VV} \rangle) + \frac{\chi_{04 22}^{(1)}}{\sigma^2} [2\langle V^2 \omega^2 \rangle + 2\langle V^2 \rangle \langle \omega^2 \rangle - \langle (\mathbf{V} \cdot \omega)^2 \rangle - \langle \mathbf{VV} \rangle : \langle \omega \omega \rangle] + \frac{\chi_{04 22}^{(2)}}{\sigma^2} [2\langle (\mathbf{V} \cdot \omega) \mathbf{V} \rangle \cdot \langle \omega \rangle - 4\langle V^2 \omega \rangle \cdot \langle \omega \rangle - \langle \mathbf{V} \cdot \omega \rangle^2 - \langle \mathbf{V} \omega \rangle : \langle \mathbf{V} \omega \rangle + 4\langle \mathbf{V} \omega \rangle : \langle \omega \mathbf{V} \rangle] + \chi_{04 04}^{(1)} \langle \omega^4 \rangle + \chi_{04 04}^{(2)} \langle \omega^2 \rangle^2 + \chi_{04 04}^{(3)} \langle \omega \omega \rangle : \langle \omega \omega \rangle + \chi_{04 04}^{(4)} \langle \omega^2 \omega \rangle \cdot \langle \omega \rangle$
$V^2 \omega^2$	$\frac{\chi_{22 40}^{(1)}}{\sigma^2} (\langle V^4 \rangle + \langle V^2 \rangle^2) + \frac{\chi_{22 40}^{(2)}}{\sigma^2} \langle \mathbf{VV} \rangle : \langle \mathbf{VV} \rangle + \chi_{22 22}^{(1)} \langle V^2 \omega^2 \rangle + \chi_{22 22}^{(2)} \langle V^2 \rangle \langle \omega^2 \rangle + \chi_{22 22}^{(3)} \langle (\mathbf{V} \cdot \omega)^2 \rangle + \chi_{22 22}^{(4)} \langle \mathbf{VV} \rangle : \langle \omega \omega \rangle + \chi_{22 22}^{(5)} \langle (\mathbf{V} \cdot \omega) \mathbf{V} \rangle \cdot \langle \omega \rangle + \chi_{22 22}^{(6)} \langle V^2 \omega \rangle \cdot \langle \omega \rangle + \chi_{22 22}^{(7)} \langle \mathbf{V} \cdot \omega \rangle^2 + \chi_{22 22}^{(8)} \langle \mathbf{V} \omega \rangle : \langle \mathbf{V} \omega \rangle + \chi_{22 22}^{(9)} \langle \mathbf{V} \omega \rangle : \langle \omega \mathbf{V} \rangle + \chi_{22 22}^{(10)} \langle \omega \mathbf{V} \rangle : \langle \omega \omega \rangle + \chi_{22 04}^{(1)} \sigma^2 (\langle \omega^4 \rangle + \langle \omega^2 \rangle^2) + \chi_{22 04}^{(2)} \sigma^2 \langle \omega \omega \rangle : \langle \omega \omega \rangle + \chi_{22 04}^{(3)} \sigma^2 \langle \omega^2 \omega \rangle \cdot \langle \omega \rangle$
$(\mathbf{V} \cdot \omega)^2$	$\overline{\chi}_{22 40}^{(1)} (\langle V^2 \rangle^2 - \langle \mathbf{VV} \rangle : \langle \mathbf{VV} \rangle) + \overline{\chi}_{22 22}^{(1)} \langle V^2 \omega^2 \rangle + \overline{\chi}_{22 22}^{(2)} \langle V^2 \rangle \langle \omega^2 \rangle + \overline{\chi}_{22 22}^{(3)} \langle (\mathbf{V} \cdot \omega)^2 \rangle + \overline{\chi}_{22 22}^{(4)} \langle \mathbf{VV} \rangle : \langle \omega \omega \rangle + \overline{\chi}_{22 22}^{(5)} \langle (\mathbf{V} \cdot \omega) \mathbf{V} \rangle \cdot \langle \omega \rangle + \overline{\chi}_{22 22}^{(6)} \langle V^2 \omega \rangle \cdot \langle \omega \rangle + \overline{\chi}_{22 22}^{(7)} \langle \mathbf{V} \cdot \omega \rangle^2 + \overline{\chi}_{22 22}^{(8)} \langle \mathbf{V} \omega \rangle : \langle \mathbf{V} \omega \rangle + \overline{\chi}_{22 22}^{(9)} \langle \mathbf{V} \omega \rangle : \langle \omega \mathbf{V} \rangle + \overline{\chi}_{22 04}^{(1)} \sigma^2 (\langle \omega^2 \rangle^2 - \langle \omega \omega \rangle : \langle \omega \omega \rangle)$

8 *Inelastic and rough Maxwell particles***Table 2** Comparison between the basic production rates in the IRHSM and IRMM with two main choices of ν_M .

Choice for ν_M	$HS \simeq M$	α	β
$\frac{5}{4}\nu_{HS}$	$\zeta_\Omega^{HS} \simeq \zeta_\Omega^M$	Arbitrary	Arbitrary
	$\zeta_r^{HS} \simeq \zeta_r^M$	Arbitrary	Arbitrary
	$\zeta_t^{HS} \simeq \zeta_t^M$	1	Arbitrary
	$\zeta^{HS} \simeq \zeta^M$	1	Arbitrary
$\frac{5}{2}\nu_{HS}$	$\zeta_t^{HS} \simeq \zeta_t^M$	Arbitrary	-1
	$\zeta^{HS} \simeq \zeta^M$	Arbitrary	± 1

$$= \frac{2\nu_M}{3} \left[\tilde{\alpha}(1 - \tilde{\alpha}) + 2\tilde{\beta}(1 - \tilde{\beta}) - \frac{2\tilde{\beta}^2}{\kappa} \theta(1 + X) \right], \quad (16b)$$

$$\zeta_r^M = \nu_M \left[\frac{\kappa}{4\theta} \chi_{02|20} + \chi_{02|02} (1 + X) \right] = \frac{4\nu_M}{3} \frac{\tilde{\beta}}{\kappa} \left[\left(1 - \frac{\tilde{\beta}}{\kappa} \right) (1 + X) - \frac{\tilde{\beta}}{\theta} \right], \quad (16c)$$

$$\begin{aligned} \zeta^M &= \frac{\nu_M}{1 + \theta} \left[\chi_{20|20} + \frac{\kappa}{4} \chi_{02|20} + \left(\frac{4}{\kappa} \chi_{20|02} + \chi_{02|02} \right) \theta (1 + X) \right] \\ &= \frac{1}{6} \frac{\nu_M}{1 + \theta} \left[1 - \alpha^2 + 2 \frac{1 - \beta^2}{1 + \kappa} \theta \left(\frac{\kappa}{\theta} + 1 + X \right) \right], \end{aligned} \quad (16d)$$

where in the second equalities we have made use of Eqs. (B1), (B2), and (B4).

Comparison with the approximate results for the IRHSM, Eqs. (13) suggests the two main choices $\nu_M = \frac{5}{4}\nu_{HS}$ and $\nu_M = \frac{5}{2}\nu_{HS}$, as summarized in Table 2. The choices $\nu_M = \frac{5}{4}\nu_{HS}$ and $\nu_M = \frac{5}{2}\nu_{HS}$ are related to the rotational and translational degrees of freedom, respectively. As can be seen, there is no unique choice for ν_M allowing for an agreement between the IRHSM and IRMM basic production rates for any α and β . As a compromise, one may take the intermediate value $\nu_M \approx 2\nu_{HS}$.

It must be remarked that the results for $\Psi(\xi) = \omega$, $\Psi(\xi) = V^2$, and $\Psi(\xi) = \omega^2$ are also valid in the case of a hybrid model where $\mathcal{F} = \text{const} \times \Theta(x)$. However, this does not happen in the case of the other quantities in Table 1, in which case the associated collisional moments cannot be expressed only in terms of velocity moments of the same or lower degree if $\mathcal{F} = \text{const} \times \Theta(x)$.

4 Application to the Homogeneous Cooling State (HCS)

In the absence of gradients or any external driving, Eq. (5) becomes

$$\frac{d}{dt}\langle \Psi \rangle = \mathcal{J}[\Psi]. \quad (17)$$

For sufficiently long times, the system asymptotically reaches the HCS, which is characterized by a uniform, isotropic, and stationary *scaled* VDF [2, 75]

$$\phi(\mathbf{c}, \mathbf{w}) = \frac{\sqrt{4T_t T_r / mI}}{n} f(\mathbf{V}, \boldsymbol{\omega}; t), \quad \mathbf{c} \equiv \frac{\mathbf{V}}{\sqrt{2T_t(t)/m}}, \quad \mathbf{w} \equiv \frac{\boldsymbol{\omega}}{\sqrt{2T_r(t)/m}}. \quad (18)$$

In particular, $X \rightarrow 0$ and $\dot{\theta} \rightarrow 0$. The latter condition implies $\zeta_t - \zeta_r \rightarrow 0$, which yields the HCS temperature ratio

$$\theta = h + \sqrt{1 + h^2}, \quad (19a)$$

$$h \equiv \frac{\kappa}{8} \frac{\chi_{02|02} - \chi_{20|20}}{\chi_{20|02}} = \frac{1 + \kappa}{4\kappa(1 + \beta)} \left[(1 + \kappa) \frac{1 - \alpha^2}{1 + \beta} - 2(1 - \kappa)(1 - \beta) \right]. \quad (19b)$$




Fig. 2 Plot of the temperature ratios **a** $T_r/T_t = \theta$ and **b** $T_r/T = 2\theta/(1 + \theta)$ as functions of β for $\kappa = \frac{2}{5}$ and $\alpha = 0.4$ (—), 0.6 (---), 0.8 (—), and 1 (···).

Figure 2 shows the β -dependence of the temperature ratios T_r/T_t and T_r/T for uniform particles ($\kappa = \frac{2}{5}$) and several representative values of the coefficient of normal restitution. We observe that, except in the case of elastic collisions ($\alpha = 1$), $T_r/T_t \rightarrow \infty$ and, hence, $T_r/T \rightarrow 2$ in the quasi-smooth limit ($\beta \rightarrow -1$).

The scaled fourth-degree moments are $\langle c^4 \rangle = \frac{9}{4} \langle V^4 \rangle / \langle V^2 \rangle^2$, $\langle w^4 \rangle = \frac{9}{4} \langle \omega^4 \rangle / \langle \omega^2 \rangle^2$, $\langle c^2 w^2 \rangle = \frac{9}{4} \langle V^2 \omega^2 \rangle / \langle V^2 \rangle \langle \omega^2 \rangle$, and $\langle (\mathbf{c} \cdot \mathbf{w})^2 \rangle = \frac{9}{4} \langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle / \langle V^2 \rangle \langle \omega^2 \rangle$. Assuming the stationary VDF $\phi(\mathbf{c}, \mathbf{w})$ has not been reached yet, the evolution equations for those moments are

$$\begin{aligned} \frac{d\langle c^4 \rangle}{\nu_M dt} &= \frac{1}{\langle V^2 \rangle^2} \frac{\mathcal{J}_M[V^4]}{\nu_M} - 2 \frac{\langle V^4 \rangle}{\langle V^2 \rangle^3} \frac{\mathcal{J}_M[V^2]}{\nu_M} \\ &= -\chi_{40|40}^{(1)} \frac{\langle V^4 \rangle}{\langle V^2 \rangle^2} - \chi_{40|40}^{(2)} - \frac{1}{3} \chi_{40|40}^{(3)} - \frac{16\theta^2}{\kappa^2} \chi_{40|04} \left(\frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^2} + \frac{5}{3} \right) - \frac{4\theta}{\kappa} \chi_{40|22}^{(1)} \\ &\quad \times \left[2 \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} + \frac{5}{3} \right] + 2 \left(\chi_{20|20} + \frac{4\theta}{\kappa} \chi_{20|02} \right) \frac{\langle V^4 \rangle}{\langle V^2 \rangle^2}, \end{aligned} \quad (20a)$$

$$\begin{aligned} \frac{d\langle w^4 \rangle}{\nu_M dt} &= \frac{1}{\langle \omega^2 \rangle^2} \frac{\mathcal{J}_M[\omega^4]}{\nu_M} - 2 \frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^3} \frac{\mathcal{J}_M[\omega^2]}{\nu_M} \\ &= -\frac{\kappa^2}{16\theta^2} \chi_{04|40} \left(\frac{\langle V^4 \rangle}{\langle V^2 \rangle^2} + \frac{5}{3} \right) - \frac{\kappa}{4\theta} \chi_{04|22}^{(1)} \left[2 \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} + \frac{5}{3} \right] \\ &\quad - \chi_{04|04}^{(1)} \frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^2} - \chi_{04|04}^{(2)} - \frac{1}{3} \chi_{04|04}^{(3)} + 2 \left(\chi_{02|02} + \frac{\kappa}{4\theta} \chi_{02|20} \right) \frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^2}, \end{aligned} \quad (20b)$$

$$\begin{aligned} \frac{d\langle c^2 w^2 \rangle}{\nu_M dt} &= \frac{1}{\langle V^2 \rangle \langle \omega^2 \rangle} \frac{\mathcal{J}_M[V^2 \omega^2]}{\nu_M} - \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle^2 \langle \omega^2 \rangle} \frac{\mathcal{J}_M[V^2]}{\nu_M} - \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle^2} \frac{\mathcal{J}_M[\omega^2]}{\nu_M} \\ &= -\frac{\kappa}{4\theta} \chi_{22|40}^{(1)} \left(\frac{\langle V^4 \rangle}{\langle V^2 \rangle^2} + 1 \right) - \frac{\kappa}{12\theta} \chi_{22|40}^{(2)} - \chi_{22|22}^{(1)} \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \chi_{22|22}^{(2)} \\ &\quad - \chi_{22|22}^{(3)} \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \frac{1}{3} \chi_{22|22}^{(4)} - \frac{4\theta}{\kappa} \chi_{22|04}^{(1)} \left(\frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^2} + 1 \right) - \frac{4\theta}{3\kappa} \chi_{22|04}^{(2)} \\ &\quad + \left(\chi_{20|20} + \frac{4\theta}{\kappa} \chi_{20|02} + \chi_{02|02} + \frac{\kappa}{4\theta} \chi_{02|20} \right) \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle}, \end{aligned} \quad (20c)$$

$$\begin{aligned} \frac{d\langle (\mathbf{c} \cdot \mathbf{w})^2 \rangle}{\nu_M dt} &= \frac{1}{\langle V^2 \rangle \langle \omega^2 \rangle} \frac{\mathcal{J}_M[(\mathbf{V} \cdot \boldsymbol{\omega})^2]}{\nu_M} - \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle^2 \langle \omega^2 \rangle} \frac{\mathcal{J}_M[V^2]}{\nu_M} - \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle^2} \frac{\mathcal{J}_M[\omega^2]}{\nu_M} \\ &= -\frac{\kappa}{6\theta} \bar{\chi}_{22|40} - \bar{\chi}_{22|22}^{(1)} \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \bar{\chi}_{22|22}^{(2)} - \bar{\chi}_{22|22}^{(3)} \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - \frac{1}{3} \bar{\chi}_{22|22}^{(4)} \\ &\quad - \frac{8\theta}{3\kappa} \bar{\chi}_{22|04} + \left(\chi_{20|20} + \frac{4\theta}{\kappa} \chi_{20|02} + \chi_{02|02} + \frac{\kappa}{4\theta} \chi_{02|20} \right) \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle}. \end{aligned} \quad (20d)$$

The departure of the HCS scaled VDF $\phi(\mathbf{c}, \mathbf{w})$ from the Maxwellian $\pi^{-3}e^{-c^2-w^2}$ can be measured by the four cumulants [73, 74, 88]

$$a_{20}^{(0)} \equiv \frac{3}{5} \frac{\langle V^4 \rangle}{\langle V^2 \rangle^2} - 1 = \frac{4}{15} \langle c^4 \rangle - 1, \quad a_{02}^{(0)} \equiv \frac{3}{5} \frac{\langle \omega^4 \rangle}{\langle \omega^2 \rangle^2} - 1 = \frac{4}{15} \langle w^4 \rangle - 1, \quad (21a)$$

$$a_{11}^{(0)} \equiv \frac{\langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} - 1 = \frac{4}{9} \langle c^2 w^2 \rangle - 1, \quad (21b)$$

$$a_{00}^{(1)} \equiv \frac{6}{5} \frac{\langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle - \frac{1}{3} \langle V^2 \omega^2 \rangle}{\langle V^2 \rangle \langle \omega^2 \rangle} = \frac{8}{15} \left[\langle (\mathbf{c} \cdot \mathbf{w})^2 \rangle - \frac{1}{3} \langle c^2 w^2 \rangle \right]. \quad (21c)$$

In terms of those cumulants, Eqs. (20) can be recast as

$$\nu_M^{-1} \frac{d}{dt} \begin{bmatrix} a_{20}^{(0)} \\ a_{02}^{(0)} \\ a_{11}^{(0)} \\ a_{00}^{(1)} \end{bmatrix} + \mathbf{M} \cdot \begin{bmatrix} a_{20}^{(0)} \\ a_{02}^{(0)} \\ a_{11}^{(0)} \\ a_{00}^{(1)} \end{bmatrix} = -\mathbf{L}, \quad (22)$$

where the elements of the matrices \mathbf{M} and \mathbf{L} are

$$M_{11} = \chi_{40|40}^{(1)} - 2\chi_{20|20} - \frac{8\theta}{\kappa} \chi_{20|02}, \quad (23a)$$

$$M_{12} = \frac{16\theta^2}{\kappa^2} \chi_{40|04}, \quad M_{13} = \frac{4\theta}{\kappa} \chi_{40|22}^{(1)}, \quad M_{14} = -\frac{2\theta}{\kappa} \chi_{40|22}^{(1)}, \quad (23b)$$

$$M_{21} = \frac{\kappa^2}{16\theta^2} \chi_{04|40}, \quad M_{22} = \chi_{04|04}^{(1)} - 2\chi_{02|02} - \frac{\kappa}{2\theta} \chi_{02|20}, \quad (23c)$$

$$M_{23} = \frac{\kappa}{4\theta} \chi_{04|22}^{(1)}, \quad M_{24} = -\frac{\kappa}{8\theta} \chi_{04|22}^{(1)}, \quad (23d)$$

$$M_{31} = \frac{\kappa}{4\theta} \chi_{22|40}^{(1)}, \quad M_{32} = \frac{4\theta}{\kappa} \chi_{22|04}^{(1)}, \quad M_{34} = \frac{1}{2} \chi_{22|22}^{(3)}, \quad (23e)$$

$$M_{33} = \frac{3}{5} \left[\chi_{22|22}^{(1)} - \chi_{20|20} - \frac{4\theta}{\kappa} \chi_{20|02} - \chi_{02|02} - \frac{\kappa}{4\theta} \chi_{02|20} \right] + \frac{1}{5} \chi_{22|22}^{(3)}, \quad (23f)$$

$$M_{41} = 0, \quad M_{42} = 0, \quad M_{43} = \bar{\chi}_{22|22}^{(1)} + \frac{2}{5} M_{44}, \quad (23g)$$

$$M_{44} = \frac{5}{6} \left[\bar{\chi}_{22|22}^{(3)} - \chi_{20|20} - \frac{4\theta}{\kappa} \chi_{20|02} - \chi_{02|02} - \frac{\kappa}{4\theta} \chi_{02|20} \right], \quad (23h)$$

$$L_1 = M_{11} + 2M_{12} + 2M_{13} + \frac{3}{5} \chi_{40|40}^{(2)} + \frac{1}{5} \chi_{40|40}^{(3)}, \quad (24a)$$

$$L_2 = 2M_{21} + M_{22} + 2M_{23} + \frac{3}{5} \chi_{04|04}^{(2)} + \frac{1}{5} \chi_{04|04}^{(3)}, \quad (24b)$$

$$L_3 = \frac{8}{5} M_{31} + \frac{8}{5} M_{32} + M_{33} + \frac{\kappa}{20\theta} \chi_{22|40}^{(2)} + \frac{3}{5} \chi_{22|22}^{(2)} + \frac{1}{5} \chi_{22|22}^{(4)} + \frac{4\theta}{5\kappa} \chi_{22|04}^{(2)}, \quad (24c)$$

$$L_4 = M_{43} + \frac{\kappa}{6\theta} \bar{\chi}_{22|40} + \bar{\chi}_{22|22}^{(2)} + \frac{1}{3} \bar{\chi}_{22|22}^{(4)} + \frac{8\theta}{3\kappa} \bar{\chi}_{22|04}. \quad (24d)$$




Fig. 3 **a** Plot of the smallest eigenvalue μ_{\min} as a function of β for $\kappa = \frac{2}{5}$ and $\alpha = 0.4$ (—), 0.6 (---), 0.8 (--) and 1 (···). The circles denote the location of the threshold value (β_0) below which $\mu_{\min} < 0$. **b** Plot of β_0 as a function of α for $\kappa = \frac{2}{5}$. The fourth-degree moments diverge in the region below the curve.

The time evolution of the cumulants is characterized by the eigenvalues of the matrix \mathbf{M} , the asymptotic behavior being governed by the smallest eigenvalue, μ_{\min} . If $\mu_{\min} > 0$, the cumulants relax to their finite stationary values $\left[a_{20}^{(0)}, a_{02}^{(0)}, a_{11}^{(0)}, a_{00}^{(1)} \right]^\dagger = -\mathbf{M}^{-1} \cdot \mathbf{L}$. On the other hand, if $\mu_{\min} < 0$ then the cumulants diverge. Figure 3a shows the β -dependence of μ_{\min} for the same representative values of α as in Fig. 2. We observe that, at a given $\alpha < 1$, there exists a threshold value β_0 such that $\mu_{\min} < 0$ if $\beta < \beta_0$. The α -dependence of β_0 is shown in Fig. 3b. For the pairs (α, β) below the curve in Fig. 3b, the cumulants, and hence the fourth-degree moments, diverge. This divergence is likely associated with an algebraic high-velocity tail of the VDF, as already present in the case of the IMM [13–17, 32, 39, 40, 89].

If the particles are perfectly smooth ($\beta = -1$), the temperature ratio θ is irrelevant and all the matrix elements vanish except $M_{11} = \frac{1}{120}(1 + \alpha)^2(5 + 6\alpha - 3\alpha^2)$, $M_{33} = \frac{1}{20}(1 + \alpha)^2$, $M_{43} = \frac{1}{36}(1 + \alpha)^2$, $M_{44} = \frac{1}{9}(1 + \alpha)^2$, and $L_1 = -\frac{1}{20}(1 - \alpha^2)^2$. This implies that $a_{02}^{(0)}$ is arbitrary, $a_{11}^{(0)} = a_{00}^{(1)} = 0$, and $a_{20}^{(0)} = 6(1 - \alpha)^2/(5 + 6\alpha - 3\alpha^2)$. The latter result agrees with a previous derivation for the IMM [21].

On the other hand, in the special case of the Pidduck gas ($\alpha = \beta = 1$), the HCS reduces to equilibrium and Eqs. (19) yield $\theta = 1$, as expected. Moreover, from Eqs. (C15) and (C16), one gets $\mathbf{L} = 0$, implying that the cumulants (21) vanish in that case.

Apart from the IMM and the equilibrium Pidduck gas, in the general case with $\alpha < 1$ and $|\beta| < 1$, the stationary values of the cumulants are given, as said above, by $\left[a_{20}^{(0)}, a_{02}^{(0)}, a_{11}^{(0)}, a_{00}^{(1)} \right]^\dagger = -\mathbf{M}^{-1} \cdot \mathbf{L}$, provided that $\beta > \beta_0$. The β -dependence of the four cumulants are displayed in Fig. 4 for the same choices of α as in Figs. 2 and 3a. The positivity of $a_{11}^{(0)}$ implies that high translational velocities are strongly correlated to high angular velocities [73, 76, 88]. In




Fig. 4 Plot of the cumulants **a** $a_{20}^{(0)}$, **b** $a_{02}^{(0)}$, **c** $a_{11}^{(0)}$, and **d** $a_{00}^{(1)}$ as functions of β for $\kappa = \frac{2}{5}$ and $\alpha = 0.4$ (—), 0.6 (---), 0.8 (- -), and 1 (···). The vertical lines denote the asymptotes at $\beta = \beta_0$.

turn, the negative values of $a_{00}^{(1)}$ mean that quasinormal orientations between the translational and angular velocity vectors tend to be favored against quasiparallel orientations (“lifted-tennis-ball” effect) [67, 73, 76, 90, 91].

5 Conclusions

In this work, we have worked out a simple Maxwell model (the IRMM) by keeping the collision rules of inelastic rough hard spheres but, on the other hand, assuming an effective mean-field collision rate independent of the relative velocity of the colliding pair. The latter assumption allows one to express a collisional moment of degree k as a bilinear combination of velocity moments of degrees $k_1 \leq k$ and $k_2 \leq k$ with $k_1 + k_2 = k$, as happens in the well-known IMM (smooth particles) [40]. Nevertheless, the derivation of exact results in the IRMM is much more complicated than in the IMM. Not only are the translational and rotational degrees of freedom entangled, thus increasing the number and structure of the moments, but also the coefficients in the bilinear combinations are functions of the coefficient of normal restitution (as in the

14 *Inelastic and rough Maxwell particles*

IMM) and, additionally, of the coefficient of tangential restitution and the reduced moment of inertia.

Specifically, we have considered the moments of first and second degree, the moments of third degree related to the heat flux, and the isotropic fourth-degree moments. The structure of the associated collisional moments was displayed in Table 1. The number of coefficients is 60, since in some cases a common coefficient factorizes more than one combination of moments. The exact expressions of those 60 coefficients in terms of the two coefficients of restitution (α and β) and of the reduced moment of inertia (κ) are presented in Appendix B. We have checked that the coefficients satisfy some consistency tests. First, they reduce to the expressions reported in the literature in the smooth limit ($\beta = -1$) [32, 40]. Next, when particularized to elastic and perfectly rough particles (Pidduck model, $\alpha = \beta = 1$), the coefficients obey a number of relations needed to allow the equilibrium VDF to be an exact solution of the model.

The knowledge of the collisional moments stemming from the IRMM is not constrained to a specific physical situation and, thus, it can be exploited in several applications. Here we have applied the results to the HCS. Specifically, the rotational-to-translational temperature ratio has been found [see Eqs. (19)] and the linear evolution equations for the fourth-degree cumulants have been obtained [see Eq. (22)]. Interestingly, we have found that those cumulants diverge in time if, at a given α , the coefficient of tangential restitution lies below a threshold value $\beta_0(\alpha)$ [see Fig. 3]. This reflects the existence of an algebraic high-velocity tail of the VDF, reminiscent of the case of the IMM [14–17, 89]. If $\beta > \beta_0(\alpha)$, the exact stationary cumulants have been obtained (see Fig. 4). In general, the departure of the HCS VDF from the equilibrium one is rather strong: (a) the (marginal) translational and angular VDFs exhibit large excess kurtoses, (b) high translational velocities are correlated to high angular velocities, and (c) the translational and angular velocities tend to adopt quasi-normal orientations (“lifted-tennis-ball” effect).

It must be stressed that, in analogy to the relationship between the IHSM and the IMM, the IRMM is intended as a simplification of the IRHSM and not necessarily as an approximation of the latter. In this respect, the predictions obtained analytically from the IRMM do not need to be interpreted strictly as a quantitative replacement for the predictions obtained either numerically or by simulations from the IRHSM. On the other hand, the IRMM provides a tractable toy model that can be used to cleanly unveil nontrivial physical properties and serve as a benchmark for numerical or simulation approaches.

As further applications of the model, we plan to apply it to the exact derivation of the Navier–Stokes transport coefficients and to the non-Newtonian rheological properties of a granular gas under simple shear flow. The results will be published elsewhere.

Acknowledgments. A.S. acknowledges financial support from Grant PID2020-112936GB-I00 funded by MCIN/AEI/10.13039/501100011033, and from grants IB20079 and GR21014 funded by Junta de Extremadura (Spain)

and by ERDF “A way of making Europe.” G.M.K. is grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for financial support through Grant No. 304054/2019-4.

Appendix A Collision rules

Let $(\mathbf{v}_1, \mathbf{v}_2)$ and $(\boldsymbol{\omega}_1, \boldsymbol{\omega}_2)$ denote the linear and the angular precollisional velocities, respectively, of two rough spherical particles of mass m , diameter σ , and moment of inertia I ; and let $(\mathbf{v}'_1, \mathbf{v}'_2)$ and $(\boldsymbol{\omega}'_1, \boldsymbol{\omega}'_2)$ denote the corresponding postcollisional velocities. The pre- and postcollisional velocities are related by

$$m\mathbf{v}'_1 = m\mathbf{v}_1 - \mathbf{Q}, \quad I\boldsymbol{\omega}'_1 = I\boldsymbol{\omega}_1 - \frac{\sigma}{2}\hat{\boldsymbol{\sigma}} \times \mathbf{Q}, \quad (\text{A1a})$$

$$m\mathbf{v}'_2 = m\mathbf{v}_2 + \mathbf{Q}, \quad I\boldsymbol{\omega}'_2 = I\boldsymbol{\omega}_2 - \frac{\sigma}{2}\hat{\boldsymbol{\sigma}} \times \mathbf{Q}, \quad (\text{A1b})$$

where \mathbf{Q} is the impulse exerted by particle 1 on particle 2 and $\hat{\boldsymbol{\sigma}} = (\mathbf{r}_2 - \mathbf{r}_1)/\sigma$ is the unit collision vector joining the centers of the two colliding spheres. Additionally, the relative velocity $\bar{\mathbf{g}}$ of the points of the spheres at contact during a collision and the center-of-mass relative velocity $\mathbf{g} = \mathbf{v}_1 - \mathbf{v}_2$ are related by

$$\bar{\mathbf{g}} = \mathbf{g} - \frac{\sigma}{2}\hat{\boldsymbol{\sigma}} \times (\boldsymbol{\omega}_1 + \boldsymbol{\omega}_2). \quad (\text{A2})$$

A similar relation holds between $\bar{\mathbf{g}}'$ and \mathbf{g}' .

Combination of Eqs. (A1) and (A2) yields

$$\bar{\mathbf{g}}' = \bar{\mathbf{g}} - \frac{2}{m} \frac{\kappa + 1}{\kappa} \mathbf{Q} + \frac{2}{m\kappa} (\hat{\boldsymbol{\sigma}} \cdot \mathbf{Q}) \hat{\boldsymbol{\sigma}}. \quad (\text{A3})$$

Here,

$$\kappa = \frac{4I}{m\sigma^2} \quad (\text{A4})$$

is a dimensionless moment of inertia. It ranges from $\kappa = 0$ (mass concentrated at the center of the sphere) to a maximum value $\kappa = \frac{2}{3}$ (mass concentrated at the spherical surface). The value $\kappa = \frac{2}{5}$ corresponds to a uniform mass distribution.

In the most basic model, the inelastic collisions of rough spherical particles are characterized by

$$\hat{\boldsymbol{\sigma}} \cdot \bar{\mathbf{g}}' = -\alpha(\hat{\boldsymbol{\sigma}} \cdot \bar{\mathbf{g}}), \quad \hat{\boldsymbol{\sigma}} \times \bar{\mathbf{g}}' = -\beta(\hat{\boldsymbol{\sigma}} \times \bar{\mathbf{g}}), \quad (\text{A5})$$

where $0 < \alpha \leq 1$ and $-1 \leq \beta \leq 1$ are the coefficients of normal and tangential restitution, respectively. For an elastic collision of perfectly smooth spheres one has $\alpha = 1$ and $\beta = -1$, while $\alpha = 1$ and $\beta = 1$ for an elastic encounter of perfectly rough spherical particles.

Insertion of Eq. (A3) into Eq. (A5) gives $\hat{\sigma} \cdot \mathbf{Q} = m\tilde{\alpha}(\hat{\sigma} \cdot \bar{\mathbf{g}})$ and $\hat{\sigma} \times \mathbf{Q} = m\tilde{\beta}(\hat{\sigma} \times \bar{\mathbf{g}})$, where

$$\tilde{\alpha} \equiv \frac{1+\alpha}{2}, \quad \tilde{\beta} \equiv \frac{1+\beta}{2} \frac{\kappa}{1+\kappa}. \quad (\text{A6})$$

Therefore, the impulse can be expressed as

$$\mathbf{Q} = m\tilde{\alpha}(\hat{\sigma} \cdot \mathbf{g})\hat{\sigma} - m\tilde{\beta}\hat{\sigma} \times \left(\hat{\sigma} \times \mathbf{g} + \sigma \frac{\omega_1 + \omega_2}{2} \right). \quad (\text{A7})$$

Equations (A1) and (A7) express the postcollisional velocities in terms of the precollisional velocities and of the collision vector [92].

Apart from the linear momentum, the total angular momentum $\mathbf{L} = m(\mathbf{r}_1 \times \mathbf{v}_1 + \mathbf{r}_2 \times \mathbf{v}_2) + I(\omega_1 + \omega_2)$ is conserved in a collision. On the other hand, the change of the total (translational plus rotational) kinetic energy is

$$\Delta K = -m \frac{1-\beta^2}{4} \frac{\kappa}{\kappa+1} \left[\hat{\sigma} \times \left(\hat{\sigma} \times \mathbf{g} + \sigma \frac{\omega_1 + \omega_2}{2} \right) \right]^2 - m \frac{1-\alpha^2}{4} (\hat{\sigma} \cdot \mathbf{g})^2. \quad (\text{A8})$$

The right-hand side vanishes only either for elastic collisions of perfectly smooth spheres ($\alpha = 1, \beta = -1$) or for elastic collisions of perfectly rough spherical particles ($\alpha = \text{beta} = 1$).

Appendix B Explicit expressions for the coefficients appearing in Table 1

The 60 coefficients appearing in Table 1 are given by

$$\varphi_{01|01} = \frac{4\tilde{\beta}}{3\kappa}, \quad (\text{B1})$$

$$\chi_{20|20} = \frac{2}{3} \left[\tilde{\alpha}(1-\tilde{\alpha}) + 2\tilde{\beta}(1-\tilde{\beta}) \right], \quad \chi_{20|02} = -\frac{\tilde{\beta}^2}{3}, \quad (\text{B2})$$

$$\psi_{20|20} = \frac{2}{15} \left(5\tilde{\alpha} - 2\tilde{\alpha}^2 - 6\tilde{\alpha}\tilde{\beta} + 10\tilde{\beta} - 7\tilde{\beta}^2 \right), \quad \psi_{20|02} = \frac{\tilde{\beta}^2}{6}, \quad (\text{B3})$$

$$\chi_{02|02} = \frac{4\tilde{\beta}}{3\kappa} \left(1 - \frac{\tilde{\beta}}{\kappa} \right), \quad \chi_{02|20} = -\frac{16\tilde{\beta}^2}{3\kappa^2}, \quad (\text{B4})$$

$$\psi_{02|02} = \frac{2\tilde{\beta}}{15\kappa} \left(10 - \frac{7\tilde{\beta}}{\kappa} \right), \quad \psi_{02|20} = \frac{8\tilde{\beta}^2}{3\kappa^2}, \quad (\text{B5})$$

$$\psi_{11|11} = \frac{1}{3} \left(\tilde{\alpha} + 2\tilde{\beta} \frac{1+\kappa}{\kappa} \right), \quad (\text{B6})$$

$$\varphi_{30|30} = \frac{1}{15} \left(15\tilde{\alpha} - 11\tilde{\alpha}^2 - 8\tilde{\alpha}\tilde{\beta} + 30\tilde{\beta} - 26\tilde{\beta}^2 \right), \quad \varphi_{30|12} = -\frac{\tilde{\beta}^2}{6}, \quad (\text{B7})$$

$$\varphi_{12|30} = -\frac{8\tilde{\beta}^2}{3\kappa^2}, \quad \varphi_{12|12}^{(1)} = \frac{1}{15} \left[5\tilde{\alpha} + 10\tilde{\beta} + \frac{2\tilde{\beta}}{\kappa} \left(10 - 4\tilde{\alpha} - 11\tilde{\beta} - \frac{5\tilde{\beta}}{\kappa} \right) \right], \quad (\text{B8a})$$

$$\varphi_{12|12}^{(2)} = \frac{2\tilde{\beta}}{15\kappa} \left(2\tilde{\alpha} + 3\tilde{\beta} \right), \quad \varphi_{12|12}^{(3)} = \frac{4\tilde{\beta}}{3\kappa} \left(1 - \frac{\tilde{\beta}}{\kappa} \right), \quad \varphi_{12|12}^{(4)} = \frac{2\tilde{\beta}^2}{3\kappa}, \quad (\text{B8b})$$

$$\overline{\varphi}_{12|12}^{(1)} = \frac{1}{15} \left[5\tilde{\alpha} + 10\tilde{\beta} + \frac{\tilde{\beta}}{\kappa} \left(20 - 3\tilde{\alpha} - 7\tilde{\beta} \frac{1+\kappa}{\kappa} \right) \right], \quad (\text{B9a})$$

$$\overline{\varphi}_{12|12}^{(2)} = \frac{\tilde{\beta}}{15\kappa} \left(\tilde{\alpha} - \tilde{\beta} \frac{1+\kappa}{\kappa} \right), \quad \overline{\varphi}_{12|12}^{(3)} = \frac{\tilde{\beta}}{15\kappa} \left(10 + 5\tilde{\alpha} + 15\tilde{\beta} - \frac{7\tilde{\beta}}{\kappa} \right), \quad (\text{B9b})$$

$$\overline{\varphi}_{12|12}^{(4)} = -\frac{2\tilde{\beta}^2}{15\kappa^2}, \quad \overline{\varphi}_{12|12}^{(5)} = \frac{\tilde{\beta}}{15\kappa} \left(10 - 5\tilde{\alpha} - 15\tilde{\beta} - \frac{7\tilde{\beta}}{\kappa} \right), \quad (\text{B9c})$$

$$\chi_{40|40}^{(1)} = \frac{2}{15} \left[10\tilde{\alpha} - 11\tilde{\alpha}^2 + 6\tilde{\alpha}^3 - 3\tilde{\alpha}^4 + 20\tilde{\beta} - 26\tilde{\beta}^2 + 16\tilde{\beta}^3 - 8\tilde{\beta}^4 - 4\tilde{\alpha}\beta(2 - \alpha - \beta + \alpha\beta) \right], \quad (\text{B10a})$$

$$\chi_{40|40}^{(2)} = -\frac{2}{15} \left[7\tilde{\alpha}^2 - 6\tilde{\alpha}^3 + 3\tilde{\alpha}^4 + 12\tilde{\beta}^2 - 16\tilde{\beta}^3 + 8\tilde{\beta}^4 - 4\tilde{\alpha}\beta(1 + \alpha + \beta - \alpha\beta) \right], \quad (\text{B10b})$$

$$\chi_{40|40}^{(3)} = -\frac{4}{15} \left[2\tilde{\alpha}^2 - 6\tilde{\alpha}^3 + 3\tilde{\alpha}^4 + 7\tilde{\beta}^2 - 16\tilde{\beta}^3 + 8\tilde{\beta}^4 + 2\alpha\beta(3 - 2\alpha - 2\beta + 2\alpha\beta) \right], \quad (\text{B10c})$$

$$\chi_{40|04} = -\frac{\tilde{\beta}^4}{15}, \quad \chi_{40|22}^{(1)} = -\frac{\tilde{\beta}^2}{15} \left[5 - 2\tilde{\alpha}(1 - \tilde{\alpha}) - 8\tilde{\beta}(1 - \tilde{\beta}) \right], \quad (\text{B10d})$$

$$\chi_{40|22}^{(2)} = \frac{2\tilde{\beta}^2}{15} \left[\tilde{\alpha}(1 - \tilde{\alpha}) + 4\tilde{\beta}(1 - \tilde{\beta}) \right], \quad (\text{B10e})$$

$$\chi_{04|40} = -\frac{256\tilde{\beta}^4}{15\kappa^4}, \quad \chi_{04|22}^{(1)} = -\frac{16\tilde{\beta}^2}{15\kappa^2} \left(5 - \frac{8\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^2}{\kappa^2} \right), \quad (\text{B11a})$$

$$\chi_{04|22}^{(2)} = -\frac{128\tilde{\beta}^3}{15\kappa^3} \left(1 - \frac{\tilde{\beta}}{\kappa} \right), \quad \chi_{04|04}^{(1)} = \frac{4\tilde{\beta}}{15\kappa} \left(10 - \frac{13\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^2}{\kappa^2} - \frac{4\tilde{\beta}^3}{\kappa^3} \right), \quad (\text{B11b})$$

$$\chi_{04|04}^{(2)} = -\frac{8\tilde{\beta}^2}{15\kappa^2} \left(3 - \frac{4\tilde{\beta}}{\kappa} + \frac{2\tilde{\beta}^2}{\kappa^2} \right), \quad \chi_{04|04}^{(3)} = -\frac{4\tilde{\beta}^2}{15\kappa^2} \left(7 - \frac{16\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^2}{\kappa^2} \right), \quad (\text{B11c})$$

$$\chi_{04|04}^{(4)} = \frac{8\tilde{\beta}}{15\kappa} \left(1 - \frac{\beta}{\kappa} \right) \left(5 - \frac{8\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^2}{\kappa^2} \right), \quad (\text{B11d})$$

$$\chi_{22|40}^{(1)} = -\frac{8\tilde{\beta}^2}{15\kappa^2} \left[5 - 2\tilde{\alpha}(1 - \tilde{\alpha}) - 8\tilde{\beta}(1 - \tilde{\beta}) \right], \quad (\text{B12a})$$

$$\chi_{22|40}^{(2)} = \frac{32\tilde{\beta}^2}{15\kappa^2} \left[\tilde{\alpha}(1 - \tilde{\alpha}) + 4\tilde{\beta}(1 - \tilde{\beta}) \right], \quad (\text{B12b})$$

$$\begin{aligned} \chi_{22|22}^{(1)} = & \frac{1}{15} \left[5\tilde{\alpha}(2 - \tilde{\alpha}) + 20\tilde{\beta} \frac{1 + \kappa}{\kappa} - 2\tilde{\beta}^2 \frac{5 + 22\kappa + 5\kappa^2}{\kappa^2} - \frac{8\tilde{\alpha}(2 - \tilde{\alpha})\tilde{\beta}}{\kappa} \right. \\ & \left. + \frac{8\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}^2}{\kappa^2} + 32\tilde{\beta}^3 \frac{1 + \kappa}{\kappa^2} - \frac{64\tilde{\beta}^4}{\kappa^2} \right], \end{aligned} \quad (\text{B12c})$$

$$\begin{aligned} \chi_{22|22}^{(2)} = & -\frac{1}{15} \left[5\tilde{\alpha}^2 + 10\tilde{\beta}^2 \frac{1 + \kappa^2}{\kappa^2} - \frac{8\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}^2}{\kappa^2} - \frac{8\tilde{\alpha}^2\tilde{\beta}}{\kappa} - 32\tilde{\beta}^3 \frac{1 + \kappa}{\kappa^2} \right. \\ & \left. + \frac{64\tilde{\beta}^4}{\kappa^2} \right], \end{aligned} \quad (\text{B12d})$$

$$\chi_{22|22}^{(3)} = \frac{4\tilde{\beta}}{15\kappa} \left[\tilde{\alpha}(2 - \tilde{\alpha}) + 3\tilde{\beta} - 4\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} - \frac{\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12e})$$

$$\chi_{22|22}^{(4)} = -\frac{4\tilde{\beta}}{15\kappa} \left[\tilde{\alpha}^2 + 4\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} + \frac{\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} - \frac{8\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12f})$$

$$\chi_{22|22}^{(5)} = \frac{4\tilde{\beta}}{15\kappa} \left[2\tilde{\alpha}(1 - \tilde{\alpha}) + 3\tilde{\beta} - 8\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} - \frac{2\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} + \frac{16\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12g})$$

$$\begin{aligned} \chi_{22|22}^{(6)} = & \frac{4\tilde{\beta}}{15\kappa} \left[5 - 4\tilde{\alpha}(1 - \tilde{\alpha}) - \tilde{\beta} \frac{5 + 11\kappa}{\kappa} + 16\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} + \frac{4\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} - \frac{32\tilde{\beta}^3}{\kappa} \right], \\ & \quad (\text{B12h}) \end{aligned}$$

$$\chi_{22|22}^{(7)} = -\frac{4\tilde{\beta}}{15\kappa} \left[\tilde{\alpha}(1 - \tilde{\alpha}) + 4\tilde{\beta} - 4\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} - \frac{\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12i})$$

$$\chi_{22|22}^{(8)} = -\frac{4\tilde{\beta}}{15\kappa} \left[\tilde{\alpha}(1 - \tilde{\alpha}) - \tilde{\beta} - 4\tilde{\beta}^2 \frac{1 + \kappa}{\kappa} - \frac{\tilde{\alpha}(1 - \tilde{\alpha})\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12j})$$

$$\chi_{22|22}^{(9)} = \frac{4\tilde{\beta}}{15\kappa} \left[4\tilde{\alpha}(1-\tilde{\alpha}) + 11\tilde{\beta} - 16\tilde{\beta}^2 \frac{1+\kappa}{\kappa} - \frac{4\tilde{\alpha}(1-\tilde{\alpha})\tilde{\beta}}{\kappa} + \frac{32\tilde{\beta}^3}{\kappa} \right], \quad (\text{B12k})$$

$$\chi_{22|04}^{(1)} = -\frac{\tilde{\beta}^2}{30} \left(5 - \frac{8\tilde{\beta}}{\kappa} + \frac{8\tilde{\beta}^2}{\kappa^2} \right), \quad \chi_{22|04}^{(2)} = \frac{8\tilde{\beta}^3}{15\kappa} \left(1 - \frac{\tilde{\beta}}{\kappa} \right), \quad (\text{B12l})$$

$$\chi_{22|04}^{(3)} = -\frac{\tilde{\beta}^2}{15} \left(5 - \frac{16\tilde{\beta}}{\kappa} + \frac{16\tilde{\beta}^2}{\kappa^2} \right), \quad (\text{B12m})$$

$$\bar{\chi}_{22|40} = -\frac{4\tilde{\beta}^2}{3\kappa^2}, \quad \bar{\chi}_{22|22}^{(1)} = -\frac{1}{15} \left(\tilde{\alpha} - \tilde{\beta} \frac{1+\kappa}{\kappa} \right)^2, \quad (\text{B13a})$$

$$\bar{\chi}_{22|22}^{(2)} = -\frac{1}{15} \left[\left(\tilde{\alpha} - \tilde{\beta} \right)^2 + \frac{\tilde{\beta}^2}{\kappa^2} \right], \quad (\text{B13b})$$

$$\bar{\chi}_{22|22}^{(3)} = \frac{1}{15} \left[2\tilde{\alpha}(5-\tilde{\alpha}) + 2\tilde{\beta}(10-3\tilde{\alpha}) \frac{1+\kappa}{\kappa} - 7\tilde{\beta}^2 \frac{(1+\kappa)^2}{\kappa^2} \right], \quad (\text{B13c})$$

$$\bar{\chi}_{22|22}^{(4)} = -\frac{1}{15} \left(2\tilde{\alpha}^2 + 6\tilde{\alpha}\tilde{\beta} + 7\tilde{\beta}^2 \frac{1+\kappa^2}{\kappa^2} \right), \quad (\text{B13d})$$

$$\bar{\chi}_{22|22}^{(5)} = \frac{2\tilde{\beta}}{15\kappa} \left(10 - 3\tilde{\alpha} - 7\tilde{\beta} \frac{1+\kappa}{\kappa} \right), \quad \bar{\chi}_{22|22}^{(6)} = \frac{2\tilde{\beta}}{15\kappa} \left(\tilde{\alpha} - \tilde{\beta} \frac{1+\kappa}{\kappa} \right), \quad (\text{B13e})$$

$$\bar{\chi}_{22|22}^{(7)} = -\frac{2\tilde{\beta}}{15\kappa} \left(\tilde{\alpha} + 4\tilde{\beta} \right), \quad \bar{\chi}_{22|22}^{(8)} = \frac{2\tilde{\beta}}{15\kappa} \left(4\tilde{\alpha} + 11\tilde{\beta} \right), \quad (\text{B13f})$$

$$\bar{\chi}_{22|22}^{(9)} = -\frac{2\tilde{\beta}}{15\kappa} \left(\tilde{\alpha} - \tilde{\beta} \right), \quad \bar{\chi}_{22|04} = -\frac{\tilde{\beta}^2}{12}. \quad (\text{B13g})$$

Appendix C Consistency tests

Tables C1–C3 display the expressions of the 60 coefficients in two interesting situations: (i) inelastic and perfectly smooth particles (i.e., the IMM, $\alpha < 1$, $\beta = -1$) and (ii) elastic and perfectly rough particles (i.e., the Pidduck model, $\alpha = \beta = 1$). Let us use both situations (i) and (ii) as tests of the results.

C.1 Inelastic Maxwell model (IMM)

In case (i), the angular velocities are unaffected by collisions and thus $\mathcal{J}_M[\Psi] = 0$ if $\Psi(\xi)$ is a function of ω only. Moreover, even though $\psi_{11|11} = \varphi_{12|12}^{(1)} = \overline{\varphi}_{12|12}^{(1)} \neq 0$, the angular and translational velocities are uncorrelated and thus $\mathcal{J}_M[V_i\omega_j] \propto \langle V_i\omega_j \rangle = \langle V_i \rangle \langle \omega_j \rangle = 0$, $\mathcal{J}_M[\omega^2 V_i] \propto \langle \omega^2 V_i \rangle = \langle \omega^2 \rangle \langle V_i \rangle = 0$, and $\mathcal{J}_M[(\mathbf{V} \cdot \boldsymbol{\omega})\omega_i] \propto \langle (\mathbf{V} \cdot \boldsymbol{\omega})\omega_i \rangle = \langle \mathbf{V} \rangle \cdot \langle \boldsymbol{\omega}\omega_i \rangle = 0$. Additionally, the following relations are fulfilled:

$$\chi_{22|22}^{(1)} + \chi_{22|22}^{(2)} = \chi_{20|20}, \quad (\text{C14a})$$

Table C1 Coefficients associated with the collisional moments of first, second, and third degree in Table 1 in the special cases of (i) inelastic and perfectly smooth particles ($\alpha < 1$, $\beta = -1$) and (ii) elastic and perfectly rough particles ($\alpha = \beta = 1$).

Coefficient	(i) Inelastic and perfectly smooth	(ii) Elastic and perfectly rough
	$(\alpha < 1, \beta = -1)$	$(\alpha = \beta = 1)$
$\varphi_{01 01}$	0	$\frac{4}{3(1+\kappa)}$
$\chi_{20 20}$	$\frac{1-\alpha^2}{6}$	$\frac{4\kappa}{3(1+\kappa)^2}$
$\chi_{20 02}$	0	$-\frac{\kappa}{4}\chi_{20 20}$
$\psi_{20 20}$	$\frac{(1+\alpha)(4-\alpha)}{15}$	$\frac{2(3+10\kappa)}{15(1+\kappa)^2}$
$\psi_{20 02}$	0	$\frac{\kappa}{8}\chi_{20 20}$
$\chi_{02 02}$	0	$\chi_{20 20}$
$\chi_{02 20}$	0	$-\frac{4}{\kappa}\chi_{20 20}$
$\psi_{02 02}$	0	$\psi_{20 20}$
$\psi_{02 20}$	0	$\frac{2}{\kappa}\chi_{20 20}$
$\psi_{11 11}$	$\frac{1+\alpha}{6}$	1
$\varphi_{30 30}$	$\frac{(1+\alpha)(19-11\alpha)}{60}$	$\frac{2(2+15\kappa)}{15(1+\kappa)^2}$
$\varphi_{30 12}$	0	$-\frac{\kappa}{8}\chi_{20 20}$
$\varphi_{12 30}$	0	$-\frac{2}{\kappa}\chi_{20 20}$
$\varphi_{12 12}^{(1)}$	$\psi_{11 11}$	$\frac{7+10\kappa+15\kappa^2}{15(1+\kappa)^2}$
$\varphi_{12 12}^{(2)}$	0	$\frac{2(2+5\kappa)}{15(1+\kappa)^2}$
$\varphi_{12 12}^{(3)}$	0	$\chi_{20 20}$
$\varphi_{12 12}^{(4)}$	0	$\frac{1}{2}\chi_{20 20}$
$\bar{\varphi}_{12 12}^{(1)}$	$\psi_{11 11}$	1
$\bar{\varphi}_{12 12}^{(2)}$	0	0
$\bar{\varphi}_{12 12}^{(3)}$	0	$\frac{2(4+15\kappa)}{15(1+\kappa)^2}$
$\bar{\varphi}_{12 12}^{(4)}$	0	$-\frac{1}{10\kappa}\chi_{20 20}$
$\bar{\varphi}_{12 12}^{(5)}$	0	$-\frac{2(1+5\kappa)}{15(1+\kappa)^2}$

Table C2 Coefficients associated with the collisional moments $\mathcal{J}_M[V^4]$ and $\mathcal{J}_M[\omega^4]$ in Table 1 in the special cases of (i) inelastic and perfectly smooth particles ($\alpha < 1$, $\beta = -1$) and (ii) elastic and perfectly rough particles ($\alpha = \beta = 1$).

Coefficient	(i) Inelastic and perfectly smooth	(ii) Elastic and perfectly rough
	$(\alpha < 1, \beta = -1)$	$(\alpha = \beta = 1)$
$\chi_{40 40}^{(1)}$	$\frac{(1 + \alpha)(45 - 29\alpha + 3\alpha^2 - 3\alpha^3)}{120}$	$\frac{4(1 + 12\kappa + 17\kappa^2 + 10\kappa^3)}{15(1 + \kappa)^4}$
$\chi_{40 40}^{(2)}$	$-\frac{(1 + \alpha)^2(19 - 6\alpha + 3\alpha^2)}{120}$	$-\frac{8(1 + 2\kappa + 3\kappa^2)}{15(1 + \kappa)^4}$
$\chi_{40 40}^{(3)}$	$\frac{(1 + \alpha)^2(1 + 6\alpha - 3\alpha^2)}{60}$	$\frac{4(1 + 2\kappa - 7\kappa^2)}{15(1 + \kappa)^4}$
$\chi_{40 04}$	0	$-\frac{15(1 + \kappa)^4}{\kappa^4}$
$\chi_{40 22}^{(1)}$	0	$-\frac{\kappa^2(5 + 2\kappa + 5\kappa^2)}{15(1 + \kappa)^4}$
$\chi_{40 22}^{(2)}$	0	$-\frac{8}{\kappa}\chi_{40 04}$
$\chi_{04 40}$	0	$\frac{256}{\kappa^4}\chi_{40 04}$
$\chi_{04 22}^{(1)}$	0	$\frac{16}{\kappa^2}\chi_{40 22}^{(1)}$
$\chi_{04 22}^{(2)}$	0	$\frac{128}{\kappa^3}\chi_{40 04}$
$\chi_{04 04}^{(1)}$	0	$\chi_{40 40}^{(1)}$
$\chi_{04 04}^{(2)}$	0	$\chi_{40 40}^{(2)}$
$\chi_{04 04}^{(3)}$	0	$\chi_{40 40}^{(3)}$
$\chi_{04 04}^{(4)}$	0	$-\frac{8}{\kappa}\chi_{40 22}^{(1)}$

$$\bar{\chi}_{22|22}^{(1)} + \bar{\chi}_{22|22}^{(2)} = \frac{1}{3} (\chi_{20|20} - \psi_{20|20}), \quad \bar{\chi}_{22|22}^{(3)} + \bar{\chi}_{22|22}^{(4)} = \psi_{20|20}. \quad (\text{C14b})$$

Equations (C14a) and (C14b) imply $\mathcal{J}_M[V^2\omega^2] = \langle \omega^2 \rangle \mathcal{J}_M[V^2]$ and $\mathcal{J}_M[(\mathbf{V} \cdot \boldsymbol{\omega})^2] = \langle \boldsymbol{\omega} \boldsymbol{\omega} \rangle : \mathcal{J}_M[\mathbf{V}\mathbf{V}]$, respectively. Finally, the coefficients $\chi_{20|20}$, $\psi_{20|20}$, $\varphi_{30|30}$, $\chi_{40|40}^{(1)}$, $\chi_{40|40}^{(2)}$, and $\chi_{40|40}^{(3)}$ agree with previous results for the IMM [2, 21, 31, 32, 39, 40].

C.2 Pidduck model

In case (ii) the total energy is conserved by collisions, i.e., $\mathcal{J}_M[mV^2 + I\omega^2] = 0$. This is guaranteed by the relations

$$\chi_{20|20} + \frac{\kappa}{4}\chi_{02|20} = 0, \quad \chi_{20|02} + \frac{\kappa}{4}\chi_{02|02} = 0. \quad (\text{C15})$$

Other tests in case (ii) correspond to the fact that $\mathcal{J}_M[\Psi] = 0$ for any $\Psi(\xi)$ if the system is at equilibrium, in which case all the anisotropic moments vanish and

Table C3 Coefficients associated with the collisional moments $\mathcal{J}_M[V^2\omega^2]$ and $\mathcal{J}_M[(\mathbf{V} \cdot \boldsymbol{\omega})^2]$ in Table 1 in the special cases of (i) inelastic and perfectly smooth particles ($\alpha < 1$, $\beta = -1$) and (ii) elastic and perfectly rough particles ($\alpha = \beta = 1$).

Coefficient	(i) Inelastic and perfectly smooth	(ii) Elastic and perfectly rough
	$(\alpha < 1, \beta = -1)$	$(\alpha = \beta = 1)$
$\chi_{22 40}^{(1)}$	0	$\frac{8}{\kappa^2} \chi_{40 22}^{(1)}$
$\chi_{22 40}^{(2)}$	0	$-\frac{128}{\kappa^3} \chi_{40 04}$
$\chi_{22 22}^{(1)}$	$\frac{(1+\alpha)(3-\alpha)}{12}$	$\frac{7+44\kappa+18\kappa^2+60\kappa^3+15\kappa^4}{15(1+\kappa)^4}$
$\chi_{22 22}^{(2)}$	$-\frac{(1+\alpha)^2}{12}$	$-\frac{7-16\kappa+26\kappa^2+15\kappa^4}{15(1+\kappa)^4}$
$\chi_{22 22}^{(3)}$	0	$\frac{4(1+2\kappa+9\kappa^2)}{15(1+\kappa)^4}$
$\chi_{22 22}^{(4)}$	0	$-\frac{4(1+7\kappa+3\kappa^2+5\kappa^3)}{15(1+\kappa)^4}$
$\chi_{22 22}^{(5)}$	0	$-\frac{4\kappa(5-6\kappa+5\kappa^2)}{15(1+\kappa)^4}$
$\chi_{22 22}^{(6)}$	0	$-2\chi_{22 22}^{(5)}$
$\chi_{22 22}^{(7)}$	0	$\frac{32}{\kappa^2} \chi_{40 04}$
$\chi_{22 22}^{(8)}$	0	$-\frac{4}{\kappa} \chi_{40 22}^{(1)}$
$\chi_{22 22}^{(9)}$	0	$-\frac{4\kappa(5-22\kappa+5\kappa^2)}{15(1+\kappa)^4}$
$\chi_{22 04}^{(1)}$	0	$\frac{1}{2} \chi_{40 22}^{(1)}$
$\chi_{22 04}^{(2)}$	0	$-\frac{8}{\kappa} \chi_{40 04}$
$\chi_{22 04}^{(3)}$	0	$\frac{\kappa}{4} \chi_{22 22}^{(5)}$
$\bar{\chi}_{22 40}$	0	$-\frac{1}{\kappa} \chi_{20 20}$
$\bar{\chi}_{22 22}^{(1)}$	$\frac{1}{5} \chi_{22 22}^{(2)}$	0
$\bar{\chi}_{22 22}^{(2)}$	$\frac{1}{5} \chi_{22 22}^{(2)}$	$-\frac{1}{10\kappa} \chi_{20 20}$
$\bar{\chi}_{22 22}^{(3)}$	$\frac{(1+\alpha)(9-\alpha)}{30}$	1
$\bar{\chi}_{22 22}^{(4)}$	$\frac{2}{5} \chi_{22 22}^{(2)}$	$-\frac{9+10\kappa+15\kappa^2}{15(1+\kappa)^2}$
$\bar{\chi}_{22 22}^{(5)}$	0	0
$\bar{\chi}_{22 22}^{(6)}$	0	0
$\bar{\chi}_{22 22}^{(7)}$	0	$\bar{\varphi}_{12 12}^{(5)}$
$\bar{\chi}_{22 22}^{(8)}$	0	$\bar{\varphi}_{12 12}^{(3)}$
$\bar{\chi}_{22 22}^{(9)}$	0	$-\frac{1}{10\kappa} \chi_{20 20}$
$\bar{\chi}_{22 04}$	0	$-\frac{\kappa}{16} \chi_{20 20}$

$\langle \omega^2 \rangle = \frac{m}{T} \langle V^2 \rangle = \frac{4}{\kappa \sigma^2} \langle V^2 \rangle$, $\langle V^2 \omega^2 \rangle = 3 \langle (\mathbf{V} \cdot \boldsymbol{\omega})^2 \rangle = \langle V^2 \rangle \langle \omega^2 \rangle$, $\langle V^4 \rangle = \frac{5}{3} \langle V^2 \rangle^2$, and $\langle \omega^4 \rangle = \frac{5}{3} \langle \omega^2 \rangle^2$. Therefore, according to Table 1, one should have

$$\chi_{20|20} + \frac{4}{\kappa} \chi_{20|02} = 0, \quad \chi_{02|20} + \frac{4}{\kappa} \chi_{02|02} = 0, \quad (\text{C16a})$$

$$5\chi_{40|40}^{(1)} + 3\chi_{40|40}^{(2)} + \chi_{40|40}^{(3)} + \frac{40}{\kappa} \chi_{40|22}^{(1)} + \frac{160}{\kappa^2} \chi_{40|04} = 0, \quad (\text{C16b})$$

$$5\chi_{04|40} + \frac{20}{\kappa} \chi_{04|22}^{(1)} + \frac{8}{\kappa^2} \left[5\chi_{04|04}^{(1)} + 3\chi_{04|04}^{(2)} + \chi_{04|04}^{(3)} \right] = 0, \quad (\text{C16c})$$

$$8\chi_{22|40}^{(1)} + \chi_{22|40}^{(2)} + \frac{4}{\kappa} \left[3\chi_{22|22}^{(1)} + 3\chi_{22|22}^{(2)} + \chi_{22|22}^{(3)} + \chi_{22|22}^{(4)} \right] \\ + \frac{16}{\kappa^2} \left[8\chi_{22|04}^{(1)} + \chi_{22|04}^{(2)} \right] = 0, \quad (\text{C16d})$$

$$\bar{\chi}_{22|40} + \frac{2}{\kappa} \left[3\bar{\chi}_{22|22}^{(1)} + 3\bar{\chi}_{22|22}^{(2)} + \bar{\chi}_{22|22}^{(3)} + \bar{\chi}_{22|22}^{(4)} \right] + \frac{16}{\kappa^2} \bar{\chi}_{22|04} = 0. \quad (\text{C16e})$$

From the expressions in Tables C1–C3 one can check that Eqs. (C16) are indeed satisfied.

References

- [1] N.V. Brilliantov, T. Pöschel, *Kinetic Theory of Granular Gases* (Oxford University Press, Oxford, 2004)
- [2] V. Garzó, *Granular Gaseous Flows. A Kinetic Theory Approach to Granular Gaseous Flows* (Springer Nature, Switzerland, 2019)
- [3] J.C. Maxwell, IV. On the dynamical theory of gases. Phil. Trans. Roy. Soc. (London) **157**, 49–88 (1867). <https://doi.org/10.1098/rstl.1867.0004>
- [4] C. Truesdell, R.G. Muncaster, *Fundamentals of Maxwell's Kinetic Theory of a Simple Monatomic Gas* (Academic Press, New York, 1980)
- [5] M.H. Ernst, Nonlinear model-Boltzmann equations and exact solutions. Phys. Rep. **78**, 1–171 (1981). [https://doi.org/10.1016/0370-1573\(81\)90002-8](https://doi.org/10.1016/0370-1573(81)90002-8)
- [6] V. Garzó, A. Santos, *Kinetic Theory of Gases in Shear Flows: Nonlinear Transport*. Fundamental Theories of Physics (Springer, Dordrecht, 2003)
- [7] A. Santos, Solutions of the moment hierarchy in the kinetic theory of Maxwell models. Cont. Mech. Thermodyn. **21**, 361–387 (2009). <https://doi.org/10.1007/s00161-009-0113-5>
- [8] G.M. Kremer, *An Introduction to the Boltzmann Equation and Transport Processes in Gases* (Springer, Berlin, 2010)

24 *Inelastic and rough Maxwell particles*

- [9] A.V. Bobylev, J.A. Carrillo, I.M. Gamba, On some properties of kinetic and hydrodynamic equations for inelastic interactions. *J. Stat. Phys.* **98**, 743–773 (2000). <https://doi.org/10.1023/A:1018627625800>
- [10] J.A. Carrillo, C. Cercignani, I.M. Gamba, Steady states of a Boltzmann equation for driven granular media. *Phys. Rev. E* **62**, 7700–7707 (2000). <https://doi.org/10.1103/PhysRevE.62.7700>
- [11] E. Ben-Naim, P.L. Krapivsky, Multiscaling in inelastic collisions. *Phys. Rev. E* **61**, R5–R8 (2000). <https://doi.org/10.1103/PhysRevE.61.R5>
- [12] C. Cercignani, Shear flow of a granular material. *J. Stat. Phys.* **102**, 1407–1415 (2001). <https://doi.org/10.1023/A:1004804815471>
- [13] A. Baldassarri, U. Marini Bettolo Marconi, A. Puglisi, Influence of correlations on the velocity statistics of scalar granular gases. *Europhys. Lett.* **58**, 14–20 (2002). <https://doi.org/10.1209/epl/i2002-00600-6>
- [14] E. Ben-Naim, P.L. Krapivsky, Scaling, multiscaling, and nontrivial exponents in inelastic collision processes. *Phys. Rev. E* **66**, 011309 (2002). <https://doi.org/10.1103/PhysRevE.66.011309>
- [15] P.L. Krapivsky, E. Ben-Naim, Nontrivial velocity distributions in inelastic gases. *J. Phys. A: Math. Gen.* **35**, L147–L152 (2002). <https://doi.org/10.1088/0305-4470/35/11/103>
- [16] M.H. Ernst, R. Brito, High-energy tails for inelastic Maxwell models. *Europhys. Lett.* **58**, 182–187 (2002). <https://doi.org/10.1209/epl/i2002-00622-0>
- [17] M.H. Ernst, R. Brito, Scaling solutions of inelastic Boltzmann equations with over-populated high energy tails. *J. Stat. Phys.* **109**, 407–432 (2002). <https://doi.org/10.1023/A:1020437925931>
- [18] E. Ben-Naim, P.L. Krapivsky, in *Granular Gas Dynamics, Lecture Notes in Physics*, vol. 624, ed. by T. Pöschel, S. Luding (Springer, Berlin, 2003), pp. 65–94
- [19] A.V. Bobylev, C. Cercignani, Self-similar asymptotics for the Boltzmann equation with inelastic and elastic interactions. *J. Stat. Phys.* **110**, 333–375 (2003). <https://doi.org/10.1023/A:1021031031038>
- [20] A.V. Bobylev, C. Cercignani, G. Toscani, Proof of an asymptotic property of self-similar solutions of the Boltzmann equation for granular materials. *J. Stat. Phys.* **111**, 403–417 (2003). <https://doi.org/10.1023/A:1022273528296>

- [21] A. Santos, Transport coefficients of d -dimensional inelastic Maxwell models. *Physica A* **321**, 442–466 (2003). [https://doi.org/10.1016/S0378-4371\(02\)01005-1](https://doi.org/10.1016/S0378-4371(02)01005-1)
- [22] A. Santos, M.H. Ernst, Exact steady-state solution of the Boltzmann equation: A driven one-dimensional inelastic Maxwell gas. *Phys. Rev. E* **68**, 011305 (2003). <https://doi.org/10.1103/PhysRevE.68.011305>
- [23] V. Garzó, Nonlinear transport in inelastic Maxwell mixtures under simple shear flow. *J. Stat. Phys.* **112**, 657–683 (2003). <https://doi.org/10.1023/A:1023828109434>
- [24] R. Brito, M.H. Ernst, in *Advances in Condensed Matter and Statistical Mechanics*, ed. by E. Korutcheva, R. Cuerno (Nova Science Publishers, New York, 2004), pp. 177–202
- [25] V. Garzó, A. Astillero, Transport coefficients for inelastic Maxwell mixtures. *J. Stat. Phys.* **118**, 935–971 (2005). <https://doi.org/10.1007/s10955-004-2006-0>
- [26] M. Bisi, J.A. Carrillo, G. Toscani, Decay rates in probability metrics towards homogeneous cooling states for the inelastic Maxwell model. *J. Stat. Phys.* **124**, 625–653 (2006). <https://doi.org/10.1007/s10955-006-9035-9>
- [27] A.V. Bobylev, I.M. Gamba, Boltzmann equations for mixtures of Maxwell gases: exact solutions and power like tails. *J. Stat. Phys.* **124**, 497–516 (2006). <https://doi.org/10.1007/s10955-006-9044-8>
- [28] F. Bolley, J.A. Carrillo, Tanaka theorem for inelastic Maxwell models. *Commun. Math. Phys.* **276**, 287–314 (2007). <https://doi.org/10.1007/s00220-007-0336-x>
- [29] J.A. Carrillo, G. Toscani, Contractive probability metrics and asymptotic behavior of dissipative kinetic equations. *Riv. Mat. Univ. Parma (7)* **6**, 75–198 (2007)
- [30] V. Garzó, Shear-rate dependent transport coefficients for inelastic Maxwell models. *J. Phys. A: Math. Theor.* **40**, 10,729–10,767 (2007). <https://doi.org/10.1088/1751-8113/40/35/002>
- [31] A. Santos, V. Garzó, Simple shear flow in inelastic Maxwell models. *J. Stat. Mech.* p. P08021 (2007). <https://doi.org/10.1088/1742-5468/2007/08/P08021>
- [32] V. Garzó, A. Santos, Third and fourth degree collisional moments for inelastic Maxwell model. *J. Phys. A: Math. Theor.* **40**, 14,927–14,943

- (2007). <https://doi.org/10.1088/1751-8113/40/50/002>
- [33] A.V. Bobylev, C. Cercignani, I.M. Gamba, *Generalized kinetic Maxwell models of granular gases* (Springer, Berlin, 2008), *Lecture Notes in Mathematics*, vol. 1937, pp. 23–58
- [34] A.V. Bobylev, C. Cercignani, I.M. Gamba, On the self-similar asymptotics for generalized non-linear kinetic Maxwell models. *Commun. Math. Phys.* **291**, 599–644 (2009). <https://doi.org/10.1007/s00220-009-0876-3>
- [35] E.A. Carlen, J.A. Carrillo, M.C. Carvalho, Strong convergence towards homogeneous cooling states for dissipative Maxwell models. *Ann. I. H. Poincaré – AN* **26**, 167–1700 (2009). <https://doi.org/10.1016/j.anihpc.2008.10.005>
- [36] V. Garzó, E. Trizac, Rheological properties for inelastic Maxwell mixtures under shear flow. *J. Non-Newton. Fluid Mech.* **165**, 932–940 (2010). <https://doi.org/10.1016/j.jnnfm.2010.01.016>
- [37] G. Furioli, A. Pulvirenti, E. Terraneo, G. Toscani, Convergence to self-similarity for the Boltzmann equation for strongly inelastic Maxwell molecules. *Ann. I. H. Poincaré – AN* **27**, 719–737 (2010). <https://doi.org/10.1016/j.anihpc.2009.11.005>
- [38] J.J. Brey, M.I. García de Soria, P. Maynar, Breakdown of hydrodynamics in the inelastic Maxwell model of granular gases. *Phys. Rev. E* **82**, 021303 (2010). <https://doi.org/10.1103/PhysRevE.82.021303>
- [39] V. Garzó, A. Santos, Hydrodynamics of inelastic Maxwell models. *Math. Model. Nat. Phenom.* **6**(4), 37–76 (2011). <https://doi.org/10.1051/mmnp/20116403>
- [40] A. Santos, V. Garzó, Collisional rates for the inelastic Maxwell model. Application to the divergence of anisotropic high-order velocity moments in the homogeneous cooling state. *Gran. Matt.* **14**, 105–110 (2012). <https://doi.org/10.1007/s10035-012-0336-1>
- [41] N. Khalil, V. Garzó, A. Santos, Hydrodynamic Burnett equations for inelastic Maxwell models of granular gases. *Phys. Rev. E* **89**, 052201 (2014). <https://doi.org/10.1103/PhysRevE.89.052201>
- [42] R. Gómez González, V. Garzó, Simple shear flow in granular suspensions: Inelastic Maxwell models and BGK-type kinetic model. *J. Stat. Mech.* **013206** (2019). <https://doi.org/10.1088/1742-5468/aaf719>
- [43] N. Khalil, V. Garzó, Unified hydrodynamic description for driven and undriven inelastic Maxwell mixtures at low density. *J. Phys. A: Math.*

- Theor. **53**, 355002 (2020). <https://doi.org/10.1088/1751-8121/ab9f72>
- [44] I. Goldhirsch, Rapid granular flows. Annu. Rev. Fluid Mech. **35**, 267–293 (2003). <https://doi.org/10.1146/annurev.fluid.35.101101.161114>
- [45] J.T. Jenkins, M.W. Richman, Kinetic theory for plane flows of a dense gas of identical, rough, inelastic, circular disks. Phys. Fluids **28**, 3485–3494 (1985). <https://doi.org/10.1063/1.865302>
- [46] C.K.K. Lun, S.B. Savage, A simple kinetic theory for granular flow of rough, inelastic, spherical particles. J. Appl. Mech. **54**, 47–53 (1987). <https://doi.org/10.1115/1.3172993>
- [47] C.S. Campbell, The stress tensor for simple shear flows of a granular material. J. Fluid Mech. **203**, 449–473 (1989). <https://doi.org/10.1017/S0022112089001540>
- [48] C.K.K. Lun, Kinetic theory for granular flow of dense, slightly inelastic, slightly rough spheres. J. Fluid Mech. **233**, 539–559 (1991). <https://doi.org/10.1017/S0022112091000599>
- [49] C.K.K. Lun, A.A. Bent, Numerical simulation of inelastic frictional spheres in simple shear flow. J. Fluid Mech. **258**, 335–353 (1994). <https://doi.org/10.1017/S0022112094003356>
- [50] A. Goldshtain, M. Shapiro, Mechanics of collisional motion of granular materials. Part 1. General hydrodynamic equations. J. Fluid Mech. **282**, 75–114 (1995). <https://doi.org/10.1017/S0022112095000048>
- [51] S. Luding, Granular materials under vibration: Simulations of rotating spheres. Phys. Rev. E **52**, 4442–4457 (1995). <https://doi.org/10.1103/PhysRevE.52.4442>
- [52] C.K.K. Lun, Granular dynamics of inelastic spheres in Couette flow. Phys. Fluids **8**, 2868–2883 (1996). <https://doi.org/10.1063/1.869068>
- [53] M. Huthmann, A. Zippelius, Dynamics of inelastically colliding rough spheres: Relaxation of translational and rotational energy. Phys. Rev. E **56**, R6275–R6278 (1997). <https://doi.org/10.1103/PhysRevE.56.R6275>
- [54] P. Zamankhan, H.V. Tafreshi, W. Polashenski, P. Sarkomaa, C.L. Hyndman, Shear induced diffusive mixing in simulations of dense Couette flow of rough, inelastic hard spheres. J. Chem. Phys. **109**, 4487–4491 (1998). <https://doi.org/10.1063/1.477076>
- [55] S. McNamara, S. Luding, Energy nonequipartition in systems of inelastic, rough spheres. Phys. Rev. E **58**, 2247–2250 (1998). <https://doi.org/10.1103/PhysRevE.58.2247>

[1103/PhysRevE.58.2247](https://doi.org/10.1103/PhysRevE.58.2247)

- [56] S. Luding, M. Huthmann, S. McNamara, A. Zippelius, Homogeneous cooling of rough, dissipative particles: Theory and simulations. *Phys. Rev. E* **58**, 3416–3425 (1998). <https://doi.org/10.1103/PhysRevE.58.3416>
- [57] O. Herbst, M. Huthmann, A. Zippelius, Dynamics of inelastically colliding spheres with Coulomb friction: relaxation of translational and rotational energy. *Granul. Matter* **2**, 211–219 (2000). <https://doi.org/10.1007/PL00010915>
- [58] T. Aspelmeier, M. Huthmann, A. Zippelius, in *Granular Gases, Lectures Notes in Physics*, vol. 564, ed. by T. Pöschel, S. Luding (Springer, Berlin, 2001), pp. 31–58
- [59] J.T. Jenkins, C. Zhang, Kinetic theory for identical, frictional, nearly elastic spheres. *Phys. Fluids* **14**, 1228–1235 (2002). <https://doi.org/10.1063/1.1449466>
- [60] W. Polashenski, P. Zamankhan, S. Mäkiharju, P. Zamankhan, Fine structures in sheared granular flows. *Phys. Rev. E* **66**, 021303 (2002). <https://doi.org/10.1103/PhysRevE.66.021303>
- [61] R. Cafiero, S. Luding, H.J. Herrmann, Rotationally driven gas of inelastic rough spheres. *Europhys. Lett.* **60**, 854–860 (2002). <https://doi.org/10.1209/epl/i2002-00295-7>
- [62] N. Mitarai, H. Hayakawa, H. Nakanishi, Collisional granular flow as a micropolar fluid. *Phys. Rev. Lett.* **88**, 174301 (2002). <https://doi.org/10.1103/PhysRevLett.88.174301>
- [63] P. Viot, J. Talbot, Thermalization of an anisotropic granular particle. *Phys. Rev. E* **69**, 051106 (2004). <https://doi.org/10.1103/PhysRevE.69.051106>
- [64] I. Goldhirsch, S.H. Noskowicz, O. Bar-Lev, Nearly smooth granular gases. *Phys. Rev. Lett.* **95**, 068002 (2005). <https://doi.org/10.1103/PhysRevLett.95.068002>
- [65] I. Goldhirsch, S.H. Noskowicz, O. Bar-Lev, Hydrodynamics of nearly smooth granular gases. *J. Phys. Chem. B* **109**, 21,449–21,470 (2005). <https://doi.org/10.1021/jp0532667>
- [66] A. Zippelius, Granular gases. *Physica A* **369**, 143–158 (2006). <https://doi.org/10.1016/j.physa.2006.04.012>

- [67] N.V. Brilliantov, T. Pöschel, W.T. Kranz, A. Zippelius, Translations and rotations are correlated in granular gases. *Phys. Rev. Lett.* **98**, 128001 (2007). <https://doi.org/10.1103/PhysRevLett.98.128001>
- [68] J. Piasecki, J. Talbot, P. Viot, Angular velocity distribution of a granular planar rotator in a thermalized bath. *Phys. Rev. E* **75**, 051307 (2007). <https://doi.org/10.1103/PhysRevE.75.051307>
- [69] F. Cornu, J. Piasecki, Granular rough sphere in a low-density thermal bath. *Physica A* **387**, 4856–4862 (2008). <https://doi.org/10.1016/j.physa.2008.03.014>
- [70] A. Santos, G.M. Kremer, V. Garzó, Energy production rates in fluid mixtures of inelastic rough hard spheres. *Prog. Theor. Phys. Suppl.* **184**, 31–48 (2010). <https://doi.org/10.1143/PTPS.184.31>
- [71] A. Santos, A Bhatnagar–Gross–Krook-like model kinetic equation for a granular gas of inelastic rough hard spheres. *AIP Conf. Proc.* **1333**, 41–48 (2011). <https://doi.org/10.1063/1.3562623>
- [72] A. Santos, Homogeneous free cooling state in binary granular fluids of inelastic rough hard spheres. *AIP Conf. Proc.* **1333**, 128–133 (2011). <https://doi.org/10.1063/1.3562637>
- [73] F. Vega Reyes, A. Santos, G.M. Kremer, Role of roughness on the hydrodynamic homogeneous base state of inelastic spheres. *Phys. Rev. E* **89**, 020202(R) (2014). <https://doi.org/10.1103/PhysRevE.89.020202>
- [74] F. Vega Reyes, A. Santos, G.M. Kremer, Properties of the homogeneous cooling state of a gas of inelastic rough particles. *AIP Conf. Proc.* **1628**, 494–501 (2014). <https://doi.org/10.1063/1.4902634>
- [75] G.M. Kremer, A. Santos, V. Garzó, Transport coefficients of a granular gas of inelastic rough hard spheres. *Phys. Rev. E* **90**, 022205 (2014). <https://doi.org/10.1103/PhysRevE.90.022205>
- [76] F. Vega Reyes, A. Santos, Steady state in a gas of inelastic rough spheres heated by a uniform stochastic force. *Phys. Fluids* **27**, 113301 (2015). <https://doi.org/10.1063/1.4934727>
- [77] F. Vega Reyes, A. Lasanta, A. Santos, V. Garzó, Thermal properties of an impurity immersed in a granular gas of rough hard spheres. *EPJ Web Conf.* **140**, 04003 (2017). <https://doi.org/10.1051/epjconf/201714004003>
- [78] F. Vega Reyes, A. Lasanta, A. Santos, V. Garzó, Energy nonequipartition in gas mixtures of inelastic rough hard spheres: The tracer limit. *Phys. Rev. E* **96**, 052901 (2017). <https://doi.org/10.1103/PhysRevE.96.052901>.

Erratum: **100**, 2274–2275 (2012)

- [79] V. Garzó, A. Santos, G.M. Kremer, Impact of roughness on the instability of a free-cooling granular gas. *Phys. Rev. E* **97**, 052901 (2018). <https://doi.org/10.1103/PhysRevE.97.052901>
- [80] A. Santos, Interplay between polydispersity, inelasticity, and roughness in the freely cooling regime of hard-disk granular gases. *Phys. Rev. E* **98**, 012904 (2018). <https://doi.org/10.1103/PhysRevE.98.012904>
- [81] A. Megías, A. Santos, Driven and undriven states of multicomponent granular gases of inelastic and rough hard disks or spheres. *Granul. Matter* **21**, 49 (2019). <https://doi.org/10.1007/s10035-019-0901-y>
- [82] A. Megías, A. Santos, Energy production rates of multicomponent granular gases of rough particles. A unified view of hard-disk and hard-sphere systems. *AIP Conf. Proc.* **2132**, 080003 (2019). <https://doi.org/10.1063/1.5119584>
- [83] A. Torrente, M.A. López-Castaño, A. Lasanta, F. Vega Reyes, A. Prados, A. Santos, Large Mpemba-like effect in a gas of inelastic rough hard spheres. *Phys. Rev. E* **99**, 060901(R) (2019). <https://doi.org/10.1103/PhysRevLett.119.148001>
- [84] R. Gómez González, V. Garzó, Non-Newtonian rheology in inertial suspensions of inelastic rough hard spheres under simple shear flow. *Phys. Fluids* **32**, 073315 (2020). <https://doi.org/10.1063/5.0015241>
- [85] A. Megías, A. Santos, Hydrodynamics of granular gases of inelastic and rough hard disks or spheres. I. Transport coefficients. *Phys. Rev. E* **104**, 034901 (2021). <https://doi.org/10.1103/PhysRevE.104.034901>
- [86] A. Megías, A. Santos, Hydrodynamics of granular gases of inelastic and rough hard disks or spheres. II. Stability analysis. *Phys. Rev. E* **104**, 034902 (2021). <https://doi.org/10.1103/PhysRevE.104.034902>
- [87] F.B. Pidduck, The kinetic theory of a special type of rigid molecule. *Proc. R. Soc. Lond. A* **101**, 101–112 (1922). <https://doi.org/10.1098/rspa.1922.0028>
- [88] A. Santos, G.M. Kremer, M. dos Santos, Sonine approximation for collisional moments of granular gases of inelastic rough spheres. *Phys. Fluids* **23**, 030604 (2011). <https://doi.org/10.1063/1.3558876>
- [89] M.H. Ernst, R. Brito, Driven inelastic Maxwell models with high energy tails. *Phys. Rev. E* **65**, 040301(R) (2002). <https://doi.org/10.1103/PhysRevE.65.040301>

- [90] W.T. Kranz, N.V. Brilliantov, T. Pöschel, A. Zippelius, Correlation of spin and velocity in the homogeneous cooling state of a granular gas of rough particles. *Eur. Phys. J. Spec. Top.* **179**, 91–111 (2009). <https://doi.org/10.1140/epjst/e2010-01196-0>
- [91] R. Rongali, M. Alam, Higher-order effects on orientational correlation and relaxation dynamics in homogeneous cooling of a rough granular gas. *Phys. Rev. E* **89**, 062201 (2014). <https://doi.org/10.1103/PhysRevE.89.062201>
- [92] For an interactive animation, see A. Santos, “Inelastic Collisions of Two Rough Spheres”, <http://demonstrations.wolfram.com/InelasticCollisionsOfTwoRoughSpheres/>, Wolfram Demonstrations Project (2010).