

PERTURBATION ANALYSIS OF RANDOMIZED SVD AND ITS APPLICATIONS TO HIGH-DIMENSIONAL STATISTICS

BY YICHI ZHANG AND MINH TANG

Department of Statistics, North Carolina State University, yzhan239@ncsu.edu; mtang8@ncsu.edu

Randomized singular value decomposition (RSVD) is a class of computationally efficient algorithms for computing the truncated SVD of large data matrices. Given an $n \times n$ symmetric matrix \mathbf{M} , the prototypical RSVD algorithm outputs an approximation of the k leading singular vectors of \mathbf{M} by computing the SVD of $\mathbf{M}^g \mathbf{G}$; here $g \geq 1$ is an integer and $\mathbf{G} \in \mathbb{R}^{n \times k}$ is a random Gaussian sketching matrix. In this paper we study the statistical properties of RSVD under a general “signal-plus-noise” framework, i.e., the observed matrix $\hat{\mathbf{M}}$ is assumed to be an additive perturbation of some true but unknown signal matrix \mathbf{M} . We first derive upper bounds for the ℓ_2 and $\ell_{2 \rightarrow \infty}$ distances between the *approximate* singular vectors of $\hat{\mathbf{M}}$ and the true singular vectors of \mathbf{M} . These upper bounds depend on the signal-to-noise ratio (SNR) and the number of power iterations g . A phase transition phenomenon is observed in which a smaller SNR requires larger values of g to guarantee convergence of the ℓ_2 and $\ell_{2 \rightarrow \infty}$ distances. We also show that the thresholds for g where these phase transitions occur are sharp whenever the noise matrices satisfy a certain trace growth condition. Finally, we derive normal approximations for the row-wise fluctuations of these approximate singular vectors and the entrywise fluctuations when $\hat{\mathbf{M}}$ is projected onto these vectors. We illustrate our theoretical results by deriving nearly-optimal performance guarantees for RSVD when applied to three statistical inference problems, namely, community detection, matrix completion, and PCA with missing data.

CONTENTS

1	Introduction	2
2	Background and settings	4
	2.1 Notation	4
	2.2 Signal-plus-noise perturbation framework	5
	2.3 Randomized SVD with repeated sampling	6
3	Theoretical results	7
	3.1 ℓ_2 error bound	9
	3.2 $\ell_{2, \infty}$ norm error bound	10
	3.3 Comparison with existing results	13
	3.4 Row-wise limiting distribution	14
4	Random graph inference	15
	4.1 Subspace perturbation error bound	16
	4.2 Lower bound and phase transition sharpness	17
	4.3 Exact recovery for stochastic blockmodels	18
	4.4 Normal approximation for $\hat{\mathbf{U}}_g$	20
5	Additional applications	21
	5.1 Matrix completion with noises	21
	5.2 PCA with missing data	25
6	Discussions	27
	Supplementary Material	27
	References	28

S.1 Numerical experiments	33
S.1.1 RSVD-based spectral clustering for SBM	33
S.1.1.1 Phase transition	33
S.1.1.2 Exact recovery	34
S.1.1.3 Limiting distribution	35
S.1.2 PCA with missing data	35
S.1.3 Distance matrix completion	38
S.2 Extension to asymmetric matrices	40
S.3 General theoretical results for any \mathbf{G}	42
S.4 Technical lemmas	43
S.5 Proofs for Section 3	49
S.5.1 Proof of Theorem S.3.1	49
S.5.2 Proof of Theorem S.3.2	50
S.5.3 Proof of Theorem S.3.3	51
S.5.4 Proof of Theorem 3.1	53
S.5.5 Proof of Theorem 3.2	54
S.5.6 Proof of Corollary 3.1	54
S.5.7 Proof of Theorem 3.3	54
S.5.8 Proof of Corollary 3.2	55
S.5.9 Proof of Theorem 3.4	58
S.6 Proofs for Section 4, Section 5, and Section S.2	61
S.6.1 Proof of Theorem 4.1	61
S.6.2 Proof of Theorem 4.2	65
S.6.3 Proof of Theorem 5.1	66
S.6.3.1 Preliminary	66
S.6.3.2 Assumption verification	66
S.6.3.3 Bounding $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$	67
S.6.3.4 Bounding $\ \hat{\mathbf{T}}_g - \mathbf{T}\ _F$	68
S.6.3.5 Bounding $\ \hat{\mathbf{T}}_g - \mathbf{T}\ _{\max}$	68
S.6.3.6 Entrywise limiting distribution and confidence interval	68
S.6.4 Proof of Theorem 5.2	74
S.6.5 Proof of Theorem S.6.1	75
S.6.6 Proof of Theorem S.2.1	78
S.7 Additional Proofs	80
S.7.1 Proof of Proposition 3.1	80
S.7.2 Proof of Proposition 3.2	80
S.7.3 Proof of Claim S.1	86
S.7.4 Proof of Claim S.2	87
S.7.5 Proof of Claim S.3	88
S.7.6 Proof of Claim S.4	88
S.7.7 Proof of Claim S.5	89

1. Introduction. Spectral methods are popular in statistics and machine learning as they provide computationally efficient algorithms with strong theoretical guarantees for a diverse number of inference problems including network analysis [91, 99, 113], matrix completion and denoising [4, 23], covariance estimation/PCA [50, 66], non-linear dimension reduction and manifold learning [12, 32, 114], ranking [26, 27], etc. A common unifying theme for many spectral algorithms is, given a data matrix $\hat{\mathbf{M}}$, first compute a factorization of $\hat{\mathbf{M}}$ using the ubiquitous singular value decomposition (SVD), then truncate this SVD to keep only the $k < \min\{m, n\}$ leading singular values and singular vectors, and finally perform inference

using the truncated SVD representation. The value k is usually chosen to be as small as possible while still preserving most of the information in $\hat{\mathbf{M}}$.

Classical algorithms for SVD, such as those based on pivotal QR decompositions and/or Householder transformations, require $\mathcal{O}(m^2n)$ floating-point operations (flops) assuming $m \leq n$ and return the full set of m singular values and vectors, even when only the leading $k < m$ of them are desired; see Sections 5.4 and 8.3 of [55]. The $\mathcal{O}(m^2n)$ flops is a severe computational burden when m is large. These algorithms also require random access to the entries of $\hat{\mathbf{M}}$ and are thus inefficient when $\hat{\mathbf{M}}$ is too large to store in RAM. These computation and memory issues limit the use of classical SVD in modern data applications.

Recently in the numerical linear algebra community, randomized SVD (RSVD) [58, 89, 101, 116, 117] had been widely studied with the aim of providing fast, memory efficient, and accurate approximations for the truncated SVD of large data matrices. In particular, suppose $\hat{\mathbf{M}}$ is an $m \times n$ matrix and we are interested in finding the k leading (left) singular vectors of $\hat{\mathbf{M}}$ for some $k \ll n$. The prototypical RSVD algorithm [58] first sketches $\hat{\mathbf{M}}$ into a smaller matrix $\mathbf{Y} = (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} \in \mathbb{R}^{m \times \tilde{k}}$ where $g \geq 1$ and $\tilde{k} \geq k$ are user-specified parameters and $\mathbf{G} \in \mathbb{R}^{n \times \tilde{k}}$ is a random matrix, and then uses the k leading left singular vectors of \mathbf{Y} as an approximation to the k leading left singular vectors of $\hat{\mathbf{M}}$. The parameter g is usually chosen to be a small integer and \tilde{k} is chosen to be equal to or slightly larger than k . There are numerous choices for \mathbf{G} including Gaussian matrices, random orthogonal matrices, Rademacher matrices, and column-sampling matrices; see [68, 83, 124] and the references therein.

The *sketch-and-solve* strategy of RSVD has several important practical advantages. Firstly, it has a computational complexity of $\mathcal{O}(mnk)$ flops and this can be further reduced to $\mathcal{O}(mn \log k)$ flops when the sketching matrix \mathbf{G} is structured. These $\mathcal{O}(mnk)$ or $\mathcal{O}(mn \log k)$ flops are also very fast as the main operations are the matrix-matrix products $\hat{\mathbf{M}}\mathbf{X}$ and $\hat{\mathbf{M}}^\top \mathbf{X}$ where \mathbf{X} is either a $m \times \tilde{k}$ or $n \times \tilde{k}$ matrix; matrix products are highly-optimized on almost all computing platforms. Secondly, it is “pass efficient” [41, 58] and only requires $(2g + 1)$ passes through the data; this dramatically reduces memory storage [55, 81]. Thirdly, RSVD also allows for data compression [33] and can be adapted to a streaming setting [116]. Many recent works have discussed replacing the classical SVD with RSVD, see [9, 36, 49, 62, 78, 95, 97, 118] for examples in covariance matrix estimation, matrix completion, network embeddings, and dynamic mode decompositions.

Existing theoretical results for RSVD, such as those in [58, 59, 81, 101, 106], had mainly focused on the setting where $\hat{\mathbf{M}}$ is assumed to be noise-free; these results either bound the approximation error between $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{U}}$ or bound the difference between the low-rank approximation error of $\|(\mathbf{I} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \hat{\mathbf{M}}\|$ and that of $\|(\mathbf{I} - \hat{\mathbf{U}} \hat{\mathbf{U}}^\top) \hat{\mathbf{M}}\|$. Here $\hat{\mathbf{U}}$ denote the true leading singular vectors of $\hat{\mathbf{M}}$ and $\|\cdot\|$ denote some unitarily invariant norm. In particular these approximation error decreases if g (the number of power iterations) and/or \tilde{k} (the sketching dimensions) increases. However, for many inference problem, the observed matrix $\hat{\mathbf{M}}$ is also noisy due to sampling and/or perturbation errors. More specifically it is often the case that $\hat{\mathbf{M}}$ is generated from a “signal-plus-noise” model $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} is assumed to be the underlying true signal matrix of $\hat{\mathbf{M}}$ with certain structure e.g., (approximately) low rank and/or sparse, and \mathbf{E} is the unobservable perturbation noise. As $\hat{\mathbf{M}}$ is a noisy realization of \mathbf{M} , $\hat{\mathbf{U}}$ is a noisy estimate of \mathbf{U} , the leading singular vectors of \mathbf{M} . Under this perspective, the main aim is now to understand the relationship between $\hat{\mathbf{U}}_g$ and \mathbf{U} , and we thus need to balance between the approximation error of $\hat{\mathbf{U}}_g$ and the estimation error of $\hat{\mathbf{U}}$, i.e., it is neither necessary nor beneficial to make the approximation error between $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{U}}$ much smaller than the estimation error between $\hat{\mathbf{U}}$ and \mathbf{U} .

Estimation errors between $\hat{\mathbf{U}}$ and \mathbf{U} is a fundamental topic in matrix perturbation theory. Classic results in matrix perturbation include Weyl’s and Lidskii’s inequalities [80, 123] for

eigenvalues and singular values, and the Davis–Kahan Theorem and Wedin’s Theorem [38, 122, 129] for subspaces; these results make minimal assumptions on the perturbation matrices \mathbf{E} . The last decade has witnessed further study of matrix perturbations from more statistical perspectives through the introduction of additional minor assumptions on \mathbf{E} and \mathbf{M} such as the entries of \mathbf{E} are independent random variables and/or the leading singular vectors of \mathbf{M} has bounded coherence. Examples include more refined matrix concentration inequalities [5, 93, 104, 115], rate-optimal unilateral singular subspace perturbation bound [17], and $\ell_{2 \rightarrow \infty}$ singular subspace perturbation bounds [3, 21, 22, 46, 51, 75, 84]. These developments in turns lead to stronger statistical guarantees for many spectral methods; see [28] for a recent and comprehensive survey.

In this paper, we will study the perturbation error between $\hat{\mathbf{U}}_g$ and \mathbf{U} under the “signal-plus-noise” model described above. In particular we analyze a repeated sampling version of RSVD (namely, *rs*-RSVD) with $\mathcal{O}(a_n g m n)$ complexity where a_n is the number of random sketches and g is the number of power iterations of $\hat{\mathbf{M}}$; see Section 2. The choice $a_n = 1$ corresponds to the `PowerRangeFinder` and `SubspacePowerMethod` described in [58, 89, 124]. Theoretical results for *rs*-RSVD, under a set of mild and typically seen assumptions for \mathbf{E} , are presented in Section 3. These results include perturbation bounds for the difference in either ℓ_2 or $\ell_{2 \rightarrow \infty}$ norm between the approximate singular vectors $\hat{\mathbf{U}}_g$ of $\hat{\mathbf{M}}$ and the true singular vectors \mathbf{U} of \mathbf{M} , normal approximations for the row-wise fluctuations of $\hat{\mathbf{U}}_g$, and entrywise concentration and normal approximations for $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M}$. The ℓ_2 and $\ell_{2 \rightarrow \infty}$ bounds exhibit a phase-transition phenomenon in that as the signal-to-noise ratio (SNR) decreases, the number of power iterations g need to increase to guarantee fast convergence rates; the phase transition thresholds are also sharp provided that the noise matrices \mathbf{E} satisfy a certain trace growth conditions. We then apply our theoretical results to three different statistical inference problems: community detection, matrix completion, and PCA with missing data. For all three problems we show that the approximate singular vectors $\hat{\mathbf{U}}_g$ achieve the same or nearly the same theoretical guarantees as that for the true singular vectors $\hat{\mathbf{U}}$; see Section 4 and Section 5. Our results thus provide a bridge between the numerical linear algebra and statistics communities. For conciseness and ease of exposition we only present results for symmetric $\hat{\mathbf{M}}$ and \mathbf{M} in the main paper. Extension of these results to the case of asymmetric $\hat{\mathbf{M}}$ and \mathbf{M} are provided in Section S.2 of the supplementary document. Numerical experiments and detailed proofs for the main paper are also included in the supplementary document.

2. Background and settings.

2.1. *Notation.* Let a be a positive integer. We then write $[a]$ to denote the set $\{1, \dots, a\}$. For two sequences $\{a_n\}_{n \geq 1}$ and $\{b_n\}_{n \geq 1}$, we write $a_n \lesssim b_n$ if there exists a constant c not depending on n such that $a_n \leq c b_n$ for all but finitely many $n \geq 1$; we write $a_n \gtrsim b_n$ if $b_n \lesssim a_n$ and we write $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $a_n \gtrsim b_n$.

The set of $d \times d'$ matrices with orthonormal columns is denoted as $\mathbb{O}_{d \times d'}$ when $d \neq d'$ and is denoted as \mathbb{O}_d otherwise. Let \mathbf{N}_1 and \mathbf{N}_2 be symmetric matrices. We write $\mathbf{N}_1 \succeq \mathbf{N}_2$ if $\mathbf{N}_1 - \mathbf{N}_2$ is positive semidefinite and we write $\mathbf{N}_1 \succ \mathbf{N}_2$ if $\mathbf{N}_1 - \mathbf{N}_2$ is positive definite. Let \mathbf{N} be an arbitrary matrix. We denote the i th row of \mathbf{N} by $[\mathbf{N}]_i$, and the ij th entry of \mathbf{N} by $[\mathbf{N}]_{ij}$. We write $\text{tr } \mathbf{N}$ and $\text{rk } \mathbf{N}$ to denote the trace and rank of a square matrix \mathbf{N} , respectively. The spectral and Frobenius norm of \mathbf{N} are denoted as $\|\mathbf{N}\|$ and $\|\mathbf{N}\|_F$, respectively. The maximum (in modulus) of the entries of \mathbf{N} is denoted as $\|\mathbf{N}\|_{\max}$. In addition we denote the $2 \rightarrow \infty$ norm of \mathbf{N} by

$$\|\mathbf{N}\|_{2 \rightarrow \infty} = \max_{\|\mathbf{x}\|=1} \|\mathbf{N}\mathbf{x}\|_\infty = \max_i \|[\mathbf{N}]_i\|,$$

i.e., $\|\mathbf{N}\|_{2 \rightarrow \infty}$ is the maximum of the ℓ_2 norms of the rows of \mathbf{N} . We have the relationships

$$n^{-1/2}\|\mathbf{N}\| \leq \|\mathbf{N}\|_{2 \rightarrow \infty} \leq \|\mathbf{N}\|, \quad \|\mathbf{N}\|_{\max} \leq \|\mathbf{N}\|_{2 \rightarrow \infty} \leq d^{1/2}\|\mathbf{N}\|_{\max},$$

where n and d are the number of rows and columns of \mathbf{N} , respectively. Given two matrices $\mathbf{U}_1 \in \mathbb{O}_{n \times d}$ and $\mathbf{U}_2 \in \mathbb{O}_{n \times d}$, we define their ℓ_2 and $\ell_{2 \rightarrow \infty}$ distances as

$$\begin{aligned} d_2(\mathbf{U}_1, \mathbf{U}_2) &:= \inf_{\mathbf{W} \in \mathbb{O}_d} \|\mathbf{U}_1 - \mathbf{U}_2 \mathbf{W}\|, \\ d_{2 \rightarrow \infty}(\mathbf{U}_1, \mathbf{U}_2) &:= \inf_{\mathbf{W} \in \mathbb{O}_d} \|\mathbf{U}_1 - \mathbf{U}_2 \mathbf{W}\|_{2 \rightarrow \infty}. \end{aligned}$$

Recall that $d_2(\mathbf{U}_1, \mathbf{U}_2) \leq \sqrt{2}\|\sin \Theta(\mathbf{U}_1, \mathbf{U}_2)\|$ where $\Theta(\mathbf{U}_1, \mathbf{U}_2)$ are the *principal* angles between \mathbf{U}_1 and \mathbf{U}_2 . Furthermore, by the relationship between the spectral and $\ell_{2 \rightarrow \infty}$ norms, $d_{2 \rightarrow \infty}$ yields finer and more uniform control of the row-wise differences (up to orthogonal transformations) between \mathbf{U}_1 and \mathbf{U}_2 . See [3, 22, 28, 34, 75] for further discussions of the $d_{2 \rightarrow \infty}$ and its uses in matrix perturbation inequalities and statistics applications.

2.2. Signal-plus-noise perturbation framework. For ease of exposition we restrict ourselves to the case when the observed matrices are symmetric in the main paper. The general case of rectangular and/or asymmetric matrices are presented in Section S.2 of the supplementary document. For a $n \times n$ symmetric matrix $\mathbf{M} = [m_{ij}]$ with $\text{rk}(\mathbf{M}) = k^*$, we consider the additive perturbation

$$\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E},$$

where \mathbf{E} is a $n \times n$ symmetric *noise* matrix; we shall generally assume that \mathbf{M} and \mathbf{E} are *unobserved*, i.e., we observed only $\hat{\mathbf{M}}$. Denote the eigendecompositions of $\hat{\mathbf{M}}$ and \mathbf{M} by

$$\begin{aligned} (2.1) \quad \mathbf{M} &:= (\mathbf{U} \ \mathbf{U}_\perp) \begin{pmatrix} \mathbf{\Lambda} & \mathbf{0} \\ \mathbf{0} & \mathbf{\Lambda}_\perp \end{pmatrix} \begin{pmatrix} \mathbf{U}^\top \\ \mathbf{U}_\perp^\top \end{pmatrix}, \\ \hat{\mathbf{M}} &:= (\hat{\mathbf{U}} \ \hat{\mathbf{U}}_\perp) \begin{pmatrix} \hat{\mathbf{\Lambda}} & \mathbf{0} \\ \mathbf{0} & \hat{\mathbf{\Lambda}}_\perp \end{pmatrix} \begin{pmatrix} \hat{\mathbf{U}}^\top \\ \hat{\mathbf{U}}_\perp^\top \end{pmatrix}, \end{aligned}$$

where $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_{k^*})$ and $\hat{\mathbf{\Lambda}} = \text{diag}(\hat{\lambda}_1, \dots, \hat{\lambda}_{k^*})$ are $k^* \times k^*$ diagonal matrices containing the k^* largest (in magnitude) nonzero eigenvalues of \mathbf{M} and $\hat{\mathbf{M}}$, respectively. Let $\psi := |\lambda_1/\lambda_{k^*}|$ denote the condition number for the first k^* eigenvalues of \mathbf{M} .

We focus mainly on the setting where \mathbf{M} has an approximately low-rank structure, i.e., both k^* and ψ are either bounded or slowly diverging; we say that a quantity z depending on n is slowly diverging if $|z| = \mathcal{O}(\log n)$ as $n \rightarrow \infty$. These low-rank assumptions appear frequently in many inference problems for high-dimensional matrix-valued data including community detection in graphs, matrix completion, and covariance matrix estimation; see e.g., [3, 16, 21, 46, 51, 88] among others. We now make an assumption on $\|\mathbf{E}\|$.

ASSUMPTION 1 (Signal-to-Noise Ratio). Let $S_n = \|\mathbf{M}\|$. There exists fixed but arbitrary constants $\beta_\Delta > 0$, $N_0 > 0$ and a quantity E_n depending on n such that $S_n/E_n \gtrsim n^{\beta_\Delta}$ and

$$(2.2) \quad \mathbb{P}[\|\mathbf{E}\| \leq E_n] \geq 1 - n^{-6},$$

whenever $n \geq N_0$. We view E_n as the noise level of \mathbf{E} and n^{β_Δ} as a lower bound on the SNR.

Assumption 1 provides a concentration inequality for $\|\mathbf{E}\|$ in terms of $\|\mathbf{M}\|$ and the signal to noise ratio (SNR) n^{β_Δ} . As $\beta_\Delta > 0$, Assumption 1 guarantees that $S_n \gg E_n$ and hence, by Weyl's inequality, there exists with high probability a sufficiently large gap between the

leading eigenvalues of $\hat{\mathbf{M}}$ (as induced by the signal \mathbf{M}) and the remaining eigenvalues of $\hat{\mathbf{M}}$ (as induced by the noise \mathbf{E}). This allows us to relate the singular vectors of the sketched matrix $\hat{\mathbf{M}}^g \mathbf{G}$ to the leading singular vectors of \mathbf{M} as g increases. We show in Section 3 that, as the SNR decreases, we need larger values of g to accurately recover the singular subspace of \mathbf{M} from that of $\hat{\mathbf{M}}^g \mathbf{G}$.

We emphasize that Assumption 1 generally holds if \mathbf{E} is a symmetric matrix whose upper triangular entries are independent mean 0 random variables. Examples include the case where the entries of \mathbf{E} are either Gaussian random variables with variances bounded by $n^{-(1+2\beta_\Delta)} S_n^2$ or bounded random variables with variances bounded by $n^{-(1+2\beta_\Delta)} S_n^2$; see e.g., Corollary 3.9 in [11] or Theorem 1.4 in [115] for a justification of these claims. Furthermore the lower bound $1 - n^{-6}$ in Eq. (2.2) is somewhat arbitrary and is chosen mainly for convenience. Indeed, the above cited results also show that for any constant $c > 0$ there exists a constant $C > 0$ depending only on c such that $\|\mathbf{E}\| \leq CE_n$ with probability at least $1 - n^{-c}$. This constant C can then be subsumed into the definition of E_n without changing the SNR n^{β_Δ} . Assumption 1 can also hold when the entries of \mathbf{E} are not mutually independent. See for example the presentation of PCA with missing data in Section 5.2, i.e., the matrix \mathbf{E} in Section 5.2 is of the form $\mathbf{E} = \mathbf{Z}\mathbf{Z}^\top - \mathbf{M}$ where \mathbf{M} is a deterministic symmetric matrix and \mathbf{Z} is a $d \times m$ matrix whose entries are independent random variables. The product $\mathbf{Z}\mathbf{Z}^\top$ creates dependence between the upper triangular entries of \mathbf{E} .

Finally we note that the noise level E_n in Assumption 1 is possibly decoupled from the signal strength S_n . This is intentional as it allows us to obtain more general theoretical results that can then be applied in diverse settings. For example in Section 4 we consider the random graphs setting wherein the entries of \mathbf{M} are binary random variables with $\mathbb{E}[\hat{m}_{ij}] = m_{ij}$. The distribution of \mathbf{E} therefore depends only on \mathbf{M} and thus E_n is also a function of S_n . In particular $n^{\beta_\Delta} \asymp E_n \asymp S_n^{1/2}$ and so as the signal level S_n decreases (for example either by decreasing the number of non-zero entries and/or their magnitudes) the SNR also decreases and inference using $\hat{\mathbf{M}}$ becomes harder. In contrast, for the problems of matrix completion and PCA with missing data discussed in Section 5, the distribution for \mathbf{E} depends on other parameters distinct from \mathbf{M} and it is thus possible to reduce both S_n and E_n without changing the SNR, thereby not affecting the general behavior of $\hat{\mathbf{M}}$.

2.3. Randomized SVD with repeated sampling. Given $\hat{\mathbf{M}}$ we wish to find the eigenvectors $\hat{\mathbf{U}}$ associated with the k^* largest eigenvalues (in magnitude) of $\hat{\mathbf{M}}$. One popular and widely used approach for finding $\hat{\mathbf{U}}$ is via randomized subspace iteration. More specifically we first sample an $n \times \tilde{k}$ matrix \mathbf{G} whose entries are iid standard normals and compute $\mathbf{Y} = \hat{\mathbf{M}}^g \mathbf{G}$ for some positive integer $g \geq 1$. Now let $\hat{\mathbf{U}}_g$ be the $n \times k^*$ matrix whose columns form an orthonormal basis for the k^* leading left singular vectors of \mathbf{Y} . The matrix $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$ is a low rank approximation to $\hat{\mathbf{M}}$ and we can take $\hat{\mathbf{U}}_g$ as an approximation to $\hat{\mathbf{U}}$. We note that \tilde{k} , the number of columns of \mathbf{G} , is often chosen to be slightly larger than k^* in order to increase the probability that the column space of \mathbf{Y} also include the column space of $\hat{\mathbf{U}}$; empirical observations suggest that $\tilde{k} = k^* + 5$ or $\tilde{k} = k^* + 10$ is sufficient for most practical applications [58, Section 1.3]. For more discussion on randomized subspace iteration, see Section 4.5 of [58], Section 11.6 of [86], Section 4.3 of [124], and [89]. Recently, [81] has purposed a data-driven bootstrap algorithm to estimate the approximation error of sketching SVD, which might also be adapted to select \tilde{k} for the general RSVD in practice.

This paper considers a variant of the above procedure in which we sample independent realizations $\mathbf{G}_1, \mathbf{G}_2, \dots, \mathbf{G}_{a_n}$ and choose the \mathbf{G} which maximizes $\sigma_k(\hat{\mathbf{M}}^g \mathbf{G}_a)$; here $\sigma_k(\cdot)$ denotes the k th largest singular value of a matrix. We term the resulting procedure as repeated-sampling randomized SVD (*rs*-RSVD); see Algorithm 1 for a formal description. The main

Algorithm 1: *rs*-RSVD

Input: $\hat{\mathbf{M}} \in \mathbb{R}^{n \times n}$, rank $k \geq 1$ and sketching dimension $\tilde{k} \geq k$. Integers $a_n, g \geq 1$.

1. Generate a_n iid standard Gaussian sketching matrices $\mathbf{G}_1, \dots, \mathbf{G}_{a_n} \in \mathbb{R}^{n \times \tilde{k}}$;
2. For $a \in [a_n]$, form $\hat{\mathbf{M}}^g \mathbf{G}_a$ by computing $\hat{\mathbf{M}} \mathbf{G}_a, \hat{\mathbf{M}}(\hat{\mathbf{M}} \mathbf{G}_a), \dots, \hat{\mathbf{M}}(\hat{\mathbf{M}}^{g-1} \mathbf{G}_a)$;
3. For $a \in [a_n]$, compute the *exact* SVD of $\hat{\mathbf{M}}^g \mathbf{G}_a$ and retain the k leading singular values $\{\sigma_i(\hat{\mathbf{M}}^g \mathbf{G}_a)\}_{i \leq k}$ of $\hat{\mathbf{M}}^g \mathbf{G}_a$ and the $n \times k$ matrix of corresponding left singular vectors $\hat{\mathbf{U}}_g^{(a)}$;
4. Choose $\tilde{a} = \operatorname{argmax}_{a \in [a_n]} \sigma_k(\hat{\mathbf{M}}^g \mathbf{G}_a)$ and $\mathbf{G} = \mathbf{G}_{\tilde{a}}$. Let $\hat{\mathbf{U}}_g = \hat{\mathbf{U}}_g^{(\tilde{a})}$;
5. Let $\tilde{\sigma}_1 \geq \dots \geq \tilde{\sigma}_k$ be the singular values of $\hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$.

Output: Estimated singular vectors $\hat{\mathbf{U}}_g$, singular values $\{\tilde{\sigma}_1, \dots, \tilde{\sigma}_k\}$, low-rank approximation $\tilde{\mathbf{M}} = \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$ or $\tilde{\mathbf{M}} = \frac{1}{2}(\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} + \hat{\mathbf{M}} \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top)$.

rationale for the repeated sampling step is that letting $a_n > 1$ allows us to choose $\tilde{k} = k^*$, which results in $\hat{\mathbf{M}}^g \mathbf{G}$ having the smallest number of columns while still obtaining an estimate $\hat{\mathbf{U}}_g$ of \mathbf{U} that is qualitatively similar to that when $\tilde{k} > k^*$; see the theoretical results in Section 3 and the numerical experiments in Sections S.1 in the Supplementary File for demonstrations of this claim. In addition, if $\hat{\mathbf{M}}$ arises from the additive model described in Section 2.2 then $\tilde{k} = k^*$ also leads to simpler theoretical results compared to $\tilde{k} > k^*$.

REMARK 2.1 (Implementation details for Algorithm 1). For ease of exposition we had written steps 1–3 of Algorithm 1 as if there were a_n different sketches $\{\hat{\mathbf{M}}^g \mathbf{G}_a\}_{a \leq a_n}$. As the computation of $\hat{\mathbf{M}}^g \mathbf{G}_a$ involves g passes through the data matrix $\hat{\mathbf{M}}$, having $a_n > 1$ is undesirable if we then require $a_n g$ passes, especially when the dimensions of $\hat{\mathbf{M}}$ are large. In practice we can combine all $a_n \geq 1$ different sketches into a single sketch, i.e., as the $\{\mathbf{G}_a\}$ are iid, we first generate a standard Gaussian matrix \mathbf{G}_* with n rows and $a_n \tilde{k}$ columns, compute a single sketched matrix $\hat{\mathbf{M}}^g \mathbf{G}_*$, and then form the $\{\hat{\mathbf{M}}^g \mathbf{G}_a\}_{a \leq a_n}$ by sequentially extracting (without replacement) k columns from $\hat{\mathbf{M}}^g \mathbf{G}_*$. The number of passes through the data is then still g . We can also replace Algorithm 1 with a more numerically stable albeit algebraically equivalent version wherein one periodically orthonormalizes $\hat{\mathbf{M}}^{g'} \mathbf{G}$ (using QR decomposition) for $g' < g$ before computing $\hat{\mathbf{M}}^{g'+1} \mathbf{G}$; see [58, Remark 4.3] and [85]. This extra orthonormalization step has no impact on the theoretical results presented subsequently.

REMARK 2.2 (Sketching with $g = 1$). If we set both $a_n = 1$ and $g = 1$ in Algorithm 1 then we get the “sketched SVD” algorithm described in [81, 83]. Sketched SVD is very useful when $\hat{\mathbf{M}}$ is too large to store in fast memory as the procedure only requires one pass through the data. However, as we will show in Section 4, setting $g = 1$ leads to poor estimates of \mathbf{U} using *rs*-RSVD unless $\tilde{k} \gg k^*$; indeed the theoretical analysis in this paper can be extended to show that if $g = 1$ then $\tilde{k} = \Omega(n)$ is possibly necessary to guarantee accurate estimation of \mathbf{U} using *rs*-RSVD; the choice $\tilde{k} = \Omega(n)$ has recently been considered by [126], which helps to establish precise asymptotically exact results in the context of sketching PCA. In practice, it is usually preferable to choose \tilde{k} as small as possible, thus we will not present theoretical results for the $g = 1, \tilde{k} = \Omega(n)$ regime as it detracts from the main message of current paper.

3. Theoretical results. We now present large-sample deviation and fluctuation results between the approximate singular vectors $\hat{\mathbf{U}}_g$ of $\hat{\mathbf{M}}$ (as obtained via *rs*-RSVD) and the true

singular vectors \mathbf{U} of the signal matrix \mathbf{M} . In particular Sections 3.1 and 3.2 give high-probability ℓ_2 and $\ell_{2 \rightarrow \infty}$ perturbation bounds (deviations) of $\hat{\mathbf{U}}_g$. The normal approximation for the row-wise fluctuations of $\hat{\mathbf{U}}_g$ is given in Section 3.4.

We first introduce some notation. Let the SVDs of $\hat{\mathbf{M}}^g \mathbf{G}$ and $\mathbf{M}^g \mathbf{G}$ be denoted as

$$(3.1) \quad \begin{aligned} \mathbf{M}^g \mathbf{G} &:= \mathbf{U}_g \boldsymbol{\Sigma}_g \mathbf{V}_g^\top; \\ \hat{\mathbf{M}}^g \mathbf{G} &:= (\hat{\mathbf{U}}_g \hat{\mathbf{U}}_{g,\perp}) \begin{pmatrix} \hat{\boldsymbol{\Sigma}}_g & \mathbf{0} \\ \mathbf{0} & \hat{\boldsymbol{\Sigma}}_{g,\perp} \end{pmatrix} \begin{pmatrix} \hat{\mathbf{V}}_g^\top \\ \hat{\mathbf{V}}_{g,\perp}^\top \end{pmatrix}. \end{aligned}$$

where $\boldsymbol{\Sigma}_g = \text{diag}(\sigma_1, \dots, \sigma_{k^*})$ and $\hat{\boldsymbol{\Sigma}}_g = \text{diag}(\hat{\sigma}_1, \dots, \hat{\sigma}_{k^*})$ contain the largest k^* singular values of $\mathbf{M}^g \mathbf{G}$ and $\hat{\mathbf{M}}^g \mathbf{G}$, respectively. Recall that we assumed $\text{rk}(\mathbf{M}) = k^*$. The following proposition shows that the leading singular vectors of $\mathbf{M}^g \mathbf{G}$ and \mathbf{M} are equivalent.

PROPOSITION 3.1. *Let $\mathbf{G} \in \mathbb{R}^{n \times \tilde{k}}$ be a Gaussian random matrix with $\tilde{k} \geq k^*$. Let g be a finite positive integer. Then $d_2(\mathbf{U}_g, \mathbf{U}) = d_{2 \rightarrow \infty}(\mathbf{U}_g, \mathbf{U}) = 0$ almost surely.*

Proposition 3.1 requires the sketching dimension \tilde{k} to be no smaller than k^* , the rank of \mathbf{M} , but allows for g to be any positive integer. This requirement also indicates that the sketching of $\hat{\mathbf{M}}$ is generally more difficult compared to that of \mathbf{M} , even when $\|\hat{\mathbf{M}} - \mathbf{M}\| = o(\|\mathbf{M}\|)$ with high probability. Indeed, $\hat{\mathbf{M}}$ is usually full-rank and hence the leading k^* singular vectors of $\hat{\mathbf{M}}^g \mathbf{G}$ need not be close to that of \mathbf{M} for any arbitrary g .

REMARK 3.1. (Known k^*) In this paper we focus on the scenario where the parameter k in Algorithm 1 is correctly specified, i.e., $k = k^*$, and thus k is used interchangeably with k^* in the following theoretical results. In practice k^* might be unknown and need to be estimated. There are a large number of methods for estimating k^* consistently and a few representative, but by no means exhaustive, references include [6, 60, 94, 127, 132]. We note, however, that all of the approaches described in these references require knowing the leading singular values of $\hat{\mathbf{M}}$ and this is possibly problematic in the context of RSVD if these singular values have to be first computed using some classical SVD algorithm. Estimation of k^* can also be done within rs -RSVD itself using similar ideas to that in Sections 4.3 and 4.4 of [58] and while we surmise that the consistency of these estimates are reasonably straightforward to establish, they nevertheless involve additional notations and derivations. We thus decided to leave the *simultaneous* estimation of k^* and \mathbf{U} using RSVD for future work.

REMARK 3.2 (Distribution for \mathbf{G}). In addition to standard Gaussian, other distributions for \mathbf{G} such as uniform and Rademacher distributions have also been studied [68, 83, 124]. We focus on the Gaussian distribution mainly for ease of exposition; any distribution for which \mathbf{G} satisfies Proposition 3.1 and Lemmas S.4.4–S.4.5 in the Supplementary File, will also lead to the same theoretical results as that presented in Theorems 3.1–3.3 of this paper. In particular if \mathbf{G} is such that $\mathbb{P}\{\text{rk}(\mathbf{U}^\top \mathbf{G}) = k^*\} = 1$ then Proposition 3.1 holds. If \mathbf{G} has iid sub-Gaussian entries then Lemmas S.4.4–S.4.5 also hold (using existing results on the smallest singular values of random matrices from [105]). Thus the main theorems of this paper remain unchanged if the entries of \mathbf{G} are uniformly distributed. The case where \mathbf{G} is Rademacher requires further analysis as $\mathbb{P}\{\text{rk}(\mathbf{U}^\top \mathbf{G}) = k^*\} < 1$ and Proposition 3.1 no longer holds.

3.1. ℓ_2 error bound. Proposition 3.1 implies the almost-sure equivalence between \mathbf{U}_g and \mathbf{U} . Thus, to obtain a ℓ_2 perturbation bound between $\hat{\mathbf{U}}_g$ and \mathbf{U} , we can alternatively study the ℓ_2 perturbation between $\hat{\mathbf{U}}_g$ and \mathbf{U}_g ; Recall that $\hat{\mathbf{U}}_g$ and \mathbf{U}_g are the left singular matrices of $\hat{\mathbf{M}}^g \mathbf{G}$ and $\mathbf{M}^g \mathbf{G}$. We first derive an upper bound for $\|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\|$ using the expansion for $\hat{\mathbf{M}}^g - \mathbf{M}^g$ introduced in Lemma S.4.2; see Theorem S.3.1. This upper bound depends on the SNR as well as several quantities that depend only on the sketching matrix \mathbf{G} . We bound the quantities depending on \mathbf{G} separately (see Lemma S.4.4 and Lemma S.4.5) and thereby obtain the following result.

THEOREM 3.1. *Let $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} and \mathbf{E} satisfy the conditions in Assumption 1. Let \mathbf{G} be generated via Algorithm 1 with a fixed $g \geq 1$, $a_n \lesssim \log^2 n$, and $\tilde{k} \geq k^*$. Denote $\vartheta := \max\{1, \tilde{k}^{-1/2} \log^{1/2} n\}$. We then have, with probability at least $1 - n^{-5}$,*

$$(3.2) \quad \|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\| \lesssim n^{1/2} E_n^g + \mathbb{I}(g \geq 2) \cdot \vartheta \tilde{k}^{1/2} S_n^{g-1} E_n.$$

Furthermore, if $g \geq 1 + (2\beta_\Delta)^{-1}$ then the above bound can be simplified to

$$(3.3) \quad \|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\| \lesssim \vartheta \tilde{k}^{1/2} S_n^{g-1} E_n.$$

By Wedin's Theorem [122], the spectral-norm concentration in Theorem 3.1 together with the lower bound for $\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})$ in Lemma S.4.5 yield the following bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$.

THEOREM 3.2. *Suppose Assumption 1 hold, and that $\hat{\mathbf{U}}_g$ is generated via Algorithm 1 for some $\tilde{k} \geq k$ satisfying either one of the following three conditions:*

- (a) both k and \tilde{k} are fixed.
- (b) k is fixed and \tilde{k} is growing with n .
- (c) k is growing with n and $\tilde{k} \geq c_{\text{gap}} k$ for any fixed $c_{\text{gap}} > 1$.

Assume the condition number ψ satisfies $\psi = o(S_n/E_n)$. Choose $a_n \geq \lceil C \tilde{k}^{-1} \log n \rceil$ where C is a constant not depending on n . Then for any $g \geq 1 + (2\beta_\Delta)^{-1}$ we have, with probability at least $1 - n^{-5}$,

$$(3.4) \quad d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \leq \sqrt{2} \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U})\| \lesssim \vartheta \psi^g \frac{E_n}{S_n}.$$

Recall ϑ is defined in Theorem 3.1.

REMARK 3.3. We now make a few technical remarks concerning Theorem 3.2.

1. If $(2\beta_\Delta)^{-1} < g < 1 + (2\beta_\Delta)^{-1}$ then a weaker bound than Eq. (3.4) still holds, namely

$$(3.5) \quad d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim n^{1/2} \tilde{k}^{-1/2} \psi^g \left(\frac{E_n}{S_n}\right)^g + \mathbb{I}(g \geq 2) \vartheta \psi^g \frac{E_n}{S_n}$$

with probability at least $1 - n^{-5}$. The upper bound in Eq. (3.5) diverges and thus becomes trivial when $g \leq (2\beta_\Delta)^{-1}$; indeed, we always have $d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \leq \sqrt{2}$.

2. The constant C in Theorem 3.2 does not depend on n but can depend on other parameters such as k, \tilde{k} and c_{gap} . More specifically the proof of Lemma S.4.5 shows that
 - C depends only on k and \tilde{k} under scenario (a).
 - $C \leq 15$ under scenario (b).
 - C depends only on c_{gap} under scenario (c) with C decreasing as c_{gap} increases.
3. The error bounds given in Theorem 3.2 also hold for $d_2(\hat{\mathbf{V}}_g, \mathbf{V}_g)$.

4. The bound in Eq.(3.4) depends on the condition number ψ and large values of ψ can lead to sub-optimal bounds. Nevertheless the bound in Eq. (3.4) is meaningful provided that $\psi = \mathcal{O}(n^\epsilon)$ for any $\epsilon < 2\beta_\Delta^2/(1 + 2\beta_\Delta)$.

Theorem 3.2 also yields a perturbation bound between the (approximate) singular values $\tilde{\sigma}_1, \dots, \tilde{\sigma}_k$ obtained from Algorithm 1 and the true singular values $\sigma_1, \dots, \sigma_k$ of \mathbf{M} .

COROLLARY 3.1. *Assume the setting of Theorem 3.2. Let $\{\tilde{\sigma}_i\}_{i=1}^k$ be the output of Algorithm 1 and $\sigma_1 \geq \dots \geq \sigma_k$ be the leading singular values of \mathbf{M} . Then for $g \geq 1 + (2\beta_\Delta)^{-1}$ we have, with probability at least $1 - n^{-5}$, $\max_{\ell \in [k]} |\tilde{\sigma}_\ell - \sigma_\ell| \lesssim \vartheta \psi^g E_n$.*

Assumption 1 together with Weyl's inequality implies that $\max_{\ell \in [k]} |\hat{\sigma}_\ell - \sigma_\ell| \lesssim E_n$ with high probability where $\hat{\sigma}_\ell$ are the (exact) leading singular values of $\hat{\mathbf{M}}$. Corollary 3.1 thus indicates that the estimation error rate of $\tilde{\sigma}_\ell$ is the same as that for $\hat{\sigma}_\ell$, up to some factors depending on the conditional number ψ and ϑ ; recall that $\vartheta = \mathcal{O}(1)$ when $\tilde{k} \asymp \log^{1/2} n$ and ψ is either bounded or, at most, slowly-diverging.

3.2. $\ell_{2,\infty}$ norm error bound. We now study the $\ell_{2 \rightarrow \infty}$ perturbation between $\hat{\mathbf{U}}_g$ and \mathbf{U} . As the $\ell_{2 \rightarrow \infty}$ norm is generally more stringent than the ℓ_2 norm, we introduce an additional assumption on the row-wise fluctuations of \mathbf{E}^g for $g \geq 2$.

ASSUMPTION 2. There exist positive constants $\delta > 1$ and $N_0 > 0$ such that, for any $n \geq N_0$, any positive integer $g \leq \log n$, any standard basis $e_i \in \mathbb{R}^n$, and any vector $\mathbf{u} \in \mathbb{R}^n$ not dependent on \mathbf{E} , we have with probability at least $1 - n^{-8}$ that

$$(3.6) \quad \max_{i \in [n]} |e_i^\top \mathbf{E}^g \mathbf{u}| \leq C_{2 \rightarrow \infty} E_n^g (\log n)^{\delta g} \|\mathbf{u}\|_{\max},$$

Here e_i is the i th basis vector in \mathbb{R}^n and $C_{2 \rightarrow \infty}$ is a finite constant depending possibly on N_0 but not on n and g .

The concentration bound in Eq. (3.6) appears frequently in the literature on $\ell_{2 \rightarrow \infty}$ perturbation, see e.g., [21, 35, 46, 84, 90, 100], and is satisfied by a large class of random matrices \mathbf{E} . More specifically, Proposition 3.2 below shows that if \mathbf{E} is a (generalized) Wigner matrix whose entries are independent mean 0 random variables with subexponential tails then \mathbf{E} satisfies Assumption 2. Proposition 3.2 is motivated by Remark 2.5 in [48] and we will use it to verify Assumption 2 for the noise matrices considered in Sections 4 and 5. We also note that the probability lower bound $1 - n^{-8}$ of Eq. (3.6) is somewhat arbitrary and is chosen mainly for convenience. Indeed, Proposition 3.2 shows that for any $c > 0$ there exists a N_0 depending on c such that the left hand side of Eq. (3.6) holds with probability at least $1 - n^{-c}$ whenever $n > N_0$. See also the discussion after Assumption 1.

PROPOSITION 3.2. *Let $\mathbf{E} = [e_{ij}]$ be a $n \times n$ matrix whose entries (or upper triangular entries if \mathbf{E} is symmetric) are independent zero-mean random variables. Suppose that there exist a constant $C_1 > 0$ not depending on n and a quantity q_n such that*

$$q_n^{g'-2} \mathbb{E} |e_{ij}/E_n|^{g'} \leq C_1^{g'} n^{-1}$$

for all $i \in [n], j \in [n]$, and integer $g' \in [2, (\log n)^{A_0 \log \log n}]$; here $A_0 \geq 10$ is a fixed constant, E_n is the quantity appearing in Assumption 1, and $q_n \rightarrow \infty$ as $n \rightarrow \infty$. Then for any fixed constant $\delta > 1$, any integer $g \leq \log n$, and any vector $\mathbf{u} \in \mathbb{R}^n$ **not** dependent on \mathbf{E} , we have

$$\max_{i \in [n]} |e_i^\top \mathbf{E}^g \mathbf{u}| \leq 2E_n^g (\log n)^{\delta g} \|\mathbf{u}\|_{\max}$$

with probability at least $1 - \exp\{-\nu(\log n)^\delta\}$, provided that $n \geq N$. Here N and ν are constants depending only on C_1, δ , and the divergence rate of q_n .

We now provide two simple examples illustrating the use of Proposition 3.2.

EXAMPLE 1 (Bernoulli Entries). Let $C_{\text{EI}} > 0$ be a finite constant not depending on n . Let $\mathbf{E} = [e_{ij}] \in \mathbb{R}^{n \times n}$ be a symmetric matrix with independent upper triangular entries such that, for all $(i, j) \in [n]^2$ we have $e_{ij} = 1 - C_{ij}\rho_n$ with probability $C_{ij}\rho_n$ and $e_{ij} = -C_{ij}\rho_n$ with probability $1 - C_{ij}\rho_n$, for some $C_{ij} < C_{\text{EI}}$. Suppose furthermore that $\rho_n \in [0, 1]$ satisfies $n\rho_n \rightarrow \infty$ as $n \rightarrow \infty$. This type of matrix \mathbf{E} appears frequently in random graph inference; see Section 4. Now let $E_n = (n\rho_n)^{1/2}$. If $g = 1$ then Eq. (3.6) follows from Bernstein's inequality with $\delta = \frac{1}{2}$ (see e.g. Lemma 16 in [3]). Now suppose that $g \geq 2$ and let $q_n = E_n$. Then, after some straightforward algebra, we have (1) $\mathbb{E}[\mathbf{E}] = \mathbf{0}$ and (2) there exist some $C_1 > 1$ such that for all $g' \geq 2$,

$$q_n^{g'-2} \mathbb{E}|e_{ij}/E_n|^{g'} \leq (n\rho_n)^{-1} \mathbb{E}|e_{ij}|^2 \leq C_1 n^{-1}$$

where the first inequality is because $|e_{ij}| < 1$ for all $(i, j) \in [n]^2$. The matrix \mathbf{E} satisfies the conditions in Proposition 3.2 and hence \mathbf{E} satisfies Assumption 2.

EXAMPLE 2 (Sub-Gaussian Entries). Let $\mathbf{E} = [e_{ij}] \in \mathbb{R}^{n \times n}$ be symmetric with independent zero-mean sub-Gaussian entries whose Orlicz-2 norms are bounded from above by σ_n^2 . Let $E_n = \sqrt{2n}\sigma_n$. If $g = 1$ then Eq. (3.6) follows from a general version of Hoeffding's inequality (see e.g., Theorem 2.6.3 in [120]) with $\delta = \frac{1}{2}$. Now suppose $g \geq 2$. We then have $\mathbb{P}(n^{1/2}|e_{ij}/E_n| \geq x) \leq \exp(-x^2)$ [120, § 2.5.2] which implies the moment condition

$$\mathbb{E}|e_{ij}/E_n|^{g'} \leq \frac{(g'/2)^{g'/2}}{n^{g'/2}}.$$

Let $A_0 = 10$, $q_n = (2n)^{1/2}\{(\log n)^{10 \log \log n}\}^{-1/2}$, and $C_1 = 2$, we have

$$q_n^{g'-2} \mathbb{E}|e_{ij}/E_n|^{g'} \leq n^{g'/2-1} (g'/2)^{-g'/2+1} \frac{(g'/2)^{g'/2}}{n^{g'/2}} = \frac{g'/2}{n} \leq C_1^{g'} n^{-1},$$

We have thus verified all conditions in Proposition 3.2 hence \mathbf{E} satisfies Assumption 2.

Given Assumption 2, the derivation of our $d_{2 \rightarrow \infty}$ bounds proceeds as follows. We first combine the Procrustean matrix decomposition in Theorem 3.1 of [22] with the series expansion for $\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}$ given in Lemma S.4.2. This yields an expression for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ that depends mainly on two type of terms, namely (1) terms of the form $\|\mathbf{E}^g \mathbf{X}\|_{2 \rightarrow \infty}$ for some matrix \mathbf{X} of dimensions $\mathbb{R}^{n \times k}$ or $\mathbb{R}^{n \times \tilde{k}}$ where \mathbf{X} does not depend on \mathbf{E} , and (2) terms of the form $\|\mathbf{X}^\top \mathbf{G}\|$ for some $\mathbf{X} \in \mathbb{O}_{n \times k}$ not depending on \mathbf{G} . The terms $\|\mathbf{E}^g \mathbf{X}\|_{2 \rightarrow \infty}$ are bounded using the concentration inequality in Assumption 2 while the terms $\|\mathbf{X}^\top \mathbf{G}\|$ are bounded in a similar manner to those in Theorem 3.2; see Theorem S.3.3 and Lemma S.4.4 to Lemma S.4.6 in the Supplementary File, for more details. In summary we obtain the following result.

THEOREM 3.3. *Assume the setting of Theorem 3.2. Further assume \mathbf{E} satisfies Assumption 2. If $g \geq 1 + (2\beta_\Delta)^{-1}$ then, with probability at least $1 - n^{-5}$,*

$$(3.7) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \Theta_n \frac{E_n}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Here $\Theta_n := (\log n)^{\delta^*} \vartheta \psi^g k^{1/2}$, and $\delta^* = \delta$ if either $g > 1 + (2\beta_\Delta)^{-1}$ or $\|\mathbf{U}\|_{2 \rightarrow \infty} \gtrsim n^{-\beta_c/2}$ for some $\beta_c \in [0, 1]$; the constant δ is defined in Assumption 2. Otherwise, if $g = 1 + (2\beta_\Delta)^{-1}$, then $\delta^* = \delta g + 1/2$.

The upper bound in Eq. (3.7) decreases with $\|\mathbf{U}\|_{2 \rightarrow \infty}$. Furthermore, as $\mathbf{U} \in \mathbb{O}_{n \times k}$, we have $k^{1/2}n^{-1/2} \leq \|\mathbf{U}\|_{2 \rightarrow \infty} \leq k^{1/2}$ always. The quantity $\frac{n}{k}\|\mathbf{U}\|_{2 \rightarrow \infty}^2$ is termed the coherence of \mathbf{U} [20]. A matrix \mathbf{U} with $n^{1/2}\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim 1$ is said to have bounded coherence; bounded coherence for \mathbf{U} is a prevalent and typically mild assumption in many high-dimensional statistics problem including matrix completion, covariance estimation, subspace estimation, and random graph inference; see e.g., [3, 18, 20, 22, 51, 75] and the examples in Section 4 and Section 5 of this paper.

REMARK 3.4. The upper bound in Eq. (3.7) requires $g \geq 1 + (2\beta_\Delta)^{-1}$ and is slightly sharper when $g > 1 + (2\beta_\Delta)^{-1}$; indeed $\delta^* = \delta$ if $g > 1 + (2\beta_\Delta)^{-1}$ while $\delta^* = \delta g + 1/2$ if $g = 1 + (2\beta_\Delta)^{-1}$ and $\|\mathbf{U}\|_{2 \rightarrow \infty} \asymp n^{-1/2}$. This difference is, however, very minor. Indeed, as g is an integer, $g = 1 + (2\beta_\Delta)^{-1}$ is possible only when $\beta_\Delta = (2\ell)^{-1}$ for integers $\ell \geq 1$. If $g < 1 + (2\beta_\Delta)^{-1}$ then a similar bound to Eq. (3.7) is available, albeit with a provably worse upper bound, i.e., with probability at least $1 - n^{-5}$,

$$(3.8) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \rho_n^{(1)}\|\mathbf{U}\|_{2 \rightarrow \infty} + \rho_n^{(2)},$$

where $\rho_n^{(1)}$ and $\rho_n^{(2)}$ are defined in Eq. (S.5.27) and Eq. (S.5.28) of the supplementary.

REMARK 3.5. For ease of exposition we have stated Eq. (3.6) in terms of a single $\delta > 1$ for all $g \in \{1, 2, \dots, \log n\}$. We can instead assume that there exists a non-decreasing and bounded sequence δ_g such that, for all $g \leq \log n$ we have

$$\max_{i \in [n]} |e_i^\top \mathbf{E}^g \mathbf{u}| \leq C_{2 \rightarrow \infty} E_n^g (\log n)^{\delta_g g} \|\mathbf{u}\|_{\max},$$

with probability at least $1 - n^{-8}$. Then Theorem 3.3 still holds under the above assumption but with $\delta = \delta_1$ and this allows for a more precise control of the factor Θ_n in Eq. (3.7). Indeed, as we see from Example 1 and Example 2 above, for many inference problem such as network analysis and matrix completion we can control $e_i^\top \mathbf{E} \mathbf{u}$ using either Bernstein's inequality or Hoeffding's inequality so that $\delta_1 = \frac{1}{2}$; the remaining δ_g are bounded from above by some finite constant $C_\delta > 1$ via Proposition 3.2. With $\delta_1 = \frac{1}{2}$ the term Θ_n in Theorem 3.3 (and Corollary 3.2 later) is simplified to $\Theta_n = (\log n)^{1/2}$ when $\tilde{k} = \Omega(\log n)$ and $g > 1 + (2\beta_\Delta)^{-1}$; this factor $\Theta_n = (\log n)^{1/2}$ is generally optimal.

REMARK 3.6. Bounds for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$ can be derived using several different approaches, including leave-one-out analysis [3, 3, 18, 27], von Neumann series expansion [24, 30, 46], holomorphic functional analysis [75], and techniques tailored towards Gaussian ensembles [71–73]. See [28] for a comprehensive survey. Our work, meanwhile, focuses on $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. As $\hat{\mathbf{U}}_g$ is obtained from the SVD of $\hat{\mathbf{M}}^g \mathbf{G}$, the bounds for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ depends on $(\hat{\mathbf{M}}^g - \mathbf{M}^g) \mathbf{G}$ which has a substantially more complicated entrywise dependency structure than that in $\mathbf{E} = \hat{\mathbf{M}} - \mathbf{M}$. As a result not all of the above cited techniques are directly applicable for bounding $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. For this paper we combine the Procrustean analysis argument from [22] with the expansion for $\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}$ given in Lemma S.4.2. This approach yields a perturbation expansion for $\hat{\mathbf{U}}_g \mathbf{W} - \mathbf{U}$ that includes auxiliary terms of the form $\|\mathbf{E}^g \mathbf{X}\|_{2 \rightarrow \infty}$ where $\mathbf{X} \in \mathbb{O}_{n \times \tilde{k}}$ does not depend on \mathbf{E} , and we leveraged Assumption 2 and Proposition 3.2 to bound these terms. We leave the problem of adapting other techniques (such as leave-one-out analysis) to control the row-wise deviation of $\hat{\mathbf{U}}_g$ for future work.

The row-wise fluctuation for $\hat{\mathbf{U}}_g$ in Theorem 3.3 also yields an entrywise concentration bound for $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M}$. In particular we have the following result.

COROLLARY 3.2. *Assume the setting of Theorem 3.3 and suppose that both k and ψ are bounded as n increases. Let $g \geq 1 + (2\beta_\Delta)^{-1}$. We then have*

$$\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M}\|_{\max} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

with probability at least $1 - n^{-5}$; here Θ_n satisfies $\Theta_n \lesssim (\log n)^\delta \vartheta$ if $g > 1 + (2\beta_\Delta)^{-1}$ and $\Theta_n \lesssim (\log n)^{\delta g + 1/2} \vartheta$ otherwise.

3.3. Comparison with existing results. We now compare the perturbation bounds for $\hat{\mathbf{U}}_g$ in ℓ_2 and $\ell_{2 \rightarrow \infty}$ norms with existing results in the literature. As we mentioned in the introduction, most theoretical analysis of RSVD focused exclusively on the setting where $\hat{\mathbf{M}}$ is assumed to be noise-free and the quantities of interest are bounds for $(\mathbf{I} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \hat{\mathbf{M}}$ in either spectral norm or Frobenius norm. For example by combining Theorem 9.2 and Theorem 10.8 in [58] we have that if $\hat{\mathbf{M}}$ is symmetric and $\tilde{k} \geq k + 2$ then

$$(3.9) \quad \|(\mathbf{I} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \hat{\mathbf{M}}\| \leq \left[\left(1 + \frac{k^{1/2}}{(\ell - 1)^{1/2}} \right) \sigma_{k+1}^g(\hat{\mathbf{M}}) + \frac{e^{\tilde{k}^{1/2}}}{\ell} \left\{ \sum_{j>k} \sigma_j^{2g}(\hat{\mathbf{M}}) \right\}^{1/2} \right]^{1/g}$$

with high probability. Here $\ell = \tilde{k} - k$ and $\sigma_j(\hat{\mathbf{M}})$ denote the j th largest singular value of $\hat{\mathbf{M}}$. If $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} and \mathbf{E} satisfy Assumption 1 then, without additional information, we can only conclude that $\max_{j>k} \sigma_j(\hat{\mathbf{M}}) \leq E_n$. Eq. (3.9) then reduces to

$$\|(\mathbf{I} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \hat{\mathbf{M}}\| \lesssim n^{1/2g} E_n,$$

with high probability, for any fixed g . Recall that $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$ is a low rank approximation of $\hat{\mathbf{M}}$ and serves as an estimate of the true but unknown signal matrix \mathbf{M} . Invoking Assumption 1 again we obtain

$$(3.10) \quad \|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\| \leq \|\hat{\mathbf{M}} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\| + \|\hat{\mathbf{M}} - \mathbf{M}\| \lesssim n^{1/2g} E_n,$$

with high probability. On the other hand, Theorem 3.2 of this paper implies

$$(3.11) \quad \|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\| \leq \|(\mathbf{U}\mathbf{U}^\top - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \mathbf{M}\| + \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top (\mathbf{M} - \hat{\mathbf{M}})\| \lesssim \vartheta \psi^g E_n,$$

with high probability. Note that Eq. (3.11) is much sharper than Eq. (3.10) when \mathbf{M} is (approximately) low-rank as assumed in the current paper. Indeed if ψ^g is bounded or slowly diverging then the multiplicative factor in Eq. (3.11) is either bounded or of order $o(n^\epsilon)$ for any arbitrary constant $\epsilon > 0$ while the multiplicative factor in Eq. (3.10) is $\Theta(n^{1/2g})$ and thus requires $g \asymp \log n$ to be competitive with the bound in Eq. (3.11); note that $g \asymp \log n$ power iterations was also used in the theoretical analysis of RSVD in [89] and Theorem 60 in [124]; as we only require $g \geq 1 + (2\beta_\Delta)^{-1}$, our bounds are more closely tailored to the setting of noisy $\hat{\mathbf{M}}$. Finally note that $\|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\| \leq 2E_n$ with high probability. Thus $\|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\|$ attains the same error rate as $\|\mathbf{M} - \hat{\mathbf{U}} \hat{\mathbf{U}}^\top \hat{\mathbf{M}}\|$, up to some (at most) logarithmic factor.

The above bounds for $\|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\|$ are based on $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$. We now consider the implication of the bounds for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. Suppose \mathbf{U} has bounded coherence, i.e., $\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim k^{1/2} n^{-1/2}$, and \mathbf{E} satisfies Assumption 2. Let $g > 1 + (2\beta_\Delta)^{-1}$ and \tilde{k}, a_n are as specified in Theorem 3.3. If k and ψ are slowly growing then Corollary 3.2 implies

$$(3.12) \quad \|\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}\|_{\max} \lesssim (\log n)^\delta \vartheta \psi^g k \frac{E_n}{n},$$

with high probability. Eq. (3.12) is, to the best of our knowledge, the first bound for entry-wise differences between the RSVD-based low-rank approximation of the noisily observed

$\hat{\mathbf{M}}$ and the underlying signal matrix \mathbf{M} , and provides much finer control for $\mathbf{M} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$. Eq. (3.12) cannot be obtained by simply combining existing bounds for $\|(\mathbf{I} - \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) \hat{\mathbf{M}}\|$ (as given in the RSVD literature) and bounds for $\|\hat{\mathbf{M}} - \mathbf{M}\|$ (as given in the matrix perturbation literature); here $\|\cdot\|$ denote any unitarily invariant norm. Finally, as we will see in Sections 5.1, the bound in Eq. (3.12) can have the same error rate as that for $\|\mathbf{M} - \hat{\mathbf{U}} \hat{\mathbf{U}}^\top \hat{\mathbf{M}}\|_{\max}$, up to some (at most) logarithmic factor.

3.4. Row-wise limiting distribution. The row-wise limiting distribution of singular vectors can be applied for uncertainty quantification of membership inference in network analysis [10, 77]. Under some appropriate probabilistic structures of \mathbf{M} and \mathbf{E} , $\hat{\mathbf{U}}$ is proved to have a row-wise limiting distribution [21]. In this section we prove that $\hat{\mathbf{U}}_g$ also has row-wise asymptotic normal distributions, provided that \mathbf{M} and \mathbf{E} satisfy the following condition.

ASSUMPTION 3. Let $\{\sigma_n\}_{n \geq 1}$ be a sequence of bounded non-negative numbers which can converge to 0 as $n \rightarrow \infty$. Assume that both of the following conditions hold as $n \rightarrow \infty$.

- (a). \mathbf{E} satisfies Assumption 1 with $E_n = \Theta(\sqrt{n}\sigma_n)$ and $\frac{S_n}{E_n} \lesssim n^{\beta_\Delta}$ for some constant $\beta_\Delta > 0$.
- (b). For any fixed $i \in [n]$, we have

$$(3.13) \quad S_n \sigma_n^{-1} \mathbf{\Gamma}_i^{-1/2} [\mathbf{E} \mathbf{U} \mathbf{\Lambda}^{-1}]_i \rightsquigarrow \mathcal{N}(0, \mathbf{I}).$$

where $\mathbf{\Gamma}_i$ is a deterministic matrix depending possibly on i and \mathbf{M} .

Assumption 3 is valid for more general \mathbf{M} and \mathbf{E} , compared with Assumption 5 in [21], which typically focuses on a random graph context. One scenario for which Assumption 3 holds is when \mathbf{E} has symmetrically independent entries and the variances of $\Theta(n^2)$ entries in \mathbf{E} are of the same asymptotic order $\Theta(\sigma_n^2)$, while the variance of the other entries are negligible. Such entrywise independent and homogenous conditions of \mathbf{E} appear frequently in many statistical problems including matrix completion, graph embeddings, and multi-dimensional scaling; see e.g., [21, 23, 79].

THEOREM 3.4. Suppose Assumptions 1–3 hold with fixed $k^* = k$ and bounded ψ . Let $\hat{\mathbf{U}}_g$ be generated via Algorithm 1 with $\tilde{k} \geq k$, $a_n \geq \lceil C \tilde{k}^{-1} \log n \rceil$ where the universal constant C is given in Remark 3.3, and $g \geq 2 + (2\beta_\Delta)^{-1}$. Now suppose that as $n \rightarrow \infty$, we have

$$(3.14) \quad \max \left\{ \sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|, n \sigma_n \ell_{\text{clt}} S_n^{-1} \right\} \|\mathbf{U}\|_{2 \rightarrow \infty} \rightarrow 0,$$

in probability, where ℓ_{clt} is a logarithmic factor such that

$$(3.15) \quad \ell_{\text{clt}} := \begin{cases} (\log n)^{g\delta+1/2} & \text{if } g = 2 + (2\beta_\Delta)^{-1}, \\ (\log n)^{2\delta} \vartheta & \text{if } g > 2 + (2\beta_\Delta)^{-1}. \end{cases}$$

Then there exist a sequence of orthogonal matrices $\mathbf{W}_{\text{clt}} = \mathbf{W}_{\text{clt}}^{(n)}$ such that for any $i \in [n]$,

$$S_n \sigma_n^{-1} \mathbf{\Gamma}_i^{-1/2} (\mathbf{W}_{\text{clt}} [\hat{\mathbf{U}}_g]_i - [\mathbf{U}]_i) \rightsquigarrow \mathcal{N}(\mathbf{0}, \mathbf{I}).$$

For each n the matrix \mathbf{W}_{clt} solves the orthogonal Procrustes problem between $\hat{\mathbf{U}}_g$ and \mathbf{U} (see Eq. (S.5.47)).

The condition $g \geq 2 + (2\beta_\Delta)^{-1}$ in Theorem 3.4 is slightly more stringent than the condition $g \geq 1 + (2\beta_\Delta)^{-1}$ in Theorem 3.1 and Theorem 3.3. The main reason for this discrepancy is that while $g \geq 1 + (2\beta_\Delta)^{-1}$ might be sufficient for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$

to achieve the optimal error rate, it does not guarantee that the row-wise fluctuations of $\hat{\mathbf{U}}_g \mathbf{W}_{\text{clt}}^\top - \mathbf{U}$ is asymptotically equivalent to the row-wise fluctuations of $\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}$ given in Assumption 3. We will present, in Section S.1.1.3 in the Supplementary File, simulation results for the row-wise Gaussian fluctuations of $\hat{\mathbf{U}}_g$ when $\hat{\mathbf{M}}$ is the adjacency matrix of a SBM random graph; our simulations indicate that the condition $g \geq 2 + (2\beta_\Delta)^{-1}$ is in fact necessary and sufficient.

REMARK 3.7. We now make a few technical remarks concerning Theorem 3.4.

1. Theorem 3.4 can be extended to the case where either (or both) of ψ and k are slowly diverging, provided that we also increase the exponent for $\log n$ in the definition of ℓ_{clt} .
2. The term $\sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}$ in Eq. (3.14) can usually be bounded by $\text{polylog}(n) \|\mathbf{U}\|_{2 \rightarrow \infty}$ with high probability. More specifically, let $\mathbf{E} = [e_{ij}]_{n \times n}$ be a symmetric matrix whose upper triangular entries are independent mean 0 random variables with $\mathbb{E}[e_{ij}^2] \leq \sigma_n^2$ and $|e_{ij}| \leq B_n$ for all i, j . Now if $B_n \lesssim \sigma_n (\log n)^{-1/2} \|\mathbf{U}\|_{2 \rightarrow \infty}^{-1}$ then, by Bernstein inequality,

$$\sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \lesssim \log^{1/2} n$$

with high probability; see Eq. (4.137) in [28] for more details. Thus $\sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}$ converges to 0 in probability whenever $\|\mathbf{U}\|_{2 \rightarrow \infty} = \mathcal{O}(\log^{-1/2} n)$. Recall that $n^{-1/2} \lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim 1$.

3. The term $n\sigma_n S_n^{-1} \ell_{\text{clt}} \|\mathbf{U}\|_{2 \rightarrow \infty}$ in Eq. (3.14) depends on the SNR and the coherence of \mathbf{U} i.e., this term is equivalent to $n^{(1/2-\beta_\Delta)} \ell_{\text{clt}} \|\mathbf{U}\|_{2 \rightarrow \infty}$. Thus larger values of $\|\mathbf{U}\|_{2 \rightarrow \infty}$ require larger SNR n^{β_Δ} in order to guarantee convergence in Eq. (3.14). If \mathbf{U} has bounded coherence, i.e., $n^{1/2} \|\mathbf{U}\|_{2 \rightarrow \infty}$ is bounded, then $n^{(1/2-\beta_\Delta)} \ell_{\text{clt}} \|\mathbf{U}\|_{2 \rightarrow \infty} \rightarrow 0$ due to our assumption that $\beta_\Delta > 0$. As we alluded to earlier, bounded coherence for \mathbf{U} is a prevalent and typically mild assumption in many high-dimensional statistics problem including matrix completion, covariance estimation, subspace estimation, and random graph inference; see e.g., [3, 18, 20, 22, 51, 75].
4. Theorem 3.4 can be used to derive normal approximations for the entries of $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$. For brevity we defer this result to Theorem S.6.1 in the Supplementary File; see Theorem 5.1 for an example of this normal approximation for noisy matrix completion.

4. Random graph inference. As an illustrative example of Theorem 3.2 and Theorem 3.3, we consider the problem of estimating the leading eigenvectors for edge-independent random graphs with low-rank edge probabilities matrices. More specifically, let $\hat{\mathbf{M}} = [\hat{m}_{ij}]$ be the adjacency matrix of a random graph on n vertices with edge probabilities $\mathbf{M} = [m_{ij}] \in [0, 1]^{n \times n}$ i.e., $\hat{\mathbf{M}}$ is a symmetric, binary matrix whose upper triangular entries are independent Bernoulli random variables with $\mathbb{P}(\hat{m}_{ij} = 1) = m_{ij}$. Next assume that \mathbf{M} is of rank k for some fixed $k \geq 1$ and has bounded condition number, i.e., $\psi \leq C_{\text{CE}}$ for some fixed constant $C_{\text{CE}} > 0$. Let $\beta \in (0, 1]$ and suppose that Assumption 1 holds with

$$(4.1) \quad S_n \asymp n^\beta \text{ and } E_n \asymp n^{\beta/2}$$

for some fixed constant $\beta \in (0, 1]$. We interpret n^β as the average expected degree for the vertices of $\hat{\mathbf{M}}$, i.e., with high probability $\hat{\mathbf{M}}$ has $\Theta(n^{1+\beta})$ non-zero entries. Smaller values of β thus implies sparser networks. We note that Eq. (4.1) is satisfied by many random graph models including Erdős–Rényi [47], SBM and its degree-corrected and/or mixed-membership variants [7, 63, 69, 108], (generalized) random dot product graphs [103, 128], as well as any edge-independent random graph whose edge probabilities are sufficiently homogeneous, e.g., $\sum_j m_{ij} \asymp n^\beta$ for all $i \in [n]$. Assumption 2 is also satisfied by the random graphs mentioned above; see e.g. [21, 48, 84] and Example 1. In addition, we also assume the bounded coherence for \mathbf{U} , $\|\mathbf{U}\|_{2 \rightarrow \infty} \asymp n^{-1/2}$. This is a prevalent and typically mild assumption for random graphs, see e.g., [3, 22, 75].

4.1. *Subspace perturbation error bound.* Now consider the case where n , the number of vertices in $\hat{\mathbf{M}}$, is large and the graph $\hat{\mathbf{M}}$ is possibly semi-sparse, and we are interested in computing the k leading singular vectors of $\hat{\mathbf{M}}$ as an estimate for the k leading singular vectors of \mathbf{M} . To save computational time, we will use Algorithm 1 to approximate the singular vectors of $\hat{\mathbf{M}}$. Now suppose we choose a fixed $g \geq 1$, and either $\tilde{k} = k$ and $a_n \asymp \log n$ or $\tilde{k} \asymp \log n$ and $a_n = 1$; recall that $\tilde{k} \geq k$ is the sketching dimension and a_n is the number of repeated sampling steps. Then from Theorems 3.2 and 3.3 (and the corresponding remarks) we have that, with high probability,

$$(4.2) \quad \begin{aligned} d_2(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \begin{cases} \text{polylog}(n) \times n^{-\beta/2} & \text{if } g \geq 1 + \beta^{-1}, \\ \text{polylog}(n) \times n^{(-g\beta+1)/2} & \text{if } \beta^{-1} < g < 1 + \beta^{-1}, \\ 1 & \text{if } 1 \leq g \leq \beta^{-1}; \end{cases} \\ d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \begin{cases} \text{polylog}(n) \times n^{-(\beta+1)/2} & \text{if } g \geq 1 + \beta^{-1}, \\ \text{polylog}(n) \times n^{-g\beta/2} & \text{if } \beta^{-1} < g < 1 + \beta^{-1}, \\ n^{-1/2} & \text{if } 1 \leq g \leq \beta^{-1}. \end{cases} \end{aligned}$$

Here $\text{polylog}(n)$ denote $\log^c n$ for some fixed $c \geq 0$. Eq. (4.2) implies a phase transition for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ as g changes. Recall that $\hat{\mathbf{U}}$ is the matrix whose columns are the (exact) leading k singular vectors of $\hat{\mathbf{M}}$ and Eq. (4.1) together with the Davis-Kahan theorem implies $d_2(\hat{\mathbf{U}}, \mathbf{U}) \lesssim n^{-\beta/2}$ with high probability. On the other hand, it has recently been shown that with high probability $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U}) \lesssim (\log n)^{1/2} n^{-(\beta+1)/2}$; see e.g., [21, 75]. Therefore if $g \geq 1 + \beta^{-1}$ then both $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ converges to 0 at the same rates as $d_2(\hat{\mathbf{U}}, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$, up to some logarithmic factors. Meanwhile, if $\beta^{-1} < g < 1 + \beta^{-1}$ then $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ converge to 0 at the slower rates of $n^{(-g\beta+1)/2}$ and $n^{-g\beta/2}$, respectively. Convergences of $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ are not guaranteed when $g \leq \beta^{-1}$. These different convergence rates can be motivated by the observation that, since $\hat{\mathbf{M}}$ is a binary matrix, the main thing that changes when β decreases is the number of non-zero entries in $\hat{\mathbf{M}}$. In other words, we are estimating the leading k singular vectors of \mathbf{M} using a noisy realization $\hat{\mathbf{M}}$ that, when β decreases, contains less information about \mathbf{M} . It is thus unlikely that any fixed value of g will work uniformly well for all values of β . Indeed, as we will see in Theorem 4.1 and Section S.1.1, both of the thresholds $g > \beta^{-1}$ and $g \geq 1 + \beta^{-1}$ in Eq. (4.2) are necessary and sufficient. We now provide three examples illustrating the relationships between g and β ; see also the visual summary in Figure 1.

- First consider the dense regime with $\beta = 1$. Then $g \geq 2$ is sufficient for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ to attain the optimal rate $n^{-1/2}$; and for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ to attain the optimal rate n^{-1} . No convergence is guaranteed when $g = 1$.
- Next consider a semi-sparse regime with $\beta = 2/3$. For $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$, the optimal rate $n^{-1/3}$ is attained when $g \geq 3$, the sub-optimal rate $n^{-1/6}$ is attained when $g = 2$. For $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$, the optimal rate $n^{-5/6}$ is attained when $g \geq 3$, the sub-optimal rate $n^{-2/3}$ is attained when $g = 2$. No convergence is guaranteed when $g = 1$.
- Finally consider a semi-sparse regime with $\beta = 1/2$. For $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$, the optimal rate $n^{-1/4}$ is attained when $g \geq 3$. For $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$, the optimal rate $n^{-3/4}$ is attained when $g \geq 3$. No convergence is guaranteed when $g = 1, 2$.

In summary, the above discussions provide theoretical justification to the well-known advice that choosing a slightly larger g and \tilde{k} are essential to the success of RSVD in practical applications [86].




FIG 1. Phase transition diagram for the error rate $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ in Eq. (4.2) as β and g changes. Recall that $\|\mathbf{M}\| \asymp n^\beta$ and $\|\mathbf{E}\| \asymp n^{\beta/2}$ and hence $n^{\beta/2}$ represents the signal to noise ratio. The different regions represent different convergence rates for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ or $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. For example, $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ converges at the optimal rate of $n^{-\beta/2}$ (blue region), converges at the slower rate of $n^{(-g\beta+1)/2}$ (green region), and has no convergence (yellow region). The solid line represents the transition threshold $g = 1 + \beta^{-1}$ and the dashed line represents the transition threshold $g = \beta^{-1}$.

4.2. *Lower bound and phase transition sharpness.* We now study the sharpness of the phase transition thresholds described in Section 4.1. More specifically, we derive lower bounds for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$, and thereby show that the conditions $g > \beta^{-1}$ and $g \geq 1 + \beta^{-1}$ are both necessary and sufficient for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ to converge to 0, at either the slow rate or the optimal rate shown in Eq. (4.2), respectively.

The lower bounds for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ depend on the following assumption on the growth rate of $\text{tr} \mathbf{E}^{2g}$, namely that for any finite $g \geq 1$ there exists some fixed constant $c_g > 0$ such that,

$$(4.3) \quad \mathbb{E} \text{tr} \mathbf{E}^{2g} \geq c_g n^{1+\beta}.$$

We now clarify Eq. (4.3). Suppose \mathbf{E} is a Wigner matrix whose entries are independent mean 0 random variables with equal variances given by $\sigma^2 \asymp n^{\beta-1}$. Then \mathbf{E} satisfy Eq. (4.3) using the well-known “trace method” combinatorial arguments from random matrix theory; see e.g. Lemma 1.5 in [15] for more details. Eq. (4.3) and Eq. (4.1) are therefore satisfied for Erdős–Rényi graphs. By adapting the trace method arguments for Wigner matrices, we can show that Eq. (4.3) continues to hold for any edge-independent random graphs with homogeneous variances, i.e., \mathbf{E} satisfies $\text{Var}[e_{ij}] \asymp n^{\beta-1}$ for all i, j , or equivalently, that $m_{ij} \asymp n^{\beta-1}$ for all i, j . In other words, Eq. (4.3) is satisfied by stochastic blockmodel graphs and their degrees-corrected and/or mixed membership variants, and by (generalized) random dot product graphs.

THEOREM 4.1. *Assume the setting of Theorem 3.2 and further suppose: (i) $S_n \asymp n^\beta$ and $E_n \asymp n^{\beta/2}$ for some fixed constant $\beta \in (0, 1]$; (ii) there exists a constant $c_g > 0$ not depending on n but can depend on g such that $\mathbb{E} \text{tr} \mathbf{E}^{2g} \geq c_g n^{1+\beta}$, (iii) $k, \psi \lesssim 1$. Choose*

$\tilde{k} = k$ and $a_n = o(\log^5 n)$. Let $p_0 \in (0, 1)$ be fixed but arbitrary. Then there exists a constant $C_{LB} > 0$ depending only on p_0 such that, with probability approaching p_0 , we have

$$d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \geq \begin{cases} C_{LB}(\log \log n)^{-1/2} \cdot n^{(-g\beta+1)/2} & \text{if } \beta^{-1} \leq g < 1 + \beta^{-1} \\ C_{LB} & \text{if } 1 \leq g < \beta^{-1} \end{cases},$$

$$d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \geq \begin{cases} C_{LB}(\log \log n)^{-1/2} \cdot n^{-g\beta/2} & \text{if } \beta^{-1} \leq g < 1 + \beta^{-1} \\ C_{LB} \cdot n^{-1/2} & \text{if } 1 \leq g < \beta^{-1} \end{cases}.$$

REMARK 4.1. For simplicity we had presented Theorem 4.1 in the case where $\tilde{k} = k$. An almost identical lower bound is available for the case where $\tilde{k} = O(\log n)$, provided that we replaced the $(\log \log n)^{-1/2}$ term in the above expressions with a $(\log n)^{-1/2}$ term.

For ease of exposition we will ignore any logarithmic factor of n in the following discussion. We see that the lower bounds in Theorem 4.1 match the upper bounds in Eq. (4.2) for both the no convergence regime where $g \leq \beta^{-1}$ and the sub-optimal regime where $\beta^{-1} < g < 1 + \beta^{-1}$; recall that if $g \geq 1 + \beta^{-1}$ then $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ has the same convergence rate as $d_2(\hat{\mathbf{U}}, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$, respectively, and are thus rate-optimal. Here we note that in Theorem 4.1, $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \gtrsim n^{-1/2}$ when $1 \leq g < \beta^{-1}$ is equivalent to $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ showing no convergence as $\|\mathbf{U}\|_{2 \rightarrow \infty} \asymp n^{-1/2}$. See Section S.1.1 in the Supplementary File for simulation results illustrating these upper and lower bounds for both $d_2(\mathbf{U}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\mathbf{U}_g, \mathbf{U})$. Finally we emphasize Theorem 4.1 is stated in a more general setting than that for random graphs, e.g., the lower bound holds whenever \mathbf{E} is a random symmetric Gaussian matrix provided that $\|\mathbf{E}\| \lesssim n^{\beta/2}$ with high probability and $\mathbb{E} \operatorname{tr} \mathbf{E}^{2g}$ satisfies Eq.(4.3).

4.3. *Exact recovery for stochastic blockmodels.* We now apply the $\ell_{2 \rightarrow \infty}$ bound for $\hat{\mathbf{U}}_g$ and \mathbf{U} given in Eq. (4.2), to the problem of community detection in stochastic blockmodel graphs [63], one of the most popular generative model for network data with an assumed intrinsic community structure. We first recall the definition of stochastic blockmodel graphs.

DEFINITION 1 (SBM). Let $K \geq 1$ be a positive integer and let $\boldsymbol{\pi} \in \mathbb{R}^K$ be positive vector satisfying $\sum_{\ell} \pi_{\ell} = 1$. Let \mathbf{B} be a symmetric $K \times K$ matrix whose entries are in $[0, 1]$. We say that $(\mathbf{A}, \boldsymbol{\tau}) \sim \text{SBM}(\mathbf{B}, \boldsymbol{\pi})$ is a K -blocks stochastic blockmodel (SBM) graph with parameters \mathbf{B} and $\boldsymbol{\pi}$, and sparsity factor ρ_n , if the following holds. First $\boldsymbol{\tau} = (\tau_1, \dots, \tau_n)$ where the τ_i are iid with $\mathbb{P}(\tau_i = \ell) = \pi_{\ell}$ for all $\ell \in [K]$. Then \mathbf{A} is a $n \times n$ symmetric binary matrix such that, conditioned on $\boldsymbol{\tau}$, for all $i \leq j$ the $\{A_{ij}\}$ are independent Bernoulli random variables with $\mathbb{E}[A_{ij}] = \rho_n B_{\tau_i, \tau_j}$.

Community detection is well-studied (see e.g., the surveys [1, 53]), with many available techniques including those based on maximizing modularity and likelihood [14, 92, 110], random walks [96, 102], semidefinite programming [2, 57], and spectral clustering [121]. In particular spectral clustering using the adjacency matrix is a simple and popular community detection algorithm wherein, given \mathbf{A} , we first choose an embedding dimension d and compute the matrix $\hat{\mathbf{U}}$ of eigenvectors corresponding to the d largest (in modulus) eigenvalues of \mathbf{A} . Next we cluster the rows of $\hat{\mathbf{U}}$ into K cluster using either the K -means or K -medians algorithms and let $\hat{\tau}_i$ be the resulting cluster membership for the i th row of $\hat{\mathbf{U}}$. The motivation behind spectral clustering is that for stochastic blockmodel graphs (1) $\hat{\mathbf{U}}$ serves as an estimate for the leading eigenvectors \mathbf{U} of the underlying edge probabilities matrix $\mathbf{P} = (\rho_n B_{\tau_i, \tau_j})$

Algorithm 2: RSVD-based Spectral Clustering

Input: Adjacency matrix \mathbf{A} , embedding dimension d , number of clusters K . Integers $a_n, g \geq 1, \tilde{k} \geq d$.

1. Obtain $\hat{\mathbf{U}}_g$ via Algorithm 1 with $\hat{\mathbf{M}} = \mathbf{A}, k = d$, sampling times a_n , power g , sampling dimension \tilde{k} ;
2. Estimate the community membership $\hat{\tau}$, by performing either K -means or K -medians on rows of $\hat{\mathbf{U}}_g$;

Output: Membership estimation $\hat{\tau}$ and singular matrix recovery $\hat{\mathbf{U}}_g$.

and (2) the rows of \mathbf{U} contain all of the necessary information for recovering τ , i.e., there exists a collection of K distinct vectors $\{\nu_1, \nu_2, \dots, \nu_K\}$ such that $[\mathbf{U}]_i = \nu_{\tau_i}$ for all $i \in [n]$.

Statistical properties of spectral clustering had been widely studied in recent years, see e.g., [3, 67, 74–76, 82, 99, 112, 113] for an incomplete list of references. In particular it is well-known that spectral clustering will, with high probability, exactly recover the community assignment τ for SBM graphs as well as its variants including degree-corrected SBM [69] and popularity adjusted SBM [108], i.e., spectral clustering yields a $\hat{\tau}$ such that with high probability there exists a permutation ς of $[K]$ for which $\hat{\tau}_i = \varsigma(\tau_i)$ for all $i \in [n]$.

In many real-world applications like social or biological networks, the number of nodes can be on the order of 10^6 , see e.g. [56]. As a result, the computation of $\hat{\mathbf{U}}$ in spectral clustering using standard SVD algorithms can be prohibitively demanding in terms of both the computational time and memory requirement. Under our *rs*-RSVD framework, by taking \mathbf{A} as the observed matrix $\hat{\mathbf{M}}$ we propose an economical spectral clustering procedure (see Algorithm 2) that replaces $\hat{\mathbf{U}}$ by its approximation $\hat{\mathbf{U}}_g$ produced by *rs*-RSVD. We note that the time complexity of Algorithm 2 is on the order of $\mathcal{O}(a_n g \times \text{nnz}(\mathbf{A}))$ operations and only require enough memory to store the $\text{nnz}(\mathbf{A})$ non-zero elements of \mathbf{A} . These time and memory requirements can be considerably smaller than that of the standard spectral clustering, especially for large and non-dense graphs. The following result shows that for a fixed g depending on the sparsity ρ_n of \mathbf{A} , Algorithm 2 will, with high probability, also exactly recovers τ .

THEOREM 4.2. *Let $(\mathbf{A}, \tau) \sim \text{SBM}(\mathbf{B}, \pi)$ be a K -blocks SBM with sparse parameter ρ_n where $\pi_\ell > 0$ for all $\ell \in [K]$ and $n\rho_n \gtrsim n^\beta$ for some $\beta \in (0, 1]$. Let $d = \text{rk}(\mathbf{B})$ and suppose that $\hat{\tau}$ is given by Algorithm 2 wherein we choose: (i) $\tilde{k} \geq d$, (ii) $a_n \geq \max\{C\tilde{k}^{-1} \log n, 1\}$ for some universal constant C and (iii) $g > \beta^{-1}$. Then for sufficiently large n , we have that $\hat{\tau}$ exactly recovers τ with probability at least $1 - n^{-5}$.*

REMARK 4.2. In practice one may consider clustering with the $(1 + \epsilon)$ -approximate K -medians algorithm when implementing Algorithm 2, instead of the exact K -median algorithm which is NP-hard [87]. See e.g. [76] for a more detailed discussion comparing the $(1 + \epsilon)$ -approximate K -medians and the exact K -medians algorithms. Theorem 4.2 still holds for approximate K -medians.

REMARK 4.3. Community detection using RSVD was also studied in [130]. In particular, Theorem 1 in [130] shows that if $n\rho_n = \omega(\log n)$ then weak recovery is possible using $\hat{\mathbf{U}}_g$, provided that one choose $\tilde{k} \geq k + 4$ and $g = \Omega(n^\epsilon)$ for any fixed but arbitrary $\epsilon > 0$. Recall that weak recovery only requires the *proportion* of mis-clustered vertices to converge to 0. Comparing the two results, we see the sparsity condition $n\rho_n = \omega(\log n)$ in [130] is less stringent than the condition $n\rho_n = \Omega(n^\epsilon)$ in Theorem 4.2 while the exact recovery with fixed g in Theorem 4.2 is a much stronger guarantee (and also is more computationally efficient) than weak recovery in [130].

The proof of Theorem 4.2 is based on the $\ell_{2 \rightarrow \infty}$ perturbation bounds given in Eq. (4.2). More specifically, if $g > \beta^{-1}$ then, with high probability,

$$(4.4) \quad \sqrt{nd}d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \rightarrow 0.$$

If Eq. (4.4) holds then the arguments for exact recovery of $\boldsymbol{\tau}$ using $\hat{\mathbf{U}}$ can be easily adapted to show exact recovery using $\hat{\mathbf{U}}_g$; see the proof of Theorem 2.6 in [82] or the proof of Theorem 5.2 in [75] for some examples of these types of arguments. Furthermore, while $g = \lceil \beta^{-1} \rceil$ is sufficient to show exact recovery using $\hat{\mathbf{U}}_g$ when β^{-1} is not an integer, in practice we recommend choosing the slightly larger value of $g = \lceil \beta^{-1} \rceil + 1$. Indeed, as shown in Section 4.2, the convergence rate for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ when $g = \lceil \beta^{-1} \rceil + 1$ is generally faster than that for $g = \lceil \beta^{-1} \rceil$, and this can lead to better finite-sample performance.

Theorem 4.2 also indicates that smaller values of β will require larger number of power iterations g and hence the computational cost of Algorithm 2 increases as the graphs become sparser. In other words, there is an inherent trade-off between the sparsity of the network and the computation cost needed to approximate its eigenvectors. In addition, as we allude to earlier in Remark 4.3, Algorithm 2 does not guarantee exact recovery of $\boldsymbol{\tau}$ in the very sparse asymptotic regime where $n\rho_n = \omega(\log n)$ and this is due entirely to the use of $\hat{\mathbf{U}}_g$, as an approximation for $\hat{\mathbf{U}}$. More specifically, if $n\rho_n = \omega(\log n)$ then [3, 75, 112] showed that exact recovery of $\boldsymbol{\tau}$ is possible using $\hat{\mathbf{U}}$. In contrast, by Theorem 4.1, we see that neither $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ nor $n^{1/2}d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ converge to 0 for any finite $g \geq 1$. Hence, to guarantee exact recovery, we will need to let g grows with n and this will require a more careful analysis of Algorithm 1 to remove the effect of the condition number ψ when $g \rightarrow \infty$. More specifically, instead of estimating all k leading singular vectors simultaneously, we might need to estimate these singular vectors in stages, starting from the first leading singular vector. After each stage we will project $\hat{\mathbf{M}}$ onto the orthogonal complement of the singular vectors estimated thus far before proceeding with the estimation of the singular vectors in the next stage. Finally, while it is certainly the case that, asymptotically, $\log n = o(n^\epsilon)$ for any $\epsilon > 0$, in practice $\log n \geq n^{0.1}$ for all $n \leq 10^{15}$. Hence, while asymptotic results in the $\omega(\log n)$ regime are interesting, they need not lead to better understanding of RSVD for practical applications.

4.4. *Normal approximation for $\hat{\mathbf{U}}_g$.* We now consider the row-wise fluctuations of $\hat{\mathbf{U}}_g$. Recall the setting at the beginning of this section where $\hat{\mathbf{M}}$ is the adjacency matrix of an edge independent random graph whose edge probabilities matrix \mathbf{M} has rank k^* matrix with k^* bounded, $S_n = \|\mathbf{M}\| \asymp n^\beta$ and $E_n \asymp n^{\beta/2}$. Suppose furthermore that the entries of \mathbf{M} are homogeneous so that $\max_{ij} m_{ij} \asymp \min_{ij} m_{ij}$. Then $\|\mathbf{M}\|_F = n\|\mathbf{M}\|_{\max}$ and hence $m_{ij} \asymp n^{\beta-1}$ for all i, j . In addition the $\{m_{ii}^{(2)}\}_{i=1}^n$ are also homogeneous; here $m_{ii}^{(2)}$ denote the diagonal elements of \mathbf{M}^2 . As $m_{ii}^{(2)} = \|[\mathbf{U}\boldsymbol{\Lambda}]_i\|^2$ and \mathbf{M} has bounded condition number, we conclude that \mathbf{U} has bounded coherence, i.e., $\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim k^{1/2}n^{-1/2}$. The above conditions are satisfied for many random graph models including stochastic blockmodels and their degree-corrected and mixed-membership stochastic variants [63, 69] as well as (generalized) random dot product graphs [103, 128].

Recall Assumption 3. As $E_n \asymp n^{\beta/2}$, take $\sigma_n = n^{(\beta-1)/2}$. We now verify Eq (3.13). Note, $S_n\sigma_n^{-1}[\mathbf{E}\mathbf{U}\boldsymbol{\Lambda}^{-1}]_i = n^{(1+\beta)/2}\boldsymbol{\Lambda}^{-1}\sum_{j=1}^n e_{ij}\mathbf{u}_j$ where e_{ij} is the ij th entry of \mathbf{E} and \mathbf{u}_j is the j th row of \mathbf{U} . For a fixed i , the $\{e_{ij}\}_{j \leq n}$ are independent mean 0 random variables and hence, by the Lindeberg-Feller central limit theorem, we have

$$S_n\sigma_n^{-1}\boldsymbol{\Gamma}_i^{-1/2}[\mathbf{E}\mathbf{U}\boldsymbol{\Lambda}^{-1}]_i = n^{(1+\beta)/2}\boldsymbol{\Lambda}^{-1}\sum_{j=1}^n e_{ij}\mathbf{u}_j \rightsquigarrow \mathcal{N}(\mathbf{0}, \mathbf{I}),$$

where Γ_i is the $k^* \times k^*$ matrix of the form

$$\Gamma_i = n^{(1+\beta)} \mathbf{\Lambda}^{-1} \sum_{j=1}^n \text{Var}[e_{ij}] \mathbf{u}_j \mathbf{u}_j^\top \mathbf{\Lambda}^{-1} = n^{(1+\beta)} \mathbf{\Lambda}^{-1} \left\{ \sum_{j=1}^n m_{ij} (1 - m_{ij}) \mathbf{u}_j \mathbf{u}_j^\top \right\} \mathbf{\Lambda}^{-1}.$$

Note that Γ_i is non-degenerate. Indeed, as \mathbf{M} is homogeneous, we have

$$\Gamma_i \succeq n^{(1+\beta)} \mathbf{\Lambda}^{-1} \min_{kl} m_{kl} \sum_j \mathbf{u}_j \mathbf{u}_j^\top \mathbf{\Lambda}^{-1} \succeq n^{2\beta} \mathbf{\Lambda}^{-2} \succeq c \mathbf{I},$$

for some fixed constant $c > 0$. In summary, Assumption 3 is satisfied and we have the following normal approximations for the row-wise fluctuations of $\hat{\mathbf{U}}$, namely there exists a sequence of orthogonal matrices \mathbf{W}_n such that for any fixed index i ,

$$S_n \sigma_n^{-1} \Gamma_i^{-1/2} (\mathbf{W}_n [\hat{\mathbf{U}}]_i - [\mathbf{U}]_i) \rightsquigarrow \mathcal{N}(0, \mathbf{I})$$

as $n \rightarrow \infty$. This result was derived previously in [21, 42, 52, 103]; see for example Theorem 2.8 in [42]. Theorem 3.4 implies the same limiting distribution for $\hat{\mathbf{U}}_g$, i.e., for any $g \geq 2 + \beta^{-1}$ there exists a sequence of orthogonal matrix \mathbf{W}_n^* such that

$$S_n \sigma_n^{-1} \Gamma_i^{-1/2} (\mathbf{W}_n^* [\hat{\mathbf{U}}_g]_i - [\mathbf{U}]_i) \rightsquigarrow \mathcal{N}(0, \mathbf{I})$$

as $n \rightarrow \infty$. See Section S.1.1.3 for a numerical illustration of this convergence.

5. Additional applications. In this section we apply our main theorems to two other high-dimensional statistical problems, namely, symmetric matrix completion and PCA with missing data. For both of these problems we show that the error rates achieved by the approximate singular vectors $\hat{\mathbf{U}}_g$ is (almost) identical to that for the exact singular vectors $\hat{\mathbf{U}}$ of the noisily observed matrices.

5.1. Matrix completion with noises. Let $\mathbf{T} \in \mathbb{R}^{n \times n}$ be a matrix whose entries are only partially and noisily observed. Such matrix occurs in many real-world applications, including the famous Netflix challenge [98]. As another example, if \mathbf{T} is an Euclidean distance matrix (EDM) between n points in \mathbb{R}^d then $\text{rk}(\mathbf{T}) \leq d + 2$ and it is commonly the case that \mathbf{T} is noisily observed [8, 40, 54, 65]. Finally, if \mathbf{T} is a signal correlation matrix between multiple remote sensors [107] then \mathbf{T} is usually only partial observed due to power constraints [31].

For this paper we assume that $\mathbf{T} \in \mathbb{R}^{n \times n}$ is symmetric, rank k , and we observed

$$(5.1) \quad \hat{\mathbf{T}} = \mathcal{P}_\Omega(\mathbf{T} + \mathbf{N}) := \Omega \circ (\mathbf{T} + \mathbf{N}).$$

Here \mathbf{N} is an unobserved $n \times n$ noise matrix, Ω is a symmetric binary matrix, and \circ denote the Hadamard product. The matrix \mathbf{N} and Ω induces noise and missingness in the entries of $\hat{\mathbf{T}}$, respectively. We shall assume, for ease of exposition, that the (upper triangle) entries of \mathbf{N} are iid $\mathcal{N}(0, \sigma^2)$ random variables while the (upper triangular) entries of Ω are iid Bernoulli random variables with success probability p . The aim of matrix completion is to recover \mathbf{T} from $\hat{\mathbf{T}}$. Since $\mathbb{E}[p^{-1} \hat{\mathbf{T}}] = \mathbf{T}$, one simple and widely used estimate for \mathbf{T} is given by $p^{-1} \hat{\mathbf{T}}_{(k)}$ where $\hat{\mathbf{T}}_{(k)}$ is the truncated rank- k SVD of $\hat{\mathbf{T}}$; see [3, 19, 23, 28, 70] and the references therein.

In many real-world applications, the dimensions of $\hat{\mathbf{T}}$ can be rather large and yet $\hat{\mathbf{T}}$ can be quite sparse compared to \mathbf{T} , i.e., the number of non-zero entries of $\hat{\mathbf{T}}$ is much smaller than n^2 . It is thus computationally attractive to approximate the left singular vectors of $\hat{\mathbf{T}}$ using randomized algorithms such as *rs*-RSVD. More specifically, let $\hat{\mathbf{U}}_g$ be the output of Algorithm 1 with $\hat{\mathbf{M}} = \hat{\mathbf{T}}$ for some properly specified choices of k, \tilde{k}, a_n and g . We view $\hat{\mathbf{U}}_g$ as an approximation for $\hat{\mathbf{U}}$, and thus as an estimate of \mathbf{U} . Given $\hat{\mathbf{U}}_g$ we compute a rank- k

Algorithm 3: RSVD-based Matrix Completion

Input: Observed matrix $\hat{\mathbf{T}}$, rank k . Sampling probability p . Integers $a_n, g \geq 1$ and $\tilde{k} \geq K$.

1. Get $\hat{\mathbf{U}}_g, \tilde{\mathbf{M}}$ via Algorithm 1 with $\hat{\mathbf{M}} = p^{-1}\hat{\mathbf{T}}$, rank k , sampling times a_n , power g and dimension \tilde{k} ;
2. Set $\hat{\mathbf{T}}_g = \tilde{\mathbf{M}}$ as either $\hat{\mathbf{T}}_g = p^{-1}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$ or $\hat{\mathbf{T}}_g = (2p)^{-1}(\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}} + \hat{\mathbf{T}}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top)$;

Output: Approximate singular vectors $\hat{\mathbf{U}}_g$ of $\hat{\mathbf{T}}$ and estimate $\hat{\mathbf{T}}_g$ of \mathbf{T} .

approximation for $\hat{\mathbf{T}}$ via $\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$. We can then take $\hat{\mathbf{T}}_g := p^{-1}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$ as an estimate for \mathbf{T} or, in the event that we desire symmetry, take $2^{-1}(\hat{\mathbf{T}}_g + \hat{\mathbf{T}}_g^\top)$ as an estimate for \mathbf{T} . The resulting algorithm, termed RSVD-based matrix completion, is presented in Algorithm 3.

Using the ℓ_2 and $\ell_{2 \rightarrow \infty}$ perturbation results in Theorem 3.2 and 3.3 and the entrywise concentration in Corollary 3.2, we derive the error bounds for the estimates $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{T}}_g$. Furthermore, by extending Theorem 3.4, we establish the general entrywise limiting distribution for the RSVD-based low-rank approximation $\tilde{\mathbf{M}}$, in Theorem S.6.1 of the supplementary document. This result allows us to construct confidence interval for any $[\hat{\mathbf{T}}_g]_{ij}$ with $(i, j) \in [n]^2$. For ease of exposition we shall assume that both k and ψ are bounded, and that p is known. These are typical assumptions in the literature. If p is unknown then, as the entries of \mathbf{T} are assumed to be missing completely at random, it can be consistently estimated by calculating the proportion of missing entries among all the entries in \mathbf{T} . The resulting \hat{p} converges to p at the rate of $n^{-1}p^{-1/2}$ and has no effect on the theoretical results in Theorem 5.1

THEOREM 5.1. *Let \mathbf{T} be a symmetric $n \times n$ matrix with $k, \psi \lesssim 1$. Let $\hat{\mathbf{T}}$ be a noisily observed version of \mathbf{T} generated according to Eq. (5.1) for some known value of $p \in (0, 1]$. Suppose there exists constants $\beta_\Delta > 0$ and $\beta \in (0, 1]$ such that*

$$(5.2) \quad np \gtrsim n^\beta \quad \text{and} \quad \frac{1}{\sqrt{np}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \lesssim n^{-\beta_\Delta}.$$

Let $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{T}}_g$ be obtained from Algorithm 3 wherein we chose: (i) $\tilde{k} \geq k$; (ii) $a_n \geq \max\{[C\tilde{k}^{-1} \log n], 1\}$ and (iii) $g \geq 1 + (2\beta_\Delta)^{-1}$. Denote $\vartheta = \max\{1, \tilde{k}^{-1/2} \log^{1/2} n\}$.

(Error Bounds) *We then have, with probability at least $1 - n^{-5}$, that*

$$(5.3) \quad \begin{aligned} d_2(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \frac{\vartheta}{\sqrt{np}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|}, \\ d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \frac{\vartheta(\log n)^{\delta^*}}{\sqrt{np}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\ \frac{1}{n} \|\hat{\mathbf{T}}_g - \mathbf{T}\|_{\text{F}} &\lesssim \frac{\vartheta}{\sqrt{np}} (\|\mathbf{T}\|_{\max} + \sigma), \\ \|\hat{\mathbf{T}}_g - \mathbf{T}\|_{\max} &\lesssim \frac{\sqrt{n}\vartheta(\log n)^{\delta^*}}{\sqrt{p}} (\|\mathbf{T}\|_{\max} + \sigma) \|\mathbf{U}\|_{2 \rightarrow \infty}^2, \end{aligned}$$

where δ^* is as defined in the statement of Theorem 3.3.

(Entrywise limiting distribution) *Suppose, in addition to the above assumptions, that \mathbf{T} is homogeneous, i.e., $\min_{k\ell} |T_{k\ell}| \asymp \|\mathbf{T}\|_{\max}$. Let $\zeta_{k\ell} = [\mathbf{U}\mathbf{U}^\top]_{k\ell}$ and denote the variance*

of $[\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}$ by

$$\begin{aligned}
 v_{ij}^* &:= \sum_{\ell \neq j} \mathbb{E}([\mathbf{E}]_{i\ell}^2) \zeta_{\ell j}^2 + \sum_{\ell \neq i} \mathbb{E}([\mathbf{E}]_{\ell j}^2) \zeta_{i\ell}^2 + \mathbb{E}([\mathbf{E}]_{ij}^2) (\zeta_{ii} + \zeta_{jj})^2 \\
 (5.4) \quad &= \frac{1}{p} \sum_{\ell \neq j} \{(1-p)T_{i\ell}^2 + \sigma^2\} \zeta_{\ell j}^2 + \frac{1}{p} \sum_{\ell \neq i} \{(1-p)T_{\ell j}^2 + \sigma^2\} \zeta_{i\ell}^2 \\
 &\quad + \frac{1}{p} \{(1-p)T_{ij}^2 + \sigma^2\} (\zeta_{ii} + \zeta_{jj})^2.
 \end{aligned}$$

Choose \tilde{k} and a_n as above and let $g \geq 2 + (2\beta_\Delta)^{-1}$. Then for any given pair $(i, j) \in \mathbb{R}^2$, we have

$$(5.5) \quad (v_{ij}^*)^{-1/2} [\hat{\mathbf{T}}_g - \mathbf{T}]_{ij} \rightsquigarrow \mathcal{N}(0, 1)$$

as $n \rightarrow \infty$. Furthermore, for any finite set of indices $\mathcal{I} = \{(i_1, j_1), \dots, (i_m, j_m)\}$, the collection of random variables $\{[\hat{\mathbf{T}}_g - \mathbf{T}]_{i_k j_k}\}_{k=1}^m$ are (asymptotically) mutually independent.

(Entrywise confidence interval) Suppose above assumptions hold. Let $\hat{\zeta}_{k\ell} = [\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top]_{k\ell}$. Define the RSVD-based estimate of v_{ij}^* by

$$(5.6) \quad \hat{v}_{ij} := \sum_{\ell \neq j} [\hat{\mathbf{E}}]_{i\ell}^2 \hat{\zeta}_{\ell j}^2 + \sum_{\ell \neq i} [\hat{\mathbf{E}}]_{\ell j}^2 \hat{\zeta}_{i\ell}^2 + [\hat{\mathbf{E}}]_{ij}^2 \{\hat{\zeta}_{ii} + \hat{\zeta}_{jj}\}^2$$

where $\hat{\mathbf{E}} := \hat{\mathbf{T}}_g - p^{-1}\hat{\mathbf{T}}$. Then for any indices pair (i, j) we have

$$(5.7) \quad (\hat{v}_{ij})^{-1/2} [\hat{\mathbf{T}}_g - \mathbf{T}]_{ij} \rightsquigarrow \mathcal{N}(0, 1).$$

We now compare Theorem 5.1 with existing results in the literature. Recall that the prototypical spectral estimate for \mathbf{T} is given by $\hat{\mathbf{T}}_S = p^{-1}\hat{\mathbf{T}}_{(k)}$ where $\hat{\mathbf{T}}_k$ is the truncated rank- k SVD of $\hat{\mathbf{T}}$; for conciseness we refer to $p^{-1}\hat{\mathbf{T}}_k$ as the deterministic SVD-based estimator for \mathbf{T} . First suppose that Eq. (5.2) holds for some constants $\beta \in (0, 1]$ and $\beta_\Delta > 0$. We then have

$$\begin{aligned}
 \|\hat{\mathbf{T}}_S - \mathbf{T}\|_F &\lesssim \frac{\|\mathbf{T}\|_{\max} + \sigma}{\sqrt{np}}, \\
 (5.8) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U}) &\lesssim \frac{(\log n)^{1/2} n(\|\mathbf{T}\|_{\max} + \sigma)}{\sqrt{np} |\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
 \|\hat{\mathbf{T}}_S - \mathbf{T}\|_{\max} &\lesssim (\log n)^{1/2} (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2,
 \end{aligned}$$

with high probability. The above bound for $\|\hat{\mathbf{T}}_S - \mathbf{T}\|_F$ is from Theorem 1.1 and Theorem 1.3 of [70] while the remaining bounds in Eq. (5.8) are from Theorem 3.4 of [3].

Now consider the bounds for $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{T}}_g$ given in Theorem 5.1. For ease of presentation we choose $\tilde{k} = \Omega(\log n)$ and $g > 1 + (2\beta_\Delta)^{-1}$. These choices for \tilde{k} and g allow us to set $\vartheta = 1$ and $\delta^* = \delta$ for any arbitrary $\delta > 1$. Eq. (5.3) then implies

$$\begin{aligned}
 \|\hat{\mathbf{T}}_g - \mathbf{T}\|_F &\lesssim \frac{\|\mathbf{T}\|_{\max} + \sigma}{\sqrt{np}}, \\
 (5.9) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \frac{(\log n)^\delta n(\|\mathbf{T}\|_{\max} + \sigma)}{\sqrt{np} |\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
 \|\hat{\mathbf{T}}_g - \mathbf{T}\|_{\max} &\lesssim (\log n)^\delta (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2,
 \end{aligned}$$

with high probability. Comparing Eq. (5.8) and Eq. (5.9) we see that the rs -RSVD estimate $\hat{\mathbf{T}}_g$ achieves the same Frobenius norm error rate as that for the deterministic SVD estimate $\hat{\mathbf{T}}_S$. Note that the Frobenius norm error for $\hat{\mathbf{T}}_S$ is rate-optimal whenever $\sigma \lesssim \|\mathbf{T}\|_{\max}$ [3, 19]. The max-norm error rate for $\hat{\mathbf{T}}_g$ is also almost identical to that of $\hat{\mathbf{T}}_S$ with the only difference being the extra $(\log n)^{1/2+\epsilon}$ factor for $\hat{\mathbf{T}}_g$; here ϵ is an arbitrary positive constant. Similarly, $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ converges to 0 at the same rate as $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$ up to the $(\log n)^{1/2+\epsilon}$ factor. These $(\log n)^{1/2+\epsilon}$ extra factors for $\hat{\mathbf{T}}_g$ can be removed with more careful book keeping; see Remark 3.5 for more details. If \tilde{k} is either bounded or diverge at a slower rate than $\log n$, or when $g = 1 + (2\beta_\Delta)^{-1}$, then Eq. (5.9) still holds but with the $(\log n)^\delta$ factor replaced by a $(\log n)^{\delta^*}$ factor for some finite $\delta^* > \delta$.

We next discuss the limiting distributions and confidence intervals given in Theorem 5.1. We had assumed that \mathbf{T} is homogeneous; this assumption was also used in Theorem 4.12 in [28] for the estimator $\hat{\mathbf{T}}_S$. The main role of the homogeneity assumption is to guarantee that the entrywise noise levels are roughly on the same order, and thereby simplify the statement of Theorem 5.1. Similar assumption are also used in the distributional theory for PCA with missing data; see e.g. Assumption 1 in [125]. Our general result, namely Theorem S.6.1, does not require \mathbf{T} to be homogenous. The condition $g \geq 2 + (2\beta_\Delta)^{-1}$ in Theorem 5.1 is then similar to that in Theorem 3.4 for the row-wise limiting distribution of $\hat{\mathbf{U}}_g$, i.e., we need $g \geq 2 + (2\beta_\Delta)^{-1}$ to guarantee that the entrywise fluctuation of $[\hat{\mathbf{T}}_g - \mathbf{T}]_{ij}$ is asymptotically equivalent to the entrywise fluctuation of $[\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}$. For any given pair $(i, j) \in \mathbb{R}^2$, we can thus quantify the uncertainty of the $[\hat{\mathbf{T}}]_{ij}$ by constructing the confidence interval:

$$([\hat{\mathbf{T}}_g]_{ij} - z_{\alpha/2} \sqrt{\hat{v}_{ij}}, [\hat{\mathbf{T}}_g]_{ij} + z_{\alpha/2} \sqrt{\hat{v}_{ij}})$$

where $z_{\alpha/2}$ is the $(1 - \alpha/2)$ quantile of $\mathcal{N}(0, 1)$. The resulting confidence interval has (asymptotic) coverage of $1 - \alpha$. We emphasize that the estimated variance \hat{v}_{ij} are computed using only the rs -RSVD outputs $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{T}}_g$; see Eq. (5.6).

REMARK 5.1. The conditions posited in Theorem 5.1 for the estimator $\hat{\mathbf{T}}_g$ is slightly more restrictive than those for $\hat{\mathbf{T}}_S$ in the current literature. Recall that from Eq. (5.2), we have $np \lesssim n^\beta$ for some arbitrary $\beta \in (0, 1]$. Eq. (5.2) also implies a lower bound for the signal-to-noise ratio of the form

$$S_n/E_n \lesssim \left\{ \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{1}{\sqrt{np}} \right\}^{-1} \lesssim n^{\beta_\Delta},$$

for some arbitrary $\beta_\Delta > 0$. In contrast, the conditions for $\hat{\mathbf{T}}_S$ given in Theorem 3.4 of [3] is that $np \lesssim \log n$ and $S_n/E_n \lesssim (\log n)^{1/2}$. The conditions $np \lesssim n^\beta$ and $S_n/E_n \lesssim n^{\beta_\Delta}$ in Theorem 5.1 is analogous to the condition $n\rho_n = \omega(\log n)$ for exact recovery of SBM using $\hat{\mathbf{U}}_g$. In particular for matrix completion using rs -RSVD to work in the $np = \omega(\log n)$ and $S_n/E_n = \omega(\log n)$ regime we will need to have g diverging with n and this requires a more careful analysis of RSVD to remove the effect of the condition number ψ as g increases. Furthermore, very large and/or diverging values of g also reduces the effectiveness of RSVD procedures for large-scale matrix computations, especially when the matrices are too large to store in fast memory as it requires a possibly excessive number of passes through the data. Finally we note that β_Δ depends on the parameters σ , $\lambda_k(\mathbf{T})$, $\|\mathbf{T}\|_{\max}$ and p . In particular, if \mathbf{T} is homogeneous (as assumed in [28]) so that $\min_{ij} |T_{ij}| \asymp \|\mathbf{T}\|_{\max}$, then $n(\|\mathbf{T}\|_{\max} + \sigma) \asymp |\lambda_k(\mathbf{T})|$ and Eq. (5.2) simplifies to the condition that $\sqrt{np} \lesssim n^{\beta_\Delta}$ and hence $\beta_\Delta = \frac{1}{2}\beta$.

Algorithm 4: RSVD-based PCA with Missing Data

Input: Partially observed data matrix $\mathcal{P}_\Omega(\mathbf{X})$. Sampling probability p . Integers $a_n \geq 1, g \geq 1$ and $\tilde{k} \geq k$.

1. Form $\mathbf{Q} = \mathcal{P}_{\text{off-diag}} \left[p^{-2} \mathcal{P}_\Omega(\mathbf{X}) \mathcal{P}_\Omega^\top(\mathbf{X}) \right]$;

2. Generate $\hat{\mathbf{U}}_g$ via Algorithm 1 with $\hat{\mathbf{M}} = \mathbf{Q}$, rank k , a_n repeated samples, power g and sketching dimension \tilde{k} ;

Output: Principal Component Recovery $\hat{\mathbf{U}}_g$.

5.2. *PCA with missing data.* We now consider principal components estimation with missing data. In particular we consider the following data generating mechanism from [18]:

$$(5.10) \quad \mathbf{X} = \mathbf{B}\mathbf{F} + \mathbf{N}.$$

Here $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_m] \in \mathbb{R}^{d \times m}$ is a data matrix with $\{\mathbf{x}_i\}$ being d -dimensional iid random vectors, \mathbf{B} is a $d \times k$ matrix, $\mathbf{F} = [\mathbf{f}_1, \dots, \mathbf{f}_m] \in \mathbb{R}^{k \times m}$ is a $k \times m$ matrix whose entries are iid $\mathcal{N}(0, 1)$ random variables, and \mathbf{N} is a $d \times m$ matrix whose entries are iid $\mathcal{N}(0, \sigma^2)$ random variables. We emphasize that \mathbf{B} , \mathbf{N} and \mathbf{F} are mutually independent. We can see that the rows of \mathbf{X} are iid random vectors with mean $\mathbf{0}$ and covariance matrix $\text{Var}[\mathbf{X}] = m\mathbf{B}\mathbf{B}^\top + \sigma^2\mathbf{I}_d$. We denote the SVD of $\mathbf{B}\mathbf{B}^\top$ by $\mathbf{B}\mathbf{B}^\top = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top$ where $\mathbf{U} \in \mathbb{O}_{d \times k}$ and $\mathbf{\Lambda} = \text{diag}(\lambda_1, \dots, \lambda_k)$; the columns of $\mathbf{U} \in \mathbb{R}^{d \times k}$ are leading principal components.

Due to sampling issues and/or privacy-preserving intention, it is often the case that only a partial subset of the entries in \mathbf{X} are observed. More specifically let Ω be a $d \times m$ binary matrix whose entries are iid Bernoulli random variables with success probability p . Then, instead of observing \mathbf{X} , we only observed $\mathcal{P}_\Omega(\mathbf{X}) = \Omega \circ \mathbf{X}$ where \circ denote the Hadamard product between matrices. Given an observed $\mathcal{P}_\Omega(\mathbf{X})$, [18] propose the following spectral procedure for recovering the principle components \mathbf{U} . First form the symmetric matrix

$$(5.11) \quad \mathbf{Q} = \mathcal{P}_{\text{off-diag}} \left[\frac{1}{p^2} \mathcal{P}_\Omega(\mathbf{X}) \mathcal{P}_\Omega^\top(\mathbf{X}) \right],$$

where $\mathcal{P}_{\text{off-diag}}[\cdot]$ sets diagonal entries of the corresponding matrix to zero. Next compute the $d \times k$ matrix $\hat{\mathbf{U}}$ whose columns are the left singular vectors corresponding to the k largest singular values of \mathbf{Q} ; $\hat{\mathbf{U}}$ serves as an estimate of \mathbf{U} . The authors of [18] then derive high-probability error bounds for $d_2(\hat{\mathbf{U}}, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$.

As the dimension d can be reasonably large compared to the number of samples m while the number of non-zero entries in $\mathcal{P}_\Omega(\mathbf{X})$ can be much smaller than md , we consider replacing the singular vectors $\hat{\mathbf{U}}$ of \mathbf{Q} by the approximate singular vectors $\hat{\mathbf{U}}_g$ computed using RSVD; see Algorithm 4. The following result shows that by choosing either $\tilde{k} \asymp \log d$ or $a_d \asymp \log d$, the approximate singular vectors $\hat{\mathbf{U}}_g$ achieve the same estimation rate as that for $\hat{\mathbf{U}}$ given in [18]. For ease of presentation, and following [18], we shall assume that \mathbf{B} has a bounded condition number, i.e., $\kappa := \lambda_1/\lambda_k \lesssim 1$.

THEOREM 5.2. *Suppose \mathbf{X} is a $d \times m$ matrix generated according to Eq. (5.10). Let $\mu = dr^{-1} \|\mathbf{U}\|_{2 \rightarrow \infty}^2$ denote the coherence parameter for \mathbf{U} and let $s_* = \log(m + d)$. Suppose there exist constants \tilde{c}_0 and \tilde{c}_1 such that $k \leq \tilde{c}_1 \frac{d}{\mu}$ and m satisfies the sample size condition*

$$(5.12) \quad m \geq \tilde{c}_0 \max \left\{ \frac{\mu^2 k^2 s_*^6}{dp^2}, \frac{\mu k s_*^3}{p}, \frac{\sigma^4 s_*^2}{\lambda_k^2 p^2}, \frac{\sigma^2 d s_*}{\lambda_k p} \right\}.$$

Define $\vartheta' = \max\{1, \tilde{k}^{-1/2} \log^{1/2} d\}$ and

$$(5.13) \quad \mathcal{E} := \frac{\mu k s_*^2}{(md)^{1/2} p} + \frac{(\mu k)^{1/2} s_*}{(md)^{1/2}} + \frac{\sigma^2 d^{1/2} s_*}{\lambda_k^2 m^{1/2} p} + \frac{\sigma d^{1/2} s_*^{1/2}}{(\lambda_k m p)^{1/2}} + \frac{\mu k}{d}.$$

Suppose that $\mathcal{E} \lesssim d^{-\beta_{\mathcal{E}}}$ for some $\beta_{\mathcal{E}} \geq 0$. Let $\hat{\mathbf{U}}_g$ be generated via Algorithm 4, wherein we choose: (i) $\tilde{k} \geq c_{\text{gap}} k$ for any fixed $c_{\text{gap}} > 1$; (ii) $a_d \geq \max\{C \tilde{k}^{-1} \log d, 1\}$ and (iii) $g \geq 1 + (2\beta_{\mathcal{E}})^{-1}$. Then with probability at least $1 - d^{-5}$, we have $d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \vartheta' \mathcal{E}$.

REMARK 5.2. The assumption of bounded κ is widely considered in the literature, see e.g., the discussion prior to Eq.(4.17) in [18] and see also [25, 29, 64, 109]. For example, $\kappa \lesssim 1$ is satisfied with high-probability when \mathbf{B} is generated from a standard Gaussian matrix (see e.g. [119]). Note that the sample size requirement in Eq. (5.12) is exactly the same as that in Eq. (4.14) of [18] for bounded κ . If κ grows with d then the sample size requirement in Eq. (5.12) depend on polynomial factors κ^c of κ and these factors can be extracted from the proof of Theorem 5.2 with a bit more careful book-keeping.

REMARK 5.3. Comparing the Theorem 5.2 with Corollary 4.3 in [18], we see that there exists a finite $g \geq 2$ such that $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_2(\hat{\mathbf{U}}, \mathbf{U})$ converges to 0 at the same rate \mathcal{E} , up to a $\log^{1/2} d$ factor; furthermore this logarithmic gap can be removed by choosing $\tilde{k} \asymp \log d$. The condition $g \geq 1 + (2\beta_{\mathcal{E}})^{-1}$ in Theorem 5.2 indicates that an appropriate value for g will depend on the relationship between p, d, σ and m , e.g., larger dimensions d (compared to m) will require larger power iterations g . In particular, suppose $\mu = \sigma = p = \lambda_1 = k = 1$ and $d \asymp m^{\beta_d}$ for some $\beta_d \in (0, 1)$; here $\beta_d < 1$ is necessary for Eq. (5.12) to hold. Then from the definition \mathcal{E} in Eq. (5.13), we have, after some straightforward algebra, that

$$\mathcal{E} \asymp d^{-(\beta_d^{-1}-1)/2} \log d + d^{-1} \lesssim \begin{cases} d^{-(\beta_d^{-1}-1)/2} \log d & \text{if } \beta_d \in [1/3, 1), \\ d^{-1} & \text{if } \beta_d \in [0, 1/3]. \end{cases}$$

To attain the rate in Eq. (5.13), we thus need $g \geq \frac{1}{1-\beta_d} + \epsilon$, where $\epsilon > 0$ is any small constant.

We note that Theorem 5.2 only provides perturbation bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and not more refined results such as perturbation bound for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. This omission is due mainly to the noise structure in the matrix \mathbf{Q} . More specifically if we take $\mathbf{M} = (\mathbf{B}\mathbf{F})(\mathbf{B}\mathbf{F})^\top$ and view \mathbf{Q} as the noisily observed version of \mathbf{M} then $\mathbf{E} = \mathbf{Q} - \mathbf{M}$ is of the form

$$\mathbf{E} = \mathcal{P}_{\text{off-diag}} \left[\frac{1}{p^2} \mathcal{P}_\Omega(\mathbf{X}) \mathcal{P}_\Omega^\top(\mathbf{X}) \right] - \mathbf{B}\mathbf{F}(\mathbf{B}\mathbf{F})^\top.$$

The (upper triangular) entries of \mathbf{E} are therefore not mutually independent and this leads to non-trivial technical challenges in deriving $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ bounds for \mathbf{Q} . In particular, Assumption 2 need not hold. In contrast, existing bounds for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$ are based on leave-two-out analysis which is a variant of leave-one-out analysis; see Appendix G in [125]. Similar to the discussion in Remark 3.6, this leave-two-out approach is not directly applicable to the analysis of $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ because $\hat{\mathbf{U}}_g$ depends on $(\hat{\mathbf{M}}^g - \mathbf{M}^g)\mathbf{G}$ and the entrywise dependency structure in $(\hat{\mathbf{M}}^g - \mathbf{M}^g)$ is more complicated than that in \mathbf{E} .

Recent work show that we can estimate the principal components \mathbf{U} by using the k leading left singular vectors of $\mathcal{P}_\Omega(\mathbf{X})$ directly (as opposed to $\hat{\mathbf{U}}$ which uses the k leading eigenvectors of \mathbf{Q}); the resulting estimate has ℓ_2 and $\ell_{2 \rightarrow \infty}$ error bounds that are comparable to those for $\hat{\mathbf{U}}$ in the regime where either $m \lesssim d$ or m is not much larger (in magnitude) compared to d ; see Section 3.2 in [125] for more details. Note that $\mathcal{P}_\Omega(\mathbf{X})$ is a rectangular matrix whose

entries, conditional on \mathbf{F} , are mutually independent. We can thus apply the results for asymmetric rs -RSVD in Section S.2 in the Supplementary File, to derive perturbation bounds for the approximate singular vectors of $\mathcal{P}_\Omega(\mathbf{X})$. A sketch of this argument is given below.

Let $\tilde{\mathbf{U}}$ be the k leading singular vectors of $\hat{\mathbf{M}} := m^{-1/2}p^{-1}\mathcal{P}_\Omega(\mathbf{X})$ and let $\tilde{\mathbf{U}}_g$ be the rs -RSVD approximation of $\tilde{\mathbf{U}}$ obtained from Algorithm S.1 for $\hat{\mathbf{M}}$. Then $\mathbb{E}_{\mathbf{N}}[\hat{\mathbf{M}}] = \mathbf{B}\mathbf{F}$ where $\mathbb{E}_{\mathbf{N}}$ denote the expectation with respect to \mathbf{N} only. Define, with a slight abuse of notation,

$$\mathbf{M} = m^{-1/2}\mathbf{B}\mathbf{F} \text{ and } \mathbf{E} = \hat{\mathbf{M}} - \mathbf{M} = m^{-1/2}p^{-1}\mathcal{P}_\Omega(\mathbf{X}) - m^{-1/2}\mathbf{B}\mathbf{F}.$$

Denote $n := \max\{d, m\}$ and $\eta := \max\{n_2, \log n\}$. Then \mathbf{E} satisfies Assumption S.1 with $E_{d,m} \asymp \sqrt{n}\sigma$; see Lemma 1 in [125]. Now suppose $S_{d,m}/E_{d,m} \asymp \eta^{\beta_\Delta}$ for some constant $\beta_\Delta > 0$. Then \mathbf{E} also satisfies Assumption S.2; this claim follows from Remark S.2.2. Theorem S.2.1 then yields upper bounds for $d_2(\tilde{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\tilde{\mathbf{U}}_g, \mathbf{U})$ when $g \geq (4\beta_\Delta)^{-1}$; these bounds are the same, up to some logarithmic factors, as those given in Lemmas 1–4 and Theorem 1 of [125] for $d_2(\tilde{\mathbf{U}}, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\tilde{\mathbf{U}}, \mathbf{U})$.

6. Discussions. In this paper we analyzed the statistical behavior of RSVD algorithms under the assumption that the observed matrix $\hat{\mathbf{M}}$ is generated according to a low-rank “signal-plus-noise” model. In particular, we derived ℓ_2 and $\ell_{2 \rightarrow \infty}$ perturbation bounds for $\hat{\mathbf{U}}_g$, entrywise concentration bounds for $\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$, and normal approximations for the row-wise fluctuations of $\hat{\mathbf{U}}_g$. There are several directions to pursue for the future research. Firstly, our upper and lower bounds for d_2 and $d_{2 \rightarrow \infty}$ include factors depending on the rank k , condition number ψ and other logarithmic terms. These factors are generally sub-optimal and can be sharpened with a more refined analysis; see for example Remark 3.5 and the discussion after Theorem 4.2. Secondly, as we shown in Section 4.2, our d_2 and $d_{2 \rightarrow \infty}$ bounds and the corresponding phase transition are sharp whenever \mathbf{E} satisfies a certain trace growth condition. This condition is satisfied in the context of random graphs inference and it is of some interest to find other statistical inference problem where this condition also holds. Thirdly, the theoretical results of this paper currently assume that g is bounded and that k^* and ψ are either bounded or slowly growing. It is thus natural to extend these results to more general settings, such as when k^* grows with n at rate n^c for some $c \in (0, 1)$, when the noise matrix \mathbf{E} has more complicated entrywise dependence structure, when the SNR is only of order $\omega(\log^c n)$ for some $c > 1$, or when $g \rightarrow \infty$. We note that these extensions are related, e.g., $k^* = \Omega(n^c)$ for some c will, in general, require $g \rightarrow \infty$ for consistent estimation of \mathbf{U} using $\hat{\mathbf{U}}_g$. Fourthly, while subspace power iterations (as considered in this paper) is one of the most popular approach for RSVD, there are other approaches such as those based on Krylov subspaces [89] and perturbation bounds in ℓ_2 and $\ell_{2 \rightarrow \infty}$ for RSVD using Krylov subspaces may require quite different techniques than those presented here. Finally, many modern dataset are represented as tensors and are analyzed using higher-order SVD; these procedures generally flatten tensor data into matrices across different dimensions and then compute the truncated SVD of the resulting flattened matrices [39, 131]. Statistical analysis of RSVD in the context of noisy tensor data is thus of some theoretical and practical interest.

SUPPLEMENTARY MATERIAL

Supplementary File for “Perturbation Analysis of Randomized SVD and its Applications to High-dimensional Statistics”

The Supplementary File contains numerical experiments, additional theoretical results, and all technical lemmas and proofs.

REFERENCES

- [1] ABBE, E. (2017). Community detection and stochastic block models: recent developments. *Journal of Machine Learning Research* **18** 6446–6531.
- [2] ABBE, E., BANDEIRA, A. S. and HALL, G. (2016). Exact recovery in the stochastic blockmodel. *IEEE Transactions on Information Theory* **62** 471–487.
- [3] ABBE, E., FAN, J., WANG, K. and ZHONG, Y. (2020). Entrywise eigenvector analysis of random matrices with low expected rank. *The Annals of Statistics* **48** 1452–1474.
- [4] ACHLIOPTAS, D. and MCSHERRY, F. (2007). Fast computation of low-rank matrix approximations. *Journal of the ACM* **54**.
- [5] AHLWEDE, R. and WINTER, A. (2002). Strong converse for identification via quantum channels. *IEEE Transactions on Information Theory* **48** 569–579.
- [6] AHN, S. C. and HORENSTEIN, A. (2013). Eigenvalue ratio test for the number of factors. *Econometrica* **81** 1203–1227.
- [7] AIROLDI, E. M., BLEI, D. M., FIENBERG, S. E. and XING, E. P. (2008). Mixed membership stochastic blockmodels. *Journal of Machine Learning Research* **9** 1981–2014.
- [8] ALFAKIH, A. Y., KHANDANI, A. and WOLKOWICZ, H. (1999). Solving Euclidean distance matrix completion problems via semidefinite programming. *Computational Optimization and Applications* **12** 13–30.
- [9] ASKHAM, T. and KUTZ, J. N. (2018). Variable projection methods for an optimized dynamic mode decomposition. *SIAM Journal on Applied Dynamical Systems* **17** 380–416.
- [10] ATHREYA, A., TANG, M., PARK, Y. and PRIEBE, C. E. (2021). On estimation and inference in latent structure random graphs. *Statistical Science* **36** 68–88.
- [11] BANDEIRA, A. S. and VAN HANDEL, R. (2016). Sharp nonasymptotic bounds on the norm of random matrices with independent entries. *Annals of Probability* **44** 2479–2506.
- [12] BELKIN, M. and NIYOGI, P. (2003). Laplacian eigenmaps for dimensionality reduction and data representation. *Neural Computation* **15** 1373–1396.
- [13] BEREND, D. and TASSA, T. (2010). Improved Bounds on Bell Numbers and on Moments of Sums of Random Variables. *Probability and Mathematical Statistics* **30** 185–205.
- [14] BICKEL, P. J. and CHEN, A. (2009). A nonparametric view of network models and Newman-Girvan and other modularities. *Proceedings of the National Academy of Sciences of the United States of America* **106** 21068–21073.
- [15] BORDENAVE, C. (2019). Lectures on random matrix theory. available online at <https://www.math.univ-toulouse.fr/~bordenave/IMPA-RMT.pdf>.
- [16] CAI, T. T., MA, Z. and WU, Y. (2013). Sparse PCA: Optimal rates and adaptive estimation. *The Annals of Statistics* **41** 3074–3110.
- [17] CAI, T. T. and ZHANG, A. (2018). Rate-optimal perturbation bounds for singular subspaces with applications to high-dimensional statistics. *The Annals of Statistics* **46** 60–89.
- [18] CAI, C., LI, G., CHI, Y., POOR, H. V. and CHEN, Y. (2021). Subspace estimation from unbalanced and incomplete data matrices: $\ell_{2,\infty}$ statistical guarantees. *The Annals of Statistics* **49** 944–967.
- [19] CANDÈS, E. J. and PLAN, Y. (2010). Matrix completion with noise. *Proceedings of the IEEE* **98** 925–936.
- [20] CANDÈS, E. J. and RECHT, B. (2009). Exact matrix completion via convex optimization. *Foundations of Computational Mathematics* **9** 717–772.
- [21] CAPE, J., TANG, M. and PRIEBE, C. E. (2019a). Signal-plus-noise matrix models: eigenvector deviations and fluctuations. *Biometrika* **106** 243–250.
- [22] CAPE, J., TANG, M. and PRIEBE, C. E. (2019b). The two-to-infinity norm and singular subspace geometry with applications to high-dimensional statistics. *The Annals of Statistics* **47** 2405–2439.
- [23] CHATTERJEE, S. (2015). Matrix estimation by universal singular value thresholding. *The Annals of Statistics* **43** 177–214.
- [24] CHEN, Y., CHENG, C. and FAN, J. (2021). Asymmetry helps: Eigenvalue and eigenvector analyses of asymmetrically perturbed low-rank matrices. *The Annals of Statistics* **49** 435–458.
- [25] CHEN, Y., GUIBAS, L. and HUANG, Q. (2014). Near-Optimal Joint Object Matching via Convex Relaxation. In *Proceedings of the 31st International Conference on Machine Learning* 100–108.
- [26] CHEN, Y. and SUH, C. (2015). Spectral MLE: Top-k rank aggregation from pairwise comparisons. In *Proceedings of the 32nd International Conference on Machine Learning* 371–380.
- [27] CHEN, Y., FAN, J., MA, C. and WANG, K. (2019). Spectral method and regularized MLE are both optimal for top- K ranking. *The Annals of Statistics* **47** 2204–2235.
- [28] CHEN, Y., CHI, Y., FAN, J. and MA, C. (2021a). Spectral Methods for Data Science: A Statistical Perspective. *Foundations and Trends® in Machine Learning* **14** 566–806.

- [29] CHEN, Y., FAN, J., MA, C. and YAN, Y. (2021b). Bridging convex and nonconvex optimization in robust PCA: Noise, outliers and missing data. *The Annals of Statistics* **49** 2948–2971.
- [30] CHENG, C., WEI, Y. and CHEN, Y. (2021). Tackling small eigen-gaps: Fine-grained eigenvector estimation and inference under heteroscedastic noise. *IEEE Transactions on Information Theory* **67** 7380–7419.
- [31] CHENG, J., YE, Q., JIANG, H., WANG, D. and WANG, C. (2012). STCDG: An efficient data gathering algorithm based on matrix completion for wireless sensor networks. *IEEE Transactions on Wireless Communications* **12** 850–861.
- [32] COIFMAN, R. and LAFON, S. (2006). Diffusion maps. *Applied and Computational Harmonic Analysis* **21** 5–30.
- [33] CORMODE, G., GAROFALAKIS, M., HAAS, P. J. and JERMAINE, C. (2011). Synopses for massive data: samples, histogram, wavelets, sketches. *Foundations and Trends in Databases* **4** 1–294.
- [34] DAMLE, A. and SUN, Y. (2020). Uniform bounds for invariant subspace perturbations. *SIAM Journal on Matrix Analysis and Applications* **41** 1208–1236.
- [35] D’ASPROMONT, A., CUCURINGU, M. and TYAGI, H. (2021). Ranking and synchronization from pairwise measurements via SVD. *Journal of Machine Learning Research* **22** 1–63.
- [36] DAVENPORT, M. A. and ROMBERG, J. (2016). An overview of low-rank matrix recovery from incomplete observations. *IEEE Journal of Selected Topics in Signal Processing* **10** 608–622.
- [37] DAVIDSON, K. R. and SZAREK, S. J. (2001). *Handbook of the geometry of Banach spaces, Vol. I* Local operator theory, random matrices and Banach spaces, 317–366. North-Holland.
- [38] DAVIS, C. and KAHAN, W. M. (1970). The rotation of eigenvectors by a perturbation. III. *SIAM Journal on Numerical Analysis* **7** 1–46.
- [39] DE LATHAUWER, L., DE MOOR, B. and VANDEWALLE, J. (2000). On the best rank-1 and rank- (r_1, r_2, \dots, r_n) approximation of higher-order tensors. *SIAM Journal on Matrix Analysis and Applications* **21** 1324–1342.
- [40] DING, Y., KRISLOCK, N., QIAN, J. and WOLKOWICZ, H. (2010). Sensor network localization. Euclidean distance matrix completions, and graph realization. *Optimization and Engineering* **11** 45–66.
- [41] DRINEAS, P., KANNAN, R. and MAHONEY, M. W. (2006). Fast Monte Carlo algorithms for matrices II: Computing a low-rank approximation to a matrix. *SIAM Journal on Computing* **36** 158–183.
- [42] DU, X. and TANG, M. (2021). Hypothesis testing for equality of latent positions in random graphs. arXiv preprint at <https://arxiv.org/abs/2105.10838>.
- [43] EDELMAN, A. (1988). Eigenvalues and condition numbers of random matrices. *SIAM Journal on Matrix Analysis and Applications* **9** 543–560.
- [44] EDELMAN, A. (1991). The distribution and moments of the smallest eigenvalue of a random matrix of Wishart type. *Linear Algebra and its Applications* **159** 55–80.
- [45] ELДАР, Y. C. and KUTYNIOK, G. (2012). *Compressed sensing: theory and applications*. Cambridge University Press.
- [46] ELDRIDGE, J., BELKIN, M. and WANG, Y. (2018). Unperturbed: spectral analysis beyond Davis-Kahan. In *Proceedings of Algorithmic Learning Theory* 321–358.
- [47] ERDŐS, P. and RÉNYI, A. (1960). On the evolution of random graphs. *Publications of the Mathematical Institute of the Hungarian Academy of Sciences* **5** 17–60.
- [48] ERDŐS, L., KNOWLES, A., YAU, H. T. and YIN, J. (2013). Spectral statistics of Erdős-Rényi graphs I: Local semicircle law. *Annals of Probability* **41** 2279–2375.
- [49] ERICHSON, N. B. and DONOVAN, C. (2016). Randomized low-rank dynamic mode decomposition for motion detection. *Computer Vision and Image Understanding* **146** 40–50.
- [50] FAN, J., LIAO, Y. and MINCHEVA, M. (2013). Large covariance estimation by thresholding principal orthogonal complements. *Journal of the Royal Statistical Society. Series B* **75** 603–647.
- [51] FAN, J., WANG, W. and ZHONG, Y. (2018). An ℓ_∞ eigenvector perturbation bound and its application to robust covariance estimation. *Journal of Machine Learning Research* **18** 1–42.
- [52] FAN, J., FAN, Y., HAN, X. and LV, J. (2022+). SIMPLE: Statistical Inference on Membership Profiles in Large Networks. *Journal of the Royal Statistical Society, Series B*.
- [53] FORTUNATO, S. (2010). Community detection in graphs. *Physics Reports* **486** 75–174.
- [54] GLUNT, W., HAYDEN, T. L. and RAYDAN, M. (1993). Molecular conformations from distance matrices. *Journal of Computational Chemistry* **14** 114–120.
- [55] GOLUB, G. H. and VAN LOAN, C. F. (2013). *Matrix computations*, 3rd ed. JHU press.
- [56] GOPALAN, P. K. and BLEI, D. M. (2013). Efficient discovery of overlapping communities in massive networks. *Proceedings of the National Academy of Sciences* **110** 14534–14539.
- [57] HAJEK, B., WU, Y. and XU, J. (2016). Achieving exact cluster recovery threshold via semidefinite programming. *IEEE Transactions on Information Theory* **62** 2788–2797.

- [58] HALKO, N., MARTINSSON, P.-G. and TROPP, J. A. (2011). Finding structure with randomness: Probabilistic algorithms for constructing approximate matrix decompositions. *SIAM Review* **53** 217–288.
- [59] HALKO, N., MARTINSSON, P.-G., SHKOLNISKY, Y. and TYGERT, M. (2011). An algorithm for the principal component analysis of large data sets. *SIAM Journal on Scientific Computing* **33** 2580–2594.
- [60] HAN, X., YANG, Q. and FAN, Y. (2019). Universal rank inference via residual subsampling with applications to large networks. arXiv preprint at <https://arxiv.org/abs/1912.11583>.
- [61] HANSON, D. L. and WRIGHT, E. T. (1971). A bound on tail probabilities for quadratic forms in independent random variables. *Annals of Mathematical Statistics* **42** 1079–1083.
- [62] HIE, B., BRYSON, B. and BERGER, B. (2019). Efficient integration of heterogeneous single-cell transcriptomes using Scanorama. *Nature Biotechnology* **37** 685–691.
- [63] HOLLAND, P. W., LASKEY, K. B. and LEINHARDT, S. (1983). Stochastic blockmodels: First steps. *Social Networks* **5** 109–137.
- [64] HUANG, Q.-X. and GUIBAS, L. (2013). Consistent shape maps via semidefinite programming. In *Proceedings of the Eleventh Eurographics/ACMSIGGRAPH Symposium on Geometry Processing* 177–186.
- [65] JAVANMARD, A. and MONTANARI, A. (2013). Localization from Incomplete Noisy Distance Measurements. *Foundations of Computational Mathematics* **13** 297–345.
- [66] JOHNSTONE, I. M. (2001). On the distribution of the largest eigenvalue in principal components analysis. *The Annals of Statistics* **29** 295–327.
- [67] JOSEPH, A. and YU, B. (2016). Impact of regularization on spectral clustering. *The Annals of Statistics* **44** 1765–1791.
- [68] KANNAN, R. and VEMPALA, S. (2017). Randomized algorithms in numerical linear algebra. *Acta Numerica* **26** 95–135.
- [69] KARRER, B. and NEWMAN, M. E. (2011). Stochastic blockmodels and community structure in networks. *Physical Review E* **83** 016107.
- [70] KESHAVAN, R. H., MONTANARI, A. and OH, S. (2010). Matrix completion from noisy entries. *Journal of Machine Learning Research* **11** 2057–2078.
- [71] KOLTCHINSKII, V., LÖFFLER, M. and NICKL, R. (2020). Efficient estimation of linear functionals of principal components. *The Annals of Statistics* **48** 464–490.
- [72] KOLTCHINSKII, V. and LOUNICI, K. (2016). Asymptotics and concentration bounds for bilinear forms of spectral projectors of sample covariance. *Annales de l’Institut Henri Poincaré, Probabilités et Statistiques* **52** 1976–2013.
- [73] KOLTCHINSKII, V. and XIA, D. (2016). Perturbation of linear forms of singular vectors under Gaussian noise. In *High Dimensional Probability VII* 397–423. Springer.
- [74] KOO, J., TROSSET, M. W. and TANG, M. (2021). Popularity adjusted blockmodels are generalized random dot product graphs. arXiv preprint at <https://arxiv.org/abs/2109.04010>.
- [75] LEI, L. (2019). Unified $\ell_{2 \rightarrow \infty}$ eigenspace perturbation theory for symmetric random matrices. arXiv preprint at <https://arxiv.org/abs/1909.04798>.
- [76] LEI, J. and RINALDO, A. (2015). Consistency of spectral clustering in stochastic block models. *The Annals of Statistics* **43** 215–237.
- [77] LEVIN, K. D., ROOSTA, F., TANG, M., MAHONEY, M. W. and PRIEBE, C. E. (2021). Limit theorems for out-of-sample extensions of the adjacency and Laplacian spectral embeddings. *Journal of Machine Learning Research* **22**.
- [78] LI, B., GOULD, J., YANG, Y., SARKIZOVA, S., TABAKA, M., ASHENBERG, O., ROSEN, Y., SLYPER, M., KOWALCZYK, M. S., VILLANI, A.-C. et al. (2020a). Cumulus provides cloud-based data analysis for large-scale single-cell and single-nucleus RNA-seq. *Nature Methods* **17** 793–798.
- [79] LI, G. K., TANG, M., CHARON, N. and PRIEBE, C. E. (2020b). Central limit theorems for classical multidimensional scaling. *Electronic Journal of Statistics* **14** 2362–2394.
- [80] LIDSKII, V. B. (1950). On the proper values of a sum and product of symmetric matrices. *Doklady Akademii Nauk SSSR* **75** 769–772.
- [81] LOPES, M., ERICHSON, N. B. and MAHONEY, M. (2020). Error estimation for sketched SVD via the bootstrap. In *Proceedings of the 37th International Conference on Machine Learning* 6382–6392.
- [82] LYZINSKI, V., SUSSMAN, D. L., TANG, M., ATHREYA, A. and PRIEBE, C. E. (2014). Perfect clustering for stochastic blockmodel graphs via adjacency spectral embedding. *Electronic Journal of Statistics* **8** 2905–2922.
- [83] MAHONEY, M. W. (2011). Randomized algorithms for matrices and data. *Foundations and Trends in Machine Learning* **3** 123–224.
- [84] MAO, X., SARKAR, P. and CHAKRABARTI, D. (2021). Estimating mixed memberships with sharp eigenvector deviations. *Journal of the American Statistical Association* **116** 1928–1940.

- [85] MARTINSSON, P. G., SZLAM, A. and TYGERT, M. (2010). Normalized power iterations for the computation of SVD. available online at <http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.683.7657>.
- [86] MARTINSSON, P. G. and TROPP, J. A. (2020). Randomized numerical linear algebra: Foundations and algorithms. *Acta Numerica* **29** 403–572.
- [87] MEGIDDO, N. and SUPOWIT, K. J. (1984). On the complexity of some common geometric location problems. *SIAM Journal on Computing* **13** 182–196.
- [88] MONTANARI, A. and VENKATARAMANAN, R. (2021). Estimation of low-rank matrices via approximate message passing. *The Annals of Statistics* **49** 321–345.
- [89] MUSCO, C. and MUSCO, C. (2015). Randomized block Krylov methods for stronger and faster approximate singular value decomposition. In *Advances in Neural Information Processing Systems* **28** 1396–1404.
- [90] NETRAPALLI, P., NIRANJAN, U., SANGHAVI, S., ANANDKUMAR, A. and JAIN, P. (2014). Non-convex Robust PCA. In *Advances in Neural Information Processing Systems* **27** 1107–1115.
- [91] NEWMAN, M. E. (2006). Finding community structure in networks using the eigenvectors of matrices. *Physical Review E* **74** 036104.
- [92] NEWMAN, M. and GIRVAN, M. (2004). Finding and evaluating community structure in networks. *Physical Review E* **69** 1–15.
- [93] OLIVEIRA, R. I. (2010). Sum of random Hermitean matrices and an inequality by Rudelson. *Electronic Communications in Probability* **15** 203–212.
- [94] ONATSKI, A. (2010). Determining the number of factors from empirical distribution in large factor models. *Review of Economics and Statistics* **92** 1004–1016.
- [95] ORÚS, R. (2019). Tensor networks for complex quantum systems. *Nature Reviews Physics* **1** 538–550.
- [96] PONS, P. and LATAPY, M. (2005). Computing communities in large networks using random walks. In *Proceedings of the 20th International Conference on Computer and Information Sciences* 284–293.
- [97] QIU, J., DONG, Y., MA, H., LI, J., WANG, C., WANG, K. and TANG, J. (2019). NetSMF: Large-scale network embedding as sparse matrix factorization. In *The World Wide Web Conference* 1509–1520.
- [98] RENNIE, J. D. and SREBRO, N. (2005). Fast maximum margin matrix factorization for collaborative prediction. In *Proceedings of the 22nd International Conference on Machine learning* 713–719.
- [99] ROHE, K., CHATTERJEE, S. and YU, B. (2011). Spectral clustering and the high-dimensional stochastic blockmodel. *The Annals of Statistics* **39** 1878–1915.
- [100] ROHE, K. and ZENG, M. (2020). Vintage factor analysis with Varimax performs statistical inference. arXiv preprint at <https://arxiv.org/abs/2004.05387>.
- [101] ROKHLIN, V., SZLAM, A. and TYGERT, M. (2010). A randomized algorithm for principal component analysis. *SIAM Journal on Matrix Analysis and Applications* **31** 1100–1124.
- [102] ROSVALL, M. and BERGSTROM, C. T. (2008). Maps of random walks on complex networks reveal community structure. *Proceedings of the National Academy of Sciences of the United States of America* **105**.
- [103] RUBIN-DELANCHY, P., CAPE, J., TANG, M. and PRIEBE, C. E. (2022+). A statistical interpretation of spectral embedding: the generalised random dot product graph. *Journal of the Royal Statistical Society, Series B*.
- [104] RUDELSON, M. (1999). Random vectors in the isotropic position. *Journal of Functional Analysis* **164** 60–772.
- [105] RUDELSON, M. and VERSHYNIN, R. (2010). Non-asymptotic theory of random matrices: extreme singular values. In *Proceedings of the International Congress of Mathematicians* **1** 1576–1602.
- [106] SAIBABA, A. K. (2019). Randomized Subspace Iterations: Analysis of Canonical Angles and Unitarily Invariant Norms. *SIAM Journal on Matrix Analysis and Applications* **40** 23–38.
- [107] SCHMIDT, R. (1986). Multiple emitter location and signal parameter estimation. *IEEE Transactions on Antennas and Propagation* **34** 276–280.
- [108] SENGUPTA, S. and CHEN, Y. (2018). A block model for node popularity in networks with community structure. *Journal of the Royal Statistical Society, Series B*. **80** 365–386.
- [109] SINGER, A. (2011). Angular synchronization by eigenvectors and semidefinite programming. *Applied and Computational Harmonic Analysis* **30** 20–36.
- [110] SNIJDERS, T. A. B. and NOWICKI, K. (1997). Estimation and Prediction for Stochastic Blockmodels for Graphs with Latent Block Structure. *Journal of Classification* **14** 75–100.
- [111] STEWART, G. W. (1977). On the perturbation of pseudo-inverses, projections and linear least squares problems. *SIAM review* **19** 634–662.
- [112] SU, L., WANG, W. and ZHANG, Y. (2019). Strong consistency of spectral clustering for stochastic block models. *IEEE Transactions on Information Theory* **66** 324–338.

- [113] SUSSMAN, D. L., TANG, M., FISHKIND, D. E. and PRIEBE, C. E. (2012). A consistent adjacency spectral embedding for stochastic blockmodel graphs. *Journal of the American Statistical Association* **107** 1119–1128.
- [114] TENENBAUM, J. B., SILVA, V. D. and LANGFORD, J. C. (2000). A Global Geometric Framework for Nonlinear Dimensionality Reduction. *Science* **290** 2319–2323.
- [115] TROPP, J. A. (2012). User-friendly tail bounds for sums of random matrices. *Foundations of Computational Mathematics* **12** 389–434.
- [116] TROPP, J. A., YURTSEVER, A., UDELL, M. and CEVHER, V. (2017). Practical sketching algorithms for low-rank matrix approximation. *SIAM Journal on Matrix Analysis and Applications* **38** 1454–1485.
- [117] TROPP, J. A., YURTSEVER, A., UDELL, M. and CEVHER, V. (2019). Streaming low-rank matrix approximation with an application to scientific simulation. *SIAM Journal on Scientific Computing* **41** A2430–A2463.
- [118] TSILIGKARIDIS, T. and HERO, A. O. (2013). Covariance estimation in high dimensions via Kronecker product expansions. *IEEE Transactions on Signal Processing* **61** 5347–5360.
- [119] VERSHYNIN, R. (2012). In *Compressed Sensing: Theory and Applications* Introduction to the non-asymptotic analysis of random matrices, 210–268. Cambridge University Press.
- [120] VERSHYNIN, R. (2018). *High-dimensional probability: An introduction with applications in data science*. Cambridge University Press.
- [121] VON LUXBURG, U. (2007). A tutorial on spectral clustering. *Statistics and Computing* **17** 395–416.
- [122] WEDIN, P.-Å. (1972). Perturbation bounds in connection with singular value decomposition. *BIT Numerical Mathematics* **12** 99–111.
- [123] WEYL, H. (1912). Das asymptotische Verteilungsgesetz der Eigenwerte linearer partieller Differentialgleichungen (mit einer Anwendung auf die Theorie der Hohlraumstrahlung). *Mathematische Annalen* **71** 441–479.
- [124] WOODRUFF, D. P. (2014). Sketching as a tool for numerical linear algebra. *Foundations and Trends in Theoretical Computer Science* **10** 1–157.
- [125] YAN, Y., CHEN, Y. and FAN, J. (2021). Inference for heteroskedastic PCA with missing data. *arXiv preprint at <https://arxiv.org/abs/2107.12365>*.
- [126] YANG, F., LIU, S., DOBRIBAN, E. and WOODRUFF, D. P. (2021a). How to reduce dimension with PCA and random projections? *IEEE Transactions on Information Theory* **67** 8154–8189.
- [127] YANG, C., PRIEBE, C. E., PARK, Y. and MARCHETTE, D. J. (2021b). Simultaneous dimensionality and complexity model selection for spectral graph clustering. *Journal of Computational and Graphical Statistics* **30** 422–441.
- [128] YOUNG, S. J. and SCHEINERMAN, E. R. (2007). Random dot product graph models for social networks. In *Proceedings of the 5th International Conference on Algorithms and Models for the Web-graph* 138–149.
- [129] YU, Y., WANG, T. and SAMWORTH, R. J. (2015). A useful variant of the Davis-Kahan theorem for statisticians. *Biometrika* **102** 315–323.
- [130] ZHANG, H., GUO, X. and CHANGE, X. (2020). Randomized spectral clustering in large-scale stochastic blockmodels. *arXiv preprint at <https://arxiv.org/abs/2002.00839>*.
- [131] ZHANG, A. and XIA, D. (2018). Tensor SVD: Statistical and computational limits. *IEEE Transactions on Information Theory* **64** 7311–7338.
- [132] ZHU, M. and GHODSI, A. (2006). Automatic dimensionality selection from the scree plot via the use of profile likelihood. *Computational Statistics & Data Analysis* **51** 918–930.

SUPPLEMENTARY FILE FOR “PERTURBATION ANALYSIS OF RANDOMIZED SVD AND ITS APPLICATIONS TO HIGH-DIMENSIONAL STATISTICS”

The Supplementary File can be divided to four parts. Section S.1 presents several numerical experiments to verify our main theoretical investigations. Section S.2 contains an extension of our main results to the case of asymmetric $\hat{\mathbf{M}}$ and \mathbf{M} . Section S.3 states more general counterparts of our main theorems when \mathbf{G} is assumed to be non-random. Sections S.4–S.7 include all the technical lemmas and proofs.

S.1. Numerical experiments. We perform simulations to illustrate our theoretical results. In particular, Section S.1.1 present results for RSVD-based subspace estimation and community detection in SBM graphs (see Section 4.3), Section S.1.2 presents results for RSVD-based PCA with missing data (see Section 5.2) and Section S.1.3 present results for matrix completion with noisy entries.

S.1.1. *RSVD-based spectral clustering for SBM.* We consider two-blocks SBM graphs with equal sized blocks and block probabilities matrix

$$\mathbf{B}_0 = \rho_n \begin{pmatrix} 0.8 & 0.3 \\ 0.3 & 0.8 \end{pmatrix}.$$

Recall that ρ_n is the sparsity scaling parameter. We consider three different values for ρ_n , namely $\rho_n = 1$ (dense setting), $\rho_n = 3n^{-1/3}$ (semi-sparse setting I), $\rho_n = 4n^{-1/2}$ (semi-sparse setting II). As the edge probability matrix is of rank 2, we set $k = 2$. Let \mathbf{U} and $\hat{\mathbf{U}} \in \mathbb{R}^{n \times 2}$ be the $n \times k$ matrix whose columns are the leading k singular vectors of \mathbf{P} and \mathbf{A} , respectively. Let $\hat{\mathbf{U}}_g$ be the approximation of $\hat{\mathbf{U}}$ computed using Algorithm 2 for some choices of a_n, g and \tilde{k} that are specified subsequently.

S.1.1.1. *Phase transition.* We first verify the convergence rate for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ as we vary g and ρ_n ; see Section 3.1 and Section 3.2. For simplicity we only consider $\rho_n = 1$ and $\rho_n = 3n^{-1/3}$, and we ignore any potential logarithmic factors in the convergence rate for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ as n increases, i.e., we ignore the polylog terms in Eq. (4.2).

We apply *rs*-RSVD with two settings of a_n, \tilde{k} , namely, (i) $a_n = \lceil \log n \rceil$ and $\tilde{k} = 12$; (ii) $a_n = 1$ and $\tilde{k} = 5 \lceil \log n \rceil$. For each combination of (ρ_n, a_n, \tilde{k}) , we numerically estimate the polynomial convergence rates of $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ as follows. For each $n = 2000, \dots, 7000$ and specific ρ_n , we generate one realization of \mathbf{A} with equal block sized and block probabilities \mathbf{B}_0 and then compute $\hat{\mathbf{U}}_g$ via Algorithm 2 for each $g \in \{1, 2, \dots, 5\}$. We then evaluate $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ and compare it against the theoretical error rate given in Section 4.1 by running a simple linear regression between the (negative logarithm) of the empirical error rate as the response variable against $\log n$ as the predictor variable. The estimated coefficient $\hat{\beta}$ are then recorded. We repeat the above simulations for 500 Monte Carlo (MC) iterations. The box-plots of the estimated convergence rate $\hat{\beta}$ over these 500 iterations are presented in Figure S.1. For comparison we had also included the (estimated) convergence rate for $d_2(\hat{\mathbf{U}}, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$ (these are labeled as “true SVD”).

The results in Figure S.1 match the theoretical results presented in Theorem 4.1 and Eq.(4.2) exactly. In particular we see no convergence when $g = 1$ and $\rho_n = 1$, slow or no convergence when $g \leq 2$ and $\rho_n = 3n^{-1/3}$, and asymptotically optimal convergence when $g \geq 2$ and $\rho_n = 1$ or $g \geq 3$ and $\rho_n = 3n^{-1/3}$.




FIG S.1. The left panel reports box-plots of simulated $\hat{\beta}$ when *rs-RSVD* is implemented with $a_n = \lceil \log n \rceil$ and $\tilde{k} = 12$. The right panel reports box-plots of simulated $\hat{\beta}$ for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ (x -axis: $g = 1, \dots, 5$) or $d_2(\hat{\mathbf{U}}, \mathbf{U})$ (x -axis: **True SVD**) and the bottom plot reports the β for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ (x -axis: $g = 1, \dots, 5$) or $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}, \mathbf{U})$ (x -axis: **True SVD**). Box-plots for dense ($\rho_n = 1$) and sparse ($\rho_n = 3n^{-1/3}$) graphs are in yellow and blue, respectively.

TABLE S.1

Proportion of times that *RSVD*-based spectral clustering ($g = 1, 2, 3$) or the original spectral clustering (*OSC*) exactly recover the memberships of all nodes; here $a_n = \lceil \log n \rceil$ and $\tilde{k} = 12$. Standard errors are reported in parentheses.

Sparsity	n	$g = 1$	$g = 2$	$g = 3$	OSC
$\rho_n \asymp 1$	1000	0.098 (0.013)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	2000	0.050 (0.010)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	3000	0.034 (0.008)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	4000	0.018 (0.006)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	5000	0.006 (0.003)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
$\rho_n \asymp n^{-1/3}$	1000	0.000 (0.000)	0.938 (0.012)	1.000 (0.000)	1.000 (0.000)
	2000	0.000 (0.000)	0.980 (0.006)	1.000 (0.000)	1.000 (0.000)
	3000	0.000 (0.000)	0.980 (0.006)	1.000 (0.000)	1.000 (0.000)
	4000	0.000 (0.000)	0.994 (0.003)	1.000 (0.000)	1.000 (0.000)
	5000	0.000 (0.000)	0.998 (0.002)	1.000 (0.000)	1.000 (0.000)
$\rho_n \asymp n^{-1/2}$	1000	0.000 (0.000)	0.000 (0.000)	0.384 (0.022)	0.906 (0.013)
	2000	0.000 (0.000)	0.000 (0.000)	0.688 (0.021)	0.986 (0.005)
	3000	0.000 (0.000)	0.000 (0.000)	0.864 (0.015)	1.000 (0.000)
	4000	0.000 (0.000)	0.000 (0.000)	0.956 (0.009)	1.000 (0.000)
	5000	0.000 (0.000)	0.000 (0.000)	0.976 (0.007)	1.000 (0.000)

S.1.1.2. *Exact recovery.* We used the same simulation setting as that described in Section S.1.1.1 with $n \in \{1000, \dots, 5000\}$ and $\rho_n \in \{1, 3n^{-1/3}, 4n^{-1/2}\}$. For each combination of (ρ_n, a_n, \tilde{k}) and for each n , we generate an adjacency matrix \mathbf{A} with equal sized blocks and block probabilities \mathbf{B}_0 . We then perform *RSVD*-based spectral clustering as described in Algorithm 2 with $g \leq 3$, where we use the `Gmedian` library in **R** to perform fast (approximate) K -medians clustering on the rows of $\hat{\mathbf{U}}_g$. For comparison, we also use `Gmedian` to perform K -medians clustering on the rows of $\hat{\mathbf{U}}$. We repeat the above steps for 500 Monte Carlo replicates. The proportion of times (among these 500 replicates) in which the K -medians clustering of either $\hat{\mathbf{U}}_g$ or $\hat{\mathbf{U}}$ correctly recover the community memberships for all nodes are reported in Table S.1 and Table S.2.

From Tables S.1 and S.2 we see that Algorithm 2 exactly recover the underlying community assignment with probabilities converging to 1 as n increases, provided that $g \geq 2$

TABLE S.2

Proportion of times that RSVD-based spectral clustering ($g = 1, 2, 3$) or the original spectral clustering (OSC) exactly recover the memberships of all nodes, among 500 MC rounds; here $a_n = 1$ and $\tilde{k} = 5\lceil \log n \rceil$. Standard errors are reported in parentheses.

Sparsity	n	$g = 1$	$g = 2$	$g = 3$	OSC
$\rho_n \asymp 1$	1000	0.536 (0.022)	0.384 (0.000)	1.000 (0.000)	1.000 (0.000)
	2000	0.616 (0.022)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	3000	0.674 (0.021)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	4000	0.600 (0.022)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	5000	0.626 (0.022)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
$\rho_n \asymp n^{-1/3}$	1000	0.000 (0.000)	0.982 (0.012)	1.000 (0.000)	1.000 (0.000)
	2000	0.000 (0.000)	0.996 (0.006)	1.000 (0.000)	1.000 (0.000)
	3000	0.000 (0.000)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	4000	0.000 (0.000)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
	5000	0.000 (0.000)	1.000 (0.000)	1.000 (0.000)	1.000 (0.000)
$\rho_n \asymp n^{-1/2}$	1000	0.000 (0.000)	0.000 (0.000)	0.604 (0.022)	0.914 (0.013)
	2000	0.000 (0.000)	0.000 (0.000)	0.892 (0.014)	1.000 (0.000)
	3000	0.000 (0.000)	0.000 (0.000)	0.974 (0.007)	1.000 (0.000)
	4000	0.000 (0.000)	0.000 (0.000)	0.996 (0.003)	1.000 (0.000)
	5000	0.000 (0.000)	0.000 (0.000)	0.998 (0.002)	1.000 (0.000)

when $\rho_n \in \{1, 3n^{-1/3}\}$ and $g \geq 3$ when $\rho_n = 4n^{-1/2}$. These empirical results are consistent with the theoretical results presented in Theorem 4.2, i.e., exact recover is guaranteed if and only if $g > \beta^{-1}$. Note that $\beta^{-1} = 1$ for $\rho_n = 1$, $\beta^{-1} = \frac{3}{2}$ when $\rho_n \asymp n^{-1/3}$ and $\beta^{-1} = 2$ when $\rho_n \asymp n^{-1/2}$. Finally we note that if $g = 1$ and $\rho_n = 1$ then RSVD-based spectral clustering occasionally recovers the community membership for all nodes. This is due to the fact that although $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_1, \mathbf{U}) = \Omega(n^{1/2})$ with high probability, it does not prevent $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_1, \mathbf{U}) \leq \zeta_*$ where ζ_* is the minimum ℓ_2 distance between any two nodes i and j belonging to different communities; exact recovery is certainly expected if $d_2(\hat{\mathbf{U}}_1, \mathbf{U}) \leq \zeta_*$.

S.1.1.3. Limiting distribution. We now illustrate the normal approximations for the row-wise fluctuations of $\hat{\mathbf{U}}_g$, as implied by Theorem 3.4. For brevity we fixed $\tilde{k} = 2$ and $a_n = \lceil \log n \rceil$. Similar results are available for $\tilde{k} \asymp \log n$ and $a_n = 1$. We generate a single realization of \mathbf{A} on $n = 10000$ vertices and $\rho_n = 4n^{-1/2}$, with equal sized blocks and block probabilities \mathbf{B}_0 . Scatter plots of the rows of $\hat{\mathbf{U}}_g$ for $g \in \{1, 2, \dots, 5\}$ as well as the rows of $\hat{\mathbf{U}}$ are presented in Figure S.2. From Figure S.2 we see that $g \geq 3$ is sufficient for the rows of $\hat{\mathbf{U}}_g$ to be clustered into two clusters but $g \geq 4$ is necessary for the empirical 95% confidence ellipses of $\hat{\mathbf{U}}_g$ to match the theoretical 95% confidence ellipses given in Theorem 3.4; the scatter plots for $g \geq 4$ are also virtually indistinguishable from that for $\hat{\mathbf{U}}$. In other words, the more stringent threshold $g \geq 2 + \beta^{-1}$ in Theorem 3.4 is both necessary and sufficient for the normal approximations of $\hat{\mathbf{U}}_g$; note that for the current setting we have $\rho_n \asymp n^{-1/2}$ and hence $\beta^{-1} = 2$.

Additional simulation results with respect to $\rho_n = 1$ or $n = 1000$, are further illustrated in Figures S.3–S.5. Since our theoretical results are valid up to some rotation, we shall orthogonally rotate the plots occasionally for the sake of better illustrations. Similar row-wise fluctuations and the corresponding phase transition phenomenon with respect to g , can also be seen in these additional simulations.

S.1.2. PCA with missing data. For this simulation study we used the same data generation mechanism as that described in section 7 of [18] but with larger values of d . More specifically we first sample a $d \times n$ matrix \mathbf{X}^* whose columns are iid multivariate normal




FIG S.2. Row-wise fluctuations for $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{U}}$ for two-blocks SBM with $n = 10000$ and $\rho_n \asymp n^{-1/2}$. (From top left to top right) scatter plots of the rows of $\hat{\mathbf{U}}_g$ when $g \leq 3$; (From bottom left to bottom right) scatter plots of the rows of $\hat{\mathbf{U}}_g$ when $g \in \{4, 5\}$, and the rows of $\hat{\mathbf{U}}$. The points are colored according to the true community assignments. The red dashed curves are the 95% empirical confidence ellipses, while the solid black curves are the 95% theoretical confidence ellipses given by Theorem 3.4




FIG S.3. Second-order simulation for two-block SBM with $n = 1000$ and $\rho_n \asymp n^{-1/2}$. (From top left to top right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 1, 2, 3$; (From bottom left to bottom right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 4, 5$ and $\hat{\mathbf{U}}$. The point color reflects the block membership of the corresponding row vectors. The red dashed curves are the 95% empirical confidence ellipses, while the solid black curves are the 95% theoretical confidence ellipses according to Theorem 3.4




FIG S.4. Second-order simulation for two-block SBM with $n = 1000$ and $\rho_n \asymp 1$. (From top left to top right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 1, 2, 3$; (From bottom left to bottom right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 4, 5$ and $\hat{\mathbf{U}}$. The point color reflects the block membership of the corresponding row vectors. The red dashed curves are the 95% empirical confidence ellipses, while the solid black curves are the 95% empirical confidence theoretical according to Theorem 3.4




FIG S.5. Second-order simulation for two-block SBM with $n = 10000$ and $\rho_n \asymp 1$. (From top left to top right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 1, 2, 3$; (From bottom left to bottom right) the scatter plots of row-wise vectors in $\hat{\mathbf{U}}_g$ with $g = 4, 5$ and $\hat{\mathbf{U}}$. The point color reflects the block membership of the corresponding row vectors. The red dashed curves are the 95% empirical confidence ellipses, while the solid black curves are the 95% theoretical confidence ellipses according to Theorem 3.4




FIG S.6. (Top Left) ℓ_2 recovery errors vs. p (here $d = 3000, n = 1000, \sigma = 1$); (Top Right) ℓ_2 recovery errors vs. d (here $n = 1000, p = 0.02, \sigma = 1$); (Bottom Left) ℓ_2 recovery errors vs. n (here $d = 3000, p = 0.02, \sigma = 1$); (Bottom Right) ℓ_2 recovery errors vs. σ (here $d = 3000, n = 1000, p = 0.02$).

random vectors with mean $\mathbf{0}$ and covariance matrix $\mathbf{U}^* \mathbf{U}^{*\top}$; here \mathbf{U}^* is a $d \times 4$ matrix whose entries are iid standard normals. We then generate $\mathcal{P}_\Omega(\mathbf{X}) = \Omega \circ (\mathbf{X}^* + \mathbf{E})$ where the entries of \mathbf{E} are iid $\mathcal{N}(0, \sigma^2)$, the entries of Ω are iid Bernoulli(p) and \circ denote the Hadamard product. The matrix $\mathcal{P}_\Omega(\mathbf{X})$ represent a noisily observed version of \mathbf{X}^* (with missing entries).

Given $\mathcal{P}_\Omega(\mathbf{X})$ we compute $\hat{\mathbf{U}}_g$ using Algorithm 4 with $k = 4, \bar{k} = 14, a_d = \lceil \log d \rceil$ and $g \in \{1, 2, \dots, 5\}$. We then record $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$; here \mathbf{U} denote the $d \times 4$ matrix whose columns are the leading eigenvectors of $\mathbf{U}^* \mathbf{U}^{*\top}$. For comparison we also record $d_2(\hat{\mathbf{U}}, \mathbf{U})$ where $\hat{\mathbf{U}}$ is the $d \times 4$ matrix of leading eigenvectors of \mathbf{Q} and \mathbf{Q} is as defined in Eq. (5.11).

Figure S.6 reports sample means for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_2(\hat{\mathbf{U}}, \mathbf{U})$ as we vary the sampling probability p , data dimension d , sample size n , and noise level σ ; these sample means are computed based on 500 Monte Carlo replicates where we resampled the matrices \mathbf{U}^*, \mathbf{E} and Ω in each replicate. Figure S.6 shows that Algorithm 4 yields nearly-optimal performance (as compared to $\hat{\mathbf{U}}$) when $g \geq 3$ for most settings of p, d, σ and n . It is only when d is large with respect to n that $\hat{\mathbf{U}}_3$ have worse performance compared to $\hat{\mathbf{U}}_4, \hat{\mathbf{U}}_5$ and $\hat{\mathbf{U}}$. Finally, $\hat{\mathbf{U}}_1$ is always sub-optimal. These observations are consistent with the theoretical results presented in Section 5.2.

S.1.3. Distance matrix completion. For this section we apply Algorithm 3 to recover the missing entries of a partially observed Euclidean distance matrix. In particular we use the `world_cities` dataset containing the locations of the 4428 most populous cities around the world; this dataset is part of the `mdsr` library in **R**. We first construct the 4428×4428 matrix $\mathbf{D} = [D_{ij}]$ whose elements are

$$D_{ij} = (\text{Lon}_i - \text{Lon}_j)^2 + (\text{Lat}_i - \text{Lat}_j)^2.$$




FIG S.7. Matrix plots for the true \mathbf{D} (Left), partially observed $\mathbf{D}_{0.8}$ (Middle), and partially observed $\mathbf{D}_{0.4}$ (Right).




FIG S.8. From left to right: matrix plots of RSVD-based estimates $\hat{\mathbf{D}}_{0.8}^{(1)}$, $\hat{\mathbf{D}}_{0.8}^{(2)}$, $\hat{\mathbf{D}}_{0.8}^{(5)}$, and exact SVD estimate $\hat{\mathbf{D}}_{0.8}$.

Here Lon_i and Lat_i represent the longitude and latitude of the i th city, respectively. We then sample a matrix $\mathbf{D}_{0.8}$ (resp. $\mathbf{D}_{0.4}$) by keeping roughly 80% (resp. 40%) of the entries in \mathbf{D} , i.e., $\mathbf{D}_{0.8} = \mathbf{\Omega} \circ \mathbf{D}$ where $\mathbf{\Omega}$ is a symmetric matrix whose upper triangular entries are iid Bernoulli(0.8). We now recover \mathbf{D} from $\mathbf{D}_{0.8}$ (resp. $\mathbf{D}_{0.4}$) using Algorithm 3. As the entries of \mathbf{D} are Euclidean distances between points in \mathbb{R}^2 , we have $\text{rk}(\mathbf{D}) \leq 4$. We therefore choose $k = 4$, $a_n = \tilde{k} = 15$ and $g \in \{1, 2, 5\}$, and let $\hat{\mathbf{D}}_{0.8}^{(g)}$ (resp. $\hat{\mathbf{D}}_{0.4}^{(g)}$) be the resulting estimate of \mathbf{D} . For comparison we also consider the spectral estimate $\hat{\mathbf{D}}_{0.8}$ (resp. $\hat{\mathbf{D}}_{0.4}$) obtained by truncating the exact SVD of $\mathbf{D}_{0.8}$ (resp. $\mathbf{D}_{0.4}$); see [70] for more details.

A plot of the true distance matrix \mathbf{D} and one random realization of the partially observed $\mathbf{D}_{0.8}$ and $\mathbf{D}_{0.4}$ are presented in Figure S.7. The corresponding RSVD-based estimates and exact SVD based estimates of \mathbf{D} are then shown in Figure S.8–S.9. Figure S.8 shows that $\hat{\mathbf{D}}_{0.8}^{(2)}$ and $\hat{\mathbf{D}}_{0.8}^{(5)}$ both have comparable accuracy to $\hat{\mathbf{D}}_{0.8}$ while Figure S.9 shows that $\hat{\mathbf{D}}_{0.4}^{(2)}$ has much worse accuracy compared to $\hat{\mathbf{D}}_{0.4}^{(5)}$ and $\hat{\mathbf{D}}_{0.4}$.

We also record the entrywise relative errors between the RSVD-based estimates $\hat{\mathbf{D}}_p^{(g)}$ (resp. the exact SVD-based estimate $\hat{\mathbf{D}}_p$) against that of \mathbf{D} . A summary of the quantile levels for these relative errors are presented in Tables S.3. The α th-quantile of the entrywise relative errors between an estimate \mathbf{Z} and the true distance \mathbf{D} is defined as the α th quantile of $\{||[\mathbf{Z} - \mathbf{D}]_{ij} / [\mathbf{D}]_{ij}| : (i, j) \in [4428] \times [4428]\}$; for example the median relative error for $\hat{\mathbf{D}}_{0.8}^{(5)}$ and $\hat{\mathbf{D}}_{0.4}^{(5)}$ are ≈ 0.029 and ≈ 0.086 , respectively. Note that the numbers in Table S.3 are averaged over 200 independent random samples of either $\mathbf{D}_{0.8}$ or $\mathbf{D}_{0.4}$. From Table S.3 we see that the relative error decreases as g increases with p fixed; indeed, the relative error of $\hat{\mathbf{D}}_{0.8}^{(5)}$ is exactly the same as that for $\hat{\mathbf{D}}_{0.8}$. Table S.3 also indicates that as p decrease we need to increase g to achieve a recovery rate close to that of $\hat{\mathbf{D}}$. These observations are consistent with the theoretical results in Theorem 5.1.




FIG S.9. From left to right: matrix plots of RSVD-based estimates $\hat{\mathbf{D}}_{0.4}^{(1)}$, $\hat{\mathbf{D}}_{0.4}^{(2)}$, $\hat{\mathbf{D}}_{0.4}^{(5)}$, and exact SVD estimate $\hat{\mathbf{D}}_{0.4}$

TABLE S.3

α th-quantiles for the relative entrywise errors of the RSVD-based estimates $\hat{\mathbf{D}}_p^{(g)}$ and exact SVD estimate $\hat{\mathbf{D}}_p$; standard errors for these quantiles are given in the parenthesis.

α	0.05	0.15	0.35	0.5	0.65	0.85	0.95
$\hat{\mathbf{D}}_{0.8}^{(1)}$	0.0169 (1.3×10^{-4})	0.0515 (3.9×10^{-4})	0.1315 (9.2×10^{-4})	0.2203 (1.4×10^{-3})	0.4232 (2.7×10^{-3})	2.4408 (1.9×10^{-2})	15.2389 (1.1×10^{-1})
$\hat{\mathbf{D}}_{0.8}^{(2)}$	0.0035 (1.4×10^{-5})	0.0107 (4.2×10^{-5})	0.0273 (1.2×10^{-4})	0.0453 (2.3×10^{-4})	0.0789 (4.6×10^{-4})	0.2798 (1.9×10^{-3})	1.5766 (1.3×10^{-2})
$\hat{\mathbf{D}}_{0.8}^{(5)}$	0.0024 (1.4×10^{-7})	0.0073 (2.8×10^{-7})	0.0182 (6.6×10^{-7})	0.0289 (1.1×10^{-6})	0.0464 (2.3×10^{-6})	0.1531 (1.0×10^{-5})	0.7823 (6.7×10^{-5})
$\hat{\mathbf{D}}_{0.8}$	0.0024	0.0073	0.0182	0.0289	0.0464	0.1530	0.7816
$\hat{\mathbf{D}}_{0.4}^{(1)}$	0.0371 (3.6×10^{-4})	0.1112 (1.0×10^{-3})	0.2644 (2.1×10^{-3})	0.4049 (2.7×10^{-3})	0.6715 (3.0×10^{-3})	3.8785 (2.0×10^{-2})	24.4391 (1.4×10^{-1})
$\hat{\mathbf{D}}_{0.4}^{(2)}$	0.0106 (3.7×10^{-5})	0.0323 (1.2×10^{-4})	0.0849 (3.5×10^{-4})	0.1472 (7.2×10^{-4})	0.2717 (1.5×10^{-3})	1.3069 (1.0×10^{-2})	8.9068 (7.0×10^{-2})
$\hat{\mathbf{D}}_{0.4}^{(5)}$	0.0070 (9.6×10^{-6})	0.0211 (3.0×10^{-5})	0.0531 (8.2×10^{-5})	0.0860 (1.5×10^{-4})	0.1454 (3.4×10^{-4})	0.5072 (1.2×10^{-3})	3.0309 (9.8×10^{-3})
$\hat{\mathbf{D}}_{0.4}$	0.0064	0.0194	0.0484	0.0774	0.1269	0.4439	2.4614

Algorithm S.1: rs -RSVD for asymmetric $\hat{\mathbf{M}}$

Input: $\hat{\mathbf{M}} \in \mathbb{R}^{n_1 \times n_2}$, rank $k \geq 1$ and sketching dimension $\tilde{k} \geq k$. Integers $a_n, g \geq 1$.

1. Generate a_n iid standard Gaussian sketching matrices $\mathbf{G}_1, \dots, \mathbf{G}_{a_n} \in \mathbb{R}^{n_2 \times \tilde{k}}$;
2. For $a \in [a_n]$, form $(\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}_a$ by computing $\hat{\mathbf{M}}\mathbf{G}_a, \hat{\mathbf{M}}^\top(\hat{\mathbf{M}}\mathbf{G}_a), \dots, (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}_a$;
3. For $a \in [a_n]$, compute the rank- k SVD of $(\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}_a$ to obtain the k th largest singular values $\sigma_k\{(\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}_a\}$ and the $n_1 \times k$ matrix of left singular vectors $\hat{\mathbf{U}}_g^{(a)}$;
4. Choose $\tilde{a} = \operatorname{argmax}_{a \in [a_n]} \sigma_k\{(\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}_a\}$ and $\mathbf{G} = \mathbf{G}_{\tilde{a}}$. Let $\hat{\mathbf{U}}_g = \hat{\mathbf{U}}_g^{(\tilde{a})}$;
5. Let $\tilde{\sigma}_1 \geq \dots \geq \tilde{\sigma}_k$ be the singular values of $\hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$.

Output: Estimated singular vectors $\hat{\mathbf{U}}_g$, singular values $\{\tilde{\sigma}_1, \dots, \tilde{\sigma}_k\}$.

S.2. Extension to asymmetric matrices. We now extend Theorems 3.1–3.3 to the setting where $\hat{\mathbf{M}}$ and \mathbf{M} are either asymmetric or rectangular. Let \mathbf{M} be a $n_1 \times n_2$ matrix with $\operatorname{rk}(\mathbf{M}) = k^*$ and consider the additive perturbation $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{E} is a $n_1 \times n_2$ noise matrix. We once again assume that both \mathbf{M} and \mathbf{E} are unobserved, i.e., we only observed $\hat{\mathbf{M}}$. Let $n = \max\{n_1, n_2\}$ and let ψ denote the condition number of \mathbf{M} . Finally denote the SVD of \mathbf{M} as $\mathbf{M} = \mathbf{U}\Sigma\mathbf{V}^\top$ where $\Sigma \in \mathbb{R}^{k^* \times k^*}$ is the diagonal matrix of non-zero singular values.

Given a matrix $\hat{\mathbf{M}}$ the rs -RSVD procedure for approximating the leading left singular vectors of $\hat{\mathbf{M}}$ is presented in Algorithm S.1. The leading right singular vectors of $\hat{\mathbf{M}}$ can be approximated by considering $\hat{\mathbf{M}}^\top$ and thus, without loss of generality, we will only focus on the left singular vectors of $\hat{\mathbf{M}}$. Note that Algorithm S.1 is equivalent to the PowerRangeFinder and SubspacePowerMethod procedures in [58, 124] when $a_n = 1$. We now introduce a few assumptions on \mathbf{M} and \mathbf{E} ; these assumptions are natural generalizations of Assumptions 1–2 for the symmetric case.

ASSUMPTION S.1 (Signal-to-noise ratio). Let $S_{n_1, n_2} = \|\mathbf{M}\|$, $n = \max\{n_1, n_2\}$, and $\eta = \max\{n_2, \log n\}$. There exist fixed but arbitrary constants $\beta_\Delta > 0$, $N_0 > 0$, and a quantity $E_{n_1, n_2} > 0$ depend on n_1, n_2 , such that $S_{n_1, n_2}/E_{n_1, n_2} \gtrsim \eta^{\beta_\Delta}$ and

$$\mathbb{P}[\|\mathbf{E}\| \leq E_{n_1, n_2}] \geq 1 - n^{-6}$$

whenever $n \geq N_0$. We view E_{n_1, n_2} as the noise level of \mathbf{E} and η^β as a lower bound on the SNR.

ASSUMPTION S.2. S.2 There exists some positive constants $\delta > 1$ such that, for any positive integer $g \leq \log n$, any standard basis $\mathbf{e}_i \in \mathbb{R}^{n_1}$, and any given vector \mathbf{u} **not** dependent on n , we have with probability at least $1 - n^{-8}$ that

$$|\mathbf{e}_i^\top \underbrace{\mathbf{E}\mathbf{E}^\top \mathbf{E} \cdots \mathbf{E}^\star}_{g \text{ terms}} \mathbf{u}| \leq C'_{2 \rightarrow \infty} E_{n_1, n_2}^g (\log n)^{\delta g} \|\mathbf{u}\|_{\max},$$

when n is sufficiently large. Here $\mathbf{u} \in \mathbb{R}^{n_1}$ and $\mathbf{E}^\star = \mathbf{E}^\top$ when g is even, and $\mathbf{u} \in \mathbb{R}^{n_2}$ and $\mathbf{E}^\star = \mathbf{E}$ otherwise. The $C'_{2 \rightarrow \infty}$ is a finite constant not depending on n and g .

REMARK S.2.1 (Divergence Regime). In contrast to the symmetric matrix case, the quantities S_{n_1, n_2} and E_{n_1, n_2} now depend on both n_1 and n_2 and hence β_Δ now depends on the growth rate of n_1 compared to n_2 . For an illustrative example, suppose that \mathbf{M} is a homogenous matrix whose entries are all of order $\Theta(1)$ and let \mathbf{E} be a standard Gaussian matrix. Assume also that $\text{rk}(\mathbf{M}) = k^*$ for some constant k^* . We then have $S_{n_1, n_2} \asymp \sqrt{n_1 n_2}$ and $E_{n_1, n_2} \asymp \sqrt{n}$. Next suppose n_1 and n_2 are growing such that $n_2 = n_1^c$ for some constant $c > 1$. We then have $\eta = n_2$, $S_{n_1, n_2}/E_{n_1, n_2} \asymp n_2^{1/(2c)}$, and $\beta_\Delta = (2c)^{-1}$.

REMARK S.2.2 (Assumption S.2 Verification). If the entries in \mathbf{E} are independent then we can verify Assumption S.2 by applying Proposition 3.2 together with a ‘‘symmetric dilation’’ argument. Consider, as an example, the case when \mathbf{E} is a standard Gaussian matrix as described in Remark S.2.1 with $E_{n_1, n_2} \asymp \sqrt{n}$. Now define

$$\mathbf{E}^+ := \begin{pmatrix} \mathbf{0} & \mathbf{E} \\ \mathbf{E}^\top & \mathbf{0} \end{pmatrix} \in \mathbb{R}^{n \times n}.$$

Then \mathbf{E}^+ is a symmetric matrix whose (upper triangular) entries are independent random variables satisfying the conditions in Proposition 3.2; see Example 2. We can therefore apply Proposition 3.2 to \mathbf{E}^+ and show, for any standard basis $\mathbf{e}_i \in \mathbb{R}^{n_1}$, any vector $\mathbf{u} \in \mathbb{R}^{n_1}$ and $g \leq \log n$ with g even, that

$$|\mathbf{e}_i^\top (\mathbf{E}\mathbf{E}^\top)^{g/2} \mathbf{u}| = |(\mathbf{e}_i^\top \mathbf{0}^\top) (\mathbf{E}^+)^g \begin{pmatrix} \mathbf{u} \\ \mathbf{0} \end{pmatrix}| \leq 2E_{n_1, n_2}^g (\log n)^{\delta g} \|\mathbf{u}\|_{\max},$$

with high probability. Bound for $\mathbf{e}_i^\top (\mathbf{E}\mathbf{E}^\top)^{(g-1)/2} \mathbf{E}\mathbf{u}$ when $g \leq \log n$ is odd is derived similarly and hence \mathbf{E} satisfies Assumption S.2.

The following bounds for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ for asymmetric $\hat{\mathbf{M}}$ follow *mutatis mutandis* from the proofs of Theorems 3.1–3.3 and the conditions posited in Assumptions S.1 and S.2; see Section S.6.6 for a sketch of the arguments. We surmise that normal approximations for the row-wise fluctuations of $\hat{\mathbf{U}}_g$ are also available in the asymmetric setting and we leave the precise formulation of this result and its conditions to the interested reader.

THEOREM S.2.1. *Suppose Assumption S.1 holds, and that $\hat{\mathbf{U}}_g$ is computed using Algorithm S.1 with $k = k^* \leq \tilde{k}$ satisfying one of the three conditions (a)–(c) as specified in Theorem 3.2. Assume also that $\psi = o(S_{n_1, n_2}/E_{n_1, n_2})$. Denote $\vartheta = \max\{1, \tilde{k}^{-1/2} \log^{1/2} n\}$. Let $g \geq (4\beta_\Delta)^{-1}$ and $a_n = \lceil C\tilde{k}^{-1} \log n \rceil$ where C is a constant not depending on n . We then have the following bounds.*

(Spectral-norm Concentration): *With probability at least $1 - n^{-5}$,*

$$(S.2.1) \quad \|(\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} - (\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G}\| \lesssim \vartheta \tilde{k}^{1/2} S_{n_1, n_2}^{2g} E_{n_1, n_2}.$$

(ℓ_2 Perturbation): *With probability at least $1 - n^{-5}$,*

$$(S.2.2) \quad d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \leq \sqrt{2} \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U})\| \lesssim \vartheta \psi^{2g+1} \frac{E_{n_1, n_2}}{S_{n_1, n_2}}.$$

($\ell_{2, \infty}$ Perturbation): *If \mathbf{E} satisfies Assumption S.2 then, with probability at least $1 - n^{-5}$,*

$$(S.2.3) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim (\log n)^{\delta^\circ} \vartheta \psi^{2g+1} k^{1/2} \frac{E_{n_1, n_2}}{S_{n_1, n_2}} \|\mathbf{U}\|_{2 \rightarrow \infty},$$

where $\delta^\circ = \delta$ if $g > (4\beta_\Delta)^{-1}$ and $\delta^\circ = (2g + 1)\delta + 1/2$ if $g = (4\beta_\Delta)^{-1}$.

Note that the parameter g in Algorithm S.1 and Theorem S.2.1 actually represents $2g + 1$ power iterations of $\hat{\mathbf{M}}$. We show in Section 4 that, in the context of random graphs inference, doing 3 or 5 power iterations is often sufficient for $\hat{\mathbf{U}}_g$ to converge to \mathbf{U} in both d_2 and $d_{2 \rightarrow \infty}$ distances. This is consistent with the recommendation of $g = 1$ or $g = 2$ provided in [58, Section 1.6].

S.3. General theoretical results for any \mathbf{G} . We first state a set of slightly more general versions of Theorem 3.1, Theorem 3.2, and Theorem 3.3. More specifically the following results provide bounds for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ conditional on \mathbf{G} , i.e., the resulting bounds depend on several quantities associated with \mathbf{G} such as $\|\mathbf{G}\|$. Given these results, we derive Theorem 3.1 through Theorem 3.3 by bounding the quantities that depend on \mathbf{G} separately through a set of technical lemmas. In particular Lemma S.4.4 and Lemma S.4.5 provide these bounds for the case where \mathbf{G} is chosen via repeated sampling of independent Gaussian matrices as described in Algorithm 1 of this paper. We note that this type of approach allow us to easily extend our results to other class of sketching matrices. Indeed, suppose \mathbf{G} is a random Rademacher matrix or a random column-subsampling matrix. Then Theorem 3.1 through Theorem 3.3 will continue to hold for this choice of \mathbf{G} provided that we can establish analogous results to that of Lemma S.4.4 and Lemma S.4.5.

THEOREM S.3.1. *Let $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} and \mathbf{E} satisfy Assumption 1. Let $g \geq 1$ be a fixed but arbitrary integer and let \mathbf{G} be any $n \times \tilde{k}$ matrix. For $\xi \geq 1$ let $\tilde{\mathbf{V}}_\xi \in \mathbb{R}^{k \times n}$ be the matrix whose columns are the right singular vectors of $\mathbf{U}^\top \mathbf{E}^\xi$ and define*

$$(S.3.1) \quad r_2(\mathbf{G}) := E_n^g \|\mathbf{G}\| + \mathbb{I}(g \geq 2) S_n^{g-1} E_n \left\{ \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| + \|\mathbf{U}^\top \mathbf{G}\| \right\}.$$

Then for sufficiently large n , we have with probability at least $1 - n^{-6}$ that

$$(S.3.2) \quad \|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\| \lesssim r_2(\mathbf{G}).$$

The bound for $\|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\|$ in Theorem S.3.1 together with the Wedin $\sin \Theta$ Theorem [122] implies the following bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_2(\hat{\mathbf{V}}_g, \mathbf{V})$.

THEOREM S.3.2. *Assume the setting in Theorem S.3.1. Then for sufficiently large n , we have with probability at least $1 - n^{-6}$ that*

$$(S.3.3) \quad \max \{d_2(\hat{\mathbf{U}}_g, \mathbf{U}), d_2(\hat{\mathbf{V}}_g, \mathbf{V}_g)\} \lesssim \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})}.$$

THEOREM S.3.3. *Assume the setting in Theorem S.3.1. Suppose furthermore that \mathbf{E} satisfies Assumption 2. Next define*

$$r_{2 \rightarrow \infty}(\mathbf{G}) := \tilde{k}^{1/2} E_n^g (\log n)^{\delta g} \|\mathbf{G}\|_{\max} \\ + \mathbb{I}(g \geq 2) (\log n)^\delta \left\{ \|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \right\} k^{1/2} S_n^{g-1} E_n \|\mathbf{U}\|_{2 \rightarrow \infty},$$

where δ is the constant appearing in Assumption 2. Then for sufficiently large n , we have with probability at least $1 - C_g n^{-6}$ that

$$d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} + \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Note $C_g > 0$ is a fixed constant only dependent on g .

S.4. Technical lemmas. Before commencing with the proofs of the main results, we first state some technical lemmas that we will repeatedly use throughout this paper. We start by listing some basic properties of the $\ell_{2 \rightarrow \infty}$ norm.

LEMMA S.4.1. *For any $\mathbf{M}_1 \in \mathbb{R}^{d_1 \times d_2}$, $\mathbf{M}_2 \in \mathbb{R}^{d_2 \times d_3}$ and $\mathbf{M}_3 \in \mathbb{R}^{d_4 \times d_1}$, we have*

$$\|\mathbf{M}_1 \mathbf{M}_2\|_{2 \rightarrow \infty} \leq \|\mathbf{M}_1\|_{2 \rightarrow \infty} \|\mathbf{M}_2\|; \\ \|\mathbf{M}_3 \mathbf{M}_1\|_{2 \rightarrow \infty} \leq \|\mathbf{M}_3\|_\infty \|\mathbf{M}_1\|_{2 \rightarrow \infty}; \\ \|\mathbf{M}_1\|_\infty \leq \sqrt{d_2} \|\mathbf{M}_1\|_{2 \rightarrow \infty}.$$

For a proof of Lemma S.4.1, see e.g., [22]. The next two results, namely Lemma S.4.2 and Lemma S.4.3, will be used for bounding $\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}$ in Theorem 3.1.

LEMMA S.4.2 (Decomposition of powered matrices difference). *Let $\hat{\mathbf{M}}$ and \mathbf{M} be two square matrices. Denote $\mathbf{E} = \hat{\mathbf{M}} - \mathbf{M}$. Then for any integer $g \geq 2$ we have the expansion*

$$\hat{\mathbf{M}}^g - \mathbf{M}^g = \mathbf{E}^g + \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi + \sum_{\xi=0}^{g-2} \mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1}.$$

PROOF OF LEMMA S.4.2. Let $\mathbf{E} = \hat{\mathbf{M}} - \mathbf{M}$. We then have the expansion

$$(S.4.1) \quad \hat{\mathbf{M}}^g - \mathbf{M}^g = \sum_{\ell=0}^{g-2} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\ell} + \hat{\mathbf{M}}^{g-1} \mathbf{E} \\ = \sum_{\ell=0}^{g-2} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\ell} + \mathbf{M}^{g-1} \mathbf{E} + (\hat{\mathbf{M}}^{g-1} - \mathbf{M}^{g-1}) \mathbf{E}.$$

If $g \geq 3$ then we can repeat the above step for $\hat{\mathbf{M}}^{g-1} - \mathbf{M}^{g-1}$ and obtain

$$\hat{\mathbf{M}}^g - \mathbf{M}^g = \sum_{\xi=0}^1 \left\{ \sum_{\ell=0}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \right\} \mathbf{E}^\xi + \sum_{\xi=0}^1 \mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} + (\hat{\mathbf{M}}^{g-2} - \mathbf{M}^{g-2}) \mathbf{E}^2.$$

In summary, by iterating the expansion in Eq. (S.4.1) a total of $g - 1$ times, we derive

$$\hat{\mathbf{M}}^g - \mathbf{M}^g = \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi + \sum_{\xi=0}^{g-2} \mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} + \mathbf{E}^g$$

as desired. \square

LEMMA S.4.3. *Let $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} and \mathbf{E} satisfy Assumption 1. Let \mathbf{G} be any given matrix. Then $\|\mathbf{M}^g \mathbf{G}\| \lesssim S_n^g \|\mathbf{U}^\top \mathbf{G}\|$. Let $g \geq 1$ be fixed but arbitrary. Then for sufficiently large n , we have with probability at least $1 - n^{-6}$ that simultaneously,*

$$\begin{aligned} \|\hat{\mathbf{M}}\| &\lesssim S_n, \\ \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| &\lesssim E_n^g \|\tilde{\mathbf{V}}_g^\top \mathbf{G}\|. \end{aligned}$$

Here $\tilde{\mathbf{V}}_\xi \in \mathbb{R}^{n \times k}$ for any $\xi \geq 1$ is the right singular matrix of $\mathbf{U}^\top \mathbf{E}^\xi \mathbf{G} \in \mathbb{R}^{k \times n}$ and $r_{2,n}(\mathbf{G})$ is as defined in Theorem S.3.1.

PROOF OF LEMMA S.4.3. By definition, we have

$$\|\mathbf{M}^g \mathbf{G}\| = \|\mathbf{U} \mathbf{\Lambda}^g \mathbf{U}^\top \mathbf{G}\| = \|\mathbf{\Lambda}^g \mathbf{U}^\top \mathbf{G}\| \leq \|\mathbf{\Lambda}^g\| \cdot \|\mathbf{U}^\top \mathbf{G}\| \asymp S_n^g \|\mathbf{U}^\top \mathbf{G}\|.$$

Next, by Assumptions 1, we have $\|\hat{\mathbf{M}}\| \leq \|\hat{\mathbf{M}} - \mathbf{M}\| + \|\mathbf{M}\| \lesssim S_n + E_n \lesssim S_n$, with probability at least $1 - n^{-6}$ as $S_n \geq C_0 n^{\beta_\Delta} E_n \gg E_n$. We now bound $\mathbf{U}^\top \mathbf{E}^g \mathbf{G}$. Since $\mathbf{U}^\top \mathbf{E}^g$ is a $k \times n$ matrix, let us denote the SVD of $\mathbf{U}^\top \mathbf{E}^g \mathbf{G}$ by

$$\mathbf{U}^\top \mathbf{E}^g \mathbf{G} = \tilde{\mathbf{U}} \tilde{\mathbf{\Sigma}} \tilde{\mathbf{V}}_g^\top,$$

where $\tilde{\mathbf{U}}$ is a $k \times k$ matrix, $\tilde{\mathbf{\Sigma}}$ is a $k \times k$ diagonal matrix, and $\tilde{\mathbf{V}}_g$ is a $n \times k$ matrix. Once again by Assumption 1 we have

$$(S.4.2) \quad \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| = \|\tilde{\mathbf{U}} \tilde{\mathbf{\Sigma}} \tilde{\mathbf{V}}_g^\top \mathbf{G}\| \leq \|\tilde{\mathbf{\Sigma}}\| \|\tilde{\mathbf{V}}_g^\top \mathbf{G}\| \leq \|\mathbf{E}\|^g \|\tilde{\mathbf{V}}_g^\top \mathbf{G}\| \lesssim E_n^g \|\tilde{\mathbf{V}}_g^\top \mathbf{G}\|$$

with probability at least $1 - n^{-6}$. Note the above upper bounds only relies on the high probability bound of $\|\mathbf{E}\|$, with probability at least $1 - n^{-6}$, and thus they hold simultaneously with probability at least $1 - n^{-6}$. \square

The following Lemma S.4.4 and Lemma S.4.5 provide a collection of bounds for the quantities depending on \mathbf{G} in the statement of Theorem S.3.1 through Theorem S.3.3, when \mathbf{G} is chosen via repeated sampling of independent Gaussian matrices as described in Algorithm 1.

LEMMA S.4.4. *Consider the setting in Lemma S.4.3 and let \mathbf{G} be the sketching matrix generated via Algorithm 1 where $a_n \lesssim (\log n)^{\beta_2}$ for some fixed $\beta_2 > 0$. Then for sufficiently large n , we have with probability at least $1 - n^{-6}$ that simultaneously,*

$$\begin{aligned} \|\mathbf{G}\|_{\max} &\leq C_u (\log n)^{1/2}, \\ \|\mathbf{G}\| &\leq C_u \sqrt{n}, \\ \|\mathbf{U}^\top \mathbf{G}\| + \max_{1 \leq \xi \leq g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| &\leq C_u \vartheta \tilde{k}^{1/2}. \end{aligned}$$

Here $\tilde{\mathbf{V}}_\xi$ is as defined in Lemma S.4.3, $\vartheta := \max\{1, \tilde{k}^{-1/2} \log^{1/2} n\}$, and C_u is a constant depending only on β_2 .

PROOF OF LEMMA S.4.4. In the following argument we always condition on \mathbf{M} and $\hat{\mathbf{M}}$. As the final bounds for $\|\mathbf{G}\|_{\max}$, $\|\mathbf{G}\|$ and $\|\mathbf{U}^\top \mathbf{G}\| + \max_{1 \leq \xi \leq g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|$ does not depend on the choice of \mathbf{M} and $\hat{\mathbf{M}}$, we can then uncondition with respect to \mathbf{M} and $\hat{\mathbf{M}}$ to obtain the desired results.

We first bound $\|\mathbf{G}\|_{\max}$. We have

$$\|\mathbf{G}\|_{\max} = \max_{a \in [a_n]} \|\mathbf{G}_a\|_{\max} = \max_{a \in [a_n]} \max_{i \in [n], j \in [\tilde{k}]} |g_{ij}^{(a)}|,$$

where $g_{ij}^{(a)}$ is ij th entry of \mathbf{G}_a . Since $g_{ij}^{(a)}$ is standard Gaussian, there exists a universal constant $K > 0$ such that $\mathbb{P}(|g_{ij}^{(a)}| < t) = 1 - 2 \exp(-t^2/K)$ for all $t > 0$; see e.g Section 2.5.1 of [120]. Let $c > 8$ be arbitrary. Then by the union bound, we have

$$\begin{aligned} \mathbb{P}\left(\|\mathbf{G}\|_{\max} \leq (cK \log n)^{1/2}\right) &= \mathbb{P}\left(\max_{i,j,a} |g_{ij}^{(a)}| \leq (cK \log n)^{1/2}\right) \\ \text{(S.4.3)} \quad &\geq 1 - 2n\tilde{k}a_n \cdot \exp\{-c \log n\} \geq 1 - n^{-6}/3. \end{aligned}$$

Note that the last inequality follows from $\tilde{k}a_n \leq n \log^{\beta_2} n$.

We next bound $\|\mathbf{G}\|$. Recall the non-asymptotic bound for the spectral norm of a Gaussian random matrix (see e.g. Corollary 5.35 in [119]). Then for a $d_1 \times d_2$ standard Gaussian matrix \mathbf{Z}_{d_1, d_2} and $t \geq 0$, we have

$$\text{(S.4.4)} \quad \|\mathbf{Z}_{d_1, d_2}\| \leq \sqrt{d_1} + \sqrt{d_2} + t,$$

with probability at least $1 - 2 \exp(-t^2/2)$. As each \mathbf{G}_a is a $n \times \tilde{k}$ matrix with iid standard Gaussian entries, we can set $d_1 = n$, $d_2 = \tilde{k} \leq d_1$ and $t = n^{1/2}$ in Eq. (S.4.4) to obtain $\|\mathbf{G}_a\| \leq 3\sqrt{n}$ with probability at least $1 - 2 \exp(-n/2)$. A union bound over all $a \in [a_n]$ yields

$$\text{(S.4.5)} \quad \mathbb{P}(\|\mathbf{G}\| \leq 3\sqrt{n}) \geq \mathbb{P}(\max_{a \in [a_n]} \|\mathbf{G}_a\| \leq 3\sqrt{n}) \geq 1 - 2a_n \exp(-n/2) \geq 1 - n^{-6}/3.$$

We finally bound $\|\mathbf{U}^\top \mathbf{G}\| + \max_{1 \leq \xi \leq g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|$. Recall that in Algorithm 1 the $\mathbf{G}_1, \dots, \mathbf{G}_{a_n}$ are generated independently of the $\hat{\mathbf{M}}$ and \mathbf{M} . Hence, conditional on $\hat{\mathbf{M}}$, the $\mathbf{U}^\top \mathbf{G}_a$ and $\tilde{\mathbf{V}}_\xi^\top \mathbf{G}_a$, for $\xi \in [g]$ and $a \in [a_n]$, are $k \times \tilde{k}$ standard Gaussian matrices. Recall also that $\tilde{k} \geq k$. By Eq. (S.4.4), we therefore have $\|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}_a\| \leq 2\tilde{k}^{1/2} + (2c \log n)^{1/2}$ with probability at least $1 - 2n^{-c}$. The same bound also holds for $\mathbf{U}^\top \mathbf{G}_a$. Now choose $c > 7$. We then have, by a union bound over $a \in [a_n]$ and $\xi \in [g]$, that when n is sufficiently large,

$$\begin{aligned} \mathbb{P}\left(\max\{\|\mathbf{U}^\top \mathbf{G}\|, \max_{\xi \in [g]} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|\} \leq 2\tilde{k}^{1/2} + (2c \log n)^{1/2} \mid \hat{\mathbf{M}}\right) &\geq 1 - 2a_n(g+1)n^{-7} \\ &\geq 1 - n^{-6}/3, \end{aligned}$$

where the last inequality follows from the fact that $a_n g = o(n)$. \square

LEMMA S.4.5. *Suppose Assumption 1 hold, and \mathbf{G} is generated via Algorithm 1 with $1 \leq k \leq \tilde{k} \leq n$ where k and \tilde{k} satisfy either one of the three following conditions*

- (a). *Both k and \tilde{k} are fixed.*
- (b). *k is fixed and \tilde{k} is growing with n .*
- (c). *k is growing with n and $\tilde{k} \geq c_{\text{gap}} k$ for some fixed $c_{\text{gap}} > 1$.*

Now choose $a_n \geq \max\{C\tilde{k}^{-1} \log n, 1\}$ for some sufficiently large but fixed constant $C > 0$; a detailed discussion for the value of C is given in Remark 3.3. Assume also that the condition number ψ satisfies $\psi = o(S_n/E_n)$. Let $g \geq 1$ be fixed but arbitrary. Then with probability at least $1 - 2n^{-6}$, we have

$$\sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} \psi^g S_n^{-g}.$$

PROOF OF LEMMA S.4.5. We recall the notations in Eq. (2.1). For any $a \in [a_n]$ we have

$$(S.4.6) \quad \mathbf{G}_a^\top \hat{\mathbf{M}}^{2g} \mathbf{G}_a = \mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a + \mathbf{G}_a^\top \hat{\mathbf{U}}_\perp \hat{\mathbf{\Lambda}}_\perp^{2g} \hat{\mathbf{U}}_\perp^\top \mathbf{G}_a.$$

Now $\mathbf{G}_a^\top \hat{\mathbf{U}}_\perp \hat{\mathbf{\Lambda}}_\perp^{2g} \hat{\mathbf{U}}_\perp^\top \mathbf{G}_a$ is positive semi-definitive and hence, by Weyl's inequality,

$$(S.4.7) \quad \sigma_k^2(\hat{\mathbf{M}}^g \mathbf{G}_a) = \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{M}}^{2g} \mathbf{G}_a) \geq \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a).$$

From the description of Algorithm 1 we have

$$(S.4.8) \quad \sigma_k^2(\hat{\mathbf{M}}^g \mathbf{G}) = \max_{a \in [a_n]} \sigma_k^2(\hat{\mathbf{M}}^g \mathbf{G}_a) \geq \max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a).$$

We now show that the term on the right-hand side of the above display can be lower-bounded by a term of the form $\tilde{k} \psi^{-2g} S_n^{2g}$.

As $\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a$ is positive semi-definitive for all a , we have

$$\begin{aligned} \max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) &\geq \lambda_k(\hat{\mathbf{\Lambda}})^{2g} \cdot \max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{U}}^\top \mathbf{G}_a) \\ &= \sigma_k^{2g}(\hat{\mathbf{M}}) \cdot \max_{a \in [a_n]} \sigma_k^2(\hat{\mathbf{U}}^\top \mathbf{G}_a). \end{aligned}$$

Note that, by Assumption 1 and Weyl's inequality, we have

$$\sigma_k(\hat{\mathbf{M}}) \geq \sigma_k(\mathbf{M}) - \|\hat{\mathbf{M}} - \mathbf{M}\| \geq \psi^{-1} c_l S_n - E_n,$$

with probability at least $1 - n^{-6}$. Recall that we had assumed $\psi = o(S_n/E_n)$ in the statement of Lemma S.4.5. Thus E_n is negligible compared to $\psi^{-1} c_l S_n$ when n increases. We therefore have

$$(S.4.9) \quad \sigma_k(\hat{\mathbf{M}}) \geq \psi^{-1} c_l S_n / 2.$$

with probability at least $1 - n^{-6}$.

Thus, to obtain a lower bound for $\sigma_k^2(\hat{\mathbf{M}}^g \mathbf{G})$ in Eq. (S.4.8), we only need to lower bound $\max_{a \in [a_n]} \sigma_k^2(\hat{\mathbf{U}}^\top \mathbf{G}_a)$ in Eq. (S.4.9). Let $\mathbf{Z}_{k, \tilde{k}} \in \mathbb{R}^{k \times \tilde{k}}$ be a random matrix whose entries are independent standard normal random variables and let $\Psi_{k, \tilde{k}}(t)$ be the cumulative distribution function (CDF) for the smallest singular value of $\mathbf{Z}_{k, \tilde{k}}$; i.e., define

$$(S.4.10) \quad \Psi_{k, \tilde{k}}(t) = \mathbb{P}(\sigma_k(\mathbf{Z}_{k, \tilde{k}}) \leq t).$$

for all $t \geq 0$. We first condition on an observed $\hat{\mathbf{M}}$. Then for any $a \in [a_n]$, the matrix $\hat{\mathbf{U}}^\top \mathbf{G}_a$ is identically distributed as $\mathbf{Z}_{k, \tilde{k}}$ and furthermore the collection of matrices $\{\hat{\mathbf{U}}^\top \mathbf{G}_a\}_{a \in [a_n]}$ are mutually independent. Thus, by the definition of $\Psi_{k, \tilde{k}}$, we have

$$\mathbb{P}\left(\max_{a \in [a_n]} \sigma_k(\hat{\mathbf{U}}^\top \mathbf{G}_a) \geq \epsilon_n \mid \hat{\mathbf{M}}\right) = 1 - (\Psi_{k, \tilde{k}}(\epsilon_n))^{a_n}.$$

for any arbitrary $\epsilon_n > 0$. As the probability on the right-hand side above does not depend on the particular choice of $\hat{\mathbf{M}}$, we can now uncondition on $\hat{\mathbf{M}}$ to obtain

$$\mathbb{P}\left(\max_{a \in [a_n]} \sigma_k(\hat{\mathbf{U}}^\top \mathbf{G}_a) \geq \epsilon_n\right) = 1 - (\Psi_{k, \tilde{k}}(\epsilon_n))^{a_n}.$$

Combining Eq.(S.4.9) and Eq. (S.4.11) we have

$$(S.4.11) \quad \max_{a \in a_n} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) \gtrsim \epsilon_n^2 \psi^{-2g} S_n^{2g},$$

with probability at least $1 - n^{-6} - (\Psi_{k, \tilde{k}}(\epsilon_n))^{a_n}$.

We now consider three scenarios, namely (a) k and \tilde{k} are both fixed, (b) k is fixed and \tilde{k} is growing with n and (c) k and \tilde{k} are both growing with n .

Case (a). Suppose both k and $\tilde{k} \geq k$ are fixed. Then for any arbitrary $\eta_0 > 0$ there exists a finite $\epsilon^* > 0$ such that $\Psi_{k, \tilde{k}}(\epsilon^*) \leq 1 - \eta_0$ (see e.g. Section 4.4 in [43]); we emphasize that ϵ^* depends on η_0 , k and \tilde{k} but does not depend on n . There thus exists a constant $C > 0$ depending only on ϵ^* such that if $a_n \geq C \log n$ then $\{\Psi_{k, \tilde{k}}(\epsilon^*)\}^{a_n} \leq n^{-6}$. Set $\epsilon_n = \epsilon^*$. Then by Eq. (S.4.11), we have $\max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) \gtrsim \psi^{-2g} S_n^{2g}$ with probability at least $1 - 2n^{-6}$.

Case (b). Suppose k is fixed and $\tilde{k} \geq k$ is growing with n . Then by the non-asymptotic bound of a ‘‘tall’’ Gaussian matrix [37] we have

$$(S.4.12) \quad \Psi_{k, \tilde{k}}(\tilde{k}^{1/2} - k^{1/2} - t) \leq 2e^{-t^2/2} = \exp(\log 2 - t^2/2).$$

Now let $t = 0.9\tilde{k}^{1/2} + C'$ and $\epsilon_n = 0.1\tilde{k}^{1/2} - k^{1/2} - C'$ for some sufficiently large but finite $C' > 0$. As k is fixed, Eq. (S.4.11) implies

$$(S.4.13) \quad \max_{1 \leq a \leq a_n} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) \gtrsim \epsilon_n^2 \psi^{-2g} S_n^{2g} \gtrsim \tilde{k} \psi^{-2g} S_n^{2g},$$

with probability at least

$$\begin{aligned} 1 - n^{-6} - \{\Psi_{k, \tilde{k}}(0.1\tilde{k}^{1/2} - k^{1/2} - C')\}^{a_n} &\geq 1 - n^{-6} - \exp\left[\frac{1}{2}a_n\{\log 2 - (0.9\tilde{k}^{1/2} + C')^2\}\right] \\ &\geq 1 - n^{-6} - \exp(-0.4a_n\tilde{k}). \end{aligned}$$

Note that the second to last inequality in the above display follows from Eq. (S.4.12), provided that $C' \geq (2 \log 2)^{1/2}$. Let $C \geq 15$. Then for $a_n \geq Ck^{-1} \log n$ we have that Eq. (S.4.13) holds with probability at least $1 - n^{-6} - \exp(-0.4C \log n) \geq 1 - 2n^{-6}$.

Case (c). Suppose $\tilde{k} \geq c_{\text{gap}} k$ for some $c_{\text{gap}} > 1$. Let $t = \frac{1}{2}(1 - c_{\text{gap}}^{-1/2})\tilde{k}^{1/2} + C'$ for some $C' \geq (2 \log 2)^{1/2}$. Then by Eq. (S.4.12), we have

$$(S.4.14) \quad \Psi_{k, \tilde{k}}\left(\frac{1}{2}(1 + c_{\text{gap}}^{-1/2})\tilde{k}^{1/2} - k^{1/2} - C'\right) \leq \exp\left(-\frac{1}{8}(1 - c_{\text{gap}}^{-1/2})^2 \tilde{k}\right).$$

Note that $1 - c_{\text{gap}}^{-1/2} > 0$ as $c_{\text{gap}} > 1$. Let $\epsilon_n = \tilde{k}^{1/2} - k^{1/2} - t$. We then have

$$(S.4.15) \quad \epsilon_n = \frac{1}{2}(1 + c_{\text{gap}}^{-1/2})\tilde{k}^{1/2} - k^{1/2} - C' \geq \frac{1}{2}(1 - c_{\text{gap}}^{-1/2})\tilde{k}^{1/2} - C' \gtrsim \tilde{k}^{1/2}.$$

Combining Eq. (S.4.11), Eq. (S.4.12), Eq. (S.4.14), and Eq. (S.4.15), we obtain

$$(S.4.16) \quad \max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) \gtrsim \epsilon_n^2 \psi^{-2g} S_n^{2g} \gtrsim \tilde{k} \psi^{-2g} S_n^{2g},$$

with probability at least $1 - n^{-6} - \exp(-\frac{1}{8}(1 - c_{\text{gap}}^{-1/2})^2 \tilde{k} a_n) \geq 1 - 2n^{-6}$, provided that $a_n \geq C\tilde{k}^{-1} \log n$ for some finite $C > 0$. We emphasize that C depends only on c_{gap} such that C decreases when c_{gap} increases.

Summarizing the results for the above three cases considered in Lemma S.4.5, we have that if $a_n \geq \max\{C\tilde{k}^{-1} \log n, 1\}$ then

$$(S.4.17) \quad \sigma_k^2(\hat{\mathbf{M}}^g \mathbf{G}) \geq \max_{a \in [a_n]} \lambda_k(\mathbf{G}_a^\top \hat{\mathbf{U}} \hat{\mathbf{\Lambda}}^{2g} \hat{\mathbf{U}}^\top \mathbf{G}_a) \gtrsim \tilde{k} \psi^{-2g} S_n^{2g},$$

with probability at least $1 - 2n^{-6}$. We therefore have

$$\sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} \psi^g S_n^{-g},$$

with probability at least $1 - 2n^{-6}$. Finally the above derivations also show that (a) C depends only on k and \tilde{k} if k and \tilde{k} are both fixed, (b) C can be any constant larger than 15, and (c) C depends only on c_{gap} when both k and \tilde{k} are growing. \square

The last technical lemma is used when bounding $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ in Theorem 3.3.

LEMMA S.4.6. *Let $\hat{\mathbf{M}} = \mathbf{M} + \mathbf{E}$ where \mathbf{M} and \mathbf{E} satisfy Assumption 2. Let \mathbf{G} be any arbitrary matrix not depending on $\hat{\mathbf{M}}$. Let g be fixed but arbitrary. Then for sufficiently large n , we have with probability at least $1 - C'_g n^{-6}$, that simultaneously*

$$\begin{aligned} \|\mathbf{E}^g \mathbf{U}\|_{2 \rightarrow \infty} &\lesssim k^{1/2} E_n^g(\log n)^{\delta g} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\ \|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} &\lesssim \tilde{k}^{1/2} E_n^g(\log n)^{\delta g} \|\mathbf{G}\|_{\max}, \\ \|\hat{\mathbf{M}}^g \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} &\lesssim k^{1/2} E_n S_n^g \|\mathbf{U}\|_{2 \rightarrow \infty}, \end{aligned}$$

where $C'_g > 0$ is some constant dependent only on g .

PROOF OF LEMMA S.4.6. Under Assumption 2, a similar argument to the section 3.5 of [22] yields

$$\begin{aligned} \|\mathbf{E}^g \mathbf{U}\|_{2 \rightarrow \infty} &\leq k^{1/2} \max_{i \in [n]} \max_{j \in [k]} |e_i^\top \mathbf{E}^g \mathbf{u}_j| \lesssim k^{1/2} E_n^g(\log n)^{\delta g} \max_{j \in [k]} \|\mathbf{u}_j\|_{\max} \\ (S.4.18) \quad &\lesssim k^{1/2} E_n^g(\log n)^{\delta g} \|\mathbf{U}\|_{2 \rightarrow \infty}, \end{aligned}$$

with probability at least $1 - (nk) \cdot n^{-8} \cdot \tilde{k}^{-1} \geq 1 - n^{-6}$. Here e_i denote the i th standard basis in \mathbb{R}^n and \mathbf{u}_j is the j th column vector of \mathbf{U} . The same argument also yields

$$\begin{aligned} \|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} &\leq \tilde{k}^{1/2} \max_{i \in [n]} \max_{j \in [\tilde{k}]} |e_i^\top \mathbf{E}^g \mathbf{g}_j| \lesssim \tilde{k}^{1/2} E_n^g(\log n)^{\delta g} \max_{j \in [\tilde{k}]} \|\mathbf{g}_j\|_{\max} \\ (S.4.19) \quad &\lesssim \tilde{k}^{1/2} E_n^g(\log n)^{\delta g} \|\mathbf{G}\|_{\max}, \end{aligned}$$

with probability at least $1 - (n\tilde{k}) \cdot n^{-8} \cdot \tilde{k}^{-1} \geq 1 - n^{-6}$. Here \mathbf{g}_j is the j th column of \mathbf{G} .

Finally we bound $\|\hat{\mathbf{M}}^g \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty}$. We start with the expansions

$$\begin{aligned} \hat{\mathbf{M}}^g \mathbf{E} \mathbf{U} &= \mathbf{E} \hat{\mathbf{M}}^{g-1} \mathbf{E} \mathbf{U} + \mathbf{M} \hat{\mathbf{M}}^{g-1} \mathbf{E} \mathbf{U} \\ (S.4.20) \quad &= \mathbf{E}^2 \hat{\mathbf{M}}^{g-2} \mathbf{E} \mathbf{U} + \mathbf{E} \mathbf{M} \hat{\mathbf{M}}^{g-2} \mathbf{E} \mathbf{U} + \mathbf{M} \hat{\mathbf{M}}^{g-1} \mathbf{E} \mathbf{U} \quad (\text{if } g \geq 2) \\ &= \dots = \mathbf{E}^{g+1} \mathbf{U} + \sum_{\xi=0}^{g-1} \mathbf{E}^\xi \mathbf{M} \hat{\mathbf{M}}^{g-1-\xi} \mathbf{E} \mathbf{U}. \end{aligned}$$

We will bound the $2 \rightarrow \infty$ norm for each individual term in the right hand side of the above display. By Eq. (S.4.18), we have

$$(S.4.21) \quad \|\mathbf{E}^{g+1} \mathbf{U}\|_{2 \rightarrow \infty} \lesssim k^{1/2} E_n^{g+1}(\log n)^{\delta(g+1)} \|\mathbf{U}\|_{2 \rightarrow \infty}$$

with probability at least $1 - n^{-6}$.

Now let $\xi = 0$. We then have, with probability at least $1 - n^{-6}$,

$$\begin{aligned}
 (S.4.22) \quad \|\mathbf{M}\hat{\mathbf{M}}^{g-1}\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{A}\mathbf{U}^\top \hat{\mathbf{M}}^{g-1}\mathbf{E}\mathbf{U}\| \\
 &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{A}\| \cdot \|\mathbf{U}^\top\| \cdot \|\hat{\mathbf{M}}\|^{g-1} \cdot \|\hat{\mathbf{M}} - \mathbf{M}\| \cdot \|\mathbf{U}\| \\
 &\lesssim S_n^g E_n \|\mathbf{U}\|_{2 \rightarrow \infty} \\
 &\lesssim k^{1/2} S_n^g E_n \|\mathbf{U}\|_{2 \rightarrow \infty},
 \end{aligned}$$

where the first inequality in the above display follows from Lemma S.4.1 and the last two inequalities follow from Lemma S.4.3 and $k \geq 1$.

Next suppose $\xi > 0$. Then using Eq. (S.4.18) we have, with probability at least $1 - 2n^{-6}$,

$$\begin{aligned}
 (S.4.23) \quad \|\mathbf{E}^\xi \mathbf{M} \hat{\mathbf{M}}^{g-1-\xi} \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}^\xi \mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{A} \mathbf{U}^\top \hat{\mathbf{M}}^{g-\xi-1} \mathbf{E} \mathbf{U}\| \\
 &\lesssim k^{1/2} E_n^\xi (\log n)^{\delta \xi} \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot S_n^{g-\xi} E_n \\
 &\lesssim k^{1/2} E_n^{\xi+1} S_n^{g-\xi} (\log n)^{\delta \xi} \|\mathbf{U}\|_{2 \rightarrow \infty},
 \end{aligned}$$

for any $\xi \leq g-1$. Recall that under Assumption 1, we have $S_n \lesssim n^{\beta_\Delta} E_n$ for some $\beta_\Delta > 0$. We therefore have

$$k^{1/2} S_n^g E_n \|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim k^{1/2} E_n^{\xi+1} S_n^{g-\xi} (\log n)^{\delta \xi} \|\mathbf{U}\|_{2 \rightarrow \infty}$$

for any $\xi > 0$. In other words, the bound in Eq. (S.4.23) when $\xi > 1$ is dominated by that when $\xi = 1$. Collecting the bounds in Eq. (S.4.21), Eq. (S.4.22) and Eq. (S.4.23), we obtain

$$\begin{aligned}
 (S.4.24) \quad \|\hat{\mathbf{M}}^g \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} &\lesssim k^{1/2} E_n^{g+1} (\log n)^{\delta(g+1)} \|\mathbf{U}\|_{2 \rightarrow \infty} + k^{1/2} S_n^g E_n \|\mathbf{U}\|_{2 \rightarrow \infty} \\
 &\lesssim k^{1/2} S_n^g E_n \|\mathbf{U}\|_{2 \rightarrow \infty}.
 \end{aligned}$$

with probability at least $1 - (C'_g - 2)n^{-6}$ with some large enough yet fixed $C'_g > 0$ when g is fixed. Note that we had used Assumption 1 to bound $\tilde{k}^{1/2} E_n^{g+1} (\log n)^{\delta(g+1)} = o(S_n^g E_n)$ as n increases. The desired results follow immediately by unifying Eq. (S.4.18), Eq. (S.4.19), and Eq. (S.4.24). \square

S.5. Proofs for Section 3.

S.5.1. *Proof of Theorem S.3.1.* If $g = 1$ then, by Assumption 1, we have

$$\|\hat{\mathbf{M}}\mathbf{G} - \mathbf{M}\mathbf{G}\| \leq \|\mathbf{E}\| \|\mathbf{G}\| \lesssim E_n \|\mathbf{G}\| \leq r_2(\mathbf{G})$$

with probability at least $1 - n^{-6}$. We next consider $g \geq 2$. Let $\Delta_g = (\hat{\mathbf{M}}^g - \mathbf{M}^g)\mathbf{G}$. Using the expansion for $\hat{\mathbf{M}}^g - \mathbf{M}^g$ in Lemma S.4.2, we have

$$(S.5.1) \quad \Delta_g = \underbrace{\mathbf{E}^g \mathbf{G}}_{\mathbf{L}_1} + \underbrace{\sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \mathbf{G}}_{\mathbf{L}_2} + \underbrace{\sum_{\xi=0}^{g-2} \mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} \mathbf{G}}_{\mathbf{L}_3}.$$

We now bound the spectral norms of each of the terms \mathbf{L}_1 through \mathbf{L}_3 .

Bounding $\|\mathbf{L}_1\|$. By Assumption 1 we have, with probability at least $1 - n^{-6}$,

$$(S.5.2) \quad \|\mathbf{L}_1\| \leq \|\mathbf{E}\|^g \|\mathbf{G}\| \lesssim E_n^g \|\mathbf{G}\|.$$

Bounding $\|\mathbf{L}_2\|$. By the triangle inequality we have

$$(S.5.3) \quad \|\mathbf{L}_2\| \leq \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \mathbf{G}\|.$$

First suppose that $\xi \geq 1$. Then with probability at least $1 - n^{-6}$, we have

$$\begin{aligned}
\|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \mathbf{G}\| &= \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-2-\xi-\ell} \mathbf{M} \mathbf{E}^\xi \mathbf{G}\| \\
&= \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-2-\xi-\ell} \mathbf{U} \mathbf{\Lambda} \mathbf{U}^\top \mathbf{E}^\xi \mathbf{G}\| \\
\text{(S.5.4)} \quad &\leq \|\hat{\mathbf{M}}\|^\ell \cdot \|\mathbf{E}\| \cdot \|\mathbf{M}\|^{g-2-\xi-\ell} \cdot \|\mathbf{\Lambda}\| \cdot \|\mathbf{U}^\top \mathbf{E}^\xi \mathbf{G}\| \\
&\lesssim S_n^{g-1-\xi} E_n^{\xi+1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|.
\end{aligned}$$

where the last inequality follows from Lemma S.4.3 and Assumption 1.

If $\xi = 0$ then, with probability at least $1 - n^{-6}$,

$$\begin{aligned}
\|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\ell} \mathbf{G}\| &= \|\hat{\mathbf{M}}\|^\ell \cdot \|\mathbf{E}\| \cdot \|\mathbf{M}\|^{g-1-\ell-1} \cdot \|\mathbf{\Lambda}\| \cdot \|\mathbf{U}^\top \mathbf{G}\| \\
\text{(S.5.5)} \quad &\lesssim S_n^{g-1} E_n \|\mathbf{U}^\top \mathbf{G}\|
\end{aligned}$$

where the last inequality follows from Lemma S.4.3.

Plugging the bounds from Eq. (S.5.4) and Eq. (S.5.5) into Eq. (S.5.3) yield

$$\begin{aligned}
\|\mathbf{L}_2\| &\lesssim \sum_{\xi=0}^{g-2} (g-1-\xi) \cdot S_n^{g-1-\xi} E_n^{\xi+1} \cdot \left\{ \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| + \|\mathbf{U}^\top \mathbf{G}\| \right\} \\
\text{(S.5.6)} \quad &\lesssim S_n^{g-1} E_n \left\{ \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| + \|\mathbf{U}^\top \mathbf{G}\| \right\}.
\end{aligned}$$

with probability at least $1 - n^{-6}$. Here we had used the fact that g is fixed and that $\|S_n^{g-1} E_n\|$ dominates $\|S_n^{g-1-\xi} E_n^{\xi+1}\|$ for all $\xi \geq 0$ as $\|E_n\| \lesssim n^{\beta_E} = o(\|S_n\|)$.

Bounding $\|\mathbf{L}_3\|$ A similar argument to that for bounding $\|\mathbf{L}_2\|$ yields,

$$\begin{aligned}
\|\mathbf{L}_3\| &\leq \sum_{\xi=0}^{g-2} \|\mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} \mathbf{G}\| \\
\text{(S.5.7)} \quad &\leq \sum_{\xi=0}^{g-2} \|\mathbf{M}^{g-2-\xi}\| \cdot \|\mathbf{\Lambda}\| \cdot \|\mathbf{U}^\top \mathbf{E}^{\xi+1} \mathbf{G}\| \\
&\lesssim \sum_{\xi=0}^{g-2} S_n^{g-1-\xi} E_n^{\xi+1} \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \lesssim S_n^{g-1} E_n \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|
\end{aligned}$$

where the penultimate inequality follows from Lemma S.4.3 and Assumption 1.

Combining Eq. (S.5.2), Eq. (S.5.6), and Eq. (S.5.7), we obtain

$$\begin{aligned}
\|(\hat{\mathbf{M}}^g - \mathbf{M}^g) \mathbf{G}\| &\leq \|\mathbf{L}_1\| + \|\mathbf{L}_2\| + \|\mathbf{L}_3\| \\
&\lesssim E_n^g \|\mathbf{G}\| + \mathbb{I}(g \geq 2) S_n^{g-1} E_n \left\{ \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| + \|\mathbf{U}^\top \mathbf{G}\| \right\},
\end{aligned}$$

with probability at least $1 - n^{-6}$, as desired. Note the only probabilistic bound used in proof is $\|\mathbf{E}\| \lesssim E_n$ with probability at least $1 - n^{-6}$, and thus, the final result's probability is still larger than or equal to $1 - n^{-6}$. \square

S.5.2. Proof of Theorem S.3.2. We note that $\hat{\mathbf{M}}^g \mathbf{G}$ is an additive perturbation of $\mathbf{M}^g \mathbf{G}$, i.e.,

$$\hat{\mathbf{M}}^g \mathbf{G} = \mathbf{M}^g \mathbf{G} + (\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}).$$

Since \mathbf{M} is rank k , we have that $\mathbf{M}^g \mathbf{G}$ is also rank k and hence $\sigma_{k+1}(\mathbf{M}^g \mathbf{G}) = 0$. Let \mathbf{U}_g and \mathbf{V}_g be the matrix of left and right singular vectors of $\mathbf{M}^g \mathbf{G}$, respectively. We thus have, by Wedin's sin- Θ theorem [122], that

$$(S.5.8) \quad \max\{\|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|, \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|\} \leq \frac{\|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G}) - \sigma_{k+1}(\mathbf{M}^g \mathbf{G})} \\ \lesssim \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \cdot r_2(\mathbf{G})$$

with probability at least $1 - n^{-6}$. Note that the last equality follows from the bound for $\|(\hat{\mathbf{M}}^g - \mathbf{M}^g) \mathbf{G}\|$ in Theorem S.3.1.

The bound for $d_2(\hat{\mathbf{V}}_g, \mathbf{V}_g)$ given in Theorem S.3.2 follows from Eq. (S.5.8) and the relationship $d_2(\hat{\mathbf{V}}_g, \mathbf{V}_g) \leq \sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|$ (see Lemma 1 in [17]). Similarly, the bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ follows from Eq. (S.5.8) and the fact that $\|\sin \Theta(\mathbf{U}, \mathbf{U}_g)\| = 0$ almost surely (see Proposition 3.1).

S.5.3. Proof of Theorem S.3.3. We denote by $\mathbf{W}_\mathbf{U}$ and $\mathbf{W}_\mathbf{V}$ the solution to the Frobenius-norm Procrustes problems

$$(S.5.9) \quad \mathbf{W}_\mathbf{U} = \operatorname{argmin}_{\mathbf{W}_k \in \mathbb{O}_k} \|\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_k\|_F, \quad \mathbf{W}_\mathbf{V} = \operatorname{argmin}_{\mathbf{W}_k \in \mathbb{O}_k} \|\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{W}_k\|_F.$$

We note that the convergence rate of $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ is equivalent to the convergence rate of $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}_g)$ since, by Proposition 3.1, $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) = 0$ almost surely. Furthermore, from the definition of $d_{2 \rightarrow \infty}$, we have $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}_g) \leq \|\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_\mathbf{U}\|_{2 \rightarrow \infty}$.

We will thus focus our effort on bounding $\|\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_\mathbf{U}\|_{2 \rightarrow \infty}$. Let $\Delta_g = \hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}$. Taking $r = k$ in Theorem 3.1 of [22], we have the decomposition

$$(S.5.10) \quad \hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_\mathbf{U} = \underbrace{(\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \Delta_g \mathbf{V}_g \mathbf{W}_\mathbf{V} \hat{\Sigma}_g^{-1}}_{\mathbf{R}_0} + \underbrace{\mathbf{U}_g (\mathbf{U}_g^\top \hat{\mathbf{U}}_g - \mathbf{W}_\mathbf{U})}_{\mathbf{R}_3} \\ + \underbrace{(\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \Delta_g (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{W}_\mathbf{V}) \hat{\Sigma}_g^{-1}}_{\mathbf{R}_1} \\ + \underbrace{(\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \mathbf{M}^g \mathbf{G} (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{V}_g^\top \hat{\mathbf{V}}_g) \hat{\Sigma}_g^{-1}}_{\mathbf{R}_2}$$

Note that if $k = \tilde{k}$ then $\mathbf{R}_2 = \mathbf{0}$ since $\mathbf{V}_g \mathbf{V}_g^\top = \mathbf{I}$, as $\mathbf{V}_g \in \mathbb{O}_k$. We now bound the $\ell_{2 \rightarrow \infty}$ norms of the three terms (or four terms when $\tilde{k} > k$) on the right-hand side of Eq. (S.5.10).

Bounding $\|\mathbf{R}_0\|_{2 \rightarrow \infty}$. We have, with probability at least $1 - n^{-6}$, that,

$$(S.5.11) \quad \|\mathbf{R}_0\|_{2 \rightarrow \infty} \leq \|\Delta_g \mathbf{V}_g \mathbf{W}_\mathbf{V} \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} + \|\mathbf{U}_g \mathbf{U}_g^\top \Delta_g \mathbf{V}_g \mathbf{W}_\mathbf{V} \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} \\ \leq \|\Delta_g\|_{2 \rightarrow \infty} \cdot \|\hat{\Sigma}_g^{-1}\| + \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\Delta_g\| \cdot \|\hat{\Sigma}_g^{-1}\| \\ = \|\Delta_g\|_{2 \rightarrow \infty} \cdot \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) + \|\Delta_g\| \cdot \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \\ \lesssim \|\Delta_g\|_{2 \rightarrow \infty} \cdot \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) + r_2(\mathbf{G}) \cdot \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}).$$

Here the last inequality uses the bound for $\|\Delta_g\|$ given in Theorem S.3.1 together with the fact that $\mathbf{U}_g = \mathbf{U} \mathbf{W}_{k,g}$ for some $\mathbf{W}_{k,g} \in \mathbb{O}_{k \times k}$ almost surely (see Proposition 3.1) and thus $\|\mathbf{U}_g\|_{2 \rightarrow \infty} = \|\mathbf{U} \mathbf{W}_{k,g}\|_{2 \rightarrow \infty} = \|\mathbf{U}\|_{2 \rightarrow \infty}$.

We now consider $\|\Delta_g\|_{2 \rightarrow \infty}$. If $g = 1$ then by Lemma S.4.6 we have

$$\|\Delta_1\|_{2 \rightarrow \infty} = \|\mathbf{E} \mathbf{G}\|_{2 \rightarrow \infty} \lesssim \tilde{k}^{1/2} E_n(\log n)^\delta \|\mathbf{G}\|_{\max}$$

with probability at least $1 - n^{-6}$. For $g \geq 2$ we first recall the expansion $\mathbf{\Delta}_g = \mathbf{L}_1 + \mathbf{L}_2 + \mathbf{L}_3$ given in Eq. (S.5.1). We now bound $\|\mathbf{L}_1\|_{2 \rightarrow \infty}$ through $\|\mathbf{L}_3\|_{2 \rightarrow \infty}$.

Bound for $\|\mathbf{L}_1\|_{2 \rightarrow \infty}$. By Lemma S.4.6 we have, with probability at least $1 - C'_g n^{-6}$, that

$$(S.5.12) \quad \|\mathbf{L}_1\|_{2 \rightarrow \infty} = \|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} \lesssim \tilde{k}^{1/2} E_n^g (\log n)^{\delta g} \|\mathbf{G}\|_{\max}.$$

Bound for $\|\mathbf{L}_2\|_{2 \rightarrow \infty}$. Using the expansion in Eq. (S.5.3) and Eq. (S.5.4), we have

$$(S.5.13) \quad \begin{aligned} \|\mathbf{L}_2\|_{2 \rightarrow \infty} &\leq \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \mathbf{G}\|_{2 \rightarrow \infty} \\ &= \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U} \mathbf{\Lambda}^{g-1-\xi-\ell} \mathbf{U}^\top \mathbf{E}^\xi \mathbf{G}\|_{2 \rightarrow \infty} \\ &\leq \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Lambda}\|^{g-1-\xi-\ell} \cdot \|\mathbf{U}^\top \mathbf{E}^\xi \mathbf{G}\| \end{aligned}$$

Recall the bound for $\|\mathbf{U}^\top \mathbf{E}^\xi \mathbf{G}\|$ given in Lemma S.4.3. Let $\varphi_g(\mathbf{G}) = \|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|$. We then have, with probability at least $1 - C'_g n^{-6}$,

$$(S.5.14) \quad \begin{aligned} \|\mathbf{L}_2\|_{2 \rightarrow \infty} &\lesssim \varphi_g(\mathbf{G}) \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} S_n^{g-1-\xi-\ell} E_n^\xi \\ &\lesssim \varphi_g(\mathbf{G}) k^{1/2} \sum_{\xi=0}^{g-2} \left(S_n^{g-1-\xi} (\log n)^\delta + \sum_{\ell=1}^{g-2-\xi} S_n^{g-1-\xi} \right) E_n^{\xi+1} \|\mathbf{U}\|_{2 \rightarrow \infty} \\ &\lesssim k^{1/2} \varphi_g(\mathbf{G}) S_n^{g-1} E_n (\log n)^\delta \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

where the penultimate inequality in the above display follows from Lemma S.4.6.

Bound for $\|\mathbf{L}_3\|_{2 \rightarrow \infty}$. From Lemma S.4.3 we have, with probability at least $1 - C'_g n^{-6}$,

$$(S.5.15) \quad \begin{aligned} \|\mathbf{L}_3\|_{2 \rightarrow \infty} &\leq \sum_{\xi=0}^{g-2} \|\mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} \mathbf{G}\|_{2 \rightarrow \infty} \\ &\leq \sum_{\xi=0}^{g-2} \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Lambda}\|^{g-1-\xi} \cdot \|\mathbf{U}^\top \mathbf{E}^{\xi+1} \mathbf{G}\| \\ &\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} S_n^{g-1} E_n \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|. \end{aligned}$$

Combining the above bounds for $\|\mathbf{L}_1\|_{2 \rightarrow \infty}$, $\|\mathbf{L}_2\|_{2 \rightarrow \infty}$ and $\|\mathbf{L}_3\|_{2 \rightarrow \infty}$, we obtain

$$(S.5.16) \quad \begin{aligned} \|\mathbf{\Delta}_g\|_{2 \rightarrow \infty} &\lesssim \tilde{k}^{1/2} E_n^g (\log n)^{\delta g} \|\mathbf{G}\|_{\max} \\ &+ \mathbb{I}(g \geq 2) k^{1/2} S_n^{g-1} E_n (\log n)^\delta \left\{ \|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \right\} \|\mathbf{U}\|_{2 \rightarrow \infty} \\ &= r_{2 \rightarrow \infty}(\mathbf{G}). \end{aligned}$$

with probability at least $1 - C'_g n^{-6}$. Recalling Eq. (S.5.11), we conclude that

$$(S.5.17) \quad \|\mathbf{R}_0\|_{2 \rightarrow \infty} \lesssim \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} + \frac{r_2(\mathbf{G}) \|\mathbf{U}\|_{2 \rightarrow \infty}}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})}$$

with probability at least $1 - (C'_g + 1)n^{-6}$.

Bounding $\|\mathbf{R}_1\|_{2 \rightarrow \infty}$. We have, with probability at least $1 - (1 + C'_g)n^{-6}$, that

$$\begin{aligned}
 \|\mathbf{R}_1\|_{2 \rightarrow \infty} &= \|(\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \mathbf{\Delta}_g (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{W}_V) \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} \\
 &\leq (\|\mathbf{\Delta}_g\|_{2 \rightarrow \infty} + \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Delta}_g\|) \cdot \frac{\sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \\
 &\lesssim \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \left\{ \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \wedge 1 \right\} + \frac{r_2(\mathbf{G}) \|\mathbf{U}\|_{2 \rightarrow \infty}}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \left\{ \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \wedge 1 \right\}
 \end{aligned}
 \tag{S.5.18}$$

Note that the last inequality in the above display follows from Theorem S.3.2 and Eq. (S.5.16) while the bound $\|\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{W}_V\| \leq \sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|$ is given in Lemma 6.8 of [22]. Finally we had used the fact that $\|\mathbf{U}_g\|_{2 \rightarrow \infty} = \|\mathbf{U}\|_{2 \rightarrow \infty}$; see the discussion after Eq. (S.5.11).

Bounding $\|\mathbf{R}_2\|_{2 \rightarrow \infty}$: Recall Eq. (3.1). We immediately have,

$$\begin{aligned}
 \mathbf{R}_2 &= (\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \mathbf{M}^g \mathbf{G} (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{V}_g^\top \hat{\mathbf{V}}_g) \hat{\Sigma}_g^{-1} \\
 &= (\mathbf{I} - \mathbf{U}_g \mathbf{U}_g^\top) \mathbf{U}_g \Sigma_g \mathbf{V}_g^\top (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{V}_g^\top \hat{\mathbf{V}}_g) \hat{\Sigma}_g^{-1} \\
 &= (\mathbf{U}_g \Sigma_g \mathbf{V}_g^\top - \mathbf{U}_g \Sigma_g \mathbf{V}_g^\top) (\hat{\mathbf{V}}_g - \mathbf{V}_g \mathbf{V}_g^\top \hat{\mathbf{V}}_g) \hat{\Sigma}_g^{-1} \\
 &= \mathbf{0}.
 \end{aligned}
 \tag{S.5.19}$$

Bounding $\|\mathbf{R}_3\|_{2 \rightarrow \infty}$. We have, by Lemma S.4.1 and Theorem S.3.2, that

$$\begin{aligned}
 \|\mathbf{R}_3\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\mathbf{U}_g^\top \hat{\mathbf{U}}_g - \mathbf{W}_U\| \\
 &\leq \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \quad (\text{Lemma 6.7 in [22]}) \\
 &\lesssim \left\{ \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \wedge 1 \right\}^2 \cdot \|\mathbf{U}\|_{2 \rightarrow \infty} \quad (\text{Eq. (S.5.8)})
 \end{aligned}
 \tag{S.5.20}$$

with probability at least $1 - n^{-6}$. The last inequality uses the fact that $\|\mathbf{U}_g\|_{2 \rightarrow \infty} = \|\mathbf{U}\|_{2 \rightarrow \infty}$ as discussed in Eq. (S.5.11).

Collecting the bounds for $\|\mathbf{R}_0\|_{2 \rightarrow \infty}$ through $\|\mathbf{R}_3\|_{2 \rightarrow \infty}$ and recalling Eq. (S.5.10), we obtain

$$d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} + \frac{r_2(\mathbf{G}) \|\mathbf{U}\|_{2 \rightarrow \infty}}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})}$$

with probability at least $1 - C_g n^{-6}$ where C_g is a constant depending only on g . Eq. (S.3.4) is thus established. \square

S.5.4. Proof of Theorem 3.1. We prove Theorem 3.1 by combining Theorem S.3.1 and Lemma S.4.4. More specifically, by Theorem S.3.1 we have

$$\|\hat{\mathbf{M}}^g - \mathbf{M}^g\| \lesssim E_n^g \|\mathbf{G}\| + \mathbb{I}(g \geq 2) \cdot S_n^{g-1} E_n \left\{ \|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \right\}.
 \tag{S.5.21}$$

with probability at least $1 - n^{-6}$, where \mathbf{G} is chosen among a_n independent standard Gaussian matrices $\mathbf{G}_1, \mathbf{G}_2, \dots, \mathbf{G}_{a_n}$ according to Algorithm 1. Substituting the bounds for $\|\mathbf{G}\|$ and $\|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi < g} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|$ from Lemma S.4.4 into Eq. (S.5.21) we obtain

$$\|\hat{\mathbf{M}}^g - \mathbf{M}^g\| \lesssim \sqrt{n} E_n^g + \mathbb{I}(g \geq 2) \cdot \vartheta_n \tilde{k}^{1/2} S_n^{g-1} E_n
 \tag{S.5.22}$$

with probability at least $1 - 2n^{-6} \geq 1 - n^{-5}$ when n is sufficiently large. The first part of Theorem 3.1 is thus established.

We now consider the case when $g \geq 1 + (2\beta_\Delta)^{-1}$ and thus $g \geq 2$. Recall that, from Assumption 1, we have $S_n/E_n \lesssim n^{\beta_\Delta}$ with $\beta_\Delta > 0$, and hence

$$(S.5.23) \quad \begin{aligned} \frac{\vartheta_n \tilde{k}^{1/2} S_n^{g-1} E_n}{\sqrt{n} E_n^g} &= \vartheta_n \tilde{k}^{1/2} n^{-1/2} S_n^{g-1} E_n^{1-g} \\ &\lesssim \vartheta_n \tilde{k}^{1/2} n^{(g-1)\beta_\Delta - 1/2} \lesssim \vartheta_n \tilde{k}^{1/2} \lesssim 1. \end{aligned}$$

Eq. (S.5.21) and Eq. (S.5.23) together imply

$$\|\hat{\mathbf{M}}^g \mathbf{G} - \mathbf{M}^g \mathbf{G}\| \lesssim \vartheta_n \tilde{k}^{1/2} S_n^{g-1} E_n$$

with probability at least $1 - 2n^{-6} \geq 1 - n^{-5}$ when n is sufficiently large, as desired. \square

S.5.5. *Proof of Theorem 3.2.* We prove Theorem 3.2 by combining Theorem 3.1, Theorem S.3.2 and Lemma S.4.5. More specifically, by Theorem S.3.2 we have

$$(S.5.24) \quad d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \leq \sqrt{2} \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U})\| \leq \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})}$$

with probability at least $1 - n^{-6}$. Recall that we assumed $\psi = o(S_n/E_n)$ in the statement of Theorem 3.2. Lemma S.4.5 then implies

$$(S.5.25) \quad \hat{\sigma}_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} \psi^g S_n^{-g},$$

with probability at least $1 - 2n^{-6}$. Now when $g \geq 1 + (2\beta_\Delta)^{-1}$, Eq. (3.3) in Theorem 3.1 implies,

$$(S.5.26) \quad r_2(\mathbf{G}) \lesssim S_n^{g-1} E_n \max\{\tilde{k}^{1/2}, \log^{1/2} n\}$$

with probability at least $1 - 2n^{-6}$ when $n \rightarrow +\infty$. Substituting Eq. (S.5.25) and Eq. (S.5.26) into Eq. (S.5.24) yields the desired result. Note when $g < 1 + (2\beta_\Delta)^{-1}$, Eq. (3.2) and Eq. (S.5.25) similarly yields a weaker bound in Eq. (3.5). \square

S.5.6. *Proof of Corollary 3.1.* Let $\mathbf{W}_n \in \mathbb{O}_k$ minimizes $\|\hat{\mathbf{U}}_g - \mathbf{U} \mathbf{W}_n\|$ over all $k \times k$ orthogonal matrices. Then $\{\sigma_\ell\}_{\ell \leq k}$ are also the singular values of $\mathbf{W}_n^\top \mathbf{U}^\top \mathbf{M}$. We thus have

$$\begin{aligned} \|\hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{W}_n^\top \mathbf{U}^\top \mathbf{M}\| &\leq \|\hat{\mathbf{U}}_g^\top (\hat{\mathbf{M}} - \mathbf{M})\| + \|(\hat{\mathbf{U}}_g - \mathbf{U} \mathbf{W}_n)^\top \mathbf{M}\| \\ &\leq \|\mathbf{E}\| + d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \|\mathbf{M}\| \\ &\lesssim E_n + \vartheta \psi^g \frac{E_n}{S_n} S_n \lesssim \vartheta \psi^g E_n, \end{aligned}$$

with probability at least $1 - n^{-5}$, where the third inequality follows from Assumption 1 and Theorem 3.2. \square

S.5.7. *Proof of Theorem 3.3.* Let $\vartheta = \max\{1, \tilde{k}^{-1/2} \log^{1/2} n\}$. Now define the quantities

$$(S.5.27) \quad \rho_n^{(1)} := \frac{\sqrt{n} \psi^g}{\tilde{k}^{1/2}} \left(\frac{E_n}{S_n}\right)^g + \mathbb{I}(g \geq 2) \vartheta \psi^g \frac{E_n}{S_n},$$

$$(S.5.28) \quad \rho_n^{(2)} := (\log n)^{\delta g + 1/2} \psi^g \left(\frac{E_n}{S_n}\right)^g + \mathbb{I}(g \geq 2) (\log n)^\delta \sqrt{k} \vartheta \psi^g \frac{E_n}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Recall Theorem 3.2, Theorem S.3.3, and the error bounds in Lemma S.4.4 and Lemma S.4.5. In particular, with probability at least $1 - n^{-5}$, we have simultaneously,

$$\begin{aligned}\sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) &\lesssim \tilde{k}^{-1/2} \psi^g S_n^{-g}, \\ r_2(\mathbf{G}) &\lesssim E_n^g n^{1/2} + \mathbb{I}(g \geq 2) S_n^{g-1} E_n \tilde{k}^{1/2} \vartheta, \\ r_{2 \rightarrow \infty}(\mathbf{G}) &\lesssim \tilde{k}^{1/2} E_n^g (\log n)^{\delta g} (\log n)^{1/2} + \mathbb{I}(g \geq 2) k^{1/2} S_n^{g-1} E_n (\log n)^\delta \vartheta \tilde{k}^{1/2} \|\mathbf{U}\|_{2 \rightarrow \infty}.\end{aligned}$$

We therefore have, with probability at least $1 - n^{-5}$,

$$\frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \lesssim \rho_n^{(1)}, \quad \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \lesssim \rho_n^{(2)}.$$

Hence, by Theorem S.3.3, we have

$$(S.5.29) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \rho_n^{(1)} \|\mathbf{U}\|_{2 \rightarrow \infty} + \rho_n^{(2)},$$

with probability at least $1 - n^{-5}$,

Now consider $g \geq 1 + (2\beta_\Delta)^{-1}$. Recalling Eq. (S.5.23), we have

$$(S.5.30) \quad \rho_n^{(1)} \asymp \vartheta \psi^g \frac{E_n}{S_n}.$$

For $\rho_n^{(2)}$, recall that $\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim n^{-\beta_c/2}$ for some $\beta_c \in [0, 1]$. We then have

$$\begin{aligned}\frac{(\log n)^\delta \sqrt{k} \vartheta \psi^g \frac{E_n}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}}{(\log n)^{\delta g + 1/2} \psi^g \left(\frac{E_n}{S_n}\right)^g} &= k^{1/2} (\log n)^{\delta - \delta g - 1/2} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty} \frac{S_n^{g-1}}{E_n^{g-1}} \\ &\lesssim k^{1/2} (\log n)^{\delta - \delta g - 1/2} \vartheta n^{(g-1)\beta_\Delta - \beta_c/2} \\ &\lesssim \begin{cases} 1 & \text{if } g > 1 + (2\beta_\Delta)^{-1} \text{ or } \beta_c < 1 \\ (\log n)^{\delta - \delta g - 1/2} & \text{otherwise,} \end{cases}\end{aligned}$$

We therefore have

$$(S.5.31) \quad \rho_n^{(2)} \lesssim (\log n)^{\delta^*} \sqrt{k} \vartheta \psi^g \frac{E_n}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty} =: \tilde{\rho}_n^{(2)},$$

where $\delta^* = \delta$ if either $g > 1 + (2\beta_\Delta)^{-1}$ or $\beta_c < 1$, and $\delta^* = \delta g + 1/2$ otherwise.

Combining Eq. (S.5.30) and Eq. (S.5.31), we can now rewrite Eq. (S.5.29) as

$$(S.5.32) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \rho_n^{(1)} \|\mathbf{U}\|_{2 \rightarrow \infty} + \tilde{\rho}_n^{(2)}$$

We now note that $\frac{\tilde{\rho}_n^{(2)}}{\rho_n^{(1)} \|\mathbf{U}\|_{2 \rightarrow \infty}} \asymp k^{1/2} (\log n)^{\delta^*} \gtrsim 1$ and hence, by using Eq. (S.5.30) and Eq. (S.5.31), we can further simply Eq. (S.5.32) to

$$d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \tilde{\rho}_n^{(2)}.$$

Eq.(3.7) is thus established. \square

S.5.8. *Proof of Corollary 3.2.* We only consider the case where $\tilde{\mathbf{M}} = \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$ as

$$\begin{aligned}\left\| \frac{1}{2} (\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} + \hat{\mathbf{M}} \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top) - \mathbf{M} \right\|_{\max} &\leq \frac{1}{2} \left\| \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M} \right\|_{\max} + \frac{1}{2} \left\| \hat{\mathbf{M}} \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{M} \right\|_{\max} \\ &= \left\| \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M} \right\|_{\max}.\end{aligned}$$

Recall Proposition 3.1. Let $\bar{\mathbf{U}}_g = \mathbf{U}_g \mathbf{W}_U$ and $\mathbf{R}_\star = \hat{\mathbf{U}}_g - \bar{\mathbf{U}}_g = \mathbf{R}_0 + \mathbf{R}_1 + \mathbf{R}_2 + \mathbf{R}_3$ where \mathbf{W}_U and \mathbf{R}_0 through \mathbf{R}_3 are the matrices appearing in Eq. (S.5.10). We then have

$$\begin{aligned}
\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M} &= \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{U} \mathbf{U}^\top \mathbf{M} \\
&= \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{U}_g \mathbf{U}_g^\top \mathbf{M} \\
&= (\bar{\mathbf{U}}_g + \mathbf{R}_\star)(\bar{\mathbf{U}}_g + \mathbf{R}_\star)^\top \hat{\mathbf{M}} - \mathbf{U}_g \mathbf{U}_g^\top \mathbf{M} \\
&= \mathbf{U} \mathbf{U}^\top \mathbf{E} + \bar{\mathbf{U}}_g \mathbf{R}_\star^\top \hat{\mathbf{M}} + \mathbf{R}_\star \bar{\mathbf{U}}_g^\top \hat{\mathbf{M}} + \mathbf{R}_\star \mathbf{R}_\star^\top \hat{\mathbf{M}}
\end{aligned}
\tag{S.5.33}$$

We now bound the max norm of the four terms on the right-hand side of Eq. (S.5.33). Recall that from the proofs of Theorem S.3.3 and Theorem 3.3 we have

$$\|\mathbf{R}_\star\|_{2 \rightarrow \infty} \lesssim \Theta_n \frac{E_n}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. We now bound $\|\mathbf{R}_\star\|$ and $\|\mathbf{E} \mathbf{R}_\star\|_{2 \rightarrow \infty}$. Our bounds for these quantities depend on bounds for $\|\Delta_g\|$, $\|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V})\|$, etc., given in Theorem 3.1, Theorem 3.2 and Lemma S.4.4 through Lemma S.4.6. For conciseness we generally omit specific references to these bounds in the following derivations.

Bounding $\|\mathbf{R}_\star\|$. Using the expressions for \mathbf{R}_0 through \mathbf{R}_3 given in the proof of Theorem S.3.3, we have (recall that $g \geq 1 + (2\beta_\Delta)^{-1}$)

$$\begin{aligned}
\|\mathbf{R}_\star\| &\leq \|\mathbf{R}_0\| + \|\mathbf{R}_1\| + \|\mathbf{R}_2\| + \|\mathbf{R}_3\| \\
&\leq \|\Delta_g\| \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) + \sqrt{2} \|\Delta_g\| \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \cdot \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\| + \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \\
&\lesssim \vartheta \psi^g \frac{E_n}{S_n} + \vartheta^2 \psi^{2g} \frac{E_n^2}{S_n^2} + \vartheta^2 \psi^{2g} \frac{E_n^2}{S_n^2} \lesssim \vartheta \psi^g \frac{E_n}{S_n},
\end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$.

Bounding $\|\mathbf{E} \mathbf{R}_\star\|_{2 \rightarrow \infty}$. We first bound $\|\mathbf{E} \Delta_g\|_{2 \rightarrow \infty}$ where $\Delta_g = \mathbf{L}_1 + \mathbf{L}_2 + \mathbf{L}_3$ and the \mathbf{L}_k are as given in Eq. (S.5.1).

- Similar to Eq. (S.5.12), with probability at least $1 - \mathcal{O}(n^{-6})$, we have

$$\|\mathbf{E} \mathbf{L}_1\|_{2 \rightarrow \infty} \leq \|\mathbf{E}^{g+1} \mathbf{G}\|_{2 \rightarrow \infty} \lesssim (\log n)^{(g+1)\delta+1/2} \cdot \tilde{k}^{1/2} E_n^{g+1}.$$

- Recall Eq. (S.4.20). We then have

$$\mathbf{E} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U} = \mathbf{E}^{\ell+2} \mathbf{U} + \sum_{\xi=1}^{\ell} \mathbf{E}^\xi \mathbf{U} \mathbf{A} \mathbf{U}^\top \mathbf{M}^{\ell-\xi} \mathbf{E} \mathbf{U}.$$

Using a similar argument to that for deriving Eq. (S.4.21) through Eq. (S.4.24), we obtain

$$\begin{aligned}
\|\mathbf{E} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} &\lesssim \|\mathbf{E}^{\ell+2} \mathbf{U}\|_{2 \rightarrow \infty} + \sum_{\xi=1}^{\ell} \|\mathbf{E}^\xi \mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{A} \mathbf{U}^\top \mathbf{M}^{\ell-\xi} \mathbf{E} \mathbf{U}\| \\
&\lesssim (\log n)^\delta k^{1/2} E_n^2 S_n^\ell \|\mathbf{U}\|_{2 \rightarrow \infty}
\end{aligned}
\tag{S.5.34}$$

with probability at least $1 - \mathcal{O}(n^{-6})$.

- Let $\varphi_g(\mathbf{G}) = \|\mathbf{U}^\top \mathbf{G}\| + \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\|$. Eq. (S.5.34) together with a similar argument to that for deriving Eq. (S.5.14) yield

$$\begin{aligned}
\|\mathbf{E} \mathbf{L}_2\|_{2 \rightarrow \infty} &\lesssim \varphi_g(\mathbf{G}) \sum_{\xi=0}^{g-2} \left(\|\mathbf{E}^2 \mathbf{U}\|_{2 \rightarrow \infty} S_n^{g-1-\xi} + \sum_{\ell=1}^{g-2-\xi} \|\mathbf{E} \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} S_n^{g-1-\xi-\ell} \right) E_n^\xi \\
&\lesssim (\log n)^{2\delta} \vartheta k^{1/2} \cdot \tilde{k}^{1/2} E_n^2 S_n^{g-1} \|\mathbf{U}\|_{2 \rightarrow \infty}.
\end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$.

- Similar to Eq. (S.5.15), with probability at least $1 - \mathcal{O}(n^{-6})$, we have

$$\begin{aligned} \|\mathbf{E}\mathbf{L}_3\|_{2 \rightarrow \infty} &\leq \sum_{\xi=0}^{g-2} \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{A}\|^{g-1-\xi} \cdot \|\mathbf{U}^\top \mathbf{E}^{\xi+1} \mathbf{G}\| \\ &\lesssim (\log n)^\delta \vartheta \tilde{k}^{1/2} k^{1/2} E_n^2 S_n^{g-1} \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

Combining the above bounds we have

$$(S.5.35) \quad \|\mathbf{E}\mathbf{\Delta}_g\|_{2 \rightarrow \infty} = f_n \tilde{k}^{1/2} k^{1/2} E_n^2 S_n^{g-1} \|\mathbf{U}\|_{2 \rightarrow \infty},$$

with probability at least $1 - \mathcal{O}(n^{-6})$; here $f_n = (\log n)^{2\delta} \vartheta + \mathbb{I}\{g = 1 + (2\beta_\Delta)^{-1}\} (\log n)^{(g+1)\delta+1/2}$ is a poly-log factor.

Using the above bounds for $\mathbf{E}\mathbf{\Delta}_g$ in spectral and $2 \rightarrow \infty$ norms, we obtain

$$\begin{aligned} \|\mathbf{E}\mathbf{R}_\star\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}\mathbf{R}_0\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{R}_1\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{R}_2\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{R}_3\|_{2 \rightarrow \infty} \\ &\leq \|\mathbf{E}\mathbf{\Delta}_g\|_{2 \rightarrow \infty} \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) + \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Delta}_g\| \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \\ &\quad + (\|\mathbf{E}\mathbf{\Delta}_g\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Delta}_g\|) \cdot \frac{\sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \\ &\quad + \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \\ &= o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}), \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. In the above derivations, we had used the assumption that $E_n/S_n \lesssim n^{-\beta_\Delta}$ which converges to 0 faster than any poly-logarithmic factor in n .

Given the above bounds for $\|\mathbf{R}_\star\|$ and $\|\mathbf{E}\mathbf{R}_\star\|_{2 \rightarrow \infty}$, we are now ready to bound the max norm of each term in the right-hand side of Eq. (S.5.33). We will frequently use the inequality $\|\mathbf{A}\mathbf{B}\|_{\max} \leq \|\mathbf{A}\|_{2 \rightarrow \infty} \|\mathbf{B}^\top\|_{2 \rightarrow \infty}$.

1. **Bounding** $\|\mathbf{U}\mathbf{U}^\top \mathbf{E}\|_{\max}$. By Lemma S.4.6 we have

$$\|\mathbf{U}\mathbf{U}^\top \mathbf{E}\|_{\max} \leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim (\log n)^\delta k^{1/2} E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

with probability at least $1 - \mathcal{O}(n^{-6})$.

2. **Bounding** $\|\bar{\mathbf{U}}_g \mathbf{R}_\star^\top \hat{\mathbf{M}}\|_{\max}$. Note that $\bar{\mathbf{U}}_g \mathbf{R}_\star^\top \hat{\mathbf{M}} = \bar{\mathbf{U}}_g \mathbf{R}_\star^\top \mathbf{E} + \bar{\mathbf{U}}_g \mathbf{R}_\star^\top \mathbf{U} \mathbf{A} \mathbf{U}^\top$. We then have, with probability at least $1 - \mathcal{O}(n^{-6})$,

$$\|\bar{\mathbf{U}}_g \mathbf{R}_\star^\top \mathbf{E}\|_{\max} \leq \|\mathbf{E}\mathbf{R}_\star\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} = o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2)$$

$$\|\bar{\mathbf{U}}_g \mathbf{R}_\star^\top \mathbf{U} \mathbf{A} \mathbf{U}^\top\|_{\max} \leq \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{A}\| \|\mathbf{R}_\star\| \|\bar{\mathbf{U}}_g\|_{2 \rightarrow \infty} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

and thus $\|\bar{\mathbf{U}}_g \mathbf{R}_\star^\top \hat{\mathbf{M}}\|_{\max} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$.

3. **Bounding** $\|\mathbf{R}_\star \bar{\mathbf{U}}_g^\top \hat{\mathbf{M}}\|_{\max}$. Note that $\mathbf{R}_\star \bar{\mathbf{U}}_g^\top \hat{\mathbf{M}} = \mathbf{R}_\star \bar{\mathbf{U}}_g^\top \mathbf{E} + \mathbf{R}_\star \bar{\mathbf{U}}_g^\top \mathbf{U} \mathbf{A} \mathbf{U}^\top$. We then have, with probability at least $1 - \mathcal{O}(n^{-6})$,

$$\|\mathbf{R}_\star \bar{\mathbf{U}}_g^\top \mathbf{E}\|_{\max} \leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{R}_\star\|_{2 \rightarrow \infty} = o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2)$$

$$\|\mathbf{R}_\star \bar{\mathbf{U}}_g^\top \mathbf{U} \mathbf{A} \mathbf{U}^\top\|_{\max} \leq \|\mathbf{R}_\star\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{A}\| \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

and thus $\|\mathbf{R}_\star \bar{\mathbf{U}}_g^\top \hat{\mathbf{M}}\|_{\max} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$.

4. **Bounding** $\|\mathbf{R}_\star \mathbf{R}_\star^\top \hat{\mathbf{M}}\|_{\max}$. Decompose $\mathbf{R}_\star \mathbf{R}_\star^\top \hat{\mathbf{M}} = \mathbf{R}_\star \mathbf{R}_\star^\top \mathbf{E} + \mathbf{R}_\star \mathbf{R}_\star^\top \mathbf{M}$. Then with probability at least $1 - \mathcal{O}(n^{-6})$

$$\|\mathbf{R}_\star \mathbf{R}_\star^\top \mathbf{E}\|_{\max} \leq \|\mathbf{R}_\star\|_{2 \rightarrow \infty} \|\mathbf{E} \mathbf{R}_\star\|_{2 \rightarrow \infty} = o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2)$$

$$\|\mathbf{R}_\star \mathbf{R}_\star^\top \mathbf{M}\|_{\max} \leq \|\mathbf{R}_\star\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{A}\| \|\mathbf{R}_\star\| = o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2)$$

and thus $\|\mathbf{R}_\star \mathbf{R}_\star^\top \hat{\mathbf{M}}\|_{\max} = o(E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2)$.

Combining the above results yield $\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{M}\|_{\max} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$ as desired. \square

S.5.9. Proof of Theorem 3.4. We first make a few remarks. For ease of presentation, and as Theorem 3.4 is a limit result, we say that an event \mathcal{E} happens with high probability (whp) if \mathcal{E} happens with probability at least $1 - Cn^{-6}$. Here $C > 0$ is a finite constant that can change from line to line. In addition, the conditions assumed in Theorem 3.4 are at least as restrictive as those assumed in Theorems 3.1 through Theorem 3.3 and Lemmas S.4.4 through Lemma S.4.6. and hence the error bounds provided in Theorems 3.1 through Theorem 3.3 and Lemmas S.4.4 through Lemma S.4.6 still hold whp. For example, when $k, \psi \lesssim 1$ and $g \geq 2 + (2\beta_\Delta)^{-1}$ we have by Eq. (S.5.30) and Eq. (S.5.31) that, whp,

$$(S.5.36) \quad \begin{aligned} \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} &\lesssim \rho_n^{(1)} \lesssim \vartheta E_n / S_n, \\ \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} &\lesssim \tilde{\rho}_n^{(2)} \lesssim \vartheta (\log n)^\delta \|\mathbf{U}\|_{2 \rightarrow \infty} E_n / S_n \end{aligned}$$

where the constant δ is specified in Assumption 2.

Our main proof strategy is as follows. First, we derive an upper bound for the $\ell_{2 \rightarrow \infty}$ norm of $\mathbf{R} := \{\hat{\mathbf{U}}_g - \mathbf{U} \mathbf{W}_\mathbf{U}\} - \{\mathbf{E} \mathbf{U} \mathbf{A}_k^{-1} \mathbf{W}_\mathbf{U}\}$. In particular we show that the row-wise fluctuations of \mathbf{R} is negligible compared to the row-wise fluctuations for $\mathbf{E} \mathbf{U} \mathbf{A}_k^{-1} \mathbf{W}_\mathbf{U}$. The row-wise limiting distributions of $\hat{\mathbf{U}}_g - \mathbf{U} \mathbf{W}_\mathbf{U}$ are thus the same as those for $\mathbf{E} \mathbf{U} \mathbf{A}_k^{-1} \mathbf{W}_\mathbf{U}$. The proof is then completed by invoking limiting distribution for $\mathbf{E} \mathbf{U} \mathbf{A}_k^{-1} \mathbf{W}_\mathbf{U}$ in Assumption 3.

We now bound $\|\mathbf{R}\|_{2 \rightarrow \infty}$. Let $\mathbf{\Delta}_g = (\hat{\mathbf{M}}^g - \mathbf{M}^g) \mathbf{G}$. Recalling Eq. (S.5.10), we have

$$(S.5.37) \quad \hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_\mathbf{U} = (\mathbf{I} - \mathbf{U} \mathbf{U}^\top) \mathbf{\Delta}_g \mathbf{V}_g \mathbf{W}_\mathbf{V} \hat{\Sigma}_g^{-1} + \mathbf{R}_1 + \mathbf{R}_2 + \mathbf{R}_3$$

where $\mathbf{R}_2 = 0$ and $\mathbf{U}_g \mathbf{U}_g^\top = \mathbf{U} \mathbf{U}^\top$ by Proposition 3.1. Then by Eq. (S.5.18) we have, whp,

$$\begin{aligned} \|\mathbf{R}_1\|_{2 \rightarrow \infty} &\lesssim \frac{r_{2 \rightarrow \infty}(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} + \left[\frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \right]^2 \|\mathbf{U}\|_{2 \rightarrow \infty} \\ &\lesssim (\log n)^\delta \vartheta^2 (E_n / S_n)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

Similarly by Eq. (S.5.20) we have, whp,

$$\|\mathbf{R}_3\|_{2 \rightarrow \infty} \lesssim \left[\frac{r_2(\mathbf{G})}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \right]^2 \|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim \vartheta^2 (E_n / S_n)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

We further decompose the term $(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\Delta_g \mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}$ in Eq. (S.5.37) as follows. By splitting the second term in the expansion for $\hat{\mathbf{M}}^g - \mathbf{M}^g$ given in Lemma S.4.2's, we have

$$(S.5.38) \quad \begin{aligned} \Delta_g &= \left(\mathbf{E}^g + \mathbf{E}\mathbf{M}^{g-1} + \sum_{\xi=1}^{g-2} \mathbf{E}\mathbf{M}^{g-1-\xi} \mathbf{E}^\xi + \sum_{\xi=0}^{g-3} \sum_{\ell=1}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E}\mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \right) \mathbf{G} \\ &\quad + \sum_{\xi=0}^{g-2} \mathbf{M}^{g-1-\xi} \mathbf{E}^{\xi+1} \mathbf{G}. \end{aligned}$$

Let $\tilde{\mathbf{G}}_g := \mathbf{G}\mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}$. Reordering the terms in Eq. (S.5.38), we obtain

$$(S.5.39) \quad \begin{aligned} (\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\Delta_g \mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1} &= (\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{E}\mathbf{M}^{g-1}\tilde{\mathbf{G}}_g \\ &\quad + \underbrace{(\mathbf{I} - \mathbf{U}\mathbf{U}^\top) \left(\sum_{\xi=1}^{g-2} \mathbf{E}\mathbf{M}^{g-1-\xi} \mathbf{E}^\xi + \sum_{\xi=0}^{g-3} \sum_{\ell=1}^{g-2-\xi} \hat{\mathbf{M}}^\ell \mathbf{E}\mathbf{M}^{g-1-\xi-\ell} \mathbf{E}^\xi \right)}_{\mathbf{R}_4} \tilde{\mathbf{G}}_g \\ &\quad + \underbrace{(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{E}^g}_{\mathbf{R}_5} \tilde{\mathbf{G}}_g. \end{aligned}$$

Note that in the above derivations we had used the fact that, by the definition of \mathbf{M} in Eq. (2.1), we have $(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{M}^{g-1-\xi} = \mathbf{0}$ for all $\xi \leq g-2$. We now show that, ignoring logarithm terms, $\|\mathbf{R}_4\|_{2 \rightarrow \infty}$ and $\|\mathbf{R}_5\|_{2 \rightarrow \infty}$ can be bounded by $(E_n/S_n)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}$ whp.

Bounding $\|\mathbf{R}_4\|_{2 \rightarrow \infty}$. The following claim provides a high-probability bound for $\|\mathbf{R}_4\|_{2 \rightarrow \infty}$. For ease of exposition we defer its proof to Section S.7.16

CLAIM S.1. *Assume the setting of Theorem 3.4, we have whp,*

$$\|\mathbf{R}_4\|_{2 \rightarrow \infty} \lesssim (\log n)^{2\delta} \vartheta(E_n/S_n)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Bounding $\|\mathbf{R}_5\|_{2 \rightarrow \infty}$. For \mathbf{R}_5 , by Lemma S.4.1, we have

$$(S.5.39) \quad \begin{aligned} \|\mathbf{R}_5\|_{2 \rightarrow \infty} &\leq \|(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{E}^g \mathbf{G}\mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} \\ &\leq \|\mathbf{E}^g \mathbf{G}\mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} + \|\mathbf{U}\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} \\ &\leq \|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} \cdot \|\hat{\Sigma}_g^{-1}\| + \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\mathbf{V}_g \mathbf{W}_g \hat{\Sigma}_g^{-1}\|. \end{aligned}$$

For the first term on the right-hand side of Eq. (S.5.39), we have whp

$$(S.5.40) \quad \begin{aligned} \frac{\|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} \cdot \|\hat{\Sigma}_g^{-1}\|}{(E_n/S_n)^2 (\log n)^{\delta^{**}} \|\mathbf{U}\|_{2 \rightarrow \infty}} &\lesssim (E_n/S_n)^{g-2} \|\mathbf{U}\|_{2 \rightarrow \infty}^{-1} (\log n)^{\delta g + 1/2 - \delta^{**}} \\ &\lesssim n^{-(g-2)\beta_\Delta + 1/2} (\log n)^{\delta g + 1/2 - \delta^{**}} \\ &\lesssim 1 \quad (\text{By } g \geq 2 + (2\beta_\Delta)^{-1}), \end{aligned}$$

by Lemma S.4.6, where we define

$$(S.5.41) \quad \delta^{**} := \begin{cases} g\delta + 1/2 & g = 2 + (2\beta_\Delta)^{-1} \\ 0 & g > 2 + (2\beta_\Delta)^{-1}, \end{cases}$$

and the second inequality is by $\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim n^{-\beta_c/2}$, $E_n/S_n \lesssim n^{\beta_\Delta}$, and $g \geq 3$. For the second term in Eq. (S.5.39), we have whp

$$\begin{aligned}
\|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G} \mathbf{V}_g \mathbf{W}_V \hat{\Sigma}_g^{-1}\| &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| \cdot \|\hat{\Sigma}_g^{-1}\| \\
&\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} (E_n/S_n)^g \|\tilde{\mathbf{V}}_g^\top \mathbf{G}\| \\
&\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} (E_n/S_n)^g (\log n)^{1/2} \\
&\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} (E_n/S_n)^2,
\end{aligned} \tag{S.5.42}$$

where the second inequality is by Lemmas S.4.3 and S.4.5, the third inequality is from the bound $\|\tilde{\mathbf{V}}_g^\top \mathbf{G}\| \lesssim (\log n)^{1/2}$ whp by Lemma S.4.4, and the last inequality is because $g \geq 3$ and $E_n = o(S_n)$. As $\delta > 1$, combining Eq. (S.5.40) and Eq. (S.5.42) yields, whp,

$$\|\mathbf{R}_5\|_{2 \rightarrow \infty} \lesssim (E_n/S_n)^2 (\log n)^{\delta^{**}} \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Summarizing the above results, we have

$$\begin{aligned}
\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_U &= (\mathbf{I} - \mathbf{U} \mathbf{U}^\top) \mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \mathbf{W}_V \hat{\Sigma}_g^{-1} + \sum_{\ell \leq 5} \mathbf{R}_\ell \\
&= (\mathbf{I} - \mathbf{U} \mathbf{U}^\top) \mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \Sigma_g^{-1} \mathbf{W}_U \\
&\quad + \underbrace{(\mathbf{I} - \mathbf{U} \mathbf{U}^\top) \mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g (\mathbf{W}_V \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{W}_U)}_{\mathbf{R}_6} + \sum_{\ell \leq 5} \mathbf{R}_\ell
\end{aligned}$$

Now for any $g \geq 2$, we have $\mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{U}^\top \mathbf{M}^g = \mathbf{M}^{g-1}$. Recall that $\mathbf{U}_g \Sigma_g \mathbf{V}_g^\top$ is the singular value decomposition of $\mathbf{M}^g \mathbf{G}$ and furthermore, $\mathbf{U} = \mathbf{U}_g \mathbf{W}_*$ for some orthogonal matrix \mathbf{W}_* . We therefore have

$$\mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \Sigma_g^{-1} = \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{W}_*^\top.$$

Substituting the above relationship into Eq.(S.5.43), we obtain

$$\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_U = \mathbf{E} \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{W}_*^\top \mathbf{W}_U - \underbrace{\mathbf{U} \mathbf{U}^\top \mathbf{E} \mathbf{U} \mathbf{\Lambda}^{-1} \mathbf{W}_*^\top \mathbf{W}_U}_{\mathbf{R}_7} + \sum_{\ell=1}^6 \mathbf{R}_\ell \tag{S.5.43}$$

We now bound $\|\mathbf{R}_6\|_{2 \rightarrow \infty}$ and $\|\mathbf{R}_7\|_{2 \rightarrow \infty}$.

Bounding $\|\mathbf{R}_6\|_{2 \rightarrow \infty}$. We first state the following claim whose proof, for ease of exposition, is deferred to Section S.7.4.

CLAIM S.2. *Assume the setting of Theorem 3.2 where $\psi \lesssim 1$ and $g \geq 1 + (2\beta_\Delta)^{-1}$ is fixed. We then have whp,*

$$\|\mathbf{W}_V \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{W}_U\| \lesssim E_n S_n^{-g-1} \vartheta \tilde{k}^{-1/2}.$$

Let $\tilde{\mathbf{R}}_6 = \mathbf{W}_V \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{W}_U$. By Claim S.2 we have

$$\begin{aligned}
\|\mathbf{R}_6\|_{2 \rightarrow \infty} &= \|(\mathbf{I} - \mathbf{U} \mathbf{U}^\top) \mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \tilde{\mathbf{R}}_6\|_{2 \rightarrow \infty} \\
&\leq \|\mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \tilde{\mathbf{R}}_6\|_{2 \rightarrow \infty} + \|\mathbf{U} \mathbf{U}^\top \mathbf{E} \mathbf{M}^{g-1} \mathbf{G} \mathbf{V}_g \tilde{\mathbf{R}}_6\|_{2 \rightarrow \infty} \\
&\leq (\|\mathbf{E} \mathbf{U}\|_{2 \rightarrow \infty} + \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|) \|\mathbf{\Lambda}\|^{g-1} \cdot \|\mathbf{U}^\top \mathbf{G}\| \cdot \|\tilde{\mathbf{R}}_6\| \\
&\lesssim (E_n (\log n)^\delta + \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|) \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot E_n S_n^{-2} \vartheta^2 \\
&= \{(\log n)^\delta \vartheta^2 (E_n/S_n)^2 + \vartheta^2 \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\| E_n S_n^{-2}\} \|\mathbf{U}\|_{2 \rightarrow \infty},
\end{aligned} \tag{S.5.44}$$

whp, where the second inequality in the above display is by Lemma S.4.1 while the third inequality is by Assumption 1, Lemma S.4.5, and Claim S.2.

Bounding $\|\mathbf{R}_7\|_{2 \rightarrow \infty}$. By definition, we have, whp,

$$\begin{aligned}
 \|\mathbf{R}_7\|_{2 \rightarrow \infty} &= \|\mathbf{U}\mathbf{U}^\top \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{W}_\mathbf{U}\|_{2 \rightarrow \infty} \\
 (S.5.45) \quad &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \cdot \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \cdot \|\mathbf{\Lambda}^{-1}\| \quad (\text{by Lemma S.4.1}) \\
 &\lesssim S_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \cdot \|\mathbf{U}\|_{2 \rightarrow \infty}.
 \end{aligned}$$

Let $\mathbf{R} = \mathbf{R}_1 + \mathbf{R}_3 + \dots + \mathbf{R}_7$. As $E_n = \mathcal{O}(n^{1/2}\sigma_n)$, we have from the above bounds for $\{\mathbf{R}_\ell\}$ that

$$\begin{aligned}
 (S.5.46) \quad S_n\sigma_n^{-1}\|\mathbf{R}\|_{2 \rightarrow \infty} &\leq S_n\sigma_n^{-1} \sum_{\ell \leq 7} \|\mathbf{R}_\ell\|_{2 \rightarrow \infty} \\
 &\lesssim \left\{ S_n\sigma_n^{-1}(E_n/S_n)^2 \ell_{\text{clt}} + \sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \right\} \|\mathbf{U}\|_{2 \rightarrow \infty} \\
 &\lesssim \left\{ n\sigma_n S_n^{-1} \ell_{\text{clt}} + \sigma_n^{-1} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \right\} \|\mathbf{U}\|_{2 \rightarrow \infty} \longrightarrow 0,
 \end{aligned}$$

in probability, where the final convergence in Eq. (S.5.46) follows from Eq. (3.14). Here ℓ_{clt} is given in Eq. (3.15) and is an upper bound for logarithmic factors appearing in \mathbf{R}_1 through \mathbf{R}_7 . Recalling Eq. (S.5.43) and the relationship $\mathbf{U} = \mathbf{U}_g \mathbf{W}_*$, we obtain

$$\hat{\mathbf{U}}_g \mathbf{W}_\mathbf{U}^\top \mathbf{W}_* - \mathbf{U} = \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1} + \mathbf{R}\mathbf{W}_\mathbf{U}^\top \mathbf{W}_*.$$

Therefore, by Slutsky's theorem, we have

$$S_n\sigma_n^{-1} \left(\mathbf{W}_*^\top \mathbf{W}_\mathbf{U} [\hat{\mathbf{U}}_g]_i - [\mathbf{U}]_i \right) = S_n\sigma_n^{-1} [\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}]_i + S_n\sigma_n^{-1} [\mathbf{R}\mathbf{W}_\mathbf{U}^\top \mathbf{W}_*]_i \rightsquigarrow \mathcal{N}_k(\mathbf{0}, \mathbf{\Gamma}_i)$$

as $n \rightarrow \infty$. Indeed, by Eq. (S.5.46), $S_n\sigma_n^{-1} [\mathbf{R}\mathbf{W}_\mathbf{U}^\top \mathbf{W}_*]_i$ is negligible in the limit while the convergence of $S_n\sigma_n^{-1} [\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}]_i$ to $\mathcal{N}_k(\mathbf{0}, \mathbf{\Gamma}_i)$ is by Assumption 3. Note since \mathbf{U} and $\mathbf{U}_g \mathbf{W}_*$ are exactly equal, by the rotation invariant of Frobenius norm, we actually have $\mathbf{W}_*^\top \mathbf{W}_\mathbf{U} = \mathbf{W}_{\text{clt}}$, where \mathbf{W}_{clt} solves Procrustes problem between $\hat{\mathbf{U}}_g$ and \mathbf{U} such that

$$(S.5.47) \quad \underset{\mathbf{W} \in \mathbb{O}_k}{\operatorname{argmin}} \|\hat{\mathbf{U}}_g - \mathbf{U}\mathbf{W}\|_{\text{F}}.$$

□

S.6. Proofs for Section 4, Section 5, and Section S.2.

S.6.1. *Proof of Theorem 4.1.* Recall that $\tilde{k} = k$ and $\hat{\mathbf{U}}_g \in \mathbb{R}^{n \times k}$ is the left singular matrix of $\hat{\mathbf{M}}^g \mathbf{G} \in \mathbb{R}^{n \times k}$. For this proof we define a sparsity scaling parameter $\rho_n \asymp n^{\beta-1}$ such that $S_n = n\rho_n$ and $E_n \asymp (n\rho_n)^{1/2}$. By Lemma S.4.5, $\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G} \in \mathbb{R}^{k \times k}$ is invertible with probability at least $1 - 2n^{-6}$. Now suppose that $\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G} \in \mathbb{R}^{k \times k}$ is invertible. Then $\hat{\mathbf{M}}^g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \hat{\mathbf{M}}^g = \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top$ and hence

$$\begin{aligned}
 d_2(\hat{\mathbf{U}}_g, \mathbf{U})^2 &\geq \frac{1}{2k} \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U}\mathbf{U}^\top\|_{\text{F}}^2 \\
 &= \frac{1}{2k} \left\{ 2k - 2\operatorname{tr}(\mathbf{U}^\top \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \mathbf{U}) \right\} = 1 - \frac{1}{k} \operatorname{tr} \mathbf{U}^\top \hat{\mathbf{M}}^g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \hat{\mathbf{M}}^g \mathbf{U}.
 \end{aligned}$$

Let $\zeta_1 = \operatorname{tr} \mathbf{U}^\top \hat{\mathbf{M}}^g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \hat{\mathbf{M}}^g \mathbf{U}$ and $\mathbf{\Delta}_g = \hat{\mathbf{M}}^g - \mathbf{M}^g$. We then have,

$$\begin{aligned}
 (S.6.1) \quad \zeta_1 &= \operatorname{tr} \mathbf{G}^\top \mathbf{M}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} + \operatorname{tr} \mathbf{G}^\top (\mathbf{M}^g \mathbf{\Delta}_g + \mathbf{\Delta}_g \mathbf{M}^g) \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \\
 &\quad + \operatorname{tr} \mathbf{U}^\top \mathbf{\Delta}_g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \mathbf{\Delta}_g \mathbf{U}.
 \end{aligned}$$

In the above derivations we used the fact $\text{tr } \mathbf{A}\mathbf{B} = \text{tr } \mathbf{B}\mathbf{A}$ for any matrices \mathbf{A} and \mathbf{B} whose products $\mathbf{A}\mathbf{B}$ and $\mathbf{B}\mathbf{A}$ are well defined, and the relationship $\mathbf{M}^g \mathbf{U} \mathbf{U}^\top = \mathbf{M}^g = \mathbf{U} \mathbf{U}^\top \mathbf{M}^g$. Next note that

$$\begin{aligned} \text{tr } \mathbf{G}^\top \mathbf{M}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} &= k - \text{tr } \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \\ &\quad - \text{tr } \mathbf{G}^\top \{ \hat{\mathbf{M}}^{2g} - \mathbf{M}^{2g} - \mathbf{E}^{2g} \} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}, \end{aligned}$$

which together with Eq. (S.6.1) yields

$$(S.6.2) \quad \begin{aligned} \zeta_1 &= k - \text{tr } \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} + \text{tr } \mathbf{U}^\top \boldsymbol{\Delta}_g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \boldsymbol{\Delta}_g \mathbf{U} \\ &\quad + \text{tr } \mathbf{G}^\top \{ \mathbf{M}^{2g} - \hat{\mathbf{M}}^{2g} + \mathbf{E}^{2g} + \mathbf{M}^g \boldsymbol{\Delta}_g + \boldsymbol{\Delta}_g \mathbf{M}^g \} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}. \end{aligned}$$

Let $\tilde{\boldsymbol{\Delta}}_g = \boldsymbol{\Delta}_g - \mathbf{E}^g = \hat{\mathbf{M}}^g - \mathbf{M}^g - \mathbf{E}^g$. We then have the identity

$$\mathbf{M}^{2g} - \hat{\mathbf{M}}^{2g} + \mathbf{E}^{2g} + \mathbf{M}^g \boldsymbol{\Delta}_g + \boldsymbol{\Delta}_g \mathbf{M}^g = \mathbf{E}^{2g} - \boldsymbol{\Delta}_g^2 = -\tilde{\boldsymbol{\Delta}}_g^2 - \tilde{\boldsymbol{\Delta}}_g \mathbf{E}^g - \mathbf{E}^g \tilde{\boldsymbol{\Delta}}_g.$$

Define the quantities

$$\begin{aligned} \mathcal{T}_1 &= \text{tr } \mathbf{U}^\top \boldsymbol{\Delta}_g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \mathbf{G}^\top \boldsymbol{\Delta}_g \mathbf{U} \\ \mathcal{T}_2 &= \text{tr } \mathbf{G}^\top \tilde{\boldsymbol{\Delta}}_g^2 \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \\ \mathcal{T}_3 &= \text{tr } \mathbf{G}^\top \tilde{\boldsymbol{\Delta}}_g \mathbf{E}^g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} = \text{tr } \mathbf{G}^\top \mathbf{E}^g \tilde{\boldsymbol{\Delta}}_g \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}. \end{aligned}$$

We now bound \mathcal{T}_1 through \mathcal{T}_3 .

Bounding \mathcal{T}_1 . Write the SVD of $\mathbf{U}^\top \boldsymbol{\Delta}_g$ as $\mathbf{U}_\Delta \boldsymbol{\Sigma}_\Delta \mathbf{V}_\Delta^\top$ where $\mathbf{U}_\Delta \in \mathbb{R}^{k \times k}$, $\boldsymbol{\Sigma}_\Delta \in \mathbb{R}^{k \times k}$ and $\mathbf{V}_\Delta \in \mathbb{R}^{n \times k}$. As \mathbf{E} and \mathbf{G} are independent, there exists a constant $C'_u > 0$ such that, with probability at least $1 - n^{-6}$,

$$(S.6.3) \quad \|\mathbf{V}_\Delta^\top \mathbf{G}\| \leq C'_u \log^{1/2} n.$$

See the proof of Lemma S.4.4 for a derivation of the above bound. We thus have, with probability at least $1 - n^{-5}$,

$$\begin{aligned} \|\mathbf{U}^\top \boldsymbol{\Delta}_g \mathbf{G}\|_F &= \|\mathbf{U}_\Delta \boldsymbol{\Sigma}_\Delta \mathbf{V}_\Delta^\top \mathbf{G}\|_F \leq \|\mathbf{U}_\Delta \boldsymbol{\Sigma}_\Delta\|_F \|\mathbf{V}_\Delta^\top \mathbf{G}\| \\ &\leq \sqrt{k} \|\boldsymbol{\Delta}_g\| \cdot \|\mathbf{V}_\Delta^\top \mathbf{G}\| \lesssim (n\rho_n)^{g-1/2} \log^{1/2} n. \end{aligned}$$

where the last inequality in the above display used the following claim whose proof can be derived using the expansion for $\boldsymbol{\Delta}_g$ given in Lemma S.4.2 together with the high probability bound for $\|\mathbf{E}\|$ from Assumption 1.

CLAIM S.3. *Assume the setting of Theorem 4.1. Choose a fixed $g \geq 1$. Then with probability at least $1 - n^{-6}$ we have $\|\boldsymbol{\Delta}_g\| \lesssim (n\rho_n)^{g-1/2}$.*

Now $\psi \lesssim 1$, $S_n/E_n \asymp (n\rho_n)^{1/2}$, and $a_n = \lceil C\tilde{k}^{-1} \log n \rceil$. Therefore by Lemma S.4.5 we have, with probability at least $1 - n^{-6}$,

$$\|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\| = \sigma_k^{-2}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim (n\rho_n)^{-2g}.$$

We therefore have, with probability at least $1 - 2n^{-6}$,

$$|\mathcal{T}_1| \leq \|\mathbf{U}^\top \boldsymbol{\Delta}_g \mathbf{G}\|_F^2 \cdot \|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\| \lesssim (n\rho_n)^{-1} \log n.$$

Bounding \mathcal{T}_2 and \mathcal{T}_3 . Recall the proof of Theorem S.3.1, in particular the bounds for \mathbf{L}_2 and \mathbf{L}_3 in Eq. (S.5.6) and Eq. (S.5.7). Now $\tilde{\boldsymbol{\Delta}}_g \mathbf{G} = \mathbf{L}_2 + \mathbf{L}_3$ and hence

$$(S.6.4) \quad \|\tilde{\boldsymbol{\Delta}}_g \mathbf{G}\| \lesssim (n\rho_n)^{g-1/2} \left\{ \max_{\xi \leq 2g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| + \|\mathbf{U}^\top \mathbf{G}\| \right\} \lesssim (n\rho_n)^{g-1/2} \log^{1/2} n$$

with probability at least $1 - 2n^{-6}$. We therefore have

$$|\mathcal{T}_2| \leq \|\tilde{\Delta}_g \mathbf{G}\|_{\mathbb{F}}^2 \cdot \|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\| \leq k \|\tilde{\Delta}_g \mathbf{G}\|^2 \|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\| \lesssim (n\rho_n)^{-1} \log n$$

with probability at least $1 - 2n^{-6}$. Next let \mathbf{U}_* be the $n \times k$ matrix whose columns are the left singular vectors of $\mathbf{E}^g \tilde{\Delta}_g \mathbf{G}$. We then have, by Claim S.3 and Eq. (S.6.4),

$$\begin{aligned} |\mathcal{T}_3| &\leq \|\mathbf{G}^\top \mathbf{E}^g \tilde{\Delta}_g \mathbf{G}\|_{\mathbb{F}} \cdot \|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\|_{\mathbb{F}} \\ &\leq k \|\mathbf{G}^\top \mathbf{U}_*\| \cdot \|\mathbf{E}^g\| \cdot \|\tilde{\Delta}_g \mathbf{G}\| \cdot \|(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\| \lesssim (n\rho_n)^{-1} \log n, \end{aligned}$$

with probability at least $1 - 6n^{-6}$. Note that we had bound $\|\mathbf{G}^\top \mathbf{U}_*\| \lesssim \log^{1/2} n$ with probability at least $1 - n^{-6}$ similar to Eq. (S.6.3). Collecting the above bounds for \mathcal{T}_1 through \mathcal{T}_3 and recalling Eq.(S.6.2), we obtain

$$(S.6.5) \quad d_2(\hat{\mathbf{U}}_1, \mathbf{U})^2 \geq 1 - \frac{1}{k} \zeta_1 = \frac{1}{k} \text{tr} \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} + \frac{r'_n}{k},$$

where $|r'_n| = \mathcal{O}\{(n\rho_n)^{-1} \log n\} = \mathcal{O}(n^{-\beta} \log n)$ with probability at least $1 - n^{-5}$.

Our last step is to lower bound $\text{tr} \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}$. As $\mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G}$ and $(\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}$ are both positive semi-definite matrices, we have

$$\begin{aligned} \text{tr} \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} &\geq \frac{\text{tr} \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G}}{\|\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G}\|} \\ &\geq \frac{\min_{a \in [a_n]} \sum_{j=1}^k \{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)}}{\|\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G}\|}, \end{aligned}$$

where $\mathbf{g}_j^{(a)}$ is the j th column of \mathbf{G}_a .

Now by Theorem 3.1, Lemma S.4.4, and Assumption 1, we have

$$(S.6.6) \quad \begin{aligned} \|\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G}\| &= \|\hat{\mathbf{M}}^g \mathbf{G}\|^2 \leq 2\|(\hat{\mathbf{M}}^g - \mathbf{M}^g) \mathbf{G}\|^2 + 2\|\mathbf{U} \Lambda^g \mathbf{U}^\top \mathbf{G}\|^2 \\ &\lesssim n^{g+1} \rho_n^g + \mathbb{I}(g \geq 2) (n\rho_n)^{2g-1} \log^{1/2} n + (n\rho_n)^{2g} \|\mathbf{U}^\top \mathbf{G}\|^2, \end{aligned}$$

with probability at least $1 - n^{-5}$. Recall that $\mathbf{U}^\top \mathbf{G}_a$ for $a \in [a_n]$ are iid $k \times k$ standard Gaussian matrices. By the spectral norm bound of a standard Gaussian matrix [45], we have

$$(S.6.7) \quad \|\mathbf{U}^\top \mathbf{G}\|^2 \leq \max_{a \in [a_n]} \|\mathbf{U}^\top \mathbf{G}_a\|^2 \leq (2\sqrt{k} + C_1 \sqrt{\log \log n})^2 \lesssim \log \log n,$$

with probability at least $1 - (\log n)^5 \exp\{-(kC_1^2/2) \cdot (\log \log n)\} \rightarrow 1$ where C_1 is some large but finite constant. Together with Eq. (S.6.6), we have with probability approaching 1,

$$(S.6.8) \quad \|\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G}\| \lesssim n^{g+1} \rho_n^g + (n\rho_n)^{2g} \log \log n,$$

noting here $(n\rho_n)^{2g-1} \log^{1/2} n$ is negligible as $n\rho_n \asymp n^\beta$ for some $\beta > 0$.

In summary, with probability approaching 1, we have

$$(S.6.9) \quad \text{tr} \mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1} \gtrsim \frac{\min_{a \in [a_n]} \sum_{j=1}^k \{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)}}{n^{g+1} \rho_n^g + (n\rho_n)^{2g} \log \log n}.$$

We now bound $\{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)}$. Let $p_0 \in (0, 1)$ be an arbitrary constant and let $\mathcal{C}_{\mathcal{E},3}$ be a finite constant such that $\mathbb{P}(\text{tr} \mathbf{E}^{2g} \geq \mathcal{C}_{\mathcal{E},3} n^{g+1} \rho_n^g) \geq p_0$. The existence of $\mathcal{C}_{\mathcal{E},3}$ follows from the assumption $\mathbb{E} \text{tr} \mathbf{E}^{2g} \geq c_{\text{low}} n^{g+1} \rho_n^g$ and Markov's inequality. Next define the event

$$\mathcal{E} = \{\|\mathbf{E}^{2g}\|_{\mathbb{F}} \leq C_{\mathcal{E},1} n^{g+1/2} \rho_n^g\} \cap \{\|\mathbf{E}^{2g}\| \leq C_{\mathcal{E},2} (n\rho_n)^g\} \cap \{\text{tr} \mathbf{E}^{2g} \geq \mathcal{C}_{\mathcal{E},3} n^{g+1} \rho_n^g\}$$

where $C_{\mathcal{E},1}$ and $C_{\mathcal{E},2}$ are constants not depending on n . Assumption 1 implies that \mathcal{E} occurs with probability approaching p_0 . Now suppose that \mathcal{E} occurs, and let \mathbf{g}_* be an n -dimensional standard Gaussian random vector that is independent of \mathbf{E} . We then have

$$(S.6.10) \quad \mathbb{E}[\mathbf{g}_*^\top \mathbf{E}^{2g} \mathbf{g}_*] = \text{tr} \mathbb{E}[\mathbf{g}_* \mathbf{g}_*^\top \mathbf{E}^{2g}] = \text{tr}(\mathbb{E}[\mathbf{g}_* \mathbf{g}_*^\top] \mathbf{E}^{2g}) = \text{tr} \mathbf{E}^{2g} \gtrsim n^{g+1} \rho_n^g,$$

where the above expectations are with respect to \mathbf{g}_* and conditional on \mathbf{E} .

Now fix an $a \in [a_n]$ and a $j \in [k]$. As $\mathbf{g}_j^{(a)}$ is a random vector whose elements are iid standard normal independent of \mathbf{E} , the Hanson-Wright inequality for quadratic forms [61] implies

$$\begin{aligned} \mathbb{P}\left\{\{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)} \geq \text{tr} \mathbf{E}^{2g} - t \mid \mathcal{E}\right\} &\leq \exp\left\{-C_2 \min\left(\frac{t^2}{\|\mathbf{E}^{2g}\|_F^2}, \frac{t}{\|\mathbf{E}^{2g}\|}\right)\right\} \\ &\leq \exp\left\{-C_3 \min\left(\frac{t^2}{n^{2g+1} \rho_n^{2g}}, \frac{t}{(n \rho_n)^g}\right)\right\}, \end{aligned}$$

for any $t \geq 0$; here $C_2 \geq 0$ and $C_3 \geq 0$ are constants not depending on n . We thus have, by a union bound over all $a \in [a_n]$ and $j \in [k]$,

$$\mathbb{P}\left\{\min_{a \in [a_n], j \in [k]} \{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)} \geq \text{tr} \mathbf{E}^{2g} - t \mid \mathcal{E}\right\} \leq k a_n \exp\left\{-C_3 \min\left(\frac{t^2}{n^{2g+1} \rho_n^{2g}}, \frac{t}{(n \rho_n)^g}\right)\right\}.$$

Recall that k is fixed and $a_n = o(\log^5 n)$. We now choose $t = C_4 n^{g+1/2} \rho_n^g \log^{1/2} n$ for some sufficiently large $C_4 \geq 0$. Then, with high probability (assuming \mathcal{E} occurs)

$$(S.6.11) \quad \min_{a \in [a_n], j \in [k]} \{\mathbf{g}_j^{(a)}\}^\top \mathbf{E}^{2g} \mathbf{g}_j^{(a)} \geq \text{tr} \mathbf{E}^{2g} - C_4 n^{g+1/2} \rho_n^g \log^{1/2} n \gtrsim n^{g+1} \rho_n^g,$$

where the last inequality follows from Eq. (S.6.10). As \mathcal{E} occurs with probability approaching p_0 , we can now uncondition with respect to \mathcal{E} and still have the bound in Eq. (S.6.11) holding with probability approaching p_0 .

In summary, combining Eq. (S.6.5), Eq. (S.6.9) and Eq. (S.6.11), we have with probability approaching p_0 that

$$\begin{aligned} d_2(\hat{\mathbf{U}}_g, \mathbf{U})^2 &\gtrsim \text{tr}\{\mathbf{G}^\top \mathbf{E}^{2g} \mathbf{G} (\mathbf{G}^\top \hat{\mathbf{M}}^{2g} \mathbf{G})^{-1}\} - |r'_n| \\ &\gtrsim \frac{n^{g+1} \rho_n^g}{n^{g+1} \rho_n^g + (n \rho_n)^{2g} \log \log n} - |r'_n| \\ &\gtrsim \begin{cases} 1 & \text{if } g < \beta^{-1}, \\ (\log \log n)^{-1} \cdot n^{1-\beta g} & \text{if } \beta^{-1} \leq g < 1 + \beta^{-1}. \end{cases} \end{aligned}$$

Note that for the above derivations if $\beta^{-1} \leq g < 1 + \beta^{-1}$ then

$$\frac{n^{g+1} \rho_n^g}{n^{g+1} \rho_n^g + (n \rho_n)^{2g} \log \log n} \gtrsim \frac{n^{g+1} \rho_n^g}{(n \rho_n)^{2g} \log \log n} = \frac{n}{(n \rho_n)^g \log \log n} \asymp \frac{n^{1-g\beta}}{\log \log n} \succ n^{-\beta} \log n,$$

and therefore r'_n is negligible, while if $g < \beta^{-1}$ then $n^{g+1} \rho_n^g$ dominates $(n \rho_n)^{2g} \log \log n$ which makes the first term $\Theta(1)$; thus r'_n is also negligible. The lower bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ is thereby established. The lower bound for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ follows immediately from the lower bound for $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and the relationship $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \geq \frac{1}{\sqrt{n}} d_2(\hat{\mathbf{U}}_g, \mathbf{U})$. \square

S.6.2. *Proof of Theorem 4.2.* Let $\hat{\mathbf{M}} = \mathbf{A}$, $\mathbf{M} = \mathbf{P}$ and $\mathbf{E} = \mathbf{A} - \mathbf{P}$. We will verify that these matrices satisfy Assumption 2. Let n_ℓ be the number of nodes in the ℓ th community. By the law of large numbers, we have for all $\ell \in [K^*]$ that,

$$(S.6.12) \quad n_\ell/n \rightarrow \pi_\ell,$$

almost surely as $n \rightarrow \infty$. We now define the normalized membership matrix

$$\bar{\Theta} := \begin{pmatrix} n_1^{-1/2} \mathbf{1}_{n_1} & \mathbf{0} & \dots & \mathbf{0} \\ \mathbf{0} & n_2^{-1/2} \mathbf{1}_{n_2} & \dots & \mathbf{0} \\ \vdots & \vdots & \ddots & \vdots \\ \mathbf{0} & \mathbf{0} & \dots & n_K^{-1/2} \mathbf{1}_{n_K} \end{pmatrix},$$

and let $\bar{\mathbf{D}} := n^{-1/2} \text{diag}(\sqrt{n_1}, \dots, \sqrt{n_K})$. Without loss of generality, e.g., by reordering the rows and columns of \mathbf{P} so that the first n_1 vertices are assigned to community 1, the next n_2 vertices are assigned to community 2, and so on, \mathbf{P} can be represented as

$$\mathbf{P} = n\rho_n \bar{\Theta} \bar{\mathbf{D}} \bar{\mathbf{B}} \bar{\mathbf{D}} \bar{\Theta}^\top.$$

By Eq. (S.6.12) and an elementwise argument, $\|\bar{\mathbf{D}} \bar{\mathbf{B}} \bar{\mathbf{D}} - \mathbf{D}^* \mathbf{B} \mathbf{D}^*\| \rightarrow 0$ almost surely, where $\mathbf{D}^* = \text{diag}(\sqrt{\pi_1}, \dots, \sqrt{\pi_K})$. Weyl's inequality then implies

$$\max_{\ell \in [d]} |\lambda_\ell(\bar{\mathbf{D}} \bar{\mathbf{B}} \bar{\mathbf{D}}) - \lambda_\ell(\mathbf{D}^* \mathbf{B} \mathbf{D}^*)| \rightarrow 0,$$

where $d = \text{rk}(\mathbf{B})$. Now $\mathbf{D}^* \mathbf{B} \mathbf{D}^*$ is a fixed matrix not depending on n and $\lambda_\ell(\mathbf{D}^* \mathbf{B} \mathbf{D}^*) > 0$ for all $\ell \in [d]$. As $\bar{\Theta}$ is an orthogonal matrix, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n\rho_n} \lambda_\ell(\mathbf{P}) = \lim_{n \rightarrow \infty} \lambda_\ell(\bar{\Theta} \bar{\mathbf{D}} \bar{\mathbf{B}} \bar{\mathbf{D}} \bar{\Theta}^\top) = \lim_{n \rightarrow \infty} \lambda_\ell(\bar{\mathbf{D}} \bar{\mathbf{B}} \bar{\mathbf{D}}) = \lambda_\ell(\mathbf{D}^* \mathbf{B} \mathbf{D}^*) > 0.$$

Therefore $\mathbf{M} = \mathbf{P}$ with $\psi \lesssim 1$, $k^* = d$ and $S_n \asymp n\rho_n$, asymptotically almost surely. Next by the definition of \mathbf{P} we have $n \max_{ij} [\mathbf{P}]_{ij} \lesssim n\rho_n$ with $n\rho_n \succ \log n$. Hence, by Theorem 5.2 in [76], there exist a constant $C_E > 0$ such that, with probability at least $1 - n^{-6}$,

$$(S.6.13) \quad \|\mathbf{A} - \mathbf{P}\| \leq C_E \sqrt{n\rho_n},$$

Assumption 1 therefore holds with $E_n = C_E \sqrt{n\rho_n}$ and $\beta_\Delta = \beta/2$. Furthermore by Proposition 3.2 we have that $\mathbf{E} = \mathbf{A} - \mathbf{P}$ satisfies Assumption 2. See Example 1 for more details.

Given Assumptions 1–2, and with \bar{k}, a_n satisfying the conditions in Theorem 3.3, we have that if $g > \beta^{-1}$ then

$$(S.6.14) \quad d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \lesssim \rho_n^{(1)} \|\mathbf{U}\|_{2 \rightarrow \infty} + \rho_n^{(2)}$$

with probability at least $1 - n^{-5}$. Here $\rho_n^{(1)}$ and $\rho_n^{(2)}$ are as defined in Eq. (S.5.27) and Eq. (S.5.28). It is well-known that the coherence is bounded: $\|\mathbf{U}\|_{2 \rightarrow \infty} \lesssim n^{-1/2}$ a.s. (see e.g. the proof of Theorem 5.2 in [75]). Therefore, since $\psi \lesssim 1$, $n\rho_n \lesssim n^\beta$, $g > \beta^{-1}$ and $\beta \in (0, 1]$, we have

$$\begin{aligned} \rho_n^{(1)} \|\mathbf{U}\|_{2 \rightarrow \infty} &\lesssim n^{-g\beta/2} + n^{-\beta/2-1/2} \log^{1/2} n = o(n^{-1/2}), \\ \rho_n^{(2)} &\lesssim (\log n)^{\delta g + 1/2} n^{-g\beta/2} + n^{-\beta/2-1/2} (\log n)^{\delta + 1/2} = o(n^{-1/2}), \end{aligned}$$

which, when combined with Eq. (S.6.14), yields

$$(S.6.15) \quad n^{1/2} d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) \rightarrow 0$$

with probability at least $1 - n^{-5}$. Eq. (S.6.15) implies that, with high probability, the ℓ_2 distance between the i th and j th rows of $\hat{\mathbf{U}}$ when $\tau_i = \tau_j$ is always smaller than the ℓ_2

distance when $\tau_i \neq \tau_j$. Hence, clustering the rows of $\hat{\mathbf{U}}$ using either the K -means or K -medians algorithm will exactly recover $\boldsymbol{\tau}$ with probability at least $1 - n^{-5}$. See for example Theorem 2.6 in [82] and Theorem 5.2 in [75] for proofs of this claim for K -means and K -medians, respectively. \square

S.6.3. Proof of Theorem 5.1.

S.6.3.1. *Preliminary.* Let $\hat{\mathbf{M}} = p^{-1}\hat{\mathbf{T}}$ and $\mathbf{M} = \mathbb{E}[p^{-1}\hat{\mathbf{T}}] = \mathbb{E}[p^{-1}\{\mathcal{P}_\Omega(\mathbf{T} + \mathbf{N})\}] = \mathbf{T}$. Now define \mathbf{U} , $\hat{\mathbf{U}}$ and $\hat{\mathbf{U}}_g$ accordingly. We have two choices for $\hat{\mathbf{T}}_g$, namely, $\frac{1}{p}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$ and $\frac{1}{2p}(\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}} + \hat{\mathbf{T}}^\top\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top)$. Since $\|\frac{1}{2p}(\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}} + \hat{\mathbf{T}}^\top\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top) - \mathbf{T}\| \leq \|\frac{1}{p}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}} - \mathbf{T}\|$ for any norm $\|\cdot\|$, we choose $\hat{\mathbf{T}}_g = \frac{1}{p}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$ and bound $\hat{\mathbf{T}}_g - \mathbf{T}$ in Frobenius norm and max-norm. Finally we also derive the entrywise limiting distribution for $\hat{\mathbf{T}}_g - \mathbf{T}$.

S.6.3.2. *Assumption verification.* We first verify that \mathbf{M} and $\mathbf{E} = \hat{\mathbf{M}} - \mathbf{M}$ satisfy Assumption 1. As \mathbf{T} has bounded condition number, we have $S_n \asymp |\lambda_k(\mathbf{T})|$. Next write

$$(S.6.16) \quad \mathbf{E} = \hat{\mathbf{M}} - \mathbf{M} = \underbrace{\frac{1}{p}\{\mathcal{P}_\Omega(\mathbf{T}) - p\mathbf{T}\}}_{\mathbf{E}_1} + \underbrace{\frac{1}{p}\mathcal{P}_\Omega(\mathbf{N})}_{\mathbf{E}_2}.$$

Now \mathbf{E}_1 is a random symmetric matrix whose upper triangular entries are independent mean 0 random variables with

$$\frac{1}{\|\mathbf{T}\|_{\max}} \max_{(i,j) \in [n]^2} |p[\mathbf{E}_1]_{ij}| = \frac{1}{p\|\mathbf{T}\|_{\max}} \max_{(i,j) \in [n]^2} \left\{ (1-2p)|[\mathbf{T}]_{ij}|, p|[\mathbf{T}]_{ij}| \right\} \leq 1.$$

Furthermore we also have

$$(S.6.17) \quad \begin{aligned} \max_i \sum_{j=1}^n \frac{\mathbb{E}|p[\mathbf{E}_1]_{ij}|^2}{\|\mathbf{T}\|_{\max}^2} &= \max_i \sum_{j=1}^n \frac{p\{[\mathbf{T}]_{ij} - p[\mathbf{T}]_{ij}\}^2 + (1-p)\{-p[\mathbf{T}]_{ij}\}^2}{\|\mathbf{T}\|_{\max}^2} \\ &\leq \max_i \sum_{j=1}^n [p(1-p)^2 + (1-p)p^2] \leq np. \end{aligned}$$

By Remark 3.13 in [11], there exists a universal constant $c > 0$ such that for $t = (np)^{1/2}$,

$$\mathbb{P}\left[\frac{p}{\|\mathbf{T}\|_{\max}} \|\mathbf{E}_1\| \geq 4(np)^{1/2} + t\right] \leq n \exp(-t^2/c),$$

which immediately implies, $\|\mathbf{E}_1\| \lesssim (n/p)^{1/2} \|\mathbf{T}\|_{\max}$ with probability at least $1 - \frac{1}{2}n^{-6}$. On the other hand, $p^{-1}\|\mathbf{E}_2\| \leq \|\mathcal{P}_\Omega(\mathbf{N})\| \lesssim \sigma(n/p)^{1/2}$, with probability at least $1 - \frac{1}{2}n^{-6}$, see e.g., Lemma 13 in [3]. Eq. (S.6.16) therefore implies

$$(S.6.18) \quad \|\mathbf{E}\| \lesssim (\sigma + \|\mathbf{T}\|_{\max}) \sqrt{\frac{n}{p}}$$

with probability at least $1 - n^{-6}$. We can thus choose $E_n = C_{\text{MC}}(\sigma + \|\mathbf{T}\|_{\max})(n/p)^{1/2}$ for some finite constant $C_{\text{MC}} > 2$. In summary we have

$$(S.6.19) \quad \frac{E_n}{S_n} \asymp \frac{1}{\sqrt{np}} \frac{n(\sigma + \|\mathbf{T}\|_{\max})}{|\lambda_k(\mathbf{T})|} \lesssim n^{-\beta_\Delta}$$

where the last inequality follows from the assumption in Eq. (5.2).

We next verify Assumption 2. Recall Proposition 3.2. Now let $A_0 = 10$ and

$$q_n = \sqrt{2np} \left[\{(\log n)^{10 \log \log n}\}^{-1/2} \wedge 1 \right]$$

Note that we require $q_n = \omega(1)$ in the statement of Proposition 3.2 and this is satisfied when $np \gtrsim n^\beta$ for some arbitrary constant $\beta > 0$ as assumed in Eq. (5.2). We then have, (S.6.20)

$$\begin{aligned} q_n^{g'-2} \mathbb{E} |[\mathbf{E}]_{ij}/E_n|^{g'} &\leq 2^{g'-1} \{q_n^{g'-2} \mathbb{E} |[\mathbf{E}_1]_{ij}/E_n|^{g'} + q_n^{g'-2} \mathbb{E} |[\mathbf{E}_2]_{ij}/E_n|^{g'}\} \\ &\leq 2^{g'-1} \left\{ (q_n^{(1)})^{g'-2} \mathbb{E} |[\mathbf{E}_1]_{ij}/E_n^{(1)}|^{g'} + (q_n^{(2)})^{g'-2} \mathbb{E} |[\mathbf{E}_2]_{ij}/E_n^{(2)}|^{g'} \right\} \end{aligned}$$

where $q_n^{(1)}, q_n^{(2)}, E_n^{(1)}$, and $E_n^{(2)}$ are defined as follows

$$\begin{aligned} q_n^{(1)} &= \sqrt{np} \geq q_n, & q_n^{(2)} &= \sqrt{np} \{(\log n)^{10 \log \log n} / 2\}^{-1/2} \geq q_n, \\ E_n^{(1)} &= C_{\text{MC}} \|\mathbf{T}\|_{\max} \sqrt{n/p} \leq E_n, & E_n^{(2)} &= C_{\text{MC}} \sigma \sqrt{n/p} \leq E_n. \end{aligned}$$

We now bound the two terms on the right-hand side of Eq. (S.6.20).

- Let $\bar{\mathbf{E}}_1 = \mathbf{E}_1 / (C_{\text{MC}} \|\mathbf{T}\|_{\max} / p)$. Each entry of $\bar{\mathbf{E}}_1$ takes on two distinct values and are uniformly bounded from above by 1. There thus exists some constant $C_1^{(1)} > 0$ such that

$$(q_n^{(1)})^{g'-2} \mathbb{E} |[\mathbf{E}_1]_{ij}/E_n^{(1)}|^{g'} = (\sqrt{np})^{g'-2} \mathbb{E} |[\bar{\mathbf{E}}_1]_{ij}/\sqrt{np}|^{g'} \leq C_1^{(1)} n^{-1}$$

for all $g' \geq 2$. See Example 1 for more details.

- Now $\mathbf{E}_2 = \mathcal{P}_\Omega(\bar{\mathbf{N}})$ where $\bar{\mathbf{N}} = p^{-1}\mathbf{N}$ is symmetric with iid $\mathcal{N}(0, p^{-2}\sigma^2)$ upper triangular entries and the random sampling $\mathcal{P}_\Omega(\cdot)$ is independent of $\bar{\mathbf{N}}$. We therefore have

$$\mathbb{E} |[\mathbf{E}_2]_{ij}/E_n^{(2)}|^{g'} = p \mathbb{E} |[\bar{\mathbf{N}}]_{ij}/E_n^{(2)}|^{g'} + (1-p) \cdot 0 = p \mathbb{E} |[\bar{\mathbf{N}}]_{ij}/E_n^{(2)}|^{g'}.$$

Recall Example 2 and note that $C_{\text{MC}} \geq 2$. There thus exists a constant $C_1^{(2)} > 0$ such that for any $g' \in [2, (\log n)^{10 \log \log n}]$ we have

$$(q_n^{(2)})^{g'-2} \mathbb{E} |[\mathbf{E}_2]_{ij}/E_n^{(2)}|^{g'} = (q_n^{(3)})^{g'-2} \mathbb{E} |[\bar{\mathbf{N}}]_{ij}/E_n^{(3)}|^{g'} \leq (C_1^{(2)})^{g'} n^{-1}$$

where we had defined $q_n^{(3)} := n^{1/2} \{(\log n)^{10 \log \log n} / 2\}^{-1/2}$ and $E_n^{(3)} := C_{\text{MC}} \sqrt{n} \sigma / p$.

In summary if $g' \in [2, (\log n)^{10 \log \log n}]$ then

$$q_n^{g'-2} \mathbb{E} |[\mathbf{E}]_{ij}/E_n|^{g'} \leq 2^{g'-1} \{(C_1^{(1)}) n^{-1} + (C_1^{(2)})^{g'} n^{-1}\} \leq C_1^{g'} n^{-1}$$

for all $(i, j) \in [n]^2$; here $C_1 > 0$ is a finite constant. As $\mathbb{E}\mathbf{E} = \mathbf{0}$, \mathbf{E} satisfies all conditions in Proposition 3.2 and hence also satisfies Assumption 2.

S.6.3.3. Bounding $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ and $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$. Recall Eq. (S.6.19). Let $g \geq 1 + (2\beta_\Delta)^{-1}$. We then have, by Theorem 3.2 and Theorem 3.3, that

$$\begin{aligned} d_2(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \vartheta \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{1}{\sqrt{np}}, \\ d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) &\lesssim \vartheta \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{(\log n)^{\delta^*}}{\sqrt{np}} \|\mathbf{U}\|_{2 \rightarrow \infty} \end{aligned} \tag{S.6.21}$$

with probability at least $1 - n^{-5}$. Here δ^* is the constant defined in Theorem 3.3.

S.6.3.4. *Bounding $\|\hat{\mathbf{T}}_g - \mathbf{T}\|_F$.* Recall that we defined $\hat{\mathbf{T}}_g = p^{-1}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$. Now by Eq. (S.6.18), Eq. (S.6.19), and the assumption $\|\mathbf{T}\| \asymp \lambda_k(\mathbf{T})$ (as \mathbf{T} has bounded condition number), we have with probability at least $1 - n^{-6}$ that

$$(S.6.22) \quad \|\hat{\mathbf{T}}\| \leq \|\mathbf{E}\| + \|\mathbf{T}\| \lesssim (\sigma + \|\mathbf{T}\|_{\max})\sqrt{\frac{n}{p}} + \|\mathbf{T}\| \asymp \|\mathbf{T}\|.$$

We now bound $\|\hat{\mathbf{T}}_g - \mathbf{T}\|_F$ for some fixed $g \geq 1 + 2(\beta_\Delta)^{-1}$. As $\mathbf{T} = \mathbf{U}\mathbf{U}^\top\mathbf{T}$ we have

$$(S.6.23) \quad \hat{\mathbf{T}}_g - \mathbf{T} = \underbrace{(\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top - \mathbf{U}\mathbf{U}^\top)\hat{\mathbf{T}}}_{\mathbf{D}_1} + \underbrace{\mathbf{U}\mathbf{U}^\top(\hat{\mathbf{T}} - \mathbf{T})}_{\mathbf{D}_2}$$

Since $\hat{\mathbf{U}}_g, \mathbf{U} \in \mathbb{R}^{n \times k}$, we have $\text{rk}(\mathbf{D}_1) \leq 2k$ and $\text{rk}(\mathbf{D}_2) \leq k$. Eq. (S.6.21) and Eq. (S.6.22) therefore imply

$$\|\mathbf{D}_1\|_F \leq \sqrt{2k}\|\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top - \mathbf{U}\mathbf{U}^\top\| \|\hat{\mathbf{T}}\| \leq 2\sqrt{2k}d_2(\hat{\mathbf{U}}_g, \mathbf{U}) \times \|\hat{\mathbf{T}}\| \lesssim \vartheta(\|\mathbf{T}\|_{\max} + \sigma)\sqrt{\frac{n}{p}}$$

with probability at least $1 - n^{-5}$. Similarly, by Eq. (S.6.18),

$$\|\mathbf{D}_2\|_F = \|\mathbf{U}\mathbf{U}^\top(\hat{\mathbf{T}} - \mathbf{T})\|_F \leq \sqrt{k}\|\mathbf{E}\| \lesssim (\|\mathbf{T}\|_{\max} + \sigma)\sqrt{\frac{n}{p}}$$

with probability at least $1 - n^{-5}$. Combining the above results with Eq. (S.6.23) we obtain

$$\frac{1}{n}\|\hat{\mathbf{T}}_g - \mathbf{T}\|_F \lesssim \vartheta(\|\mathbf{T}\|_{\max} + \sigma)\frac{1}{\sqrt{np}}.$$

with probability at least $1 - n^{-5}$.

S.6.3.5. *Bounding $\|\hat{\mathbf{T}}_g - \mathbf{T}\|_{\max}$.* Fix a $g \geq 1 + (2\beta_\Delta)^{-1}$. Then under the condition of Theorem 5.1 we have by Corollary 3.2 that

$$(S.6.24) \quad \|\hat{\mathbf{T}}_g - \mathbf{T}\|_{\max} \lesssim \Theta_n E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \vartheta(\log n)^{\delta^*} (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2,$$

with probability at least $1 - n^{-5}$.

S.6.3.6. *Entrywise limiting distribution and confidence interval.* Fix a $g \geq 2 + (2\beta_\Delta)^{-1}$. We will derive the limiting distribution of $[\hat{\mathbf{T}}_g]_{ij}$ for any $(i, j) \in [n]^2$ where $\hat{\mathbf{T}}_g = p^{-1}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}}$; the same result also holds for $\hat{\mathbf{T}}_g = (2p)^{-1}(\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top\hat{\mathbf{T}} + \hat{\mathbf{T}}\hat{\mathbf{U}}_g\hat{\mathbf{U}}_g^\top)$. For ease of exposition we say that an event \mathcal{E} happens with high probability (whp) if \mathcal{E} happens with probability at least $1 - Cn^{-6}$. Here $C > 0$ is an arbitrary constant that can change from line to line.

Compared with the entrywise concentration bound in Eq. (S.6.24), the entrywise limiting distribution requires substantially more detailed analysis. For clarity we first state a general entrywise CLT result which does not depend on the matrix completion setting and then apply this result to derive the normal approximations in Eq. (5.5) and Eq. (5.7).

THEOREM S.6.1. *Suppose Assumptions 1–2 hold with fixed k^* and bounded ψ . Let $\tilde{\mathbf{M}}$ be generated via Algorithm 1 with $\tilde{k} \geq k$, $a_n \geq \lceil C\tilde{k}^{-1} \log n \rceil$ where the universal constant C is given in Remark 3.3, and $g \geq 2 + (2\beta_\Delta)^{-1}$. Define the random variable,*

$$\alpha_n := \ell_{\text{clt}}\left(\frac{E_n}{S_n}\right) E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \frac{\|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|^2}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}^2,$$

where ℓ_{clt} is defined in Theorem 3.4. Fix an $(i, j) \in [n]^2$ and suppose that there exists a quantity $v_{ij} > 0$ depending possibly on n such that

$$\frac{[\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}}{\sqrt{v_{ij}}} \rightsquigarrow \mathcal{N}(0, 1).$$

Then, provided that $(v_{ij})^{-1/2} \alpha_n \rightarrow 0$ in probability, we also have

$$\frac{[\tilde{\mathbf{M}} - \mathbf{M}]_{ij}}{\sqrt{v_{ij}}} \rightsquigarrow \mathcal{N}(0, 1).$$

The proof of Theorem S.6.1 is deferred to Section S.6.5. To apply Theorem S.6.1, we first derive an upper bound for $\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|$ (see Claim S.4). We next compute $v_{ij} = \text{Var}([\mathbf{U}\mathbf{U}^\top \tilde{\mathbf{E}}]_{ij} + [\tilde{\mathbf{E}}\mathbf{U}\mathbf{U}^\top]_{ij})$ where $\tilde{\mathbf{E}}$ is obtained by truncating the entries of \mathbf{E} . Using this v_{ij} we show that $\{[\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}\} / \sqrt{v_{ij}} \rightsquigarrow \mathcal{N}(0, 1)$ and $\alpha_n / \sqrt{v_{ij}} \rightarrow 0$ in probability. Theorem S.6.1 then implies $\{[\hat{\mathbf{T}}_g - \mathbf{T}]_{ij}[\mathbf{T}]_{ij}\} / \sqrt{v_{ij}} \rightsquigarrow \mathcal{N}(0, 1)$. Finally we show that the quantities v_{ij}^* and \hat{v}_{ij} defined in Eq.(5.6) satisfy $v_{ij}^*/v_{ij} \rightarrow 1$ and $\hat{v}_{ij}/v_{ij} \rightarrow 1$ in probability. Our derivation of the limiting distribution for $\{[\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}\} / \sqrt{v_{ij}}$ follows the same ideas as that presented in [28] but our derivations are for the RSVD estimates $\hat{\mathbf{U}}_g$ while the derivations in [28] are for $\hat{\mathbf{U}}$. Nevertheless, for completeness we will present most of the technical details with references to [28] when appropriate.

Before getting into the proof note that by Eq. (5.2) we have

$$(S.6.25) \quad C_n^{(1)} := \ell_{\text{clt}} \frac{n^{1/2}(\|\mathbf{T}\|_{\max} + \sigma)}{p^{1/2}|\lambda_k(\mathbf{T})|} \rightarrow 0,$$

$$(S.6.26) \quad C_n^{(2)} := \vartheta(\log n)^{\delta^* + 1/2} \frac{n^{1/2}(\|\mathbf{T}\|_{\max} + \sigma)}{p^{1/2}|\lambda_k(\mathbf{T})|} \rightarrow 0.$$

where both of the above convergence are in probability.

We first consider a truncated version of $\tilde{\mathbf{T}}$. Recall $[\mathbf{N}]_{ij} = \eta_{ij}$ where $\eta_{ij} \sim \mathcal{N}(0, \sigma^2)$. Let $\tilde{\eta}_{ij} := \eta_{ij} \mathbb{I}\{|\eta_{ij}| \leq 5\sigma\sqrt{\log n}\}$ and let $\tilde{\mathbf{N}}$ be the matrix whose entries are the $\tilde{\eta}_{ij}$. Now define

$$(S.6.27) \quad \tilde{\mathbf{T}} := \mathcal{P}_\Omega(\mathbf{T} + \tilde{\mathbf{N}}) \text{ and } \tilde{\mathbf{E}} = \tilde{\mathbf{T}}/p - \mathbf{T}.$$

Now by the tail bounds for Gaussian distribution, we have

$$\mathbb{P}\left(\max_{ij} |\eta_{ij}| \leq 5\sigma\sqrt{\log n}\right) \geq 1 - n^2 \mathbb{P}(|\eta_{ij}| > 5\sigma\sqrt{\log n}) \geq 1 - n^{-10}.$$

See [28, Section 3.2.3] for more details. We thus have $\tilde{\mathbf{E}} = \mathbf{E}$ and $\tilde{\mathbf{T}} = \hat{\mathbf{T}}$ whp. The following result uses $\tilde{\mathbf{E}}$ to bound $\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|$; see Section S.7.6 for a proof.

CLAIM S.4. *Assume the setting of Theorem 5.1. We then have whp that*

$$\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \lesssim \sqrt{\frac{\log n}{p}} (\|\mathbf{T}\|_{\max} + \sigma) + \frac{\log n}{p} (\|\mathbf{T}\|_{\max} + \sigma\sqrt{\log n}) \|\mathbf{U}\|_{2 \rightarrow \infty}^2.$$

Claim S.4 implies

$$\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \lesssim \sqrt{\frac{\log n}{n}} E_n + \frac{\log^{3/2} n}{\sqrt{np}} E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \frac{\log^{3/2} n}{\sqrt{np}} E_n,$$

whp and hence $\|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|/S_n = (\log n)^{3/2} (np)^{-1/2} (E_n/S_n) \rightarrow 0$ in probability. Recalling the definition of α_n in Theorem S.6.1 we obtain

$$(S.6.28) \quad \begin{aligned} \alpha_n &\lesssim \ell_{\text{clt}} \frac{n(\|\mathbf{T}\|_{\max}^2 + \sigma^2)}{p\lambda_k(\mathbf{T})} \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \sqrt{\frac{\log n}{p}} (\|\mathbf{T}\|_{\max} + \sigma) \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \\ &\quad + \frac{\log n}{p} (\|\mathbf{T}\|_{\max} + \sigma \sqrt{\log n}) \|\mathbf{U}\|_{2 \rightarrow \infty}^4. \end{aligned}$$

with high probability.

Let v_{ij} be the variance of $[\mathbf{U} \mathbf{U}^\top \tilde{\mathbf{E}}]_{ij} + [\tilde{\mathbf{E}} \mathbf{U} \mathbf{U}^\top]_{ij}$. We first bound v_{ij} from below using a similar argument to the proof of Theorem 4.12 in [28]. Using this bound we then show that α_n/v_{ij} is negligible whp. Denote $\zeta_{k\ell} = [\mathbf{U} \mathbf{U}^\top]_{k\ell}$ and note that

$$\zeta_{k\ell}^2 = ([\mathbf{U}]_k^\top [\mathbf{U}]_\ell)^2 \leq \|[\mathbf{U}]_k\|^2 \times \|[\mathbf{U}]_\ell\|^2$$

for all $(k, \ell) \in [n^2]$. Following the proof of Lemma 4.19 in [28] we have

$$(S.6.29) \quad \begin{aligned} v_{ij} &= \sum_{\ell \neq j} \mathbb{E}([\tilde{\mathbf{E}}]_{i\ell}^2) \zeta_{k\ell}^2 + \sum_{\ell \neq i} \mathbb{E}([\tilde{\mathbf{E}}]_{\ell j}^2) \zeta_{i\ell}^2 + \mathbb{E}([\tilde{\mathbf{E}}]_{ij}^2) \{\zeta_{ii} + \zeta_{jj}\}^2 \\ &\geq \left\{ \min_{(k, \ell) \in [n^2]} \mathbb{E}([\tilde{\mathbf{E}}]_{k\ell}^2) \right\} \left\{ \|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2 \right\}. \end{aligned}$$

As the distribution of η_{ij} is symmetric around 0, the distribution of $\tilde{\eta}_{ij}$ is also symmetric around 0 and $\mathbb{E}[\tilde{\eta}_{ij}] = 0$. We therefore have

$$(S.6.30) \quad \begin{aligned} \mathbb{E}([\tilde{\mathbf{E}}]_{ij}^2) &= p^{-1} \mathbb{E}\{([\mathbf{T}]_{ij} - p[\mathbf{T}]_{ij} + [\tilde{\mathbf{N}}]_{ij})^2\} + (1-p)[\mathbf{T}]_{ij}^2 \\ &= \frac{1-p}{p} [\mathbf{T}]_{ij}^2 + \frac{1}{p} \mathbb{E}[\tilde{\eta}_{ij}^2] \geq \frac{1-p}{p} \min_{k\ell} |T_{k\ell}|^2 + \frac{1}{2p} \sigma^2, \end{aligned}$$

provided that n is sufficiently large. The inequality in the above display is derived as follows. Let $t_n = 5\sqrt{\log n}$. By the Cauchy–Schwarz inequality we have

$$(S.6.31) \quad \frac{1}{\sigma^2} \mathbb{E}[\eta_{ij}^2 \mathbb{I}(|\eta_{ij}| > \sigma t_n)] \leq \left(\mathbb{E}[(\eta_{ij}/\sigma)^4] \mathbb{P}[|\eta_{ij}/\sigma| > t_n] \right)^{1/2} = o(1),$$

and hence

$$\mathbb{E}([\tilde{\mathbf{N}}]_{ij}^2/\sigma^2) = \sigma^{-2} \mathbb{E}[\eta_{ij}^2 \{1 - \mathbb{I}(|\eta_{ij}| > \sigma t_n)\}] = 1 + o(1).$$

Under the conditions of Theorem 5.1 we have that p is bounded away from 1 and $\min_{k\ell} |T_{k\ell}| \asymp \|\mathbf{T}\|_{\max}$, and thus

$$(S.6.32) \quad \sqrt{v_{ij}} \gtrsim \left(\frac{\|\mathbf{T}\|_{\max}}{\sqrt{p}} + \frac{\sigma}{\sqrt{p}} \right) \sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}.$$

We therefore have, by Eq. (S.6.28), that

$$(S.6.33) \quad \begin{aligned} \frac{\alpha_n}{\sqrt{v_{ij}}} &\lesssim \ell_{\text{clt}} \left\{ \frac{n\|\mathbf{T}\|_{\max}}{\sqrt{p}\lambda_k(\mathbf{T})} + \frac{n\sigma}{\sqrt{p}\lambda_k(\mathbf{T})} \right\} \frac{\|\mathbf{U}\|_{2 \rightarrow \infty}^2}{\sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}} \\ &\quad + \frac{(\log n)^{1/2} \|\mathbf{U}\|_{2 \rightarrow \infty}^2}{\sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}} + \frac{(\log n)^{3/2} \|\mathbf{U}\|_{2 \rightarrow \infty}^4}{\sqrt{p} \sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}} \end{aligned}$$

We now show that $\alpha_n/\sqrt{v_{ij}} \rightarrow 0$ in probability. As $\min_{k\ell} |T_{k\ell}| \asymp \|\mathbf{T}\|_{\max}$ and $k, \psi \lesssim 1$, we have $n\|\mathbf{T}\|_{\max} \asymp \lambda_k(\mathbf{T})$ and thus

$$\frac{(\log n)^{1/2} \|\mathbf{U}\|_{2 \rightarrow \infty}^2}{\sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}} \lesssim \ell_{\text{clt}} \frac{n\|\mathbf{T}\|_{\max}}{\sqrt{p}\lambda_k(\mathbf{T})} \frac{\|\mathbf{U}\|_{2 \rightarrow \infty}^2}{\sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}}.$$

Also, since \mathbf{T} is homogeneous, the $\{[\mathbf{T}^2]_{ii}\}_{i=1}^n$ are homogeneous. As $[\mathbf{T}^2]_{ii} = \|[\mathbf{U}\mathbf{\Lambda}]_i\|^2$ and \mathbf{T} have bounded condition number, the $\{\|[\mathbf{U}]_i\|\}_{i=1}^n$ are also homogeneous, i.e., $\min_i \|[\mathbf{U}]_i\|^2 \asymp \max_i \|[\mathbf{U}]_i\|^2$ and hence

$$(S.6.34) \quad \sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2} \asymp \|\mathbf{U}\|_{2 \rightarrow \infty} \asymp n^{-1/2}$$

Combining Eq.(S.6.33), Eq. (S.6.34), and Eq. (S.6.25), and recalling the assumption $np = n^\beta$ for some constant $\beta > 0$, (see (5.2)) we conclude that $\alpha_n/\sqrt{v_{ij}} \rightarrow 0$ in probability.

Now we show the limiting distribution of $\tilde{L}_{ij} := ([\mathbf{U}\mathbf{U}^\top \tilde{\mathbf{E}}]_{ij} + [\tilde{\mathbf{E}}\mathbf{U}\mathbf{U}^\top]_{ij})/\sqrt{v_{ij}}$. Write

$$\tilde{L}_{ij} = \frac{1}{\sqrt{v_{ij}}} \left[\sum_{\ell \neq j} [\tilde{\mathbf{E}}]_{i\ell} \zeta_{\ell j} + \sum_{\ell \neq i} [\tilde{\mathbf{E}}]_{\ell j} \zeta_{i\ell} + [\tilde{\mathbf{E}}]_{ij} \{\zeta_{ii} + \zeta_{jj}\} \right].$$

The summands on the right-hand side of the above display are independent mean 0 random variables. Similar to the proof of Lemma 4.19 in [28], we invoke the Berry–Esseen Theorem to show the desired limiting distribution. We first check the third-order moment condition for \tilde{L}_{ij} , i.e.,

$$\begin{aligned} \gamma_n &:= \frac{1}{(v_{ij})^{3/2}} \sum_{j \neq \ell} \mathbb{E} |[\tilde{\mathbf{E}}]_{i\ell}|^3 |\zeta_{\ell j}|^3 + \sum_{\ell \neq i} \mathbb{E} |[\tilde{\mathbf{E}}]_{\ell j}|^3 |\zeta_{i\ell}|^3 + \mathbb{E} |[\tilde{\mathbf{E}}]_{ij}|^3 |\zeta_{ii} + \zeta_{jj}|^3 \\ &\leq \frac{2\|\mathbf{U}\mathbf{U}^\top\|_{\max} \|\tilde{\mathbf{E}}\|_{\max}}{|v_{ij}|^{3/2}} \left(\sum_{\ell \neq j} \mathbb{E} |[\tilde{\mathbf{E}}]_{i\ell}|^2 \zeta_{\ell j}^2 + \sum_{\ell \neq i} \mathbb{E} |[\tilde{\mathbf{E}}]_{\ell j}|^2 \zeta_{i\ell}^2 + \mathbb{E} |[\tilde{\mathbf{E}}]_{ij}|^2 (\zeta_{ii} + \zeta_{jj})^2 \right) \\ &= \frac{2\|\mathbf{U}\mathbf{U}^\top\|_{\max} \|\tilde{\mathbf{E}}\|_{\max}}{\sqrt{v_{ij}}} \leq \frac{2\|\mathbf{U}\|_{2 \rightarrow \infty}^2 \|\tilde{\mathbf{E}}\|_{\max}}{\sqrt{v_{ij}}} \lesssim \sqrt{\frac{\log n}{p}} \frac{\|\mathbf{U}\|_{2 \rightarrow \infty}^2}{\sqrt{\|[\mathbf{U}]_i\|^2 + \|[\mathbf{U}]_j\|^2}} \rightarrow 0 \end{aligned}$$

in probability. Note that the penultimate inequality in the above derivations follows from Eq. (S.6.32) and the bound

$$(S.6.35) \quad \|\tilde{\mathbf{E}}\|_{\max} = \|\tilde{\mathbf{T}}/p - \mathbf{T}\|_{\max} \lesssim \frac{1}{p} \left\{ \|\mathbf{T}\|_{\max} + \sigma \sqrt{\log n} \right\}$$

with high probability, and the last inequality follows from Eq. (S.6.34) and Eq. (S.6.25).

Let $\Phi(\cdot)$ be the CDF of $\mathcal{N}(0, 1)$ and denote $L_{ij} = ([\mathbf{U}\mathbf{U}^\top \mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij})/\sqrt{v_{ij}}$. Then by the Berry–Esseen bound (see e.g. Theorem 4.21 in [28]), we have

$$\sup_{z \in \mathbb{R}} |\mathbb{P}(\tilde{L}_{ij} \leq z) - \Phi(z)| \leq 10\gamma_n \rightarrow 0.$$

as $n \rightarrow \infty$. Now $L_{ij} = \tilde{L}_{ij}$ whenever $\mathbf{E} = \tilde{\mathbf{E}}$ and hence

$$\begin{aligned} \sup_{z \in \mathbb{R}} |\mathbb{P}(L_{ij} \leq z) - \Phi(z)| &\leq \sup_{z \in \mathbb{R}} |\mathbb{P}[\{L_{ij} \leq z\} \cap \{\tilde{\mathbf{E}} = \mathbf{E}\}] - \Phi(z)| + \mathbb{P}(\tilde{\mathbf{E}} \neq \mathbf{E}) \\ &= \sup_{z \in \mathbb{R}} |\mathbb{P}[\{\tilde{L}_{ij} \leq z\} \cap \{\tilde{\mathbf{E}} = \mathbf{E}\}] - \Phi(z)| + \mathbb{P}(\tilde{\mathbf{E}} \neq \mathbf{E}) \\ &\leq \sup_{z \in \mathbb{R}} |\mathbb{P}[\tilde{L}_{ij} \leq z] - \Phi(z)| + 2\mathbb{P}(\tilde{\mathbf{E}} \neq \mathbf{E}) \\ &\leq \gamma_n + 2\mathbb{P}[\tilde{\mathbf{E}} \neq \mathbf{E}] \rightarrow 0 \end{aligned}$$

as $n \rightarrow \infty$. In summary we have $L_{ij} \rightsquigarrow \mathcal{N}(0, 1)$. Recall that $\alpha_n/\sqrt{v_{ij}} \rightarrow 0$ in probability and hence, by Theorem S.6.1, we have

$$(S.6.36) \quad \frac{[\hat{\mathbf{T}}_g - \mathbf{T}]_{ij}}{\sqrt{v_{ij}}} \rightsquigarrow \mathcal{N}(0, 1).$$

We next show that $v_{ij}/v_{ij}^* \rightarrow 1$ in probability and hence v_{ij} can be replaced by v_{ij}^* in without changing the limit result in Eq.(S.6.36); here v_{ij}^* is defined in (5.4). Recall that $\mathbf{E} = p^{-1}\mathcal{P}_\Omega(\mathbf{T} + \mathbf{N}) - \mathbf{T}$ and hence

$$\mathbb{E}([\mathbf{E}]_{k\ell}^2) = \frac{1-p}{p}[\mathbf{T}]_{k\ell}^2 + \frac{1}{p}\mathbb{E}[\eta_{ij}^2].$$

Then, by Eq. (S.6.30) and Eq. (S.6.31), we have

$$|\mathbb{E}[\mathbf{E}]_{k\ell}^2 - \mathbb{E}[\tilde{\mathbf{E}}]_{k\ell}^2| = \left| \frac{1}{p}\mathbb{E}[\eta_{kl}^2] - \frac{1}{p}\mathbb{E}[\tilde{\eta}_{kl}^2] \right| \leq \frac{\sigma^2}{p} \left(\mathbb{E}[(\eta_{kl}/\sigma)^4] \mathbb{P}(|\eta_{kl}| > 5\sigma\sqrt{\log n}) \right)^{1/2}$$

and thus

$$(S.6.37) \quad \max_{(k,\ell) \in [n]^2} |\mathbb{E}[\mathbf{E}]_{k\ell}^2 - \mathbb{E}[\tilde{\mathbf{E}}]_{k\ell}^2| = o(p^{-1}\sigma^2).$$

Now recall the definitions of v_{ij} and v_{ij}^* . Then by Eq. (S.6.32) and Eq. (S.6.37), we have

$$\begin{aligned} \left| \frac{v_{ij}^* - v_{ij}}{v_{ij}} \right| &\leq \left\{ \max_{(k,\ell) \in [n]^2} |\mathbb{E}[\mathbf{E}]_{k\ell}^2 - \mathbb{E}[\tilde{\mathbf{E}}]_{k\ell}^2| \right\} \cdot \frac{\left[\sum_{\ell \neq j} \zeta_{\ell j}^2 + \sum_{\ell \neq i} \zeta_{i\ell}^2 + \{\zeta_{ii} + \zeta_{jj}\}^2 \right]}{v_{ij}} \\ &\leq \frac{2 \max_{(k,\ell) \in [n]^2} |\mathbb{E}[\mathbf{E}]_{k\ell}^2 - \mathbb{E}[\tilde{\mathbf{E}}]_{k\ell}^2| \cdot \left\{ \|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2 \right\}}{\min_{(k,\ell) \in [n]^2} \mathbb{E}([\tilde{\mathbf{E}}]_{k\ell}^2) \cdot \left\{ \|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2 \right\}} = o(1), \end{aligned}$$

where the last equality is because $\min_{(k,\ell) \in [n]^2} \mathbb{E}([\tilde{\mathbf{E}}]_{k\ell}^2) \geq (2p)^{-1}\sigma^2$; see Eq. (S.6.30). In summary we have $v_{ij}^*/v_{ij} \rightarrow 1$ in probability, as desired.

We now derive the confidence interval for $[\hat{\mathbf{T}}_g]_{ij}$ given in Eq. (5.7). This is equivalent to showing that $\hat{v}_{ij}/v_{ij} \rightarrow 1$ in probability where \hat{v}_{ij} is defined in Eq. (5.6). We first define

$$\tilde{v}_{ij} := \sum_{\ell \neq j} [\tilde{\mathbf{E}}]_{i\ell}^2 \zeta_{\ell j}^2 + \sum_{\ell \neq i} [\tilde{\mathbf{E}}]_{\ell j}^2 \zeta_{i\ell}^2 + [\tilde{\mathbf{E}}]_{ij}^2 (\zeta_{ii} + \zeta_{jj})^2.$$

Using the same argument as that for deriving Eq. (4.171) in [28], we have

$$(S.6.38) \quad \begin{aligned} |\tilde{v}_{ij} - v_{ij}| &\lesssim \frac{(\|\mathbf{T}\|_{\max} + \sigma)^2 \log n}{(np)^{3/2}} + \frac{(\|\mathbf{T}\|_{\max} + \sigma)^2 (\log n)^2}{(np)^2} \\ &\asymp \frac{(\|\mathbf{T}\|_{\max} + \sigma)^2 \log n}{(np)^{3/2}}. \end{aligned}$$

with high probability. We now bound $|\tilde{v}_{ij} - \hat{v}_{ij}|$. Recall that we had defined $\hat{\mathbf{E}} = \hat{\mathbf{T}}_g - p^{-1}\hat{\mathbf{T}}$ in the statement of Theorem 5.1. Now let $\check{\mathbf{E}} := \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \tilde{\mathbf{T}}/p - p^{-1}\hat{\mathbf{T}}$ where $\tilde{\mathbf{T}}$ is defined in Eq. (S.6.27) and let

$$\check{v}_{ij} = \sum_{\ell \neq j} [\check{\mathbf{E}}]_{i\ell}^2 \zeta_{\ell j}^2 + \sum_{\ell \neq i} [\check{\mathbf{E}}]_{\ell j}^2 \zeta_{i\ell}^2 + [\check{\mathbf{E}}]_{ij}^2 (\zeta_{ii} + \zeta_{jj})^2.$$

Then whp $\check{\mathbf{E}} = \hat{\mathbf{E}}$ and $\check{v}_{ij} = \hat{v}_{ij}$. As $\tilde{\mathbf{E}} = \mathbf{E}$ whp, we have by Eq. (S.6.24) that

$$\|\check{\mathbf{E}} - \tilde{\mathbf{E}}\|_{\max} = \|\hat{\mathbf{E}} - \mathbf{E}\|_{\max} = \|p^{-1}\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{T}} - \mathbf{T}\|_{\max} \lesssim \vartheta(\log n)^{\delta^*} (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2.$$

whp. Now recall Eq. (S.6.34) and Eq. (S.6.35). We then have

$$\begin{aligned} \|\check{\mathbf{E}}\|_{\max} &\lesssim \|\tilde{\mathbf{E}}\|_{\max} + \|\check{\mathbf{E}} - \tilde{\mathbf{E}}\|_{\max} \\ &\lesssim \vartheta(\log n)^{\delta^*} (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \frac{1}{p} \left\{ \|\mathbf{T}\|_{\max} + \sigma \sqrt{\log n} \right\} \\ &\lesssim \frac{1}{p} \left\{ \|\mathbf{T}\|_{\max} + \sigma \sqrt{\log n} \right\}, \end{aligned}$$

whp. Furthermore, we have by Eq. (S.6.21) that

$$\|\hat{\mathbf{U}}_g\|_{2 \rightarrow \infty} \leq \|\mathbf{U}\|_{2 \rightarrow \infty} + d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) = (1 + o(1)) \|\mathbf{U}\|_{2 \rightarrow \infty}$$

whp and hence

$$\begin{aligned} \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U} \mathbf{U}^\top\|_{\max} &\leq d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U}) (\|\hat{\mathbf{U}}_g\|_{2 \rightarrow \infty} + \|\mathbf{U}\|_{2 \rightarrow \infty}) \\ &\lesssim \vartheta \frac{n(\|\mathbf{T}\|_{\max} + \sigma) (\log n)^{\delta^*}}{|\lambda_k(\mathbf{T})| \sqrt{np}} \|\mathbf{U}\|_{2 \rightarrow \infty}^2 = o(\|\mathbf{U}\|_{2 \rightarrow \infty}^2), \end{aligned}$$

whp. In summary we have whp that

$$\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} \leq \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U} \mathbf{U}^\top\|_{\max} + \|\mathbf{U} \mathbf{U}^\top\|_{\max} \lesssim \|\mathbf{U}\|_{2 \rightarrow \infty}^2.$$

We can now use the same arguments as that presented in Eq. (4.176)–Eq. (4.177) of [28], but with terms depending on $\hat{\mathbf{U}}$ replaced by terms depending on $\hat{\mathbf{U}}_g$. More specifically let $\hat{\zeta}_{kl} = [\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top]_{kl}$. We then have, after some straightforward but tedious algebra, that (S.6.39)

$$\begin{aligned} |\hat{v}_{ij} - \tilde{v}_{ij}| &= |\check{v}_{ij} - \tilde{v}_{ij}| \\ &\leq \left| \sum_{\ell \neq j} ([\check{\mathbf{E}}]_{i\ell}^2 \hat{\zeta}_{\ell j}^2 - [\tilde{\mathbf{E}}]_{i\ell}^2 \zeta_{\ell j}^2) \right| + \left| \sum_{\ell \neq i} ([\check{\mathbf{E}}]_{\ell j}^2 \hat{\zeta}_{i\ell}^2 - [\tilde{\mathbf{E}}]_{\ell j}^2 \zeta_{i\ell}^2) \right| + 2 \left| [\check{\mathbf{E}}]_{ij}^2 \hat{\zeta}_{ii} \hat{\zeta}_{jj} - [\tilde{\mathbf{E}}]_{ij}^2 \zeta_{ii} \zeta_{jj} \right| \\ &\lesssim (\|\tilde{\mathbf{E}}\|_{\max} + \|\check{\mathbf{E}}\|_{\max}) \|\check{\mathbf{E}} - \tilde{\mathbf{E}}\|_{\max} \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} \\ &\quad + (\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} + \|\mathbf{U} \mathbf{U}^\top\|_{\max}) \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U} \mathbf{U}^\top\|_{\max} \|\tilde{\mathbf{E}}\|^2 \\ &\quad + (\|\tilde{\mathbf{E}}\|_{\max} + \|\check{\mathbf{E}}\|_{\max}) \|\check{\mathbf{E}} - \tilde{\mathbf{E}}\|_{\max} \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max}^2 \\ &\quad + \|\tilde{\mathbf{E}}\|_{\max}^2 (\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} + \|\mathbf{U} \mathbf{U}^\top\|_{\max}) \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U} \mathbf{U}^\top\|_{\max} \\ &\lesssim \underbrace{\vartheta(\log n)^{\delta^*+1/2} \left(\frac{n}{p}\right)^{3/2} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^3}{|\lambda_k(\mathbf{T})|}}_{\zeta_n} \|\mathbf{U}\|_{2 \rightarrow \infty}^4, \end{aligned}$$

whp. In particular the last inequality in the above display is derived as follows.

(i) Let $\delta_1 = (\|\tilde{\mathbf{E}}\|_{\max} + \|\check{\mathbf{E}}\|_{\max}) \|\check{\mathbf{E}} - \tilde{\mathbf{E}}\|_{\max} \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max}$. Then

$$\begin{aligned} \delta_1 &\lesssim \frac{1}{p} \left\{ \|\mathbf{T}\|_{\max} + \sigma \sqrt{\log n} \right\} \cdot \vartheta(\log n)^{\delta^*} (\|\mathbf{T}\|_{\max} + \sigma) \sqrt{\frac{n}{p}} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \\ &\leq \vartheta(\log n)^{\delta^*+1/2} \left(\frac{n}{p}\right)^{3/2} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^2 \|\mathbf{T}\|_{\max}}{n \|\mathbf{T}\|_{\max}} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \lesssim \zeta_n \end{aligned}$$

whp. Note that we had used the fact that $|\lambda_k(\mathbf{T})| \asymp n \|\mathbf{T}\|_{\max}$ in the above derivations.

(ii) Next let $\delta_2 = (\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} + \|\mathbf{U}\mathbf{U}^\top\|_{\max}) \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U}\mathbf{U}^\top\|_{\max} \|\tilde{\mathbf{E}}\|^2$. Then

$$\begin{aligned} \delta_2 &\lesssim \vartheta \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{(\log n)^{\delta^*}}{\sqrt{np}} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \|\tilde{\mathbf{E}}\|^2 \\ &\lesssim \vartheta (\log n)^{\delta^*} (n/p)^{3/2} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^3}{|\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \prec \zeta_n \end{aligned}$$

whp where we used the fact that $\|\tilde{\mathbf{E}}\|$ has the same upper bound as $\|\mathbf{E}\|$ given in (S.6.18).

(iii) Finally let $\delta_3 = \|\mathbf{E}\|_{\max}^2 (\|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top\|_{\max} + \|\mathbf{U}\mathbf{U}^\top\|_{\max}) \|\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top - \mathbf{U}\mathbf{U}^\top\|_{\max}$. Then

$$\begin{aligned} \delta_3 &\lesssim \vartheta (\log n)^{1+\delta^*} \frac{1}{p^2} \sqrt{\frac{n}{p}} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^3}{|\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \\ &\leq \frac{np}{p^2} \sqrt{\frac{n}{p}} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^3}{|\lambda_k(\mathbf{T})|} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \lesssim \zeta_n \end{aligned}$$

whp where we had used the fact that $np \lesssim n^\beta = \omega(\vartheta(\log n)^{1+\delta^*})$.

Note that $\delta_1 + \delta_2$ and $\delta_1 + \delta_3$ are upper bounds for the terms denoted by α_1 and α_3 in Eq. (4.176) and Eq. (4.177) of [28], respectively. Combining Eq. (S.6.29), Eq. (S.6.38), and Eq. (S.6.39), we have whp that

$$\begin{aligned} \frac{|\hat{v}_{ij} - v_{ij}|}{v_{ij}} &\lesssim \frac{(\|\mathbf{T}\|_{\max} + \sigma)^2 \log n}{(np)^{3/2} v_{ij}} + \vartheta (\log n)^{\delta^*+1/2} \left(\frac{n}{p}\right)^{3/2} \frac{(\|\mathbf{T}\|_{\max} + \sigma)^3}{|\lambda_k(\mathbf{T})| v_{ij}} \|\mathbf{U}\|_{2 \rightarrow \infty}^4 \\ &\lesssim \frac{\log n}{\sqrt{np} \cdot n(\|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2)} + \vartheta (\log n)^{\delta^*+1/2} \sqrt{\frac{n}{p}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{\|\mathbf{U}\|_{2 \rightarrow \infty}^4}{\|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2} \\ &\lesssim \frac{\log n}{\sqrt{np}} + \vartheta (\log n)^{\delta^*+1/2} \sqrt{\frac{n}{p}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma)}{|\lambda_k(\mathbf{T})|} \frac{\|\mathbf{U}\|_{2 \rightarrow \infty}^4}{\|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2} \\ &\lesssim \vartheta (\log n)^{\delta^*+1/2} \sqrt{\frac{n}{p}} \frac{n(\|\mathbf{T}\|_{\max} + \sigma) \|\mathbf{U}\|_{2 \rightarrow \infty}^4}{|\lambda_k(\mathbf{T})| (\|\mathbf{U}\|_i^2 + \|\mathbf{U}\|_j^2)} \\ &= o(1), \end{aligned}$$

where the last equality is by Eq. (S.6.34) and Eq. (S.6.26). In summary we have $\sqrt{v_{ij}/\hat{v}_{ij}} \rightarrow 1$ in probability and hence, by Slutsky's Theorem and Eq. (S.6.36), we have

$$\frac{[\hat{\mathbf{T}}_g - \mathbf{T}]_{ij}}{\sqrt{\hat{v}_{ij}}} = \sqrt{\frac{v_{ij}}{\hat{v}_{ij}}} \cdot \frac{[\hat{\mathbf{T}}_g - \mathbf{T}]_{ij}}{\sqrt{v_{ij}}} \rightsquigarrow \mathcal{N}(0, 1),$$

as desired.

S.6.4. Proof of Theorem 5.2. The proof uses the same ideas as that presented in the proof of Theorem 3.3 and thus, for conciseness, we only sketch some of the main steps here. Define $\mathbf{Q}^* = \mathbf{B}\mathbf{F}(\mathbf{B}\mathbf{F})^\top$. Then with probability at least $1 - d^{-6}$, we have

$$(S.6.40) \quad |\lambda_i(\mathbf{Q}^*)| \asymp m\lambda_i \text{ for } i \in [k].$$

and conditional on given \mathbf{Q}^* , with probability at least $1 - d^{-6}$,

$$\|\mathbf{Q} - \mathbf{Q}^*\| \lesssim m\lambda_k \mathcal{E}.$$

See Section 8.2 and Section 10.2 in the supplementary material of [18] for derivations of these bounds. Then we can condition on a high-probability event such that given \mathbf{Q}^* satisfying $|\lambda_i(\mathbf{Q}^*)| \asymp m\lambda_i$ for all $i \in [k]$. Recall that we have assumed bounded condition number

among $\lambda_1, \dots, \lambda_k$. So we then take $\mathbf{M} = \mathbf{Q}^*$ and $\hat{\mathbf{M}} = \mathbf{Q}$ in Assumption 1 with bounded ψ and $S_n = C_1 m \lambda_k$ for some fixed constant $C_1 \geq 0$. By (S.6.40), we further have $E_n = m \lambda_k \mathcal{E}$. Then by Theorem 3.2 and the fact that \mathbf{G} is independent with \mathbf{Q} and \mathbf{Q}^* , we can uncondition the high-probability event of $\mathbf{M} = \mathbf{Q}^*$, and the desired results follow immediately. \square

S.6.5. *Proof of Theorem S.6.1.* We first recall the decomposition of $\hat{\mathbf{U}}_g - \mathbf{U}_g \mathbf{W}_{\text{clt}}$ in Eq. (S.5.43), i.e.,

$$(S.6.41) \quad \hat{\mathbf{U}}_g \mathbf{W}_{\text{clt}}^\top - \mathbf{U} = \mathbf{E} \mathbf{U} \mathbf{A}^{-1} + [\mathbf{R}_1 + \dots + \mathbf{R}_7] \mathbf{W}_{\text{clt}}^\top = \mathbf{E} \mathbf{U} \mathbf{A}^{-1} + \mathbf{R} \mathbf{W}_{\text{clt}}^\top,$$

where $\mathbf{R}_1, \dots, \mathbf{R}_3$ are defined in the proof of Theorem S.3.3 in Section S.5.3, and $\mathbf{R}_4, \dots, \mathbf{R}_7$, \mathbf{W}_{clt} are defined in the proof of Theorem 3.4 in Section S.5.9; in particular \mathbf{W}_{clt} solves the orthogonal Procrustes problem between $\hat{\mathbf{U}}_g$ and $\hat{\mathbf{U}}$ (see Eq. (S.5.47)). We emphasize that Eq. (S.6.41) holds for general matrices $\hat{\mathbf{M}}$ and \mathbf{M} , and does not require Assumption 3 and Eq.(3.14) to be satisfied. For this proof we only consider $\tilde{\mathbf{M}} = \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}}$ and note that the same arguments also apply when $\tilde{\mathbf{M}} = 2^{-1}(\hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} + \hat{\mathbf{M}} \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top)$. For ease of exposition we say that an event \mathcal{E} happens with high probability (whp) if \mathcal{E} happens with probability at least $1 - Cn^{-6}$ where $C > 0$ is a finite constant that can change from line to line.

Let $\tilde{\mathbf{R}} := \mathbf{R} \mathbf{W}_{\text{clt}}^\top$ and note that $\mathbf{E} \mathbf{U} \mathbf{A}^{-1} \mathbf{U}^\top \hat{\mathbf{M}} = \mathbf{E} \mathbf{U} \mathbf{U}^\top + \mathbf{E} \mathbf{U} \mathbf{A}^{-1} \mathbf{U}^\top \mathbf{E}$. Then from Eq. (S.6.41) we have

$$(S.6.42) \quad \begin{aligned} \tilde{\mathbf{M}} - \mathbf{M} &= \hat{\mathbf{U}}_g \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{U} \mathbf{U}^\top \mathbf{M} = \hat{\mathbf{U}}_g \mathbf{W}_{\text{clt}}^\top \mathbf{W}_{\text{clt}} \hat{\mathbf{U}}_g^\top \hat{\mathbf{M}} - \mathbf{U} \mathbf{U}^\top \mathbf{M} \\ &= (\mathbf{U} + \mathbf{E} \mathbf{U} \mathbf{A}^{-1} + \tilde{\mathbf{R}}) (\mathbf{U} + \mathbf{E} \mathbf{U} \mathbf{A}^{-1} + \tilde{\mathbf{R}})^\top \hat{\mathbf{M}} - \mathbf{U} \mathbf{U}^\top \mathbf{M} \\ &= \mathbf{U} \mathbf{U}^\top \mathbf{E} + \mathbf{U} \mathbf{A}^{-1} \mathbf{U}^\top \mathbf{E} \hat{\mathbf{M}} + \mathbf{U} \tilde{\mathbf{R}}^\top \hat{\mathbf{M}} + \mathbf{E} \mathbf{U} \mathbf{A}^{-1} \mathbf{U}^\top \mathbf{E} + \mathbf{E} \mathbf{U} \mathbf{U}^\top \\ &\quad + \mathbf{E} \mathbf{U} \mathbf{A}^{-2} \mathbf{U}^\top \mathbf{E} \hat{\mathbf{M}} + \mathbf{E} \mathbf{U} \mathbf{A}^{-1} \tilde{\mathbf{R}}^\top \hat{\mathbf{M}} + \tilde{\mathbf{R}} \mathbf{U}^\top \hat{\mathbf{M}} \\ &\quad + \tilde{\mathbf{R}} \mathbf{A}^{-1} \mathbf{U}^\top \mathbf{E} \hat{\mathbf{M}} + \tilde{\mathbf{R}} \tilde{\mathbf{R}}^\top \hat{\mathbf{M}}, \end{aligned}$$

The following claim provides upper bound for $\|\mathbf{R}\|_{2 \rightarrow \infty}$, $\|\mathbf{R}\|$ and $\|\mathbf{E} \mathbf{R}\|_{2 \rightarrow \infty}$; we deferred its proof to Section S.7.7. As \mathbf{W}_{clt} is orthogonal, the same bounds also hold for $\|\tilde{\mathbf{R}}\|_{2 \rightarrow \infty}$, $\|\tilde{\mathbf{R}}\|$ and $\|\mathbf{E} \tilde{\mathbf{R}}\|_{2 \rightarrow \infty}$.

CLAIM S.5. *Assume the setting of Theorem 3.2 with k, ψ bounded. Let $g \geq 2 + (2\beta_\Delta)^{-1}$. We then have whp,*

$$(S.6.43) \quad \begin{aligned} \|\mathbf{R}\|_{2 \rightarrow \infty} &\lesssim \ell_{\text{clt}} \left(\frac{E_n}{S_n} \right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty} + \frac{\|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\ \|\mathbf{R}\| &\lesssim \max \{ \vartheta^2, (\log n)^{1/2} \} \left(\frac{E_n}{S_n} \right)^2 + \frac{\|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|}{S_n}, \\ \|\mathbf{E} \mathbf{R}\|_{2 \rightarrow \infty} &\lesssim \ell_{\text{clt}} (\log n)^\delta \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty} + (\log n)^\delta \frac{E_n \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}, \end{aligned}$$

where ℓ_{clt} is specified in Theorem 3.4.

Recall that

$$\alpha_n = \ell_{\text{clt}} \left(\frac{E_n}{S_n} \right) E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \|\mathbf{U}^\top \mathbf{E} \mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \frac{\|\mathbf{U}^\top \mathbf{E} \mathbf{U}\|^2}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}^2.$$

We now bound the max norm of each term on the right-hand side of Eq. (S.6.42) by α_n . We will repeatedly use the observations that (1) $E_n/S_n \lesssim n^{-\beta_\Delta}$ for some $\beta_\Delta > 0$ and thus

decreases to 0 faster than any $\text{polylog}(n)$ factor and (2) $\|\mathbf{M}_1\mathbf{M}_2^\top\|_{\max} \leq \|\mathbf{M}_1\|_{2 \rightarrow \infty} \times \|\mathbf{M}_2\|_{2 \rightarrow \infty}$.

1. $\|\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max}$. As $\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}} = \mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}^2 + \mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\mathbf{M}$ and thus

$$\|\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}^2\|_{\max} \leq \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}^{-1}\| \|\mathbf{E}^2\mathbf{U}\|_{2 \rightarrow \infty} \lesssim (\log n)^{2\delta} \left(\frac{E_n}{S_n}\right) E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

$$\begin{aligned} \|\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top\|_{\max} &\leq \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \|\mathbf{\Lambda}^{-1}\| \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{\Lambda}\| \\ &\lesssim \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \end{aligned}$$

Each of the above terms are less than α_n and hence $\|\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

2. $\|\mathbf{U}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max}$. We decompose $\mathbf{U}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}} = \mathbf{U}\tilde{\mathbf{R}}^\top\mathbf{E} + \mathbf{U}\tilde{\mathbf{R}}^\top\mathbf{M}$. First we have

$$\begin{aligned} \|\mathbf{U}\tilde{\mathbf{R}}^\top\mathbf{E}\|_{\max} &\leq \|\mathbf{E}\tilde{\mathbf{R}}\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \\ &\lesssim \ell_{\text{clt}} (\log n)^\delta \left(\frac{E_n}{S_n}\right)^2 E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + (\log n)^\delta \frac{E_n \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\|}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \alpha_n. \end{aligned}$$

Furthermore we have

$$\begin{aligned} \|\mathbf{U}\tilde{\mathbf{R}}^\top\mathbf{M}\|_{\max} &\leq \|\mathbf{M}\mathbf{R}\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \leq \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \|\mathbf{\Lambda}\| \|\mathbf{R}\| \\ &\lesssim \max\{\vartheta^2, (\log n)^{1/2}\} \left(\frac{E_n}{S_n}\right) E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \alpha_n. \end{aligned}$$

In summary, we have $\|\mathbf{U}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

3. $\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\|_{\max}$. We have whp,

$$\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\|_{\max} \leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty}^2 \|\mathbf{\Lambda}^{-1}\| \lesssim (\log n)^{2\delta} \left(\frac{E_n}{S_n}\right) E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \lesssim \alpha_n$$

4. $\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max}$. We have $\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}} = \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}^2 + \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}\mathbf{M}$. First, we have

$$\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}^2\|_{\max} \leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{E}^2\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}^{-2}\| \lesssim (\log n)^{3\delta} \left(\frac{E_n}{S_n}\right)^2 E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

whp. Second, we have whp,

$$\begin{aligned} \|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}\mathbf{M}\|_{\max} &\leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}\| \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{\Lambda}^{-2}\| \\ &\lesssim (\log n)^\delta \left(\frac{E_n}{S_n}\right) \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2. \end{aligned}$$

In summary, we have $\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-2}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

5. $\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max}$. We decompose $\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}} = \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\mathbf{E} + \mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\mathbf{M}$. First we have

$$\begin{aligned} \|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\mathbf{E}\|_{\max} &\leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{E}\tilde{\mathbf{R}}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}^{-1}\| \\ &\lesssim \ell_{\text{clt}} (\log n)^{2\delta} \left(\frac{E_n}{S_n}\right)^3 E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + (\log n)^{2\delta} \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \end{aligned}$$

whp. Second we have whp

$$\begin{aligned} \|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\mathbf{M}\|_{\max} &\leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}^{-1}\| \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}\| \|\tilde{\mathbf{R}}\| \\ &\lesssim \vartheta (\log n)^\delta \left(\frac{E_n}{S_n}\right)^2 E_n \|\mathbf{U}\|_{2 \rightarrow \infty}^2 + (\log n)^\delta \frac{E_n}{S_n} \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2 \rightarrow \infty}^2. \end{aligned}$$

In summary, we have $\|\mathbf{E}\mathbf{U}\mathbf{\Lambda}^{-1}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

6. $\|\tilde{\mathbf{R}}\mathbf{U}^\top\hat{\mathbf{M}}\|_{\max}$. We decompose $\tilde{\mathbf{R}}\mathbf{U}^\top\hat{\mathbf{M}} = \tilde{\mathbf{R}}\mathbf{U}^\top\mathbf{E} + \tilde{\mathbf{R}}\mathbf{U}^\top\mathbf{M} = \tilde{\mathbf{R}}\mathbf{U}^\top\mathbf{E} + \tilde{\mathbf{R}}\mathbf{\Lambda}\mathbf{U}^\top$. First we have

$$\begin{aligned}\|\tilde{\mathbf{R}}\mathbf{U}^\top\mathbf{E}\|_{\max} &= \|\mathbf{E}\mathbf{U}\|_{2\rightarrow\infty}\|\tilde{\mathbf{R}}\|_{2\rightarrow\infty} \\ &\lesssim \ell_{\text{clt}}(\log n)^\delta \left(\frac{E_n}{S_n}\right)^2 E_n \|\mathbf{U}\|_{2\rightarrow\infty}^2 + (\log n)^\delta \frac{E_n}{S_n} \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2\end{aligned}$$

Second we have

$$\|\tilde{\mathbf{R}}\mathbf{\Lambda}\mathbf{U}^\top\|_{\max} \leq \|\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\mathbf{U}\mathbf{\Lambda}\|_{2\rightarrow\infty} \lesssim \ell_{\text{clt}}\left(\frac{E_n}{S_n}\right) E_n \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2$$

whp. In summary we have $\|\tilde{\mathbf{R}}\mathbf{U}^\top\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

7. $\|\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max}$. We decompose $\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}} = \tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}^2 + \tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\mathbf{M}$. First we have

$$\begin{aligned}\|\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}^2\|_{\max} &\leq \|\mathbf{E}^2\mathbf{U}\|_{2\rightarrow\infty}\|\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\mathbf{\Lambda}^{-1}\| \\ &\lesssim \ell_{\text{clt}}(\log n)^{2\delta} \left(\frac{E_n}{S_n}\right)^3 E_n \|\mathbf{U}\|_{2\rightarrow\infty}^2 + (\log n)^{2\delta} \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2\end{aligned}$$

whp. Second we have whp

$$\begin{aligned}\|\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\mathbf{M}\|_{\max} &\leq \|\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\|_{2\rightarrow\infty}\|\mathbf{M}\mathbf{E}\mathbf{U}\|_{2\rightarrow\infty} \\ &\leq \|\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\mathbf{\Lambda}^{-1}\| \|\mathbf{U}\|_{2\rightarrow\infty}\|\mathbf{\Lambda}\| \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \\ &\lesssim \ell_{\text{clt}}\left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \frac{\|\mathbf{U}^\top\mathbf{E}\mathbf{U}\|^2}{S_n} \|\mathbf{U}\|_{2\rightarrow\infty}^2.\end{aligned}$$

In summary we have $\|\tilde{\mathbf{R}}\mathbf{\Lambda}^{-1}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

8. $\|\tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max}$. We decompose $\tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}} = \tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\mathbf{E} + \tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\mathbf{M}$. First whp

$$\begin{aligned}\|\tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\mathbf{E}\|_{\max} &\leq \|\mathbf{E}\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\tilde{\mathbf{R}}\|_{2\rightarrow\infty} \\ &\lesssim \left(\frac{E_n}{S_n}\right)^3 E_n \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \frac{\|\mathbf{U}^\top\mathbf{E}\mathbf{U}\|^2}{S_n} \|\mathbf{U}\|_{2\rightarrow\infty}^2,\end{aligned}$$

Second we have whp,

$$\begin{aligned}\|\tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\mathbf{M}\|_{\max} &\leq \|\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top\tilde{\mathbf{R}}\|_{2\rightarrow\infty} \\ &\leq \|\tilde{\mathbf{R}}\|_{2\rightarrow\infty}\|\mathbf{U}\|_{2\rightarrow\infty}\|\mathbf{\Lambda}\|\|\tilde{\mathbf{R}}\| \\ &\lesssim \left(\frac{E_n}{S_n}\right)^2 E_n \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \|\mathbf{U}^\top\mathbf{E}\mathbf{U}\| \|\mathbf{U}\|_{2\rightarrow\infty}^2 + \frac{\|\mathbf{U}^\top\mathbf{E}\mathbf{U}\|^2}{S_n} \|\mathbf{U}\|_{2\rightarrow\infty}^2.\end{aligned}$$

In summary we have $\|\tilde{\mathbf{R}}\tilde{\mathbf{R}}^\top\hat{\mathbf{M}}\|_{\max} = \mathcal{O}(\alpha_n)$ whp.

Substituting the above bounds into Eq. (S.6.42) we have, for any $(i, j) \in [n]^2$, that

$$[\tilde{\mathbf{M}}]_{ij} - [\mathbf{M}]_{ij} = [\mathbf{U}\mathbf{U}^\top\mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij} + \mathcal{R}_{ij},$$

where $|\mathcal{R}_{ij}| \lesssim \alpha_n$ whp. Then, under the condition of Theorem S.6.1, we have

$$\frac{[\tilde{\mathbf{M}}]_{ij} - [\mathbf{M}]_{ij}}{\sqrt{v_{ij}}} = \frac{[\mathbf{U}\mathbf{U}^\top\mathbf{E}]_{ij} + [\mathbf{E}\mathbf{U}\mathbf{U}^\top]_{ij}}{\sqrt{v_{ij}}} + \frac{\mathcal{R}_{ij}}{\sqrt{v_{ij}}} \rightsquigarrow \mathcal{N}(0, 1),$$

by Slutsky's theorem. \square

S.6.6. *Proof of Theorem S.2.1.* Theorem S.2.1 follows *mutatis mutandis* from the proofs of Theorem 3.2 and Theorem 3.3, with the conditions in Assumptions 1 and 2 replaced by those in Assumptions S.1 and S.2. For brevity we only present proof sketches for the bounds $\|(\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G} - (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G}\|$ and $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$ given in Eq. (S.2.1) and Eq. (S.2.2). The bound for $d_{2 \rightarrow \infty}(\hat{\mathbf{U}}_g, \mathbf{U})$ in Eq.(S.2.3) is also straightforward but requires more tedious book-keeping and is thus omitted.

Let \mathbf{U}_g be the left singular vectors of $(\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G} = \mathbf{U}\Sigma^{2g+1}\mathbf{U}^\top \mathbf{G}$ where \mathbf{G} is $n \times \tilde{k}$ matrix whose entries are iid standard normals. As $\text{rk}(\mathbf{U}^\top \mathbf{G}) = k$ almost surely, \mathbf{U}_g is equivalent to \mathbf{U} , i.e.,

$$(S.6.44) \quad d_2(\mathbf{U}_g, \mathbf{U}) = \|\sin \Theta(\mathbf{U}_g, \mathbf{U})\| = d_{2 \rightarrow \infty}(\mathbf{U}_g, \mathbf{U}) = 0$$

almost surely. See also the proof of Proposition 3.1 in Section S.7.1.

We now introduce some notations. Let J be any positive integer and $\mathbf{M}_1, \dots, \mathbf{M}_J$ be arbitrary matrices of dimensions $n_1 \times n_2$. First define

$$\langle \mathbf{M}_1 \cdots \mathbf{M}_J \rangle := \mathbf{M}_1 \mathbf{M}_2^\top \mathbf{M}_3 \mathbf{M}_4^\top \cdots \mathbf{M}_J^*$$

where $\mathbf{M}_J^* = \mathbf{M}_J^\top$ if J is even and $\mathbf{M}_J^* = \mathbf{M}_J$ otherwise. Next, for $\ell_1, \dots, \ell_J > 0$, define

$$\langle \mathbf{M}_1^{\ell_1} \cdots \mathbf{M}_J^{\ell_J} \rangle := \underbrace{\langle \mathbf{M}_1 \cdots \mathbf{M}_1 \rangle}_{\ell_1 \text{ terms}} \cdots \underbrace{\langle \mathbf{M}_J \cdots \mathbf{M}_J \rangle}_{\ell_J \text{ terms}}.$$

Finally, define

$$\Delta_g = (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} - (\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G} = \{ \langle \hat{\mathbf{M}}^{2g+1} \rangle - \langle \mathbf{M}^{2g+1} \rangle \} \mathbf{G}.$$

We say that an event \mathcal{A} happens with probability at least $1 - \mathcal{O}(n^{-6})$ if $\mathbb{P}(\mathcal{A}) \geq 1 - Cn^{-6}$ where C is some fixed constant not depending on n .

We now derive the bound for $\|\Delta_g\|$ given in Eq. (S.2.1). Similar to Lemma S.4.2, we have

$$\Delta_g = \underbrace{\langle \mathbf{E}^{2g+1} \rangle \mathbf{G}}_{\mathbf{L}_1} + \underbrace{\sum_{\xi=0}^{2g-1} \sum_{\ell=0}^{2g-1-\xi} \langle \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \rangle \mathbf{G}}_{\mathbf{L}_2} + \underbrace{\sum_{\xi=0}^{2g-1} \langle \mathbf{M}^{2g-\xi} \mathbf{E}^{\xi+1} \rangle \mathbf{G}}_{\mathbf{L}_3}.$$

Bounding $\|\mathbf{L}_1\|$: Similar to Eq. (S.5.2), we have with probability at least $1 - \mathcal{O}(n^{-6})$,

$$\|\mathbf{L}_1\| \leq \|\langle \mathbf{E}^{2g+1} \rangle\| \times \|\mathbf{G}\| \leq \|\mathbf{E}\|^{2g+1} \times \|\mathbf{G}\| \lesssim E_{n_1, n_2}^{2g+1} \|\mathbf{G}\|.$$

Then similar to Eq. (S.4.4) and Eq. (S.4.5), we have

$$\|\mathbf{G}\| \lesssim \max\{n_2^{1/2}, \tilde{k}^{1/2}, \log^{1/2} n\} = \max\{n_2^{1/2}, \log^{1/2} n\},$$

with probability at least $1 - \mathcal{O}(n^{-6})$. In summary, $\|\mathbf{L}_1\| \lesssim E_{n_1, n_2}^{2g+1} \max\{n_2^{1/2}, \log^{1/2} n\}$ with probability at least $1 - \mathcal{O}(n^{-6})$.

Bounding $\|\mathbf{L}_2\|$: First we note that

$$\|\mathbf{L}_2\| \leq \sum_{\xi=0}^{2g-1} \sum_{\ell=0}^{2g-1-\xi} \|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \mathbf{G}\|.$$

We now bound $\|\hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \mathbf{G}\|$. Suppose that ξ is odd (the case when ξ is even follows the same identical argument). We then have

$$\langle \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \rangle \mathbf{G} = \langle \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell-1} \rangle \mathbf{V} \Sigma \mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E} \mathbf{G}.$$

Then, similar to Eq. (S.5.3), we have,

$$\begin{aligned} \|\langle \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \rangle \mathbf{G}\| &\leq \|\hat{\mathbf{M}}\|^\ell \times \|\mathbf{E}\| \times \|\mathbf{M}\|^{2g-\xi-\ell-1} \times \|\boldsymbol{\Sigma}\| \times \|\mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E} \mathbf{G}\| \\ &\lesssim S_{n_1, n_2}^{2g-\xi} E_{n_1, n_2} \|\mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E} \mathbf{G}\| \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. Similar to Lemma S.4.3, let $\bar{\mathbf{V}}_\xi \in \mathbb{R}^{n_2 \times k}$ be the right singular vectors of $\mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E}$. We then have

$$\begin{aligned} \|\mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E} \mathbf{G}\| &\leq \|\mathbf{U}^\top (\mathbf{E} \mathbf{E}^\top)^{(\xi-1)/2} \mathbf{E}\| \times \|\bar{\mathbf{V}}_\xi^\top \mathbf{G}\| \\ (S.6.45) \quad &\lesssim E_{n_1, n_2}^\xi \max\{\tilde{k}^{1/2}, \log^{1/2} n\}, \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. Therefore, with probability at least $1 - \mathcal{O}(n^{-6})$,

$$\|\langle \hat{\mathbf{M}}^\ell \mathbf{E} \mathbf{M}^{2g-\xi-\ell} \mathbf{E}^\xi \rangle \mathbf{G}\| \lesssim S_{n_1, n_2}^{2g-\xi} E_{n_1, n_2}^{\xi+1} \max\{\tilde{k}^{1/2}, \log^{1/2} n\},$$

and thus $\|\mathbf{L}_2\| \lesssim S_{n_1, n_2}^{2g} E_{n_1, n_2} \max\{\tilde{k}^{1/2}, \log^{1/2} n\}$.

Bounding $\|\mathbf{L}_3\|$: Once again we focus on the case where ξ is odd. First note that

$$\langle \mathbf{M}^{2g-\xi} \mathbf{E}^{\xi+1} \rangle \mathbf{G} = \langle \mathbf{M}^{2g-\xi-1} \rangle \mathbf{U} \boldsymbol{\Sigma} \mathbf{V}^\top (\mathbf{E}^\top \mathbf{E})^{(\xi+1)/2} \mathbf{G}.$$

Next, similar to Eq. (S.5.7) and Eq. (S.6.45), we have

$$\begin{aligned} \|\langle \mathbf{M}^{2g-\xi} \mathbf{E}^{\xi+1} \rangle \mathbf{G}\| &\leq \|\mathbf{M}\|^{2g-\xi-1} \times \|\boldsymbol{\Sigma}\| \times \|\mathbf{V}^\top (\mathbf{E}^\top \mathbf{E})^{(\xi+1)/2} \mathbf{G}\| \\ (S.6.46) \quad &\lesssim S_{n_1, n_2}^{2g-\xi} E_{n_1, n_2}^{\xi+1} \max\{\tilde{k}^{1/2}, \log^{1/2} n\} \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. In summary, similar to Eq. (S.5.7), we have

$$\|\mathbf{L}_3\| \leq \sum_{\xi=0}^{2g-1} \|\langle \mathbf{M}^{2g-\xi} \mathbf{E}^{\xi+1} \rangle \mathbf{G}\| \lesssim S_{n_1, n_2}^{2g} E_{n_1, n_2} \max\{\tilde{k}^{1/2}, \log^{1/2} n\},$$

with probability at least $1 - \mathcal{O}(n^{-6})$.

Combining the upper bounds above for $\|\mathbf{L}_1\|$, $\|\mathbf{L}_2\|$ and $\|\mathbf{L}_3\|$ we obtain

$$\begin{aligned} \|\boldsymbol{\Delta}_g\| &\leq \|\mathbf{L}_1\| + \|\mathbf{L}_2\| + \|\mathbf{L}_3\| \\ &\lesssim E_{n_1, n_2}^{2g+1} \eta^{1/2} + S_{n_1, n_2}^{2g} E_{n_1, n_2} \max\{\tilde{k}^{1/2}, \log^{1/2} n\}, \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$, where $\eta = \max\{n_2, \log n\}$. Now recall Assumption S.1. Then by choosing $g \geq (4\beta_\Delta)^{-1}$, we have

$$(S.6.47) \quad \|\boldsymbol{\Delta}_g\| \lesssim S_{n_1, n_2}^{2g} E_{n_1, n_2} \max\{\tilde{k}^{1/2}, \log^{1/2} n\},$$

with probability at least $1 - \mathcal{O}(n^{-6})$. Eq. (S.2.1) is thereby established.

Bounding $\sigma_k^{-1}\{(\hat{\mathbf{M}} \hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}} \mathbf{G}\}$: Denote the SVD of $\hat{\mathbf{M}}$ as

$$\hat{\mathbf{M}} = \hat{\mathbf{U}} \hat{\boldsymbol{\Sigma}} \hat{\mathbf{V}}^\top + \hat{\mathbf{U}}_\perp \hat{\boldsymbol{\Sigma}}_\perp \hat{\mathbf{V}}_\perp^\top,$$

where $\hat{\boldsymbol{\Sigma}} \in \mathbb{R}^{k \times k}$ is the diagonal matrix formed by the k largest singular values of $\hat{\mathbf{M}}$ and $\hat{\boldsymbol{\Sigma}}_\perp \in \mathbb{R}^{(n-k) \times (n-k)}$ is the diagonal matrix for the remaining singular values. Similar to Eq. (S.4.6)–Eq. (S.4.9), we have for any $a \in [a_n]$ that

$$\mathbf{G}_a^\top \hat{\mathbf{M}}^\top (\hat{\mathbf{M}} \hat{\mathbf{M}}^\top)^{2g} \hat{\mathbf{M}} \mathbf{G}_a = \mathbf{G}_a^\top \hat{\mathbf{V}} \hat{\boldsymbol{\Sigma}}^{4g+2} \hat{\mathbf{V}}^\top \mathbf{G}_a + \mathbf{G}_a^\top \hat{\mathbf{V}}_\perp \hat{\boldsymbol{\Sigma}}_\perp^{4g+2} \hat{\mathbf{V}}_\perp^\top \mathbf{G}_a.$$

Therefore, by Weyl's inequality,

$$\begin{aligned} \sigma_k^2\{(\hat{\mathbf{M}} \hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}} \mathbf{G}\} &= \max_{a \in [a_n]} \lambda_k\{\mathbf{G}_a^\top \hat{\mathbf{M}}^\top (\hat{\mathbf{M}} \hat{\mathbf{M}}^\top)^{2g} \hat{\mathbf{M}} \mathbf{G}_a\} \\ &\geq \max_{a \in [a_n]} \lambda_k\{\mathbf{G}_a^\top \hat{\mathbf{V}} \hat{\boldsymbol{\Sigma}}^{4g+2} \hat{\mathbf{V}}^\top \mathbf{G}_a\} \geq \sigma_k^{4g+2}(\hat{\mathbf{M}}) \max_{a \in [a_n]} \sigma_k^2\{\hat{\mathbf{V}}^\top \mathbf{G}_a\}. \end{aligned}$$

When $\psi = o(S_{n_1, n_2}/E_{n_1, n_2})$, similar to Eq. (S.4.9), we have $\sigma_k(\hat{\mathbf{M}}) \lesssim \psi^{-1} S_{n_1, n_2}$ with probability at least $1 - n^{-6}$. Next, using the same arguments as that for deriving Eq. (S.4.11)–Eq. (S.4.17), we obtain

$$\max_{a \in [a_n]} \sigma_k^2 \{ \hat{\mathbf{V}}^\top \mathbf{G}_a \} \gtrsim \tilde{k},$$

with probability at least $1 - \mathcal{O}(n^{-6})$, provided that k and \tilde{k} satisfy one of the conditions (a)–(c) as specified in Theorem 3.2 and $a_n \geq C\tilde{k}^{-1} \log n$ for some universal constant C . In summary we have, with probability at least $1 - \mathcal{O}(n^{-6})$,

$$(S.6.48) \quad \sigma_k^{-1} \{ (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} \} \lesssim \tilde{k}^{-1/2} \psi^{2g+1} S_{n_1, n_2}^{-(1+2g)}.$$

Bounding $d_2(\hat{\mathbf{U}}_g, \mathbf{U})$: We apply Wedin’s sin- Θ theorem [122], i.e.,

$$\begin{aligned} \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\| &\leq \frac{\|\Delta_g\|}{\sigma_k \{ (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} \} - \sigma_{k+1} \{ (\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G} \}} \\ &= \|\Delta_g\| \sigma_k^{-1} \{ (\hat{\mathbf{M}}\hat{\mathbf{M}}^\top)^g \hat{\mathbf{M}}\mathbf{G} \} \lesssim \psi \psi^{2g+1} \frac{E_{n_1, n_2}}{S_{n_1, n_2}}, \end{aligned}$$

with probability at least $1 - \mathcal{O}(n^{-6})$. The equality in the above display is because $\text{rk}\{(\mathbf{M}\mathbf{M}^\top)^g \mathbf{M}\mathbf{G}\} \leq k$ while the last inequality follows from Eq. (S.6.47) and Eq. (S.6.48). Eq. (S.2.2) is thereby established. \square

S.7. Additional Proofs.

S.7.1. Proof of Proposition 3.1. When $\text{rk}(\mathbf{M}) = k^*$, we have $\mathbf{M} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^\top$ by Eq. (2.1). Then $\mathbf{M}^g \mathbf{G} = \mathbf{U}\mathbf{\Lambda}^g \mathbf{U}^\top \mathbf{G}$ where $\mathbf{\Lambda}^g$ is a $k^* \times k^*$ diagonal matrix with full rank. In addition, since \mathbf{G} is chosen from many standard Gaussian matrix \mathbf{G}_a with $a \in [a_n]$, and $\mathbf{U} \in \mathbb{O}_{n \times k}$, it is easy to see each $\mathbf{U}^\top \mathbf{G}_a$ is a $k^* \times \tilde{k}$ standard Gaussian matrix. Recall that $\tilde{k} \geq k^*$.

Now if $\mathbf{U}^\top \mathbf{G}$ has rank k^* then $\mathbf{\Lambda}^g \mathbf{U}^\top \mathbf{G}$ is $k^* \times \tilde{k}$ matrix with rank k^* and thus, as a result, the column space of $\mathbf{M}^g \mathbf{G} = \mathbf{U} \cdot \mathbf{\Lambda}^g \mathbf{U}^\top \mathbf{G}$ coincides with the column space of \mathbf{U} . As $\mathbf{U}_g \in \mathbb{O}_{n \times k^*}$ is the left singular matrix of $\mathbf{M}^g \mathbf{G}$ and $\mathbf{U} \in \mathbb{O}_{n \times k^*}$, there exists an orthogonal $\mathbf{W}_{k, g} \in \mathbb{O}_{k^*}$ such that $\mathbf{U}_g = \mathbf{U}\mathbf{W}_{k^*, g}$. By the definition of d_* for $* \in \{2, 2 \rightarrow \infty\}$, we have

$$(S.7.1) \quad 0 \leq d_*(\mathbf{U}_g, \mathbf{U}) = \inf_{\mathbf{W}_{k^*} \in \mathbb{O}_{k^*}} \|\mathbf{U}_g - \mathbf{U}\mathbf{W}_{k^*}\|_* \leq \|\mathbf{U}_g - \mathbf{U}\mathbf{W}_{k^*, g}\|_* = 0.$$

Finally, since $\mathbb{P}(\text{rk}(\mathbf{U}^\top \mathbf{G}_a) = k^*) = 1$ for any $a \in [a_n]$ (see e.g. [44]) and \mathbf{G} is chosen from $\mathbf{G}_1, \dots, \mathbf{G}_{a_n}$, we immediately have $\mathbb{P}(\text{rk}(\mathbf{U}^\top \mathbf{G}) = k^*) = 1$. Therefore, we conclude that Eq. (S.7.1) holds almost surely. \square

S.7.2. Proof of Proposition 3.2. The ideas of this proof are adapted from the proof of Lemmas 6.5 and 7.10 in [48]. We first assume that \mathbf{E} is entrywise independent and then extend it to the case where \mathbf{E} is symmetric with independent (upper-triangular) entries. Fix a particular basis vector \mathbf{e}_{i^*} and a vector $\mathbf{u} \in \mathbb{R}^n$ not depending on \mathbf{E} . Our target event can be written as

$$(S.7.2) \quad \mathcal{A} = \left\{ \mathbf{E}: |\mathbf{e}_{i^*}^\top \mathbf{E}^g \mathbf{u}| \leq 1.5 E_n^g (\log n)^{g\delta} \|\mathbf{u}\|_{\max} \right\} = \left\{ \mathbf{H}: |\mathbf{e}_{i^*}^\top \mathbf{H}^g \bar{\mathbf{u}}| \leq 1.5 (\log n)^{g\delta} \right\},$$

where $\mathbf{H} := \mathbf{E}/E_n = (h_{ij})_{n \times n}$ and $\bar{\mathbf{u}} := \mathbf{u}/\|\mathbf{u}\|_{\max} = (\bar{u}_1, \dots, \bar{u}_n)^\top$. We now derive an upper bound for $\mathbb{P}(\mathcal{A})$. Denote, for brevity, $\mathcal{X} := \mathbf{e}_{i^*}^\top \mathbf{H}^g \bar{\mathbf{u}}$.

Case a. ($g = 1$) When $g = 1$, we have

$$\mathcal{X} = \mathbf{e}_i^\top \mathbf{H} \bar{\mathbf{u}} = \sum_{i=1}^n h_{i^*i} \bar{u}_i.$$

By the moment conditions of e_{ij} , we know for any $2 \leq g' \leq (\log n)^{A_0 \log \log n}$,

$$(S.7.3) \quad \mathbb{E}|h_{ii^*}|^{g'} = \mathbb{E}[|e_{ij}/E_n|^{g'}] \leq \frac{C_1^{g'}}{nq_n^{g'-2}}.$$

In addition, $h_{i^*1}, \dots, h_{i^*n}$ are independent mean zero random variables. Then, by taking $\alpha = 1, \beta = -2, \gamma = 1$ in Lemma 3.8 in [48], we have when $\delta > 1$ is fixed,

$$\begin{aligned} |\mathcal{X}| &= \left| \sum_{i=1}^n h_{i^*i} \bar{u}_i \right| \leq (\log n)^\delta \left[\frac{\sup_i |\bar{u}_i|}{q_n} + \left\{ \frac{1}{n} \sum_{i=1}^n \bar{u}_i^2 \right\}^{1/2} \right] \\ &\leq (\log n)^\delta \left[\frac{1}{q_n} + 1 \right] \leq 1.1(\log n)^\delta \end{aligned}$$

with probability at least $1 - \exp\{-\nu(\log n)^\delta\}$, provided that $n \geq N(C_1, \delta, A_0, \{q_n\}_{n \geq 1})$. Here $N(C_1, \delta, A_0, \{q_n\}_{n \geq 1})$ is a constant depending only on C_1, δ, A_0 , and the divergence rate of q_n , while ν is a constant depending only on C_1 . Eq. (S.7.3) can be rewritten as $|\mathbf{e}_i^\top \mathbf{E} \mathbf{u}| \leq 1.1 E_n (\log n)^\delta \|\mathbf{u}\|_{\max}$ which implies the desired result.

Case b. ($2 \leq g \leq \log n$) We first note the following two deterministic bounds: By the moment condition of e_{ij} , we have when $g' = 2$,

$$\mathbb{E}|h_{ij}|^2 = \mathbb{E}[|e_{ij}|^2]/E_n^2 \leq C_1^2/n,$$

for any $i, j \in [n]$. Next, if $3 \leq g' \leq (\log n)^{\delta+2}$ and n is sufficiently large such that $(\log n)^{A_0 \log \log n} \geq (\log n)^{\delta+2}$ and $q_n > 3 \log C_1$, then

$$\begin{aligned} \mathbb{E}|h_{ij}|^{g'} &\leq \frac{C_1^{g'}}{nq_n^{g'-2}} = \frac{1}{n} \exp \left\{ g'(\log C_1 - \log q_n) + 2 \log q_n \right\}. \\ &\leq \frac{1}{n} \exp \left\{ 3(\log C_1 - \log q_n) + 2 \log q_n \right\} \leq \frac{C_1^2}{n}. \end{aligned}$$

In other words, there exists a constant $N_1(C_1, A_0, \{q_n\}_{n \geq 1})$ (henceforth N_1) depending only on $C_1, A_0 > 0$, and the divergence rate of q_n such that for all $n \geq N_1$, for any $g' \in [2, (\log n)^{\delta+2}]$, and for all $i, j \in [n]$ we have

$$(S.7.4) \quad \mathbb{E}|h_{ij}|^{g'} \leq \frac{C_1^2}{n}.$$

We next introduce some additional notation. Let α denote a generic pair (i, j) with $i, j \in [n]$. The s th coordinate of α (with $s \in \{1, 2\}$) is denoted as $\alpha^{(s)}$. For two pairs α and $\tilde{\alpha}$, define $\chi_{\alpha\tilde{\alpha}} = \mathbf{1}\{\alpha^{(2)} = \tilde{\alpha}^{(1)}\}$.

We now bound the ι th moment of \mathcal{X} for integer $\iota > 0$. By definition, we have

$$(S.7.5) \quad \begin{aligned} \mathcal{X} &= \sum_{(i_1, \dots, i_g) \in [n]^g} h_{i^*i_1} h_{i_1i_2} \cdots h_{i_{g-1}i_g} \bar{u}_{i_g} \\ &= \sum_{(\alpha_1, \dots, \alpha_g) \in ([n]^2)^g} \chi_{\alpha_0\alpha_1} \chi_{\alpha_1\alpha_2} \cdots \chi_{\alpha_{g-1}\alpha_g} h_{\alpha_1} \cdots h_{\alpha_g} \bar{u}_{\alpha_g^{(2)}} := \sum_{\beta \in ([n]^2)^g} \zeta_\beta, \end{aligned}$$

where (i_1, \dots, i_g) denote a collection of g integers (not necessarily distinct) from $[n]$, $\alpha_0 := (0, i^*)$, and $(\alpha_1, \dots, \alpha_g)$ denote a collection of g pairs (not necessarily distinct) from $[n] \times [n]$. For brevity we use β to denote a generic $(\alpha_1, \dots, \alpha_g)$ and $\zeta_\beta := \chi_{\alpha_0 \alpha_1} \chi_{\alpha_1 \alpha_2} \cdots \chi_{\alpha_{g-1} \alpha_g} h_{\alpha_1} \cdots h_{\alpha_g} \bar{u}_{\alpha_g^{(2)}}$. Then, for any $\iota \geq 1$,

$$(S.7.6) \quad \mathcal{X}^\iota = \sum_{(\beta_1, \beta_2, \dots, \beta_\iota)} \zeta_{\beta_1} \zeta_{\beta_2} \cdots \zeta_{\beta_\iota},$$

where $(\beta_1, \beta_2, \dots, \beta_\iota)$ now runs over all possible combinations of $\beta_1 \dots \beta_\iota \in ([n]^2)^g$. For a particular $(\beta_1, \beta_2, \dots, \beta_\iota)$, we denote $\beta_\ell = (\alpha_{1,\ell}, \dots, \alpha_{g,\ell})$ for $\ell \in [\iota]$. We denote a generic $(\beta_1, \dots, \beta_\iota)$ by \mathbf{B} .

Suppose \mathbf{B} is given. We can then construct a partition $\Gamma_{\mathbf{B}}$ over $[\iota] \times [g]$ (induced by \mathbf{B} as follows). $\Gamma_{\mathbf{B}} = \{\gamma_{\zeta, \mathbf{B}}\}$ is a collection of disjoint sets such that for any $(\ell, \tau) \in [\iota] \times [g]$ and $(\ell', \tau') \in [\iota] \times [g]$, (ℓ, τ) and (ℓ', τ') belong to the same partition $\gamma_{\zeta, \mathbf{B}} \in \Gamma_{\mathbf{B}}$ if and only if,

$$(S.7.7) \quad \beta_\ell^{(\tau)} := \alpha_{\tau, \ell} = \alpha_{\tau', \ell'} =: \beta_{\ell'}^{(\tau')}.$$

The above property for $\gamma_{\zeta, \mathbf{B}}$ implies the existence of a $\alpha \in [n]^2$ such that $\alpha_{\tau, \ell} = \alpha$ for all $(\ell, \tau) \in \gamma_{\zeta, \mathbf{B}}$. We denote this α by $\alpha_{\gamma_{\zeta, \mathbf{B}}}$. There is then a bijection between the set of distinct $\mathbf{B} = (\beta_1, \dots, \beta_\iota)$ and the set of partition $\Gamma_{\mathbf{B}}$ and α associated with $\Gamma_{\mathbf{B}}$, i.e., a bijection between \mathbf{B} and $\bar{\Gamma}_{\mathbf{B}} := \{(\gamma_{\zeta, \mathbf{B}}, \alpha_{\gamma_{\zeta, \mathbf{B}}})\}_{\zeta=1}^{|\bar{\Gamma}_{\mathbf{B}}|}$. We denote the mapping from \mathbf{B} to $\bar{\Gamma}_{\mathbf{B}}$ by $\Upsilon(\cdot)$, i.e., $\Upsilon(\mathbf{B}) = \bar{\Gamma}_{\mathbf{B}}$ and the inverse mapping by Υ^{-1} . More specifically, given a partition Γ of $[\iota] \times [g]$ together with its α -assignment, i.e.,

$$(S.7.8) \quad \bar{\Gamma} = \{(\gamma_\zeta, \alpha_{\gamma_\zeta})\}_{\zeta=1}^{|\bar{\Gamma}|},$$

there exists a unique $\mathbf{B}_{\bar{\Gamma}}$ such that

$$(S.7.9) \quad \mathbf{B}_{\bar{\Gamma}} = (\beta_{1, \bar{\Gamma}}, \dots, \beta_{\iota, \bar{\Gamma}}) := \Upsilon^{-1}(\bar{\Gamma}).$$

For a $\bar{\Gamma}$ of the form in Eq. (S.7.8), we can construct a multigraph $G_{\bar{\Gamma}} = [V(G_{\bar{\Gamma}}), E(G_{\bar{\Gamma}})]$ with vertex set $V(G_{\bar{\Gamma}})$ and edge set $E(G_{\bar{\Gamma}})$ as follows.

- $V(G_{\bar{\Gamma}}) = \{\gamma_0, \gamma_1, \dots, \gamma_{|\bar{\Gamma}|}\}$ where γ_0 is an auxiliary node with $\alpha_{\gamma_0} = (0, i^*)$. Recall that we currently assume a fixed i^* and hence γ_0 does not depend on $\bar{\Gamma}$.
- For any two indices $\zeta \in \{0, 1, \dots, |\bar{\Gamma}|\}$ and $\zeta' \in \{0, \dots, |\bar{\Gamma}|\}$, the pair $(\gamma_\zeta, \gamma_{\zeta'})$ belongs to $E(G_{\bar{\Gamma}})$ if and only if $\chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}$ is a factor in one of $\zeta_{\beta_{\ell, \bar{\Gamma}}}$ for some $\beta_{\ell, \bar{\Gamma}}$ in $\mathbf{B}_{\bar{\Gamma}} = \Upsilon^{-1}(\bar{\Gamma})$, defined in Eq. (S.7.9).

REMARK S.7.1. The graph $G_{\bar{\Gamma}}$ can also be defined using only the $\Gamma = \{\gamma_1, \dots, \gamma_{|\bar{\Gamma}|}\}$ without explicit reference to the $\{\alpha_{\gamma_i}\}$. More specifically, for any $\gamma_\zeta, \gamma_{\zeta'}$, the term $\chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}$ is a factor in one of the $\zeta_{\beta_{\ell, \bar{\Gamma}}}$ if and only if:

- for $\gamma_\zeta \neq \gamma_0$ and $\gamma_{\zeta'} \neq \gamma_0$, there exist some τ, ℓ such that $(\ell, \tau) \in \gamma_\zeta$ and $(\ell, \tau + 1) \in \gamma_{\zeta'}$;
- for $\gamma_\zeta = \gamma_0$ and $\gamma_{\zeta'} \neq \gamma_0$, there exists some ℓ such that $(\ell, 1) \in \gamma_{\zeta'}$.

Given the above bijection between $\mathbf{B} = (\beta_1, \dots, \beta_g)$ and $\bar{\Gamma}_{\mathbf{B}}$, we can rewrite Eq. (S.7.6) as

$$\begin{aligned} \mathcal{X}^\iota &= \sum_{(\beta_1, \beta_2, \dots, \beta_\iota)} \zeta_{\beta_1} \zeta_{\beta_2} \cdots \zeta_{\beta_\iota} = \sum_{\bar{\Gamma}_{\mathbf{B}}} \zeta_{\beta_{1, \bar{\Gamma}_{\mathbf{B}}}} \zeta_{\beta_{2, \bar{\Gamma}_{\mathbf{B}}}} \cdots \zeta_{\beta_{\iota, \bar{\Gamma}_{\mathbf{B}}}} \\ &= \sum_{\bar{\Gamma}_{\mathbf{B}}} \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_{\mathbf{B}}})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_{\mathbf{B}}}) \setminus \{\gamma_0\}} h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|} \cdot \left(\prod_{\ell=1}^{\iota} \bar{u}_{\alpha_{g, \ell, \bar{\Gamma}_{\mathbf{B}}}}^{(2)} \right), \end{aligned}$$

where $\alpha_{g,\ell,\bar{\Gamma}_B}$ denote the g th α in $\beta_{\ell,\bar{\Gamma}_B}$.

Now let $\iota = \frac{(\log n)^\delta}{C_B C_1 s g}$ for some fixed $s, \delta > 0$. The constant C_B is specified later in Eq. (S.7.14). Without loss of generality we shall assume that ι is even (otherwise we let $\iota = \lceil \frac{(\log n)^\delta}{C_B C_1 s g} \rceil$ or $\iota = \lceil \frac{(\log n)^\delta}{C_B C_1 s g} \rceil + 1$ without affecting the subsequent argument). Now there exists a constant $N_2(C_1, \delta)$ (henceforth N_2) such that for any $n \geq N_2$ we have (as $\gamma_\zeta \subseteq [\iota] \times [g]$)

$$(S.7.10) \quad |\gamma_\zeta| \leq \iota g \leq (\log n)^{\delta+1} / (C_B C_1 s g) \leq (\log n)^{\delta+2},$$

We then have, for $n \geq \max\{N_1, N_2\}$,

$$(S.7.11) \quad \begin{aligned} |\mathbb{E}\mathcal{X}^\iota| &= \left| \sum_{\bar{\Gamma}_B} \mathbb{E} \left[\left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|} \cdot \left(\prod_{\ell=1}^{\iota} \bar{u}_{\alpha_{g,\ell,\bar{\Gamma}_B}}^{(2)} \right) \right] \right| \\ &= \left| \sum_{\bar{\Gamma}_B} \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \left(\prod_{\ell=1}^{\iota} \bar{u}_{\alpha_{g,\ell,\bar{\Gamma}_B}}^{(2)} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} \mathbb{E}[h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}] \right| \\ &= \left| \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \left(\prod_{\ell=1}^{\iota} \bar{u}_{\alpha_{g,\ell,\bar{\Gamma}_B}}^{(2)} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} \mathbb{E}[h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}] \right| \\ &\leq \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} \left| \mathbb{E}[h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}] \right| \\ &\leq \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} \mathbb{E}[|h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}|] \\ &\leq \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \prod_{\gamma_\zeta \in V(G_{\bar{\Gamma}_B}) \setminus \{\gamma_0\}} C_1^2/n \\ &\leq \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) (C_1^2/n)^{|\bar{\Gamma}_B|} \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \end{aligned}$$

where $\mathbb{I}(\bar{\Gamma}_B) = 1$ if $|\gamma_\zeta| \geq 2$ for all $\zeta \in [|\bar{\Gamma}_B|]$ and $\mathbb{I}(\bar{\Gamma}_B) = 0$ otherwise. Here $\gamma_1, \dots, \gamma_{|\bar{\Gamma}_B|}$ are the components of $\bar{\Gamma}_B$ (see Eq. (S.7.8)). The third equality in the above display is because, for a given $\bar{\Gamma}_B$, if there exists a γ_ζ such that $|\gamma_\zeta| = 1$ then $\mathbb{E}[h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}] = \mathbb{E}[h_{\alpha_{\gamma_\zeta}}] = 0$ as the entries of \mathbf{E} have mean 0. The first inequality is because $|\bar{u}_i| \leq 1$ for all $i \in [n]$ while the third inequality follows from Eq. (S.7.4); indeed, the non-zero summands in $\mathbb{E}[\mathcal{X}^\iota]$ all include terms of the form $\mathbb{E}[h_{\alpha_{\gamma_\zeta}}^{|\gamma_\zeta|}]$ for some $|\gamma_\zeta| \in [2, (\log n)^{\delta+2}]$ (see Eq. (S.7.10)).

It remains to bound

$$\sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) (C_1^2/n)^{|\bar{\Gamma}_B|} \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right)$$

Recall Eq. (S.7.8), i.e., any $\bar{\Gamma}_B$ include two components, namely the partition $\{\gamma_{\zeta, B}\}$ of $[\iota] \times [g]$ and the $\{\alpha_{\gamma_\zeta}\}$ associated with $\{\gamma_\zeta\}$. We therefore have

$$\begin{aligned} |\mathbb{E}\mathcal{X}^\iota| &\leq \sum_{\bar{\Gamma}_B} \mathbb{I}(\bar{\Gamma}_B) (C_1^2/n)^{|\bar{\Gamma}_B|} \left(\prod_{(\gamma_\zeta, \gamma_{\zeta'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_\zeta}, \alpha_{\gamma_{\zeta'}}} \right) \\ &= \sum_{\Gamma_B} \sum_{\{\alpha_{\gamma_\zeta, B}\}} \mathbb{I}(\Gamma_B, \{\alpha_{\gamma_\zeta, B}\}) (C_1^2/n)^{|\Gamma_B|} \prod_{(\gamma_{\zeta, B}, \gamma_{\zeta', B}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_{\zeta, B}}, \alpha_{\gamma_{\zeta', B}}} \end{aligned}$$

where the first sum is over all possible partitions $\Gamma_B = \{\gamma_{\varsigma, B}\}$ of $[l] \times [g]$ while the second sum is over all possible $\{\alpha_{\gamma_{\varsigma, B}}\}$ of α -assignments of Γ_B . We now bound

$$\sum_{\{\alpha_{\gamma_{\varsigma, B}}\}} \prod_{(\gamma_{\varsigma, B}, \gamma_{\varsigma', B}) \in E(G_{\Gamma_B})} \chi_{\alpha_{\gamma_{\varsigma, B}}, \alpha_{\gamma_{\varsigma', B}}}$$

for some arbitrary but fixed Γ_B . By Remark S.7.1, the $(\gamma_1, \dots, \gamma_{|\bar{\Gamma}_B|})$ corresponds to a multi-graph $G_{\bar{\Gamma}_B}$ and furthermore $G_{\bar{\Gamma}_B}$ is connected for any Γ_B . To see this, take γ_0 as the root node. Then any γ_{ς_1} for which $(\ell, 1) \in \gamma_{\varsigma_1}$ for some $\ell \in [l]$ will have an edge with γ_0 . Any γ_{ς_2} for which $(\ell, 2) \in \gamma_{\varsigma_2}$ for some $\ell \in [l]$ will have an edge to γ_{ς_1} . Continuing this argument we obtain a path between any $\gamma_{\varsigma} \in \{\gamma_1, \dots, \gamma_{|\bar{\Gamma}_B|}\}$ and $\gamma_{\varsigma'} \in \{\gamma_1, \dots, \gamma_{|\bar{\Gamma}_B|}\}$. There thus exists a spanning tree $\mathcal{T}_{\bar{\Gamma}_B} \subseteq G_{\bar{\Gamma}_B}$ rooted at α_0 . Now $\mathcal{T}_{\bar{\Gamma}_B} \subseteq G_{\bar{\Gamma}_B}$ satisfies

$$(S.7.12) \quad \prod_{(\gamma_{\varsigma}, \gamma_{\varsigma'}) \in E(G_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_{\varsigma}}, \alpha_{\gamma_{\varsigma'}}} = 1 \quad \text{only if} \quad \prod_{(\gamma_{\varsigma}, \gamma_{\varsigma'}) \in E(\mathcal{T}_{\bar{\Gamma}_B})} \chi_{\alpha_{\gamma_{\varsigma}}, \alpha_{\gamma_{\varsigma'}}} = 1.$$

We now study the behavior of $\chi_{\alpha_{\gamma_{\varsigma}}, \alpha_{\gamma_{\varsigma'}}$ for an arbitrary $\mathcal{T}_{\bar{\Gamma}}$. For a given ancestor γ_{ς_a} in $\mathcal{T}_{\bar{\Gamma}}$ with $\alpha_{\gamma_{\varsigma_a}}$, there are at most n choices of α -assignment for any of its descendants, say γ_{ς_d} , that can make $\chi_{\alpha_{\gamma_{\varsigma_a}}, \alpha_{\gamma_{\varsigma_d}}} = 1$. This is because $\alpha_{\gamma_{\varsigma_d}}^{(1)}$ must be $\alpha_{\gamma_{\varsigma_a}}^{(2)}$, and $\alpha_{\gamma_{\varsigma_d}}^{(2)}$ can be any of $1, \dots, n$. Now we count from the root node γ_0 in $\mathcal{T}_{\bar{\Gamma}}$. For each of γ_0 's descendants in $\mathcal{T}_{\bar{\Gamma}}$, there are at most n different α -assignment that make the corresponding $\chi_{\alpha_{\gamma_0}, \alpha} = 1$. Same argument also holds for the descendants of γ_0 's descendants. There are thus at most $n^{|\bar{\Gamma}|}$ different $(\alpha_1, \dots, \alpha_{|\bar{\Gamma}|})$ for which $\prod_{(\gamma_{\varsigma}, \gamma_{\varsigma'}) \in E(\mathcal{T}_{\bar{\Gamma}})} \chi_{\alpha_{\gamma_{\varsigma}}, \alpha_{\gamma_{\varsigma'}}} = 1$. Eq. (S.7.12) therefore implies

$$(S.7.13) \quad \sum_{\{\alpha_{\gamma_{\varsigma, B}}\}} \prod_{(\gamma_{\varsigma, B}, \gamma_{\varsigma', B}) \in E(G_{\Gamma_B})} \chi_{\alpha_{\gamma_{\varsigma, B}}, \alpha_{\gamma_{\varsigma', B}}} \leq \sum_{\{\alpha_{\gamma_{\varsigma, B}}\}} \prod_{(\gamma_{\varsigma, B}, \gamma_{\varsigma', B}) \in E(\mathcal{T}_{\Gamma_B})} \chi_{\alpha_{\gamma_{\varsigma, B}}, \alpha_{\gamma_{\varsigma', B}}} \leq n^{|\bar{\Gamma}_B|}.$$

Combining Eq. (S.7.4), Eq. (S.7.11) and Eq. (S.7.13), we have

$$|\mathbb{E}\mathcal{X}^\iota| \leq \sum_{\Gamma_B} \mathbb{I}(\Gamma_B, \{\alpha_{\gamma_{\varsigma, B}}\}) (C_1^2/n)^{|\bar{\Gamma}_B|} \cdot n^{|\bar{\Gamma}_B|} \leq \sum_{\Gamma_B} \mathbb{I}(\Gamma_B, \{\alpha_{\gamma_{\varsigma, B}}\}) (C_1 \log \log n)^{\iota g},$$

where the last inequality is because $C_1^2 \leq C_1^2 (\log \log n)^2$ when n is large. The number of all distinct partitions $(\gamma_1, \dots, \gamma_{|\bar{\Gamma}|})$, of $\{1, \dots, \iota\} \times \{1, \dots, g\}$ (under permutation equivalence) is the Bell number $\mathbb{B}_{g\iota}$. In particular we have

$$(S.7.14) \quad \mathbb{B}_{g\iota} < \left(\frac{C_B g \iota}{\log(g\iota + 1)} \right)^{g\iota}$$

where $C_B = 0.792$; see [13] for more details. We therefore have

$$|\mathbb{E}\mathcal{X}^\iota| \leq \left\{ \frac{C_B g \iota}{\log(g\iota + 1)} \right\}^{g\iota} \cdot \{C_1 (\log \log n)\}^{\iota g}.$$

Recall that we let $\iota = \frac{(\log n)^\delta}{C_B C_1 s g}$ for any fixed $s, \delta > 0$. Summarizing the results above, we have,

$$\begin{aligned} |\mathbb{E}\mathcal{X}^\iota| &= \mathbb{E}|\mathcal{X}^\iota| \leq \left\{ \frac{C_B g (\log n)^\delta / (C_B C_1 s g)}{\log(g\iota + 1)} \right\}^{\iota g} \cdot \{C_1 (\log \log n)\}^{\iota g} \\ &= \left\{ (\log n)^\delta / s \right\}^{\iota g} \cdot \left\{ \frac{\log \log n}{\log(g\iota + 1)} \right\}^{\iota g} \\ &\leq (\log n)^{\iota g \delta} / s^{\iota g}, \end{aligned}$$

provided that ι is assumed to be an even number without loss of generality, $n \geq N(C_1, \delta, A_0, \{q_n\}_{n \geq 1})$ (henceforth N) where N is a constant depending only on C_1, δ, A_0 and the divergence rate of the $\{q_n\}$. Note that the last inequality in the above display is because

$$\frac{\log \log n}{\log(g\nu + 1)} \leq \frac{\log \log n}{\log\{(\log n)^\delta / (C_B C_1 s) + 2\}} < 1,$$

for sufficiently large but finite n (as C_B, s, δ are fixed and $\delta > 1$). We therefore have, from Markov's inequality, that if $n \geq N$ then

$$\mathbb{P}[|\mathcal{X}| > (\log n)^{g\delta}] \leq \frac{\mathbb{E}|\mathcal{X}|^\iota}{(\log n)^{\iota g\delta}} \leq s^{-\iota g} = s^{-\frac{(\log n)^\delta}{C_B C_1 s}}.$$

Let $s = e$ and $\nu = 1/(C_B C_1 e)$, we derives, when $n \geq N$,

$$(S.7.15) \quad \begin{aligned} \mathbb{P}\left[|\mathbf{e}_{i^*}^\top \mathbf{E}^g \mathbf{u}| \leq (n\rho_n)^{g/2} (\log n)^{g\delta} \|\mathbf{u}\|_{\max}\right] &= 1 - \mathbb{P}[|\mathcal{X}| > (\log n)^{g\delta}] \\ &\geq 1 - \exp\{-\nu(\log n)^\delta\}. \end{aligned}$$

Now consider when \mathbf{E} is symmetrically independent. When $g = 1$, the argument is exactly same with the elementwise independent case. When $g \in [2, \log n]$, we can let $\mathbf{H} = \mathbf{H}_u + \mathbf{H}_d$ where ij th entry in \mathbf{H}_u is same with \mathbf{H} when $i \leq j$ and 0 otherwise; ij th entry in \mathbf{H}_d is same with \mathbf{H} when $i > j$ and 0 otherwise. We decompose

$$\mathbf{e}_{i^*}^\top \mathbf{H}^g \mathbf{u} = \mathbf{e}_{i^*}^\top \mathbf{H}_u^g \mathbf{u} + \mathbf{e}_{i^*}^\top \mathbf{H}_d \mathbf{H}_u^{g-1} \mathbf{u} + \cdots + \mathbf{e}_{i^*}^\top \mathbf{H}_d^g \mathbf{u},$$

where there are 2^g terms on the right-hand side above. Now it is easy to see \mathbf{H}_u is a random matrix with entrywise independence and the corresponding $\mathbf{E} = \mathbf{H}_u$ still satisfies (1)–(3) in Lemma 3.2. So we can bound $\mathbf{e}_{i^*}^\top \mathbf{H}_u^g \mathbf{u}$ similar to Eq. (S.7.15) with a modified upper bound $(n\rho_n)^{g/2} (\log n)^{g\delta} \|\mathbf{u}\|_{\max} / 2^g$. All other terms can be treated similarly. Consider, for example, the term $\mathbf{e}_{i^*}^\top \mathbf{H}_d \mathbf{H}_u^{g-1} \mathbf{u}$. Then, similar to Eq. (S.7.5), we have

$$\begin{aligned} \mathbf{e}_{i^*}^\top \mathbf{H}_d \mathbf{H}_u^{g-1} \mathbf{u} &= \sum_{(\alpha_1, \dots, \alpha_g) \in [n]^2 \times \cdots \times [n]^2} \chi_{\alpha_0 \alpha_1} \chi_{\alpha_1 \alpha_2} \cdots \chi_{\alpha_{g-1} \alpha_g} h'_{\alpha_1} h_{\alpha_2} \cdots h_{\alpha_g} \bar{u}_{\alpha_g^{(2)}} \\ &= \sum_{(\alpha_1, \dots, \alpha_g) \in [n]^2 \times \cdots \times [n]^2} \chi_{\alpha_0 \alpha_1} \chi_{\alpha_1 \alpha_2} \cdots \chi_{\alpha_{g-1} \alpha_g} h_{\bar{\alpha}_1} h_{\alpha_2} \cdots h_{\alpha_g} \bar{u}_{\alpha_g^{(2)}} \end{aligned}$$

where h'_{ij} and h_{ij} are the entries of \mathbf{H}_d and \mathbf{H}_u , respectively, and $\bar{\alpha}_1 = (\alpha_1^{(2)}, \alpha_1^{(1)})$ as $h'_{ij} = h_{ji}$ and hence $h'_{\alpha_1} = h_{\bar{\alpha}_1}$. We can thus bound $\mathbf{e}_{i^*}^\top \mathbf{H}_d \mathbf{H}_u^{g-1} \mathbf{u}$ using the same ‘‘partition-and-moment-bounding’’ arguments as that for the entrywise independent case described above (c.f. Eq. (S.7.15)). We only need a small modification of the criteria used for constructing a partition Γ_B over $[\iota] \times [g]$ in (S.7.7), i.e., for any $(\ell, \tau) \in [\iota] \times [g]$ and $(\ell', \tau') \in [\iota] \times [g]$, we say (ℓ, τ) and (ℓ', τ') belong to the same partition $\gamma_{\zeta, B} \in \Gamma_B$ if and only if,

$$\tilde{\alpha}_{\tau, \ell} = \tilde{\alpha}_{\tau', \ell'},$$

where $\tilde{\alpha}_{\tau, 1} := (\alpha_{\tau, 1}^{(2)}, \alpha_{\tau, 1}^{(1)})$ and $\tilde{\alpha}_{\tau, \ell} = \alpha_{\tau, \ell}$ when $\ell \geq 2$. Finally, recall each term has an upper bound of $(n\rho_n)^{g/2} (\log n)^{g\delta} \|\mathbf{u}\|_{\max} / 2^g$. Since $2^g \leq 2^{\log n} = n$, the bound for $\mathbf{e}_{i^*}^\top \mathbf{H}^g \mathbf{u}$ is the same as that in Eq. (S.7.15) even after taking a union bound over these 2^g terms, provided that we modify ν, N , and s accordingly. \square

S.7.3. *Proof of Claim S.1.* If $g = 2$ then $\mathbf{R}_4 = 0$. We thus consider only $g \geq 3$. Let $\tilde{\mathbf{G}}_g := \mathbf{G}\mathbf{V}_g\mathbf{W}_V\hat{\Sigma}_g^{-1}$. By the triangle inequality we have

$$(S.7.16) \quad \begin{aligned} \|\mathbf{R}_4\|_{2 \rightarrow \infty} &= \underbrace{\sum_{\xi=1}^{g-2} \|(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{E}\mathbf{M}^{g-1-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty}}_{R_{4,1}} \\ &+ \underbrace{\sum_{\xi=0}^{g-3} \sum_{\ell=1}^{g-2-\xi} \|(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\hat{\mathbf{M}}^\ell\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty}}_{R_{4,2}} \end{aligned}$$

We now bound $R_{4,1}$. We first observe that whp,

$$(S.7.17) \quad \begin{aligned} R_{4,1} &\leq \sum_{\xi=1}^{g-2} \|\mathbf{E}\mathbf{M}^{g-1-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} + \sum_{\xi=1}^{g-2} \|\mathbf{U}\mathbf{U}^\top\mathbf{E}\mathbf{M}^{g-1-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} \\ &\lesssim (E_n/S_n)^2(\log n)^{\delta+1/2}\|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

To see this, note that $g-1-\xi \geq 1$ for all $\xi \in [g-2]$ and hence, whp

$$(S.7.18) \quad \begin{aligned} \|\mathbf{E}\mathbf{M}^{g-1-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} &= \|\mathbf{E}\mathbf{U}\mathbf{A}^{g-1-\xi}\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\mathbf{V}_g\mathbf{W}_V\hat{\Sigma}_g^{-1}\|_{2 \rightarrow \infty} \\ &\leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{A}^{g-1-\xi}\| \|\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\| \|\hat{\Sigma}_g^{-1}\| \\ &\lesssim E_n(\log n)^\delta \|\mathbf{U}\|_{2 \rightarrow \infty} S_n^{g-1-\xi} E_n^\xi \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \tilde{k}^{-1/2} S_n^{-g} \\ &\lesssim (E_n/S_n)^2(\log n)^\delta \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}, \end{aligned}$$

where the first inequality follow from Lemma S.4.1, the second inequality follows from Lemmas S.4.3 and S.4.5, and the last inequality follows from Lemma S.4.4. Similarly, we have whp,

$$\begin{aligned} \|\mathbf{U}\mathbf{U}^\top\mathbf{E}\mathbf{M}^{g-1-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{E}\| \|\mathbf{A}^{g-1-\xi}\| \|\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\| \|\hat{\Sigma}_g^{-1}\| \\ &\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} (E_n/S_n)^{\xi+1} \tilde{k}^{-1/2} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \\ &\lesssim (E_n/S_n)^2 \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

We now bound $R_{4,2}$. For a given $0 \leq \xi \leq g-3$ and $1 \leq \ell \leq g-2$, denote

$$r_{4,2}(\ell, \xi) := (\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\hat{\mathbf{M}}^\ell\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g$$

As $(\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{M} = 0$, we have

$$(S.7.19) \quad \begin{aligned} r_{4,2}(\ell, \xi) &= (\mathbf{I} - \mathbf{U}\mathbf{U}^\top)(\mathbf{M} + \mathbf{E})\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \\ &= (\mathbf{I} - \mathbf{U}\mathbf{U}^\top)\mathbf{E}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \\ &= \hat{\mathbf{E}}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g - \mathbf{U}\mathbf{U}^\top\hat{\mathbf{E}}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g, \end{aligned}$$

In particular $r_{4,2}(1, \xi) = \mathbf{E}^2\mathbf{M}^{g-2-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g - \mathbf{U}\mathbf{U}^\top\mathbf{E}^2\mathbf{M}^{g-2-\xi}\mathbf{E}^\xi\tilde{\mathbf{G}}_g$.

If $\ell \geq 2$ then, by replacing $\mathbf{E}\hat{\mathbf{M}}^{\ell-1}$ with $\mathbf{E}^2\hat{\mathbf{M}}^{\ell-2} + \mathbf{E}\mathbf{M}\hat{\mathbf{M}}^{\ell-2}$, we obtain

$$\begin{aligned} r_{4,2}(\ell, \xi) &= \mathbf{E}\mathbf{M}\hat{\mathbf{M}}^{\ell-2}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g + \mathbf{E}^2\hat{\mathbf{M}}^{\ell-2}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \\ &\quad - \mathbf{U}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \\ &= \dots \\ &= \sum_{\zeta=1}^{\ell-1} \left\{ \mathbf{E}^\zeta\mathbf{M}\hat{\mathbf{M}}^{\ell-\zeta-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \right\} + \mathbf{E}^\ell\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g \\ &\quad - \mathbf{U}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g. \end{aligned}$$

We now bound the $\ell_{2 \rightarrow \infty}$ norm of $r_{4,2}(\ell, \xi)$ for each $\xi \leq g-3$ and $2 \leq \ell \leq g-2-\xi$. For any $\zeta \leq \ell-1$, using the same argument as that for deriving Eq. (S.7.18), we have (S.7.20)

$$\begin{aligned} \|\mathbf{E}^\zeta\mathbf{M}\hat{\mathbf{M}}^{\ell-\zeta-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}^\zeta\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}\| \|\hat{\mathbf{M}}\|^{\ell-\zeta-1} \|\mathbf{E}\| \|\mathbf{\Lambda}^{g-1-\xi-\ell}\| \|\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\| \|\hat{\Sigma}_g^{-1}\| \\ &\lesssim (E_n/S_n)^{\zeta+\xi+1} (\log n)^{\delta\zeta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty} \\ &\lesssim (E_n/S_n)^2 (\log n)^{\delta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}, \end{aligned}$$

whp, where the last inequality is because $\zeta = 1$ and $\xi = 0$ results in the largest upper bound for $(E_n/S_n)^{\zeta+\xi+1} (\log n)^{\delta\zeta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}$; indeed, other combinations of ξ, ζ lead to smaller order terms as $E_n/S_n \lesssim n^{-\beta_\Delta}$ for some $\beta_\Delta > 0$ under Assumption 1 which decays to 0 faster than the logarithmic factor $(\log n)^{\delta\zeta}$. Similarly, we also have

$$\begin{aligned} \|\mathbf{E}^{\ell+1}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}^{\ell+1}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{\Lambda}\|^{g-1-\xi-\ell} \|\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\| \|\hat{\Sigma}_g^{-1}\| \\ (S.7.21) \quad &\lesssim (E_n/S_n)^2 (\log n)^{2\delta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

Finally, for any $0 \leq \xi \leq g-3$ and $1 \leq \ell \in g-2-\xi$, we have

$$\begin{aligned} \|\mathbf{U}\mathbf{U}^\top\mathbf{E}\hat{\mathbf{M}}^{\ell-1}\mathbf{E}\mathbf{M}^{g-1-\xi-\ell}\mathbf{E}^\xi\tilde{\mathbf{G}}_g\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{E}\| \|\hat{\mathbf{M}}\|^{\ell-1} \|\mathbf{E}\| \|\mathbf{\Lambda}\|^{g-1-\xi-\ell} \|\mathbf{U}^\top\mathbf{E}^\xi\mathbf{G}\| \|\hat{\Sigma}_g^{-1}\| \\ (S.7.22) \quad &\lesssim \|\mathbf{U}\|_{2 \rightarrow \infty} (E_n/S_n)^{\xi+2} \vartheta \\ &\lesssim (E_n/S_n)^2 \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}. \end{aligned}$$

Combining Eq. (S.7.20), Eq(S.7.21), and Eq. (S.7.22), we have

$$\|r_{4,2}(\ell, \xi)\|_{2 \rightarrow \infty} \lesssim (E_n/S_n)^2 (\log n)^{2\delta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty}$$

whp. We therefore obtain

$$(S.7.23) \quad R_{4,2} \leq \sum_{\xi=0}^{g-3} \sum_{\ell=1}^{g-2-\xi} \|r_{4,2}(\ell, \xi)\|_{2 \rightarrow \infty} \lesssim (E_n/S_n)^2 (\log n)^{2\delta} \vartheta \|\mathbf{U}\|_{2 \rightarrow \infty},$$

whp. The desired bound for $\|\mathbf{R}_4\|$ follows directly from Eq. (S.7.17) and Eq. (S.7.23). \square

S.7.4. Proof of Claim S.2. Let $\tilde{\mathbf{R}}_6 = \mathbf{W}_\mathbf{V}\hat{\Sigma}_g^{-1} - \Sigma_g^{-1}\mathbf{W}_\mathbf{U}$. We then have

$$\begin{aligned} \|\tilde{\mathbf{R}}_6\| &\leq \|\mathbf{V}_g^\top\hat{\mathbf{V}}_g\hat{\Sigma}_g^{-1} - \Sigma_g^{-1}\mathbf{U}_g^\top\hat{\mathbf{U}}_g\| + \|(\mathbf{W}_\mathbf{V} - \mathbf{V}_g^\top\hat{\mathbf{V}}_g)\hat{\Sigma}_g^{-1}\| + \|\Sigma_g^{-1}(\mathbf{U}_g^\top\hat{\mathbf{U}}_g - \mathbf{W}_\mathbf{U})\| \\ &\leq \|\mathbf{V}_g^\top\hat{\mathbf{V}}_g\hat{\Sigma}_g^{-1} - \Sigma_g^{-1}\mathbf{U}_g^\top\hat{\mathbf{U}}_g\| + \|\hat{\Sigma}_g^{-1}\| \|\sin\Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|^2 + \|\Sigma_g^{-1}\| \|\sin\Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \end{aligned}$$

where the last inequality follows from Lemma 6.7 of [22]. Now recall Theorem 3.2 and Remark 3.3. Then for $g \geq 1 + (2\beta_\Delta)^{-1}$ and \tilde{k}, ψ are bounded, we have whp

$$(S.7.24) \quad \max \left\{ \|\sin\Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2, \|\sin\Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|^2 \right\} \lesssim (E_n/S_n)^2 \vartheta^2,$$

Furthermore, by Lemma S.4.5 we have

$$(S.7.25) \quad \|\hat{\Sigma}_g^{-1}\| = \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} S_n^{-g}, \quad \|\Sigma_g^{-1}\| = \sigma_k^{-1}(\mathbf{M}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} S_n^{-g},$$

whp. Combining Eq. (S.7.24)–Eq. (S.7.25), we have whp,

$$(S.7.26) \quad \|\mathbf{W}_V \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{W}_U\| \lesssim \|\mathbf{V}_g^\top \hat{\mathbf{V}}_g \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{U}_g^\top \hat{\mathbf{U}}_g\| + E_n^2 S_n^{-g-2} \vartheta^2 \tilde{k}^{-1/2}.$$

We now bound $\mathbf{V}_g^\top \hat{\mathbf{V}}_g \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{U}_g^\top \hat{\mathbf{U}}_g$. As $\hat{\mathbf{U}}_g^\top \hat{\mathbf{U}}_g = \mathbf{I}$ and $\mathbf{V}_g^\top \mathbf{V}_g = \mathbf{I}$, we have

$$\begin{aligned} \mathbf{V}_g^\top \hat{\mathbf{V}}_g \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{U}_g^\top \hat{\mathbf{U}}_g &= \mathbf{V}_g^\top (\hat{\mathbf{V}}_g \hat{\Sigma}_g^{-1} \hat{\mathbf{U}}_g^\top - \mathbf{V}_g \Sigma_g^{-1} \mathbf{U}_g^\top) \hat{\mathbf{U}}_g \\ &= \mathbf{V}_g^\top \left((\hat{\mathbf{M}}^g \mathbf{G})^\dagger - (\mathbf{M}^g \mathbf{G})^\dagger \right) \hat{\mathbf{U}}_g, \end{aligned}$$

where $(\hat{\mathbf{M}}^g \mathbf{G})^\dagger := \hat{\mathbf{V}}_g \hat{\Sigma}_g^{-1} \hat{\mathbf{U}}_g^\top$ and $(\mathbf{M}^g \mathbf{G})^\dagger = \mathbf{V}_g \Sigma_g^{-1} \mathbf{U}_g^\top$ are the Moore-Penrose pseudoinverse of $\hat{\mathbf{M}}^g \mathbf{G}$ and $\mathbf{M}^g \mathbf{G}$, respectively. By Theorem 3.2 in [111], we have,

$$(S.7.27) \quad \begin{aligned} \|(\hat{\mathbf{M}}^g \mathbf{G})^\dagger - (\mathbf{M}^g \mathbf{G})^\dagger\| &= \max \{ \|\hat{\Sigma}_g^{-1}\|^2, \|\Sigma_g^{-1}\|^2 \} \|\mathbf{M}^g \mathbf{G} - \hat{\mathbf{M}}^g \mathbf{G}\| \\ &\lesssim \tilde{k}^{-1} S_n^{-2g} \cdot \vartheta \tilde{k}^{1/2} S_n^{g-1} E_n = E_n S_n^{-g-1} \vartheta \tilde{k}^{-1/2}. \end{aligned}$$

Finally, combining Eq. (S.7.26) and Eq. (S.7.27), we have whp,

$$\|\mathbf{W}_V \hat{\Sigma}_g^{-1} - \Sigma_g^{-1} \mathbf{W}_U\| \lesssim E_n S_n^{-g-1} \vartheta \tilde{k}^{-1/2} + E_n^2 S_n^{-g-2} \vartheta^2 \tilde{k}^{-1/2} \asymp E_n S_n^{-g-1} \vartheta \tilde{k}^{-1/2},$$

where $E_n^2 S_n^{-g-2} \vartheta^2 \tilde{k}^{-1/2}$ is negligible as S_n/E_n is polynomially divergent. \square

S.7.5. *Proof of Claim S.3.* By Lemma S.4.2, we have, with probability at least $1 - n^{-6}$,

$$\begin{aligned} \|\Delta_g\| &= \|\mathbf{E}\|^g + \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} \|\hat{\mathbf{M}}\|^\ell \|\mathbf{E}\| \|\mathbf{M}\|^{g-1-\xi-\ell} \|\mathbf{E}\|^\xi + \sum_{\xi=0}^{g-2} \|\mathbf{M}\|^{g-1-\xi} \|\mathbf{E}\|^{\xi+1} \\ &\lesssim (n\rho_n)^{g/2} + \sum_{\xi=0}^{g-2} \sum_{\ell=0}^{g-2-\xi} (n\rho_n)^{g-1/2-\xi/2} + \sum_{\xi=0}^{g-2} (n\rho_n)^{g-1/2-\xi/2} \lesssim (n\rho_n)^{g-1/2}, \end{aligned}$$

where the first inequality follows from Lemma S.4.3.

S.7.6. *Proof of Claim S.4.* The current proof follows the same argument as that presented for the proof of Eq. (4.137) in [28]. First recall the matrix $\tilde{\mathbf{E}}$ defined in Eq. (S.6.27). Let $\mathbf{Z}_{ij} = [\tilde{\mathbf{E}}]_{ij} \left\{ [\mathbf{U}]_i [\mathbf{U}]_j^\top + [\mathbf{U}]_j [\mathbf{U}]_i^\top \right\}$. We then have

$$\mathbf{U}^\top \tilde{\mathbf{E}} \mathbf{U} = \sum_{i \geq j} \sum_{i \geq j} \mathbf{Z}_{ij}.$$

Next note that $(\mathbf{A} + \mathbf{A}^\top)^2 \lesssim 2\mathbf{A}\mathbf{A}^\top + 2\mathbf{A}^\top \mathbf{A}$ (where \lesssim denote the Lowner ordering for positive semidefinite matrices) and

$$(S.7.28) \quad \begin{aligned} \max_{i,j} \mathbb{E}[\tilde{\mathbf{E}}]_{ij}^2 &= p^{-2} \mathbb{E} \left\{ p([\mathbf{T}]_{ij} - p[\mathbf{T}]_{ij} + [\tilde{\mathbf{N}}]_{ij})^2 \right\} + (1-p)([\mathbf{T}]_{ij})^2 \\ &\leq p^{-1} \|\mathbf{T}\|_{\max}^2 + p^{-1} \mathbb{E}[\tilde{\mathbf{N}}]_{ij}^2 \leq p^{-1} (\|\mathbf{T}\|_{\max}^2 + \sigma^2). \end{aligned}$$

We therefore have

$$\begin{aligned} \mathbb{E}[\mathbf{Z}_{ij}^2] &\leq 2\mathbb{E}[\tilde{\mathbf{E}}]_{ij}^2 \left(\|\mathbf{U}\|_i^2 \|\mathbf{U}\|_j \|\mathbf{U}\|_j^\top + \|\mathbf{U}\|_j^2 \|\mathbf{U}\|_i \|\mathbf{U}\|_i^\top \right) \\ &\leq 2p^{-1} (\|\mathbf{T}\|_{\max}^2 + \sigma^2) \left(\|\mathbf{U}\|_i^2 \|\mathbf{U}\|_j \|\mathbf{U}\|_j^\top + \|\mathbf{U}\|_j^2 \|\mathbf{U}\|_i \|\mathbf{U}\|_i^\top \right) \end{aligned}$$

and hence

$$(S.7.29) \quad \begin{aligned} \left\| \sum_{i \geq j} \mathbb{E}[\mathbf{Z}_{ij}^2] \right\| &\leq 2p^{-1} (\|\mathbf{T}\|_{\max}^2 + \sigma^2) \left\| \sum_i \sum_j \|\mathbf{U}_i\|^2 \|\mathbf{U}_j\| \mathbf{U}_j^\top \right\| \\ &= 2kp^{-1} (\|\mathbf{T}\|_{\max}^2 + \sigma^2). \end{aligned}$$

On the other hand, by the definition of $\tilde{\mathbf{E}}$,

$$(S.7.30) \quad \begin{aligned} \max_{(i,j) \in [n]^2} \|\mathbf{Z}_{ij}\| &\leq 2 \max_{(i,j) \in [n]^2} |[\tilde{\mathbf{E}}]_{ij}| \cdot \max_{(i,j) \in [n]^2} \|\mathbf{U}_i\| \|\mathbf{U}_j\| \\ &\leq 2\|\tilde{\mathbf{E}}\|_{\max} \|\mathbf{U}\|_{2 \rightarrow \infty}^2 \leq \frac{2}{p} (\|\mathbf{T}\|_{\max} + 5\sigma\sqrt{\log n}) \|\mathbf{U}\|_{2 \rightarrow \infty}^2. \end{aligned}$$

We now apply the matrix Bernstein inequality [115, Theorem 1.4] and obtain

$$\|\mathbf{U}^\top \tilde{\mathbf{E}} \mathbf{U}\| = \left\| \sum_{i \geq j} \mathbf{Z}_{ij} \right\| \lesssim \sqrt{\frac{\log n}{p}} (\|\mathbf{T}\|_{\max} + \sigma) + \frac{\log n}{p} (\|\mathbf{T}\|_{\max} + \sigma\sqrt{\log n}) \|\mathbf{U}\|_{2 \rightarrow \infty}^2$$

whp. As $\mathbf{E} = \tilde{\mathbf{E}}$ whp, the above bound also holds for \mathbf{E} whp. \square

S.7.7. Proof of Claim S.5. We will bound $\|\mathbf{R}_i\|_{2 \rightarrow \infty}$, $\|\mathbf{R}_i\|$, and $\|\mathbf{E}\mathbf{R}_i\|_{2 \rightarrow \infty}$ for $1 \leq i \leq 7$. Combining these bound yield the desired results for $\mathbf{R} = \sum_i \mathbf{R}_i$. We first summarize a few frequently used bounds that were derived previously in the proof of Theorems 3.1–3.2. For any $g \geq 2 + (2\beta_\Delta)^{-1}$ we have whp

$$(S.7.31) \quad \|\mathbf{G}\| \lesssim \sqrt{n}, \quad \|\mathbf{G}\|_{\max} \lesssim \log^{1/2} n \quad \sigma_k^{-1}(\hat{\mathbf{M}}^g \mathbf{G}) \lesssim \tilde{k}^{-1/2} S_n^{-g}$$

$$(S.7.32) \quad \|\Delta_g\| \lesssim \vartheta \tilde{k}^{1/2} S_n^{g-1} E_n, \quad \|\Delta_g\|_{2 \rightarrow \infty} \lesssim (\log n)^\delta \vartheta \tilde{k}^{1/2} S_n^{g-1} E_n \|\mathbf{U}\|_{2 \rightarrow \infty}$$

$$(S.7.33) \quad \max \left\{ \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|, \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\| \right\} \lesssim \vartheta \frac{E_n}{S_n}$$

$$(S.7.34) \quad \max \left\{ \|\mathbf{U}^\top \mathbf{G}\|, \max_{\xi \leq g-1} \|\tilde{\mathbf{V}}_\xi^\top \mathbf{G}\| \right\} \lesssim \vartheta \tilde{k}^{1/2}$$

$$(S.7.35) \quad \|\mathbf{U}^\top \mathbf{E}^{g'} \mathbf{G}\| \lesssim \vartheta \tilde{k}^{1/2} E_n^{g'}, \quad \text{for all } g' \geq 1$$

$$(S.7.36) \quad \|\mathbf{E}^{g'} \mathbf{G}\|_{2 \rightarrow \infty} \lesssim (\log n)^{\delta g' + 1/2} \tilde{k}^{1/2} E_n^{g'}, \quad \text{for all } g' \leq g$$

$$(S.7.37) \quad \|\mathbf{E}^{g'} \mathbf{U}\|_{2 \rightarrow \infty} = \|\mathbf{E}^{g'} \mathbf{U}_g\|_{2 \rightarrow \infty} \lesssim (\log n)^{\delta g'} E_n^{g'} \|\mathbf{U}\|_{2 \rightarrow \infty}, \quad \text{for all } g' \leq g$$

$$(S.7.38) \quad \|\mathbf{E} \Delta_g\|_{2 \rightarrow \infty} \lesssim (\log n)^{2\delta} \vartheta \tilde{k}^{1/2} E_n^2 S_n^{g-1} \|\mathbf{U}\|_{2 \rightarrow \infty}.$$

Eq. (S.7.31) is from Lemma S.4.5 and Lemma S.4.4, while Eq. (S.7.32) is from Theorem 3.1 and Eq. (S.5.16). Eq. (S.7.33) is from Theorem 3.2 and Remark 3.3, while Eq. (S.7.34) and Eq. (S.7.35) are from Lemma S.4.4 and Lemma S.4.3, respectively. Eq. (S.7.36) and Eq. (S.7.37) follow from Eq. (3.6) and Proposition 3.1, and finally Eq. (S.7.38) follows from Eq. (S.5.35),

We now proceed with the bounds for \mathbf{R}_i and $\mathbf{E}\mathbf{R}_i$.

R₁: Recall Eq. (S.5.18). We then have whp

$$\begin{aligned}
\|\mathbf{R}_1\|_{2 \rightarrow \infty} &\leq (\|\mathbf{\Delta}_g\|_{2 \rightarrow \infty} + \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Delta}_g\|) \cdot \frac{\sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \\
&\lesssim (\log n)^\delta \vartheta^2 \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
\|\mathbf{R}_1\| &\leq \|\mathbf{\Delta}_g\| \cdot \frac{\sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \lesssim \vartheta^2 \left(\frac{E_n}{S_n}\right)^2, \\
\|\mathbf{ER}_1\|_{2 \rightarrow \infty} &\leq (\|\mathbf{E}\mathbf{\Delta}_g\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\mathbf{\Delta}_g\|) \cdot \frac{\sqrt{2} \|\sin \Theta(\hat{\mathbf{V}}_g, \mathbf{V}_g)\|}{\sigma_k(\hat{\mathbf{M}}^g \mathbf{G})} \\
&\lesssim (\log n)^{2\delta} \vartheta^2 \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty}.
\end{aligned}$$

R₂: $\|\mathbf{R}_2\| = \|\mathbf{R}_2\|_{2 \rightarrow \infty} = \|\mathbf{ER}_2\|_{2 \rightarrow \infty} = \mathbf{0}$; see Eq. (S.5.19).

R₃: Recall Eq. (S.5.20). We then have whp

$$\begin{aligned}
\|\mathbf{R}_3\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \lesssim \vartheta^2 \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
\|\mathbf{R}_3\| &\leq \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \lesssim \vartheta^2 \left(\frac{E_n}{S_n}\right)^2, \\
\|\mathbf{ER}_3\|_{2 \rightarrow \infty} &\lesssim \|\mathbf{E}\mathbf{U}_g\|_{2 \rightarrow \infty} \cdot \|\sin \Theta(\hat{\mathbf{U}}_g, \mathbf{U}_g)\|^2 \lesssim (\log n)^\delta \vartheta^2 \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty}.
\end{aligned}$$

R₄: The bound for $\|\mathbf{R}_4\|_{2 \rightarrow \infty}$ is given in Claim S.1 while bounds for $\|\mathbf{R}_4\|_{2 \rightarrow \infty}$ and $\|\mathbf{ER}_4\|_{2 \rightarrow \infty}$ can be extracted from the proof of Claim S.1 (see below). For brevity we omit the details.

$$\begin{aligned}
\|\mathbf{R}_4\|_{2 \rightarrow \infty} &\lesssim (\log n)^{2\delta} \vartheta \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
\|\mathbf{R}_4\| &\lesssim \vartheta \left(\frac{E_n}{S_n}\right)^2, \quad \|\mathbf{ER}_4\|_{2 \rightarrow \infty} \lesssim (\log n)^{3\delta} \vartheta \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty}.
\end{aligned}$$

R₅: Recall Eq. (S.5.39) and Eq. (S.5.40). We then have whp

$$\begin{aligned}
\|\mathbf{R}_5\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}^g \mathbf{G}\|_{2 \rightarrow \infty} \|\hat{\mathbf{\Sigma}}_g^{-1}\| + \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| \|\hat{\mathbf{\Sigma}}_g^{-1}\| \lesssim (\log n)^{\delta^{**}} \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
\|\mathbf{R}_5\| &\leq \|\mathbf{E}^g \mathbf{G}\| \|\hat{\mathbf{\Sigma}}_g^{-1}\| + \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| \|\hat{\mathbf{\Sigma}}_g^{-1}\| \lesssim (\log n)^{1/2} \left(\frac{E_n}{S_n}\right)^2, \\
\|\mathbf{ER}_5\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}^{g+1} \mathbf{G}\|_{2 \rightarrow \infty} \|\hat{\mathbf{\Sigma}}_g^{-1}\| + \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}^g \mathbf{G}\| \|\hat{\mathbf{\Sigma}}_g^{-1}\| \lesssim (\log n)^{\delta^{**} + \delta} \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty},
\end{aligned}$$

where δ^{**} is defined in Eq. (S.5.41).

R₆: Recall Eq. (S.5.44) and Claim S.2. We then have whp

$$\begin{aligned}
 \|\mathbf{R}_6\|_{2 \rightarrow \infty} &\leq (\|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} + \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|) \|\mathbf{\Lambda}\|^{g-1} \|\mathbf{U}^\top \mathbf{G}\| \|\tilde{\mathbf{R}}_6\| \\
 &\lesssim (\log n)^\delta \vartheta^2 \left(\frac{E_n}{S_n}\right)^2 \|\mathbf{U}\|_{2 \rightarrow \infty} + \vartheta^2 \frac{E_n \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty}, \\
 \|\mathbf{R}_6\| &\leq (\|\mathbf{E}\mathbf{U}\| + \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|) \|\mathbf{\Lambda}\|^{g-1} \|\mathbf{U}^\top \mathbf{G}\| \|\tilde{\mathbf{R}}_6\| \lesssim \vartheta^2 \left(\frac{E_n}{S_n}\right)^2 + \vartheta^2 \frac{E_n \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n^2}, \\
 \|\mathbf{E}\mathbf{R}_6\|_{2 \rightarrow \infty} &\leq (\|\mathbf{E}^2 \mathbf{U}\|_{2 \rightarrow \infty} + \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|) \|\mathbf{\Lambda}\|^{g-1} \|\mathbf{U}^\top \mathbf{G}\| \|\tilde{\mathbf{R}}_6\| \\
 &\lesssim (\log n)^{2\delta} \vartheta^2 \frac{E_n^3}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty} + (\log n)^\delta \vartheta^2 \frac{E_n^2 \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n^2} \|\mathbf{U}\|_{2 \rightarrow \infty}.
 \end{aligned}$$

R₇: Recall Eq. (S.5.45). We then have whp

$$\begin{aligned}
 \|\mathbf{R}_7\|_{2 \rightarrow \infty} &\leq \|\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \|\mathbf{\Lambda}^{-1}\| \lesssim \frac{\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty} \\
 \|\mathbf{R}_7\| &\leq \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \|\mathbf{\Lambda}^{-1}\| \lesssim \frac{\|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n} \\
 \|\mathbf{E}\mathbf{R}_7\|_{2 \rightarrow \infty} &\leq \|\mathbf{E}\mathbf{U}\|_{2 \rightarrow \infty} \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\| \|\mathbf{\Lambda}^{-1}\| \lesssim (\log n)^\delta \frac{E_n \|\mathbf{U}^\top \mathbf{E}\mathbf{U}\|}{S_n} \|\mathbf{U}\|_{2 \rightarrow \infty}. \quad \square
 \end{aligned}$$