ON WALL-CROSSING INVARIANCE OF CERTAIN SUMS OF WELSCHINGER NUMBERS

S. FINASHIN, V. KHARLAMOV

ABSTRACT. We continue our quest for real enumerative invariants not sensitive to changing the real structure and extend the construction we uncovered previously for counting curves of anti-canonical degree ≤ 2 on del Pezzo surfaces with $K^2 = 1$ to curves of any anti-canonical degree and on any del Pezzo surfaces of degree $K^2 \leq 3$.

The obvious answer is always overlooked

Known as Whitehead's Law

1. INTRODUCTION

1.1. **Problem formulation.** Up to 2000s it was a rather common opinion that there can not exist any reasonable integer valued enumerative geometry over the reals, and that it is a prerogative of geometry over the field of complex numbers and other algebraically closed fields. The situation is radically changed in 2003, when J.-Y. Welschinger invented an integer signed count of point-constrained real rational curves on real rational surfaces that respects the necessary invariance property to be preserved under equivariant deformations and equivariant isomorphisms.

Those Welschinger invariants, like most of their subsequent analogs, turned out to be sensitive to changes that a real structure experiences under *wall-crossing*. However, quite soon after Welschinger's discovery, some examples appeared where under an appropriate counting scheme the invariance under wall-crossing holds. Initially, it was a signed count of real lines on real projective hypersurfaces (see [S1], [Wal], [O-T], [FK-1]), which includes as the starting case the count of real lines on real cubic surfaces in real projective 3-space.

It is the latter signed count on cubic surfaces that was recently extended by us to counting curves of anticanonical degree 1 and 2 on del Pezzo surfaces of degree $K^2 = 1$ (see [FK-2], [FK-3]). The invariance under wall-crossing was achieved there by combining the original Welschinger invariants with a certain intrinsic Pin⁻-structure we attributed to real loci of these surfaces.

This led us to a general question: What kind of real rational surfaces X carry similar Pin^- -structures on $X_{\mathbb{R}}$ and what are (anticanonical) degrees for which the real rational curves in X can be counted invariantly under wall-crossing?

In this paper we answer this question for del Pezzo surfaces of degree $K^2 \leq 3$.

²⁰²⁰ Mathematics Subject Classification. Primary: 14N10. Secondary: 14P25, 14J26, 14N15, 53D45.

Key words and phrases. Real del Pezzo surfaces, Pin-structures, Real enumerative geometry, Welschinger invariants, Wall-crossing.

1.2. Main results. Let X be a real del Pezzo surface of degree $K^2 \leq 3$ and $\operatorname{conj} : X \to X$ the complex conjugation. Denote by $\operatorname{Eff}(X) \subset \operatorname{Pic}(X) = H_2(X)$ the semigroup of effective divisor classes, and put $\operatorname{Eff}_{\mathbb{R}}(X) = \operatorname{Eff}(X) \cap \ker(1 + \operatorname{conj}_*)$. If $X_{\mathbb{R}} \neq \emptyset$ (which is always the case if K^2 is odd), the latter semigroup coincides with the semigroup of divisor classes that can be realized by a real effective divisor.

For del Pezzo surfaces, Eff(X) and $\text{Eff}_{\mathbb{R}}(X)$ are both finitely generated. For $K^2 = 2$ and 3, they are generated by *lines*, which are, by definition, embedded (-1)-curves of genus 0 (defined over the reals if we speak of *real lines*). For $K^2 = 1$, these semigroups have one additional generator, -K.

We split both Eff(X) and $\text{Eff}_{\mathbb{R}}(X)$, into subsets called *layers*:

$$\mathcal{L}^{m}(X) = \{ \alpha \in \text{Eff}(X) \mid -\alpha K = m \}, \quad \mathcal{L}^{m}_{\mathbb{R}}(X) = \{ \alpha \in \text{Eff}_{\mathbb{R}}(X) \mid -\alpha K = m \}.$$

All the layers are finite sets. They are empty if m is not a positive integer.

For each pair of integers m, k with $0 \leq k \leq m-1, k = m-1 \mod 2$, and a collection $\mathbf{x} \subset X$ of m-1 points including k real ones and $\frac{1}{2}(m-k-1)$ pairs of complex conjugate imaginary points, we consider the set $\mathcal{C}_{\mathbb{R}}(m, k, \mathbf{x})$ of real rational irreducible reduced curves $A, [A] \in \mathcal{L}^m_{\mathbb{R}}(X), \mathbf{x} \subset A$ that pass through \mathbf{x} . The sets $\mathcal{C}_{\mathbb{R}}(m, k, \mathbf{x})$ are all finite for a generic choice of \mathbf{x} .

The main object of this paper is the following double sequence of integers,

(1.2.1)
$$N_{m,k} = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x})} i^{\hat{q}([A]) - m^2} w(A),$$

where $\hat{q} : \text{Eff}_{\mathbb{R}}(X) \to \mathbb{Z}/4$ is a certain quadratic function that X inherits from its natural embeddings in an appropriate 3-fold (see Section 2) and w(A) is the modified Welschinger number $w(A) = (-1)^{c_A}$ in which c_A denotes the number of cross-point real nodes in the real locus of A.

As is known, for any $\alpha \in \mathcal{L}^m_{\mathbb{R}}(X)$ a partial sum $W_{\alpha,k} = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x}), [A]=\alpha} w(A)$, which

we call the modified *Welschinger invariant*, does not depend on a generic choice of \mathbf{x} (see [Br]) and is invariant under real deformations of X (see [Wel1]). The same invariance of $N_{m,k}$ now follows from the real deformation invariance of $\hat{\mathbf{q}}$ (see Section 2).

Some precaution must be however taken in the case $K^2 = 1$, m = 1. Namely, to have such invariance for numbers $W_{-K,0}$ and thus, for $N_{1,0}$, we need, in addition to genericness of \mathbf{x} , assume also genericness of X (see [IKS-3]).

Recall that over \mathbb{C} all del Pezzo surfaces with a fixed K^2 are deformation equivalent to each other, while over \mathbb{R} two real del Pezzo surfaces having the same K^2 are real deformation equivalent if, and only if, their real structures conj : $X \to X$ are diffeomorphic.

The goal of this paper is to prove that the sequence $N_{m,k}$ has the following strong invariance property.

1.2.1. **Theorem.** The double sequence $N_{m,k}$ is the same for all real del Pezzo surfaces X with $X_{\mathbb{R}} \neq \emptyset$ having a given degree $K^2 \leq 3$.

Since for any $k \ge 2$ and any nonsingular real rational surface X with disconnected real locus $X_{\mathbb{R}}$ the Welschinger invariants $W_{\alpha,k}$ vanish (see [Br]), and since for any $K^2 \in \{1, 2, 3\}$ there exist real del Pezzo surfaces with disconnected $X_{\mathbb{R}}$, Theorem 1.2.1 implies the following vanishing for $N_{m,k}$.

1

1.2.2. Corollary. For all real del Pezzo surfaces as in Theorem 1.2.1, each of the numbers $N_{m,k}$ with $k \ge 2$ is equal to 0.

The numbers $N_{m,k}$ happen to have remarkable properties. In particular, for k = 1 they turn out to be related in a "magic way" with Gromov-Witten invariants.

1.2.3. **Theorem.** For any $m \in \mathbb{Z}_{\geq 1}$ and any real surface X as in Theorem 1.2.1,

(1.2.2)
$$N_{2m,1} = 2^{m-3} \sum_{\alpha \in \mathcal{L}^m(X)} (e\alpha)^2 G W_{\alpha}$$

where e is an arbitrary class $e \in K^{\perp}$ with $e^2 = -2$ and GW_{α} states for the Gromov-Witten invariant which counts rational irreducible reduced curves in class $\alpha \in \mathcal{L}^m(X)$ passing through a generic collection of m-1 points.

Besides giving a way to calculate the numbers $N_{2m,1}$, this result plays a crucial role in deriving simple recursive formulas governing the both sequences, $N_{2m,1}$ and $N_{2m+1,0}$.

1.2.4. **Theorem.** For any $n \in \mathbb{Z}_{\geq 1}$ and any real surface X as in Theorem 1.2.1,

$$mK^{2}N_{m,0} = 2\sum_{j=1}^{n} \binom{n-1}{n-j} j(m-2j)^{2} N_{m-2j,0} N_{2j,1} \quad for \ m = 2n+1,$$

$$mK^{2}N_{m,1} = 2\sum_{j=1}^{n} \binom{n-1}{n-j} j(m-2j)^{2} N_{m-2j,1} N_{2j,1} \quad for \ m = 2n+2.$$

These recursion relations allow to reconstruct the numbers $N_{m,k}$ from the initial values (see Section 5.5) and to observe their non-vanishing and positivity (except the case of $N_{2n+1,0} = 0$ for $K^2 = 2$).

1.2.5. *Remark.* In the case $K^2 = 2$, the numbers $N_{m,k}$ vanish for odd m (see Proposition 2.1.2), and so in this case the first formula in Theorem 1.2.4 holds for trivial reasons.

In Theorem 5.4.1, we solve the above recurrence relations and get the following explicit formulas:

$$N_{2n+1,0} = \frac{1}{4} N_{1,0} b^n \left(n + \frac{1}{2}\right)^{n-2}, \qquad N_{2n+2,1} = N_{2,1} b^n \left(n + 1\right)^{n-2}, \qquad b = \frac{4N_{2,1}}{K^2}$$

As an immediate consequence of these expressions, we evaluate the growth rate of the sequences $N_{2n+1,0}$ and $N_{2n+2,1}$ and compare it with the growth rate of an analogous sequence of Gromov-Witten numbers (see Corollary 6.2.2).

1.2.6. Remark. In the case of X with $K^2 = 2$ and $X_{\mathbb{R}} = \emptyset$, one can prove that $\mathcal{C}_{\mathbb{R}}(m,k,x) = \emptyset$ for any m,k, and so, for such X, all the numbers $N_{m,k}$ vanish. This shows that Theorem 1.2.1 can not be extended to this case.

1.3. Plan of the paper. Section 2 starts from constructing of natural, basic for our counting scheme, Pin⁻-structures and establishing their main properties. We conclude this preliminary section by a discussion of wall-crossing and precise the limit behavior of the curves involved into counting of $N_{2m,1}$ in the case of contracting a spherical component of $X_{\mathbb{R}}$. Section 3 is devoted to a proof of the central result, Theorem 1.2.1. Theorem 1.2.3 is proved in Section 4. In Section 5 we prove theorems 1.2.4, 5.4.1 and discuss a few simple concrete applications. In Section 6 we are making few remarks on generating functions, on comparison of our count with a similar count for Gromov-Witten invariants, and on a situation with other, $K^2 > 3$, del Pezzo surfaces.

1.4. Acknowledgements. The idea of this work arose during a stay of the second author at the Max-Planck Institute for Mathematics in Bonn in spring 2021 and took its shape during a RIP-stay of the authors at the Mathematisches Forschungsinstitut Oberwolfach in summer 2021. We thank the both institutions for hospitality and excellent (despite a complicated pandemic situation) working conditions.

Our special thanks go to R. Rasdeaconu, discussions with whom were among the motivations for this study, and to J. Solomon, for encouragement and helpful remarks.

The second author was partially supported by the grant ANR-18-CE40-0009 of French Agence Nationale de Recherche and the grant by Ministry of Science and Higher Education of Russia under the contract 075-15-2019-1620 with St. Petersburg Department of Steklov Mathematical Institute.

2. Preliminaries

From now on, we denote the anticanonical degree K^2 by d.

2.1. **Basic** Pin⁻ structures. Recall that each del Pezzo surface X of degree d = 3 has a natural anticanonical embedding $X \hookrightarrow \mathbb{P}^3$ representing it as a non-singular cubic surface. For del Pezzo surfaces of degree d = 2, the anticanonical map is a double covering $X \to \mathbb{P}^2$ branched along a non-singular quartic curve, and its deck transformation $\gamma: X \to X$ is called *Geiser involution*. This covering lifts naturally to an embedding $X \hookrightarrow \mathbb{P}(1, 1, 1, 2)$ into the weighted projective space $\mathbb{P}(1, 1, 1, 2)$. For del Pezzo surfaces of degree d = 1, the bi-anticanonical map $X \to \mathbb{P}^3$ represents X as a double covering of a non-degenerate quadratic cone $Q \subset \mathbb{P}^3$ branched at its vertex and along a non-singular sextic $C \subset Q$ (traced on Q by a transversal cubic surface). The deck transformation $\tau: X \to X$ in this case is known as the *Bertini involution*. The above covering $X \to Q$ lifts to an embedding $X \hookrightarrow \mathbb{P}(1, 1, 2, 3)$ provided by the graded anti-canonical ring $R = \sum_{m \ge 0} H^0(X; -mK)$. In each case d = 1, 2, 3, these embeddings of X are defined uniquely up to automorphisms of the ambient 3-space (see, for example, [Do]).

These constructions are exhaustive, functorial, and work equally well over \mathbb{C} and \mathbb{R} . In particular, over \mathbb{R} we obtain a natural embedding of the real locus $X_{\mathbb{R}}$ into \mathbb{RP}^3 if d = 3, into $\mathbb{RP}^2 \times \mathbb{R} = \mathbb{P}_{\mathbb{R}}(1, 1, 1, 2) \setminus \mathfrak{v}_q, \mathfrak{v}_q = (0, 0, 0, 1)$ if d = 2, and into $\mathbb{RP}^2 \times \mathbb{R} = \mathbb{P}_{\mathbb{R}}(1, 1, 2, 3) \setminus \mathfrak{v}_p, \mathfrak{v}_p = (0, 0, 1, 0)$ if d = 1 (see [FK-2]).

Therefore, in each case d = 1, 2, 3, the real locus $X_{\mathbb{R}}$ inherits from the ambient 3-space some natural Pin⁻-structures. Namely, if d = 3, we have a pair of Spinstructures in \mathbb{RP}^3 which differ by a shift by $h \in H^1(\mathbb{RP}^3; \mathbb{Z}/2), h \neq 0$. They induce on $X_{\mathbb{R}} \subset \mathbb{RP}^3$ a pair of Pin⁻-structures, θ^X and $\theta^X + w_1$, where $w_1 = w_1(X_{\mathbb{R}})$ is the pull-back of h. If d = 1 or 2, the above embeddings $X_{\mathbb{R}} \subset \mathbb{RP}^2 \times \mathbb{R}$ are twosided, and thus a pair of Pin⁻-structures on $\mathbb{RP}^2 \times \mathbb{R}$ (which differ by a generator in $H^1(\mathbb{RP}^2 \times \mathbb{R}; \mathbb{Z}/2)$) descends to a pair of Pin⁻-structures on $X_{\mathbb{R}}$, also denoted θ^X and $\theta^X + w_1$ (since also differ by a shift by $w_1 = w_1(X_{\mathbb{R}})$).

Recall that there is a canonical correspondence between Pin⁻-structures θ on $X_{\mathbb{R}}$ and *quadratic functions* $q_{\theta} : H_1(X_{\mathbb{R}}; \mathbb{Z}/2) \to \mathbb{Z}/4$, that is the functions satisfying $q_{\theta}(x+y) = q_{\theta}(x) + q_{\theta}(y) + 2(x,y) \mod 4$. These functions can be viewed as $\mathbb{Z}/4$ liftings of w_1 seen as a homomorphism $w_1 : H_1(X; \mathbb{Z}/2) \to \mathbb{Z}/2$. In particular, it implies that $q_{\theta+w_1} = -q_{\theta}$ (cf. [FK-2, Lemma 2.4.1]).

Thus, in the case d = 1 or 3, we distinguish the Pin⁻-structure θ^X from $\theta^X + w_1$ by requiring that θ^X is monic that is $q_{\theta^X}(w_1^*) = 1$, where $w_1^* \in H_1(X_{\mathbb{R}}; \mathbb{Z}/2)$ is dual to w_1 (then $q_{\theta^X+w_1} = -q_{\theta^X}$ takes on w_1^* value -1). We call such θ^X basic Pin⁻-structure on $X_{\mathbb{R}}$. In the case d = 2, there is no natural way to distinguish θ^X from $\theta^X + w_1$ and we call basic both of them. Note that our definition of basic structures is independent of the choice of a graded anticanonical embedding of $X_{\mathbb{R}}$.

As in [FK-2], we consider also the function

$$\hat{q}_{\theta}: H_2^-(X) \to \mathbb{Z}/4, \quad \hat{q}_{\theta} = q_{\theta} \circ \Upsilon$$

where $H_2^-(X) = \ker(1 + \operatorname{conj}_* : H_2(X) \to H_2(X))$ and $\Upsilon : H_2^-(X) \to H_1(X_{\mathbb{R}}; \mathbb{Z}/2)$ is the Viro homomorphism (see [DIK, Chapter 1] or [FK-2]). This homomorphism respects the intersection form, so that \hat{q}_{θ} inherits from q_{θ} the property

$$\hat{q}_{\theta}(x+y) = \hat{q}_{\theta}(x) + \hat{q}_{\theta}(y) + 2(x,y) \mod 4.$$

The Viro homomorphism Υ induces an isomorphism

$$H_1(X_{\mathbb{R}}; \mathbb{Z}/2) \cong H_2^-(X)/(1 - \operatorname{conj}_*)H_2(X),$$

and has a simple interpretation in differential topology setting. Namely, for any real del Pezzo surface (and, more generally, for any compact complex surface Xwith a real structure and $H_1(X; \mathbb{Z}/2) = 0$) each class $\alpha \in H_2^-(X)$ can be realized by a conj-invariant smoothly embedded oriented 2-manifold $F \subset X$ and then it is $F \cap X_{\mathbb{R}}$, which is a collection (may be empty) of smooth circles in $X_{\mathbb{R}}$, that realizes the class $\Upsilon(\alpha)$. Furthermore, if $\Upsilon(\alpha) = 0$ then α can be represented by a conj-invariant smoothly embedded oriented 2-manifold disjoint from $X_{\mathbb{R}}$, and if $\Upsilon(\alpha) \neq 0$ then $F \subset X$ can be chosen in such a way that each of the components of $F \cap X_{\mathbb{R}}$ represents a non-zero element in $H_1(X_{\mathbb{R}}; \mathbb{Z}/2)$.

2.1.1. **Theorem.**

- Real automorphisms and real deformations preserve the basic Pin⁻-structure (resp. the pair of basic Pin⁻-structures) of del Pezzo surfaces of degree 1 and 3 (resp. of degree 2).
- (2) The quadratic functions $q_{\theta X}$ of basic Pin⁻-structures θ^X vanish on each real vanishing cycle in $H_1(X_{\mathbb{R}}; \mathbb{Z}/2)$.

Proof. (1) holds because real automorphisms and real deformations preserve w_1 and any real automorphism of X is induced by a real automorphism of the ambient 3-space. Property (2) is a special case of [FK-2, Lemma 2.4.2].

2.1.2. **Proposition.** For d = 2 and any of the two basic Pin^- -structures θ^X , the following holds:

(1) The functions \hat{q} and q associated with θ^X are skew-symmetric with respect to γ and its restriction $\gamma_{\mathbb{R}}: X_{\mathbb{R}} \to X_{\mathbb{R}}$, correspondingly. That is

$$q \circ (\gamma_{\mathbb{R}})_* = -q, \qquad \hat{q} \circ (\gamma)_* = -\hat{q}.$$

(2) For each pair $(m,k) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{>0}$, $k = m-1 \mod 2$, the number $N_{m,k}$ does not depend on the choice of θ^X . If, in addition, m is odd, then $N_{m,k} = 0$.

Proof. (1) The reflection map $\mathbb{RP}^2 \times \mathbb{R} \to \mathbb{RP}^2 \times \mathbb{R}$ given by $(x, y, z, t) \mapsto (x, y, z, -t)$ shifts each of Pin⁻-structures on $\mathbb{RP}^2 \times \mathbb{R}$ by $w_1(\mathbb{RP}^2 \times \mathbb{R})$ (cf. [K-T, Lemma 1.10]). Since $\gamma_{\mathbb{R}}$ is the restriction of this reflection to $X_{\mathbb{R}}$, it interchanges the two basic Pin⁻-structures induced on $X_{\mathbb{R}}$, and hence transforms q into -q.

Skew-symmetry of \hat{q} follows from that of q.

(2) Suppose first that m = 2n + 1. Let **x** be a generic collection of k real points and $\frac{1}{2}(m - k - 1)$ pairs of complex conjugate imaginary points and let A, $[A] \in \mathcal{L}^m_{\mathbb{R}}(X)$, be a real rational curve passing through **x**. Note that A can not be γ -invariant, since otherwise $\hat{q}([A]) = -\hat{q}([\gamma(A)]) = -\hat{q}([A]) \in \mathbb{Z}/4$, which implies that $\hat{q}([A])$ is even and thus contradicts to the congruences $\hat{q}([A]) \equiv [A]^2 \equiv -AK \equiv 2n + 1 \mod 2$. Therefore, the set of such curves A splits into a finite union of pairs, $\{A, \gamma(A)\}$, and there remains to notice that $i^{\hat{q}([A])-1} + i^{\hat{q}([\gamma(A)])-1} = 0$ due to $\hat{q}([A]) + \hat{q}([\gamma(A)]) = 0, \hat{q}([A]) = \pm 1$.

In the case m = 2n, it is sufficient to notice that then $\hat{q}(\alpha) = -\hat{q}(\alpha) \in \mathbb{Z}/4$ for any $\alpha \in \mathcal{L}_{\mathbb{R}}^m$, since (similar to the above) $\hat{q}(\alpha) \equiv \alpha^2 \equiv -\alpha K \equiv 2n \mod 2$. \Box

2.2. Three auxiliary surfaces. To simplify proving Theorem 1.2.4 (see Section 5) we pick some particular real del Pezzo surface X for each of the degrees d = 1, 2, 3.

2.2.1. Proposition.

- (1) For d = 1, pick X with $X_{\mathbb{R}} = \mathbb{RP}^2 \perp \mathbb{I}3\mathbb{S}^2$. Then :
 - H₂⁻(X) ≅ Z² is generated by K and a real root vector e ∈ K[⊥], while the first real layer L_R¹(X) consists of -K and the divisor classes of 2 real lines, -K±e.
 The Bertini involution acts on H₂⁻(X) preserving the basic quadratic form q̂ and -K, but permuting the divisor classes of lines.
- (2) For d = 2, pick X with $X_{\mathbb{R}} = \mathbb{K} \perp \mathbb{S}^2$ (\mathbb{K} denotes a Klein bottle). Then :
 - $H_2^-(X) \cong \mathbb{Z}^4$ is generated by K, three pairwise orthogonal real root vectors $e_1, e_2, e_3 \in K^{\perp}$, and $\frac{1}{2}(-K e_1 e_2 e_3)$, while the first real layer $\mathcal{L}^1_{\mathbb{R}}(X)$ consists of divisor classes of 8 real lines, $\frac{1}{2}(-K \pm e_1 \pm e_2 \pm e_3)$.

• X can be chosen so that its group of automorphisms contains $\mathbb{Z}/2 \times \mathfrak{D}_4$, where $\mathbb{Z}/2$ is generated by the Geiser involution γ and \mathfrak{D}_4 is the dihedral group of a square.

• $\mathbb{Z}/2 \times \mathfrak{D}_4$ acts transitively on the above 8 lines, and, in particular, one of the \mathfrak{D}_4 -orbits $\{\frac{1}{2}(-K-e_1-e_2-e_3), \frac{1}{2}(-K+e_1+e_2-e_3), \frac{1}{2}(-K+e_1-e_2+e_3), \frac{1}{2}(-K-e_1+e_2+e_3)\}$ is permuted by γ -action with the other \mathfrak{D}_4 -orbit. The \mathfrak{D}_4 -action preserves each of the basic quadratic functions $\pm \hat{q}$, while γ_* acts as $(-1)^n$ on $\hat{q}|_{\mathcal{L}^n_*}$.

• The multiples of K are the only γ_* -invariant classes and the only ones invariant under the \mathfrak{D}_4 -action.

- (3) For d = 3, pick X with $X_{\mathbb{R}} = \mathbb{RP}^2 \sqcup \mathbb{S}^2$. Then :
 - $H_2^-(X)$ is generated by the divisor classes of the 3 real lines $L_1, L_2, L_3 \subset X$.

• X can be chosen so that its automorphism group contains the symmetric group S_3 acting by permutations on the above lines.

- The S₃-action in $H_2^-(X)$ preserves the basic quadratic function \hat{q} .
- The multiples of K are the only classes invariant under this action.

Proof. In all 3 cases we use the 1–1 correspondence between real lines and divisor classes $\alpha \in \mathcal{L}^1_{\mathbb{R}}$ with $\alpha^2 = -1$ and apply the Lefschetz fixed point theorem, $\operatorname{rk} H_2^+(X) - \operatorname{rk} H_2^-(X) = \chi(X_{\mathbb{R}}) - 2$, which together with $\operatorname{rk} H_2^+(X) + \operatorname{rk} H_2^-(X) = -1$

 $\operatorname{rk} H_2(X) = 10 - d$ calculates the ranks of $H_2^{\pm}(X) = \operatorname{ker}(1 \mp \operatorname{conj}_*)$. We use also that the lattices $H_2^{\pm}(X) = \operatorname{ker}(1 \mp \operatorname{conj}_*)$ have 2-periodic discriminant groups of rank $\frac{1}{2}(10 - d - \operatorname{rk} H^*(X_{\mathbb{R}}))$ (see, f.e., [DIK, Proposition 8.3.3]). Recall besides that both Bertini (for d = 1) and Geiser (for d = 2) involutions act on $H_2(X)$ as a reflection against the line spanned by K.

(1) In this case, $K^{\perp} \cap H_2^- \cong \langle -2 \rangle$. The divisor classes of real lines $\alpha \in \mathcal{L}^1_{\mathbb{R}}$ split as $\alpha = -K \pm e$, where e is a generator of $\langle -2 \rangle$, and the Bertini involution permutes these lines. Preserving of \hat{q} by its action is due to Theorem 2.1.1(1).

(2) In this case, $\operatorname{rk} H_2^{\pm} = 4$ and $\operatorname{rk}(K^{\perp} \cap H_2^{-}) = 3$. Since X is an (M-2)-surface of type I, the discriminant form of H_2^+ is even 2-periodic of rank 2. Thus, we conclude that $H_2^+ \cong D_4$. By Nikulin's gluing theorem [N], the discriminant group discr $(K^{\perp} \cap H_2^-)$ is also 2-periodic and has rank 1 or 3 (since $K^{\perp} \cap H_2^-$ is complementary to H_2^+ in $K^{\perp} = E_7$). In the case of discriminant rank 1, the Brown invariant would be $\pm 1 \in \mathbb{Z}/8$, which contradicts to Brown's congruence $\operatorname{Br}(\operatorname{discr}(K^{\perp} \cap H_2^-)) = \sigma(K^{\perp} \cap H_2^-) = -3 \mod 8$. Thus, the discriminant rank is 3, which implies $K^{\perp} \cap H_2^- \cong 3A_1$. Arithmetical description of the real lines is then straightforward.

For constructing X with the required \mathfrak{D}_4 -symmetry, it is sufficient to pick a \mathfrak{D}_4 -symmetric perturbation of a \mathfrak{D}_4 -symmetric pair of conics, as shown on Fig. 1. Note that the above \mathfrak{D}_4 -action is realized by projective transformations $\mathbb{P}^2 \to$

FIGURE 1. Degree 2 del Pezzo surface with \mathfrak{D}_4 -symmetry lifted to a double plane branched along a \mathfrak{D}_4 -symmetric quartic curve



The thick/dotted coloring of bitangents shows how they are lifted so that \mathfrak{D}_4 -symmetry is preserved in the double covering.

 \mathbb{P}^2 , and hence has a "cylindrical" lifting to $\mathbb{P}(1,1,1,2)$ which preserves X. Such transformations $\mathbb{P}(1,1,1,2) \to \mathbb{P}(1,1,1,2)$ preserve each of the two Pin⁻-structures on $\mathbb{P}_{\mathbb{R}}(1,1,1,2)$ and send outward vector fields on $X_{\mathbb{R}}$ to outward fields. Therefore, each of \mathfrak{D}_4 -symmetries preserves each of the basic Pin⁻-structures on $X_{\mathbb{R}}$ and thus preserves \hat{q} . Proposition 2.1.2 gives also the required skew-symmetry of the γ -action on \hat{q} .

(3) In this case, $\operatorname{rk} H_2^-(X) = 3$ and its discriminant group has order 4. Since the intersection matrix $\begin{bmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{bmatrix}$ of the three real lines lying on X (the only existing ones) has the same determinant 4, these lines must generate the whole lattice $H_2^-(X)$.

An example of X with a required S_3 -action can be given by an equation of the form $x_0(x_1^2 + x_2^2 + x_3^2 - x_0^2) = \varepsilon x_1 x_2 x_3$. This action preserves \hat{q} due to Theorem 2.1.1(1). The last claim about invariant classes follows from linear independence of divisor classes of the lines L_1, L_2, L_3 and the relation $[L_1] + [L_2] + [L_3] = -K$. \Box

2.3. Wall-crossing. We call a complex analytic family $X(z), z \in \mathbb{D} = \{z \in \mathbb{C} \mid |z| < 1\}$ a Morse-Lefschetz family, if : $\mathcal{X} = \bigcup_z X(z)$ is a non-singular 3-fold; each X(z) with $z \neq 0$ is non-singular while X(0) is uninodal; the projection $\pi : \mathcal{X} \to \mathbb{D}$ is a proper map and it is a submersion at every point except the nodal point of X(0); at the nodal point of X(0) in appropriate local coordinates z_1, z_2, z_3 the projection can be written as $z = \sum_i z_i^2$. If \mathcal{X} is equipped with a real structure sending X(z) to $X(\bar{z})$ for each $z \in \mathbb{D}$, the family is called *real*.

The following is well known (for del Pezzo surfaces with $K^2 = 1$, see, for example, [FK-3]).

2.3.1. **Proposition.** Any pair of real del Pezzo surfaces of same degree $d \leq 3$ can be connected either by a real deformation, or a finite sequence of real Morse-Lefschetz families of del Pezzo surfaces embedded into the corresponding projective or weighted projective space. If the real locus of the both surfaces is non-empty the sequence of real Morse-Lefschetz families can be chosen not to involve surfaces with empty real locus.

Every real Morse-Lefschetz family $\pi : \mathcal{X} \to \mathbb{D}$ being restricted to the punctured upper half disc $\mathbb{D}^+ = \{z \in \mathbb{D}, \operatorname{Im} z \ge 0, z \ne 0\}$ and considered as a fibration of smooth 4-manifolds is trivial. This provides a natural identification of lattices $H_2(X(t)), t \in \mathbb{R}^*, |t| < 1$, so that for these values of t the involutions $c_t : H_2(X(t)) \to H_2(X(t))$ and $c_{-t} : H_2(X(-t)) \to H_2(X(-t))$ induced by the complex conjugation become related by the Picard-Lefschetz transformation:

(2.3.1) $c_t = s_e \circ c_{-t} \quad s_e(v) = v + (ev)e,$

where $e \in H_2(X(t))$ is the vanishing class of the underlying nodal degeneration.

2.3.2. **Proposition.** Let $\pi : \mathcal{X} \to \mathbb{D}$ be a real Morse-Lefschetz family of del Pezzo surfaces of degree $K^2 \leq 3$, embedded into the corresponding projective or weighted projective space, and let the direction of the wall-crossing is selected so that $\chi(X_{\mathbb{R}}(-t)) < \chi(X_{\mathbb{R}}(t))$ for t > 0. Then, $H_2^-(X(t))$ is naturally identified with $H_2^-(X(-t)) \cap e^{\perp}$ and with respect to this identification $\hat{q}_{\theta^{X(t)}}$ is the restriction of $\hat{q}_{\theta^{X(-t)}}$, where in the case $K^2 = 2$ we pick arbitrary Pin⁻-structure on the ambient $\mathbb{P}(1, 1, 1, 2)$ and choose that basic Pin⁻-structures which are induced by means of outward vector fields.

Proof. To identify the lattices $H_2(X(t)), t \in \mathbb{R}^*, |t| < 1$ we use, as it is described above, the trivialization of the family over the upper disc. Then, under assumption $\chi(X_{\mathbb{R}}(-t)) < \chi(X_{\mathbb{R}}(t))$ for t > 0, the identity $c_t = s_e \circ c_{-t}$ implies that $e \in$ $H_2^-(X(-t))$ and $H_2^-(X(t)) = H_2^-(X(-t)) \cap e^{\perp}$. Also, $e \in H_2^+(X(t)) = \ker(1 - \operatorname{conj}_*)$. Next, we follow the geometric interpretation of the Viro homomorphism Υ (see Section 2.1).

We fix $t_0 > 0$ and pick a class $\alpha \in H_2^-(X(t_0))$. If $\Upsilon \alpha \neq 0$, then we realize it by a conj-invariant smooth embedded oriented 2-manifold $F(t_0) \subset X(t_0)$ disjoint from $X_{\mathbb{R}}(t_0)$. If $\Upsilon(\alpha) \neq 0$, we choose $F(t_0) \subset X(t_0)$ in such a way that each of the components of $F(t_0) \cap X_{\mathbb{R}}(t_0)$ represents a non-zero element in $H_1(X_{\mathbb{R}}(t_0); \mathbb{Z}/2)$. In both cases, if one of spherical components of $X_{\mathbb{R}}(t_0)$ is a vanishing sphere $S^2(t_0)$, then $F(t_0)$ does not intersect $S^2(t_0)$ (since the circle components of intersection should be homologically non-trivial, while a presence of an isolated intersection point contradicts to smoothness of $F(t_0)$ and its conj-invariance). If a vanishing sphere $S^2(t_0)$, chosen conj-equivariant, is not contained in $X_{\mathbb{R}}(t_0)$, then, due to our assumption on the direction of the wall crossing, $S^2(t_0) \cap X_{\mathbb{R}}(t_0)$ consists of 2 points, while the intersection number $e \cdot \alpha$ is 0 (since e and α belong to opposite eighenspaces for the action of conj in $H_2(X(t_0))$). Thus, we can eliminate, respecting conj-invariance, all geometric intersection points between $F(t_0)$ and $S^2(t_0)$ that might appear when $S^2(t_0) \cap X_{\mathbb{R}}(t_0) \neq \emptyset$, since in such a case any 2 complex conjugate intersection points are of opposite intersection index and, hence, can be eliminated by a surgery of $F(t_0)$ along a conj-invariant path joining them along $S^2(t_0)$.

As soon as $F(t_0) \cap S^2(t_0) = \emptyset$, there is no more obstruction for extending $F(t_0)$ up to a locally trivial conj-invariant family $F(t) \subset X(t), t \in \mathbb{D}_{\mathbb{R}}$, avoiding the node of X(0). Now, the invariance of $q_{\theta^{X_{\mathbb{R}}(t)}}([F(t) \cap X_{\mathbb{R}}(t)])$ follows from its continuity (note that $q^{X_{\mathbb{R}}(0)}$ is well defined on H_1 of the smooth part of $X_{\mathbb{R}}(0)$). \Box

Recall that untwisting of a given real Morse-Lefschetz family X(z) induced by the substitution $z = t^2$ (see [FK-3]) and followed by resolution of the nodal singularity defines a complex analytic fibered 3-fold $\hat{\pi} : \hat{\mathcal{X}} \to \mathbb{D}$ equipped with a real structure. This new family $\hat{\mathcal{X}}(t) = \hat{\pi}^{-1}(t), t \in \mathbb{D}$ is formed by $\hat{\mathcal{X}}(t) = X(t^2)$ for $t \neq 0$ and a reducible fiber $\hat{\mathcal{X}}(0) = \hat{X}(0) \cup \mathcal{E}$, where $\hat{X}(0)$ is the resolution of the nodal surface $X(0), \mathcal{E}$ is a non-singular projective quadric surface, and their intersection $E = X(0) \cap \mathcal{E}$ can be considered at the same time as the exceptional divisor of the blow-down $\hat{X}(0) \to X(0)$ and as a hyperplane section of \mathcal{E} viewed as a projective quadric.

The following lemma, which will be used in Section 4, requires a more flexible symplectic setting. So, we endow \mathcal{X} and then $\hat{\mathcal{X}}$ with a Kähler structure ω respected by the complex conjugation conj and then take a small real perturbation of the (almost) complex structure in $\hat{\mathcal{X}}$ to make it generic in the class of *admissible almost-complex structures* on a *symplectic fibration* ($\hat{\pi} : \hat{\mathcal{X}} \to \mathbb{D}, \omega$) (see [T] for definitions and existence).

In this lemma we consider a very special type of wall-crossing, contraction of a spherical component, meaning by this that the node of $X_{\mathbb{R}}(0)$ is solitary and give birth to an S^2 -component in $X_{\mathbb{R}}(t)$, t > 0, or equivalently that $\mathcal{E}_{\mathbb{R}} = S^2$ and $E_{\mathbb{R}} = \emptyset$.

2.3.3. Lemma. Let $\hat{\pi} : \hat{\mathcal{X}} \to \mathbb{D}$ be a described above real symplectic fibration (equipped with an admissible generic almost-complex structure) contracting a spherical component, and let $\mathbf{x}(t) = (x(t), z_1(t), w_1(t), \dots, z_{m-1}(t), w_{m-1}(t))$ be a generic conj-equivariant family of point-constraints such that x(0) is real and belongs to the sphere $S^2 = \mathcal{E}_{\mathbb{R}}$, while $w_1(0) = \operatorname{conj} z_1(0), \dots, w_{m-1}(0) = \operatorname{conj} z_{m-1}(0)$ are imaginary and belong to $\hat{X}(0)$. Then:

- (1) For any selection $p_1 \in \{z_1(0), w_1(0)\}, \ldots, p_{m-1} \in \{z_{m-1}(0), w_{m-1}(0)\},$ each of J-holomorphic rational curves $C \subset \hat{X}(0)$ having $-K \cdot C = m$ and passing through p_1, \ldots, p_{m-1} is irreducible nodal and intersects E transversally. The number of such curves C in a given divisor class does not depend on the selection of points p_1, \ldots, p_{m-1} .
- (2) For any continuous family $A(t) \in C_{\mathbb{R}}(2m, 1, \mathbf{x}(t))$, the limit-curve $\lim_{t\to 0^+} A(t)$ is a curve-configuration

$$C \cup \operatorname{conj} C \cup L_1 \cup \operatorname{conj} L_1 \cup \cdots \cup L_{n-1} \cup \operatorname{conj} L_{n-1} \cup H_n \subset \mathcal{X}(0)$$

with the following properties: C is a J-holomorphic rational curve which lies in $\hat{X}(0)$ and passes through a selection $p_1 \in \{z_1(0), w_1(0)\}, \ldots, p_{m-1} \in \{z_{m-1}(0), w_{m-1}(0)\}; -K \cdot C = m; n = C \cdot E; C \cap E$ consists of n distinct points $q_1, \ldots, q_n; L_1, \ldots, L_{n-1}$ are \mathbb{P}^1 -generators of $\mathcal{E} \cong \mathbb{P}^1 \times \mathbb{P}^1$ and pass through a selection of n-1 points between $q_1, \ldots, q_n; H_n$ is a real hyperplane section of $\mathcal{E} \cong \mathbb{P}^1 \times \mathbb{P}^1$ that passes through the remaining point and the point x(0).

(3) For any sufficiently small t > 0, passing to the limits as in item (2) establishes a bijection between the set C_ℝ(2m, 1, **x**(t)) and the set I of curve-configurations described in item (2). The set I is a disjoint union of 2^{m-2} subsets I_{**p**,conj**p**} where {**p**, conj**p**} is pair of complex conjugate selections and I_{{**p**,conj**p**}} consists of configurations with **p** ⊂ C, conj**p** ⊂ conjC. Each I_{**p**,conj**p**} is itself a disjoint union of subsets I_{C,conjC} consisting of configurations sharing the same pair {C, conjC}. Each I_{C,conjC} contains n2ⁿ⁻¹ elements.

Proof. Claim (2) and the first part of Claim (1) follow from [Br-P, Proposition 3.7] (the option of tangency between H and E is excluded, since $E_{\mathbb{R}} = \emptyset$). Due to the transversality of curves C in Claim (1), their number is nothing but a relative Gromov-Witten invariant of pair ($\hat{X}(0), E$) (without constraints on E). The bijectivity in Claim (3) follows from Symplectic Sum Formula in its simplest, transversal intersection, case. The rest of Claim (3) is a straightforward consequence of bijectivity.

3. Proof of Theorem 1.2.1

In this section we consider real del Pezzo surfaces X of degree $d \leq 3$, with a basic Pin⁻-structure θ^X (cf. Theorem 2.1.1) and the associated with it quadratic function $\hat{q}_{\theta^X} : H_2^-(X) \to \mathbb{Z}/4$. In Introduction we omitted, for shortness, indicating X and θ^X . In this section we need to use a full notation, since X will be varying. So, let us adjust our notation and rewrite our main definition (1.2.1) correspondingly:

$$N_{m,k}^X = \sum_{\alpha \in \mathcal{L}_{\mathbb{R}}^m(X)} \mathcal{N}_{\alpha,k}^X, \ \mathcal{N}_{\alpha,k}^X = i^{\hat{q}_{\theta^X}(\alpha) - m^2} W_{\alpha,k}^X, \ W_{\alpha,k}^X = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x}), [A] = \alpha} w(A).$$

Recall that the *traditional Welschinger invariants* are defined similarly as

$$\widetilde{W}_{\alpha,k}^{X} = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x}), [A] = \alpha} \widetilde{w}(A), \quad \text{with} \quad \widetilde{w}(A) = (-1)^{s_{A}}$$

where s_A stands for the number of solitary real nodal points of A. Since $w(A) = (-1)^{c_A}$ and $c_A + s_A$ is the arithmetic genus $g_a(\alpha)$, where $\alpha = [A]$ (and c_A is the number of cross-point real nodes), we have

$$\widetilde{W}_{\alpha,k}^X = (-1)^{g_a(\alpha)} W_{\alpha,k}^X.$$

3.1. **Push-forward formula.** Assume that real del Pezzo surfaces X and Y are related by wall-crossing, and consider a real Morse-Lefschetz family $\pi : \mathcal{X} \to \mathbb{D}$ such that $X = X(-t_0)$ and $Y = X(t_0)$ for a fixed $0 < t_0 < 1$. Following the exposition in Section 2.3, we identify the lattices $H_2(X) = H_2(Y)$ using a smooth trivialization of the fibration $\pi : \mathcal{X} \to \mathbb{D}$ over the upper half disc.

We choose the direction of wall-crossing so that $\chi(X_{\mathbb{R}}) < \chi(Y_{\mathbb{R}})$, which implies $H_2^-(Y) = \{ \alpha \in H_2^-(X) \mid \alpha e = 0 \}$ where $e \in H_2^-(X)$ is the vanishing class,

and consider the orthogonal projection

(3.1.1)
$$p^e: H_2^-(X) \to H_2^-(Y) \otimes \mathbb{Q} \qquad v \mapsto v + \frac{1}{2}(ev)e.$$

Our aim is to treat the push-forward by p^e of the function $\mathcal{N}_k^X : H_2^-(X) \to \mathbb{Z}$ defined by $\mathcal{N}_k^X(\alpha) = \mathcal{N}_{\alpha,k}^X$ if α is an effective divisor class and $\mathcal{N}_k^X(\alpha) = 0$ otherwise.

Note that, due to Theorem 2.1.1(1) and Proposition 2.1.2(2), none of the above ingredients, including p^e and \mathcal{N}_k^X , depends on a choice of t_0 .

3.1.1. **Proposition.** If $X_{\mathbb{R}}$ and $Y_{\mathbb{R}}$ are non-empty, then: (1) the push-forward $p_*^e(\mathcal{N}_k^X)$ is well-defined, (2) it is supported in the integer part $H_2^-(Y) = H_2^-(Y) \otimes 1 \subset H_2^-(Y) \otimes \mathbb{Q}$, and (3) being restricted to this integer part, $p_*^e(\mathcal{N}_k^X) = \mathcal{N}_k^Y$ on $H_2^-(Y)$.

Proof. By definition, $p_*^e(\mathcal{N}_k^X)(\alpha) = \sum_{n \in \mathbb{Z}} \mathcal{N}_k^X(\alpha + n e)$, and (1) means finiteness of the sum. But if |n| is sufficiently large, then $(\alpha + ne)^2 < -2$ and thus, $\alpha + ne$ can not be realized by a reduced irreducible rational curve, which gives $\mathcal{N}_k^X(\alpha + ne) = 0$.

According to the projection formula (3.1.1), $p_*^e(\mathcal{N}_k^X)$ is supported in $H_2^-(Y) \otimes \frac{1}{2}\mathbb{Z}$, and (2) means that

$$\sum_{n \in \mathbb{Z}} \mathcal{N}_k^X(\alpha + ne) = 0 \quad \text{ for each } \alpha \in H_2^-(X) \text{ with } r = \alpha e \text{ odd}$$

To prove that this sum is zero, we group the summands by pairs $\alpha + ne$, $\alpha + (r-n)e$ and notice that the reflection s_e permutes the elements in each of these pairs. As is known (see for example [IKS-2, Theorem 4.3(1)]) the traditional Welschinger invariants are preserved under such a reflection, so that $\widetilde{W}_{\alpha+ne,k}^X = \widetilde{W}_{\alpha+(r-n)e,k}^X$ As it follows from adjunction formula, the curves in the divisor class $\alpha + ne$ and the curves in the divisor class $\alpha + (d - n)e$ have the same parity of the arithmetic genus. Hence, $W_{\alpha+ne,k}^X$ and $W_{\alpha+(r-n)e,k}^X$ are also equal. Thus, there remains to notice, that $(\alpha + ne)K = \alpha K = (\alpha + (r - n)e)K$, and that

$$i^{\hat{q}_{\theta X}(\alpha+(r-n)e)} = i^{\hat{q}_{\theta X}(\alpha+ne)+\hat{q}_{\theta X}((r-2n)e)+2(\alpha+ne)((r-2n)e)}$$
$$= i^{\hat{q}_{\theta X}(\alpha+ne)+2r(r-2n)} = -i^{\hat{q}_{\theta X}(\alpha+ne)}$$

due to $\hat{q}_{\theta^X}(e) = 0$ (see Theorem 2.1.1) and $r = 1 \mod 2$.

To prove (3) we apply Theorem 2.1 from [Br]. According to this theorem, if $\alpha e = 0$, then

$$\widetilde{W}_{\alpha,k}^Y = \sum_{n \in \mathbb{Z}} (-1)^n \widetilde{W}_{\alpha+n\,e,k}^X.$$

This implies

$$W^Y_{\alpha,k} = \sum_{n \in \mathbb{Z}} W^X_{\alpha+ne,k},$$

since, due to the adjunction formula, $g_a(\alpha + ne) = g_a(\alpha) + n \mod 2$ as soon as $\alpha e = 0$. Finally, it is left to notice that $\hat{q}_{\theta X}(\alpha + ne) = \hat{q}_{\theta X}(\alpha)$ for all $n \in \mathbb{Z}$, and that $\hat{q}_{\theta Y}(\alpha) = \hat{q}_{\theta X}(\alpha)$ due to Proposition 2.3.2.

3.2. **Proof of Theorem 1.2.1.** Due to deformation invariance of \hat{q} (see Theorem 2.1.1), and due to independence of \mathcal{N}_k^X from a choice of a basic Pin⁻-structure when $K^2 = 2$ (see Proposition 2.1.2(2)), it is left to consider sequences of real Morse-Lefschetz families as in Proposition 2.3.1 and to apply Proposition 3.1.1 (noticing that the projection p^e preserves the layers).

WALL-CROSSING INVARIANCE

4. Proof of Theorem 1.2.3

Throughout this section we fix a real del Pezzo surface X of degree $d = K^2 \leq 3$ and use notation $\hat{q}: H_2(X) \to \mathbb{Z}/4$ for the quadratic function associated to a basic Pin⁻-structure on $X_{\mathbb{R}}$.

4.1. Switch to open Gromov-Witten invariants. In the computations below we essentially rely on Solomon's recursion relations for open Gromov-Witten invariants (see [H-S]). In accordance with notation in [H-S], for each $m \ge 1$, $0 \le k \le m-1, k = m-1 \mod 2$, and each $v \in K^{\perp} \cap \ker(1 + \operatorname{conj}_*)$, we put

(4.1.1)

$$N_{m,v,k} = \sum_{\substack{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x}) \\ d[A] = -mK - v}} i^{\hat{q}([A]) - m^2} w(A),$$

$$\Gamma_{m,v,k} = -2^{1-l} \sum_{\substack{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x}) \\ d[A] = -mK - v}} i^{\hat{q}([A]) - m^2} w(A), \qquad l = \frac{1}{2}(m - k - 1)$$

where **x** is a generic collection of k real points and $l = \frac{1}{2}(m-k-1)$ pairs of complex conjugate imaginary points, and define

$$\Gamma_{m,k} = \sum_{v \in K^{\perp} \cap \ker(1 + \operatorname{conj}_{*})} \Gamma_{m,v,k}.$$

Note that:

- If -mK v is not divisible by d, then the sums involved in (4.1.1) are void and the numbers $N_{m,v,k}$, $\Gamma_{m,v,k}$ are zero by definition.
- Numbers $N_{m,v,k}$ and $\Gamma_{m,v,k}$ does not depend on a choice of **x** and are preserved under equivariant deformations and equivariant isomorphisms (*cf.* discussion in Introduction).
- In accordance with (1.2.1) and above definitions we have

(4.1.2)
$$N_{m,k} = -2^{l-1} \sum_{v \in K^{\perp} \cap \ker(1 + \operatorname{conj}_{*})} \Gamma_{m,v,k} = -2^{l-1} \Gamma_{m,k},$$

while Theorem 1.2.1 implies that both $N_{m,k}$ and $\Gamma_{m,k}$ do not depend even on a choice of a real del Pezzo surface of given degree.

• If d is odd, we have

$$\hat{q}([A]) - m^2 = \hat{q}(d[A]) - m^2 = \hat{q}(-mK - v) - m^2 = \hat{q}(v),$$

so that the above definitions can be rewritten as follows

$$N_{m,v,k} = \sum_{\substack{A \in C_{\mathbb{R}}(m,k,\mathbf{x}) \\ K^{2}[A] = -mK - v}} i^{\hat{q}(v)} w(A),$$

$$\Gamma_{m,v,k} = -2^{1-l} \sum_{\substack{A \in C_{\mathbb{R}}(m,k,\mathbf{x}) \\ d[A] = -mK - v}} i^{\hat{q}(v)} w(A) \qquad l = \frac{1}{2}(m-k-1)$$

• Theorem 1.2.4 is reformulated as

(4.1.3)
$$2md\Gamma_{m,0} + \sum_{j=1}^{n} \binom{n-1}{n-j} j(m-2j)^{2}\Gamma_{m-2j,0}\Gamma_{2j,1} \quad \text{for } m = 2n+1,$$
$$2md\Gamma_{m,1} + \sum_{j=1}^{n} \binom{n-1}{n-j} j(m-2j)^{2}\Gamma_{m-2j,1}\Gamma_{2j,1} \quad \text{for } m = 2n+2.$$

4.2. Proof of Theorem 1.2.3.

4.2.1. Lemma. For any $n \in \mathbb{N}$, we have $n2^n = \sum_{k=0}^n (n-2k)^2 {n \choose k}$.

Proof. The identities $\sum_{k=0}^{n} \binom{n}{k} = 2^{n}$, $\sum_{k=0}^{n} k\binom{n}{k} = n2^{n-1}$, and $\sum_{k=0}^{n} k^{2}\binom{n}{k} = (n+n^{2})2^{n-2}$ imply $\sum_{k=0}^{n} (n-2k)^{2}\binom{n}{k} = n^{2}2^{n} - 4n^{2}2^{n-1} + 4(n+n^{2})2^{n-2} = n2^{n}$. \Box

Proof of Theorem 1.2.3. Due to Theorem 1.2.1, it is enough to check the formula (1.2.2) for one particular del Pezzo surface X of each degree $d \leq 3$. We make choice by picking $X = X(t_0), 0 < t_0 \ll 1$, from a real Morse-Lefschetz family X(t) contracting a spherical component $S^2 \subset X_{\mathbb{R}}(t_0)$ (see Sec. 2.3).

First, we interpret the right-hand side of (1.2.2) as a weighted count of complex rational curves C on the resolution $\hat{X}(0)$ of the nodal surface X(0) that (1) belong to the *m*-th level $\mathcal{L}^m(\hat{X}(0))$, (2) pass through a fixed generic collection \mathbf{p} of m-1points, and (3) have a non-trivial intersection, $C \cdot E > 0$, with the (-2)-curve $E \subset \hat{X}(0)$ representing the node. Namely, we observe that, in accordance with the Abramovich-Bertram-Vakil formula (see [IKS-2, Proposition 4.1] and [Br-P, Theorem 2.5]), the input of each of such curves C into the right-hand side is equal to $\sum_{k=0}^{n} (n-2k)^2 {n \choose k}$ where $n = C \cdot E$. Thus, in accordance with Lemma 4.2.1, the right-hand side of (1.2.2) can be seen as the weighted count of the above curves $C \subset \hat{X}(0)$ with weights $n2^n$.

To treat the left-hand side we consider the untwisted family $\hat{\mathcal{X}}(t)$ and apply Lemma 2.3.3 choosing the constraint $\mathbf{x}(t) = (x(t), z_1(t), w_1(t), \ldots, z_{m-1}(t), w_{m-1}(t))$ as indicated there. First, we note that the input of each of the curves $A(t_0) \in \mathcal{C}(2m, 1, \mathbf{x}(t_0))$ into $N_{2m,1}$ is equal to 1. This is because \hat{q} vanishes on the spherical component of $X_{\mathbb{R}}(t_0)$ containing $x(t_0)$ and curves $A_{\mathbb{R}}(t)$ have no real cross-point nodes for all sufficiently small t > 0 (the latter follows from the explicit description of the limit curves $\hat{A}_{\mathbb{R}}(0)$ in Lemma 2.3.3(2)). So, $N_{2m,1}$ is just the cardinality of $\mathcal{C}(2m, 1, \mathbf{x}(t_0))$, which in its turn coincides with the cardinality of the set I that counts the limit curve-configurations see Lemma 2.3.3(3).

To compare this cardinal count of the limit curve-configurations with the weighted count we made at the beginning, let us choose as \mathbf{p} a selection $p_1 \in \{z_1(0), w_1(0)\}, \dots, p_{m-1} \in \{z_{m-1}(0), w_{m-1}(0)\}$. Restricting the cardinal count to any particular selection \mathbf{p} is equivalent to dividing $N_{2m,1}$ by 2^{m-2} (in accord with subsets $I_{\{\mathbf{p}, \operatorname{conj} \mathbf{p}\}}$ in Lemma 2.3.3(3)). On the other hand, forgetting the line components in the curve configuration is equivalent to counting with the weight $n2^{n-1}$. Thus, we conclude that

$$\sum_{\alpha \in \mathcal{L}^m} (e\alpha)^2 G W_\alpha = 2 \cdot 2^{-(m-2)} N_{2m,1} = 2^{3-m} N_{2m,1}.$$

Due to (4.1.2), the result of Theorem 1.2.3 can be rewritten as follows.

4.2.2. Corollary. In the same setting as in Theorem 1.2.3, we have

(4.2.1)
$$-2\Gamma_{2m,1} = \sum_{\alpha \in \mathcal{L}^m} (e\alpha)^2 G W_{\alpha} . \square$$

5. Proof of Theorem 1.2.4

5.1. Preparation for proving Theorem 1.2.4. Apart of Theorem 1.2.1 and Proposition 4.2.2, the proof is based on the following Solomon's recursion rule (cf. [H-S, (OGW3)] and [Ch, Th.1.1(RWDVV3)]) which is a corollary of an analog for WDVV-equation designed by Solomon [S2] in the framework of open strings.

5.1.1. Theorem. ¹ Let X be a real del Pezzo surface equipped with a basic Pin^{-} structure, and let $H_1, H_2, H_3 \in H_2^-(X)$ be fixed elements with $H_1H_3 = 0$. Then, for each pair $(m,k) \in \mathbb{Z}_{>0} \times \mathbb{Z}_{\geq 0}$ with $l = \frac{1}{2}(m-1-k) \geq 1$ and any $B \in \mathcal{L}^m_{\mathbb{R}}$, the following relation holds:

$$(H_{1}H_{2})(H_{3}B)\Gamma_{B,k} = \Theta_{B,k}^{(1)} + \Theta_{B,k}^{(2)}, \text{ where}$$

$$\Theta_{B,k}^{(1)} = \frac{1}{4} \sum_{\substack{B_{1}+B_{2}=B\\k_{1}+k_{2}=k+1}} (H_{1}B_{1})((H_{3}B_{1})(H_{2}B_{2}) - (H_{2}B_{1})(H_{3}B_{2})) \binom{l-1}{l_{1}}\binom{k}{k_{1}}\Gamma_{B_{1},k_{1}}\Gamma_{B_{2},k_{2}}$$

$$\Theta_{B,k}^{(2)} = \frac{1}{2} \sum_{\substack{B_{F}-\operatorname{conj}_{*},B_{F}\\+B_{U}=B}} (B_{U}B_{F})(H_{1}B_{F})((H_{3}B_{F})(H_{2}B_{U}) - (H_{2}B_{F})(H_{3}B_{U}))\binom{l-1}{l_{U}}GW_{B_{F}}\Gamma_{B_{U},k}$$

In this theorem, notation $\Gamma_{B,k}$ stands for $\Gamma_{m,v,k}$ such that $B = \frac{1}{d}(-mK - v)$ (and similar for B_i , m_i , v_i , etc.). We let also $l = \frac{1}{2}(m-1-k)$ (and similar for $m_i, k_i, l_i,$ etc.). For shortness, we adopt the convention to put $\Gamma_{\alpha,k} = 0$ for any $\alpha \notin H_2^-(X).$

Theorem 1.2.1 allows us to give the proof only for one particular real del Pezzo surface X for each degree d = 1, 2, 3. We take X as in Proposition 2.2.1 and denote by G_d its automorphism group described there, that is:

$$G_d = \begin{cases} \mathbb{Z}/2\mathbb{Z} \text{ generated by the Bertini involution } \tau, \text{ if } d = 1, \\ \mathbb{Z}/2\mathbb{Z} \times \mathfrak{D}_4 \text{ where } \mathbb{Z}/2 \text{ is generated by the Geiser involution } \gamma, \text{ if } d = 2, \\ S_3, \text{ if } d = 3. \end{cases}$$

To apply Theorem 5.1.1, we choose:

- $H_1 \in \mathcal{L}^1_{\mathbb{R}} \setminus \{-K\}$, so that $H_1^2 = KH_1 = -1$, $H_2 = -K$, so that $H_1H_2 = 1$,
- $H_3 = H_1 + H_2$, so that $H_1H_3 = 0$ as required in Theorem 5.1.1,

which gives

(5.1.1)
$$(H_3B)\Gamma_{B,k} = \Theta_{B,k}^{(1)} + \Theta_{B,k}^{(2)}$$
, where

¹In fact, this theorem, with appropriate definitions for the numbers $\Gamma_{B,k}$, holds for all real rational surfaces X with any Pin⁻-structure on $X_{\mathbb{R}}$.

$$\begin{split} \Theta_{B,k}^{(1)} &= \frac{1}{4} \sum_{\substack{m_1 + m_2 = m, m_1, m_2 \ge 1 \\ B_1 + B_2 = B, B_i \in \mathcal{L}_{\mathbb{R}}^{m_i} \\ k_1 + k_2 = k + 1}} (H_1 B_1) ((H_1 B_1) m_2 - m_1 (H_1 B_2)) \binom{l-1}{l_1} \binom{k}{k_1} \Gamma_{B_1, k_1} \Gamma_{B_2, k_2} \\ \Theta_{B,k}^{(2)} &= \frac{1}{2} \sum_{\substack{m_F + M_U = m, m_F, m_U \ge 1 \\ B_F \in \mathcal{L}^{m_F}, B_U \in \mathcal{L}_{\mathbb{R}}^{m_U}, \\ B_F - \operatorname{conj}_* B_F + B_U = B}} (B_U B_F) (H_1 B_F) ((H_1 B_F) m_U - m_F (H_1 B_U)) \binom{l-1}{l_U} G W_{B_F} \Gamma_{B_U, k_U}$$

Our aim is to perform summation of these identities over all $B \in \mathcal{L}_{\mathbb{R}}^m$ and all $H_1 \in \mathcal{L}_{\mathbb{R}}^1 \setminus \{-K\}$. As a preliminary step, we prove a few auxiliary identities which then will be used repeatedly. In what follows (similarly to above) we write H_1 and $B_i \in \mathcal{L}^{m_i}$ in a form

$$H_1 = \frac{1}{d}(-K - w), \quad B_i = \frac{1}{d}(-m_i K - v_i), \text{ where } w, v_i \in K^{\perp},$$

so that $w^2 = -d(1+d)$ and both -K - w, $-m_iK - v_i$ are divisible by d in $H_2(X)$. We let also

$$\mathfrak{W}_{\mathbb{R}} = \{ w \in K^{\perp} \cap H_2^{-}(X) \mid w^2 = -d(1+d) \text{ and } -K - w \in dH_2^{-}(X) \}.$$

5.1.2. **Proposition.** For surfaces X as in Proposition 2.2.1, the following holds. (1) If either $d \in \{1,3\}$, or d = 2 and m is even, then

- $$\begin{split} \Gamma_{m,v,k} &= \Gamma_{m,gv,k} \quad \text{for any} \quad g \in G_d \quad \text{and any} \quad v \in K^{\perp}.\\ \text{If } d &= 2, \ m \ \text{is odd, and} \ v \in K^{\perp}, \ \text{then} \quad \Gamma_{m,v,k} &= -\Gamma_{m,\gamma v,k} \quad \text{and} \\ \Gamma_{m,v,k} &= \Gamma_{m,gv,k} \quad \text{for any} \quad g \in \mathfrak{D}_4. \end{split}$$
- (2) $\sum_{w \in \mathfrak{W}_{\mathbb{R}}} w = 0$ and $\sum_{H \in \mathcal{L}_{\mathbb{R}}^1 \setminus \{-K\}} H = -\frac{1}{d} \operatorname{card} \{\mathfrak{W}_{\mathbb{R}}\} K.$
- (3) If either $d \in \{1,3\}$ or m is even, then for any $H \in \mathcal{L}^1_{\mathbb{R}} \setminus \{-K\}$ we have

$$\sum_{B \in \mathcal{L}_{\mathbb{R}}^m} \Gamma_{B,k} B = -\frac{m}{d} \Gamma_{m,k} K, \ \sum_{B \in \mathcal{L}_{\mathbb{R}}^m} (HB) \Gamma_{B,k} = \frac{m}{d} \Gamma_{m,k}.$$

- If d = 2 and m is odd, then the above sums vanish.
- (4) ² For any $v \in K^{\perp}$, $v \neq 0$, we have

$$\sum_{B \in \mathcal{L}^m} B(vB) GW_B = \frac{1}{v^2} \sum_{B \in \mathcal{L}^m} v(vB)^2 GW_B.$$

(5) For every $u, v \in K^{\perp}$, we have

$$u^2 \sum_{B \in \mathcal{L}^m} (vB)^2 GW_B = v^2 \sum_{B \in \mathcal{L}^m} (uB)^2 GW_B.$$

Proof. Proof of the first claim is a direct combination of preservation of the arithmetic genus g_a and Welschinger numbers by the action of $g \in G_d$ with the properties of their action on \hat{q} described in Proposition 2.2.1.

To prove Claims (2) and (3), except the vanishing statement in (3), it is sufficient to notice that the expressions $\sum w$, $\sum H$, and $\sum \Gamma_{B,k}B$ are invariant under the action of G_d , hence proportional to K (see Proposition 2.2.1), and that to determine the coefficient of proportionality it remains to take scalar product with K. For proving the vanishing statement, we notice that, as it follows from Claim (1), if

²Properties (4) and (5) hold without any assumption on a real structure of X.

for an arbitrary chosen $B \in \mathcal{L}_{\mathbb{R}}^m$ the number $\Gamma_{m,k}$ is not zero, then the orbit of B under the action of $\mathbb{Z}/2 \times \mathfrak{D}_4$ splits into 2 orbits of \mathfrak{D}_4 that are interchanged by γ . This implies (using the same arguments as above) that

$$\sum_{g \in \mathbb{Z}/2 \times \mathfrak{D}_4} \Gamma_{gB,k} B = \sum_{g \in \mathfrak{D}_4} \Gamma_{gB,k} gB + \sum_{g \in \mathfrak{D}_4} \Gamma_{\gamma gB,k} \gamma gB = \Gamma_{B,k} (\sum_{g \in \mathfrak{D}_4} gB - \gamma (\sum_{g \in \mathfrak{D}_4} gB)) = \lambda K - \gamma \lambda K = 0.$$

To prove Claims (4) and (5) we notice that due to the invariance of Gromov-Witten numbers GW_B under the Weyl group action on $K^{\perp} = E_{9-d}$ (which follows from to the monodromy invariance of Gromov-Witten numbers, and interpretation of the Weyl group as monodromy), these claims become straightforward consequences of the fact that the function $K^{\perp} \to \mathbb{R}$ given by $\alpha \mapsto \alpha^2$ is the only, up to scalar factor, quadratic function invariant under the action of the Weyl group (as it follows from the irreducibility of the action, see [Bo, §2.1, Prop. 1]).

Next, we note that in the left-hand side of (5.1.1) we have $(H_3B)\Gamma_{B,k} = ((\frac{1}{d} + 1)m - \frac{1}{d}wB)\Gamma_{B,k}$. Then, denoting by $\Theta_{m,k}^{(i)}$ the sum of $\Theta_{B,k}^{(i)}$ over all $H_1 \in \mathcal{L}^1_{\mathbb{R}} \setminus \{-K\}$ and $B \in \mathcal{L}^m_{\mathbb{R}}$, we conclude that

(5.1.2)
$$(\frac{1}{d}+1)\operatorname{card}(\mathcal{L}^1_{\mathbb{R}}\smallsetminus\{-K\})m\Gamma_{m,k} = \Theta_{m,k}^{(1)} + \Theta_{m,k}^{(2)}$$

since the terms $(wB)\Gamma_{B,k}$ vanish after summation due to Proposition 5.1.2(2).

5.2. **Proof of Theorem 1.2.4 for odd** $\mathbf{m} = 2\mathbf{n} + \mathbf{1} \ge \mathbf{3}$. Due to Proposition 2.1.2(2), this part of Theorem 1.2.4 holds trivially for d = 2. Therefore, here we may assume (for simplicity) that $d \ne 2$.

We let k = 0 and have

$$\Theta_{m,0}^{(1)} = \frac{1}{4} \sum_{\substack{m_1+m_2=m, m_i \geqslant 1\\ B_i \in \mathcal{L}_{\mathbb{R}}^{m_i}, H_1 \in \mathcal{L}_{\mathbb{R}}^1}} {\binom{l-1}{l_1} (H_1B_1)((H_1B_1)m_2 - (H_1B_2)m_1)\Gamma_{B_1,0}\Gamma_{B_2,1}}$$
$$= \frac{1}{4} \sum_{H_1 \in \mathcal{L}_{\mathbb{R}}^1} \sum_{\substack{m_1+m_2=m\\ m_1,m_2 \geqslant 1}} \sum_{B_2 \in \mathcal{L}_{\mathbb{R}}^{m_2}} {\binom{l-1}{l_1}} m_2\Gamma_{B_2,1} \sum_{B_1 \in \mathcal{L}_{\mathbb{R}}^{m_1}} (H_1B_1)^2\Gamma_{B_1,0}$$
$$- \frac{1}{4} \sum_{H_1 \in \mathcal{L}_{\mathbb{R}}^1} \sum_{\substack{m_1+m_2=m\\ m_1,m_2 \geqslant 1}} {\binom{l-1}{l_1}} m_1 (\sum_{B_1 \in \mathcal{L}_{\mathbb{R}}^m} (H_1B_1)\Gamma_{B_1,0} \sum_{B_2 \in \mathcal{L}_{\mathbb{R}}^{m_2}} (H_1B_2)\Gamma_{B_2,1}),$$

$$\begin{aligned} \Theta_{m,0}^{(2)} &= \frac{1}{2} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1 \\ B_F \in \mathcal{L}^{m_F}, B_U \in \mathcal{L}_{\mathbb{R}}^{m_U}, H_1 \in \mathcal{L}_{\mathbb{R}}^1 \\ B_F \in \mathcal{L}^{m_F}, B_U \in \mathcal{L}_{\mathbb{R}}^{m_U}, H_1 \in \mathcal{L}_{\mathbb{R}}^1 \\ &= \frac{1}{2} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1 \\ H_1 \in \mathcal{L}_{\mathbb{R}}^1}} \left(\sum_{B_U \in \mathcal{L}^{m_U}} m_U B_U \binom{l-1}{l_U} \Gamma_{B_U, 0} \cdot \sum_{B_F \in \mathcal{L}^{m_F}} B_F (H_1 B_F)^2 G W_{B_F} \right) \\ &- \frac{1}{2} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1 \\ H_1 \in \mathcal{L}_{\mathbb{R}}^1}} \left(\sum_{B_U \in \mathcal{L}_{\mathbb{R}}^{m_U}} B_U (H_1 B_U) \binom{l-1}{l_U} \Gamma_{B_U, 0} \cdot \sum_{B_F \in \mathcal{L}^{m_F}} m_F B_F (H_1 B_F) G W_{B_F} \right). \end{aligned}$$

In the last relation, symbol "." is used to denote the intersection index in $H_2(X)$.

Substituting $H_1 = \frac{1}{d}(-K - w)$ and observing the vanishing (due to Proposition 5.1.2(2)) of summands where the factors (wB_i) enter linearly, we obtain

$$\Theta_{m,0}^{(1)} = \frac{1}{4d^2} \sum_{w \in \mathfrak{W}} \Big(\sum_{\substack{m_1 + m_2 = m \\ m_i \ge 1, B_i \in \mathcal{L}_{\mathbb{R}}^{m_i}}} \binom{l-1}{l_1} \Big) \Big(m_2 (wB_1)^2 - m_1 (wB_1) (wB_2) \Big) \Gamma_{B_1,0} \Gamma_{B_2,1} \Big).$$

Next, we note that the terms with a factor $(wB_2)\Gamma_{B_2,1}$ vanish after summation over B_2 , since by Proposition 5.1.2(3), $\sum B_2\Gamma_{B_2,k}$ is collinear with K, which is orthogonal to ω . Thus, we get

(5.2.1)
$$\Theta_{m,0}^{(1)} = \frac{1}{4d^2} \sum_{\substack{m_1+m_2=m\\m_1,m_2 \ge 1}} \left(\binom{l-1}{l_1} m_2 \Gamma_{m_2,1} \sum_{\substack{B_1 \in \mathcal{L}_{\mathbb{R}}^{m_1}\\w \in \mathfrak{W}_{\mathbb{R}}}} (wB_1)^2 \Gamma_{B_1,0} \right).$$

Similarly, after the same substitution for H_1 into the first summand of $\Theta_{m,0}^{(2)}$, we apply Proposition 5.1.2(3) to the factor containing $\sum B_U \Gamma_{B_U,0}$ and obtain

$$\frac{1}{2d^3} \sum_{\substack{w \in \mathfrak{W}_{\mathbb{R}} \\ 2m_F + m_U = m \\ m_F, m_U \geqslant 1}} {\binom{l-1}{l_U}} m_U^2 m_F \Gamma_{m_U,0} \sum_{B_F \in \mathcal{L}^{m_F}} (m_F - (wB_F))^2 GW_{B_F}.$$

Then, cancelation in $(m_F - (wB_F))^2$ of linear in w term (since $\sum_{w \in \mathfrak{M}_{\mathbb{R}}} w = 0$), gives

$$\frac{1}{2d^3} \sum_{\substack{w \in \mathfrak{M}_{\mathbb{R}} \\ 2m_F + m_U = m \\ m_F, m_U \geqslant 1}} {\binom{l-1}{l_U}} m_U^2 \Gamma_{m_U,0} \left(m_F^3 N_{m_F}^{GW} + m_F \sum_{B_F \in \mathcal{L}^{m_F}} (wB_F)^2 GW_{B_F} \right).$$

In the second summand of $\Theta_{m,0}^{(2)}$, after the same substitution for H_1 and cancelation of linear in w terms, we apply Proposition 5.1.2(4) (with the choice v = w) and obtain

$$-\frac{1}{2d^{3}}\sum_{\substack{w\in\mathfrak{W}_{\mathbb{R}}\\2m_{F}+m_{U}=m\\m_{F},m_{U}\geqslant1}} \binom{l-1}{l_{U}}m_{F}^{3}m_{U}^{2}\Gamma_{m_{U},0}N_{m_{F}}^{GW}$$

$$+\frac{1}{2d^{3}(1+d)}\sum_{\substack{w\in\mathfrak{W}_{\mathbb{R}}\\2m_{F}+m_{U}=m\\m_{F},m_{U}\geqslant1}} (\sum_{B_{U}\in\mathcal{L}_{\mathbb{R}}} \binom{l-1}{l_{U}}(wB_{U})^{2}\Gamma_{B_{U},0}\sum_{B_{F}\in\mathcal{L}^{m_{F}}}m_{F}(wB_{F})^{2}GW_{B_{F}}).$$

Afterwards we cancel two opposite terms in the sum and conclude that

$$\Theta_{m,0}^{(2)} = \frac{1}{2d^3} \sum_{\substack{w \in \mathfrak{W}_{\mathbb{R}} \\ 2m_F + m_U = m \\ m_F, m_U \geqslant 1}} (\binom{l-1}{l_U} m_F m_U^2 \Gamma_{m_U,0} \sum_{B_F \in \mathcal{L}^{m_F}} (wB_F)^2 GW_{B_F}) +$$

$$(5.2.2) \qquad \frac{1}{2d^3(1+d)} \sum_{\substack{w \in \mathfrak{W}_{\mathbb{R}} \\ B_U \in \mathcal{L}_{\mathbb{R}}^{m_U}}} (\sum_{l_U} \binom{l-1}{l_U} (wB_U)^2 \Gamma_{B_U,0} \sum_{B_F \in \mathcal{L}^{m_F}} m_F (wB_F)^2 GW_{B_F}).$$

Now, we substitute the expressions obtained in (5.2.1) and (5.2.2) into (5.1.2) and observe that, as it follows from Proposition 4.2.2 where we transform e into w in accordance with Proposition 5.1.2(5), the second term in (5.2.2) cancels (5.2.1). In this way we get

$$\frac{1+d}{d}\operatorname{card}(\mathcal{L}^{1}_{\mathbb{R}}\smallsetminus\{-K\})m\Gamma_{m,0} = \frac{d(1+d)}{4d^{3}}\operatorname{card}\mathfrak{W}_{\mathbb{R}}\sum_{\substack{2m_{F}+m_{U}=m\\m_{F},m_{U}\geqslant1}} \left(m_{F}m_{U}^{2}\binom{l-1}{l_{U}}\right)\Gamma_{m_{U},0}\sum_{B_{F}\in\mathcal{L}^{m_{F}}} (eB_{F})^{2}GW_{B_{F}}\right).$$

Finally, division by $\frac{1+d}{2d^3} \operatorname{card}(\mathcal{L}^1_{\mathbb{R}} \setminus \{-K\}) = \frac{1+d}{2d^3} \operatorname{card} \mathfrak{W}_{\mathbb{R}}$ and using once more Proposition 4.2.2 gives

$$2dm\Gamma_{m,0} = \frac{1}{2} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \ge 1}} \left(m_F m_U^2 \binom{l-1}{l_U} \Gamma_{m_U,0}(-2\Gamma_{2m_F,1}) \right)$$

which proves the first relation of Theorem 1.2.4 (cf. its reformulation (4.1.3)).

5.3. Proof of Theorem 1.2.4 for $\mathbf{m} = 2\mathbf{n} + 2 \ge 4$. We let k = 1 and have:

(5.3.1)
$$\frac{1+d}{d}\operatorname{card}(\mathcal{L}^{1}_{\mathbb{R}}\smallsetminus\{-K\})m\Gamma_{m,1} = \Theta_{m,1}^{(1)} + \Theta_{m,1}^{(2)},$$

(5.3.2)
$$\Theta_{m,1}^{(1)} = \frac{1}{4} \sum_{\substack{m_1+m_2=m,m_i \ge 1 \\ B_i \in \mathcal{L}_{\mathbb{R}}^{m_i}, H_1 \in \mathcal{L}_{\mathbb{R}}^1 \\ B_i \in \mathcal{L}_{\mathbb{R}}^{m_i}, H_1 \in \mathcal{L}_{\mathbb{R}}^1}} {\binom{l-1}{l_1} (H_1 B_1) ((H_1 B_1) m_2 - (H_1 B_2) m_1) \Gamma_{B_1,1} \Gamma_{B_2,1}}{\frac{1}{4d^4} \sum_{\substack{m_1+m_2=m,m_i \ge 1 \\ m_1+m_2=m,m_i \ge 1}} {\binom{l-1}{l_1} m_2 \Gamma_{m_2,1}} \sum_{\substack{w \in \mathfrak{W}_{\mathbb{R}}, B_1 \in \mathcal{L}_{\mathbb{R}}^{m_1}}} (w B_1)^2 \Gamma_{B_1,1} \right),$$

$$\Theta_{m,1}^{(2)} = \frac{1}{2} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1 \\ B_F \in \mathcal{L}^{m_1, B_U} \in \mathcal{L}_{\mathbb{R}}^{m_2} \\ H_1 \in \mathcal{L}_{\mathbb{R}}^{\frac{1}{2}}}} (B_U B_F) (H_1 B_F) ((H_1 B_F) m_U - (H_1 B_U) m_F) \binom{l-1}{l_U} G W_{B_F} \Gamma_{B_U,1} \\ (5.3.3) = \frac{d}{2d^6} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1}} \binom{l-1}{l_U} m_F m_U^2 \Gamma_{m_U,1} \sum_{w \in \mathfrak{W}_{\mathbb{R}}, v_F \in K^{\perp}} (wv_F)^2 G W_{B_F} - \frac{1}{2d^6} \sum_{\substack{2m_F + m_U = m \\ m_F, m_U \geqslant 1}} \binom{l-1}{l_U} (v_F v_U) (wv_F) (wv_U) G W_{B_F} \Gamma_{B_U,1}.$$

We substitute (5.3.2) and (5.3.3) into (5.3.1), apply Proposition 5.1.2(5) where we choose u = w and v = e with $e^2 = -2, e \in K^{\perp}$ to perform a transformation

$$\sum_{v_F \in K^{\perp}} (wv_F)^2 GW_{B_F} = d^2 \sum_{B_F \in \mathcal{L}^{m_F}} (wB_F)^2 GW_{B_F} = \frac{d^3(1+d)}{2} \sum_{B_F \in \mathcal{L}^{m_F}} (eB_F)^2 GW_{B_F} \stackrel{\text{Prop.4.2.2}}{=} -d^3(1+d)\Gamma_{2m_F,1}$$

cancel similar terms in $\Theta_{m,1}^{(1)}+\Theta_{m,1}^{(2)}$ and get

(5.3.4)
$$\frac{1+d}{d}m\Gamma_{m,1} = -\frac{d^3(1+d)}{2\cdot d^5} \sum_{\substack{2m_F+m_U=m\\m_F,m_U \ge 1}} \binom{l-1}{l_U} m_F m_U^2 \Gamma_{m_U,1} \Gamma_{2m_F,1}$$

wherefrom the required recursion relation (cf. (4.1.3))

$$2md\Gamma_{m,1} + \sum_{j=1}^{n} \left(\binom{n-1}{n-j} j(m-2j)^2 \Gamma_{m-2j,1} \Gamma_{2j,1} = 0 \quad \text{for } m = 2n+2. \quad \Box$$

5.4. **Explicit formulas.** The recursive formulas of Theorem 1.2.4 have a surprisingly simple explicit solution.

5.4.1. Theorem. For each d = 1, 2, 3 and any $n \in \mathbb{Z}_{\geq 0}$, we have

$$N_{2n+1,0} = \frac{1}{4} N_{1,0} b^n \left(n + \frac{1}{2}\right)^{n-2}, \quad N_{2n+2,1} = N_{2,1} b^n \left(n + 1\right)^{n-2} \quad with \quad b = \frac{4N_{2,1}}{d}$$

Proof. Both relations stated in Theorem 1.2.4 hold trivially for these values if n = 0. To check the second of these two relations for n > 0 we substitute there the given values of $N_{2k,1}$, $k \leq n + 1$ and get

$$(2n+2)dN_{2,1}b^{n}(n+1)^{n-2} = 2\sum_{j=1}^{n} \binom{n-1}{n-j} j(2n+2-2j)^{2}N_{2,1}b^{n-j}(n-j+1)^{n-j-2}N_{2,1}b^{j-1}j^{j-3}$$

division by $2d N_{2,1}b^n = 8N_{2,1}^2b^{n-1}$ it gives

$$(n+1)^{n-1} = \sum_{j=1}^{n} \binom{n-1}{n-j} (n-j+1)^{n-j} j^{j-2}$$

which is a special case of Abel's binomial theorem (see, for example, [Co])

$$\frac{(x+y)^m}{x} = \sum_{k=0}^m \binom{m}{k} (x-kz)^{k-1} (y+kz)^{m-k}$$

where we put x = -z = 1, y = n, m = n - 1, k = j - 1.

Substituting the given values into the first relation we obtain

$$(2n+1)d\frac{1}{4}N_{1,0}b^{n}(n+\frac{1}{2})^{n-2} = 2\sum_{j=1}^{n} \binom{n-1}{n-j} j(2n+1-2j)^{2}\frac{1}{4}N_{1,0}b^{n-j}(n-j+\frac{1}{2})^{n-j-2}N_{2,1}b^{j-1}j^{j-3}.$$

Division by $\frac{1}{2}dN_{1,0}b^n = 2N_{1,0}b^{n-1}N_{2,1}$ turns it into

$$(n+\frac{1}{2})^{n-1} = \sum_{j=1}^{n} \binom{n-1}{n-j} (n-j+\frac{1}{2})^{n-j} j^{j-2}$$

which is Abel's binomial relation for $x = -z = 1, y = n - \frac{1}{2}, m = n - 1, k = j - 1$. \Box

5.4.2. Corollary. For real del Pezzo surfaces X of degree d = 1 or d = 2 whose real locus $X_{\mathbb{R}}$ has the maximal number of connected components, the invariant $N_{dm,k}$ coincides with the Welschinger invariant $W_{-mK,k}$. In particular, for d = 1 and $X_{\mathbb{R}} = \mathbb{RP}^2 \perp 4S^2$ we have

$$W_{-(2n+1)K,0} = 2 (120)^n (n + \frac{1}{2})^{n-2},$$

$$W_{-(2n+2)K,1} = 30 (120)^n (n+1)^{n-2},$$

and for d = 2 and $X_{\mathbb{R}} = \bot 4S^2$

$$W_{-(n+1)K,1} = 6 (12)^n (n+1)^{n-2}.$$

Proof. This is immediate from Theorem 5.4.1 and knowledge of values of $N_{1,0}, N_{2,1}$ (see Table below), since under the assumptions imposed on $X_{\mathbb{R}}$ the eighenspace $\ker(1 + \operatorname{conj}_*) \subset H_2(X)$ is generated by K and, in addition, $\hat{q}(mK) - m^2 = m^2 \hat{q}(K) - m^2 = 0$.

5.5. Few first values and simple applications. To begin, we list the values of $N_{m,k}$ for $m \leq 6$ obtained by means of the formulas established above. The initial values $N_{1,0}, N_{2,1}$ we use there:

- for $K^2 = 1$, were obtained in [FK-2], [FK-3] via some "lattice calculation",
- for $K^2 = 2$, can be obtained in a similar way,
- for $K^2 = 3$, the value $N_{1,0} = 3$ was explained in [FK-1], and $N_{2,1} = N_{1,0}$ follows from a natural one-to-one correspondence $\mathcal{L}^1_{\mathbb{R}} \to \mathcal{L}^2_{\mathbb{R}}$ assigning to a line the divisor class of residual conics of hyperplane sections containing this line.

$N_{m,k}$				$ \Gamma_{m,k} $				
(m,k)	$K^2 = 1$	$K^2=2$	$K^2 = 3$		(m,k)	$K^2 = 1$	$K^2 = 2$	$K^2 = 3$
(1,0)	8	0	3		(1,0)	16	0	6
(2,1)	30	6	3		(2,1)	60	12	6
(3,0)	160	0	2		(3,0)	160	0	2
(4,1)	1800	36	6		(4,1)	1800	36	6
(5,0)	28800	0	12		(5,0)	14400	0	6
(6,1)	432000	864	48		(6,1)	216000	432	24

5.5.1. **Proposition.** For any nonsingular real cubic surface X:

- The signed count of 2-points-constrained real twisted cubics is equal to $3 \chi(X_{\mathbb{R}})$, independently on the number of real points in the points-constraint. When the surface is maximal, it contains h = 40 hyperbolic (with $\hat{q} = 1$) and e = 32 elliptic (with $\hat{q} = -1$) twisted cubics.
- The signed count of 3-points-constrained real non-singular rational quartic curves is equal to $9 3\chi(X_{\mathbb{R}})$, independently on the number of real points in the points-constraint. When the surface is maximal, it contains h = 120 hyperbolic (with $\hat{q} = 0$) and e = 96 elliptic (with $\hat{q} = 2$) quartics.

Proof. The third layer is formed by -K and divisor classes of twisted cubics. Therefore, the first statement follows from $N_{3,k} = 2 - k$ and $W_{-K,k} = \chi(X_{\mathbb{R}}) - (k+1)$. For a separate count of hyperbolic and elliptic twisted cubics, it is sufficient to notice, in addition, that their number is equal to the number of roots, 72, in E_6 .

The fourth layer is formed by divisor classes representable by non-singular rational quartic curves (that is by divisor classes of form -2K - L' - L'' where L', L'' any pair of disjoint lines) and genus 1 quartic curves (that is by divisor classes -K + Lwhere L is any line). Therefore, the second statement follows from $N_{4,k} = 9 - 4K$, $W_{-K+L,k} = \chi(X_{\mathbb{R}}) - k$ and $\hat{q}(-K+L) = \hat{q}(L) - 1$. For a separate count of hyperbolic and elliptic quartics, we note that the number of pairs of disjoint real lines on a maximal cubic surface is equal to $\frac{27\cdot16}{2} = 216$.

5.5.2. *Remark.* In the case of cubic surfaces, there are natural bijections: between the set of divisor classes of twisted cubics (resp. non-singular rational quartic curves) and the set of isomorphism classes of presentations of the surface as 6-blowup of \mathbb{P}^2 (resp. as 5-blowup of $\mathbb{P}^1 \times \mathbb{P}^1$). Therefore, cited counts can be interpreted as signed counts of the corresponding blow-up models.

Considering real del Pezzo surfaces X with $K^2 = 2$ as double coverings of \mathbb{P}^2 branched along real non-singular quartic curves $A \subset \mathbb{P}^2$ and using skew-invariance of \hat{q} under the deck transformation, we translate our signed count of curves belonging to even layers $\mathcal{L}_{\mathbb{R}}^{2n}$ into a signed count of real rational point-constrained curves of degree *n* tangent to *A* at 2n - 1 points plus a signed count of real rational pointconstrained curves of degree 2n tangent to *A* at 4n points. In particular, such a point-constrained curve of degree 2n is called *hyperbolic* (resp. *elliptic*), if \hat{q} takes value 0 (resp. 2) on its lifts to X.

5.5.3. **Proposition.** Let $A \subset \mathbb{P}^2$ be a real non-singular quartic curve and Ω one of two halves of \mathbb{RP}^2 bounded by $A_{\mathbb{R}}$. Then:

• The signed count of real conics 4-tangent to A and constrained by a point in Ω gives $8 - 2\chi(\Omega)$. If A is maximal and Ω is non-orientable, then this count involves h = 70 hyperbolic and e = 56 elliptic conics.

WALL-CROSSING INVARIANCE

 If A is maximal and Ω is non-orientable, then signed count of real nonsingular rational quartic curves 8-tangent to with k real point-constraints chosen in Ω gives 336 if k = 1 and 896 if k = 3.

Proof. The second layer, \mathcal{L}^2 , is formed by -K and the divisor classes of lifts of 4-tangent conics. Therefore, the first statement follows from $N_{2,1} = 6$ and $W_{-K,1} = \chi(X_{\mathbb{R}}) - 2 = 2\chi(\Omega) - 2$. In the maximal case, $8 - 2\chi(\Omega) = 14$ and the number of 4-tangent conics is the number of roots, 126, in E_7 .

The fourth layer is formed by -2K and the divisor classes of lifts of 8-tangent quartics. Here we obtain the required number of quartics, $4 \cdot 224$ if k = 3 and 2(132 + 36) if k = 1, due to $N_{4,k} = 0$, $W_{-2K,k} = -224$ for k = 3 and $N_{4,k} = 36$, $W_{-2K,k} = -132$ for k = 1 (see [IKS-2] for these values of $W_{-2K,k}$).

Considering real del Pezzo surfaces X with $K^2 = 1$ as double coverings $\pi : X \to Q$ of a real quadric cone $Q \subset \mathbb{P}^2$ branched along real non-singular sextics $C \subset Q$ and using invariance of \hat{q} under the deck transformation, we translate our signed count of curves belonging to the second layer $\mathcal{L}^2_{\mathbb{R}}$ into a signed count of real rational 1-point-constrained hyperplane sections tangent to C at 2 points plus a signed count of real 1-point-constrained quartics tangent to C at 6 points (and obtained as transversal sections of Q by quadrics). In particular, such a point-constrained quartic is called *hyperbolic* (resp. *elliptic*), if \hat{q} takes value 0 (resp. 2) on its lifts to X.

5.5.4. **Proposition.** For a real non-singular sextic $C \subset Q$, the signed count of real quartics 6-tangent to C constrained by a point in $\pi(X_{\mathbb{R}})$ gives $6 + \frac{\chi^2(X_{\mathbb{R}})-1}{2}$. If X is maximal and $X_{\mathbb{R}}$ is connected, then this count involves h = 1192 hyperbolic and e = 1208 elliptic quartics.

Proof. Follows from
$$N_{2,1} = 30$$
 and $W_{-2K,1} = 6 + \frac{\chi^2(X_{\mathbb{R}}) - 1}{2}$ (see [FK-3]).

6. Concluding Remarks

6.1. Generating functions. Recall the tree function $T(x) = \sum_{n \ge 1} n^{n-1} \frac{x^n}{n!}$.

6.1.1. **Proposition.** For each d = 1, 2 and 3, the functions

$$N^{even}(x) = \sum_{n \ge 0} N_{2n+2,1} \frac{x^n}{n!}, \quad N^{odd}(x) = \sum_{n \ge 0} N_{2n+1,0} \frac{x^n}{n!}$$

can be expressed through T(x) as

$$N^{even}(x) = N_{2,1}(bx)^{-1}(T(bx) - \frac{1}{2}T^2(bx)), \quad N^{odd}(x) = N_{1,0}(bx)^{-\frac{1}{2}}(T^{\frac{1}{2}}(bx) - \frac{1}{3}T^{\frac{3}{2}}(bx))$$

where b is 120, 12 and 4 for $d = 1, 2, 3$ respectively.

 $a_{1} = 1, 2, 0$ respectively.

Proof. After substitution of the values of $N_{2n+2,1}$ from Theorem 5.4.1 we obtain

$$N^{\text{even}}(x) = N_{2,1} \sum_{n \ge 0} (n+1)^{n-1} \frac{(bx)^n}{(n+1)!} = N_{2,1} G(bx), \text{ where}$$
$$G(x) = \sum_{n \ge 1} n^{n-2} \frac{x^{n-1}}{n!} = \frac{1}{x} (T(x) - \frac{1}{2} T^2(x)).$$

For the latter identity, see, *i.e.*, [M]. For values of $b = \frac{4N_{2,1}}{d}$ see Section 5.5.

A similar substitution for $N_{2n+1,1}$ gives

$$N^{\text{odd}}(x) = \sum_{n \ge 0} \frac{1}{4} N_{1,0} b^n \left(n + \frac{1}{2}\right)^{n-2} \frac{x^n}{n!} = N_{1,0} Q(bx) \quad \text{with}$$
$$Q(x) = \sum_{n \ge 0} \frac{1}{4} \left(n + \frac{1}{2}\right)^{n-2} \frac{x^n}{n!} = x^{-\frac{1}{2}} \int_0^x \frac{1}{2} x^{-\frac{1}{2}} e^{\frac{1}{2}T(x)} dx = x^{-\frac{1}{2}} \left(T^{\frac{1}{2}}(x) - \frac{1}{3}T^{\frac{3}{2}}(x)\right)$$

where the last equalities follow from the following relations (see, f.e., [CGHJK])

$$\sum \frac{1}{2} (n + \frac{1}{2})^{n-1} \frac{x^n}{n!} = e^{\frac{1}{2}T(x)} = \left(\frac{T(x)}{x}\right)^{\frac{1}{2}}.$$

6.2. On Gromov-Witten side. It is thought-provoking that over the complex field, for any del Pezzo surface, the sums of genus-0 Gromov-Witten invariants over layers \mathcal{L}^m satisfy essentially the same recursion relation as genus-0 Gromov-Witten invariants of projective plane.

6.2.1. **Proposition.** For every del Pezzo surface of degree $1 \le d \le 6$ and every $m \ge 4$,

(6.2.1)
$$d^2 N_m^{GW} = \sum_{\substack{m_1 + m_2 = m \\ m_1, m_2 \ge 1}} N_{m_1}^{GW} N_{m_2}^{GW} m_1^2 m_2 \left(m_2 \binom{m-4}{m_1-2} - m_1 \binom{m-4}{m_1-1} \right)$$

where N_m^{GW} stands for $\sum_{\alpha \in \mathcal{L}^m(X)} GW(\alpha)$. The initial values of N_m^{GW} for m = 1, 2, 3 and $1 \leq d \leq 6$ are shown in the table below.

	d=1	d=2	d=3	d=4	d=5	d = 6
N_1^{GW}	252	56	27	16	10	6
$N_2^{\overline{GW}}$	5130	138	27	10	5	3
$N_3^{\overline{GW}}$	446400	344	84	16	5	2

It is not difficult to show that this implies the following square root rule.

6.2.2. Corollary. For any $1 \leq d \leq 6$ one has $\log N_{2m}^{GW} = 2m \log m + O(m)$, while $\log N_{2m,1}$ for d = 1, 2, 3 and $\log N_{2m+1,0}$ for d = 1, 3 are of type $m \log m + O(m)$ (recall that $N_{2m+1,0} = 0$ for d = 2).

In the proof of Proposition 6.2.1 we use (in a similar, but much simpler, way as in out proof of recursion relations for $N_{m,k}$) the fact that in the case of $d \leq 6$ the action of the Weyl group on K^{\perp} has no any nonzero invariant element. Note that, as an elementary check shows, the relation (6.2.1) does not hold for d = 7 and 8. When d = 9, it turns (after replacing m_i by $3d_i$) into celebrated Kontsevich-Manin recursion relation.

The next formula is an immediate consequence of Proposition 4.2.2 combined with Theorem 5.4.1.

6.2.3. **Proposition.** $\sum_{\alpha \in \mathcal{L}^n} (e\alpha)^2 GW_{\alpha} = 2da^n n^{n-3}$ where *e* is an arbitrary class $e \in K^{\perp}$ with $e^2 = -2$ and *a* equals 60 if d = 1, 6 if d = 2, and 2 if d = 3.

6.3. Other del Pezzo surfaces. There are two more kinds of del Pezzo surfaces to which our approach applies almost literally. One of them is real nonsingular quadrics $X \subset P^3$ with $X_{\mathbb{R}} \neq \emptyset$. Such non-empty quadrics form two real deformation classes: one with $X_{\mathbb{R}} = S^2$ and another with $X_{\mathbb{R}} = S^1 \times S^1$. Pick a Spin-structure on $X_{\mathbb{R}}$ induced from a Spin-structure on \mathbb{RP}^3 , and let q: $H_1(X_{\mathbb{R}}; \mathbb{Z}/2) \to \mathbb{Z}/2$ denote the associated quadratic function. We put in this case $N_{m,k} = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x})} (-1)^{q(A_{\mathbb{R}})} w(A)$ and obtain the following results.

- The numbers $N_{m,k}$ are independent of the choice of a real quadric X with $X_{\mathbb{R}} \neq \emptyset$ and a Spin-structure on \mathbb{RP}^3 .
- The numbers $N_{m,k}$ vanish unless m = 4n, while $N_{4n,k}$ is equal to the The numbers N_{m,k} vanish timess mⁱ = 4n, while N_{4n,k} is equal to the Welschinger invariant W_{nh,k}, h = -¹/₂K, of X with X_ℝ = S² (see [Wel2], [IKS-1], [Br-P], [Ch-Z] for various methods of calculation of W_{nh,k}).
 The relation N_{4n,1} = 2²ⁿ⁻³ Σ_{α∈L²ⁿ}(αe)²GW_α (where e is an arbitrary class e ∈ K[⊥] with e² = -2) holds for any n ≥ 1.

Another example is provided by del Pezzo surfaces X of degree 4 presented as double coverings of real non-singular quadrics $Q \subset \mathbb{P}^3$ branched along real nonsingular curves representing the doubled hyperplane section class. The covering $X \to Q$ is naturally embedded into the quadratic cone $Z \subset \mathbb{P}^4$ over Q. Thus, we can pick a Pin⁻-structure on $X_{\mathbb{R}}$ induced from a Pin⁻-structure on the non-singular part of $Z_{\mathbb{R}}$ and put $N_{m,k} = \sum_{A \in \mathcal{C}_{\mathbb{R}}(m,k,\mathbf{x})} i^{\hat{q}([A])-m^2} w(A)$, as usual.

- The numbers $N_{m,k}$ are preserved under change of the branching curve.
- The numbers $N_{m,k}$ vanish if either m is odd or k > 1.
- The relation $N_{2n,1} = 2^{n-3} \sum_{\alpha \in \mathcal{L}^n} (\alpha e)^2 GW_{\alpha}$ (where e is an arbitrary class $e \in K^{\perp}$ with $e^2 = -2$) holds for any $n \ge 1$.

References

- [Bo] N. BOURBAKI, Groupes et Algèbres de Lie, Ch. 4, 5 et 6. Hermann, Paris, 1968.
- [Br] E. BRUGALLÉ, On the invariance of Welschinger invariants. Algebra i Analiz, 32 (2020), 1 - 20.
- [Br-P] E. BRUGALLÉ, N. PUIGNAU On Welschinger invariants of symplectic 4-manifolds. Comment. Math. Helv. 90 (2015), 905 - 938.
- [Ch] X. CHEN, Steenrod pseudocycles, lifted cobordisms, and Solomon's relations for Welschinger's invariants. arXiv: 1809.089v2 (2019), 70 pages.
- [Ch-Z] X. CHEN, A. ZINGER WDVV-type relations for Welschinger invariants: Applications. Kyoto J. Math. bf 61 (2021), 339 - 376.
- [Co] L. COMTET, Advanced Combinatorics: The Art of Finite and Infinite Expansions. D. Reidel Publishing Company, Dordrecht, 1974.
- [CGHJK] R.M.CORLESS, G.H.GONNET, D.E.G. HARE, D.J. JEFFREY, D.E. KNUTH, On the Lambert W function. Advances in Comput. Math., 5 (1996), 329 - 359.
- [DIK] A. DEGTYAREV, I. ITENBERG, V. KHARLAMOV, Real Enriques Surfaces. Springer, Lecture Notes in Mathematics 1746, 2000.
- [Do] I.V. DOLGACHEV, Classical algebraic geometry. A modern view. Cambridge University Press, Cambridge, 2012.
- [FK-1] S. FINASHIN, V. KHARLAMOV, Abundance of real lines on real projective hypersurfaces. Int. Math. Res. Not. IMRN 2013, 46.
- [FK-2] S. FINASHIN, V. KHARLAMOV, Two kinds of real lines on real del Pezzo surfaces of degree 1. Selecta Math. (N.S.) 27 (2021), no. 5, Paper No. 83, 23 pp.
- [FK-3] S. FINASHIN, V. KHARLAMOV, Combined count of real rational curves of canonical degree 2 on real del Pezzo surfaces with $K^2 = 1$. arXiv:2107.06988, 21 pp.
- [G-P] L. GÖTTSCHE, R.PANDHARIPANDE The quantum cohomology of blow-ups of \mathbb{P}^2 and enumerative geometry J. Differential Geom. 48 (1998), no. 1, 61 - 90.

- [H-S] A. HOREV, J. P. SOLOMON The open Gromov-Witten-Welschinger theory of blowups of projective plane. arXiv:1210.4034, 34 pages.
- [IKS-1] I. ITENBERG, E. SHUSTIN, V. KHARLAMOV, Welschinger invariants of small non-toric Del Pezzo surfaces. J. Eur. Math. Soc. (JEMS) 15 (2013), no. 2, 539–594.
- [IKS-2] I. ITENBERG, E. SHUSTIN, V. KHARLAMOV, Welschinger invariants of real del Pezzo surfaces of degree ≥ 2. International J. Math. 26 (2015), no. 6., 1550060, 63 pp.
- [IKS-3] I. ITENBERG, E. SHUSTIN, V. KHARLAMOV, Welschinger invariants revisited. in book "Analysis meets geometry: A Tribute to Mikael Passare", Springer International Publishing, Trends in Math., 2017, 239 – 260,
- [K-T] R. KIRBY, L. TAYLOR, Pin-Structures on Low-Dimensional Manifolds. Geometry of Lowdimensional Manifolds, 2. Ed. S.K. Donaldson, C.B. Thomas, London Math. Soc. Lect. Notes (151), Cambridge Univ. Press, 1991
- [K-M] M. KONTSEVICH, YU. MANIN, Gromov-Witten classes, Quantum Cohomology, and Enumerative Geometry. Commun. Math. Physics, 164(1994), 525 - 562.
- [M] J.W. MOON, Counting labelled trees. Canadian Mathematical Monographs 1, 1970. 113 pp.
- [N] V. V. NIKULIN, Integer symmetric bilinear forms and some of their geometric applications. Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 1, 111–177.
- [O-T] C. OKONEK, A. TELEMAN, Intrinsic signs and lower bounds in real algebraic geometry. J. Reine Ang. Math. 688 (2014), 219 - 241.
- [S1] J. P. SOLOMON Intersection theory on the moduli space of holomorphic curves with Lagrangian boundary conditions. arXiv:math/0606429 2006, 79 pages.
- [S2] J. P. SOLOMON A differential equation for the open Gromov-Witten potential. preprint 2007, 66 pages.
- [T] F. TEHRANI Open Gromov-Witten theory on symplectic manifolds and symplectic cutting. Advances in Mathematics 232 (2013), 238 – 270.
- [Wal] J. WALCHER, Opening mirror symmetry on the quintic. Communications in Mathematical Physics **276** (2007), no. 3, 671 689.
- [Wel1] J.-Y. WELSCHINGER, Invariants of real symplectic 4-manifolds and lower bounds in real enumerative geometry. Invent. Math. 162 (2005), no. 1, 195–234.
- [Wel2] J.-Y. WELSCHINGER, Optimalité, congruences et calculs d'invariants des variétés symplectiques réelles de dimension quatre. arXiv:0707.4317, 42 pages.