

# Linear Convergence of Generalized Proximal Point Algorithms for Monotone Inclusion Problems

Hui Ouyang\*

March 25, 2022

## Abstract

We focus on the linear convergence of generalized proximal point algorithms for solving monotone inclusion problems. Under the assumption that the associated monotone operator is metrically subregular or that the inverse of the monotone operator is Lipschitz continuous, we provide  $Q$ -linear and  $R$ -linear convergence results on generalized proximal point algorithms. Comparisons between our results and related ones in the literature are presented in remarks of this work.

**2020 Mathematics Subject Classification:** Primary 47J25, 47H05; Secondary 65J15, 90C25, 90C30.

**Keywords:** Proximal point algorithm, monotone inclusion problems, resolvent, firmly nonexpansiveness, linear convergence

## 1 Introduction

Throughout this work,

$\mathcal{H}$  is a real Hilbert space,

with inner product  $\langle \cdot, \cdot \rangle$  and induced norm  $\|\cdot\|$ .

Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ . Denote the set of all nonnegative integers by  $\mathbb{N} := \{0, 1, 2, \dots\}$ . The iteration sequence of the *generalized proximal point algorithm* is generated by conforming to the iteration scheme:

$$(\forall k \in \mathbb{N}) \quad x_{k+1} = (1 - \lambda_k) x_k + \lambda_k J_{c_k A} x_k + \eta_k e_k,$$

where  $x_0 \in \mathcal{H}$  is the *initial point* and  $(\forall k \in \mathbb{N}) \lambda_k \in [0, 2]$  and  $\eta_k \in \mathbb{R}_+$  are the *relaxation coefficients*,  $c_k \in \mathbb{R}_{++}$  is the *regularization coefficient*, and  $e_k \in \mathcal{H}$  is the *error term*.

*The goal of this work is to investigate the linear convergence of the generalized proximal point algorithm for solving the associated monotone inclusion problem, that is, finding a point  $\bar{x}$  in  $\text{zer } A := \{x \in \mathcal{H} : 0 \in Ax\}$ .*

Many celebrated optimization algorithms are actually specific cases of the generalized proximal point algorithm when the operator  $A$  is specified accordingly; such algorithms include the projected gradient method [18], the extragradient method [11], the forward-backward splitting algorithm [14], the Peaceman-Rachford splitting algorithm [17], the Douglas-Rachford splitting algorithm [5, 14, 19], the split inexact Uzawa method [24], and so on; in addition, the augmented Lagrangian method (i.e., the method of multipliers) [9] and the alternating direction method of multipliers [7] are instances of the generalized proximal point algorithm applied to dual problems (see, e.g., [3, 6, 10, 23] for exposition). Hence, studying the linear convergence of the generalized proximal point algorithm helps us deduce corresponding results on the linear convergence of algorithms mentioned above.

Main results in this work are summarized as follows.

---

\*Mathematics, University of British Columbia, Kelowna, B.C. V1V 1V7, Canada. E-mail: [hui.ouyang@alumni.ubc.ca](mailto:hui.ouyang@alumni.ubc.ca).

- R1:** We show  $R$ -linear convergence results on generalized proximal point algorithms in [Proposition 5.1](#) and [Theorem 5.12](#) under the assumption of metrical subregularity.
- R2:**  $Q$ -linear convergence results of generalized proximal point algorithms are presented in [Theorem 5.6](#) and [Proposition 5.8](#) under the assumption of metrical subregularity.
- R3:** In [Theorem 5.9](#) and [Proposition 5.10](#), under the assumption of the Lipschitz continuity of the inverse of the related monotone operator, we obtain  $Q$ -linear convergence results of generalized proximal point algorithms.

The rest of this work is organized as follows. We collect some basic definitions, fundamental facts, and auxiliary results in [Section 2](#). To facilitate proofs in subsequent sections, we work on the metrical subregularity of set-valued operators in [Section 3](#). In [Section 4](#), we consider the inexact version of the non-stationary Krasnosel'skiĭ-Mann iterations. In particular, we establish a  $R$ -linear convergence result on the non-stationary Krasnosel'skiĭ-Mann iterations, which will be used to deduce the corresponding result on the generalized proximal point algorithm in [Section 5](#). Our main results on the linear convergence of generalized proximal point algorithms are presented in [Section 5](#). In the last section [Section 6](#), we summarize this work and list some possible future work.

We now turn to the notation used in this work.  $\text{Id}$  stands for the *identity mapping*. Denote by  $\mathbb{R}_+ := \{\lambda \in \mathbb{R} : \lambda \geq 0\}$  and  $\mathbb{R}_{++} := \{\lambda \in \mathbb{R} : \lambda > 0\}$ . Let  $\bar{x}$  be in  $\mathcal{H}$  and let  $r \in \mathbb{R}_+$ .  $B[\bar{x}; r] := \{y \in \mathcal{H} : \|y - \bar{x}\| \leq r\}$  is the *closed ball centered at  $\bar{x}$  with radius  $r$* . Let  $C$  be a nonempty set of  $\mathcal{H}$ . Then  $(\forall x \in \mathcal{H}) d(x, C) = \inf_{y \in C} \|x - y\|$ . If  $C$  is nonempty closed and convex, then the *projector* (or *projection operator*) onto  $C$  is the operator, denoted by  $P_C$ , that maps every point in  $\mathcal{H}$  to its unique projection onto  $C$ , that is,  $(\forall x \in \mathcal{H}) \|x - P_C x\| = d(x, C)$ . Let  $\mathcal{D}$  be a nonempty subset of  $\mathcal{H}$  and let  $T : \mathcal{D} \rightarrow \mathcal{H}$ .  $\text{Fix } T := \{x \in \mathcal{D} : x = T(x)\}$  is the *set of fixed points of  $T$* . Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a set-valued operator. Then  $A$  is characterized by its *graph*  $\text{gra } A := \{(x, u) \in \mathcal{H} \times \mathcal{H} : u \in A(x)\}$ . The *inverse* of  $A$ , denoted by  $A^{-1}$ , is defined through its graph  $\text{gra } A^{-1} := \{(u, x) \in \mathcal{H} \times \mathcal{H} : (x, u) \in \text{gra } A\}$ . The *domain*, *range*, and *set of zeros* of  $A$  are defined by  $\text{dom } A := \{x \in \mathcal{H} : Ax \neq \emptyset\}$ ,  $\text{ran } A := \{y \in \mathcal{H} : \exists x \in \mathcal{H} \text{ s.t. } y \in Ax\}$ , and  $\text{zer } A := \{x \in \mathcal{H} : 0 \in Ax\}$ , respectively. Let  $(y_k)_{k \in \mathbb{N}}$  be a sequence in  $\mathcal{H}$  and let  $\bar{y}$  be in  $\mathcal{H}$ .  $\Omega((y_k)_{k \in \mathbb{N}})$  stands for the *set of all weak sequential clusters of the sequence  $(y_k)_{k \in \mathbb{N}}$* . If  $(y_k)_{k \in \mathbb{N}}$  converges (strongly) to  $\bar{y}$ , then we denote by  $y_k \rightarrow \bar{y}$ .  $(y_k)_{k \in \mathbb{N}}$  converges weakly to  $\bar{y}$  if, for every  $u \in \mathcal{H}$ ,  $\langle y_k, u \rangle \rightarrow \langle \bar{y}, u \rangle$ ; in symbols,  $y_k \rightharpoonup \bar{y}$ . Suppose that  $(y_k)_{k \in \mathbb{N}}$  converges to  $\bar{y}$ . Then  $(y_k)_{k \in \mathbb{N}}$  is  $R$ -linearly convergent (or converges  $R$ -linearly) to  $\bar{y}$  if  $\limsup_{k \rightarrow \infty} (\|y_k - \bar{y}\|)^{\frac{1}{k}} < 1$ ; when  $(\forall k \in \mathbb{N}) y_k \neq \bar{y}$ , we say  $(y_k)_{k \in \mathbb{N}}$  is  $Q$ -linearly convergent (or converges  $Q$ -linearly) to  $\bar{y}$  if  $\limsup_{k \rightarrow \infty} \frac{\|y_{k+1} - \bar{y}\|}{\|y_k - \bar{y}\|} < 1$ . For other notation not explicitly defined here, we refer the reader to [1].

## 2 Preliminaries

The definitions, facts, and lemmas gathered in this section are fundamental to our analysis in the subsequent sections.

### 2.1 Nonexpansive operators

All algorithms considered in this work are based on nonexpansive operators.

**Definition 2.1.** [1, Definition 4.1] Let  $D$  be a nonempty subset of  $\mathcal{H}$  and let  $T : D \rightarrow \mathcal{H}$ . Then  $T$  is

- (i) *nonexpansive* if it is Lipschitz continuous with constant 1, i.e.,  $(\forall x \in D) (\forall y \in D) \|Tx - Ty\| \leq \|x - y\|$ ;
- (ii) *firmly nonexpansive* if  $(\forall x \in D) (\forall y \in D) \|Tx - Ty\|^2 + \|(\text{Id} - T)x - (\text{Id} - T)y\|^2 \leq \|x - y\|^2$ .

Although [1, Definition 4.33] considers only the case  $\alpha \in ]0, 1[$ , we extend the definition to  $\alpha \in ]0, 1]$ . Clearly, by Definition 2.1(i) and Definition 2.2,  $T$  is 1-averaged if and only if  $T$  is nonexpansive. This extension will facilitate our future statements. It is clear that both firmly nonexpansive and averaged operators must be nonexpansive.

**Definition 2.2.** [1, Definition 4.33] Let  $D$  be a nonempty subset of  $\mathcal{H}$ , let  $T : D \rightarrow \mathcal{H}$  be nonexpansive, and let  $\alpha \in ]0, 1]$ . Then  $T$  is *averaged with constant  $\alpha$* , or  *$\alpha$ -averaged*, if there exists a nonexpansive operator  $R : D \rightarrow \mathcal{H}$  such that  $T = (1 - \alpha) \text{Id} + \alpha R$ .

**Fact 2.3.** [16, Proposition 2.7(ii)] Let  $\alpha \in ]0, 1]$  and let  $T : \mathcal{H} \rightarrow \mathcal{H}$  be an  $\alpha$ -averaged operator with  $\text{Fix } T \neq \emptyset$ . Let  $x$  and  $e$  be in  $\mathcal{H}$  and let  $\lambda$  and  $\eta$  be in  $\mathbb{R}_+$ . Define

$$y_x := (1 - \lambda)x + \lambda Tx \quad \text{and} \quad z_x := (1 - \lambda)x + \lambda Tx + \eta e = y_x + \eta e.$$

Then for every  $\bar{x} \in \text{Fix } T$ ,

$$\|y_x - \bar{x}\|^2 \leq \|x - \bar{x}\|^2 - \lambda \left( \frac{1}{\alpha} - \lambda \right) \|x - Tx\|^2; \quad (2.1a)$$

$$\|z_x - \bar{x}\|^2 \leq \|x - \bar{x}\|^2 - \lambda \left( \frac{1}{\alpha} - \lambda \right) \|x - Tx\|^2 + \eta \|e\| (2 \|y_x - \bar{x}\| + \eta \|e\|). \quad (2.1b)$$

**Lemma 2.4.** Let  $\alpha \in ]0, 1]$  and let  $T : \mathcal{H} \rightarrow \mathcal{H}$  be an  $\alpha$ -averaged operator with  $\text{Fix } T \neq \emptyset$ . Let  $x$  and  $e$  be in  $\mathcal{H}$ , let  $\lambda \in \mathbb{R}$ , and let  $\eta \in \mathbb{R}_+$ . Define

$$y_x := (1 - \lambda)x + \lambda Tx \quad \text{and} \quad z_x := (1 - \lambda)x + \lambda Tx + \eta e = y_x + \eta e.$$

Let  $\varepsilon$  and  $\beta$  be in  $\mathbb{R}_+$  with  $\eta\varepsilon \in [0, 1[$  and let  $\bar{x} \in \mathcal{H}$ . Suppose that  $\|e\| \leq \varepsilon \|x - z_x\|$  and  $\|y_x - \bar{x}\| \leq \beta \|x - \bar{x}\|$ . Then

$$\|z_x - \bar{x}\| \leq \frac{\beta + \eta\varepsilon}{1 - \eta\varepsilon} \|x - \bar{x}\|.$$

*Proof.* Apply the assumptions  $\|e\| \leq \varepsilon \|x - z_x\|$  and  $\|y_x - \bar{x}\| \leq \beta \|x - \bar{x}\|$  in the following second inequality to force that

$$\|z_x - \bar{x}\| \leq \|y_x - \bar{x}\| + \eta \|e\| \leq \beta \|x - \bar{x}\| + \eta\varepsilon \|x - z_x\| \leq \beta \|x - \bar{x}\| + \eta\varepsilon (\|x - \bar{x}\| + \|\bar{x} - z_x\|),$$

which implies directly that  $\|z_x - \bar{x}\| \leq \frac{\beta + \eta\varepsilon}{1 - \eta\varepsilon} \|x - \bar{x}\|$ . ■

## 2.2 Resolvent of monotone operators

**Definition 2.5.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a set-valued operator. Then we say

- (i) [1, Definition 20.1]  $A$  is *monotone* if  $(\forall (x, u) \in \text{gra } A) (\forall (y, v) \in \text{gra } A) \langle x - y, u - v \rangle \geq 0$ ;
- (ii) [1, Definition 20.20] a monotone operator  $A$  is *maximally monotone* (or *maximal monotone*) if there exists no monotone operator  $B : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  such that  $\text{gra } B$  properly contains  $\text{gra } A$ , i.e., for every  $(x, u) \in \mathcal{H} \times \mathcal{H}$ ,

$$(x, u) \in \text{gra } A \Leftrightarrow (\forall (y, v) \in \text{gra } A) \langle x - y, u - v \rangle \geq 0.$$

**Definition 2.6.** [1, Definition 23.1] Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  and let  $\gamma \in \mathbb{R}_{++}$ . The *resolvent* of  $A$  is

$$J_A = (\text{Id} + A)^{-1}.$$

**Fact 2.7** illustrates that the resolvent of a maximally monotone operator is single-valued, full domain, and firmly nonexpansive, which is essential to our study on generalized proximal point algorithms in subsequent sections.

**Fact 2.7.** [1, Proposition 23.10] *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be such that  $\text{dom } A \neq \emptyset$ , set  $D := \text{ran } A$ , and set  $T = J_A|_D$ . Then  $A$  is maximally monotone if and only if  $T$  is firmly nonexpansive and  $D = \mathcal{H}$ .*

**Lemma 2.8.** [1, Proposition 23.38] *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone and let  $\gamma \in \mathbb{R}_{++}$ . Then  $J_{\gamma A}$  is  $\frac{1}{2}$ -averaged and*

$$\text{zer}(\text{Id} - J_{\gamma A}) = \text{Fix } J_{\gamma A} = \text{zer } A. \quad (2.2)$$

*Proof.* Because  $A$  is maximally monotone, due to [1, Proposition 20.22], **Fact 2.7**, and [1, Remark 4.34(iii)], we know that  $\gamma A$  is also maximally monotone and that  $J_{\gamma A}$  is  $\frac{1}{2}$ -averaged, which, combined with [1, Proposition 23.38], yields (2.2). ■

**Fact 2.9.** [1, Proposition 23.2(ii)] *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone. Then*

$$(x \in \mathcal{H}) (\gamma \in \mathbb{R}_{++}) \quad \left( J_{\gamma A} x, \frac{1}{\gamma} (x - J_{\gamma A} x) \right) \in \text{gra } A.$$

**Fact 2.10.** [16, Lemma 2.12] *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ . Let  $x$  and  $e$  be in  $\mathcal{H}$ , let  $\lambda$  and  $\eta$  be in  $\mathbb{R}_+$ , and let  $\gamma \in \mathbb{R}_{++}$ . Define*

$$y_x := (1 - \lambda)x + \lambda J_{\gamma A} x \quad \text{and} \quad z_x := (1 - \lambda)x + \lambda J_{\gamma A} x + \eta e.$$

*Then  $(\forall \bar{x} \in \text{zer } A) \|y_x - \bar{x}\|^2 \leq \|x - \bar{x}\|^2 - \lambda(2 - \lambda) \|x - J_{\gamma A} x\|^2$ .*

**Fact 2.11.** [16, Lemma 2.15] *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$  and let  $\gamma \in \mathbb{R}_{++}$ . Then  $(\forall x \in \mathcal{H})(\forall z \in \text{zer } A) \|J_{\gamma A} x - z\|^2 + \|(\text{Id} - J_{\gamma A}) x\|^2 \leq \|x - z\|^2$ .*

### 2.3 Miscellany

Results presented in this subsection will facilitate some proofs later.

The identity shown in **Fact 2.12** below is essentially given on [8, Page 4] to illustrate the linear convergence of a special instance of the exact version of the generalized proximal point algorithm.

**Fact 2.12.** [8, Page 4] *Let  $t$  be in  $\mathbb{R} \setminus \{-1\}$  and let  $\lambda$  be in  $\mathbb{R}$ . Then*

$$\left(1 - \frac{\lambda}{t+1}\right)^2 - \left(1 - \lambda(2 - \lambda) \frac{1}{1+t^2}\right) = \frac{2t\lambda}{(1+t^2)(t+1)^2} (1 - \lambda - t^2).$$

The following result will be used to compare convergence rates of generalized proximal point algorithms later.

**Corollary 2.13.** *Let  $t$  be in  $\mathbb{R}_+$ . Then  $(1 - \frac{1}{t+1})^2 \leq (1 - \frac{1}{1+t^2})$ .*

*Proof.* Apply **Fact 2.12** with  $\lambda = 1$  to deduce that

$$\left(1 - \frac{1}{t+1}\right)^2 - \left(1 - \frac{1}{1+t^2}\right) = \frac{2t}{(1+t^2)(t+1)^2} (1 - 1 - t^2) = -t^2 \frac{2t}{(1+t^2)(t+1)^2} \leq 0,$$

which leads to the required inequality. ■

**Lemma 2.14** will play a critical role in the proof of **Theorem 3.10(i)** later.

**Lemma 2.14.** Let  $u$  and  $v$  be in  $\mathcal{H}$  and let  $t \in \mathbb{R}_{++}$  such that

$$\|v\| \leq t \|u - v\|. \quad (2.3)$$

Then the following hold.

$$(i) \quad (\forall \lambda \in [0, 1]) \quad \|(1 - \lambda)u + \lambda v\|^2 \leq \left(1 - \frac{\lambda}{t+1}\right)^2 \|u\|^2 + \lambda(t^2 + \lambda - 1) \left\| \frac{\sqrt{t}}{1+t}u - \frac{1}{\sqrt{t}}v \right\|^2.$$

Moreover, the equality holds if and only if  $\|v\| = t \|u - v\|$ .

(ii) Suppose that  $0 \leq \lambda \leq 1 - t^2$ . Then

$$\|(1 - \lambda)u + \lambda v\|^2 \leq \left(1 - \frac{\lambda}{t+1}\right)^2 \|u\|^2 \leq \|u\|^2.$$

*Proof.* (i): Let  $\lambda \in [0, 1]$ . Set  $\zeta := \frac{\lambda(1-\lambda)}{t} + \frac{\lambda^2}{1+t}$ . Since  $\lambda \in [0, 1]$  and  $t \in \mathbb{R}_{++}$ , we know that  $\zeta \geq 0$ . This combined with (2.3) guarantees that

$$\|(1 - \lambda)u + \lambda v\|^2 \quad (2.4a)$$

$$\leq \|(1 - \lambda)u + \lambda v\|^2 + \zeta (t^2 \|u - v\|^2 - \|v\|^2) \quad (2.4b)$$

$$= (1 - \lambda)^2 \|u\|^2 + \lambda^2 \|v\|^2 + 2\lambda(1 - \lambda) \langle u, v \rangle + \zeta t^2 (\|u\|^2 - 2 \langle u, v \rangle + \|v\|^2) - \zeta \|v\|^2 \quad (2.4c)$$

$$= \left(1 - \frac{\lambda}{t+1}\right)^2 \|u\|^2 + \left( (1 - \lambda)^2 - \left(1 - \frac{\lambda}{t+1}\right)^2 + \zeta t^2 \right) \|u\|^2 + 2(\lambda(1 - \lambda) - \zeta t^2) \langle u, v \rangle + (\lambda^2 + \zeta(t^2 - 1)) \|v\|^2. \quad (2.4d)$$

On the other hand, by some elementary algebra, it is easy to establish that

$$(1 - \lambda)^2 - \left(1 - \frac{\lambda}{t+1}\right)^2 + \zeta t^2 = \frac{t\lambda}{(1+t)^2} (t^2 + \lambda - 1); \quad (2.5a)$$

$$\lambda(1 - \lambda) - \zeta t^2 = -\frac{\lambda}{1+t} (t^2 + \lambda - 1); \quad (2.5b)$$

$$\lambda^2 + \zeta(t^2 - 1) = \frac{\lambda}{t} (t^2 + \lambda - 1). \quad (2.5c)$$

Combine (2.4) and (2.5) to ensure that

$$\begin{aligned} \|(1 - \lambda)u + \lambda v\|^2 &\leq \left(1 - \frac{\lambda}{t+1}\right)^2 \|u\|^2 + \lambda(t^2 + \lambda - 1) \left( \frac{t}{(1+t)^2} \|u\|^2 - \frac{2}{1+t} \langle u, v \rangle + \frac{1}{t} \|v\|^2 \right) \\ &= \left(1 - \frac{\lambda}{t+1}\right)^2 \|u\|^2 + \lambda(t^2 + \lambda - 1) \left\| \frac{\sqrt{t}}{1+t}u - \frac{1}{\sqrt{t}}v \right\|^2. \end{aligned}$$

In addition, based on our proof above, it is clear that the inequality above turns to an equality if and only if  $\|v\| = t \|u - v\|$ .

(ii): Inasmuch as  $\lambda \geq 0$  and  $t > 0$ , we have that

$$\lambda \leq 1 - t^2 \Leftrightarrow t^2 + \lambda - 1 \leq 0; \quad (2.6a)$$

$$\left(1 - \frac{\lambda}{t+1}\right)^2 \leq 1 \Leftrightarrow 0 \leq \frac{\lambda}{t+1} \leq 2 \Leftrightarrow \lambda \leq 2(t+1). \quad (2.6b)$$

Notice that  $0 \leq \lambda \leq 1 - t^2$  and that  $\lambda \leq 1 - t^2 \leq 1 \leq 2(t+1)$ . Hence, as a consequence of (i), the first and second inequalities in (ii) follow directly from (2.6a) and (2.6b), respectively.  $\blacksquare$

**Remark 2.15.** Lemma 2.14 is inspired by the second part of the proof of [8, Theorem 3.1] which works on the convergence rate of a special instance of the exact version of the generalized proximal point algorithm in  $\mathbb{R}^n$ . We assume there is a tiny typo,  $\frac{t_k}{t_k^2+1}$  should be  $\frac{t_k}{(t_k+1)^2}$ , in the equation (3.10) of the proof of [8, Theorem 3.1]. Hence, the part after (3.10) in the proof of [8, Theorem 3.1] could have been simplified.

**Lemma 2.16.** Let  $(\varepsilon_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$  and let  $(\rho_k)_{k \in \mathbb{N}}$  be in  $[0, 1]$ . Define

$$(\forall k \in \mathbb{N}) \quad \chi_k := \prod_{i=0}^k \rho_i \text{ and } \zeta_k := \sum_{i=0}^k \left( \prod_{j=i+1}^k \rho_j \right) \varepsilon_i.$$

Suppose that  $\sum_{k \in \mathbb{N}} \varepsilon_k < \infty$  and that  $\limsup_{k \rightarrow \infty} (\chi_k)^{\frac{1}{k}} < 1$ . Then the following hold.

- (i)  $\sum_{k \in \mathbb{N}} \chi_k < \infty$  and  $\lim_{k \rightarrow \infty} \chi_k = 0$ .
- (ii) Suppose that  $(\forall k \in \mathbb{N}) \rho_{k+1} \leq \rho_k$ . Then  $\sum_{k \in \mathbb{N}} \zeta_k < \infty$ .

*Proof.* (i): Because  $\chi := \limsup_{k \rightarrow \infty} (\chi_k)^{\frac{1}{k}} < 1$ , there exists  $\varepsilon > 0$  and  $K \in \mathbb{N}$  such that  $\chi + \varepsilon < 1$  and

$$(\forall k \geq K) \quad (\chi_k)^{\frac{1}{k}} < \chi + \varepsilon, \quad \text{that is, } \chi_k < (\chi + \varepsilon)^k. \quad (2.7)$$

Hence,

$$(\forall k \geq K) \quad \sum_{i=0}^k \chi_i = \sum_{i=0}^K \chi_i + \sum_{i=K+1}^k \chi_i \stackrel{(2.7)}{\leq} \sum_{i=0}^K \chi_i + \sum_{i \in \mathbb{N}} (\chi + \varepsilon)^i = \sum_{i=0}^K \chi_i + \frac{1}{1 - (\chi + \varepsilon)} < \infty,<sup>1</sup>$$

which yields that  $\sum_{k \in \mathbb{N}} \chi_k < \infty$  and  $\lim_{k \rightarrow \infty} \chi_k = 0$ .

(ii): Suppose that there exists  $\bar{k} \in \mathbb{N}$  such that  $\rho_{\bar{k}} = 0$ . Then, by the assumption  $(\forall k \in \mathbb{N}) \rho_{k+1} \leq \rho_k$  and  $\rho_k \geq 0$ , we know that  $(\forall k \geq \bar{k}) \rho_k = 0$ . So  $(\forall k \geq \bar{k}) (\forall i \in \{0, 1, \dots, k\}) \prod_{j=i+1}^k \rho_j = 0$ , which implies that  $(\forall k \geq \bar{k}) \zeta_k = 0$ . Hence, in this case, it is trivial that  $\sum_{k \in \mathbb{N}} \zeta_k < \infty$ .

Suppose that  $(\forall k \in \mathbb{N}) \rho_k > 0$ . Notice that

$$(\forall k \in \mathbb{N}) \quad \zeta_k = \sum_{i=0}^k \left( \prod_{j=i+1}^k \rho_j \right) \varepsilon_i = \sum_{i=0}^k \left( \left( \prod_{j=0}^{k-i} \rho_j \right) \varepsilon_i \cdot \frac{\left( \prod_{j=i+1}^k \rho_j \right)}{\left( \prod_{j=0}^{k-i} \rho_j \right)} \right)$$

and that for every  $k \in \mathbb{N}$  and  $i \in \{0, 1, \dots, k\}$

$$\begin{aligned} \frac{\prod_{j=i+1}^k \rho_j}{\prod_{j=0}^{k-i} \rho_j} &= \begin{cases} \frac{\prod_{j=i+1}^k \rho_j}{\prod_{j=0}^{k-i} \rho_j} & \text{if } k-i \leq i+1; \\ \frac{\prod_{j=k-i+1}^k \rho_j}{\prod_{j=0}^i \rho_j} & \text{if } k-i > i+1, \end{cases} \\ &= \begin{cases} \frac{1}{\rho_0} \prod_{j=1}^{k-i} \frac{\rho_{j+i}}{\rho_j} & \text{if } k-i \leq i+1; \\ \frac{1}{\rho_0} \prod_{j=1}^i \frac{\rho_{j+k-i}}{\rho_j} & \text{if } k-i > i+1, \end{cases} \\ &\leq \frac{1}{\rho_0}, \end{aligned}$$

where we invoke the assumption  $(\forall k \in \mathbb{N}) \rho_{k+1} \leq \rho_k$  in the last inequality.

<sup>1</sup>In the whole work, we use the empty sum convention. Hence, if  $k = K$  in this inequality, then  $\sum_{i=K+1}^k \chi_i = 0$ .

On the other hand, due to [21, Page 80], the Cauchy product  $\sum_{k \in \mathbb{N}} \sum_{i=0}^k \chi_{k-i} \varepsilon_i$  of the two absolutely convergent series  $\sum_{k \in \mathbb{N}} \varepsilon_k$  and  $\sum_{k \in \mathbb{N}} \chi_k$  is absolutely convergent, that is,

$$\sum_{k \in \mathbb{N}} \sum_{i=0}^k \chi_{k-i} \varepsilon_i < \infty. \quad (2.8)$$

Taking all results obtained above into account, we derive that

$$\sum_{k \in \mathbb{N}} \xi_k \leq \frac{1}{\rho_0} \sum_{k \in \mathbb{N}} \sum_{i=0}^k \left( \prod_{j=0}^{k-i} \rho_j \right) \varepsilon_i = \frac{1}{\rho_0} \sum_{k \in \mathbb{N}} \sum_{i=0}^k \chi_{k-i} \varepsilon_i \stackrel{(2.8)}{<} \infty,$$

which reaches the required inequality in (ii). ■

### 3 Metrical Subregularity

Metrical subregularity is a critical assumption for the linear convergence of algorithms studied in this work. In this section, we exhibit results related to metrical subregularity.

**Definition 3.1.** [4, Pages 183 and 184] Let  $F : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a set-valued operator.  $F$  is called *metrically subregular at  $\bar{x}$  for  $\bar{y}$*  if  $(\bar{x}, \bar{y}) \in \text{gra } F$  and there exist  $\kappa \in \mathbb{R}_+$  and a neighborhood  $U$  of  $\bar{x}$  such that

$$(\forall x \in U) \quad d(x, F^{-1}(\bar{y})) \leq \kappa d(\bar{y}, F(x)).$$

The constant  $\kappa$  is called *constant of metric subregularity*. The infimum of all  $\kappa$  for which the inequality above holds is the *modulus of metric subregularity*, denoted by  $\text{subreg}(F; \bar{x} | \bar{y})$ . The absence of metric subregularity is signaled by  $\text{subreg}(F; \bar{x} | \bar{y}) = \infty$ .

**Fact 3.2.** [16, Lemma 2.19] Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ , let  $\bar{x} \in \text{zer } A$ , and let  $\gamma \in \mathbb{R}_{++}$ . Then  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$  if and only if  $\text{Id} - J_{\gamma A}$  is metrically subregular at  $\bar{x}$  for  $0 = (\text{Id} - J_{\gamma A})\bar{x}$ . In particular, if  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d(x, A^{-1}0) \leq \kappa d(0, Ax),$$

then  $\text{Id} - J_{\gamma A}$  is metrically subregular at  $\bar{x}$  for  $0 = (\text{Id} - J_{\gamma A})\bar{x}$ ; more specifically,

$$(\forall x \in B[\bar{x}; \delta]) \quad d(x, (\text{Id} - J_{\gamma A})^{-1}0) \leq \left(1 + \frac{\kappa}{\gamma}\right) d(0, (\text{Id} - J_{\gamma A})x).$$

Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\bar{x} \in \text{zer } A$ , let  $\gamma$  and  $\delta$  be in  $\mathbb{R}_{++}$ , and let  $x \in \mathcal{H}$ . If  $x \in B[\bar{x}; \delta]$ , then due to **Fact 2.11**,  $J_{\gamma A} x \in B[\bar{x}; \delta]$ . Hence, applying **Lemma 3.3**, we easily deduce [12, Theorem 3.1] and [23, Lemma 5.3]. In fact, the proofs of **Lemma 3.3**, [12, Theorem 3.1], and [23, Lemma 5.3] are similar.

**Lemma 3.3.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$  and let  $\gamma \in \mathbb{R}_{++}$ . Assume that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d(x, A^{-1}0) \leq \kappa d(0, Ax). \quad (3.1)$$

Then for every  $x \in \mathcal{H}$  with  $J_{\gamma A} x \in B[\bar{x}; \delta]$ ,

$$d(J_{\gamma A} x, A^{-1}0) \leq \frac{1}{\sqrt{1 + \frac{\gamma^2}{\kappa^2}}} d(x, A^{-1}0). \quad (3.2)$$

*Proof.* As a consequence of [Fact 2.9](#),

$$(\forall x \in \mathcal{H}) \quad \frac{1}{\gamma} (x - J_{\gamma A} x) \in A (J_{\gamma A} x). \quad (3.3)$$

Due to [Fact 2.11](#),

$$(\forall x \in \mathcal{H})(\forall z \in \text{zer } A) \quad \|J_{\gamma A} x - z\|^2 + \|(\text{Id} - J_{\gamma A}) x\|^2 \leq \|x - z\|^2. \quad (3.4)$$

By virtue of the maximal monotonicity of  $A$  and via [\[1, Proposition 23.39\]](#), we know that  $\text{zer } A$  is closed and convex. So  $P_{\text{zer } A} x \in \text{zer } A$  is well-defined. Substitute  $z = P_{\text{zer } A} x$  in [\(3.4\)](#) to establish that

$$(\forall x \in \mathcal{H}) \quad \|x - J_{\gamma A} x\|^2 \leq \|x - P_{\text{zer } A} x\|^2 - \|J_{\gamma A} x - P_{\text{zer } A} x\|^2. \quad (3.5)$$

Let  $x \in \mathcal{H}$  with  $J_{\gamma A} x \in B[\bar{x}; \delta]$ . Applying [\(3.1\)](#) with  $x = J_{\gamma A} x$  in the first inequality below and employing  $\|J_{\gamma A} x - P_{\text{zer } A} (J_{\gamma A} x)\| \leq \|J_{\gamma A} x - P_{\text{zer } A} x\|$  in the fourth inequality, we deduce that

$$\begin{aligned} d^2 (J_{\gamma A} x, A^{-1}0) &\leq \kappa^2 d^2 (0, A (J_{\gamma A} x)) \\ &\stackrel{(3.3)}{\leq} \frac{\kappa^2}{\gamma^2} \|x - J_{\gamma A} x\|^2 \\ &\stackrel{(3.5)}{\leq} \frac{\kappa^2}{\gamma^2} (\|x - P_{\text{zer } A} x\|^2 - \|J_{\gamma A} x - P_{\text{zer } A} x\|^2) \\ &\leq \frac{\kappa^2}{\gamma^2} \|x - P_{\text{zer } A} x\|^2 - \frac{\kappa^2}{\gamma^2} \|J_{\gamma A} x - P_{\text{zer } A} (J_{\gamma A} x)\|^2 \\ &= \frac{\kappa^2}{\gamma^2} d^2 (x, A^{-1}0) - \frac{\kappa^2}{\gamma^2} d^2 (J_{\gamma A} x, A^{-1}0), \end{aligned}$$

which yields [\(3.2\)](#) easily, since it is clear that  $\frac{\frac{\kappa^2}{\gamma^2}}{1 + \frac{\kappa^2}{\gamma^2}} = \frac{1}{1 + \frac{\gamma^2}{\kappa^2}}$ . ■

**Theorem 3.4.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ . Let  $x \in \mathcal{H}$ , let  $e \in \mathcal{H}$  and  $\eta \in \mathbb{R}_+$ , let  $(\gamma, \varepsilon) \in \mathbb{R}_{++}^2$ , and let  $\lambda \in ]0, 2[$ . Define

$$y_x := (1 - \lambda) x + \lambda J_{\gamma A} x \quad \text{and} \quad z_x := (1 - \lambda) x + \lambda J_{\gamma A} x + \eta e = y_x + \eta e.$$

Suppose that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d(x, A^{-1}0) \leq \kappa d(0, Ax). \quad (3.6)$$

Set  $\rho := \left(1 - \lambda(2 - \lambda) \frac{1}{(1 + \frac{\kappa}{\gamma})^2}\right)^{\frac{1}{2}}$ . Suppose that  $J_{\gamma A} x \in B[\bar{x}; \delta]$ . Then the following statements hold.

- (i)  $\rho \in ]0, 1[$ .
- (ii)  $\|y_x - P_{\text{zer } A} x\| \leq \rho \|x - P_{\text{zer } A} x\|$ .
- (iii) Suppose that  $\|e\| \leq \varepsilon \|x - z_x\|$  and  $\eta\varepsilon \in [0, 1[$ . Then

$$\|z_x - P_{\text{zer } A} x\| \leq \frac{\rho + \eta\varepsilon}{1 - \eta\varepsilon} \|x - P_{\text{zer } A} x\|.$$



*Proof.* (i): This is clear from  $\lambda \in ]0, 2[$ ,  $\lambda(2 - \lambda) = -(1 - \lambda)^2 + 1 \leq 1$ , and  $\frac{1}{(1 + \frac{\kappa}{\gamma})^2} \in ]0, 1[$ .

(ii): Applying (3.6) with  $x = J_{\gamma A} x$  in the first inequality and employing Fact 2.9 in the second inequality, we know that

$$\|J_{\gamma A} x - P_{\text{zer } A}(J_{\gamma A} x)\| = d(J_{\gamma A} x, A^{-1}0) \leq \kappa d(0, A(J_{\gamma A} x)) \leq \frac{\kappa}{\gamma} \|x - J_{\gamma A} x\|. \quad (3.7)$$

Hence,

$$\|x - P_{\text{zer } A} x\| \leq \|x - P_{\text{zer } A}(J_{\gamma A} x)\| \quad (3.8a)$$

$$\leq \|x - J_{\gamma A} x\| + \|J_{\gamma A} x - P_{\text{zer } A}(J_{\gamma A} x)\| \quad (3.8b)$$

$$\stackrel{(3.7)}{\leq} \left(1 + \frac{\kappa}{\gamma}\right) \|x - J_{\gamma A} x\|. \quad (3.8c)$$

Applying Fact 2.10 with  $\bar{x} = P_{\text{zer } A} x$  in the first inequality below, we observe that

$$\begin{aligned} \|y_x - P_{\text{zer } A} x\|^2 &\leq \|x - P_{\text{zer } A} x\|^2 - \lambda(2 - \lambda) \|x - J_{\gamma A} x\|^2 \\ &\stackrel{(3.8)}{\leq} \left(1 - \lambda(2 - \lambda) \frac{1}{\left(1 + \frac{\kappa}{\gamma}\right)^2}\right) \|x - P_{\text{zer } A} x\|^2, \end{aligned}$$

which guarantees the desired inequality in (ii).

(iii): Based on Lemma 2.8,  $J_{\gamma A}$  is  $\frac{1}{2}$ -averaged and  $\text{Fix } J_{\gamma A} = \text{zer } A \neq \emptyset$ . Employing (ii) and applying Lemma 2.4 with  $T = J_{\gamma A}$ ,  $\alpha = \frac{1}{2}$ ,  $\beta = \rho$ , and  $\bar{x} = P_{\text{zer } A} x$ , we deduce (iii). ■

**Remark 3.5.** We uphold the assumptions of Theorem 3.4. By some easy algebra, it is not difficult to get that

$$\max \left\{ \left(1 - \frac{\lambda}{\frac{\kappa}{\gamma} + 1}\right)^2, \left(1 - \lambda(2 - \lambda) \frac{1}{1 + \frac{\kappa^2}{\gamma^2}}\right) \right\} < 1 - \lambda(2 - \lambda) \frac{1}{\left(1 + \frac{\kappa}{\gamma}\right)^2}.$$

Note that if  $\text{zer } A$  is a singleton, then  $P_{\text{zer } A} x = P_{\text{zer } A}(J_{\gamma A} x)$ . Therefore, based on Theorem 3.12(i) below, if  $\text{zer } A$  is a singleton, then the coefficient  $\rho$  in Theorem 3.4(ii) can be decreased.

**Definition 3.6.** [20, Page 885] Let  $F : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a set-valued operator. We say  $F^{-1}$  is *Lipschitz continuous at 0* (with modulus  $\alpha \geq 0$ ) if there is a unique solution  $\bar{z}$  to  $0 \in F(z)$  (i.e.  $F^{-1}(0) = \{\bar{z}\}$ ), and for some  $\tau > 0$  we have

$$\|z - \bar{z}\| \leq \alpha \|w\| \quad \text{whenever } z \in F^{-1}(w) \text{ and } \|w\| \leq \tau.$$

Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be a maximally monotone operator with  $\text{zer } A \neq \emptyset$ . It was claimed in [12, Page 684] and [22, Page 5] without proof that the assumption that  $A^{-1}$  is Lipschitz continuous at 0 is stronger than that  $A$  is metrically subregular. For completeness, we provide a detailed proof for this claim below.

**Fact 3.7.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ . Suppose that  $A^{-1}$  is Lipschitz continuous at 0 with modulus  $\alpha > 0$ , i.e.,  $A^{-1}(0) = \{\bar{x}\}$  and there exists  $\tau > 0$  such that

$$\left(\forall (w, x) \in \text{gra } A^{-1} \text{ with } w \in B[0; \tau]\right) \quad \|x - \bar{x}\| \leq \alpha \|w\|. \quad (3.9)$$

Set  $\delta := \alpha\tau$ . Then  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$  with  $\text{subreg}(A; \bar{x}|\bar{y}) \leq \alpha$ ; more precisely,

$$\left(\forall x \in B[\bar{x}; \delta]\right) \quad d(x, A^{-1}0) \leq \alpha d(0, Ax). \quad (3.10)$$

*Proof.* Let  $x \in B[\bar{x}; \delta]$ . Because  $A^{-1}0 = \{\bar{x}\}$ , we know that

$$d(x, A^{-1}0) = \|x - \bar{x}\|. \quad (3.11)$$

If  $Ax = \emptyset$ , then, via the convention  $\inf \emptyset = \infty$ ,  $d(x, A^{-1}0) \leq \infty = \alpha d(0, Ax)$ .

Assume that  $Ax \neq \emptyset$ . Then, invoking the maximal monotonicity of  $A$  and employing [1, Proposition 23.39], we know that  $Ax$  is nonempty closed and convex. So, via [1, Theorem 3.16],  $y := P_{Ax} 0$  is a well-defined point in  $Ax$ . Then we have exactly the following two cases.

*Case 1:*  $y \in B[0; \tau]$ . Now  $(y, x) \in \text{gra } A^{-1}$  with  $y \in B[0; \tau]$ . Bearing (3.9) and (3.11) in mind, we observe that

$$d(x, A^{-1}0) = \|x - \bar{x}\| \leq \alpha \|y\| = \alpha \|P_{Ax} 0\| = \alpha d(0, Ax).$$

*Case 2:*  $y \notin B[0; \tau]$ . Then  $\|y\| = \|P_{Ax} 0\| > \tau$ . Hence, it is easy to see that

$$d(x, A^{-1}0) = \|x - \bar{x}\| \leq \delta = \alpha \tau \leq \alpha \|P_{Ax} 0\| = \alpha d(0, Ax).$$

Altogether, (3.10) is true in both cases and the proof is complete. ■

Naturally, one may have the following question.

**Question 3.8.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone. Suppose that  $\text{zer } A = \{\bar{x}\}$ . Are the following two statements equivalent?

- (i)  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ .
- (ii)  $A^{-1}$  is Lipschitz continuous at 0 with a positive modulus.

Based on Fact 3.7, we know that (ii)  $\Rightarrow$  (i). Therefore, (i)  $\Leftrightarrow$  (ii) if and only if (i)  $\Rightarrow$  (ii).

Example 3.9 illustrates that for all continuous and monotone function  $f : \mathbb{R} \rightarrow \mathbb{R}$  with  $f^{-1}(0) = \{\bar{x}\}$ , the metrical subregularity of  $f$  is equivalent to the Lipschitz continuity of  $f^{-1}$  at 0 with a positive modulus. In other words, we provide a specific example showing the equivalence of the two statements in Question 3.8.

**Example 3.9.** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be continuous and monotone and let  $f^{-1}(0) = \{\bar{x}\}$ . Then the following hold.

- (i)  $f$  is maximally monotone.
- (ii) Let  $\epsilon \in \mathbb{R}_{++}$ . Then there exists  $\delta \in \mathbb{R}_{++}$  such that

$$f(x) \in B[0; \delta] \Rightarrow x \in B[\bar{x}; \epsilon]. \quad (3.12)$$

- (iii)  $f$  is metrically subregular at  $\bar{x}$  for  $0 = f(\bar{x})$  if and only if  $f^{-1}$  is Lipschitz continuous at 0 with a positive modulus.

*Proof.* (i): Because  $f$  is continuous and monotone, via [1, Corollary 20.28],  $f$  is maximally monotone.

(ii): Suppose to the contrary that (3.12) is not true. Then

$$(\forall k \in \mathbb{N} \setminus \{0\}) \text{ there exists } x_k \in \mathbb{R} \text{ such that } f(x_k) \in B\left[0; \frac{1}{k}\right] \text{ and } x_k \notin B[\bar{x}; \epsilon],$$

which forces that

$$f(x_k) \rightarrow 0 \quad \text{and} \quad \Omega((x_k)_{k \in \mathbb{N}}) \cap B\left[\bar{x}; \frac{\epsilon}{2}\right] = \emptyset, \quad (3.13)$$

where  $\Omega((x_k)_{k \in \mathbb{N}})$  is the set of all sequential cluster points of  $(x_k)_{k \in \mathbb{N}}$ .

We have exactly the following cases.

*Case 1:*  $(x_k)_{k \in \mathbb{N}}$  is bounded. Then this together with (3.13) implies that there exists a subsequence  $(x_{k_i})_{i \in \mathbb{N}}$  of  $(x_k)_{k \in \mathbb{N}}$  such that

$$f(x_{k_i}) \rightarrow 0 \quad \text{and} \quad x_{k_i} \rightarrow \hat{x} \notin B\left[\bar{x}; \frac{\epsilon}{2}\right].$$

On the other hand, the continuity of  $f$  implies that

$$f(\hat{x}) = f\left(\lim_{i \rightarrow \infty} x_{k_i}\right) = \lim_{i \rightarrow \infty} f(x_{k_i}) = 0,$$

which contradicts that  $f^{-1}(0) = \{\bar{x}\}$  and  $\hat{x} \neq \bar{x}$ .

*Case 2:*  $(x_k)_{k \in \mathbb{N}}$  is not bounded. Without loss of generality, we assume that there exists a subsequence  $(x_{k_i})_{i \in \mathbb{N}}$  of  $(x_k)_{k \in \mathbb{N}}$  such that

$$x_{k_i} \rightarrow \infty. \tag{3.14}$$

Let  $\tilde{x} > \bar{x}$ . (If  $x_{k_i} \rightarrow -\infty$ , then we choose  $\tilde{x} < \bar{x}$  and the remaining proof is similar to the following proof.) Inasmuch as  $f^{-1}(0) = \{\bar{x}\}$ , we have exactly the following two subcases.

*Subcase 2.1:*  $f(\tilde{x}) > 0$ . Combine this with (3.13) and (3.14) to deduce that there exists  $N \in \mathbb{N}$  such that

$$(\forall i \geq N) \quad f(x_{k_i}) < f(\tilde{x}) \quad \text{and} \quad x_{k_i} > \tilde{x},$$

which entails that

$$\langle \tilde{x} - \bar{x}, f(\tilde{x}) - f(\bar{x}) \rangle > 0 \quad \text{and} \quad \langle x_{k_N} - \tilde{x}, f(x_{k_N}) - f(\tilde{x}) \rangle < 0.$$

This contradicts the monotonicity of  $f$ .

*Subcase 2.2:*  $f(\tilde{x}) < 0$ . Similarly, as a consequence of (3.13) and (3.14), there exists  $N \in \mathbb{N}$  such that

$$(\forall i \geq N) \quad f(x_{k_i}) > f(\tilde{x}) \quad \text{and} \quad x_{k_i} > \tilde{x}.$$

This necessitates that

$$\langle \tilde{x} - \bar{x}, f(\tilde{x}) - f(\bar{x}) \rangle < 0 \quad \text{and} \quad \langle x_{k_N} - \tilde{x}, f(x_{k_N}) - f(\tilde{x}) \rangle > 0,$$

which contradicts the monotonicity of  $f$  as well.

Altogether, (ii) holds in all cases.

(iii): Suppose that  $f$  is metrically subregular at  $\bar{x}$  for  $0 = f(\bar{x})$ . Then

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d\left(x, f^{-1}(0)\right) \leq \kappa d(0, f(x)), \quad \text{that is, } |x - \bar{x}| \leq \kappa |f(x)|. \tag{3.15}$$

As a result of Fact 3.7, it remains to prove that  $f^{-1}$  is Lipschitz continuous at 0 with a positive modulus.

Replace  $\epsilon = \delta$  in (ii) above to see that there exists  $\delta' \in \mathbb{R}_{++}$  such that

$$f(x) \in B[0; \delta'] \Rightarrow x \in B[\bar{x}; \delta]. \tag{3.16}$$

Now, invoke (3.15) and (3.16) to obtain that

$$\left(\forall (f(x), x) \in \text{gra } f^{-1} \text{ with } f(x) \in B[0; \delta']\right) \quad |x - \bar{x}| \leq \kappa |f(x)|,$$

which, via Definition 3.6, ensures that  $f^{-1}$  is Lipschitz continuous at 0 with modulus  $\kappa > 0$ .

Altogether, the proof is complete. ■

[Theorem 3.10](#) together with [Theorem 3.4](#) will play a critical role to prove the linear convergence of generalized proximal point algorithms later.

**Theorem 3.10.** *Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ , let  $x \in \mathcal{H}$ , let  $\gamma \in \mathbb{R}_{++}$ , and let  $\lambda \in [0, 2]$ . Define*

$$y_x := (1 - \lambda)x + \lambda J_{\gamma A} x. \quad (3.17)$$

Let  $z \in \text{zer } A$  and let  $t \in \mathbb{R}_{++}$ . Suppose that

$$\|J_{\gamma A} x - z\| \leq t \|x - J_{\gamma A} x\|. \quad (3.18)$$

Then the following statements hold.

(i) Suppose that  $\lambda \in [0, 1]$ . Then

$$\|y_x - z\|^2 \leq \begin{cases} \left(1 - \frac{\lambda}{t+1}\right)^2 \|x - z\|^2 & \text{if } t^2 + \lambda - 1 \leq 0; \\ \left(1 - \lambda(2 - \lambda) \frac{1}{1+t^2}\right) \|x - z\|^2 & \text{if } t^2 + \lambda - 1 > 0. \end{cases}$$

(ii) Suppose that  $\lambda \in [1, 2]$ . Then

$$\|y_x - z\|^2 \leq \left(1 - \lambda(2 - \lambda) \frac{1}{1+t^2}\right) \|x - z\|^2.$$

(iii) Define

$$\rho := \begin{cases} \left(1 - \frac{\lambda}{t+1}\right)^2 & \text{if } 0 \leq \lambda \leq 1 - t^2; \\ 1 - \lambda(2 - \lambda) \frac{1}{1+t^2} & \text{if } 1 - t^2 < \lambda \leq 2. \end{cases} \quad (3.20)$$

Suppose that  $\lambda \in ]0, 2[$ . Then

$$\rho = \max \left\{ \left(1 - \frac{\lambda}{t+1}\right)^2, 1 - \lambda(2 - \lambda) \frac{1}{1+t^2} \right\} \in ]0, 1[. \quad (3.21)$$

Moreover,

$$\|y_x - z\|^2 \leq \rho \|x - z\|^2.$$

*Proof.* Invoke [Fact 2.11](#) in the following second inequality to entail that

$$\left(1 + \frac{1}{t^2}\right) \|J_{\gamma A} x - z\|^2 \stackrel{(3.18)}{\leq} \|J_{\gamma A} x - z\|^2 + \|x - J_{\gamma A} x\|^2 \leq \|x - z\|^2,$$

which necessitates that

$$\|J_{\gamma A} x - z\|^2 \leq \frac{1}{1 + \frac{1}{t^2}} \|x - z\|^2. \quad (3.22)$$

(i): If  $\lambda = 0$ , then  $\|y_x - z\|^2 = \|x - z\|^2 = \left(1 - \frac{\lambda}{t+1}\right)^2 \|x - z\|^2 = \left(1 - \lambda(2 - \lambda) \frac{1}{1+t^2}\right) \|x - z\|^2$ . Hence, the result in (i) is trivial.

Suppose that  $\lambda \in ]0, 1]$ . Then

$$\left( \lambda - \lambda(1 - \lambda) \frac{1}{t^2} \right) \leq 0 \Leftrightarrow t^2 + \lambda - 1 \leq 0. \quad (3.23)$$

It is clear that

$$(1 - \lambda) + \left( \lambda - \lambda(1 - \lambda) \frac{1}{t^2} \right) \frac{1}{1 + \frac{1}{t^2}} = 1 + \lambda \left( \frac{1}{1 + \frac{1}{t^2}} - 1 \right) - \lambda(1 - \lambda) \frac{1}{t^2} \frac{1}{1 + \frac{1}{t^2}} \quad (3.24a)$$

$$= 1 - \lambda \frac{1}{1 + t^2} - \lambda(1 - \lambda) \frac{1}{1 + t^2} \quad (3.24b)$$

$$= 1 - \lambda(2 - \lambda) \frac{1}{1 + t^2}. \quad (3.24c)$$

Employing [1, Corollary 2.15] in the following second equality and invoking both (3.22) and (3.23) in the second inequality, we observe that

$$\begin{aligned} \|y_x - z\|^2 &\stackrel{(3.17)}{=} \|(1 - \lambda)(x - z) + \lambda(J_{\gamma A}x - z)\|^2 \\ &= (1 - \lambda)\|x - z\|^2 + \lambda\|J_{\gamma A}x - z\|^2 - \lambda(1 - \lambda)\|x - J_{\gamma A}x\|^2 \\ &\stackrel{(3.18)}{\leq} (1 - \lambda)\|x - z\|^2 + \lambda\|J_{\gamma A}x - z\|^2 - \lambda(1 - \lambda)\frac{1}{t^2}\|J_{\gamma A}x - z\|^2 \\ &= (1 - \lambda)\|x - z\|^2 + \left( \lambda - \lambda(1 - \lambda) \frac{1}{t^2} \right) \|J_{\gamma A}x - z\|^2 \\ &\leq \begin{cases} (1 - \lambda)\|x - z\|^2 & \text{if } t^2 + \lambda - 1 \leq 0; \\ \left( (1 - \lambda) + (\lambda - \lambda(1 - \lambda) \frac{1}{t^2}) \frac{1}{1 + \frac{1}{t^2}} \right) \|x - z\|^2 & \text{if } t^2 + \lambda - 1 > 0. \end{cases} \end{aligned}$$

This combined with (3.24) guarantees that

$$\|y_x - z\|^2 \leq \begin{cases} (1 - \lambda)\|x - z\|^2 & \text{if } t^2 + \lambda - 1 \leq 0; \\ \left( (1 - \lambda(2 - \lambda) \frac{1}{1 + t^2}) \right) \|x - z\|^2 & \text{if } t^2 + \lambda - 1 > 0. \end{cases} \quad (3.25)$$

On the other hand, if  $t^2 + \lambda - 1 \leq 0$ , then utilizing (3.18) and applying Lemma 2.14(ii) with  $u = x - z$  and  $v = J_{\gamma A}x - z$ , we know that

$$\|y_x - z\|^2 \leq \left( 1 - \frac{\lambda}{t + 1} \right)^2 \|x - z\|^2. \quad (3.26)$$

Notice that, by some easy algebra,  $(1 - \frac{\lambda}{t+1})^2 \leq 1 - \lambda \Leftrightarrow t^2 + \lambda - 1 \leq 0$ . Hence, combining (3.25) and (3.26), we deduce (i).

(ii): Because  $\lambda \in [1, 2]$ , we have that

$$\lambda(1 - \lambda) \leq 0. \quad (3.27)$$

Due to Fact 2.9,  $(J_{\gamma A}x, \frac{1}{\gamma}(x - J_{\gamma A}x)) \in \text{gra } A$ . This combined with  $(z, 0) \in \text{gra } A$  and the monotonicity of  $A$  entails that

$$\langle J_{\gamma A}x - z, x - J_{\gamma A}x \rangle \geq 0. \quad (3.28)$$

Invoke (3.27) and (3.28) in the following first inequality to derive that

$$\begin{aligned}
& \|y_x - z\|^2 \\
(3.17) \quad &= \|(1 - \lambda)(x - z) + \lambda(J_{\gamma A}x - z)\|^2 \\
&= (1 - \lambda)^2 \|x - z\|^2 + \lambda^2 \|J_{\gamma A}x - z\|^2 + 2\lambda(1 - \lambda) \langle x - z, J_{\gamma A}x - z \rangle \\
&= (1 - \lambda)^2 \|x - z\|^2 + \lambda^2 \|J_{\gamma A}x - z\|^2 + 2\lambda(1 - \lambda) \|J_{\gamma A}x - z\|^2 + 2\lambda(1 - \lambda) \langle x - J_{\gamma A}x, J_{\gamma A}x - z \rangle \\
&\leq (1 - \lambda)^2 \|x - z\|^2 + \lambda(2 - \lambda) \|J_{\gamma A}x - z\|^2 \\
(3.22) \quad &\leq (1 - \lambda)^2 \|x - z\|^2 + \lambda(2 - \lambda) \frac{1}{1 + t^2} \|x - z\|^2 \\
&= \left(1 - \lambda(2 - \lambda) \frac{1}{1 + t^2}\right) \|x - z\|^2,
\end{aligned}$$

where the last equality follows from

$$(1 - \lambda)^2 + \lambda(2 - \lambda) \frac{1}{1 + t^2} = 1 + \lambda(2 - \lambda) \left(-1 + \frac{1}{1 + t^2}\right) = 1 - \lambda(2 - \lambda) \frac{1}{1 + t^2}.$$

(iii): Inasmuch as  $\lambda \in [0, 2]$  and  $t \in \mathbb{R}_{++}$ , it is easy to get that

$$\begin{aligned}
\left(1 - \frac{\lambda}{t + 1}\right)^2 &\in ]0, 1[ \Leftrightarrow \lambda < 2(t + 1); \\
1 - \lambda(2 - \lambda) \frac{1}{1 + t^2} &\in ]0, 1[ \Leftrightarrow \lambda(2 - \lambda) > 0.
\end{aligned}$$

Hence,  $\lambda \in ]0, 2[$  and  $t \in \mathbb{R}_{++}$  lead to

$$\max \left\{ \left(1 - \frac{\lambda}{t + 1}\right)^2, 1 - \lambda(2 - \lambda) \frac{1}{1 + t^2} \right\} \in ]0, 1[.$$

Combine this with Fact 2.12 and (3.20) to yield (3.21).

Furthermore, the last assertion in (iii) is clear from (i) and (ii) above. ■

The inequality (3.29) presented in Remark 3.11 will be used to compare convergence rates of generalized proximal point algorithms later.

**Remark 3.11.** Let  $\lambda \in \mathbb{R}_+$  and let  $t \in \mathbb{R}_{++}$  such that  $\lambda \leq 1 - t^2$ . Then

$$\begin{aligned}
1 - \lambda \frac{1}{1 + t^2} - \left(1 - \frac{\lambda}{t + 1}\right)^2 &= \frac{2\lambda}{t + 1} - \lambda \frac{1}{1 + t^2} - \frac{\lambda^2}{(t + 1)^2} \\
&= \frac{\lambda}{(1 + t^2)(t + 1)^2} (2(t + 1)(1 + t^2) - (t + 1)^2 - \lambda(1 + t^2)) \\
&\geq \frac{\lambda}{(1 + t^2)(t + 1)^2} (2(t + 1)(1 + t^2) - (t + 1)^2 - (1 - t^2)(1 + t^2)) \\
&= \frac{\lambda}{(1 + t^2)(t + 1)^2} (2t^3 + t^2 + t^4) = \frac{\lambda t^2}{1 + t^2} \geq 0,
\end{aligned}$$

where we use the assumption  $\lambda \leq 1 - t^2$  in the first inequality above.

Hence, based on (3.21) in Theorem 3.10(iii), we know that

$$\max \left\{ \left(1 - \frac{\lambda}{t + 1}\right)^2, 1 - \lambda(2 - \lambda) \frac{1}{1 + t^2} \right\} \leq \max \left\{ 1 - \lambda \frac{1}{1 + t^2}, 1 - \lambda(2 - \lambda) \frac{1}{1 + t^2} \right\}. \quad (3.29)$$

**Theorem 3.12.** Let  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  be maximally monotone with  $\text{zer } A \neq \emptyset$ . Let  $x$  and  $e$  be in  $\mathcal{H}$ , let  $\bar{x} \in \text{zer } A$ , let  $\eta$  and  $\varepsilon$  be in  $\mathbb{R}_+$ , let  $\gamma \in \mathbb{R}_{++}$ , and let  $\lambda \in ]0, 2[$ . Define

$$y_x := (1 - \lambda)x + \lambda J_{\gamma A} x \quad \text{and} \quad z_x := (1 - \lambda)x + \lambda J_{\gamma A} x + \eta e. \quad (3.30)$$

Then the following statements hold.

(i) Suppose that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d(x, A^{-1}0) \leq \kappa d(0, Ax). \quad (3.31)$$

Set  $\rho := \max \left\{ \left(1 - \frac{\lambda}{\frac{\kappa}{\gamma} + 1}\right)^2, \left(1 - \lambda(2 - \lambda) \frac{1}{1 + \frac{\kappa^2}{\gamma^2}}\right) \right\}^{\frac{1}{2}}$ . Suppose that  $J_{\gamma A} x \in B[\bar{x}; \delta]$ . Then the following hold.

- (a)  $\rho \in ]0, 1[$ .
- (b)  $\|y_x - P_{\text{zer } A}(J_{\gamma A} x)\| \leq \rho \|x - P_{\text{zer } A}(J_{\gamma A} x)\|$ .
- (c) If  $\text{zer } A = \{\bar{x}\}$ , then  $\|y_x - \bar{x}\| \leq \rho \|x - \bar{x}\|$ .
- (d) Suppose that  $\|e\| \leq \varepsilon \|x - z_x\|$  and that  $\eta\varepsilon \in [0, 1[$ . Then

$$\|z_x - P_{\text{zer } A}(J_{\gamma A} x)\| \leq \frac{\rho + \eta\varepsilon}{1 - \eta\varepsilon} \|x - P_{\text{zer } A}(J_{\gamma A} x)\|.$$

In addition, if  $\text{zer } A = \{\bar{x}\}$ , then

$$\|z_x - \bar{x}\| \leq \frac{\rho + \eta\varepsilon}{1 - \eta\varepsilon} \|x - \bar{x}\|.$$

(ii) Suppose that  $A^{-1}$  is Lipschitz continuous at 0 with modulus  $\alpha > 0$ , i.e.,  $A^{-1}(0) = \{\bar{x}\}$  and there exists  $\tau > 0$  such that

$$(\forall (w, x) \in \text{gra } A^{-1} \text{ with } w \in B[0; \tau]) \quad \|x - \bar{x}\| \leq \alpha \|w\|. \quad (3.32)$$

Set  $\rho := \max \left\{ \left(1 - \frac{\lambda}{\frac{\alpha}{\gamma} + 1}\right)^2, \left(1 - \lambda(2 - \lambda) \frac{1}{1 + \frac{\alpha^2}{\gamma^2}}\right) \right\}^{\frac{1}{2}}$ . Suppose that  $\frac{1}{\gamma}(x - J_{\gamma A} x) \in B[0; \tau]$ . Then the following hold.

- (a)  $\rho \in ]0, 1[$ .
- (b)  $\|y_x - \bar{x}\| \leq \rho \|x - \bar{x}\|$ .
- (c) Suppose that  $\|e\| \leq \varepsilon \|x - z_x\|$  and that  $\eta\varepsilon \in [0, 1[$ . Then

$$\|z_x - \bar{x}\| \leq \frac{\rho + \eta\varepsilon}{1 - \eta\varepsilon} \|x - \bar{x}\|.$$

*Proof.* (i): Because  $J_{\gamma A} x \in B[\bar{x}; \delta]$ , adopt (3.31) with  $x = J_{\gamma A} x$  in the first inequality to derive that

$$\|J_{\gamma A} x - P_{\text{zer } A}(J_{\gamma A} x)\| = d(J_{\gamma A} x, A^{-1}0) \leq \kappa d(0, A(J_{\gamma A} x)) \leq \frac{\kappa}{\gamma} \|x - J_{\gamma A} x\|, \quad (3.33)$$

where we utilize Fact 2.9 in the last inequality. In view of  $P_{\text{zer } A}(J_{\gamma A} x) \in \text{zer } A$ , employing (3.33) and applying Theorem 3.10(iii) with  $t = \frac{\kappa}{\gamma}$  and  $z = P_{\text{zer } A}(J_{\gamma A} x)$ , we establish (i)(a) and (i)(b).

Notice that if  $\text{zer } A = \{\bar{x}\}$ , then  $P_{\text{zer } A}(J_{\gamma A} x) = \bar{x}$ . Hence, (i)(c) is clear from (i)(b).

In addition, according to Lemma 2.8, we know that  $J_{\gamma A}$  is  $\frac{1}{2}$ -averaged and  $\text{Fix } J_{\gamma A} = \text{zer } A \neq \emptyset$ . Hence, combine (i)(b)&(i)(c) with Lemma 2.4 to guarantee (i)(d).

(ii): Employing Fact 2.9 again, we get that  $(J_{\gamma A} x, \frac{1}{\gamma}(x - J_{\gamma A} x)) \in \text{gra } A$ . Taking the assumption,  $\frac{1}{\gamma}(x - J_{\gamma A} x) \in B[0; \tau]$ , and (3.32) into account, we establish that

$$\|J_{\gamma A} x - \bar{x}\| \leq \frac{\alpha}{\gamma} \|x - J_{\gamma A} x\|,$$

which, applying Theorem 3.10(iii) with  $t = \frac{\alpha}{\gamma}$  and  $z = \bar{x}$ , ensures (ii)(a) and (ii)(b).

At last, similarly with the proof of (i)(d) above, (ii)(b) and Lemma 2.4 entail (ii)(c) directly.  $\blacksquare$

**Remark 3.13.** (i) Suppose  $\text{zer } A = \{\bar{x}\}$ . If the assumption that  $A^{-1}$  is Lipschitz continuous at 0 with a positive modulus is strictly stronger than that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , then Theorem 3.12(i) is more interesting than Theorem 3.12(ii).

(ii) The idea of Theorem 3.12(ii)(b) is essentially presented in [8, Theorem 3.1] on the linear convergence rate of the exact version of the generalized proximal point algorithm with the relaxation coefficient being a constant in  $\mathbb{R}^n$ . Notice that the proof of Theorem 3.12(ii)(b) is closely related to Theorem 3.10(i)&(ii), that Lemma 2.14 is critical to the proof of Theorem 3.10(i), and that Lemma 2.14 is inspired by [8, Theorem 3.1] (see Remark 2.15 for details). But the proof of Theorem 3.12(ii)(b) is more natural and easier to understand than that of [8, Theorem 3.1]. In addition, actually we shall use Theorem 3.12 to investigate the linear convergence of the inexact version of generalized proximal point algorithms later.

(iii) It is not difficult to see that Theorem 3.12(ii)(b) can actually also be obtained by employing Fact 3.7 and applying Theorem 3.12(i)(c) with an extra assumption on the distance from  $x$  to  $\text{zer } A$ .

## 4 Inexact Version of the Non-stationary Krasnosel'skiĭ-Mann iterations

In the whole section, we suppose that  $(\alpha_k)_{k \in \mathbb{N}}$  is in  $]0, 1]$ , that  $(\forall k \in \mathbb{N}) \lambda_k \in [0, \frac{1}{\alpha_k}]$ , and that

$$(\forall k \in \mathbb{N}) \quad T_k : \mathcal{H} \rightarrow \mathcal{H} \text{ is } \alpha_k\text{-averaged with } \bigcap_{i \in \mathbb{N}} \text{Fix } T_i \neq \emptyset.$$

Let  $x_0$  and  $(e_k)_{k \in \mathbb{N}}$  be in  $\mathcal{H}$  and let  $(\eta_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$ . In this section, we investigate the *inexact non-stationary Krasnosel'skiĭ-Mann iterations* generated by following the iteration scheme

$$(\forall k \in \mathbb{N}) \quad x_{k+1} = (1 - \lambda_k)x_k + \lambda_k T_k x_k + \eta_k e_k. \quad (4.1)$$

Note that the generalized proximal point algorithm studied in this work is actually a special case of the iteration sequence generated by (4.1). Some results obtained in this section will be applied to generalized proximal point algorithms in the following section.

**Fact 4.1.** [16, Theorem 4.3] *The following statements hold.*

(i)  $(\forall \bar{x} \in \bigcap_{k \in \mathbb{N}} \text{Fix } T_k) (\forall k \in \mathbb{N}) \|x_{k+1} - \bar{x}\| \leq \|x_0 - \bar{x}\| + \sum_{i=0}^k \eta_i \|e_i\|.$

(ii) *Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ . Then the following hold.*

(a)  $\sum_{k \in \mathbb{N}} \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) \|x_k - T_k x_k\|^2 < \infty.$



(b) Suppose that  $\liminf_{k \rightarrow \infty} \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) > 0$  (e.g.,  $\liminf_{k \rightarrow \infty} \lambda_k > 0$  and  $\limsup_{k \rightarrow \infty} \lambda_k < \frac{1}{\limsup_{k \rightarrow \infty} \alpha_k} < \infty$ ). Then  $\sum_{k \in \mathbb{N}} \|x_k - T_k x_k\|^2 < \infty$ . Consequently,  $\lim_{k \rightarrow \infty} \|x_k - T_k x_k\| = 0$ .

**Lemma 4.2.** Denote by  $C := \bigcap_{k \in \mathbb{N}} \text{Fix } T_k$ . Define

$$(\forall k \in \mathbb{N}) \quad y_k := (1 - \lambda_k)x_k + \lambda_k T_k x_k \text{ and } \varepsilon_k := \eta_k \|e_k\| (2 \|y_k - P_C x_k\| + \eta_k \|e_k\|).$$

Then

$$(\forall k \in \mathbb{N}) \quad d^2(x_{k+1}, C) \leq d^2(x_k, C) - \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) \|x_k - T_k x_k\|^2 + \varepsilon_k.$$

*Proof.* Inasmuch as  $(\forall k \in \mathbb{N}) T_k$  is nonexpansive, via [1, Proposition 4.23(ii)],  $C = \bigcap_{k \in \mathbb{N}} \text{Fix } T_k$  is closed and convex. So, by [1, Theorem 3.16],  $(\forall x \in \mathcal{H}) P_C x$  is a well-defined point in  $C$ .

For every  $k \in \mathbb{N}$ , applying (2.1b) in Fact 2.3 with  $T = T_k$ ,  $\alpha = \alpha_k$ ,  $x = x_k$ ,  $y_x = y_k$ ,  $z_x = x_{k+1}$ ,  $\eta = \eta_k$ ,  $e = e_k$ , and  $\bar{x} = P_C x_k$  in the second inequality below, we derive that

$$\begin{aligned} d^2(x_{k+1}, C) &\leq \|x_{k+1} - P_C x_k\|^2 \\ &\leq \|x_k - P_C x_k\|^2 - \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) \|x_k - T_k x_k\|^2 + \varepsilon_k \\ &= d^2(x_k, C) - \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) \|x_k - T_k x_k\|^2 + \varepsilon_k. \end{aligned}$$

Altogether, the proof is complete. ■

**Lemma 4.3.** Let  $\bar{x} \in \bigcap_{k \in \mathbb{N}} \text{Fix } T_k$ . Suppose that  $(\forall k \in \mathbb{N}) \text{Id} - T_k$  is metrically subregular at  $\bar{x}$  for  $0 \in (\text{Id} - T_k) \bar{x}$ , i.e.,

$$(\exists \gamma_k > 0)(\exists \delta_k > 0)(\forall x \in B[\bar{x}; \delta_k]) \quad d(x, \text{Fix } T_k) \leq \gamma_k \|x - T_k x\|. \quad (4.2)$$

Suppose that  $\delta := \inf_{k \in \mathbb{N}} \delta_k > 0$  and that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \delta$ . Let  $0 < \tau \leq \delta - \sum_{k \in \mathbb{N}} \eta_k \|e_k\|$  and let  $x_0 \in B[\bar{x}; \tau]$ . Then the following hold.

(i)  $(\forall k \in \mathbb{N}) x_k \in B[\bar{x}; \delta]$ .

(ii) Suppose that  $(\forall k \in \mathbb{N}) C := \text{Fix } T_k \neq \emptyset$ , that  $\gamma := \sup_{k \in \mathbb{N}} \gamma_k < \infty$ , and that  $\liminf_{k \rightarrow \infty} \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) > 0$  (e.g.,  $\liminf_{k \rightarrow \infty} \lambda_k > 0$  and  $\limsup_{k \rightarrow \infty} \lambda_k < \frac{1}{\limsup_{k \rightarrow \infty} \alpha_k} < \infty$ ). Then  $\sum_{k \in \mathbb{N}} d^2(x_k, C) < \infty$ .

Consequently,  $d(x_k, C) \rightarrow 0$ .

*Proof.* (i): Due to Fact 4.1(i) and the assumptions that  $x_0 \in B[\bar{x}; \tau]$  and  $\tau + \sum_{k \in \mathbb{N}} \eta_k \|e_k\| \leq \delta$ , we easily get (i).

(ii): Based on the assumptions and Fact 4.1(ii)(b),  $\sum_{k \in \mathbb{N}} \|x_k - T_k x_k\|^2 < \infty$ . Bearing (i) in mind and for every  $k \in \mathbb{N}$ , applying (4.2) with  $x = x_k$ , we observe that

$$\sum_{k \in \mathbb{N}} d^2(x_k, C) = \sum_{k \in \mathbb{N}} d^2(x_k, \text{Fix } T_k) \leq \gamma^2 \sum_{k \in \mathbb{N}} \|x_k - T_k x_k\|^2 < \infty.$$

Altogether, the proof is complete. ■

The following result is inspired by the proof of [2, Proposition 4.2(iii)] which is on iteration sequences generated by  $\mathcal{T}$ -class operators (see [2, Page 3] for a detailed definition).

**Proposition 4.4.** *Suppose that  $(\forall k \in \mathbb{N}) \lambda_k \in \left[0, \frac{1}{\alpha_k}\right]$ , that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ , that  $\limsup_{k \rightarrow \infty} \alpha_k > 0$ , and that  $\limsup_{k \rightarrow \infty} \lambda_k < \frac{1}{\limsup_{k \rightarrow \infty} \alpha_k}$ . Define  $M := \limsup_{k \rightarrow \infty} \frac{\lambda_k}{\frac{1}{\alpha_k} - \lambda_k}$ . Then the following hold.*

(i)  $M < \infty$ .

(ii)  $(\exists K \in \mathbb{N}) (\forall k \geq K) \|x_{k+1} - x_k\|^2 \leq 2(M+1)\lambda_k \left(\frac{1}{\alpha_k} - \lambda_k\right) \|x_k - T_k x_k\|^2 + 2\eta_k^2 \|e_k\|^2$ .

(iii)  $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < \infty$ . Consequently,  $\|x_{k+1} - x_k\| \rightarrow 0$ .

*Proof.* (i): Inasmuch as  $\limsup_{k \rightarrow \infty} \alpha_k > 0$  and  $(\forall k \in \mathbb{N}) \alpha_k \in ]0, 1]$ , we observe that  $\limsup_{k \rightarrow \infty} \alpha_k \in ]0, 1]$ , which, connected with  $\limsup_{k \rightarrow \infty} \lambda_k < \frac{1}{\limsup_{k \rightarrow \infty} \alpha_k}$ , ensures that

$$M = \limsup_{k \rightarrow \infty} \frac{\lambda_k}{\frac{1}{\alpha_k} - \lambda_k} \leq \frac{\limsup_{k \rightarrow \infty} \lambda_k}{\frac{1}{\limsup_{k \rightarrow \infty} \alpha_k} - \limsup_{k \rightarrow \infty} \lambda_k} < \infty.$$

(ii): Due to (i), there exists  $K \in \mathbb{N}$  such that  $(\forall k \geq K) \frac{\lambda_k}{\frac{1}{\alpha_k} - \lambda_k} \leq M + 1$ . Using this in the last inequality below, we observe that for every  $k \geq K$ ,

$$\begin{aligned} \|x_{k+1} - x_k\|^2 &\stackrel{(4.1)}{=} \|\lambda_k (T_k x_k - x_k) + \eta_k e_k\|^2 \\ &\leq 2\lambda_k^2 \|T_k x_k - x_k\|^2 + 2\eta_k^2 \|e_k\|^2 \\ &= 2\frac{\lambda_k}{\frac{1}{\alpha_k} - \lambda_k} \lambda_k \left(\frac{1}{\alpha_k} - \lambda_k\right) \|x_k - T_k x_k\|^2 + 2\eta_k^2 \|e_k\|^2 \\ &\leq 2(M+1)\lambda_k \left(\frac{1}{\alpha_k} - \lambda_k\right) \|x_k - T_k x_k\|^2 + 2\eta_k^2 \|e_k\|^2. \end{aligned}$$

(iii): Combining (ii) with Fact 4.1(ii)(a) and the assumption that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ , we derive that  $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < \infty$ , which is followed immediately by the last assertion that  $\|x_{k+1} - x_k\| \rightarrow 0$ . ■

Proposition 4.5 is inspired by [13, Theorem 3]. In particular, it generalizes [13, Theorem 3] by replacing the nonexpansive operator  $T$  therein with a sequence of averaged operators  $(T_k)_{k \in \mathbb{N}}$ . In Proposition 4.5 below, we consider the convergence of the sequence  $(d(x_k, C))_{k \in \mathbb{N}}$ , where  $(x_k)_{k \in \mathbb{N}}$  is generated by (4.1) with  $(\forall k \in \mathbb{N}) C := \text{Fix } T_k$ . Proposition 4.5 will be applied to deduce a  $R$ -linear convergence result on generalized proximal point algorithms in the next section.

**Proposition 4.5.** *Suppose that  $(\forall k \in \mathbb{N}) C := \text{Fix } T_k \neq \emptyset$ . Let  $\bar{x} \in C$ . Suppose that  $(\forall k \in \mathbb{N}) \text{Id} - T_k$  is metrically subregular at  $\bar{x}$  for  $0 \in (\text{Id} - T_k) \bar{x}$ , i.e.,*

$$(\exists \gamma_k > 0)(\exists \delta_k > 0)(\forall x \in B[\bar{x}; \delta_k]) \quad d(x, \text{Fix } T_k) \leq \gamma_k \|x - T_k x\|. \quad (4.3)$$

Suppose that  $\delta := \inf_{k \in \mathbb{N}} \delta_k > 0$  and that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \delta$ . Let  $0 < \tau \leq \delta - \sum_{k \in \mathbb{N}} \eta_k \|e_k\|$  and let  $x_0 \in B[\bar{x}; \tau]$ . Define for every  $k \in \mathbb{N}$

$$\begin{aligned} y_k &:= (1 - \lambda_k)x_k + \lambda_k T_k x_k, \quad \varepsilon_k := \eta_k \|e_k\| (2\|y_k - P_C x_k\| + \eta_k \|e_k\|), \\ \beta_k &:= \frac{\lambda_k \left(\frac{1}{\alpha_k} - \lambda_k\right)}{\gamma_k^2}, \quad \text{and} \quad \rho_k := \begin{cases} 1 - \beta_k & \text{if } \beta_k \leq 1; \\ \frac{1}{1 + \beta_k} & \text{if } \beta_k > 1. \end{cases} \end{aligned}$$

Then the following assertions hold.

(i)  $(\forall k \in \mathbb{N}) d^2(x_{k+1}, C) \leq (1 - \beta_k) d^2(x_k, C) + \varepsilon_k$ .

(ii)  $(\forall k \in \mathbb{N}) \rho_k \in [0, 1]$  and  $d^2(x_{k+1}, C) \leq \rho_k d^2(x_k, C) + \varepsilon_k$ .

(iii)  $(\forall k \in \mathbb{N}) d^2(x_{k+1}, C) \leq \left(\prod_{i=0}^k \rho_i\right) d^2(x_0, C) + \sum_{i=0}^k \left(\prod_{j=i+1}^k \rho_j\right) \varepsilon_i$ . Moreover, the following hold.

(a) Suppose that  $\limsup_{k \rightarrow \infty} \left(\prod_{i=0}^k \rho_i\right)^{\frac{1}{k}} < 1$  and  $(\forall k \in \mathbb{N}) \rho_{k+1} \leq \rho_k$ . Then  $\sum_{k \in \mathbb{N}} d^2(x_k, C) < \infty$ .  
Consequently,  $d(x_k, C) \rightarrow 0$ .

(b) Suppose that  $\rho := \sup_{k \in \mathbb{N}} \rho_k < 1$ . Then

$$(\forall k \in \mathbb{N}) \quad d^2(x_{k+1}, C) \leq \rho^k \left( \rho d^2(x_0, C) + \sum_{i=0}^k \frac{\varepsilon_i}{\rho^i} \right).$$

Consequently, if  $\sum_{k \in \mathbb{N}} \frac{\varepsilon_k}{\rho^k} < \infty$ , then  $\left(d^2(x_k, C)\right)_{k \in \mathbb{N}}$  converges R-linearly to 0.

*Proof.* (i): Based on Lemma 4.3(i), we know that  $(\forall k \in \mathbb{N}) x_k \in B[\bar{x}; \delta]$ . For every  $k \in \mathbb{N}$ , apply (4.3) with  $x = x_k$  to deduce that

$$d(x_k, C) = d(x_k, \text{Fix } T_k) \leq \gamma_k \|x_k - T_k x_k\|. \quad (4.4)$$

Apply Lemma 4.2 in the first inequality below to get that

$$\begin{aligned} d^2(x_{k+1}, C) &\leq d^2(x_k, C) - \lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right) \|x_k - T_k x_k\|^2 + \varepsilon_k \\ &\stackrel{(4.4)}{\leq} d^2(x_k, C) - \frac{\lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right)}{\gamma_k^2} d^2(x_k, C) + \varepsilon_k \\ &\leq \left( 1 - \frac{\lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right)}{\gamma_k^2} \right) d^2(x_k, C) + \varepsilon_k, \end{aligned}$$

which guarantees (i).

(ii): Note that  $(\forall k \in \mathbb{N}) (1 - \beta_k) \leq \frac{1}{1 + \beta_k} \Leftrightarrow 1 - \beta_k^2 \leq 1$ . So we know that  $(\forall k \in \mathbb{N}) 1 - \beta_k \leq \rho_k$ . Hence, (ii) is clear from (i) above.

(iii): Applying (ii), by induction, we easily establish that

$$(\forall k \in \mathbb{N}) \quad d^2(x_{k+1}, C) \leq \left(\prod_{i=0}^k \rho_i\right) d^2(x_0, C) + \sum_{i=0}^k \left(\prod_{j=i+1}^k \rho_j\right) \varepsilon_i. \quad (4.5)$$

For every  $k \in \mathbb{N}$ , applying (2.1a) in Fact 2.3 with  $x = x_k$ ,  $y_x = y_k$ ,  $T = T_k$ ,  $\lambda = \lambda_k$ ,  $\alpha = \alpha_k$ , and  $\bar{x} = P_C x_k$ , we observe that

$$\|y_k - P_C x_k\| \leq \|x_k - P_C x_k\| = d(x_k, C),$$

which, combined with [16, Corollary 4.4(iii)], implies that  $(\|y_k - P_C x_k\|)_{k \in \mathbb{N}}$  is bounded. This together with the assumption  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \delta < \infty$  necessitates that  $\sum_{k \in \mathbb{N}} \varepsilon_k < \infty$ .

(iii)(a): As a consequence of Lemma 2.16(i)&(ii), our assumptions imply that

$$\sum_{k \in \mathbb{N}} \left(\prod_{i=0}^k \rho_i\right) < \infty \quad \text{and} \quad \sum_{k \in \mathbb{N}} \sum_{i=0}^k \left(\prod_{j=i+1}^k \rho_j\right) \varepsilon_i < \infty.$$

This combined with (4.5) ensures that  $\sum_{k \in \mathbb{N}} d^2(x_k, C) < \infty$ , which forces that  $d(x_k, C) \rightarrow 0$ .

(iii)(b): This is immediate from (4.5). ■

In [Corollary 4.6\(i\)](#) below, by applying [Proposition 4.5\(i\)](#), we deduce the main result of [[16](#), [Theorem 4.9](#)] again.

**Corollary 4.6.** *Suppose that  $(\forall k \in \mathbb{N}) C := \text{Fix } T_k \neq \emptyset$  and that  $(\forall k \in \mathbb{N}) e_k \equiv 0$  and  $\eta_k \equiv 0$  in [\(4.1\)](#). Let  $\bar{x} \in C$ . Suppose that  $(\forall k \in \mathbb{N}) \text{Id} - T_k$  is metrically subregular at  $\bar{x}$  for  $0 \in (\text{Id} - T_k) \bar{x}$ , i.e.,*

$$(\exists \gamma_k > 0)(\exists \delta_k > 0)(\forall x \in B[\bar{x}; \delta_k]) \quad d(x, \text{Fix } T_k) \leq \gamma_k \|x - T_k x\|.$$

Suppose that  $\delta := \inf_{k \in \mathbb{N}} \delta_k > 0$ . Let  $x_0 \in B[\bar{x}; \delta]$ . Define

$$(\forall k \in \mathbb{N}) \quad \beta_k := \frac{\lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right)}{\gamma_k^2} \quad \text{and} \quad \rho_k := 1 - \beta_k.$$

Then the following hold.

- (i)  $(\forall k \in \mathbb{N}) \rho_k \in [0, 1]$  and  $d(x_{k+1}, C) \leq \rho_k^{\frac{1}{2}} d(x_k, C)$ . Consequently,  $\lim_{k \rightarrow \infty} d(x_k, C)$  exists.
- (ii) Suppose that  $0 < \underline{\lambda} := \inf_{k \in \mathbb{N}} \lambda_k \leq \bar{\lambda} := \sup_{k \in \mathbb{N}} \lambda_k < \frac{1}{\alpha}$  where  $\alpha := \sup_{k \in \mathbb{N}} \alpha_k > 0$  and that  $\gamma := \sup_{k \in \mathbb{N}} \gamma_k \in \mathbb{R}_{++}$ . Define  $\rho := \sup_{k \in \mathbb{N}} \rho_k$ . Then there exists  $\hat{x} \in C$  such that

$$(\forall k \in \mathbb{N}) \quad \|x_k - \hat{x}\| \leq 2\rho^{\frac{k}{2}} d(x_0, C). \quad (4.6)$$

Consequently,  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in C$ .

*Proof.* We uphold the notation used in [Proposition 4.5](#) above. Notice that if  $(\forall k \in \mathbb{N}) e_k \equiv 0$  and  $\eta_k \equiv 0$ , then, by definition,  $(\forall k \in \mathbb{N}) \varepsilon_k \equiv 0$  in [Proposition 4.5](#). Then [Proposition 4.5\(i\)](#) forces that  $(\forall k \in \mathbb{N}) 1 - \beta_k \in \mathbb{R}_+$ , that is,  $\beta_k \leq 1$ . Hence, we have that  $(\forall k \in \mathbb{N}) \rho_k = 1 - \beta_k \in [0, 1]$  in [Proposition 4.5](#).

(i): According to our analysis above, this is immediate from [Proposition 4.5\(ii\)](#).

(ii): Because  $(\forall k \in \mathbb{N}) \rho_k = 1 - \beta_k \in [0, 1]$  and  $0 < \underline{\lambda} \leq \bar{\lambda} < \frac{1}{\alpha} < \infty$ , we observe that

$$\rho = \sup_{k \in \mathbb{N}} \rho_k = 1 - \inf_{k \in \mathbb{N}} \frac{\lambda_k \left( \frac{1}{\alpha_k} - \lambda_k \right)}{\gamma_k^2} \leq 1 - \frac{\underline{\lambda} \left( \frac{1}{\alpha} - \bar{\lambda} \right)}{\gamma} \in [0, 1[. \quad (4.7)$$

Notice that [[1](#), [Proposition 4.23\(ii\)](#)] implies that  $C$  is nonempty closed and convex and that our assumptions necessitate  $(\forall k \in \mathbb{N}) 0 < \inf_{i \in \mathbb{N}} \lambda_i \leq \lambda_k \leq \sup_{i \in \mathbb{N}} \lambda_i < \frac{1}{\sup_{i \in \mathbb{N}} \alpha_i} \leq \frac{1}{\alpha}$ . Hence, [\(4.7\)](#) combined with (i) above leads to

$$(\forall k \in \mathbb{N}) \quad d(x_{k+1}, C) \leq \rho^{\frac{1}{2}} d(x_k, C),$$

which, connecting with [[16](#), [Theorem 4.9\(i\)](#)] and [[1](#), [Theorem 5.12](#)], ensures [\(4.6\)](#). ■

## 5 Generalized Proximal Point Algorithms

Throughout this section,  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  is maximally monotone with  $\text{zer } A \neq \emptyset$  and

$$(\forall k \in \mathbb{N}) \quad x_{k+1} = (1 - \lambda_k) x_k + \lambda_k J_{c_k A} x_k + \eta_k e_k, \quad (5.1)$$

where  $x_0 \in \mathcal{H}$  is the *initial point* and  $(\forall k \in \mathbb{N}) \lambda_k \in [0, 2]$  and  $\eta_k \in \mathbb{R}_+$  are the *relaxation coefficients*,  $c_k \in \mathbb{R}_{++}$  is the *regularization coefficient*, and  $e_k \in \mathcal{H}$  is the *error term*.

Generalized proximal point algorithms generate the iteration sequence by conforming to the scheme [\(5.1\)](#). The classic proximal point algorithm generates the iteration sequence by following [\(5.1\)](#) with  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$ ,  $e_k \equiv 0$ , and  $\eta_k \equiv 0$ . In this section, we investigate the linear convergence of generalized proximal point algorithms for solving the monotone inclusion problem, i.e., finding a zero of the associated monotone operator.

## 5.1 Auxiliary results

**Proposition 5.1** is motivated by [12, Theorem 3.1] and [22, Theorem 3.1]. In particular, **Proposition 5.1(i)(a)** and **Proposition 5.1(ii)(a)** reduce to [12, Theorem 3.1] and [22, Theorem 3.1], respectively, when  $(\forall k \in \mathbb{N}) c_k \equiv c \in \mathbb{R}_{++}$ . **Proposition 5.1(i)** works on the local convergence of the exact version of the proximal point algorithm. Note that if we restrict  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$  in **Proposition 5.1(ii)(a)**, the convergence rate in **Proposition 5.1(ii)(b)** is better than that of **Proposition 5.1(ii)(a)** since  $\frac{1}{1+\frac{c_k^2}{\kappa^2}} = 1 - \frac{1}{1+\frac{\kappa^2}{c_k^2}} \leq 1 - \frac{1}{\left(1+\frac{\kappa}{c_k}\right)^2}$ .

**Proposition 5.1.** *Suppose that  $(\forall k \in \mathbb{N}) e_k \equiv 0$  and  $\eta_k \equiv 0$  in (5.1). Let  $\bar{x}$  be in  $\text{zer } A$ . Suppose that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,*

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad d(x, A^{-1}0) \leq \kappa d(0, Ax). \quad (5.2)$$

Set  $c := \inf_{k \in \mathbb{N}} c_k$ . Then the following assertions hold.

(i) Suppose that  $x_0 \in B[\bar{x}; \delta]$  and that  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$ . Then

(a)  $(\forall k \in \mathbb{N}) d(x_{k+1}, \text{zer } A) \leq \frac{1}{\sqrt{1+\frac{c_k^2}{\kappa^2}}} d(x_k, \text{zer } A);$

(b) if  $c > 0$ , then  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in \text{zer } A$ .

(ii) Suppose that  $x_k \rightarrow \bar{x}$ . Then the following hold.

(a) Set  $(\forall k \in \mathbb{N}) \rho_k := \left(1 - \lambda_k(2 - \lambda_k) \frac{1}{\left(1+\frac{\kappa}{c_k}\right)^2}\right)^{\frac{1}{2}}$ . Then

i. there exists  $K \in \mathbb{N}$  such that  $(\forall k \geq K) d(x_{k+1}, \text{zer } A) \leq \rho_k d(x_k, \text{zer } A);$

ii. if  $c > 0$  and  $0 < \underline{\lambda} := \inf_{k \in \mathbb{N}} \lambda_k \leq \bar{\lambda} := \sup_{k \in \mathbb{N}} \lambda_k < 2$ , then  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in \text{zer } A$ .

(b) Suppose that  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$ . Then

i. there exists  $K \in \mathbb{N}$  such that  $(\forall k \geq K) d(x_{k+1}, \text{zer } A) \leq \frac{1}{\sqrt{1+\frac{c_k^2}{\kappa^2}}} d(x_k, \text{zer } A);$

ii. if  $c > 0$ , then  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in \text{zer } A$ .

*Proof.* (i): For every  $k \in \mathbb{N}$ , applying **Fact 2.10** with  $x = x_k$ ,  $y_x = x_{k+1}$ ,  $\lambda = \lambda_k$ , and  $\gamma = c_k$ , we know that  $\|x_{k+1} - \bar{x}\| \leq \|x_k - \bar{x}\|$ ; and employing **Fact 2.11** with  $x = x_k$ ,  $\gamma = c_k$ , and  $z = \bar{x}$ , we observe that  $\|J_{c_k A} x_k - \bar{x}\| \leq \|x_k - \bar{x}\|$ . Combine these results with  $x_0 \in B[\bar{x}; \delta]$  to deduce that

$$x_k \in B[\bar{x}; \delta] \quad \text{and} \quad J_{c_k A} x_k \in B[\bar{x}; \delta]. \quad (5.3)$$

Let  $k \in \mathbb{N}$ . Taking (5.3) into account and applying **Lemma 3.3** with  $x = x_k$  and  $\gamma = c_k$ , we obtain that

$$d(x_{k+1}, \text{zer } A) = d(J_{c_k A} x_k, \text{zer } A) \leq \frac{1}{\sqrt{1+\frac{c_k^2}{\kappa^2}}} d(x_k, \text{zer } A).$$

Suppose that  $c = \inf_{k \in \mathbb{N}} c_k > 0$ . Then  $\rho := \sup_{k \in \mathbb{N}} \frac{1}{\sqrt{1+\frac{c_k^2}{\kappa^2}}} = \frac{1}{\sqrt{1+\frac{c^2}{\kappa^2}}} \in [0, 1[$  and

$$(\forall k \in \mathbb{N}) \quad d(x_{k+1}, \text{zer } A) \leq \rho d(x_k, \text{zer } A). \quad (5.4)$$

Notice that, via [1, Proposition 23.39] and by virtue of the maximal monotonicity of  $A$ ,  $\text{zer } A$  is closed and convex. Similarly with the proof of [Corollary 4.6\(ii\)](#), combining [\(5.4\)](#) with [16, Theorem 5.6(i)] and [1, Theorem 5.12], we obtain that  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in \text{zer } A$ .

(ii): Because  $x_k \rightarrow \bar{x}$ , we know that there exists  $K \in \mathbb{N}$  such that  $(\forall k \geq K) x_k \in B[\bar{x}; \delta]$ , which, connected with [Fact 2.11](#), ensures that

$$(\forall k \geq K) \quad J_{c_k A} x_k \in B[\bar{x}; \delta]. \quad (5.5)$$

(ii)(a): Let  $k \in \mathbb{N}$  such that  $k \geq K$ . Bearing [\(5.5\)](#) in mind and applying [Theorem 3.4\(ii\)](#) with  $x = x_k$ ,  $y_x = x_{k+1}$ ,  $\gamma = c_k$ ,  $\lambda = \lambda_k$ , and  $\rho = \rho_k$ , we deduce that

$$d(x_{k+1}, \text{zer } A) \leq \|x_{k+1} - P_{\text{zer } A} x_k\| \leq \rho_k \|x_k - P_{\text{zer } A} x_k\| = \rho_k d(x_k, \text{zer } A). \quad (5.6)$$

Suppose that  $0 < \underline{\lambda} = \inf_{k \in \mathbb{N}} \lambda_k \leq \bar{\lambda} = \sup_{k \in \mathbb{N}} \lambda_k < 2$  and  $c = \inf_{k \in \mathbb{N}} c_k > 0$ . Then  $\rho := \sup_{k \in \mathbb{N}} \rho_k \leq \left(1 - \underline{\lambda} (2 - \bar{\lambda}) \frac{1}{(1 + \frac{\bar{\lambda}}{c})^2}\right)^{\frac{1}{2}} \in [0, 1[$ . Moreover, due to [\(5.6\)](#),

$$(\forall k \geq K) \quad d(x_{k+1}, \text{zer } A) \leq \rho^{k-K+1} d(x_K, \text{zer } A),$$

which, combined with [16, Theorem 5.6(i)] and [1, Theorem 5.12], guarantees that  $(x_k)_{k \in \mathbb{N}}$  converges  $R$ -linearly to a point  $\hat{x} \in \text{zer } A$ .

(ii)(b): Invoking [\(5.5\)](#) and employing almost the same arguments used in the proof of [\(i\)\(a\)](#) and [\(ii\)\(a\)ii.](#) above, we obtain [\(ii\)\(b\)](#).  $\blacksquare$

**Corollary 5.2.** *Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ . Set  $c := \inf_{k \in \mathbb{N}} c_k$ . Then the following hold.*

- (i) *Suppose that  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Then  $\sum_{k \in \mathbb{N}} \|x_k - J_{c_k A} x_k\|^2 < \infty$ . Moreover, if  $c > 0$ , then  $\frac{1}{c_k} (x_k - J_{c_k A} x_k) \rightarrow 0$ .*
- (ii) *Suppose that  $\sup_{k \in \mathbb{N}} \lambda_k < 2$ . Then  $\sum_{k \in \mathbb{N}} \|x_k - x_{k+1}\|^2 < \infty$ . Moreover, if  $c > 0$  and  $\lambda := \inf_{k \in \mathbb{N}} \lambda_k > 0$ , then  $\frac{1}{c_k} (x_k - J_{c_k A} x_k) \rightarrow 0$ .*

*Proof.* According to [Lemma 2.8](#),  $(\forall k \in \mathbb{N}) J_{c_k A}$  is  $\frac{1}{2}$ -averaged operator and  $\text{Fix } J_{c_k A} = \text{zer } A \neq \emptyset$ .

(i): Applying [Fact 4.1\(ii\)\(b\)](#) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$  and  $\alpha_k = \frac{1}{2}$ , we deduce  $\sum_{k \in \mathbb{N}} \|x_k - J_{c_k A} x_k\|^2 < \infty$ , which forces  $\|x_k - J_{c_k A} x_k\| \rightarrow 0$ .

In addition, if  $c = \inf_{k \in \mathbb{N}} c_k > 0$ , then

$$\left\| \frac{1}{c_k} (x_k - J_{c_k A} x_k) \right\| \leq \frac{1}{c} \|x_k - J_{c_k A} x_k\| \rightarrow 0.$$

(ii): It is not difficult to verify that  $\sup_{k \in \mathbb{N}} \lambda_k < 2$  if and only if  $\limsup_{k \rightarrow \infty} \lambda_k < 2$  and  $(\forall k \in \mathbb{N}) \lambda_k < 2$ . Apply [Proposition 4.4\(iii\)](#) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$  and  $\alpha_k = \frac{1}{2}$  to yield that  $\sum_{k \in \mathbb{N}} \|x_{k+1} - x_k\|^2 < \infty$  and  $\|x_{k+1} - x_k\| \rightarrow 0$ . Suppose that  $c = \inf_{k \in \mathbb{N}} c_k > 0$  and  $\lambda = \inf_{k \in \mathbb{N}} \lambda_k > 0$ . Based on [\(5.1\)](#),

$$\left\| \frac{1}{c_k} (x_k - J_{c_k A} x_k) \right\| = \left\| \frac{1}{c_k \lambda_k} (x_k - x_{k+1} + \eta_k e_k) \right\| \leq \frac{1}{c \lambda} (\|x_k - x_{k+1}\| + \eta_k \|e_k\|) \rightarrow 0.$$

Altogether, the proof is complete.  $\blacksquare$

**Proposition 5.3.** *Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$  and that  $\sum_{k \in \mathbb{N}} \left|1 - \frac{c_{k+1}}{c_k}\right| < \infty$ . Then the following hold.*

- (i)  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\|$  exists.

(ii) Assume that  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$ . Then  $x_k - J_{c_k A} x_k \rightarrow 0$ .

*Proof.* (i): Because  $\sum_{k \in \mathbb{N}} \left| 1 - \frac{c_{k+1}}{c_k} \right| < \infty$  and  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ , due to [16, Lemma 5.2(ii)] and [1, Lemma 5.31], we know that  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\|$  exists in  $\mathbb{R}_+$ .

(ii): Applying Fact 4.1(ii)(a) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$  and  $\alpha_k = \frac{1}{2}$  and employing the assumption  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$ , we establish that  $\liminf_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\| = 0$ . This as well as (i) yields  $x_k - J_{c_k A} x_k \rightarrow 0$ . ■

Theorem 5.4 generalizes [23, Theorem 4.5] by replacing the constant  $\lambda \in ]0, 2[$  therein with a sequence  $(\lambda_k)_{k \in \mathbb{N}}$  in  $[0, 2]$ . To do this generalization, we can require that  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$  like [6, Theorem 3] or that  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$  and  $\sum_{k \in \mathbb{N}} \left| 1 - \frac{c_{k+1}}{c_k} \right| < \infty$  like our Theorem 5.4(ii). In [6, Theorem 3], to obtain the required weak convergence of the sequence generated by (5.1) with  $(\forall k \in \mathbb{N}) \eta_k = \lambda_k$ , the authors require that  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$ , which is stronger than our assumption  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$  in Theorem 5.4, but our assumption  $\sum_{k \in \mathbb{N}} \left| 1 - \frac{c_{k+1}}{c_k} \right| < \infty$  in Theorem 5.4(ii) is not needed for [6, Theorem 3].

**Theorem 5.4.** Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$  and  $\inf_{k \in \mathbb{N}} c_k > 0$ . Then the following assertions hold.

(i) Suppose that  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\| = 0$ . Then  $(x_k)_{k \in \mathbb{N}}$  converges weakly to a point in  $\text{zer } A$ .

(ii) Suppose that  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$  and  $\sum_{k \in \mathbb{N}} \left| 1 - \frac{c_{k+1}}{c_k} \right| < \infty$ . Then  $(x_k)_{k \in \mathbb{N}}$  converges weakly to a point in  $\text{zer } A$ .

*Proof.* (i): Taking  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\| = 0$ ,  $\inf_{k \in \mathbb{N}} c_k > 0$ , and [15, Proposition 2.16(i)] into account, we know that  $\Omega((x_k)_{k \in \mathbb{N}}) \subseteq \text{zer } A$ . On the other hand, due to [16, Lemma 5.1(iii)(b)], we have that  $(\forall z \in \text{zer } A) \lim_{k \rightarrow \infty} \|x_k - z\|$  exists in  $\mathbb{R}_+$ . These results combined with [1, Lemma 2.47] entail that  $(x_k)_{k \in \mathbb{N}}$  converges weakly to a point in  $\text{zer } A$ .

(ii): Combine our assumption with Proposition 5.3(ii) to get that  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\| = 0$ . Hence, the desired weak convergence is clear from (i) above. ■

The convergence result of Corollary 5.5 is consistent with that of [6, Theorem 3] except that the assumption  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$  in [6, Theorem 3] is replaced by  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$  in Corollary 5.5.

**Corollary 5.5.** Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ , that  $\inf_{k \in \mathbb{N}} c_k > 0$ , and that  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Then  $(x_k)_{k \in \mathbb{N}}$  converges weakly to a point in  $\text{zer } A$ .

*Proof.* Clearly, the assumption  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$  entails that  $\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) = \infty$ . Moreover, apply Fact 4.1(ii)(a) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$  and  $\alpha_k = \frac{1}{2}$  to get that

$$\sum_{k \in \mathbb{N}} \lambda_k (2 - \lambda_k) \|x_k - J_{c_k A} x_k\|^2 < \infty,$$

which, combined with  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ , guarantees that  $\sum_{k \in \mathbb{N}} \|x_k - J_{c_k A} x_k\|^2 < \infty$  and hence,  $\lim_{k \rightarrow \infty} \|x_k - J_{c_k A} x_k\| = 0$ .

Therefore, by Theorem 5.4(i), we obtain the required weak convergence. ■

## 5.2 Linear convergence of generalized proximal point algorithms

In this subsection, we consider the linear convergence of generalized proximal point algorithms.

Theorem 5.6 shows a  $Q$ -linear convergence result of generalized proximal point algorithms.

**Theorem 5.6.** Let  $\bar{x} \in \text{zer } A$ . Suppose that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad \mathbf{d}(x, A^{-1}0) \leq \kappa \mathbf{d}(0, Ax).$$

Let  $(\varepsilon_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$  such that  $\eta_k \varepsilon_k \rightarrow 0$ . Suppose that  $(\forall k \in \mathbb{N}) \|e_k\| \leq \varepsilon_k \|x_k - x_{k+1}\|$  and  $\lambda_k \in ]0, 2[$ , that  $J_{c_k A} x_k \rightarrow \bar{x}$ , and that  $c := \liminf_{k \rightarrow \infty} c_k > 0$  and  $0 < \underline{\lambda} := \liminf_{k \rightarrow \infty} \lambda_k \leq \bar{\lambda} := \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Set

$$(\forall k \in \mathbb{N}) \quad \rho_k := \max \left\{ \left( 1 - \frac{\lambda_k}{\frac{\kappa}{c_k} + 1} \right)^2, \left( 1 - \lambda_k (2 - \lambda_k) \frac{1}{1 + \frac{\kappa^2}{c_k^2}} \right) \right\}^{\frac{1}{2}}.$$

Then the following statements hold.

(i) For every  $k$  large enough, we have that  $\rho_k \in ]0, 1[$  and that

$$\|x_{k+1} - P_{\text{zer } A}(J_{c_k A} x_k)\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - P_{\text{zer } A}(J_{c_k A} x_k)\|.$$

Moreover, there exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - P_{\text{zer } A}(J_{c_k A} x_k)\| \leq \mu \|x_k - P_{\text{zer } A}(J_{c_k A} x_k)\| \leq \mu^{k-K+1} \|x_K - P_{\text{zer } A}(J_{c_K A} x_K)\|.$$

(ii) Suppose that  $\text{zer } A = \{\bar{x}\}$ . Then for every  $k$  large enough,

$$\|x_{k+1} - \bar{x}\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|. \quad (5.7)$$

Moreover, there exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - \bar{x}\| \leq \mu \|x_k - \bar{x}\| \leq \mu^{k-K+1} \|x_K - \bar{x}\|.$$

*Proof.* (i): Inasmuch as  $J_{c_k A} x_k \rightarrow \bar{x}$  and  $\eta_k \varepsilon_k \rightarrow 0$ , there exists  $K_1 \in \mathbb{N}$  such that

$$(\forall k \geq K_1) \quad J_{c_k A} x_k \in B[\bar{x}; \delta] \quad \text{and} \quad \eta_k \varepsilon_k \in \left[0, \frac{1}{2}\right]. \quad (5.8)$$

For every  $k \geq K_1$ , applying [Theorem 3.12\(i\)\(a\)&\(i\)\(d\)](#) with  $x = x_k$ ,  $z_x = x_{k+1}$ ,  $\gamma = c_k$ ,  $\lambda = \lambda_k$ ,  $\eta = \eta_k$ ,  $e = e_k$ , and  $\varepsilon = \varepsilon_k$ , and employing [\(5.8\)](#), we get  $\rho_k \in ]0, 1[$  and

$$\|x_{k+1} - P_{\text{zer } A}(J_{c_k A} x_k)\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - P_{\text{zer } A}(J_{c_k A} x_k)\|. \quad (5.9)$$

In addition, because  $c = \liminf_{k \rightarrow \infty} c_k > 0$  and  $0 < \underline{\lambda} = \liminf_{k \rightarrow \infty} \lambda_k \leq \bar{\lambda} = \limsup_{k \rightarrow \infty} \lambda_k < 2$ , we observe that  $\limsup_{k \rightarrow \infty} \left| 1 - \frac{\lambda_k}{\frac{\kappa}{c_k} + 1} \right| \leq \max \left\{ \left| 1 - \frac{\underline{\lambda}}{\frac{\kappa}{c} + 1} \right|, |\bar{\lambda} - 1| \right\}$  and that

$$\rho := \limsup_{k \rightarrow \infty} \rho_k \leq \max \left\{ \left| 1 - \frac{\underline{\lambda}}{\frac{\kappa}{c} + 1} \right|, |\bar{\lambda} - 1|, \left( 1 - \underline{\lambda} (2 - \bar{\lambda}) \frac{1}{1 + \frac{\kappa^2}{c^2}} \right)^{\frac{1}{2}} \right\} < 1.$$

Since  $\eta_k \varepsilon_k \rightarrow 0$ , we have that  $\limsup_{k \rightarrow \infty} \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} = \rho < 1$ , which necessitates that there exist  $K \geq K_1$  and  $\mu \in ]\rho, 1[$  such that

$$(\forall k \geq K) \quad \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \leq \mu.$$



This combined with (5.9) deduces the last assertion in (i).

(ii): Notice that the assumption  $\text{zer } A = \{\bar{x}\}$  forces that

$$(\forall k \in \mathbb{N}) \quad \text{P}_{\text{zer } A} (J_{c_k A} x_k) \equiv \bar{x}.$$

Therefore, (ii) is immediate from (i). ■

**Remark 5.7.** We uphold the assumption and notation of Theorem 5.6. Taking Corollary 2.13 into account, we observe that if  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$ , then

$$(\forall k \in \mathbb{N}) \quad \rho_k = \left(1 - \frac{1}{1 + \frac{\kappa^2}{c_k^2}}\right)^{\frac{1}{2}} = \frac{1}{\sqrt{1 + \frac{c_k^2}{\kappa^2}}}.$$

Note that this convergence rate is consistent with that of Proposition 5.1(i)(a).

**Proposition 5.8.** Suppose that  $\mathcal{H} = \mathbb{R}^n$ . Suppose that  $\bar{x} \in \text{zer } A$  and that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad \text{d} \left( x, A^{-1}0 \right) \leq \kappa \text{d} (0, Ax).$$

Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ . Let  $(\varepsilon_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$  such that  $\eta_k \varepsilon_k \rightarrow 0$ . Suppose that  $(\forall k \in \mathbb{N}) \|e_k\| \leq \varepsilon_k \|x_k - x_{k+1}\|$  and  $\lambda_k \in ]0, 2[$ , and that  $\liminf_{k \rightarrow \infty} c_k > 0$  and  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Set

$$(\forall k \in \mathbb{N}) \quad \rho_k := \max \left\{ \left(1 - \frac{\lambda_k}{\frac{\kappa}{c_k} + 1}\right)^2, \left(1 - \lambda_k (2 - \lambda_k) \frac{1}{1 + \frac{\kappa^2}{c_k^2}}\right) \right\}^{\frac{1}{2}}.$$

Then the following statements hold.

(i) For every  $k$  large enough, we have that

$$\|x_{k+1} - \text{P}_{\text{zer } A} (J_{c_k A} x_k)\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \text{P}_{\text{zer } A} (J_{c_k A} x_k)\|.$$

Moreover, there exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - \text{P}_{\text{zer } A} (J_{c_k A} x_k)\| \leq \mu \|x_k - \text{P}_{\text{zer } A} (J_{c_k A} x_k)\| \leq \mu^{k-K+1} \|x_K - \text{P}_{\text{zer } A} (J_{c_K A} x_K)\|.$$

(ii) Suppose that  $\text{zer } A = \{\bar{x}\}$ . Then for every  $k$  large enough,

$$\|x_{k+1} - \bar{x}\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|.$$

Moreover, there exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - \bar{x}\| \leq \mu \|x_k - \bar{x}\| \leq \mu^{k-K+1} \|x_K - \bar{x}\|.$$

*Proof.* Because  $(c_k)_{k \in \mathbb{N}}$  is in  $\mathbb{R}_{++}$ , it is not difficult to verify that  $\liminf_{k \rightarrow \infty} c_k > 0 \Leftrightarrow \inf_{k \in \mathbb{N}} c_k > 0$ . Moreover, bearing  $\mathcal{H} = \mathbb{R}^n$ ,  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ , and  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$  in mind, and employing Corollary 5.5, we know that  $x_k \rightarrow \bar{x}$ . Notice that, via Fact 2.11,  $(\forall k \in \mathbb{N}) \|J_{c_k A} x_k - \bar{x}\| \leq \|x_k - \bar{x}\|$ . These results entail that

$$J_{c_k A} x_k \rightarrow \bar{x}.$$

Therefore, in view of Theorem 5.6, we obtain the required results. ■

**Theorem 5.9.** Suppose that  $A^{-1}$  is Lipschitz continuous at 0 with modulus  $\alpha > 0$ , i.e.,  $A^{-1}(0) = \{\bar{x}\}$  and there exists  $\tau > 0$  such that

$$\left( \forall (w, x) \in \text{gra } A^{-1} \text{ with } w \in B[0; \tau] \right) \quad \|x - \bar{x}\| \leq \alpha \|w\|.$$

Suppose that  $\frac{1}{c_k} (x_k - J_{c_k A} x_k) \rightarrow 0$ . Let  $(\varepsilon_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$  such that  $\eta_k \varepsilon_k \rightarrow 0$ . Suppose that  $(\forall k \in \mathbb{N}) \|e_k\| \leq \varepsilon_k \|x_k - x_{k+1}\|$  and  $\lambda_k \in ]0, 2[$ , and that  $\liminf_{k \rightarrow \infty} c_k > 0$  and  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Set

$$(\forall k \in \mathbb{N}) \quad \rho_k := \max \left\{ \left( 1 - \frac{\lambda_k}{\frac{\alpha}{c_k} + 1} \right)^2, \left( 1 - \lambda_k (2 - \lambda_k) \frac{1}{1 + \frac{\alpha^2}{c_k^2}} \right) \right\}^{\frac{1}{2}}.$$

Then the following statements hold.

(i) For every  $k$  large enough, we have that  $\rho_k \in ]0, 1[$  and that

$$\|x_{k+1} - \bar{x}\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|.$$

(ii) There exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - \bar{x}\| \leq \mu \|x_k - \bar{x}\| \leq \mu^{k-K+1} \|x_K - \bar{x}\|. \quad (5.10)$$

*Proof.* (i): In view of  $\frac{1}{c_k} (x_k - J_{c_k A} x_k) \rightarrow 0$  and  $\eta_k \varepsilon_k \rightarrow 0$ , there exists  $K_1 \in \mathbb{N}$  such that

$$(\forall k \geq K_1) \quad \frac{1}{c_k} (x_k - J_{c_k A} x_k) \in B[0; \tau] \quad \text{and} \quad \eta_k \varepsilon_k \in \left[ 0, \frac{1}{2} \right]. \quad (5.11)$$

For every  $k \geq K_1$ , applying [Theorem 3.12\(ii\)\(a\)&\(ii\)\(c\)](#) with  $x = x_k$ ,  $z_x = x_{k+1}$ ,  $\gamma = c_k$ ,  $\lambda = \lambda_k$ ,  $\eta = \eta_k$ ,  $e = e_k$ , and  $\varepsilon = \varepsilon_k$ , and employing (5.11), we derive  $\rho_k \in ]0, 1[$  and

$$\|x_{k+1} - \bar{x}\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|. \quad (5.12)$$

(ii): Similarly with the proof of the last part of [Theorem 5.6\(i\)](#), there exist  $K \geq K_1$  and  $\mu \in [0, 1[$  such that

$$(\forall k \geq K) \quad \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \leq \mu,$$

which, combined with (5.12), guarantees (5.10). ■

**Proposition 5.10.** Suppose that  $A^{-1}$  is Lipschitz continuous at 0 with modulus  $\alpha > 0$ , i.e.,  $A^{-1}(0) = \{\bar{x}\}$  and there exists  $\tau > 0$  such that

$$\left( \forall (w, x) \in \text{gra } A^{-1} \text{ with } w \in B[0; \tau] \right) \quad \|x - \bar{x}\| \leq \alpha \|w\|.$$

Suppose that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \infty$ . Let  $(\varepsilon_k)_{k \in \mathbb{N}}$  be in  $\mathbb{R}_+$  such that  $\eta_k \varepsilon_k \rightarrow 0$ . Suppose that  $(\forall k \in \mathbb{N}) \|e_k\| \leq \varepsilon_k \|x_k - x_{k+1}\|$  and  $\lambda_k \in ]0, 2[$ , and that  $\liminf_{k \rightarrow \infty} c_k > 0$  and  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . Set

$$(\forall k \in \mathbb{N}) \quad \rho_k := \max \left\{ \left( 1 - \frac{\lambda_k}{\frac{\alpha}{c_k} + 1} \right)^2, \left( 1 - \lambda_k (2 - \lambda_k) \frac{1}{1 + \frac{\alpha^2}{c_k^2}} \right) \right\}^{\frac{1}{2}}.$$

Then the following hold.

(i) For every  $k$  large enough, we have that

$$\|x_{k+1} - \bar{x}\| \leq \frac{\rho_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|. \quad (5.13)$$

(ii) There exist  $\mu \in [0, 1[$  and  $K \in \mathbb{N}$  such that

$$(\forall k \geq K) \quad \|x_{k+1} - \bar{x}\| \leq \mu \|x_k - \bar{x}\| \leq \mu^{k-K+1} \|x_K - \bar{x}\|.$$

*Proof.* Similarly with the proof of [Proposition 5.8](#), the assumptions  $\liminf_{k \rightarrow \infty} c_k > 0$  and  $(\forall k \in \mathbb{N}) c_k > 0$  ensure that  $\inf_{k \in \mathbb{N}} c_k > 0$ . Then, via [Corollary 5.2\(i\)](#), we establish that  $\frac{1}{c_k} (x_k - J_{c_k A} x_k) \rightarrow 0$ .

Hence, as a consequence of [Theorem 5.9](#), we obtain the required results.  $\blacksquare$

**Remark 5.11.** We uphold the notation used in [Proposition 5.10](#).

(i) [Proposition 5.10](#) improves [[23](#), Theorem 4.7] from the following two aspects.

- Note that  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$  if and only if  $(\forall k \in \mathbb{N}) \lambda_k \in ]0, 2[$  and  $0 < \liminf_{k \rightarrow \infty} \lambda_k \leq \limsup_{k \rightarrow \infty} \lambda_k < 2$ . We see clearly that the constant  $\lambda \in ]0, 2[$  in [[23](#), Theorem 4.7] is replaced by a sequence  $(\lambda_k)_{k \in \mathbb{N}}$  satisfying  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$  in [Proposition 5.10](#).
- Notice that the linear convergence result provided in [[23](#), Theorem 4.7] is essentially that for every  $k$  large enough,  $\|x_{k+1} - \bar{x}\| \leq \frac{\zeta_k + \eta_k \varepsilon_k}{1 - \eta_k \varepsilon_k} \|x_k - \bar{x}\|$ , where

$$(\forall k \in \mathbb{N}) \quad \zeta_k := \max \left\{ \left( 1 - \lambda \frac{1}{1 + \frac{\alpha^2}{c_k^2}} \right), \left( 1 - \lambda (2 - \lambda) \frac{1}{1 + \frac{\alpha^2}{c_k^2}} \right) \right\}^{\frac{1}{2}} \quad \text{with } \lambda \in ]0, 2[.$$

Bearing [Remark 3.11](#) in mind, we conclude that the convergence rate given in (5.13) of our [Proposition 5.10\(i\)](#) is better than the corresponding rate in [[23](#), Theorem 4.7].

(ii) Taking [Corollary 2.13](#) into account, we observe that if  $(\forall k \in \mathbb{N}) \lambda_k \equiv 1$ , then  $(\forall k \in \mathbb{N})$

$$\rho_k = \left( 1 - \frac{1}{1 + \frac{\alpha^2}{c_k^2}} \right)^{\frac{1}{2}} = \frac{1}{\sqrt{1 + \frac{\alpha^2}{c_k^2}}} = \frac{\alpha}{(\alpha^2 + c_k^2)^{\frac{1}{2}}}. \quad \text{Therefore, we see that [Proposition 5.10](#) extends [[20](#),$$

[Theorem 2\]](#) from  $\lambda \equiv 1$  to a sequence  $(\lambda_k)_{k \in \mathbb{N}}$  satisfying  $0 < \inf_{k \in \mathbb{N}} \lambda_k \leq \sup_{k \in \mathbb{N}} \lambda_k < 2$ .

[Theorem 5.12](#) below provides  $R$ -linear convergence results on generalized proximal point algorithms. It is an application of [Proposition 4.5](#) which shows a  $R$ -linear convergence result on the non-stationary Krasnosel'skiĭ-Mann iterations.

**Theorem 5.12.** Let  $\bar{x} \in \text{zer } A$ . Suppose that  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ , i.e.,

$$(\exists \kappa > 0)(\exists \delta > 0)(\forall x \in B[\bar{x}; \delta]) \quad \text{d} \left( x, A^{-1}0 \right) \leq \kappa \text{d} (0, Ax). \quad (5.14)$$

Suppose that  $c := \inf_{k \in \mathbb{N}} c_k > 0$  and that  $\sum_{k \in \mathbb{N}} \eta_k \|e_k\| < \delta$ . Let  $0 < \tau \leq \delta - \sum_{k \in \mathbb{N}} \eta_k \|e_k\|$  and let  $x_0 \in B[\bar{x}; \tau]$ . Define

$$(\forall k \in \mathbb{N}) \quad y_k := (1 - \lambda_k)x_k + \lambda_k T_k x_k \quad \text{and} \quad \varepsilon_k := \eta_k \|e_k\| (2 \|y_k - P_{\text{zer } A} x_k\| + \eta_k \|e_k\|).$$

Set  $(\forall k \in \mathbb{N}) \gamma_k := \left( 1 + \frac{\kappa}{c_k} \right)$ ,  $\beta_k := \frac{\lambda_k(2 - \lambda_k)}{\gamma_k^2}$ , and  $\rho_k := 1 - \beta_k$ . Denote by  $\rho := \sup_{k \in \mathbb{N}} \rho_k$ ,  $\underline{\lambda} := \inf_{k \in \mathbb{N}} \lambda_k$ , and  $\bar{\lambda} := \sup_{k \in \mathbb{N}} \lambda_k$ . Then the following statements hold.

(i)  $(\forall k \in \mathbb{N}) \rho_k \in [0, 1]$  and  $d^2(x_{k+1}, \text{zer } A) \leq \rho_k d^2(x_k, \text{zer } A) + \varepsilon_k$ .

(ii)  $(\forall k \in \mathbb{N}) d^2(x_{k+1}, \text{zer } A) \leq \left(\prod_{i=0}^k \rho_i\right) d^2(x_0, \text{zer } A) + \sum_{i=0}^k \left(\prod_{j=i+1}^k \rho_j\right) \varepsilon_i$ .

(iii) Suppose that  $0 < \underline{\lambda} \leq \bar{\lambda} < 2$ . Then the following hold.

(a)  $0 \leq \rho \leq 1 - \frac{\underline{\lambda}(2-\bar{\lambda})}{(1+\frac{\kappa}{c})^2} < 1$ .

(b)  $(\forall k \in \mathbb{N}) d^2(x_{k+1}, \text{zer } A) \leq \rho^k \left(\rho d^2(x_0, \text{zer } A) + \sum_{i=0}^k \frac{\varepsilon_i}{\rho^i}\right)$ . Consequently, if  $\sum_{k \in \mathbb{N}} \frac{\varepsilon_k}{\rho^k} < \infty$ , then  $\left(d^2(x_k, \text{zer } A)\right)_{k \in \mathbb{N}}$  converges R-linearly to 0.

(iv) Suppose that  $0 < \underline{\lambda} \leq \bar{\lambda} < 2$  and that  $(\forall k \in \mathbb{N}) e_k \equiv 0$  and  $\eta_k \equiv 0$ . Then the following hold.

(a)  $(\forall k \in \mathbb{N}) d(x_{k+1}, \text{zer } A) \leq \rho_k^{\frac{1}{2}} d(x_k, \text{zer } A)$ .

(b) There exists a point  $\hat{x} \in \text{zer } A$  such that

$$(\forall k \in \mathbb{N}) \quad \|x_k - \hat{x}\| \leq 2\rho_k^{\frac{1}{2}} d(x_0, \text{zer } A).$$

Consequently,  $(x_k)_{k \in \mathbb{N}}$  converges R-linearly to a point  $\hat{x} \in \text{zer } A$ .

*Proof.* In view of [Lemma 2.8](#),  $(\forall k \in \mathbb{N}) J_{c_k A}$  is  $\frac{1}{2}$ -averaged operator and

$$(\forall k \in \mathbb{N}) \quad (\text{Id} - J_{c_k A})^{-1} 0 = \text{Fix } J_{c_k A} = \text{zer } A.$$

Moreover, employing [\(5.14\)](#) and for every  $k \in \mathbb{N}$ , applying [Fact 3.2](#) with  $\gamma = c_k$ , we know that  $(\forall k \in \mathbb{N}) \text{Id} - J_{c_k A}$  is metrically subregular at  $\bar{x}$  for  $0 = (\text{Id} - J_{c_k A}) \bar{x}$ ; more precisely,

$$(\forall x \in B[\bar{x}; \delta]) \quad d(x, \text{Fix } J_{c_k A}) \leq \left(1 + \frac{\kappa}{c_k}\right) \|x - J_{c_k A} x\|. \quad (5.15)$$

**(i)&(ii):** Inasmuch as  $(\forall k \in \mathbb{N}) \lambda_k \in [0, 2]$  and  $\gamma_k > 1$ , we have that  $(\forall k \in \mathbb{N}) \lambda_k(2 - \lambda_k) \in [0, 1]$  and  $\beta_k = \frac{\lambda_k(2-\lambda_k)}{\gamma_k^2} \in [0, 1]$ . Hence,  $(\forall k \in \mathbb{N}) \rho_k \in [0, 1]$ .

Apply [Proposition 4.5\(ii\)&\(iii\)](#) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$ ,  $C = \text{zer } A$ ,  $\gamma_k = \left(1 + \frac{\kappa}{c_k}\right)$ ,  $\delta_k \equiv \delta$ , and  $\alpha_k \equiv \frac{1}{2}$  to derive results in **(i)&(ii)**.

**(iii)(a):** Note that

$$\begin{aligned} \sup_{k \in \mathbb{N}} \gamma_k &= \sup_{k \in \mathbb{N}} \left(1 + \frac{\kappa}{c_k}\right) = 1 + \frac{\kappa}{\inf_{k \in \mathbb{N}} c_k} = 1 + \frac{\kappa}{c} \\ \Rightarrow \rho &= \sup_{k \in \mathbb{N}} \rho_k = 1 - \inf_{k \in \mathbb{N}} \beta_k \leq 1 - \frac{\underline{\lambda}(2-\bar{\lambda})}{\left(1 + \frac{\kappa}{c}\right)^2} \in ]0, 1[. \end{aligned}$$

Hence, **(iii)(a)** is true.

**(iii)(b):** Employing **(iii)(a)** above and applying [Proposition 4.5\(iii\)\(b\)](#) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$ ,  $C = \text{zer } A$ ,  $\gamma_k = \left(1 + \frac{\kappa}{c_k}\right)$ ,  $\delta_k \equiv \delta$ , and  $\alpha_k \equiv \frac{1}{2}$ , we establish the required results in **(iii)(b)**.

**(iv):** Taking **(iii)(a)** above into account and applying [Corollary 4.6](#) with  $(\forall k \in \mathbb{N}) T_k = J_{c_k A}$ ,  $C = \text{zer } A$ ,  $\gamma_k = \left(1 + \frac{\kappa}{c_k}\right)$ ,  $\delta_k \equiv \delta$ , and  $\alpha_k \equiv \frac{1}{2}$ , we directly obtain the desired results in **(iv)**.  $\blacksquare$

## 6 Conclusion and Future Work

In this work, we considered the metrical subregularity of set-valued operators, which is a popular assumption for the linear convergence of optimization algorithms. We also provided an  $R$ -linear convergence result on the non-stationary Krasnosel'skiĭ-Mann iterations. Because the generalized proximal point algorithm is a special instance of the non-stationary Krasnosel'skiĭ-Mann iterations, the result on the non-stationary Krasnosel'skiĭ-Mann iterations was applied to the generalized proximal point algorithm. In addition, we showed some  $Q$ -linear convergence results on the generalized proximal point algorithm under the assumption that the associated monotone operator is metrically subregular or that the inverse of the monotone operator is Lipschitz continuous with a positive modulus.

Given a maximally monotone operator  $A : \mathcal{H} \rightarrow 2^{\mathcal{H}}$  with  $\text{zer } A = \{\bar{x}\}$ , as stated in [Question 3.8](#) and [Remark 3.13\(i\)](#), it is interesting to know if the following two statements are equivalent.

- (i)  $A$  is metrically subregular at  $\bar{x}$  for  $0 \in A\bar{x}$ .
- (ii)  $A^{-1}$  is Lipschitz continuous at 0 with a positive modulus.

We stated in [Fact 3.7](#) that (ii) implies (i) and we also found in [Example 3.9](#) a specific  $A : \mathbb{R} \rightarrow \mathbb{R}$  suggesting a positive answer for the equivalence. In the future, we shall try either proving (i)  $\Rightarrow$  (ii) or finding a counterexample for it. In addition, as we presented in the introduction, many popular optimization algorithms are instances of the generalized proximal point algorithm when the associated monotone operator is specified accordingly. We shall also apply our linear convergence results to some particular examples of the generalized proximal point algorithm.

## References

- [1] H. H. BAUSCHKE AND P. L. COMBETTES: *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*, second edition, Springer, 2017.
- [2] P. L. COMBETTES: Quasi-Fejérian analysis of some optimization algorithms, in *Inherently parallel algorithms in feasibility and optimization and their applications (Haifa, 2000)* North-Holland, Amsterdam, 2001, pp. 115–152.
- [3] E. CORMAN AND X. YUAN: A generalized proximal point algorithm and its convergence rate, *SIAM Journal on Optimization* 24 (2014), pp. 1614–1638.
- [4] A. L. DONTCHEV AND R. T. ROCKAFELLAR: *Implicit Functions and Solution Mappings*, second edition, Springer, 2014.
- [5] J. DOUGLAS AND H. H. RACHFORD: On the numerical solution of heat conduction problems in two and three space variables, *Transactions of the American Mathematical Society* 82 (1956), pp. 421–439.
- [6] J. ECKSTEIN AND D. P. BERTSEKAS: On the Douglas-Rachford splitting method and the proximal point algorithm for maximal monotone operators, *Mathematical Programming* 55 (1992), pp. 293–318.
- [7] D. GABAY AND B. MERCIER: A dual algorithm for the solution of nonlinear variational problems via finite element approximation, *Computers and Mathematics with Applications* 2 (1976), pp. 17–40.
- [8] Y. GU AND J. YANG: On the optimal linear convergence factor of the relaxed proximal point algorithm for monotone inclusion problems, *arXiv Preprint: 1905.04537* (2019), pp. 1–9.

- [9] M. R. HESTENES: Multiplier and gradient methods, *Journal of Optimization Theory and Applications* 4 (1969), pp. 303–320.
- [10] D. KIM: Accelerated proximal point method for maximally monotone operators, *Mathematical Programming* 190 (2021), pp. 57–87.
- [11] G. M. KORPELEVIČ: An extragradient method for finding saddle points and for other problems, *Ėkonomika i Matematicheskie Metody* 12 (1976), pp. 747–756.
- [12] D. LEVENTHAL: Metric subregularity and the proximal point method, *Journal of Mathematical Analysis and Applications* 360 (2009), pp. 681–688.
- [13] J. LIANG, J. FADILI, AND G. PEYRÉ: Convergence rates with inexact non-expansive operators, *Mathematical Programming* 159 (2016), pp. 403–434.
- [14] P. L. LIONS AND B. MERCIER: Splitting algorithms for the sum of two nonlinear operators, *SIAM Journal on Numerical Analysis* 16 (1979), pp. 964–979.
- [15] H. OUYANG: Weak and strong convergence of generalized proximal point algorithms, *arXiv preprint: 2110.07015* (2021), pp. 1–30.
- [16] H. OUYANG: On the Stability of Krasnosel’skiĭ-Mann Iterations, *arXiv preprint: 2203.04527* (2022), pp. 1–30.
- [17] D. W. PEACEMAN AND H. H. RACHFORD: The numerical solution of parabolic and elliptic differential equations, *Journal of the Society for Industrial and Applied Mathematics* 3 (1955), pp. 28–41.
- [18] B. T. POLYAK: *Introduction to Optimization*, Translations Series in Mathematics and Engineering, Optimization Software, Inc., Publications Division, New York, 1978.
- [19] M. J. D. POWELL: A method for nonlinear constraints in minimization problems, in *Optimization (Sympos., Univ. Keele, Keele, 1968)* Academic Press, London, 1969, pp. 283–298.
- [20] T. R. ROCKAFELLAR: Monotone operators and the proximal point algorithm, *SIAM Journal on Control and Optimization* 14 (1976), pp. 877–898.
- [21] W. RUDIN: *Principles of Mathematical Analysis*, International Series in Pure and Applied Mathematics, McGraw-Hill Book Co., New York-Auckland-Düsseldorf, third ed., 1976. Optimization Software, Inc., Publications Division, New York, 1978.
- [22] L. SHEN AND S. PAN: Linear convergence of the generalized PPA and several splitting methods for the composite inclusion problem, *arXiv preprint: 1508.05156* (2016), pp. 1–22.
- [23] M. TAO AND X. YUAN: On the optimal linear convergence rate of a generalized proximal point algorithm, *Journal of Scientific Computing* 74 (2018), pp. 826–850.
- [24] X. ZHANG, M. BURGER, X. BRESSON, AND S. OSHER: Bregmanized nonlocal regularization for deconvolution and sparse reconstruction, *SIAM Journal on Imaging Sciences* 3 (2010), pp. 253–276.