K-STABILITY OF GORENSTEIN FANO GROUP COMPACTIFICATIONS WITH RANK TWO

JAE-HYOUK LEE, KYEONG-DONG PARK, AND SUNGMIN YOO

ABSTRACT. We give a classification of Gorenstein Fano bi-equivariant compactifications of semisimple complex Lie groups with rank two, and determine which of them are equivariant K-stable and admit (singular) Kähler–Einstein metrics. As a consequence, we obtain several explicit examples of K-stable Fano varieties admitting (singular) Kähler–Einstein metrics. We also compute the greatest Ricci lower bounds, equivalently the delta invariants for K-unstable varieties. This gives us three new examples on which each solution of the Kähler–Ricci flow is of type II.

1. INTRODUCTION

The Yau–Tian–Donaldson conjecture for the Fano case states that a Fano manifold admits a Kähler–Einstein metric if and only if it satisfies the algebraic geometric condition, called the K-polystability. This conjecture for the Fano case were completely solved by Chen–Donaldson–Sun [CDS15a, CDS15b, CDS15c] and Tian [Tia15]. Recently, Li [Li22] generalized this conjecture to Q-Fano varieties with singular Kähler–Einstein metrics using the notion of uniform K-stability. In general, however, it is difficult to verify the K-stability condition to show the existence of a Kähler–Einstein metric since one should consider infinite number of possible degenerations (test configurations).

Despite this issue, when the manifold has large symmetry, we can reduce the problem to checking only finite number of degenerations. In fact, Datar–Székelyhidi [DS16] proved that if a Fano manifold X is equivariant K-stable, i.e., K-stable with respect to special degenerations that are G-equivariant for some reductive subgroup G of $\operatorname{Aut}_0(X)$, then it admits a Kähler–Einstein metric. Using this theorem, they could recover the theorem of Wang–Zhu which says that a toric Fano manifold admits a Kähler–Einstein metric if and only if the Futaki invariant vanishes [WZ04]. By Mabuchi's theorem [Mab87], this is equivalent to that the barycenter of the moment polytope locates at the origin.

The spherical varieties consist of an important and large class of highly symmetric varieties which include toric varieties and (more generally) bi-equivariant compactifications of reductive Lie groups, called the *group compactifications*. In [Del17], Delcroix proved that a smooth Fano group compactification admits a Kähler–Einstein metric if and only if the barycenter of the corresponding moment polytope translated by -2ρ locates in the relative interior of the cone generated by positive roots, where 2ρ denotes the sum of all positive roots of G. He also proved in [Del20] that for Q-Fano spherical varieties, this combinatorial condition is equivalent to the equivariant K-stability.

Very recently, Delcroix's theorem for smooth Fano group compactifications is generalized to \mathbb{Q} -Fano group compactifications by Li–Tian–Zhu [LTZ23]. They proved that the same combinatorial criterion is also applicable to check the existence of singular Kähler–Einstein metric using a variational approach. Combining this with the above result by Delcroix, we have the following

²⁰¹⁰ Mathematics Subject Classification. Primary: 14M27, 32Q20, Secondary: 32M12, 53C55.

Key words and phrases. singular Kähler–Einstein metrics, equivariant K-stability, Gorenstein Fano group compactifications, moment polytopes, greatest Ricci lower bounds, Kähler–Ricci flow.

Theorem 1.1 (Delcroix [Del17, Del20], Li–Tian–Zhu [LTZ23]). Let X be a \mathbb{Q} -Fano variety which is a bi-equivariant compactification of reductive complex Lie group G. Then the followings are equivalent.

- (1) X admits a singular Kähler–Einstein metric.
- (2) X is K-stable with respect to special degenerations that are G-equivariant.
- (3) $bar_{DH}(\Delta) \in 2\rho + \Xi$, where $bar_{DH}(\Delta)$ is the barycenter of the moment polytope Δ with respect to the Duistermaat-Heckman measure and Ξ is the relative interior of the cone generated by positive roots of G.

In this paper, we study Gorenstein Fano group compactifications. Recall that a complete normal variety X is called *Gorenstein Fano* if its anticanonical divisor $-K_X$ is Cartier and ample. For toric varieties, it is well-known that classifying Gorenstein Fano toric varieties is equivalent to classifying reflexive lattice polytopes; see e.g. [CLS11, Theorem 8.3.4]. In particular, Gorenstein Fano toric surfaces correspond to 16 equivalence classes of reflexive lattice polygons in the plane up to lattice equivalence (cf. [CLS11, Theorem 8.3.7]).

By applying theorems for generalizing the results for the toric case, we classify all Gorenstein Fano bi-equivariant compactifications of semisimple complex Lie groups with rank two, and determine the equivariant K-stability so as to conclude the admittance of (singular) Kähler–Einstein metrics.

Lie type	dima Y	G	Gorenstein Fano	K -stable (\exists KE)	K -unstable (\nexists KE)
The type	me type dime A	U	total(smooth, singular)	total(smooth, singular)	total(smooth, singular)
		$\operatorname{SL}_2(\mathbb{C}) \times \operatorname{SL}_2(\mathbb{C})$	15(2,13)	3(2,1)	$12 (\cdot, 12)$
$\Lambda_{\perp} \times \Lambda_{\perp}$	$A_1 \times A_1 = 6$	$\mathrm{PSL}_2(\mathbb{C}) \times \mathrm{PSL}_2(\mathbb{C})$	7(2,5)	5(2,3)	$2(\cdot, 2)$
$A_1 \times A_1$	0	$\operatorname{SL}_2(\mathbb{C}) \times \operatorname{PSL}_2(\mathbb{C})$	14(2,12)	4(1,3)	10(1,9)
		$\mathrm{SO}_4(\mathbb{C})$	6(3,3)	2(1,1)	4(2,2)
٨	Λο	$\mathrm{SL}_3(\mathbb{C})$	5(3,2)	2(1,1)	3(2,1)
A 2	0	$\mathrm{PSL}_3(\mathbb{C})$	$3~(3,\cdot)$	$3~(3,\cdot)$	
Ba	10	$\operatorname{Sp}_4(\mathbb{C})$	4(3,1)	$2 (2, \cdot)$	2(1,1)
D ₂ 10	$\mathrm{SO}_5(\mathbb{C})$	4(2,2)	4(2,2)		
G ₂	14	G_2	$2 (2, \cdot)$	$2 (2, \cdot)$	•
	t	otal	60(22,38)	27 (16, 11)	33 (6, 27)

Theorem 1.2. There are exactly 60 Gorenstein Fano (22 smooth, 38 singular) varieties which are bi-equivariant compactifications of semisimple complex Lie groups with rank two. Among them, 27 (16 smooth, 11 singular) varieties are equivariant K-stable and admit Kähler–Einstein metrics.

We note that Delcroix [Del20] already obtained the related results for the case of smooth $SO_4(\mathbb{C})$ compactifications, and Li–Tian–Zhu [LTZ23] gave the full description of Gorenstein Fano $SO_4(\mathbb{C})$ compactifications including K-stabilities. We also note that in [Del15], the case of all smooth
toroidal simple group compactifications was treated.

One advantage of the combinatorial approach is that we can explicitly compute the value of the greatest Ricci lower bound R(X) (also called Tian's β -invariant), which is a measurement how far the Fano manifold X is from being K-stable. More precisely, Delcroix [Del17] proved that for a smooth Fano group compactification without a Kähler–Einstein metric, the greatest Ricci lower bound satisfies

$$R(X) = \sup\left\{t < 1 \colon \frac{1}{1-t}(2\rho - t \cdot \mathbf{bar}_{DH}(\Delta)) \in -\Xi + \Delta\right\}$$

Recently, it is known that for a K-unstable Fano manifold X, the greatest Ricci lower bound R(X) is equal to the delta invariant $\delta(X)$, which is defined by Fujita–Odaka [FO18] using the log canonical threshold [BBJ21, CRZ19]. Using this fact for the K-unstable smooth Fano varieties in the above theorem, we also obtain the following.

Theorem 1.3. Let X be a K-unstable smooth Fano variety which is a bi-equivariant compactification of a semisimple complex Lie group G with rank two. Denote its Picard number by $\rho(X)$. Then the greatest Ricci lower bound R(X), equivalently the delta invariant $\delta(X)$ is as follows:

G	$SL_2 \times PSL_2$	$\mathrm{SO}_4(\mathbb{C})$		SL	$_{3}(\mathbb{C})$	$\operatorname{Sp}_4(\mathbb{C})$		
$\rho(X)$	3	2	3	2	3	3		
R(X)	$\frac{8869}{9333} \approx 0.95$	$\frac{49}{51} \approx 0.96$	$\frac{75257}{99843} \approx 0.75$	$\frac{1419621}{1493483} \approx 0.95$	$\frac{21100419}{28437901} \approx 0.74$	$\frac{1046175339}{1236719713} \approx 0.84$		

The greatest Ricci lower bound of the last case was already computed in [Del17, Example 6.9]. We remark that by a result of Golota [Gol20], one can also compute the delta invariants of K-unstable singular Fano varieties applying the same formula.

The above theorem induces interesting facts in Kähler–Ricci flow. For a Fano manifold X with a Kähler form $\omega_0 \in 2\pi c_1(X)$, it is proved by Cao [Cao85] that the normalized Kähler–Ricci flow

$$\frac{d\omega(t)}{dt} = -\operatorname{Ric}(\omega(t)) + \omega(t), \quad \omega(0) = \omega_0,$$

has solution for all $t \geq 0$. Moreover, Perelman [Per02] proved that if X has the unique Kähler– Einstein metric ω_{KE} , then $\omega(t)$ smoothly converges to ω_{KE} . However, in general, it may not have the limit. It is proved in [LTZ18] that if a smooth Fano equivariant compactification X of a semisimple complex Lie group admits no Kähler–Einstein metrics, then any solution of the Kähler–Ricci flow is of type II, that is, the curvature of $\omega(t)$ is not uniformly bounded. As a result, Li–Tian–Zhu [LTZ18] obtain three examples having type II solutions of the Kähler–Ricci flow from smooth Fano equivariant compactifications of SO₄(\mathbb{C}) and Sp₄(\mathbb{C}). We get three additional such examples from the above classification.

Corollary 1.4. The manifolds in Theorem 1.3 have type II solutions of the Kähler–Ricci flows.

This paper is organized as follows. In Section 2, we recall the theory of spherical varieties by focusing on equivariant group compactifications. In Section 3, we discuss a combinatorial criterion for the existence of a singular Kähler–Einstein metric on a Q-Fano group compactification and its greatest Ricci lower bound in terms of the barycenter of its moment polytope with respect to the Duistermaat–Heckman measure. In Section 4, we give a classification of Gorenstein Fano bi-equivariant compactifications of semisimple complex Lie groups with rank two, and determine which of them are K-stable and admit (singular) Kähler–Einstein metrics.

Acknowledgements. Jae-Hyouk Lee was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2019R1F1A1058962). Kyeong-Dong Park was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2019R1A2C3010487, NRF-2021R1C1C2092610). Sungmin Yoo was supported by the Institute for Basic Science (IBS-R032-D1).

Kyeong-Dong Park would like to sincerely thank Michel Brion for his kind answers to several questions related to moment polytopes. The authors thank the anonymous referee for the careful reading and many valuable suggestions.

2. Spherical varieties and algebraic moment polytopes

Let G be a connected reductive algebraic group over \mathbb{C} .

2.1. Spherical varieties and colors. Bi-equivariant compactifications of G are spherical G-varieties. In the following, we review general notions and results about spherical varieties. We refer [Kno91], [Tim11] and [Gan18] as references for spherical varieties.

Definition 2.1. A normal variety X equipped with an action of G is called *spherical* if a Borel subgroup B of G acts on X with an open orbit.

Let G/H be an open G-orbit of a spherical variety X and T be a maximal torus of B. For a character $\chi \in \mathfrak{X}(B) = \mathfrak{X}(T)$, let $\mathbb{C}(G/H)_{\chi}^{(B)} = \{f \in \mathbb{C}(G/H) : b \cdot f = \chi(b)f$ for all $b \in B\}$ be the set of B-semi-invariant functions in $\mathbb{C}(G/H)$ associated to χ , where $\mathbb{C}(G/H) = \mathbb{C}(X)$ denotes the function field of G/H.

The spherical weight lattice \mathcal{M} of G/H is defined as a subgroup of the character group $\mathfrak{X}(T)$ such that each element $\chi \in \mathcal{M}$ has the non-zero set of *B*-semi-invariant functions, that is,

$$\mathcal{M} = \{ \chi \in \mathfrak{X}(T) : \mathbb{C}(G/H)_{\chi}^{(B)} \neq 0 \}$$

Note that every function f_{χ} in $\mathbb{C}(G/H)_{\chi}^{(B)}$ is determined by its weight χ up to constant. This is because any *B*-invariant rational function on G/H is constant, that is, $\mathbb{C}(G/H)^B = \mathbb{C}$. The spherical weight lattice \mathcal{M} is a free abelian group of finite rank. We define the rank of G/H as the rank of the lattice \mathcal{M} . Let $\mathcal{N} = \text{Hom}(\mathcal{M}, \mathbb{Z})$ denote its dual lattice together with the natural pairing $\langle \cdot, \cdot \rangle \colon \mathcal{N} \times \mathcal{M} \to \mathbb{Z}$.

As the open B-orbit of a spherical variety X is an affine variety, its complement has pure codimension one and is a finite union of B-stable prime divisors.

Definition 2.2. For a spherical homogeneous space G/H, *B*-stable prime divisors in G/H are called *colors* of G/H. We denote by $\mathfrak{D} = \{D_1, \dots, D_k\}$ the set of colors of G/H.

Every discrete \mathbb{Q} -valued valuation ν of the function field $\mathbb{C}(G/H)$ induces a homomorphism $\rho(\nu) \colon \mathcal{M} \to \mathbb{Q}$ which is defined by $\langle \rho(\nu), \chi \rangle = \nu(f_{\chi})$, where $f_{\chi} \in \mathbb{C}(G/H)_{\chi}^{(B)} \setminus \{0\}$. Hence, we get a map

 ρ : {discrete \mathbb{Q} -valued valuations on G/H} $\to \mathcal{N} \otimes \mathbb{Q}$.

Luna and Vust [LV83] showed that the restriction of ρ to the set \mathcal{V} of *G*-invariant discrete valuations on *G*/*H* is injective. Since the map ρ is injective on \mathcal{V} , we may consider \mathcal{V} as a subset of $\mathcal{N} \otimes \mathbb{Q}$ via ρ . It is known that \mathcal{V} is a full-dimensional (co)simplicial cone of $\mathcal{N} \otimes \mathbb{Q}$, which is called the *valuation cone* of *G*/*H*. For a *B*-stable divisor *D* in *X*, we have a discrete valuation ν_D associated to *D*, that is, $\nu_D(f)$ is the vanishing order of *f* along *D*. For the sake of simplicity, we simplify the notation $\rho(\nu_D)$ as $\rho(D)$.

Remark 2.3. The normal equivariant embeddings of a given spherical homogeneous space are classified by combinatorial objects called *colored fans*, which generalize the fans appearing in the classification of toric varieties. In a brief way, a colored fan is a finite collection of colored cones, which is a pair $(\mathcal{C}, \mathfrak{F})$ consisting of $\mathfrak{F} \subset \mathfrak{D}$ and a strictly convex cone $\mathcal{C} \subset \mathcal{N} \otimes \mathbb{Q}$ generated by $\rho(\mathfrak{F})$ and finitely many elements in the valuation cone \mathcal{V} (see [LV83] and [Kno91] for details).

2.2. Algebraic moment polytopes and anticanonical line bundles. Let L be a G-linearized ample line bundle on a spherical G-variety X. By the multiplicity-free property of spherical varieties, the algebraic moment polytope $\Delta(X, L)$ encodes the structure of representation of G in the spaces of multi-sections of tensor powers of L.

Definition 2.4. The algebraic moment polytope $\Delta(X, L)$ of L with respect to B is defined as the closure of $\bigcup_{k \in \mathbb{N}} \Delta_k / k$ in $\mathfrak{X}(T) \otimes \mathbb{R}$, where Δ_k is a finite set consisting of (dominant) weights λ such that

$$H^0(X, L^{\otimes k}) = \bigoplus_{\lambda \in \Delta_k} V_G(\lambda).$$

Here, $V_G(\lambda)$ means the irreducible representation of G with highest weight λ .

The algebraic moment polytope $\Delta(X, L)$ for a polarized (spherical) *G*-variety *X* was introduced by Brion in [Bri87] as a purely algebraic version of the Kirwan polytope. This is indeed the convex hull of finitely many points in $\mathfrak{X}(T) \otimes \mathbb{R}$ (see [Bri87]).

For a spherical G-variety X, consider the open B-orbit. As B stabilizes the open B-orbit, the stabilizer P under the G-action of the open B-orbit is a parabolic subgroup of G. Let ρ_P be the

half sum of roots of P. Then, by [GH15, Remark 4.3] the *B*-weight ξ of a *B*-semi-invariant global section $s \in \Gamma(X, K_X^{-1})$ of the anticanonical bundle K_X^{-1} is equal to $2\rho_P$. From [Bri97, Theorem 4.2] and [Lun97, Section 3.6], we can get an explicit expression of the anticanonical divisor $-K_X$ for a spherical variety X. Furthermore, based on the works of Brion [Bri89, Bri97], Gagliardi and Hofscheier [GH15, Section 9] described the (algebraic) moment polytope of the anticanonical line bundle on a Gorenstein Fano spherical variety.

Proposition 2.5. Let X be a Gorenstein Fano embedding of a spherical homogeneous space G/H. If a B-stable Weil divisor $-K_X = \sum_{i=1}^k m_i D_i + \sum_{j=1}^\ell E_j$ represents the anticanonical line bundle K_X^{-1} for colors D_i and G-stable divisors E_j in X, the moment polytope $\Delta(X, K_X^{-1})$ is $2\rho_P + Q_X^*$, where the polytope Q_X is the convex hull of the set

$$\left\{\frac{\rho(D_i)}{m_i}: i=1,\cdots,k\right\} \cup \left\{\rho(E_j): j=1,\cdots,\ell\right\}$$

in $\mathcal{N} \otimes \mathbb{R}$ and its dual polytope Q_X^* is defined as $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle n, m \rangle \geq -1 \text{ for every } n \in Q_X\}.$

2.3. Gorenstein Fano group compactifications. We refer the reader to [Tim03], [AB04], [AK05], [Tim11] and [Del17] as references for group compactifications. Any reductive algebraic group G is isomorphic to a symmetric homogeneous space $(G \times G)/\text{diag}(G)$ under the action of the group $G \times G$ for the involution $\theta(g_1, g_2) = (g_2, g_1), g_1, g_2 \in G$. Indeed, G is spherical with respect to the action of $G \times G$ by left and right multiplication from the Bruhat decomposition. If T is a maximal torus of G, then $T \times T$ is a maximal torus of $G \times G$ and we get the spherical weight lattice

$$\mathcal{M} = \mathfrak{X}((T \times T)/\operatorname{diag}(T)) = \{(\lambda, -\lambda) : \lambda \in \mathfrak{X}(T)\}.$$

Thus, \mathcal{M} can be identified with the character lattice $\mathfrak{X}(T)$ of T by the projection to the first coordinate. Under this identification, we also identify the dual lattice \mathcal{N} of \mathcal{M} with the dual character lattice $\mathfrak{X}(T)^{\vee}$.

We now describe the colors of G. As $B^-B \subset G$ is open for the opposite Borel B^- of B, by the Bruhat decomposition the colors of G coincide with the Schubert divisors $D_{\alpha} := \overline{B^- s_{\alpha} B}$ for simple root α , where $s_{\alpha} \in N_G(T)$ is a representative for the simple reflection associated to α . Therefore, the colors are identified with the simple roots of G and the image of D_{α} under the map ρ is equal to the corresponding coroot α^{\vee} , that is, $\rho(D_{\alpha}) = \alpha^{\vee}$ (see e.g. [Gan18, Example 3.6]). Recall that the coroot α^{\vee} of a root α is defined as the unique element in $[\mathfrak{g},\mathfrak{g}] \cap \mathfrak{t}$ such that $\alpha(x) = \frac{2 \kappa(x, \alpha^{\vee})}{\kappa(\alpha^{\vee}, \alpha^{\vee})}$ for all $x \in \mathfrak{t}$, where κ is the Cartan–Killing form on the Lie algebra \mathfrak{g} of G and \mathfrak{t} denotes the Lie algebra of T.

Corollary 2.6. Let X be a Gorenstein Fano equivariant compactification of a reductive group G. If E_j are G-stable divisors in X for $1 \le j \le \ell$, then the moment polytope $\Delta(X, K_X^{-1})$ is a polytope of which facets lie in any Weyl wall and the affine hyperplanes $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle \rho(E_j), m - 2\rho \rangle = -1\}$ for $1 \le j \le \ell$, where $2\rho = \sum_{\alpha \in \Phi^+} \alpha$ is the sum of all positive roots of G.

Proof. Suppose that the rank of G is r. For colors D_i and G-stable divisors E_j in X, a Weil divisor $-K_X = \sum_{i=1}^r 2D_i + \sum_{j=1}^\ell E_j$ represents the anticanonical line bundle K_X^{-1} by [AB04, Section 5.2]. Then, for a color D_i associated to a simple root α_i , the image $\rho(D_i) = \alpha_i^{\vee}$ gives an inequality

$$\left\langle \frac{\alpha_i^{\vee}}{2}, m - 2\rho \right\rangle \ge -1 \quad \iff \quad \left\langle \alpha_i^{\vee}, m \right\rangle \ge 0 \quad \text{for all } 1 \le i \le r$$

Thus, the images of all colors D_1, \dots, D_r determine the positive restricted Weyl chamber, and the result follows from Proposition 2.5.

Conversely, given reductive algebraic group G a lattice polytope in $\mathcal{M} \otimes \mathbb{R}$ with one vertex as the origin and Weyl walls as facets determine a Gorenstein Fano group compactification if the other facets lie on affine hyperplanes $\{m \in \mathcal{M} \otimes \mathbb{R} : \langle \nu, m - 2\rho \rangle = -1\}$ defined by primitive elements ν in \mathcal{N} . In general, Gagliardi and Hofscheier [GH15] classify the Gorenstein Fano spherical embeddings of a spherical homogeneous space G/H in terms of G/H-reflexive polytopes.

Example 2.7 (Gorenstein Fano compactifications of $SL_2(\mathbb{C})$ and $PSL_2(\mathbb{C})$). As $SL_2(\mathbb{C})$ is simplyconnected, the spherical weight lattice \mathcal{M} of $SL_2(\mathbb{C})$ is spanned by the fundamental weight ϖ_1 and its dual lattice \mathcal{N} is spanned by the coroot α_1^{\vee} . The only possible lattice moment polytope is $[0, 3\varpi_1]$ from the inequality $\langle -\alpha_1^{\vee}, (x-2)\varpi_1 \rangle \geq -1$. Indeed, this is the moment polytope of the 3-dimensional hyperquadric \mathbb{Q}^3 which is obtained by an equivariant open embedding

 $\operatorname{SL}_2(\mathbb{C}) \hookrightarrow X := \{ [x,t] : \operatorname{det}(x) = t^2 \} \subset \mathbb{P}(\operatorname{Mat}_{2 \times 2}(\mathbb{C}) \oplus \mathbb{C}), x \mapsto [x,1].$

Note that we have the anticanonical divisor $-K_{\mathbb{Q}^3} = 2D + E$ with $\rho(E) = -\alpha_1^{\vee}$, where $D = \overline{B^- s_{\alpha_1} B}$ and $E = \mathbb{Q}^3 \setminus \mathrm{SL}_2(\mathbb{C})$.

On the other hand, since $\text{PSL}_2(\mathbb{C}) = \text{SL}_2(\mathbb{C})/\{\pm I\} \cong \text{SO}_3(\mathbb{C})$ is of adjoint type, the spherical weight lattice \mathcal{M} of $\text{PSL}_2(\mathbb{C})$ is spanned by the simple root $\alpha_1 = 2\varpi_1$ and its dual lattice \mathcal{N} is spanned by the coweight ϖ_1^{\vee} . In this case, the only possible lattice moment polytope is $[0, 2\alpha_1] = [0, 4\varpi_1]$ from the inequality $\langle -\varpi_1^{\vee}, (x-1)\alpha_1 \rangle \geq -1$. Indeed, this is the moment polytope of the 3-dimensional projective space \mathbb{P}^3 which is an example of *wonderful compactifications* constructed by De Concini and Procesi [DCP83] (see also [LPY21, Example 2.4]).

3. Singular Kähler–Einstein metrics and greatest Ricci lower bounds

Let X be a Q-Fano variety, i.e., a normal projective complex variety such that $-K_X$ is Q-Cartier and ample.

3.1. Singular Kähler–Einstein metrics. By Kodaira's embedding theorem, we can embed a \mathbb{Q} -Fano variety X into a projective space \mathbb{P}^N using a basis of $H^0(X, K_X^{-k})$ for some k > 0. Let ω_{FS} be the Fubini–Study metric of \mathbb{P}^N . We choose a reference Kähler form ω_0 on X by

$$\omega_0 := \frac{1}{k} \omega_{FS}|_X \in 2\pi c_1(X).$$

Let h be the normalized Ricci potential of ω_0 such that $\operatorname{Ric}(\omega_0) - \omega_0 = \sqrt{-1}\partial\bar{\partial}h$ on X_{reg} , where X_{reg} means the complement of the singular locus of X.

Definition 3.1. Let X be a Q-Fano variety with log terminal singularities. A singular Kähler-Einstein metric on X is a current ω_{φ} satisfying the complex Monge-Ampère equation

$$\omega_{\varphi}^n = e^{h - \varphi} \omega_0^n$$

where $\omega_{\varphi} = \omega_0 + \sqrt{-1}\partial\bar{\partial}\varphi$ and $\varphi \in PSH(X,\omega_0)$ has full Monge-Ampère mass. (For precise definitions, we refer [BBEGZ19].)

Remark 3.2. If ω_{φ} is a singular Kähler–Einstein metric, then $\omega_{\varphi} \in C^{\infty}(X_{reg})$ and it satisfies

$$\operatorname{Ric}(\omega_{\varphi}) = \omega_{\varphi}$$
 on X_{reg} .

Li, Tian and Zhu [LTZ23, Theorem 1.2] proved a criterion for the existence of singular Kähler– Einstein metrics on Q-Fano group compactifications as a generalization of Delcroix's result [Del17] for smooth Fano group compactifications (see also [Del20]). Let X be a bi-equivariant compactification of a reductive group G, and let $\Phi = \Phi(G, T)$ be the root system of G with respect to a maximal torus T. Fix a set of positive roots Φ^+ , and denote by $2\rho = \sum_{\alpha \in \Phi^+} \alpha$ the sum of positive roots and by \mathcal{C}^+ the cone generated by positive roots in Φ^+ . Let Ξ be the relative interior of \mathcal{C}^+ . **Theorem 3.3** ([Del17, LTZ23]). For a Q-Fano compactification X of a reductive group G with the moment polytope $\Delta := \Delta(X, K_X^{-1})$, X admits a singular Kähler–Einstein metric if and only if

$$bar_{DH}(\Delta) \in 2\rho + \Xi,$$

where $bar_{DH}(\Delta)$ is the barycenter of the moment polytope with respect to the Duistermaat–Heckman measure $\prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 dp$ and $\kappa(\cdot, \cdot)$ is the Cartan–Killing inner product.

3.2. Greatest Ricci lower bounds of Fano manifolds. Let X be a Fano manifold, that is, a compact complex manifold whose anticanonical line bundle K_X^{-1} is ample. We define the greatest Ricci lower bound R(X) of X as

 $R(X) := \sup\{0 \le t \le 1: \text{ there exists a Kähler form } \omega \in c_1(X) \text{ with } \operatorname{Ric}(\omega) > t \omega\}.$

Theorem 3.4 (Székelyhidi [Szé11]). Let $\omega \in 2\pi c_1(X)$.

 $R(X) = \sup\{0 \le t \le 1: \text{ there exists a smooth solution } \varphi_t \text{ of } \omega_{\varphi_t}^n = e^{h - t\varphi} \omega^n\}.$

In other words, R(X) is the supremum of the existence of Aubin's continuity path:

$$\operatorname{Ric}(\omega_{\varphi_t}) = t \,\omega_{\varphi_t} + (1-t)\omega$$

Remark 3.5. R(X) is an invariant of the Fano manifold X in the sense that this is independent of choice of $\omega \in 2\pi c_1(X)$. This invariant was first studied by Tian [Tia92], and was explicitly defined by Rubinstein [Rub08, Rub09], where it was called Tian's β -invariant.

Remark 3.6. By definition, if X admits a Kähler–Einstein metric, then the greatest Ricci lower bound R(X) is equal to 1. If X does not admit a Kähler–Einstein metric, R(X) measures somehow a Fano manifold X fails to be Kähler–Einstein. However, R(X) = 1 does not guarantee X to be Kähler–Einstein. For example, Tian [Tia97] constructed the unstable deformation of the Mukai– Umemura 3-fold having R(X) = 1 (see [Szé11, Section 3]).

Theorem 3.7 (Li [Li17]). R(X) = 1 if and only if X is K-semistable.

The greatest Ricci lower bound R(X) of a Fano manifold X is closely related with Tian's α -invariant (or the global log canonical threshold). If X does not admit a Kähler–Einstein metric then by [Tia87, Theorem 2.1] we have a lower bound of R(X) in terms of the alpha invariant

$$R(X) \ge \alpha(X) \cdot \frac{\dim X + 1}{\dim X}$$

On the other hand, the greatest Ricci lower bound R(X) is also related with the δ -invariant $\delta(X, -K_X)$, defined by Fujita and Odaka [FO18] using log canonical thresholds of basis type divisors. In fact, for a K-unstable Fano manifold X, the greatest Ricci lower bound R(X) is equal to the delta invariant $\delta(X, -K_X)$. More precisely, we have

Theorem 3.8 ([BBJ21, CRZ19]). $R(X) = \min\{1, \delta(X, -K_X)\}.$

For any smooth Fano equivariant compactification of complex Lie groups, Delcroix [Del17] obtained an explicit formula for the greatest Ricci lower bound in terms of the moment polytope as follows. Note that this is a generalization of Li's results for toric manifolds [Li11], since the root system of this case is trivial so that the cone generated by positive roots is restricted to the origin, the Duistermaat–Heckman measure is the Lebesgue measure, and $2\rho = 0$.

Theorem 3.9 (Delcroix [Del17]). For a smooth Fano group compactification X of G,

$$R(X) = \sup\left\{t \in (0,1) : 2\rho + \frac{t}{1-t}(2\rho - \boldsymbol{bar}_{DH}(\Delta)) \in \operatorname{Relint}(\Delta - \mathcal{C}^+)\right\},\$$

where $\operatorname{Relint}(\Delta - C^+)$ means the relative interior of the Minkovski sum of the moment polytope Δ and the negative $-C^+$ of the cone generated by positive roots in Φ^+ . We immediately get an elementary geometric expression for R(X) by Proposition 3.9 as follows:

Corollary 3.10. Let A be the point corresponding to 2ρ in $\mathfrak{X}(T)$ and C be the barycenter $bar_{DH}(\Delta)$ of the moment polytope $\Delta(X, K_X^{-1})$ with respect to the Duistermaat–Heckman measure. If Q is the point at which the half-line starting from the barycenter C in the direction of A intersects the

boundary of
$$\Delta - \mathcal{C}^+$$
, then we have $R(X) = \frac{|AQ|}{|\overrightarrow{CQ}|}$

4. GORENSTEIN FANO GROUP COMPACTIFICATIONS WITH RANK TWO

In this section, we give a classification of Gorenstein Fano bi-equivariant compactifications of semisimple complex Lie groups with rank two, and check their K-stability as follows:

First, we classify all possible Gorenstein Fano bi-equivariant compactifications of each semisimple complex Lie group G using Corollary 2.6. In [Ruz12, Section 2.4], Ruzzi classified the locally factorial Fano symmetric varieties of rank two in terms of colored fans. As bi-equivariant group caompactifications are characterized by their moment polytopes, we can give a more explicit description of Gorenstein Fano group compactifications via moment polytopes. Second, for each compactification, we use the following criterion due to Alexeev–Katzarkov to check the singularity of the corresponding projective varieties. Recall that a polytope $\Delta \subset \mathcal{M} \otimes \mathbb{R}$ is called *Delzant* if the integral generators of the edges at each vertex form a basis of the lattice \mathcal{M} .

Proposition 4.1 (Proposition 2.5 of [AK05]). Let X be a polarized bi-equivariant compactification of a reductive group G and Δ^{toric} be its toric polytope formed by the Weyl group action from the moment polytope Δ , that is, $\Delta^{toric} = \bigcup_{w \in W} w\Delta$ for the Weyl group W of G.

- (1) If X is smooth, then Δ^{toric} satisfies the Delzant condition.
- (2) If Δ^{toric} is Delzant and no vertex of Δ^{toric} lies in a Weyl wall, then X is smooth.

Lastly, we determine which of them admit (singular) Kähler–Einstein metrics by computing the barycenter of each moment polytope with respect to the Duistermaat–Heckman measure (Theorem 3.3). In the case of K-unstable smooth Fano manifolds, we compute the greatest Ricci lower bounds applying Theorem 3.9 and Corollary 3.10.

Before we begin the above procedure, we first recall the classification of semisimple complex Lie groups G with rank two. Note that a connected semisimple Lie group G is determined uniquely up to an isomorphism by its Dynkin diagram and the character lattice $\mathfrak{X}(T)$ of a maximal torus $T \subset G$ (see e.g. [OV90, Theorem 4.3.9 and Theorem 4.3.10]).

Proposition 4.2. Let $Q \subset P$ be the root and weight lattice for a reduced root system Φ . For a lattice L such that $Q \subset L \subset P$, there exist a connected semisimple Lie group G, its maximal torus T and a root system isomorphism $\Phi_G \to \Phi$ mapping $\mathfrak{X}(T)$ to L.

As the group L/Q is isomorphic to the center Z(G) of the semisimple Lie group G corresponding to L by [OV90, Theorem 4.3.7], the classification of connected semisimple complex Lie groups can be given in terms of subgroups of the fundamental group P/Q of a reduced root system Φ . The fundamental groups of classical complex Lie groups are well-known (see [OV90, Chapter 1, § 3]):

Lemma 4.3. The complex Lie goups $SL_n(\mathbb{C})$ and $Sp_{2n}(\mathbb{C})$ are simply-connected, but $\pi_1(SO_n(\mathbb{C})) = \mathbb{Z}_2$ for $n \geq 3$.

As a result, we obtain a classification of semisimple complex Lie groups G with rank two as shown in Table 1.

Now, we start the classification of Gorenstein Fano bi-equivariant compactifications of each semisimple complex Lie group with rank two. Because of its complexity, we will deal with the cases of $A_1 \times A_1$ type in the last part of this paper. We start with the cases of A_2 type.

Lie type	Lie group G
$A_1 \times A_1$	$\operatorname{SL}_2(\mathbb{C}) \times \operatorname{SL}_2(\mathbb{C}), \operatorname{SL}_2(\mathbb{C}) \times \operatorname{PSL}_2(\mathbb{C}), \operatorname{PSL}_2(\mathbb{C}) \times \operatorname{PSL}_2(\mathbb{C}), \operatorname{SO}_4(\mathbb{C})$
A ₂	$\mathrm{SL}_3(\mathbb{C}),\mathrm{PSL}_3(\mathbb{C})$
$B_2=C_2$	$\operatorname{Sp}_4(\mathbb{C}) \cong \operatorname{Spin}_5(\mathbb{C}), \operatorname{SO}_5(\mathbb{C})$
G ₂	the simply-connected complex Lie group G_2

TABLE 1. Classification of semisimple complex Lie groups with rank two.

4.1. A₂-type.

4.1.1. Gorenstein Fano group compactifications of $SL_3(\mathbb{C})$. As $SL_3(\mathbb{C})$ is simply-connected, the spherical weight lattice \mathcal{M} is spanned by the fundamental weights $\varpi_1 = (1,0)$ and $\varpi_2 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$, and its dual lattice \mathcal{N} is spanned by the coroots $\alpha_1^{\vee} = \left(1, -\frac{\sqrt{3}}{3}\right)$ and $\alpha_2^{\vee} = \left(0, \frac{2\sqrt{3}}{3}\right)$. The Weyl walls are given by $W_1 := \{y = 0\}$ and $W_2 := \{y = \sqrt{3}x\}$. The sum of positive roots is $2\rho = 2\varpi_1 + 2\varpi_1 = (3, \sqrt{3})$. Choosing a realization of the root system A_2 in the Euclidean plane \mathbb{R}^2 with $\varpi_1 = (1, 0)$ and $\varpi_2 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$, for p = (x, y) we obtain the Duistermaat–Heckman measure

$$\prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp = \left(\frac{3}{2}x - \frac{\sqrt{3}}{2}y\right)^2 \left(\frac{3}{2}x + \frac{\sqrt{3}}{2}y\right)^2 (\sqrt{3}y)^2 \, dxdy$$

because A₂ has 3 positive roots: $\Phi^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2\} = \left\{ \left(\frac{3}{2}, -\frac{\sqrt{3}}{2}\right), (0, \sqrt{3}), \left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right) \right\}.$

Theorem 4.4. There are five Gorenstein Fano equivariant compactifications of $SL_3(\mathbb{C})$ up to isomorphism: three smooth compactifications and two singular compactifications. Their moment polytopes are given in the following Table 2 and Figure 1. Among them only one smooth Fano compactification of $SL_3(\mathbb{C})$ and a unique singular one admit singular Kähler–Einstein metrics.

X	$\Delta(K_X^{-1})$	Case	Edges (except Weyl walls)	Vertices	Smooth?	KE
$\overline{\mathrm{SL}}_3(1)$	Δ_1	II	$x + \frac{1}{\sqrt{3}}y = 5$	$\left(\frac{5}{2},\frac{5}{2}\sqrt{3}\right),(5,0)$	smooth	Yes
$\overline{\mathrm{SL}}_3(2)$	Δ_2	I.2	$x + \sqrt{3}y = 7, \ x + \frac{1}{\sqrt{3}}y = 5$	$\left(\frac{7}{4},\frac{7}{4}\sqrt{3}\right),(4,\sqrt{3}),(5,0)$	smooth	No
$\overline{\mathrm{SL}}_3(3)$	Δ_3	I.3.2	$x + \sqrt{3}y = 7, \ x + \frac{1}{\sqrt{3}}y = 5,$ $2x + \frac{4}{\sqrt{3}}y = 11$	$ \begin{pmatrix} \frac{7}{4}, \frac{7}{4}\sqrt{3} \\ (5,0) \end{pmatrix}, \begin{pmatrix} \frac{5}{2}, \frac{3}{2}\sqrt{3} \\ (5,0) \end{pmatrix}, \begin{pmatrix} \frac{9}{2}, \frac{\sqrt{3}}{2} \\ (5,0) \end{pmatrix}, $	smooth	No
$\overline{\mathrm{SL}}_3(4)$	Δ_4	I.1	$x + \sqrt{3}y = 7$	$\left(\frac{7}{4},\frac{7}{4}\sqrt{3}\right),(7,0)$	singular	Yes
$\overline{\mathrm{SL}}_3(5)$	Δ_5	I.3.1	$x + \sqrt{3}y = 7, \ 3x + \frac{5}{\sqrt{3}}y = 15$	$\left(\frac{7}{4}, \frac{7}{4}\sqrt{3}\right), \left(\frac{5}{2}, \frac{3}{2}\sqrt{3}\right), (5, 0)$	singular	No

TABLE 2. Gorenstein Fano equivariant compactifications of $SL_3(\mathbb{C})$.

Proof. Using Corollary 2.6 (cf. [LTZ23, Lemma 3.1]), we construct all possible Gorenstein Fano biequivariant compactifications of $SL_3(\mathbb{C})$. Let F_i be the facet of the moment polytope with primitive outer normal vector $m_i \alpha_1^{\vee} + n_i \alpha_2^{\vee} = \left(m_i, \frac{-m_i + 2n_i}{3}\sqrt{3}\right) \in \mathcal{N}$. Then F_i can be written as

$$F_i = \Big\{ (x,y) \in \mathcal{M} \otimes \mathbb{R} : m_i x + \frac{-m_i + 2n_i}{\sqrt{3}} y = 2m_i + 2n_i + 1 \Big\},\$$

and $0 \leq \frac{-m_i+2n_i}{3m_i} \leq 1$ by the convexity of the moment polytope. We start with the edge F_1 , the facet which intersects the Weyl wall W_2 . We divide this into two cases: **Case-I.** F_1 is orthogonal to W_2 ; **Case-II.** F_1 is not orthogonal to W_2 .

• Case-I: F_1 is orthogonal to W_2 .

In this case, $F_1 \perp W_2$ implies that $\frac{-m_1+2n_1}{3m_1} = 1$ so that $(m_1, n_1) = (1, 2)$. Thus

$$F_1 = \{x + \sqrt{3}y = 7\},\$$

and $F_1 \cap W_2 = \left(\frac{7}{4}, \frac{7}{4}\sqrt{3}\right)$. Then the first vertex point $A_1 := F_1 \cap F_2 \in \mathcal{M}$ is given by

$$A_1 = \left(7 - \frac{3}{2} \cdot \frac{5m_2 - 2n_2 - 1}{2m_2 - n_2}, \frac{\sqrt{3}}{2} \cdot \frac{5m_2 - 2n_2 - 1}{2m_2 - n_2}\right)$$

The convexity and primitive conditions of the moment polytope imply that

$$A_1 = 7\varpi_1 \text{ or } 3\varpi_1 + 2\varpi_2 \text{ or } \varpi_1 + 3\varpi_2.$$

- Case-I.1: $A_1 = 7\varpi_1 = (7,0)$ In this case, F_1 intersects the Weyl wall W_1 so that the corresponding moment polytope is Δ_4 (Figure 1(4)).
- Case-I.2: $A_1 = 3\varpi_1 + 2\varpi_2 = (4, \sqrt{3})$ implies that $(m_2, n_2) = (1, 1)$ so that we have

$$F_2 = \left\{ x + \frac{\sqrt{3}}{3}y = 5 \right\},$$

and it must intersect with W_1 at the vertex $A_2 = F_2 \cap W_1 = (5, 0)$. The corresponding moment polytope is Δ_2 (Figure 1(2)).

- **Case-I.3:** $A_1 = \varpi_1 + 3\varpi_2 = \left(\frac{5}{2}, \frac{3}{2}\sqrt{3}\right)$ implies that $m_2 + 1 = n_2$ so that we have

$$F_2 = \left\{ m_2 x + \frac{m_2 + 2}{\sqrt{3}}y = 4m_2 + 3 \right\}.$$

By convexity and primitive condition, $A_2 = 5\varpi_1$ or $A_2 = 4\varpi_1 + \varpi_2$.

- * **Case-I.3.1:** $A_2 = 5\varpi_1 \Rightarrow (m_2, n_2) = (3, 4)$ so that $F_2 = \left\{ 3x + \frac{5}{\sqrt{3}}y = 15 \right\}$. In this case, F_2 intersects W_1 at A_2 . The moment polytope is Δ_5 (Figure 1(5)).
- * **Case-I.3.2:** $A_2 = 4\varpi_1 + \varpi_2 \Rightarrow (m_2, n_2) = (2, 3)$ so that $F_2 = \left\{2x + \frac{4}{\sqrt{3}}y = 11\right\}$. In this case, $F_3 = \left\{x + \frac{\sqrt{3}}{3}y = 5\right\}$ intersects W_1 at the vertex $A_3 = 5\varpi_1$. The corresponding moment polytope is Δ_3 (Figure 1(3)).
- Case-II: F_1 is not orthogonal to W_2 .

In this case, $F_1 \cap W_2$ is a vertex of the polytope so that $A_1 := F_1 \cap W_2 \in \mathcal{M}$. This implies that $A_1 = \left(\frac{2m_1+2n_1+1}{2n_1}, \frac{2m_1+2n_1+1}{2n_1}\sqrt{3}\right)$ and $0 \leq \frac{-m_1+2n_1}{3m_1} < 1$ by the convexity of the polytope. Therefore, $A_1 = 5\varpi_2 = \left(\frac{5}{2}, \frac{5}{2}\sqrt{3}\right)$ and $2m_1 + 1 = 3n_1$ so that

$$F_1 = \left\{ (3n_1 - 1)x + \frac{n_1 + 1}{\sqrt{3}}y = 10n_1 \right\}.$$

Then the second vertex $A_2 = 5\omega_1$ or $3\omega_1 + \omega_2$ or $2\omega_1 + 3\omega_2$ or $\omega_1 + 4\omega_2$.

Removing duplications (up to isomorphisms), one can check that we have only one case $A_2 = 5\omega_1$. The corresponding moment polytope is Δ_1 (Figure 1(1)).

(1) Case-II. Let $\overline{\mathrm{SL}_3}(1)$ be the equivariant compactification of $\mathrm{SL}_3(\mathbb{C})$ whose moment polytope Δ_1 is the convex hull of three points 0, $5\varpi_1$, $5\varpi_2$ in $\mathcal{M} \otimes \mathbb{R}$. Then $\overline{\mathrm{SL}_3}(1)$ is a smooth projective symmetric variety with Picard number one studied by Ruzzi [Ruz10]. Using the above formula for

10

the Duistermaat–Heckman measure, we get the barycenter of Δ_1 with respect to the Duistermaat–Heckman measure

$$\mathbf{bar}_{DH}(\Delta_1) = (\bar{x}, \bar{y}) = \frac{1}{\mathrm{Vol}_{DH}(\Delta_1)} \left(\int_{\Delta_1} x \prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp, \int_{\Delta_1} y \prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp \right)$$
$$= \left(\frac{10}{3}, \frac{10}{9}\sqrt{3} \right) = \frac{10}{9} \times 2\rho.$$

Since $2\rho = (3,\sqrt{3})$ and the cone \mathcal{C}^+ is generated by the vectors $\left(\frac{3}{2}, -\frac{\sqrt{3}}{2}\right)$ and $(0,\sqrt{3})$, the barycenter **bar**_{DH}(Δ_1) is in the relative interior of the translated cone $2\rho + \mathcal{C}^+$. Therefore, $\overline{\mathrm{SL}}_3(1)$ admits a Kähler–Einstein metric by Theorem 3.3.

(2) Case-I.2. Let $\overline{\operatorname{SL}}_3(2)$ be the equivariant compactification of $\operatorname{SL}_3(\mathbb{C})$ whose moment polytope Δ_2 is the convex hull of three points 0, $5\varpi_1$, $3\varpi_1 + 2\varpi_2$, $\frac{7}{2}\varpi_2$ in $\mathcal{M} \otimes \mathbb{R}$. Recall from [Bri94] that for smooth projective spherical varieties of rank two, the blow-ups can be described and every birational equivariant morphism is a composition of blow-ups along smooth centers. As a result, $\overline{\operatorname{SL}}_3(2)$ is the blow-up of $\overline{\operatorname{SL}}_3(1)$ along the closed orbit corresponding to a vertex A_1 in Figure 1(1). We know that

$$\mathbf{bar}_{DH}(\Delta_2) = \left(\frac{156038947}{45427872}, \frac{16309243}{19469088}\sqrt{3}\right) \approx (3.435, 0.838 \times \sqrt{3}).$$

Since $\mathbf{bar}_{DH}(\Delta_2)$ is not in the relative interior of the translated cone $2\rho + \mathcal{C}^+$, $\overline{\mathrm{SL}}_3(2)$ does not admit any Kähler–Einstein metric by Theorem 3.3. Let Q be the point at which the half-line starting from the barycenter $C = \mathbf{bar}_{DH}(\Delta_2)$ in the direction of $A = (3, \sqrt{3})$ intersects the boundary of $(\Delta_2 - \mathcal{C}^+)$. Considering the line $x + \sqrt{3}y = 7$ giving a part of $\partial(\Delta_2 - \mathcal{C}^+)$ and the half-line \overrightarrow{CA} , we can compute

$$Q = \left(-\frac{12664579}{2363584}, \frac{29209667}{7090752}\right) \approx (-5.358, 4.119).$$

By Corollary 3.10, the greatest Ricci lower bound of $\overline{\text{SL}}_3(2)$ is

$$R(\overline{\mathrm{SL}}_3(2)) = \frac{AQ}{\overline{CQ}} = \frac{1419621}{1493483} \approx 0.9505.$$

(3) Case-I.3.2. As before, the variety $\overline{\mathrm{SL}}_3(3)$ is the blow-up of the smooth Fano compactification $\overline{\mathrm{SL}}_3(2)$ along the closed orbit corresponding to a vertex A_1 in Figure 1(2). Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_3) = \left(\frac{2234103775}{675213408}, \frac{527459083}{675213408}\sqrt{3}\right) \approx (3.309, 0.781 \times \sqrt{3})$$

of the moment polytope Δ_3 with respect to the Duistermaat-Heckman measure is not in the relative interior of the translated cone $2\rho + C^+$, $\overline{\mathrm{SL}}_3(3)$ does not admit any Kähler-Einstein metric by Theorem 3.3. Let Q be the point at which the half-line starting from the barycenter $C = \mathbf{bar}_{DH}(\Delta_3)$ in the direction of $A = (3, \sqrt{3})$ intersects the boundary of $(\Delta_3 - C^+)$. Considering the line $x + \sqrt{3}y = 7$ giving a part of $\partial(\Delta_3 - C^+)$ and the half-line \overrightarrow{CA} , we can compute

$$Q = \left(\frac{495934721}{234799424}, \frac{382553749}{234799424}\right) \approx (2.112, 1.629).$$

By Corollary 3.10, the greatest Ricci lower bound of $\overline{SL_3}(3)$ is

$$R(\overline{\mathrm{SL}}_{3}(3)) = \frac{\overline{AQ}}{\overline{CQ}} = \frac{21100419}{28437901} \approx 0.74198.$$

(4) Case-I.1. Let $\overline{\mathrm{SL}}_3(4)$ be the equivariant compactification of $\mathrm{SL}_3(\mathbb{C})$ whose moment polytope Δ_4 is the convex hull of three points 0, $7\varpi_1$, $\frac{7}{2}\varpi_2$ in $\mathcal{M} \otimes \mathbb{R}$. As the toric polytope formed by the

Weyl group action from Δ_4 is not a Delzant polytope, $\overline{SL_3}(4)$ is singular by Proposition 4.1. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_4) = \left(\frac{11183}{288}, \frac{203}{288}\sqrt{3}\right) \approx (4.108, 0.705 \times \sqrt{3})$$

of the moment polytope Δ_4 with respect to the Duistermaat–Heckman measure is in the relative interior of the translated cone $2\rho + C^+$, $\overline{SL_3}(4)$ admits a singular Kähler–Einstein metric by Theorem 3.3.

(5) Case-I.3.1. As the toric polytope formed by the Weyl group action from the moment polytope Δ_5 is not a Delzant polytope, $\overline{SL}_3(5)$ is singular by Proposition 4.1. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_5) = \left(\frac{1580795359}{507050784}, \frac{402732299}{507050784}\sqrt{3}\right) \approx (3.118, 0.794 \times \sqrt{3})$$

of the moment polytope Δ_5 with respect to the Duistermaat–Heckman measure is not in the relative interior of the translated cone $2\rho + C^+$, $\overline{SL}_3(5)$ does not admit any singuar Kähler–Einstein metric by Theorem 3.3.



FIGURE 1. Moment polytopes of Gorenstein Fano compactifications of $SL_3(\mathbb{C})$.

4.1.2. Gorenstein Fano group compactification of $PSL_3(\mathbb{C})$. As $PSL_3(\mathbb{C})$ is of adjoint type, the spherical weight lattice \mathcal{M} is spanned by the simple roots $\alpha_1 = \left(\frac{3}{2}, -\frac{\sqrt{3}}{2}\right)$ and $\alpha_2 = (0, \sqrt{3})$, and its dual lattice \mathcal{N} is spanned by the fundamental coweights $\varpi_1^{\vee} = \begin{pmatrix} 2\\ 3\\ \end{pmatrix}$ and $\varpi_2^{\vee} = \begin{pmatrix} \frac{1}{3}, \frac{\sqrt{3}}{3} \end{pmatrix}$. The Weyl walls are given by $W_1 = \{y = 0\}$ and $W_2 = \{y = \sqrt{3}x\}$. The sum of positive roots is $2\rho = (3, \sqrt{3})$.

Theorem 4.5. There are three Gorenstein Fano equivariant compactifications of $PSL_3(\mathbb{C})$. Their moment polytopes are given in the following Table 3 and Figure 2. They all are smooth and admit Kähler–Einstein metrics.

X	$\Delta(K_X^{-1})$	Case	Edges (except Weyl walls)	Vertices	Smoothness	KE
$\overline{\mathrm{PSL}_3}(1)$	Δ_1	I.3	$x + \sqrt{3}y = 9$	$\left(\frac{9}{4},\frac{9}{4}\sqrt{3}\right),(9,0)$	smooth	Yes
$\overline{\mathrm{PSL}_3}(2)$	Δ_2	I.2	$x + \sqrt{3}y = 9, \ x = \frac{9}{2}$	$\left(\frac{9}{4},\frac{9}{4}\sqrt{3}\right), \left(\frac{9}{2},\frac{3}{2}\sqrt{3}\right), \left(\frac{9}{2},0\right)$	smooth	Yes
$\overline{\mathrm{PSL}}_3(3)$	Δ_3	I.1	$ x + \sqrt{3}y = 9, \ x = \frac{9}{2}, 3x + \sqrt{3}y = 15 $	$ \begin{pmatrix} \frac{9}{4}, \frac{9}{4}\sqrt{3} \\ (3, 2\sqrt{3}) \end{pmatrix}, \begin{pmatrix} \frac{9}{2}, \frac{\sqrt{3}}{2} \\ \frac{9}{2}, 0 \end{pmatrix}, $	smooth	Yes

TABLE 3. Gorenstein Fano equivariant compactifications of $PSL_3(\mathbb{C})$.

Proof. Let $m_i \varpi_1^{\vee} + n_i \varpi_2^{\vee} = \left(\frac{2m_i + n_i}{3}, \frac{n_i}{3}\sqrt{3}\right) \in \mathcal{N}$ be the primitive outer normal vector of the facet $F_i = \{(2m_i + n_i)x + n_i\sqrt{3}y = 3 + 6m_i + 6n_i\},\$

where $m_i \ge 0, n_i \ge 0$ by the convexity of the polytope. Let F_1 be the facet which intersects the Weyl wall W_2 .

• Case-I: F_1 is orthogonal to W_2 .

$$F_1 \perp W_2 \Rightarrow (m_1, n_1) = (0, 1) \Rightarrow F_1 = \{x + \sqrt{3}y = 9\}, F_1 \cap W_2 = \left(\frac{9}{4}, \frac{9}{4}\sqrt{3}\right).$$

Then the vertex point $A_1 := F_1 \cap F_2 \in \mathcal{M}$ is given by

$$A_1 = \left(3 + \frac{3}{2} \left(\frac{1 - n_2}{m_2}\right), \left\{2 - \frac{1}{2} \left(\frac{1 - n_2}{m_2}\right)\right\} \sqrt{3}\right).$$

The convexity and primitive conditions imply that $A_1 = 2\alpha_1 + 3\alpha_2$ or $3\alpha_1 + 3\alpha_2$ or $4\alpha_1 + 3\alpha_2$ or $5\alpha_1 + 3\alpha_2$ or $6\alpha_1 + 3\alpha_2$. But if $A_1 = 4\alpha_1 + 3\alpha_2$ or $5\alpha_1 + 3\alpha_2$, there is no lattice polytope.

- Case-I.1: $A_1 = 2\alpha_1 + 3\alpha_2 = (3, 2\sqrt{3}) \Rightarrow n_2 = 1 \Rightarrow F_2 = \{(2m_2 + 1)x + \sqrt{3}y = 9 + 6m_2\}.$ By convexity and primitive condition, $A_2 = 3\alpha_1 + 2\alpha_2$ or $4\alpha_1 + 2\alpha_2$. If $A_2 = 4\alpha_1 + 2\alpha_2 = (6,0) \Rightarrow 6m_2 = 3$ so that this is not Gorenstein.

- If $A_2 = 3\alpha_1 + 2\alpha_2 = (\frac{9}{2}, \frac{\sqrt{3}}{2}) \Rightarrow (m_2, n_2) = (1, 1)$ so that $F_2 = \{3x + \sqrt{3}y = 15\}$. In this case, $(m_3, n_3) = (1, 0)$ so that $F_3 = \{x = \frac{9}{2}\} \perp W_1$ (Δ_3 , Figure 2(3)).
- Case-I.2: $A_1 = 3\alpha_1 + 3\alpha_2 = \left(\frac{9}{2}, \frac{3}{2}\sqrt{3}\right) \Rightarrow \frac{1-n_2}{m_2} = 1 \Rightarrow (m_2, n_2) = (1, 0)$ so that $F_2 = \left\{ x = \frac{9}{2} \right\}$ and $F_2 \perp W_1$ (Δ_2 , Figure 2(2)).

- Case-I.3:
$$A_1 = 6\alpha_1 + 3\alpha_2 = 9\varpi_1 (\Delta_1, \text{ Figure 2(1)}).$$

• Case-II: F_1 is not orthogonal to W_2 . $F_1 \cap W_2 =: A_1 = (\frac{2m_1 + 2n_1 + 1}{m_1 + 2n_1} \cdot \frac{3}{2}, \frac{2m_1 + 2n_1 + 1}{m_1 + 2n_1} \cdot \frac{3}{2}\sqrt{3}) \in \mathcal{M} \Rightarrow \frac{2m_1 + 2n_1 + 1}{m_1 + 2n_1} = 1 + \frac{m_1 + 1}{m_1 + 2n_1} \Rightarrow (m_1, n_1) = (1, 0).$ This is isomorphic to Case-I.3.

(1) Case-I.3. By [Ruz10, Theorem 5], the variety $\overline{\text{PSL}_3}(1)$ is isomorphic to $\mathbb{P}(\text{Mat}_{3\times 3}(\mathbb{C})) = \mathbb{P}^8$ as $PSL_3(\mathbb{C}) \times PSL_3(\mathbb{C})$ -variety. This has three orbits and the closed orbit consists of rank one 3×3 matrices which is isomorphic to $\mathbb{P}^2 \times \mathbb{P}^2$. As $\overline{PSL_3}(1)$ is a homogeneous variety, it naturally admits a Kähler–Einstein metric (see [Mat72, Section 5]).

(2) Case-I.2. The variety $\overline{\text{PSL}_3}(2)$ is the blow-up of the smooth Fano compactification $\overline{\text{PSL}_3}(1)$ along the closed orbit corresponding to a vertex A_1 in Figure 2(1). Indeed, $\overline{\text{PSL}_3}(2)$ is the wonderful compactification of $\text{PSL}_3(\mathbb{C})$. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_2) = \left(\frac{24641}{6592}, \frac{24641}{19776}\sqrt{3}\right) = \frac{24641}{19776} \times 2\rho \approx (3.738, 1.246 \times \sqrt{3})$$

of the moment polytope Δ_2 with respect to the Duistermaat–Heckman measure is in the relative interior of the translated cone $2\rho + C^+$, $\overline{PSL_3}(2)$ admit a Kähler–Einstein metric by Theorem 3.3.

(3) Case-I.1. The variety $\overline{\text{PSL}_3}(3)$ is the blow-up of $\overline{\text{PSL}_3}(2)$ along the closed orbit corresponding to a vertex A_1 in Figure 2(2). Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_3) = \left(\frac{189565091}{57005952}, \frac{189565091}{171017856}\sqrt{3}\right) = \frac{189565091}{171017856} \times 2\rho \approx (3.325, 1.108 \times \sqrt{3})$$

of the moment polytope Δ_3 with respect to the Duistermaat–Heckman measure is in the relative interior of the translated cone $2\rho + C^+$, $\overline{PSL_3}(3)$ admit a Kähler–Einstein metric by Theorem 3.3. \Box



FIGURE 2. Moment polytopes of Gorenstein Fano compactifications of $PSL_3(\mathbb{C})$.

4.2. B₂-type.

4.2.1. Gorenstein Fano group compactifications of $\operatorname{Sp}_4(\mathbb{C}) \cong \operatorname{Spin}_5(\mathbb{C})$. As $\operatorname{Sp}_4(\mathbb{C})$ is simply-connected, the spherical weight lattice \mathcal{M} is spanned by $\varpi_1 = \left(\frac{1}{2}, \frac{1}{2}\right)$ and $\varpi_2 = (0, 1)$, and its dual lattice \mathcal{N} is spanned by $\alpha_1^{\vee} = (2, 0)$ and $\alpha_2^{\vee} = (-1, 1)$. The Weyl walls are given by $W_1 = \{y = x\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (1, 3)$. Choosing a realization of the root system C_2 in the Euclidean plane \mathbb{R}^2 with $\alpha_1 = (1, 0)$ and $\alpha_2 = (-1, 1)$, for p = (x, y) we obtain the Duistermaat–Heckman measure

$$\prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp = x^2 (-x+y)^2 y^2 (x+y)^2 \, dx dy$$

because C_2 has 4 positive roots: $\Phi^+ = \{\alpha_1, \alpha_2, \alpha_1 + \alpha_2, 2\alpha_1 + \alpha_2\} = \{(1, 0), (-1, 1), (0, 1), (1, 1)\}.$

Theorem 4.6. There are four Gorenstein Fano equivariant compactifications of $\text{Sp}_4(\mathbb{C})$ up to isomorphism: three smooth compactifications and one singular compactification. Their moment polytopes are given in the following Table 4 and Figure 3. Among them only two smooth Fano compactifications admit Kähler–Einstein metrics.

X	$\Delta(K_X^{-1})$	Case	Edges (except Weyl walls)	Vertices	Smoothness	KE
$\overline{\mathrm{Sp}_4}(1)$	Δ_1	I.3	$y = \frac{7}{2}$	$\left(0,\frac{7}{2}\right), \left(\frac{7}{2},\frac{7}{2}\right)$	smooth	Yes
$\overline{\mathrm{Sp}_4}(2)$	Δ_2	I.2	$y = \frac{7}{2}, \ x + y = 5$	$\left(0,\frac{7}{2}\right), \left(\frac{3}{2},\frac{7}{2}\right), \left(\frac{5}{2},\frac{5}{2}\right)$	smooth	Yes
$\overline{\mathrm{Sp}_4}(3)$	Δ_3	I.1.2	$y = \frac{7}{2}, x + y = 5,$ x + 3y = 11	$(0, \frac{7}{2}), (\frac{1}{2}, \frac{7}{2}), (2, 3), (\frac{5}{2}, \frac{5}{2})$	smooth	No
$\overline{\mathrm{Sp}_4}(4)$	Δ_4	I.1.1	$y = \frac{7}{2}, \ 2x + 4y = 15$	$\left(0,\frac{7}{2}\right), \left(\frac{1}{2},\frac{7}{2}\right), \left(\frac{5}{2},\frac{5}{2}\right)$	singular	No

TABLE 4. Gorenstein Fano equivariant compactifications of $\text{Sp}_4(\mathbb{C})$.

Proof. Let $m_i \alpha_1^{\vee} + n_i \alpha_2^{\vee} = (2m_i - n_i, n_i) \in \mathcal{N}$ be the primitive outer normal vector of the facet $F_i = \{(2m_i - n_i)x + n_iy = 2m_i + 2n_i + 1\},$

where $n_i \ge 2m_i - n_i \ge 0$ by the convexity of the polytope.

Let F_1 be the facet which intersects the Weyl wall W_2 , then we have two cases:

- Case-I: F_1 is orthogonal to W_2 .
- $$\begin{split} F_1 \perp W_2 \Rightarrow (m_1, n_1) &= (1, 2) \Rightarrow F_1 = \left\{ y = \frac{7}{2} \right\}, \ F_1 \cap W_2 = \left(0, \frac{7}{2}\right). \ \text{Let } A_1 := F_1 \cap F_2. \\ A_1 &= \left(\frac{4m_2 3n_2 + 2}{4m_2 2n_2}, \frac{7}{2}\right) \in \mathcal{M} \Rightarrow A_1 = \varpi_1 + 3\varpi_2 \text{ or } 3\varpi_1 + 2\varpi_2 \text{ or } 5\varpi_1 + \varpi_2 \text{ or } 7\varpi_1. \\ \text{But if } A_1 &= 5\varpi_1 + 1\varpi_2 = \left(\frac{5}{2}, \frac{7}{2}\right), \text{ there is no lattice polytope.} \\ &- \mathbf{Case-I.1:} \ A_1 = \varpi_1 + 3\varpi_2 = \left(\frac{1}{2}, \frac{7}{2}\right) \Rightarrow m_2 + 1 = n_2, \ F_2 = \{(m_2 1)x + (m_2 + 1)y = 4m_2 + 3\}. \ \text{By convexity and primitive condition, } A_2 = 5\varpi_1 \text{ or } 4\varpi_1 + \varpi_2. \\ &* \mathbf{Case-I.1.1:} \ A_2 = 5\varpi_1 \Rightarrow (m_2, n_2) = (3, 4), \ F_2 = \{2x + 4y = 15\} \\ &(\Delta_4, \text{ Figure 3(4)}). \\ &* \mathbf{Case-I.1.2:} \ A_2 = 4\varpi_1 + \varpi_2 \Rightarrow (m_2, n_2) = (2, 3), \ F_2 = \{x + 3y = 11\} \\ &\text{In this case, } (m_3, n_3) = (1, 1) \text{ so that } F_3 = \{x + y = 5\} \perp W_1 \ (\Delta_3, \text{ Figure 3(3)}). \\ &- \mathbf{Case-I.2:} \ A_1 = 3\varpi_1 + 2\varpi_2 = \left(\frac{3}{2}, \frac{7}{2}\right) \Rightarrow (m_2, n_2) = (1, 1), \ F_2 = \{x + y = 5\} \\ &(\Delta_2, \text{ Figure 3(2)}). \\ &- \mathbf{Case-I.3:} \ A_1 = 7\varpi_1 = (\frac{7}{2}, \frac{7}{2}) \ (\Delta_1, \text{ Figure 3(1)}). \end{split}$$
 Case-II: F_1 is not orthogonal to W_2.
 F_1 \cap W_2 =: A_1 = (0, \frac{2m_1 + 2n_1 + 1}{n_1}) \in \mathcal{M} \Rightarrow 3 + \frac{1}{n_1} \leq \frac{2m_1 + 2n_1 + 1}{n_1} < 4 + \frac{1}{n_1} \Rightarrow \frac{2m_1 + 2n_1 + 1}{n_1} = 4 \\ \text{ In this case, there is no primitive vector.} \end{split}

(1) Case-I.3. The variety $\overline{\text{Sp}_4}(1)$ is isomorphic to the 10-dimensional Lagrangian Grassmannian Lag(4,8) parametrizing 4-dimensional isotropic subspaces in the 8-dimensional symplectic vector space (see [Ruz10, Theorem 5]). Since $\overline{\text{Sp}_4}(1)$ is homogeneous, it admits a Kähler–Einstein metric.

(2) Case-I.2. The variety $\text{Sp}_4(2)$ is the blow-up of the smooth Fano compactification $\text{Sp}_4(1)$ along the closed orbit corresponding to a vertex A_1 in Figure 3(1). Indeed, $\overline{\text{Sp}_4}(2)$ is the wonderful compactification of $\text{Sp}_4(\mathbb{C})$. Since $2\rho = (1,3)$ and the cone \mathcal{C}^+ is generated by the vectors (1,0) and (-1,1), the barycenter

$$\mathbf{bar}_{DH}(\Delta_2) = (\frac{135148980025}{104829824704}, \frac{5019760035}{1637966011}) \approx (1.289, 3.065)$$

of the moment polytope Δ_2 with respect to the Duistermaat–Heckman measure is in the relative interior of the translated cone $2\rho + C^+$. By Theorem 3.3, $\overline{\text{Sp}_4}(2)$ admits a Kähler–Einstein metric.

(3) Case-I.1.2. The variety $\text{Sp}_4(3)$ is the blow-up of the wonderful compactification $\text{Sp}_4(2)$ along the unique closed orbit corresponding to a vertex A_1 in Figure 3(2). Since the barycenter

 $\mathbf{bar}_{DH}(\Delta_3) = (\frac{27756440595}{22318407232}, \frac{3043253830}{1046175339}) \approx (1.244, 2.909)$

of the moment polytope Δ_3 with respect to the Duistermaat–Heckman measure is not in the relative interior of the translated cone $2\rho + C^+$, $\overline{\text{Sp}_4}(3)$ does not admit any Kähler–Einstein metric as already showed in [Del17, Example 5.4].

(4) Case-I.1.1. By Proposition 4.1, $\overline{\text{Sp}_4}(4)$ is singular. Since the barycenter

 $\mathbf{bar}_{DH}(\Delta_4) = \left(\frac{53741124025}{47717371328}, \frac{192699595}{67780357}\right) \approx (1.262, 2.843)$

of the moment polytope Δ_4 with respect to the Duistermaat–Heckman measure is not in the relative interior of the translated cone $2\rho + C^+$, $\overline{\text{Sp}_4}(4)$ does not admit any Kähler–Einstein metric.



FIGURE 3. Moment polytopes of Gorenstein Fano compactifications of $\text{Sp}_4(\mathbb{C})$.

4.2.2. Gorenstein Fano group compactification of $PSp_4(\mathbb{C}) \cong SO_5(\mathbb{C})$. As $PSp_4(\mathbb{C})$ is of adjoint type, the spherical weight lattice \mathcal{M} is spanned by $\alpha_1 = (1,0)$ and $\alpha_2 = (-1,1)$, and its dual lattice \mathcal{N} is spanned by $\varpi_1^{\vee} = (1,1)$ and $\varpi_2^{\vee} = (0,1)$. The Weyl walls are given by $W_1 = \{y = x\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (1,3)$.

Theorem 4.7. There are four Gorenstein Fano equivariant compactifications of $SO_5(\mathbb{C})$: two smooth compactifications and two singular compactifications. Their moment polytopes are given in the following Table 5 and Figure 4. They all admit singular Kähler–Einstein metrics.

X	$\Delta(K_X^{-1})$	Case	Edges (except Weyl walls)	Vertices	Smoothness	KE
$\overline{SO_5}(1)$	Δ_1	I.2	y = 4	(0,4),(4,4)	smooth	Yes
$\overline{SO_5}(2)$	Δ_2	I.1	$y = 4, \ x + y = 5$	$(0,4),(1,4),(\frac{5}{2},\frac{5}{2})$	smooth	Yes
$\overline{SO_5}(3)$	Δ_3	II.1	x + y = 5	$(0,5), (\frac{5}{2}, \frac{5}{2})$	singular	Yes
$\overline{SO_5}(4)$	Δ_4	II.2	$x + 2y = 8, \ x + y = 5$	$(0,4),(2,3),(\frac{5}{2},\frac{5}{2})$	singular	Yes

TABLE 5.	Gorenstein	Fano e	equivariant	compactifications	of	SO_5	(\mathbb{C})).
							`	

Proof. Let $m_i \varpi_1^{\vee} + n_i \varpi_2^{\vee} = (m_i, m_i + n_i) \in \mathcal{N}$ be the primitive outer normal vector of the facet $F_i = \{m_i x + (m_i + n_i)y = 4m_i + 3n_i + 1\},$

where $m_i \ge m_i + n_i \ge 0$ by the convexity of the polytope.

Let F_1 be the facet which intersects the Weyl wall W_2 , then we have two cases: • Case-I: $F_1 \perp W_2 \Rightarrow (m_1, n_1) = (0, 1)$ so that $F_1 = \{y = 4\}, F_1 \cap W_2 = (0, 4)$. Note that $F_1 \cap F_2 =: A_1 = \left(\frac{-n_2+1}{m_2}, 4\right) \in \mathcal{M}$. $\Rightarrow A_1 = 5\alpha_1 + 4\alpha_2 \text{ or } 6\alpha_1 + 4\alpha_2 \text{ or } 7\alpha_1 + 4\alpha_2 \text{ or } 8\alpha_1 + 4\alpha_2.$ If $A_1 = 6\alpha_1 + 4\alpha_2$ or $7\alpha_1 + 4\alpha_2$, there is no lattice polytope. - Case-I.1: $A_1 = 5\alpha_1 + 4\alpha_2 = (1, 4) \Rightarrow m_2 + n_2 = 1$ $\Rightarrow F_2 = \{m_2 x + y = m_2 + 4\}$ Convexity and primitive condition $\Rightarrow m_2 = 1$ so that $F_2 = \{x + y = 5\}$ $(\Delta_2, \text{ Figure 4(2)}).$ • Case-II: $F_1 \cap W_2 =: A_1 = \left(0, \frac{4m_1 + 3n_1 + 1}{m_1 + n_1}\right) \in \mathcal{M}.$ By convexity $m_1 + n_1 \ge m_1 > 0 \Rightarrow A_1 = (0, 5)$ or (0, 4). - Case-II.1: $A_1 = (0,5) \Rightarrow (m_1, n_1) = (1,0) \Rightarrow F_1 = \{x + y = 5\} \perp W_1$ $(\Delta_3, \text{ Figure 4(3)}).$ - Case-II.2: $A_1 = (0, 4) \Rightarrow n_1 = 1$ $\Rightarrow F_1 = \{m_1 x + (m_1 + 1)y = 4(m_1 + 1)\}\$ Convexity and primitive conditions \Rightarrow $(m_1, n_1) = (1, 1)$ so that $F_1 = \{x + 2y = 8\}$ $A_2 := F_1 \cap F_2 = 4\varpi_1 + 1\varpi_2 = (2,3)$ and $F_2 = \{x + y = 5\} \perp W_1$ (Δ_4 , Figure 4(4)).

(1) Case-I.2. The variety $\overline{SO_5}(1)$ is the 10-dimensional spinor variety $\mathbb{S}_5 \subset \mathbb{P}^{15}$ which is one component of the variety parametrizing 5-dimensional isotropic subspaces in the 10-dimensional orthogonal vector space (see [Ruz10, Theorem 5]). As $\overline{SO_5}(1)$ is a homogeneous variety, it naturally admits a Kähler–Einstein metric.

(2) Case-I.1. The variety $\overline{SO_5}(2)$ is the blow-up of the smooth Fano compactification $\overline{SO_5}(1)$ along the closed orbit corresponding to a vertex A_1 in Figure 4(1). Indeed, $\overline{SO_5}(2)$ is the wonderful compactification of $SO_5(\mathbb{C})$. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_2) = \left(\frac{6332682925}{5547479872}, \frac{18534175955}{5547479872}\right) \approx (1.415, 3.341)$$

of the moment polytope Δ_2 with respect to the Duistermaat-Heckman measure is in the relative interior of the translated cone $2\rho + C^+$, $\overline{SO_5}(2)$ admit a Kähler–Einstein metric by Theorem 3.3.

(3) Case-II.1. As the toric polytope Δ_3^{toric} formed by the Weyl group action from the moment polytope Δ_3 is not a Delzant polytope, $\overline{SO_5}(3)$ is singular. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_3) = \left(\frac{725}{704}, \frac{225}{64}\right) \approx (1.030, 3.516)$$

is in the relative interior of the cone $2\rho + C^+$, $\overline{SO_5}(3)$ admits a singular Kähler–Einstein metric. (4) Case-II.2. As the toric polytope Δ_4^{toric} formed by the Weyl group action from the moment polytope Δ_4 is not a Delzant polytope, $\overline{SO_5}(4)$ is singular. Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_4) = \left(\frac{1110073615}{943350848}, \frac{8612750675}{2830052544}\right) \approx (1.177, 3.043)$$

is in the relative interior of the cone $2\rho + \mathcal{C}^+$, $\overline{SO_5}(4)$ admits a singular Kähler–Einstein metric. \Box



FIGURE 4. Moment polytopes of Gorenstein Fano compactifications of $SO_5(\mathbb{C})$.

4.3. Gorenstein Fano group compactification of G_2 . As the root lattice and weight lattice of G_2 coincide, the simply-connected complex Lie group of type G_2 is of adjoint type. Thus, the spherical weight lattice \mathcal{M} is spanned by the fundamental weights $\varpi_1 = \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right)$ and $\varpi_2 = (0, \sqrt{3})$, and its dual lattice \mathcal{N} is spanned by $\alpha_1^{\vee} = (2, 0)$ and $\alpha_2^{\vee} = (-1, \frac{1}{\sqrt{3}})$. The Weyl walls are given by $W_1 = \{y = \sqrt{3}x\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (1, 3\sqrt{3})$. Note that the complex Lie group G_2 has 6 positive roots:

$$\Phi^{+} = \{\alpha_{1}, \alpha_{2}, \alpha_{1} + \alpha_{2}, 2\alpha_{1} + \alpha_{2}, 3\alpha_{1} + \alpha_{2}, 3\alpha_{1} + 2\alpha_{2}\} \\ = \{(1,0), \left(-\frac{3}{2}, \frac{\sqrt{3}}{2}\right), \left(-\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{1}{2}, \frac{\sqrt{3}}{2}\right), \left(\frac{3}{2}, \frac{\sqrt{3}}{2}\right), (0, \sqrt{3})\}.$$

Thus, the Duistermaat–Heckman measure is given as

$$\prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp = x^2 \Big(-\frac{3}{2}x + \frac{\sqrt{3}}{2}y \Big)^2 \Big(-\frac{1}{2}x + \frac{\sqrt{3}}{2}y \Big)^2 \Big(\frac{1}{2}x + \frac{\sqrt{3}}{2}y \Big)^2 \Big(\frac{3}{2}x + \frac{\sqrt{3}}{2}y \Big)^2 (\sqrt{3}y)^2 \, dxdy.$$

Theorem 4.8. There are only two Gorenstein Fano equivariant compactifications of G_2 . Their moment polytopes are given in the following Table 6 and Figure 5. They all are smooth and admit Kähler-Einstein metrics.

X	$\Delta(K_X^{-1})$	Case	Edges (except Weyl walls)	Vertices	Smoothness	KE
$\overline{G_2}(1)$	Δ_1	I.1	$y = \frac{7}{2}\sqrt{3}, \ x + \sqrt{3}y = 11$	$\left(0,\frac{7}{2}\sqrt{3}\right), \left(\frac{1}{2},\frac{7}{2}\sqrt{3}\right), \left(\frac{11}{4},\frac{11}{4}\sqrt{3}\right)$	smooth	Yes
$\overline{G_2}(2)$	Δ_2	I.2	$y = \frac{7}{2}\sqrt{3}$	$\left(\frac{7}{2},\frac{7}{2}\sqrt{3}\right)$	smooth	Yes

TABLE 6. Gorenstein Fano equivariant compactifications of G_2 .

Proof. Let $m_i \alpha_1^{\vee} + n_i \alpha_2^{\vee} = \left(2m_i - n_i, \frac{n_i}{\sqrt{3}}\right) \in \mathcal{N}$ be the primitive outer normal vector of the facet

$$F_i = \left\{ (2m_i - n_i)x + \frac{n_i}{\sqrt{3}}y = 2m_i + 2n_i + 1 \right\},\$$

where $\frac{n_i}{3} \ge 2m_i - n_i \ge 0$ by the convexity of the polytope. Let F_1 be the facet which intersects the Weyl wall W_2 , we have two cases:

- Case-I: $F_1 \perp W_2 \Rightarrow F_1 = \left\{ y = \frac{7}{2}\sqrt{3} \right\}$ $\Rightarrow A_1 = \varpi_1 + 3\varpi_2 \text{ or } 3\varpi_1 + 2\varpi_2 \text{ or } 5\varpi_1 + \varpi_2 \text{ or } 7\varpi_1$ If $A_1 = 3\varpi_1 + 2\varpi_2 \text{ or } 5\varpi_1 + \varpi_2$, there is no lattice polytope. - Case-I.1: $A_1 = \varpi_1 + 3\varpi_2 = \left(\frac{1}{2}, \frac{7}{2}\sqrt{3}\right) \Rightarrow (m_2, n_2) = (2, 3) \text{ and } F_2 = \{x + \sqrt{3}y = 11\}$ (Δ_1 , Figure 5(1)) - Case-I.2: $A_1 = 7\varpi_1 = \left(\frac{7}{2}, \frac{7}{2}\sqrt{3}\right)$ (Δ_2 , Figure 5(2))
- Case-II: F_1 is not orthogonal to W_2 . $F_1 \cap W_2 = \left(0, \left(\frac{2m_1+2n_1+1}{n_1}\right)\sqrt{3}\right) \in \mathcal{M} \Rightarrow \frac{4}{3} + \frac{1}{n_1} \ge \frac{2m_1+1}{n_1} > 1 + \frac{1}{n_1} \Rightarrow \text{ No lattice polytope}\right)$
- (1) Case-I.1. This variety $\overline{G_2}(1)$ is the wonderful compactification of G_2 . Computation shows

$$\operatorname{Vol}_{DH}(\Delta_1) = \int_0^{\frac{11\sqrt{3}}{4}} \int_0^{\frac{y}{\sqrt{3}}} \prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp + \int_{\frac{11\sqrt{3}}{4}}^{\frac{7\sqrt{3}}{2}} \int_0^{-\sqrt{3}y+11} \prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp = \frac{107945390367459}{8830976} \sqrt{3}.$$

This implies that

$$\mathbf{bar}_{DH}(\Delta_1) = \left(\frac{32567112922303267}{27292859194142720}, \frac{247470028273390111}{81878577582428160}\sqrt{3}\right) \approx (1.193, 3.022 \times \sqrt{3}).$$

Since it is in the relative interior of the translated cone $2\rho + C^+$, $\overline{G_2}(1)$ admits a Kähler–Einstein metric.

(2) Case-I.2. This variety $\overline{G_2}(2)$ is the smooth Fano compactification of G_2 with Picard number one called the *double Cayley Grassmannian* (see [Man21]). In this case, the barycenter

$$\mathbf{bar}_{DH}(\Delta_2) = \left(\frac{139601}{79360}, \frac{49}{15}\sqrt{3}\right) \approx (1.759, 3.267 \times \sqrt{3})$$

is in the relative interior of the translated cone $2\rho + C^+$ so that $\overline{G_2}(2)$ admits a Kähler–Einstein metric (see [LPY21, Section 3.6]).



FIGURE 5. Moment polytopes of Gorenstein Fano compactifications of G_2 .

4.4. $A_1 \times A_1$ -type.

4.4.1. Gorenstein Fano group compactification of $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$. As $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$ is simplyconnected, the spherical weight lattice \mathcal{M} is spanned by $\varpi_1 = (1,0)$ and $\varpi_2 = (0,1)$, and its dual lattice \mathcal{N} is spanned by $\alpha_1^{\vee} = (1,0)$ and $\alpha_2^{\vee} = (0,1)$. The Weyl walls are given by $W_1 = \{y = 0\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (2,2)$. Choosing a realization of the root system $A_1 \times A_1$ in the Euclidean plane \mathbb{R}^2 with $\varpi_1 = (1,0)$ and $\varpi_2 = (0,1)$, for p = (x,y) we obtain the Duistermaat–Heckman measure

$$\prod_{\alpha \in \Phi^+} \kappa(\alpha, p)^2 \, dp = (2x)^2 (2y)^2 \, dx dy$$

because $A_1 \times A_1$ has 2 positive roots: $\Phi^+ = \{\alpha_1, \alpha_2\} = \{(2, 0), (0, 2)\}.$

Theorem 4.9. There are 15 Gorenstein Fano equivariant compactifications of $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$ up to isomorphism: two smooth compactifications and 13 singular compactifications. Their moment polytopes are given in the following Table 7 and Figure 6. Among them only two smooth Fano compactifications and a unique singular one admit Kähler–Einstein metrics.

X	Case	Edges (except Weyl walls)	Vertices	Smooth?	KE	$\mathbf{bar}(\Delta)$
$\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(1)$	I.1.1	$y = 3, \ x + 2y = 7$	(0,3),(1,3),(7,0)	singular	No	$\left(\frac{11427}{3794}, \frac{5661}{3794}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{SL}_2(2)$	I.1.1.a	y = 3, x + 2y = 7, x + y = 5	(0,3), (1,3), (3,2), (5,0)	singular	No	$\left(\frac{174753}{72086}, \frac{119919}{72086}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{SL}_2(3)$	I.1.1.b	y = 3, x + 2y = 7, x = 3	(0,3), (1,3), (3,2), (3,0)	singular	No	$\left(\frac{106833}{52486}, \frac{4233}{2282}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(4)$	I.1.2	y = 3, 2x + 3y = 11, x + y = 5	(0, 3), (1, 3), (4, 1), (5, 0)	singular	No	$\left(\frac{130089}{55657}, \frac{87441}{55657}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(5)$	I.1.3	$y = 3, \ 3x + 4y = 15$	(0,3),(1,3),(5,0)	singular	No	$\left(\frac{243}{112}, \frac{177}{112}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(6)$	I.2.1	$y = 3, \ x + y = 5$	(0, 3), (2, 3), (5, 0)	singular	No	$\left(\frac{1569}{665}, \frac{1233}{665}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(7)$	I.2.1.a	y = 3, x + y = 5, x = 3	(0, 3), (2, 3), (3, 2), (3, 0)	smooth	Yes	$\left(\frac{28779}{14035}, \frac{28779}{14035}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(8)$	I.2.2	$y = 3, \ 3x + y = 9$	(0, 3), (2, 3), (3, 0)	singular	No	$\left(\frac{2817}{1631}, \frac{3411}{1631}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(9)$	I.3	y = 3, x = 3	(0, 3), (3, 3)(3, 0),	smooth	Yes	$\left(\frac{9}{4},\frac{9}{4}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(10)$	II.1	x + y = 5	(0,5), (5,0)	singular	Yes	$\left(\frac{15}{7},\frac{15}{7}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{SL}_2(11)$	II.2.1	x + 3y = 9	(0,3), (9,0)	singular	No	$\left(\frac{27}{7},\frac{9}{7}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{SL}_2(12)$	II.2.1.a	$x + 3y = 9, \ x + 2y = 7$	(0,3), (3,2), (7,0)	singular	No	$\left(\frac{293793}{93989}, \frac{131547}{93989}\right)$
$\overline{\mathrm{SL}}_2 \times \overline{\mathrm{SL}}_2(13)$	II.2.1.b	$x + 3y = 9, \ x + y = 5$	(0,3), (3,2), (5,0)	singular	No	$\left(\frac{160017}{63637}, \frac{98619}{63637}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(14)$	II.2.2	$2x + 5y = 15, \ x + 2y = 7$	(0,3), (5,1), (7,0)	singular	No	$\left(\frac{245313}{77896}, \frac{101469}{77896}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{SL}_2(15)$	II.2.3	3x + 7y = 21	(0,3), (7,0)	singular	No	$\left(3,\frac{9}{7}\right)$

TABLE 7. Gorenstein Fano equivariant compactifications of $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$.

Proof. Let $m_i \alpha_1^{\vee} + n_i \alpha_2^{\vee} = (m_i, n_i) \in \mathcal{N}$ be the primitive outer normal vector of the facet $F_i = \{m_i x + n_i y = 2m_i + 2n_i + 1\},$

where $m_i > 0, n_i \ge 0$ by the convexity of the polytope.

• Case-I: F_1 is orthogonal to W_2 . $F_1 \perp W_2 \Rightarrow F_1 = \{y = 3\}$ $F_1 \cap F_2 =: A_1 = \left(\frac{2m_2 - n_2 + 1}{m_2}, 3\right) = \left(2 + \frac{-n_2 + 1}{m_2}, 3\right) \in \mathcal{M} \Rightarrow A_1 = (1,3) \text{ or } (2,3) \text{ or } (3,3)$ $- \text{Case-I.1:} A_1 = (1,3) \Rightarrow m_2 + 1 = n_2$ $* \text{Case-I.1.1:} (m_2, n_2) = (1,2) \text{ and } F_2 = \{x + 2y = 7\}, A_2 = (7,0)$ $(\Delta_1, \text{ Figure 6(1)}).$

- * Case-I.1.1.a: $A_2 = (3, 2)$ and $F_3 = \{x + y = 5\}$ (Δ_2 , Figure 6(2)).
- * Case-I.1.1.b: $A_2 = (3, 2)$ and $F_3 = \{x = 3\}$ (Δ_3 , Figure 6(3)).
- * Case-I.1.2: $(m_2, n_2) = (2, 3)$ and $F_2 = \{2x + 3y = 11\}, A_2 = (4, 1)$ and $F_3 = \{x + y = 5\}$ (Δ_4 , Figure 6(4)).
- * Case-I.1.3: $(m_2, n_2) = (3, 4)$ and $F_2 = \{3x + 4y = 15\}, A_2 = (5, 0)$ $(\Delta_5, \text{ Figure } 6(5)).$
- Case-I.2: $A_1 = (2,3) \Rightarrow n_2 = 1$

If $(m_2, n_2) = (2, 1)$ so that $F_2 = \{2x + y = 7\}$, then $A_2 = (3, 1)$ and $F_3 = \{x = 3\}$. In this case, the polytope is isomorphic to Case-I.1.1.b.

- * Case-I.2.1: $(m_2, n_2) = (1, 1)$ and $F_2 = \{x + y = 5\}, A_2 = (5, 0)$ $(\Delta_6, \text{ Figure } 6(6)).$
- * Case-I.2.1.a: $A_2 = (3, 2)$ and $F_3 = \{x = 3\}$ (Δ_7 , Figure 6(7)).
- * Case-I.2.2: $(m_2, n_2) = (3, 1)$ and $F_2 = \{3x + y = 9\}, A_2 = (3, 0)$ $(\Delta_8, \text{ Figure 6(8)}).$

- Case-I.3: $A_1 = (3,3) \Rightarrow (m_2, n_2) = (1,0)$ so that $F_2 = \{x = 3\}$ (Δ_9 , Figure 6(9)). • Case-II: F_1 is not orthogonal to W_2 .

- $F_1 \cap W_2 =: A_1 = (0, \frac{2m_1 + 2n_1 + 1}{n_1}) \text{ where } n_1 \ge m_1 > 0 \text{ (Convexity and symmetry)} \\ A_1 \in \mathcal{M} \Rightarrow \frac{2m_1 + 1}{n_1} \in \mathbb{N}_+ \Rightarrow \frac{2m_1 + 1}{n_1} = 1(A_1 = (0, 3)) \text{ or } \frac{2m_1 + 1}{n_1} = 3(A_1 = (0, 5)) \text{ (primitive)} \\ \text{Case-III.1: } \frac{2m_1 + 1}{n_1} = 3 \Rightarrow (m_1, n_1) = (1, 1) \text{ and } F_1 = \{x + y = 5\}, A_2 = (5, 0)$

$$(\Delta_{10}, \, {
m Figure} \, \, {
m 6(10)}).$$

- **Case-II.2**: $\frac{2m_1+1}{n_1} = 1 \Rightarrow F_1 = \{m_1x + (2m_1+1)y = 6m_1+3\} \Rightarrow m_1 = 1, 2, 3$
 - * Case-II.2.1: $(m_1, n_1) = (1, 3)$ and $F_1 = \{x + 3y = 9\}, A_2 = (9, 0) (\Delta_{11}, \dots, \Delta_{11})$ Figure 6(11)).
 - * Case-II.2.1.a: $A_2 = (3, 2)$ and $F_2 = \{x + 2y = 7\}$ (Δ_{12} , Figure 6(12)).
 - * Case-II.2.1.b: $A_2 = (3, 2)$ and $F_2 = \{x + y = 5\}$ (Δ_{13} , Figure 6(13)).
 - * Case-II.2.2: $(m_1, n_1) = (2, 5)$ and $F_1 = \{2x + 5y = 15\}, A_2 = (5, 1)$ and $F_2 = \{x + 2y = 7\}$ (Δ_{14} , Figure 6(14)).
 - * Case-II.2.3: $(m_1, n_1) = (3, 7)$ and $F_1 = \{3x + 7y = 21\}, A_2 = (7, 0)$ $(\Delta_{15}, \text{ Figure 6(15)}).$

In the Case-I.3, the variety $\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(9)$ is isomorphic to $\mathbb{Q}^3 \times \mathbb{Q}^3$ from Example 2.7. As $SL_2 \times SL_2(9)$ is a homogeneous variety, it naturally admits a Kähler–Einstein metric.

Applying Theorem 3.3, we can determine which of them are equivariant K-stable and admit (singular) Kähler-Einstein metric. For instance, in the Case-I.2.1.a, the variety $\overline{SL_2 \times SL_2(7)}$ is the blow-up of the smooth Fano compactification $\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(9)$ along the closed orbit which is a diagonal embedding of \mathbb{Q}^3 . Since the barycenter

$$\mathbf{bar}_{DH}(\Delta_7) = \left(\frac{28779}{14035}, \frac{28779}{14035}\right) \approx (2.051, 2.051)$$

of the moment polytope Δ_7 with respect to the Duistermaat-Heckman measure is in the relative interior of the translated cone $2\rho + \mathcal{C}^+$, $\overline{\mathrm{SL}_2 \times \mathrm{SL}_2}(7)$ admits a Kähler–Einstein metric by Theorem 3.3. JAE-HYOUK LEE, KYEONG-DONG PARK, AND SUNGMIN YOO



FIGURE 6. Moment polytopes of Gorenstein Fano compactifications of $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$ -(i).



FIGURE 6. Moment polytopes of Gorenstein Fano compactifications of $SL_2(\mathbb{C}) \times SL_2(\mathbb{C})$ -(ii).

4.4.2. Gorenstein Fano group compactification of $PSL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$. As $PSL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ is of adjoint type, the spherical weight lattice \mathcal{M} is spanned by $\alpha_1 = (2,0)$ and $\alpha_2 = (0,2)$, and its dual lattice \mathcal{N} is spanned by $\varpi_1^{\vee} = \left(\frac{1}{2}, 0\right)$ and $\varpi_2^{\vee} = \left(0, \frac{1}{2}\right)$. The Weyl walls are given by $W_1 = \{y = 0\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (2, 2)$.

Theorem 4.10. There are seven Gorenstein Fano equivariant compactifications of $PSL_2(\mathbb{C}) \times$ $PSL_2(\mathbb{C})$ up to isomorphism: two smooth compactifications and five singular compactifications. Their moment polytopes are given in the following Table 8 and Figure 7. Among them only two smooth compactifications and three singular compactifications admit Kähler-Einstein metrics.

X	Case	Edges (except Weyl walls)	Vertices	Smooth?	KE	$\mathbf{bar}(\Delta)$
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(1)$	I.1.1	$y = 4, \ x + y = 6$	(0,4), (2,4), (6,0)	singular	Yes	$\left(\frac{783}{287}, \frac{678}{287}\right)$
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(2)$	I.1.1.a	y = 4, x + y = 6, x = 4	(0,4), (2,4), (4,2), (4,0)	smooth	Yes	$\left(\frac{10254}{4081}, \frac{10254}{4081}\right)$
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(3)$	I.1.2	$y = 4, \ 2x + y = 8$	(0,4), (2,4), (4,0)	singular	Yes	$\left(\frac{99}{49}, \frac{128}{49}\right)$
$\overline{\mathrm{PSL}}_2 \times \mathrm{PSL}_2(4)$	I.2	y = 4, x = 4	(0,4), (4,4), (4,0)	smooth	Yes	(3,3)
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(5)$	II.1	x + y = 6	(0, 6), (6, 0)	singular	Yes	$\left(\frac{18}{7},\frac{18}{7}\right)$
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(6)$	II.2.1	x + 2y = 8	(0,4),(8,0)	singular	No	$\left(\frac{24}{7},\frac{12}{7}\right)$
$\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(7)$	II.2.1.a	$x + 2y = 8, \ x + y = 6$	(0, 4), (4, 2), (6, 0)	singular	No	$\left(\frac{8418}{2863}, \frac{5346}{2863}\right)$

TABLE 8.	Gorenstein	Fano	equivariant	compactifications	of PSL_2	$(\mathbb{C}$	$) \times PSL_2$	(\mathbb{C})	ļ
			1	1		`	/ -	· ·	

Proof. Let $m_i \varpi_1^{\vee} + n_i \varpi_2^{\vee} = \left(\frac{m_i}{2}, \frac{n_i}{2}\right) \in \mathcal{N}$ be the primitive outer normal vector of the facet $F_i = \{m_i x + n_i y = 2(m_i + n_i + 1)\},\$

where $m_i \ge 0, n_i \ge 0$ by the convexity of the polytope.

For the facet F_1 which intersects the Weyl wall W_2 , we have two cases:

- Case-I: F_1 is orthogonal to W_2 . $\Rightarrow F_1 = \{y = 4\}, F_1 \cap F_2 =: A_1 = \left(2 + 2\frac{-n_2+1}{m_2}, 4\right) \in \mathcal{M}$ - Case-I.1: $A_1 = (2, 4) \Rightarrow n_2 = 1$
 - * Case-I.1.1: $(m_2, n_2) = (1, 1)$ and $F_2 = \{x + y = 6\}, A_2 = (6, 0)$ $(\Delta_1, \text{ Figure 7(1)}).$
 - * Case-I.1.1.a: $A_2 = (4, 2)$ and $F_3 = \{x = 4\}$ (Δ_2 , Figure 7(2)).
 - * Case-I.1.2: $(m_2, n_2) = (2, 1)$ and $F_2 = \{2x + y = 8\}, A_2 = (4, 0)$
 - $(\Delta_3, \text{ Figure 7(3)}).$

- Case-I.2:
$$A_1 = (4,4) \Rightarrow (m_2, n_2) = (1,0)$$
 so that $F_2 = \{x = 4\}$ (Δ_4 , Figure 7(4)).

- **Case-II:** F_1 is not orthogonal to W_2 . $A_1 = (0, 2\frac{m_1 + n_1 + 1}{n_1}) \in \mathcal{M} \Rightarrow \frac{m_1 + n_1 + 1}{n_1} = 2(A_1 = (0, 4)) \text{ or } \frac{m_1 + n_1 + 1}{n_1} = 3(A_1 = (0, 6))$ -**Case-II.1:** $\frac{m_1 + n_1 + 1}{n_1} = 3 \Rightarrow (m_1, n_1) = (1, 1) \text{ and } F_1 = \{x + y = 6\}, A_2 = (6, 0)$
 - $(\Delta_5, \text{ Figure 7(5)})$
 - Case-II.2: $\frac{m_1+n_1+1}{n_1} = 2 \Rightarrow m_1 + 1 = n_1 \Rightarrow F_1 = \{m_1x + (m_1+1)y = 4(m_1+1)\}$
 - * Case-II.2.1: $m_1 = 1 \Rightarrow F_1 = \{x + 2y = 8\}, A_2 = (8,0) \ (\Delta_6, \text{ Figure 7(6)}).$
 - * Case-II.2.1.a: $A_2 = (4, 2)$ and $F_2 = \{x + y = 6\}$ (Δ_7 , Figure 7(7)).

In the Case-I.2, the variety $\overline{\text{PSL}_2 \times \text{PSL}_2}(4)$ is isomorphic to $\mathbb{P}^3 \times \mathbb{P}^3$ from Example 2.7. As $\overline{PSL_2} \times \overline{PSL_2}(4)$ is a homogeneous variety, it naturally admits a Kähler–Einstein metric.

Applying Theorem 3.3, we can determine which of them are equivariant K-stable and admit (singular) Kähler-Einstein metric. For instance, in the Case-I.1.1.a, the variety $\overline{PSL_2 \times PSL_2}(2)$ is the blow-up of the smooth Fano compactification $\overline{PSL_2 \times PSL_2}(4)$ along the closed orbit which is a diagonal embedding of \mathbb{P}^3 . Since the barycenter $\mathbf{bar}_{DH}(\Delta_2) = \left(\frac{10254}{4081}, \frac{10254}{4081}\right) \approx (2.513, 2.513)$ is in the relative interior of the cone $2\rho + \mathcal{C}^+$, $\overline{\mathrm{PSL}_2 \times \mathrm{PSL}_2}(2)$ admits a Kähler–Einstein metric. \Box



FIGURE 7. Moment polytopes of Gorenstein Fano compactifications of $PSL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$.

4.4.3. Gorenstein Fano group compactification of $SL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$. The spherical weight lattice \mathcal{M} is spanned by $\varpi_1 = (1,0)$ and $\alpha_2 = (0,2)$, and its dual lattice \mathcal{N} is spanned by $\alpha_1^{\vee} = (1,0)$ and $\varpi_2^{\vee} = \left(0, \frac{1}{2}\right)$. The Weyl walls are given by $W_1 = \{y = 0\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (2,2)$.

Theorem 4.11. There are 14 Gorenstein Fano equivariant compactifications of $SL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ up to isomorphism: two smooth compactifications and 12 singular compactifications. Their moment polytopes are given in the following Table 9 and Figure 8. Among them only one smooth compactification and three singular compactifications admit singular Kähler–Einstein metrics.

X	Case	Edges (except Weyl walls)	Vertices	Smooth?	KE	$\mathbf{bar}(\Delta)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(1)$	I.1.1	$y = 4, \ x + y = 5$	(0, 4), (1, 4), (5, 0)	singular	Yes	$\left(\frac{243}{112}, \frac{59}{28}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(2)$	I.1.1.a	$y = 4, x + y = 5, x + \frac{1}{2}y = 4$	(0, 4), (1, 4), (3, 2), (4, 0)	singular	Yes	$\left(\frac{12069}{6034}, \frac{13383}{6034}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(3)$	I.1.1.b	y = 4, x + y = 5, x = 3	(0, 4), (1, 4), (3, 2), (3, 0)	singular	No	$\left(\frac{2067}{1099}, \frac{2550}{1099}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(4)$	I.1.2	$y = 4, \ 4x + 3y = 16$	(0, 4), (1, 4), (4, 0)	singular	No	$\left(\frac{1797}{1022}, \frac{1128}{511}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(5)$	I.2.1	$y = 4, \ x + \frac{1}{2}y = 4$	(0,4), (2,4), (4,0)	singular	Yes	$\left(\frac{99}{49}, \frac{128}{49}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(6)$	I.2.1.a	$y = 4, \ x + \frac{1}{2}y = 4, \ x = 3$	(0, 4), (2, 4), (3, 2), (3, 0)	smooth	No	$\left(\frac{17274}{8869}, \frac{23943}{8869}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(7)$	I.2.2	$y = 4, \ 4x + y = 12$	(0, 4), (2, 4), (3, 0)	singular	No	$\left(\frac{2817}{1631}, \frac{4548}{1631}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(8)$	I.3	y = 4, x = 3	(0, 4), (3, 4), (3, 0)	smooth	Yes	$\left(\frac{9}{4},3\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(9)$	II.1	4x + y = 12	(0, 12), (3, 0)	singular	No	$\left(\frac{9}{7},\frac{36}{7}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{PSL}_2}(10)$	II.2.1	2x + y = 8	(0,8),(4,0)	singular	No	$\left(\frac{12}{7},\frac{24}{7}\right)$
$\overline{\mathrm{SL}}_2 \times \mathrm{PSL}_2(11)$	II.2.1.a	$2x + y = 8, \ x = 3$	(0, 8), (3, 2), (3, 0)	singular	No	$\left(\frac{846}{511}, \frac{1797}{511}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{PSL}_2}(12)$	II.2.1.b	$2x + y = 8, \ 4x + y = 12$	(0,8), (3,2), (3,0)	singular	No	$\left(\frac{4209}{2863}, \frac{10692}{2863}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{PSL}_2}(13)$	II.2.1.c	$2x + y = 8, \ 3x + y = 9$	(0,8),(1,6),(3,0)	singular	No	$\left(\frac{10887}{8288}, \frac{30927}{8288}\right)$
$\overline{\mathrm{SL}_2 \times \mathrm{PSL}_2}(14)$	II.2.2	8x + 3y = 24	(0,8),(3,0)	singular	No	$\left(\frac{9}{7}, \frac{24}{7}\right)$

TABLE 9. Gorenstein Fano equivariant compactifications of $SL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$.

Proof. Let $m_i \varpi_1^{\vee} + n_i \alpha_2^{\vee} = (m_i, 2n_i) \in \mathcal{N}$ be the primitive outer normal vector of the facet

$$F_i = \left\{ m_i x + \frac{n_i}{2} y = 2m_i + n_i + 1 \right\},\,$$

where $m_i \ge 0, n_i \ge 0$ by the convexity of the polytope. For the facet F_1 which intersects the Weyl wall W_2 , we have two cases:

• Case-I: F_1 is orthogonal to W_2 . $F_1 \perp W_2 \Rightarrow F_1 = \{y = 4\}$ $F_1 \cap F_2 =: A_1 = \left(\frac{2m_2 - n_2 + 1}{m_2}, 4\right) = \left(2 + \frac{-n_2 + 1}{m_2}, 4\right) \in \mathcal{M}$ -Case-I.1: $A_1 = (1, 4) \Rightarrow m_2 + 1 = n_2$ * Case-I.1.1: $(m_2, n_2) = (1, 2)$ and $F_2 = \{x + y = 5\}, A_2 = (5, 0)$ $(\Delta_1,$ Figure 8(1)). * Case-I.1.1.a: $A_2 = (3, 2)$ and $F_3 = \{x + \frac{1}{2}y = 4\}$ (Δ_2 , Figure 8(2)). * Case-I.1.1.b: $A_2 = (3, 2)$ and $F_3 = \{x = 3\}$ (Δ_3 , Figure 8(3)). * Case-I.1.2: $(m_2, n_2) = (2, 3)$ and $F_2 = \{4x + 3y = 16\}, A_2 = (4, 0)$ $(\Delta_4,$ Figure 8(4)). - Case-I.2: $A_1 = (2, 4) \Rightarrow n_2 = 1$

- * **Case-I.2.1**: $(m_2, n_2) = (1, 1)$ and $F_2 = \left\{ x + \frac{1}{2}y = 4 \right\}, A_2 = (4, 0)$ (Δ_5 , Figure 8(5)).
- * Case-I.2.1.a: $A_2 = (3, 2)$ and $F_3 = \{x = 3\}$ (Δ_6 , Figure 8(6)).
- * Case-I.2.2: $(m_2, n_2) = (2, 1)$ and $F_2 = \{4x + y = 12\}, A_2 = (3, 0)$ (Δ_7 , Figure 8(7)).
- Case-I.3: $A_1 = (3,4) \Rightarrow m_2 + n_2 = 1 \Rightarrow (m_2, n_2) = (1,0)$ and $F_2 = \{x = 3\}$ (Δ_8 , Figure 8(8)).
- Case-II: F_1 is not orthogonal to W_2 .

$$F_{1} \cap W_{2} =: A_{1} = \left(0, 2^{\frac{2m_{1}+n_{1}+1}{n_{1}}}\right) = \left(0, 2\left(1+\frac{2m_{1}+1}{n_{1}}\right)\right) \in \mathcal{M} \Rightarrow \frac{2m_{1}+1}{n_{1}} \in \mathbb{N}_{+}$$
Note that $\emptyset \neq F_{i} \cap W_{1} = \left(0, 2+\frac{n_{i}+1}{m_{i}}\right) \in \mathcal{M} \Rightarrow \frac{n_{i}+1}{m_{i}} \ge 1 \Rightarrow n_{i} + 1 \ge m_{i}$

$$- \operatorname{Case-II.1:} \frac{2m_{1}+1}{n_{1}} = 5 \Rightarrow (m_{1}, n_{1}) = (2, 1) \text{ and } F_{1} = \{4x + y = 12\}, A_{2} = (3, 0)$$

$$(\Delta_{9}, \operatorname{Figure} 8(9)).$$

$$- \operatorname{Case-II.2:} \frac{2m_{1}+1}{n_{1}} = 3 \Rightarrow (m_{1}, n_{1}) = (1, 1) \text{ or } (m_{1}, n_{1}) = (4, 3)$$

$$* \operatorname{Case-II.2.1:} (m_{1}, n_{1}) = (1, 1) \text{ and } F_{1} = \{2x + y = 8\}, A_{2} = (4, 0)$$

$$(\Delta_{10}, \operatorname{Figure} 8(10)).$$

$$* \operatorname{Case-II.2.1.a:} A_{2} = (3, 2) \text{ and } F_{2} = \{x = 3\} (\Delta_{11}, \operatorname{Figure} 8(11)).$$

$$* \operatorname{Case-II.2.1.b:} A_{2} = (2, 4) \text{ and } F_{2} = \{4x + y = 12\} (\Delta_{12}, \operatorname{Figure} 8(12)).$$

$$* \operatorname{Case-II.2.1.c:} A_{2} = (1, 6) \text{ and } F_{2} = \{3x + y = 9\} (\Delta_{3}, \operatorname{Figure} 8(13)).$$

$$* \operatorname{Case-II.2.1.c:} A_{2} = (1, 6) \text{ and } F_{2} = \{8x + 3y = 24\}, A_{2} = (3, 0)$$

* Case-II.2.2: $(m_1, n_1) = (4, 3)$ and $F_1 = \{8x + 3y = 24\}, A_2 = (3, 0)$ (Δ_{14} , Figure 8(14)).

In the Case-I.2.1.a, the toric polytope Δ_6^{toric} formed by the Weyl group action from the moment polytope Δ_6 is a Delzant polytope and no vertices of Δ_6^{toric} lie in Weyl walls, hence $\overline{SL_2 \times PSL_2}(6)$ is smooth by Proposition 4.1. Since the barycenter of the moment polytope Δ_6 with respect to the Duistermaat–Heckman measure is given by

$$\mathbf{bar}_{DH}(\Delta_6) = \left(\frac{17274}{8869}, \frac{23943}{8869}\right) \approx (1.948, 2.700),$$

it is not in the relative interior of the translated cone $2\rho + C^+$. Thus $\overline{\text{SL}_2 \times \text{PSL}_2}(6)$ does not admit any singular Kähler–Einstein metric by Theorem 3.3.

Let Q be the point at which the half-line starting from the barycenter $C = \mathbf{bar}_{DH}(\Delta_6)$ in the direction of A = (2, 2) intersects the boundary of $(\Delta_6 - \mathcal{C}^+)$. Considering the line x = 3 giving a part of $\partial(\Delta_6 - \mathcal{C}^+)$ and the half-line \overrightarrow{CA} , by Corollary 3.10, the greatest Ricci lower bound is

$$R(\overline{\mathrm{SL}_2 \times \mathrm{PSL}_2}(6)) = \frac{\overline{AQ}}{\overline{CQ}} = \frac{1}{3 - \frac{17274}{8869}} = \frac{8869}{9333} \approx 0.9503.$$

In the Case-I.3, the variety $\overline{\text{SL}_2 \times \text{PSL}_2(8)}$ is isomorphic to $\mathbb{Q}^3 \times \mathbb{P}^3$ from Example 2.7. As $\overline{\text{SL}_2 \times \text{PSL}_2(8)}$ is a homogeneous variety, it naturally admits a Kähler–Einstein metric.

JAE-HYOUK LEE, KYEONG-DONG PARK, AND SUNGMIN YOO



FIGURE 8. Moment polytopes of Gorenstein Fano compactifications of $SL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ -(i).



FIGURE 8. Moment polytopes of Gorenstein Fano compactifications of $SL_2(\mathbb{C}) \times PSL_2(\mathbb{C})$ -(ii).

4.4.4. Gorenstein Fano equivariant compactifications of $SO_4(\mathbb{C})$. The spherical weight lattice \mathcal{M} of $SO_4(\mathbb{C})$ is spanned by $\varpi_1 + \varpi_2 = (1, 1)$ and $-\varpi_1 + \varpi_2 = (-1, 1)$, and its dual lattice \mathcal{N} is spanned by $\varpi_1^{\vee} + \varpi_2^{\vee} = \left(\frac{1}{2}, \frac{1}{2}\right)$ and $-\varpi_1^{\vee} + \varpi_2^{\vee} = \left(-\frac{1}{2}, \frac{1}{2}\right)$. The Weyl walls are given by $W_1 = \{y = 0\}$ and $W_2 = \{x = 0\}$. The sum of positive roots is $2\rho = (2, 2)$.

In [LTZ23, Section 7.1], Li–Tian–Zhu obtained the following results. Note that they used different bases for \mathcal{M} and \mathcal{N} . In fact, taking coordinate changes via the linear transformation given by a matrix $\begin{pmatrix} \frac{1}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{1}{2} \end{pmatrix}$, the bases $\varpi_1 + \varpi_2 = (1, 1)$ and $-\varpi_1 + \varpi_2 = (-1, 1)$ of \mathcal{M} can be transformed as (1, 0) and (0, 1), respectively, as in [LTZ23, Section 7.1].

Theorem 4.12. [LTZ23, Section 7.1] There are six Gorenstein Fano equivariant compactifications of $SO_4(\mathbb{C})$ up to isomorphism: three smooth compactifications and three singular compactifications. Their moment polytopes are given in the following Table 10 and Figure 9. Among them, the only one compactification in each cases admits a Kähler–Einstein metric.

X	Case	Edges (except Weyl walls)	Vertices	Smooth?	KE	$\mathbf{bar}(\Delta)$
$\overline{\mathrm{SO}_4}(1)$	I.1.1	y = 3, x + 3y = 10	(0,3),(1,3),(10,0)	singular	No	$\left(\frac{8229}{1918}, \frac{1368}{959}\right)$
$\overline{\mathrm{SO}_4}(2)$	I.1.2	y = 3, x + 3y = 10, x + y = 6	(0,3), (1,3), (4,2), (6,0)	smooth	No	$\left(\frac{467331}{150514}, \frac{125928}{75257}\right)$
$\overline{\mathrm{SO}_4}(3)$	I.1.3	$y = 3, \ 3x + 5y = 18$	(0,3),(1,3),(6,0)	singular	No	$\left(\frac{1341}{518}, \frac{396}{259}\right)$
$\overline{\mathrm{SO}_4}(4)$	I.2.1	$y = 3, \ x + y = 6$	(0,3),(3,3),(6,0)	smooth	No	$\left(\frac{297}{98},\frac{96}{49}\right)$
$\overline{\mathrm{SO}_4}(5)$	I.2.2	y = 3, x = 3	(0,3),(3,3),(3,0)	singular	Yes	$\left(\frac{9}{4},\frac{9}{4}\right)$
$\overline{\mathrm{SO}_4}(6)$	II	x + y = 6	(0, 6), (6, 0)	smooth	Yes	$\left(\frac{18}{7},\frac{18}{7}\right)$

TABLE 10. Gorenstein Fano equivariant compactifications of $SO_4(\mathbb{C})$.

Proof of Theorem 1.3 for K-unstable smooth compactifications of $SO_4(\mathbb{C})$. (1) In the Case I.1.2 (Case 1.1.2 in [LTZ23, Section 7.1]), $\overline{SO_4}(2)$ is a smooth compactification of $SO_4(\mathbb{C})$ corresponding to Figure 9(2) and the barycenter of its moment polytope is

$$(\bar{x}, \bar{y}) = \left(\frac{467331}{150514}, \frac{125928}{75257}\right) \approx (3.1049, 0.7538).$$

Let Q be the point at which the half-line starting from the barycenter C in the direction of A = (2, 2)intersects the boundary of the moment polytope. Considering the line y = 3 and the half-line \overrightarrow{CA} , we can compute $Q = \left(-\frac{67959}{49172}, 3\right)$. Thus, we obtain the greatest Ricci lower bound $R = \frac{\overrightarrow{AQ}}{\overrightarrow{CQ}} = \frac{75257}{99843} \approx 0.7538$ by Corollary 3.10.

(2) In the Case I.2.1 (Case 1.2.1 in [LTZ23, Section 7.1]), $\overline{SO_4}(4)$ is a smooth compactification of $SO_4(\mathbb{C})$ corresponding to Figure 9(4) and the barycenter of its moment polytope is

$$(\bar{x}, \bar{y}) = \left(\frac{297}{98}, \frac{96}{49}\right) \approx (3.0306, 1.9592).$$

Let Q be the point at which the half-line starting from the barycenter C in the direction of A = (2, 2)intersects the boundary of the moment polytope. Considering the line y = 3 and the half-line \overrightarrow{CA} , we compute $Q = \left(-\frac{93}{4},3\right)$. Thus, we obtain the greatest Ricci lower bound $R = \frac{\overrightarrow{AQ}}{\overrightarrow{CQ}} = \frac{49}{51} \approx 0.9608$ by Corollary 3.10.



FIGURE 9. Moment polytopes of Gorenstein Fano compactifications of $SO_4(\mathbb{C})$.

References

- [AB04] Valery Alexeev and Michel Brion, Stable reductive varieties II: projective case, Adv. Math. 184 (2004), no. 2, 380–408.
- [AK05] Valery Alexeev and Ludmil Katzarkov, On K-stability of reductive varieties, Geom. Funct. Anal. 15 (2005), 297–310.
- [BBEGZ19] Robert Berman, Sebastien Boucksom, Philippe Eyssidieux, Vincent Guedj and Ahmed Zeriahi, Kähler-Einstein metrics and the Kähler-Ricci flow on log Fano varieties, J. Reine Angew. Math. 751 (2019), 27–89.
- [BBJ21] Robert Berman, Sébastien Boucksom and Mattias Jonsson, A variational approach to the Yau-Tian-Donaldson conjecture, J. Amer. Math. Soc. 34 (2021), no. 3, 605–652.
- [Bri87] Michel Brion, Sur l'image de l'application moment, Séminaire d'algèbre Paul Dubreil et Marie-Paule Malliavin (Paris, 1986), Lecture Notes in Math., vol. 1296, Springer, Berlin, 1987, pp. 177–192.
- [Bri89] _____, Groupe de Picard et nombres caractéristiques des variétés sphériques, Duke Math. J. 58 (1989), no. 2, 397–424.
- [Bri94] _____, Factorisation de certains morphismes birationnels, Compos. Math. 91 (1994), no. 1, 57–66.
- [Bri97] _____, Curves and divisors in spherical varieties, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 21–34.
- [Cao85] Huai-Dong Cao, Deformation of Kähler metrics to Kähler–Einstein metrics on compact Kähler manifolds, Invent. Math. 81 (1985), no. 2, 359–372.
- [CDS15a] Xiuxiong Chen, Simon Donaldson, and Song Sun, K\u00e4hler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities, J. Amer. Math. Soc. 28 (2015), no. 1, 183–197.
- [CDS15b] _____, Kähler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π, J. Amer. Math. Soc. 28 (2015), no. 1, 199–234.
- [CDS15c] _____, Kähler–Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof, J. Amer. Math. Soc. **28** (2015), no. 1, 235–278.
- [CLS11] David Cox, John Little and Hal Schenck, Toric Varieties, Vol. 124. American Mathematical Soc., 2011.
- [CRZ19] Ivan Cheltsov, Yanir Rubinstein and Kewei Zhang, Basis log canonical thresholds, local intersection estimates, and asymptotically log del Pezzo surfaces, Selecta Math. 25 (2019), Article number 34.
- [DS16] Ved Datar and Gábor Székelyhidi, Kähler-Einstein metrics along the smooth continuity method, Geom. Funct. Anal. 26 (2016), 975–1010.
- [DCP83] Corrado De Concini and Claudio Procesi, Complete symmetric varieties, Invariant theory (Montecatini, 1982), Lecture Notes in Math., vol. 996, Springer, Berlin, 1983, pp. 1–44.
- [Del15] Thibaut Delcroix, *Métriques de Kähler–Einstein sur les compactifications de groupes*, PhD thesis, Université Grenoble Alpes, October 2015.
- [Del17] _____, Kähler–Einstein metrics on group compactifications, Geom. Funct. Anal. 27 (2017), no. 1, 78–129.
- [Del20] _____, K-stability of Fano spherical varieties, Ann. Sci. Éc. Norm. Supér. (4) 53 (2020), no. 3, 615–662.
- [FO18] Kento Fujita and Yuji Odaka, On the K-stability of Fano varieties and anticanonical divisors, Tohoku Math. J. 70 (2018), no. 4, 511–521.
- [Gan18] Jacopo Gandini, Embeddings of spherical homogeneous spaces, Acta Math. Sin. (Engl. Ser.) 34 (2018), no. 3, 299–340.
- [GH15] Giuliano Gagliardi and Johannes Hofscheier, Gorenstein spherical Fano varieties, Geom. Dedicata 178 (2015), 111–133.
- [Gol20] Aleksei Golota, Delta-invariants for Fano varieties with large automorphism groups, Internat. J. Math. 31 (2020), no. 10, 2050077.
- [Kno91] Friedrich Knop, The Luna-Vust theory of spherical embeddings, Proceedings of the Hyderabad Conference on Algebraic Groups (Hyderabad, 1989), Manoj Prakashan, Madras, 1991, pp. 225–249.
- [Li11] Chi Li, Greatest lower bounds on Ricci curvature for toric Fano manifolds, Adv. Math. 226 (2011), no. 6, 4921–4932.
- [Li17] _____, Yau-Tian-Donaldson correspondence for K-semistable Fano manifolds, J. Reine Angew. Math. 733 (2017), 55–85.
- [Li22] _____, G-uniform stability and Kähler-Einstein metrics on Fano varieties, Invent. Math. 227 (2022), no. 2, 661–744.
- [LPY21] Jae-Hyouk Lee, Kyeong-Dong Park and Sungmin Yoo, Kähler–Einstein metrics on smooth Fano symmetric varieties with Picard number one, Mathematics 9 (2021), no. 1, 102.
- [LTZ18] Yan Li, Gang Tian and Xiaohua Zhu, Singular limits of Kähler-Ricci flow on Fano G-manifolds, preprint (2021), arXiv:1807.09167v5.
- [LTZ23] _____, Singular Kähler–Einstein metrics on Q-Fano compactifications of Lie groups, Math. Eng., 5 (2023), no. 2, 1–43.

- [Lun97] Domingo Luna, Grosses cellules pour les variétés sphériques, Algebraic groups and Lie groups, Austral. Math. Soc. Lect. Ser., vol. 9, Cambridge Univ. Press, Cambridge, 1997, pp. 267–280.
- [LV83] Domingo Luna and Thierry Vust, Plongements d'espaces homogènes, Comment. Math. Helv. 58 (1983), no. 2, 186–245.
- [Mab87] Toshiki Mabuchi, Einstein-Kähler forms, Futaki invariants and convex geometry on toric Fano varieties, Osaka J. Math. 24 (1987), no. 4, 705–737.
- [Man21] Laurent Manivel, The double Cayley Grassmannian, Math. Res. Lett. 28 (2021), no. 6, 1765–1792.
- [Mat72] Yozo Matsushima, Remarks on Kähler-Einstein manifolds, Nagoya Math. J. 46 (1972), 161–173.
- [OV90] Arkadij L. Onishchik and Ernest B. Vinberg, Lie groups and algebraic groups, Translated from the Russian
- and with a preface by D. A. Leites, Springer Series in Soviet Mathematics, Springer-Verlag, Berlin, 1990. [Per02] Grisha Perelman, *The entropy formula for the Ricci flow and its geometric applications*, preprint (2002), arXiv:0211159.
- [Rub08] Yanir A. Rubinstein, Some discretizations of geometric evolution equations and the Ricci iteration on the space of Kähler metrics, Adv. Math. 218 (2008), no. 5, 1526–1565.
- [Rub09] _____, On the construction of Nadel multiplier ideal sheaves and the limiting behavior of the Ricci flow, Trans. Amer. Math. Soc. **361** (2009), no. 11, 5839–5850.
- [Ruz10] Alessandro Ruzzi, Geometrical description of smooth projective symmetric varieties with Picard number one, Transform. Groups 15 (2010), no. 1, 201–226.
- [Ruz12] _____, Fano symmetric varieties with low rank, Publ. Res. Inst. Math. Sci. 48 (2012), no. 2, 235–278.
- [Szé11] Gábor Székelyhidi, Greatest lower bounds on the Ricci curvature of Fano manifolds, Compos. Math. 147, no. 1, 319–331.
- [Tia87] Gang Tian, On Kähler–Einstein metrics on certain Kähler manifolds with $C_1(M) > 0$, Invent. Math. 89 (1987), no. 2, 225–246.
- [Tia92] _____, On stability of the tangent bundles of Fano varieties, Internat. J. Math. 3 (1992), no. 3, 401–413.
- [Tia97] _____, Kähler–Einstein metrics with positive scalar curvature, Invent. Math. 130 (1997), no. 1, 1–37.
- [Tia15] _____, K-stability and Kähler–Einstein metrics, Commun. Pure Appl. Math. 68 (2015), no. 7, 1085–1156.
- [Tim03] Dmitry A. Timashev, Equivariant compactifications of reductive groups, Sbornik: Mathematics 194 (2003), no. 4, 589–616.
- [Tim11] _____, Homogeneous spaces and equivariant embeddings, Encyclopaedia of Mathematical Sciences, vol. 138, Springer, Heidelberg, 2011, Invariant Theory and Algebraic Transformation Groups, 8.
- [WZ04] Xu-Jia Wang and Xiaohua Zhu, Kähler-Ricci solitons on toric manifolds with positive first Chern class, Adv. Math. 188 (2004), no. 1, 87–103.

DEPARTMENT OF MATHEMATICS, EWHA WOMANS UNIVERSITY, SEODAEMUN-GU, SEOUL 03760, REPUBLIC OF KOREA

Email address: jaehyoukl@ewha.ac.kr

DEPARTMENT OF MATHEMATICS AND RESEARCH INSTITUTE OF NATURAL SCIENCE, GYEONGSANG NATIONAL UNIVERSITY, JINJU 52828, REPUBLIC OF KOREA

Email address: kdpark@gnu.ac.kr

DEPARTMENT OF MATHEMATICS, INCHEON NATIONAL UNIVERSITY, YEONSU-GU, INCHEON 22012, REPUBLIC OF KOREA

Email address: sungminyoo.math@inu.ac.kr