

NLO QCD Renormalization Group Evolution for Non-Leptonic $\Delta F = 2$ Transitions in the SMEFT

Jason Aebischer^a, Andrzej J. Buras^b, Jacky Kumar^b

^aPhysik-Institut, Universität Zürich, CH-8057 Zürich, Switzerland

^bTUM Institute for Advanced Study, Lichtenbergstr. 2a, D-85747 Garching, Germany

Abstract

We present for the first time NLO QCD Renormalization Group (RG) evolution matrices for non-leptonic $\Delta F = 2$ transitions in the Standard Model Effective Field Theory (SMEFT). To this end we transform first the known two-loop QCD anomalous dimension matrices (ADMs) of the BSM operators in the so-called BMU basis into the ones in the common Weak Effective Theory (WET) basis (the so-called JMS basis) for which tree-level and one-loop matching to the SMEFT are already known. This allows us subsequently to find the two-loop QCD ADMs for the SMEFT non-leptonic $\Delta F = 2$ operators in the Warsaw basis. Having all these ingredients we investigate the impact of these NLO QCD effects on the QCD RG evolution of SMEFT Wilson coefficients for non-leptonic $\Delta F = 2$ transitions from the new physics scale Λ down to the electroweak scale μ_{ew} . The main benefit of these new contributions is that they allow to remove renormalization scheme dependences present both in the one-loop matchings between the WET and SMEFT and also between SMEFT and a chosen UV completion. But the NLO QCD effects, calculated here in the NDR scheme, turn out to be small, in the ballpark of a few percent but larger than one-loop Yukawa top effects when only the $\Delta F = 2$ operators are considered. The technology developed in our paper allows to obtain the ADMs in the SMEFT from the ones of the BMU basis also for non-leptonic $\Delta F = 1$ decays and the results of this more involved analysis will be presented soon in another publication.

1 Introduction

Non-leptonic $\Delta F = 2$ transitions, represented by $K^0 - \bar{K}^0$, $D^0 - \bar{D}^0$ and $B_{d,s}^0 - \bar{B}_{d,s}^0$ mixings play very important roles in the tests of the Standard Model (SM) and of the New Physics (NP) beyond it [1]. In order to increase the precision of these tests it is necessary to go beyond the leading order (LO) analyses both in the Weak Effective Theory (WET) and also in the Standard Model Effective Field Theory (SMEFT). To this end it is mandatory to include first in the renormalization group (RG) analyses in these theories the one-loop matching contributions, both between these two theories as well as when passing thresholds at which heavy particles are integrated out. But this is not the whole story, a fact which is forgotten in some recent SMEFT analyses present in the literature. To complete a NLO analysis and remove various renormalization scheme (RS) dependences in the one-loop matching also two-loop anomalous dimensions of all operators in the WET and SMEFT have to be included.

The present status of these efforts in the case of non-leptonic meson $\Delta F = 1$ decays and $\Delta F = 2$ quark mixing processes is as follows:

- The matchings in question are known by now both at tree-level [2] and one-loop level [3]¹.
- The one-loop ADMs relevant for the RG in WET [9, 10] and SMEFT [11–13] are also known.
- The two-loop QCD ADMs relevant for RG evolutions for both $\Delta F = 1$ and $\Delta F = 2$ transitions in WET are also known [14–16].

The main goal of the present paper is to extend the QCD RG evolution in the SMEFT for $\Delta F = 2$ transitions beyond the leading order. In fact at first sight this is straightforward because the $SU(3)_c$ symmetry remains unbroken in the SMEFT and in the absence of electroweak interactions one could just use the NLO QCD BSM analysis of [14] up to the new physics scale Λ in the so-called BMU operator basis that is useful for NLO QCD calculations. However, in the presence of electroweak interactions, the so-called SMEFT Warsaw basis [17] is more suitable and it is necessary to perform the QCD renormalization group analysis within the SMEFT in that basis. Therefore we present here for the first time two-loop ADMs for $\Delta F = 2$ four-quark operators of the SMEFT. We will demonstrate that they can be obtained from the two-loop ADMs in the BMU basis [14]. As an intermediate step we will present the ADM relevant for the WET in the so-called JMS basis [2] which for the matching of the WET to the SMEFT is more useful than the BMU basis.

The main technology presented here can be extended to non-leptonic $\Delta F = 1$ transitions but due to the large number of operators involved [14, 15] the corresponding analysis is much more complicated and will be presented in due time in a separate publication.

Having all these ingredients we investigate numerically the impact of the NLO corrections calculated here on the LO RG evolutions of SMEFT Wilson coefficients for $\Delta F = 2$ transitions from the new physics scale Λ down to the electroweak scale μ_{ew} presented by us in [16]. Including also the effects of top Yukawa couplings at the one-loop level, taken already into account in the latter paper, we find that the main benefit from the present analysis is the removal of QCD renormalization scheme dependences present currently both in the one-loop matchings between the WET and SMEFT and also between SMEFT and a chosen UV

¹Previous partial results can be found, for example, in [4–8].

completion. But the NLO QCD effects in RG QCD evolution, calculated here in the NDR scheme [18], turn out to be small, in the ballpark of a few percent but larger than one-loop Yukawa top effects when only the $\Delta F = 2$ operators are considered.

Our paper is organized as follows. In Section 2 we derive the general relation between QCD RG evolutions in two different operator bases as well as the corresponding relations for one-loop and two-loop ADMs. This presentation is valid for any number of operators and can be extended to other processes, in particular $\Delta F = 1$ non-leptonic transitions. Using these formulae we summarize in Section 3 the one-loop and two-loop ADMs for $\Delta F = 2$ transitions in BMU, JMS and SMEFT bases as well as the matching matrices between these bases. Subsequently in Section 4 we analyze numerically the size of the NLO QCD corrections calculated here by presenting the RG evolution matrices in the WET and SMEFT both at LO and NLO in QCD. We summarize in Section 5

2 Basic Formulae for NLO QCD RG Evolution

2.1 Evolution Matrix

The RG evolution matrix for non-leptonic transitions in the BMU basis for SM and BSM operators is known including one-loop and two-loop QCD contributions. It is given as follows

$$\hat{U}_{\text{BMU}}(\mu_{\text{had}}, \mu_{\text{ew}}) = \left[1 + \hat{J}_{\text{BMU}} \frac{\alpha_s(\mu_{\text{had}})}{4\pi} \right] \hat{U}_{\text{BMU}}^{(0)}(\mu_{\text{had}}, \mu_{\text{ew}}) \left[1 - \hat{J}_{\text{BMU}} \frac{\alpha_s(\mu_{\text{ew}})}{4\pi} \right], \quad (1)$$

where $\hat{U}_{\text{BMU}}^{(0)}$ is the RS-independent LO evolution matrix. On the other hand, \hat{J}_{BMU} stems from the RS-dependent two-loop ADMs, which makes them sensitive to the renormalization scheme considered. This scheme dependence is cancelled by the one of the matching at μ_{ew} and by the one of the hadronic matrix elements at μ_{had} . Explicit general expressions for $\hat{U}_i^{(0)}$ and \hat{J}_i in terms of the coefficients of the one-loop and two-loop perturbative expansions for the ADM $\hat{\gamma}$ and the QCD β -function can be found including their derivations in Chapter 5 of [1]. They will be listed in the next subsection. Unless explicitly stated the following discussion applies to both $\Delta F = 2$ and $\Delta F = 1$ transitions which are characterized by different operators and their corresponding ADMs.

Our goal is to obtain an analogous expression in the SMEFT, that is

$$\hat{U}_{\text{SMEFT}}(\mu_{\text{ew}}, \Lambda) = \left[1 + \hat{J}_{\text{SMEFT}} \frac{\alpha_s(\mu_{\text{ew}})}{4\pi} \right] \hat{U}_{\text{SMEFT}}^{(0)}(\mu_{\text{ew}}, \Lambda) \left[1 - \hat{J}_{\text{SMEFT}} \frac{\alpha_s(\Lambda)}{4\pi} \right]. \quad (2)$$

To this end we notice that this evolution depends only on \hat{J}_{SMEFT} and $\hat{U}_{\text{SMEFT}}^{(0)}$ without any explicit dependence on one- and two-loop anomalous dimensions of the SMEFT operators. This gives us a hint that it should be possible to obtain \hat{J}_{SMEFT} and $\hat{U}_{\text{SMEFT}}^{(0)}$ from the known \hat{J}_{BMU} and $\hat{U}_{\text{BMU}}^{(0)}$ without knowing explicitly one-loop and two-loop ADMs in the SMEFT. It turns out that this is indeed possible but as an intermediate step we should first find \hat{J}_{JMS} and $\hat{U}_{\text{JMS}}^{(0)}$ from \hat{J}_{BMU} and $\hat{U}_{\text{BMU}}^{(0)}$. We will perform this intermediate step in what follows.

In order to reach this goal we first present the general formula which allows to obtain \hat{J}_{B} and $\hat{U}_{\text{B}}^{(0)}$ in a given basis B from \hat{J}_{A} and $\hat{U}_{\text{A}}^{(0)}$ in the basis A for which these two objects, like in the BMU basis, are already known.

Defining the transformation matrix \hat{R} between these two operator bases through

$$\vec{Q}_B = \hat{R}\vec{Q}_A, \quad \vec{C}_A = \hat{R}^T\vec{C}_B, \quad \hat{R} = \hat{R}_0 + \frac{\alpha_s(\mu)}{4\pi}\hat{R}_1, \quad (3)$$

a straightforward calculation results in the relations we were looking for:

$$\boxed{\hat{J}_B = (\hat{R}_0^{-1})^T \hat{J}_A \hat{R}_0^T - (\hat{R}_0^{-1})^T \hat{R}_1^T, \quad \hat{U}_B^{(0)}(\mu_1, \mu_2) = (\hat{R}_0^{-1})^T \hat{U}_A^{(0)}(\mu_1, \mu_2) \hat{R}_0^T,} \quad (4)$$

Let us apply these formulae to the two cases of interest.

Example 1

Here

$$A=BMU, \quad B=JMS, \quad \hat{R}^T \equiv \hat{M}_{JMS} = \hat{M}_0 + \frac{\alpha_s(\mu)}{4\pi}\hat{M}_1 \quad (5)$$

and we find

$$\boxed{\hat{J}_{JMS} = \hat{M}_0^{-1} \hat{J}_{BMU} \hat{M}_0 - \hat{M}_0^{-1} \hat{M}_1, \quad \hat{U}_{JMS}^{(0)}(\mu_1, \mu_2) = \hat{M}_0^{-1} \hat{U}_{BMU}^{(0)}(\mu_1, \mu_2) \hat{M}_0.} \quad (6)$$

Example 2

Here

$$A=JMS, \quad B=SMEFT, \quad \hat{R}^T \equiv \hat{K} = \hat{K}_0 + \frac{\alpha_s(\mu)}{4\pi}\hat{K}_1. \quad (7)$$

and we find

$$\boxed{\hat{J}_{SMEFT} = \hat{K}_0^{-1} \hat{J}_{JMS} \hat{K}_0 - \hat{K}_0^{-1} \hat{K}_1, \quad \hat{U}_{SMEFT}^{(0)} = \hat{K}_0^{-1} \hat{U}_{JMS}^{(0)} \hat{K}_0,} \quad (8)$$

where we suppressed the scales in the last relation. The reason is that SMEFT and JMS are valid at different energy scales. Therefore in order to use the second relation in the equation above in the basic formula (2) one should properly adjust the scales in $\hat{U}_{JMS}^{(0)}$. To test it one can use the one-loop ADMs of [11–13].

In summary starting with the known RG NLO evolution in the BSM basis one can not only find the corresponding NLO evolution in WET in the JMS basis but also the NLO evolution in the SMEFT. To this end the matrices \hat{M}_{JMS} and \hat{K} including one-loop contributions have to be calculated.

This formulation is general but in the case at hand significant simplifications take place:

$$\hat{M}_{JMS} = \hat{M}_0, \quad \hat{K}_0 = \hat{1}. \quad (9)$$

Both are related to the absence of evanescent contributions in the relevant basis transformations [16]. But the second one demonstrates the advantage of the JMS basis over the BMU basis when the matching to SMEFT is concerned.

2.2 One-Loop and Two-Loop ADMs in SMEFT

Despite the possibility of finding the NLO RG evolution matrix in the basis B from the one in the basis A without knowing anomalous dimensions in basis B, it is useful to find these anomalous dimensions if one wants to include subsequently the effects of Yukawa couplings.

In order to reach this goal we first present the general formula which allows to obtain $\hat{\gamma}_B$ in a given basis B from $\hat{\gamma}_A$ in the basis A which is, like in the BMU basis, already known.

The general ADM is defined as [1]

$$\hat{\gamma}(g) = \hat{Z}^{-1} \frac{d\hat{Z}}{d\ln\mu}, \quad (10)$$

where \hat{Z} is the operator renormalization matrix which depends on the basis considered. The ADM has the following perturbative expansion

$$\hat{\gamma}(\alpha_s) = \frac{\alpha_s}{4\pi} \hat{\gamma}^{(0)} + \left(\frac{\alpha_s}{4\pi}\right)^2 \hat{\gamma}^{(1)} + \mathcal{O}(\alpha_s^3). \quad (11)$$

Using the general notation for the transformation matrix \hat{R} in (3), the renormalization matrices in the two bases A and B are related through

$$\hat{Z}_B = \hat{Z}_A \left(\hat{R}_0 - \frac{\alpha_s(\mu)}{4\pi} \hat{R}_1 \right). \quad (12)$$

In [1] the derivation of the relations we are seeking for has been presented for the case $R_0 = \hat{1}$ and $\hat{R}_1 \equiv \hat{s}$ with the result given in (5.180) of that book. Generalizing it to the case at hand we find

$$\boxed{\hat{\gamma}_B^{(0)} = \hat{R}_0^{-1} \hat{\gamma}_A^{(0)} \hat{R}_0, \quad \hat{\gamma}_B^{(1)} = \hat{R}_0^{-1} \hat{\gamma}_A^{(1)} \hat{R}_0 - \hat{R}_0^{-1} \hat{\gamma}_A^{(0)} \hat{R}_1 + \hat{R}_1 \hat{\gamma}_A^{(0)} \hat{R}_0 + 2\beta_0 \hat{R}_0^{-1} \hat{R}_1.} \quad (13)$$

In obtaining the last term in the second equation above we used the relation

$$\frac{d\alpha_s}{d\ln\mu} = -2\beta_0 \frac{\alpha_s^2}{4\pi}, \quad (14)$$

where $\beta_0 = (11N_c - 2N_f)/3$ represents the one-loop QCD β function. For $R_0 = \hat{1}$ and $\hat{R}_1 = \hat{s}$ we confirm (5.180) in [1].

Starting then with the ADMs in the BMU basis one can find the ADMs in the JMS basis and subsequently ADMs in the SMEFT. However, there is one subtlety to be taken care of. The anomalous dimension matrix $\hat{\gamma}$ as defined in (10) and in the most literature including [1] and in particular in the BMU basis in [14] governs the RG evolution of the matrix elements of the operators involved while the evolution of the corresponding Wilson coefficients is governed by the transposed matrix $\hat{\gamma}^T$. But the authors that introduced the JMS and SMEFT bases decided to denote by $\hat{\gamma}$ the one which usually would be called $\hat{\gamma}^T$.

This is unfortunate and in what follows we will within QCD use the standard notation so that the RG evolution of operator matrix elements will be governed by $\hat{\gamma}$ and the evolution of Wilson coefficients by $\hat{\gamma}^T$. Therefore when listing ADMs and \hat{J} in the JMS and SMEFT bases we will use the standard notation used also in the BMU basis in [14]. The resulting QCD RG evolution matrices at LO and NLO obviously do not depend on this choice. We will also see at the end of the next section that the LO Yukawa effects in the RG evolution can also be incorporated in the presence of NLO QCD corrections in a straightforward manner.

3 One-Loop and Two-Loop ADMs in BMU, JMS and SMEFT Bases

3.1 BMU

We begin with the BMU basis [14] for which the complete ADMs at NLO in QCD have been calculated in [14]. The BMU basis consists in full generality of $(5 + 3) = 8$ physical

operators belonging to the five distinct sectors (VLL, SLL, LR, VRR, SRR). However, SLL and SRR operators, violating hypercharge conservation are not allowed within the SMEFT at dimension-six level and we will not consider them in what follows. Adopting the `WCxf` convention [19], the remaining four operators are ($ij = sd, db, sb, cu$)²

$$\begin{aligned} Q_{\text{VLL}}^{ij} &= [\bar{d}_i \gamma_\mu P_L d_j][\bar{d}_i \gamma^\mu P_L d_j], & Q_{\text{VRR}}^{ij} &= [\bar{d}_i \gamma_\mu P_R d_j][\bar{d}_i \gamma^\mu P_R d_j], \\ Q_{\text{LR},1}^{ij} &= [\bar{d}_i \gamma_\mu P_L d_j][\bar{d}_i \gamma^\mu P_R d_j], & Q_{\text{LR},2}^{ij} &= [\bar{d}_i P_L d_j][\bar{d}_i P_R d_j], \end{aligned} \quad (15)$$

which are built exclusively out of colour-singlet currents $[\bar{d}_i^\alpha \dots d_j^\alpha][\bar{d}_i^\beta \dots d_j^\beta]$, where α, β denote colour indices. This feature is very useful for calculations in DQCD [20, 21], because their matrix elements in the large- N_c limit can be obtained directly without using Fierz identities.

The one-loop and two-loop anomalous dimensions are given as follows

$$\hat{\gamma}_{\text{BMU}}^{(0)} = \begin{pmatrix} 6 - \frac{6}{N_c} & 0 & 0 & 0 \\ 0 & 6 - \frac{6}{N_c} & 0 & 0 \\ 0 & 0 & \frac{6}{N_c} & 12 \\ 0 & 0 & 0 & -6N_c + \frac{6}{N_c} \end{pmatrix}, \quad \hat{\gamma}_{\text{BMU}}^{(1)} = \begin{pmatrix} \hat{\gamma}_{\text{VLL}}^{(1)} & 0 & 0 & 0 \\ 0 & \hat{\gamma}_{\text{VRR}}^{(1)} & 0 & 0 \\ 0 & 0 & (\hat{\gamma}_{\text{LR}}^{(1)})_{11} & (\hat{\gamma}_{\text{LR}}^{(1)})_{12} \\ 0 & 0 & (\hat{\gamma}_{\text{LR}}^{(1)})_{21} & (\hat{\gamma}_{\text{LR}}^{(1)})_{22} \end{pmatrix}. \quad (16)$$

Here

$$\hat{\gamma}_{\text{VLL}}^{(1)} = \hat{\gamma}_{\text{VRR}}^{(1)} = -\frac{19}{6}N_c - \frac{22}{3} + \frac{39}{N_c} - \frac{57}{2N_c^2} + \frac{2}{3}N_f - \frac{2}{3N_c}N_f, \quad (17)$$

and

$$(\hat{\gamma}_{\text{LR}}^{(1)})_{11} = \frac{137}{6} + \frac{15}{2N_c^2} - \frac{22}{3N_c}N_f, \quad (18)$$

$$(\hat{\gamma}_{\text{LR}}^{(1)})_{12} = \frac{200}{3}N_c - \frac{6}{N_c} - \frac{44}{3}N_f, \quad (19)$$

$$(\hat{\gamma}_{\text{LR}}^{(1)})_{21} = \frac{71}{4}N_c + \frac{9}{N_c} - 2N_f, \quad (20)$$

$$(\hat{\gamma}_{\text{LR}}^{(1)})_{22} = -\frac{203}{6}N_c^2 + \frac{479}{6} + \frac{15}{2N_c^2} + \frac{10}{3}N_cN_f - \frac{22}{3N_c}N_f. \quad (21)$$

Here N_c is the number of colours with $N_c = 3$ in QCD. N_f is the number of quark flavours, $N_f = 3, 4, 5$ in the WET and $N_f = 6$ in the SMEFT. The numerical solutions for evolution matrices for $ij = sd, db, sb$ are given in [22].

3.2 JMS

The JMS basis has been introduced to facilitate the classification of the complete WET operator basis [2] for the purpose of matching from SMEFT onto WET. The relevant $\Delta F = 2$ operators are

²We use the ordering of flavours as in [16] but different papers use different conventions, which has to be taken into account.

$$\begin{aligned}
[Q_{dd}^{VLL}]_{ijij} &= Q_{VLL}^{ij}, & [Q_{dd}^{VRR}]_{ijij} &= Q_{VRR}^{ij}, & [Q_{dd}^{V1,LR}]_{ijij} &= Q_{LR,1}^{ij}, \\
[Q_{dd}^{V8,LR}]_{ijij} &= [\bar{d}_i \gamma_\mu P_L T^A d_j][\bar{d}_i \gamma^\mu P_R T^A d_j] = -\frac{1}{2N_c} Q_{LR,1}^{ij} - Q_{LR,2}^{ij},
\end{aligned} \tag{22}$$

where T^A are $SU(3)_c$ colour generators of the fundamental representation.

As demonstrated in [16], using the method described in [23], the matching matrix \hat{R}_1 vanishes in this case and consequently only \hat{R}_0 enters the basis transformation. Using the relations above one finds

$$\hat{R}_0 = \hat{R}_0^{-1} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -\frac{1}{2N_c} & -1 \end{pmatrix}. \tag{23}$$

Using (13) we find one-loop and two-loop ADMs in the JMS basis

$$\hat{\gamma}_{\text{JMS}}^{(0)} = \begin{pmatrix} 6 - \frac{6}{N_c} & 0 & 0 & 0 \\ 0 & 6 - \frac{6}{N_c} & 0 & 0 \\ 0 & 0 & 0 & -12 \\ 0 & 0 & -\frac{6}{N_c} C_F & -6N_c + \frac{12}{N_c} \end{pmatrix}, \quad \hat{\gamma}_{\text{JMS}}^{(1)} = \begin{pmatrix} (\hat{\gamma}_{\text{JMS}}^{(1)})_{11} & 0 & 0 & 0 \\ 0 & (\hat{\gamma}_{\text{JMS}}^{(1)})_{22} & 0 & 0 \\ 0 & 0 & (\hat{\gamma}_{\text{JMS}}^{(1)})_{33} & (\hat{\gamma}_{\text{JMS}}^{(1)})_{34} \\ 0 & 0 & (\hat{\gamma}_{\text{JMS}}^{(1)})_{43} & (\hat{\gamma}_{\text{JMS}}^{(1)})_{44} \end{pmatrix}, \tag{24}$$

where we have still used the BMU notation for $\hat{\gamma}$ so that in contrast to [9]

$$\mu \frac{d}{d\mu} \vec{C}_{\text{JMS}} = \hat{\gamma}_{\text{JMS}}^T \vec{C}_{\text{JMS}}, \tag{25}$$

as emphasized above. Here,

$$(\hat{\gamma}_{\text{JMS}}^{(1)})_{11} = (\hat{\gamma}_{\text{JMS}}^{(1)})_{22} = \frac{(N_c - 1)(171 - 19N_c^2 - N_c(63 - 4N_f))}{6N_c^2}, \tag{26}$$

$$(\hat{\gamma}_{\text{JMS}}^{(1)})_{33} = \frac{21}{2} \left(-1 + \frac{1}{N_c^2}\right), \tag{27}$$

$$(\hat{\gamma}_{\text{JMS}}^{(1)})_{34} = \frac{6}{N_c} - \frac{200N_c}{3} + \frac{44N_f}{3}, \tag{28}$$

$$(\hat{\gamma}_{\text{JMS}}^{(1)})_{43} = \frac{(N_c^2 - 1)(9 - 208N_c^2 + 22N_c N_f)}{6N_c^3}, \tag{29}$$

$$(\hat{\gamma}_{\text{JMS}}^{(1)})_{44} = \frac{27 + 679N_c^2 - 203N_c^4 - 88N_c N_f + 20N_c^3 N_f}{6N_c^2}, \tag{30}$$

and $C_F = (N_c^2 - 1)/(2N_c)$. We have checked that the obtained $\hat{\gamma}_{\text{JMS}}^{(0)}$ agrees after transposition with the results in [9].

3.3 SMEFT

3.3.1 Operators

In the SMEFT there are five operators which can contribute to $\Delta F = 2$ processes at the dimension-six level

$$\begin{aligned}\mathcal{O}_{qq}^{(1)} &= (\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t), & \mathcal{O}_{qq}^{(3)} &= (\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t), \\ \mathcal{O}_{dd} &= (\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t), & \mathcal{O}_{qd}^{(1)} &= (\bar{q}_p \gamma_\mu q_r)(\bar{d}_s \gamma^\mu d_t), \\ \mathcal{O}_{qd}^{(8)} &= (\bar{q}_p \gamma_\mu T^A q_r)(\bar{d}_s \gamma^\mu T^A d_t).\end{aligned}\tag{31}$$

The relevant SMEFT Wilson coefficients are

$$B = \left\{ \mathcal{C}_{qq}^{(1)} + \mathcal{C}_{qq}^{(3)}, \quad \mathcal{C}_{aa}, \quad \mathcal{C}_{qa}^{(1)}, \quad \mathcal{C}_{qa}^{(8)} \right\},\tag{32}$$

in the down ($a = d$) and up ($a = u$) sector, respectively [4]. At tree-level for $B_{s,d} - \bar{B}_{s,d}$ and $K^0 - \bar{K}^0$ mixing one finds the following matching conditions at μ_{ew} in the **down-basis**

$$\begin{aligned}[C_{dd}^{V,LL}]_{ijij} &= -[\mathcal{C}_{qq}^{(1)}]_{ijij} - [\mathcal{C}_{qq}^{(3)}]_{ijij}, & [C_{dd}^{V,RR}]_{ijij} &= -[\mathcal{C}_{dd}]_{ijij}, \\ [C_{dd}^{V1,LR}]_{ijij} &= -[\mathcal{C}_{qd}^{(1)}]_{ijij}, & [C_{dd}^{V8,RR}]_{ijij} &= -[\mathcal{C}_{qd}^{(8)}]_{ijij},\end{aligned}\tag{33}$$

and for $D^0 - \bar{D}^0$ mixing in the **up-basis**

$$\begin{aligned}[C_{uu}^{V,LL}]_{ijij} &= -[\hat{\mathcal{C}}_{qq}^{(1)}]_{ijij} - [\hat{\mathcal{C}}_{qq}^{(3)}]_{ijij}, & [C_{uu}^{V,RR}]_{ijij} &= -[\hat{\mathcal{C}}_{uu}]_{ijij}, \\ [C_{uu}^{V1,LR}]_{ijij} &= -[\hat{\mathcal{C}}_{qu}^{(1)}]_{ijij}, & [C_{uu}^{V8,RR}]_{ijij} &= -[\hat{\mathcal{C}}_{qu}^{(8)}]_{ijij},\end{aligned}\tag{34}$$

where we have neglected contributions from modified Z -coupling operators. Note that we use the Hamiltonian for WET to define Wilson coefficients contrary to [2, 3], who use the Lagrangian, in consequence minus signs are present in the matching conditions.

3.3.2 QCD anomalous dimensions

The ADMs in this case are the same as for the JMS basis

$$\hat{\gamma}_{\text{SMEFT}}^{(0)} = \hat{\gamma}_{\text{JMS}}^{(0)}, \quad \hat{\gamma}_{\text{SMEFT}}^{(1)} = \hat{\gamma}_{\text{JMS}}^{(1)}.\tag{35}$$

Here $\hat{\gamma}_{\text{JMS}}^{(0)}$ and $\hat{\gamma}_{\text{JMS}}^{(1)}$ are given by (24)-(30).

Inspecting the RG evolution for $[\hat{\mathcal{C}}_{qq}^{(1)}]_{ijij}$ and $[\hat{\mathcal{C}}_{qq}^{(3)}]_{ijij}$ one finds that the sum for dd or uu indices from these two Wilson coefficients evolves without being affected by other operators and only this sum matches on the VLL operator in the JMS basis. While this can be verified explicitly by using the RG equations (35) and (36) in [16], the inclusion of flavour diagonal gluon exchanges at the two-loop level cannot change this property.

3.3.3 Top Yukawa anomalous dimension

In this subsection we report the LO ADM resulting from top Yukawa interactions. It reads for a given sector with flavour indices $ij = sd, db, sb, cu$ in the down-basis [4]:

$$\hat{\gamma}_{y_t}^{(0)} = \frac{y_t^2}{16\pi^2} \begin{pmatrix} r_{ij} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2}r_{ij} & 0 \\ 0 & 0 & 0 & \frac{1}{2}r_{ij} \end{pmatrix}, \quad (36)$$

with the combination $r_{ij} = |V_{ti}|^2 + |V_{tj}|^2$. In the up-basis the ADM is zero and the corresponding mixing effects only come into play when considering back-rotation [24] at the EW scale.

Indeed this anomalous dimension matrix can be extracted from formulae (30)-(33) in [16]. It is the coefficient of

$$L = \frac{1}{(4\pi)^2} \ln\left(\frac{\mu_{\text{ew}}}{\Lambda}\right) \quad (37)$$

in the solution of RG equations retaining only the first leading logarithm and neglecting the μ -dependence of y_t . Defining

$$[\mathcal{C}_{qq}^{(1+3)}]_{ijij} \equiv [\mathcal{C}_{qq}^{(1)}]_{ijij} + [\mathcal{C}_{qq}^{(3)}]_{ijij}, \quad (38)$$

these equations read in **the down-basis**

$$[\mathcal{C}_{qq}^{(1+3)}]_{ijij}(\mu_{\text{ew}}) = [\mathcal{C}_{qq}^{(1+3)}]_{ijij} + y_t^2 \left[\lambda_t^{ik} [\mathcal{C}_{qq}^{(1+3)}]_{kjiij} + \lambda_t^{kj} [\mathcal{C}_{qq}^{(1+3)}]_{ikij} \right] L, \quad (39)$$

$$[\mathcal{C}_{qd}^{(1)}]_{ijij}(\mu_{\text{ew}}) = [\mathcal{C}_{qd}^{(1)}]_{ijij} + y_t^2 \left[\frac{\lambda_t^{ik}}{2} [\mathcal{C}_{qd}^{(1)}]_{kjiij} + \frac{\lambda_t^{kj}}{2} [\mathcal{C}_{qd}^{(1)}]_{ikij} \right] L, \quad (40)$$

$$[\mathcal{C}_{qd}^{(8)}]_{ijij}(\mu_{\text{ew}}) = [\mathcal{C}_{qd}^{(8)}]_{ijij} + y_t^2 \left[\frac{\lambda_t^{ik}}{2} [\mathcal{C}_{qd}^{(8)}]_{kjiij} + \frac{\lambda_t^{kj}}{2} [\mathcal{C}_{qd}^{(8)}]_{ikij} \right] L, \quad (41)$$

where a summation over k is implied and where we only considered the $\Delta F = 2$ operators in (31). We have suppressed the argument of the NP scale Λ in the Wilson coefficients and SM parameters on the r.h.s to simplify the notation. The μ dependence of y_t will be included in the next subsection.

3.4 Explicit Expression for the Evolution Matrix

3.4.1 Pure QCD

We can now find the NLO QCD evolution matrix in any of the bases considered by us using the general known expressions that we recall here in the case of SMEFT for completeness. Simplifying the notation by dropping the subscript SMEFT we have:

$$\hat{U}(\mu_{\text{ew}}, \Lambda) = \left[1 + \hat{J} \frac{\alpha_s(\mu_{\text{ew}})}{4\pi} \right] \hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda) \left[1 - \hat{J} \frac{\alpha_s(\Lambda)}{4\pi} \right]. \quad (42)$$

Here $\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)$ denotes the usual LO RG evolution matrix that is explicitly given as follows

$$\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda) = \hat{V} \left(\left[\frac{\alpha_s(\Lambda)}{\alpha_s(\mu_{\text{ew}})} \right]^{\frac{\hat{\gamma}^{(0)}}{2\beta_0}} \right)_D \hat{V}^{-1}, \quad (43)$$

where \hat{V} diagonalizes $\hat{\gamma}^{(0)T}$

$$\hat{\gamma}_D^{(0)} = \hat{V}^{-1} \hat{\gamma}^{(0)T} \hat{V}, \quad (44)$$

and $\vec{\gamma}^{(0)}$ is the vector containing the diagonal elements of the diagonal matrix $\hat{\gamma}_D^{(0)}$.

The NLO matrix \hat{J} is given by

$$\hat{J} = \hat{V} \hat{H} \hat{V}^{-1} \quad (45)$$

with

$$(H)_{ij} = \delta_{ij} (\gamma^{(0)})_i \frac{\beta_1}{2\beta_0^2} - \frac{G_{ij}}{2\beta_0 + (\gamma^{(0)})_i - (\gamma^{(0)})_j}, \quad (46)$$

where

$$\hat{G} = \hat{V}^{-1} \hat{\gamma}^{(1)T} \hat{V}, \quad (47)$$

with the two-loop matrix $\hat{\gamma}^{(1)}$ found using (13) and $\beta_1 = \frac{34}{3}N_c^2 - \frac{10}{3}N_cN_f - 2C_F N_f$ [25].

Setting $N_c = 3$ and $N_f = 6$ we find

$$\hat{J}_{\text{SMEFT}}^{(6)} = \hat{J}_{\text{JMS}}^{(6)} = \begin{pmatrix} 1.37 & 0 & 0 & 0 \\ 0 & 1.37 & 0 & 0 \\ 0 & 0 & 1.47 & 2.75 \\ 0 & 0 & 16.60 & 6.92 \end{pmatrix}, \quad \hat{J}_{\text{BMU}}^{(6)} = \begin{pmatrix} 1.37 & 0 & 0 & 0 \\ 0 & 1.37 & 0 & 0 \\ 0 & 0 & -1.30 & -1.38 \\ 0 & 0 & -16.60 & 9.69 \end{pmatrix}. \quad (48)$$

The corresponding matrices for $N_f = 4$ and $N_f = 5$ can be found in Appendix A.

3.4.2 Including Top Yukawa Effects

Until now we succeeded to find the NLO QCD RG evolution in the SMEFT but also the evolution due to the top Yukawa has to be taken into account. But the two-loop ADM for $\Delta F = 2$ operators including top Yukawa couplings is not known at present. Therefore we can only combine the known LO evolution due to Yukawa couplings with the NLO QCD evolution just found. The full evolution is then given by

$$[\hat{U}(\mu_{\text{ew}}, \Lambda)]_{\text{QCD}+y_t} = \left[\hat{1} + \hat{J} \frac{\alpha_s(\mu_{\text{ew}})}{4\pi} \right] [\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)]_{\text{QCD}+y_t} \left[\hat{1} - \hat{J} \frac{\alpha_s(\Lambda)}{4\pi} \right], \quad (49)$$

where the label QCD + y_t indicates that besides QCD also Yukawa contributions have been taken into account. Note that \hat{J} contains only QCD contributions. As NLO corrections due to top Yukawa effects are unknown, it is legitimate to proceed in this manner.

There are two routes to find the LO evolution matrix in this formula. If one is interested only in the numerical result one can simply replace the LO QCD evolution matrix in (42) by the one present in the usual computer codes like `wilson` [26] or `DsixTools` [27, 28] for LO RG evolution in the SMEFT.

However, as demonstrated in [29] a very accurate analytic formula for the RG evolution including scale dependence of y_t can be found. To this end we note that

$$[\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)]_{\text{QCD}+y_t} = [\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)]_{\text{QCD}} [\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)]_{y_t} \quad (50)$$

with the QCD evolution matrix given in (43). To find the second matrix one can simply follow the *case C* in Appendix E of [1] that is a particular case of the general formulae in [29]. We find

$$[\hat{U}^{(0)}(\mu_{\text{ew}}, \Lambda)]_{y_t} = \exp \left[\frac{1}{8\pi} \frac{\hat{\gamma}_t}{\gamma_m^{(0)} - \beta_0} \left([\alpha_s(\Lambda)]^{\frac{\gamma_m^{(0)}}{\beta_0} - 1} - [\alpha_s(\mu_{\text{ew}})]^{\frac{\gamma_m^{(0)}}{\beta_0} - 1} \right) \right], \quad (51)$$

where the diagonal matrix $\hat{\gamma}_t$

$$\hat{\gamma}_t = \hat{b} y_t^2(\mu_0) [\alpha_s(\mu_0)]^{-\frac{\gamma_m^{(0)}}{\beta_0}}, \quad \gamma_m^{(0)} = 8, \quad (52)$$

with the diagonal matrix $\hat{b} = \text{diag}(1, 0, 1/2, 1/2)$ deduced from (36) and an arbitrary scale μ_0 , which we choose to be 160 GeV in our numerical analysis.

4 Numerical Analysis

In order to illustrate the importance of NLO QCD corrections within the SMEFT with respect to the LO ones and top quark Yukawa effects as well as LO and NLO QCD effects within the WET we derive in the following numerical expressions for the evolution matrices in the WET and SMEFT. The results in this section have been obtained using the analytic expressions derived in the previous section.

4.1 WET

Setting $\mu_{\text{had}} = 1.3$ GeV and $\mu_{\text{ew}} = 160$ GeV and using the threshold scale $\mu_5 = 4.2$ GeV between $N_f = 5$ and $N_f = 4$ we find

$$[\hat{U}_{\text{JMS}}^{(0)}]_{\text{QCD}} = \begin{pmatrix} 0.76 & 0 & 0 & 0 \\ 0 & 0.76 & 0 & 0 \\ 0 & 0 & 1.10 & 0.31 \\ 0 & 0 & 1.38 & 2.71 \end{pmatrix}, \quad [\hat{U}_{\text{JMS}}]_{\text{QCD}} = \begin{pmatrix} 0.76 & 0 & 0 & 0 \\ 0 & 0.76 & 0 & 0 \\ 0 & 0 & 1.24 & 0.57 \\ 0 & 0 & 2.02 & 3.59 \end{pmatrix}. \quad (53)$$

Here $\hat{U}_{\text{JMS}}^{(0)}$ and \hat{U}_{JMS} are LO and NLO evolution matrices in the JMS basis, respectively. We observe that in the LR sector the NLO effects are large and it is mandatory to include them in any phenomenological analysis.

4.2 SMEFT

Here we study the evolution between $\mu_{\text{ew}} = 160$ GeV and $\Lambda = 10$ TeV for various cases in the $N_f = 6$ flavour theory.

4.2.1 Pure QCD Evolution

$$[\hat{U}_{\text{SMEFT}}^{(0)}]_{\text{QCD}} = \begin{pmatrix} 0.89 & 0 & 0 & 0 \\ 0 & 0.89 & 0 & 0 \\ 0 & 0 & 1.02 & 0.10 \\ 0 & 0 & 0.43 & 1.52 \end{pmatrix}, \quad [\hat{U}_{\text{SMEFT}}]_{\text{QCD}} = \begin{pmatrix} 0.89 & 0 & 0 & 0 \\ 0 & 0.89 & 0 & 0 \\ 0 & 0 & 1.02 & 0.12 \\ 0 & 0 & 0.46 & 1.57 \end{pmatrix}. \quad (54)$$

As expected, due to a much slower running of α_s and its smaller value than in WET, QCD effects are significantly smaller and this applies in particular to NLO QCD effects.

4.2.2 Pure Yukawa Evolution

When only the Yukawa running in the SMEFT at one-loop is considered, the resulting evolution matrix reads

$$[\hat{U}_{\text{SMEFT}}^{(0)}]_{y_t} = \begin{pmatrix} 0.98 & 0 & 0 & 0 \\ 0 & 1.00 & 0 & 0 \\ 0 & 0 & 0.99 & 0 \\ 0 & 0 & 0 & 0.99 \end{pmatrix} \quad (55)$$

where we have used $y_t(\mu = 160\text{GeV}) = 0.94$.

4.2.3 QCD+Yukawa Evolution

In this subsection we consider the combination of QCD and Yukawa running effects at LO and NLO. The corresponding evolution matrices read

$$[\hat{U}_{\text{SMEFT}}^{(0)}]_{\text{QCD}+y_t} = \begin{pmatrix} 0.87 & 0 & 0 & 0 \\ 0 & 0.89 & 0 & 0 \\ 0 & 0 & 1.01 & 0.10 \\ 0 & 0 & 0.43 & 1.51 \end{pmatrix}, \quad [\hat{U}_{\text{SMEFT}}]_{\text{QCD}+y_t} = \begin{pmatrix} 0.88 & 0 & 0 & 0 \\ 0 & 0.89 & 0 & 0 \\ 0 & 0 & 1.01 & 0.11 \\ 0 & 0 & 0.44 & 1.54 \end{pmatrix}, \quad (56)$$

where for both matrices we have used eq. (49), setting \hat{J} to zero in the LO case, and keeping \hat{J} in the computation of \hat{U}_{SMEFT} .

We observe that when only $\Delta F = 2$ operators are considered the impact of top-Yukawa effects is very small. It is significantly larger when also $\Delta F = 1$ operators are included in the analysis. A detailed discussion of these effects has been presented in [16].

5 Conclusions

The main results of our paper are as follows.

- General formulae for the relation of QCD RG evolution matrices at LO and NLO between two different operator bases, given in eq. (4).

- General formulae for the relation of QCD one-loop and two-loop anomalous dimension matrices between two different operator bases, reported in eq. (13).
- The two-loop QCD ADMs for $\Delta F = 2$ operators in the JMS WET basis and in the SMEFT Warsaw basis. They are given in eqs. (24)-(30) and eq. (35).
- Master formulae for QCD RG evolution matrices for the Wilson coefficients of $\Delta F = 2$ operators in the SMEFT Warsaw basis at the NLO were derived. They are given in eq. (42).
- Generalization of these formulae to include top Yukawa effects at the one-loop level.

These findings allow for a general and scheme-independent QCD analysis of non-leptonic $\Delta F = 2$ processes in the SMEFT and WET at NLO. In a given UV completion in which the Wilson coefficients have been calculated at a NP scale Λ , our master formulae allow to calculate them at the μ_{ew} scale. The inclusion of NLO QCD corrections in the RG evolution in the WET from the hadronic scale μ_{had} to the electroweak scale μ_{ew} allows a correct matching of Wilson coefficients to the matrix elements calculated by lattice QCD (LQCD) or other non-perturbative methods sensitive to renormalization scheme dependences. The use of the JMS basis on the other hand allows to generalize this formula to the SMEFT, because in this basis the tree-level matching of SMEFT onto WET [2] and the one-loop matching [3, 4] are known.

The main messages from the numerical analysis in Section 4 are as follows:

- The NLO QCD corrections to the $\Delta F = 2$ RG evolution matrices within WET are substantial.
- The NLO QCD corrections to the $\Delta F = 2$ RG evolution matrices within SMEFT are small, in the ballpark of a few percent.
- Even smaller are top-Yukawa effects if only $\Delta F = 2$ operators are included in the analysis.

The small NLO QCD corrections to the $\Delta F = 2$ matrices within the SMEFT could be considered at first sight disappointing. But it should be realized that they have been calculated in the NDR scheme but could be larger in a different RS. This also does not preclude significant one-loop matching contributions, which in principle could be larger than the QCD effects found here. However, to combine these one-loop matching conditions with the NLO effects calculated here they have to be calculated in the NDR scheme. Only then RS independent results can be obtained.

Acknowledgements

We thank Christoph Bobeth for useful discussions and comments on the document. J. A. acknowledges financial support from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme under grant agreement 833280 (FLAY), and from the Swiss National Science Foundation (SNF) under contract 200020-204428. A.J.B acknowledges financial support from the Excellence Cluster ORIGINS, funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC-2094 – 390783311. J.K. is financially supported by the Alexander von Humboldt Foundation's postdoctoral research fellowship.

A $\hat{J}_{\text{JMS}}^{(N_f)}$ and $\hat{J}_{\text{BMU}}^{(N_f)}$

For convenience we report in this appendix the \hat{J} matrices for $N_f = 5, 4$ flavours in the JMS and BMU basis, obtained from eq. (45).

$$\hat{J}_{\text{JMS}}^{(5)} = \begin{pmatrix} 1.63 & 0 & 0 & 0 \\ 0 & 1.63 & 0 & 0 \\ 0 & 0 & 1.67 & 2.44 \\ 0 & 0 & 17.04 & 5.12 \end{pmatrix}, \quad \hat{J}_{\text{JMS}}^{(4)} = \begin{pmatrix} 1.79 & 0 & 0 & 0 \\ 0 & 1.79 & 0 & 0 \\ 0 & 0 & 2.43 & 2.10 \\ 0 & 0 & 21.15 & 3.18 \end{pmatrix}, \quad (57)$$

$$\hat{J}_{\text{BMU}}^{(5)} = \begin{pmatrix} 1.63 & 0 & 0 & 0 \\ 0 & 1.63 & 0 & 0 \\ 0 & 0 & -1.17 & -1.39 \\ 0 & 0 & -17.04 & 7.96 \end{pmatrix}, \quad \hat{J}_{\text{BMU}}^{(4)} = \begin{pmatrix} 1.79 & 0 & 0 & 0 \\ 0 & 1.79 & 0 & 0 \\ 0 & 0 & -1.10 & -1.39 \\ 0 & 0 & -21.15 & 6.71 \end{pmatrix}. \quad (58)$$

References

- [1] A. J. Buras, *Gauge Theory of Weak Decays*. Cambridge University Press, 6, 2020.
- [2] E. E. Jenkins, A. V. Manohar, and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Operators and Matching*, *JHEP* **03** (2018) 016, [[arXiv:1709.04486](#)].
- [3] W. Dekens and P. Stoffer, *Low-energy effective field theory below the electroweak scale: matching at one loop*, *JHEP* **10** (2019) 197, [[arXiv:1908.05295](#)].
- [4] J. Aebischer, A. Crivellin, M. Fael, and C. Greub, *Matching of gauge invariant dimension-six operators for $b \rightarrow s$ and $b \rightarrow c$ transitions*, *JHEP* **05** (2016) 037, [[arXiv:1512.02830](#)].
- [5] C. Bobeth, A. J. Buras, A. Celis, and M. Jung, *Yukawa enhancement of Z -mediated new physics in $\Delta S = 2$ and $\Delta B = 2$ processes*, *JHEP* **07** (2017) 124, [[arXiv:1703.04753](#)].
- [6] T. Hurth, S. Renner, and W. Shepherd, *Matching for FCNC effects in the flavour-symmetric SMEFT*, *JHEP* **06** (2019) 029, [[arXiv:1903.00500](#)].
- [7] M. Endo, T. Kitahara, and D. Ueda, *SMEFT top-quark effects on $\Delta F = 2$ observables*, *JHEP* **07** (2019) 182, [[arXiv:1811.04961](#)].
- [8] B. Grzadkowski and M. Misiak, *Anomalous Wtb coupling effects in the weak radiative B -meson decay*, *Phys. Rev.* **D78** (2008) 077501, [[arXiv:0802.1413](#)].
- [9] E. E. Jenkins, A. V. Manohar, and P. Stoffer, *Low-Energy Effective Field Theory below the Electroweak Scale: Anomalous Dimensions*, *JHEP* **01** (2018) 084, [[arXiv:1711.05270](#)].
- [10] J. Aebischer, M. Fael, C. Greub, and J. Virto, *B physics Beyond the Standard Model at One Loop: Complete Renormalization Group Evolution below the Electroweak Scale*, *JHEP* **09** (2017) 158, [[arXiv:1704.06639](#)].

- [11] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators I: Formalism and lambda Dependence*, *JHEP* **10** (2013) 087, [[arXiv:1308.2627](#)].
- [12] E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators II: Yukawa Dependence*, *JHEP* **01** (2014) 035, [[arXiv:1310.4838](#)].
- [13] R. Alonso, E. E. Jenkins, A. V. Manohar, and M. Trott, *Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology*, *JHEP* **04** (2014) 159, [[arXiv:1312.2014](#)].
- [14] A. J. Buras, M. Misiak, and J. Urban, *Two loop QCD anomalous dimensions of flavor changing four quark operators within and beyond the standard model*, *Nucl. Phys.* **B586** (2000) 397–426, [[hep-ph/0005183](#)].
- [15] J. Aebischer, C. Bobeth, A. J. Buras, J. Kumar, and M. Misiak, *General non-leptonic $\Delta F = 1$ WET at the NLO in QCD*, *JHEP* **11** (2021) 227, [[arXiv:2107.10262](#)].
- [16] J. Aebischer, C. Bobeth, A. J. Buras, and J. Kumar, *SMEFT ATLAS of $\Delta F = 2$ transitions*, *JHEP* **12** (2020) 187, [[arXiv:2009.07276](#)].
- [17] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek, *Dimension-Six Terms in the Standard Model Lagrangian*, *JHEP* **1010** (2010) 085, [[arXiv:1008.4884](#)].
- [18] A. J. Buras and P. H. Weisz, *QCD Nonleading Corrections to Weak Decays in Dimensional Regularization and 't Hooft-Veltman Schemes*, *Nucl. Phys.* **B333** (1990) 66–99.
- [19] J. Aebischer et al., *WCxf: an exchange format for Wilson coefficients beyond the Standard Model*, *Comput. Phys. Commun.* **232** (2018) 71–83, [[arXiv:1712.05298](#)].
- [20] A. J. Buras and J.-M. Gérard, *Dual QCD Insight into BSM Hadronic Matrix Elements for $K^0 - \bar{K}^0$ Mixing from Lattice QCD*, [arXiv:1804.02401](#).
- [21] J. Aebischer, A. J. Buras, and J.-M. Gérard, *BSM hadronic matrix elements for ϵ'/ϵ and $K \rightarrow \pi\pi$ decays in the Dual QCD approach*, *JHEP* **02** (2019) 021, [[arXiv:1807.01709](#)].
- [22] A. J. Buras, S. Jäger, and J. Urban, *Master formulae for $\Delta F = 2$ NLO QCD factors in the standard model and beyond*, *Nucl. Phys.* **B605** (2001) 600–624, [[hep-ph/0102316](#)].
- [23] J. Aebischer, A. J. Buras, and J. Kumar, *Simple Rules for Evanescent Operators in One-Loop Basis Transformations*, [arXiv:2202.01225](#).
- [24] J. Aebischer and J. Kumar, *Flavour Violating Effects of Yukawa Running in SMEFT*, *JHEP* **09** (2020) 187, [[arXiv:2005.12283](#)].
- [25] D. R. T. Jones, *Two Loop Diagrams in Yang-Mills Theory*, *Nucl. Phys. B* **75** (1974) 531.
- [26] J. Aebischer, J. Kumar, and D. M. Straub, *Wilson: a Python package for the running and matching of Wilson coefficients above and below the electroweak scale*, *Eur. Phys. J.* **C78** (2018), no. 12 1026, [[arXiv:1804.05033](#)].
- [27] A. Celis, J. Fuentes-Martin, A. Vicente, and J. Virto, *DsixTools: The Standard Model Effective Field Theory Toolkit*, *Eur. Phys. J.* **C77** (2017), no. 6 405, [[arXiv:1704.04504](#)].
- [28] J. Fuentes-Martin, P. Ruiz-Femenia, A. Vicente, and J. Virto, *DsixTools 2.0: The Effective Field Theory Toolkit*, *Eur. Phys. J. C* **81** (2021), no. 2 167, [[arXiv:2010.16341](#)].

-
- [29] A. J. Buras and M. Jung, *Analytic inclusion of the scale dependence of the anomalous dimension matrix in Standard Model Effective Theory*, *JHEP* **06** (2018) 067, [[arXiv:1804.05852](#)].