

DEFORMATIONS OF SOME LOCAL CALABI-YAU MANIFOLDS

ROBERT FRIEDMAN AND RADU LAZA

ABSTRACT. We study deformations of certain crepant resolutions of isolated rational Gorenstein singularities. After a general discussion of the deformation theory, we specialize to dimension three and consider examples which are good (log) resolutions as well as the case of small resolutions. We obtain some partial results on the classification of canonical threefold singularities that admit good crepant resolutions. Finally, we study a non-crepant example, the blowup of a small resolution whose exceptional set is a smooth curve.

INTRODUCTION

This paper is part of a series [FL22, FL24, FL23] investigating the deformation theory of singular Calabi-Yau varieties, i.e. compact analytic spaces Y with isolated Gorenstein canonical singularities such that $\omega_Y \cong \mathcal{O}_Y$, building on and generalizing previously known results due to Kawamata, Namikawa, Namikawa-Steenbrink, and others [Fri86], [Kaw92], [NS95], [Nam94, Nam02]. The deformation theory of Y as studied in [FL22] has both a local and a global aspect. Locally, if $x \in Y$ is a singular point, we can study the deformation functor $\mathbf{Def}_{(Y,x)}$ of the germ (Y, x) . In particular, the tangent space to this functor, or equivalently the “first order deformations,” i.e. the deformations over the dual numbers $\text{Spec } \mathbb{C}[\varepsilon]$, are classified by a finite dimensional vector space $T_{Y,x}^1$, or equivalently by the corresponding skyscraper sheaf supported at x . Globally, for a compact analytic space Y , there is the deformation functor \mathbf{Def}_Y , whose tangent space is a finite dimensional vector space that we denote by \mathbb{T}_Y^1 . There is a corresponding sheaf T_Y^1 , which is supported on the singular locus of Y . In case the singularities of Y are isolated, \mathbb{T}_Y^1 is a skyscraper sheaf supported at the singular points and the stalk of \mathbb{T}_Y^1 at x is the vector space $T_{Y,x}^1$. There is a natural morphism of deformation functors $\mathbf{Def}_Y \rightarrow \prod_{x \in Y_{\text{sing}}} \mathbf{Def}_{(Y,x)}$. Note that \mathbf{Def}_Y and $\mathbf{Def}_{(Y,x)}$ are pro-represented by germs of analytic spaces and the morphism of functors corresponds to a morphism of germs of analytic spaces. On Zariski tangent spaces, the differential of this morphism of functors or germs of spaces becomes a homomorphism $\mathbb{T}_Y^1 \rightarrow H^0(Y; T_Y^1) = \bigoplus_{x \in Y_{\text{sing}}} T_{Y,x}^1$. For $\dim Y \geq 3$, this morphism is almost never surjective. It is thus important to identify interesting tangent directions in $\bigoplus_{x \in Y_{\text{sing}}} T_{Y,x}^1$ and try to lift these to \mathbb{T}_Y^1 . If the deformations of Y are unobstructed, such first order deformations of Y will come from actual deformations.

Taking a local point of view, let (X, x) be the germ of an isolated Gorenstein canonical singularity (or equivalently an isolated rational Gorenstein singularity [KM98], [Kol97, (11.1)]). We will usually take X to be a *good Stein representative* for the germ (X, x) , i.e. a contractible Stein representative with a unique singular point x . Let $\pi: \widehat{X} \rightarrow X$ be a *good resolution* or a *log resolution*, i.e. $\pi^{-1}(x) = E$ is a divisor with simple normal crossings. The assumption of Gorenstein canonical singularities means that $K_{\widehat{X}} = \pi^* \omega_X \otimes \mathcal{O}_{\widehat{X}}(D)$ for some effective divisor D on \widehat{X} . If $D = 0$, we say that π is a *good crepant resolution* of X . Typical examples of singularities that admit good crepant resolutions are the O_{16} singularities, i.e. affine cones over a smooth cubic surface in \mathbb{P}^3 . More general examples are singularities which are analytically isomorphic to cones over Fano manifolds embedded via the anticanonical line bundle, for example the cone over a smooth hypersurface of degree n in \mathbb{P}^n . A related case is that where there exists a *small resolution* $\pi': X' \rightarrow X$, i.e. a

Research of the second author is supported in part by NSF grant DMS-2101640.

resolution where the fiber of π' over x has dimension 1 (more generally one could also consider the case where the exceptional set has dimension less than $\dim X - 1$). However, at least when X is a local complete intersection and $\dim X \geq 3$, such resolutions can only exist for $\dim X = 3$ (see Remark 4.1). Note that small resolutions $\pi': X' \rightarrow X$ are automatically crepant, in the sense that $K_{X'} \cong \mathcal{O}_{X'}$. Examples of such singularities are A_{2k-1} singularities in dimension 3: these are locally defined by the equation $x^2 + y^2 + z^2 + w^{2k}$. A singularity X admitting a small resolution is terminal, but not (locally) \mathbb{Q} -factorial. However, most canonical singularities do not admit good crepant or small resolutions. For instance, terminal \mathbb{Q} -factorial singularities, such as the A_{2k} singularities in dimension 3 (i.e. those given by $x^2 + y^2 + z^2 + w^{2k+1}$, $k \geq 1$), do not have either a good crepant or a small resolution.

There is a natural subspace of $H^0(X; T_X^1)$ defined as follows: Let $U = X - \{x\} = \widehat{X} - E$, where as above $\pi: \widehat{X} \rightarrow X$ is a good resolution and $E = \pi^{-1}(x)$. In case $\dim X \geq 3$, a theorem of Schlessinger implies that $H^0(X; T_X^1) \cong H^1(U; T_U)$, where T_U is the tangent bundle of U [Sch71, Theorem 2]. By Wahl's theory [Wah76], there is a morphism of functors $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_X$ where the induced map on tangent spaces is the natural restriction map

$$H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(U; T_U) \cong H^0(X; T_X^1).$$

Informally, we can think of the image of $\mathbf{Def}_{\widehat{X}}$ as the *simultaneous resolution locus*, i.e. as the deformations of X which lift to deformations of the resolution \widehat{X} . If $\pi: \widehat{X} \rightarrow X$ is a good crepant resolution, then $\mathbf{Def}_{\widehat{X}}$ is unobstructed by a theorem of Gross [Gro97, Proposition 3.4]. If $\pi': X' \rightarrow X$ is a small resolution, then $\mathbf{Def}_{X'}$ is unobstructed by [Fri86, Proposition 2.1]. If X is a local complete intersection singularity, then \mathbf{Def}_X is always unobstructed (cf. [Loo84, §6]).

In trying to understand the image of the morphism $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_X$, and more concretely the image of $H^1(\widehat{X}; T_{\widehat{X}})$ in $H^0(X; T_X^1)$, and their relevance in the Calabi-Yau case, there are two major obstacles:

- (1) In the local setting, the image of $H^1(\widehat{X}; T_{\widehat{X}})$ in $H^0(X; T_X^1)$, is not a birational invariant, i.e. is not independent of the choice of a good resolution. The possible naturally occurring birationally invariant subspaces of $H^0(X; T_X^1)$ are rather the images of $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E))$ or $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E))$. These images are studied in [FL22, Theorem 2.1(iii)], where we prove that the image of $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1})$ is the same as that of $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E))$, at least in the local complete intersection case or if $\dim X = 3$, and is thus independent of the choice of resolution. However, this image does not seem to have an obvious deformation-theoretic interpretation. One important case where such an interpretation exists is when \widehat{X} is a good crepant resolution of X : in this case $\Omega_{\widehat{X}}^{n-1} \cong T_{\widehat{X}}$ and hence the image of $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E))$ agrees with that of $H^1(\widehat{X}; T_{\widehat{X}})$. A similar result holds in the case of a small resolution (Proposition 4.13(i)).
- (2) In the global setting, where Y is a Calabi-Yau variety, it seems difficult to lift deformations arising in this manner to global deformations of Y . For example, in dimension 3 and for small resolutions Y' , this issue is connected with Clemens' conjecture about smooth rational curves in Y' in the case where Y' is a quintic threefold [Cle87], which is still open and where we have nothing new to add. For another example, the deformations of a quintic threefold Y with an O_{16} singularity $x \in Y$ are versal for the deformations of the isolated singularity at x , i.e. the map $\mathbb{T}_Y^1 \rightarrow H^0(Y; T_Y^1)$ is surjective. Hence, if $\widehat{Y} \rightarrow Y$ is the natural crepant resolution and \widehat{X} is a good Stein representative of the germ (Y, x) with crepant resolution \widehat{X} , the image of $\mathbb{T}_Y^1 \rightarrow H^0(Y; T_Y^1)$ contains the image of $H^1(\widehat{X}; T_{\widehat{X}})$. However, the analogous result fails for hypersurfaces of degree $n + 2$ in \mathbb{P}^{n+1} containing an isolated singularity isomorphic to the cone over a hypersurface of degree n in \mathbb{P}^n for

$n \geq 4$, and in this case “most” of the image of $H^1(\widehat{X}; T_{\widehat{X}})$ fails to lift to \mathbb{T}_Y^1 . Thus, the strategy of [FL22] is to work modulo the deformations induced from a resolution, i.e. with $K = \text{Coker} \left(H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E)) \rightarrow H^0(X; T_X^1) \right) \cong H_E^2(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E))$ (see [FL22, Theorem 2.1(v)]); this has the virtue of globalizing well in the Calabi-Yau and Fano case [FL22, §4, §5].

Nonetheless, the study of $\mathbf{Def}_{\widehat{X}}$ and of $H^1(\widehat{X}; T_{\widehat{X}})$ involves a lot of interesting geometry, and the goal of this paper is to investigate some of this geometry. For the reasons outlined above, we restrict to the local case and (mostly) either to the good crepant case or the case of a small resolution. Finally, for most of our results, we restrict to dimension $n = 3$, where good structural results for canonical singularities exist, via the deformation theory of canonical and log canonical surface singularities. Specifically, by [KSB88], [Kaw88], [KM98, Theorem 5.35], an isolated threefold canonical Gorenstein singularity, not necessarily admitting a crepant resolution, can be realized as the total space of a one parameter deformation of a rational double point or a simple elliptic or cusp surface singularity (or equivalently, an isolated log canonical Gorenstein surface singularity). In particular, as discussed below, semistable models for deformations of cusp singularities are a plentiful source of examples of good crepant resolutions and can be obtained systematically by the methods of [FM83], [Eng18], [EF21].

The contents of this paper are as follows. Section 1 analyzes the deformation theory of good crepant resolutions $\pi: \widehat{X} \rightarrow X$, where $\dim X \geq 3$. The main result is Theorem 1.6, which gives some very general results about the first order deformations of \widehat{X} and their relation to first order deformations of E . In this case, the tangent space to the deformation functor \mathbf{Def}_E is the vector space \mathbb{T}_E^1 and there is a corresponding sheaf T_E^1 . Then Theorem 1.6 relates the tangent space to $\mathbf{Def}_{\widehat{X}}$ and the corresponding obstruction space to the cohomology of the exceptional divisor E and its components E_i . Among other things, we show:

Theorem 0.1. *Let $\pi: \widehat{X} \rightarrow X$ be a good crepant resolution of the isolated rational singularity X , with $n = \dim X \geq 3$, and let $E = \bigcup_{i=1}^r E_i$ be the exceptional divisor of π . Then the maps $\mathbb{T}_E^1 \rightarrow H^0(E; T_E^1)$ and $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^0(E; T_E^1)$ are surjective. In particular, all first order smoothing directions for E are realized via first order deformations of \widehat{X} . The map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective $\iff H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$. In this case, all first order locally trivial deformations of E are realized by first order deformations of \widehat{X} and the divisors E_1, \dots, E_r .*

More precise results require better control of the structure of E , which in turn leads to restricting to the case $\dim X = 3$, the running assumption starting with Section 2.

In dimension 2, Gorenstein canonical singularities are du Val singularities, also called rational double point (RDP), simple, or ADE singularities. Their structure is well understood, as is their deformation theory. The purpose of Sections 2 and 3 is to give a partial classification of the isolated Gorenstein canonical threefold singularities (X, x) which admit good crepant resolutions, 3-dimensional analogues of the ADE case, and to discuss their associated deformation theory. As noted above, there is a close connection between isolated threefold canonical Gorenstein singularities and one parameter smoothings of simple elliptic or cusp singularities. Such smoothings are in turn closely related to degenerations of K3 surfaces (cf. for example [FM83]). This leads us to define *divisors of Type II, Type III₁, and Type III₂* (Definition 2.5 and Figures 1, 2, 3). We then obtain a partial classification of the threefold singularities admitting good crepant resolutions, a result of independent interest:

Theorem 0.2 (= Theorems 3.1 and 3.6). *Let (X, x) be an isolated Gorenstein canonical singularity of dimension 3 with a good crepant resolution $\pi: \widehat{X} \rightarrow X$ and let $E = \pi^{-1}(x)$ be the reduced exceptional divisor.*

- (i) If the general hypersurface section of X passing through x is a simple elliptic singularity, then E is of Type II.
- (ii) If the general hypersurface section of X passing through x is a cusp and ω_E^{-1} is nef and big, then E is of Type III₁ or Type III₂.
- (ii)' If the general hypersurface section S of X passing through x is a cusp and the full inverse image $\pi^{-1}(S)$ has normal crossings, then after a sequence of flops (elementary modifications of type 2), ω_E^{-1} becomes nef and big, hence E is of Type III₁ or Type III₂.

Combining Theorem 0.2 with Theorem 1.6, we are able to obtain a deeper understanding of the various deformation theoretic invariants, especially in the Type II and Type III₁ cases. In particular, the following is a somewhat less precise formulation of Proposition 2.11, Proposition 2.14, and Remark 2.15:

Theorem 0.3. *If E is of Type II and irreducible or if E is of Type III₁, then $H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$ and hence the map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective. However, if E is of Type II and reducible, then the map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is never surjective, and if E is of Type III₂, then the map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is not in general surjective.*

In Section 4, we switch our attention to the case of singularities which admit small resolutions $p: X' \rightarrow X$, technically a much simpler case. Here, we relate $H^1(X'; T_{X'})$ to the birational invariants $H^1(\widehat{X}; \Omega_{\widehat{X}}^2(\log E)(-E))$, $H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$, and $H^1(\widehat{X}; \Omega_{\widehat{X}}^2(\log E))$ arising from a good resolution; these invariants are controlled by the Du Bois invariants $b^{p,q}$ and link invariants $\ell^{p,q}$ introduced by Steenbrink [Ste97]. In particular, we recover some results of Steenbrink regarding the dimension of the versal deformation spaces for such singularities (Remark 4.16) and discuss some interesting examples (Examples 4.17 and 4.18). After the first version of this paper was posted, Sz-Sheng Wang sent us a preprint (now [Wan22]) which has substantial overlap with the material in Section 4.

In the final Section 5, we discuss a non-crepant example of a very special type, the blowup \widehat{X} of a smooth curve C which is the exceptional set of a small resolution $\pi: X' \rightarrow X$. In this case, X is a threefold A_{2n-1} singularity, i.e. defined locally by the equation $x^2 + y^2 + z^2 + w^{2n}$, where we assume $n \geq 2$. The question here is to relate the deformations of \widehat{X} to those of X' and X . In particular, we show the following (Theorem 5.7):

Theorem 0.4. *For the above example, let $(S_{\widehat{X}}, 0)$ and $(S_{X'}, 0)$ be the germs prorepresenting the functors $\mathbf{Def}_{\widehat{X}}$ and $\mathbf{Def}_{X'}$ respectively. Then the induced morphism $S_{\widehat{X}} \rightarrow S_{X'}$ is finite of degree n and its differential at the origin has a one-dimensional kernel and cokernel.*

This kind of example is also relevant to the study of deformations of \mathbb{Q} -factorial terminal threefold singularities such as the A_{2n} singularities in dimension 3. While this example is very specific, it helps to illustrate the difference between the image of $H^1(\widehat{X}; T_{\widehat{X}})$ and the birationally invariant image of $H^1(X'; T_{X'})$. It is also interesting from the perspective of the minimal model program. Generally speaking, the analysis of this paper shadows the steps of the minimal program. Namely, Sections 1–3 roughly parallel the fact that a canonical threefold singularity has a partial crepant (divisorial) resolution with terminal singularities ([KM98, Theorem 6.23]). Similarly, Section 4 is the deformation theoretic counterpart of the statement that a terminal singularity admits a small partial resolution to a terminal \mathbb{Q} -factorial singularity ([KM98, Theorem 6.25]), which cannot be further improved. There is however an important difference between the deformation theoretic point of view and that of the minimal model program: in our arguments we need the partial resolutions to be actual resolutions, i.e. we only consider crepant partial resolutions of a canonical singularity which are smooth, not just terminal. Nonetheless, we believe that the discussion here captures some important general phenomena for these classes of singularities, which in turn will

help to better understand the geometry of the moduli spaces of Calabi-Yau varieties, especially in dimension 3.

Notation and conventions. Throughout the paper, we work with the notations and assumptions made above: (X, x) denotes the germ of an isolated singularity, X is a *good Stein representative* for the germ (X, x) , i.e. a contractible Stein representative with a unique singular point x (cf. [Loo84, §2]), and $\pi: \widehat{X} \rightarrow X$ is a *good resolution*, i.e. $\pi^{-1}(x) = E$ is a divisor with simple normal crossings. Unless otherwise specified, all singular cohomology (including local cohomology) is with \mathbb{C} -coefficients.

Acknowledgement. We would like to thank the referee for a careful reading of our paper, and for extensive comments which helped us to improve it.

1. DEFORMATION THEORY IN THE GOOD CREPANT CASE

We begin with a general definition.

Definition 1.1. Let $\pi: \widehat{X} \rightarrow X$ be a resolution of X . Then π is *equivariant* if $R^0\pi_*T_{\widehat{X}} \cong T_X^0$. By e.g. [Fri86, Lemma 3.1], a small resolution is equivariant. Note that, for $q > 0$, $R^q\pi_*T_{\widehat{X}}$ is a torsion sheaf supported on x .

The resolution $\pi: \widehat{X} \rightarrow X$ is:

- (1) a *good resolution* (sometimes called a *log resolution*) if $\pi^{-1}(x) = E$ is a divisor with simple normal crossings.
- (2) a *good equivariant resolution* if π is good and equivariant. By resolution of singularities, good equivariant resolutions always exist.
- (3) *crepant* if $K_{\widehat{X}} = \pi^*\omega_X$ and hence $K_{\widehat{X}} \cong \mathcal{O}_{\widehat{X}}$, and is a *good crepant resolution* if π is also a good resolution. Thus, with these definitions, a small resolution is crepant but not a good crepant resolution. (Note: If the isolated normal singularity (X, x) admits a not necessarily good resolution $\pi: \widehat{X} \rightarrow X$ with $K_{\widehat{X}} \cong \mathcal{O}_{\widehat{X}}$, then (X, x) is automatically a rational Gorenstein singularity, cf. [Kol97, (11.9)]).

Given a good resolution $\pi: \widehat{X} \rightarrow X$, let $E = \pi^{-1}(x) = \bigcup_{i=1}^r E_i$, where the E_i are smooth divisors in \widehat{X} .

Proposition 1.2. *A crepant resolution is equivariant.*

Proof. We begin by showing:

Lemma 1.3. *Let $\pi: \widehat{X} \rightarrow X$ be a crepant resolution and let $\pi': X' \rightarrow X$ be an arbitrary resolution. Then there is a closed analytic subset \widehat{V} of \widehat{X} of codimension at least 2 and a proper analytic subset V' of X' such that the birational map $X' \dashrightarrow \widehat{X}$ restricts on $X' - V'$ to a surjective morphism $\nu: X' - V' \rightarrow \widehat{X} - \widehat{V}$.*

Proof. By Hironaka's theorem, there is a blowup $f: \widetilde{X} \rightarrow \widehat{X}$ which dominates X' . We can further assume that all centers of blowups lie over the inverse image of the singular point x . Let \widehat{V} be the image in \widehat{X} of the centers of the blowups, hence f is an isomorphism from $\widetilde{X} - f^{-1}(\widehat{V})$ to $\widehat{X} - \widehat{V}$. Moreover, $K_{\widetilde{X}} = \mathcal{O}_{\widetilde{X}}(G)$, where $G = \sum_i n_i G_i$ is a divisor with $n_i > 0$ for all i such that $f(G) \subseteq \widehat{V}$. Since $K_{X'} \cong \mathcal{O}_{X'}(D)$ for an effective divisor D whose image in X is the point x , it follows easily that all of the exceptional divisors for the morphism $\widetilde{X} \rightarrow X'$ are of the form G_i for some i . Thus, if V' is the closure of the union of the images of the G_i which are exceptional for the morphism $\widetilde{X} \rightarrow X'$, there is a surjective morphism $X' - V' \rightarrow \widehat{X} - \widehat{V}$. \square

Remark 1.4. The argument of Lemma 1.3 proves the standard fact that, if in addition X' is also a crepant resolution of X , then X' and \widehat{X} are isomorphic outside a set of codimension two.

Continuing with the proof of Proposition 1.2, we must show that the natural injective map $R^0\pi_*T_{\widehat{X}} \rightarrow T_X^0$ is surjective. Choose an equivariant resolution $\pi': X' \rightarrow X$. If ξ is a local section of T_X^0 , then ξ lifts to a section of $T_{X'}$ over the inverse image of the open set and thus defines a section of $T_{\widehat{X}}$ over the complement of V in the notation of Lemma 1.3. By Hartogs, this section extends to a section $\hat{\xi}$ of $T_{\widehat{X}}$. The image of $\hat{\xi}$ in T_X^0 is then ξ . Thus $R^0\pi_*T_{\widehat{X}} \rightarrow T_X^0$ is surjective and therefore an isomorphism. \square

Suppose now that \widehat{X} is a good, not necessarily crepant resolution of X . There is an exact sequence

$$0 \rightarrow T_{\widehat{X}}(-\log E) \rightarrow T_{\widehat{X}} \rightarrow \bigoplus_i N_{E_i/\widehat{X}} \rightarrow 0.$$

Here $H^1(\widehat{X}; T_{\widehat{X}}(-\log E))$ is the tangent space to the functor of deformations of \widehat{X} keeping the divisors E_i . Let \mathcal{D}^\bullet be the complex given by

$$T_{\widehat{X}} \rightarrow N_{E/\widehat{X}}$$

(in degrees 0, 1 respectively). Then $\mathbb{H}^1(\widehat{X}; \mathcal{D}^\bullet)$ is the tangent space to deformations of \widehat{X} keeping the divisor E (as an effective Cartier divisor). Also, let \mathcal{C}^\bullet be the complex

$$T_{\widehat{X}}|E \rightarrow N_{E/\widehat{X}}.$$

Thus \mathcal{C}^\bullet is the dual complex to the complex $I_E/I_E^2 \rightarrow \Omega_{\widehat{X}}^1|E$, which is quasi-isomorphic to Ω_E^1 , the sheaf of Kähler differentials on E . It follows that $\mathcal{H}^i(\mathcal{C}^\bullet) = T_E^i$, $i = 0, 1$, and $\mathbb{H}^i(E; \mathcal{C}^\bullet) = \text{Ext}^i(\Omega_E^1, \mathcal{O}_E) = \mathbb{T}_E^i$. There is a commutative diagram

$$\begin{array}{ccccccc} & & 0 & & 0 & & \\ & & \downarrow & & \downarrow & & \\ & & T_{\widehat{X}}(-E) & \longequal{\quad} & T_{\widehat{X}}(-E) & & \\ & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & T_{\widehat{X}}(-\log E) & \longrightarrow & T_{\widehat{X}} & \longrightarrow & \bigoplus_i N_{E_i/\widehat{X}} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T_E^0 & \longrightarrow & T_{\widehat{X}}|E & \longrightarrow & N_{E/\widehat{X}} \longrightarrow T_E^1 \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \\ & & 0 & & 0 & & \end{array}$$

In particular, there is always an exact sequence

$$0 \rightarrow \bigoplus_i N_{E_i/\widehat{X}} \rightarrow N_{E/\widehat{X}} \rightarrow T_E^1 \rightarrow 0.$$

Moreover, $\mathcal{H}^0(\mathcal{D}^\bullet) = T_{\widehat{X}}(-\log E)$ and $\mathcal{H}^1(\mathcal{D}^\bullet) = \mathcal{H}^1(\mathcal{C}^\bullet) = T_E^1$. Also, from the hypercohomology spectral sequences, there are exact sequences

$$\begin{aligned} H^0(\widehat{X}; T_{\widehat{X}}) &\rightarrow H^0(E; N_{E/\widehat{X}}) \rightarrow \mathbb{H}^1(\widehat{X}; \mathcal{D}^\bullet) \rightarrow H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(E; N_{E/\widehat{X}}); \\ H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) &\rightarrow \mathbb{H}^1(\widehat{X}; \mathcal{D}^\bullet) \rightarrow H^0(E; T_E^1) \rightarrow H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) \end{aligned}$$

as well as the usual exact sequences

$$\begin{aligned} H^0(E; T_{\widehat{X}}|E) &\rightarrow H^0(E; N_{E/\widehat{X}}) \rightarrow \mathbb{T}_E^1 \rightarrow H^1(E; T_{\widehat{X}}|E) \rightarrow H^1(E; N_{E/\widehat{X}}); \\ 0 &\rightarrow H^1(E; T_E^0) \rightarrow \mathbb{T}_E^1 \rightarrow H^0(E; T_E^1) \rightarrow H^2(E; T_E^0) \rightarrow \mathbb{T}_E^2 \rightarrow H^1(E; T_E^1) \rightarrow 0. \end{aligned}$$

Most of the homomorphisms in the above exact sequences have a geometric meaning.

Now suppose \widehat{X} is a good crepant resolution of (X, x) . Then $T_{\widehat{X}}(-\log E)$ is isomorphic to $\Omega_{\widehat{X}}^{n-1}(\log E)(-E)$, so, by the vanishing theorem of Guillén, Navarro Aznar, Pascual-Gainza, Puerta and Steenbrink (see e.g. [PS08, p. 181]), for $p \geq 2$,

$$H^p(\widehat{X}; T_{\widehat{X}}(-\log E)) = H^p(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E)) = 0.$$

Also, $N_{E_i/\widehat{X}} = \omega_{E_i} = K_{E_i}$ and $N_{E/\widehat{X}} = \omega_E$. Moreover, as previously noted, $\mathbf{Def}_{\widehat{X}}$ is unobstructed.

Finally, we have the following result of Steenbrink [Ste83, Lemma 2.14]:

Theorem 1.5 (Steenbrink). *In the above notation, if X has an isolated rational singularity, then $H^i(E; \mathcal{O}_E) = 0$ for $i > 0$.* \square

With these preliminaries, we turn now to the main result of this section:

Theorem 1.6. *Let $\pi: \widehat{X} \rightarrow X$ be a good crepant resolution of the isolated rational singularity X , with $n = \dim X \geq 3$.*

- (i) $\mathbb{H}^1(\widehat{X}; \mathcal{D}^\bullet) \cong H^1(\widehat{X}; T_{\widehat{X}})$. In other words, the first order deformations of \widehat{X} are exactly the first order deformations keeping the effective divisor E .
- (ii) $H^0(E; T_E^1) \cong \bigoplus_i H^1(E_i; N_{E_i/\widehat{X}}) = \bigoplus_i H^1(E_i; \omega_{E_i})$. Thus $H^0(E; T_E^1) = 0 \iff h^{0, n-2}(E_i) = 0$ for every i . (The condition $H^0(E; T_E^1) = 0$ is the condition that all first order deformations of \widehat{X} induce locally trivial first order deformations of E .) More generally, $\dim H^0(E; T_E^1) = \sum_i h^{0, n-2}(E_i)$.
- (iii) If $\dim X = 3$, then $H^1(E; T_E^1)$ has dimension $r - 1$, where r is the number of components of E , and is more intrinsically dual to the cokernel of $H^0(E; \mathcal{O}_E) \rightarrow \bigoplus_i H^0(E_i; \mathcal{O}_{E_i})$. If $\dim X > 3$, then $\dim H^1(E; T_E^1) = \sum_i \dim H^{n-3}(E_i; \mathcal{O}_{E_i})$.
- (iv) $H^2(\widehat{X}; T_{\widehat{X}}) \cong \bigoplus_i H^2(E_i; N_{E_i/\widehat{X}})$ and, if $H^3(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$ (which is always satisfied if $\dim X = 3$), then $H^2(E; T_E^0) = 0$, i.e. all locally trivial first order deformations of E are unobstructed. Thus, in this case,

$$\dim \mathbb{T}_E^2 = \dim H^1(E; T_E^1) = \begin{cases} r - 1, & \text{if } \dim X = 3; \\ \sum_i \dim H^{n-3}(E_i; \mathcal{O}_{E_i}), & \text{if } \dim X > 3. \end{cases}$$

- (v) The maps $\mathbb{T}_E^1 \rightarrow H^0(E; T_E^1)$ and $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^0(E; T_E^1)$ are surjective. In particular, all first order smoothing directions for E are realized via first order deformations of \widehat{X} .
- (vi) The map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective $\iff H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^1(E; T_E^0)$ is surjective $\iff H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$. In this case, all first order locally trivial deformations of E are realized by first order deformations of \widehat{X} and the divisors E_1, \dots, E_r .

Proof. (i) By adjunction and Serre duality, $H^0(E; N_{E/\widehat{X}}) = H^0(E; \omega_E)$ is dual to $H^{n-1}(E; \mathcal{O}_E) = 0$. Likewise $H^1(E; N_{E/\widehat{X}}) = H^1(E; \omega_E)$ is dual to $H^{n-2}(E; \mathcal{O}_E) = 0$. Hence $\mathbb{H}^1(\widehat{X}; \mathcal{D}^\bullet) \cong H^1(\widehat{X}; T_{\widehat{X}})$.

(ii) There is an exact sequence

$$H^0(E; N_{E/\widehat{X}}) \rightarrow H^0(E; T_E^1) \rightarrow \bigoplus_i H^1(E_i; N_{E_i/\widehat{X}}) \rightarrow H^1(E; N_{E/\widehat{X}}).$$

As in (i), $H^0(E; N_{E/\widehat{X}}) = H^1(E; N_{E/\widehat{X}}) = 0$. Thus $H^0(E; T_E^1) \cong \bigoplus_i H^1(E_i; N_{E_i/\widehat{X}})$. But $H^1(E_i; N_{E_i/\widehat{X}}) = H^1(E_i; \omega_{E_i})$ is Serre dual to $H^{n-2}(E_i; \mathcal{O}_{E_i})$ and therefore has dimension $h^{0, n-2}(E_i)$. In particular, $H^0(E; T_E^1) = 0 \iff h^{0, n-2}(E_i) = 0$ for all i .

(iii) Continuing with the above exact sequence, we have

$$0 = H^1(E; N_{E/\widehat{X}}) \rightarrow H^1(E; T_E^1) \rightarrow \bigoplus_i H^2(E_i; N_{E_i/\widehat{X}}) \rightarrow H^2(E; N_{E/\widehat{X}}) \rightarrow 0.$$

If $\dim X = 3$, then $H^1(E; T_E^1)$ is dual to the cokernel of $H^0(E; \mathcal{O}_E) \rightarrow \bigoplus_i H^0(E_i; \mathcal{O}_{E_i})$ and hence has dimension $r - 1$. If $\dim X > 3$, then $H^2(E_i; N_{E_i/\widehat{X}}) = H^2(E_i; \omega_{E_i})$ is Serre dual to $H^{n-3}(E_i; \mathcal{O}_{E_i})$ and $H^2(E; N_{E/\widehat{X}})$ is Serre dual to $H^{n-3}(E; \mathcal{O}_E) = 0$. Thus $\dim H^1(E; T_E^1) = \sum_i \dim H^{n-3}(E_i; \mathcal{O}_{E_i})$ if $\dim X > 3$.

(iv) The statement about $H^2(\widehat{X}; T_{\widehat{X}})$ follows from the exact sequence

$$H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0 \rightarrow H^2(\widehat{X}; T_{\widehat{X}}) \rightarrow \bigoplus_i H^2(E_i; N_{E_i/\widehat{X}}) \rightarrow H^3(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0.$$

The long exact sequence associated to

$$0 \rightarrow T_{\widehat{X}}(-E) \rightarrow T_{\widehat{X}}(-\log E) \rightarrow T_E^0 \rightarrow 0$$

gives rise to an exact sequence

$$H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^2(E; T_E^0) \rightarrow H^3(\widehat{X}; T_{\widehat{X}}(-E)).$$

As we have seen, $H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$, and $H^3(\widehat{X}; T_{\widehat{X}}(-E)) = 0$ if $\dim X = 3$ for dimension reasons. Thus $H^2(E; T_E^0) = 0$.

(v) There is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) & \longrightarrow & H^1(\widehat{X}; T_{\widehat{X}}) & \longrightarrow & \bigoplus_i H^1(N_{E_i/\widehat{X}}) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \cong \\ 0 & \longrightarrow & H^1(E; T_E^0) & \longrightarrow & \mathbb{T}_E^1 & \longrightarrow & H^0(E; T_E^1) \longrightarrow H^2(E; T_E^0). \end{array}$$

(Here, the top right hand arrow is surjective because $H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$.) Thus the induced map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^0(E; T_E^1)$ is surjective, and therefore the map $\mathbb{T}_E^1 \rightarrow H^0(E; T_E^1)$ is surjective as well.

(vi) From the diagram in (v), the map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective $\iff H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^1(E; T_E^0)$ is surjective. Since $H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$, the cokernel of $H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^1(E; T_E^0)$ is $H^2(\widehat{X}; T_{\widehat{X}}(-E))$. Thus $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective $\iff H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$. \square

Remark 1.7. (i) In the situation of (v), it follows from [FL22, Theorem 2.1(iii)] that, for a crepant isolated rational singularity,

$$H^1(\widehat{X}; T_{\widehat{X}}) \cong H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}) \cong H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E)) \oplus H_E^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}).$$

Moreover, $H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}(\log E)(-E)) \cong H^1(\widehat{X}; T_{\widehat{X}}(-\log E))$. The induced map

$$H_E^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}) \rightarrow H^1(\widehat{X}; \Omega_{\widehat{X}}^{n-1}) \rightarrow H^1(E; \Omega_E^{n-1}/\tau_E^{n-1}) \cong \bigoplus_i H^1(E_i; \omega_{E_i}),$$

is an isomorphism, and thus by (v) there is a splitting

$$H^1(\widehat{X}; T_{\widehat{X}}) \cong H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \oplus H^0(E; T_E^1).$$

Note that, while $H^1(\widehat{X}; T_{\widehat{X}}(-\log E))$ and $H^1(\widehat{X}; T_{\widehat{X}})$ have the same image in $H^0(X; T_X^1)$, this is just a statement about the differential of the corresponding morphism of deformation functors $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_X$, and it is reasonable to ask if the actual morphism of deformation functors is finite (meaning that the corresponding morphism of the analytic germs which prorepresent them is finite).

(ii) If $\dim X = 3$ and E is smooth, we will show in the next section that $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ is surjective. However, it is not in general an isomorphism, for example in case X is an O_{16} singularity, the cone over a smooth cubic surface E . A calculation shows that, in this case, $\dim H^0(X; T_X^1) = 16$ and that $\ell = 6$, where $\ell = \ell^{1,1}$ is the link invariant of [Ste97]. By [Wah76], $H^0(X; R^1\pi_*T_{\widehat{X}}) \cong H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$ is the nonnegative part of the deformations of O_{16} and is easily computed to have dimension 5. But $\dim H^1(E; T_E) = 4$, so that $H^0(X; R^1\pi_*T_{\widehat{X}}) \rightarrow H^1(E; T_E)$ is not injective. Here, the weight zero piece of $H^0(X; T_X^1)$ has dimension four and corresponds to deformations of the cubic surface, hence gives all of the first order deformations of E . However, there is also a weight one piece of dimension one. Starting with the cone over the Fermat cubic surface E , for example, the general weight one deformation is given by $f_t = x^3 + y^3 + z^3 + w^3 + txyzw$, defining the singularity X_t . A calculation shows that, for $t \neq 0$, $\dim H^0(X; T_{X_t}^1) = 15$, and hence that the invariant a of [FL22, Theorem 2.1(iv)] is not 0 in this case. In fact, $a = 1$, and the spectral sequence with E_1 page $E_1^{p,q} = H_E^q(\widehat{X}_t; \Omega_{\widehat{X}_t}^p) \implies H_E^{p+q}(\widehat{X}_t)$ does not degenerate at E_2 for $t \neq 0$.

Finally, if E is not smooth, then $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow \mathbb{T}_E^1$ typically fails to be surjective. We will give a geometric explanation for this failure in the next section.

2. THE GOOD CREPANT CASE: SOME CLASSES OF EXAMPLES

In this section, $\pi: \widehat{X} \rightarrow X$ denotes a good crepant resolution of the rational, Gorenstein, isolated singularity X , with $\dim X = 3$. Denote the exceptional divisor $E = \pi^{-1}(x) = \bigcup_{i \geq 1} E_i$. We fix the following notation: $D_{ij} = E_i \cap E_j$, $D_i = E_i \cap \left(\bigcup_{j \neq i} E_j \right) = \bigcup_{j \neq i} D_{ij}$, and $D = \bigcup_i D_i = E_{\text{sing}}$. Our goal is to describe in more detail the case where E looks like a K -trivial semistable degeneration of $K3$ surfaces with one component supporting a negative definite anticanonical divisor removed, or more generally a K -trivial semistable resolution of a one parameter smoothing of a simple elliptic or cusp singularity, minus the component containing the resolution of the germ of the singularity. Note that by [FM83], [Eng18], [EF21], there is a procedure for generating all such examples. There is also an easy criterion for determining the multiplicity and hence for deciding when such examples are good crepant resolutions of complete intersection or even hypersurface singularities.

Before we begin our discussion, we record the following, which holds for an isolated rational singularity X of dimension 3, not necessarily crepant:

Proposition 2.1. *Let $\pi: \widehat{X} \rightarrow X$ be a good resolution of the isolated rational singularity (X, x) of dimension 3, with exceptional divisor $E = \pi^{-1}(x) = \bigcup_{i \geq 1} E_i$, $D_{ij} = E_i \cap E_j$, and triple points t_{ijk} .*

- (i) *For $i > 0$, $H^i(E; \mathcal{O}_E) = 0$ and $H^i(|\Gamma|) = 0$ for $i > 0$, where Γ is the dual complex of E .*
- (ii) *The two homomorphisms $\bigoplus_i H^1(E_i; \mathcal{O}_{E_i}) \rightarrow \bigoplus_{i < j} H^1(D_{ij}; \mathcal{O}_{D_{ij}})$ and $\bigoplus_i H^0(E_i; \Omega_{E_i}^1) \rightarrow \bigoplus_{i < j} H^0(D_{ij}; \Omega_{D_{ij}}^1)$ are isomorphisms.*
- (iii) *Let Ω_E^1 be the sheaf of Kähler differentials and τ_E^1 the subsheaf of torsion differentials. Then $H^0(E; \Omega_E^1/\tau_E^1) = 0$ and $H^1(E) = 0$.*
- (iv) *The map $\bigoplus_i H^2(E_i) \rightarrow \bigoplus_{i < j} H^2(D_{ij})$ is surjective and its kernel is $H^2(E)$. Hence $b_2(E) = \sum_{i=1}^r b_2(E_i) - \#\{\text{double curves}\}$.*
- (v) *Let L be the link of the singularity (X, x) and let ℓ be the dimension of $H^2(L)$ or equivalently $H^3(L)$. Then $H^2(L)$ is a pure Hodge structure and $H^2(L) = H^{1,1}(L)$, so that $\ell = \ell^{1,1}$, where*

$\ell^{1,1}$ is the link invariant of [Ste97]. Finally,

$$\ell = b_2(E) - r = \sum_i b_2(E_i) - r - \#\{\text{double curves}\} = \sum_i (b_2(E_i) - 1) - \#\{\text{double curves}\}.$$

Proof. We shall just sketch the proof. By Theorem 1.5, $H^i(E; \mathcal{O}_E) = 0$ for $i > 0$. The weight spectral sequence for E degenerates at E_2 , and hence so does the Mayer-Vietoris spectral sequence for \mathcal{O}_E . This degeneration, along with the fact that $H^i(E; \mathcal{O}_E) = 0$, $i = 1, 2$, implies that $H^i(|\Gamma|) = 0$ for $i > 0$ and that $\bigoplus_i H^1(E_i; \mathcal{O}_{E_i}) \rightarrow \bigoplus_{i < j} H^1(D_{ij}; \mathcal{O}_{D_{ij}})$ is an isomorphism. Then $\bigoplus_i H^0(E_i; \Omega_{E_i}^1) \rightarrow \bigoplus_{i < j} H^0(D_{ij}; \Omega_{D_{ij}}^1)$ is an isomorphism as well, as follows by taking complex conjugation. This proves (i) and (ii). There is an exact sequence

$$0 \rightarrow \Omega_E^1 / \tau_E^1 \rightarrow \bigoplus_i \Omega_{E_i}^1 \rightarrow \bigoplus_{i < j} \Omega_{D_{ij}}^1 \rightarrow 0.$$

Thus, by (ii), $H^0(E; \Omega_E^1 / \tau_E^1) = 0$. There is a spectral sequence converging to $H^{p+q}(E)$ with $E_1^{p,q} = H^q(E; \Omega_E^p / \tau_E^p)$. As $H^0(E; \Omega_E^1 / \tau_E^1) = H^1(E; \mathcal{O}_E) = 0$, $H^1(E) = 0$, proving (iii). The semipurity theorem (see e.g. [Ste83, Corollary 1.12]) implies that the mixed Hodge structure on $H^k(E)$ is pure for $k \geq 3$. An examination of the weight spectral sequence shows that this forces the map $\bigoplus_i H^2(E_i) \rightarrow \bigoplus_{i < j} H^2(D_{ij})$ to be surjective and identifies $H^2(E)$ with its kernel. Thus $b_2(E) = \sum_{i=1}^r b_2(E_i) - \#\{\text{double curves}\}$, which is (iv). By the link exact sequence and semipurity ([Ste83, Corollary 1.12] again), there is an exact sequence of mixed Hodge structures

$$0 \rightarrow H_E^2(\widehat{X}) \rightarrow H^2(E) \rightarrow H^2(L) \rightarrow 0,$$

where as in the statement of (v) L is the link of the isolated singularity. By duality, $\dim H_E^2(\widehat{X}) = \dim H^4(E) = \sum_i H^4(E_i) = r$. Thus

$$\ell = \dim H^2(L) = b_2(E) - r = \sum_{i=1}^r b_2(E_i) - \#\{\text{double curves}\} - r,$$

using (iv). The above exact sequence also shows that $H^2(L)$ is a pure Hodge structure and that $H^2(L) = H^{1,1}(L)$, hence $\ell = \ell^{1,1}$. This establishes all of the statements of (v). \square

Remark 2.2. Kawamata [Kaw88, p. 97] introduced an invariant $\sigma(Y)$, the *defect* of Y , for a normal projective variety measuring the failure of \mathbb{Q} -factoriality for Y . For a rational singularity (X, x) in dimension 3, there is a local analogue $\sigma(x) = \dim H^2(\widehat{X}) / \sum_i \mathbb{C}[E_i]$ defined by Namikawa-Steenbrink [NS95, (3.9)]. It is immediate to see that $\sigma(x) = \ell$ via the link exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_E^2(\widehat{X}) & \longrightarrow & H^2(E) & \longrightarrow & H^2(L) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow \cong & & \\ & & \bigoplus_i H_4(E_i) & \longrightarrow & H^2(\widehat{X}) & & \end{array}$$

The relationship between the local and global invariants is as follows. If Y is a compact complex threefold with isolated rational singularities, for each $y \in Y_{\text{sing}}$ we have the link L_y and the local invariant $\ell_y = \dim H^2(L_y) = \dim H^3(L_y)$. Let T be the kernel of the natural map $\bigoplus_{y \in Y_{\text{sing}}} H^3(L_y) \rightarrow H^4(Y)$ and let $\sigma(Y)$ be the dimension of the image of $\bigoplus_{y \in Y_{\text{sing}}} H^3(L_y) \rightarrow H^4(Y)$. Thus clearly

$$\dim T = \sum_{y \in Y_{\text{sing}}} \ell_y - \sigma(Y).$$

A straightforward argument with Mayer-Vietoris and semipurity shows that

$$\sigma(Y) = b_4(Y) - b_2(Y).$$

Thus the defect measures the extent to which Poincaré duality fails on Y .

For the remainder of this section, we return to the case where X has a crepant resolution $\pi: \widehat{X} \rightarrow X$. Following Reid [Rei80, Definition 2.5], we make the following definition:

Definition 2.3. Let (X, x) be the germ of an isolated singularity. We say that *Property P holds for a general $t \in \mathfrak{m}_x$, or for the general hypersurface section of X defined by t* , if there exists a finite dimensional subspace V of \mathfrak{m}_x , mapping surjectively onto $\mathfrak{m}_x/\mathfrak{m}_x^2$, such that Property P holds for all t in a nonempty Zariski open subset of V .

With that said, we have the following, essentially due to Kollár-Shepherd-Barron [KSB88] and Kawamata [Kaw88]:

Theorem 2.4. *Let (X, x) be an isolated rational Gorenstein singularity of dimension 3 and let S be a hypersurface section of X passing through x defined by a general $t \in \mathfrak{m}_x$ in the sense of Definition 2.3. Then S has a du Val, a simple elliptic or a cusp singularity.* \square

Since an isolated compound du Val singularity is terminal, the du Val case is excluded if X has a crepant resolution. Define E_0 to be the proper transform of S on \widehat{X} . Thus $E_0 \rightarrow S$ is birational but the inverse image of the singular point is $\bigcup_{i \geq 1} (E_0 \cap E_i)$. If S is defined by the function $t \in \mathfrak{m}_{X,x}$, then as divisors $(\pi^*t) = E_0 + \sum_{i \geq 1} a_i E_i$, where a_i is an integer ≥ 1 . Hence

$$\mathcal{O}_{\widehat{X}}\left(E_0 + \sum_{i \geq 1} a_i E_i\right) \cong \mathcal{O}_{\widehat{X}}.$$

By the crepant assumption, $K_{E_i} = \mathcal{O}_{\widehat{X}}(E_i)|_{E_i} = \mathcal{O}_{E_i}(E_i)$, including the case $i = 0$ where we might have to replace K_{E_0} by ω_{E_0} , and $\omega_E = \mathcal{O}_{\widehat{X}}(E)|_E = \mathcal{O}_E(E)$.

Motivated by the description of semistable smoothings of simple elliptic and cusp singularities (cf. [FM83] and Remark 2.6 below), we make the following definition:

Definition 2.5. A compact analytic surface $E = \bigcup_{i \geq 1} E_i$ with simple normal crossings is *Type II* (Figure 1) if, after relabeling, $E = \bigcup_{i=1}^r E_i$ with:

- (i) E_1, \dots, E_{r-1} are elliptic ruled (not necessarily minimally ruled) and E_r is rational, with $-K_{E_r}$ nef and big. (Here a smooth surface S is a not necessarily minimally ruled surface over a base curve D if there exists a morphism $S \rightarrow D$ whose generic fiber is \mathbb{P}^1 , and it is elliptic ruled if D is an elliptic curve.)
- (ii) The dual complex is a line segment with adjacent vertices E_1, \dots, E_r , or a single point if $r = 1$.
- (iii) With $D_{i,i+1} = E_i \cap E_{i+1}$, $1 \leq i \leq r-1$ as above, $D_{i,i+1}$ is a smooth elliptic curve.
- (iv) There exists a smooth elliptic curve $D_0 \subseteq E_1$, disjoint from D_{12} , such that, in case $r > 1$, $K_{E_1} = \mathcal{O}_{E_1}(-D_0 - D_{12})$, $K_{E_i} = \mathcal{O}_{E_i}(-D_{i-1,i} - D_{i,i+1})$ for $1 \leq i \leq r-1$, and $K_{E_r} = \mathcal{O}_{E_r}(-D_{r-1,r})$, and $K_{E_1} = \mathcal{O}_{E_1}(-D_0)$ in case $r = 1$.
- (v) $\mathcal{O}_E(E) = \omega_E = \mathcal{O}_E(-D_0)$.

Note that (v) implies that $\mathcal{O}_{D_{i,i+1}}(E) \cong \mathcal{O}_{D_{i,i+1}}$, and thus $T_E^1 \cong \bigoplus_i \mathcal{O}_{\widehat{X}}(E)|_{D_{i,i+1}} \cong \mathcal{O}_D$, where $D = \bigcup_{i=1}^{r-1} D_{i,i+1} = E_{\text{sing}}$.

The compact analytic surface $E = \bigcup_{i=1}^r E_i$ with simple normal crossings is *Type III₁* (Figure 2) if, possibly after relabeling:

- (i) All of the E_i are smooth rational.
- (ii) $E_i \cap E_j = \emptyset$ unless $j = \pm i$. In particular, the dual complex of E is a point (if $r = 1$) or a line segment (if $r \geq 2$).




FIGURE 1. Exceptional divisor E of Type II

- (iii) Assume that $r \geq 2$. Let $D_{i,i+1} = E_i \cap E_{i+1}$. Then $D_{i,i+1}$ is a smooth rational curve. For $i = 1, r$, there exist connected curves C_1, C_r on E_1, E_r respectively, where C_1 and C_r are chains of smooth rational curves meeting transversally, with $(C_1 \cdot D_{1,2})_{E_1} = (C_r \cdot D_{r-1,r})_{E_r} = 2$, and we have: $K_{E_1} = \mathcal{O}_{E_1}(-C_1 - D_{1,2})$ and $K_{E_r} = \mathcal{O}_{E_r}(-C_r - D_{r-1,r})$. For $2 \leq i \leq r-1$, there exist two curves C'_i and C''_i , both chains of smooth rational curves, supporting effective divisors of self-intersection 0, and not containing $D_{i-1,i}$ or $D_{i,i+1}$, with

$$(C'_i \cdot D_{i-1,i})_{E_i} = (C''_i \cdot D_{i-1,i})_{E_i} = (C'_i \cdot D_{i,i+1})_{E_i} = (C''_i \cdot D_{i,i+1})_{E_i} = 1,$$

such that

$$K_{E_i} = \mathcal{O}_{E_i}(-D_{i-1,i} - C'_i - D_{i,i+1} - C''_i).$$

Moreover, $C = C_1 + C'_2 + C''_2 + \cdots + C'_{r-1} + C''_{r-1} + C_r$ is a Cartier divisor on E , where $C' = \sum_{i=2}^{r-1} C'_i$ and $C'' = \sum_{i=2}^{r-1} C''_i$ are connected.

- (iv) In all cases, $\mathcal{O}_E(E) = \omega_E = \mathcal{O}_E(-C)$, where, if $r = 1$, C is a cycle of smooth rational curves or an irreducible nodal curve.



FIGURE 2. Exceptional divisor E of Type III_1

Note that (iv) implies that there exist points p'_i, p''_i in $D_{i,i+1}$ for $1 \leq i \leq r-1$, which are smooth points on D , and such that $T_E^1 \cong \mathcal{O}_D(-\sum_i(p'_i + p''_i))$.

The compact analytic surface $E = \bigcup_{i=1}^r E_i$ with simple normal crossings is Type III_2 (Figure 3) if $r \geq 2$ and, possibly after relabeling:

- (i) All of the E_i are smooth rational.
- (ii) There are distinguished components E_1, \dots, E_s such that $E_i \cap E_j = \emptyset$ unless $j = \pm i \bmod s$. In particular, for $s \geq 3$, the dual complex of E_1, \dots, E_s is a circle (and similarly for $r = 2$

where the dual complex has 2 vertices joined by 2 edges). Moreover, there exist chains of smooth rational curves C_i on E_i , $1 \leq i \leq s$ such that C_i and D_i have no component in common, and $C_i \cap E_j \neq \emptyset \iff 1 \leq j \leq s$ and $j = \pm 1 \pmod s$. In this case, $(C_i \cdot D'_{i,i+1})_{E_i} = 1$ for every component $D'_{i,i+1}$ of $D_{i,i+1}$ meeting E_0 , with the convention that $D_{s,s+1} = D_{1,s}$. Moreover, $C = C_1 + \dots + C_s$ is a Cartier divisor on E contained in $\bigcup_{i=1}^s E_i$, more precisely $C_i \cap E_{i+1} = C_{i+1} \cap E_i$.

- (iii) For $1 \leq i \leq s$, $C_i + D_i$ is a cycle of smooth rational curves, and $K_{E_i} = \mathcal{O}_{E_i}(-C_i - D_i)$. For $i > s$, D_i is a cycle of smooth rational curves, and $K_{E_i} = \mathcal{O}_{E_i}(-D_i)$.
- (iv) The dual complex of E is a semi-simplicial triangulation of the 2-disk, and E_1, \dots, E_r are the boundary vertices.
- (v) $\mathcal{O}_E(E) = \omega_E = \mathcal{O}_E(-\sum_{i=1}^s C_i)$.




FIGURE 3. Exceptional divisor E of Type III_2

Note that (v) implies the following (keeping the convention that $D_{s,s+1} = D_{1,s}$): Suppose $D_{i,i+1}$ is irreducible. Then there exist points p_i in $D_{i,i+1}$ for $1 \leq i \leq s$, not in D_{sing} , such that $T_E^1 \cong \mathcal{O}_D(-\sum_i p_i)$. An analogous statement holds if $D_{i,i+1}$ is reducible, where such points exist in every component of $D_{i,i+1}$ meeting C_i .

Remark 2.6. The above terminology is modeled on the corresponding terminology for semistable degenerations of $K3$ surfaces due to Persson, Kulikov and others. More precisely, consider a smoothing $\rho: X \rightarrow (\Delta, 0)$, where $X = (X, x)$ is the germ of a simple elliptic or cusp singularity over the unit disk (i.e. ρ is a flat morphism of germs, $\rho^{-1}(0)$ is a simple elliptic or cusp singularity and $\rho^{-1}(t)$ is smooth for $t \neq 0$). By [FM83, Theorem 2.5], possibly after a base change, there exists a semistable crepant resolution $\pi: \widehat{X} \rightarrow X$. In other words, \widehat{X} is smooth and induces a minimal resolution of $\rho^{-1}(0)$ with exceptional divisor D_0 , $K_{\widehat{X}} \cong \mathcal{O}_{\widehat{X}}$, and the induced morphism $f = \rho \circ \pi: \widehat{X} \rightarrow \Delta$ satisfies: $f^{-1}(0)$ is a reduced divisor with simple normal crossings. Then the

dual complex of $f^{-1}(0)$ is a line segment in the simple elliptic case (Type II) or a triangulation of S^2 in the cusp case (Type III). Thus, the dual complex of $E = \pi^{-1}(0)$ is again a line segment in the simple elliptic case, and hence is of Type II in the sense of Definition 2.5, or it obtained by deleting a vertex in a triangulation of S^2 and hence is either a line segment or a disk, i.e. is of Type III₁ or Type III₂ in the sense of Definition 2.5. See also Theorem 3.6 for a partial converse to this picture.

Remark 2.7. As noted in greater generality in [Kaw88], the number r of divisors in the exceptional set E of the crepant resolution $\pi: \widehat{X} \rightarrow X$ is independent of the choice of crepant resolution. This follows easily from the fact that any two crepant resolutions are isomorphic in codimension one (see e.g. Remark 1.4). The invariant $\ell = \dim H^2(L)$ defined in Proposition 2.1(v) can also be computed directly in this case, at least for a semistable smoothing:

Proposition 2.8. (i) *Suppose that \widehat{X} is the semistable model for a smoothing of a simple elliptic singularity, of multiplicity $m = -D_0^2$. Then*

$$\ell = 9 - m.$$

(ii) *Suppose that \widehat{X} is the semistable model for a smoothing of a cusp singularity, of multiplicity $m = -D_0^2$ and length s , the number of components of the cusp. Then*

$$\ell = 9 - m + s.$$

Proof. By Proposition 2.1(v), $\ell = \sum_{i=1}^r b_2(E_i) - \#\{\text{double curves}\} - r$, where r is the number of components of E . In the simple elliptic case, after standard birational operations (flops or Type II elementary modifications), we can assume that E_1, \dots, E_{r-1} are minimal elliptic ruled surfaces and that E_r is a generalized del Pezzo surface of degree $-(D_{r-1,r})_{E_{r-1}}^2 = (D_{r-1,r})_{E_r}^2 = K_{E_r}^2 = m$. Thus there are $r - 1$ double curves, $r - 1$ elliptic ruled components E_i with $b_2(E_i) = 2r$, and the remaining component E_r satisfies: $b_2(E_r) = 10 - m$. Putting this together gives

$$\ell = 2(r - 1) + 10 - m - 2r + 1 = 9 - m.$$

In the cusp case, we shall just write down the proof for E of Type III₂ (the proof in the Type III₁ case is similar but simpler). Let e be the number of double curves of E and let t be the number of triple points. By taking Euler characteristics,

$$r - e + t = 1.$$

Each surface E_i satisfies: $-K_{E_i} = \mathcal{O}_{E_i}(D_i + C_i)$ or $-K_{E_i} = \mathcal{O}_{E_i}(D_i)$, depending on whether E_i meets E_0 , and in this case C_i is irreducible since by assumption \widehat{X} is semistable. Set $\widehat{D}_i = D_i$, if E_i does not meet E_0 (i.e. $i > s$), and $\widehat{D}_i = D_i + C_i$, if E_i meets E_0 (i.e. $1 \leq i \leq s$). If s_i is the number of components of \widehat{D}_i , then $\sum_i s_i = 2e + s$. Every triple point is contained in three edges, and every edge contains two triple points except for the edges corresponding to the double curves $E_i \cap E_{i+1}$, $1 \leq i \leq s$, which just meet one triple point. Thus $2e = 3t + s$.

The surfaces E_i , $i > 0$, are rational surfaces with an anticanonical cycle \widehat{D}_i . Following [FM83, Definition 3.1], define the *charge* $Q(E_i, \widehat{D}_i)$ by:

$$Q(E_i, \widehat{D}_i) = 12 - (\widehat{D}_i)^2 - s_i.$$

Note that this definition makes sense for E_0 as well, where we set $\widehat{D}_0 = D_0 = \sum_{i=1}^r C_i$ and $Q(E_0, \widehat{D}_0) = 12 + m - s$. For $i > 0$, by [Fri15, Lemma 1.2], $b_2(E_i) = Q(E_i, \widehat{D}_i) - 2 + s_i$. Then

$$\begin{aligned} \ell &= \sum_{i \geq 1} b_2(E_i) - e - r = \sum_{i \geq 1} Q(E_i, \widehat{D}_i) - 3r + \sum_{i \geq 1} s_i - e \\ &= \sum_{i \geq 1} Q(E_i, \widehat{D}_i) - 3(1 + e - t) + 2e + s - e \\ &= \sum_{i \geq 1} Q(E_i, \widehat{D}_i) - 3 + (3t - 2e + s) = \sum_{i \geq 1} Q(E_i, \widehat{D}_i) - 3. \end{aligned}$$

The principle of ‘‘conservation of charge’’ [FM83, Proposition 3.7] implies that

$$\sum_{i \geq 1} Q(E_i, \widehat{D}_i) + Q(E_0, \widehat{D}_0) = 24.$$

Thus $\sum_{i \geq 1} Q(E_i, \widehat{D}_i) = 12 - m + s$ and so finally

$$\ell = \sum_{i \geq 1} Q(E_i, \widehat{D}_i) - 3 = 9 - m + s. \quad \square$$

Remark 2.9. Proposition 2.8 implies the well-known results that $m \leq 9$ for a smoothable simple elliptic singularity and $m \leq 9 + s$ for a smoothable cusp singularity.

In the Type II and Type III₁ cases, T_E^1 is uniquely specified as noted in the definition. For the Type III₂ case, T_E^1 is also uniquely specified by Condition (v):

Lemma 2.10. *Suppose that E is of Type III₂. For every i , $1 \leq i \leq s$, and every component $\Gamma_{i\alpha}$ of $D_{i,i+1}$ meeting C_i , choose points $q_{i\alpha} \in \Gamma_{i\alpha}$, not in E_j for $j \neq i$. Then $\mathcal{O}_D(-\sum_{i,\alpha} q_{i\alpha}) \cong T_E^1$.*

Proof. Let $D = D' \cup D''$, where D'' consists of the components of $D_{i,i+1}$ meeting C_i , $1 \leq i \leq s$ with the usual conventions, and D' is the union of all of the other components. For simplicity, we just consider the case where $D_{i,i+1} \cong \mathbb{P}^1$ is irreducible for every i and write q_i instead of $q_{i\alpha}$. Let L be a line bundle on D such that $L|_{D'} \cong \mathcal{O}_{D'}$ and $L|_{D''} \cong \mathcal{O}_{D''}$. We claim that $L \cong \mathcal{O}_D$. Applying this to $\mathcal{O}_D(-\sum_i q_i) \otimes T_E^1 \cong \mathcal{O}_D(p_i - \sum_i q_i)$ then proves the lemma.

More precisely, we shall show that $\text{Pic } D \cong \text{Pic } D' \times \text{Pic } D'' \cong \text{Pic } D' \times \mathbb{Z}^s$, and hence $p_a(D) = p_a(D')$. We have an exact sequence

$$0 \rightarrow \mathcal{O}_D^* \rightarrow \mathcal{O}_{D'}^* \times \mathcal{O}_{D''}^* \rightarrow \prod_{i=1}^s \mathbb{C}_{p_i}^* \rightarrow 0,$$

where $D' \cap D'' = \{p_1, \dots, p_s\}$ and $\mathbb{C}_{p_i}^*$ is the skyscraper sheaf at p_i with stalk \mathbb{C}^* . But $H^0(D''; \mathcal{O}_{D''}^*) \cong (\mathbb{C}^*)^s$ and the induced map

$$H^0(D''; \mathcal{O}_{D''}^*) \rightarrow H^0(D; \prod_{i=1}^s \mathbb{C}_{p_i}^*) \cong (\mathbb{C}^*)^s$$

coming from the above exact sequence is an isomorphism. Thus

$$H^1(D; \mathcal{O}_D^*) = \text{Pic } D \cong H^1(D'; \mathcal{O}_{D'}^*) \times H^1(D''; \mathcal{O}_{D''}^*) = \text{Pic } D' \times \text{Pic } D'',$$

completing the proof. \square

Next we describe the cokernel of $H^1(\widehat{X}; T_{\widehat{X}})$ in $H^1(E; T_E^0)$. As we have seen in the proof of Theorem 1.6(vi), this cokernel is $H^2(\widehat{X}; T_{\widehat{X}}(-E))$, and it measures the failure of a first order locally trivial deformation of E to be realized by a deformation of \widehat{X} and the divisors E_1, \dots, E_r .

Proposition 2.11. *Suppose that E is Type II. Then $\dim H^2(\widehat{X}; T_{\widehat{X}}(-E)) \geq r - 1$, where r is the number of components of E , and $H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$ if $r = 1$, i.e. E is irreducible.*

Remark 2.12. The geometric interpretation of Proposition 2.11 is as follows. By Theorem 1.6, $H^2(\widehat{X}; T_{\widehat{X}}(-E))$ is the obstruction to realizing all first order locally trivial deformations of E by deforming \widehat{X} . We have seen that, in the Type II case, where $E \subseteq \widehat{X}$, we have $T_E^1 \cong \mathcal{O}_D$, where $D = E_{\text{sing}}$. For a general deformation of E , the line bundle T_E^1 on D has degree 0, but is not necessarily the trivial line bundle. Here, $r - 1$ is the number of conditions on the deformation required to keep T_E^1 trivial.

Proof of Proposition 2.11. By the formal functions theorem,

$$H^2(\widehat{X}; T_{\widehat{X}}(-E)) = \varprojlim_n H^2(\widehat{X}; T_{\widehat{X}}(-E)|nE).$$

Since $H^3(\widehat{X}; T_{\widehat{X}}(-2E)) = 0$, $H^2(\widehat{X}; T_{\widehat{X}}(-E)) \rightarrow H^2(E; T_{\widehat{X}}(-E)|E)$ is surjective. For $n \geq 1$, we have the exact sequence

$$0 \rightarrow T_{\widehat{X}}(-E)|E \otimes \mathcal{O}_E(-nE) \rightarrow T_{\widehat{X}}(-E)|(n+1)E \rightarrow T_{\widehat{X}}(-E)|nE \rightarrow 0.$$

By duality (where we also allow the cases $n = 0, -1$), $H^2(E; T_{\widehat{X}}|E \otimes \mathcal{O}_E(-(n+1)E))$ is dual to

$$H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+1)E) \otimes \omega_E) = H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E)).$$

By Theorem 3.1(i), $\mathcal{O}_E(E) = K_E = \mathcal{O}_E(-D_0)$ for an effective nonzero divisor D_0 on E , disjoint from the singular locus, and similarly $\mathcal{O}_E(mE) = \mathcal{O}_E(-mD_0)$. In particular, for $m \geq 0$, there is an inclusion $\mathcal{O}_E(mE) \rightarrow \mathcal{O}_E$ which only vanishes along the divisor D_0 for $m > 0$. Now use the conormal sequence

$$0 \rightarrow I_E/I_E^2 \rightarrow \Omega_{\widehat{X}}^1|E \rightarrow \Omega_E^1 \rightarrow 0.$$

Since $I_E/I_E^2 = \mathcal{O}_E(-E)$, $I_E/I_E^2 \otimes \mathcal{O}_E((n+2)E) = \mathcal{O}_E((n+1)E)$. First assume $r = 1$, i.e. E is smooth. From the conormal sequence, we get an exact sequence

$$H^0(E; \mathcal{O}_E((n+1)E)) \rightarrow H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E)) \rightarrow H^0(E; \Omega_E^1 \otimes \mathcal{O}_E((n+2)E)).$$

Since $H^0(E; \mathcal{O}_E((n+1)E)) = 0$ for all $n \geq 0$, there is a sequence of inclusions

$$H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E)) \rightarrow H^0(E; \Omega_E^1 \otimes \mathcal{O}_E((n+2)E)) \rightarrow H^0(E; \Omega_E^1) = 0.$$

Hence $H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E)) = 0$ for all $n \geq 0$, and therefore $H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$.

Now assume $r \geq 2$. We freely use the notation of Definition 2.5. With notation as in Definition 2.5, $\mathcal{O}_E(mE)|D_{i,i+1} \cong \mathcal{O}_{D_{i,i+1}}$, $\mathcal{O}_E(mE)|E_i \cong \mathcal{O}_{E_i}$ if $i > 1$, and $\mathcal{O}_E(mE)|E_1 \cong \mathcal{O}_{E_1}(-mD_0)$. Using the exact sequence (from the normalization)

$$0 \rightarrow \mathcal{O}_E(mE) \rightarrow \bigoplus_i \mathcal{O}_E(mE)|E_i \rightarrow \bigoplus_i \mathcal{O}_{D_{i,i+1}} \rightarrow 0,$$

the map $\bigoplus_i H^0(E_i; \mathcal{O}_E(mE)|E_i) \rightarrow \bigoplus_i H^0(D_{i,i+1}; \mathcal{O}_{D_{i,i+1}})$ is an isomorphism. Thus, as in the case $r = 1$, $H^0(E; \mathcal{O}_E(mE)) = 0$ for all $m > 0$.

As in the case $r = 1$, we want to analyze $H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E))$. By the above,

$$H^0(E; \Omega_{\widehat{X}}^1|E \otimes \mathcal{O}_E((n+2)E)) \subseteq H^0(E; \Omega_E^1 \otimes \mathcal{O}_E((n+2)E)).$$

Now we have the exact sequence

$$0 \rightarrow \tau_E^1 \rightarrow \Omega_E^1 \rightarrow \Omega_E^1/\tau_E^1 \rightarrow 0.$$

Also, $H^0(E; \Omega_E^1/\tau_E^1) = 0$ by Proposition 2.1(iii). Since there is an inclusion

$$H^0(E; (\Omega_E^1/\tau_E^1) \otimes \mathcal{O}_E((n+2)E)) \subseteq H^0(E; \Omega_E^1/\tau_E^1),$$

we conclude that $H^0(E; (\Omega_E^1/\tau_E^1) \otimes \mathcal{O}_E((n+2)E)) = 0$. The remaining term is

$$H^0(E; \tau_E^1 \otimes \mathcal{O}_E((n+2)E)).$$

As D is smooth and the curve $D_{i,i+1}$ appears twice in $\coprod_i E_i$, by [Fri83, Prop. 1.10(2)], there is an exact sequence

$$0 \rightarrow \mathcal{O}_{\widehat{X}}(-E)|_D \rightarrow \bigoplus_i (\mathcal{O}_{\widehat{X}}(-E)|_{D_{i,i+1}})^2 \rightarrow \tau_E^1 \rightarrow 0.$$

Since D is a disjoint union of the smooth components D_i and $\mathcal{O}_{\widehat{X}}(-E)|_D \cong \mathcal{O}_D$, this says that $\tau_E^1 \cong \mathcal{O}_D = \bigoplus_{i=1}^{r-1} \mathcal{O}_{D_{i,i+1}}$. Hence

$$H^0(E; \tau_E^1 \otimes \mathcal{O}_E((n+2)E)) \cong H^0(D; \mathcal{O}_D) = \mathbb{C}^{r-1}.$$

In particular, taking $n = 1$, we get $\dim H^2(E; T_{\widehat{X}}(-E)|_E) = r - 1$. Thus $\dim H^2(\widehat{X}; T_{\widehat{X}}(-E)) \geq r - 1$. \square

Remark 2.13. If we wanted to fully calculate $\dim H^2(\widehat{X}; T_{\widehat{X}}(-E))$, we would have to understand the coboundary

$$H^1(nE; T_{\widehat{X}}(-E)|_nE) \rightarrow H^2(E; T_{\widehat{X}}(-E)|_E \otimes \mathcal{O}_E(-nE)),$$

where the last part of the proof shows that

$$\dim H^2(E; T_{\widehat{X}}(-E)|_E \otimes \mathcal{O}_E(-nE)) = r - 1$$

for all $n \geq 1$. This seems difficult, and most likely depends on the higher infinitesimal neighborhoods of E in \widehat{X} .

In the Type III₁ case (and also the Type III₂ case if every component of D meets C) then in fact $H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$:

Proposition 2.14. *In the Type III₁ case, $H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$.*

Proof. The proof is similar to the proof of Proposition 2.11, but in this case $\tau_E^1 \otimes \mathcal{O}_E((n+2)E)$ is a line bundle of negative degree on every component of $D = E_{\text{sing}}$. Thus $H^0(E; \tau_E^1 \otimes \mathcal{O}_E((n+2)E)) = 0$. \square

Remark 2.15. A similar (but also ultimately inconclusive) analysis along the lines of Proposition 2.11 is possible in the Type III₂ case: one can show that, in this case,

$$\dim H^0(E; \tau_E^1 \otimes \mathcal{O}_E((n+2)E)) \geq p_a(D),$$

and in particular that $\dim H^2(\widehat{X}; T_{\widehat{X}}(-E)) \geq p_a(D)$. It is easy to see that $p_a(D)$ is not 0 in general. More precisely, using [EF21, Lemma 2.25(iv)], one can check that $p_a(D) = r - s$, which is typically nonzero.

3. THE GOOD CREPANT CASE: A PARTIAL CLASSIFICATION

Let (X, x) be an isolated singularity of dimension 3 with a good crepant resolution $\pi: \widehat{X} \rightarrow X$. It is natural to ask if the (reduced) exceptional divisor E of π is Type II, Type III₁, or Type III₂. We have some partial results along these lines, inspired by the arguments of Shepherd-Barron [SB83a, SB83b]:

Theorem 3.1. *With notation and assumptions as above, let t be a general element of $\mathfrak{m}_{X,x}$ in the sense of Definition 2.3.*

(i) *With $(\pi^*t) = E_0 + \sum_{i \geq 1} a_i E_i$, we have $a_i = 1$ for all i . Hence*

$$\mathcal{O}_{\widehat{X}}\left(\sum_{i \geq 1} E_i\right) \cong \mathcal{O}_{\widehat{X}}(-E_0).$$

- (ii) If the hypersurface section of X defined by t is simple elliptic, then E is of Type II. In particular, all components of E are rational or elliptic ruled.
- (iii) If the hypersurface section of X defined by t is a cusp, then all components of E are rational and every component of a double curve D_{ij} is a smooth rational curve. (The proof gives much more detailed information about this case.)

Proof. We begin with two lemmas.

Lemma 3.2. *Let S be the germ of a simple elliptic singularity or a cusp singularity, with singular point x , and let $p: \tilde{S} \rightarrow S$ be a birational morphism such that \tilde{S} is normal, $p^{-1}(x)$ is a curve, i.e. has dimension one, and $K_{\tilde{S}} = \mathcal{O}_{\tilde{S}}(-D_0)$, where D_0 is an effective nonzero Weil divisor whose support is contained in $p^{-1}(x)$.*

- (i) *If S is simple elliptic, D_0 is a (reduced) smooth elliptic curve, S is Gorenstein, and D_0 is a Cartier divisor with $D_0^2 < 0$.*
- (ii) *If S is a cusp, then D_0 is a reduced cycle of smooth rational curves, or an irreducible nodal curve.*

Proof. First consider the simple elliptic case. There exists a resolution of singularities $T \rightarrow S$ which dominates \tilde{S} , and hence T is the blowup of a minimal resolution of singularities of S , say T_0 with $K_{T_0} = \mathcal{O}_{T_0}(-C)$. Thus T is obtained by successively blowing up points on T_0 . All blowups at points not on the proper transform of C correspond to components of K_T occurring with positive coefficients: as Cartier divisors, $K_T = -C' + \sum_i a_i e_i$, where the e_i are proper transforms of exceptional curves corresponding to blowups at points not on the proper transform of C , and $a_i > 0$. Let D'_0 be the nonzero effective Cartier divisor on T which is the proper transform of D_0 . Then

$$K_T = -C' + \sum_i a_i e_i = -D'_0 + \text{a sum of exceptional fibers of the morphism } T \rightarrow \tilde{S}.$$

The only way this is possible is if all of the e_i are fibers of the morphism $T \rightarrow \tilde{S}$ and C' is not an exceptional fiber of the morphism. It follows that \tilde{S} is dominated by a surface T , a blowup of T_0 where all blowups are at points on the proper transform of C , and C itself is not blown down. Hence \tilde{S} has at worst A_n singularities, so is Gorenstein, and $K_{\tilde{S}} = \omega_{\tilde{S}} = \mathcal{O}_{\tilde{S}}(-D_0)$, where D_0 is the image of the proper transform of C . Thus D_0 is a (reduced) smooth elliptic curve and $D_0^2 < 0$.

A similar argument in the cusp case shows that, if T_0 is the minimal resolution and $K_{T_0} = \mathcal{O}_{T_0}(-\sum_k \Gamma_k)$, where $\sum_k \Gamma_k$ is a cycle of rational curves on T_0 , then \tilde{S} is dominated by a surface T , a blowup of T_0 where all blowups are at points on the proper transform of $\sum_k \Gamma_k$, and $\sum_k \Gamma_k$ itself is not entirely blown down. In particular, the image D_0 of $\sum_k \Gamma_k$ is a reduced cycle of smooth rational curves or an irreducible nodal curve. \square

Lemma 3.3. *Let T be a smooth algebraic surface such that $-K_T = \sum_i a_i D_i$ where the D_i are irreducible curves, $a_i > 0$, and the sum is nonempty. Then either T is an elliptic ruled surface and $-K_T = \sigma' + \sigma''$, where σ', σ'' are disjoint smooth elliptic curves, or $\bigcup_i D_i$ is connected.*

Proof. First, if $\rho: T \rightarrow \bar{T}$ is the blowdown of an exceptional curve, so that \bar{T} is smooth, then it is easy to check that $-K_{\bar{T}} = \sum_i a_i \rho_*(D_i)$ and that T is a blowup of \bar{T} at a point of $\bigcup_i \rho_*(D_i)$. Then $\bigcup_i D_i$ is connected $\iff \bigcup_i \rho_*(D_i)$ is connected. Thus, we may as well assume that T is minimal, and hence is either \mathbb{P}^2 , where the result is automatic, or the blowup of a ruled surface over a curve of genus g . In this case, using [Har77] as a general reference, let e be the invariant of the ruled surface, i.e. $-e$ is the minimal self-intersection of a curve on T , and let σ_0 be a curve of self-intersection $-e$. In particular, σ_0 is a section of the ruling and

$$K_T \equiv -2\sigma_0 + (2g - 2 - e)f,$$

where f is the numerical equivalence class of a fiber. Note also that either $e > 0$ and σ_0 is the unique curve on T with negative self-intersection, or $e \leq 0$ and every curve on T has nonnegative self-intersection.

If $-K_T = D' + D''$, where D' and D'' are disjoint and nonempty effective divisors, then at most one of D' , D'' can have negative self-intersection. Thus we can assume that $(D'')^2 \geq 0$. If $(D'')^2 > 0$, then $(D')^2 < 0$ by the Hodge index theorem, hence σ_0 is a component of D' . Then every component of D'' is disjoint from σ_0 , hence is numerically equivalent to a positive multiple of $\sigma_0 + ef$. It follows that D'' is numerically equivalent to $\sigma_0 + ef$ and $D' = \sigma_0$. Moreover, $-K_T = D' + D''$ is a union of two disjoint smooth sections. The argument also shows that, if $(D'')^2 = 0$, then $(D')^2 = 0$ as well and hence D' and D'' are numerically equivalent. In particular, since $K_T \equiv -2\sigma_0 + (2g - 2 - e)f$, there exist disjoint sections $\sigma' \subseteq \text{Supp } D'$ and $\sigma'' \subseteq \text{Supp } D''$, and all remaining components of D' , D'' are fibers. Since D' and D'' are disjoint, we must have $\sigma' = D'$ and $\sigma'' = D''$. In all cases, if $\bigcup_i D_i$ is not connected, $-K_T = \sigma' + \sigma''$, where σ', σ'' are disjoint sections of T . Then

$$2g(\sigma') - 2 = 2g(\sigma'') - 2 = K_S \cdot \sigma' + (\sigma')^2 = -(\sigma')^2 + (\sigma')^2 = 0.$$

Hence σ', σ'' are disjoint smooth elliptic curves and T is elliptic ruled. \square

Returning to the proof of Theorem 3.1, let S be a general hypersurface section through x . Then (S, x) is either simple elliptic or a cusp.

The case where (S, x) is simple elliptic:

Let E_0 be the proper transform of S . First we claim that E_0 is normal. In any case, E_0 is Gorenstein, and $\omega_{E_0} = \mathcal{O}_{E_0}(-D_0)$, where $D_0 = \sum_{i \geq 1} a_i(E_0 \cap E_i)$. Let $\nu: \widetilde{E}_0 \rightarrow E_0$ be the normalization (necessarily Cohen-Macaulay), and suppose that ν is not an isomorphism. Then $\omega_{\widetilde{E}_0} = \omega_{\widetilde{E}_0/E_0} \otimes \nu^* \omega_{E_0}$. Since E_0 is Gorenstein, it must fail to be normal in codimension one. Then $\omega_{\widetilde{E}_0/E_0}$ is an ideal sheaf (the conductor ideal sheaf) of $\mathcal{O}_{\widetilde{E}_0}$, hence $\omega_{\widetilde{E}_0/E_0}$ is a rank one reflexive sheaf and $\omega_{\widetilde{E}_0/E_0} \cong \mathcal{O}_{\widetilde{E}_0}(-F)$ for some effective nonzero divisor F on \widetilde{E}_0 . Since the set of non-normal points of E_0 is contained in $E_0 \cap E$, every component of F is also a component of $\nu^{-1}D_0$. In particular, some component of $-K_{\widetilde{E}_0}$ is non-reduced. But this contradicts Lemma 3.2.

Thus, E_0 is normal. By Lemma 3.2, E_0 is Gorenstein and $\omega_{E_0} = \mathcal{O}_{E_0}(-D_0)$, where D_0 is smooth elliptic and $(D_0)_{E_0}^2 < 0$. In particular,

$$\mathcal{O}_{E_0}(E_0) = \mathcal{O}_{E_0}(-D_0) = \mathcal{O}_{\widehat{X}}(-\sum_{i \geq 1} a_i E_i)|_{E_0}.$$

Then E_0 meets $\bigcup_{i \geq 1} E_i$ for a unique i , say $i = 1$, and $a_1 = 1$. Thus

$$K_{E_1} = \mathcal{O}_{E_1}(E_1) = \mathcal{O}_{\widehat{X}}(-E_0 - \sum_{i \geq 2} a_i E_i)|_{E_0}.$$

Moreover, $(D_0)_{E_1}^2 = -(D_0)_{E_0}^2 > 0$. By Lemma 3.3, there are two possibilities:

- (i) $D_0 + \sum_{i \geq 2} a_i(E_1 \cap E_i)$ is connected. Then $E_1 \cap E_i = \emptyset$ for $i > 1$, so that $r = 1$, $E = E_1$, and $K_{E_1} = \mathcal{O}_{E_1}(-D_0)$.
- (ii) E_1 is elliptic ruled and $K_{E_1} = \mathcal{O}_{E_1}(-D_0 - \Gamma)$ for some smooth elliptic curve Γ disjoint from D_0 with $\Gamma^2 < 0$ by the Hodge index theorem.

In the first case, E_1 is rational since $K_{E_1} = (D_0)_{E_1}^2 > 0$. In the second case, E_1 meets $\bigcup_{i \geq 2} E_i$ for a unique i , say $i = 2$, $a_2 = 1$, and $\Gamma = D_{12}$ with $(D_{12})_{E_2}^2 = -(D_{12})_{E_1}^2 > 0$. Then we can repeat this analysis. Eventually this process must terminate with a rational E_r . Moreover, $K_{E_r} = \mathcal{O}_{E_r}(-D_{r-1,r})$, with $(D_{r-1,r})_{E_r}^2 > 0$, so that K_{E_r} is nef and big. Hence we have shown that

E satisfies (i)–(iv) of the Type II case of Definition 2.5. Condition (v) follows since $\sum_{i \geq 0} E_i = (\pi^*t)$ is pulled back from X . Note that $i \geq 1$,

$$\begin{aligned} T_E^1|D_{i,i+1} &= N_{D_{i,i+1}/E_i} \otimes N_{D_{i,i+1}/E_{i+1}} = \mathcal{O}_{\widehat{X}}(E_i + E_{i+1})|D_{i,i+1} \\ &= \mathcal{O}_{\widehat{X}}\left(\sum_{i \geq 0} E_i\right)|D_{i,i+1} = \mathcal{O}_{D_{i,i+1}}, \end{aligned}$$

as remarked after the definition of Type II.

The case where (S, x) is a cusp:

Let E_0 be the proper transform of S . By the same arguments as in the simple elliptic case, E_0 is normal and Gorenstein, and $\omega_{E_0} = \mathcal{O}_{E_0}(-\sum_{k=1}^n \Gamma_k)$, where either each Γ_k is smooth rational and the dual graph of the Γ_k is a cycle, or $n = 1$ and $C = \Gamma_1$ is irreducible. Moreover, E_0 is dominated by a smooth surface \widetilde{E}_0 which is a blowup of a minimal resolution of the cusp, and we can further assume that no fibers of $\widetilde{E}_0 \rightarrow E_0$ are exceptional curves, i.e. that $\widetilde{E}_0 \rightarrow E_0$ is a minimal resolution. Thus E_0 is obtained from \widetilde{E}_0 by contracting chains of curves of self-intersection -2 . We have

$$\mathcal{O}_{E_0}(E_0) = \mathcal{O}_{E_0}\left(-\sum_{k=1}^n \Gamma_k\right) = \mathcal{O}_{\widehat{X}}\left(-\sum_{i \geq 1} a_i E_i\right)|E_0.$$

After reindexing, we can assume that the components E_i of E meeting E_0 are E_1, \dots, E_s , and the above shows that $a_i = 1$ for such i . Define $E_i \cap E_0 = C_i$. Then $E_i \cap E_0 = C_i$ is a union of some of the Γ_k , and the connected components of C_i are either chains of rational curves, or C_i is a cycle of rational curves or an irreducible curve of arithmetic genus one. By Lemma 3.3, the last two cases can only arise if $r = 1$ and $E = E_1$ is irreducible, and necessarily a rational surface. If $r \geq 2$, then C_i is a disjoint union of chains of smooth rational curves, and each E_i with $1 \leq i \leq s$ meets at least one other component E_j for which $C_i \cap C_j \neq \emptyset$. In fact, we can reorder the E_i so that, for $1 \leq i \leq s-1$, E_i meets E_{i+1} with $E_i \cap E_{i+1} \cap E_0 \neq \emptyset$. Note that, as C_i is reduced, no component of C_i is contained in E_j for $j \neq i$, and thus D_{ij} is not contained in C_i for $j \neq i$.

Taking for example $i = 1$, E_1 meets E_2 at a point of $C_1 \cap C_2$. Then there exists a component G_2 of D_{12} meeting the chain C_1 , necessarily at an end component of the chain. Now

$$-K_{E_1} = C_1 + D_{12} + \sum_{j \neq 1,2} a_j D_{1j}$$

and the support $C_1 \cup D_{12} \cup \bigcup_{j \neq 1,2} D_{1j}$ is connected by Lemma 3.3. Hence

$$(K_{E_1} + G_2) \cdot G_2 = -(C_1 \cdot G_2) - \sum_{j \neq 1,2} a_j (D_{1j} \cdot G_2) < 0.$$

Thus $(K_{E_1} + G_2) \cdot G_2 = -2$ and G_2 is a smooth rational curve. Also, $(C_1 \cdot G_2)$ is either 1 or 2. If it is 2, then $G_2 = D_{12}$, C_1 is connected, and E_1 is a rational surface with $(D_{1j} \cdot D_{12}) = 0$ for $j \neq 1, 2$. This implies that $E_1 \cap E_j = \emptyset$ for $j \neq 1, 2$. Also, by the connectedness of C_1 , it follows that $E_2 \cap E_j = \emptyset$ for $j \neq 1, 2$, and we are in the case $s = 2$. So we can assume that $(C_1 \cdot G) = 1$, there is a unique $k \neq 1, 2$, say $k = 3$, such that $(D_{13} \cdot G_2) \neq 0$, and $(D_{13} \cdot G_2) = 1$ and $a_3 = 1$.

Let G_3 be the unique component of D_{13} meeting G_2 . Repeating this argument with G_3 , we see that G_3 is smooth rational and that $(C_1 + D_{12}) \cdot G_3$ is either 1 or 2. If it is 2, then the only possibility is that G_3 meets C_1 at the other end of the chain from G_2 . In this case, $E_1 \cap E_j = \emptyset$ for $j \neq 1, 2, 3$. Otherwise we can continue this process with an E_j and with a smooth component G_j of D_{1j} such that $G_j \cdot G_3 \neq 0$. Eventually the curves C_1 and the G_j must close up (although it is possible for some E_j to be equal to E_ℓ at an intermediate stage). We can do this analysis for all E_i , $1 \leq i \leq s$: every component of D_{ij} is a smooth rational curve, and $a_j = 1$ for every j such that $E_i \cap E_j \neq \emptyset$. Moreover, the scheme-theoretic intersection of E_0 , E_i , and $E_{i \pm 1}$ is a reduced point,

and hence is a smooth point of E_0 . In particular, $C = \bigcup_i C_i$ is a cycle of smooth rational curves, hence has arithmetic genus 1.

Now let E_k be a component of E with $k > s$ such that $E_k \cap E_1 \neq \emptyset$, say, and let G be a component of D_{1k} . Then $a_k = 1$ and G is a smooth rational curve. Moreover, there exists an $\ell \neq i, j$ such that $G \cap E_\ell \neq \emptyset$ as well. Thus $E_1 \cap E_k \cap E_\ell \neq \emptyset$, say, and

$$-K_{E_k} = D_{1k} + D_{k\ell} + \sum_{t \neq 1, k, \ell} a_t D_{kt}.$$

Let G' be a component of $D_{k\ell}$ meeting G . Arguments as above show that G is smooth rational and that $a_t = 1$ for every t such that $E_t \cap G' \neq \emptyset$. Continuing in this way and using the connectedness of E , it follows that $a_i = 1$ for every i and that every component of D_{ij} is a smooth rational curve. Moreover, for every i , there exists a curve in $|-K_{E_i}|$ whose components are rational curves. Hence E_i is rational. Thus, E satisfies the conditions of the second statement in Theorem 3.1(iii). \square

Remark 3.4. The proof shows that, in case the hypersurface section S is a cusp and using the notation of the proof, $-K_{E_i}$ is effective and is a cycle of smooth rational curves which contains C_i if $E_i \cap E_0 \neq \emptyset$.

Remark 3.5. (i) It seems quite possible that, in general, E might not be of Type III₁ or Type III₂, even after making some flops. In particular, from the point of view of the classification of algebraic varieties, it is reasonable to allow E to have more complicated singularities than normal crossings, namely dlt singularities.

(ii) In the Type II case, $\dim H^0(E; T_E^1) = r - 1$ is the number of elliptic ruled components. In the Type III₁ or Type III₂ cases, $H^0(E; T_E^1) = 0$, so that all first order deformations of E are locally trivial in these cases. This follows more generally in case the general hypersurface section of X passing through x is a cusp, by Theorem 3.1(iii) and Theorem 1.6(ii).

(iii) It is easy to see that, if $|\Gamma|$ is the dual complex of E , then $H^i(|\Gamma|) = 0$ for $i > 0$, and indeed a theorem of [dFKX17] says that $|\Gamma|$ is contractible. In the Type II and Type III₁ case, $|\Gamma|$ is a point or a line segment, and in the Type III₂ case $|\Gamma|$ is a disk. However, in the general case, without making flops, the topological type of the dual complex can be more complicated than a line segment or a disk. For example, it can be a disk meeting a line segment at a point.

On the positive side, there is the following:

Theorem 3.6. *Let (X, x) be an isolated singularity of dimension 3 with a good crepant resolution $\pi: \widehat{X} \rightarrow X$ and let $E = \pi^{-1}(x)$ be the reduced exceptional divisor.*

- (i) *If the general hypersurface section of X passing through x is a cusp and ω_E^{-1} is nef and big, then E is of Type III₁ or Type III₂.*
- (ii) *If the general hypersurface section S of X passing through x is a cusp and the full inverse image $\pi^{-1}(S)$ has normal crossings, then after a sequence of flops (elementary modifications of type 2), ω_E^{-1} becomes nef and big, hence E is of Type III₁ or Type III₂.*

Proof. First assume that ω_E^{-1} is nef and big. In the notation of the proof of Theorem 3.1, $C = \bigcup_{i=1}^s C_i = E \cap E_0$ is a cycle of rational curves and $\mathcal{O}_{E_0}(-C) = \omega_{E_0} = \mathcal{O}_{E_0}(-\sum_i \Gamma_i)$. Likewise, $\omega_E = \mathcal{O}_E(-C) = \mathcal{O}_E(E)$. Since $(C)_E^2 = -(C)_{E_0}^2 > 0$, the total degree of $\mathcal{O}_C(C) = \omega_E^{-1}|_C$ is positive. Then general results on line bundles on cycles of rational curves (cf. [Fri15, Lemma 1.7]) imply either that $(C)_E^2 \geq 2$ and $\mathcal{O}_C(C)$ has no base points or $(C)_E^2 = 1$ and $\mathcal{O}_C(C)$ has a single base point at a smooth point of C . From the exact sequence

$$0 \rightarrow \mathcal{O}_E \rightarrow \mathcal{O}_E(C) \rightarrow \mathcal{O}_C(C) \rightarrow 0,$$

there exists a section of $\mathcal{O}_E(C)$ vanishing at C^* , where $C^* = \bigcup_{i=1}^s C_i^*$ and C_i^* is smooth for all i . Since $\mathcal{O}_E(-E)|_C$ is nef and has nonnegative degree on every component and positive total degree, $H^1(E; \mathcal{O}_E(-NE)) = 0$ for all $N \geq 0$. By induction, it is then easy to see that $H^1(nE; \mathcal{O}_{\widehat{X}}(-NE)|_{nE}) = 0$ for all $N \geq 0$ and all $n > 0$. Thus $R^1\pi_*\mathcal{O}_{\widehat{X}}(-NE) = 0$ for all $N \geq 0$ by the formal functions theorem. In particular, $R^1\pi_*\mathcal{O}_{\widehat{X}}(-2E) = 0$. By applying $R^i\pi_*$ to the exact sequence

$$0 \rightarrow \mathcal{O}_{\widehat{X}}(-2E) \rightarrow \mathcal{O}_{\widehat{X}}(-E) \rightarrow \mathcal{O}_E(C) \rightarrow 0,$$

it follows that the natural map

$$R^0\pi_*\mathcal{O}_{\widehat{X}}(-E) \rightarrow H^0(E; \mathcal{O}_E(C))$$

is surjective. Hence, there exists an element $t \in R^0\pi_*\mathcal{O}_{\widehat{X}}(-E) \subseteq \mathfrak{m}_x$ which lifts to a function π^*t whose restriction to E is C^* . Thus $S^* = \{t = 0\}$ defines a cusp singularity on X , the proper transform E_0^* of S^* is a resolution of singularities of S^* and $E \cup E_0^*$ has simple normal crossings. It follows from the classification of Type III degenerations of K3 surfaces that $E \cup E_0^*$ is a Type III anticanonical pair, in the notation of [FM83, p. 192] or [EF21, §2], except that the Hirzebruch-Inoue component has been replaced by the local surface E_0^* . The assumption that ω_E^{-1} is nef and big is then equivalent to the assumption that $(C'_i)_{E_i}^2 \geq 0$ for every component C'_i of $C_i^* = E_0^* \cap E_i$.

On the other hand, under the assumption of (ii), $E \cup E_0$ is again a Type III anticanonical pair as above. Then every component C'_i of $C_i = E_0 \cap E_i$ is a smooth rational curve and $(C'_i)_{E_0}^2 < 0$ for every i . By the triple point formula, $(C'_i)_{E_i}^2 \geq 0$ unless $(C'_i)_{E_0}^2 = (C'_i)_{E_i}^2 = -1$. In this case, the standard flop (type 2 modification) eliminates C'_i but does not alter the assumption that $\pi^{-1}(S)$ has normal crossings. In this process, the total number i such that $(C'_i)_{E_0}^2 = -1$ decreases, so it must ultimately terminate at a stage where $(C'_i)_{E_0}^2 \leq -2$ for every i . Thus we can assume in (i) and in (ii) that $(C'_i)_{E_i}^2 \geq 0$ for every component C'_i of $C_i = E_0 \cap E_i$.

In this case, suppose that there exists a component E_1 such that $C_1 = E_0 \cap E_1$ is disconnected. Then every component of C_1 has nonnegative square on E_1 . By the Hodge index theorem, every component of C_1 has square 0 on E_1 and E_1 is a minimal rational ruled surface with the remaining double curves sections of the ruling. If E_2 is a component meeting E_0 and E_1 , then the same analysis shows that either every component of C_2 has square 0 on E_2 and E_2 is a minimal rational ruled surface with the remaining double curves sections of the ruling, or C_2 is connected, hence an irreducible smooth rational curve, and C_1 has just two components. Continuing in this way, we see that E is of Type III₁. In the remaining case, E_0 meets every component of E in an irreducible smooth rational curve. It is easy to see in this case that the dual complex of E triangulates a 2-disk and thus that E is of Type III₂. \square

4. THE CASE OF A SMALL RESOLUTION

In this section, we consider the case of a small resolution $\pi': X' \rightarrow X$, i.e. (X, x) is the germ of an isolated Gorenstein singularity of dimension 3 with a good Stein representative X , and $p: X' \rightarrow X$ is a small resolution with exceptional set C .

Remark 4.1. There is no real limitation to restricting to dimension 3, at least in case the singularity (X, x) is a local complete intersection. Indeed, such resolutions can only exist for $\dim X = 3$: By the Grothendieck-Lefschetz theorem, the local ring $\mathcal{O}_{X,x}$ of an isolated local complete intersection singularity is a UFD for $\dim X \geq 4$. The proof of the “easy case” of Zariski’s Main Theorem shows that, in case $\mathcal{O}_{X,x}$ is a UFD and $\pi: \widehat{X} \rightarrow X$ is a resolution of singularities, there exists a divisor D on \widehat{X} such that $\text{codim } \overline{\pi(D)} \geq 2$. Thus, if (X, x) is an isolated local complete intersection singularity and $\pi': X' \rightarrow X$ is a resolution such that the exceptional set $(\pi')^{-1}(x)$ has dimension $\leq \dim X - 2$, then $\dim X = 3$.

Conversely, in dimension 3, suppose that (X, x) is the germ of an isolated singularity with a small resolution $\pi': X' \rightarrow X$. If X is Cohen-Macaulay, then it follows from results of Laufer, Reid, and Pinkham (see for example [Pin83, §8]) that X is a compound du Val singularity, i.e. that the general hyperplane section of X in the sense of Definition 2.3 is a rational double point. In particular, X is a hypersurface singularity and $(\pi')^{-1}(x) = C = \bigcup_{i=1}^r C_i$, where the C_i are smooth rational curves meeting (pairwise) transversally (but three C_i can meet at a point).

For the case of a small resolution, the functor $\mathbf{Def}_{X'}$ has a more than purely formal meaning: By [Lau80, Theorem 2], there is a deformation of a neighborhood of the exceptional curve C over the smooth germ $(H^1(X'; T_{X'}), 0)$ for which the Kodaira-Spencer map is an isomorphism. As previously noted, X' is a crepant resolution of X , and hence $K_{X'} \cong \mathcal{O}_{X'}$. Also, the resolution $p: X' \rightarrow X$ is equivariant (cf. Definition 1.1).

We will use the following standard fact about local cohomology:

Lemma 4.2. *If \mathcal{F} is a locally free sheaf on X' , then $H_C^1(X'; \mathcal{F}) = 0$.*

Proof. Using the Mayer-Vietoris sequence [Har77, Ex. III (2.4) p. 212], it suffices to show that $H_{C_i}^1(X'; \mathcal{F}) = 0$ for every irreducible component C_i of C and that $H_p^2(X'; \mathcal{F}) = 0$ for every point $p \in X'$. By [Har67, (1.4)], there is a spectral sequence with E_1 term $E_1^{p,q} = H^p(X'; \mathcal{H}_{C_i}^q(\mathcal{F}))$ converging to $H_{C_i}^{p+q}(X'; \mathcal{F})$, where $\mathcal{H}_{C_i}^q(\mathcal{F})$ is the associated local cohomology sheaf. Since C_i is smooth, it is a local complete intersection. Hence, by [Har67, (3.7), (3.8)] $\mathcal{H}_{C_i}^q(\mathcal{F}) = 0$ for $i = 0, 1$. Thus $H_{C_i}^1(X'; \mathcal{F}) = 0$. The vanishing of $H_p^2(X'; \mathcal{F})$ is similar. \square

Since the fibers of p have dimension one, $R^2 p_* T_{X'} = 0$ and hence $H_x^0(X; R^2 p_* T_{X'}) = 0$. Applying the Leray spectral in local cohomology to the morphism p and the sheaf $T_{X'}$ and using Lemma 4.2 to see that $H_C^1(X'; T_{X'}) = 0$ gives:

Lemma 4.3. *There is an exact sequence*

$$0 \rightarrow H^0(X; R^1 p_* T_{X'}) \rightarrow H^0(X; T_X^1) \rightarrow H_C^2(X'; T_{X'}) \rightarrow 0. \quad \square$$

Note that $H_x^0(X; R^1 p_* T_{X'}) = H^0(X; R^1 p_* T_{X'})$ and that $H_x^2(X; T_X^0) \cong H^0(X; T_X^1)$. Since $K_{X'}$ is trivial, $T_{X'} \cong \Omega_{X'}^2$, and hence $H_C^2(X'; T_{X'}) \cong H_C^2(X'; \Omega_{X'}^2)$.

The (singular) local cohomology groups $H_C^k(X')$ can be described via duality:

$$H_C^k(X') \cong H_{6-k}(C) = \begin{cases} 0, & \text{if } k \neq 4, 6; \\ \bigoplus_i H_2(C_i), & \text{if } k = 4; \\ \mathbb{C}, & \text{if } k = 6. \end{cases}$$

Moreover, there is a spectral sequence

$$E_1^{p,q} = H_C^q(X'; \Omega_{X'}^p) \implies \mathbb{H}_C^{p+q}(X'; \Omega_{X'}^\bullet) = H_C^{p+q}(X').$$

Many of the terms in the E_1 page of the spectral sequence are zero:

Lemma 4.4. *If $q = 0, 1$, then $H_C^q(X'; \Omega_{X'}^p) = 0$ for all p , and $H_C^2(X'; \Omega_{X'}^3) = H_C^2(X'; \mathcal{O}_{X'}) = 0$.*

Proof. The first statement follows from Lemma 4.2. The second follows by considering the Leray spectral sequence with $E_2^{p,q} = H_x^p(X; R^q p_* \mathcal{O}_{X'}) \implies H_C^{p+q}(X'; \mathcal{O}_{X'})$. Here, $R^q p_* \mathcal{O}_{X'} = 0$ for $q > 0$ since (X, x) is a rational singularity and

$$H_x^p(X; R^0 p_* \mathcal{O}_{X'}) = H_x^p(X; \mathcal{O}_X) = 0$$

for $p < 3$ because $\text{depth } \mathcal{O}_{X,x} = 3$. \square

Thus we have the following picture for the $E_1^{p,q}$ page of the spectral sequence converging to $\mathbb{H}_C^*(X'; \Omega_{X'}^\bullet) = H_C^{p+q}(X')$:

$H_C^3(X'; \mathcal{O}_{X'})$	$H_C^3(X'; \Omega_{X'}^1)$	$H_C^3(X'; \Omega_{X'}^2)$	$H_C^3(X'; \Omega_{X'}^3)$
	$H_C^2(X'; \Omega_{X'}^1)$	$H_C^2(X'; \Omega_{X'}^2)$	

Lemma 4.5. *The differential $d: H_C^2(X'; \Omega_{X'}^1) \rightarrow H_C^2(X'; \Omega_{X'}^2)$ is injective.*

Proof. This is clear since the kernel of $d: H_C^2(X'; \Omega_{X'}^1) \rightarrow H_C^2(X'; \Omega_{X'}^2)$ would inject into $H_C^3(X') = 0$. \square

By examining the above spectral sequence, there is a homomorphism $H_C^2(X'; \Omega_{X'}^2) \rightarrow H_C^4(X')$.

Proposition 4.6. *The map $H_C^2(X'; \Omega_{X'}^2) \rightarrow H_C^4(X')$ is surjective and split by the fundamental class map. Thus, if K'_x denotes the kernel of $H_C^2(X'; \Omega_{X'}^2) \rightarrow H_C^4(X')$, we have a direct sum decomposition*

$$H_C^2(X'; \Omega_{X'}^2) \cong K'_x \oplus H_C^4(X') \cong K'_x \oplus \left(\bigoplus_{i=1}^r \mathbb{C}[C_i] \right).$$

Proof. First note that the fundamental classes of the C_i are a basis for $H_C^4(X') \cong H_2(C)$. On the other hand, for every i one can construct a fundamental class $[C_i] \in H_C^2(X'; \Omega_{X'}^2)$ which maps to the fundamental class $[C_i] \in H_C^4(X')$. We also have the Mayer-Vietoris sequence

$$0 = \bigoplus_{z \in C_{\text{sing}}} H_z^2(X'; \Omega_{X'}^2) \rightarrow \bigoplus_i H_{C_i}^2(X'; \Omega_{X'}^2) \rightarrow H_C^2(X'; \Omega_{X'}^2),$$

and hence we can view the $[C_i]$ as linearly independent elements of $H_C^2(X'; \Omega_{X'}^2)$. The image of $H_C^2(X'; \Omega_{X'}^2)$ in $H_C^4(X')$ therefore contains the vector space spanned by the fundamental classes of the components of C , and hence is equal to $H_C^4(X')$. Thus $H_C^2(X'; \Omega_{X'}^2) \rightarrow H_C^4(X')$ is surjective and the subspace spanned by the fundamental classes of the components of C is a complement to the kernel. \square

Define $S_x \cong \mathbb{C}^r \subseteq H_C^2(\Omega_{X'}^2)$ to be the image of the fundamental class map. Thus $H_C^2(X'; \Omega_{X'}^2) \cong K'_x \oplus S_x$. By Proposition 4.6, we can identify S_x with $H_C^4(X')$. It can also be identified with $H^3(L)$, where $L = X - \{x\} = X' - C$ is the link of the singularity. This follows from the exact sequence

$$0 = H^3(X') \rightarrow H^3(U) \rightarrow H_C^4(X') \rightarrow H^4(X') = 0.$$

In particular, the mixed Hodge structure on $S_x \cong H^3(L)$ is pure weight 4 and of type $(2, 2)$.

To say more about K'_x , we have:

Lemma 4.7. *Let A_x be the kernel of $d: H_C^3(X'; \mathcal{O}_{X'}) \rightarrow H_C^3(X'; \Omega_{X'}^1)$ and let $a = \dim A_x$. Then there is an exact sequence*

$$0 \rightarrow H_C^2(X'; \Omega_{X'}^1) \rightarrow K'_x \rightarrow A_x \rightarrow 0.$$

Thus $\dim K'_x = \dim H_C^2(X'; \Omega_{X'}^1) + a$. If $A_x = 0$, then $d: H_C^2(X'; \Omega_{X'}^1) \rightarrow K'_x$ is an isomorphism.

Proof. By examining the spectral sequence, we see that $d_2: A_x \rightarrow H_C^2(X'; \Omega_{X'}^2)/d(H_C^2(X'; \Omega_{X'}^1))$ is injective and that its image is $K'_x/d(H_C^2(X'; \Omega_{X'}^1))$. Hence $A_x \cong K'_x/d(H_C^2(X'; \Omega_{X'}^1))$. The remaining statements of the lemma are clear. \square

Corollary 4.8. *In the above notation, there is an exact sequence*

$$0 \rightarrow (K'_x)^\vee/A_x^\vee \rightarrow H^0(X; T_X^1) \rightarrow K'_x \oplus S_x \rightarrow 0.$$

Proof. By duality, arguing as in [Kar86],

$$H^0(X; R^1 p_* T_{X'}) \cong H_C^2(X'; \Omega_{X'}^1)^\vee \cong (K'_x)^\vee/A_x^\vee.$$

The proof then follows from Proposition 4.6 and Lemma 4.7. \square

Corollary 4.9. *Let $b = \dim K'_x$. Then*

- (i) $\dim H_C^2(X'; \Omega_{X'}^2) = b + r$.
- (ii) $\dim H_C^2(X'; \Omega_{X'}^1) = \dim H^0(X; R^1 p_* T_{X'}) = b - a$.
- (iii) $\dim H^0(X; T_X^1) = 2b - a + r$. Thus $b + r \leq \dim H^0(X; T_X^1) \leq 2b + r$.

Proof. These follows from Proposition 4.6, Lemma 4.7 and Corollary 4.8 (and its proof). \square

Theorem 4.10. *In the above situation, the following are equivalent:*

- (i) C is smooth, i.e. $r = 1$, and the normal bundle $N_{C/X'} \cong \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. In other words, (X, x) is the germ of an ordinary double point.
- (ii) $\dim H_C^2(X'; \Omega_{X'}^2) = r$, i.e. $K'_x = 0$, or equivalently $b = 0$.
- (iii) $R^1 p_* T_{X'} = 0$, or equivalently $b = a$.

Proof. (i) \implies (ii): In this case, $\dim H^0(X; T_X^1) = 1 = 2b - a + 1$. Since $b \geq 0$ and $b - a = \dim H_C^2(X'; \Omega_{X'}^1) \geq 0$, we must have $b = 0$, and hence $K'_x = 0$.

(ii) \implies (iii): By Corollary 4.9(ii), $\dim H^0(X; R^1 p_* T_{X'}) = b - a$. If $K'_x = 0$, then $b = \dim K'_x = 0$, hence $a = 0$ and $R^1 p_* T_{X'} = 0$.

(iii) \implies (i): Following the discussion in [Fri86, pp. 678–679], there exists a small deformation of X' to X'_t , where the exceptional curve C splits up into a union of δ disjoint copies of \mathbb{P}^1 with normal bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$, and such a deformation blows down to a deformation of X to a union of δ ordinary double points. But if $R^1 p_* T_{X'} = 0$, then the only deformations of X' are locally trivial. Hence the exceptional curve C on X' is already a single \mathbb{P}^1 with normal bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$, and (X, x) is the germ of an ordinary double point. \square

Proposition 4.11. *Suppose that there is a small deformation of X' to a space with exactly δ compact curves, and all of these have normal bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. Then $\dim H_C^2(X'; \Omega_{X'}^2) = \delta$.*

Proof. Let (S, s_0) be the germ of a smooth analytic space prorepresenting \mathbf{Def}_X and let (S', s'_0) prorepresent $\mathbf{Def}_{X'}$. The morphism of functors $\mathbf{Def}_{X'} \rightarrow \mathbf{Def}_X$ then induces a morphism of germs $S' \rightarrow S$ which is an immersion by [Fri86, Proposition 2.1]. By Lemma 4.3, $H_C^2(X'; \Omega_{X'}^2)$ is the normal bundle to this immersion, so it will suffice to prove that the image of S' has codimension δ . As noted in the proof of Theorem 4.10, there is an open dense subset of S' corresponding to a germ with exactly δ singularities, all of which are ordinary double points. By openness of versality, the codimension of the image of S' in S then has codimension δ as claimed. \square

Corollary 4.12. (Compare [Nam02, Lemma 1.9].) We have $b + r = \delta$. Thus $\delta \geq r$, with equality $\iff (X, x)$ is an ordinary double point. \square

We now compare the above discussion with the case of a good resolution, as described in [FL22, Theorem 2.1]. By successively blowing up the curves C_i in some order, we obtain a good resolution $\pi: \widehat{X} \rightarrow X$ which is an iterated blowup of X' . Let $\rho: \widehat{X} \rightarrow X'$ be the blowup morphism, and $E = \bigcup_i E_i$ the exceptional divisor of ρ or of π . To distinguish groups on X' and on \widehat{X} , we denote the latter with a “ $\widehat{}$ ”. Then we have the group \widehat{K}'_x defined in [FL22, Theorem 2.1(vi)]:

$$\widehat{K}'_x = \text{Ker}\{H_E^2(\widehat{X}; \Omega_{\widehat{X}}^2) \rightarrow H^2(\widehat{X}; \Omega_{\widehat{X}}^2)\}.$$

We also have $\text{Gr}_F^2 H^3(L) = H^3(L)$ and $\widehat{A}_x = \text{Ker}\{d: H^3(\widehat{X}; \mathcal{O}_{\widehat{X}}) \rightarrow H_E^3(\widehat{X}; \Omega_{\widehat{X}}^1)\}$. (In [FL22, Corollary 1.8], \widehat{A}_x is defined to be the kernel of $d: H^3(\widehat{X}; \mathcal{O}_{\widehat{X}}) \rightarrow H_E^3(\widehat{X}; \Omega_{\widehat{X}}^1(\log E))$), but it is easy to check that in our case $H_E^3(\widehat{X}; \Omega_{\widehat{X}}^1) \rightarrow H_E^3(\widehat{X}; \Omega_{\widehat{X}}^1(\log E))$ is injective.) There is an exact sequence [FL22, Theorem 2.1(v) and (vi)]

$$0 \rightarrow H^1(\widehat{X}; \Omega_{\widehat{X}}^2(\log E)(-E)) \rightarrow H^0(X; T_X^1) \rightarrow \widehat{K}'_x \oplus H^3(L) \rightarrow 0,$$

with $H^1(\widehat{X}; \Omega_{\widehat{X}}^2(\log E)(-E))^\vee \cong H_E^2(\widehat{X}; \Omega_{\widehat{X}}^2(\log E))$, and by [FL22, Corollary 1.8 and Theorem 2.1(iv)] an exact sequence

$$0 \rightarrow H_E^2(\widehat{X}; \Omega_{\widehat{X}}^2(\log E)) \rightarrow \widehat{K}'_x \rightarrow \widehat{A}_x \rightarrow 0.$$

Hence $H^1(\widehat{X}; \Omega_{\widehat{X}}^2(\log E)(-E)) \cong (\widehat{K}'_x)^\vee / \widehat{A}_x^\vee$.

Proposition 4.13. *Let $\rho: \widehat{X} \rightarrow X$ be a good resolution of X which is obtained by successively blowing up the curves C_i in some order.*

- (i) *There is are isomorphisms $H^1(\widehat{X}; \Omega_{\widehat{X}}^2) \cong H^1(X'; \Omega_{X'}^2) \cong H^1(X'; T_{X'})$, compatible with the natural homomorphisms to $H^1(U; T_U) \cong H^0(X; T_X^1)$.*
- (ii) *Let \widehat{K}'_x , $H^3(L)$, and \widehat{A}_x be the groups described above. Then $\widehat{K}'_x \cong K'_x$, $H^3(L) \cong S_x$, and $\widehat{A}_x \cong A_x$, so that the following diagram commutes:*

$$\begin{array}{ccccccc} 0 & \longrightarrow & (\widehat{K}'_x)^\vee / \widehat{A}_x^\vee & \longrightarrow & H^0(X; T_X^1) & \longrightarrow & \widehat{K}'_x \oplus H^3(L) \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & (K'_x)^\vee / A_x^\vee & \longrightarrow & H^0(X; T_X^1) & \longrightarrow & K'_x \oplus S_x \longrightarrow 0. \end{array}$$

Proof. (i): By standard results, $R^0 \rho_* \Omega_{\widehat{X}}^k = \Omega_{X'}^k$, $R^i \rho_* \mathcal{O}_{X'} = 0$ for $i > 0$, and $R^i \rho_* \Omega_{\widehat{X}}^k = 0$ for $i \geq 2$. As for $R^1 \rho_* \Omega_{\widehat{X}}^k$, first suppose that C is smooth. Then, by a standard argument (cf. [Gro85, V(1.2.1)]), $R^1 \rho_* \Omega_{\widehat{X}}^1 \cong \mathcal{O}_C$ and $R^1 \rho_* \Omega_{\widehat{X}}^2 \cong \Omega_C^1$. In the general case, a successive application of the Leray spectral sequence for the iterated blowup shows that $R^1 \rho_* \Omega_{\widehat{X}}^1$ has a filtration whose successive quotients are \mathcal{O}_{C_i} , $1 \leq i \leq r$, and similarly $R^1 \rho_* \Omega_{\widehat{X}}^2$ has a filtration whose successive quotients are $\Omega_{C_i}^1$, $1 \leq i \leq r$. In particular, $H^0(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) = 0$. Thus, by the Leray spectral sequence, $H^1(\widehat{X}; \Omega_{\widehat{X}}^2) \cong H^1(X'; R^0 \rho_* \Omega_{X'}^2) \cong H^1(X'; \Omega_{X'}^2)$, and the remaining statements in (i) are clear.

(ii): By the proof of (i), $H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^1) = H_C^0(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) = 0$, and $\dim H_C^0(X'; R^1 \rho_* \Omega_{\widehat{X}}^1) = \dim H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) = r$. Similarly, $R^0 \rho_* \mathbb{C}_{\widehat{X}} = \mathbb{C}_{X'}$, $R^2 \rho_* \mathbb{C}_{\widehat{X}}$ is a successive extension of the \mathbb{C}_{C_i} , and otherwise $R^i \rho_* \mathbb{C}_{\widehat{X}} = 0$. In particular, $H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) \cong H_C^2(X'; R^2 \rho_* \mathbb{C}_{\widehat{X}})$.

The Leray spectral sequence in local cohomology gives a spectral sequence with

$$E_2^{a,b} = H_C^a(X'; R^b \rho_* \Omega_{\widehat{X}}^2) \implies H_E^{a+b}(\widehat{X}; \Omega_{\widehat{X}}^2).$$

The only nonzero terms are $H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^2)$ and $H_C^2(X'; R^0 \rho_* \Omega_{\widehat{X}}^2) = H_C^2(X'; \Omega_{X'}^2)$. It follows that $H_E^1(\widehat{X}; \Omega_{\widehat{X}}^2) = 0$ and there is a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_C^2(X'; \Omega_{X'}^2) & \longrightarrow & H_E^2(\widehat{X}; \Omega_{\widehat{X}}^2) & \longrightarrow & H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) \longrightarrow 0 \\ & & \parallel & & \parallel & & \downarrow \cong \\ & & K'_x \oplus \mathbb{C}^r & \longrightarrow & \widehat{K}'_x \oplus H_E^4(\widehat{X}) & \longrightarrow & \mathbb{C}^r \end{array}$$

Here, we use the Leray spectral sequence to conclude also that there is an exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_C^4(X') & \longrightarrow & H_E^4(\widehat{X}) & \longrightarrow & H_C^2(X'; R^2 \rho_* \mathbb{C}_{\widehat{X}}) \longrightarrow 0 \\ & & \parallel & & \parallel & & \downarrow \cong \\ & & \bigoplus_i \mathbb{C}[C_i] & & & & H_C^1(X'; R^1 \rho_* \Omega_{\widehat{X}}^2) \end{array}$$

Tracing through the identifications gives $\widehat{K}'_x \cong K'_x$. By definition, $H^3(L) \cong \bigoplus_i \mathbb{C}[C_i] = S_x$. The remaining identification of \widehat{A}_x with A_x is similar, using the Leray spectral sequence to conclude that $H^3_E(\widehat{X}; \mathcal{O}_{\widehat{X}}) \cong H^3_C(X'; \mathcal{O}_{X'})$ and $H^3_E(\widehat{X}; \Omega^1_{\widehat{X}}) \cong H^3_C(X'; \Omega^1_{X'})$, compatibly with d . \square

Recall that a \mathbb{C}^* action on the germ of an analytic space (X, x) is *good* if the weights of the induced action on the Zariski tangent space of the fixed point x are all positive. Applying the case of a good (divisorial) resolution [FL22, Theorem 2.1(iv)] and using the fact that X is a hypersurface singularity, hence a local complete intersection singularity, we have the following corollary:

Corollary 4.14. *The germ (X, x) has a good \mathbb{C}^* action $\iff A_x = 0 \iff a = 0 \iff$ the spectral sequence with $E_1^{p,q} = H^q_C(X'; \Omega^p_{X'}) \implies \mathbb{H}^{p+q}_C(X'; \Omega^\bullet_{X'}) = H^{p+q}_C(X')$ degenerates at E_2 . \square*

The invariants b and a can also be described in terms of the Du Bois invariants $b^{p,q}(X, x)$ of the singularity X , using [FL22, Theorem 2.1(iv), (vi)]:

Corollary 4.15. $b = b^{1,1}(X, x)$ and $b - a = b^{2,1}(X, x)$. \square

Remark 4.16. In particular, by Corollary 4.9(iii) and the fact that $\ell = \ell^{2,1} = r$, we recover the result of Steenbrink [Ste97, Theorem 4] that

$$\dim H^0(X; T^1_X) = b^{1,1}(X, x) + b^{2,1}(X, x) + \ell^{2,1},$$

the inequalities due to Namikawa [Nam97, Theorem 1]

$$b^{1,1}(X, x) + \ell^{2,1} \leq \dim H^0(X; T^1_X) \leq 2b^{1,1}(X, x) + \ell^{2,1},$$

as well as the statement that $\dim H^0(X; T^1_X) = 2b^{1,1}(X, x) + \ell^{2,1} \iff (X, x)$ is weighted homogeneous (since (X, x) is a local complete intersection). As shown in the papers of Steenbrink and Namikawa cited above and [FL22, Theorem 2.1(iv)], these results hold more generally for isolated rational Gorenstein singularities of dimension 3.

Example 4.17. Consider the A_{2n-1} singularity $x^2 + y^2 + z^2 + w^{2n}$ (a compound A_1 singularity). Here, $T^1_X \cong \mathbb{C}[w]/(w^{2n-1})$ has dimension $2n - 1$ and $r = 1$. A calculation shows that $\dim H^0(X; R^1 p_* T_{X'}) = n - 1$. Hence $\dim H^2_C(X'; \Omega^2_{X'}) = n$ by Lemma 4.3. Note also that we can deform \widehat{X} so that the exceptional curve C breaks up into a union of n curves with normal bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. The this deformation then blows down to a deformation of X to the union of n ordinary double points, i.e. $\delta = n$ in the notation of Proposition 4.11.

Example 4.18. Consider the compound A_{n-1} singularity $x^2 + y^2 + f(z, w)$, where $f(z, w) = \prod_{i=1}^n (z + \lambda_i w)$ defines a plane curve which is the union of distinct n lines meeting at the origin. An easily computable example is $f(z, w) = z^n - w^n$. Thus

$$T^1_X \cong \mathbb{C}[z, w]/(z^{n-1}, w^{n-1}).$$

A calculation shows that $\dim T^1_X = (n - 1)^2$. Note that, for $n \geq 4$, $(X, 0)$ has nontrivial equisingular deformations (there are local moduli). This is also reflected in the fact that there are nontrivial weight zero deformations for $n \geq 4$ (and nontrivial positive weight deformations for $n \geq 5$). By Corollary 4.9, since $a = 0$ and $r = n - 1$, $\dim H^0(X; R^1 p_* T_{X'}) = (n - 1)(n - 2)/2$ and $\dim H^1_C(X'; \Omega^2_{X'}) = n(n - 1)/2$.

The surface X_0 defined by $w = 0$ is an A_{n-1} singularity and the inverse image X'_0 in X' is a resolution of singularities. Moreover all of the components C_i in X' have normal bundle $\mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-1)$. By general results (e.g. [Bri71], [Art74], or the paper by Pinkham [DPT80, *Résolution simultanée de points doubles rationels*]), there is a morphism of functors $\mathbf{Def}_{X'_0} \rightarrow \mathbf{Def}_{X_0}$ with the following property: If T is the analytic germ prorepresenting the functor \mathbf{Def}_{X_0} and \widetilde{T} is the germ prorepresenting $\mathbf{Def}_{X'_0}$, then the induced morphism $\widetilde{T} \rightarrow T$ is a Galois cover of smooth germs

with Galois group the Weyl group of the corresponding root system, in this case A_{n-1} . The inverse image in \widetilde{T} of the discriminant locus in T consists of δ hyperplanes, corresponding to keeping one of the δ positive roots the class of an irreducible effective curve.

This is in agreement with Proposition 4.11, because in this case $\delta = \binom{n}{2}$ (one can deform the union of n concurrent lines to a union of n lines meeting transversally).

Specializing to the case $n = 5$, and hence $r = 4$, we have $\dim H^0(X; T_X^1) = 16$. We can deform the singularity $x^2 + y^2 + z^5 - w^5$ in the weight one direction to X_t which is defined by

$$x^2 + y^2 + z^5 - w^5 + tz^3w^3.$$

A calculation shows that $\dim H^0(X_t; T_{X_t}^1) = 15$ if $t \neq 0$. In particular, $a \neq 0$ in this case (in fact $a = 1$), and the spectral sequence with E_1 page $H_C^q(X'_t; \Omega_{X'_t}^p) \implies H_C^{p+q}(X'_t)$ does not degenerate at E_2 for $t \neq 0$.

5. A NON-CREPANT EXAMPLE

In this final section, we consider a non-crepant example \widehat{X} , the blowup of a small resolution with exceptional set a smooth curve C along the curve C . Following the discussion of the introduction, there are homomorphisms $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(U, T_U) \cong H^0(X; T_X^1)$ and $H^1(\widehat{X}; \Omega_{\widehat{X}}^2) \rightarrow H^1(U, T_U) \cong H^0(X; T_X^1)$. The image of $H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$ is a birational invariant, i.e. is independent of the choice of a good resolution, and is identified with the image of $H^1(X'; T_{X'})$ by Proposition 4.13(i). On the other hand, the image of $H^1(\widehat{X}; T_{\widehat{X}})$ has also geometric meaning (it is the tangent space to the “simultaneous resolution locus” for the resolution $\pi: \widehat{X} \rightarrow X$) and the map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^0(X; T_X^1)$ factors through the natural map $H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$. Our goal in this section is to explicitly compare the image of $H^1(\widehat{X}; T_{\widehat{X}})$ in $H^0(X; T_X^1)$ with that of $H^1(\widehat{X}; \Omega_{\widehat{X}}^2) \cong H^1(X'; T_{X'})$. More generally we compare $\mathbf{Def}_{\widehat{X}}$ and $\mathbf{Def}_{X'}$. While this example is somewhat special, similar techniques will handle other examples, such as the natural good resolution of the A_2 singularity defined by $x^2 + y^2 + z^2 + w^3$.

We begin with a general result:

Lemma 5.1. *Let $g: \widehat{X} \rightarrow X'$ be the blowup of a smooth threefold X' along a smooth compact curve C , with exceptional divisor E . Then $R^1 g_* T_{\widehat{X}} = R^1 g_* T_{\widehat{X}}(-E) = 0$, and there is an exact sequence*

$$0 \rightarrow R^0 g_* T_{\widehat{X}} \rightarrow T_{X'} \rightarrow N_{C/X'} \rightarrow 0.$$

Proof. For the first statement, we must show that $(R^1 g_* T_{\widehat{X}})_t = 0$ for all $t \in X'$, and we may as well assume that $t \in C$. By the formal functions theorem, it suffices to show that

$$\varprojlim_n H^1(nf; T_{\widehat{X}}|nf) = 0,$$

where f is a fiber of $g: \widehat{X} \rightarrow X'$ over a point $t \in C$, I_f is the ideal sheaf defining the reduced scheme f , and nf is the scheme defined by I_f^n . From the exact sequence

$$0 \rightarrow N_{f/E} \rightarrow N_{f/\widehat{X}} \rightarrow N_{E/\widehat{X}}|f \rightarrow 0,$$

and the fact that $N_{f/E} \cong \mathcal{O}_f$ and $N_{E/\widehat{X}}|f \cong \mathcal{O}_f(-1)$, we see that

$$N_{f/\widehat{X}} \cong \mathcal{O}_f \oplus \mathcal{O}_f(-1).$$

Because f is a local complete intersection, there is an exact sequence

$$0 \rightarrow \mathrm{Sym}^n(I_f/I_f^2) \rightarrow \mathcal{O}_{(n+1)f} \rightarrow \mathcal{O}_{nf} \rightarrow 0,$$

where I_f/I_f^2 is the conormal bundle. Hence

$$\mathrm{Sym}^n(I_f/I_f^2) = \mathcal{O}_f \oplus \mathcal{O}_f(1) \oplus \cdots \oplus \mathcal{O}_f(n).$$

Then from the normal bundle sequence

$$0 \rightarrow T_f \rightarrow T_{\widehat{X}}|_f \rightarrow N_{f/\widehat{X}} \rightarrow 0,$$

and the fact that $T_f \cong \mathcal{O}_f(2)$, it follows easily that $H^1(nf; T_{\widehat{X}}|_f) = 0$ for all $n \geq 1$, hence that $R^1g_*T_{\widehat{X}} = 0$. The proof that $R^1g_*T_{\widehat{X}}(-E) = 0$ is similar, using $\mathcal{O}_{\widehat{X}}(-E)|_f = \mathcal{O}_f(1)$.

To see the statement about $R^0g_*T_{\widehat{X}}$, there is an exact sequence

$$0 \rightarrow T_{\widehat{X}} \rightarrow g^*T_{X'} \rightarrow i_*T_{E/C}(E) \rightarrow 0,$$

where $i: E \rightarrow \widehat{X}$ is the inclusion. Hence there is an exact sequence

$$(*) \quad 0 \rightarrow g_*T_{\widehat{X}} \rightarrow g_*g^*T_{X'} \rightarrow g_*i_*T_{E/C}(E) \rightarrow R^1g_*T_{\widehat{X}} = 0.$$

Then $g_*g^*T_{X'} = T_{X'}$ and $g_*i_*T_{E/C}(E) = r_*T_{E/C}(E)$, where $r: E \rightarrow C$ is the projection. Also, we have the Euler exact sequence

$$0 \rightarrow \mathcal{O}_E \rightarrow r^*N_{C/X'}(1) \rightarrow T_{E/C} \rightarrow 0.$$

Thus, using $\mathcal{O}_E(E) = \mathcal{O}_E(-1)$, we get:

$$0 \rightarrow \mathcal{O}_E(E) \rightarrow r^*N_{C/X'} \rightarrow T_{E/C}(E) \rightarrow 0.$$

Taking r_* and using $R^1r_*\mathcal{O}_E(E) = 0$ gives $r_*T_{E/C}(E) = N_{C/X'}$. Thus $(*)$ becomes

$$0 \rightarrow g_*T_{\widehat{X}} \rightarrow T_{X'} \rightarrow N_{C/X'} \rightarrow 0,$$

as claimed. \square

Corollary 5.2. *Suppose as above that $g: \widehat{X} \rightarrow X'$ is the blowup of a smooth threefold along a smooth compact curve C , and that moreover $p: X' \rightarrow X$ is a resolution of an isolated singular point x , with $C \subseteq p^{-1}(x)$. Suppose in addition either that $p: X' \rightarrow X$ is a small resolution of X or that it is a good resolution with $\deg N_{C/E_i} < 0$ for every component E_i of $E = p^{-1}(x)$ containing C . Then there is an exact sequence*

$$0 \rightarrow H^0(C; N_{C/X'}) \rightarrow H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(X'; T_{X'}) \rightarrow H^1(C; N_{C/X'}).$$

Hence, if $p: X' \rightarrow X$ is an equivariant resolution of X , for example if p is a small resolution, then so is $\pi = p \circ g: \widehat{X} \rightarrow X$.

Proof. By the Leray spectral sequence and Lemma 5.1, $H^i(\widehat{X}; T_{\widehat{X}}) = H^i(X'; R^0g_*T_{\widehat{X}})$ for all i . The exact sequence of the statement then follows from the exact sequence for $R^0g_*T_{\widehat{X}}$ in Lemma 5.1, except for the injectivity on the left. If $H^0(C; N_{C/X'}) \rightarrow H^1(\widehat{X}; T_{\widehat{X}})$ is not injective, then the map $H^0(X'; T_{X'}) \rightarrow H^0(C; N_{C/X'})$ is nonzero. Thus there exists a nonzero element of $H^0(C; N_{C/X'})$ which lifts to $\theta \in H^0(X'; T_{X'})$. Exponentiating the vector field θ , we see that C moves in a one parameter family in X' . However, given the contraction $p: X' \rightarrow X$, every such family must be contained in the exceptional set of p . This is clearly impossible if p is a small resolution, or if C does not move in a family in some component E_i of the exceptional set E . In particular, if $\deg N_{C/E_i} < 0$ for every component E_i of E containing C , then C does not move inside any E_i .

To see the last statement, the above shows that the map $H^0(\widehat{X}; T_{\widehat{X}}) \rightarrow H^0(X'; T_{X'})$ is surjective. Equivalently, $R^0\pi_*T_{\widehat{X}} \rightarrow R^0p_*T_{X'}$ is surjective, and it is clearly injective, hence an isomorphism. Since by assumption $R^0p_*T_{X'} = T_X^0$, this says that $R^0\pi_*T_{\widehat{X}} = T_X^0$, i.e. π is equivariant. \square

For the rest of this section, (X, x) is a threefold A_{2n-1} singularity, $p: X' \rightarrow X$ is a small resolution with exceptional curve C , and $g: \widehat{X} \rightarrow X'$ is the blowup of the curve C , with exceptional divisor E . Then $\pi = p \circ g: \widehat{X} \rightarrow X$ is a non-crepant resolution of X , and it is equivariant by Corollary 5.2. If $n = 1$, then $H^i(\widehat{X}; T_{\widehat{X}}) = H^i(X'; T_{X'}) = 0$ for $i = 1, 2$, and both $\mathbf{Def}_{\widehat{X}}$ and $\mathbf{Def}_{X'}$ are (represented by) a single point. Thus, we shall always assume that $n \geq 2$, so that $N_{C/X'} \cong \mathcal{O}_C \oplus \mathcal{O}_C(-2)$ and $E \cong \mathbb{F}_2$. Note that $K_{\widehat{X}} = \mathcal{O}_{\widehat{X}}(E)$ and $K_E = K_{\widehat{X}} \otimes \mathcal{O}_{\widehat{X}}(E)|_E = \mathcal{O}_{\widehat{X}}(2E)|_E$. As $K_E = \mathcal{O}_E(-2\sigma - 4f)$, where σ is the negative section on E and f is the class of a fiber, $\mathcal{O}_{\widehat{X}}(E)|_E = \mathcal{O}_E(-\sigma - 2f)$. In particular, $N_{E/\widehat{X}}^\vee = \mathcal{O}_{\widehat{X}}(-E)|_E$ is effective, nef, and big, and $H^i(E; N_{E/\widehat{X}}) = 0$ for all i since

$$H^2(E; N_{E/\widehat{X}})^\vee = H^0(E; K_E \otimes \mathcal{O}_{\widehat{X}}(-E)|_E) = H^0(E; \mathcal{O}_E(-\sigma - 2f)) = 0.$$

Corollary 5.3. *Under the above assumptions,*

- (i) $H^2(\widehat{X}; T_{\widehat{X}}) = H^2(\widehat{X}; T_{\widehat{X}}(-E)) = 0$. In particular, $\mathbf{Def}_{\widehat{X}}$ is unobstructed of dimension $\dim H^1(\widehat{X}; T_{\widehat{X}})$.
- (ii) $H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \cong H^1(\widehat{X}; T_{\widehat{X}})$ and the natural map

$$H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^1(E; T_E)$$

is surjective. Hence E is a stable submanifold of \widehat{X} , deformations of \widehat{X} are versal for deformations of E , and there exist small deformations of \widehat{X} for which E deforms to \mathbb{F}_0 .

- (iii) For all i , $H^i(\widehat{X}; T_{\widehat{X}}) = H^i(X'; R^0 g_* T_{\widehat{X}})$, and there is an exact sequence

$$0 \rightarrow \mathbb{C} = H^0(C; N_{C/X'}) \rightarrow H^1(\widehat{X}; T_{\widehat{X}}) \rightarrow H^1(X'; T_{X'}) \rightarrow H^1(C; N_{C/X'}) \rightarrow 0.$$

Thus $\dim H^1(\widehat{X}; T_{\widehat{X}}) = \dim H^1(X'; T_{X'}) = n - 1$.

Proof. (i) To see that $H^2(\widehat{X}; T_{\widehat{X}}) = 0$, it suffices to show that $R^2 \pi_* T_{\widehat{X}} = 0$. In the Leray spectral sequence with $E_2^{a,b} = R^a p_* R^b g_* T_{\widehat{X}} \implies R^{a+b} \pi_* T_{\widehat{X}}$, all possible terms contributing to $R^2 \pi_* T_{\widehat{X}}$ are 0, either for dimension reasons or because $R^1 p_* R^1 g_* T_{\widehat{X}} = 0$ by Lemma 5.1. Thus $R^2 \pi_* T_{\widehat{X}} = 0$, and the proof for $H^2(\widehat{X}; T_{\widehat{X}}(-E))$ is similar.

- (ii) From the exact sequence

$$0 \rightarrow T_{\widehat{X}}(-\log E) \rightarrow T_{\widehat{X}} \rightarrow N_{E/\widehat{X}} \rightarrow 0,$$

and the fact that $H^i(E; N_{E/\widehat{X}}) = 0$ for all i (apply Leray to the morphism $r: E \rightarrow C$), we have $H^i(\widehat{X}; T_{\widehat{X}}(-\log E)) \cong H^i(\widehat{X}; T_{\widehat{X}})$ for all i . Thus in particular $H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$ and $H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \cong H^1(\widehat{X}; T_{\widehat{X}})$. Finally, from the exact sequence

$$0 \rightarrow T_{\widehat{X}}(-E) \rightarrow T_{\widehat{X}}(-\log E) \rightarrow T_E \rightarrow 0$$

and the vanishing of $H^2(\widehat{X}; T_{\widehat{X}}(-E))$, we see that $H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \rightarrow H^1(E; T_E)$ is surjective.

- (iii) This follows from Corollary 5.2 and the fact that $H^2(\widehat{X}; T_{\widehat{X}}) = 0$. \square

Remark 5.4. (i) By (iii) above, the images of $H^1(\widehat{X}; T_{\widehat{X}})$ and $H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$ in $H^0(X; T_X^1)$ are different, since by Proposition 4.13 the image of $H^1(\widehat{X}; \Omega_{\widehat{X}}^2)$ is that of $H^1(X'; \Omega_{X'}^2) = H^1(X'; T_{X'})$, and this image is strictly larger than that of $H^1(\widehat{X}; T_{\widehat{X}})$.

(ii) The functors $\mathbf{Def}_{\widehat{X}}$ and $\mathbf{Def}_{X'}$ are both smooth of dimension $n - 1$, but the differential of the corresponding morphism of functors $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_{X'}$, i.e. the induced map on Zariski tangent spaces, is not an isomorphism at 0: it has a one-dimensional kernel and cokernel. We will describe the morphism $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_{X'}$ explicitly.

First, let $\mathbf{Def}_{\widehat{X},E}$ denote the functor of deformations of the pair (X, E) : for a germ (S, s_0) , an element of $\mathbf{Def}_{\widehat{X},E}(S, s_0)$ consists of a deformation $\widehat{\mathcal{X}}$ of \widehat{X} over S , together with an effective Cartier divisor \mathcal{E} of $\widehat{\mathcal{X}}$, flat over S and restricting to E over s_0 . The functor $\mathbf{Def}_{X',C}$ is defined similarly; the objects over S are pairs $(\mathcal{X}', \mathcal{C})$ where \mathcal{C} is flat over S and restricts to the reduced subscheme C of X' . In particular, as we are only considering germs of spaces, \mathcal{C} is smooth over S with all fibers irreducible.

Proposition 5.5. $\mathbf{Def}_{\widehat{X}} \cong \mathbf{Def}_{\widehat{X},E} \cong \mathbf{Def}_{X',C}$, and the morphism $\mathbf{Def}_{\widehat{X}} \rightarrow \mathbf{Def}_{X'}$ is the same under the above identification as the forgetful morphism $\mathbf{Def}_{X',C} \rightarrow \mathbf{Def}_{X'}$.

Proof. It is a standard result that the tangent space to $\mathbf{Def}_{\widehat{X},E}$ is $H^1(\widehat{X}; T_{\widehat{X}}(-\log E))$ (cf. §1) and the obstruction space is $H^2(\widehat{X}; T_{\widehat{X}}(-\log E)) = 0$. Thus, $\mathbf{Def}_{\widehat{X},E}$ is smooth by Corollary 5.3(ii), and the first statement of the proposition is the isomorphism $H^1(\widehat{X}; T_{\widehat{X}}(-\log E)) \cong H^1(\widehat{X}; T_{\widehat{X}})$. For the second, given a pair $(\widehat{\mathcal{X}}, \mathcal{E})$ in $\mathbf{Def}_{\widehat{X},E}(S, s_0)$, it is easy to check that the morphism $r: E \rightarrow C$ extends to a morphism $\mathcal{E} \rightarrow \mathcal{C} \cong C \times S$ (note that $C \cong \mathbb{P}^1$ is rigid), and that \mathcal{E} can be blown down to a subspace $\mathcal{C} \cong C \times S \subseteq \mathcal{X}'$. Conversely, given a pair $(\mathcal{X}', \mathcal{C})$ over S , let $\widehat{\mathcal{X}}$ be the blowup of \mathcal{X}' along \mathcal{C} and let \mathcal{E} be the exceptional divisor. This gives two morphisms of functors $\mathbf{Def}_{\widehat{X},E} \rightarrow \mathbf{Def}_{X',C}$ and $\mathbf{Def}_{X',C} \rightarrow \mathbf{Def}_{\widehat{X},E}$ which are clearly inverses. Hence $\mathbf{Def}_{\widehat{X},E} \cong \mathbf{Def}_{X',C}$. \square

To put the above in more manageable form, we give an explicit description of $\mathbf{Def}_{X',C}$. First, we recall the basics about deformations of X and X' . Let Z be the germ of the standard ordinary double point in dimension 2, given by $x^2 + y^2 + z^2 = 0$, and let Z' be the resolution of singularities of Z , with $C \subseteq Z'$ the exceptional set. Then \mathbf{Def}_Z is represented by the germ $(\mathbb{C}, 0)$, with coordinate t and universal family $\mathcal{Z} \rightarrow (\mathbb{C}, 0)$ given by $x^2 + y^2 + z^2 + t = 0$. Likewise, $\mathbf{Def}_{Z'}$ is represented by the germ $(\mathbb{C}, 0)$, with coordinate u , and universal family $\mathcal{Z}' \rightarrow (\mathbb{C}, 0)$ given by a choice for a small resolution of $x^2 + y^2 + z^2 + u^2 = 0$. The threefold X' is isomorphic in a neighborhood of C to f^*Z' , where $f: (\mathbb{C}, 0) \rightarrow (\mathbb{C}, 0)$ is given by $u = f(w) = w^n$. A deformation \mathcal{X}' of X' over a germ (S, s_0) corresponds to a morphism $F: (\mathbb{C}, 0) \times S \rightarrow (\mathbb{C}, 0)$ with $F(w, s_0) = f(w) = w^n$, with $\mathcal{X}' = F^*Z'$. A polynomial F restricting to w^n is analytically equivalent to one of the form $w^n + b_{n-2}(s)w^{n-1} + \dots + b_0(s)$. In particular, $\mathbf{Def}_{X'}$ is represented by the germ of the affine space $(\mathbb{C}^{n-1}, 0)$, with coordinates (b_{n-2}, \dots, b_0) . Note also that $F^{-1}(0) \rightarrow S$ is a finite cover of degree n .

Lemma 5.6. *There is an isomorphism of functors from $\mathbf{Def}_{X',C}$ to \mathbf{F} , where, for (S, s_0) the germ of an analytic space, $\mathbf{F}(S)$ is the set of pairs (F, σ) , where as above $F: (\mathbb{C}, 0) \times S \rightarrow (\mathbb{C}, 0)$ is a morphism such that $F(w, s_0) = f(w) = w^n$, and $\sigma \subseteq F^{-1}(0)$ is a section of the finite cover $F^{-1}(0) \rightarrow S$, or equivalently a morphism $\lambda: S \rightarrow (\mathbb{C}, 0)$ such that $F(\lambda(s), s)$ is identically 0. Moreover, via this isomorphism, the morphism $\mathbf{Def}_{X',C} \rightarrow \mathbf{Def}_{X'}$ corresponds the forgetful map $(F, \sigma) \in \mathbf{F}(S) \mapsto F$.*

Proof. Given an object (F, σ) of $\mathbf{F}(S)$, the morphism F defines \mathcal{X}' in $\mathbf{Def}_{X'}(S)$ corresponding to the morphism $F: (\mathbb{C}, 0) \times S \rightarrow (\mathbb{C}, 0)$ as above, and a Cartesian diagram

$$\begin{array}{ccc} \mathcal{X}' & \longrightarrow & Z' \\ \downarrow & & \downarrow \\ (\mathbb{C}, 0) \times S & \xrightarrow{F} & (\mathbb{C}, 0). \end{array}$$

Note that $\mathcal{X}'|_{F^{-1}(0)} = Z' \times F^{-1}(0) \subseteq \mathcal{X}'$, and thus

$$C \times F^{-1}(0) \subseteq \mathcal{X}'|_{F^{-1}(0)} \subseteq \mathcal{X}',$$

compatibly with the projection to S . Given the section $\sigma \subseteq F^{-1}(0)$, define

$$\mathcal{C} = C \times \sigma \subseteq C \times F^{-1}(0) \subseteq \mathcal{X}'.$$

Thus the pair (F, σ) defines a deformation of (X', C) over S , and hence an element of $\mathbf{Def}_{X', C}$.

Conversely, suppose that we are given a pair $(\mathcal{X}', \mathcal{C}) \in \mathbf{Def}_{X', C}(S)$, and let $F: (\mathbb{C}, 0) \times S \rightarrow (\mathbb{C}, 0)$ be the morphism corresponding to \mathcal{X}' in $\mathbf{Def}_{X'}(S)$. Note that C does not deform in \mathcal{Z}' , even to first order. Thus $\mathcal{C} \cong C \times S \subseteq C \times F^{-1}(0)$, compatibly with the projection to S , so that the projection of \mathcal{C} onto the second factor $F^{-1}(0)$ defines a section σ of the morphism $F^{-1}(0) \rightarrow S$. Clearly, the two constructions $\mathbf{F}(S) \rightarrow \mathbf{Def}_{X', C}(S)$ and $\mathbf{Def}_{X', C}(S) \rightarrow \mathbf{F}(S)$ are mutual inverses and are functorial under pullback. This defines the isomorphism of functors, and the final statement is clear from the construction. \square

Explicitly, with $P(w; \mathbf{b}) = w^n + \sum_{i=0}^{n-2} b_i w^i$, where $\mathbf{b} = (b_{n-2}, \dots, b_0)$, the universal deformation \mathcal{U}' of X' is given as a small resolution of $x^2 + y^2 + z^2 + (P(w; \mathbf{b}))^2 = 0$. Consider

$$\{(\lambda, b_{n-2}, \dots, b_0) : P(\lambda; \mathbf{b}) = 0\} \subseteq (\mathbb{C} \times \mathbb{C}^{n-1}, 0).$$

Note that, if $w - \lambda$ is a factor of $P(w; \mathbf{b})$, then

$$P(w; \mathbf{b}) = (w - \lambda)(w^{n-1} + \lambda w^{n-2} + t_{n-3} w^{n-3} + \dots + t_1 w + t_0) = (w - \lambda)Q(w; \lambda, \mathbf{t}),$$

say, where $Q(w; \lambda, \mathbf{t}) = w^{n-1} + \lambda w^{n-2} + t_{n-3} w^{n-3} + \dots + t_1 w + t_0$. Solving explicitly for the coefficients b_i , we see:

$$\begin{aligned} b_{n-2} &= -\lambda^2 + t_{n-3}; \\ b_i &= -\lambda t_i + t_{i-1}, \quad 1 \leq i \leq n-3; \\ b_0 &= -\lambda t_0. \end{aligned}$$

If $\lambda, t_{n-3}, \dots, t_0$ are coordinates on \mathbb{C}^{n-1} , this defines a morphism $\Phi: \mathbb{C}^{n-1} \rightarrow \mathbb{C}^{n-1}$ by:

$$\Phi(\lambda, t_{n-3}, \dots, t_0) = (-\lambda^2 + t_{n-3}, -\lambda t_{n-3} + t_{n-4}, \dots, -\lambda t_0) = (b_{n-2}, \dots, b_0).$$

Solving for t_i in terms of λ and the b_i gives

$$\begin{aligned} t_i &= \lambda^{n-i-1} + b_{n-2} \lambda^{n-i-3} + \dots + b_{i+1}, \quad 1 \leq i \leq n-3; \\ -b_0 &= \lambda^n + b_{n-2} \lambda^{n-2} + \dots + \lambda b_1, \end{aligned}$$

recovering the fact that λ is a root of $P(w; \mathbf{b})$ (and there are exactly n such roots). Thus the morphism $\pi_1 \times \Phi: \mathbb{C}^{n-1} \rightarrow \mathbb{C} \times \mathbb{C}^{n-1}$ defined by

$$(\lambda, t_{n-3}, \dots, t_0) \mapsto (\lambda, \Phi(\lambda, t_{n-3}, \dots, t_0))$$

is an isomorphism from \mathbb{C}^{n-1} to $\{(\lambda, b_{n-2}, \dots, b_0) : P(\lambda; \mathbf{b}) = 0\}$. The germ $(\mathbb{C}^{n-1}, 0)$, with coordinates $\lambda, t_{n-3}, \dots, t_0$, together with the family which is a small resolution of $\Phi^* \mathcal{U}'$, represents $\mathbf{Def}_{X', C}$, and hence after blowing up represents $\mathbf{Def}_{\hat{X}}$. Moreover Φ corresponds to the forgetful morphism $\mathbf{Def}_{X', C} \rightarrow \mathbf{Def}_{X'}$.

By the above, Φ is finite of degree n and surjective, and is ramified exactly where $Q(\lambda; \lambda, \mathbf{t}) = 0$, i.e. where $P(w; \mathbf{b})$ has λ as a double root (or where the discriminant of $P(w; \mathbf{b})$ vanishes). In fact,

$$\det \begin{pmatrix} \frac{\partial b_{n-2}}{\partial \lambda} & \dots & \frac{\partial b_0}{\partial \lambda} \\ \frac{\partial b_{n-2}}{\partial t_{n-3}} & \dots & \frac{\partial b_0}{\partial t_{n-3}} \\ \vdots & \vdots & \vdots \\ \frac{\partial b_{n-2}}{\partial t_0} & \dots & \frac{\partial b_0}{\partial t_0} \end{pmatrix} = \pm Q(\lambda; \lambda, \mathbf{t}).$$

Summarizing the above discussion, then, we have:

Theorem 5.7. *Let $(S_{\widehat{X}}, 0)$ and $(S_{X'}, 0)$ be the germs prorepresenting the functors $\mathbf{Def}_{\widehat{X}}$ and $\mathbf{Def}_{X'}$ respectively. Then the induced morphism $S_{\widehat{X}} \rightarrow S_{X'}$ is finite of degree n and its differential at the origin has a one-dimensional kernel and cokernel. \square*

REFERENCES

- [Art74] M. Artin, *Algebraic construction of Brieskorn’s resolutions*, J. Algebra **29** (1974), 330–348.
- [Bri71] E. Brieskorn, *Singular elements of semi-simple algebraic groups*, Actes du Congrès International des Mathématiciens (Nice, 1970), Tome 2, Gauthier-Villars Éditeur, Paris, 1971, pp. 279–284.
- [Cle87] H. Clemens, *Curves on higher-dimensional complex projective manifolds*, Proceedings of the International Congress of Mathematicians, Vol. 1, 2 (Berkeley, Calif., 1986), Amer. Math. Soc., Providence, RI, 1987, pp. 634–640.
- [dFKX17] T. de Fernex, J. Kollár, and C. Xu, *The dual complex of singularities*, Higher dimensional algebraic geometry—in honour of Professor Yujiro Kawamata’s sixtieth birthday, Adv. Stud. Pure Math., vol. 74, Math. Soc. Japan, Tokyo, 2017, pp. 103–129.
- [DPT80] M. Demazure, H. Pinkham, and B. Teissier (eds.), *Séminaire sur les Singularités des Surfaces*, Lecture Notes in Mathematics, vol. 777, Springer, Berlin, 1980, Held at the Centre de Mathématiques de l’École Polytechnique, Palaiseau, 1976–1977.
- [EF21] P. Engel and R. Friedman, *Smoothings and rational double point adjacencies for cusp singularities*, J. Differential Geom. **118** (2021), no. 1, 23–100.
- [Eng18] P. Engel, *Looijenga’s conjecture via integral-affine geometry*, J. Differential Geom. **109** (2018), no. 3, 467–495.
- [FL22] R. Friedman and R. Laza, *Deformations of singular Fano and Calabi-Yau varieties*, arXiv:2203.04823, 2022.
- [FL23] ———, *Deformations of Calabi-Yau varieties with isolated log canonical singularities*, arXiv:2306.03755, 2023.
- [FL24] ———, *Deformations of Calabi-Yau varieties with k -liminal singularities*, Forum Math. Sigma **12** (2024), Paper No. e59.
- [FM83] R. Friedman and R. Miranda, *Smoothing cusp singularities of small length*, Math. Ann. **263** (1983), no. 2, 185–212.
- [Fri83] R. Friedman, *Global smoothings of varieties with normal crossings*, Ann. of Math. (2) **118** (1983), no. 1, 75–114.
- [Fri86] ———, *Simultaneous resolution of threefold double points*, Math. Ann. **274** (1986), no. 4, 671–689.
- [Fri15] ———, *On the geometry of anticanonical pairs*, arXiv:1502.02560, 2015.
- [Gro85] M. Gros, *Classes de Chern et classes de cycles en cohomologie de Hodge-Witt logarithmique*, Mém. Soc. Math. France (N.S.) (1985), no. 21, 87.
- [Gro97] M. Gross, *Deforming Calabi-Yau threefolds*, Math. Ann. **308** (1997), no. 2, 187–220.
- [Har67] R. Hartshorne, *Local cohomology*, Lecture Notes in Mathematics, vol. No. 41, Springer-Verlag, Berlin-New York, 1967, A seminar given by A. Grothendieck, Harvard University, Fall, 1961.
- [Har77] ———, *Algebraic geometry*, Graduate Texts in Mathematics, No. 52, Springer-Verlag, New York-Heidelberg, 1977.
- [Kar86] U. Karras, *Local cohomology along exceptional sets*, Math. Ann. **275** (1986), no. 4, 673–682.
- [Kaw88] Y. Kawamata, *Crepan blowing-up of 3-dimensional canonical singularities and its application to degenerations of surfaces*, Ann. of Math. (2) **127** (1988), no. 1, 93–163.
- [Kaw92] ———, *Unobstructed deformations. A remark on a paper of Z. Ran*, J. Algebraic Geom. **1** (1992), no. 2, 183–190.
- [KM98] J. Kollár and S. Mori, *Birational geometry of algebraic varieties*, Cambridge Tracts in Mathematics, vol. 134, Cambridge University Press, Cambridge, 1998.
- [Kol97] J. Kollár, *Singularities of pairs*, Algebraic geometry—Santa Cruz 1995, Proc. Sympos. Pure Math., vol. 62, Amer. Math. Soc., Providence, RI, 1997, pp. 221–287.
- [KSB88] J. Kollár and N. I. Shepherd-Barron, *Threefolds and deformations of surface singularities*, Invent. Math. **91** (1988), no. 2, 299–338.
- [Lau80] H. B. Laufer, *Versal deformations for two-dimensional pseudoconvex manifolds*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **7** (1980), no. 3, 511–521.
- [Loo84] E. J. N. Looijenga, *Isolated singular points on complete intersections*, London Mathematical Society Lecture Note Series, vol. 77, Cambridge University Press, Cambridge, 1984.
- [Nam94] Y. Namikawa, *On deformations of Calabi-Yau 3-folds with terminal singularities*, Topology **33** (1994), no. 3, 429–446.

- [Nam97] ———, *Deformation theory of Calabi-Yau threefolds and certain invariants of singularities*, J. Algebraic Geom. **6** (1997), no. 4, 753–776.
- [Nam02] ———, *Stratified local moduli of Calabi-Yau threefolds*, Topology **41** (2002), no. 6, 1219–1237.
- [NS95] Y. Namikawa and J. H. M. Steenbrink, *Global smoothing of Calabi-Yau threefolds*, Invent. Math. **122** (1995), no. 2, 403–419.
- [Pin83] H. C. Pinkham, *Factorization of birational maps in dimension 3*, Singularities, Part 2 (Arcata, Calif., 1981), Proc. Sympos. Pure Math., vol. 40, Amer. Math. Soc., Providence, RI, 1983, pp. 343–371. MR 713260
- [PS08] C. A. M. Peters and J. H. M. Steenbrink, *Mixed Hodge structures*, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, vol. 52, Springer-Verlag, Berlin, 2008.
- [Rei80] M. Reid, *Canonical 3-folds*, Journées de Géométrie Algébrique d’Angers, Juillet 1979/Algebraic Geometry, Angers, 1979, Sijthoff & Noordhoff, Alphen aan den Rijn—Germantown, Md., 1980, pp. 273–310.
- [SB83a] N. I. Shepherd-Barron, *Degenerations with numerically effective canonical divisor*, The birational geometry of degenerations (Cambridge, Mass., 1981), Progr. Math., vol. 29, Birkhäuser Boston, Boston, MA, 1983, pp. 33–84.
- [SB83b] ———, *Extending polarizations on families of K3 surfaces*, The birational geometry of degenerations (Cambridge, Mass., 1981), Progr. Math., vol. 29, Birkhäuser, Boston, Mass., 1983, pp. 135–171.
- [Sch71] M. Schlessinger, *Rigidity of quotient singularities*, Invent. Math. **14** (1971), 17–26.
- [Ste83] J. H. M. Steenbrink, *Mixed Hodge structures associated with isolated singularities*, Singularities, Part 2 (Arcata, Calif., 1981), Proc. Sympos. Pure Math., vol. 40, Amer. Math. Soc., Providence, RI, 1983, pp. 513–536.
- [Ste97] ———, *Du Bois invariants of isolated complete intersection singularities*, Ann. Inst. Fourier (Grenoble) **47** (1997), no. 5, 1367–1377.
- [Wah76] J. M. Wahl, *Equisingular deformations of normal surface singularities. I*, Ann. of Math. (2) **104** (1976), no. 2, 325–356.
- [Wan22] S.-S. Wang, *A remark on deformations of 1-convex manifolds with exceptional curves*, Bull. Sci. Math. **178** (2022), Paper No. 103144, 11.

COLUMBIA UNIVERSITY, DEPARTMENT OF MATHEMATICS, NEW YORK, NY 10027
Email address: rf@math.columbia.edu

STONY BROOK UNIVERSITY, DEPARTMENT OF MATHEMATICS, STONY BROOK, NY 11794
Email address: radu.laza@stonybrook.edu